

Knot Floer homology in cyclic branched covers

J ELISENDA GRIGSBY

In this paper, we introduce a sequence of invariants of a knot K in S^3 : the knot Floer homology groups $\widehat{HFK}(\Sigma^m(K); \widetilde{K}, i)$ of the preimage of K in the m -fold cyclic branched cover over K . We exhibit $\widehat{HFK}(\Sigma^m(K); \widetilde{K}, i)$ as the categorification of a well-defined multiple of the Turaev torsion of $\Sigma^m(K) - \widetilde{K}$ in the case where $\Sigma^m(K)$ is a rational homology sphere. In addition, when K is a two-bridge knot, we prove that $\widehat{HFK}(\Sigma^2(K); \widetilde{K}, \mathfrak{s}_0) \cong \widehat{HFK}(S^3; K)$ for \mathfrak{s}_0 the spin Spin^c structure on $\Sigma^2(K)$. We conclude with a calculation involving two knots with identical $\widehat{HFK}(S^3; K, i)$ for which $\widehat{HFK}(\Sigma^2(K); \widetilde{K}, i)$ differ as \mathbb{Z}_2 -graded groups.

57R58, 57M27; 57M05

1 Introduction

Let Y be a closed, connected, oriented 3-manifold and \mathfrak{s} a Spin^c structure on Y . In [13], Ozsváth and Szabó assign to the pair $(Y; \mathfrak{s})$ a graded abelian group, denoted $\widehat{HF}(Y; \mathfrak{s})$.

The additional data of an oriented, nullhomologous link L in Y induces a filtration on the chain complex used to compute $\widehat{HF}(Y; \mathfrak{s})$ for each Spin^c structure \mathfrak{s} [11], [16]. The filtered chain homotopy type of this complex is an invariant of the oriented link L in Y . One can, in particular, calculate the associated graded object of this filtration, yielding a sequence of graded abelian groups $\widehat{HFK}(Y; L, \mathfrak{s}, i)$, called the knot Floer homology groups of L in Y .

Now consider $\Sigma^m(K)$, the m -fold cyclic branched cover of S^3 branched along K . Let $p: \Sigma^m(K) \rightarrow S^3$ denote the associated projection map and $\widetilde{K} = p^{-1}(K)$ denote the preimage of K in $\Sigma^m(K)$. Consideration of \widetilde{K} in each cyclic branched cover, $\Sigma^m(K)$, yields a sequence of invariants of the original knot K in S^3 . Namely, for each $m \in \mathbb{Z}^+$ we have:

Definition 1.1 $\widehat{HFK}(\Sigma^m(K); \widetilde{K}) = \bigoplus_{\mathfrak{s}, i} \widehat{HFK}(\Sigma^m(K); \widetilde{K}, \mathfrak{s}, i)$, the knot Floer homology groups of $\widetilde{K} \subset \Sigma^m(K)$.

Our aim here is to study this sequence of invariants, focusing on the case where K is a two-bridge knot and $m = 2$. Then $\Sigma^2(K)$ is a lens space (Chapter 12 in [2]) with $H_1(\Sigma^2(K)) = \mathbb{Z}_n$, n an odd integer. Our main result, stated more precisely in Section 4, is:

Theorem 4.3 *For K a two-bridge knot in S^3 , there exists a Spin^c structure, \mathfrak{s}_0 , on $\Sigma^2(K)$ for which*

$$\widehat{HFK}(\Sigma^2(K); \widetilde{K}; \mathfrak{s}_0) \cong \widehat{HFK}(S^3; K).$$

This result falsely suggests that the groups $\widehat{HFK}(\Sigma^m(K); \widetilde{K})$ contain no more information than the groups $\widehat{HFK}(S^3; K)$. In fact, there are knot pairs K_1, K_2 in S^3 for which

$$\widehat{HFK}(S^3; K_1) \cong \widehat{HFK}(S^3; K_2)$$

but for which

$$\widehat{HFK}(\Sigma^2(K); \widetilde{K}_1) \not\cong \widehat{HFK}(\Sigma^2(K); \widetilde{K}_2)$$

as \mathbb{Z}_2 -graded groups. Such a pair (the two-bridge knots $K(15, 4)$ and $K(15, 7)$) is discussed in detail in Section 4.1.

We also show that for K a nullhomologous knot in a rational homology sphere Y , $\widehat{HFK}(Y; K)$ is a categorification of a multiple of the Turaev torsion of $Y - K$. The connection, established by Kirk and Livingston in [5], between the Casson–Gordon invariant of K and various torsions of $\Sigma^2(K)$ bears further examination, particularly since it may yield new obstructions to K being slice.

The paper is laid out as follows:

In Section 2 we recall the relevant definitions and theorems in Heegaard Floer homology as well as describe and develop notation for certain natural handlebody decompositions and Heegaard diagrams associated to $K \subset S^3$ and $\widetilde{K} \subset \Sigma^m(K)$.

In Section 3 we discuss torsions of chain complexes and prove that $\widehat{HFK}(Y; K)$ is the categorification of a multiple of the Turaev torsion of $Y - K$ in the case where Y is a rational homology sphere.

In Section 4 we study the invariant $\widehat{HFK}(\Sigma^2(K); \widetilde{K})$ for the case where K is a two-bridge knot in S^3 . We also compute $\widehat{HFK}(\Sigma^2(K); \widetilde{K})$ in a few Spin^c structures for the two-bridge knots $K(15, 7)$ and $K(15, 4)$, whose double branched covers are the lens spaces $-L(15, 7)$ and $-L(15, 4)$, respectively.

Acknowledgments I am grateful to Peter Ozsváth for recommending this problem and for numerous indispensable conversations, to Rob Kirby for his guidance and support and for a careful reading of the manuscript, and to the referee for many useful comments.

2 Background and conventions

We begin by reminding the reader of the Floer homology setup for $\widehat{HFK}(Y)$. For details, see Ozsváth and Szabó [11; 12; 13]. For the knot Floer homology refinements, see also Rasmussen [16].

2.1 Heegaard Floer homology background

Let K be a nullhomologous knot in a closed, oriented, connected 3-manifold Y . Although Ozsváth and Szabó's theory assigns homology groups more generally to nullhomologous links in Y , we will focus on knots in this paper.

In [11], Ozsváth and Szabó present the data of a knot K in Y by means of a *doubly-pointed Heegaard diagram* compatible with K . More specifically, they construct a handlebody decomposition of Y arising from a generic self-indexing Morse function

$$f: Y \rightarrow \mathbb{R}$$

with a single index 0 and 3 critical point and g index 1 and 2 critical points. This decomposition yields a Heegaard diagram for Y . The data of two points on the Heegaard surface, S , specifies the knot, K .

Definition 2.1 A doubly-pointed Heegaard diagram for a pair (Y, K) is a tuple $(S, \vec{\alpha}, \vec{\beta}, w, z)$ where

- $\vec{\alpha}$ is the g -tuple of co-attaching circles for the g 1-handles
- $\vec{\beta}$ is the g -tuple of attaching circles for the g 2-handles
- $w, z \in S - \vec{\alpha} - \vec{\beta}$
- K is the isotopy class of $-\gamma_w \cup \gamma_z$, where γ_w and γ_z are gradient flow lines from the index 3 to index 0 critical points associated to any generic metric on Y , intersecting S at z and w , respectively.

We gather the standard definitions and notation here for the reader's convenience:

- $\text{Sym}^g(S) = S^{\times g} / \Sigma_g$ is the g -fold symmetric product of the Heegaard surface, S .

- $\mathbb{T}_\alpha = \alpha_1 \times \dots \times \alpha_g$ (resp. \mathbb{T}_β) is the half-dimensional torus of co-attaching (resp. attaching) circles of the 1–handles (resp. 2–handles) inside $\text{Sym}^g(S)$.
- $V_z = \{z\} \times \text{Sym}^{g-1}(S)$ (resp. V_w) is the codimension \mathbb{C} 1 subvariety of $\text{Sym}^g(S)$ consisting of g –tuples where one point is constrained to lie at z (resp. w).
- $n_z = (-) \cap V_z$ (resp. $n_w = (-) \cap V_w$) is the algebraic intersection number of a class with V_z (resp. V_w) in $\text{Sym}^g(S)$.

This data gives rise to a \mathbb{Z} –filtered chain complex, \widehat{CF} whose

- generators are elements $\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$,
- differential is given by:

$$\widehat{\partial} \mathbf{x} = \sum_{\mathbf{y} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta} \sum_{\{\phi \in \pi_2(\mathbf{x}, \mathbf{y}) \mid \mu(\phi) = 1, n_w(\phi) = 0\}} \#(\widehat{\mathcal{M}}(\phi)) \mathbf{y}$$

- where $\widehat{\mathcal{M}}(\phi)$ is the moduli space of holomorphic maps of the standard unit disk into $\text{Sym}^g(S)$ with boundary conditions as in [13], in the homotopy class of ϕ , modded out by the standard \mathbb{R} action (for the sake of simplicity, count $\# \widehat{\mathcal{M}}(\phi)$ with \mathbb{Z}_2 coefficients),
- $\mu(\phi)$ is the expected dimension of the moduli space $\mathcal{M}(\phi)$ (before we mod out by the \mathbb{R} action), given by the Maslov index of ϕ ,
- and the relative \mathbb{Z} filtration on generators is given by

$$\mathcal{F}(\mathbf{x}) - \mathcal{F}(\mathbf{y}) = n_z(\phi) - n_w(\phi)$$

where $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$.

The relative \mathbb{Z} filtration is improved to an absolute \mathbb{Z} filtration by requiring that the Euler characteristic of the associated bi-graded complex is the symmetrized Alexander polynomial of K . We will have more to say on this point in Section 3. See also Section 2.3 of [11].

The homology groups of the associated graded object of this \mathbb{Z} –filtered complex are Ozsváth and Szabó’s knot Floer homology groups; ie, $\widehat{HFK}(Y; K, j)$ is the homology of the chain complex $\mathcal{F}_j / \mathcal{F}_{j-1}$.

For convenience, we introduce a couple more pieces of notation:

- $s(\mathbf{x}, \mathbf{y}) = n_z(\phi) - n_w(\phi)$ is the \mathbb{Z} –filtration difference between \mathbf{x} and \mathbf{y} in \widehat{CFK} (where $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$).
- $m(\mathbf{x}, \mathbf{y}) = \mu(\phi) - 2n_w(\phi)$ is the relative homological grading of \mathbf{x} and \mathbf{y} in \widehat{CFK} (where, again, $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$).

2.2 Natural handlebody decompositions

In what follows, whenever we refer to a handlebody decomposition for $S^3 - K$, we will always mean one with

- a single 0–handle h_0 ,
- g 1–handles $h_{\alpha_1}, \dots, h_{\alpha_g}$,
- $(g - 1)$ 2–handles $h_{\beta_1}, \dots, h_{\beta_{g-1}}$,
- no 3–handles.

We will also specify an oriented meridian, μ , for K (along which the final 2–handle, h_{β_g} , will be attached to build S^3) such that the attaching circle of h_{β_g} goes over one of the h_{α_i} (for definiteness, h_{α_g}) geometrically once and over all of the other 1–handles geometrically 0 times. μ generates $H_1(S^3 - K)$ and specifies an orientation.

We will use the notation $\text{hb}(S^3 - K; \mu)$ to denote such a handlebody decomposition.

Similarly, $\text{hb}(S^3; K; \mu)$ will denote the extension of $\text{hb}(S^3 - K; \mu)$ to a handlebody decomposition for S^3 . In particular,

$$\text{hb}(S^3; K; \mu) = \text{hb}(S^3 - K; \mu) \cup h_\mu \cup h_3.$$

Accordingly, we construct a doubly-pointed Heegaard diagram $\text{hd}(S^3; K; \mu)$ for the pair (S^3, K) by choosing an oriented arc δ on S meeting μ transversely in a single intersection point and having 0 geometric intersection with all other co-attaching (attaching) circles for the 1–handles (2–handles). Our two basepoints z and w are then the initial and final points, respectively, of δ . See Figure 1.

$\text{hd}(S^3; K; \mu)$ has the properties:

- $\vec{\alpha} = \alpha_1 \cup \dots \cup \alpha_g$ are the coattaching circles associated to the 1–handles, $h_{\alpha_1}, \dots, h_{\alpha_g}$,
- $\vec{\beta} = \beta_1 \cup \dots \cup \beta_{g-1} \cup (\beta_g = \mu)$ are the attaching circles associated to the 2–handles $h_{\beta_1}, \dots, h_{\beta_{g-1}}, h_{\beta_g = \mu}$,
- The orientation convention for K given in Definition 2.1 has the property that if $\lambda \subset S$ is a longitude which agrees with the orientation on K , then $\lambda \cap \mu = \delta \cap \mu$.

A natural handlebody decomposition and doubly-pointed Heegaard diagram can be constructed for the m –fold cyclic branched cover $\Sigma^m(K)$ as follows:




Figure 1: Example of a doubly-pointed Heegaard diagram

- (1) Begin with the natural \mathbb{Z}_m -equivariant handlebody decomposition of $\Sigma^m(K) - \tilde{K}$ which associates to each handle, h , in the handlebody decomposition of $S^3 - K$, m handles in the handlebody decomposition of $\Sigma^m(K) - \tilde{K}$ consisting of a chosen lift \tilde{h} of h and $m - 1$ translates $\tau_m(\tilde{h}), \tau_m^2(\tilde{h}), \dots, \tau_m^{m-1}(\tilde{h})$ of \tilde{h} . We now have a \mathbb{Z}_m -equivariant handlebody decomposition for $\Sigma^m(K) - \tilde{K}$ but too many 0-handles (the theory requires a handlebody decomposition for $\Sigma^m(K)$ with a single 0 and 3 handle).

To correct this, recall that μ is the core circle for a single one of the 1-handles, h_{α_g} . Use $m - 1$ of the lifts of h_{α_g} : $\tau_m(\tilde{h}_{\alpha_g}), \dots, \tau_m^{m-1}(\tilde{h}_{\alpha_g})$ to cancel the extra 0-handles $\tau_m(\tilde{h}_0), \dots, \tau_m^{m-1}(\tilde{h}_0)$. This new handlebody decomposition is still \mathbb{Z}_m -equivariant with respect to the projection map (only now the action on \tilde{h}_0 and \tilde{h}_{α_g} are trivial). We denote this handlebody decomposition by $\text{hb}(\Sigma^m(K) - \tilde{K}; \tilde{\mu})$.

- (2) We extend this to a \mathbb{Z}_m -equivariant handlebody decomposition of $\Sigma^m(K)$ by adding one more 2-handle attached along $\tilde{\mu}$, which will be our choice of meridian for the knot \tilde{K} in $\Sigma^m(K)$, and a 3-handle to fill in the rest of the solid torus neighborhood of \tilde{K} . We denote this handlebody decomposition by $\text{hb}(\Sigma^m(K); \tilde{K}; \tilde{\mu})$.
- (3) Associated to this $\text{hb}(\Sigma^m(K); \tilde{K}; \tilde{\mu})$ is the corresponding doubly-pointed Heegaard diagram $\text{hd}(\Sigma^m(K); \tilde{K}; \tilde{\mu})$ with basepoints w and z on either side of $\tilde{\mu}$.

An example should make everything concrete.

2.3 Example: $K =$ right handed trefoil

Consider the genus 2 Heegaard diagram for S^3 compatible with $K =$ the right handed trefoil pictured in Figure 2. This is what we have been calling $\text{hd}(S^3; K; \mu = \beta_2)$. Notice that if we remove the 3–handle and h_{β_2} we get a handlebody decomposition for $S^3 - K$.




Figure 2: Genus 2 Heegaard diagram for the RH Trefoil

A presentation for $\pi_1(S^3 - K)$ is generated by $h_{\alpha_1}, h_{\alpha_2}$ with the single relation given by the attaching map of the 2–handle h_{β_1} . This relation can be read off by traversing β_1 and keeping track of the intersections with α_1 and α_2 :

$$\pi_1(S^3 - K) = \langle h_{\alpha_1}, h_{\alpha_2} \mid \partial(h_{\beta_1}) = h_{\alpha_1} h_{\alpha_2} h_{\alpha_1}^{-1} h_{\alpha_2}^{-1} h_{\alpha_1}^{-1} \rangle$$

The associated Heegaard diagram for $\Sigma^2(K)$ that we have been calling $\text{hd}(\Sigma^2(K); \tilde{K}; \tilde{\beta}_2)$ has the following properties:

- $\tilde{\alpha}_1, \tau_2(\tilde{\alpha}_1), \tilde{\alpha}_2$ are the co-attaching circles for the lifts of the 1–handles (note that we have used $\tau_2(\tilde{\alpha}_2)$ to cancel the extra 0–handle)
- $\tilde{\beta}_1, \tau_2(\tilde{\beta}_1), \tilde{\beta}_2$ are the attaching circles for the lifts of the 2–handles (note that $\tilde{\beta}_2$ still intersects $\tilde{\alpha}_2$ geometrically once and intersects no other α curves)

One obtains a handlebody decomposition for $\Sigma^2(K) - \tilde{K}$ by removing the 3–handle and the 2–handle, \tilde{h}_{β_2} .

A presentation for $\pi_1(\Sigma^2(K) - \tilde{K})$ is therefore generated by $\tilde{h}_{\alpha_1}, \tilde{h}_{\tau_2(\alpha_1)}, \tilde{h}_{\alpha_2}$ with the following relations:

$$\begin{aligned}\partial(\tilde{h}_{\beta_1}) &= \tilde{h}_{\alpha_1} \cdot \tilde{h}_{\alpha_1} \cdot \tilde{h}_{\alpha_2}^{-1} \cdot \tau_2(\tilde{h}_{\alpha_1})^{-1} \cdot \tilde{h}_{\alpha_2}^{-1} \\ \partial(\tau_2(\tilde{h}_{\beta_1})) &= \tau_2(\tilde{h}_{\alpha_1}) \cdot \tilde{h}_{\alpha_2} \cdot \tau_2(\tilde{h}_{\alpha_1}) \cdot \tilde{h}_{\alpha_1}^{-1}\end{aligned}$$

3 $\widehat{HFK}(Y)$ and $\check{\tau}(Y - K)$

The aim of this section is to understand $\widehat{HFK}(Y; K)$ as the categorification of a well-defined multiple of a version of the Turaev torsion of $Y - K$ in the case where Y is a rational homology sphere. K , as before, is a nullhomologous knot in Y .

To understand $\widehat{HFK}(Y; K)$ as a categorification, we must first recall that the chain complex for $\widehat{CFK}(Y; K)$ splits as a sum of chain complexes, naturally indexed by elements of $\text{Spin}^c(Y_0(K))$, where $Y_0(K)$ denotes the canonical 0 surgery on K . The set, $\text{Spin}^c(Y_0(K))$, is often referred to as the set of relative Spin^c structures of the pair (Y, K) and denoted $\underline{\text{Spin}}^c(Y, K)$. In Section 2.3 of [11] (see also Section 2.6 of [13]), Ozsváth and Szabó describe, given a doubly-pointed Heegaard diagram, how to construct a map

$$\mathbb{T}_\alpha \cap \mathbb{T}_\beta \rightarrow \underline{\text{Spin}}^c(Y, K)$$

and a splitting

$$\underline{\text{Spin}}^c(Y, K) \cong \text{Spin}^c(Y) \times \mathbb{Z}.$$

Taking the Euler characteristic of each summand in this splitting will yield a formal polynomial in two variables, one of which indexes the $\text{Spin}^c(Y)$ structure and one of which indexes the \mathbb{Z} factor.

In brief, we obtain such a formal two-variable polynomial as a multiple of the Reidemeister torsion of the maximal abelian cover of $Y - K$. Specifically, computation of the Reidemeister torsion yields a rational expression in $\mathbb{Q}(H_1(Y))(T)$ which depends on the choice of a lift of a \mathbb{Z} -basis for $C_*(Y - K)$ to a $\mathbb{Z}[H_1(Y)][T, T^{-1}]$ -basis for $C_*(Y - K)^{[H_1(Y-K)]}$, where $(Y - K)^{[H_1(Y-K)]}$ is the maximal abelian cover of $Y - K$. A multiple of this rational expression yields a well-defined formal two-variable polynomial in $\mathbb{Z}[\text{Spin}^c(Y)][T, T^{-1}]$ once we use Turaev's correspondence between Spin^c structures on 3-manifolds and lifts of \mathbb{Z} -module bases to $\mathbb{Z}[H_1(Y)]$ bases of the maximal abelian cover of Y .

Before launching into a formal discussion of these ideas, we state our main result.

Theorem 3.1 *Let Y be an oriented rational homology sphere and K an oriented, nullhomologous knot in Y .*

Let $e: \pi_1(Y - K) \rightarrow \mathbb{Z}$ be given by $e(\gamma) = lk(\gamma, K)$. Let $r: \pi_1(Y - K) \rightarrow H_1(Y)$ be the projection onto $H_1(Y)$.

Let $\check{\tau}(Y - K) \in \mathbb{Q}(\text{Spin}^c(Y)(T))$ be the variant of the Reidemeister torsion of the knot complement given in Definition 3.3. Then

$$(\check{\tau}(Y - K)) \cdot (T - 1) = \sum_{\mathfrak{s} \in \text{Spin}^c(Y)} p_{\mathfrak{s}}(T) \cdot \mathfrak{s}$$

where

$$p_{\mathfrak{s}}(T) = \sum_i \chi(\widehat{HFK}(Y; K, \mathfrak{s}, i)) \cdot T^i.$$

Recall that

$$\chi(\widehat{HFK}(Y; K, \mathfrak{s}, i)) = \sum_{\{d \in d_0 + \mathbb{Z}\}} (-1)^{d-d_0} rk(\widehat{HFK}_d(Y; K, \mathfrak{s}, i)).$$

In the above, d is the absolute homological grading of a generator, defined in [14]. We can, however, define $\chi(\widehat{HFK}(Y; K, \mathfrak{s}, i))$ without reference to this absolute grading for a rational homology sphere by using the relative \mathbb{Z}_2 homological grading on generators induced by comparing the local intersection numbers of \mathbb{T}_α and \mathbb{T}_β at two generators \mathbf{x} and \mathbf{y} (see Theorem 3.7).

We then lift this relative \mathbb{Z}_2 grading to an absolute \mathbb{Z}_2 grading by making the choice which insures

$$\sum_j (-1)^j rk(\widehat{HF}_j(Y)) = |H_1(Y; \mathbb{Z})|$$

for $j \in \mathbb{Z}_2$.

One should think of $\check{\tau}(Y - K)$ as a rational function in the formal variables $\{\mathfrak{s} | \mathfrak{s} \in \text{Spin}^c(Y)\}$ and T . In the numerator of a particular form of this rational expression, the \mathfrak{s} term records the Spin^c structure and the exponent on the T variable records the filtration level of a generator in $\widehat{HFK}(Y)$.

3.1 Background on torsions of chain complexes

We start by recalling some definitions. The classical references for this material are the papers of Milnor [7], [8]. The particular version of interest to us is developed in Turaev [19].

First, recall that the torsion can be defined for a finite acyclic chain complex

$$0 \longrightarrow C_m \xrightarrow{\partial_m} \cdots \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \xrightarrow{\partial_0} 0$$

of vector spaces over a field \mathbb{F} with fixed \mathbb{F} -bases $\{c_q\}$ for each C_q . One then chooses, for each q , a collection $\{b_q\}$ of elements whose images form a basis for $im(\partial_q)$ in C_{q-1} .

Given two bases $\{b\}$ and $\{c\}$ for a given vector space, let $[b/c]$ denote the determinant of the change of basis matrix (ie, the nonsingular matrix $A = (a_{ij})$ where $b_i = \sum_{j=1}^n a_{ij}c_j$).

Then

Definition 3.2 The torsion of C_* is defined as

$$\tau(C_*) = \prod_{q=0}^m [\{\partial_{q+1}(b_{q+1}), b_q\} / \{c_q\}]^{(-1)^{q+1}}.$$

It is well-known that $\tau(C_*)$ depends only upon the original choices of bases $\{c_q\}$ for C_q .

Classically, we have been interested in torsions of chain complexes arising as covers. For example, the Alexander polynomial of a knot is (a multiple of) the torsion of the infinite cyclic cover of the knot complement.

In general, we start with a finite chain complex of \mathbb{Z} -modules and construct a cover of X via a surjective homomorphism $\pi_1(X) \rightarrow G$. Call such a cover $X^{[G]}$. $C_*(X^{[G]})$ is a free $\mathbb{Z}[G]$ -module with a basis given by a choice of lift of the \mathbb{Z} -module basis downstairs. If G is an abelian group, we can construct the field of fractions $\mathbb{Q}(G)$ of $\mathbb{Z}[G]$ by inverting all non-zero-divisors. The free $\mathbb{Z}[G]$ -module basis we chose for $C_*(X^{[G]})$ then becomes a free $\mathbb{Q}(G)$ -basis for $C_*(X^{\mathbb{Q}(G)}) := \mathbb{Q}(G) \otimes_{\mathbb{Z}[G]} C_*(X^{[G]})$. If $C_*(X^{[G]})$ is an acyclic complex, we can compute its torsion.

3.2 Torsion of $(Y - K)_{\mathbb{Q}(G \times \mathbb{Z})}$

Let K be a nullhomologous, oriented knot in an oriented rational homology sphere, Y , and let $\pi = \pi_1(Y - K)$. We are interested in the torsion of the chain complex arising from the surjective Hurewicz homomorphism

$$\pi \rightarrow H_1(Y - K).$$

Notice that a choice of oriented meridian μ for the knot yields a splitting

$$\varphi_\mu = r \times e: H_1(Y - K) \xrightarrow{\cong} H_1(Y) \times \mathbb{Z} \cong G \times \mathbb{Z}$$

specified by

$$\varphi_\mu(h) = (h - e(h)\mu, e(h)).^1$$

Here,

$$e: \pi \rightarrow \mathbb{Z}$$

is given by linking number with K :

$$e(h) = lk(h, K)$$

and

$$r: \pi \rightarrow H_1(Y) \cong G$$

is the composition $\pi_1(Y - K) \rightarrow \pi_1(Y)$ with $\pi_1(Y) \rightarrow H_1(Y)$.

Let $\epsilon: \mathbb{Z}[\pi] \rightarrow \mathbb{Z}[\mathbb{Z}]$ denote the \mathbb{Z} -ring extension of e and let $\rho: \mathbb{Z}[\pi] \rightarrow \mathbb{Z}[G]$ denote the \mathbb{Z} -ring extension of r .

Then let $C_*(\widetilde{Y - K})$ denote the \mathbb{Z} -linear chain complex of the universal cover, $\widetilde{Y - K}$, of $Y - K$ and form the free $\mathbb{Z}[G \times \mathbb{Z}]$ -module $\mathbb{Z}[G \times \mathbb{Z}] \otimes_{\rho \otimes \epsilon} C_*(\widetilde{Y - K})$ and denote it by

$$C_*((Y - K)^{[G \times \mathbb{Z}]})$$

At this point, we can form $\mathbb{Q}(G \times \mathbb{Z})$, the ring of quotients of $\mathbb{Z}[G \times \mathbb{Z}]$ and construct

$$C_{\mathbb{Q}(G \times \mathbb{Z})} := \mathbb{Q}(G \times \mathbb{Z}) \otimes_{\mathbb{Z}[G \times \mathbb{Z}]} C_*((Y - K)^{[G \times \mathbb{Z}]})$$

Now, to compute $\tau(C_{\mathbb{Q}(G \times \mathbb{Z})})$, we fix a handlebody decomposition for $Y - K$ with

- a single 0-handle h_0 ,
- g 1-handles $h_{\alpha_1}, \dots, h_{\alpha_g}$, where, again, we are choosing h_{α_g} to be μ , a meridian, (hence, $e(h_{\alpha_g}) = 1$ and $r(h_{\alpha_g}) = 0$),
- $(g - 1)$ 2-handles $h_{\beta_1}, \dots, h_{\beta_{g-1}}$,
- no 3-handles.

which yields a \mathbb{Z} -module basis for $C_*(Y - K)$.

The boundary maps

$$\partial_*: C_*((Y - K)^{[G \times \mathbb{Z}]}) \rightarrow C_{*-1}((Y - K)^{[G \times \mathbb{Z}]})$$

¹Note that I am abusing notation here (and will throughout), denoting an element of H_1 or π_1 by a loop representing it, and vice versa.

are most easily expressed using Fox calculus (see [4]). Specifically ∂_2 is the $(g-1) \times g$ matrix

$$(\partial_2)_{ij} = (\rho \otimes \epsilon) \left(\frac{\partial h_{\beta_i}}{\partial h_{\alpha_j}} \right) \in \mathbb{Z}[G \times \mathbb{Z}].$$

and ∂_1 is the $g \times 1$ matrix

$$(\partial_1)_i = (\rho \otimes \epsilon)(h_{\alpha_i} - 1) \in \mathbb{Z}[G \times \mathbb{Z}].$$

After verifying that the chain complex $C_{\mathbb{Q}(G \times \mathbb{Z})}$ is acyclic (addressed by Lemma 3.4, whose proof we give in Section 3.4), we can pick lifts $\tilde{h}_0, \tilde{h}_{\alpha_i}, \tilde{h}_{\beta_j}$ of the the \mathbb{Z} -module bases downstairs and compute the torsion by comparing that lift with

- $b_2 = \{\tilde{h}_{\beta_1}, \dots, \tilde{h}_{\beta_{g-1}}\}$
- $\partial_2(b_2) = \{\partial(\tilde{h}_{\beta_1}), \dots, \partial(\tilde{h}_{\beta_{g-1}})\}$
- $b_1 = \{\tilde{h}_{\alpha_g}\}$
- $\partial(b_1) = \{\partial(\tilde{h}_{\alpha_g})\}$
- $b_0 = \{\}$

Then,

- $[b_2/c_2] = 1$
- $[\partial_2(b_2)b_1/c_1] = \rho \otimes \epsilon([\partial_2^g])$, where ∂_2^g is the $g \times g$ matrix obtained by inserting the tuple representing the g th basis element of the chosen basis $\{h_{\alpha_1}, \dots, h_{\alpha_g}\}$ into the g th row of ∂_2 .
- $[b_1/c_0] = \rho \otimes \epsilon([\partial_1^g])$, where ∂_1^g is the 1×1 matrix obtained by deleting the first $g-1$ rows from ∂_1 . In other words, $[b_1/c_0] = \rho \otimes \epsilon(h_{\alpha_g} - 1) = T^1 - T^0$.

We obtain the torsion of $C_{\mathbb{Q}(G \times \mathbb{Z})}$ by computing the rational expression

$$(\rho \otimes \epsilon) \left(\frac{[\partial_2^g]}{[\partial_1^g]} \right) = \frac{\rho \otimes \epsilon([\partial_2^g])}{T^1 - T^0}$$

with respect to the initial choices $\tilde{h}_0, \tilde{h}_{\alpha_i}, \tilde{h}_{\beta_j}$ of lifts.

3.3 Euler chains and bases for chain complexes

All we lack in the above is a nice way of specifying a $\mathbb{Q}(G \times \mathbb{Z})$ -basis for the chain complex $C_{\mathbb{Q}(G \times \mathbb{Z})}$, ie, a lift of a particular \mathbb{Z} -module basis for $C_*(Y - K)$. Without such a lift, τ has an indeterminacy coming from this choice.

It turns out that Turaev gives us exactly the tools we need to specify such a lift. In brief, he explains how to associate to an *Euler chain* (defined in Section 2.5 in [19]) on $Y - K$ a $\mathbb{Z}[H_1(Y - K)]$ -module basis for the maximal abelian cover, $(Y - K)^{[H_1(Y - K)]}$.

Furthermore, given a doubly-pointed Heegaard diagram for Y compatible with a knot K , Ozsváth and Szabó construct a map

$$\mathbb{T}_\alpha \cap \mathbb{T}_\beta \rightarrow \text{Eul}(Y_0(K)) \rightarrow \text{Eul}(Y - K).$$

There is a natural splitting of $\text{Eul}(Y - K) \cong \text{Eul}(Y) \times \mathbb{Z}$ which, via Turaev’s identification of Euler chains and Spin^c structures, allows us to realize the torsion of the chain complex of the maximal abelian cover as a formal element of $\mathbb{Q}(\text{Spin}^c(Y))(T)$.

So, choosing a lift of a \mathbb{Z} -module basis for $C_*(Y - K)$ to a $\mathbb{Z}[H_1(Y - K)]$ -module basis for $C_*(Y - K)^{[H_1(Y - K)]}$ is just a matter of specifying an element of $\text{Eul}(Y - K)$.

We specify such an element by using Ozsváth and Szabó’s identification [11]

$$\mathbb{T}_\alpha \cap \mathbb{T}_\beta \rightarrow \underline{\text{Spin}}^c(Y, K) \leftrightarrow \text{Eul}(Y_0(K)).$$

Their map comes complete with a natural map

$$\text{Eul}(Y_0(K)) \rightarrow \text{Eul}(Y - K)$$

induced by forgetting the final two and three handle (and corresponding arcs in the spider-like Euler chain) along with a splitting

$$p_1 \times p_2: \text{Eul}(Y - K) \rightarrow \text{Eul}(Y) \times \mathbb{Z}.$$

Here, the map $p_1: \text{Eul}(Y - K) \rightarrow \text{Eul}(Y)$ is obtained via the unique extension of an Euler chain for $Y - K$ to one for Y .²

The map $p_2: \text{Eul}(Y - K) \rightarrow \mathbb{Z}$ is defined as follows. Let $\xi \in \text{Eul}(Y - K)$ be the restriction of $\xi' \in \text{Eul}(Y_0(K))$ and $\mathfrak{s}_{\xi'}$ be the element of $\text{Spin}^c(Y_0(K))$ associated to ξ' via Turaev’s identification. Then

$$p_2(\xi) = \frac{1}{2} \langle c_1(\mathfrak{s}_{\xi'}), [\widehat{F}] \rangle,$$

²Recall that in going from $Y - K$ to Y we add a 2-handle along a meridian μ for the knot and a 3-handle, and μ has a unique intersection point with a single α curve. The arc connecting h_0 to h_3 is uniquely specified by the basepoint, w .

where $[\widehat{F}]$ is the homology class of a capped-off Seifert surface for K in $Y_0(K)$.

Notice that this splitting is defined so that it respects the natural μ -induced splitting

$$\varphi_\mu: H_1(Y - K) \xrightarrow{\cong} (H_1(Y) \times \mathbb{Z})$$

specified by $\varphi_\mu(h) = (h - e(h)\mu, e(h))$, where $e: H_1(Y - K) \rightarrow \mathbb{Z}$ is defined by

$$e(h) = lk(h, K) = \langle h', [\widehat{F}] \rangle,$$

where $h' \in H_1(Y_0(K))$ is the induced image under the inclusion map

$$H_1(Y - K) \rightarrow H_1(Y_0(K))$$

(again, we are assuming that K is an oriented knot).

We are finally ready to define the variant of Reidemeister torsion for which $\widehat{HFK}(Y, K)$ is a categorification in the case when Y is a rational homology sphere.

Definition 3.3 Let Y be a rational homology sphere, K a nullhomologous, oriented knot in Y , μ a choice of meridian for K .

Let $\text{hd}(Y; K, \mu)$ be a doubly-pointed Heegaard diagram compatible with K .

Then consider the summands of the formal determinant of the matrix

$$\begin{array}{c} \alpha_1 \\ \vdots \\ \alpha_{g-1} \end{array} \begin{pmatrix} \beta_1 & \dots & \beta_{g-1} \\ & & \\ & & \end{pmatrix}$$

where the entries of the matrix are formal sums of intersection points between the appropriate α and β curves, and each intersection point in the matrix is assigned a ± 1 according to its local intersection number (see Lemma 3.6).

By acting on this formal sum by Ozsváth and Szabó's map

$$f: \mathbb{T}_\alpha \cap \mathbb{T}_\beta \rightarrow \underline{\text{Spin}}^c(Y, K)$$

composed with the splitting

$$p_1 \times p_2: \underline{\text{Spin}}^c(Y, K) \rightarrow \text{Spin}^c(Y) \times \mathbb{Z}$$

we get a formal element of $\mathbb{Z}[\text{Spin}^c(Y)][T, T^{-1}]$; ie, a formal polynomial, $p_s(T)$, in the variables $\{\mathfrak{s} | \mathfrak{s} \in \text{Spin}^c(Y)\}$ and T, T^{-1} .

We define:

$$\check{r}(Y - K) = \frac{p_s(T)}{T - 1}.$$

Remark Note that $\check{\tau}(Y - K)$ is actually the Reidemeister torsion of the maximal abelian cover of $Y - K$, where the usual indeterminacy coming from a choice of basis has been eliminated.

More precisely, let $e: \pi_1(Y - K) \rightarrow \mathbb{Z}$ be given by

$$e(\gamma) = lk(\gamma, K),$$

$$r: \pi_1(Y - K) \rightarrow H_1(Y)$$

be the projection onto $H_1(Y)$, and

$$\epsilon: \mathbb{Z}[\pi_1(Y - K)] \rightarrow \mathbb{Z}[T, T^{-1}],$$

$$\rho: \mathbb{Z}[\pi_1(Y - K)] \rightarrow \mathbb{Z}[H_1(Y)]$$

the natural group ring extensions.

Then

$$(\rho \otimes \epsilon) \begin{pmatrix} [\partial_2^g] \\ [\partial_1^g] \end{pmatrix} = \frac{\rho \otimes \epsilon([\partial_2^g])}{T - 1}$$

yields the Reidemeister torsion associated to the maximal abelian cover of $Y - K$. If the basis of the chain complex for the maximal abelian cover is specified by Ozsváth–Szabó’s map to Spin^c structures, then we arrive at $\check{\tau}(Y - K)$.

$\check{\tau}(Y - K)$ is the variant of Reidemeister torsion which will (when multiplied by $(T - 1)$) be the categorification of the knot Floer homology for a rational homology sphere.

3.4 $C_{\mathbb{Q}(G \times \mathbb{Z})}$ is an acyclic complex

We return now to the proof of an important point which we left unresolved in an earlier part of this section.

Lemma 3.4 *The chain complex $C_{\mathbb{Q}(G \times \mathbb{Z})}$ is acyclic.*

Proof of Lemma 3.4 We need only show that the determinants of the matrices $(\rho \otimes \epsilon)(\partial_2^g)$ and $(\rho \otimes \epsilon)(\partial_1^g)$ are units in $\mathbb{Q}(G)(T)$.

$(\rho \otimes \epsilon)[\partial_1^g] = T - 1$ is clearly a nonzerodivisor in $\mathbb{Z}[G][[T, T^{-1}]]$ and hence a unit in $\mathbb{Q}(G)(T)$.

To see that $[\partial_2^g]$ must be a unit in $\mathbb{Q}(G)(T)$, consider the ring homomorphism

$$\varphi: \mathbb{Q}(G)(T) \rightarrow \mathbb{Q}$$

which sends T and all $h \in G$ to 1. If $\varphi(p) = q$ is a unit in \mathbb{Q} , then p must be a unit in $\mathbb{Q}(G)(T)$.

But $\varphi((\rho \otimes \epsilon)(\partial_2^g))$ is exactly the relation matrix for $H_1(Y; \mathbb{Q})$. $H_1(Y; \mathbb{Q}) = 0$ then implies that $\varphi((\rho \otimes \epsilon)[\partial_2^g])$ is a unit in \mathbb{Q} . \square

3.5 Proof of Theorem 3.1

The relationship between generators of the knot Floer homology and summands of the determinant of the differential

$$\partial_2: (C_{\mathbb{Q}(G \times \mathbb{Z})})_2 \rightarrow (C_{\mathbb{Q}(G \times \mathbb{Z})})_1$$

used to compute the Reidemeister torsion is clear, since the generators of \widehat{CFK} are themselves summands of an analogous formal determinant (see, eg, the discussion in the proof of Proposition 4.4).

Under this correspondence, the relative element of $\text{Spin}^c(Y, K)$ specified by the difference between a pair of generators is an element of $\underline{H}_1(Y_0(K)) \cong H_1(Y - K)$.

Furthermore, by the naturality of the H_1 action on all of these sets, the splitting

$$H_1(Y - K) \rightarrow H_1(Y) \times \mathbb{Z}$$

matches the splitting

$$\underline{\text{Spin}}^c(Y, K) \cong \text{Spin}^c(Y) \times \mathbb{Z}.$$

We need only verify that $s(\mathbf{x}, \mathbf{y})$, the relative filtration grading of two generators, \mathbf{x} and \mathbf{y} , defined by $n_z(\phi) - n_w(\phi)$ (for $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$), matches up with the relative T exponent of the corresponding summands and that $m(\mathbf{x}, \mathbf{y})$, the relative Maslov grading of the generators agrees mod 2 with the relative sign of the summands in the determinant.

Lemma 3.5 *Suppose K is an oriented, nullhomologous knot in a closed, connected, oriented 3-manifold Y and $\text{hd}(Y; K, \mu)$ is a doubly-pointed Heegaard diagram as in Section 2. Then consider the map $e: \pi \rightarrow \mathbb{Z}$ given by $e(\gamma) = lk(\gamma, K)$. Let \mathbf{x}, \mathbf{y} be Floer homology generators and let $\gamma_{\mathbf{x}}, \gamma_{\mathbf{y}} \in \pi$ be their corresponding summands in the Fox determinant, $[\partial_2^g]$. Then*

$$e(\gamma_{\mathbf{x}}) - e(\gamma_{\mathbf{y}}) = s(\mathbf{x}, \mathbf{y}).$$

Proof of Lemma 3.5 First, recall that the filtration difference between two generators \mathbf{x} and \mathbf{y} is well-defined whenever \mathbf{x} and \mathbf{y} are in the same Spin^c structure. Then there exists a topological disk $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$ and $s(\mathbf{x}, \mathbf{y}) = n_z(\phi) - n_w(\phi)$.

Recall also that in $\text{hd}(Y; K)$ we have a distinguished β circle, μ , which is a meridian for K , and next to which we place the two basepoints as shown in Figure 1. Recall (Section 2.13 of [13]) that we can represent disks in $\pi_2(\mathbf{x}, \mathbf{y})$ uniquely as \mathbb{Z} -linear combinations of *fundamental domains* \mathcal{D}_i , which are the closures of the connected components of $S - \alpha_1 - \dots - \alpha_g - \beta_1 - \dots - (\beta_g = \mu)$.

Let \mathcal{D}_z and \mathcal{D}_w be the fundamental domains containing the basepoints z and w , respectively. Given a representation of a disk ϕ as a \mathbb{Z} -linear combination of fundamental domains, n_z and n_w are then the coefficients on \mathcal{D}_z and \mathcal{D}_w , respectively.

Now suppose \mathbf{x} and \mathbf{y} are two generators in the same Spin^c structure, $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$ is a disk connecting them, and $\sum_i \mathcal{D}_i$ is the linear combination of fundamental domains representing ϕ . Note that

$$\partial(\mathcal{D}_w) = -\mu + (\text{other stuff}),$$

and

$$\partial(\mathcal{D}_z) = \mu + (\text{other stuff}).$$

Therefore,

$$\begin{aligned} \partial(\phi) &= n_z(\partial\mathcal{D}_z) + n_w(\partial\mathcal{D}_w) + (\text{other stuff}) \\ &= (n_z - n_w)\mu + (\text{other stuff}) \end{aligned}$$

Call this (other stuff) γ . Note that γ is exactly the image under the Hurewicz map $\pi_1(Y - K) \rightarrow H_1(Y - K)$ of $\gamma_{\mathbf{x}}^{-1} \cdot \gamma_{\mathbf{y}}$, where $\gamma_{\mathbf{x}}$ and $\gamma_{\mathbf{y}}$ are the summands corresponding to \mathbf{x} and \mathbf{y} in the Fox determinant, $[\partial_2^g]$. In other words,

$$e(\gamma) = e(\gamma_{\mathbf{x}}^{-1} \cdot \gamma_{\mathbf{y}}) = -(e(\gamma_{\mathbf{x}}) - e(\gamma_{\mathbf{y}})).$$

Furthermore,

$$(n_z - n_w)\mu = -\gamma$$

in $H_1(Y - K)$. Therefore,

$$\begin{aligned} e(\gamma_{\mathbf{x}}) - e(\gamma_{\mathbf{y}}) &= -e(\gamma) \\ &= (n_z(\phi) - n_w(\phi)) \cdot e(\mu) \\ &= n_z(\phi) - n_w(\phi) \\ &= s(\mathbf{x}, \mathbf{y}), \end{aligned}$$

as desired. □

Lemma 3.6 *Let \mathbf{x} and \mathbf{y} be two elements of $\widehat{CFK}(Y)$ in the same $Spin^c$ structure. Let $\gamma_{\mathbf{x}}, \gamma_{\mathbf{y}} \in \mathbb{Z}[\pi]$ be the corresponding summands in the Fox determinant, $[\partial_2^g]$. Then*

$$(-1)^{m(\mathbf{x}, \mathbf{y})} = \text{sgn}(\gamma_{\mathbf{x}}) \cdot \text{sgn}(\gamma_{\mathbf{y}}).$$

Proof of Lemma 3.6 Recall the following standard fact from Lagrangian Intersection Floer theory.

Theorem 3.7 (Floer, Robbin–Salamon [17]) *Let L_1 and L_2 be two Lagrangian submanifolds in a symplectic manifold X . Given $\mathbf{x}, \mathbf{y} \in L_1 \cap L_2$ and $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$ a pseudoholomorphic disk connecting them, we have*

$$(-1)^{\mu(\phi)} = \text{deg}(\mathbf{x}) \cdot \text{deg}(\mathbf{y})$$

Here deg denotes the local intersection number of L_1 and L_2 at \mathbf{x} and $\mu(\phi)$ is the Maslov index of ϕ .

In our setting, \mathbb{T}_α and \mathbb{T}_β play the role of Lagrangians in the symplectic manifold $\text{Sym}^g(S)$. Since the mod 2 Maslov index difference depends only on the local intersection degrees of the two intersection points \mathbf{x} and \mathbf{y} in $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$, we need only prove that $\text{sgn}(\gamma_{\mathbf{x}}) \cdot \text{sgn}(\gamma_{\mathbf{y}}) = \text{deg}(\mathbf{x}) \cdot \text{deg}(\mathbf{y})$, a straightforward calculation in local coordinates on $\text{Sym}^g(S)$. \square

3.6 Relationship to twisted Alexander polynomials

We mention some closely-related constructions developed in Wada [20], Kirk and Livingston [5] and Kitano [6].

As usual, we assume Y is an oriented rational homology sphere, and K in Y is an oriented, nullhomologous knot.

Then we have an isomorphism

$$f: H_1(Y) \rightarrow \mathbb{Z}_{a_1} \times \dots \times \mathbb{Z}_{a_k}$$

where the elements a_i are well defined under the added condition that a_i divides a_{i+1} for all $1 \leq i < k$. Let c_i denote $(p_i \circ f)(c)$, where

$$p_i: (\mathbb{Z}_{a_1} \times \dots \times \mathbb{Z}_{a_k}) \rightarrow \mathbb{Z}_{a_i}$$

denotes the projection onto the i th component. Then for each i we have a character $\phi_i: H_1(Y) \rightarrow S^1 \subset \mathbb{C}$ which lands in the cyclotomic field $\mathbb{Q}(\zeta_{a_i})$ for ζ_{a_i} a primitive

a_i th root of unity. By multiplying characters, we obtain the 1–dimensional tensor product representation:

$$\begin{aligned} \phi(c)(z) &= (\phi_1 \otimes \dots \otimes \phi_k)(c)(z) \\ &= \zeta_{a_1}^{c_1} \dots \zeta_{a_k}^{c_k}(z) \end{aligned}$$

Now we can form what is known as the ϕ –twisted Alexander polynomial of K by using $e: \pi \rightarrow \mathbb{Z}$ coming from the intersection number with a Seifert surface and its \mathbb{Z} –ring extension $\epsilon: \mathbb{Z}[\pi] \rightarrow \mathbb{Z}[\mathbb{Z}]$.

Definition 3.8 (Wada) Let Y, K, ϵ be as above. Let \mathbb{F} be a field and $\phi: \mathbb{Z}[\pi] \rightarrow Gl_n(\mathbb{F})$ a representation. Then the ϕ –twisted Alexander polynomial of K in Y is the rational expression

$$\Delta_{K,\phi}(T) = \left(\frac{(\phi \otimes \epsilon)[\partial_2^g]}{T - 1} \right)$$

where ∂_2 is the Fox matrix associated to the presentation

$$\langle h_{\alpha_1}, \dots, h_{\alpha_g} | h_{\beta_1}, \dots, h_{\beta_{g-1}} \rangle$$

of $\pi_1(Y - K)$ and ∂_2^g is ∂_2 with the g th column removed. Here, the presentation is again assumed to have the property that h_{α_g} is a meridian of the knot, implying $e(h_{\alpha_g}) = 1$ and $f(h_{\alpha_g}) = 0$.

Kitano proves, in [6], that Wada’s ϕ –twisted polynomial is the Reidemeister torsion of the chain complex associated to $\phi \otimes \epsilon$. Since this chain complex is exactly the chain complex $C_{\mathbb{Q}(G \times \mathbb{Z})}$ (the only difference is an extra map $\mathbb{Q}(G) \rightarrow \mathbb{Q}(\zeta_{a_1}, \dots, \zeta_{a_k}) \subset \mathbb{C}$, yielding an element of $\mathbb{Q}(\zeta_{a_k})[T, T^{-1}]$),³ Wada’s ϕ –twisted polynomial is the Reidemeister torsion in a slightly different form.

Kirk and Livingston, in [5], define yet another version of a ϕ –twisted Alexander polynomial, which differs slightly from Wada’s definition. They again begin with homomorphisms $e: \pi \rightarrow \mathbb{Z}$ and $f: \pi \rightarrow \mathbb{Z}_{a_1} \times \dots \times \mathbb{Z}_{a_k}$ and form the $\mathbb{Q}(\zeta_{a_k})[T, T^{-1}]$ chain complex

$$C_*(Y - K; \mathbb{Q}(\zeta_{a_k})[T, T^{-1}]_\phi) := \mathbb{Q}(\zeta_{a_1}, \dots, \zeta_{a_k}) \otimes_\phi C_*(\widetilde{Y - K})$$

where here, the T action is given (once a μ with $e(\mu) = 1$ is chosen) by

$$T^n(g \otimes c) = (g \cdot \phi(\mu^{-n})) \otimes (\mu^n \cdot c)$$

extended linearly.

³Note that, since $a_i | a_k$ for all i , $\mathbb{Q}(\zeta_{a_1}, \dots, \zeta_{a_k}) = \mathbb{Q}(\zeta_{a_k})$.

Definition 3.9 (Kirk–Livingston) The i th ϕ -twisted Alexander polynomial, denoted Δ_i , is the order of the torsion of the i th homology of

$$C_*(Y - K; \mathbb{Q}(\zeta_{a_k})[T, T^{-1}]_\phi),$$

considered as a $\mathbb{Q}(\zeta_{a_k})[T, T^{-1}]$ -module.

They go on to prove that Wada's invariant, labeled W , is related to Δ_0 and Δ_1 by the simple formula

$$W = \frac{\Delta_1}{\Delta_0}.$$

4 \widehat{HFK} for double-branched covers of two-bridge knots

We now turn to exploring $\widehat{HFK}(\Sigma^m(K); \widetilde{K})$ in the case $m = 2$ and K a two-bridge knot. Our main result is Theorem 4.3.

We start by recalling a few standard facts about two-bridge knots. A good reference is Chapter 12 of [2].

First, there is a one-to-one correspondence between isotopy classes of two-bridge knots and lens spaces arising as their double branched covers.

Theorem 4.1 [18; 3] *A two bridge knot K in S^3 with twist numbers*

$$(c_1, -c_2, c_3, -c_4, \dots, c_n)$$

(see Figure 3) has double branched covering $-L(p, q)$ where $\frac{p}{q}$ is the continued fraction expansion

$$\frac{p}{q} = c_1 + \frac{1}{c_2 + \frac{1}{\dots + \frac{1}{c_n}}}.$$

We will denote the two-bridge knot whose double branched cover is $-L(p, q)$ by $K(p, q)$.

A particularly useful projection of a two-bridge knot for our purposes is the Schubert normal form. We construct the Schubert normal form of the knot $K(p, q)$ as a union of 4 segments on S^2 : 2 straight “underbridges” U_1 and U_2 and two curvy “overbridges” O_1 and O_2 (All of the following is explained very nicely in [15]).

- (1) A neighborhood of U_1 looks like Figure 4 and a neighborhood of U_2 looks like the mirror image of U_1 reflected across a central vertical axis as in Figure 5.




Figure 3: A two-bridge knot with twist numbers $(c_1, -c_2, c_3, -c_4, \dots, c_n)$




Figure 4: A neighborhood of U_1




Figure 5: A neighborhood of U_2

O_1 and O_2 are formed by connecting $a_i \pmod{2p}$ to $b_{(i-q) \pmod{2p}}$. See Figure 6 for the example of $K(3, 1)$.




Figure 6: Schubert normal form $(U_1 \cup U_2 \cup O_1 \cup O_2)$ for $K(3, 1) =$ the right-handed trefoil

Using Proposition 2.2 in [15], we get a genus 2 handlebody decomposition for $K(p, q)$ that extends to a Heegaard decomposition of S^3 . Schematically, we can draw this handlebody decomposition for $S^3 - K$ by

- (1) placing the feet of two 1–handles at a_0, a_p and b_0, b_p , respectively,
- (2) letting $\alpha_1 = U_1 +$ core of the 1 handle whose feet are at a_0 and a_p , pushed out to the Heegaard surface,
- (3) similarly letting $\alpha_2 = U_2 +$ core of the 1–handle whose feet are at b_0 and b_p , pushed out to the Heegaard surface,
- (4) letting $\beta_1 =$ boundary of a regular neighborhood of either O_1 or O_2 .

Figure 7 provides an illustration of this for $K =$ right-handed trefoil.

Hence, Schubert normal form for a two-bridge knot K yields a genus 2 doubly-pointed Heegaard diagram $\text{hd}(S^3; K; \mu = \beta_2 = h_{\alpha_2})$. One easily checks that $h_{\alpha_1}, h_{\alpha_2}$ are both primitive in $H_1(S^3 - K)$.

As described in Section 2, this handlebody decomposition lifts to one for $\Sigma^2(K) - \tilde{K}$ with an action of \mathbb{Z}_2 . If we denote the non-trivial element of \mathbb{Z}_2 by τ_2 , then $\text{hb}(\Sigma^2(K) - \tilde{K}; \tilde{\mu})$ has

- one 0–handle \tilde{h}_0
- three 1–handles $\tilde{h}_{\alpha_1}, \tau_2(\tilde{h}_{\alpha_1}), \tilde{h}_{\alpha_2}$
- two 2–handles $\tilde{h}_{\beta_1}, \tau_2(\tilde{h}_{\beta_1})$.




Figure 7: A handlebody decomposition for $S^3 - K(3, 1)$

$\text{hb}(\Sigma^2(K); \tilde{K}; \tilde{\mu} = \tilde{h}_{\alpha_2})$ and $\text{hd}(\Sigma^2(K); \tilde{K})$ are as described in Section 2.

We begin our $\widehat{HFK}(\Sigma^2(K); \tilde{K})$ calculation by splitting generators into Spin^c classes. The following (very easy) observation will help us:

Lemma 4.2 *Suppose h is a 1-cycle in S^3 . Let \tilde{h} and $\tau_2(\tilde{h})$ be its two lifts in $\Sigma^2(K)$. Then $\tilde{h} + \tau_2(\tilde{h}) = 0$ in $H_1(\Sigma^2(K))$.*

In particular, if \tilde{h} and $\tau_2(\tilde{h})$ are themselves 1-cycles in $\Sigma^2(K)$, then $\tilde{h} = -\tau_2(\tilde{h})$ in $H_1(\Sigma^2(K))$.

Proof of Lemma 4.2 Let

$$p_!(h) = \sum_{g \in \mathbb{Z}_2} g(\tilde{h})$$

(\tilde{h} is a choice of lift of h) be the transfer map on chains (see, eg, Defn. 11.2 in [1]) associated to the map

$$p: \Sigma^2(K) \rightarrow S^3.$$

Since $h = 0 \in H_1(S^3)$ and the transfer map is a homomorphism on homology, $p_!(h) = \tilde{h} + \tau_2(\tilde{h}) = 0 \in H_1(\Sigma^2(K))$, as desired. \square

Theorem 4.3 *Given any knot K in S^3 and a particular doubly-pointed Heegaard diagram $\text{hd}(S^3; K)$, we can construct $\text{hd}(\Sigma^2(K); \tilde{K})$. Then with respect to these particular Heegaard diagrams there is a natural map on chains*

$$f: \widehat{CFK}(S^3; K) \rightarrow \widehat{CFK}(\Sigma^2(K); \tilde{K})$$

given by

$$f(\mathbf{x}) = (\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$$

for $\mathbf{x} \in \widehat{CFK}(S^3; K)$.

All generators in the image of f lie in the same Spin^c structure, which we will denote \mathfrak{s}_0 .

If K is a two-bridge knot and $\text{hd}(S^3; K)$ is the Heegaard diagram associated to the Schubert normal form of K , then f is a chain map and induces an isomorphism on homology; ie,

$$f_*: \widehat{HFK}(S^3; K) \rightarrow \widehat{HFK}(\Sigma^2(K); \tilde{K}, \mathfrak{s}_0)$$

is an isomorphism.

Here \mathbf{x} is a g -tuple of intersection points of α and β curves in $\text{hd}(S^3; K)$, $\tilde{\mathbf{x}}$ is a choice of lift of that g -tuple in $\text{hd}(\Sigma^2(K); \tilde{K})$, and $\tau_2(\tilde{\mathbf{x}})$ is the image of that g -tuple under the non-trivial deck transformation.

Remark It is worthwhile to mention that in the case where K is a two-bridge knot in S^3 , \mathfrak{s}_0 is the unique spin element of $\text{Spin}^c(\Sigma^2(K))$.

More precisely, consider the first Chern class map $c_1: \text{Spin}^c(Y) \rightarrow H^2(Y; \mathbb{Z})$:

$$c_1(\mathfrak{s}) = \mathfrak{s} - \bar{\mathfrak{s}},$$

where $\bar{\mathfrak{s}}$ is the conjugate Spin^c structure (see [13], Section 2.6). In our situation, the map τ_2 is the conjugation map on Spin^c structures: $\tau_2(\mathfrak{s}) = \bar{\mathfrak{s}}$ for all $\mathfrak{s} \in \text{Spin}^c(Y)$.

But it is clear that $\mathfrak{s}_0 = \tau_2(\mathfrak{s}_0)$, so

$$c_1(\mathfrak{s}_0) = c_1(\bar{\mathfrak{s}}_0);$$

ie, \mathfrak{s}_0 is spin.

Furthermore, \mathfrak{s}_0 is the unique Spin^c structure with this property, for if $\mathfrak{s} \neq \mathfrak{s}_0$, we have $\mathfrak{s} \neq \tau_2(\mathfrak{s})$, implying $c_1(\mathfrak{s}) \neq 0$. But $H^2(Y; \mathbb{Z}) \cong \mathbb{Z}_k$ (k odd) has no 2-torsion, so

$$c_1(\mathfrak{s}) - c_1(\tau_2(\mathfrak{s})) = 2c_1(\mathfrak{s}) \neq 0,$$

so $c_1(\mathfrak{s}) \neq c_1(\bar{\mathfrak{s}})$.

Proof of Theorem 4.3 We begin by showing that all generators of the form $f(\mathbf{x}) = (\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ in $\widehat{CFK}(\Sigma^2(K); \tilde{K})$ are in the same Spin^c structure. Recall [13] that two generators \mathbf{x} and \mathbf{y} of $\widehat{HFK}(Y)$ lie in the same Spin^c structure iff there exists some

path $\mathbf{x} \rightarrow \mathbf{y}$ along α curves and some path $\mathbf{y} \rightarrow \mathbf{x}$ along β curves such that the union represents the 0 element in $H_1(Y)$.

In our situation, we have two generators, $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ and $(\tilde{\mathbf{y}}, \tau_2(\tilde{\mathbf{y}}))$, and we wish to show that we can find a 1-cycle as above representing $0 \in H_1(\Sigma^2(K))$.

But now note that for any two generators \mathbf{x} and \mathbf{y} of $\widehat{HFK}(S^3)$ we can find a path γ_α traveling from \mathbf{x} to \mathbf{y} along α curves and a path γ_β traveling from \mathbf{y} to \mathbf{x} along β curves in the Heegaard diagram for S^3 . The union, $\gamma = \gamma_\alpha \cup \gamma_\beta$, is a closed 1-cycle representing the trivial (only) element in $H_1(S^3)$.

But $p_!(\gamma)$ (where, again, $p_!$ is the transfer map) in the Heegaard diagram $\text{hd}(\Sigma^2(K); \tilde{K}; \alpha_2)$ exactly gives a path from $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ to $(\tilde{\mathbf{y}}, \tau_2(\tilde{\mathbf{y}}))$ along α curves and a path from $(\tilde{\mathbf{y}}, \tau_2(\tilde{\mathbf{y}}))$ to $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ along β curves.

Since $p_!(\gamma) = 0$ in $H_1(\Sigma^2(K))$, all generators of the form $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ lie in the same Spin^c structure, which we have called \mathfrak{s}_0 .

We now turn to showing that, in the case of a two-bridge knot,

$$\widehat{HFK}(\Sigma^2(K); \tilde{K}, \mathfrak{s}_0) \cong \widehat{HFK}(S^3; K).$$

We will do so by showing that the lifted Heegaard diagram associated to the Schubert normal form for K , $\text{hd}(\Sigma^2(K); \tilde{K}; \tilde{h}_{\alpha_2})$, has the property that no other generators lie in \mathfrak{s}_0 . The differentials and filtrations in this central Spin^c structure will match the differentials and filtrations downstairs.

Proposition 4.4 *Let K be a two-bridge knot, $\text{hd}(S^3; K, h_{\alpha_2})$ be the genus 2 Heegaard diagram for K obtained from Schubert normal form, and $\text{hd}(\Sigma^2(K); \tilde{K}, \tilde{h}_{\alpha_2})$ be the lifted genus 3 Heegaard diagram for \tilde{K} in $\Sigma^2(K)$.*

Then the map $f: \widehat{CFK}(S^3; K) \rightarrow \widehat{CFK}(\Sigma^2(K); \tilde{K}; \mathfrak{s}_0)$ described in Theorem 4.3 is a bijection of sets. In particular, all elements of $\widehat{CFK}(\Sigma^2(K); \mathfrak{s}_0)$ are of the form $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ for $\mathbf{x} \in \widehat{CFK}(S^3; K)$.

Proof of Proposition 4.4 We have already shown that all of the τ_2 -invariant generators (those of the form $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$) are in \mathfrak{s}_0 . We now need only show that no non- τ_2 -invariant generators are in \mathfrak{s}_0 .

The proof will rely on the fact that for a two-bridge knot,

$$rk(\widehat{CFK}(S^3; K)) = |H_1(\Sigma^2(K))|,$$

which will imply that no non- τ_2 -invariant generators can appear in \mathfrak{s}_0 .⁴

First, note that for K a two-bridge knot with the standard genus 2 Heegaard decomposition given by Schubert normal form, all generators in $\widehat{CFK}(\Sigma^2(K); \widetilde{K})$ for the genus 3 Heegaard diagram constructed as the lift of this genus 2 Heegaard diagram (as described in Section 2.2) are naturally of the form $(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}))$, where \mathbf{x} and \mathbf{y} are generators in $\widehat{CFK}(S^3)$, and $\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}) \in \mathbb{T}_{\widetilde{\mathbf{f}}_1} \cap \mathbb{T}_{\widetilde{\mathbf{f}}_2}$ are lifts of \mathbf{x} and \mathbf{y} .

More precisely, note that the generators in $\widehat{CFK}(S^3; K)$ are naturally identified with intersection points of α_1 with β_1 (since β_2 , the chosen meridian of the knot, intersects only α_2). Similarly, the generators of $\widehat{CFK}(\Sigma^2(K); \widetilde{K})$ in the lifted Heegaard diagram for $\Sigma^2(K)$, are in one-to-one correspondence with formal summands of the determinant

$$\begin{matrix} & \widetilde{\beta}_1 & \tau_2(\widetilde{\beta}_1) \\ \widetilde{\alpha}_1 & \left(\begin{matrix} & \\ & \end{matrix} \right) \\ \tau_2(\widetilde{\alpha}_1) & & \end{matrix}$$

where the entries in the matrix above are formal sums of intersection points of the corresponding α and β curves. A summand of this matrix can be thought of as a pair $(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}))$ where \mathbf{x}, \mathbf{y} are generators of $\widehat{CFK}(S^3; K)$ and $\widetilde{\mathbf{x}}$ and $\widetilde{\mathbf{y}}$ are the lifts which lie on the $\widetilde{\alpha}_1$ curve.⁵

We can then measure the Spin^c structure of a non- τ_2 -invariant generator $(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}))$ by comparing it to the τ_2 -invariant generator $(\widetilde{\mathbf{y}}, \tau_2(\widetilde{\mathbf{y}}))$. More precisely:

Lemma 4.5 *If $(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}))$ is a generator of $\widehat{CFK}(\Sigma^2(K), \widetilde{K})$, then its corresponding element of $\text{Spin}^c(\Sigma^2(K))$ relative to \mathfrak{s}_0 , ie,*

$$\mathfrak{s}(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}})) - \mathfrak{s}(\widetilde{\mathbf{y}}, \tau_2(\widetilde{\mathbf{y}})),$$

thought of as an element of $H_1(\Sigma^2(K))$, is represented by the lift, $\widetilde{\gamma}$, of the word γ in $\pi_1(S^3 - K)$ read off as we travel from \mathbf{x} to \mathbf{y} along β_1 .

Proof of Lemma 4.5 $\mathfrak{s}(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}})) - \mathfrak{s}(\widetilde{\mathbf{y}}, \tau_2(\widetilde{\mathbf{y}}))$, as an element of $H_1(\Sigma^2(K))$, is represented by the cycle obtained by connecting $(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}))$ to $(\widetilde{\mathbf{y}}, \tau_2(\widetilde{\mathbf{y}}))$ along α curves and $(\widetilde{\mathbf{y}}, \tau_2(\widetilde{\mathbf{y}}))$ to $(\widetilde{\mathbf{x}}, \tau_2(\widetilde{\mathbf{y}}))$ along β curves (see Definition 2.4 and Section 2.6 of [13]; also see [19]).

We construct such a path as the product of:

⁴This condition on $rk(\widehat{CFK}(S^3; K))$ holds for a wider class of knots (eg, alternating knots), and therefore, similar results may hold in wider generality.

⁵Such an identification of generators of $\widehat{CFK}(\Sigma^2(K); \widetilde{K})$ with pairs of generators of $\widehat{CFK}(S^3; K)$ is possible for a general choice of Heegaard diagram for K in S^3 , but different generators will require different lifts of \mathbb{T}_α and \mathbb{T}_β in order to make the identification.

- the appropriate lift of a loop between \mathbf{x} and \mathbf{y} (along α_1 and back along β_1) in $\text{hd}(S^3; K)$ to a loop between $\tilde{\mathbf{x}}$ and $\tilde{\mathbf{y}}$ in $\text{hd}(\Sigma^2(K); \tilde{K})$,
- the constant path from $\tau_2(\tilde{\mathbf{y}})$ to $\tau_2(\tilde{\mathbf{y}})$,
- a path from the lone intersection point between $\tilde{\beta}_2$ and $\tilde{\alpha}_2$ to itself along $\tilde{\beta}_2$.

The element of $H_1(\Sigma^2(K))$ represented by this path is represented by the word in $\pi_1(\Sigma^2(K))$ read off as we travel from $\tilde{\mathbf{x}}$ to $\tilde{\mathbf{y}}$ along $\tilde{\beta}_1$, which is the lift, $\tilde{\gamma}$, of the word γ in $\pi_1(S^3 - K)$ read off as we travel from \mathbf{x} to \mathbf{y} along β_1 . \square

But now I claim that, since $rk(\widehat{CFK}(S^3; K)) = |H_1(\Sigma^2(K))|$ for the Heegaard diagram associated to Schubert normal form for K a two-bridge knot (see [16], [15], [10]), $\tilde{\gamma} \neq 0$ unless $\mathbf{x} = \mathbf{y}$.

Lemma 4.6 *Let K a two-bridge knot in S^3 and \mathbf{x}, \mathbf{y} two generators of $\widehat{CFK}(S^3; K)$ associated to the handlebody decomposition coming from Schubert normal form. Let $\gamma(\mathbf{x}, \mathbf{y})$ be the word in $\pi_1(S^3 - K)$ read off as we travel from \mathbf{x} to \mathbf{y} along β_1 and $\tilde{\gamma}(\mathbf{x}, \mathbf{y})$ a lift of γ to a word in $\pi_1(\Sigma^2(K) - \tilde{K})$.*

Then $\mathbf{x} \neq \mathbf{y}$ implies that $\tilde{\gamma}(\mathbf{x}, \mathbf{y}) \neq 0$ as an element of $H_1(\Sigma^2(K))$.

Proof of Lemma 4.6 The crucial observation is that the Fox matrix associated to the homomorphism

$$\pi_1(S^3 - K) \rightarrow \mathbb{Z}_2$$

is a presentation matrix for $H_1(\Sigma^2(K))$, and its summands are in natural one-to-one correspondence with the generators of $\widehat{CFK}(S^3; K)$. See Section 3.5 for a more detailed discussion of Fox calculus.

Fix a generator, \mathbf{x} , of $\widehat{CFK}(S^3; K)$ and begin reading off the relation corresponding to the boundary of β_1 , beginning at \mathbf{x} . Let

$$t: \pi_1(S^3 - K) \rightarrow \mathbb{Z}_2$$

be the homomorphism inducing the branched double cover of K and

$$\tau: \mathbb{Z}[\pi_1(S^3 - K)] \rightarrow \mathbb{Z}[\mathbb{Z}_2]$$

the group-ring extension.

Recall that the generators of $\widehat{CFK}(S^3; K)$ correspond one-to-one with the intersection points of α_1 and β_1 . For convenience, label the generators of $\widehat{CFK}(S^3; K)$ by \mathbf{y}_i , according to the order in which we encounter them as we travel along β_1 from \mathbf{x} .

Then

$$\frac{\partial \beta_1}{\partial \alpha_1} = \sum_{\mathbf{y}_i \in \widehat{CFK}(S^3; K)} \pm \gamma(\mathbf{x}, \mathbf{y}_i);$$

ie, the Fox derivative of β_1 by α_1 is the formal sum of the words connecting \mathbf{x} to each of the other generators of $\widehat{CFK}(S^3; K)$, with signs in the sum given by the local intersection number of $\alpha_1 \cap \beta_1$ at \mathbf{y}_i . Notice that there are exactly $rk(\widehat{CFK}(S^3; K))$ summands in the Fox determinant.

Furthermore, $\tau(\frac{\partial \beta_1}{\partial \alpha_1})$ is a presentation matrix for $H_1(\Sigma^2(K))$, after replacing the formal elements of \mathbb{Z}_2 by square roots of unity (± 1).

But $rk(\widehat{CFK}(S^3; K)) = |H_1(\Sigma^2(K))|$, so all summands must have the same sign once we replace the formal elements of \mathbb{Z}_2 with ± 1 and take into account local intersection multiplicities.

Since the element of $H_1(\Sigma^2(K))$ represented by $\tilde{\gamma}(\mathbf{x}, \mathbf{y}_k)$ is precisely

$$\sum_{i=1}^k \tau(\pm \gamma(\mathbf{x}, \mathbf{y}_i)) \cdot h_{\tilde{\alpha}_1},$$

and $h_{\tilde{\alpha}_1}$ is a primitive generator of $H_1(\Sigma^2(K))$, we conclude that $\tilde{\gamma}(\mathbf{x}, \mathbf{y})$ cannot be 0 in $H_1(\Sigma^2(K))$ for any $\mathbf{y} \neq \mathbf{x}$. \square

The following two lemmas prove that the map

$$f: \widehat{CFK}(S^3; K) \rightarrow \widehat{CFK}(\Sigma^2(K); \tilde{K}; \mathfrak{s}_0)$$

given above also preserves the relative filtration and homological gradings in the case that K is a two-bridge knot. Hence, f is a chain map respecting the filtration, so $f_*: \widehat{HFK}(S^3; K) \rightarrow \widehat{HFK}(\Sigma^2(K); \tilde{K}; \mathfrak{s}_0)$ is an isomorphism.

Lemma 4.7 For K any knot in S^3 ,

$$s(\mathbf{x}, \mathbf{y}) = s(f(\mathbf{x}), f(\mathbf{y}))$$

for all pairs of generators \mathbf{x} and \mathbf{y} in $\widehat{HFK}(S^3; K)$.

Proof of Lemma 4.7 Let \mathbf{x} and \mathbf{y} be two generators in $\widehat{HFK}(S^3; K)$ and $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$ a topological disk with $n_z(\phi) - n_w(\phi) = 1$.

We have already observed in the proof of Theorem 4.3 that if γ is the boundary of the image of ϕ in the Heegaard surface, S , then $\tilde{\gamma} = p_1(\gamma)$ will be the boundary of the image of $\tilde{\phi} \in \pi_2(f(\mathbf{x}), f(\mathbf{y}))$ in \tilde{S} .

Lemma 3.5 asserts that $s(\mathbf{x}, \mathbf{y})$ is equal to the coefficient on μ , our choice of meridian, in γ .

But by the way we have defined $\text{hd}(\Sigma^2(K); \tilde{K}, \tilde{\mu})$, the coefficient on $\tilde{\mu}$ in $\tilde{\gamma}$ is equal to the coefficient on μ in γ . \square

Lemma 4.8 For K a two-bridge knot in S^3 and $\text{hd}(S^3; K)$ a Heegaard diagram for K coming from Schubert normal form as before,

$$m(\mathbf{x}, \mathbf{y}) = m(f(\mathbf{x}), f(\mathbf{y})).$$

Proof of Lemma 4.8 Consider $\text{hd}(S^3; K)$ and $\text{hd}(\Sigma^2(K); \tilde{K}, \tilde{h}_{\alpha_2})$ obtained from Schubert normal form. It will be convenient to destabilize each of these Heegaard diagrams once by canceling the $h_{\alpha_2}, h_{\beta_2}$ pair and the $\tilde{h}_{\alpha_2}, \tilde{h}_{\beta_2}$ pair, respectively. Call these destabilized Heegaard diagrams $\text{hd}^\circ(S^3; K)$ and $\text{hd}^\circ(\Sigma^2(K); \tilde{K}, \tilde{h}_{\alpha_2})$, respectively.

Note that there is a natural one-to-one correspondence between \widehat{CFK} generators corresponding to hd and hd° (see Prop. 6.1 in [11]) which induces the isomorphism on \widehat{HFK} corresponding to the (de)stabilization.

This destabilized Heegaard diagram for K coming from Schubert normal form is particularly nice because we can find relative Maslov gradings between all generators just by looking at disks coming from “finger moves.”

Namely, we know that we obtain $\text{hd}^\circ(S^3; K)$ from the standard genus 1 Heegaard diagram for S^3 by performing finger moves of the curve β across the curve α .

In fact, one can check that for each generator, there is a natural disk connecting it to at least one adjacent generator by a finger. Each of these disks has either $n_w = 1, n_z = 0$ or $n_w = 0, n_z = 1$, and these “finger disks” are enough to determine the relative Maslov grading of any two generators.

Now, suppose that \mathbf{x} and \mathbf{y} are two generators in $\widehat{CFK}(S^3; K)$ connected by a finger disk. Then $(\tilde{\mathbf{x}}, \tau_2(\tilde{\mathbf{x}}))$ and $(\tilde{\mathbf{y}}, \tau_2(\tilde{\mathbf{y}}))$ are connected by the lift of the finger disk, which is a quadrilateral. Such a quadrilateral always represents a holomorphic disk in $\text{Sym}^2(S)$ of Maslov index 1.

Therefore,

$$m(f(\mathbf{x}), f(\mathbf{y})) = m(\mathbf{x}, \mathbf{y})$$

whenever \mathbf{x} and \mathbf{y} are connected by a finger disk.

Now we claim that every generator is connected to every other generator by some series of finger disks. We can see this as follows:

Notice that β_1 in $\text{hd}^\circ(S^3; K)$ is the image, after destabilizing the $\text{hd}(S^3; K)$ coming from Schubert normal form, of a regular neighborhood of either one of the overbridges (note that a regular neighborhood of O_1 is isotopic to a regular neighborhood of O_2). Similarly, α_1 is the union of one of the underbridges (whichever one was not canceled in the destabilization) – U_1 , say – with the core of the 1–handle whose feet are placed at a_0 and a_p .

Now, focus on any two adjacent generators of the Heegaard Floer homology (see Figure 8). These two adjacent generators correspond to where β_1 , considered either as $N(O_1)$ or $N(O_2)$, intersects α_1 . For definiteness, say that the two adjacent generators we are considering are associated to $N(O_1)$. Note that O_1 can be decomposed as the union of

- a small overbridge O_{small} , which exits a neighborhood of U_1 at a_i and a_{2p-i}
- an arc, O_a , connecting one of a_i and a_{2p-i} with an endpoint of U_1 (either a_0 or a_p), and
- an arc, O_b , connecting the other endpoint (a_{2p-i} or a_i) with an endpoint of U_2 (either b_0 or b_p).

Since b_p and b_0 are where the basepoints w and z are positioned in the destabilized Heegaard diagram, a regular neighborhood of the arc connecting a_i (or a_{2p-i}) to b_0 (or b_p) will be a finger disk connecting these two adjacent generators.

In other words, every two adjacent generators is connected by a finger disk. So every pair of generators is connected by some sequence of finger disks.

Therefore,

$$m(f(\mathbf{x}), f(\mathbf{y})) = m(\mathbf{x}, \mathbf{y})$$

for all pairs of generators \mathbf{x}, \mathbf{y} in $\widehat{CFK}(S^3; K)$, as desired. \square

4.1 Examples: $K(15, 7)$ and $K(15, 4)$

$\widehat{HFK}(\Sigma^2(K); \widetilde{K})$ distinguishes between knots with the same knot Floer homology. In [10] Ozsváth and Szabó prove that for K an alternating knot in S^3 , the knot Floer homology is determined completely by the Alexander polynomial and the signature.

$K(15, 7)$ and $K(15, 4)$ are two-bridge knots with the same Alexander polynomial and signature but differing \mathbb{Z}_2 –graded knot Floer homologies in the double branched cover.⁶

⁶The double-branched covers of $K(15, 7)$ and $K(15, 4)$ are different 3-manifolds (the lens spaces $-L(15, 7)$ and $-L(15, 4)$, respectively). In fact, $\widehat{HF}(\Sigma^2(K(15, 7))) \not\cong \widehat{HF}(\Sigma^2(K(15, 4)))$ as \mathbb{Q} –graded groups (See Proposition 4.8 in [9] for an inductive formula for the \mathbb{Q} grading of generators in lens spaces). This immediately implies that $\widehat{HFK}(\Sigma^2(K(15, 7)); \widetilde{K}(15, 7)) \not\cong \widehat{HFK}(\Sigma^2(K(15, 4)); \widetilde{K}(15, 4))$ as \mathbb{Q} –graded groups. We will show the stronger statement that $\widehat{HFK}(\Sigma^2(K(15, 7)); \widetilde{K}(15, 7)) \not\cong \widehat{HFK}(\Sigma^2(K(15, 4)); \widetilde{K}(15, 4))$ as \mathbb{Z}_2 –graded groups.




Figure 8: Adjacent generators in a genus 1 (destabilized) Heegaard diagram compatible with a two-bridge knot




Figure 9: \mathbb{Z} -filtered chain complex for $\widehat{CFK}(S^3; K(15, 7))$

The computations of both $\widehat{HFK}(S^3; K)$ and $\widehat{HFK}(\Sigma^2(K); \widetilde{K})$ for these two knots are given below.

Computation for $K(15, 7)$

We start by computing $\widehat{HFK}(S^3; K)$ for $K = K(15, 7)$. In the genus 1 Heegaard diagram compatible with K given in Figure 10, we have 15 generators, which we label




Figure 10: Genus 1 (destabilized) Heegaard diagram for $K(15, 7)$ in S^3

\mathfrak{s}_0	$\mathfrak{s}_{\pm 1}$	$\mathfrak{s}_{\pm 2}$	$\mathfrak{s}_{\pm 3}$	$\mathfrak{s}_{\pm 4}$
$(x_1, x_1), (y_1, y_1)$	(x_1, x_8)	$(x_1, x_2), (y_1, y_2)$	(x_1, x_7)	$(x_1, x_3), (y_1, y_3)$
$(x_2, x_2), (y_2, y_2)$		$(x_2, x_3), (y_2, y_3)$	(x_2, x_8)	$(x_2, x_4), (y_2, y_4)$
$(x_3, x_3), (y_3, y_3)$		$(x_3, x_4), (y_3, y_4)$	(y_1, y_7)	$(x_3, x_5), (y_3, y_5)$
$(x_4, x_4), (y_4, y_4)$		$(x_4, x_5), (y_4, y_5)$		$(x_4, x_6), (y_4, y_6)$
$(x_5, x_5), (y_5, y_5)$		$(x_5, x_6), (y_5, y_6)$		$(x_5, x_7), (y_5, y_7)$
$(x_6, x_6), (y_6, y_6)$		$(x_6, x_7), (y_6, y_7)$		(x_6, x_8)
$(x_7, x_7), (y_7, y_7)$		$(x_7, x_8), (y_7, y_8)$		
$(x_8, x_8),$				

$\mathfrak{s}_{\pm 5}$	$\mathfrak{s}_{\pm 6}$	$\mathfrak{s}_{\pm 7}$
(x_1, x_6)	$(x_1, x_4), (y_1, y_4)$	$(x_1, x_5), (y_1, y_5)$
(y_1, y_6)	$(x_2, x_5), (y_2, y_5)$	$(x_2, x_6), (y_2, y_6)$
(x_2, x_7)	$(x_3, x_6), (y_3, y_6)$	$(x_3, x_7), (y_3, y_7)$
(y_2, y_7)	$(x_4, x_7), (y_4, y_7)$	(x_4, x_8)
(x_3, x_8)	(x_5, x_8)	

Table 1: Spin^c structures $\mathfrak{s}_0, \dots, \mathfrak{s}_{\pm 7}$ for $\Sigma^2(K(15, 7))$




Figure 11: Genus 2 (destabilized) Heegaard diagram for $\tilde{K}(15, 7)$ in $L(15, 7)$

x_1, \dots, x_8 and y_1, \dots, y_7 . This Heegaard diagram for S^3 compatible with K was obtained by taking the handlebody decomposition of $S^3 - K$ coming from the Schubert normal form for K and destabilizing once.

The differential counts maps of the disk into $\text{Sym}^1(S) = S$. See Figure 10 for an example.

The filtered chain complex for \widehat{CF} is pictured in Figure 9. The i -th vertical slice of this filtered chain complex is the chain complex $\mathcal{F}_i/\mathcal{F}_{i-1} = \widehat{CFK}(S^3; K, i)$.

We construct a Heegaard diagram for $\Sigma^2(K)$ compatible with \tilde{K} by taking the branched double cover of the Heegaard surface Σ around the two basepoints w and z . The α and β curves of our original Heegaard diagram then lift to two α curves and two β curves, and w and z lift to the two basepoints for the doubly-pointed Heegaard diagram for $\Sigma^2(K)$.

All generators of $\widehat{HFK}(\Sigma^2(K); \tilde{K})$ are of the form $(\tilde{x}_i, \tau_2(\tilde{x}_j))$ or $(\tilde{y}_i, \tau_2(\tilde{y}_j))$, where τ_2 is the non-trivial element of \mathbb{Z}_2 . Under the \mathbb{Z}_2 action on the α and β curves, we get a natural \mathbb{Z}_2 action on these generators:

$$\tau_2(\tilde{x}_i, \tau_2(\tilde{x}_j)) = (\tau_2(\tilde{x}_i), \tilde{x}_j).$$




Figure 12: Two holomorphic disks, one in $\pi_2(x_5, y_4)$ and the other in $\pi_2(y_5, x_4)$

We shall refer to such a pair of generators as a *conjugate pair*, since they are in conjugate Spin^c structures with respect to the central Spin^c structure, \mathfrak{s}_0 .

τ_2 -invariant generators

The τ_2 -invariant generators are those of the form $(\tilde{x}_i, \tau_2(\tilde{x}_i))$ or $(\tilde{y}_i, \tau_2(\tilde{y}_i))$. Furthermore, for every topological bigon downstairs we see a corresponding topological quadrilateral upstairs (as detailed in the proof of Lemma 4.8). See the shaded disk in $\pi_2(y_5, x_4)$ in Figure 10 and its lift in $\pi_2((\tilde{y}_5, \tau_2(\tilde{y}_5)), (\tilde{x}_4, \tau_2(\tilde{x}_4)))$ in Figure 11.

τ_2 -non-invariant generators

We will first state how the generators split up according to Spin^c structures, then explicitly compute $\widehat{HFK}(\Sigma^2(K); \tilde{K}; \mathfrak{s}_{\pm 2})$. Here we use $\mathfrak{s}_{\pm 2}$ to denote the conjugate Spin^c structures on $\Sigma^2(K)$ corresponding to $\pm 2 \in H_1(\Sigma^2(K)) \cong \mathbb{Z}_{15}$.

Each column in Table 1 gives the generators in two conjugate Spin^c structures simultaneously (except in the case of the “central” Spin^c structure, \mathfrak{s}_0 , which is its own conjugate). When we write (x_1, x_8) , for example, we refer to two generators simultaneously: $(\tilde{x}_1, \tau_2(\tilde{x}_8)) \in \mathfrak{s}_1$ and $(\tau_2(\tilde{x}_1), \tilde{x}_8) \in \mathfrak{s}_{-1}$.

Now let’s more closely examine one of the Spin^c structures, $\mathfrak{s}_{\pm 2}$.




Figure 13: Two holomorphic disks, one in $\pi_2((\tilde{y}_5, \tau_2(\tilde{y}_4)), (\tilde{x}_4, \tau_2(\tilde{x}_5)))$ and one in $\pi_2((\tilde{y}_4, \tau_2(\tilde{y}_5)), (\tilde{x}_5, \tau_2(\tilde{x}_4)))$

To see that these are the generators in the two conjugate Spin^c structures $\mathfrak{s}_{\pm 2}$, connect, for example, $(\tilde{x}_1, \tau_2(\tilde{x}_1))$ to $(\tilde{x}_1, \tau_2(\tilde{x}_2))$ by the path in $\text{Sym}^2(\tilde{S})$ which is the product of the constant path $\tilde{x}_1 \rightarrow \tilde{x}_1$ with the path $\tau_2(\tilde{x}_1) \rightarrow \tau_2(\tilde{x}_2)$ along the $\tau_2(\tilde{\beta})$ curve. Close the path to a loop γ by taking a path $\tau_2(\tilde{x}_1) \rightarrow \tau_2(\tilde{x}_2)$ along the $\tau_2(\tilde{\alpha})$ curve. Since

$$\tilde{\alpha} \cdot \tilde{\beta} = \tau_2(\tilde{\alpha}) \cdot \tau_2(\tilde{\beta}) = 1,$$

and

$$\tilde{\alpha} \cdot \tau_2(\tilde{\beta}) = \tau_2(\tilde{\alpha}) \cdot \tilde{\beta} = -1,$$

we see that if we assert that $\tilde{\alpha}$ is the positive generator of $H_1(\Sigma^2(K))$, then the loop γ represents the element $2 \in H_1(\Sigma^2(K))$.

We see this because a pushoff of γ has one intersection with $\tau_2(\tilde{\alpha})$ with local multiplicity -1 (at $\tau_2(\tilde{y}_1)$) and one intersection with $\tilde{\alpha}$ with local multiplicity 1 (at \tilde{x}_2). Recall that $\tau_2(\tilde{\alpha}) = -\tilde{\alpha}$ in $H_1(\Sigma^2(K))$.

The same type of calculation can be performed to verify that all of the other generators are in the stated Spin^c structures.




Figure 14: Two holomorphic disks, one in $\pi_2(x_5, y_4)$ and the other in $\pi_2(x_6, y_3)$

Now focus on, for example, $(\tilde{x}_4, \tau_2(\tilde{x}_5))$ and $(\tilde{y}_5, \tau_2(\tilde{y}_4))$ (and the corresponding pair $(\tau_2(\tilde{x}_4), \tilde{x}_5), (\tau_2(\tilde{y}_5), \tilde{y}_4)$ in the conjugate Spin^c structure).

Notice that in the Heegaard diagram for $(S^3, K(15, 7))$ pictured in Figure 12 we see a topological disk in $\pi_2(x_5, y_4)$ and a topological disk in $\pi_2(y_5, x_4)$, both with $n_w = 0$ and $n_z = 1$.

If we lift both of these to the Heegaard diagram for $(\Sigma^2(K); \tilde{K})$ we get two quadrilaterals, one representing a disk in $\pi_2((\tilde{x}_5, \tau_2(\tilde{x}_5)), (\tilde{y}_4, \tau_2(\tilde{y}_4)))$ and one in

$$\pi_2((\tilde{y}_5, \tau_2(\tilde{y}_5)), (\tilde{x}_4, \tau_2(\tilde{x}_4))).$$

Taking the difference of these two disks yields two disks, one in

$$\pi_2((\tilde{y}_4, \tau_2(\tilde{y}_5)), (\tilde{x}_5, \tau_2(\tilde{x}_4)))$$

and one in $\pi_2((\tilde{y}_5, \tau_2(\tilde{y}_4)), (\tilde{x}_4, \tau_2(\tilde{x}_5)))$. Both of these disks have $n_z = n_w = 0$ and Maslov index 1. Both of these disks are holomorphic because they are topological quadrilaterals (see Figure 13).

A similar argument shows that there is another pair of disks, one in

$$\pi_2((\tilde{y}_4, \tau_2(\tilde{y}_3)), (\tilde{x}_4, \tau_2(\tilde{x}_5)))$$




Figure 15: A holomorphic disk in $\pi_2((\tilde{x}_6, \tau_2(\tilde{x}_5)), (\tau_2(\tilde{y}_4), \tilde{y}_3))$

and one in

$$\pi_2((\tau_2(\tilde{y}_4), \tilde{y}_3), (\tau_2(\tilde{x}_4), \tilde{x}_5)),$$

both with $n_z = n_w = 0$.

There also exist topological disks in $\text{Sym}^2(\tilde{S})$ which are the sum of a pair of lifts of disks. Focus, for example, on the pair $(\tilde{x}_6, \tau_2(\tilde{x}_5))$ and $(\tau_2(\tilde{y}_4), \tilde{y}_3)$. In the Heegaard diagram downstairs, we see a disk in $\pi_2(x_6, y_3)$ with $n_w = 0$ and $n_z = 1$. We also see a disk in $\pi_2(x_5, y_4)$. See Figure 14.

The sum of the lifts of these two disks to the Heegaard diagram for $\Sigma^2(K)$ again breaks up into two disks. One of the disks is in

$$\pi_2((\tilde{x}_6, \tau_2(\tilde{x}_5)), (\tau_2(\tilde{y}_4), \tilde{y}_3))$$

and the other is between the conjugate generators, ie, in

$$\pi_2((\tau_2(\tilde{x}_6), \tilde{x}_5), (\tilde{y}_4, \tau_2(\tilde{y}_3))).$$

Both of these disks have $n_w = 0$ and $n_z = 1$. See Figure 15 for one of the two disks.

Using similarly obtained disks, we easily calculate the relative filtration levels and Maslov grading of all generators in $\mathfrak{s}_{\pm 2}$. The generators of the \widehat{CFK} complex for




Figure 16: $C\widehat{FK}(\Sigma^2(K), \widetilde{K}(15, 7), \mathfrak{s}_{\pm 2})$




Figure 17: \mathbb{Z} -filtered chain complex for $C\widehat{FK}(S^3; K(15, 4))$

$K(15, 7)$ as well as arrows corresponding to some of the differentials are pictured in Figure 16. We omit the \sim 's and the τ_2 's, since we are thinking of this as the chain complex for both \mathfrak{s}_2 and \mathfrak{s}_{-2} .




Figure 18: Genus 1 (destabilized) Heegaard diagram for $K(15, 4)$ in S^3

\mathfrak{s}_0	$\mathfrak{s}_{\pm 1}$	$\mathfrak{s}_{\pm 2}$	$\mathfrak{s}_{\pm 3}$	$\mathfrak{s}_{\pm 4}$
$(x_1, x_1), (y_1, y_1)$	$(x_2, x_3), (y_2, y_3)$	(x_1, x_2)	$(x_3, x_4), (y_3, y_4)$	$(x_2, x_4), (y_2, y_4)$
$(x_2, x_2), (y_2, y_2)$	$(x_4, x_5), (y_4, y_5)$		$(x_5, x_6), (y_5, y_6)$	$(x_3, x_5), (y_3, y_5)$
$(x_3, x_3), (y_3, y_3)$	$(x_6, x_7), (y_6, y_7)$		$(x_7, x_8), (y_1, y_2)$	$(x_4, x_6), (y_4, y_6)$
$(x_4, x_4), (y_4, y_4)$	(x_1, x_8)		$(x_2, x_8), (y_1, y_7)$	$(x_5, x_7), (y_5, y_7)$
$(x_5, x_5), (y_5, y_5)$			(x_1, x_3)	$(x_6, x_8), (y_1, y_3)$
$(x_6, x_6), (y_6, y_6)$				$(x_3, x_8), (y_1, y_6)$
$(x_7, x_7), (y_7, y_7)$				(x_1, x_7)
(x_8, x_8)				

$\mathfrak{s}_{\pm 5}$	$\mathfrak{s}_{\pm 6}$	$\mathfrak{s}_{\pm 7}$
$(x_2, x_5), (y_2, y_5)$	$(x_2, x_7), (y_2, y_7)$	$(x_1, x_5), (y_1, y_5)$
$(x_4, x_7), (y_4, y_7)$	(x_1, x_4)	$(x_2, x_6), (y_2, y_6)$
(x_1, x_6)		$(x_3, x_7), (y_3, y_7)$
		$(x_3, x_6), (y_3, y_6)$
		$(x_5, x_8), (y_1, y_4)$
		(x_4, x_8)

Table 2: Spin^c structures $\mathfrak{s}_0, \dots, \mathfrak{s}_{\pm 7}$ for $\Sigma^2(K(15, 4))$




Figure 19: Genus 2 (destabilized) Heegaard diagram for $\tilde{K}(15, 4)$ in $L(15, 4)$




Figure 20: $C\hat{F}K(\Sigma^2(K), \tilde{K}(15, 4), \mathfrak{s}_{\pm 1})$




Figure 21: $C\widehat{FK}(\Sigma^2(K), \widetilde{K}(15, 4), \mathfrak{s}_{\pm 2})$




Figure 22: $C\widehat{FK}(\Sigma^2(K), \widetilde{K}(15, 4), \mathfrak{s}_{\pm 4})$

Note that we have made no claims about whether there are any more holomorphic disks than the ones described.⁷ However, just based on the information contained in Figure 16 we can see that $\widehat{HFK}(\Sigma^2(K); \widetilde{K}; \mathfrak{s}_{\pm 2})$ must have support in three different filtration levels, with

$$\widehat{HFK}(\Sigma^2(K); \widetilde{K}, \mathfrak{s}_{\pm 2}, i - 1) \cong \widehat{HFK}(\Sigma^2(K); \widetilde{K}, \mathfrak{s}_{\pm 2}, i + 1) \cong \mathbb{Z}_2^3.$$

⁷In fact, we cannot possibly have listed all of the d_1 differentials, for $(d_0 + d_1)^2 \neq 0$ in Figure 16.




Figure 23: $C\widehat{FK}(\Sigma^2(K), \widetilde{K}(15, 4), \mathfrak{s}_{\pm 7})$

Computation for $K(15, 4)$

We now turn to computations of $\widehat{HFK}(S^3; K)$ and $\widehat{HFK}(\Sigma^2(K); \widetilde{K})$ for $K = K(15, 4)$. Recall that our aim is to prove that

$$\widehat{HFK}(\Sigma^2(K); \widetilde{K}(15, 7)) \not\cong \widehat{HFK}(\Sigma^2(K); \widetilde{K}(15, 4))$$

as \mathbb{Z}_2 -graded groups. We will do so by showing that $\widehat{HFK}(\Sigma^2(K); \widetilde{K}(15, 4), \mathfrak{s}_k)$ does not look like $\widehat{HFK}(\Sigma^2(K); \widetilde{K}(15, 7), \mathfrak{s}_{\pm 2})$ (computed in the previous subsection) for any $k \in \mathbb{Z}_{15}$ with order 15.

In the genus 1 Heegaard diagram compatible with K given in Figure 18, we have 15 generators, which we label x_1, \dots, x_8 and y_1, \dots, y_7 . See Figure 17 for the filtered chain complex.

We construct a genus 2 Heegaard diagram for $\Sigma^2(K)$ compatible with \widetilde{K} in exactly the same way we did before (see Figure 19).

τ_2 -non-invariant generators

The generators split up according to Spin^c structures as detailed in Table 2.

Using the methods described in the previous subsection, we find relative filtration and Maslov gradings for $\mathfrak{s}_{\pm 1}, \mathfrak{s}_{\pm 2}, \mathfrak{s}_{\pm 4}$, and $\mathfrak{s}_{\pm 7}$. The generators of the \widehat{CFK} complex for $K(15, 4)$ in these Spin^c structures as well as arrows corresponding to some of the differentials are pictured in Figures 20 - 23.

Again, we make no claims about whether there are any more holomorphic disks than the ones described. However, we have enough information about the homology of the associated graded chain complexes to determine that

$$\widehat{HFK}(\Sigma^2(K); \tilde{K}(15, 7), \mathfrak{s}_{\pm 2}) \not\cong \widehat{HFK}(\Sigma^2(K); \tilde{K}(15, 4), \mathfrak{s}_k)$$

for any k relatively prime to 15.

References

- [1] **G E Bredon**, *Topology and geometry*, volume 139 of *Graduate Texts in Mathematics*, Springer-Verlag, New York (1997) MR1700700 Corrected third printing of the 1993 original
- [2] **G Burde, H Zieschang**, *Knots*, de Gruyter Studies in Mathematics 5, Walter de Gruyter & Co., Berlin (2003) MR1959408
- [3] **J H Conway**, *An enumeration of knots and links, and some of their algebraic properties*, from: “Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)”, Pergamon, Oxford (1970) 329–358 MR0258014
- [4] **R H Fox**, *A quick trip through knot theory*, from: “Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)”, Prentice-Hall, Englewood Cliffs, N.J. (1962) 120–167 MR0140099
- [5] **P Kirk, C Livingston**, *Twisted Alexander invariants, Reidemeister torsion, and Casson-Gordon invariants*, *Topology* 38 (1999) 635–661
- [6] **T Kitano**, *Twisted Alexander polynomial and Reidemeister torsion*, *Pacific J. Math.* 174 (1996) 431–442 MR1405595
- [7] **J Milnor**, *Whitehead torsion*, *Bull. Amer. Math. Soc.* 72 (1966) 358–426 MR0196736
- [8] **J W Milnor**, *Infinite cyclic coverings*, from: “Conference on the Topology of Manifolds (Michigan State Univ., E. Lansing, Mich., 1967)”, Prindle, Weber & Schmidt, Boston (1968) 115–133 MR0242163
- [9] **P Ozsváth, Z Szabó**, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, *Adv. Math.* 173 (2003) 179–261 MR1957829
- [10] **P Ozsváth, Z Szabó**, *Heegaard Floer homology and alternating knots*, *Geom. Topol.* 7 (2003) 225–254 MR1988285
- [11] **P Ozsváth, Z Szabó**, *Holomorphic disks and knot invariants*, *Adv. Math.* 186 (2004) 58–116 MR2065507
- [12] **P Ozsváth, Z Szabó**, *Holomorphic disks and three-manifold invariants: properties and applications*, *Ann. of Math. (2)* 159 (2004) 1159–1245 MR2113020
- [13] **P Ozsváth, Z Szabó**, *Holomorphic disks and topological invariants for closed three-manifolds*, *Ann. of Math. (2)* 159 (2004) 1027–1158 MR2113019

- [14] **P Ozsváth, Z Szabó**, *Holomorphic triangle invariants and the topology of symplectic four-manifolds*, *Duke Math. J.* 121 (2004) 1–34 MR2031164
- [15] **J A Rasmussen**, *Floer homology of surgeries on two-bridge knots*, *Algebr. Geom. Topol.* 2 (2002) 757–789 MR1928176
- [16] **J A Rasmussen**, *Floer homology and knot complements*, PhD thesis, Harvard University (2003) arXiv:math.GT/0306378
- [17] **J Robbin, D Salamon**, *The Maslov index for paths*, *Topology* 32 (1993) 827–844 MR1241874
- [18] **H Schubert**, *Knoten mit zwei Brücken*, *Math. Z.* 65 (1956) 133–170 MR0082104
- [19] **V Turaev**, *Torsion invariants of Spin^c -structures on 3-manifolds*, *Math. Res. Lett.* 4 (1997) 679–695 MR1484699
- [20] **M Wada**, *Twisted Alexander polynomial for finitely presentable groups*, *Topology* 33 (1994) 241–256 MR1273784

Evans Hall, University of California, Berkeley
Berkeley, CA 94720, USA

egrigsby@math.berkeley.edu

Received: 9 September 2005