

A note on spaces of asymptotic dimension one

KOJI FUJIWARA

KEVIN WHYTE

Let X be a geodesic metric space with $H_1(X)$ uniformly generated. If X has asymptotic dimension one then X is quasi-isometric to an unbounded tree. As a corollary, we show that the asymptotic dimension of the curve graph of a compact, oriented surface with genus $g \geq 2$ and one boundary component is at least two.

51F99; 20F69, 54F45, 57M50

1 Introduction

Our main results show that, under fairly weak restrictions, a geodesic metric space of asymptotic dimension one is quasi-isometric to a tree. Before stating precise results we review some definitions.

If X is a set and $X = \cup_i O_i$ a covering, we say that the *multiplicity* of the covering is at most n if any point $x \in X$ is contained in at most n elements of $\{O_i\}_i$.

Recall that the *covering dimension* of a topological space X is the minimal d such that every open covering has a refinement with multiplicity at most $d + 1$. The notion of asymptotic dimension is a coarse analogue introduced by Gromov in [8]:

Let X be a metric space, and $X = \cup_i O_i$ a covering. For $D \geq 0$, we say that the *D–multiplicity* of the covering is at most n if for any $x \in X$, the closed D –ball centered at x intersects at most n elements of $\{O_i\}_i$. The multiplicity is exactly the 0–multiplicity.

The *asymptotic dimension* of the metric space X is at most n if for any $D \geq 0$, there exists a covering $X = \cup_i O_i$ such that the diameter of O_i is uniformly bounded (that is, there exists C such that for all i , $\text{diam } O_i \leq C$), and the D –multiplicity of the covering is at most $n + 1$. We say that the asymptotic dimension of X , $\text{asdim } X$, is n if the asymptotic dimension of X is at most n , but it is not at most $n - 1$. If such n does not exist, then we define the asymptotic dimension of X to be infinite.

It is easy to see that if two metric spaces are quasi-isometric, then they have the same asymptotic dimension. A geodesic metric space has asymptotic dimension zero if and

only if it is bounded. Since unbounded trees have asymptotic dimension one, all spaces quasi-isometric to unbounded trees also have asymptotic dimension one [8].

Let X be a geodesic metric space. We say $H_1(X)$ is *uniformly generated* if there is an $L > 0$ so that $H_1(X)$ is generated by loops of length at most L .

Theorem 1.1 *Let X be a geodesic metric space with $H_1(X)$ uniformly generated. If X has asymptotic dimension one then X is quasi-isometric to an unbounded tree.*

Remark This theorem relies on a result of Manning [11] characterizing spaces quasi-isometric to trees, see Theorem 2.2. The restriction on $H_1(X)$ is essential, see Example 2.4.

Let G be a finitely generated group, and Γ the Cayley graph with respect to a finite generating set. Since the asymptotic dimension is invariant by quasi-isometry, the asymptotic dimension of G , $\text{asdim } G$, is defined to be the asymptotic dimension of Γ .

Corollary 1.2 *Let G be a finitely presented group. The asymptotic dimension of G is one if and only if G contains a non-trivial free group as a subgroup of finite index.*

The assumption that G is finitely presented is necessary. The following example is suggested by D Osin to us. We are informed by Dranishnikov that Corollary 1.2 has been known in work of Januszkiewicz and Swiatkowski [10] (see also Dranishnikov [4]) and that this example is also in Gentimis [7].

Example 1.3 Let $G = A \wr \mathbb{Z}$ be the wreath product such that A is a non-trivial finite group. Then, $\text{asdim } G = 1$ but G does not contain a free group as a subgroup of finite index. G is finitely generated, but not finitely presented.

By definition, G is the semidirect product of $A^{\mathbb{Z}}$ and \mathbb{Z} , $A^{\mathbb{Z}} \rtimes \mathbb{Z}$, such that the action of \mathbb{Z} is the obvious action of shifting the indexes of the direct product, $N = A^{\mathbb{Z}}$, of countable copies of A . There is an exact sequence $1 \rightarrow N \rightarrow G \rightarrow \mathbb{Z} \rightarrow 0$. Note that N is a locally finite (that is, any finitely generated subgroup is finite) countable group, which is not finitely generated. Therefore G does not contain a free group as a subgroup of finite index. We want to show that $\text{asdim } G = 1$. Indeed, the notion of asymptotic dimension is extended to a countable group, and it is shown that $\text{asdim } G = \sup \text{asdim } F$, where \sup is taken over all finitely generated subgroups $F < G$, see Dranishnikov and Smith [5]. Since N is locally finite, $\text{asdim } N = 0$. The Hurewicz type formula for asymptotic dimension is also extended [5] and we get $\text{asdim } G \leq \text{asdim } N + \text{asdim } \mathbb{Z} = 1$. Also, since $\mathbb{Z} < G$, $1 \leq \text{asdim } G$. It is easy to

see that if K and L are finitely generated, then the wreath product $K \wr L$ is finitely generated. If both K, L are finitely presented, $K \wr L$ is finitely presented if and only if K is trivial or L is finite, see Baumslag [1]. Therefore, G is finitely generated, but not finitely presented.

It is natural to ask the following question:

Question 1.4 Suppose G is a finitely generated group which is torsion free. If $\text{asdim } G = 1$, then is G a free group?

Corollary 1.5 Let X be geodesic metric space which is Gromov hyperbolic. If X has asymptotic dimension one then X is quasi-isometric to an unbounded tree.

This result does not require that X be proper. One important class of examples which are not proper are curve complexes. Let $S = S_{g,p}$ be a compact, orientable surface such that g is the genus and p is the number of the connected components of the boundary of S . We assume that $3g - 4 + p > 0$. The *curve complex* of S , defined by Harvey [9], are the flag complexes with 1-skeleton the *curve graph* of S , $C(S)$. The curve graph is the graph whose vertices are isotopy classes of essential, nonperipheral, simple closed curves in S , with two distinct vertices joined by an edge if the corresponding curves can be realized by disjoint curves. We remark that the curve complex of S is quasi-isometric to the curve graph of S , so that they have same asymptotic dimension.

Masur and Minsky [12] show the remarkable result that $C(S)$ is Gromov hyperbolic.

It is known that $C(S)$ has finite asymptotic dimension (see Bell and Fujiwara [2]). Since $C(S)$ is unbounded it cannot have asymptotic dimension zero. No upper bound or non-trivial lower bound is known in general. In the case $p = 1$, Schleimer shows in [14] that $C(S)$ is one-ended, from which it follows that it is not quasi-isometric to a tree. Thus we can improve the lower bound on asymptotic dimension here:

Corollary 1.6 Let S be a compact, oriented surface with genus $g \geq 2$ and one boundary component. Let $C(S)$ be the curve graph of S . Then the asymptotic dimension of $C(S)$ is at least two.

Acknowledgements

The first author would like to thank Greg Bell for discussion. We appreciate comments and information by Alexander Dranishnikov. We are grateful to Denis Osin, who shared his ideas with us.

This work was done during the first author's visits to the University of Illinois in Chicago and the Max Planck Institute in Bonn. He would like to thank both institutions for their hospitality. The second author was partially supported by an NSF Career award (DMS-0204576).

2 Proofs

Theorem 1.1 follows from two key ingredients. The first is the classical argument that surfaces have covering dimension two (see, for instance, Munkres [13, Section 55]). The fact we need is the following:

Lemma 2.1 *Let S be a compact surface with boundary S^1 . Let A , B , and C be points on ∂S , dividing it into arcs AB , AC , and BC . If $\{O_i\}_i$ is an open cover of S with multiplicity two then there is some i for which O_i intersects all three segments AB , AC , and BC .*

The second ingredient is a characterization of spaces quasi-isometric to trees due to Manning:

Theorem 2.2 (Manning [11, Theorem 4.6]) *Let Y be a geodesic space. Suppose that there exists a constant K with the following property:*

Let a, b be points in Y , γ a geodesic from a to b , and α a path from a to b . Then the K -ball at the midpoint m of the geodesic γ has non-empty intersection with α .

Then Y is quasi-isometric to a simplicial tree.

Before we start proving Theorem 1.1, we show one lemma. This lemma, essentially contained in Block and Weinberger [3], explains the role of the uniform generation assumption:

Lemma 2.3 *Let X be a complete geodesic metric space. The following are equivalent:*

- *X has uniformly generated H_1 .*
- *X is quasi-isometric to a complete geodesic metric space, Y , with $H_1(Y) = 0$.*

(These are also equivalent to the condition $H_1^{uf}(X) = 0$ which is the form which appears in [3].)

Proof Let Y be a complete geodesic metric space with $H_1(Y) = 0$, and let $f : X \rightarrow Y$ be a (K, C) -quasi-isometry. Suppose l is any loop in X . Divide l into segments $[a_0, a_1], [a_1, a_2], \dots, [a_{n-1}, a_n]$, where $a_0 = a_n$ and the length of each segment is at most 1.

Let $b_i = f(a_i)$. We have $d(b_i, b_{i+1}) \leq K + C$. For each i choose a geodesic segment in Y connecting b_i to b_{i+1} . Call the resulting loop l' . By assumption, l' bounds, so we have $l' = \partial S$ for some surface S mapping to Y . Triangulate S so that the image of each simplex has diameter at most 1 in Y . For each vertex v in S let $\alpha(v)$ be a point of X which maps close to the image of v in Y . For each edge e of S with endpoints u and v , choose a geodesic segment, $\alpha(e)$, in X connecting $\alpha(u)$ to $\alpha(v)$.

Thus we have α mapping the one-skeleton of S to X . For each simplex σ of S we have a loop in X of length at most D (depending only on K and C). The boundary component of S is mapped to a loop l'' which is within Hausdorff distance D of l . The map α exhibits that l'' is a sum of classes of length at most D , and the condition on Hausdorff distance implies the same for $l - l''$. Thus l is a sum of loops of length at most D , proving that $H_1(X)$ is uniformly generated.

Conversely, suppose $H_1(X)$ is generated by loops of length at most D . Let A be a maximal collection of points in X with $d(a, a') \geq D$ for all $a \neq a'$. Let $R = 3D$, and let Y be the space:

$$X \cup_{a \in A} \text{cone}(B(a, R))$$

In words, Y is X with each R -ball centered at $a \in A$ coned to a point. We give Y the induced path metric where each cone line has length R . The inclusion of X is then isometric, and has coarsely dense image, and so is a quasi-isometry. By construction, any loop l in X of length at most D is contained $B(a, R)$ for some a , and hence is null homotopic in Y . Since these generate $H_1(X)$, we have $H_1(X) \rightarrow H_1(Y)$ is the zero map.

Let l be any loop in Y . By compactness of the circle, l only passes through finitely many cone points. By pushing l off of the cones whose cone points it misses, we may arrange for l to visit the interior of only finitely many cones. For each such cone $\text{cone}(B(a, R))$, let u and v be the first and last places l visits this cone. By a further homotopy we may assume l travels along the cone line from u to the cone point and then along the cone line back from the cone point to v . Choose a path p_a of length at most $2R$ in $B(a, R)$ connecting u and v . Let l' be the loop with all the trips into the cones cut out and replaced with the p_a . The disks $c(p_a)$ give a homotopy between l and l' in Y . Since $l' \subset X$ we know $[l] = [l'] = 0$ in Y . Thus $H_1(Y) = 0$ as claimed. \square

Remark The above proof goes through verbatim to show that uniform generation of $\pi_1(X)$ is equivalent to X quasi-isometric to a simply connected space.

Proof of Theorem 1.1 Using Lemma 2.3, we assume X is quasi-isometric to a geodesic space Y with $H_1(Y) = 0$. Since Y has asymptotic dimension one there is a cover O_i of Y with 1–multiplicity two, and all O_i of diameter at most some $D < \infty$. Let U_i be the open subset of Y consisting of the points within a distance of 1 of O_i . By definition this cover has multiplicity at most two, and each U_i has diameter at most $D + 2$.

We check that Manning’s condition holds in Y with $K = \frac{3}{2}(D + 2)$. Let γ , a , b , m , and α be as in the statement of Theorem 2.2. Let l be the loop $\gamma \circ \alpha^{-1}$. Since $H_1(Y) = 0$ there is a surface S with $\partial S = S^1$ and a continuous map $f : S \rightarrow Y$ with $f|_{\partial S} = l$. Applying Lemma 2.1 to the covering $f^{-1}(U_i)$ shows that there is some i such that U_i has non-trivial intersection with α and with the segments \overline{am} and \overline{mb} of γ . Since U_i has diameter at most $D + 2$ this means there are points p on α and s and t on \overline{am} and \overline{mb} which are pairwise within a distance $D + 2$. Since γ is a geodesic, one of s or t must be within $\frac{1}{2}(D + 2)$ of m , so α must pass through the K ball around m as desired. We showed that Y , therefore X as well, is quasi-isometric to a tree, which is unbounded. \square

Remark The proof only uses that X has a cover by sets of bounded diameter with L_0 –multiplicity two for L_0 large enough to generate all of $H_1(X)$. Since it follows that X is quasi-isometric to a tree, we know that there are covers by uniformly bounded sets of L –multiplicity two for all L .

Theorem 1.1 does not hold without the assumption on $H_1(X)$:

Example 2.4 Let X be a graph with vertex set \mathbb{N} and two edges connecting n and $n+1$, both of length 2^n . Thus X is a string of circles of increasing radii, each touching the next at a single point. It is clear that $H_1(X)$ is not uniformly generated and in particular that X is not quasi-isometric to a tree.

For any $D > 0$, choose n such that $2^n \gg D$. Let U be the union of the vertices $\{1, 2, \dots, n\}$ of X and all adjacent edges except for the $\frac{1}{2}D$ –ball around the vertex $n+1$. For $i > 0$ let A_i be the ball of radius D around the vertex $n+i$, let B_i and C_i be the subsets of the two edges from $n+i$ to $n+i+1$ not within $\frac{1}{2}D$ of a vertex. It is easy to see that collection $\{U, A_i, B_i, C_i\}$ is a covering of X with D –multiplicity two. By subdividing the B_i and C_i to a collection of intervals of length $2D$ covering each, one gets a covering with D –multiplicity two by sets of diameter at most 2^n . Thus X has asymptotic dimension one.

Proof of Corollary 1.2 A well known corollary of work of Stallings [15] and Dunwoody [6] is that a group quasi-isometric to a tree is virtually free. Thus Corollary 1.2 follows immediately from Theorem 1.1 since the fundamental group of the Cayley graph is generated by translates of the relators, and so is generated by loops of length at most the length of the longest relator. \square

Remark In fact, we get something a priori stronger: if $\Gamma = \langle F | R \rangle$ where R_{ab} is a finitely generated Γ -module and Γ has asymptotic dimension one then Γ is virtually free. However, since every virtually free group is finitely presented, this does not include any new examples.

Proof of Corollary 1.5 To prove Corollary 1.5, we need to check that $H_1(X)$ is uniformly generated. For any graph G we have that $H_1(G)$, and, indeed, $\pi_1(G)$, are generated by isometrically embedded loops. If we divide such a loop into three equal pieces we get a geodesic triangle whose sides come close only at the corners. If G is δ -hyperbolic, this implies any such loop can have length at most 6δ , which proves the corollary. \square

References

- [1] **G Baumslag**, *Wreath products and finitely presented groups*, Math. Z. 75 (1960/1961) 22–28 MR0120269
- [2] **G Bell, K Fujiwara**, *The asymptotic dimension of a curve graph is finite*, J. Lond. Math. Soc. (to appear) arXiv:math.GT/0509216
- [3] **J Block, S Weinberger**, *Large scale homology theories and geometry*, from: “Geometric topology (Athens, GA, 1993)”, AMS/IP Stud. Adv. Math. 2, Amer. Math. Soc., Providence, RI (1997) 522–569 MR1470747
- [4] **A N Dranishnikov**, *Cohomological approach to asymptotic dimension* arXiv:math.MG/0608215
- [5] **A Dranishnikov, J Smith**, *Asymptotic dimension of discrete groups*, Fund. Math. 189 (2006) 27–34 MR2213160
- [6] **M J Dunwoody**, *The accessibility of finitely presented groups*, Invent. Math. 81 (1985) 449–457
- [7] **A Gentimis**, *Asymptotic dimension of finitely presented group* arXiv:math.AT/0610541
- [8] **M Gromov**, *Asymptotic invariants of infinite groups*, from: “Geometric group theory, Vol. 2 (Sussex, 1991)”, London Math. Soc. Lecture Note Ser. 182, Cambridge Univ. Press, Cambridge (1993) 1–295 MR1253544

- [9] **W J Harvey**, *Boundary structure of the modular group*, from: “Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)”, Ann. of Math. Stud. 97, Princeton Univ. Press, Princeton, N.J. (1981) 245–251 MR624817
- [10] **T Januszkiewicz, J Swiatkowski**, *Filling invariants of systolic complexes and groups*, Geom. Topol. 11 (2007) 727–758
- [11] **J F Manning**, *Geometry of pseudocharacters*, Geom. Topol. 9 (2005) 1147–1185 MR2174263
- [12] **H A Masur, Y N Minsky**, *Geometry of the complex of curves. I. Hyperbolicity*, Invent. Math. 138 (1999) 103–149 MR1714338
- [13] **J R Munkres**, *Topology*, second edition, Prentice Hall (2000)
- [14] **S Schleimer**, *The end of the curve complex* arXiv:math.GT/0608505
- [15] **J R Stallings**, *On torsion-free groups with infinitely many ends*, Ann. of Math. (2) 88 (1968) 312–334 MR0228573

Graduate School of Information Sciences, Tohoku University
Sendai 980-8579, Japan

Department of Mathematics, University of Illinois at Chicago
Chicago IL 60607-7045, USA

fujiwara@math.is.tohoku.ac.jp, kwhyte@math.uic.edu

Received: 30 November 2006 Revised: 1 May 2007