

## Amalgamations of Heegaard splittings in 3–manifolds without some essential surfaces

GUOQIU YANG  
FENGCHUN LEI

Let  $M$  be a compact, orientable,  $\partial$ –irreducible 3–manifold and  $F$  be a connected closed essential surface in  $M$  with  $g(F) \geq 1$  which cuts  $M$  into  $M_1$  and  $M_2$ . In the present paper, we show the following theorem: Suppose that there is no essential surface with boundary  $(Q_i, \partial Q_i)$  in  $(M_i, F)$  satisfying  $\chi(Q_i) > 2 + g(F) - 2g(M_i)$ ,  $i = 1, 2$ . Then  $g(M) = g(M_1) + g(M_2) - g(F)$ . As a consequence, we further show that if  $M_i$  has a Heegaard splitting  $V_i \cup_{S_i} W_i$  with distance  $D(S_i) \geq 2g(M_i) - g(F)$ ,  $i = 1, 2$ , then  $g(M) = g(M_1) + g(M_2) - g(F)$ .

The main results follow from a new technique which is a stronger version of Schultens' Lemma.

57M99, 57N10; 57M27

### 1 Introduction

Let  $M$  be a compact, orientable,  $\partial$ –irreducible 3–manifold and  $F$  be a connected closed essential surface in  $M$  with  $g(F) \geq 1$  which cuts  $M$  into  $M_1$  and  $M_2$ . Suppose  $V_i \cup_{S_i} W_i$  is a Heegaard splitting of  $M_i$ ,  $i = 1, 2$ . Then  $V_1 \cup_{S_1} W_1$  and  $V_2 \cup_{S_2} W_2$  induce a natural Heegaard splitting  $V \cup_S W$  of  $M$ , which is called the amalgamation of  $V_1 \cup_{S_1} W_1$  and  $V_2 \cup_{S_2} W_2$  along  $F$ , and  $g(S) = g(S_1) + g(S_2) - g(F)$ ; see Schultens [15]. Thus  $g(M) \leq g(M_1) + g(M_2) - g(F)$ .

There exist examples which show that an amalgamation of two minimal genus Heegaard splittings of  $M_1$  and  $M_2$  may be stabilized (refer to Bachman, Schleimer and Sedgwick [1], Kobayashi, Qiu, Rieck and Wang [7], Schultens and Weidmann [17] and others). On the other hand, it has been shown that under some conditions on the manifolds, or the gluing maps, the equality  $g(M) = g(M_1) + g(M_2) - g(F)$  holds; see Kobayashi and Qiu [6], Lackenby [8], Lei and Yang [9], Li [10], Souto [18], Yang and Lei [19] and others.

In the present paper, we show the following result:

**Theorem 4.2** Suppose that there is no essential surface with boundary  $(Q_i, \partial Q_i)$  in  $(M_i, F)$  satisfying  $\chi(Q_i) > 2 + g(F) - 2g(M_i)$ ,  $i = 1, 2$ . Then  $g(M) = g(M_1) + g(M_2) - g(F)$ .

As a consequence, we further show:

**Theorem 4.3** If  $M_i$  has a Heegaard splitting  $V_i \cup_{S_i} W_i$  with distance  $D(S_i) \geq 2g(M_i) - g(F)$ ,  $i = 1, 2$ , then  $g(M) = g(M_1) + g(M_2) - g(F)$ .

The paper is organized as follows. In Section 2, we review some preliminaries and notation which will be used later. In Section 3, we will prove two technical lemmas. The statements and proofs of the main results are given in Section 4. The lemmas in Section 3 play an important role in our proof of Theorem 4.2.

## 2 Preliminaries

The concepts and terminologies not defined in the paper are standard; for example, see, Hempel [3] and Jaco [5].

Suppose  $F$  is a subsurface of  $\partial M$  or a surface properly embedded in a 3-manifold  $M$ . If there is a disk  $D \subset M$  such that  $D \cap F = \partial D$  and  $\partial D$  is an essential loop in  $F$ , then we say that  $F$  is *compressible* in  $M$ . Such a disk  $D$  is called a *compressing disk*. We say that  $F$  is *incompressible* in  $M$  if  $F$  is not compressible in  $M$ . If  $\partial M$  is incompressible, then  $M$  is said to be  *$\partial$ -irreducible*. If  $F$  is an incompressible surface in  $M$  and not parallel to a subsurface of  $\partial M$ , then  $F$  is an *essential surface* in  $M$ .

A 3-manifold  $C$  is called a *compression body* if there exists a connected closed orientable surface  $S$  such that  $C$  is obtained from  $S \times I$  by attaching 2-handles along mutually disjoint loops in  $S \times \{0\} \subset S \times I$  and capping off any resulting 2-sphere boundary components with 3-handles. We denote  $S \times \{1\}$  by  $\partial_+ C$  and  $\partial C - \partial_+ C$  by  $\partial_- C$ . An essential disk in  $C$  is a compressing disk of  $\partial_+ C$  in  $C$ .

A *Heegaard splitting* of a 3-manifold  $M$  is a triplet  $(C_1, C_2; S)$ , where  $C_1$  and  $C_2$  are compression bodies with  $C_1 \cup C_2 = M$  and  $C_1 \cap C_2 = \partial_+ C_1 = \partial_+ C_2 = S$ . The surface  $S$  is called a *Heegaard surface* and the genus of a Heegaard splitting is defined by the genus of the Heegaard surface. We use  $g(M)$  to denote the *Heegaard genus* of  $M$ , which is equal to the minimal genus of all Heegaard splittings of  $M$ . A Heegaard splitting  $C_1 \cup_S C_2$  for  $M$  is *minimal* if  $g(S) = g(M)$ . A Heegaard splitting  $C_1 \cup_S C_2$  is *trivial* if  $\partial_- C_1 \cong \partial_+ C_1$  or  $\partial_- C_2 \cong \partial_+ C_2$ .

Let  $C_1 \cup_S C_2$  be a Heegaard splitting for  $M$ .  $C_1 \cup_S C_2$  is said to be *reducible* (or *weakly reducible*) if there are essential disks  $D_1 \subset C_1$  and  $D_2 \subset C_2$  with  $\partial D_1 = \partial D_2$  (or  $\partial D_1 \cap \partial D_2 = \emptyset$ ). The splitting  $C_1 \cup_S C_2$  is said to be *irreducible* if it is not reducible; and the splitting  $C_1 \cup_S C_2$  is said to be *strongly irreducible* if it is not weakly reducible.

Scharlemann and Thompson showed in [13] that any irreducible and  $\partial$ -irreducible Heegaard splitting  $C_1 \cup_S C_2$  can be broken up into a series of strongly irreducible Heegaard splittings. That is, we can begin with the handle structure determined by  $C_1 \cup_S C_2$  and rearrange the order of adding the 1- and 2-handles, so that ultimately

$$M = (C_1^1 \cup_{S_1} C_2^1) \cup_{F_1} (C_1^2 \cup_{S_2} C_2^2) \cup_{F_2} \cdots \cup_{F_{m-1}} (C_1^m \cup_{S_m} C_2^m),$$

such that each  $C_1^i \cup_{S_i} C_2^i$  is a strongly irreducible Heegaard splitting with intersections  $\partial_- C_1^i \cap \partial_- C_1^{i+1} = F_i$ ,  $1 \leq i \leq m-1$  and  $\partial_- C_1^1 = \partial_- C_1$ ,  $\partial_- C_2^m = \partial_- C_2$ . For each  $i$ , each component of  $F_i$  is a closed incompressible surface of positive genus, and only one component of  $M_i = C_1^i \cup_{S_i} C_2^i$  is not a product. None of the compression bodies  $C_1^i, C_2^i$ ,  $1 \leq i \leq m$  is trivial. Such a rearrangement of handles is called an *untelescoping* of the Heegaard splitting  $C_1 \cup_S C_2$ . Then it is easy to see  $\chi(S) \leq \chi(S_i), \chi(F_i)$  for each  $i$ , and when  $m \geq 2$ ,  $\chi(S) < \chi(S_i), \chi(F_i)$  for each  $i$ .

Let  $C_1 \cup_S C_2$  be a Heegaard splitting,  $\alpha$  and  $\beta$  two essential simple closed curves in  $S$ . The *distance*  $d(\alpha, \beta)$  of  $\alpha$  and  $\beta$  is the smallest integer  $n \geq 0$  such that there is a sequence of essential simple closed curves  $\alpha = \alpha_0, \alpha_1, \dots, \alpha_n = \beta$  in  $S$  with  $\alpha_{i-1} \cap \alpha_i = \emptyset$ , for  $1 \leq i \leq n$ . The *distance* of the Heegaard splitting  $C_1 \cup_S C_2$  is defined to be  $\min \{d(\alpha, \beta)\}$ , where  $\alpha$  bounds an essential disk in  $C_1$  and  $\beta$  bounds an essential disk in  $C_2$ , and is denoted by  $D(S)$ .

The concept of Heegaard distance of a Heegaard splitting was first defined by Hempel [4]. It is clear that  $C_1 \cup_S C_2$  is reducible if and only if  $D(S) = 0$ ,  $C_1 \cup_S C_2$  is weakly reducible if and only if  $D(S) = 1$ . Its relations to the genus of the Heegaard splitting have been discussed by Hartshorn [2], Hempel [4], Scharlemann and Tomova [14] and others.

The following Lemmas are some well-known basic facts and results:

**Lemma 2.1** [12] *Suppose  $(Q, \partial Q) \subset (M, \partial M)$  is an essential surface and  $Q'$  is the result of  $\partial$ -compressing  $Q$ . Then  $Q'$  is essential.*

**Lemma 2.2** [16] *Let  $V$  be a compression body and  $F$  be a properly embedded incompressible surface in  $V$  with  $\partial F \subset \partial_+ V$ , then each component of  $V \setminus F$  is a compression body.*

**Lemma 2.3** [11] *Let  $M = V \cup_S W$  be a strongly irreducible Heegaard splitting. If  $\alpha$  is an essential simple loop in  $S$  which bounds a disk  $D$  in  $M$  such that  $D$  is transverse to  $S$ , then  $\alpha$  bounds an essential disk in  $V$  or  $W$ .*

**Lemma 2.4** [2] *Let  $V \cup_S W$  be a Heegaard splitting of  $M$  and  $F$  be a properly embedded incompressible surface (maybe not connected) in  $M$ . Then any component of  $F$  is parallel to  $\partial M$  or  $D(S) \leq 2 - \chi(F)$ .*

In the rest of this paper, we use  $M \setminus N$  to denote the manifold obtained by cutting  $M$  along  $M \cap N$  and  $N(F, M)$  the compact regular neighborhood of the submanifold  $F$  in the manifold  $M$ .

### 3 Two technical lemmas

**Definition 3.1** Let  $F$  be a 2-sided surface properly embedded in  $M$ ,  $F \times [0, 1]$  a regular neighborhood of  $F$  in  $M$  with  $F \times \{\frac{1}{2}\} = F$ . If there are two compressing disks  $D$  and  $E$  of  $F$  such that  $D \cap (F \times \{0\}) = \emptyset$  and  $E \cap (F \times \{1\}) = \emptyset$ , then  $F$  is called *bicompressible* and  $(D, E)$  is called a *bicompressing disk pair* of  $F$ . If  $F$  is bicompressible and any bicompressing disk pair  $(D, E)$  satisfies  $\partial D \cap \partial E \neq \emptyset$ , then  $F$  is called *strongly irreducible*.

**Definition 3.2** Two surfaces  $F_1$  and  $F_2$  embedded in a 3-manifold are almost transverse if they have exactly one nontransverse intersection point, and it is a saddle point.

The following Lemma 3.3 is a stronger version of Schultens's lemma [16] as well as Lemma 3.3 of Bachman, Schleimer and Sedgwick [1]. Lemma 3.3 is essential in our proof of Theorem 4.2.

**Lemma 3.3** *Let  $M = V \cup_S W$  be a strongly irreducible Heegaard splitting and  $F$  be a 2-sided essential surface (not a disk or 2-sphere) in  $M$ . Then  $F$  can be isotoped such that at least one of the following conclusions holds:*

- (1)  *$F$  is transverse to  $S$  and any component of  $S \setminus F$  is incompressible in the respective submanifold of  $M \setminus F$  except for exactly one strongly irreducible component;*
- (2)  *$F$  is almost transverse to  $S$  and any component of  $S \setminus N(F, M)$  is incompressible in the respective submanifold of  $M \setminus N(F, M)$ .*

**Proof** From Schultens's lemma in [16], we may assume that each component of  $S \cap F$  is an essential loop in both  $F$  and  $S$ , and  $|S \cap F|$  is minimal. If  $S \setminus F$  is bicompressible in  $M \setminus F$ , then (1) is true. In the following arguments, we suppose (1) is not true. So  $S \setminus F$  is incompressible in  $V$  or  $W$ , say  $V$ . Then  $F$  satisfies the following three conditions: (i) each component of  $S \cap F$  is an essential loop in both  $F$  and  $S$ ; (ii)  $S \setminus F$  is incompressible in  $V$ ; (iii) any component of  $F \cap V$  is essential in  $V$ .

We can take  $F$  such that  $-\chi(F \cap V)$  is minimal among all surfaces isotopic to  $F$  which satisfies conditions: (i), (ii) and (iii). If  $F \cap V$  is not boundary compressible, then any component of  $F \cap V$  is spanning annulus by the properties of compression body. This means that  $S \setminus F$  is compressible in  $V$ , a contradiction. So  $F \cap V$  is boundary compressible.

We claim that there exists a boundary compressing disk  $\Delta$  of  $F \cap V$  in  $V$  such that any component of the result of  $\partial$ -compressing  $F \cap V$  along  $\Delta$  is essential in  $V$ .

Let  $\Delta_1$  be a boundary compressing disk of  $F \cap V$  in  $V$ ,  $\alpha_1$  the corresponding essential arc in  $F \cap V$  and  $\beta_1$  the corresponding essential arc in  $\partial V$  with  $\alpha_1 \cup \beta_1 = \partial \Delta_1$ . Denote the component of  $F \cap V$  which contains  $\alpha_1$  by  $P$ . Obviously,  $\chi(P) \leq 0$ .

If  $\chi(P) = 0$ , then  $P$  is an essential annulus in  $V$ . By performing  $\partial$ -compression to  $P$  along  $\Delta_1$ , we get an essential disk  $E$  with  $E \cap F = \emptyset$  in  $V$ . This means that  $S \setminus F$  is compressible in  $V$ , a contradiction.

So  $\chi(P) \leq -1$ . Denoted the result of  $\partial$ -compressing  $P$  along  $\Delta_1$  by  $P'$ . If any component of  $P'$  is essential in  $V$ , we take  $\Delta = \Delta_1$ ,  $\alpha = \alpha_1$  and  $\beta = \partial \Delta \setminus \alpha$ . If one component  $P^*$  of  $P'$  is parallel to a subsurface  $Q$  of  $\partial V$  in  $V$  with  $\partial Q = \partial P^*$ , then  $\alpha_1$  is separating in  $P$  and the other component of  $P'$  is essential in  $V$  by Lemma 2.1. Since  $\partial \beta_1 \cap \partial Q = \emptyset$ ,  $\beta_1 \subset Q$  or  $\beta_1 \cap Q = \emptyset$ . If  $\beta_1 \subset Q$ , then  $P$  is compressible in  $V$ , a contradiction. Hence  $\beta_1 \cap Q = \emptyset$  and this means that there exists a nonseparating essential arc  $\alpha$  of both  $P$  and  $P'$  and an essential arc  $\beta \subset \partial V$  which satisfies  $\alpha \cap \beta = \partial \alpha = \partial \beta$  and  $\alpha \cup \beta$  bounding a disk  $\Delta$  with  $\Delta \cap (F \cap V) = \alpha$ . So  $\Delta$  is a boundary compressing disk of  $F \cap V$  in  $V$ . By Lemma 2.1, any component of the result of  $\partial$ -compressing  $F \cap V$  along  $\Delta$  is essential in  $V$ .

Perform  $\partial$ -compression to  $F \cap V$  along  $\Delta$  to get  $F^*$ , which is an isotopy of  $F$ . Then  $F^*$  satisfies conditions (i) and (iii). Obviously,  $-\chi(F^* \cap V) < -\chi(F \cap V)$ . Since  $-\chi(F \cap V)$  is minimal among all surfaces which are isotopic to  $F$  and satisfy conditions: (i), (ii) and (iii),  $S \setminus F^*$  is compressible in  $V$ . If  $S \setminus F^*$  is compressible in  $W$ , then  $S \setminus F$  is bicompressible in  $M \setminus F$ , ie (1) is true, a contradiction. Hence,  $S \setminus F^*$  is incompressible in  $W$ . From  $S \setminus F$  to  $S \setminus F^*$ , only cut the band  $B_1 = N(\beta, S \setminus F)$  from  $S \setminus F$  along one pair opposite edges of  $B_1$  and paste it to  $S \setminus F$  along the other

pair opposite edges of  $B_1$ . Then only the component of  $S \setminus F^*$  which contains  $\beta$  is compressible in  $V$  and any component of  $S \setminus F^*$  is incompressible in  $W$ . Furthermore, only the component of  $S \setminus F^*$  which contains  $\beta$  is compressible in  $M \setminus F^*$  and the band  $B_1$  intersects with the boundary of any compressing disk  $D$  of  $S \setminus F^*$ ; see Figure 1 below. Otherwise, there is a compressing disk  $D_1$  for some component of  $S \setminus F^*$  with  $\partial D_1 \cap B_1 = \emptyset$ , by Lemma 2.3,  $\partial D_1$  must bound a compressing disk of  $S \setminus F$  in  $V$ , a contradiction. Hence, at most one of  $S \setminus F$  is compressible in  $W$ . If one component of  $S \setminus F$  is compressible in  $W$ , then the component must contain  $\beta$ . Otherwise,  $S \setminus F^*$  is bicompressible, ie (1) is true, a contradiction again. So at most the component of  $S \setminus F$  which contains  $\beta$  is compressible in  $M \setminus F$  and any compressing disk  $E$  of  $S \setminus F$  satisfies  $\partial E \cap B_1 \neq \emptyset$ ; see Figure 1.




Figure 1:  $\partial E \cap B_1$  and  $\partial D \cap B_1$

Push  $F^*$  slightly off both  $F$  and  $F^*$  to get  $F'$ , which is a parallel copy of  $F^*$  such that the disk  $\Delta$  lies in the parallelism  $N$  bounded by  $F$  and  $F'$ ; see Figure 2.



Figure 2:  $F$  and  $F^*$

Then  $B_1 \subset N$  and any component of  $S \setminus N$  is incompressible in the respective submanifold of  $M \setminus N$ ; see Figure 3.




Figure 3:  $F, F'$  and  $N$

So  $F$  can be isotoped into  $F_1$  which is almost transverse to  $S$  with a saddle point belong to both  $\alpha$  and  $\beta$  with  $N(F_1, M) = N$ ; see Figure 4.




Figure 4:  $F \cap S$  and  $F_1 \cap S$

Thus (2) is true.

This completes the proof. □

**Lemma 3.4** *Let  $N$  be a compact orientable 3-manifold and not a compression body,  $F$  a component of  $\partial N$ . Suppose  $Q$  is a properly embedded connected separating surface in  $N$  with  $\partial Q \subset F$  and any component of  $\partial Q$  is essential in  $F$ , and  $Q$  cuts  $N$  into two compression bodies  $N_1$  and  $N_2$  with  $Q = \partial_+ N_1 \cap \partial_+ N_2$ . If  $Q$  can be compressed to  $Q_i^*$  in some  $N_i$  such that any component of  $Q_i^*$  is parallel to a subsurface of  $\partial N$ , then  $g(N) \leq 1 - \frac{1}{2}\chi(F \cap N_{3-i}) - \frac{1}{2}\chi(Q)$ .*

**Proof** By assumption,  $Q^*$  and  $F \setminus (Q \cap F)$  have no disk components, and each component of  $Q^*$  is parallel into either a subsurface of  $F$  or a component of  $\partial N$ .

We may assume that  $Q$  is compressed to  $Q^*$  in  $N_1$  by cutting  $Q$  open along a collection  $\mathcal{D} = \{D_1, \dots, D_n\}$  of pairwise disjoint compressing disks in  $N_1$ .  $N'_1 = N_1 - \eta(\mathcal{D})$  where  $\eta(\mathcal{D})$  is the open regular neighborhood of  $\mathcal{D}$ . Let  $A_1, \dots, A_r$  and  $A_{r+1}, \dots, A_{r+s}$  be all the components of  $Q^*$ , where  $A_i$  is a component with  $\partial A_i \subset F$  for  $1 \leq i \leq r$  and  $A_{r+i}$  is a closed surface for  $1 \leq i \leq s$ . Suppose that each  $A_i$  is parallel to a subsurface  $A'_i$  of  $\partial N$ , where  $i = 1, 2, \dots, r + s$ . We divide it into two cases to discuss:

If there exist two components  $A'_{i_0}$  and  $A'_{j_0}$  with  $A'_{i_0} \subset A'_{j_0}$ . Then we set  $\mathcal{A}_1 = \{A_i : A'_i \subset A'_{j_0}, 1 \leq i \leq r+s, i \neq j_0\}$  and  $\mathcal{A}_2 = \{A_i : A'_i \cap A'_{j_0} = \emptyset, 1 \leq i \leq r+s\}$ . We claim that  $\mathcal{A}_2 = \emptyset$ . Otherwise, since  $Q$  is connected, there must exist  $A_{i_1} \in \mathcal{A}_1$ ,  $A_{i_2} \in \mathcal{A}_2$ , and  $D_{p_1}, D_{p_2} \in \mathcal{D}$  such that in the compression, the two copies (obtained by compressing  $Q$  along  $\mathcal{D}$ ) of  $D_{p_k}$  lie in  $A_{i_k}$  and  $A_{j_0}$  respectively,  $k = 1, 2$ . But this contradicts the assumption that  $Q$  is separating in  $N$ . Thus  $\mathcal{A}_2 = \emptyset$ . Let  $N''_1$  be the component of  $N'_1$  which contain  $A_{j_0}$ , then  $A_{j_0} \subset \partial_+ N''_1$ . Since  $A_{j_0}$  is parallel to  $A'_{j_0}$ ,  $N''_1$  is a handlebody, the other components of  $N'_1$  are  $A_{r+i} \times I$ .  $N$  is homeomorphic to  $N_2 \cup N(\mathcal{D}) \cup N'_1$ . So  $N$  is a compression body with  $F = \partial_+ N$ , a contradiction.

So for any two components  $A_i, A_j$  of  $Q^*$ ,  $A'_i \cap A'_j = \emptyset$ . Then  $N'_1 = \bigcup_{i=1}^{r+s} A_i \times I$ , where  $A_i \times \{0\} = A'_i$  and  $A_i \times \{1\} = A_i$ ,  $i = 1, 2, \dots, r+s$ . Let  $B_1, \dots, B_t$  be the components of  $F \setminus \bigcup_{i=1}^r A'_i$ . Take a small regular neighborhood  $B_i \times I$  of  $B_i$  in  $N_2$ , where  $B_i \times \{0\} = B_i$  and  $i = 1, 2, \dots, t$ . Set  $C_1 = N_1 \cup \bigcup_{i=1}^t B_i \times I$  and  $C_2 = N \setminus C_1$ . Then  $C_1$  is obtained from  $F \times I$  and  $\bigcup_{i=1}^s A_{r+i} \times I$  by adding 1–handles whose cores are  $\mathcal{D}$ , so  $C_1$  is a compression body. Note that  $C_2 = N_2 \setminus \bigcup_{i=1}^t B_i \times I \cong N_2$ ,  $C_2$  is a compression body. Let  $S = \partial_+ C_1$ , then by assumption  $Q \subset \partial_+ N_2$  and  $S = \partial_+ C_2$ . Thus,  $S$  is a Heegaard surface of  $N$ . Now  $2 - \chi(S) = 2 - \chi(F \cap N_2) - \chi(Q)$ , so we have  $g(N) \leq g(S) = \frac{1}{2}(2 - \chi(S)) \leq 1 - \frac{1}{2}\chi(F \cap N_2) - \frac{1}{2}\chi(Q)$ , as required.  $\square$

## 4 The essential surfaces and amalgamations

**Proposition 4.1** *Let  $M$  be a compact, orientable 3–manifold and  $F$  be an essential closed surface which cuts  $M$  into  $M_1$  and  $M_2$ . Suppose that there is no essential surface with boundary  $(Q_i, \partial Q_i)$  in  $(M_i, F)$  satisfying  $\chi(Q_i) > 2 + g(F) - 2g(M_i)$ ,  $i = 1, 2$ . Then for any closed incompressible surface  $F^*$  in  $M$  with  $g(F^*) < g(M_1) + g(M_2) - g(F) - 1$ , we can isotope  $F$  in  $M$  such that after isotopy,  $F \cap F^* = \emptyset$ .*

**Proof** Since  $F$  and  $F^*$  are both incompressible, we can isotope  $F$  in  $M$  such that each component of  $F \cap F^*$  is essential in both  $F$  and  $F^*$ , and  $|F \cap F^*|$  is minimal.

If  $|F \cap F^*| > 0$ , by the minimality of  $|F \cap F^*|$ , each component of  $F^* \cap M_i$  is essential in  $M_i$ ,  $i = 1, 2$ . By assumption, we have  $\chi(F^* \cap M_1) \leq 2 + g(F) - 2g(M_1)$  and  $\chi(F^* \cap M_2) \leq 2 + g(F) - 2g(M_2)$ . So

$$\chi(F^*) = \chi(F^* \cap M_1) + \chi(F^* \cap M_2) \leq 4 + 2g(F) - 2g(M_1) - 2g(M_2).$$

Then  $g(F^*) = \frac{1}{2}(2 - \chi(F^*)) \geq g(M_1) + g(M_2) - g(F) - 1$ , a contradiction to the assumption.

Thus  $F \cap F^* = \emptyset$ .  $\square$

Now we come to the main result of the paper.

**Theorem 4.2** *Let  $M$  be a compact, orientable,  $\partial$ -irreducible 3-manifold and  $F$  be an essential closed surface which cuts  $M$  into  $M_1$  and  $M_2$ . Suppose that there is no essential surface with boundary  $(Q_i, \partial Q_i)$  in  $(M_i, F)$  satisfying  $\chi(Q_i) > 2 + g(F) - 2g(M_i)$ ,  $i = 1, 2$ . Then  $g(M) = g(M_1) + g(M_2) - g(F)$ .*

**Proof** We may assume that  $M$  is irreducible. If  $M$  is reducible, we have that at least one of  $M_1$  and  $M_2$ , say  $M_i$ , is reducible. By the Prime Decomposing Theorem of 3-manifold, we can put  $M_i = M' \# M'_1$  such that  $F \subset \partial M'_1$  and  $M'_1$  is irreducible. By Haken's Lemma,  $g(M'_1) = g(M_i) - g(M')$  and  $g(M'_1 \cup_F M_2) = g(M) - g(M')$ . Since any essential surface with boundary  $(Q_1, \partial Q_1)$  in  $(M'_1, F)$  is an essential surface with boundary  $(Q_1, \partial Q_1)$  in  $(M_i, F)$  and by our assumption,  $\chi(Q_1) \leq 2 + g(F) - 2g(M_i) \leq 2 + g(F) - 2g(M'_1)$ . Then the conclusion follows immediately from the irreducible case.

So  $M$  is irreducible and  $\partial$ -irreducible.

It is easy to see that  $g(M) \leq g(M_1) + g(M_2) - g(F)$ . Suppose that  $g(M) = g(M_1) + g(M_2) - g(F)$  does not hold. Thus there exists a minimal Heegaard splitting  $V \cup_S W$  of  $M$  with

$$(1) \quad g(S) \leq g(M_1) + g(M_2) - g(F) - 1.$$

We divide it into the following two cases to discuss.

**Case 1** The Heegaard splitting  $V \cup_S W$  is strongly irreducible.

In this case, by Lemma 3.3, there are only the following two subcases.

**Subcase 1**  $F$  is transverse to  $S$  and any component of  $S \setminus F$  is incompressible in the respective submanifold of  $M \setminus F$  except for exactly one strongly irreducible component.

If one incompressible component of  $S \cap M_1$  and  $S \cap M_2$  is inessential. Then we can isotope  $S$  to  $S'$  such that  $|S \cap F| > |S' \cap F|$ , one component of  $S' \cap M_1$  and  $S' \cap M_2$  is bicompressible and any other component is incompressible. So we may assume one of  $S \cap M_1$ , say  $P$  is bicompressible and any incompressible component of  $S \cap M_1$  and  $S \cap M_2$  is essential in  $M_1$  or  $M_2$  without loss generality.

Since  $F$  is essential in  $M$  and there is no closed essential surface in a compression body,  $F \cap S \neq \emptyset$ . Then  $\chi(P) \leq -2$ . If otherwise,  $\chi(P) \geq -1$ ,  $P$  is either a disk, an annulus, twice-punctured disk, or a once-punctured torus, in each case we can conclude that a component of  $\partial P$  bounds a disk in  $M_1$ , therefore  $F$  is compressible in  $M_1$ , a contradiction. By assumption,  $\chi(S \cap M_2) \leq 2 + g(F) - 2g(M_2)$ .

If  $S \cap M_1 \neq P$ , then at least one component of  $S \cap M_1$  is essential in  $M_1$ , by assumption  $\chi(S \cap M_1) \leq 2 + g(F) - 2g(M_1) + \chi(P) \leq g(F) - 2g(M_1)$ . Thus

$$\begin{aligned}\chi(S) &= \chi(S \cap M_1) + \chi(S \cap M_2) \\ &\leq g(F) - 2g(M_1) + 2 + g(F) - 2g(M_2) \\ &\leq 2 - 2(g(M_1) + g(M_2) - g(F))\end{aligned}$$

and  $g(S) \geq g(M_1) + g(M_2) - g(F)$ , a contradiction.

So  $S \cap M_1 = P$ . By Lemma 2.2,  $P$  divides  $M_1$  into two compression bodies  $V \cap M_1$  and  $W \cap M_1$ . Since  $P$  is bicompressible in  $M_1$ , we compress  $P$  into  $P_V$  ( $P_W$ , resp.) in  $V \cap M_1$  ( $W \cap M_1$ , resp.) as possible as. Since  $V \cup_S W$  is strongly irreducible and by Lemma 2.3, any component of  $P_V$  and  $P_W$  is incompressible in  $M_1$ . If one component of  $P_V$  or  $P_W$ , say  $P_V$ , is essential in  $M_1$ , then by assumption,  $\chi(S \cap M_1) \leq \chi(P_V) - 2 \leq 2 + g(F) - 2g(M_1) - 2 \leq g(F) - 2g(M_1)$ . By the same arguments as above,  $g(S) \geq g(M_1) + g(M_2) - g(F)$ , a contradiction again. So any component of  $P_V$  or  $P_W$  is  $\partial$ -parallel in  $M_1$ . Then by Lemma 3.4,  $g(M_1) \leq 1 - \frac{1}{2}\chi(F \cap W) - \frac{1}{2}\chi(P)$  and  $g(M_1) \leq 1 - \frac{1}{2}\chi(F \cap V) - \frac{1}{2}\chi(P)$ . Note that  $\chi(F) = \chi(F \cap V) + \chi(F \cap W)$ , so  $2g(M_1) \leq 2 - \frac{1}{2}\chi(F) - \chi(P) = 1 + g(F) - \chi(P)$ . Then  $\chi(S \cap M_1) = \chi(P) \leq 1 + g(F) - 2g(M_1)$  and

$$\begin{aligned}\chi(S) &= \chi(S \cap M_1) + \chi(S \cap M_2) \\ &\leq 1 + g(F) - 2g(M_1) + 2 + g(F) - 2g(M_2) \\ &\leq 3 - 2(g(M_1) + g(M_2) - g(F)),\end{aligned}$$

so  $g(S) \geq g(M_1) + g(M_2) - g(F) - \frac{1}{2}$ , a contradiction to (1).

This finishes the proof of Subcase 1.

**Subcase 2**  $F$  is almost transverse to  $S$  and any component of  $S \setminus N(F)$  is incompressible in the respective submanifold of  $M \setminus N(F)$ .

For  $i = 1, 2$ , it is easy to see that  $S \cap M_i$  is essential in  $M_i$ , then by assumption,  $\chi(S \cap M_i) \leq 2 + g(F) - 2g(M_i)$ . Clearly,  $\chi(S \cap N(F)) = -1$ . So

$$\begin{aligned}\chi(S) &= \chi(S \cap M_1) + \chi(S \cap M_2) - 1 \\ &\leq 2 + g(F) - 2g(M_1) + 2 + g(F) - 2g(M_2) - 1 \\ &\leq 3 - 2(g(M_1) + g(M_2) - g(F)),\end{aligned}$$

hence  $g(S) \geq g(M_1) + g(M_2) - g(F) - \frac{1}{2}$ , a contradiction again.

This finishes the proof of Subcase 2.

**Case 2** The Heegaard splitting  $V \cup_S W$  is weakly reducible.

Since  $M$  is irreducible, by Haken’s lemma,  $V \cup_S W$  is irreducible. By the result of Scharlemann and Thompson [13],  $V \cup_S W$  is an amalgamation of  $n$  strongly irreducible Heegaard splitting  $V \cup_S W = (V_1 \cup_{S_1} W_1) \cup_{F_1} (V_2 \cup_{S_2} W_2) \cup_{F_2} \cdots \cup_{F_{n-1}} (V_n \cup_{S_n} W_n)$ . We may further assume that no component of  $F_i$ ,  $1 \leq i \leq n - 1$ , is  $\partial$ -parallel in  $M$ . Obviously,  $g(F_i) < g(S_i) < g(S)$  and  $g(F_i) \leq g(S) - 2$ . Then by (1) we have

$$\begin{aligned} g(F_i) &\leq g(S) - 2 \\ &\leq g(M_1) + g(M_2) - g(F) - 3 \\ &< g(M_1) + g(M_2) - g(F) - 1. \end{aligned}$$

By Proposition 4.1, we can isotope  $F$  such that  $(\cup F_i) \cap F = \emptyset$ . So  $F$  lies in the nontrivial component  $V_j^* \cup_{S_j^*} W_j^*$  of  $V_j \cup_{S_j} W_j$  for some  $1 \leq j \leq n$ .

If  $F$  is parallel to some component, say  $F^*$ , of  $\cup F_i$ , then  $F^*$  cuts  $M$  into two parts  $M_1$  and  $M_2$ .  $V \cup_S W$  is an amalgamation of two Heegaard splittings of  $M_1$  and  $M_2$  and  $g(S) \geq g(M_1) + g(M_2) - g(F)$ , a contradiction. Hence  $F$  can not be parallel to any component of  $\cup F_i$ . So  $F$  is an essential closed surface in  $V_j^* \cup_{S_j^*} W_j^*$ . Denote  $M^* = V_j^* \cup_{S_j^*} W_j^*$ . Then there are the following two subcases.

**Subcase 3**  $F$  is almost transverse to  $S_j^*$  and any component of  $S_j^* \setminus N(F)$  is incompressible in the respective submanifold of  $M^* \setminus N(F)$ ;

Then any component of  $S_j^* \setminus N(F)$  is incompressible in the respective submanifold of  $M \setminus N(F)$ . Furthermore,  $S_j^* \setminus N(F)$  is essential in  $M \setminus N(F)$  since there is no essential closed surface in a compression body. Then  $\chi(S_j^* \cap N(F)) = -1$  and by assumption,  $\chi(S_j^* \cap M_i) \leq 2 + g(F) - 2g(M_i)$  for  $i = 1, 2$ . So

$$\begin{aligned} \chi(S) &\leq \chi(S_j^*) - 2 \\ &= \chi(S_j^* \cap M_1) + \chi(S_j^* \cap M_2) + \chi(S_j^* \cap N(F)) - 2 \\ &\leq 2 + g(F) - 2g(M_1) + 2 + g(F) - 2g(M_2) - 3 \\ &\leq 1 - 2(g(M_1) + g(M_2) - g(F)). \end{aligned}$$

Hence  $g(S) \geq g(S_j^*) \geq g(M_1) + g(M_2) - g(F) + \frac{1}{2}$ , a contradiction.

This finishes the proof of Subcase 3.

**Subcase 4**  $F$  is transverse to  $S_j^*$  and any component of  $S_j^* \setminus F$  is incompressible in the respective submanifold of  $M^* \setminus F$  except for exactly one strongly irreducible component.

Since  $(\bigcup F_i)$  is incompressible in  $M$ , any component of  $S_j^* \setminus F$  is incompressible in the respective submanifold of  $M \setminus F$  except for exactly one strongly irreducible component. We may assume that any component of  $S_j^* \setminus F$  is incompressible in  $M_1$ , and denote the compressible component of  $S_j^* \cap M_2$  by  $Q'$ . Then  $S_j^* \cap M_1$  is essential in  $M_1$  and by assumption  $\chi(S_j^* \cap M_1) \leq 2 + g(F) - 2g(M_1)$ . It is easy to see that  $\chi(Q') \leq -2$ .

We compress  $Q'$  as much as possible in  $V_j^*$  ( $W_j^*$ , resp.) to obtain subsurfaces  $Q'_V$  ( $Q'_W$ , resp.), then any component of  $Q'_V$  and  $Q'_W$  is incompressible in  $V_j^* \cup_{S_j^*} W_j^*$ . Furthermore,  $Q'_V$  and  $Q'_W$  is incompressible in  $M_2$  since  $\bigcup F_i$  is incompressible in  $M$ . If there is one component of  $S_j^* \cap M_2$ ,  $Q'_V$  and  $Q'_W$  which is essential in  $V_j^* \cup_{S_j^*} W_j^*$ , then by assumption  $\chi(S_j^* \cap M_2) \leq 2 + g(F) - 2g(M_2) - 2$ . Hence  $g(S) \geq g(S_j^*) \geq g(M_1) + g(M_2) - g(F)$ , a contradiction.

So any component of  $Q'_V$  and  $Q'_W$  is parallel to a subsurface of  $F$  or a component of  $\partial_- V_j^*$  or  $\partial_- W_j^*$ . Any component of  $F \cap V_j^*$  is incompressible in  $V_j^*$ , by Lemma 2.2 any component of  $V_j^* \setminus F$  is a compression body. By the same reasons as above, any component of  $W_j^* \setminus F$  is a compression body. We amalgamate the Heegaard splitting  $V_j^* \cup_{S_j^*} W_j^*$  and the Heegaard splittings contained in  $M_2$  of the Heegaard sequence  $V_1 \cup_{S_1} W_1, V_2 \cup_{S_2} W_2, \dots, V_{n-1} \cup_{S_{n-1}} W_{n-1}, V_n \cup_{S_n} W_n$  along the components contained in  $M_2$  of  $\bigcup F_i$  to obtain a Heegaard splitting  $V' \cup_{S'} W'$  such that the following conditions are satisfied:

- (1)  $V' \cap M_1 = V_j^* \cap M_1$  and  $W' \cap M_1 = W_j^* \cap M_1$ .
- (2)  $S' \cap F = S_j^* \cap F$  and  $S' \cap M_1 = S_j^* \cap M_1$ .
- (3) Only one component of  $S' \cap M_2$  is compressible in  $V'$ , denoted by  $Q''$  and other incompressible components are just the components of  $S_j^* \cap M_2$ .
- (4)  $Q''$  can be compressed into  $Q''_V$  ( $Q''_W$ , resp.) in  $V'$  ( $W'$ , resp.) such that the incompressible components of  $Q''_V$  ( $Q''_W$ , resp.) with boundary are the same as the incompressible components of  $Q'_V$  ( $Q'_W$ , resp.) with boundary.

Then  $Q''$  satisfies the conditions of Lemma 3.4, so  $g(M_2) \leq 1 - \frac{1}{2}\chi(F \cap W') - \frac{1}{2}\chi(Q'')$  and  $g(M_2) \leq 1 - \frac{1}{2}\chi(F \cap V') - \frac{1}{2}\chi(Q'')$ . Note that  $\chi(F) = \chi(F \cap V') + \chi(F \cap W')$ , so  $2g(M_2) \leq 2 - \frac{1}{2}\chi(F) - \chi(Q'') = 1 + g(F) - \chi(Q'')$ . Then  $\chi(S' \cap M_2) \leq \chi(Q'') \leq 1 + g(F) - 2g(M_2)$ . Notice that  $\chi(S' \cap M_1) = \chi(S_j^* \cap M_1) \leq 2 + g(F) - 2g(M_1)$ . Hence,

$$\begin{aligned} \chi(S') &= \chi(S' \cap M_1) + \chi(S' \cap M_2) \\ &\leq 1 + g(F) - 2g(M_1) + 2 + g(F) - 2g(M_2) \\ &\leq 3 - 2(g(M_1) + g(M_2) - g(F)). \end{aligned}$$

So  $g(S') \geq g(M_1) + g(M_2) - g(F) - \frac{1}{2}$  and  $g(S) \geq g(S') \geq g(M_1) + g(M_2) - g(F)$ , a contradiction.

This finishes the proof of Subcase 4.

Therefore,  $g(M) = g(M_1) + g(M_2) - g(F)$  holds. This completes the proof of Theorem 4.2.  $\square$

In the following, we give an application of Theorem 4.2.

**Theorem 4.3** *Let  $M$  be a compact, orientable 3-manifold and  $F$  be an essential closed surface which cuts  $M$  into  $M_1$  and  $M_2$ . If  $M_i$  has a Heegaard splitting  $V_i \cup_{S_i} W_i$  with distance  $D(S_i) \geq 2g(M_i) - g(F)$ ,  $i = 1, 2$ . Then  $g(M) = g(M_1) + g(M_2) - g(F)$ .*

**Proof** For  $i = 1, 2$ , if  $Q_i$  is an essential surface in  $M_i$ . By assumptions and Lemma 2.4,  $2 - \chi(Q_i) \geq D(S_i) \geq 2g(M_i) - g(F)$ . Then  $\chi(Q_i) \leq 2 + 2g(M_i) - g(F)$ ,  $i = 1, 2$ . And by Theorem 4.2, we have  $g(M) = g(M_1) + g(M_2) - g(F)$ .  $\square$

**Remark** The condition in Theorem 4.2 is weaker than that in the main results of Kobayashi and Qiu [6] and Yang and Lei [19].

**Acknowledgements** The work is supported by the Programs of Cultivating Outstanding Youth Teachers in Harbin Institute of Technology. The authors would like to thank the referee for his valuable comments and suggestions.

## References

- [1] **D Bachman, S Schleimer, E Sedgwick**, *Sweepouts of amalgamated 3-manifolds*, *Algebr. Geom. Topol.* 6 (2006) 171–194 MR2199458
- [2] **K Hartshorn**, *Heegaard splittings of Haken manifolds have bounded distance*, *Pacific J. Math.* 204 (2002) 61–75 MR1905192
- [3] **J Hempel**, *3-manifolds*, *Ann. of Math. Studies* 86, Princeton Univ. Press (1976) MR0415619
- [4] **J Hempel**, *3-manifolds as viewed from the curve complex*, *Topology* 40 (2001) 631–657 MR1838999
- [5] **W Jaco**, *Lectures on three-manifold topology*, *CBMS Regional Conference Series in Math.* 43, Amer. Math. Soc. (1980) MR565450
- [6] **T Kobayashi, R Qiu**, *The amalgamation of high distance Heegaard splittings is always efficient*, *Math. Ann.* 341 (2008) 707–715 MR2399167

- [7] **T Kobayashi, R Qiu, Y Rieck, S Wang**, *Separating incompressible surfaces and stabilizations of Heegaard splittings*, Math. Proc. Cambridge Philos. Soc. 137 (2004) 633–643 MR2103921
- [8] **M Lackenby**, *The Heegaard genus of amalgamated 3-manifolds*, Geom. Dedicata 109 (2004) 139–145 MR2113191
- [9] **F Lei, G Yang**, *A lower bound of genus of amalgamations of Heegaard splittings*, Math. Proc. Cambridge Philos. Soc. 146 (2009) 615–623 MR2496347
- [10] **T Li**, *On the Heegaard splittings of amalgamated 3-manifolds*, from: “Workshop on Heegaard Splittings”, (C Gordon, Y Moriah, editors), Geom. Topol. Monogr. 12 (2007) 157–190 MR2408246
- [11] **M Scharlemann**, *Local detection of strongly irreducible Heegaard splittings*, Topology Appl. 90 (1998) 135–147 MR1648310
- [12] **M Scharlemann**, *Proximity in the curve complex: boundary reduction and bicompressible surfaces*, Pacific J. Math. 228 (2006) 325–348 MR2274524
- [13] **M Scharlemann, A Thompson**, *Thin position for 3-manifolds*, from: “Geometric topology (Haifa, 1992)”, (C Gordon, Y Moriah, B Wajnryb, editors), Contemp. Math. 164, Amer. Math. Soc. (1994) 231–238 MR1282766
- [14] **M Scharlemann, M Tomova**, *Alternate Heegaard genus bounds distance*, Geom. Topol. 10 (2006) 593–617 MR2224466
- [15] **J Schultens**, *The classification of Heegaard splittings for (compact orientable surface)  $\times S^1$* , Proc. London Math. Soc. (3) 67 (1993) 425–448 MR1226608
- [16] **J Schultens**, *Additivity of tunnel number for small knots*, Comment. Math. Helv. 75 (2000) 353–367 MR1793793
- [17] **J Schultens, R Weidmann**, *Destabilizing amalgamated Heegaard splittings*, from: “Workshop on Heegaard Splittings”, (C Gordon, Y Moriah, editors), Geom. Topol. Monogr. 12 (2007) 319–334 MR2408253
- [18] **J Souto**, *Distances in the curve complex and Heegaard genus*, Preprint (2003) Available at <http://www-personal.umich.edu/~jsouto/papers.html>
- [19] **G Yang, F Lei**, *On amalgamations of Heegaard splittings with high distance*, Proc. Amer. Math. Soc. 137 (2009) 723–731 MR2448595

*Department of Mathematics, Harbin Institute of Technology  
Harbin 150001, China*

*School of Mathematical Sciences, Dalian University of Technology  
Dalian 116024, China*

*klyang0707@yahoo.com.cn, ffcclei@yahoo.com.cn*

Received: 28 May 2009      Revised: 3 September 2009