

Symmetric homology of algebras

SHAUN V AULT

The symmetric homology of a unital algebra A over a commutative ground ring k is defined using derived functors and the symmetric bar construction of Fiedorowicz. For a group ring $A = k[\Gamma]$, the symmetric homology is related to stable homotopy theory via $HS_*(k[\Gamma]) \cong H_*(\Omega\Omega^\infty S^\infty(B\Gamma); k)$. Two chain complexes that compute $HS_*(A)$ are constructed, both making use of a symmetric monoidal category ΔS_+ containing ΔS . Two spectral sequences are found that aid in computing symmetric homology. The second spectral sequence is defined in terms of a family of complexes, $\text{Sym}_*^{(p)}$. $\text{Sym}^{(p)}$ is isomorphic to the suspension of the cycle-free chessboard complex Ω_{p+1} of Vrećica and Živaljević, and so recent results on the connectivity of Ω_n imply finite-dimensionality of the symmetric homology groups of finite-dimensional algebras. Some results about the $k\Sigma_{p+1}$ -module structure of $\text{Sym}^{(p)}$ are developed. A partial resolution is found that allows computation of $HS_1(A)$ for finite-dimensional A and some concrete computations are included.

55N35; 13D03, 18G10

1 Introduction and definitions

The theory of symmetric homology, in which the symmetric groups Σ_k^{op} , for $k \geq 0$, play the role that the cyclic groups do in cyclic homology, begins with the definition of the category ΔS , containing the simplicial category Δ as subcategory. Indeed, ΔS is an example of *crossed simplicial group* (see Fiedorowicz and Loday [10]). One develops a notion of a bar resolution over crossed simplicial groups by analogy with the cyclic bar resolution. However, the naïve contravariant symmetric bar resolution produces trivial results. Fiedorowicz found that a *covariant* symmetric bar construction, B_*^{sym} , produces a non-trivial and important homology theory (see Fiedorowicz [8]). One defines the symmetric homology of an algebra A by

Definition 1 $HS_*(A) = \text{Tor}_*^{\Delta S}(\underline{k}, B_*^{\text{sym}} A)$.

Almost 20 years ago, Fiedorowicz found but did not publish the following remarkable result [8].

Theorem 2 (Fiedorowicz’s Theorem) *If Γ is a group, then*

$$HS_*(k[\Gamma]) \cong H_*(\Omega\Omega^\infty S^\infty(B\Gamma); k).$$

This formula shows in particular that HS_* is an important and non-trivial theory. While it is true that $H_*(\Omega^\infty S^\infty X) = H_*(QX)$ is well understood, the same cannot be said of the homology of $\Omega\Omega^\infty S^\infty X$. Indeed, $H_*(QX)$ has been studied extensively by Cohen, Lada and May [4]. $H_*(QX)$ may be regarded as the free allowable AR -Hopf algebra with conjugation generated by $H_*(X)$. On the other hand, results about the homology of $\Omega\Omega^\infty S^\infty X$ are fewer and farther between. Cohen and Peterson [5] computed $H_*(\Omega\Omega^\infty S^\infty)$ (that is, the case when the space X is the zero-sphere, S^0), but there is little hope of extending this result to arbitrary X using the same methods.

We find that adjoining a “unit” to ΔS results in a permutative category ΔS_+ . This step is necessary in order to prove [Theorem 2](#) and related theorems.

By reducing the standard resolution that computes $HS_*(A)$ to one that involves only epimorphisms of ΔS_+ , we develop two spectral sequences abutting to $HS_*(A)$. The first spectral sequence is based on the work of Słomińska [27] on $E-I$ -categories, and relates symmetric homology to the homology of the symmetric groups.

The second spectral sequence makes use of a family of complexes, $\text{Sym}_*^{(p)}$ which distill the relevant combinatorial data of the nerve of $\text{Epi}_{\Delta S}$.

Theorem 3 *If A has an augmentation ideal I which is free as k -module, with countable basis X , then there is a spectral sequence $E_{p,q}^1 \Rightarrow \tilde{H}S_{p+q}(A)$, with*

$$E_{p,q}^1 \cong \bigoplus_{u \in X^{p+1}/\Sigma_{p+1}} H_{p+q}(E_*G_u \otimes_{G_u} \text{Sym}_*^{(p)}; k),$$

where G_u is the isotropy subgroup for the chosen representative of $u \in X^{p+1}/\Sigma_{p+1}$ and the symbol \otimes_{G_u} stands for the chain complex analog of equivariant half-smash product for spaces, \times_{G_u} .

The multiplicative structure of A becomes encapsulated in the differential $d_{p,q}^1$ on E^1 , and so results about $HS_*(A)$ for general algebras A will follow from the structure of $\text{Sym}_*^{(p)}$. There is an isomorphism of complexes, $k[S\Omega_{p+1}^+] \xrightarrow{\cong} \text{Sym}_*^{(p)}$, where Ω_n^+ is the augmented cycle-free $(n \times n)$ -chessboard complex of Vrećica and Živaljević [30], which produces the following immediate important corollary:

Corollary 4 *If A is finitely-generated over a Noetherian ground ring k , then $HS_*(A)$ is finitely-generated over k in each degree.*

In the final section, we develop a partial resolution of the trivial ΔS^{op} -module \underline{k} by projective modules, leading to the following:

Theorem 5 $HS_i(A)$ for $i = 0, 1$ may be computed as the degree 0 and degree 1 homology groups of the (partial) chain complex

$$0 \longleftarrow A \xleftarrow{\partial_1} A \otimes A \otimes A \xleftarrow{\partial_2} (A \otimes A \otimes A \otimes A) \oplus A,$$

where

$$\begin{aligned} \partial_1: a \otimes b \otimes c &\mapsto abc - cba, \\ \partial_2: \begin{cases} a \otimes b \otimes c \otimes d &\mapsto ab \otimes c \otimes d + d \otimes ca \otimes b \\ &\quad + bca \otimes 1 \otimes d + d \otimes bc \otimes a, \\ a &\mapsto 1 \otimes a \otimes 1. \end{cases} \end{aligned}$$

In particular, we see that $HS_0(A) = A/[A, A]$, where $[A, A]$ is the ideal generated by the commutator submodule $[A, A]$. In other words, $HS_0(A)$ is the symmetrization of A as an algebra. Compare with the zeroth Hochschild or Cyclic homology of A , which is $A/[A, A]$.

1.1 The category ΔS

Let ΔS be the category that has as objects, the ordered sets $[n] = \{0, 1, \dots, n\}$ for $n \geq 0$, and as morphisms, pairs (ϕ, g) , where $\phi: [n] \rightarrow [m]$ is a non-decreasing map of sets (that is, a morphism in Δ), and $g \in \Sigma_{n+1}^{\text{op}}$ (the opposite group of the symmetric group acting on $[n]$). The element g represents an automorphism of $[n]$, and as a set map, takes $i \in [n]$ to $g^{-1}(i)$. Equivalently, a morphism in ΔS is a morphism in Δ together with a total ordering of the domain $[n]$. Composition of morphisms is achieved as in Fiedorowicz–Loday [10], namely, $(\phi, g) \circ (\psi, h) = (\phi \cdot g^*(\psi), \psi^*(g) \cdot h)$. Observe that the properties of $g^*(\phi)$ and $\phi^*(g)$ stated in [10, Proposition 1.6] are formally rather similar to the properties of exponents (except for properties 2.h and 2.v). Indeed, the notation $g^\phi \stackrel{\text{def}}{=} \phi^*(g)$, $\phi^g \stackrel{\text{def}}{=} g^*(\phi)$ will generally be used in lieu of the original notation in what follows. For reference, we restate [10, Proposition 1.6] using the “exponent” notation:

Proposition 6 If G_* is a crossed simplicial group, for $g, h \in G_n$, $\phi \in \text{Mor}_\Delta([m], [n])$, $\psi \in \text{Mor}_\Delta([p], [m])$,

$$\begin{aligned} (1.h)' \quad g^\phi \psi &= (g^\phi) \psi & (1.v)' \quad \phi^{gh} &= (\phi^g)^h \\ (2.h)' \quad (\phi \psi)^g &= \phi^g \psi^{(g^\phi)} & (2.v)' \quad (gh)^\phi &= g^\phi h^{(\phi^g)} \\ (3.h)' \quad g^{\text{id}_n} &= g \quad \text{and} \quad 1^\phi = 1 & (3.v)' \quad \phi^1 &= \phi \quad \text{and} \quad \text{id}_n^g = \text{id} \end{aligned}$$

It is often helpful to represent morphisms of ΔS as diagrams of points and lines, indicating images of set maps. Using these diagrams, we may see clearly how ψ^g and g^ψ are related to ψ and g (see Figure 1).




Figure 1: Morphisms of ΔS : $g \cdot \psi = (g^*(\psi), \psi^*(g)) = (\psi^g, g^\psi)$

An equivalent characterization of ΔS comes from Pirashvili, as the category $\mathcal{F}(\text{as})$ of *non-commutative sets* [22]. The objects are sets $\underline{n} \stackrel{\text{def}}{=} \{1, 2, \dots, n\}$ for $n \geq 0$. By convention, $\underline{0}$ is the empty set. A morphism in $\text{Mor}_{\mathcal{F}(\text{as})}(\underline{n}, \underline{m})$ consists of a set map $f: \underline{n} \rightarrow \underline{m}$ together with a total ordering on each preimage set $f^{-1}(j)$. There is an obvious inclusion of categories, $\Delta S \hookrightarrow \mathcal{F}(\text{as})$, taking $[n]$ to $\underline{n+1}$, but there is no object of ΔS that maps to $\underline{0}$. It will be useful to define $\Delta S_+ \supset \Delta S$ which is isomorphic to $\mathcal{F}(\text{as})$:

Definition 7 ΔS_+ is the category consisting of all objects and morphisms of ΔS , with the additional object $[-1]$, representing the empty set, and a unique morphism $\iota_n: [-1] \rightarrow [n]$ for each $n \geq -1$.

Remark 8 Pirashvili’s construction is a special case of a more general construction due to May and Thomason [20]. This construction associates to any topological operad $\{\mathcal{C}(n)\}_{n \geq 0}$ a topological category $\hat{\mathcal{C}}$ together with a functor $\hat{\mathcal{C}} \rightarrow \mathcal{F}$, where \mathcal{F} is the category of finite sets, such that the inverse image of any function $f: \underline{m} \rightarrow \underline{n}$ is the space $\prod_{i=1}^n \mathcal{C}(\#f^{-1}(i))$. Composition in $\hat{\mathcal{C}}$ is defined using the composition of the operad. May and Thomason refer to $\hat{\mathcal{C}}$ as the *category of operators* associated to \mathcal{C} .

They were interested in the case of an E_∞ operad, but their construction evidently works for any operad. The category of operators associated to the discrete A_∞ operad $\mathcal{A}ss$, which parametrizes monoid structures, is precisely Pirashvili's construction of $\mathcal{F}(as)$, that is, ΔS_+ .

One very useful advantage in enlarging our category to ΔS to ΔS_+ is the added structure inherent in ΔS_+ .

Proposition 9 ΔS_+ is a permutative category.

Proof Define the monoid product on objects by $[n] \odot [m] \stackrel{\text{def}}{=} [n + m + 1]$, (disjoint union of sets), and on morphisms $(\phi, g): [n] \rightarrow [n']$, $(\psi, h): [m] \rightarrow [m']$, by $(\phi, g) \odot (\psi, h) = (\eta, k): [n + m + 1] \rightarrow [n' + m' + 1]$, where (η, k) is just the morphism (ϕ, g) acting on the first $n + 1$ points of $[n + m + 1]$, and (ψ, h) acting on the remaining points.

The unit object will be $[-1] = \emptyset$. \odot is clearly associative, and $[-1]$ acts as two-sided identity. Finally, define the transposition transformation $\gamma_{n,m}: [n] \odot [m] \rightarrow [m] \odot [n]$ to be the identity on objects, and on morphisms to be precomposition with the block transposition that switches the first block of size $n + 1$ with the second block of size $m + 1$. \square

Remark 10 The fact that ΔS_+ is permutative shall be exploited to prove that $HS_*(A)$ admits homology operations in a forthcoming paper.

Remark 11 It will become convenient to include the object $[-1] = \emptyset$ in Δ as well. Denote the enlarged category by Δ_+ .

For the purposes of computation, a morphism $\alpha: [n] \rightarrow [m]$ of ΔS may be conveniently represented as a tensor product of monomials in the formal non-commuting variables $\{x_0, x_1, \dots, x_n\}$. Let $\alpha = (\phi, g)$, with $\phi \in \text{Mor}_\Delta([n], [m])$ and $g \in \Sigma_{n+1}^{\text{op}}$. The tensor representation of α will have $m + 1$ tensor factors. Each x_i will occur exactly once, in the order $x_{g(0)}, x_{g(1)}, \dots, x_{g(n)}$. The i th tensor factor consists of the product of $\#\phi^{-1}(i - 1)$ variables, with the convention that the empty product will be denoted 1. Thus, the i th tensor factor records the total ordering of $\phi^{-1}(i)$. As an example, the tensor representation of the morphism depicted in [Figure 2](#) is $x_1 x_0 \otimes x_3 x_4 \otimes 1 \otimes x_2$. With this notation, the composition of two morphisms $\alpha = X_0 \otimes X_1 \otimes \dots \otimes X_m: [n] \rightarrow [m]$ and $\beta = Y_1 \otimes Y_2 \otimes \dots \otimes Y_n: [p] \rightarrow [n]$ is given by, $\alpha\beta = Z_0 \otimes Z_1 \otimes \dots \otimes Z_m$, where Z_i is determined by replacing each variable in the monomials $X_i = x_{j_1} \dots x_{j_s}$ by the corresponding monomials Y_{j_k} in β . So, $Z_i = Y_{j_1} \dots Y_{j_s}$.




Figure 2: Morphisms of ΔS in tensor notation

1.2 Homological algebra of functors

Recall, for a category \mathcal{C} , a \mathcal{C} -module is covariant functor $F: \mathcal{C} \rightarrow k\text{-Mod}$. Similarly, a \mathcal{C}^{op} -module is a contravariant functor $G: \mathcal{C} \rightarrow k\text{-Mod}$. Let M be a \mathcal{C}^{op} -module and N be a \mathcal{C} -module. Following Mac Lane [17], define the tensor product of functors as a coend, $M \otimes_{\mathcal{C}} N = \int^X (MX) \otimes (NX)$. That is,

$$M \otimes_{\mathcal{C}} N = \bigoplus_{X \in \text{Obj} \mathcal{C}} M(X) \otimes_k N(X) / \approx,$$

where the equivalence \approx is generated by $y \otimes f_*(x) \approx f^*(y) \otimes x$ for $f \in \text{Mor}_{\mathcal{C}}(X, Y)$, $x \in N(X)$ and $y \in M(Y)$.

Remark 12 Note, the existing literature based on the work of Connes, Loday and Quillen consistently defines the categorical tensor product in the reverse sense: $N \otimes_{\mathcal{C}} M$ is the direct sum of copies of $NX \otimes_k MX$ modded out by the equivalence $x \otimes f^*(y) \approx f_*(x) \otimes y$ for all \mathcal{C} -morphisms $f: X \rightarrow Y$. In this context, N is covariant, while M is contravariant. I chose to follow the convention of Pirashvili and Richter [23] in writing tensor products as $M \otimes_{\mathcal{C}} N$ so that the equivalence $\xi: \mathcal{C}\text{-Mod} \rightarrow k[\text{Mor } \mathcal{C}]\text{-Mod}$ passes to tensor products in a straightforward way: $\xi(M \otimes_{\mathcal{C}} N) = \xi(M) \otimes_{k[\text{Mor } \mathcal{C}]} \xi(N)$.

The *trivial* \mathcal{C} -module, resp. \mathcal{C}^{op} -module, denoted by \underline{k} (for either variance), is the functor taking each object to k and each morphism to the identity. As noted in Fiedorowicz–Loday [10], the category of \mathcal{C} -modules is abelian and has enough projectives. For any \mathcal{C}^{op} -module M , the functor $N \mapsto M \otimes_{\mathcal{C}} N$ is right-exact and so admits derived functors $\text{Tor}_n^{\mathcal{C}}(M, -)$ such that $\text{Tor}_0^{\mathcal{C}}(M, N) = M \otimes_{\mathcal{C}} N$.

1.3 The symmetric bar construction

Now that we have defined the category ΔS , the next step should be to define an appropriate bar construction. Recall that the *cyclic bar construction* is a functor $B_*^{\text{cyc}} A: \Delta C^{\text{op}} \rightarrow k\text{-Mod}$. One then takes the groups $\text{Tor}_n^{\Delta C}(B_*^{\text{cyc}} A, \underline{k})$ as the definition of $HC_n(A)$ for $n \geq 0$. However the results of [10] show that the cyclic bar construction does not extend to a functor $\Delta S^{\text{op}} \rightarrow k\text{-Mod}$. Furthermore, for any functor $F: \Delta S^{\text{op}} \rightarrow k\text{-Mod}$, the groups $\text{Tor}_n^{\Delta S}(F, \underline{k})$ simply compute the homology of the underlying simplicial module of F (given by restricting F to Δ^{op}). However, Fiedorowicz discovered that there is a natural extension of the cyclic bar construction not to a *contravariant* functor on ΔS , but to a *covariant* functor [8].

Definition 13 Let A be an associative, unital algebra over a commutative ground ring k . Define a ΔS -module (that is, a functor $\Delta S \rightarrow k\text{-Mod}$), $B_*^{\text{sym}} A$ by

$$B_n^{\text{sym}} A = B_*^{\text{sym}} A[n] \stackrel{\text{def}}{=} A^{\otimes(n+1)}$$

$$B_*^{\text{sym}} A(\alpha): (a_0 \otimes a_1 \otimes \dots \otimes a_n) \mapsto \alpha(a_0, \dots, a_n),$$

where $\alpha: [n] \rightarrow [m]$ is represented in tensor notation, and evaluation at (a_0, \dots, a_n) simply amounts to substituting each a_i for x_i and multiplying the resulting monomials in A . If the pre-image $\alpha^{-1}(i)$ is empty, then the unit of A is inserted. Observe that $B_*^{\text{sym}} A$ is natural in A .

Note that there are natural inclusions $\Delta \hookrightarrow \Delta C \hookrightarrow \Delta S$. The second inclusion is induced from inclusions of groups $C_{n+1} \hookrightarrow \Sigma_{n+1}$. To be precise, for each n , let τ_n be the $(n+1)$ -cycle $(0, n, n-1, \dots, 1) \in \Sigma_{n+1}$. τ_n generates a subgroup isomorphic to C_{n+1} . $B_*^{\text{sym}} A$ may be regarded as a simplicial k -module via the chain of functors, $\Delta^{\text{op}} \hookrightarrow \Delta C^{\text{op}} \xrightarrow{\cong} \Delta C \hookrightarrow \Delta S$. Here, the isomorphism $D: \Delta C^{\text{op}} \rightarrow \Delta C$ is the standard duality (see Loday [16]). Note that the *cyclic bar construction* can be recovered from the covariant symmetric bar construction by $B_*^{\text{cyc}} A = B_*^{\text{sym}} A \circ D$.

Definition 14 The *symmetric homology* of an associative, unital k -algebra A is denoted $HS_*(A)$, and is defined as

$$HS_*(A) \stackrel{\text{def}}{=} \text{Tor}_*^{\Delta S}(\underline{k}, B_*^{\text{sym}} A)$$

Remark 15 Since $\underline{k} \otimes_{\Delta S} M \cong \text{colim}_{\Delta S} M$, for any ΔS -module M , we can alternatively describe symmetric homology as derived functors of the colimit

$$HS_n(A) = \text{colim}_{\Delta S} {}^{(n)} B_*^{\text{sym}} A.$$

In the language of Gabriel–Zisman [11, Appendix II.3], $HS_*(A) = H_*(\Delta S, B_*^{\text{sym}} A)$. Moreover, a simplicial module whose homology computes $H_*(\Delta S, B_*^{\text{sym}} A)$ is readily available [11, page 153]. In order to describe this simplicial module, we recall that the *nerve* of a category \mathcal{C} is the simplicial set $N\mathcal{C}$ such that n –chains are sequences of n composable morphisms of \mathcal{C} , 0 –chains are simply objects of \mathcal{C} , and face maps are given by deletion of objects along with the composition of the corresponding morphisms. If $F: \mathcal{C} \rightarrow \mathcal{D}$ is a morphism and $\chi \in N\mathcal{C}$ is an n –chain,

$$\chi_n \xleftarrow{f_n} \chi_{n-1} \xleftarrow{f_{n-1}} \cdots \xleftarrow{f_2} \chi_1 \xleftarrow{f_1} \chi_0,$$

then let $F\chi = F\chi_0$. The following simplicial module computes $H_*(\Delta S, B_*^{\text{sym}} A)$:

$$\{C_n(\Delta S, B_*^{\text{sym}} A)\}_{n \geq 0} = \left\{ \bigoplus_{\chi \in N_n \Delta S} B_*^{\text{sym}} A\chi \right\}_{n \geq 0},$$

with face maps and degeneracies defined by:

$$d_i(\chi, x) = \begin{cases} (d_0\chi, (f_1)_*(x)), & i = 0, \\ (d_i\chi, x), & 1 \leq i \leq n \end{cases}$$

$$s_i(\chi, x) = (s_i\chi, x)$$

In this formula, we have $\chi \in N_n \Delta S$ and $x \in B_*^{\text{sym}} A\chi$.

$C_n(\Delta S, B_*^{\text{sym}} A)$ may be interpreted in terms of the categorical tensor product construction mentioned in Section 1.2, using the *under-category* functor, $(-\setminus \Delta S): \Delta S^{\text{op}} \rightarrow \mathbf{cat}$, which associates to each object $[p]$ of ΔS , the category $[p] \setminus \Delta S$ of objects under $[p]$ (see Quillen [24], Mac Lane [18], etc). The n –chains of $N([p] \setminus \Delta S)$ may be identified with $(n+1)$ –chains of $N\Delta S$ which have domain $[p]$. Let $k[-]$ be the functor that associates to any simplicial set the corresponding simplicial k –module. We may identify

$$C_*(\Delta S, B_*^{\text{sym}} A) = k[N(-\setminus \Delta S)] \otimes_{\Delta S} B_*^{\text{sym}} A.$$

Indeed, observe that every element of $k[N(-\setminus \Delta S)] \otimes_{\Delta S} B_*^{\text{sym}} A$ of the form

$$\left(\begin{array}{c} [p] \\ \swarrow \quad \searrow \quad \downarrow \alpha \\ [q_n] \xleftarrow{\beta_n} \cdots \xleftarrow{\beta_2} [q_1] \xleftarrow{\beta_1} [q_0] \end{array} \right) \otimes x$$

is equivalent to one in which the incoming morphism α becomes an identity:

$$\left(\begin{array}{ccccc} & & & & [q_0] \\ & & & & \downarrow \text{id} \\ & & & & [q_0] \\ & & & \swarrow \beta_1 & \\ & & [q_1] & \longleftarrow \beta_1 & \\ & & \cdots & \longleftarrow \beta_2 & \\ & [q_n] & \longleftarrow \beta_n & & \end{array} \right) \otimes \alpha_*(x),$$

the latter element being identified with

$$([q_n] \xleftarrow{\beta_n} \cdots \xleftarrow{\beta_2} [q_1] \xleftarrow{\beta_1} [q_0], \alpha_*(x)) \in C_n(\Delta S, B_*^{\text{sym}} A).$$

Proposition 16 For an associative, unital k -algebra A ,

$$HS_*(A) = H_*(k[N(- \setminus \Delta S)] \otimes_{\Delta S} B_*^{\text{sym}} A; k).$$

Remark 17 By duality of ΔC , it is clear that the related complex $k[N(- \setminus \Delta C)] \otimes_{\Delta C} B_*^{\text{sym}} A$ computes $HC_*(A)$, where we understand that the functor $B_*^{\text{sym}} A$ is restricted to $\Delta C \hookrightarrow \Delta S$.

1.4 Symmetric homology of the ground ring

We now have enough tools to compute $HS_*(k)$. First, we need to show:

Lemma 18 $N(\Delta S)$ is contractible.

Proof Define a functor $\mathcal{F}: \Delta S \rightarrow \Delta S$ on objects by $\mathcal{F}[n] = [0] \odot [n]$, and on morphisms by $\mathcal{F}f = \text{id}_{[0]} \odot f$, using the monoid multiplication \odot defined in Proposition 9. There is a natural transformation $\text{id}_{\Delta S} \rightarrow \mathcal{F}$ given by the following commutative diagram for each $f: [m] \rightarrow [n]$:

$$\begin{array}{ccc} [m] & \xrightarrow{f} & [n] \\ \delta_0 \downarrow & & \downarrow \delta_0 \\ [m+1] & \xrightarrow{\text{id} \odot f} & [n+1] \end{array}$$

Here, $\delta_j^{(k)}: [k-1] \rightarrow [k]$ is the Δ morphism that misses the point $j \in [k]$.

Consider the constant functor $\Delta S \xrightarrow{[0]} \Delta S$ that sends all objects to $[0]$ and all morphisms to $\text{id}_{[0]}$. There is a natural transformation $[0] \rightarrow \mathcal{F}$ given by the following commutative

diagram for each $f: [m] \rightarrow [n]$.

$$\begin{array}{ccc}
 [0] & \xrightarrow{\text{id}} & [0] \\
 \downarrow 0_0 & & \downarrow 0_0 \\
 [m+1] & \xrightarrow{\text{id} \odot f} & [n+1]
 \end{array}$$

Here, $0_j^{(k)}: [0] \rightarrow [k]$ is the morphism that sends the point 0 to $j \in [k]$.

Natural transformations induce homotopy equivalences (see Segal [26] or Quillen [24, Proposition 1.2]), so in particular, the identity map on $N(\Delta S)$ is homotopic to the map that sends $N(\Delta S)$ to the nerve of a trivial category. Thus, $N(\Delta S)$ is contractible. \square

Corollary 19 *The symmetric homology of the ground ring k is isomorphic to k , concentrated in degree 0.*

Proof $HS_*(k)$ is the homology of the chain complex generated (freely) over k by the chains

$$\left\{ [q_n] \xleftarrow{\beta_n} \dots \xleftarrow{\beta_2} [q_1] \xleftarrow{\beta_1} [q_0] \otimes (1 \otimes \dots \otimes 1) \right\},$$

where $\beta_i \in \text{Mor}_{\Delta S}([q_{i-1}], [q_i])$. Each such chain may be identified with the chain

$$[q_n] \xleftarrow{\beta_n} \dots \xleftarrow{\beta_2} [q_1] \xleftarrow{\beta_1} [q_0]$$

of $N(\Delta S)$, and this defines a chain isomorphism to $N(\Delta S)$. The result now follows from Lemma 18. \square

2 Symmetric homology with coefficients and the UCT

Following the conventions for Hochschild and cyclic homology in Loday [16], when we need to indicate explicitly the ground ring k over which we compute symmetric homology of A , we shall use the notation: $HS_*(A | k)$. On the other hand, for a k -module M , $HS_*(A; M)$ will denote the homology of the complex $C_*(\Delta S, B_*^{\text{sym}} A) \otimes_k M$.

Two easy propositions are collected here for reference.

Proposition 20

- (1) *If M is flat over k , then $HS_*(A; M) \cong HS_*(A) \otimes_k M$.*
- (2) *If B is a commutative k -algebra, then $HS_*(A \otimes_k B | B) \cong HS_*(A; B)$.*

Proof Left to the reader. \square

Theorem 21 (Universal Coefficient Theorem) *If A is a flat k -algebra, and B is a commutative k -algebra, then there is a spectral sequence with*

$$E_2^{p,q} = \text{Tor}_p^k(HS_q(A|k), B) \Rightarrow HS_*(A|B).$$

Proof Let $T_q: k\text{-Mod} \rightarrow k\text{-Mod}$ be the functor $HS_q(A; -)$. Observe, since A is flat, $\{T_q\}$ is a long exact sequence of additive covariant functors (see Dold [6, Definition 1.1] and also McCleary [21, Section 12.1] for details). $T_q = 0$ for sufficiently small q (indeed, for $q < 0$) and T_q commutes with arbitrary direct sums. Hence, by the Universal Coefficient Theorem [6, Theorem 2.12] (see also [21, Theorem 12.11]), there is a spectral sequence with $E_2^{p,q} = \text{Tor}_p^k(T_q(k), B) \Rightarrow T_*(B)$. \square

As an immediate consequence, we have the following result.

Corollary 22 *If $f: A \rightarrow A'$ is a k -algebra map between flat algebras inducing an isomorphism $HS_*(A) \xrightarrow{\cong} HS_*(A')$ in symmetric homology, then for a commutative k -algebra B , the chain map $f \otimes \text{id}_B$ induces an isomorphism $HS_*(A; B) \xrightarrow{\cong} HS_*(A'; B)$.*

Under stronger hypotheses, the universal coefficient spectral sequence reduces to short exact sequences.

Corollary 23 *If k has weak global dimension ≤ 1 , then the spectral sequence of Theorem 21 reduces to short exact sequences,*

$$0 \longrightarrow HS_n(A|k) \otimes_k B \longrightarrow HS_n(A; B) \longrightarrow \text{Tor}_1^k(HS_{n-1}(A|k), B) \longrightarrow 0.$$

Moreover, if k is hereditary and A is projective over k , then these sequences split (unnaturally). (See Itô [13] for precise definitions of weak global dimension and hereditary.)

Proof Assume first that k has weak global dimension ≤ 1 . So $\text{Tor}_p^k(HS_q(A|k), B) = 0$ for all $p > 1$. Following Dold's argument [6, Corollary 2.13], we obtain the required exact sequences. Assume further that k is hereditary and A is projective. Then Rotman [25, Theorem 8.22] gives us the desired splitting. \square

Remark 24 The proof given above also proves UCT for cyclic homology. A partial result along these lines exists in Loday [16, 2.1.16]. There, he shows $HC_*(A|k) \otimes_k K \cong HC_*(A|K)$ and $HH_*(A|k) \otimes_k K \cong HH_*(A|K)$ in the case that K is a localization of k , and A is a K -module, flat over k . I am not aware of a statement of UCT for cyclic or Hochschild homology in its full generality in the literature.

3 Integral symmetric homology and a Bockstein spectral sequence

We shall obtain a converse to [Corollary 22](#) in the case $k = \mathbb{Z}$.

Theorem 25 *Let $f: A \rightarrow A'$ be an algebra map between torsion-free \mathbb{Z} -algebras. Suppose that for $B = \mathbb{Q}$ or $B = \mathbb{Z}/p\mathbb{Z}$ (for any prime p) the map*

$$f \otimes \text{id}_B: C_*(\Delta S, B_*^{\text{sym}} A) \otimes_k B \longrightarrow C_*(\Delta S, B_*^{\text{sym}} A') \otimes_k B$$

induces an isomorphism $HS_(A; B) \longrightarrow HS_*(A'; B)$. Then f also induces an isomorphism $HS_*(A) \xrightarrow{\cong} HS_*(A')$.*

First, note that if A is flat over k , one can construct the Bockstein homomorphisms $\beta_n: HS_n(A; \mathbb{Z}) \rightarrow HS_{n-1}(A; X)$ associated to a short exact sequence of k -modules, $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$. These Bockstein maps are natural in A , since $HS_*(A; -)$ is a long exact sequence of functors (see [Section 2](#)). Moreover if the induced map $f_*: HS_*(A; W) \rightarrow HS_*(A'; W)$ is an isomorphism for any two of $W = X$, $W = Y$ and $W = Z$, then it is an isomorphism for the third – an easy exercise for the reader. We shall now proceed with the proof of [Theorem 25](#). All tensor products will be over \mathbb{Z} in what follows.

Proof Let A and A' be torsion-free \mathbb{Z} -modules. Let $f: A \rightarrow A'$ be an algebra map inducing isomorphism in symmetric homology with coefficients in \mathbb{Q} and also in $\mathbb{Z}/p\mathbb{Z}$ for any prime p . For $m \geq 2$, there is a short exact sequence,

$$0 \longrightarrow \mathbb{Z}/p^{m-1}\mathbb{Z} \xrightarrow{p} \mathbb{Z}/p^m\mathbb{Z} \longrightarrow \mathbb{Z}/p\mathbb{Z} \longrightarrow 0.$$

A straightforward induction argument shows that the maps induced by f are isomorphisms,

$$(1) \quad f_*: HS_*(A; \mathbb{Z}/p^m\mathbb{Z}) \xrightarrow{\cong} HS_*(A'; \mathbb{Z}/p^m\mathbb{Z})$$

Denote $\mathbb{Z}/p^\infty\mathbb{Z} = \varinjlim \mathbb{Z}/p^m\mathbb{Z}$. Note, this is a *direct limit* in the sense that it is a colimit over a directed system. The direct limit functor is exact (see Stenström [[28](#), Proposition 5.3]), so the maps $HS_n(A; \mathbb{Z}/p^\infty\mathbb{Z}) \rightarrow HS_n(A'; \mathbb{Z}/p^\infty\mathbb{Z})$ induced by f are isomorphisms, given by the following chain of isomorphisms:

$$\begin{aligned} HS_n(A; \mathbb{Z}/p^\infty\mathbb{Z}) &\cong \varinjlim H_*(k[C_*(\Delta S, B_*^{\text{sym}} A)] \otimes \mathbb{Z}/p^m\mathbb{Z}) \\ &\xrightarrow{f_*} \varinjlim H_*(k[C_*(\Delta S, B_*^{\text{sym}} A')] \otimes \mathbb{Z}/p^m\mathbb{Z}) \cong HS_n(A'; \mathbb{Z}/p^\infty\mathbb{Z}) \end{aligned}$$

(Note that f_* here stands for $\varinjlim H_n(k[C_*(\Delta S, B_*^{\text{sym}} f)] \otimes \text{id})$.)

Finally, consider the short exact sequence of abelian groups,

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Q} \longrightarrow \bigoplus_{p \text{ prime}} \mathbb{Z}/p^\infty \mathbb{Z} \longrightarrow 0$$

The isomorphism $f_*: HS_*(A; \mathbb{Z}/p^\infty \mathbb{Z}) \rightarrow HS_*(A'; \mathbb{Z}/p^\infty \mathbb{Z})$ passes to direct sums, giving isomorphisms

$$f_*: HS_n\left(A; \bigoplus_p \mathbb{Z}/p^\infty \mathbb{Z}\right) \xrightarrow{\cong} HS_n\left(A'; \bigoplus_p \mathbb{Z}/p^\infty \mathbb{Z}\right)$$

for each n . Since $HS_*(A; \mathbb{Q}) \rightarrow HS_*(A'; \mathbb{Q})$ is an isomorphism, we have the required isomorphism in symmetric homology, $f_*: HS_n(A | \mathbb{Z}) \xrightarrow{\cong} HS_n(A' | \mathbb{Z})$. \square

Remark 26 [Theorem 25](#) may be useful for determining integral symmetric homology, since rational computations are generally simpler, and computations mod p may be made easier due to the presence of additional structure, such as homology operations (see the author’s forthcoming paper).

Finally, we state a result along the lines of McCleary [[21](#), Theorem 10.3]. Denote the torsion submodule of a graded module H_* by $\tau(H_*)$.

Theorem 27 (Bockstein spectral sequence) *Suppose A is free of finite rank over \mathbb{Z} . Then there is a singly-graded spectral sequence with*

$$E_*^1 = HS_*(A; \mathbb{Z}/p\mathbb{Z}) \Rightarrow HS_*(A)/\tau(HS_*(A)) \otimes \mathbb{Z}/p\mathbb{Z},$$

with differential map $d^1 = \beta$, the standard Bockstein map associated to

$$0 \rightarrow \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{Z}/p^2\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z} \rightarrow 0.$$

Moreover, the convergence is strong.

Proof The proof McCleary gives [[21](#), page 459] carries over to our case intact. All that is required for this proof is that each $HS_n(A)$ be a finitely-generated abelian group. The hypothesis that A is finitely-generated, coupled with the result of [Corollary 78](#) of [Section 9.4](#), guarantees this. \square

4 Symmetric homology using ΔS_+

In this section, we shall show that replacing ΔS by ΔS_+ in an appropriate way does not affect the computation of HS_* . We shall define a functor $B_*^{\text{sym}+} A: \Delta S_+ \rightarrow k\text{-Mod}$ such that $H_*(\Delta S_+, B_*^{\text{sym}+} A) \cong H_*(\Delta S, B_*^{\text{sym}} A)$.

4.1 Extension of the bar construction to ΔS_+

Recall for a functor $\mathcal{F}: \mathcal{D} \rightarrow \mathcal{C}$ of small categories, the induced functor from the category of \mathcal{C} -modules to \mathcal{D} -modules given by $M \mapsto M \circ \mathcal{F}$ admits a left adjoint functor, \mathcal{F}^* (see Kan [14] or Gabriel–Zisman [11, Appendix II.3]). In particular, the inclusion $i: \Delta S \hookrightarrow \Delta S_+$ induces a functor $i^*: \Delta S\text{-Mod} \rightarrow \Delta S_+\text{-Mod}$. It is instructive to analyze the functor $i^*(B_*^{\text{sym}} A): \Delta S_+ \rightarrow k\text{-Mod}$. Note, $i^*(B_*^{\text{sym}} A)$ is just the left Kan extension of $B_*^{\text{sym}} A$ along i . By definition, $i^*(B_*^{\text{sym}} A)$ is the functor, $[n] \mapsto \text{colim}_{(i/[n])} B_*^{\text{sym}} A \circ \pi_1$, where $\pi_1: (i/[n]) \rightarrow \Delta S$ is the projection $(i[m] \xrightarrow{\phi} [n]) \mapsto [m]$. Note, for $n = -1$, $\text{colim}_{(i/[-1])} B_*^{\text{sym}} A \circ \pi_1 \cong 0$, the initial object of $k\text{-Mod}$, since the category $(i/[-1])$ is empty. For $n \geq 0$,

$$(2) \quad \text{colim}_{(i/[n])} B_*^{\text{sym}} \circ \pi_1 = \left(\bigoplus_{([n] \leftarrow [m]) \in \text{Mor } \Delta S} B_m^{\text{sym}} A \right) / (\phi \psi, x) \approx (\phi, \psi_*(x)) \cong B_n^{\text{sym}} A$$

The last isomorphism results from the “identity trick” (see, for example, Section 1.3), namely $(\phi, x) \approx (\text{id}, \phi_*(x))$. For any morphism $\beta \in \text{Mor } \Delta S$, the morphism induced by $i^*(B_*^{\text{sym}} A)$ is simply β itself. On the other hand, the morphism $\iota_n: [-1] \rightarrow [n]$ of ΔS_+ becomes the zero map $0 \rightarrow A^{\otimes(n+1)}$ under $i^*(B_*^{\text{sym}} A)$.

Definition 28 Given an associative, unital algebra A over a commutative ground ring k , we let $B_*^{\text{sym}^0_+} A$ be the functor $i^*(B_*^{\text{sym}} A): \Delta S_+ \rightarrow k\text{-Mod}$ defined above.

Lemma 29 $H_*(\Delta S_+, B_*^{\text{sym}^0_+} A) \cong H_*(\Delta S, B_*^{\text{sym}} A)$.

Proof By Gabriel–Zisman [11, Theorem 3.6], there is a spectral sequence with

$$(3) \quad E_{p,q}^2 = \text{colim}_{\Delta S_+}^{(p)} (L_q i^*)(B_*^{\text{sym}} A) \Rightarrow \text{colim}_{\Delta S}^{(p+q)} B_*^{\text{sym}} A,$$

where i is the inclusion $\Delta S \hookrightarrow \Delta S_+$ as above and $L_q i^*$ is the q th left-derived functor of the right-exact functor i^* . The target $\text{colim}_{\Delta S}^{(p+q)} B_*^{\text{sym}} A$ is the same as $HS_{p+q}(A)$. I claim that i^* is in fact left-exact.

Let $0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$ be an exact sequence of ΔS -modules. By the same argument displayed in (2), we have

$$\text{colim}_{(i/[n])} M \circ \pi_1 \cong \begin{cases} 0, & \text{if } n = -1 \\ M_n, & \text{if } n \geq 0 \end{cases}$$

Similar results follow for N and P . Thus, the sequence $0 \rightarrow i^* M \rightarrow i^* N \rightarrow i^* P \rightarrow 0$ is exact, since each sequence of k -modules, $0 \rightarrow M_n \rightarrow N_n \rightarrow P_n \rightarrow 0$ is exact, and

certainly the sequence of 0 modules corresponding to $n = -1$ is exact. Now, since i^* is shown to be an exact functor, its left-derived functors $L_q i^*$ must be trivial for $q \geq 1$. The spectral sequence (3) is trivial in columns $q \geq 1$, and thus strongly converges

$$\operatorname{colim}_{\Delta S_+}^{(p)} i^* B_*^{\operatorname{sym}} A = H_p(\Delta S_+, B_*^{\operatorname{sym}^0} A) \Rightarrow HS_p(A). \quad \square$$

4.2 A more useful extension

Unfortunately, the ΔS_+ -module $B_*^{\operatorname{sym}^0} A$ does not seem to reflect the useful properties of ΔS_+ . In particular, the module corresponding to the object $[-1]$ is trivial in $B_*^{\operatorname{sym}^0} A$. Define a related functor which is non-trivial at the object $[-1]$:

Definition 30 For an associative, unital algebra, A , over a commutative ground ring k , define a functor $B_*^{\operatorname{sym}^+} A: \Delta S_+ \rightarrow k\text{-Mod}$ by

$$\begin{cases} B_n^{\operatorname{sym}^+} A &= A^{\otimes(n+1)}, \text{ for } n \geq 0 \\ B_{-1}^{\operatorname{sym}^+} A &= k, \end{cases}$$

Morphisms are mapped in the same way as for $B_*^{\operatorname{sym}^0} A$ with the exception that ι_n gets mapped to the inclusion $k \hookrightarrow A^{\otimes(n+1)}$ as algebras.

One very useful consequence of defining symmetric in terms of the homology groups $H_*(\mathcal{C}, F)$ for an appropriate category \mathcal{C} and functor F , is that the functor $H_*(\mathcal{C}, -)$, takes short exact sequences of \mathcal{C} -modules to long exact sequences in homology (see Gabriel–Zisman [11]). We use this fact to prove:

Theorem 31 For an associative, unital k -algebra A , $HS_*(A) = H_*(\Delta S_+, B_*^{\operatorname{sym}^+} A)$.

Proof Consider the following sequence of ΔS_+ -modules:

$$(4) \quad 0 \rightarrow B_*^{\operatorname{sym}^0} A \rightarrow B_*^{\operatorname{sym}^+} A \rightarrow Q \rightarrow 0$$

Here, Q is the quotient $B_*^{\operatorname{sym}^+} A / B_*^{\operatorname{sym}^0} A$, or in other words, Q is the functor $\Delta S_+ \rightarrow k\text{-Mod}$ taking $[-1] \mapsto k$ and $[n] \mapsto 0$ for all $n \geq 0$, which has no dependence on the choice of algebra A . Indeed, with $A = k$, we have the exact sequence:

$$0 \rightarrow B_*^{\operatorname{sym}^0} k \rightarrow B_*^{\operatorname{sym}^+} k \rightarrow Q \rightarrow 0$$

The corresponding long exact sequence in homology is rather easy to analyze. Using Lemma 29,

$$\begin{aligned} H_p(\Delta S_+, B_*^{\operatorname{sym}^0} k) &= HS_p(k) = \begin{cases} k, & p = 0 \\ 0, & p > 0 \end{cases} \\ &\cong H_*(N(\Delta S_+)) \cong H_*(\Delta S_+, B_*^{\operatorname{sym}^+} k) \end{aligned}$$

The last isomorphism is obtained in the same way as in [Corollary 19](#). Thus it follows that $H_*(\Delta S_+, Q) = 0$. The 5–lemma applied to the long exact sequence corresponding to diagram (4) then proves $H_n(\Delta S_+, B_*^{\text{sym}^0_+} A) \cong H_n(\Delta S_+, B_*^{\text{sym}^m_+} A)$ in every degree. [Lemma 29](#) provides the link to symmetric homology. \square

5 Tensor algebra decomposition and Fiedorowicz’s theorems

For a general k –algebra A , finding $HS_*(A)$ using the standard resolution is often too difficult. The multiplicative structure of A is hard to control. On the other hand, tensor algebras may be easier to deal with since their multiplicative structure is so clean. Indeed, tensor algebra arguments are also key in the proof of Fiedorowicz’s Theorem about the symmetric homology of group algebras.

5.1 Resolution of algebras using the 2–sided bar construction

Let $T: k\text{–Mod} \rightarrow k\text{–Alg}$ be the functor sending a k –module to the tensor algebra generated by that k –module. In other words, $TM = \bigoplus_{n \geq 0} M^{\otimes n}$. There is an algebra homomorphism $\theta_A: TA \rightarrow A$, defined by multiplying tensor factors, $\theta_A(a_0 \otimes a_1 \otimes \cdots \otimes a_k) = a_0 a_1 \cdots a_k$. By abuse of notation, let T also stand for the functor $k\text{–Mod} \rightarrow k\text{–Mod}$ obtained by composing with the forgetful functor $k\text{–Alg} \rightarrow k\text{–Mod}$. T becomes a monad with multiplication $\theta_{TA}: T^2 \rightarrow T$ and unit transformation $h: \text{id} \rightarrow T$, defined by sending the module M identically onto the summand corresponding to $n = 1$. The homomorphism θ_A then expresses A as a T –algebra.

Observe that May’s 2–sided bar construction $B_*(T, T, A)$ (see May [19, Chapter 9]) gives a resolution of A by tensor algebras:

$$(5) \quad 0 \longleftarrow A \xleftarrow{\theta_A} TA \xleftarrow{\theta_1} T^2 A \xleftarrow{\theta_2} \dots$$

The boundary maps θ_n are defined in the standard way: $\theta_n \stackrel{\text{def}}{=} \sum_{i=0}^n (-1)^i T^{n-i} \theta_{T^i A}$. If we denote by A_* the simplicial algebra consisting of A all degrees and whose faces and degeneracies are all id_A , then there is a strong deformation retract $B_*(T, T, A) \rightarrow A_*$ [19].

5.2 Monoid algebras

In the case that $A = k[M]$ for a monoid, M , we find a remarkable interpretation of the complex $C_*(\Delta S_+, B_*^{\text{sym}^m_+} TA)$ as an E_∞ –algebra. First, define a set-valued variant of the symmetric bar construction:

Definition 32 Let M be a monoid. Define a functor $B_*^{\text{sym}}M: \Delta S \rightarrow \mathbf{Sets}$ by:

$$B_n^{\text{sym}}M = B_*^{\text{sym}}M[n] \stackrel{\text{def}}{=} M^{n+1}, \quad (\text{set product})$$

$$B_*^{\text{sym}}M(\alpha): (m_0, \dots, m_n) \mapsto \alpha(m_0, \dots, m_n), \quad \text{for } \alpha \in \text{Mor } \Delta S.$$

where $\alpha: [n] \rightarrow [k]$ is represented in tensor notation, and evaluation at (m_0, \dots, m_n) is as in Definition 13.

The functor $B_*^{\text{sym}+}M: \Delta S_+ \rightarrow \mathbf{Sets}$ will be defined by the above as well as by setting $B_{-1}^{\text{sym}+}M \stackrel{\text{def}}{=} \emptyset$, and $B_*^{\text{sym}+}M \iota_n$ is the inclusion $\emptyset \hookrightarrow M^{n+1}$.

Regarding $N(- \setminus \Delta S_+)$ as a contravariant functor to **sets**, define $N(- \setminus \Delta S_+) \times_{\Delta S_+} B_*^{\text{sym}+}M$ as a coend construction similar to that in Section 1.2. It is immediate that this construction is a simplicial set whose homology computes $HS_*(k[M])$.

Now, in the context of monoids, the James construction is the appropriate analog of the the tensor algebra construction of Section 5.1. If M is a free monoid on a generating set X , that is, $M = JX_+$, then $k[M] = k[JX_+] = T(k[X])$ is the free tensor algebra over k on the set X with disjoint basepoint. In this case, we have the following:

Lemma 33

$$(6) \quad HS_*(k[JX_+]) \cong H_* \left(\coprod_{n \geq -1} \tilde{X}_n; k \right),$$

where

$$\tilde{X}_n = \begin{cases} N(\Delta S_+), & n = -1 \\ N([n] \setminus \Delta S_+) \times_{\Sigma_{n+1}^{\text{op}}} X^{n+1}, & n \geq 0 \end{cases}$$

Proof Using the simplicial set $N(- \setminus \Delta S_+) \times_{\Delta S_+} B_*^{\text{sym}+}JX_+$, a typical generator has the form,

$$(7) \quad \left(\begin{array}{ccccccc} & & & & & & [p] \\ & & & & & & \downarrow \alpha \\ & & & & & & [q_0] \\ & & & & \swarrow \beta_1 & & \\ & & & [q_1] & \longleftarrow \beta_2 & \cdots & \longleftarrow \beta_n \\ & & & [q_n] & & & \end{array} \right) \otimes u$$

where

$$u = \left(\bigotimes_{x \in X_0} x \right) \otimes \left(\bigotimes_{x \in X_1} x \right) \otimes \cdots \otimes \left(\bigotimes_{x \in X_p} x \right),$$

and X_0, X_1, \dots, X_p are finite ordered lists of elements of X . The idea is to “expand” the element u into a tensor of individual elements of X , and at the same time to get rid of the units that may be present. Indeed, each X_j may be thought of as an element of the set product X^{m_j} for some m_j . If $X_j = \emptyset$, then set $m_j = 0$. We use the convention that an empty tensor product is equal to 1_k (think of this as the disjoint basepoint appended to X), and say that the corresponding tensor factor is *trivial*. Now let $m = (\sum m_j) - 1$. Let $\pi: X^{m_0} \times X^{m_1} \times \dots \times X^{m_p} \rightarrow X^{m+1}$ be the evident isomorphism. Let $X_m = \pi(X_0, X_1, \dots, X_p)$.

Case 1 If u is non-trivial (that is, $X_m \neq \emptyset$), then construct the element

$$u' = \bigotimes_{x \in X_m} x$$

Next, construct a Δ -morphism $\zeta_u: [m] \rightarrow [p]$ as follows: For each j , ζ_u maps the points

$$\sum_{i=0}^{j-1} m_i, \left(\sum_{i=0}^{j-1} m_i \right) + 1, \dots, \left(\sum_{i=0}^j m_i \right) - 1 \mapsto j$$

Observe, $(\zeta_u)_*(u') = u$. Under ΔS_+ -equivalence, expression (7) is equivalent to

$$(8) \quad \left(\begin{array}{c} [m] \\ \swarrow \quad \downarrow \alpha \zeta_u \\ [q_n] \xleftarrow{\beta_n} \dots \xleftarrow{\beta_2} [q_1] \xleftarrow{\beta_1} [q_0] \end{array} \right) \otimes u'$$

The choice of u' and ζ_u is well-defined with respect to the ΔS_+ -equivalence up to isomorphism of $[m]$ (an element of Σ_{m+1}^{op}). This shows that any such non-trivial chain may be written uniquely as an element of $N([m] \setminus \Delta S_+) \times_k \Sigma_{m+1}^{op} X^{m+1}$. An example should clarify this process. If $u = (x_{i_0} \otimes x_{i_1}) \otimes 1 \otimes (x_{i_2} \otimes x_{i_3}) \in (T(k[X]))^{\otimes 3}$, then $u' = x_{i_0} \otimes x_{i_1} \otimes x_{i_2} \otimes x_{i_3}$ and $\zeta_u = x_0 x_1 \otimes 1 \otimes x_2$ (written in tensor notation – these x_i 's are formal variables, not elements of X). Clearly, $\zeta_u(u') = u$.

Case 2 If u is trivial, then $u = 1_k^{\otimes(p+1)}$, and we have

$$([q_n] \leftarrow \dots \leftarrow [q_0] \xleftarrow{\alpha} [p]) \otimes u \approx ([q_n] \leftarrow \dots \leftarrow [q_0] \xleftarrow{\iota_{q_0}} [-1]) \otimes 1_k.$$

This element can be identified uniquely with the chain $[q_n] \leftarrow \dots \leftarrow [q_0] \in N(\Delta S_+)$.

Thus, the isomorphism (6) is proven. Note that the total number of non-trivial tensor factors is preserved under ΔS morphisms. This shows that the differential respects the direct sum decomposition. □

Remark 34 This proof works the same using ΔS rather than ΔS_+ , by replacing the argument of Case 2 with the “identity trick”,

$$([q_n] \leftarrow \cdots \leftarrow [q_0] \xleftarrow{\alpha} [p]) \otimes u \approx ([q_n] \leftarrow \cdots \leftarrow [q_0] \xleftarrow{\text{id}} [q_0]) \otimes 1_k^{\otimes(q_0+1)}.$$

However the extra structure of ΔS_+ will be necessary to prove the E_∞ -structure needed for the results in the next section.

Breaking normal convention, let X^0 stand for the one-point set, $\{*\}$, (or $\{1_k\}$, which signifies the purpose of this set). Then the content of Lemma 33 can be more concisely,

$$(9) \quad HS_*(k[JX_+]) \cong \bigoplus_{n \geq -1} H_*(N([n] \setminus \Delta S_+) \times_{\Sigma_{n+1}^{\text{op}}} X^{n+1}; k)$$

Remark 35 It will be useful to work with a resolution of the monoid M by free monoids. We obtain an equivalence, $B_*(J, J, M) \rightarrow M_*$, where M_* is the simplicial monoid consisting of M in all degrees and whose faces and degeneracies are all id_M .

The next lemma will provide an essential link in the computation of $HS_*(k[M])$.

Lemma 36 Let $\mathcal{N} \stackrel{\text{def}}{=} N(- \setminus \Delta S_+) \times_{\Delta S_+} B_*^{\text{sym}+} J$. There is an equivalence of functors $\Theta: \mathcal{N} \xrightarrow{\cong} C_\infty$, where C_∞ is the monad associated to the E_∞ -operad, C_∞ . Moreover, Θ induces an equivalence

$$(10) \quad \Theta_*: B_*(\mathcal{N}, J, M) \rightarrow B_*(C_\infty, J, M)$$

Remark 37 \mathcal{N} itself turns out not to be a full-fledged operad, since it fails the right-unit condition. However, in order to prove Fiedorowicz’s Theorems below, all that we require is an equivalence of functors which preserves the simplicial maps of the bar construction.

Proof By definition,

$$\mathcal{N}X = \coprod_{n \geq 0} N([n-1] \setminus \Delta S_+) \times_{\Sigma_n^{\text{op}}} X^n,$$

which has been re-indexed to begin at $n = 0$. Let \mathcal{D} denote the operad with $\mathcal{D}(n) = E\Sigma_n$. Since \mathcal{D} is an E_∞ -operad, and the monads associated to different E_∞ -operads are equivalent, it suffices to show the equivalence, $\mathcal{N}X \simeq \mathcal{D}X$.

Note, the 0-chains (objects) of $N([n-1] \setminus \Delta S_+)$ are pairs $(\phi, g) \in \text{Mor}_{\Delta_+}([n-1], [m]) \times \Sigma_n^{\text{op}}$, so the Σ_n -action is free. Since $N([n-1] \setminus \Delta S_+)$ is a contractible

simplicial set, there is an equivalence induced on 0–chains by the map sending $(\phi, g) \mapsto g$:

$$B([n - 1] \setminus \Delta S_+) \xrightarrow{\cong} E\Sigma_n^{\text{op}},$$

where $B\mathcal{C}$ is the classifying space of a category \mathcal{C} . The map $g \rightarrow g^{-1}$ provides an equivalence $E\Sigma_n^{\text{op}} \simeq E\Sigma_n$.

These equivalences induce

$$(11) \quad \coprod_{n \geq 0} N([n - 1] \setminus \Delta S_+) \times_{\Sigma_n^{\text{op}}} X^n \simeq \coprod_{n \geq 0} E_*\Sigma_n^{\text{op}} \times_{\Sigma_n^{\text{op}}} X^n \simeq \coprod_{n \geq 0} E_*\Sigma_n \times_{\Sigma_n} X^n.$$

Equation (11) gives the equivalence of functors, $\mathcal{N} \rightarrow C_\infty$. For the second part of the lemma, we need only consider the 0–faces of each bar construction in (10), since the other face maps and all of the degeneracies only depend on the monad J and the monoid M . The 0–faces correspond to the transformations $\mathcal{N}J \rightarrow \mathcal{N}$ on the one hand, and $\mathcal{D}J \rightarrow \mathcal{D}$ on the other. Since the map $\mathcal{N}X \rightarrow \mathcal{D}X$ is natural in X , the diagram below commutes:

$$\begin{array}{ccc} \mathcal{N}JX & \xrightarrow{\cong} & \mathcal{D}JX \\ \downarrow & & \downarrow \\ \mathcal{N}X & \xrightarrow{\cong} & \mathcal{D}X \end{array}$$

Thus, the equivalence passes to the bar construction with no problem. □

5.3 Fiedorowicz’s theorems

The results of this section have been known for some time now, but not previously published. Fiedorowicz studied the symmetric homology of monoid algebras and group algebras [8]. All proofs of this section are either based on or inspired by the corresponding proofs given in that preprint. The following theorem is the first major result, which serves as a stepping stone to the group-algebra case.

Theorem 38 $HS_*(k[M]) \cong H_*(B(C_\infty, C_1, M); k)$, where C_1 is the little 1–cubes monad and C_∞ is the little ∞ –cubes monad.

Proof The arguments become clearer if we use homotopy colimits throughout. The link is provided by the following fact of homological algebra:

$$\text{colim}_{\mathcal{C}}^{(*)} F = H_*\left(\text{hocolim}_{\mathcal{C}} F\right)$$

In our case, we are interested in $\mathcal{C} = \Delta S_+$ and $F = B_*^{\text{sym}+} M$. By Section 5.1, we can replace M by its resolution $B_*(J, J, M)$. So there is an equivalence,

$$\text{hocolim}_{\Delta S_+} B_*^{\text{sym}+} M \simeq \text{hocolim}_{\Delta S_+} B_*^{\text{sym}+} B_*(J, J, M)$$

Now using May [19, Lemma 9.7],

$$\text{hocolim}_{\Delta S_+} B_*^{\text{sym}+} B_*(J, J, M) \cong B_*(\text{hocolim}_{\Delta S_+} B_*^{\text{sym}+} J, J, M) = B_*(\mathcal{N}, J, M)$$

By Lemma 36 the latter is equivalent to $B_*(C_\infty, J, M)$. The proof is completed by using the equivalence $J \simeq C_1$. □

As we turn our attention to group algebras, the following lemma will be necessary.

Lemma 39 *For any topological monoid, M , there is a natural equivalence,*

$$B(\Omega^\infty S^\infty, C_1, M) \simeq \Omega \Omega^\infty S^\infty(BM)$$

Proof Consider the bar construction $B_*(S, C_1, M)$, where S is suspension. May [19, Lemma 9.7] gives the isomorphism

$$B_*(\Omega^\infty S^\infty S, C_1, M) \cong \Omega^\infty S^\infty B_*(S, C_1, M).$$

Looping on both sides and taking geometric realization, we get

$$B(\Omega \Omega^\infty S^\infty S, C_1, M) \cong \Omega \Omega^\infty S^\infty B(S, C_1, M).$$

Now $B(\Omega \Omega^\infty S^\infty S, C_1, M) \simeq B(\Omega^\infty S^\infty, C_1, M)$. Finally, by Thomason [29] and Fiedorowicz [9], $B(S, C_1, M)$ is naturally equivalent to the bar construction BM , which yields the equivalence, $\Omega \Omega^\infty S^\infty B(S, C_1, M) \simeq \Omega \Omega^\infty S^\infty(BM)$. □

When the monoid is a group, we obtain the following important consequence.

Corollary 40 *If Γ is a group, then $HS_*(k[\Gamma]) \cong H_*(\Omega \Omega^\infty S^\infty(B\Gamma); k)$.*

Proof From Theorem 38, $HS_*(k[\Gamma]) \cong H_*(B(C_\infty, C_1, \Gamma); k)$. Since Γ is a group, $B(C_\infty, C_1, \Gamma)$ is grouplike, so $B(C_\infty, C_1, \Gamma) \xrightarrow{\simeq} B(\Omega^\infty S^\infty, C_1, \Gamma)$. Lemma 39 then applies. □

The analysis of $HS_*(k[\Gamma])$ then becomes interesting in its own right. We shall have more specific to say about $HS_1(k[\Gamma])$ in Section 11.5.

6 The category $\text{Epi}\Delta S$ and a smaller resolution

The complexes for computing $H_*(\Delta S, B_*^{\text{sym}} A)$ or $H_*(\Delta S_+, B_*^{\text{sym}+} A)$ are extremely large and unwieldy for computation. Fortunately, when the algebra A is equipped with an augmentation, $\epsilon: A \rightarrow k$, we may use a more manageable subcomplex of $C_*(\Delta S_+, B_*^{\text{sym}+} A)$, related to the subcategory of ΔS_+ consisting only of epimorphisms.

6.1 Basic and reduced tensors

Recall, if A has an augmentation ϵ , then there is an augmentation ideal I and the exact sequence $0 \rightarrow I \rightarrow A \xrightarrow{\epsilon} k \rightarrow 0$ splits as k -modules. So $A \cong I \oplus k$, and every $x \in A$ can be decomposed uniquely as $x = a + \lambda$ for some $a \in I, \lambda \in k$.

Definition 41 Define $B_{-1, \emptyset} A = k$. For $n \geq 0$, if $J \subseteq [n]$, define

$$B_{n, J} A \stackrel{\text{def}}{=} B_0^J \otimes B_1^J \otimes \cdots \otimes B_n^J, \quad \text{where } B_j^J = \begin{cases} I & \text{if } j \in J \\ k & \text{if } j \notin J \end{cases}$$

Remark 42 For each $n \geq -1$, there is a direct sum decomposition of k -modules, $B_n^{\text{sym}+} A \cong \bigoplus_{J \subseteq [n]} B_{n, J} A$.

Definition 43 A *basic tensor* is any tensor $w_0 \otimes w_1 \otimes \cdots \otimes w_n$, where each w_j is in I or is equal to the unit of A . Call a tensor factor w_j *trivial* if it is the unit of A . If all factors of a basic tensor are trivial, then the tensor is called *trivial*, and if no factors are trivial, the tensor is called *reduced*.

For a basic tensor $Y \in B_n^{\text{sym}+} A$, we shall define a map $\delta_Y \in \text{Mor } \Delta_+$ as follows: If Y is trivial, let $\delta_Y = \iota_n$. Otherwise, Y has $\bar{n} + 1$ non-trivial factors for some $\bar{n} \geq 0$. Define $\delta_Y: [\bar{n}] \rightarrow [n]$ to be the unique injective map that sends each point $0, 1, \dots, \bar{n}$ to a point $p \in [n]$ such that Y is non-trivial at factor p . Let \bar{Y} be the tensor obtained from Y by omitting all trivial factors if such exist, or $\bar{Y} = 1_k$ if Y is trivial. Note, \bar{Y} is the unique basic tensor such that $(\delta_Y)_*(\bar{Y}) = Y$.

Proposition 44 Any chain

$$([q_n] \leftarrow [q_{n-1}] \leftarrow \cdots \leftarrow [q_0] \leftarrow [q]) \otimes Y \in k[N(- \setminus \Delta S_+)] \otimes_{\Delta S_+} B_*^{\text{sym}+} A,$$

where Y is a basic tensor, is equivalent to a chain

$$([q_n] \leftarrow [q_{n-1}] \leftarrow \cdots \leftarrow [q_0] \leftarrow [\bar{q}]) \otimes \bar{Y},$$

where either \bar{Y} is reduced or $\bar{Y} = 1_k$ and $\bar{q} = -1$.

Proof Let δ_Y and \bar{Y} be defined as above, and let $[\bar{q}]$ be the domain of δ_Y . Then $Y = (\delta_Y)_*(\bar{Y})$, and

$$\left([q_n] \leftarrow [q_{n-1}] \leftarrow \cdots \leftarrow [q_0] \xleftarrow{\phi} [q] \right) \otimes Y \approx \left([q_n] \leftarrow [q_{n-1}] \leftarrow \cdots \leftarrow [q_0] \xleftarrow{\phi \delta_Y} [\bar{q}] \right) \otimes \bar{Y} \quad \square$$

6.2 Reducing to epimorphisms

We now turn our attention to the morphisms in the chains. Our goal is to reduce to those chains that involve only epimorphisms.

Definition 45 Let \mathcal{C} be a category. The category $\text{Epi}\mathcal{C}$ (resp. $\text{Mono}\mathcal{C}$) is the subcategory of \mathcal{C} consisting of the same objects as \mathcal{C} and only those morphisms $f \in \text{Mor}\mathcal{C}$ that are epic (resp. monic). The set of morphisms of $\text{Epi}\mathcal{C}$ from X to Y may be denoted $\text{Epi}_{\mathcal{C}}(X, Y)$. Similarly, the set of morphisms of $\text{Mono}\mathcal{C}$ from X to Y may be denoted $\text{Mono}_{\mathcal{C}}(X, Y)$.

Remark 46 A morphism $\alpha = (\phi, g) \in \text{Mor}\Delta S_+$ is epic (resp. monic) if and only if ϕ is epic (resp. monic) as morphism in Δ_+ .

Proposition 47 Any morphism $\alpha \in \text{Mor}\Delta S_+$ decomposes uniquely as $(\eta, \text{id}) \circ \gamma$, where $\gamma \in \text{Mor}(\text{Epi}\Delta S_+)$ and $\eta \in \text{Mor}(\text{Mono}\Delta_+)$.

Proof Suppose α has source $[-1]$ and target $[n]$. Then $\alpha = \iota_n$ is the only possibility, and this decomposes as $\iota_n \circ \text{id}_{[-1]}$. Now suppose the source of α is $[p]$ for some $p \geq 0$. Write $\alpha = (\phi, g)$, with $\phi \in \text{Mor}_{\Delta}([p], [r])$ and $g \in \Sigma_{p+1}^{\text{op}}$. Suppose ϕ hits $q + 1$ distinct points in $[r]$. Then there is a morphism $\pi: [p] \rightarrow [q]$ induced by ϕ by maintaining the order of the points hit. Let η be the obvious order-preserving monomorphism $[q] \rightarrow [r]$ so that $\eta\pi = \phi$ as morphisms in Δ . To get the required decomposition in ΔS , use: $\alpha = (\eta, \text{id}) \circ (\pi, g)$.

Now, if $(\xi, \text{id}) \circ (\psi, h)$ is also a decomposition of α , with ξ monic and ψ epic, then $(\xi, \text{id}) \circ (\psi, h) = (\eta, \text{id}) \circ (\pi, g)$ implies $(\xi\psi, g^{-1}h) = (\eta\pi, \text{id})$, proving $g = h$. Uniqueness will then follow from uniqueness of such decompositions entirely within the category Δ . The latter follows from Loday [16, Theorem B.2]. □

This decomposition will be written

$$[p] \longrightarrow [r] = [p] \twoheadrightarrow \text{im}([p] \longrightarrow [r]) \hookrightarrow [r],$$

and we call the *epimorphism construction*, the rule

$$\mathcal{E}_p([p] \rightarrow [r]) = [p] \twoheadrightarrow \text{im}([p] \rightarrow [r]).$$

Proposition 48 *The epimorphism construction is a functor $\mathcal{E}_p: [p] \setminus \Delta S_+ \rightarrow [p] \setminus \text{Epi} \Delta S_+$.*

Proof If $[p] \xrightarrow{\beta} [r_1] \xrightarrow{\alpha} [r_2]$, then there is an induced map

$$\text{im}([p] \rightarrow [r_1]) \xrightarrow{\bar{\alpha}} \text{im}([p] \rightarrow [r_2])$$

making the following diagram commute:

(12)

$$\begin{array}{ccccc}
 [r_1] & \xrightarrow{\alpha} & [r_2] & & \\
 & \swarrow \beta & \nearrow \alpha\beta & & \\
 & [p] & & & \\
 & \swarrow \pi_1 & \searrow \pi_2 & & \\
 \text{im}([p] \rightarrow [r_1]) & \xrightarrow{\bar{\alpha}} & \text{im}([p] \rightarrow [r_2]) & & \\
 \uparrow \eta_1 & & \uparrow \eta_2 & & \\
 \text{im}([p] \rightarrow [r_1]) & & \text{im}([p] \rightarrow [r_2]) & &
 \end{array}$$

$\bar{\alpha}$ is the epimorphism induced from the map $\alpha\eta_1$. Furthermore, for morphisms

$$[p] \rightarrow [r_1] \xrightarrow{\alpha_1} [r_2] \xrightarrow{\alpha_2} [r_3],$$

we have $\overline{\alpha_2\alpha_1} = \bar{\alpha}_2 \circ \bar{\alpha}_1$. □

Remark 49 If $\alpha: [p] \rightarrow [r]$ is an epimorphism of ΔS_+ , then $\mathcal{E}_p(\alpha) = \alpha$.

Define a variant of the symmetric bar construction using $\text{Epi} \Delta S$:

Definition 50 $B_*^{\text{sym}+} I: \text{Epi} \Delta S_+ \rightarrow k\text{-Mod}$ is the functor defined by

$$\begin{cases} B_n^{\text{sym}+} I = I^{\otimes(n+1)}, & n \geq 0, \\ B_{-1}^{\text{sym}+} I = k, \end{cases}$$

and

$$B_*^{\text{sym}+} I(\alpha): (a_0 \otimes a_1 \otimes \cdots \otimes a_n) \mapsto \alpha(a_0, \dots, a_n), \text{ for } \alpha \in \text{Mor}(\text{Epi} \Delta S_+)$$

This definition makes sense since I is an ideal, and α is required to be epimorphic. Note, the simple tensors $w_0 \otimes \cdots \otimes w_n$ in $B_n^{\text{sym}+} I$ are by definition reduced. Consider the simplicial k -module:

$$(13) \quad C_*(\text{Epi} \Delta S_+, B_*^{\text{sym}+} I) = k[N(- \setminus \text{Epi} \Delta S_+)] \otimes_{\text{Epi} \Delta S_+} B_*^{\text{sym}+} I$$

There is an obvious inclusion

$$f: C_*(\text{Epi}\Delta S_+, B_*^{\text{sym}+} I) \longrightarrow C_*(\Delta S_+, B_*^{\text{sym}+} A).$$

Define a chain map g in the opposite direction as follows. First, by [Proposition 44](#) and observations above, we only need to define g on the chains

$$([q_n] \longleftarrow \cdots \longleftarrow [q_0] \longleftarrow [q]) \otimes Y$$

where Y is reduced (or $Y = 1_k$). In this case, define component maps $g(q) = N(\mathcal{E}_q) \otimes \text{id}$. A priori, this definition is well-defined only when the tensor product is over k . We would like to assemble the maps $g(q)$ into a chain map g . In order to do this, we must show that the maps are compatible under ΔS_+ -equivalence.

Suppose

$$v = ([p_n] \longleftarrow \cdots \longleftarrow [p_0] \xleftarrow{\phi\psi} [p]) \otimes Z$$

and

$$v' = ([p_n] \longleftarrow \cdots \longleftarrow [p_0] \xleftarrow{\phi} [q]) \otimes \psi_*(Z),$$

where $\psi \in \text{Mor}_{\Delta S_+}([p], [q])$. If Z is a basic tensor, then so is $\psi_*(Z)$. In order to apply g to v or v' , each must first be put into a reduced form.

Case 1 Suppose Z is trivial. Then v and v' both reduce to

$$([p_n] \longleftarrow \cdots \longleftarrow [p_0] \longleftarrow [-1]) \otimes 1,$$

hence $g(v) = g(v')$.

Case 2 Suppose Z is non-trivial. For the sake of clean notation, let $W = \psi_*(Z)$. Construct $\delta_Z, \bar{Z}, \delta_W$ and \bar{W} such that $Z = (\delta_Z)_*(\bar{Z})$ and $W = (\delta_W)_*(\bar{W})$ as in [Proposition 44](#), and reduce both chains:

$$(14) \quad \begin{array}{ccc} ([p_n] \longleftarrow \cdots \longleftarrow [p_0] \xleftarrow{\phi\psi} [p]) \otimes Z & \approx & ([p_n] \longleftarrow \cdots \longleftarrow [p_0] \xleftarrow{\phi} [q]) \otimes W \\ \text{reduce} \downarrow \approx & & \text{reduce} \downarrow \approx \\ ([p_n] \longleftarrow \cdots \longleftarrow [p_0] \xleftarrow{\phi\psi\delta_Z} [\bar{p}]) \otimes \bar{Z} & & ([p_n] \longleftarrow \cdots \longleftarrow [p_0] \xleftarrow{\phi\delta_W} [\bar{q}]) \otimes \bar{W} \end{array}$$

Observe that number of distinct points hit by $\psi\delta_Z$ is exactly $\bar{q} + 1$; indeed, $W = \psi_*(Z)$ has $\bar{q} + 1$ non-trivial factors. Thus, $[\bar{q}] = \text{im}([\bar{p}] \rightarrow [q])$. Now, [Proposition 47](#) implies that

there is precisely one ΔS -epimorphism $\gamma: [\bar{p}] \rightarrow [\bar{q}]$ making Diagram (15) commute.

$$(15) \quad \begin{array}{ccc} [p] & \xrightarrow{\psi} & [q] \\ \delta_Z \uparrow & \nearrow \psi \delta_Z & \uparrow \delta_W \\ [\bar{p}] & \xrightarrow{\quad} & [\bar{q}] \\ & \searrow \gamma & \\ & & \text{im}([\bar{p}] \rightarrow [q]) \end{array}$$

That is to say, there exists an epimorphism γ such that $\gamma_*(\bar{Z}) = \bar{W}$ and $\psi \delta_Z = \delta_W \gamma$. So we may replace the first morphism of the chain in the lower left of Diagram (14) with $\phi \delta_W \gamma$. Then when we apply g to the chain, the first morphism becomes $\mathcal{E}_{\bar{p}}(\phi \delta_W \gamma) = \mathcal{E}_{\bar{p}}(\phi \delta_W) \circ \gamma$, since γ is epic. Let $\pi \stackrel{\text{def}}{=} \mathcal{E}_{\bar{p}}(\phi \delta_W)$. Then the result of applying g to each side of Diagram (14) is shown below:

$$\begin{array}{c} ([p_n] \leftarrow \dots \leftarrow [p_0] \xleftarrow{\phi \psi \delta_Z} [\bar{p}]) \otimes \bar{Z} \\ \downarrow g(\bar{p}) \\ (\text{im}([\bar{p}] \rightarrow [p_n]) \leftarrow \dots \leftarrow \text{im}([\bar{p}] \rightarrow [p_0]) \xleftarrow{\pi \gamma} [\bar{p}]) \otimes \bar{Z} \\ \\ ([p_n] \leftarrow \dots \leftarrow [p_0] \xleftarrow{\phi \delta_W} [\bar{q}]) \otimes \bar{W} \\ \downarrow g(\bar{q}) \\ (\text{im}([\bar{q}] \rightarrow [p_n]) \leftarrow \dots \leftarrow \text{im}([\bar{q}] \rightarrow [p_0]) \xleftarrow{\pi} [\bar{q}]) \otimes \bar{W} \end{array}$$

Observe that there is equality of objects and morphisms up to the morphisms $\pi \gamma$ on the left and π on the right. Since γ is epic, the $\text{Epi} \Delta S_+$ -equivalence allows us to transport the morphism γ to the right of the tensor, showing that $g(v) \approx g(v')$, hence g is well-defined.

Proposition 51 *If A has augmentation ideal I , then*

$$HS_*(A) = H_*(\text{Epi} \Delta S_+, B_*^{\text{sym}+} I) = H_*(k[N(- \setminus \text{Epi} \Delta S_+)] \otimes_{\text{Epi} \Delta S_+} B_*^{\text{sym}+} I; k).$$

Proof Clearly $gf = \text{id}$, where f is the inclusion mentioned above. It remains to show that $fg \simeq \text{id}$. Assume Y is a basic tensor in $B_q^{\text{sym}+} I$. Define maps $h_j^{(n)}$ as follows:

$$h_j^{(n)}: ([q_n] \leftarrow \cdots \leftarrow [q_0] \leftarrow [q]) \otimes Y \mapsto ([q_n] \leftarrow \cdots \leftarrow [q_j] \leftarrow \text{im}([q] \rightarrow [q_j]) \leftarrow \cdots \leftarrow \text{im}([q] \rightarrow [q_0]) \leftarrow [q]) \otimes Y.$$

$h_j^{(n)}$ is well-defined by the functorial properties of the epimorphism construction, and a routine, but tedious, verification shows that h defines a presimplicial homotopy from fg to id . □

6.3 Reduced symmetric homology

Denote by $B_*^{\text{sym}} I$, the restriction of $B_*^{\text{sym}+} I$ to ΔS . That is, $B_n^{\text{sym}} I \stackrel{\text{def}}{=} I^{\otimes(n+1)}$ for all $n \geq 0$. Then there is a splitting of the chain complex,

$$C_*(\text{Epi}\Delta S_+, B_n^{\text{sym}+} I) = C_*(\text{Epi}\Delta S, B_n^{\text{sym}} I) \oplus k[N(*)],$$

where $*$ is the full subcategory of $\text{Epi}\Delta S_+$ consisting only of the object $[-1]$. Indeed, since there are no epimorphisms $[-1] \rightarrow [n]$ or $[n] \rightarrow [-1]$ for $n \geq 0$, we may think of $[-1]$ as a disconnected basepoint. Hence, we have $HS_*(A) \cong H_*(\text{Epi}\Delta S, B_n^{\text{sym}} I) \oplus k_0$, where k_0 is the graded k -module consisting of k concentrated in degree 0.

Definition 52 The *reduced symmetric homology* of A is defined,

$$\tilde{H}S_*(A) \stackrel{\text{def}}{=} H_*(\text{Epi}\Delta S, B_n^{\text{sym}} I).$$

Remark 53 Reduction to epimorphisms seems to hinge on the property that A has an augmentation ideal. This condition may be lifted (as Richter conjectures), if it can be shown that $N(\text{Epi}\Delta S)$ is contractible. As partial progress along these lines, it can be shown that $N(\text{Epi}\Delta S)$ is simply-connected.

7 Interpretation of $HS_*(A)$ in terms of group homology

$\text{Epi}\Delta S$ is an E-I-A-category, that is, a category in which all endomorphisms are isomorphisms and also each isomorphism class of an object has only a single member (all isomorphism are automorphisms). In this section, we use results of Słomińska [27] about E-I-A-categories to provide an interpretation of symmetric homology as the homology of products of symmetric groups with coefficients in certain modules. As a corollary, we find that when the ground ring is a field of characteristic 0, $HS_*(A)$ can be computed as the coinvariants of a group action.

Fix a unital associative algebra A over commutative ground ring k , and assume A has an augmentation, with augmentation ideal I . By Proposition 51 and Definition 52,

$$\tilde{H}S_*(A) = H_*(\text{Epi}\Delta S, B_*^{\text{sym}} I) = \text{colim}_{\text{Epi}\Delta S}^{(*)} B_*^{\text{sym}} I,$$

where $\text{colim}^{(*)}$ represents the left derived functors of the colimit.

Let \mathcal{S}_0 be the category whose objects are ordered tuples of non-negative integers, $(q_n < q_{n-1} < \dots < q_0)$, with a unique morphism $(q_n < \dots < q_0) \rightarrow (q'_n < \dots < q'_0)$ if $\{q'_n, \dots, q'_0\} \subseteq \{q_n, \dots, q_0\}$. Let Aut be the functor (\mathcal{S}_0 -group) defined by

$$\text{Aut}(q_n < \dots < q_0) = \prod_{i=0}^n \Sigma_{q_i+1}^{\text{op}} = \prod_{i=0}^n \text{Aut}_{\text{Epi}S}([q_i])$$

Morphisms of \mathcal{S}_0 map under Aut to the evident projections. Let $\mathcal{E}pi$ be the functor $\mathcal{S}_0 \rightarrow \mathbf{Set}$ defined by

$$\mathcal{E}pi(q_n < \dots < q_0) = \begin{cases} \{q_0\}, & \text{if } n = 0 \\ \prod_{i=1}^n \text{Epi}_{\Delta S}([q_{i-1}], [q_i]), & \text{if } n \geq 1 \end{cases}$$

If $\{q'_n, \dots, q'_0\} \subseteq \{q_n, \dots, q_0\}$, then the set map

$$\mathcal{E}pi(q_n < \dots < q_0) \longrightarrow \mathcal{E}pi(q'_n < \dots < q'_0)$$

is defined by the appropriate projections and compositions of morphisms, if $n' \geq 1$, and by the unique set map into $\{q'_0\}$ if $n = 0$.

There is an action of $\text{Aut}(q_n < \dots < q_0)$ on $\mathcal{E}pi(q_n < \dots < q_0)$ given by

$$\mu: (g_n, \dots, g_0) \otimes (\phi_n, \dots, \phi_1) \mapsto (g_n \phi_n g_{n-1}^{-1}, \dots, g_1 \phi_1 g_0^{-1})$$

The maps induced by $(q_n < \dots < q_0) \rightarrow (q'_n < \dots < q'_0)$ are equivariant with respect to this action in the sense that the following diagram commutes:

$$(16) \quad \begin{array}{ccc} (g_n, \dots, g_0) \otimes (\phi_n, \dots, \phi_1) & \xrightarrow{\mu} & (g_n \phi_n g_{n-1}^{-1}, \dots, g_1 \phi_1 g_0^{-1}) \\ \downarrow & & \downarrow \\ (g'_n, \dots, g'_0) \otimes (\phi'_n, \dots, \phi'_1) & \xrightarrow{\mu} & (g'_n \phi'_n (g'_{n-1})^{-1}, \dots, g'_1 \phi'_1 (g'_0)^{-1}) \end{array}$$

Now, given the augmentation ideal I of A , define a related functor, $\mathcal{E}pi_I: \mathcal{S}_0 \rightarrow k\text{-Mod}$ by

$$(q_n < \dots < q_0) \mapsto k[\mathcal{E}pi(q_n < \dots < q_0)] \otimes_k B_{q_0}^{\text{sym}} I$$

The effect of $\mathcal{E}pi_I$ on morphisms of \mathcal{S}_0 requires some explanation. If

$$(q_n < \dots < q_0) \rightarrow (q'_{n'} < \dots < q'_0)$$

is a morphism of \mathcal{S}_0 , then the map

$$\mathcal{E}pi_I(q_n < \dots < q_0) \rightarrow \mathcal{E}pi_I(q'_{n'} < \dots < q'_0)$$

is defined on generators $(\phi_n, \dots, \phi_1) \otimes x$ by applying the map

$$\mathcal{E}pi_I((q_n < \dots < q_0) \rightarrow (q'_{n'} < \dots < q'_0)) \otimes \phi_r \cdots \phi_1,$$

if $q'_0 = q_r$. That is, morphisms “on the left” are thrown out; morphisms “in the middle” are composed; and morphisms “on the right” are applied to the element of $B_{q_0}^{\text{sym}} I$ to produce an element of $B_{q'_0}^{\text{sym}} I$. Each $k[\mathcal{E}pi(q_n < \dots < q_0)] \otimes B_{q_0}^{\text{sym}} I$ becomes an $\text{Aut}(q_n < \dots < q_0)$ -module in the obvious way. Diagram (16) implies that there are well-defined functors, $H_q(\text{Aut}-, \mathcal{E}pi_I-): \mathcal{S}_0 \rightarrow k\text{-Mod}$.

Theorem 54 *With Aut and $\mathcal{E}pi_I$ defined as above, there is a spectral sequence*

$$\text{colim}_{\mathcal{S}_0}^{(p)} H_q(\text{Aut}-, \mathcal{E}pi_I-) \Rightarrow \tilde{H}S_{p+q}(A)$$

Proof This is essentially Słomińska [27, Proposition 1.6], reinterpreted in terms of the derived functors of colim . Compare also [27, Proposition 1.8], where variance of the functor N is opposite, and cohomology is computed. Let Q stand for $(q_n < \dots < q_0)$. For each Q , let $\mathcal{G}Q$ be the category consisting of a single object whose endomorphism set is the group $\text{Aut}(Q)$. Then we obtain

$$\begin{aligned} \text{colim}_{Q \in \mathcal{S}_0}^{(p)} H_q(\text{Aut}(Q), \mathcal{E}pi_I(Q)) &= \text{colim}_{Q \in \mathcal{S}_0}^{(p)} \text{colim}_{\mathcal{G}Q}^{(q)} (\mathcal{E}pi_I(Q)) \\ &\implies \text{colim}_{\text{Epi}\Delta S}^{(p+q)} B_*^{\text{sym}} I \cong \tilde{H}S_{p+q} A \end{aligned}$$

which completes the proof. □

7.1 Implications in characteristic 0

If k is a field of characteristic 0, then for any finite group G and kG -module M , $H_q(G, M) = 0$ for all $q > 0$ (see Brown [1], for example). Thus, the spectral sequence of Theorem 54, would collapse giving the following

Corollary 55 *If k is a field of characteristic 0, and A is a k -algebra, then*

$$\tilde{H}S_p(A) = \text{colim}_{Q \in \mathcal{S}_0}^{(p)} \mathcal{E}pi_I(Q)_{\text{Aut}(Q)},$$

where $\mathcal{E}pi_I(Q)_{Aut(Q)}$ is the k -space of coinvariants of $\mathcal{E}pi_I(Q)$ under the action of $Aut(Q)$.

8 A second spectral sequence

In Section 7, a spectral sequence was found that essentially separates the symmetric group elements from the morphisms comprising the chains of the resolution $C_*(Epi\Delta S, B_*^{sym}I)$. What remains is still rather cumbersome. The following can be viewed as an attempt to separate the combinatorial structure inherent in the category $Epi\Delta S$ from the more complicated structure generally found in $B_*^{sym}I$. Indeed, the E^1 term of our spectral sequence isolates the multiplicative structure of $B_*^{sym}I$ into the action of the differential.

8.1 Filtering by degree

Consider a filtration \mathcal{F}_* of $C_*(Epi\Delta S, B_*^{sym}I)$ given by

$$\mathcal{F}_p C_q(Epi\Delta S, B_*^{sym}I) = \bigoplus_{p \geq m_0 \geq \dots \geq m_q} k \left[\prod_{i=1}^q Epi_{\Delta S}([m_{i-1}], [m_i]) \right] \otimes B_{m_0}^{sym}I$$

\mathcal{F}_* filters the complex by the length of $x \in B_*^{sym}I$. Here, length refers to the number of tensor components of x . Tensors of $B_{m_0}^{sym}I$ are considered to have length $m_0 + 1$. The only face map that can potentially change the length of x is d_0 , and since all morphisms are epic, d_0 can only reduce the length. Thus, \mathcal{F}_* is compatible with the differential. The filtration quotients are easily described. The non-zero chains are those corresponding to $p = m_0$. In other words, the length of x is exactly $p + 1$. E^0 splits into a direct sum based on the product of x_i 's in $u = (x_0, \dots, x_p) \in X^{p+1}$. For $u \in X^{p+1}$, let P_u be the set of all distinct permutations of u . Then,

$$E_{p,q}^0 = \bigoplus_{u \in X^{p+1} / \Sigma_{p+1}} \bigoplus_{w \in P_u} \left(\bigoplus_{p = m_0 \geq \dots \geq m_q} k \left[\prod_{i=1}^q Epi_{\Delta S}([m_{i-1}], [m_i]) \right] \otimes w \right).$$

8.2 The Categories $\tilde{\mathcal{S}}_p, \tilde{\mathcal{S}}'_p, \mathcal{S}_p$ and \mathcal{S}'_p

Before proceeding with the main theorem of this section, we must define four related categories. In the definitions that follow, let $\{z_0, z_1, z_2, \dots, z_p\}$ be a set of formal non-commuting indeterminates.

Definition 56 $\tilde{\mathcal{S}}_p$ is the category with objects formal tensor products $Z_0 \otimes \cdots \otimes Z_s$, where each Z_i is a non-empty product of z_i 's, and every one of z_0, z_1, \dots, z_p occurs exactly once in the tensor product. There is a unique morphism $Z_0 \otimes \cdots \otimes Z_s \rightarrow Z'_0 \otimes \cdots \otimes Z'_t$, if and only if the tensor factors of the latter are products of the factors of the former in some order. In such a case, there is a unique $\beta \in \text{Epi}\Delta S$ so that $\beta_*(Z_0 \otimes \cdots \otimes Z_s) = Z'_0 \otimes \cdots \otimes Z'_t$.

$\tilde{\mathcal{S}}_p$ has initial objects $\sigma_*(z_0 \otimes z_1 \otimes \cdots \otimes z_p)$, for $\sigma \in \Sigma_{p+1}^{\text{op}}$, so $N\tilde{\mathcal{S}}_p$ is a contractible complex. Let $\tilde{\mathcal{S}}'_p$ be the full subcategory of $\tilde{\mathcal{S}}_p$ with all objects $\sigma_*(z_0 \otimes \cdots \otimes z_p)$ deleted.

Let \mathcal{S}_p be a skeletal category equivalent to $\tilde{\mathcal{S}}_p$. In fact, we may make \mathcal{S}_p the quotient category, identifying each object $Z_0 \otimes \cdots \otimes Z_s$ with any permutation of its tensor factors, and identifying morphisms ϕ and ψ if their source and target are equivalent. This category has nerve $N\mathcal{S}_p$ homotopy-equivalent to $N\tilde{\mathcal{S}}_p$. Now, \mathcal{S}_p is a poset with unique initial object, $z_0 \otimes \cdots \otimes z_p$. Let \mathcal{S}'_p be the full subcategory (subposet) of \mathcal{S}_p obtained by deleting the object $z_0 \otimes \cdots \otimes z_p$. Clearly, \mathcal{S}'_p is a skeletal category equivalent to $\tilde{\mathcal{S}}'_p$.

8.3 Main theorem

Theorem 57 *There is spectral sequence converging (weakly) to $\tilde{H}S_*(A)$ with*

$$E^1_{p,q} \cong \bigoplus_{u \in X^{p+1} / \Sigma_{p+1}} H_{p+q}(EG_u \times_{G_u} |N\mathcal{S}_p / N\mathcal{S}'_p|; k),$$

where G_u is the isotropy subgroup for the chosen representative of $u \in X^{p+1} / \Sigma_{p+1}$.

Recall, for a group G , right G -space X , and left G -space Y , $X \times_G Y$ denotes the *equivariant half-smash product*. If $*$ is a chosen basepoint for Y having trivial G -action, then $X \times_G Y \stackrel{\text{def}}{=} (X \times_G Y) / (X \times_G *) = X \times Y / \approx$, with equivalence relation defined by $(x.g, y) \approx (x, g.y)$ and $(x, *) \approx (x', *)$ for all $x, x' \in X$, $y \in Y$ and $g \in G$ (see Bruner, May, McClure and Steinberger [3]). In our case, X is of the form EG , with canonical underlying complex E_*G , equipped with a right G -action, $(g_0, g_1, \dots, g_n).g = (g_0, g_1, \dots, g_n g)$.

Observe, both $N\tilde{\mathcal{S}}_p$ and $N\tilde{\mathcal{S}}'_p$ carry a left Σ_{p+1} -action (hence also a G_u -action). The action is defined on 0-chains $Z_0 \otimes \cdots \otimes Z_s$ by permutation of the individual indeterminates, z_0, z_1, \dots, z_p . This action extends to n -chains in the straightforward manner.

Define for each $u \in X^{p+1}/\Sigma_{p+1}$, the following subcomplex of $E_{p,q}^0$:

$$\mathcal{M}_u \stackrel{\text{def}}{=} \bigoplus_{w \in P_u} \left(\bigoplus_{p=m_0 \geq \dots \geq m_q} k \left[\prod_{i=1}^q \text{Epi}_{\Delta S}([m_{i-1}], [m_i]) \right] \otimes w \right)$$

Lemma 58 *There is a chain-isomorphism, $(N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p)/G_u \xrightarrow{\cong} \mathcal{M}_u$.*

Proof Let C_j denote objects of $\tilde{\mathcal{S}}_p$. As above, we may view each C_j as a morphism of ΔS . By abuse of notation, let C_j also represent a 0-cell of $N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p$. Denote the chosen representative of u again by u (We view $u = (x_{i_0}, x_{i_1}, \dots, x_{i_p}) \in X^{p+1}/\Sigma_{p+1}$ as represented by a $(p+1)$ -tuple whose indices are in non-decreasing order).

First define a map $N\tilde{\mathcal{S}}_p \rightarrow \mathcal{M}_u$ on n -cells by

$$(17) \quad \alpha_*: \left(C_q \xleftarrow{\phi_q} \dots \xleftarrow{\phi_1} C_0 \right) \mapsto (\phi_q, \dots, \phi_1) \otimes C_0(u).$$

The notation $C_0(u)$ is used in place of the more correct $(C_0)_*(u)$ in order to avoid clutter. I claim α_* factors through $N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p$. Indeed, if

$$C_q \xleftarrow{\phi_q} \dots \xleftarrow{\phi_1} C_0 \in N\tilde{\mathcal{S}}'_p,$$

then we cannot have $C_0 = \sigma(z_0 \otimes \dots \otimes z_p)$ for any symmetric group element σ . That is, C_0 , viewed as a morphism, must be strictly epic. Then the length of $C_0(u)$ is strictly less than the length of u , which would make $(\phi_q, \dots, \phi_1) \otimes C_0(u)$ trivial in $E_{p,q}^0$.

The map α_* then factors through $(N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p)/G_u$, since if $\gamma \in G_u$, then γ corresponds to an automorphism $g \in \Sigma_{p+1}^{\text{op}}$, and by definition, we have

$$\begin{aligned} \gamma \cdot \left(C_q \xleftarrow{\phi_q} \dots \xleftarrow{\phi_1} C_0 \right) &= \left(C_q g \xleftarrow{\phi_q} \dots \xleftarrow{\phi_1} C_0 g \right) \mapsto \\ &(\phi_q, \dots, \phi_1) \otimes C_0(g(u)) = (\phi_q, \dots, \phi_1) \otimes C_0(u) \end{aligned}$$

(Note, $g(u) = u$ follows from the fact that $\gamma \in G_u$, the isotropy subgroup for u).

For the map in the opposite direction, consider $(\phi_q, \dots, \phi_1) \otimes w$ for $w \in P_u$. Let $t \in \Sigma_{p+1}^{\text{op}}$ so that $w = t(u)$. Define a map sending

$$(18) \quad \beta_*: (\phi_q, \dots, \phi_1) \otimes w \mapsto \left((\phi_q \dots \phi_1 t) \xleftarrow{\phi_q} \dots \xleftarrow{\phi_2} \phi_1 t \xleftarrow{\phi_1} t \right)$$

We must check that the definition of β_* does not depend on choice of t . Indeed, if $w = s(u)$ also, then $u = s^{-1}t(u)$, hence $s^{-1}t \in G_u^{\text{op}}$. Thus

$$\begin{aligned} \left((\phi_q \cdots \phi_1 s) \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_2} \phi_1 s \xleftarrow{\phi_1} s \right) &\approx (s^{-1}t) \cdot \left((\phi_q \cdots \phi_1 s) \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_2} \phi_1 s \xleftarrow{\phi_1} s \right) \\ &= \left((\phi_q \cdots \phi_1 t) \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_2} \phi_1 t \xleftarrow{\phi_1} t \right) \end{aligned}$$

The maps α_* and β_* are clearly inverse to one another. All that remains is to verify that they are chain maps. We need only check compatibility with the zeroth face maps in either case, since the i th face maps (for $i > 0$) simply compose the morphisms ϕ_{i+1} and ϕ_i in either chain complex. The zeroth face maps of either complex will be denoted d_0 .

First consider the map α_* .

$$\begin{array}{ccc} (C_q \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_1} C_0) & \xrightarrow{d_0} & (C_q \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_2} C_1) \\ \alpha_* \downarrow & & \alpha_* \downarrow \\ (\phi_q, \dots, \phi_1) \otimes C_0(u) & \xrightarrow{d_0} & (\phi_q, \dots, \phi_2) \otimes \phi_1 C_0(u) = (\phi_q, \dots, \phi_2) \otimes C_1(u) \end{array}$$

The equality in the lower right of the diagram is simply a restatement that $C_0 \xrightarrow{\phi_1} C_1$ is a morphism of \tilde{S}_p .

For the reverse direction, assume $w = t(u)$ as above. The following diagram commutes.

$$\begin{array}{ccc} (\phi_q, \dots, \phi_1) \otimes w & \xrightarrow{d_0} & (\phi_q, \dots, \phi_2) \otimes \phi_1(w) \\ \beta_* \downarrow & & \beta_* \downarrow \\ ((\phi_q \cdots \phi_1 t) \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_2} \phi_1 t \xleftarrow{\phi_1} t) & \xrightarrow{d_0} & ((\phi_q \cdots \phi_1 t) \xleftarrow{\phi_q} \cdots \xleftarrow{\phi_2} \phi_1 t) \end{array}$$

The lower righthand corner deserves some explanation. Since the target of β_* should be $(N\tilde{S}_p/N\tilde{S}'_p)/G_u$, we see that the lower righthand term is trivial unless ϕ_1 is an isomorphism. In other words, we only need to prove the diagram commutes when $\phi_1 \in \Sigma_{p+1}^{\text{op}}$. As β_* is applied to the chain on the upper righthand corner, the clear choice for group element to begin the chain is $\phi_1 t$, since $\phi_1(w) = \phi_1 t(u)$. \square

Using Lemma 58, we identify \mathcal{M}_u with the orbit complex $(N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p)/G_u$. Now, the complex $N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p$ is a free G_u -complex, so we have an isomorphism,

$$H_*((N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p)/G_u) \cong H_*^{G_u}(N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p),$$

where $H_*^{G_u}$ is G_u -equivariant homology. See Brown [1] for details). Then, by definition,

$$H_*^{G_u}(N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p) = H_*(G_u, N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p),$$

which may be computed using the free resolution, E_*G_u of k as right G_u -module. The resulting complex $k[E_*G_u] \otimes_{kG_u} k[N\tilde{\mathcal{S}}_p]/k[N\tilde{\mathcal{S}}'_p]$ is a double complex isomorphic to the quotient

$$\begin{aligned} (k[E_*G_u] \otimes_{kG_u} k[N\tilde{\mathcal{S}}_p]) / (k[E_*G_u] \otimes_{kG_u} k[N\tilde{\mathcal{S}}'_p]) \\ \cong k[(E_*G_u \times_{G_u} N\tilde{\mathcal{S}}_p) / (E_*G_u \times_{G_u} N\tilde{\mathcal{S}}'_p)]. \end{aligned}$$

This last complex may be identified with the simplicial complex of the space,

$$(EG_u \times_{G_u} |N\tilde{\mathcal{S}}_p|) / (EG_u \times_{G_u} |N\tilde{\mathcal{S}}'_p|) \cong EG_u \times_{G_u} |N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p|.$$

The last piece of the puzzle involves simplifying the spaces $|N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p|$. Since \mathcal{S} is a skeletal subcategory of $\tilde{\mathcal{S}}$, there is an equivalence of categories $\tilde{\mathcal{S}} \simeq \mathcal{S}$, inducing a homotopy equivalence of complexes (hence also of spaces) $|N\tilde{\mathcal{S}}| \simeq |N\mathcal{S}|$. Note that $N\mathcal{S}$ inherits a G_u -action from $N\tilde{\mathcal{S}}$, and the map $\tilde{\mathcal{S}} \rightarrow \mathcal{S}$ is G_u -equivariant.

Proposition 59 *There are weak equivalences, $EG_u \times_{G_u} |N\tilde{\mathcal{S}}_p| \rightarrow EG_u \times_{G_u} |N\mathcal{S}_p|$ and $EG_u \times_{G_u} |N\tilde{\mathcal{S}}'_p| \rightarrow EG_u \times_{G_u} |N\mathcal{S}'_p|$, inducing a weak equivalence $EG_u \times_{G_u} |N\tilde{\mathcal{S}}_p/N\tilde{\mathcal{S}}'_p| \rightarrow EG_u \times_{G_u} |N\mathcal{S}_p/N\mathcal{S}'_p|$.*

Proof The case $p > 2$ will be handled first. As long as the spaces are path connected, we can use the fibration sequences associated to $X \rightarrow EG_u \times_{G_u} X \rightarrow BG_u$ to show that a G_u -equivariant homotopy equivalence $X \simeq Y$ induces $EG_u \times_{G_u} X \simeq EG_u \times_{G_u} Y$. $|N\tilde{\mathcal{S}}_p|$ and $|N\mathcal{S}_p|$ are path-connected because they are contractible. It suffices to show $|N\mathcal{S}'_p|$ path connected, since $|N\tilde{\mathcal{S}}'_p| \simeq |N\mathcal{S}'_p|$.

let $W_0 = z_0z_1 \otimes z_2 \otimes \dots \otimes z_p$. This represents a vertex of $N\mathcal{S}'_p$. Suppose

$$W = Z_0 \otimes \dots \otimes Z'_i z_0 z_1 Z''_i \otimes \dots \otimes Z_s.$$

Then there is a morphism $W_0 \rightarrow W$, hence an edge between W_0 and W .

Next, suppose $W = Z_0 \otimes \cdots \otimes Z'_i z_0 Z''_i z_1 Z'''_i \otimes \cdots \otimes Z_s$. There is a path

$$\begin{array}{c}
 Z_0 \otimes \cdots \otimes Z'_i z_0 Z''_i z_1 Z'''_i \otimes \cdots \otimes Z_s \\
 \downarrow \\
 Z_0 Z_1 \dots Z'_i z_0 Z''_i z_1 Z'''_i \dots Z_s \\
 \uparrow \\
 Z_0 Z_1 \dots Z'_i \otimes z_0 \otimes Z''_i z_1 Z'''_i \dots Z_s \\
 \downarrow \\
 z_0 \otimes Z_0 Z_1 \dots Z'_i Z''_i z_1 Z'''_i \dots Z_s \\
 \uparrow \\
 z_0 \otimes Z_0 Z_1 \dots Z'_i Z''_i \otimes z_1 Z'''_i \dots Z_s \\
 \downarrow \\
 z_0 z_1 Z'''_i \dots Z_s \otimes Z_0 Z_1 \dots Z'_i Z''_i \\
 \uparrow \\
 W_0.
 \end{array}$$

Similarly, if $W = Z_0 \otimes \cdots \otimes Z'_i z_1 Z''_i z_0 Z'''_i \otimes \cdots \otimes Z_s$, there is a path to W_0 . Finally, if $W = Z_0 \otimes \cdots \otimes Z_s$ with z_0 occurring in Z_i and z_1 occurring in Z_j for $i \neq j$, there is an edge to some W' in which $Z_i Z_j$ occurs, and thus a path to W_0 .

The cases $p = 0, 1$ and 2 are handled individually:

Observe that $|N\tilde{S}'_0|$ and $|NS'_0|$ are empty spaces, since \tilde{S}'_0 has no objects. Hence

$$EG_u \times_{G_u} |N\tilde{S}'_0| = EG_u \times_{G_u} |NS'_0| = \emptyset.$$

Furthermore, any group G_u must be trivial. Thus there is a chain of homotopy equivalences

$$EG_u \times_{G_u} |N\tilde{S}_0/N\tilde{S}'_0| \simeq |N\tilde{S}_0| \simeq |NS_0| \simeq EG_u \times_{G_u} |NS_p/NS'_p|.$$

Next, since $|N\tilde{S}'_1|$ is homeomorphic to $|NS'_1|$, each space consisting of the two discrete points $z_0 z_1$ and $z_1 z_0$ with the same group action, the proposition is true for $p = 1$ as well.

For $p = 2$, observe that $|N\tilde{S}'_2|$ has two connected components, \tilde{U}_1 and \tilde{U}_2 that are interchanged by any odd permutation $\sigma \in \Sigma_3$. Similarly, $|NS'_2|$ consists of two connected components, U_1 and U_2 , interchanged by any odd permutation of Σ_3 . Now, restricted to the alternating group, A_3 , we certainly have weak equivalences for any subgroup $H_u \subseteq A_3$,

$$EH_u \times_{H_u} \tilde{U}_1 \xrightarrow{\simeq} EH_u \times_{H_u} U_1 \quad \text{and} \quad EH_u \times_{H_u} \tilde{U}_2 \xrightarrow{\simeq} EH_u \times_{H_u} U_2.$$

The action of an odd permutation induces equivariant homeomorphisms $\tilde{U}_1 \xrightarrow{\cong} \tilde{U}_2$ and $U_1 \xrightarrow{\cong} U_2$, and so if we have a subgroup $G_u \subseteq \Sigma_3$ generated by $H_u \subseteq A_3$ and a transposition, then the two connected components are identified in an $A\Sigma_3$ -equivariant manner. Thus, if G_u contains a transposition,

$$EG_u \times_{G_u} |N\tilde{S}'_2| \cong EH_u \times_{H_u} \tilde{U}_1 \simeq EH_u \times_{H_u} U_1 \cong EG_u \times_{G_u} |NS'_2|.$$

This completes the case $p = 2$ and the proof of [Proposition 59](#). □

[Proposition 59](#) coupled with [Lemma 58](#) produces the required isomorphism in homology, hence proving [Theorem 57](#):

$$\begin{aligned} E_{p,q}^1 &= \bigoplus_{u \in X^{p+1}/\Sigma_{p+1}} H_{p+q}(\mathcal{M}_u) \cong \bigoplus_{u \in X^{p+1}/\Sigma_{p+1}} H_{p+q}(EG_u \times_{G_u} |N\tilde{S}_p/N\tilde{S}'_p|; k) \\ &\cong \bigoplus_{u \in X^{p+1}/\Sigma_{p+1}} H_{p+q}(EG_u \times_{G_u} |NS_p/NS'_p|; k). \end{aligned}$$

Corollary 60 *If the augmentation ideal of A satisfies $I^2 = 0$, then*

$$\tilde{H}S_n(A) \cong \bigoplus_{p \geq 0} \bigoplus_{u \in X^{p+1}/\Sigma_{p+1}} H_n(EG_u \times_{G_u} NS_p/NS'_p; k).$$

Proof This follows from consideration of the original E^0 term of the spectral sequence. E^0 is generated by chains $(\phi_n, \dots, \phi_1) \otimes w$ with induced differential d^0 , agreeing with the differential d of $C_*(\text{Epi}\Delta S, B_*^{\text{sym}}I)$ when ϕ_0 is an isomorphism. When ϕ_0 is a strict epimorphism, however, the zeroth face map of d maps the generator to: $(\phi_n, \dots, \phi_1) \otimes (\phi_0)_*(Y) = 0$, since $(\phi_0)_*(Y)$ would have at least one tensor factor that is the product of two or more elements of I . Thus, d^0 also agrees with d in the case that ϕ_0 is strictly epic. Hence, the spectral sequence collapses at level 1. □

8.4 The complex $\text{Sym}_*^{(p)}$

Note, for $p > 0$, there are homotopy equivalences $|NS_p/N\mathcal{S}'_p| \simeq |NS_p| \vee S|N\mathcal{S}'_p| \simeq S|N\mathcal{S}'_p|$, since $|NS_p|$ is contractible. $|NS_p|$ is a disjoint union of $(p + 1)!$ p -cubes, identified along certain faces. Geometric analysis of $S|N\mathcal{S}'_p|$, however, seems quite difficult. Fortunately, there is an even smaller chain complex homotopic to $NS_p/N\mathcal{S}'_p$.

Definition 61 Let $p \geq 0$ and impose an equivalence relation on $k[\text{Epi}_{\Delta S}([p], [q])]$ generated by

$$Z_0 \otimes \cdots \otimes Z_i \otimes Z_{i+1} \otimes \cdots \otimes Z_q \approx (-1)^{ab} Z_0 \otimes \cdots \otimes Z_{i+1} \otimes Z_i \otimes \cdots \otimes Z_q,$$

where $Z_0 \otimes \cdots \otimes Z_q$ is a morphism expressed in tensor notation, and $a = \text{deg}(Z_i)$ and $b = \text{deg}(Z_{i+1})$, where $\text{deg}(Z) \stackrel{\text{def}}{=} |Z| - 1$, one less than the number of tensor factors of Z . The complex $\text{Sym}_*^{(p)}$ is then defined by

$$\text{Sym}_i^{(p)} \stackrel{\text{def}}{=} k[\text{Epi}_{\Delta S}([p], [p - i])]/ \approx .$$

The face maps will be defined recursively. On monomials,

$$d_i(z_{j_0} \dots z_{j_s}) = \begin{cases} 0, & i < 0, \\ z_{j_0} \dots z_{j_i} \otimes z_{j_{i+1}} \dots z_{j_s}, & 0 \leq i < s, \\ 0, & i \geq s. \end{cases}$$

Then, extend d_i to tensor products via

$$(19) \quad d_i(W \otimes V) = d_i(W) \otimes V + W \otimes d_{i-\text{deg}(W)}(V),$$

where W and V are formal tensors in $k[\text{Epi}_{\Delta S}([p], [q])]$, and

$$\text{deg}(W) = \text{deg}(W_0 \otimes \cdots \otimes W_t) \stackrel{\text{def}}{=} \sum_{k=0}^t \text{deg}(W_k).$$

The boundary map $\text{Sym}_n^{(p)} \rightarrow \text{Sym}_{n-1}^{(p)}$ is then $d = \sum_{i=0}^n (-1)^i d_i = \sum_{i=0}^{n-1} (-1)^i d_i$.

Remark 62 There is an action $\Sigma_{p+1} \times \text{Sym}_i^{(p)} \rightarrow \text{Sym}_i^{(p)}$, given by permuting the formal indeterminates z_i . Furthermore, this action is compatible with the differential.

Lemma 63 $\text{Sym}_*^{(p)}$ is homotopy-equivalent to $k[NS_p]/k[N\mathcal{S}'_p]$.

Proof Let v_0 represent the common initial vertex of the p -cubes making up NS_p . Then, as cell-complex, NS_p consists of v_0 together with all corners of the various p -cubes and i -cells for each i -face of the cubes. Thus, NS_p consists of $(p + 1)!$

p -cells with attaching maps $\partial I^p \rightarrow (NS_p)^{p-1}$ defined according to the face maps for NS_p given above. Note that a chain of NS_p is non-trivial in NS_p/NS'_p if and only if the initial vertex v_0 is included. Thus, any (cubical) k -cell is uniquely determined by the label of the vertex opposite v_0 .

Label each top-dimensional cell with the permutation induced on the set $\{0, 1, \dots, p\}$ by the order of indeterminates in its final vertex, $z_{i_0}z_{i_1} \dots z_{i_p}$. On a given p -cell, for each vertex $Z_0 \otimes \dots \otimes Z_s$, there is an ordering of the tensor factors so that $Z_0 \otimes \dots \otimes Z_s \rightarrow z_{i_0}z_{i_1} \dots z_{i_p}$ preserves the order of formal indeterminates z_i . Rewrite each vertex of this p -cell in this order. Now, any p -chain $(z_{i_0} \otimes z_{i_1} \otimes \dots \otimes z_{i_p}) \rightarrow \dots \rightarrow z_{i_0}z_{i_1} \dots z_{i_p}$ is obtained by choosing the order in which to combine the factors. In fact, the p -chains for this cube are in bijection with the elements of the symmetric group S_p , as in the standard decomposition of a p -cube into $p!$ simplices. A given permutation $\{1, 2, \dots, p\} \mapsto \{j_1, j_2, \dots, j_p\}$ will represent the chain obtained by first combining $z_{j_0} \otimes z_{j_1}$ into $z_{j_0}z_{j_1}$, then combining $z_{j_1} \otimes z_{j_2}$ into $z_{j_1}z_{j_2}$. In effect, we “erase” the tensor product symbol between $z_{j_{r-1}}$ and z_{j_r} for each j_r in order given by the list above.

We shall declare that the *natural* order of combining the factors will be the one that always combines the last two:

$$(z_{i_0} \otimes \dots \otimes z_{i_{p-1}} \otimes z_{i_p}) \longrightarrow (z_{i_0} \otimes \dots \otimes z_{i_{p-1}}z_{i_p}) \longrightarrow (z_{i_0} \otimes \dots \otimes z_{i_{p-2}}z_{i_{p-1}}z_{i_p}) \longrightarrow \dots \longrightarrow (z_{i_0} \dots z_{i_p}).$$

This corresponds to a permutation $\rho \stackrel{\text{def}}{=} \{1, \dots, p\} \mapsto \{p, p-1, \dots, 2, 1\}$, and this chain will be regarded as *positive*. A chain C_σ , corresponding to another permutation, σ , will be regarded as positive or negative depending on the sign of the permutation $\sigma\rho^{-1}$. Finally, the entire p -cell should be identified with the sum $\sum_{\sigma \in S_p} \text{sgn}(\sigma\rho^{-1})C_\sigma$. It is this sign convention that permits the inner faces of the cube to cancel appropriately in the boundary maps. Thus we have a map on the top-dimensional chains:

$$(20) \quad \theta_p: \text{Sym}_p^{(p)} \rightarrow (k[NS_p]/k[NS'_p])_p$$

Extend the definition θ_* to arbitrary k -cells by sending the k -chain $Z_0 \otimes \dots \otimes Z_{p-k}$ to the sum of k -length chains with source $z_0 \otimes \dots \otimes z_p$ and target $Z_0 \otimes \dots \otimes Z_{p-k}$ with signs determined by the natural order of erasing tensor product symbols of $z_0 \otimes \dots \otimes z_p$, excluding those tensor product symbols that never get erased. The following example should clarify the point. Let $W = z_3z_0 \otimes z_1 \otimes z_2z_4$. This is a 2-cell of $\text{Sym}_*^{(4)}$. W is obtained from $z_0 \otimes z_1 \otimes z_2 \otimes z_3 \otimes z_4 = z_3 \otimes z_0 \otimes z_1 \otimes z_2 \otimes z_4$ by combining factors in some order. There are only 2 erasable tensor product symbols in this example. The natural order (last to first) corresponds to the chain, $z_3 \otimes z_0 \otimes z_1 \otimes z_2 \otimes z_4 \rightarrow$

$z_3 \otimes z_0 \otimes z_1 \otimes z_2 z_4 \rightarrow z_3 z_0 \otimes z_1 \otimes z_2 z_4$. So, this chain shows up in $\theta_*(W)$ with positive sign, whereas the chain $z_3 \otimes z_0 \otimes z_1 \otimes z_2 \otimes z_4 \rightarrow z_3 z_0 \otimes z_1 \otimes z_2 \otimes z_4 \rightarrow z_3 z_0 \otimes z_1 \otimes z_2 z_4$ shows up with a negative sign.

Now, θ_* is easily seen to be a chain map $\text{Sym}_*^{(p)} \rightarrow k[N\mathcal{S}_p]/k[N\mathcal{S}'_p]$. Geometrically, θ_* has the effect of subdividing a cell-complex (defined with cubical cells) into a simplicial space, so θ_* is a homotopy-equivalence. Furthermore, θ_* is equivariant with respect to the action of the symmetric group. \square

Remark 64 As an example, consider $|N\mathcal{S}_2|$. There are 6 2-cells, each represented by a copy of I^2 . The 2-cell labelled by the permutation $\{0, 1, 2\} \mapsto \{1, 0, 2\}$ consists of the chains $z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 \otimes z_0 z_2 \rightarrow z_1 z_0 z_2$ and $-(z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 z_0 \otimes z_2 \rightarrow z_1 z_0 z_2)$. Hence, the boundary is the sum of 1-chains:

$$\begin{aligned} & [(z_1 \otimes z_0 z_2 \rightarrow z_1 z_0 z_2) - (z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 z_0 z_2) + (z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 \otimes z_0 z_2)] \\ & - [(z_1 z_0 \otimes z_2 \rightarrow z_1 z_0 z_2) - (z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 z_0 z_2) + (z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 z_0 \otimes z_2)] \\ & = (z_1 \otimes z_0 z_2 \rightarrow z_1 z_0 z_2) + (z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 \otimes z_0 z_2)(z_1 z_0 \otimes z_2 \rightarrow z_1 z_0 z_2) \\ & - (z_1 \otimes z_0 \otimes z_2 \rightarrow z_1 z_0 \otimes z_2). \end{aligned}$$

These 1-chains correspond to the 4 edges of the square. Thus, in our example this 2-cell of $|N\mathcal{S}_p|$ will correspond to $z_1 z_0 z_2 \in \text{Sym}_2^{(2)}$, and its boundary in $|N\mathcal{S}_p/N\mathcal{S}'_p|$ will consist of the two edges adjacent to the vertex labeled $z_0 \otimes z_1 \otimes z_2$, with appropriate signs: $(z_0 \otimes z_1 \otimes z_2 \rightarrow z_1 \otimes z_0 z_2) - (z_0 \otimes z_1 \otimes z_2 \rightarrow z_1 z_0 \otimes z_2)$. The corresponding boundary in $\text{Sym}_1^{(2)}$ will be $(z_1 \otimes z_0 z_2) - (z_1 z_0 \otimes z_2)$, matching the boundary map already defined on $\text{Sym}_*^{(p)}$. See Figures 3 and 4.




Figure 3: $|N\mathcal{S}_2|$ consists of six squares, grouped into two hexagons that share a common center vertex




Figure 4: $\text{Sym}^{(2)} \simeq NS_2 / NS'_2$. The center of each hexagon is $z_0 \otimes z_1 \otimes z_2$.

Now, with one piece of new notation, we may re-interpret [Theorem 57](#).

Definition 65 Let G be a group. Let k_0 be the chain complex consisting of k concentrated in degree 0, with trivial G -action. If X_* is a right G -complex, Y_* is a left G -complex with $k_0 \hookrightarrow Y_*$ as a G -subcomplex, then define the *equivariant half-smash tensor product* of the two complexes:

$$X_* \otimes_G Y_* \stackrel{\text{def}}{=} (X_* \otimes_{kG} Y_*) / (X_* \otimes_{kG} k_0)$$

Corollary 66 There is spectral sequence converging (weakly) to $\tilde{H}S_*(A)$ with

$$E_{p,q}^1 \cong \bigoplus_{u \in X^{p+1} / \Sigma_{p+1}} H_{p+q}(E_* G_u \otimes_{G_u} \text{Sym}_*^{(p)}; k),$$

where G_u is the isotropy subgroup for the chosen representative of $u \in X^{p+1} / \Sigma_{p+1}$.

9 Properties of the complex $\text{Sym}_*^{(p)}$

9.1 Algebra structure of Sym_*

We may consider $\text{Sym}_* \stackrel{\text{def}}{=} \bigoplus_{p \geq 0} \text{Sym}_*^{(p)}$ as a bigraded differential algebra, where $\text{bideg}(W) = (p + 1, i)$ for $W \in \text{Sym}_i^{(p)}$. The product

$$\boxtimes: \text{Sym}_i^{(p)} \otimes \text{Sym}_j^{(q)} \rightarrow \text{Sym}_{i+j}^{(p+q+1)}$$

is defined by $W \boxtimes V \stackrel{\text{def}}{=} W \otimes V'$, where V' is obtained from V by replacing each formal indeterminate z_r by z_{r+p+1} for $0 \leq r \leq q$. Equation (19) then implies

$$(21) \quad d(W \boxtimes V) = d(W) \boxtimes V + (-1)^{\text{bideg}(W)_2} W \boxtimes d(V),$$

where $\text{bideg}(W)_2$ is the second component of $\text{bideg}(W)$.

Proposition 67 *The product \boxtimes is well-defined on the level of homology. Furthermore, this product (on both the chain level and homology level) is skew commutative in a twisted sense: $W \boxtimes V = (-1)^{ij} \tau(V \boxtimes W)$, where $\text{bideg}(W) = (p + 1, i)$, $\text{bideg}(V) = (q + 1, j)$, and τ is the permutation sending*

$$\{0, 1, \dots, q, q + 1, q + 2, \dots, p + q, p + q + 1\} \mapsto \{p + 1, p + 2, \dots, p + q + 1, 0, 1, \dots, p - 1, p\}.$$

Proof Equation (21) implies the product passes to homology classes. Now, suppose $W = Y_0 \otimes Y_1 \otimes \dots \otimes Y_{p-i} \in \text{Sym}_i^{(p)}$ and $V = Z_0 \otimes Z_1 \otimes \dots \otimes Z_{q-j} \in \text{Sym}_j^{(p)}$.

$$(22) \quad V \boxtimes W = V \otimes W' = (-1)^\alpha W' \otimes V,$$

where W' is related to W by replacing each z_r by z_{r+q+1} . The exponent $\alpha = \text{deg}(V) \text{deg}(W) = ij$ arises from the relations in $\text{Sym}_{i+j}^{(p+q+1)}$. (The fact that $\text{deg}(V) = i$ and $\text{deg}(W) = j$ may be made clear by observing that the degree of a formal tensor product in $\text{Sym}_*^{(s)}$ is equal to the number of *cut points*, that is, the number of places where a tensor product symbol may be inserted.) Next, apply the block transformation τ to (22) to obtain $\tau(V \boxtimes W) = (-1)^\alpha \tau(W' \otimes V) = (-1)^\alpha W \otimes V' = (-1)^\alpha W \boxtimes V$, where V' is obtained by replacing z_r by z_{r+p+1} in V . □

9.2 Computer calculations

In principle, the homology of $\text{Sym}_*^{(p)}$ may be found by using a computer. In fact, we have the following results up to $p = 7$:

Theorem 68 *For $0 \leq p \leq 7$, the groups $H_*(\text{Sym}_*^{(p)})$ are free abelian and have Poincaré polynomials $P_p(t) \stackrel{\text{def}}{=} P(H_*(\text{Sym}_*^{(p)}); t)$:*

$$\begin{aligned} P_0(t) &= 1, & P_1(t) &= t, \\ P_2(t) &= t + 2t^2, & P_3(t) &= 7t^2 + 6t^3, \\ P_4(t) &= 43t^3 + 24t^4, & P_5(t) &= t^3 + 272t^4 + 120t^5, \\ P_6(t) &= 36t^4 + 1847t^5 + 720t^6, & P_7(t) &= 829t^5 + 13710t^6 + 5040t^7. \end{aligned}$$

Proof These computations were performed using scripts written for the computer algebra systems GAP [12] and Octave [7]. □

We conjecture that the $H_*(\text{Sym}_*^{(p)})$ is always free abelian.

9.3 Representation theory of $H_*(\text{Sym}_*^{(p)})$

By Remark 62, the groups $H_i(\text{Sym}_*^{(p)}; k)$ carry the structure of $k\Sigma_{p+1}$ -modules, so it seems natural to investigate the irreducible representations comprising these modules.

Proposition 69 *Let $C_{p+1} \hookrightarrow \Sigma_{p+1}$ be the cyclic group of order $p + 1$, embedded into the symmetric group as the subgroup generated by the permutation*

$$\tau_p \stackrel{\text{def}}{=} (0, p, p - 1, \dots, 1).$$

Then there is a Σ_{p+1} -isomorphism $H_p(\text{Sym}_^{(p)}) \cong AC_{p+1} \uparrow \Sigma_{p+1}$, that is, the alternating representation of the cyclic group, induced up to the symmetric group. Note, for p even, AC_{p+1} coincides with the trivial representation IC_{p+1} .*

Moreover, $H_p(\text{Sym}_^{(p)})$ is generated by the elements $\sigma(b_p)$, for the distinct cosets σC_{p+1} , where $b_p \stackrel{\text{def}}{=} \sum_{j=0}^p (-1)^{j p} \tau_p^j(z_0 z_1 \dots z_p)$.*

Proof Let w be a general element of $\text{Sym}_p^{(p)}$, $w = \sum_{\sigma \in \Sigma_{p+1}} c_\sigma \sigma(z_0 z_1 \dots z_p)$, where c_σ are constants in k . $H_p(\text{Sym}_*^{(p)})$ consists of those w such that $d(w) = 0$. That is,

$$(23) \quad 0 = \sum_{\sigma \in \Sigma_{p+1}} \sum_{i=0}^{p-1} (-1)^i c_\sigma \sigma(z_0 \dots z_i \otimes z_{i+1} \dots z_p).$$

Now for each σ , the terms corresponding to $\sigma(z_0 \dots z_i \otimes z_{i+1} \dots z_p)$ occur in pairs in the above formula. The obvious term of the pair is $(-1)^i c_\sigma \sigma(z_0 \dots z_i \otimes z_{i+1} \dots z_p)$. Not so obviously, the second term of the pair is

$$(-1)^{(p-i-1)i} (-1)^{p-i-1} c_\rho \rho(z_0 \dots z_{p-i-1} \otimes z_{p-i} \dots z_p),$$

where $\rho = \sigma \tau_p^{p-i}$. Thus, if $d(w) = 0$, then we must have

$$(-1)^i c_\sigma + (-1)^{(p-i-1)(i+1)} c_\rho = 0,$$

or $c_\rho = (-1)^{(p-i)(i+1)} c_\sigma$. Set $j = p - i$, so that $c_\rho = (-1)^{j(p-j+1)} c_\sigma = (-1)^{j p} c_\sigma$. This shows that the only restrictions on the coefficients c_σ are that the absolute values of coefficients corresponding to $\sigma, \sigma \tau_p, \sigma \tau_p^2, \dots$ must be the same, and their corresponding signs in w alternate if and only if p is odd; otherwise, they have the same signs.

Clearly, the elements $\sigma(b_p)$ for distinct cosets σC_{p+1} represents an independent set of generators over k for $H_p(\text{Sym}_*^{(p)})$.

Observe that b_p is invariant under the action of $\text{sgn}(\tau_p)\tau_p$, and so b_p generates an alternating representation AC_{p+1} over k . Induced up to Σ_{p+1} , we obtain the representation $AC_{p+1} \uparrow \Sigma_{p+1}$ of dimension $(p+1)!/(p+1) = p!$, generated by the elements $\sigma(b_p)$ as in the proposition. \square

Definition 70 For a given proper partition $\lambda = [\lambda_0, \lambda_1, \lambda_2, \dots, \lambda_s]$ of the $p+1$ integers $\{0, 1, \dots, p\}$, an element W of $\text{Sym}_*^{(p)}$ will be designated as *type* λ if it is equivalent to $\pm(Y_0 \otimes Y_1 \otimes Y_2 \otimes \dots \otimes Y_s)$ with $\text{deg}(Y_i) = \lambda_i - 1$. That is, each Y_i has λ_i factors. The notation $\text{Sym}_\lambda^{(p)}$ or Sym_λ will denote the k -submodule of $\text{Sym}_{p-s}^{(p)}$ generated by all elements of type λ .

In what follows, $|\lambda|$ will refer to the number of components of λ . The action of Σ_{p+1} leaves Sym_λ invariant for any given λ , so there is a decomposition as $k\Sigma_{p+1}$ -module:

$$\text{Sym}_{p-s}^{(p)} = \bigoplus_{\lambda \vdash (p+1), |\lambda|=s+1} \text{Sym}_\lambda.$$

Proposition 71 For a given proper partition $\lambda \vdash (p+1)$,

- (a) Sym_λ contains exactly one alternating representation $A\Sigma_{p+1}$ iff λ contains no repeated components.
- (b) Sym_λ contains exactly one trivial representation $I\Sigma_{p+1}$ iff λ contains no repeated even components.

Proof Sym_λ is a quotient of the regular representation, since it is the image of the Σ_{p+1} -map $\pi_\lambda: k\Sigma_{p+1} \rightarrow \text{Sym}_\lambda$, $\sigma \mapsto \psi_\lambda \sigma$, where $s \in \Sigma_{p+1}^{\text{op}}$ is the ΔS -automorphism of $[p]$ corresponding to σ and ψ_λ is a Δ -morphism $[p] \rightarrow [|\lambda|]$ that sends the points $0, \dots, \lambda_0 - 1$ to 0, the points $\lambda_0, \dots, \lambda_0 + \lambda_1 - 1$ to 1, and so on. Hence, there can be at most 1 copy of $A\Sigma_{p+1}$ and at most 1 copy of $I\Sigma_{p+1}$ in Sym_λ .

Let W be the “standard” element of Sym_λ . That is, the indeterminates z_i occur in W in numerical order and the degrees of monomials of W are in decreasing order. $A\Sigma_{p+1}$ exists in Sym_λ iff the element $V = \sum_{\sigma \in \Sigma_{p+1}} \text{sgn}(\sigma)\sigma(W)$ is non-zero. Suppose that some component of λ is repeated, say $\lambda_i = \lambda_{i+1} = \ell$. If $W = Y_0 \otimes Y_1 \otimes \dots \otimes Y_s$, then $\text{deg}(Y_i) = \text{deg}(Y_{i+1}) = \ell - 1$. Now, we know that $W = (-1)^{\text{deg}(Y_i)\text{deg}(Y_{i+1})} Y_0 \otimes \dots \otimes Y_{i+1} \otimes Y_i \otimes \dots \otimes Y_s = -(-1)^\ell \alpha(W)$, for the permutation $\alpha \in \Sigma_{p+1}$ that exchanges the indices of indeterminates in Y_i with those in Y_{i+1} in an order-preserving way.

In V , the term $\alpha(W)$ shows up with sign $\text{sgn}(\alpha) = (-1)^\ell$, thus cancelling with W . Hence, $V = 0$, and no alternating representation exists.

If, on the other hand, no component of λ is repeated, then no term $\pm\alpha(W)$ can be equivalent to W for $\alpha \neq \text{id}$, so V survives as the generator of $A\Sigma_{p+1}$ in Sym_λ .

A similar analysis applies for trivial representations. This time, we examine $U = \sum_{\sigma \in \Sigma_{p+1}} \sigma(W)$, which would be a generator for $I\Sigma_{p+1}$ if it were non-zero. As before, if there is a repeated component, $\lambda_i = \lambda_{i+1} = \ell$, then $W = (-1)^{\ell-1}\alpha(W)$. However, this time, W cancels with $\alpha(W)$ only if $\ell-1$ is odd. That is, $|\lambda_i| = |\lambda_{i+1}|$ is even. If $\ell-1$ is even, or if all λ_i are distinct, then the element U must be non-zero. \square

Proposition 72 $H_i(\text{Sym}_*^{(p)})$ contains an alternating representation for each partition $\lambda \vdash (p+1)$ with $|\lambda| = p-i$ such that no component of λ is repeated.

Proof This proposition will follow from the fact that $d(V) = 0$ for any generator V of an alternating representation in Sym_λ . Then, by Schur’s Lemma, the alternating representation must survive at the level of homology.

Let $V = \sum_{\sigma \in \Sigma_{p+1}} \text{sgn}(\sigma)\sigma(W)$ be the generator mentioned in Proposition 71. $d(V)$ consists of individual terms $d_j(\sigma(W)) = \sigma(d_j(W))$ along with appropriate signs. For a given j , $d_j(W)$ is identical to W except at some monomial Y_i , where a tensor product symbol is inserted. We will introduce some notation to make the argument a little cleaner. If $Y = z_{i_0}z_{i_1} \dots z_{i_r}$ is a monomial, then the notation $Y\{s, \dots, t\}$ refers to the monomial $z_{i_s}z_{i_{s+1}} \dots z_{i_t}$, assuming $0 \leq s \leq t \leq r$. Now, we write

$$(24) \quad d_j(W) = (-1)^{a+\ell} Y_0 \otimes \dots \otimes Y_i\{0, \dots, \ell\} \otimes Y_i\{\ell + 1, \dots, m\} \otimes \dots \otimes Y_s,$$

where $a = \text{deg}(Y_0) + \dots + \text{deg}(Y_{i-1})$. Use the relations in Sym_* to rewrite (24):

$$(25) \quad (-1)^{(a+\ell)+\ell(m-\ell-1)} Y_0 \otimes \dots \otimes Y_i\{\ell + 1, \dots, m\} \otimes Y_i\{0, \dots, \ell\} \otimes \dots \otimes Y_s$$

Let α be the permutation that relabels indices in such a way that $Y_i\{0, \dots, m-\ell-1\} \mapsto Y_i\{\ell + 1, \dots, m\}$ and $Y_i\{m-\ell, \dots, m\} \mapsto Y_i\{0, \dots, \ell\}$, so that the following is equivalent to (25).

$$(26) \quad (-1)^{a+m\ell-\ell^2} \alpha(Y_0 \otimes \dots \otimes Y_i\{0, \dots, m-\ell-1\} \otimes Y_i\{m-\ell, \dots, m\} \otimes \dots \otimes Y_s)$$

Now, (26) also occurs in $d_{j'}(\text{sgn}(\alpha)\alpha(W))$ for some j' . This term looks like

$$(27) \quad \text{sgn}(\alpha)(-1)^{a+m-\ell-1} \alpha(Y_0 \otimes \dots \otimes Y_i\{0, \dots, m-\ell-1\} \otimes Y_i\{m-\ell, \dots, m\} \otimes \dots \otimes Y_s)$$

which equals

$$(28) \quad (-1)^{m\ell - \ell^2 + a - 1} \alpha(Y_0 \otimes \cdots \otimes Y_i\{0, \dots, m - \ell - 1\} \otimes Y_i\{m - \ell, \dots, m\} \otimes \cdots \otimes Y_s)$$

Comparing the signs of (28) and (26), we verify the two terms cancel each other out in the sum $d(V)$. □

By Proposition 72, it is clear that if $p + 1$ is a triangular number – that is, $p + 1$ is of the form $r(r + 1)/2$ for some positive integer r , then the lowest dimension in which an alternating representation may occur is $p + 1 - r$, corresponding to the partition $\lambda = [r, r - 1, \dots, 2, 1]$. A little algebra yields the following statement for any p .

Corollary 73 $H_i(\text{Sym}_*^{(p)})$ contains an alternating representation in degree $p + 1 - r$, where $r = \lfloor \sqrt{2p + 9/4} - 1/2 \rfloor$. Moreover, there are no alternating representations present for $i \leq p - r$.

There is not much known about the other irreducible representations occurring in the homology groups of $\text{Sym}_*^{(p)}$, however computational evidence shows that $H_i(\text{Sym}_*^{(p)})$ contains no trivial representation, $I\Sigma_{p+1}$, for $i \leq p - r$ (r as in the conjecture above) up to $p = 50$.

9.4 Connectivity of $\text{Sym}_*^{(p)}$

Quite recently, Vrećica and Živaljević [30] observed that the complex $\text{Sym}_*^{(p)}$ is isomorphic to the suspension of the cycle-free chessboard complex Ω_{p+1} (in fact, the isomorphism takes the form $k[S\Omega_{p+1}^+] \rightarrow \text{Sym}_*^{(p)}$, where Ω_{p+1}^+ is the augmented complex).

The m -chains of the complex Ω_n are generated by ordered lists,

$$L = \{(i_0, j_0), (i_1, j_1), \dots, (i_m, j_m)\},$$

where $1 \leq i_0 < i_1 < \dots < i_m \leq n$, all $1 \leq j_s \leq n$ are distinct integers, and the list L is cycle-free. It may be easier to say what it means for L not to be cycle free: L is not cycle-free if there exists a subset $L_c \subseteq L$ and ordering of L_c so that

$$L_c = \{(\ell_0, \ell_1), (\ell_1, \ell_2), \dots, (\ell_{t-1}, \ell_t), (\ell_t, \ell_0)\}.$$

The face maps are defined by deletion of elements of L :

$$d_s(\{(i_0, j_0), \dots, (i_m, j_m)\}) \stackrel{\text{def}}{=} \{(i_0, j_0), \dots, (i_{s-1}, j_{s-1}), (i_{s+1}, j_{s+1}), \dots, (i_m, j_m)\}$$

For completeness, an explicit isomorphism shall be provided as part of the proof of the following proposition.

Proposition 74 Let Ω_n^+ denote the augmented cycle-free $(n \times n)$ -chessboard complex, where the unique (-1) -chain is represented by the empty $n \times n$ chessboard, and the boundary map on 0 -chains takes a vertex to the unique (-1) -chain. For each $p \geq 0$, there is a chain isomorphism, $\omega_*: k[S\Omega_{p+1}^+] \rightarrow \text{Sym}_*^{(p)}$.

Proof Note that we may define the generating m -chains of $k[\Omega_{p+1}]$ as cycle-free lists L , with no requirement on the order of L , under the equivalence relation:

$$\sigma.L \stackrel{\text{def}}{=} \{(i_{\sigma^{-1}(0)}, j_{\sigma^{-1}(0)}), \dots, (i_{\sigma^{-1}(m)}, j_{\sigma^{-1}(m)})\} \approx \text{sgn}(\sigma)L,$$

for $\sigma \in \Sigma_{m+1}$. Suppose L is an $(m + 1)$ -chain of $S\Omega_{p+1}^+$ (that is an m -chain of Ω_{p+1}^+). Call a subset $L' \subseteq L$ a *queue* if there is a reordering of L' such that

$$L' = \{(\ell_0, \ell_1), (\ell_1, \ell_2), \dots, (\ell_{t-1}, \ell_t)\}.$$

L' is called a *maximal queue* if it is not properly contained in any other queue. Since L is supposed to be cycle-free, we can partition L into some number of maximal queues, L'_1, L'_2, \dots, L'_q . Let σ be a permutation representing the reordering of L into maximal ordered queues.

Now, each maximal ordered queue L'_i will correspond to a monomial of formal indeterminates z_i as follows.

$$(29) \quad L'_s = \{(\ell_0, \ell_1), (\ell_1, \ell_2), \dots, (\ell_{t-1}, \ell_t)\} \mapsto z_{\ell_0-1}z_{\ell_1-1} \cdots z_{\ell_{t-1}}.$$

For each maximal ordered queue, L'_s , denote the monomial obtained by formula (29) by Z_s . Let k_1, k_2, \dots, k_u be the numbers in $\{0, 1, 2, \dots, p\}$ such that $k_r + 1$ does not appear in any pair $(i_s, j_s) \in L$. Now we may define ω_* on $L = L'_1 \cup L'_2 \cup \dots \cup L'_q$.

$$(30) \quad \omega_{m+1}(L) \stackrel{\text{def}}{=} Z_1 \otimes Z_2 \otimes \cdots \otimes Z_q \otimes z_{k_1} \otimes z_{k_2} \otimes \cdots \otimes z_{k_u}.$$

Observe, if $L = \emptyset$ is the (-1) -chain of Ω_{p+1}^+ , then there are no maximal queues in L , and so $\omega_0(\emptyset) = z_0 \otimes z_1 \otimes \cdots \otimes z_p$.

ω_* is a (well-defined) chain map with inverse given by essentially reversing the process. To each monomial $Z = z_{i_0}z_{i_1} \cdots z_{i_t}$ with $t > 0$, there is an associated ordered queue $L' = \{(i_0 + 1, i_1 + 1), (i_1 + 1, i_2 + 1), \dots, (i_{t-1} + 1, i_t + 1)\}$. If the monomial is a singleton, $Z = z_{i_0}$, the associated ordered queue will be the empty set. Now, given a generator $Z_1 \otimes Z_2 \otimes \cdots \otimes Z_q \in \text{Sym}_*^{(p)}$, map it to the list $L \stackrel{\text{def}}{=} L'_1 \cup L'_2 \cup \dots \cup L'_q$, preserving the original order of indices. □

Theorem 75 $\text{Sym}_*^{(p)}$ is $\lfloor \frac{2}{3}(p - 1) \rfloor$ -connected.

Proof See Vrećica and Živaljević [30, Theorem 10]. □

This remarkable fact yields the following useful corollaries:

Corollary 76 *The spectral sequences of Theorem 57 and Corollary 66 converge strongly to $\tilde{H}S_*(A)$.*

Proof The connectivity of the complexes $\text{Sym}_*^{(p)}$ is a non-decreasing function of p . So for large enough p , we have $H_n(E_*G_u \otimes_{G_u} \text{Sym}_*^{(p)}) = 0$. For fixed n , the spectral sequence is guaranteed to collapse at a finite level. \square

Corollary 77 *For each $i \geq 0$, there is a positive integer N_i so that if $p \geq N_i$, there is an isomorphism $\tilde{H}S_i(A) \cong H_i(\mathcal{F}_p C_*(\text{Epi} \Delta S, B_*^{\text{sym}} I))$.*

Corollary 78 *If A is finitely-generated over a Noetherian ground ring k , then $HS_*(A)$ is finitely-generated over k in each degree.*

The bounds on connectivity are conjectured to be tight. This is certainly true for $p \equiv 1 \pmod{3}$, based on Vrećica and Živaljević [30, Theorem 16]. [30, Corollary 12] establishes that either $H_{2k}(\text{Sym}_*^{(3k-1)}) \neq 0$ or $H_{2k}(\text{Sym}_*^{(3k)}) \neq 0$. For $k \leq 2$, both statements are true. When the latter condition is true, this gives a tight bound on connectivity for $p \equiv 0 \pmod{3}$. When the former is true, there is not enough information for a tight bound, since we are more interested in proving that $H_{2k-1}(\text{Sym}_*^{(3k-1)})$ is non-zero, since for $k = 1, 2$, we have computed the integral homology, $H_1(\text{Sym}_*^{(2)}) = \mathbb{Z}$ and $H_3(\text{Sym}_*^{(5)}) = \mathbb{Z}$.

10 A partial resolution

As before, k is a commutative ground ring. In this section, we find an explicit partial resolution of the trivial ΔS^{op} -module \underline{k} by projective modules, allowing the computation of $HS_0(A)$ and $HS_1(A)$ for a unital associative k -algebra A . The resolution will be constructed through a number of technical lemmas.

For any small category \mathcal{C} and object $X \in \text{Obj} \mathcal{C}$, the \mathcal{C} -modules $k[\text{Mor}_{\mathcal{C}}(X, -)]$ are projective (as \mathcal{C} -module), as are the \mathcal{C}^{op} -modules $k[\text{Mor}_{\mathcal{C}}(-, X)]$. In particular, each $k[\text{Mor}_{\Delta S}(-, [q])]$ is a projective ΔS^{op} -module. In proving exactness, it suffices to examine the individual sub- k -modules, $k[\text{Mor}_{\Delta S}([n], [q])]$.

Lemma 79 *For each $n \geq 0$, the sequence*

$$0 \longleftarrow k \xleftarrow{\epsilon} k[\text{Mor}_{\Delta S}([n], [0])] \xleftarrow{\rho} k[\text{Mor}_{\Delta S}([n], [2])]$$

is exact, where ϵ is defined by $\epsilon(\phi) = 1$ for any morphism $\phi: [n] \rightarrow [0]$, and ρ is defined by $\rho(\psi) = (x_0x_1x_2) \circ \psi - (x_2x_1x_0) \circ \psi$ for any morphism $\psi: [n] \rightarrow [2]$ ($x_0x_1x_2$ and $x_2x_1x_0$ are ΔS morphisms $[2] \rightarrow [0]$ written in tensor notation).

Proof Clearly, ϵ is surjective. Now, $\epsilon\rho = 0$, since $\rho(\psi)$ consists of two morphisms with opposite signs. Let $\phi_0 = x_0x_1 \dots x_n: [n] \rightarrow [0]$. The kernel of ϵ is spanned by elements $\phi - \phi_0$ for $\phi \in \text{Mor}_{\Delta S}([n], [0])$. So, it suffices to show that the submodule of $k[\text{Mor}_{\Delta S}([n], [0])]$ generated by $(x_0x_1x_2)\psi - (x_2x_1x_0)\psi$ for $\psi: [n] \rightarrow [2]$ contains all of the elements $\phi - \phi_0$. In other words, it suffices to find a sequence $\phi = \phi_k, \phi_{k-1}, \dots, \phi_2, \phi_1, \phi_0$ so that each ϕ_i is obtained from ϕ_{i+1} by reversing the order of 3 (possibly empty) blocks, $XYZ \rightarrow ZYX$. Let $\phi = x_{i_0}x_{i_1} \dots x_{i_n}$. If $\phi = \phi_0$, we may stop here. Otherwise, we may produce a sequence ending in ϕ_0 by way of a certain family of rearrangements:

k -rearrangement $x_{i_0}x_{i_1} \dots x_{i_{k-1}}x_{i_k}x_{k+1} \dots x_n \rightsquigarrow x_{k+1} \dots x_nx_{i_k}x_{i_0}x_{i_1} \dots x_{i_{k-1}}$ such that $i_k \neq k$. That is, a k -rearrangement only applies to those monomials that agree with ϕ_0 in the final $n - k$ indeterminates, but not in the final $n - k + 1$ indeterminates. If $k = n$, then this rearrangement reduces to the cyclic rearrangement, $x_{i_0}x_{i_1} \dots x_{i_n} \rightsquigarrow x_{i_n}x_{i_0}x_{i_1} \dots x_{i_{n-1}}$.

Beginning with ϕ , perform n -rearrangements until the final indeterminate is x_n . For convenience of notation, let this new monomial be $x_{j_0}x_{j_1} \dots x_{j_n}$. (Of course, $j_n = n$.) If $j_k = k$ for all $k = 0, 1, \dots, n$, then we are done. Otherwise, there will be a number k such that $j_k \neq k$ but $j_{k+1} = k + 1, \dots, j_n = n$. Perform a k -rearrangement followed by enough n -rearrangements so that the final indeterminate is again x_n . The net result of these rearrangements is that the ending block $x_{k+1}x_{k+2} \dots x_n$ remains fixed while the beginning block $x_{j_0}x_{j_1} \dots x_{j_k}$ becomes cyclically permuted to $x_{j_k}x_{j_0} \dots x_{j_{k-1}}$. It is clear that applying this combination of rearrangements repeatedly will finally obtain a monomial $x_{\ell_0}x_{\ell_1} \dots x_{\ell_{k-1}}x_kx_{k+1} \dots x_n$. Now repeat the process, until after a finite number of steps, we finally obtain ϕ_0 . □

Let $\mathcal{B}_n \stackrel{\text{def}}{=} \{x_{i_0}x_{i_1} \dots x_{i_{k-1}} \otimes x_{i_k} \otimes x_{k+1}x_{k+2} \dots x_n \mid 1 \leq k \leq n, i_k \neq k\}$. Then $k[\mathcal{B}_n]$ is a free submodule of $k[\text{Mor}_{\Delta S}([n], [2])]$ of size $(n + 1)! - 1$.

Corollary 80 When restricted to $k[\mathcal{B}_n]$, the map ρ of Lemma 79 is surjective onto the kernel of ϵ .

Proof In the proof of Lemma 79, the n -rearrangements correspond to the image of elements $x_{i_0} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1$, with $i_n \neq n$. For $k < n$, k -rearrangements correspond to the image of elements $x_{i_0} \dots x_{i_{k-1}} \otimes x_{i_k} \otimes x_{k+1} \dots x_n$, with $i_k \neq k$. □

Lemma 81 $k[\text{Mor}_{\Delta S}([n], [m])]$ is a free k -module of rank $(m + n + 1)!/m!$.

Proof A morphism $\phi: [n] \rightarrow [m]$ of ΔS is nothing more than an assignment of $n + 1$ objects into $m + 1$ compartments, along with a total ordering of the original $n + 1$ objects, hence $\#\text{Mor}_{\Delta S}([n], [m]) = \binom{m+n+1}{m}(n + 1)! = \frac{(m+n+1)!}{m!}$. \square

Lemma 82 $\rho|_{k[\mathcal{B}_n]}$ is an isomorphism $k[\mathcal{B}_n] \cong \ker \epsilon$.

Proof Since the rank of $k[\text{Mor}_{\Delta S}([n], [0])]$ is $(n + 1)!$, the rank of the $\ker \epsilon$ is $(n + 1)! - 1$. The isomorphism then follows from [Corollary 80](#). \square

Lemma 83 The relations of the form

$$(31) \quad XY \otimes Z \otimes W + W \otimes ZX \otimes Y + YZX \otimes 1 \otimes W + W \otimes YZ \otimes X \approx 0 \quad \text{and}$$

$$(32) \quad 1 \otimes X \otimes 1 \approx 0$$

collapse $k[\text{Mor}_{\Delta S}([n], [2])]$ onto $k[\mathcal{B}_n]$.

Proof This proof proceeds in multiple steps.

Step 1: Degeneracy Relations $X \otimes Y \otimes 1 \approx X \otimes 1 \otimes Y \approx 1 \otimes X \otimes Y$.

First, observe that letting $X = Y = W = 1$ in (31) yields $Z \otimes 1 \otimes 1 \approx 0$, since $1 \otimes Z \otimes 1 \approx 0$. Then, letting $X = Z = W = 1$ in (31) produces $1 \otimes 1 \otimes Y \approx 0$. Thus, any formal tensor with two trivial factors is equivalent to 0. Next, let $Z = W = 1$ in (31). Then using the above observation, we obtain $1 \otimes X \otimes Y + 1 \otimes Y \otimes X \approx 0$, that is, $1 \otimes X \otimes Y \approx -(1 \otimes Y \otimes X)$. Then, if we let $X = W = 1$, we obtain $Y \otimes Z \otimes 1 + 1 \otimes Z \otimes Y \approx 0$, which is equivalent to $Y \otimes Z \otimes 1 - 1 \otimes Y \otimes Z \approx 0$. Finally, let $X = Y = 1$ in (31). The expression reduces to $Z \otimes 1 \otimes W - 1 \otimes Z \otimes W \approx 0$.

Step 2: Sign Relation $X \otimes Y \otimes Z \approx -(Z \otimes Y \otimes X)$.

Let $Y = 1$ in (31), and use the degeneracy relations to rewrite the result as $X \otimes Z \otimes W + 1 \otimes W \otimes ZX + 1 \otimes ZX \otimes W + W \otimes Z \otimes X \approx 0$. Since $1 \otimes ZX \otimes W \approx -(1 \otimes W \otimes ZW)$, the desired result follows: $X \otimes Z \otimes W + W \otimes Z \otimes X \approx 0$.

Step 3: Hochschild Relation $XY \otimes Z \otimes 1 - X \otimes YZ \otimes 1 + ZX \otimes Y \otimes 1 \approx 0$. This relation is named after the similar relation, $XY \otimes Z - X \otimes YZ + ZX \otimes Y$, that arises in the Hochschild complex.

Let $W = 1$ in (31), and use the degeneracy and sign relations to obtain the desired result.

Step 4: Cyclic Relation $\sum_{j=0}^n \tau_n^j (x_{i_0} x_{i_1} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1) \approx 0$, where $\tau_n \in \Sigma_{n+1}$ is the $(n + 1)$ -cycle $(0, n, n - 1, \dots, 2, 1)$, which acts by permuting the indices.

For $n = 0$, there are no such relations (indeed, no relations at all). For $n = 1$, the cyclic relation takes the form $x_0 \otimes x_1 \otimes 1 + x_1 \otimes x_0 \otimes 1 \approx 0$, which follows from degeneracy and sign relations.

Assume now that $n \geq 2$. For each $k = 1, 2, \dots, n - 1$, define

$$\begin{aligned} A_k &\stackrel{\text{def}}{=} x_{i_0} x_{i_1} \dots x_{i_{k-1}}, \\ B_k &\stackrel{\text{def}}{=} x_{i_k}, \\ C_k &\stackrel{\text{def}}{=} x_{i_{k+1}} \dots x_{i_n}. \end{aligned}$$

By the Hochschild relation, $0 \approx \sum_{k=1}^{n-1} (A_k B_k \otimes C_k \otimes 1 - A_k \otimes B_k C_k \otimes 1 + C_k A_k \otimes B_k \otimes 1)$. But for $k \leq n - 2$, $A_k B_k \otimes C_k \otimes 1 = A_{k+1} \otimes B_{k+1} C_{k+1} \otimes 1$. Thus, after some cancellation,

$$(33) \quad 0 \approx -A_1 \otimes B_1 C_1 \otimes 1 + A_{n-1} B_{n-1} \otimes C_{n-1} \otimes 1 + \sum_{k=1}^{n-1} C_k A_k \otimes B_k \otimes 1.$$

Now observe that sign and degeneracy relations imply that $-A_1 \otimes B_1 C_1 \otimes 1 \approx x_{i_1} \dots x_{i_n} \otimes x_{i_0} \otimes 1$. The term $A_{n-1} B_{n-1} \otimes C_{n-1} \otimes 1$ is equal to $x_{i_0} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1$, and for $1 \leq k \leq n - 1$, $C_k A_k \otimes B_k \otimes 1 = x_{i_{k+1}} \dots x_{i_n} x_{i_0} \dots x_{i_{k-1}} \otimes x_{i_k} \otimes 1$. Thus, (33) can be rewritten as the cyclic relation,

$$0 \approx (x_{i_0} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1) + \sum_{k=0}^{n-1} x_{i_{k+1}} \dots x_{i_n} x_{i_0} \dots x_{i_{k-1}} \otimes x_{i_k} \otimes 1.$$

Step 5 Every element of the form $X \otimes Y \otimes 1$ is equivalent to a linear combination of elements of \mathcal{B}_n .

To prove this, we shall induct on the size of Y . Suppose Y consists of a single indeterminate. That is, $X \otimes Y \otimes 1 = x_{i_0} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1$. Now, if $i_n \neq n$, we are done. Otherwise, we use the cyclic relation to write $x_{i_0} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1 \approx -\sum_{j=1}^n \tau_n^j (x_{i_0} \dots x_{i_{n-1}} \otimes x_{i_n} \otimes 1)$.

Now suppose $k \geq 1$ and any element $Z \otimes W \otimes 1$ with $|W| = k$ is equivalent to an element of $k[\mathcal{B}_n]$. Consider $X \otimes Y \otimes 1 = x_{i_0} \dots x_{i_{n-k-1}} \otimes x_{i_{n-k}} \dots x_{i_n} \otimes 1$. Let

$$\begin{aligned} A &= x_{i_0} x_{i_1} \dots x_{i_{n-k-1}}, \\ B &= x_{i_{n-k}} \dots x_{i_{n-1}}, \\ C &= x_{i_n}. \end{aligned}$$

Then, by the Hochschild relation, $X \otimes Y \otimes 1 = A \otimes BC \otimes 1 \approx AB \otimes C \otimes 1 + CA \otimes B \otimes 1$. But since C has one indeterminate and B has k indeterminates, this last expression is equivalent to an element of $k[\mathcal{B}_n]$ by inductive hypothesis.

Step 6: Modified Hochschild Relation $XY \otimes Z \otimes W - X \otimes YZ \otimes W + ZX \otimes Y \otimes W \approx 0$, modulo $k[\mathcal{B}_n]$.

First, we show that $X \otimes Y \otimes W + Y \otimes X \otimes W \approx 0 \pmod{k[\mathcal{B}_n]}$. Indeed, if we let $Z = 1$ in (31), then sign and degeneracy relations yield: $X \otimes Y \otimes W + Y \otimes X \otimes W \approx XY \otimes W \otimes 1 + YX \otimes W \otimes 1$, that is, by step 5,

$$(34) \quad X \otimes Y \otimes W \approx -(Y \otimes X \otimes W) \pmod{k[\mathcal{B}_n]}.$$

Now, using sign and degeneracy relations, (31) can be re-expressed:

$$XY \otimes Z \otimes W - Y \otimes ZX \otimes W + YZX \otimes W \otimes 1 - X \otimes YZ \otimes W \approx 0.$$

Using (34), we then arrive at the modified Hochschild relation,

$$(35) \quad XY \otimes Z \otimes W - X \otimes YZ \otimes W + ZX \otimes Y \otimes W \approx 0 \pmod{k[\mathcal{B}_n]}.$$

Step 7 Modified Cyclic Relation $\sum_{j=0}^k \tau_k^j(x_{i_0}x_{i_1} \dots x_{i_{k-1}} \otimes x_{i_k} \otimes x_{i_{k+1}} \dots x_{i_n}) \approx 0$, modulo $k[\mathcal{B}_n]$. Note, the $(k + 1)$ -cycle τ_k permutes the indices i_0, i_1, \dots, i_k , and fixes the rest.

Equation (34) proves the modified cyclic relations for $k = 1$. The modified cyclic relation for $k \geq 2$ follows from the modified Hochschild relation in the same manner as in step 4. Of course, this time all equivalences are taken modulo $k[\mathcal{B}_n]$.

Step 8 Every element of the form $X \otimes Y \otimes x_n$ is equivalent to an element of $k[\mathcal{B}_n]$.

We shall use the modified cyclic and modified Hochschild relations in a similar way as cyclic and Hochschild relations were used in step 5. Again we induct on the size of Y . If $|Y| = 1$, then $X \otimes Y \otimes x_n = x_{i_0} \dots x_{i_{n-2}} \otimes x_{i_{n-1}} \otimes x_n$. If $i_{n-1} \neq n - 1$, then we are done. Otherwise, use the modified cyclic relation to re-express $X \otimes Y \otimes x_n$ as a sum of elements of $k[\mathcal{B}_n]$.

Next, suppose $k \geq 1$ and any element $Z \otimes W \otimes x_n$ with $|W| = k$ is equivalent to an element of $k[\mathcal{B}_n]$. Consider an element $X \otimes Y \otimes x_n$ with $|Y| = k + 1$. Write $Y = BC$ with $|B| = k$ and $|C| = 1$ and use the modified Hochschild relation to rewrite $X \otimes BC \otimes x_n$ in terms of two elements whose middle tensor factors are either B or C (modulo $k[\mathcal{B}_n]$). By inductive hypothesis, the rewritten expression must lie in $k[\mathcal{B}_n]$.

Step 9 Every element of $k[\text{Mor}_{\Delta S}([n], [2])]$ is equivalent to a linear combination of elements from the following set:

$$(36) \quad \mathcal{C}_n \stackrel{\text{def}}{=} \{X \otimes x_{i_n} \otimes 1 \mid i_n \neq n\} \cup \{X \otimes x_{i_{n-1}} \otimes x_n \mid i_{n-1} \neq n-1\} \\ \cup \{X \otimes Y \otimes Z x_n \mid |Z| \geq 1\}$$

Note, the k -module generated by \mathcal{C}_n contains $k[\mathcal{B}_n]$.

Let $X \otimes Y \otimes Z$ be an arbitrary element of $k[\text{Mor}_{\Delta S}([n], [2])]$. If $|X| = 0$, $|Y| = 0$, or $|Z| = 0$, then the degeneracy relations and step 5 imply that $X \otimes Y \otimes Z$ is equivalent to an element of $k[\mathcal{B}_n]$.

Suppose now that $|X|, |Y|, |Z| \geq 1$. If x_n occurs in X , use the relation $X \otimes Y \otimes W \approx -(Y \otimes X \otimes W) \pmod{k[\mathcal{B}_n]}$ to ensure that x_n occurs in the middle factor. If x_n occurs in Z , use the sign relation and the above relation to put x_n into the middle factor. In any case, it suffices to assume our element has the form: $X \otimes U x_n V \otimes Z$. Using the modified Hochschild relation, $X \otimes U x_n V \otimes Z \approx -(Z \otimes V \otimes X U x_n) + Z \otimes V X \otimes U x_n, \pmod{k[\mathcal{B}_n]}$. The first term is certainly in $k[\mathcal{C}_n]$, since $|X| \geq 1$. If $|U| > 0$, the second term also lies in $k[\mathcal{C}_n]$. If, on the other hand, $|U| = 0$, then step 8 implies that $Z \otimes V X \otimes x_n$ is an element of $k[\mathcal{B}_n]$.

Observe that Step 9 proves Lemma 83 for $n = 0, 1, 2$, since in these cases, any elements that fall within the set $\{X \otimes Y \otimes Z x_n \mid |Z| \geq 1\}$ must have either $|X| = 0$ or $|Y| = 0$, hence are equivalent via the degeneracy relation to elements of $k[\{X \otimes x_{i_n} \otimes 1 \mid i_n \neq n\}]$. In what follows, assume $n \geq 3$.

Step 10 Every element of $k[\text{Mor}_{\Delta S}([n], [2])]$ is equivalent, modulo $k[\mathcal{B}_n]$, to a linear combination of elements from the following set:

$$(37) \quad \mathcal{D}_n \stackrel{\text{def}}{=} \{X \otimes x_{i_{n-2}} \otimes x_{n-1} x_n \mid i_{n-2} \neq n-2\} \cup \{X \otimes Y \otimes Z x_{n-1} x_n \mid |Z| \geq 1\}.$$

First, we require a relation that transports x_n from the end of a tensor:

$$(38) \quad W \otimes Z \otimes X x_n \approx W \otimes x_n Z \otimes X \pmod{k[\mathcal{B}_n]}.$$

Letting $Y = x_n$ in (31), and making use of the sign relation, we have: $W \otimes Z \otimes X x_n \approx W \otimes Z X \otimes x_n + x_n Z X \otimes W \otimes 1 + W \otimes x_n Z \otimes X$. By steps 5 and 8, $W \otimes Z \otimes X x_n \approx W \otimes x_n Z \otimes X$, modulo elements of $k[\mathcal{B}_n]$.

Now, let $X \otimes Y \otimes Z$ be an arbitrary element of $k[\text{Mor}_{\Delta S}([n], [2])]$. Locate x_{n-1} and use the techniques of Step 9 to re-express $X \otimes Y \otimes Z$ as a linear combination of terms of the form: $X_j \otimes Y_j \otimes Z_j x_{n-1}$, modulo $k[\mathcal{B}_n]$. Our goal is to re-express each term as a linear combination of vectors in which x_n occurs only in the second tensor factor.

If x_n occurs in X_j , then observe $X_j \otimes Y_j \otimes Z_j x_{n-1} \approx -(Y_j \otimes X_j \otimes Z_j x_{n-1})$, (mod $k[\mathcal{B}_n]$).

If x_n occurs in Z_j , then first substitute $Y = x_{n-1}$ into (31), obtaining the relation:

$$\begin{aligned}
 & Xx_{n-1} \otimes Z \otimes W + W \otimes ZX \otimes x_{n-1} + x_{n-1}ZX \otimes 1 \otimes W + W \otimes x_{n-1}Z \otimes X \approx 0 \\
 & \Rightarrow W \otimes Z \otimes Xx_{n-1} \approx W \otimes ZX \otimes x_{n-1} + W \otimes x_{n-1}Z \otimes X \pmod{k[\mathcal{B}_n]}
 \end{aligned}$$

By the modified Hochschild relation, $W \otimes x_{n-1}Z \otimes X \approx Wx_{n-1} \otimes Z \otimes X + ZW \otimes x_{n-1} \otimes X$, (mod $k[\mathcal{B}_n]$), then using sign relations etc., we obtain:

$$\begin{aligned}
 & W \otimes Z \otimes Xx_{n-1} \approx \\
 & W \otimes ZX \otimes x_{n-1} + Z \otimes X \otimes Wx_{n-1} - ZW \otimes X \otimes x_{n-1}, \pmod{k[\mathcal{B}_n]}.
 \end{aligned}$$

Thus, we can express our original element $X \otimes Y \otimes Z$ as a linear combination of elements of the form $X' \otimes U'x_nV' \otimes Z'x_{n-1}$, (mod $k[\mathcal{B}_n]$). Then using modified Hochschild etc., rewrite each such term as follows:

$$\begin{aligned}
 & X' \otimes U'x_nV' \otimes Z'x_{n-1} \approx \\
 & X'U' \otimes x_nV' \otimes Z'x_{n-1} - U' \otimes x_nV'X' \otimes Z'x_{n-1}, \pmod{k[\mathcal{B}_n]}.
 \end{aligned}$$

By (38), we transport x_n to the end of each term, so $X' \otimes U'x_nV' \otimes Z'x_{n-1} \approx X'U' \otimes V' \otimes Z'x_{n-1}x_n - U' \otimes V'X' \otimes Z'x_{n-1}x_n$, modulo elements of $k[\mathcal{B}_n]$. If $|Z'| \geq 1$, then we are done. Otherwise, we have some elements of the form $X'' \otimes Y'' \otimes x_{n-1}x_n$. Use an induction argument analogous to that in step 8 to re-express this type of element as a linear combination of elements of the form $U \otimes x_{i_{n-2}} \otimes x_{n-1}x_n$ such that $i_{n-2} \neq n-2$, modulo elements of $k[\mathcal{B}_n]$.

Step 11 Every element of $k[\text{Mor}_{\Delta S}([n], [2])]$ is equivalent to an element of $k[\mathcal{B}_n]$.

We shall use an iterative re-writing procedure. First of all, define sets:

$$\begin{aligned}
 \mathcal{B}_n^j & \stackrel{\text{def}}{=} \{A \otimes x_{i_{n-j}} \otimes x_{n-j+1} \dots x_n \mid i_{n-j} \neq n-j\}, \\
 \mathcal{C}_n^j & \stackrel{\text{def}}{=} \{A \otimes B \otimes Cx_{n-j+1} \dots x_n \mid |C| \geq 1\}.
 \end{aligned}$$

Now clearly, $\mathcal{B}_n = \bigcup_{j=0}^{n-1} \mathcal{B}_n^j$. In what follows, ‘reduced’ will always mean reduced modulo elements of $k[\mathcal{B}_n]$. By steps 9 and 10, we can reduce an arbitrary element $X \otimes Y \otimes Z$ to linear combinations of elements in $\mathcal{B}_n^0 \cup \mathcal{B}_n^1 \cup \mathcal{B}_n^2 \cup \mathcal{C}_n^2$. Suppose now that we have reduced elements to linear combinations of elements from the set $\mathcal{B}_n^0 \cup \mathcal{B}_n^1 \cup \dots \cup \mathcal{B}_n^j \cup \mathcal{C}_n^j$, for some $j \geq 2$. I claim any element of \mathcal{C}_n^j can be re-expressed as a linear combination of elements from the set $\mathcal{B}_n^0 \cup \mathcal{B}_n^1 \cup \dots \cup \mathcal{B}_n^{j+1} \cup \mathcal{C}_n^{j+1}$. Indeed, let $X \otimes Y \otimes Zx_{n-j+1} \dots x_n$, with $|Z| \geq 1$. Let $w \stackrel{\text{def}}{=} x_{n-j+1} \dots x_n$. We may now

think of $X \otimes Y \otimes Z w$ as consisting of the ‘indeterminates’ $x_0, x_1, \dots, x_{n-j}, w$, hence, by step 10, we may reduce this element to a linear combination of elements from the set $\{X \otimes x_{i_{n-j-1}} \otimes x_{n-j} w \mid i_{n-j-1} \neq n-j-1\} \cup \{X \otimes Y \otimes Z x_{n-j} w \mid |Z| \geq 1\}$. This implies the element may be written as a linear combination of elements from the set $\mathcal{B}_n^{j+1} \cup \mathcal{C}_n^{j+1}$, modulo elements of the form $A \otimes B \otimes 1$ and $A \otimes B \otimes x_{n-j+1} \dots x_n$. Since $\{A \otimes B \otimes x_{n-j+1} x_{n-j+2} \dots x_n\} \subseteq \mathcal{C}_n^{j-1}$, the inductive hypothesis ensures that there is set containment $\{A \otimes B \otimes x_{n-j+1} \dots x_n\} \subseteq \mathcal{B}_n^0 \cup \dots \cup \mathcal{B}_n^j$. This completes the inductive step.

After a finite number of iterations, then, we can re-express any element $X \otimes Y \otimes Z$ as a linear combination from the set $\mathcal{B}_n^0 \cup \dots \cup \mathcal{B}_n^{n-1} \cup \mathcal{C}_n^{n-1} = \mathcal{B}_n \cup \mathcal{C}_n^{n-1}$. But $\mathcal{C}_n^{n-1} = \{A \otimes B \otimes C x_2 \dots x_n \mid |C| \geq 1\}$. Any element from this set has either $|A| = 0$ or $|B| = 0$, therefore is equivalent to an element of $k[\mathcal{B}_n]$ already. \square

Corollary 84 *If $\frac{1}{2} \in k$, then the four-term relation $XY \otimes Z \otimes W + W \otimes ZX \otimes Y + YZX \otimes 1 \otimes W + W \otimes YZ \otimes X \approx 0$ is sufficient to collapse $k[\text{Mor}_{\Delta S}([n], [2])]$ onto $k[\mathcal{B}_n]$.*

Proof We only need to modify step 1 of the previous proof. We will establish that $X \otimes 1 \otimes 1 \approx 1 \otimes X \otimes 1 \approx 1 \otimes 1 \otimes X \approx 0$.

Setting three variables at a time equal to 1 in (31) we obtain,

$$(39) \quad 2(W \otimes 1 \otimes 1) + 2(1 \otimes 1 \otimes W) \approx 0, \quad \text{when } X = Y = Z = 1.$$

$$(40) \quad Z \otimes 1 \otimes 1 + 3(1 \otimes Z \otimes 1) \approx 0, \quad \text{when } X = Y = W = 1.$$

$$(41) \quad 2(Y \otimes 1 \otimes 1) + 1 \otimes Y \otimes 1 + 1 \otimes 1 \otimes Y \approx 0, \quad \text{when } X = Z = W = 1.$$

Equivalently, we have a system of linear equations,

$$\begin{bmatrix} 2 & 0 & 2 \\ 1 & 3 & 0 \\ 2 & 1 & 1 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} = 0,$$

where $z_1 = X \otimes 1 \otimes 1$, $z_2 = 1 \otimes X \otimes 1$, and $z_3 = 1 \otimes 1 \otimes X$. Since the determinant of the coefficient matrix is -4 , the matrix is invertible in the ring k as long as $1/2 \in k$. \square

Define for each $m \geq 0$, $P_m \stackrel{\text{def}}{=} k[\text{Mor}_{\Delta S}(-, [m])]$ and $P_m^n = P_m([n])$. Lemma 83 together with Lemmas 82 and 79 show the following sequence of k -modules is exact for each $n \geq 0$:

$$(42) \quad 0 \longleftarrow k \xleftarrow{\epsilon} P_0^n \xleftarrow{\rho} P_2^n \xleftarrow{(\alpha, \beta)} P_3^n \oplus P_0^n$$

where $\alpha: P_3^n \rightarrow P_2^n$ is given by composition with the ΔS morphism

$$x_0x_1 \otimes x_2 \otimes x_3 + x_3 \otimes x_2x_0 \otimes x_1 + x_1x_2x_0 \otimes 1 \otimes x_3 + x_3 \otimes x_1x_2 \otimes x_0,$$

and $\beta: P_0^n \rightarrow P_2^n$ is induced by $1 \otimes x_0 \otimes 1$. This holds for all $n \geq 0$, so we have constructed a partial resolution of \underline{k} by projective ΔS^{op} -modules:

$$(43) \quad 0 \longleftarrow k \xleftarrow{\epsilon} P_0 \xleftarrow{\rho} P_2 \xleftarrow{(\alpha, \beta)} P_3 \oplus P_0$$

11 Using the partial resolution for low degree computations

Let A be a unital associative algebra over k . Since (43) is a partial resolution of \underline{k} , it can be used to find $HS_i(A)$ for $i = 0, 1$.

11.1 Main theorem

Theorem 85 *$HS_i(A)$ for $i = 0, 1$ may be computed as the degree 0 and degree 1 homology groups of the following (partial) chain complex:*

$$(44) \quad 0 \longleftarrow A \xleftarrow{\partial_1} A \otimes A \otimes A \xleftarrow{\partial_2} (A \otimes A \otimes A \otimes A) \oplus A,$$

where

$$\partial_1: a \otimes b \otimes c \mapsto abc - cba,$$

$$\partial_2: \begin{cases} a \otimes b \otimes c \otimes d & \mapsto ab \otimes c \otimes d + d \otimes ca \otimes b + bca \otimes 1 \otimes d + d \otimes bc \otimes a, \\ a & \mapsto 1 \otimes a \otimes 1. \end{cases}$$

Proof Tensoring the complex (43) with $B_*^{\text{sym}} A$ over ΔS , we obtain a complex that computes $HS_0(A)$ and $HS_1(A)$. The statement of the theorem then follows from isomorphisms induced by the evaluation map

$$k [\text{Mor}_{\Delta S}(-, [p])] \otimes_{\Delta S} B_*^{\text{sym}} A \xrightarrow{\cong} B_p^{\text{sym}} A. \quad \square$$

11.2 Degree 0 symmetric homology

Theorem 86 *For a unital associative algebra A over commutative ground ring k , $HS_0(A) \cong A/([A, A])$, where $([A, A])$ is the ideal generated by the commutator submodule $[A, A]$.*

Proof By Theorem 5, $HS_0(A) \cong A/k[\{abc - cba\}]$ as a k -module. But $k[\{abc - cba\}]$ is an ideal of A . Now clearly $[A, A] \subseteq k[\{abc - cba\}]$. On the other hand, $k[\{abc - cba\}] \subseteq ([A, A])$ since $abc - cba = a(bc - cb) + a(cb) - (cb)a$. \square

Corollary 87 *If A is commutative, then $HS_0(A) \cong A$.*

Remark 88 [Theorem 86](#) implies that symmetric homology does not preserve Morita equivalence, since for $n > 1$, $HS_0(M_n(A)) = M_n(A) / ([M_n(A), M_n(A)]) = 0$, while in general $HS_0(A) = A / ([A, A]) \neq 0$.

11.3 Degree 1 symmetric homology

Using GAP, we have made the following explicit computations of degree 1 integral symmetric homology. See [Section 13](#) for a discussion of how computer algebra systems were used in symmetric homology computations.

A	$HS_1(A \mathbb{Z})$
$\mathbb{Z}[t]/(t^2)$	$\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$
$\mathbb{Z}[t]/(t^3)$	$\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$
$\mathbb{Z}[t]/(t^4)$	$(\mathbb{Z}/2\mathbb{Z})^4$
$\mathbb{Z}[t]/(t^5)$	$(\mathbb{Z}/2\mathbb{Z})^4$
$\mathbb{Z}[t]/(t^6)$	$(\mathbb{Z}/2\mathbb{Z})^6$
$\mathbb{Z}[C_2]$	$\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$
$\mathbb{Z}[C_3]$	0
$\mathbb{Z}[C_4]$	$(\mathbb{Z}/2\mathbb{Z})^4$
$\mathbb{Z}[C_5]$	0
$\mathbb{Z}[C_6]$	$(\mathbb{Z}/2\mathbb{Z})^6$

Based on these calculations, we conjecture:

Conjecture 89

$$HS_1(k[t]/(t^n)) = \begin{cases} (k/2k)^n, & \text{if } n \geq 0 \text{ is even.} \\ (k/2k)^{n-1} & \text{if } n \geq 1 \text{ is odd.} \end{cases}$$

Remark 90 The computations of $HS_1(\mathbb{Z}[C_n])$ are consistent with those of Brown and Loday [\[2\]](#). See [Section 11.5](#) for a more detailed treatment of HS_1 for group rings.

Additionally, HS_1 has been computed for the following examples. These computations were done using GAP in some cases and in others, Fermat [\[15\]](#) computations on sparse matrices were used in conjunction with the GAP scripts. (for example, when the algebra has dimension greater than 6 over \mathbb{Z}).

A	$HS_1(A \mathbb{Z})$
$\mathbb{Z}[t, u]/(t^2, u^2)$	$\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^{11}$
$\mathbb{Z}[t, u]/(t^3, u^2)$	$\mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^{11} \oplus \mathbb{Z}/6\mathbb{Z}$
$\mathbb{Z}[t, u]/(t^3, u^2, t^2u)$	$\mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^{10}$
$\mathbb{Z}[t, u]/(t^3, u^3)$	$\mathbb{Z}^4 \oplus (\mathbb{Z}/2\mathbb{Z})^7 \oplus (\mathbb{Z}/6\mathbb{Z})^5$
$\mathbb{Z}[t, u]/(t^2, u^4)$	$\mathbb{Z}^3 \oplus (\mathbb{Z}/2\mathbb{Z})^{20} \oplus \mathbb{Z}/4\mathbb{Z}$
$\mathbb{Z}[t, u, v]/(t^2, u^2, v^2)$	$\mathbb{Z}^6 \oplus (\mathbb{Z}/2\mathbb{Z})^{42}$
$\mathbb{Z}[t, u]/(t^4, u^3)$	$\mathbb{Z}^6 \oplus (\mathbb{Z}/2\mathbb{Z})^{19} \oplus \mathbb{Z}/6\mathbb{Z} \oplus (\mathbb{Z}/12\mathbb{Z})^2$
$\mathbb{Z}[t, u, v]/(t^2, u^2, v^3)$	$\mathbb{Z}^{11} \oplus (\mathbb{Z}/2\mathbb{Z})^{45} \oplus (\mathbb{Z}/6\mathbb{Z})^4$
$\mathbb{Z}[i, j, k], i^2 = j^2 = k^2 = ijk = -1$	$(\mathbb{Z}/2\mathbb{Z})^8$
$\mathbb{Z}[C_2 \times C_2]$	$(\mathbb{Z}/2\mathbb{Z})^{12}$
$\mathbb{Z}[C_3 \times C_2]$	$(\mathbb{Z}/2\mathbb{Z})^6$
$\mathbb{Z}[C_3 \times C_3]$	$(\mathbb{Z}/3\mathbb{Z})^9$
$\mathbb{Z}[S_3]$	$(\mathbb{Z}/2\mathbb{Z})^2$

11.4 Splittings of the partial complex

Under certain circumstances, the partial complex in [Theorem 5](#) splits as a direct sum of smaller complexes. This observation becomes increasingly important as the dimension of the algebra increases. Indeed, some of the computations of the previous section were done using splittings.

Definition 91 For a commutative k -algebra A and $u \in A$, define the k -modules:

$$(A^{\otimes n})_u \stackrel{\text{def}}{=} \{a_1 \otimes a_2 \otimes \dots \otimes a_n \in A^{\otimes n} \mid a_1 a_2 \dots a_n = u\}$$

Proposition 92 If $A = k[M]$ for a commutative monoid M , then complex (44) splits as a direct sum of complexes

$$(45) \quad 0 \longleftarrow (A)_u \xleftarrow{\partial_1} (A \otimes A \otimes A)_u \xleftarrow{\partial_2} (A \otimes A \otimes A \otimes A)_u \oplus (A)_u,$$

where u ranges over the elements of M . Thus, for $i = 0, 1$, we have $HS_i(A) \cong \bigoplus_{u \in M} HS_i(A)_u$.

Proof Since M is a commutative monoid, there are direct sum decompositions as k -module: $A^{\otimes n} = \bigoplus_{u \in M} (A^{\otimes n})_u$. The boundary maps ∂_1 and ∂_2 preserve the products of tensor factors, so the inclusions $(A^{\otimes n})_u \hookrightarrow A^{\otimes n}$ induce maps of complexes, hence the complex itself splits as a direct sum. □

Definition 93 For each u , the homology groups of complex (45) will be called the u -layered symmetric homology of A , denoted $HS_i(A)_u$.

We may use layers to investigate the symmetric homology of $k[t]$. This algebra is monoidal, generated by the monoid $\{1, t, t^2, t^3, \dots\}$. Now, the t^m -layer symmetric homology of $k[t]$ will be the same as the t^m -layer symmetric homology of $k[M_{m+1}^{m+2}]$, where M_q^p denotes the cyclic monoid generated by an indeterminate s with the property that $s^p = s^q$. Using this observation and subsequent computation, we conjecture:

Conjecture 94

$$HS_1(k[t])_{t^m} = \begin{cases} 0 & m = 0, 1 \\ k/2k, & m \geq 2 \end{cases}$$

This conjecture has been verified up to $m = 18$, in the case $k = \mathbb{Z}$.

11.5 2-torsion in HS_1

The occurrence of 2-torsion in $HS_1(A)$ for the examples considered in Sections 11.3 and 11.4 comes as no surprise, based on Theorem 2. First consider the following chain of isomorphisms:

$$\begin{aligned} \pi_2^s(B\Gamma) &= \pi_2(\Omega^\infty S^\infty(B\Gamma)) \cong \pi_1(\Omega\Omega^\infty S^\infty(B\Gamma)) \\ &\cong \pi_1(\Omega_0\Omega^\infty S^\infty(B\Gamma)) \xrightarrow{h} H_1(\Omega_0\Omega^\infty S^\infty(B\Gamma)) \end{aligned}$$

Here, $\Omega_0\Omega^\infty S^\infty(B\Gamma)$ denotes the component of the constant loop, and h is the Hurewicz homomorphism, which is an isomorphism since $\Omega_0\Omega^\infty S^\infty(B\Gamma)$ is path-connected and π_1 is abelian (since it is actually π_2 of a space).

On the other hand, by Theorem 2,

$$HS_1(k[\Gamma]) \cong H_1(\Omega\Omega^\infty S^\infty(B\Gamma); k) \cong H_1(\Omega_0\Omega^\infty S^\infty(B\Gamma)) \otimes k.$$

Note, all tensor products will be over \mathbb{Z} in this section. Now $\Omega\Omega^\infty S^\infty(B\Gamma)$ consists of disjoint homeomorphic copies of $\Omega_0\Omega^\infty S^\infty(B\Gamma)$, one for each element of $\Gamma/[\Gamma, \Gamma]$, (where $[\Gamma, \Gamma]$ is the commutator subgroup of Γ), so we may write

$$H_1(\Omega\Omega^\infty S^\infty(B\Gamma)) \otimes k \cong H_1(\Omega_0\Omega^\infty S^\infty(B\Gamma)) \otimes k [\Gamma/[\Gamma, \Gamma]]$$

to obtain the following result:

Proposition 95 *If Γ is a group, then $HS_1(k[\Gamma]) \cong \pi_2^s(B\Gamma) \otimes k [\Gamma/[\Gamma, \Gamma]]$.*

As an immediate corollary, if Γ is abelian, then $HS_1(k[\Gamma]) \cong \pi_2^s(B\Gamma) \otimes k[\Gamma]$. Moreover, by results of Brown and Loday [2], if Γ is abelian, then $\pi_2^s(B\Gamma)$ is the reduced tensor square, $\Gamma \tilde{\wedge} \Gamma = (\Gamma \otimes \Gamma) / \approx$, where $g \otimes h \approx -h \otimes g$ for all $g, h \in \Gamma$.

Proposition 96

$$HS_1(k[C_n]) = \begin{cases} (\mathbb{Z}/2\mathbb{Z})^n & n \text{ even,} \\ 0 & n \text{ odd.} \end{cases}$$

Proof $\pi_2^s(BC_n) = \mathbb{Z}/2\mathbb{Z}$ if n is even, and 0 if n is odd. The result then follows from Proposition 95, as $k[C_n/[C_n, C_n]] \cong k[C_n] \cong k^n$, as k -module. \square

12 Relations to cyclic homology

The relation between the symmetric bar construction and the cyclic bar construction arising from the inclusions $\Delta C \hookrightarrow \Delta S$ gives rise to a natural map $HC_*(A) \rightarrow HS_*(A)$. Indeed, by Remark 17, we may define cyclic homology thus: $HC_*(A) = \text{Tor}_*^{\Delta C}(\underline{k}, B_*^{\text{sym}} A)$, where we understand $B_*^{\text{sym}} A$ as the restriction of the functor to ΔC .

Using the partial complex of Theorem 5, and an analogous one for computing cyclic homology (see Loday [16, page 59]), the map $HC_*(A) \rightarrow HS_*(A)$ for degrees 0 and 1 is induced by the following partial chain map:

$$\begin{array}{ccccccc} 0 & \longleftarrow & A & \xleftarrow{\partial_1^C} & A \otimes A & \xleftarrow{\partial_2^C} & A^{\otimes 3} \oplus A \\ & & \downarrow \gamma_0 = \text{id} & & \downarrow \gamma_1 & & \downarrow \gamma_2 \\ 0 & \longleftarrow & A & \xleftarrow{\partial_1^S} & A^{\otimes 3} & \xleftarrow{\partial_2^S} & A^{\otimes 4} \oplus A \end{array}$$

In this diagram, the boundary maps in the upper row are defined as follows:

$$\begin{aligned} \partial_1^C: a \otimes b &\mapsto ab - ba \\ \partial_2^C: \begin{cases} a \otimes b \otimes c &\mapsto ab \otimes c - a \otimes bc + ca \otimes b \\ a &\mapsto 1 \otimes a - a \otimes 1 \end{cases} \end{aligned}$$

The boundary maps in the lower row are defined as in Theorem 5.

$$\begin{aligned} \partial_1^S: a \otimes b \otimes c &\mapsto abc - cba \\ \partial_2^S: \begin{cases} a \otimes b \otimes c \otimes d &\mapsto ab \otimes c \otimes d - d \otimes ca \otimes b + bca \otimes 1 \otimes d + d \otimes bc \otimes a \\ a &\mapsto 1 \otimes a \otimes 1 \end{cases} \end{aligned}$$

The partial chain map is given in degree 1 by $\gamma_1(a \otimes b) \stackrel{\text{def}}{=} a \otimes b \otimes 1$. In degree 2, γ_2 is defined on the summand $A^{\otimes 3}$ via

$$a \otimes b \otimes c \mapsto (a \otimes b \otimes c \otimes 1 - 1 \otimes a \otimes bc \otimes 1 + 1 \otimes ca \otimes b \otimes 1 + 1 \otimes 1 \otimes abc \otimes 1 - b \otimes ca \otimes 1 \otimes 1, -2abc - cab),$$

and on the summand A via

$$a \mapsto (-1 \otimes 1 \otimes a \otimes 1, 4a).$$

12.1 Examples

To provide some examples, consider the maps $\gamma_1: HC_1(\mathbb{Z}[t]/(t^n)) \rightarrow HS_1(\mathbb{Z}[t]/(t^n))$.

It can be shown (for example, by direct computation) that $HC_1(\mathbb{Z}[t]/(t^2)) \cong \mathbb{Z}/2\mathbb{Z}$ is generated by the 1-chain $t \otimes t$. $\gamma_1(t \otimes t) = t \otimes t \otimes 1 \in HS_1(\mathbb{Z}[t]/(t^2))$ is a non-trivial element of $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ (which may be verified by direct computation as well).

The map $HC_1(\mathbb{Z}[t]/(t^3)) \rightarrow HS_1(\mathbb{Z}[t]/(t^3))$ may be similarly analyzed. Here, the chain $t \otimes t + t \otimes t^2$ is a generator of $HC_1(\mathbb{Z}[t]/(t^3)) \cong \mathbb{Z}/6\mathbb{Z}$, which gets sent by γ_1 to $t \otimes t \otimes 1 + t \otimes t^2 \otimes 1$, a non-trivial element of $HS_1(\mathbb{Z}[t]/(t^3)) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$.

The case $n = 4$ is bit more interesting. Here, $HC_1(\mathbb{Z}[t]/(t^4)) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$, generated by $t \otimes t$ and $t \otimes t^2 + t \otimes t^3$, respectively. The image of the map γ_1 in $HS_1(\mathbb{Z}[t]/(t^4))$ is $(\mathbb{Z}/2\mathbb{Z})^2 \subseteq (\mathbb{Z}/2\mathbb{Z})^4$.

13 Using computer algebra systems for computing symmetric homology

The computer algebra systems GAP, Octave and Fermat were used to verify proposed theorems and also to obtain some concrete computations of symmetric homology for some small algebras. A tar-file of the scripts that were created and used for this work is available at <http://arxiv.org/e-print/0807.4521v1/>. This tar-file contains the following files:

- Basic.g – Some elementary functions, necessary for some functions in DeltaS.g
- HomAlg.g – Homological Algebra functions, such as computation of homology groups for chain complexes.
- Fermat.g – Functions necessary to invoke Fermat for fast sparse matrix computations.

- `fermattogap`, `gaptofermat` – Auxiliary text files for use when invoking Fermat from GAP.
- `DeltaS.g` – This is the main repository of scripts used to compute various quantities associated with the category ΔS , including $HS_1(A)$ for finite-dimensional algebras A .

In order to use the functions of `DeltaS.g`, simply copy the above files into the working directory (such as `~/gap/`), invoke GAP, then read in `DeltaS.g` at the prompt. The dependent modules will automatically be loaded (hence they must be present in the same directory as `DeltaS.g`). Note that most of the computations involving homology require substantial memory to run. I recommend calling GAP with the command line option “`-o mem`”, where *mem* is the amount of memory to be allocated to this instance of GAP. All computations done in this dissertation can be accomplished by allocating 20 gigabytes of memory. The following provides a few examples of using the functions of `DeltaS.g`

```
[ault@math gap]$ gap -o 20g

gap> Read("DeltaS.g");
gap>
gap> ## Number of morphisms [6] --> [4]
gap> SizeDeltaS( 6, 4 );
1663200
gap>
gap> ## Generate the set of morphisms of Delta S, [2] --> [2]
gap> EnumerateDeltaS( 2, 2 );
[[ [ 0, 1, 2 ], [ ], [ ] ], [ [ 0, 2, 1 ], [ ], [ ] ],
 [ [ 1, 0, 2 ], [ ], [ ] ], [ [ 1, 2, 0 ], [ ], [ ] ],
 [ [ 2, 0, 1 ], [ ], [ ] ], [ [ 2, 1, 0 ], [ ], [ ] ],
 [ [ 0, 1 ], [ 2 ], [ ] ], [ [ 0, 2 ], [ 1 ], [ ] ],
 [ [ 1, 0 ], [ 2 ], [ ] ], [ [ 1, 2 ], [ 0 ], [ ] ],
 [ [ 2, 0 ], [ 1 ], [ ] ], [ [ 2, 1 ], [ 0 ], [ ] ],
 [ [ 0, 1 ], [ ], [ 2 ] ], [ [ 0, 2 ], [ ], [ 1 ] ],
 [ [ 1, 0 ], [ ], [ 2 ] ], [ [ 1, 2 ], [ ], [ 0 ] ],
 [ [ 2, 0 ], [ ], [ 1 ] ], [ [ 2, 1 ], [ ], [ 0 ] ],
 [ [ 0 ], [ 1, 2 ], [ ] ], [ [ 0 ], [ 2, 1 ], [ ] ],
 [ [ 1 ], [ 0, 2 ], [ ] ], [ [ 1 ], [ 2, 0 ], [ ] ],
 [ [ 2 ], [ 0, 1 ], [ ] ], [ [ 2 ], [ 1, 0 ], [ ] ],
 [ [ 0 ], [ 1 ], [ 2 ] ], [ [ 0 ], [ 2 ], [ 1 ] ], [ [ 1 ], [ 0 ], [ 2 ] ],
 [ [ 1 ], [ 2 ], [ 0 ] ], [ [ 2 ], [ 0 ], [ 1 ] ], [ [ 2 ], [ 1 ], [ 0 ] ],
 [ [ 0 ], [ ], [ 1, 2 ] ], [ [ 0 ], [ ], [ 2, 1 ] ],
 [ [ 1 ], [ ], [ 0, 2 ] ], [ [ 1 ], [ ], [ 2, 0 ] ],
 [ [ 2 ], [ ], [ 0, 1 ] ], [ [ 2 ], [ ], [ 1, 0 ] ],
 [ [ ], [ 0, 1, 2 ], [ ] ], [ [ ], [ 0, 2, 1 ], [ ] ],
```

```

[[ ], [1, 0, 2], [ ]], [[ ], [1, 2, 0], [ ]],
[[ ], [2, 0, 1], [ ]], [[ ], [2, 1, 0], [ ]],
[[ ], [0, 1], [2]], [[ ], [0, 2], [1]],
[[ ], [1, 0], [2]], [[ ], [1, 2], [0]],
[[ ], [2, 0], [1]], [[ ], [2, 1], [0]],
[[ ], [0], [1, 2]], [[ ], [0], [2, 1]],
[[ ], [1], [0, 2]], [[ ], [1], [2, 0]],
[[ ], [2], [0, 1]], [[ ], [2], [1, 0]],
[[ ], [ ], [0, 1, 2]], [[ ], [ ], [0, 2, 1]],
[[ ], [ ], [1, 0, 2]], [[ ], [ ], [1, 2, 0]],
[[ ], [ ], [2, 0, 1]], [[ ], [ ], [2, 1, 0]] ] ]

gap>
gap> ## Generate only the epimorphisms [2] --> [2]
gap> EnumerateDeltaS( 2, 2 : epi );
[[ [ 0 ], [ 1 ], [ 2 ] ], [ [ 0 ], [ 2 ], [ 1 ] ] ],
[[ [ 1 ], [ 0 ], [ 2 ] ], [ [ 1 ], [ 2 ], [ 0 ] ] ],
[[ [ 2 ], [ 0 ], [ 1 ] ], [ [ 2 ], [ 1 ], [ 0 ] ] ] ]

gap>
gap> ## Compose two morphisms of Delta S.
gap> a := Random(EnumerateDeltaS(4,3));
[[ 0 ], [ 2, 4, 1 ], [ ], [ 3 ] ]
gap> b := Random(EnumerateDeltaS(3,2));
[[ ], [ 3, 0, 2 ], [ 1 ] ]
gap> MultDeltaS(b, a);
[[ ], [ 3, 0 ], [ 2, 4, 1 ] ]
gap> MultDeltaS(a, b);
Maps incomposeable
[ ]

gap>
gap> ## Examples of using morphisms of Delta S to act on simple tensors
gap> A := TruncPolyAlg([3,2]);
<algebra of dimension 6 over Rationals>
gap> ## TruncPolyAlg is defined in Basic.g
gap> ## TruncPolyAlg([i_1, i_2, ..., i_n]) is generated by
gap> ## x_1, x_2, ..., x_n, under the relation (x_j)^(i_j) = 0.
gap> g := GeneratorsOfLeftModule(A);
[ X^[ 0, 0 ], X^[ 0, 1 ], X^[ 1, 0 ], X^[ 1, 1 ], X^[ 2, 0 ], X^[ 2, 1 ] ]
gap> x := g[2]; y := g[3];
X^[ 0, 1 ]
X^[ 1, 0 ]
gap> v := [ x*y, 1, y^2 ];
gap> ## v represents the simple tensor xy \otimes 1 \otimes y^2.
[ X^[ 1, 1 ], 1, X^[ 2, 0 ] ]
gap> ActByDeltaS( v, [[2], [], [0], [1]] );
[ X^[ 2, 0 ], 1, X^[ 1, 1 ], 1 ]
gap> ActByDeltaS( v, [[2], [0,1]] );
[ X^[ 2, 0 ], X^[ 1, 1 ] ]
gap> ActByDeltaS( v, [[2,0], [1]] );

```

```

[ 0*X^[ 0, 0 ], 1 ]
gap>
gap> ## Symmetric monoidal product on DeltaS_+
gap> a := Random(EnumerateDeltaS(4,2));
[ [ ], [ 2, 1, 0 ], [ 3, 4 ] ]
gap> b := Random(EnumerateDeltaS(3,3));
[ [ ], [ ], [ ], [ 1, 3, 2, 0 ] ]
gap> MonoidProductDeltaS(a, b);
[ [ ], [ 2, 1, 0 ], [ 3, 4 ], [ ], [ ], [ ], [ 6, 8, 7, 5 ] ]
gap> MonoidProductDeltaS(b, a);
[ [ ], [ ], [ ], [ 1, 3, 2, 0 ], [ ], [ 6, 5, 4 ], [ 7, 8 ] ]
gap> MonoidProductDeltaS(a, []);
[ [ ], [ 2, 1, 0 ], [ 3, 4 ] ]
gap>
gap> ## Symmetric Homology of the algebra A, in degrees 0 and 1.
gap> SymHomUnitalAlg(A);
[ [ 0, 0, 0, 0, 0, 0 ], [ 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 6, 0, 0 ] ]
gap> ## '0' represents a factor of Z, while a non-zero p represents
gap> ## a factor of Z/pZ.
gap>
gap> ## Using layers to compute symmetric homology
gap> C2 := CyclicGroup(2);
<pc group of size 2 with 1 generators>
gap> A := GroupRing(Rationals, DirectProduct(C2, C2));
<algebra-with-one over Rationals, with 2 generators>
gap> ## First, a direct computation without layers:
gap> SymHomUnitalAlg(A);
[ [ 0, 0, 0, 0 ], [ 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2 ] ]
gap> ## Next, compute HS_0(A)_u and HS_1(A)_u for each generator u.
gap> g := GeneratorsOfLeftModule(A);
[ (1)*<identity> of ..., (1)*f2, (1)*f1, (1)*f1*f2 ]
gap> SymHomUnitalAlgLayered(A, g[1]);
[ [ 0 ], [ 2, 2, 2 ] ]
gap> SymHomUnitalAlgLayered(A, g[2]);
[ [ 0 ], [ 2, 2, 2 ] ]
gap> SymHomUnitalAlgLayered(A, g[3]);
[ [ 0 ], [ 2, 2, 2 ] ]
gap> SymHomUnitalAlgLayered(A, g[4]);
[ [ 0 ], [ 2, 2, 2 ] ]
gap> ## Computing HS_1( Z[t] ) by layers:
gap> SymHomFreeMonoid(0,10);
HS_1(k[t])_{t^0} : [ ]
HS_1(k[t])_{t^1} : [ ]
HS_1(k[t])_{t^2} : [ 2 ]
HS_1(k[t])_{t^3} : [ 2 ]
HS_1(k[t])_{t^4} : [ 2 ]
HS_1(k[t])_{t^5} : [ 2 ]
HS_1(k[t])_{t^6} : [ 2 ]

```

```

HS_1(k[t]_{t^7}) : [ 2 ]
HS_1(k[t]_{t^8}) : [ 2 ]
HS_1(k[t]_{t^9}) : [ 2 ]
HS_1(k[t]_{t^10}) : [ 2 ]
gap> ## Poincare polynomial of \Sym_*^{(p)} for small p.
gap> ## There is a check for torsion, using a call to Fermat
gap> ## to find Smith Normal Form of the differential matrices.
gap> PoincarePolynomialSymComplex(2);
C_0 Dimension: 1
C_1 Dimension: 6
C_2 Dimension: 6
D_1
SNF(D_1)
D_2
SNF(D_2)
2*t^2+t
gap> PoincarePolynomialSymComplex(5);
C_0 Dimension: 1
C_1 Dimension: 30
C_2 Dimension: 300
C_3 Dimension: 1200
C_4 Dimension: 1800
C_5 Dimension: 720
D_1
SNF(D_1)
D_2
SNF(D_2)
D_3
SNF(D_3)
D_4
SNF(D_4)
D_5
SNF(D_5)
120*t^5+272*t^4+t^3

```

References

- [1] **KS Brown**, *Cohomology of groups*, Graduate Texts in Mathematics 87, Springer, New York (1982) [MR672956](#)
- [2] **R Brown, J-L Loday**, *Van Kampen theorems for diagrams of spaces*, Topology 26 (1987) 311–335 [MR899052](#) With an appendix by M Zisman
- [3] **RR Bruner, JP May, JE McClure, M Steinberger**, *H_∞ ring spectra and their applications*, Lecture Notes in Mathematics 1176, Springer, Berlin (1986) [MR836132](#)
- [4] **FR Cohen, T J Lada, JP May**, *The homology of iterated loop spaces*, Lecture Notes in Mathematics 533, Springer, Berlin (1976) [MR0436146](#)

- [5] **FR Cohen, FP Peterson**, *On the homology of certain spaces looped beyond their connectivity*, Israel J. Math. 66 (1989) 105–131 [MR1017156](#)
- [6] **A Dold**, *Universelle Koeffizienten*, Math. Z. 80 (1962) 63–88 [MR0140558](#)
- [7] **JW Eaton**, *GNU Octave Manual* (2002) Available at <http://www.gnu.org/software/octave/doc/interpreter/>
- [8] **Z Fiedorowicz**, *The symmetric bar construction*, preprint Available at <http://www.math.ohio-state.edu/~fiedorow/>
- [9] **Z Fiedorowicz**, *Classifying spaces of topological monoids and categories*, Amer. J. Math. 106 (1984) 301–350 [MR737777](#)
- [10] **Z Fiedorowicz, J-L Loday**, *Crossed simplicial groups and their associated homology*, Trans. Amer. Math. Soc. 326 (1991) 57–87 [MR998125](#)
- [11] **P Gabriel, M Zisman**, *Calculus of fractions and homotopy theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete 35, Springer New York, New York (1967) [MR0210125](#)
- [12] **TG Group**, *GAP – Groups, Algorithms and Programming*, version 4.4.10 (2007) Available at <http://www.gap-system.org/>
- [13] **K Itô**, *Encyclopedic Dictionary of Mathematics* (1993)
- [14] **DM Kan**, *Adjoint functors*, Trans. Amer. Math. Soc. 87 (1958) 294–329 [MR0131451](#)
- [15] **RH Lewis**, *Fermat computer algebra system* (2008) Available at <http://home.bway.net/lewis>
- [16] **J-L Loday**, *Cyclic homology*, second edition, Grundlehren der Mathematischen Wissenschaften 301, Springer, Berlin (1998) [MR1600246](#) Appendix E by María O Ronco, Chapter 13 by the author in collaboration with Teimuraz Pirashvili
- [17] **S Mac Lane**, *The Milgram bar construction as a tensor product of functors*, from: “The Steenrod Algebra and its Applications (Proc. Conf. to Celebrate N. E. Steenrod’s Sixtieth Birthday, Battelle Memorial Inst., Columbus, Ohio, 1970)”, Lecture Notes in Mathematics 168, Springer, Berlin (1970) 135–152 [MR0273618](#)
- [18] **S MacLane**, *Categories for the working mathematician*, Graduate Texts in Mathematics 5, Springer, New York (1971) [MR0354798](#)
- [19] **JP May**, *The geometry of iterated loop spaces*, Lectures Notes in Mathematics 271, Springer, Berlin (1972) [MR0420610](#)
- [20] **JP May, R Thomason**, *The uniqueness of infinite loop space machines*, Topology 17 (1978) 205–224 [MR508885](#)
- [21] **J McCleary**, *A user’s guide to spectral sequences*, second edition, Cambridge Studies in Advanced Mathematics 58, Cambridge University Press, Cambridge (2001) [MR1793722](#)

- [22] **T Pirashvili**, *On the PROP corresponding to bialgebras*, Cah. Topol. Géom. Différ. Catég. 43 (2002) 221–239 [MR1928233](#)
- [23] **T Pirashvili, B Richter**, *Hochschild and cyclic homology via functor homology*, *K–Theory* 25 (2002) 39–49 [MR1899698](#)
- [24] **D Quillen**, *Higher algebraic K–theory I*, from: “Algebraic K–theory, I: Higher K–theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)”, Springer, Berlin (1973) 85–147. Lecture Notes in Math., Vol. 341 [MR0338129](#)
- [25] **J J Rotman**, *An introduction to homological algebra*, Pure and Applied Mathematics 85, Academic Press [Harcourt Brace Jovanovich Publishers], New York (1979) [MR538169](#)
- [26] **G Segal**, *Classifying spaces and spectral sequences*, Inst. Hautes Études Sci. Publ. Math. (1968) 105–112 [MR0232393](#)
- [27] **J Słomińska**, *Homotopy colimits on E-I-categories*, from: “Algebraic topology Poznań 1989”, Lecture Notes in Math. 1474, Springer, Berlin (1991) 273–294 [MR1133907](#)
- [28] **B Stenström**, *Rings of quotients*, Die Grundlehren der Mathematischen Wissenschaften 217, Springer, New York (1975) [MR0389953](#)
- [29] **R W Thomason**, *Uniqueness of delooping machines*, *Duke Math. J.* 46 (1979) 217–252 [MR534053](#)
- [30] **S T Vrećica, R T Živaljević**, *Cycle-free chessboard complexes and symmetric homology of algebras*, *European J. Combin.* 30 (2009) 542–554 [MR2489249](#)

Department of Mathematics, Fordham University
Bronx NY 10461, USA

ault@fordham.edu

<http://fordham.academia.edu/ShawnAult/>

Received: 6 January 2010