

Expanders and property A

ANA KHUKHRO
NICK J WRIGHT

We give a cohomological characterisation of expander graphs, and use it to give a direct proof that expander graphs do not have Yu’s property A.

20F65, 20F69, 51F99, 55N20; 05C80, 43A07, 55B20, 68R10

1 Introduction

Property A, first introduced by Yu [6], is a coarse geometric analogue of amenability.

Definition 1.1 A discrete bounded geometry metric space X has *property A* if for each $x \in X$ and each $n \in \mathbb{N}$, there is an element $f_n(x) \in \text{Prob}(X)$ with

- (1) a sequence S_n such that $\text{supp}(f_n(x)) \subseteq B_{S_n}(x)$, and
- (2) for any $R > 0$, $\|f_n(x_1) - f_n(x_0)\|_{\ell^1} \rightarrow 0$ as $n \rightarrow \infty$ uniformly on the set $\{(x_0, x_1) : d(x_0, x_1) \leq R\}$.

Yu [6] proved that if a metric space has property A then it is uniformly embeddable into Hilbert space. Indeed, this was the original motivation behind this definition, since a result of the same paper [6] states that the coarse Baum–Connes conjecture holds for discrete bounded geometry metric spaces which admit a uniform embedding into Hilbert space.

There are few known examples of metric spaces which do not have property A. One such family of examples is provided by *expander graphs* (cf Lubotzky [3] and Margulis [4]). Informally, an expander is a sequence of highly connected graphs which have bounded valency. Expander graphs are used in computer science due to their high connectivity. They are also of theoretical interest as they provide counterexamples to the coarse Baum–Connes conjecture; see Higson, Lafforgue and Skandalis [2].

It is well-known that expander graphs do not uniformly embed into Hilbert space (see for example Roe [5]). It follows that expanders cannot have property A. By using nonembeddability into Hilbert space, the existing proof of this fact obscures the

relationship between these two notions. In this paper we give a new, direct proof that expanders do not have property A, making the connection between the two properties explicit. Our proof is based on the observation that both the expander condition and property A can be expressed in terms of a coboundary operator which, roughly speaking, measures the size of the (co)boundary of a set of vertices. The cohomological description of property A was given in [1], while the cohomological description of the expander condition is introduced in this paper.

2 Expanders and cohomology

Let $\{\Gamma_i\}$ be a sequence of finite graphs with uniformly bounded valencies. Abusing notation, we will also denote the vertex set by Γ_i and the edges by E_i . We take the edges to be directed, with an edge connecting x to y if and only if there is an edge connecting y to x . The *Cheeger constant* of the graph Γ_i is defined by $h(\Gamma_i) = \frac{1}{2} \inf(|\partial F|/|F|)$, where F ranges over the nonempty subsets of Γ_i such that $|F| \leq \frac{1}{2}|\Gamma_i|$ and ∂F denotes the coboundary¹ of F , ie the set of edges of Γ_i with exactly one end point in F . The factor of $\frac{1}{2}$ compensates for the doubling arising from the use of directed edges.

Definition 2.1 A finite graph Γ is a (k, ε) -*expander* if each vertex of Γ has valency at most k , and $h(\Gamma) \geq \varepsilon$.

A sequence of finite graphs $\{\Gamma_i\}$ is called an *expander sequence* if $|\Gamma_i| \rightarrow \infty$ and there exists k, ε such that each Γ_i is a (k, ε) -expander.

It is not obvious that such sequences exist. Their existence was first proved by Pinsker, in a nonconstructive way. Margulis [4] was the first to give explicit examples of expanders, using discrete groups with property (T).

Let Γ be a finite graph and let E denote its set of directed edges. We view \mathbb{C} as the subspace of $\ell^1(\Gamma)$ consisting of constant functions, and write \bar{f} for the class in $\ell^1(\Gamma)/\mathbb{C}$ represented by $f \in \ell^1(\Gamma)$. The norm on $\ell^1(\Gamma)/\mathbb{C}$ is the quotient norm defined by $\|\bar{f}\|_{\ell^1/\mathbb{C}} = \inf_{c \in \mathbb{C}} \|f + c\|_{\ell^1}$. We will write $\ell_0^1(E)$ for the subspace of $\ell^1(E)$ consisting of functions whose sum is zero. The norm on $\ell_0^1(E)$ is the usual ℓ_1 norm. Define a coboundary map

$$d: \ell^1(\Gamma)/\mathbb{C} \longrightarrow \ell_0^1(E)$$

by $d\bar{f}(e) = f(e^+) - f(e^-)$ where e^- is the starting vertex and e^+ is the end vertex of the directed edge e .

¹This is usually referred to as the boundary of F , however as the map goes from vertices to edges, homologically it is a coboundary.

Lemma 2.2 *The Cheeger constant $h(\Gamma)$ is at least $\varepsilon/2$ if and only if $\|d\bar{f}\|_{\ell^1} \geq \varepsilon\|\bar{f}\|_{\ell^1/\mathbb{C}}$ for every $\bar{f} \in \ell^1(\Gamma)/\mathbb{C}$.*

Proof Suppose $\|d\bar{f}\|_{\ell^1} \geq \varepsilon\|\bar{f}\|_{\ell^1/\mathbb{C}}$ for every $\bar{f} \in \ell^1(\Gamma)/\mathbb{C}$. Then in particular, for any subset $F \subset \Gamma$ such that $|F| \leq \frac{1}{2}|\Gamma|$ we have $\|d\bar{\chi}_F\|_{\ell^1} \geq \varepsilon\|\bar{\chi}_F\|_{\ell^1/\mathbb{C}}$, where χ_F denotes the characteristic function of F . It is clear that $\|d\bar{\chi}_F\|_1$ is equal to $|\partial F|$, the coboundary of the set F (recall that we are taking our edges to be directed). Also, since $|F| \leq \frac{1}{2}|\Gamma|$, we have

$$\sum_{\gamma \in \Gamma} |\chi_F(\gamma) + c| = \sum_{\gamma \in F} |1 + c| + \sum_{\gamma \notin F} |c| \geq \sum_{\gamma \in F} 1 - \sum_{\gamma \in F} |c| + \sum_{\gamma \notin F} |c| \geq \sum_{\gamma \in F} 1.$$

From this, we can see that the infimum over $c \in \mathbb{C}$ of $\sum_{\gamma \in \Gamma} |\chi_F(\gamma) + c|$ is achieved when $c = 0$ and so we have $\|\bar{\chi}_F\|_{\ell^1/\mathbb{C}} = |F|$. Hence for every F with $|F| \leq \frac{1}{2}|\Gamma|$, we have $|\partial F| \geq \varepsilon|F|$ and so $h(\Gamma) \geq \varepsilon/2$.

Suppose now that $h(\Gamma)$ is at least $\varepsilon/2$. Given $\bar{f} \in \ell^1(\Gamma)/\mathbb{C}$, pick an $f' \in \ell^1(\Gamma)$ which takes positive values on each element of Γ and such that $\bar{f}' = \bar{f}$. We can write f' as $\sum a_j \chi_{F_j}$ for some nested collection of subsets $F_1 \subset F_2 \subset \dots \subset F_n$ of Γ and coefficients $a_j > 0$. Now $\|d\bar{f}\|_{\ell^1} = \|d\bar{f}'\|_{\ell^1}$ is equal to $\sum a_j \|d\bar{\chi}_{F_j}\|_{\ell^1}$ since the F_j are nested. Hence

$$\|d\bar{f}\|_{\ell^1} = \sum_j a_j \|d\bar{\chi}_{F_j}\|_{\ell^1} = \sum_j a_j |\partial F_j|.$$

Let F_j^c denote the complement of F_j in Γ . Since $h(\Gamma) \geq \varepsilon/2$, when $|F_j| \leq \frac{1}{2}|\Gamma|$ we have $|\partial F_j| \geq \varepsilon|F_j| = \|\bar{\chi}_{F_j}\|_{\ell^1/\mathbb{C}}$, while for $|F_j| > \frac{1}{2}|\Gamma|$ we have

$$|\partial F_j| = |\partial F_j^c| \geq \varepsilon|F_j^c| = \varepsilon\|\bar{\chi}_{F_j^c}\|_{\ell^1/\mathbb{C}} = \varepsilon\|\bar{1} - \bar{\chi}_{F_j}\|_{\ell^1/\mathbb{C}} = \varepsilon\|\bar{\chi}_{F_j}\|_{\ell^1/\mathbb{C}},$$

and so $\|d\bar{f}\|_{\ell^1} \geq \varepsilon \sum_j a_j \|\bar{\chi}_{F_j}\|_{\ell^1/\mathbb{C}} \geq \varepsilon \left\| \sum_j a_j \chi_{F_j} \right\|_{\ell^1/\mathbb{C}} = \varepsilon\|\bar{f}\|_{\ell^1/\mathbb{C}}$.

This completes the proof. □

The map $\ell^1(\Gamma) \rightarrow \ell^1_0(\Gamma)$ taking a function $f \in \ell^1(\Gamma)$ to $g = f - (1/|\Gamma|)\sum_{\beta \in \Gamma} f(\beta)$ has kernel \mathbb{C} , and hence induces an isomorphism from $\ell^1(\Gamma)/\mathbb{C}$ to $\ell^1_0(\Gamma)$. This map has norm at most 2 since

$$\|g\|_{\ell^1} = \sum_{\gamma \in \Gamma} |f(\gamma) - \frac{1}{|\Gamma|} \sum_{\beta \in \Gamma} f(\beta)| \leq \sum_{\gamma \in \Gamma} |f(\gamma)| + \sum_{\beta \in \Gamma} |f(\beta)| = 2\|f\|_{\ell^1},$$

while the inverse is given by the inclusion of $\ell^1_0(\Gamma)$ in $\ell^1(\Gamma)$ which has norm 1. Hence identifying $\ell^1(\Gamma)/\mathbb{C}$ with $\ell^1_0(\Gamma)$, the norms differ by a factor of at most 2.

We now move on to the definition of the cohomology which detects expander sequences. Let $\{\Gamma_i\}_{i \in \mathbb{N}}$ be a sequence of graphs. We denote by $\prod_{i \in \mathbb{N}}^\infty \ell^1(\Gamma_i)$ the space of bounded elements of the direct product. That is, $\prod_{i \in \mathbb{N}}^\infty \ell^1(\Gamma_i)$ is the space of functions from $\coprod_i \Gamma_i$ to \mathbb{C} , such that the sup- ℓ^1 -norm

$$\|f\| = \sup_{i \in \mathbb{N}} \|f|_{\Gamma_i}\|_{\ell^1}$$

is finite. We define a summation map $\sigma_0: \prod_{i \in \mathbb{N}}^\infty \ell^1(\Gamma_i) \rightarrow \ell^\infty(\mathbb{N})$ by $\sigma_0(f)(i) = \sum_{x \in \Gamma_i} f(x)$. Similarly $\prod_{i \in \mathbb{N}}^\infty \ell^1(E_i)$ is the space of functions on $\coprod_i E_i$ with finite sup- ℓ^1 -norm, and we define a map $\sigma_1: \prod_{i \in \mathbb{N}}^\infty \ell^1(E_i) \rightarrow \ell^\infty(\mathbb{N})$ by $\sigma_1(f)(i) = \sum_{x \in E_i} f(x)$.

We define $C^0(\{\Gamma_i\}) = \ker(\sigma_0)$, $C^1(\{\Gamma_i\}) = \ker(\sigma_1)$.

Note that $C^0(\{\Gamma_i\})$ consists of functions whose restriction to each Γ_i lies in $\ell^1_0(\Gamma_i)$, and $C^1(\{\Gamma_i\})$ consists of functions whose restriction to each E_i is in $\ell^1_0(E_i)$. Therefore combining the coboundary maps on each component yields a coboundary map $d: C^0(\{\Gamma_i\}) \rightarrow C^1(\{\Gamma_i\})$, and it is easy to see that this is bounded. In the spirit of [1], our cohomological description of the expander condition is given by completing this cochain complex.

Definition 2.3 [1, Definition 3.1] The *quotient completion* of a pre-Fréchet space V (a space equipped with a countable family of seminorms $\|\cdot\|_j$) is the space $V_Q = \ell^\infty(\mathbb{N}, V)/c_0(\mathbb{N}, V)$ of bounded sequences in V modulo sequences vanishing at infinity.

For simplicity we suppose that the seminorms are monotonic, that is $\|\cdot\|_i \leq \|\cdot\|_j$ for $i < j$. We note the following useful property of this completion.

Lemma 2.4 Let $T: V \rightarrow W$ be a bounded map from a normed spaced V to a pre-Fréchet space W . Then T is bounded below if and only if the induced map $T^Q: V_Q \rightarrow W_Q$ [1, Proposition 3.3] is injective.

Proof One direction is obvious: if T is bounded below then T^Q is also bounded below hence injective. For the converse suppose that T is not bounded below. This means that for each seminorm $\|\cdot\|_{j,W}$ for W and all $\varepsilon > 0$ there exists v in V with $\|Tv\|_{j,W} < \varepsilon\|v\|_V$. Hence we can find a sequence $v_n \in V$ with $\|v_n\|_V = 1$ and $\|Tv_n\|_{n,W} < \frac{1}{n}$. As the sequence v_n is bounded, it determines an element v of V_Q . Its image under T^Q is given by the sequence Tv_n , and since for $n \geq j$ we have $\|Tv_n\|_{j,W} \leq \|Tv_n\|_{n,W} < \frac{1}{n}$, we have $Tv_n \in c_0(\mathbb{N}, W)$. Hence $T^Qv = 0$, so T^Q is not injective. \square

We remark that the lemma is not true in general if V is a pre-Fréchet space. Whilst for T not bounded below there still exists a sequence v_n not tending to zero such that $Tv_n \rightarrow 0$, there may be no *bounded* sequence with these properties.

We now give our cohomological description of the expander condition. Let $C_Q^p(\{\Gamma_i\})$ denote the quotient completion of $C^p(\{\Gamma_i\})$ for $p = 0, 1$. The extension of the coboundary map d to the completion we again denote by d .

Definition 2.5 The Cheeger cohomology of a sequence of graphs $\{\Gamma_i\}$, denoted $H_h^*(\{\Gamma_i\})$ is the cohomology of the cochain complex $(C_Q^p(\{\Gamma_i\}), d)$.

We remark that $C_Q^p(\{\Gamma_i\})$ is the kernel of the induced map σ_p^Q , since the quotient completion preserves exactness (cf [1]).

Theorem 2.6 Let $\{\Gamma_i\}_{i \in \mathbb{N}}$ be a sequence of finite graphs with bounded valency. Then $\{\Gamma_i\}$ is an expander sequence if and only if $H_h^0(\{\Gamma_i\})$ vanishes.

Proof Using Lemma 2.2 and the identification of $\ell^1(\Gamma_i)/\mathbb{C}$ with $\ell_0^1(\Gamma_i)$, the graphs $\{\Gamma_i\}$ form an expander sequence if and only if there exists $\varepsilon > 0$ such that for each graph Γ_i the coboundary map $d: \ell_0^1(\Gamma_i) \rightarrow \ell_0^1(E_i)$ is ε -bounded below. The individual coboundary maps are bounded below by a common ε if and only if the map $d: C^0(\{\Gamma_i\}) \rightarrow C^1(\{\Gamma_i\})$ is bounded below. By Lemma 2.4 this is equivalent to injectivity of the coboundary map $d: C_Q^0(\{\Gamma_i\}) \rightarrow C_Q^1(\{\Gamma_i\})$ on the completed complex. Hence the graphs $\{\Gamma_i\}$ form an expander sequence if and only if $H_h^0(\{\Gamma_i\}) = 0$. \square

3 Symmetrisation of property A

In this section we recall one of the cohomological characterisations of property A from [1], and prove a symmetrisation result. Throughout this section, let X denote a metric space. At certain points we will require X to be a discrete, bounded geometry space, that is, for each $R > 0$ there exists N such that for all $x \in X$ the ball of radius R about x contains at most N points.

Definition 3.1 An X -module is a triple $\mathcal{V} = (V, \|\cdot\|, \text{supp})$, where V is a Banach space with norm $\|\cdot\|$ and supp is a function from V to the power set of X such that

- (1) $\text{supp}(v) = \emptyset$ if $v = 0$,
- (2) $\text{supp}(v + w) \subseteq \text{supp}(v) \cup \text{supp}(w)$ for every $v, w \in V$,
- (3) $\text{supp}(\lambda v) = \text{supp}(v)$ for every $v \in V$ and every $\lambda \neq 0$,
- (4) if v_n is a sequence converging to v then $\text{supp}(v) \subseteq \overline{\bigcup_n \text{supp}(v_n)}$.

Let $\mathcal{E}^p(X, \mathcal{V})$ denote the space of functions ϕ from X^{p+1} to V such that for all $R > 0$ the function ϕ is bounded on

$$\Delta_R^{p+1} = \{(x_0, \dots, x_p) \in X^{p+1} : d(x_i, x_j) \leq R \text{ for all } i, j\}$$

and there exists $S > 0$ such that if $\mathbf{x} = (x_0, \dots, x_p) \in \Delta_R^{p+1}$ then $\text{supp}(\phi(\mathbf{x})) \subseteq B_S(x_i)$ for all i .

The space $\mathcal{E}^p(X, V)$ is equipped with the family of seminorms

$$\|\phi\|_R = \sup\{\|\phi(x)\|_{\mathcal{V}} : x \in \Delta_R^{p+1}\}.$$

In [1] this is denoted by $\mathcal{E}^{p,-1}(X, \mathcal{V})$, being part of a bicomplex, however for simplicity we drop the -1 from our notation. We note that $\mathcal{E}^0(X, \mathcal{V})$ is a normed space, since in dimension zero the norms are independent of R .

Let $\mathcal{E}_Q^p(X, \mathcal{V})$ denote the quotient completion of $\mathcal{E}^p(X, \mathcal{V})$. The usual formula

$$D\phi(x_0, \dots, x_{p+1}) = \sum_{i=0}^{p+1} (-1)^i \phi(x_0, \dots, \hat{x}_i, \dots, x_{p+1})$$

yields a coboundary map from $\mathcal{E}^p(X, \mathcal{V})$ to $\mathcal{E}^{p+1}(X, \mathcal{V})$, and the extension of D to the completion we again denote by D .

The *controlled cohomology* $H_Q^*(X, \mathcal{V})$ is the cohomology of the completed complex $(\mathcal{E}_Q^p(X, \mathcal{V}), D)$.

Let $\mathbf{1}_Q$ denote the constant function 1 on X , viewed as a 0-cocycle in $\mathcal{E}_Q^0(X, \mathbb{C})$, and let $\pi_*: H_Q^0(X, \ell^1(X)) \rightarrow H_Q^0(X, \mathbb{C})$ be the map on cohomology induced by the summation map $\pi: \ell^1(X) \rightarrow \mathbb{C}$. By [1, Theorem 7.2] the space X has property A if and only if the class $[\mathbf{1}_Q] \in H_Q^0(X, \mathbb{C})$ is in the image of the map π_* . Here the module $\ell^1(X)$ is equipped with the usual support function, while all elements of \mathbb{C} are defined to have empty support.

We now compare ℓ^1 and ℓ^2 coefficients. We define maps $\alpha: \ell^1(X) \rightarrow \ell^2(X)$ and $\beta: \ell^2(X) \rightarrow \ell^1(X)$ by

$$\alpha(\eta)(x) = \sqrt{|\eta(x)|} \text{ for } \eta \in \ell^1(X), \quad \beta(\xi)(x) = |\xi(x)|^2 \text{ for } \xi \in \ell^2(X).$$

Note that $\|\alpha(\eta)\|_{\ell^2}^2 = \|\eta\|_{\ell^1}$ and $\|\beta(\xi)\|_{\ell^1} = \|\xi\|_{\ell^2}^2$.

Lemma 3.2 *Let α, β be defined as above. Then the compositions with α and β yield maps $\mathcal{E}^p(X, \ell^1(X)) \rightarrow \mathcal{E}^p(X, \ell^2(X))$ and $\mathcal{E}^p(X, \ell^2(X)) \rightarrow \mathcal{E}^p(X, \ell^1(X))$ which extend in the natural way to maps α_*, β_* on the quotient completions. Moreover these maps take 0-cocycles to 0-cocycles.*

Proof The identity $\|\alpha(\eta)\|_{\ell^2}^2 = \|\eta\|_{\ell^1}$ shows that for ϕ_n a bounded sequence in $\mathcal{E}^p(X, \ell^1(X))$, the sequence $\alpha \circ \phi_n \in \mathcal{E}^p(X, \ell^2(X))$ is also bounded. Hence, as composition with α preserves supports, $\alpha \circ \phi_n$ defines an element in the quotient completion. We note that the inequalities

$$|\sqrt{|\eta(z)|} - \sqrt{|\eta'(z)|}| \leq \sqrt{||\eta(z)| - |\eta'(z)||} \leq \sqrt{|\eta(z) - \eta'(z)|}$$

imply that $\|\alpha(\eta) - \alpha(\eta')\|_{\ell^2}^2 \leq \|\eta - \eta'\|_{\ell^1}$. It follows that if ϕ'_n is another bounded sequence in $\mathcal{E}^p(X, \ell^1(X))$ such that $\|\phi_n - \phi'_n\|_R \rightarrow 0$, then $\|\alpha \circ \phi_n - \alpha \circ \phi'_n\|_R \rightarrow 0$, and so the element of $\mathcal{E}^p_Q(X, \ell^2(X))$ obtained by composition with α is independent of the choice of representative of element of $\mathcal{E}^p_Q(X, \ell^1(X))$. Thus we have a well-defined map $\alpha_*: \mathcal{E}^p_Q(X, \ell^1(X)) \rightarrow \mathcal{E}^p_Q(X, \ell^2(X))$.

The estimate $\|\alpha(\eta) - \alpha(\eta')\|_{\ell^2}^2 \leq \|\eta - \eta'\|_{\ell^1}$ also yields

$$\begin{aligned} \|D\alpha(\phi_n)(x_0, x_1)\|_{\ell^2}^2 &= \|\alpha(\phi_n(x_1)) - \alpha(\phi_n(x_0))\|_{\ell^2}^2 \\ &\leq \|\phi_n(x_1) - \phi_n(x_0)\|_{\ell^1} \\ &= \|D\phi_n(x_0, x_1)\|_{\ell^1} \end{aligned}$$

for ϕ_n a bounded sequence in $\mathcal{E}^0_Q(X, \ell^1(X))$. Hence α_* takes 0-cocycles to 0-cocycles.

The argument for β_* is similar, using the identity $\|\beta(\xi)\|_{\ell^1} = \|\xi\|_{\ell^2}^2$ and the estimate $\|\beta(\xi) - \beta(\xi')\|_{\ell^1} \leq \|\xi - \xi'\|_{\ell^2} (\|\xi\|_{\ell^2} + \|\xi'\|_{\ell^2})$ which follows from

$$\begin{aligned} ||\xi(x)|^2 - |\xi'(x)|^2| &= ||\xi(x)| - |\xi'(x)|| (|\xi(x)| + |\xi'(x)|) \\ &\leq |\xi(x) - \xi'(x)| (|\xi(x)| + |\xi'(x)|) \end{aligned}$$

by the Cauchy-Schwartz inequality. □

We now prove a symmetrisation result. Note that we will omit norm subscripts where this does not cause confusion.

For an element ϕ of $\mathcal{E}^0_Q(X, \ell^1(X))$ or $\mathcal{E}^0_Q(X, \ell^2(X))$ we say ϕ is *symmetric* if it can be represented by a sequence ϕ_n such that $\phi(x)(z)$ is real and $\phi_n(x)(z) = \phi_n(z)(x)$ for all $x, z \in X$. We say that ϕ is *everywhere unital* if $\lim_{n \rightarrow \infty} \|\phi_n(x)\| = 1$ for all $x \in X$ (note that this limit is independent of the choice of representative sequence).

Theorem 3.3 *Let X be a bounded geometry metric space. The following are equivalent:*

- (1) X has property A.
- (2) There is a cocycle $\phi \in \mathcal{E}^0_Q(X, \ell^1(X))$ such that $\pi_*(\phi) = \mathbf{1}_Q$.

- (3) There is a symmetric cocycle $\phi \in \mathcal{E}_Q^0(X, \ell^1(X))$ such that $\pi_*(\phi) = \mathbf{1}_Q$.
- (4) There is a symmetric cocycle $\psi \in \mathcal{E}_Q^0(X, \ell^2(X))$ such that ψ is everywhere unital.

Proof The equivalence of (1) and (2) is [1, Theorem 7.2].

First we prove (2) \Rightarrow (4). Suppose there exists a cocycle $\phi \in \mathcal{E}_Q^0(X, \ell^1(X))$ such that $\pi_*(\phi) = \mathbf{1}_Q$. We consider $\alpha_*\phi$. Choosing a representative sequence ϕ_n for ϕ we note that $\|\alpha(\phi_n(x))\|^2 = \|\phi_n(x)\| \geq 1$ for all x since $\pi(\phi(x)) = 1$. Let

$$\theta_n(x) = \frac{1}{\|\alpha(\phi_n(x))\|} \alpha(\phi_n(x)).$$

We know that $\alpha_*\phi$ is a cocycle. The estimate

$$\begin{aligned} \left\| \frac{1}{\|\xi\|} \xi - \frac{1}{\|\xi'\|} \xi' \right\| &\leq \frac{\|\xi - \xi'\|}{\|\xi\|} + \left| \frac{1}{\|\xi\|} - \frac{1}{\|\xi'\|} \right| \|\xi'\| = \frac{\|\xi - \xi'\| + \left| \|\xi'\| - \|\xi\| \right|}{\|\xi\|} \\ &\leq 2 \|\xi - \xi'\| \end{aligned}$$

for $\xi \in \ell^2(X)$ with $\|\xi\| \geq 1$, shows that $D\theta_n \rightarrow 0$, ie θ again determines a cocycle.

Consider the operators $T_n: \ell^2(X) \rightarrow \ell^2(X)$ defined by

$$(T_n \xi)(y) = \sum_{x \in X} \theta_n(x)(y) \xi(x).$$

The support condition on θ_n provides an $S_n > 0$ such that $\theta_n(x)$ is supported in $B_{S_n}(x)$, and bounded geometry gives a bound N_n on the size of these balls, hence the operators T_n are bounded. The support condition also shows that these operators have finite propagation, and thus they are elements of the uniform Roe algebra of X (see [5, Section 4.4]). Consider $T'_n = (T_n^* T_n)^{1/2}$. This lies in the uniform Roe algebra since T_n does, and hence for each n we can find another self-adjoint operator T''_n with T''_n of finite propagation and $\|T''_n - T'_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Define $\psi_n(x) = T''_n(\delta_x)$. For $\xi \in \ell^2(X)$ we have

$$\langle T_n \xi, T_n \xi \rangle = \langle T_n^* T_n \xi, \xi \rangle = \langle (T'_n)^2 \xi, \xi \rangle = \langle T'_n \xi, T'_n \xi \rangle$$

so $\|T_n \xi\| = \|T'_n \xi\|$ for all ξ . We have $\|T'_n(\delta_x)\| = \|T_n(\delta_x)\| = \|\theta_n(x)\| = 1$. Hence $\|\psi_n(x)\| = \|T''_n(\delta_x)\| \rightarrow 1$ as $n \rightarrow \infty$. Finite propagation of T''_n provides the support condition for ψ_n and so ψ_n gives an everywhere unital element of $\mathcal{E}_Q^0(X, \ell^2(X))$. To see that ψ is a cocycle note that $\|D\theta_n(x_0, x_1)\| = \|T_n(\delta_{x_1} - \delta_{x_0})\| = \|T'_n(\delta_{x_1} - \delta_{x_0})\|$ and $\|D\psi_n(x_0, x_1)\| = \|T''_n(\delta_{x_1} - \delta_{x_0})\|$. As $T''_n - T'_n \rightarrow 0$, $D\theta_n \rightarrow 0$ implies $D\psi_n \rightarrow 0$.

As T_n'' is self-adjoint, we have $\psi_n(x)(z) = \langle T_n'' \delta_x, \delta_z \rangle = \langle \delta_x, T_n'' \delta_z \rangle = \overline{\psi_n(z)(x)}$. To make ψ_n symmetric it therefore suffices to ensure that $\psi_n(x)(z)$ is real. For an operator $T: \ell^2(X) \rightarrow \ell^2(X)$, let \bar{T} denote the operator defined by $\bar{T}\xi = T\bar{\xi}$ where $\bar{\xi}$ denotes the entry-wise complex conjugate of ξ . As θ_n is real, it follows that $\bar{T}_n = T_n$, and so $\bar{T}_n^* T_n = \bar{T}_n^* \bar{T}_n = T_n^* T_n$, hence as $T_n^* T_n = T_n'^2$ we have $\bar{T}_n'^2 = \bar{T}_n'^2 = T_n^* T_n$. Since the positive square-root T_n' of $T_n^* T_n$ is unique we have $\bar{T}_n' = T_n'$. Without loss of generality we may assume that $\bar{T}_n'' = T_n''$, since replacing T_n'' with its real part $\frac{1}{2}(T_n'' + \bar{T}_n'')$ reduces the distance from T_n' . Hence we have $\psi_n(x)(z) = \langle T_n'' \delta_x, \delta_z \rangle$ real, so we have proved (4).

(4) \Rightarrow (3) is immediate from Lemma 3.2: given ψ , we take $\phi = \beta_* \psi$. Symmetry is preserved and as ψ is everywhere unital, the same holds for ϕ . So, as ϕ is nonnegative, we have $\pi_* \phi = \mathbf{1}_Q$.

(3) \Rightarrow (2) is trivial. □

4 Expanders do not have property A

Let Γ be a disjoint union of graphs $\{\Gamma_i\}_{i \in \mathbb{N}}$ equipped with a proper metric such that the restriction to each component Γ_i is the graph metric on Γ_i , and such that the distance between Γ_i and its complement Γ_i^c tends to infinity as $i \rightarrow \infty$. If Γ has property A then there is a cocycle $\phi \in \mathcal{E}_Q^0(\Gamma, \ell^1(\Gamma))$ with $\pi_*(\phi) = \mathbf{1}_Q$, while if $\{\Gamma_i\}$ is an expander sequence then $H_h^0(\{\Gamma_i\})$ is zero. We will show that these two cohomological conditions are contradictory, that is, expanders cannot have property A.

Theorem 4.1 *Let Γ be a disjoint union of graphs Γ_i with bounded valency, such that $d(\Gamma_i, \Gamma_i^c) \rightarrow \infty$ and $|\Gamma_i| \rightarrow \infty$ as $i \rightarrow \infty$. If there exists a cocycle $\phi \in \mathcal{E}_Q^0(\Gamma, \ell^1(\Gamma))$ such that $\pi_*(\phi) = \mathbf{1}_Q$ then $H_h^0(\{\Gamma_i\})$ is nonzero.*

Proof Suppose there exists a cocycle $\phi \in \mathcal{E}_Q^0(\Gamma, \ell^1(\Gamma))$ such that $\pi_*(\phi) = \mathbf{1}_Q$. We will use this to construct a nonzero cocycle in $C_Q^0(\{\Gamma_i\})$ thus proving that $H_h^0(\{\Gamma_i\})$ is nonzero. By Theorem 3.3 we may assume that ϕ is a symmetric cocycle.

For each $n \in \mathbb{N}$ the controlled support condition provides an $S_n > 0$ such that for each $x \in \Gamma$, the support of $\phi_n(x)$ lies in $B_{S_n}(x)$. As the distance between components tends to ∞ , if i is sufficiently large then the distance between Γ_i and the other components of Γ exceeds S_n . Hence there exists j_n such that if $i \geq j_n$ then $\phi_n(x)$ is supported in Γ_i for all $x \in \Gamma_i$.

For each i, n , we choose a vertex $e_n^i \in \Gamma_i$ so that the infimum of

$$\sum_{(x_0, x_1) \in E_i} |D\phi_n(x_0, x_1)(z)|$$

over all $z \in \Gamma_i$ is realised at $z = e_n^i$, where E_i denotes the set of edges of Γ_i . Note that the infimum is actually a minimum, since each Γ_i is finite, and so such an e_n^i exists. For $i \geq j_n$ we define $f_n^i \in \ell^1 \Gamma_i$ by $f_n^i(x) = \phi_n(x)(e_n^i) - 1/|\Gamma_i|$, and for $i < j_n$ we define f_n^i to be 0. By symmetry of ϕ_n , when $i \geq j_n$ we have

$$\sum_{x \in \Gamma_i} |f_n^i(x)| = \sum_{x \in \Gamma_i} \left| \phi_n(e_n^i)(x) - \frac{1}{|\Gamma_i|} \right| \leq \|\phi_n(e_n^i)\|_{\ell^1} + 1.$$

This is bounded in i, n , hence $f_n = (f_n^1, f_n^2, \dots)$ defines an element f in the quotient completion of $\prod_{i \in \mathbb{N}} \ell^1(\Gamma_i)$. We will show that this is a nonzero cocycle in $C_Q^0(\{\Gamma_i\})$.

For $i < j_n$ we have $\sigma_0(f_n)(i) = \sum_{x \in \Gamma_i} f_n^i(x) = 0$, while for $i \geq j_n$ we have

$$\sum_{x \in \Gamma_i} f_n^i(x) = \sum_{x \in \Gamma_i} \left(\phi_n(x)(e_n^i) - \frac{1}{|\Gamma_i|} \right) = \sum_{x \in \Gamma_i} \left(\phi_n(e_n^i)(x) - \frac{1}{|\Gamma_i|} \right) = \pi_*(\phi_n)(e_n^i) - 1$$

by symmetry of ϕ_n . Since $\pi_*(\phi) = \mathbf{1}_Q$, the sequence $\pi_*(\phi_n)(e_n^i) - 1$ tends to zero (uniformly in i) as $n \rightarrow \infty$. Thus $\sigma_0^Q(f) = 0$, so f is an element of $C_Q^0(\{\Gamma_i\})$.

Recalling that the valencies of the Γ_i are uniformly bounded, we have a bound N_n on the cardinality of the balls $B_{S_n}(e_n^i)$. As $\phi_n(e_n^i)(x) = 0$ outside $B_{S_n}(e_n^i)$, when $i \geq j_n$ we have the following lower bound for the ℓ^1 -norm of f_n^i :

$$\|f_n^i\|_{\ell^1} \geq \sum_{x \in \Gamma_i \setminus B_{S_n}(e_n^i)} \frac{1}{|\Gamma_i|} \geq \frac{|\Gamma_i| - N_n}{|\Gamma_i|} = 1 - \frac{N_n}{|\Gamma_i|}.$$

Hence $\|f_n\| = \sup_{i \in \mathbb{N}} \|f_n^i\|_{\ell^1} \geq 1$ for all n . In particular $\|f_n\|$ does not tend to zero, so f is a *nonzero* element of $C_Q^0(\{\Gamma_i\})$.

It remains to verify that f is a cocycle. We apply the coboundary operator d to f_n^i . This clearly vanishes when $i < j_n$, while for $i \geq j_n$ we have

$$df_n^i(x_0, x_1) = f_n^i(x_1) - f_n^i(x_0) = D\phi(x_0, x_1)(e_n^i).$$

Our choice of e_n^i now comes into play. Let k be an upper bound on the valency of the graphs, so that $|E_i|/|\Gamma_i| \leq k$ for all i . Then we have

$$\begin{aligned} \|df_n^i\|_{\ell^1} &\leq \sum_{(x_0, x_1) \in E_i} |D\phi(x_0, x_1)(e_n^i)| = \frac{1}{|\Gamma_i|} \sum_{z \in \Gamma_i} \sum_{(x_0, x_1) \in E_i} |D\phi(x_0, x_1)(e_n^i)| \\ &\leq \frac{1}{|\Gamma_i|} \sum_{z \in \Gamma_i} \sum_{(x_0, x_1) \in E_i} |D\phi(x_0, x_1)(z)| \\ &\leq k \|D\phi_n\|_{R=1} \end{aligned}$$

as $\sum_{z \in \Gamma_i} |D\phi(x_0, x_1)(z)| \leq \|D\phi_n\|_{R=1}$. This tends to zero as $n \rightarrow \infty$ since ϕ is a cocycle. Hence $df = 0$, so f is a nonzero cocycle and $H_h^0(\{\Gamma_i\})$ is nonzero. \square

Since property A is equivalent to existence of a cocycle $\phi \in \mathcal{E}_Q^0(X, \ell^1(X))$ such that $\pi_*(\phi) = \mathbf{1}_Q$, and a sequence of graphs is an expander if and only if $H_h^0(\{\Gamma_i\})$ vanishes we obtain the following immediate corollary to Theorem 4.1.

Corollary 4.2 *Let Γ be the disjoint union of an expander sequence, with metric as above. Then Γ does not have property A.*

References

- [1] **J Brodzki, G A Niblo, N Wright**, *A cohomological characterisation of Yu's Property A for metric spaces*, *Geom. Topol.* 16 (2012) 391–432
- [2] **N Higson, V Lafforgue, G Skandalis**, *Counterexamples to the Baum–Connes conjecture*, *Geom. Funct. Anal.* 12 (2002) 330–354 MR1911663
- [3] **A Lubotzky**, *Discrete groups, expanding graphs and invariant measures*, *Progress in Math.* 125, Birkhäuser Verlag, Basel (1994) MR1308046 With an appendix by JD Rogawski
- [4] **G A Margulis**, *Explicit group-theoretic constructions of combinatorial schemes and their applications in the construction of expanders and concentrators*, *Problemy Peredachi Informatsii* 24 (1988) 51–60 MR939574 In Russian
- [5] **J Roe**, *Lectures on coarse geometry*, *Univ. Lecture Series* 31, Amer. Math. Soc. (2003) MR2007488
- [6] **G Yu**, *The coarse Baum–Connes conjecture for spaces which admit a uniform embedding into Hilbert space*, *Invent. Math.* 139 (2000) 201–240 MR1728880

*School of Mathematics, University of Southampton
University Road, Southampton, SO17 1BJ, UK*

ak11g08@soton.ac.uk, wright@soton.ac.uk

Received: 22 August 2011 Revised: 3 October 2011

