

# Proof of a stronger version of the AJ Conjecture for torus knots

ANH T TRAN

For a knot  $K$  in  $S^3$ , the  $\mathfrak{sl}_2$ -colored Jones function  $J_K(n)$  is a sequence of Laurent polynomials in the variable  $t$  that is known to satisfy non-trivial linear recurrence relations. The operator corresponding to the minimal linear recurrence relation is called the recurrence polynomial of  $K$ . The AJ Conjecture (see Garoufalidis [4]) states that when reducing  $t = -1$ , the recurrence polynomial is essentially equal to the  $A$ -polynomial of  $K$ . In this paper we consider a stronger version of the AJ Conjecture, proposed by Sikora [14], and confirm it for all torus knots.

57N10; 57M25

## 1 Introduction

### 1.1 The AJ Conjecture

For a knot  $K$  in  $S^3$ , let  $J_K(n) \in \mathbb{Z}[t^{\pm 1}]$  be the colored Jones polynomial of  $K$  colored by the  $n$ -dimensional simple  $\mathfrak{sl}_2$ -representation (Jones [8], and Reshetikhin and Turaev [13]), normalized so that for the unknot  $U$ ,

$$J_U(n) = [n] := \frac{t^{2n} - t^{-2n}}{t^2 - t^{-2}}.$$

The color  $n$  can be assumed to take negative integer values by setting  $J_K(-n) = -J_K(n)$ . In particular,  $J_K(0) = 0$ . It is known that  $J_K(1) = 1$ , and  $J_K(2)$  is the ordinary Jones polynomial.

Define two operators  $L, M$  acting on the set of discrete functions  $f: \mathbb{Z} \rightarrow \mathcal{R} := \mathbb{C}[t^{\pm 1}]$  by

$$(Lf)(n) := f(n+1), \quad (Mf)(n) := t^{2n} f(n).$$

It is easy to see that  $LM = t^2 ML$ . Besides, the inverse operators  $L^{-1}, M^{-1}$  are well-defined. One can consider  $L, M$  as elements of the quantum torus

$$\mathcal{T} := \mathcal{R}\langle L^{\pm 1}, M^{\pm 1} \rangle / (LM - t^2 ML),$$

which is not commutative, but almost commutative.

Let

$$\mathcal{A}_K = \{P \in \mathcal{T} \mid PJ_K = 0\},$$

which is a left-ideal of  $\mathcal{T}$ , called the *recurrence ideal* of  $K$ . It was proved by Garoufalidis and Lê in [5] that for every knot  $K$ , the recurrence ideal  $\mathcal{A}_K$  is non-zero. An element in  $\mathcal{A}_K$  is called a recurrence relation for the colored Jones polynomials of  $K$ .

The ring  $\mathcal{T}$  is not a principal left-ideal domain, ie, not every left-ideal of  $\mathcal{T}$  is generated by one element. By adding all inverses of polynomials in  $t, M$  to  $\mathcal{T}$ , one gets a principal left-ideal domain  $\tilde{\mathcal{T}}$ ; cf [4]. The ring  $\tilde{\mathcal{T}}$  can be formally defined as follows. Let  $\mathcal{R}(M)$  be the fractional field of the polynomial ring  $\mathcal{R}[M]$ . Let  $\tilde{\mathcal{T}}$  be the set of all Laurent polynomials in the variable  $L$  with coefficients in  $\mathcal{R}(M)$ ,

$$\tilde{\mathcal{T}} = \left\{ \sum_{j \in \mathbb{Z}} f_j(M)L^j \mid f_j(M) \in \mathcal{R}(M), f_j = 0 \text{ almost everywhere} \right\},$$

and define the product in  $\tilde{\mathcal{T}}$  by  $f(M)L^k \cdot g(M)L^l = f(M)g(t^{2k}M)L^{k+l}$ .

The left-ideal extension  $\tilde{\mathcal{A}}_K := \tilde{\mathcal{T}}\mathcal{A}_K$  of  $\mathcal{A}_K$  in  $\tilde{\mathcal{T}}$  is then generated by a polynomial

$$\alpha_K(t; M, L) = \sum_{j=0}^d \alpha_{K,j}(t, M)L^j,$$

where  $d$  is assumed to be minimal and all the coefficients  $\alpha_{K,j}(t, M) \in \mathbb{Z}[t^{\pm 1}, M]$  are assumed to be co-prime. That  $\alpha_K$  can be chosen to have integer coefficients follows from the fact that  $J_K(n) \in \mathbb{Z}[t^{\pm 1}]$ . The polynomial  $\alpha_K$  is defined up to a polynomial in  $\mathbb{Z}[t^{\pm 1}, M]$ . Moreover, one can choose  $\alpha_K \in \mathcal{A}_K$ , ie, it is a recurrence relation for the colored Jones polynomials. We will call  $\alpha_K$  the *recurrence polynomial* of  $K$ .

Let  $\varepsilon$  be the map reducing  $t = -1$ . Garoufalidis [4] formulated the following conjecture (see also Frohman, Gelca and Lofaro [3], and Gelca [6]).

**Conjecture 1 (AJ Conjecture)** *For every knot  $K$ ,  $\varepsilon(\alpha_K)$  is equal to the  $A$ -polynomial, up to a polynomial depending on  $M$  only.*

The  $A$ -polynomial of a knot was introduced by Cooper, Culler, Gillet, Long and Shalen [1]; it describes the  $SL_2(\mathbb{C})$ -character variety of the knot complement as viewed from the boundary torus. Here in the definition of the  $A$ -polynomial, we also allow the factor  $L - 1$  coming from the abelian component of the character variety of the knot group. Hence the  $A$ -polynomial in this paper is equal to  $L - 1$  times the  $A$ -polynomial defined in [1].

The AJ Conjecture was verified for the trefoil and figure 8 knots by Garoufalidis [4], and was partially checked for all torus knots by Hikami [7]. It was established for some classes of two-bridge knots and pretzel knots, including all twist knots and  $(-2, 3, 6n \pm 1)$ -pretzel knots, by Lê and the author [9; 10]. Here we provide a full proof of the AJ Conjecture for all torus knots. Moreover, we show that a stronger version of the conjecture, due to Sikora, holds true for all torus knots.

### 1.2 Main results

For a finitely generated group  $G$ , let  $\chi(G)$  denote the  $SL_2(\mathbb{C})$ -character variety of  $G$ ; see eg Culler and Shalen [2], and Lubotzky and Magid [11]. For a manifold  $Y$  we use  $\chi(Y)$  also to denote  $\chi(\pi_1(Y))$ . Suppose  $G = \mathbb{Z}^2$ , the free abelian group with 2 generators. Every pair of generators  $\mu, \lambda$  will define an isomorphism between  $\chi(G)$  and  $(\mathbb{C}^*)^2/\tau$ , where  $(\mathbb{C}^*)^2$  is the set of non-zero complex pairs  $(M, L)$  and  $\tau$  is the involution  $\tau(M, L) := (M^{-1}, L^{-1})$ , as follows: Every representation is conjugate to an upper diagonal one, with  $M$  and  $L$  being the upper left entries of  $\mu$  and  $\lambda$ , respectively. The isomorphism does not change if one replaces  $(\mu, \lambda)$  by  $(\mu^{-1}, \lambda^{-1})$ .

For an algebraic set  $V$  (over  $\mathbb{C}$ ), let  $\mathbb{C}[V]$  denote the ring of regular functions on  $V$ . For example,  $\mathbb{C}[(\mathbb{C}^*)^2/\tau] = \mathfrak{t}^\sigma$ , the  $\sigma$ -invariant subspace of  $\mathfrak{t} := \mathbb{C}[M^{\pm 1}, L^{\pm 1}]$ , where  $\sigma(M^k L^l) := M^{-k} L^{-l}$ .

Let  $K$  be a knot in  $S^3$  and  $X = S^3 \setminus K$  its complement. The boundary of  $X$  is a torus whose fundamental group is free abelian of rank two. An orientation of  $K$  will define a unique pair of an oriented meridian  $\mu$  and an oriented longitude  $\lambda$  such that the linking number between the longitude and the knot is zero. The pair provides an identification of  $\chi(\partial X)$  and  $(\mathbb{C}^*)^2/\tau$  that actually does not depend on the orientation of  $K$ .

The inclusion  $\partial X \hookrightarrow X$  induces an algebra homomorphism

$$\theta: \mathbb{C}[\chi(\partial X)] \cong \mathfrak{t}^\sigma \longrightarrow \mathbb{C}[\chi(X)].$$

We will call the kernel  $\mathfrak{p}$  of  $\theta$  the  $A$ -ideal of  $K$ ; it is an ideal of  $\mathfrak{t}^\sigma$ . The  $A$ -ideal was first introduced in [3]; it determines the  $A$ -polynomial of  $K$ . In fact  $\mathfrak{p} = (A_K \cdot \mathfrak{t})^\sigma$ , the  $\sigma$ -invariant part of the ideal  $A_K \cdot \mathfrak{t} \subset \mathfrak{t}$  generated by the  $A$ -polynomial  $A_K$ .

The involution  $\sigma$  acts on the quantum torus  $\mathcal{T}$  also by  $\sigma(M^k L^l) = M^{-k} L^{-l}$ . Let  $\mathcal{A}_K^\sigma$  be the  $\sigma$ -invariant part of the recurrence ideal  $\mathcal{A}_K$ ; it is an ideal of  $\mathcal{T}^\sigma$ . Sikora [14] proposed the following conjecture.

**Conjecture 2** *Suppose  $K$  is a knot. Then  $\sqrt{\varepsilon(\mathcal{A}_K^\sigma)} = \mathfrak{p}$ .*

Here  $\sqrt{\varepsilon(\mathcal{A}_K^\sigma)}$  denotes the radical of the ideal  $\varepsilon(\mathcal{A}_K^\sigma)$  in the ring  $t^\sigma = \varepsilon(\mathcal{T}^\sigma)$ .

It is easy to see that [Conjecture 2](#) implies the AJ Conjecture. [Conjecture 2](#) was verified for the unknot and the trefoil knot by Sikora [14]. In the present paper we confirm it for all torus knots.

**Theorem 1** [Conjecture 2](#) holds true for all torus knots.

### 1.3 Plan of the paper

We provide a full proof of the AJ Conjecture for all torus knots in [Section 2](#) and prove [Theorem 1](#) in [Section 2](#).

### 1.4 Acknowledgements

This paper was done when the author was a graduate student in the School of Mathematics, Georgia Institute of Technology. The author would like to thank T T Q Lê for his guidance, S Garoufalidis for helpful discussions, and the referee for suggestions.

## 2 Proof of the AJ Conjecture for torus knots

We will always assume that knots have framings 0.

Let  $T(a, b)$  denote the  $(a, b)$ -torus knot. We consider the two cases,  $a, b > 2$  and  $a = 2$ , separately. Lemmas [2.1](#) and [2.5](#) below were first proved in [7] using formulas for the colored Jones polynomials and the Alexander polynomial of torus knots given in Morton [12]. We present direct proofs here.

### 2.1 The case $a, b > 2$

**Lemma 2.1** One has

$$J_{T(a,b)}(n+2) = t^{-4ab(n+1)} J_{T(a,b)}(n) + t^{-2ab(n+1)} \frac{t^{2\lambda_{(a+b)(n+1)}} - t^{-2\lambda_{(a-b)(n+1)}}}{t^2 - t^{-2}},$$

where  $\lambda_k := t^{2k} + t^{-2k}$ .

**Proof** By [12], we have

$$(1) \quad J_{T(a,b)}(n) = t^{-ab(n^2-1)} \sum_{j=-(n-1)/2}^{(n-1)/2} t^{4bj(aj+1)} [2aj + 1],$$

where  $[k] := (t^{2k} - t^{-2k}) / (t^2 - t^{-2})$ . Hence:

$$\begin{aligned}
 & J_{T(a,b)}(n+2) \\
 &= t^{-ab((n+2)^2-1)} \sum_{j=-(n+1)/2}^{(n+1)/2} t^{4bj(aj+1)} [2aj+1] \\
 &= t^{-ab((n+2)^2-1)} \sum_{j=-(n-1)/2}^{(n-1)/2} t^{4bj(aj+1)} [2aj+1] + t^{-ab((n+2)^2-1)} \\
 &\quad \times (t^{b(n+1)(a(n+1)+2)} [a(n+1)+1] - t^{b(n+1)(a(n+1)-2)} [a(n+1)-1]) \\
 &= t^{-4ab(n+1)} J_{T(a,b)}(n) + t^{-2ab(n+1)} \frac{t^{2\lambda_{(a+b)(n+1)}} - t^{-2\lambda_{(a-b)(n+1)}}}{t^2 - t^{-2}} \quad \square
 \end{aligned}$$

**Lemma 2.2** *The colored Jones function of  $T(a, b)$  is annihilated by the operator  $F_{a,b} = c_3 L^3 + c_2 L^2 + c_1 L + c_0$  where:*

$$\begin{aligned}
 c_3 &:= t^2(t^{2(a+b)} M^{a+b} + t^{-2(a+b)} M^{-(a+b)}) \\
 &\quad - t^{-2}(t^{2(a-b)} M^{a-b} + t^{-2(a-b)} M^{-(a-b)}) \\
 c_2 &:= -t^{-2ab}(t^2(t^{4(a+b)} M^{a+b} + t^{-4(a+b)} M^{-(a+b)}) \\
 &\quad + t^{-2}(t^{4(a-b)} M^{a-b} + t^{-4(a-b)} M^{-(a-b)})) \\
 c_1 &:= -t^{-8ab} M^{-2ab} c_3 \\
 c_0 &:= -t^{-4ab} M^{-2ab} c_2
 \end{aligned}$$

**Proof** It is easy to check that  $c_3 t^{-4ab(n+2)} + c_1 = c_2 t^{-4ab(n+1)} + c_0 = 0$  and

$$c_3 (t^{2\lambda_{(a+b)(n+2)}} - t^{-2\lambda_{(a-b)(n+2)}}) + c_2 t^{2ab} (t^{2\lambda_{(a+b)(n+1)}} - t^{-2\lambda_{(a-b)(n+1)}}) = 0.$$

Hence, by Lemma 2.1,  $F_{a,b}J_{T(a,b)}(n)$  is equal to:

$$\begin{aligned}
 & c_3 J_{T(a,b)}(n+3) + c_2 J_{T(a,b)}(n+2) + c_1 J_{T(a,b)}(n+1) + c_0 J_{T(a,b)}(n) \\
 &= c_3 \left( t^{-4ab(n+2)} J_{T(a,b)}(n+1) + t^{-2ab(n+2)} \frac{t^{2\lambda(a+b)(n+2)} - t^{-2\lambda(a-b)(n+2)}}{t^2 - t^{-2}} \right) \\
 &\quad + c_2 \left( t^{-4ab(n+1)} J_{T(a,b)}(n) + t^{-2ab(n+1)} \frac{t^{2\lambda(a+b)(n+1)} - t^{-2\lambda(a-b)(n+1)}}{t^2 - t^{-2}} \right) \\
 &\quad\quad\quad + c_1 J_{T(a,b)}(n+1) + c_0 J_{T(a,b)}(n) \\
 &= (c_3 t^{-4ab(n+2)} + c_1) J_{T(a,b)}(n+1) + (c_2 t^{-4ab(n+1)} + c_0) J_{T(a,b)}(n) \\
 &\quad\quad\quad + t^{-2ab(n+1)} \left( c_3 \frac{t^{2\lambda(a+b)(n+2)} - t^{-2\lambda(a-b)(n+2)}}{t^2 - t^{-2}} \right. \\
 &\quad\quad\quad \left. + c_2 t^{2ab} \frac{t^{2\lambda(a+b)(n+1)} - t^{-2\lambda(a-b)(n+1)}}{t^2 - t^{-2}} \right) \\
 &= 0
 \end{aligned}$$

This proves Lemma 2.2.  $\square$

Recall that  $\alpha_{T(a,b)}$  is the recurrence polynomial of  $T(a, b)$ .

**Proposition 2.3** For  $a, b > 2$ , one has  $\alpha_{T(a,b)} = F_{a,b}$ .

**Proof** By Lemma 2.2 it suffices to show that if an operator  $P = P_2 L^2 + P_1 L + P_0$ , where the  $P_j$  are polynomials in  $\mathbb{C}[t^{\pm 1}, M]$ , annihilates the colored Jones polynomials of  $T(a, b)$  then  $P = 0$ .

Indeed, suppose  $P J_{T(a,b)}(n) = 0$ . Then, by Lemma 2.1:

$$\begin{aligned}
 0 &= P_2 J_{T(a,b)}(n+2) + P_1 J_{T(a,b)}(n+1) + P_0 J_{T(a,b)}(n) \\
 &= P_2 \left( t^{-4ab(n+1)} J_{T(a,b)}(n) + t^{-2ab(n+1)} \frac{t^{2\lambda(a+b)(n+1)} - t^{-2\lambda(a-b)(n+1)}}{t^2 - t^{-2}} \right) \\
 &\quad\quad\quad + P_1 J_{T(a,b)}(n+1) + P_0 J_{T(a,b)}(n) \\
 &= (t^{-4ab(n+1)} P_2 + P_0) J_{T(a,b)}(n) + P_1 J_{T(a,b)}(n+1) \\
 &\quad\quad\quad + P_2 t^{-2ab(n+1)} \frac{t^{2\lambda(a+b)(n+1)} - t^{-2\lambda(a-b)(n+1)}}{t^2 - t^{-2}}
 \end{aligned}$$

Let  $P'_2 = t^{-4ab(n+1)} P_2 + P_0$  and

$$P'_0 = P_2 t^{-2ab(n+1)} \frac{t^{2\lambda(a+b)(n+1)} - t^{-2\lambda(a-b)(n+1)}}{t^2 - t^{-2}}.$$

Then,

$$(2) \quad P'_2 J_{T(a,b)}(n) + P_1 J_{T(a,b)}(n+1) + P'_0 = 0.$$

Note that  $P'_2$  and  $P'_0$  are polynomials in  $\mathbb{C}[t^{\pm 1}, M]$ . We need the following lemma.

**Lemma 2.4** *The lowest degree in  $t$  of  $J_{T(a,b)}(n)$  is*

$$l_n = -abn^2 + ab + \frac{1}{2}(1 - (-1)^{n-1})(a-2)(b-2).$$

**Proof** From (1), it follows easily that  $l_n = -abn^2 + ab$  if  $n$  is odd, and  $l_n = (-abn^2 + ab) + (ab - 2b - 2a + 4)$  if  $n$  is even.  $\square$

Let us complete the proof of Proposition 2.3. Suppose  $P'_2, P_1 \neq 0$ . Let  $r_n$  and  $s_n$  be the lowest degrees (in  $t$ ) of  $P'_2$  and  $P_1$  respectively. Note that, when  $n$  is large enough,  $r_n$  and  $s_n$  are polynomials in  $n$  of degrees at most 1. Equation (2) then implies that  $r_n + l_n = s_n + l_{n+1}$ , ie:

$$r_n - s_n = l_{n+1} - l_n = -ab(2n+1) - (-1)^n(a-2)(b-2)$$

This cannot happen since the LHS is a polynomial in  $n$ , when  $n$  is large enough, while the RHS is not (since  $(a-2)(b-2) > 0$ ). Hence  $P'_2 = P_1 = P'_0 = 0$ , which means  $P = 0$ .  $\square$

It is easy to see that  $\varepsilon(\alpha_{T(a,b)}) = M^{-2ab}(M^a - M^{-a})(M^b - M^{-b})A_{T(a,b)}$  where  $A_{T(a,b)} = (L-1)(L^2M^{2ab} - 1)$  is the  $A$ -polynomial of  $T(a,b)$  when  $a, b > 2$ . This means the AJ Conjecture holds true for  $T(a,b)$  when  $a, b > 2$ .

## 2.2 The case $a = 2$

**Lemma 2.5** *One has*

$$J_{T(2,b)}(n+1) = -t^{-(4n+2)b} J_{T(2,b)}(n) + t^{-2nb}[2n+1].$$

**Proof** By (1), we have

$$J_{T(2,b)}(n) = t^{-2b(n^2-1)} \sum_{j=-(n-1)/2}^{(n-1)/2} t^{4bj(2j+1)}[4j+1].$$

Hence

$$J_{T(2,b)}(n+1) = t^{-2b((n+1)^2-1)} \sum_{k=-n/2}^{n/2} t^{4bk(2k+1)}[4k+1].$$

Set  $k = -(j + \frac{1}{2})$ . Then:

$$\begin{aligned}
 J_{T(2,b)}(n+1) &= t^{-2b((n+1)^2-1)} \sum_{j=(n-1)/2}^{-(n+1)/2} t^{4bj(2j+1)}[-(4j+1)] \\
 &= t^{-2b((n+1)^2-1)} \left( - \sum_{j=-(n-1)/2}^{(n-1)/2} t^{4bj(2j+1)}[4j+1] + t^{2bn(n+1)}[2n+1] \right) \\
 &= -t^{-(4n+2)b} J_{T(2,b)}(n) + t^{-2nb}[2n+1]
 \end{aligned}$$

This proves [Lemma 2.5](#). □

**Lemma 2.6** *The colored Jones function of  $T(2, b)$  is annihilated by the operator  $G_{2,b} = d_2 L^2 + d_1 L + d_0$  where*

$$\begin{aligned}
 d_2 &:= t^2 M^2 - t^{-2} M^{-2}, \\
 d_1 &:= t^{-2b} (t^{-4b} M^{-2b} (t^2 M^2 - t^{-2} M^{-2}) - (t^6 M^2 - t^{-6} M^{-2})), \\
 d_0 &:= -t^{-4b} M^{-2b} (t^6 M^2 - t^{-6} M^{-2}).
 \end{aligned}$$

**Proof** From [Lemma 2.5](#) we have

$$\begin{aligned}
 J_{T(2,b)}(n+1) &= -t^{-(4n+2)b} J_{T(2,b)}(n) + t^{-2nb}[2n+1], \\
 J_{T(2,b)}(n+2) &= t^{-8(n+1)b} J_{T(2,b)}(n) - t^{-6(n+1)b}[2n+1] + t^{-2(n+1)b}[2n+3].
 \end{aligned}$$

It is easy to check that

$$\begin{aligned}
 t^{-8(n+1)b} d_2 - t^{-(4n+2)b} d_1 + d_0 &= 0, \\
 d_2 (-t^{-6(n+1)b}[2n+1] + t^{-2(n+1)b}[2n+3]) + d_1 t^{-2nb}[2n+1] &= 0.
 \end{aligned}$$

Hence:

$$\begin{aligned}
 G_{2,b} J_{T(2,b)}(n) &= d_2 J_{T(2,b)}(n+2) + d_1 J_{T(2,b)}(n+1) + d_0 J_{T(2,b)}(n) \\
 &= d_2 (t^{-8(n+1)b} J_{T(2,b)}(n) - t^{-6(n+1)b}[2n+1] + t^{-2(n+1)b}[2n+3]) \\
 &\quad + d_1 (-t^{-(4n+2)b} J_{T(2,b)}(n) + t^{-2nb}[2n+1]) + d_0 J_{T(2,b)}(n) \\
 &= (t^{-8(n+1)b} d_2 - t^{-(4n+2)b} d_1 + d_0) J_{T(2,b)}(n) \\
 &\quad + d_2 (-t^{-6(n+1)b}[2n+1] + t^{-2(n+1)b}[2n+3]) + d_1 t^{-2nb}[2n+1] \\
 &= 0
 \end{aligned}$$



This proves [Lemma 2.6](#). □

**Proposition 2.7** *One has  $\alpha_{T(2,b)} = G_{2,b}$ .*

**Proof** By [Lemma 2.6](#), it suffices to show that if an operator  $P = P_1L + P_0$ , where the  $P_j$  are polynomials in  $\mathbb{C}[t^{\pm 1}, M]$ , annihilates the colored Jones polynomials of  $T(2, b)$  then  $P = 0$ .

Indeed, suppose  $PJ_{T(2,b)}(n) = 0$ . Then:

$$\begin{aligned} 0 &= P_1 J_{T(2,b)}(n+1) + P_0 J_{T(2,b)}(n) \\ &= P_1 (-t^{-(4n+2)b} J_{T(2,b)}(n) + t^{-2nb} [2n+1]) + P_0 J_{T(2,b)}(n) \\ &= (-t^{-(4n+2)b} P_1 + P_0) J_{T(2,b)}(n) + t^{-2nb} [2n+1] P_1 \end{aligned}$$

Let  $P'_1 = -t^{-(4n+2)b} P_1 + P_0$  and  $P'_0 = t^{-2nb} [2n+1] P_1$ . Then  $P'_1, P'_0$  are polynomials in  $\mathbb{C}[t^{\pm 1}, M]$  and  $P'_1 J(n) + P'_0 = 0$ . This implies that  $P'_1 = P'_0 = 0$  since the lowest degree in  $t$  of  $J_{T(2,b)}(n)$  is  $-2bn^2 + 2b$ , which is quadratic in  $n$ , by [Lemma 2.4](#). Hence  $P = 0$ . □

It is easy to see that  $\varepsilon(\alpha_{T(2,b)}) = M^{-2b}(M^2 - M^{-2})A_{T(2,b)}$  where  $A_{T(2,b)} = (L-1)(LM^{2b} + 1)$  is the  $A$ -polynomial of  $T(2, b)$ . This means the AJ Conjecture holds true for  $T(2, b)$ .

### 3 Proof of Theorem 1

As in the previous section, we consider the two cases,  $a, b > 2$  and  $a = 2$ , separately.

#### 3.1 The case $a, b > 2$

We claim that:

**Proposition 3.1** *The colored Jones function of  $T(a, b)$  is annihilated by the operator  $PQ$  where:*

$$\begin{aligned} P &:= t^{-10ab}(L^3 M^{2ab} + L^{-3} M^{-2ab}) \\ &\quad - (t^{2(a-b)} + t^{2(b-a)})t^{-4ab}(L^2 M^{2ab} + L^{-2} M^{-2ab}) + t^{2ab}(LM^{2ab} + L^{-1} M^{-2ab}) \\ &\quad \quad \quad - (t^{2ab} + t^{-2ab})(L + L^{-1}) + (t^{2(a-b)} + t^{2(b-a)})(t^{4ab} + t^{-4ab}) \\ Q &:= t^{-6ab}(L^3 M^{2ab} + L^{-3} M^{-2ab}) \\ &\quad - (t^{2(a+b)} + t^{-2(a+b)})q^{-ab}(L^2 M^{2ab} + L^{-2} M^{-2ab}) + t^{-2ab}(LM^{2ab} + L^{-1} M^{-2ab}) \\ &\quad \quad \quad - (t^{2ab} + t^{-2ab})(L + L^{-1}) + 2(t^{2(a+b)} + t^{-2(a+b)}) \end{aligned}$$

**Proof** We first prove the following two lemmas.

**Lemma 3.2** *One has*

$$QJ_{T(a,b)}(n) = t^{4ab-2}(\lambda_{a+b} - \lambda_{a-b}) \frac{t^{2abn}\lambda_{(a-b)(n+1)} - t^{-2abn}\lambda_{(a-b)(n-1)}}{t^2 - t^{-2}}.$$

**Proof** Let

$$g(n) := t^{-2abn} \frac{t^2\lambda_{(a+b)n} - t^{-2}\lambda_{(a-b)n}}{t^2 - t^{-2}}.$$

Then, by Lemma 2.1,  $J_{T(a,b)}(n+2) = t^{-4ab(n+1)}J_{T(a,b)}(n) + g(n+1)$ . Hence:

$$\begin{aligned} & QJ_{T(a,b)}(n) \\ &= t^{-6ab}(t^{4ab(n+3)}J_{T(a,b)}(n+3) + t^{-4ab(n-3)}J_{T(a,b)}(n-3)) \\ &\quad - (t^{2(a+b)} + t^{-2(a+b)})t^{-4ab}(t^{4ab(n+2)}J_{T(a,b)}(n+2) + t^{-4ab(n-2)}J_{T(a,b)}(n-2)) \\ &\quad\quad + t^{-2ab}(t^{4ab(n+1)}J_{T(a,b)}(n+1) + t^{-4ab(n-1)}J_{T(a,b)}(n-1)) \\ &\quad\quad\quad - (t^{2ab} + t^{-2ab})(J_{T(a,b)}(n+1) + J_{T(a,b)}(n-1)) \\ &\quad\quad\quad\quad + 2(t^{2(a+b)} + t^{-2(a+b)})J_{T(a,b)}(n) \\ &= t^{-6ab}(t^{4ab}(J_{T(a,b)}(n+1) + J_{T(a,b)}(n-1)) \\ &\quad\quad\quad + t^{2ab(n+5)}g(n+2) - t^{-2ab(n-5)}g(n-2)) \\ &\quad - (t^{2(a+b)} + t^{-2(a+b)})t^{-4ab}(2t^{4ab}J_{T(a,b)}(n) \\ &\quad\quad\quad + t^{2ab(n+4)}g(n+1) - t^{-2ab(n-4)}g(n-1)) \\ &\quad + t^{-2ab}(t^{4ab}(J_{T(a,b)}(n-1) + J_{T(a,b)}(n+1)) + (t^{2ab(n+3)} - t^{-2ab(n-3)})g(n) \\ &\quad\quad\quad - (t^{2ab} + t^{-2ab})(J_{T(a,b)}(n+1) + J_{T(a,b)}(n-1)) \\ &\quad\quad\quad\quad + 2(t^{2(a+b)} + t^{-2(a+b)})J_{T(a,b)}(n) \\ &= t^{-6ab}(t^{2ab(n+5)}g(n+2) - t^{-2ab(n-5)}g(n-2)) \\ &\quad - (t^{2(a+b)} + t^{-2(a+b)})t^{-4ab}(t^{2ab(n+4)}g(n+1) - t^{-2ab(n-4)}g(n-1)) \\ &\quad\quad\quad + t^{-2ab}(t^{2ab(n+3)} - t^{-2ab(n-3)})g(n) \end{aligned}$$

Using the definition of  $g(n)$ , we get:

$$\begin{aligned} QJ_{T(a,b)}(n) = t^{4ab} & \left( t^{2abn} \frac{t^2\lambda_{(a+b)(n+2)} - t^{-2}\lambda_{(a-b)(n+2)}}{t^2 - t^{-2}} \right. \\ & \left. - t^{-2abn} \frac{t^2\lambda_{(a+b)(n-2)} - t^{-2}\lambda_{(a-b)(n-2)}}{t^2 - t^{-2}} \right) \end{aligned}$$

$$\begin{aligned}
 & -(t^{2(a+b)} + t^{-2(a+b)})t^{4ab} \times \\
 & \left( t^{2abn} \frac{t^2 \lambda_{(a+b)(n+1)} - t^{-2} \lambda_{(a-b)(n+1)}}{t^2 - t^{-2}} \right. \\
 & \qquad \qquad \qquad \left. - t^{-2abn} \frac{t^2 \lambda_{(a+b)(n-1)} - t^{-2} \lambda_{(a-b)(n-1)}}{t^2 - t^{-2}} \right) \\
 & + t^{4ab} (t^{2abn} - t^{-2abn}) \frac{t^2 \lambda_{(a+b)n} - t^{-2} \lambda_{(a-b)n}}{t^2 - t^{-2}}
 \end{aligned}$$

Now applying the equality  $\lambda_{k+l} + \lambda_{k-l} = \lambda_k \lambda_l$ , we then obtain

$$Q J_{T(a,b)}(n) = t^{4ab-2} (\lambda_{a+b} - \lambda_{a-b}) \frac{t^{2abn} \lambda_{(a-b)(n+1)} - t^{-2abn} \lambda_{(a-b)(n-1)}}{t^2 - t^{-2}}.$$

This proves [Lemma 3.2](#). □

Let  $h(n) := t^{2abn} \lambda_{(a-b)(n+1)} - t^{-2abn} \lambda_{(a-b)(n-1)}$ .

**Lemma 3.3** *The function  $h(n)$  is annihilated by the operator  $P$ , ie,  $Ph(n) = 0$ .*

**Proof** Let  $c = a - b$ . Then:

$$\begin{aligned}
 Ph(n) &= t^{-10ab} (t^{4ab(n+3)} h(n+3) + t^{-4ab(n-3)} h(n-3)) \\
 & - (t^{2(a-b)} + t^{2(b-a)}) t^{-4ab} (t^{4ab(n+2)} h(n+2) + t^{-4ab(n-2)} h(n-2)) \\
 & + t^{2ab} (t^{4ab(n+1)} h(n+1) + t^{-4ab(n-1)} h(n-1)) \\
 & - (t^{2ab} + t^{-2ab}) (h(n+1) + h(n-1)) + (t^{2(a-b)} + t^{2(b-a)}) (t^{4ab} + t^{-4ab}) h(n) \\
 & = (t^{2ab(3n+4)} \lambda_{c(n+4)} - t^{2ab(n-2)} \lambda_{c(n+2)} \\
 & \qquad \qquad \qquad + t^{-2ab(n+2)} \lambda_{c(n-2)} - t^{-2ab(3n-4)} \lambda_{c(n-4)}) \\
 & - \lambda_c (t^{2ab(3n+4)} \lambda_{c(n+3)} - t^{2abn} \lambda_{c(n+1)} + t^{-2abn} \lambda_{c(n-1)} - t^{-2ab(3n-4)} \lambda_{c(n-3)}) \\
 & + (t^{2ab(3n+4)} \lambda_{c(n+2)} - t^{2ab(n+2)} \lambda_{cn} + t^{-2ab(n-2)} \lambda_{cn} - t^{-2ab(3n-4)} \lambda_{c(n-2)}) \\
 & - (t^{2ab} + t^{-2ab}) (t^{2ab(n+1)} \lambda_{c(n+2)} - t^{-2ab(n+1)} \lambda_{cn} \\
 & + t^{2ab(n-1)} \lambda_{cn} - t^{-2ab(n-1)} \lambda_{c(n-2)}) \\
 & + \lambda_c (t^{4ab} + t^{-4ab}) (t^{2abn} \lambda_{c(n+1)} - t^{-2abn} \lambda_{c(n-1)})
 \end{aligned}$$

Note that  $\lambda_{k+l} + \lambda_{k-l} = \lambda_k \lambda_l$ . Hence:

$$\begin{aligned}
 Ph(n) &= (-t^{2ab(n-2)}\lambda_{c(n+2)} + t^{-2ab(n+2)}\lambda_{c(n-2)}) \\
 &\quad - \lambda_c(-t^{2abn}\lambda_{c(n+1)} + t^{-2abn}\lambda_{c(n-1)}) \\
 &\quad + (-t^{2ab(n+2)}\lambda_{cn} + t^{-2ab(n-2)}\lambda_{cn}) \\
 &\quad - (t^{2ab} + t^{-2ab})(t^{2ab(n+1)}\lambda_{c(n+2)} - t^{-2ab(n+1)}\lambda_{cn} \\
 &\quad \quad + t^{2ab(n-1)}\lambda_{cn} - t^{-2ab(n-1)}\lambda_{c(n-2)}) \\
 &\quad + \lambda_c(t^{4ab} + t^{-4ab})(t^{2abn}\lambda_{c(n+1)} - t^{-2abn}\lambda_{c(n-1)}) \\
 &= -(t^{4ab} + t^{-4ab} + 1)t^{2abn}\lambda_{c(n+2)} + (t^{4ab} + t^{-4ab} + 1)t^{-2abn}\lambda_{c(n-2)} \\
 &\quad - (t^{4ab} + t^{-4ab} + 1)(t^{2abn} - t^{-2abn})\lambda_{cn} \\
 &\quad \quad + \lambda_c(t^{4ab} + t^{-4ab} + 1)(t^{2abn}\lambda_{c(n+1)} - t^{-2abn}\lambda_{c(n-1)}) \\
 &= -(t^{4ab} + t^{-4ab} + 1)t^{2abn}(\lambda_{c(n+2)} + \lambda_{cn} - \lambda_c\lambda_{c(n+1)}) \\
 &\quad + (t^{4ab} + t^{-4ab} + 1)t^{-2abn}(\lambda_{c(n-2)} + \lambda_{cn} - \lambda_c\lambda_{c(n-1)}) \\
 &= 0
 \end{aligned}$$

This proves [Lemma 3.3](#). □

[Proposition 3.1](#) follows directly from [Lemmas 3.2](#) and [3.3](#). □

### 3.2 The case $a = 2$

We claim that:

**Proposition 3.4** *The colored Jones function of  $T(2, b)$  is annihilated by the operator*

$$\begin{aligned}
 R &= t^{-4b}(L^2M^{2b} + L^{-2}M^{-2b}) + (t^{2b} + t^{-2b})(L + L^{-1}) \\
 &\quad - (t^4 + t^{-4})t^{-2b}(LM^{2b} + L^{-1}M^{-2b}) + (M^{2b} + M^{-2b}) - 2(t^4 + t^{-4}).
 \end{aligned}$$

**Proof** From [Lemma 2.5](#) we have

$$\begin{aligned}
 J_{T(2,b)}(n+1) &= -t^{-(4n+2)b}J_{T(2,b)}(n) + t^{-2nb}[2n+1], \\
 J_{T(2,b)}(n+2) &= t^{-8(n+1)b}J_{T(2,b)}(n) - t^{-6(n+1)b}[2n+1] + t^{-2(n+1)b}[2n+3], \\
 J_{T(2,b)}(n-1) &= -t^{(4n-2)b}J_{T(2,b)}(n) + t^{2nb}[2n-1], \\
 J_{T(2,b)}(n-2) &= t^{8(n-1)b}J_{T(2,b)}(n) - t^{6(n-1)b}[2n-1] + t^{2(n-1)b}[2n-3].
 \end{aligned}$$

Hence

$$\begin{aligned}
 RJ_{T(2,b)}(n) &= t^{-4b} (t^{4(n+2)b} J_{T(2,b)}(n+2) + t^{-4(n-2)b} J_{T(2,b)}(n-2)) \\
 &\quad + (t^{2b} + t^{-2b})(J_{T(2,b)}(n+1) + J_{T(2,b)}(n-1)) \\
 &\quad - (t^4 + t^{-4})t^{-2b} (t^{4(n+1)b} J_{T(2,b)}(n+1) + t^{-4(n-1)b} J_{T(2,b)}(n-1)) \\
 &\quad + ((t^{4nb} + t^{-4nb}) - 2(t + t^{-4}))J_{T(2,b)}(n) \\
 &= t^{-4b} (-t^{-2(n-1)b}[2n+1] + t^{2(n+3)b}[2n+3] \\
 &\quad - t^{2(n+1)b}[2n-1] + t^{-2(n-3)b}[2n-3]) \\
 &\quad + (t^{2b} + t^{-2b})(t^{-2nb}[2n+1] + t^{2nb}[2n-1]) \\
 &\quad - (t^4 + t^{-4})t^{-2b} (t^{2(n+4)b}[2n+1] + t^{-(2n+4)b}[2n-1]) \\
 &\quad - (t + t^{-4})t^{2b} (t^{2nb}[2n+1] + t^{-2nb}[2n-1]) \\
 &= t^{2b} t^{2nb} ([2n+3] + [2n-1] - (t^4 + t^{-4})[2n+1]) \\
 &\quad + t^{2b} t^{-2nb} ([2n-3] + [2n+1] - (t^4 + t^{-4})[2n-1]) \\
 &= 0,
 \end{aligned}$$

since  $[k+l] + [k-l] = (t^{2l} + t^{-2l})[k]$ . □

### 3.3 Proof of Theorem 1

We first note that the  $A$ -ideal  $\mathfrak{p}$ , the kernel of  $\theta: \mathfrak{t}^\sigma \rightarrow \mathbb{C}[\chi(X)]$ , is radical, ie,  $\sqrt{\mathfrak{p}} = \mathfrak{p}$ , since the character ring  $\mathbb{C}[\chi(X)]$  is reduced, ie, has nil-radical 0, by definition.

**Lemma 3.5** *Suppose  $\delta(t, M, L) \in \mathcal{A}_K$ . Then there are polynomials  $g(t, M) \in \mathbb{C}[t^{\pm 1}, M]$  and  $\gamma(t, M, L) \in \mathcal{T}$  such that*

$$(3) \quad \delta(t, M, L) = \frac{1}{g(t, M)} \gamma(t, M, L) \alpha_K(t, M, L).$$

Moreover,  $g(t, M)$  and  $\gamma(t, M, L)$  can be chosen so that  $\varepsilon(g) \neq 0$ .

**Proof** By definition  $\alpha_K$  is a generator of  $\tilde{\mathcal{A}}_K$ , the extension of  $\mathcal{A}_K$  in the principal left-ideal domain  $\tilde{\mathcal{T}}$ . Since  $\delta \in \mathcal{A}_K$ , it is divisible by  $\alpha_K$  in  $\tilde{\mathcal{T}}$ . Hence (3) follows.

We can assume that  $t+1$  does not divide both  $g(t, M)$  and  $\gamma(t, M, L)$  simultaneously. If  $\varepsilon(g) = 0$  then  $g$  is divisible by  $t+1$ , and hence  $\gamma$  is not. But then from the equality  $g\delta = \gamma\alpha_K$ , it follows that  $\alpha_K$  is divisible by  $t+1$ , which is impossible, since all the coefficients of powers of  $L$  in  $\alpha_K$  are supposed to be co-prime. □

Showing  $\sqrt{\varepsilon(\mathcal{A}_K^\sigma)} \subset \mathfrak{p}$  For torus knots, by Section 1, we have  $\varepsilon(\alpha_K) = f(M)A_K$ , where  $f(M) \in \mathbb{C}[M^{\pm 1}]$ . For every  $\delta \in \mathcal{A}_K$ , by Lemma 3.5, there exist  $g(t, M) \in \mathbb{C}[t^{\pm 1}, M]$  and  $\gamma \in \mathcal{T}$  such that  $\delta = \frac{1}{g(t, M)} \gamma \alpha_K$  and  $\varepsilon(g) \neq 0$ . It implies that

$$(4) \quad \varepsilon(\gamma) = \frac{1}{\varepsilon(g(M))} \varepsilon(\gamma) \varepsilon(\alpha_K) = \frac{1}{\varepsilon(g(M))} \varepsilon(\gamma) f(M) A_K.$$

The  $A$ -polynomial of a torus knot does not contain any non-trivial factor depending on  $M$  only. Since  $\varepsilon(\gamma) \in \mathfrak{t} = \mathbb{C}[L^{\pm 1}, M^{\pm 1}]$ , equation (4) implies that

$$h := \frac{1}{\varepsilon(g(M))} \varepsilon(\gamma) f(M)$$

is an element of  $\mathfrak{t}$ . Hence  $\varepsilon(\gamma) \in A_K \cdot \mathfrak{t}$ , the ideal of  $\mathfrak{t}$  generated by  $A_K$ . It follows that  $\varepsilon(\mathcal{A}_K) \subset A_K \cdot \mathfrak{t}$  and thus  $\varepsilon(\mathcal{A}_K^\sigma) \subset (A_K \cdot \mathfrak{t})^\sigma = \mathfrak{p}$ . Hence  $\sqrt{\varepsilon(\mathcal{A}_K^\sigma)} \subset \sqrt{\mathfrak{p}} = \mathfrak{p}$ .

Showing  $\mathfrak{p} \subset \sqrt{\varepsilon(\mathcal{A}_K^\sigma)}$  For  $a, b > 2$ , by Proposition 3.1 the colored Jones function of  $T(a, b)$  is annihilated by the operator  $PQ$ . Note that

$$\begin{aligned} \varepsilon(PQ) &= (L + L^{-1} - 2)^2 (L^2 M^{2ab} + L^{-2} M^{-2ab} - 2)^2 \\ &= L^{-2} (L^{-1} M^{-ab} (L - 1) (L^2 M^{2ab} - 1))^4. \end{aligned}$$

If  $u \in \mathfrak{p}$  then  $u = v A'_{T(a,b)}$ , where

$$A'_{T(a,b)} := L^{-1} M^{-ab} (L - 1) (L^2 M^{2ab} - 1) = L^{-1} M^{-ab} A_{T(a,b)}$$

and  $v \in \mathbb{C}[M^{\pm 1}, L^{\pm 1}]$ . It is easy to see that  $\sigma(v) = Lv$ , since  $\sigma(u) = u$  and  $\sigma(A'_{T(a,b)}) = L^{-1} A'_{T(a,b)}$ . This implies that  $\sigma(v^2 L) = \sigma(v)^2 L^{-1} = v^2 L$ . We then have

$$u^4 = v^4 A'^4_{T(a,b)} = \varepsilon(v^4 L^2 PQ) \in \varepsilon(\mathcal{A}_K^\sigma),$$

hence  $u \in \sqrt{\varepsilon(\mathcal{A}_K^\sigma)}$ .

For  $a = 2$ , by Proposition 3.4 the colored Jones function of  $T(2, b)$  is annihilated by the operator  $R$ . Note that  $\sigma(R) = R$  and

$$\varepsilon(R) = (L + L^{-1} - 2)(LM^{2b} + L^{-1} M^{-2b} + 2) = (L^{-1} M^{-b} (L - 1)(LM^{2b} + 1))^2.$$

If  $u \in \mathfrak{p}$  then  $u = v A'_{T(2,b)}$ , where

$$A'_{T(2,b)} := L^{-1} M^{-b} (L - 1)(LM^{2b} + 1) = L^{-1} M^{-b} A_{T(2,b)}$$

and  $v \in \mathbb{C}[M^{\pm 1}, L^{\pm 1}]$ . It is easy to see that  $\sigma(v) = -v$  and hence  $\sigma(v^2) = \sigma(v)\sigma(v) = v^2$ . We then have

$$u^2 = v^2 A'^2_{T(2,b)} = \varepsilon(v^2 R) \in \varepsilon(\mathcal{A}_K^\sigma),$$

hence  $u \in \sqrt{\varepsilon(\mathcal{A}_K^\sigma)}$ .

In both cases  $\mathfrak{p} \subset \sqrt{\varepsilon(\mathcal{A}_K^\sigma)}$ . Hence  $\sqrt{\varepsilon(\mathcal{A}_K^\sigma)} = \mathfrak{p}$  for all torus knots.

## References

- [1] **D Cooper, M Culler, H Gillet, D D Long, P B Shalen**, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. 118 (1994) 47–84 [MR1288467](#)
- [2] **M Culler, P B Shalen**, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. 117 (1983) 109–146 [MR683804](#)
- [3] **C Frohman, R Gelca, W Lofaro**, *The A-polynomial from the noncommutative viewpoint*, Trans. Amer. Math. Soc. 354 (2002) 735–747 [MR1862565](#)
- [4] **S Garoufalidis**, *On the characteristic and deformation varieties of a knot*, from: “Proceedings of the Casson Fest”, (C Gordon, Y Rieck, editors), Geom. Topol. Monogr. 7 (2004) 291–309 [MR2172488](#)
- [5] **S Garoufalidis, T T Q Lê**, *The colored Jones function is q-holonomic*, Geom. Topol. 9 (2005) 1253–1293 [MR2174266](#)
- [6] **R Gelca**, *On the relation between the A-polynomial and the Jones polynomial*, Proc. Amer. Math. Soc. 130 (2002) 1235–1241 [MR1873802](#)
- [7] **K Hikami**, *Difference equation of the colored Jones polynomial for torus knot*, Internat. J. Math. 15 (2004) 959–965 [MR2106155](#)
- [8] **V F R Jones**, *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc. 12 (1985) 103–111 [MR766964](#)
- [9] **T T Q Lê**, *The colored Jones polynomial and the A-polynomial of knots*, Adv. Math. 207 (2006) 782–804 [MR2271986](#)
- [10] **T Lê, A Tran**, *On the AJ conjecture for knots*, to appear in Transactions of the AMS [arXiv:1111.5258](#)
- [11] **A Lubotzky, A R Magid**, *Varieties of representations of finitely generated groups*, Mem. Amer. Math. Soc. 58 (1985) xi+117 [MR818915](#)
- [12] **H R Morton**, *The coloured Jones function and Alexander polynomial for torus knots*, Math. Proc. Cambridge Philos. Soc. 117 (1995) 129–135 [MR1297899](#)
- [13] **N Y Reshetikhin, V G Turaev**, *Ribbon graphs and their invariants derived from quantum groups*, Comm. Math. Phys. 127 (1990) 1–26 [MR1036112](#)
- [14] **A Sikora**, *Quantizations of character varieties and quantum knot invariants* [arXiv:0807.0943](#)

*Department of Mathematics, The Ohio State University  
100 Math Tower, 231 West 18th Avenue, Columbus OH 43210, USA*

[tran.350@osu.edu](mailto:tran.350@osu.edu)

Received: 22 November 2011