

Factorization rules in quantum Teichmüller theory

JULIEN ROGER

For a punctured surface S , a point of its Teichmüller space $\mathcal{T}(S)$ determines an irreducible representation of its quantization $\mathcal{T}^q(S)$. We analyze the behavior of these representations as one goes to infinity in $\mathcal{T}(S)$, or in the moduli space $\mathcal{M}(S)$ of the surface. The main result of this paper states that an irreducible representation of $\mathcal{T}^q(S)$ limits to a direct sum of representations of $\mathcal{T}^q(S_\gamma)$, where S_γ is obtained from S by pinching a multicurve γ to a set of nodes. The result is analogous to the factorization rule found in conformal field theory.

57M50; 32G15, 20G42

Let S be an oriented surface of genus g obtained from a closed compact surface \bar{S} by removing s punctures v_1, \dots, v_s . The *Teichmüller space* $\mathcal{T}(S)$ of S is the space of isotopy classes of complete hyperbolic metrics on S with finite area. It comes equipped with a natural Kähler metric, called the Weil–Petersson metric, which is invariant under the action of the mapping class group $\text{MCG}(S)$ on $\mathcal{T}(S)$. A quantization of the Teichmüller space was successfully described by L Chekhov and V V Fock in [10], and, in a slightly different setting, by R Kashaev in [12]. In the work of Chekhov and Fock, the main geometric ingredient is the notion of shear coordinates on the enhanced Teichmüller space $\tilde{\mathcal{T}}(S)$, which were introduced by W Thurston [19]. On the algebraic side, they make use of the quantum dilogarithm as described by L Faddeev and R Kashaev [8].

In the physics literature, the interest for the quantization of Teichmüller theory can be traced back to the work of E Verlinde and H Verlinde [21; 20]. In particular, H Verlinde conjectured in [20] that quantum Teichmüller theory should give rise to a family of representations of the mapping class groups which could be identified with a modular functor obtained from Liouville conformal field theory. The existence of such a modular functor associated to the quantum Teichmüller space was conjectured also by Fock [9] and was studied further by J Teschner [18]. A generalized version of this conjecture was made by Fock and A B Goncharov [11] in their study of (quantum) higher Teichmüller theory.

The goal of this paper is to investigate a similar question in the context of the exponential version of the quantum Teichmüller space studied by H Bai [3], F Bonahon and

X Liu [7; 14]. Given a parameter $q \in \mathbb{C}^*$, the quantum Teichmüller space $\mathcal{T}^q(S)$ is a non-commutative algebra, a deformation of an algebra of functions on $\mathcal{T}(S)$. For q a root of unity, Bonahon and Liu [7] described a complete classification of the finite-dimensional irreducible representations $\rho: \mathcal{T}^q(S) \rightarrow \text{End}(V)$. In particular, in the case when q is a primitive N^{th} root of unity with N odd and fixing weights $p_1, \dots, p_s \in \{0, \dots, N-1\}$ labelling the punctures v_1, \dots, v_s of S , one can associate to every hyperbolic metric $m \in \mathcal{T}(S)$ a unique irreducible representation $\rho_m: \mathcal{T}^q(S) \rightarrow \text{End}(V)$. Using a construction similar to the one described in [4], one can then construct a projective vector bundle $\mathcal{K}^q = \mathcal{K}^q(p_1, \dots, p_s)$ over $\mathcal{T}(S)$ with fiber $\mathbb{P}V$, where the fiber at $m \in \mathcal{T}(S)$ is endowed with the action of the irreducible representation ρ_m of $\mathcal{T}^q(S)$. This construction behaves well under the action of the mapping class group $\text{MCG}(S)$ and we obtain a projective vector bundle $\widetilde{\mathcal{K}}^q$ over the moduli space $\mathcal{M}(S) = \mathcal{T}(S)/\text{MCG}(S)$.

In the spirit of conformal field theory, one can then ask if this bundle extends to the Deligne–Mumford compactification $\overline{\mathcal{M}}(S)$ of the moduli space. To study this question, we analyze how the representation ρ_m of $\mathcal{T}^q(S)$ breaks down when the metric m approaches a point of $\overline{\mathcal{M}}(S) \setminus \mathcal{M}(S)$, that is, when the lengths of a finite number of geodesics of S tend to 0 for this metric. The result we obtain can be interpreted as a *factorization rule* for this theory.

More precisely, let λ be an *ideal triangulation* of S , that is, a triangulation of S with vertices at the punctures. Following the construction of Chekhov and Fock in [10], Bonahon and Liu [7; 14] associate to S , the triangulation λ and a parameter $q \in \mathbb{C}^*$, an algebra $\mathcal{T}_\lambda^q(S)$ called the *Chekhov–Fock algebra*. It is the skew-commutative algebra over \mathbb{C} with generators $X_1^{\pm 1}, \dots, X_n^{\pm 1}$ associated to the edges of λ and relations $X_i X_j = q^{2\sigma_{ij}} X_j X_i$, where the $\sigma_{ij} \in \{-2, -1, 0, 1, 2\}$ are the coefficients of the Weil–Petersson Poisson structure on the enhanced Teichmüller space $\widetilde{\mathcal{T}}(S)$, parametrized using Thurston’s shear coordinates.

Approaching a point in the boundary of $\overline{\mathcal{M}}(S)$ corresponds to shrinking a finite number of non-intersecting geodesics of S to points. Hence we give ourselves a finite union of non-intersecting, non-homotopic, essential simple closed curves $\gamma = \bigcup_i \gamma_i \subset S$ and we consider the surface $S_\gamma = S \setminus \gamma$. One should think of S_γ as being obtained from S by pinching the multicurve γ to a set of nodes and removing them. It is a possibly disconnected surface with two new punctures for each curve removed. The ideal triangulation λ on S induces an ideal triangulation λ_γ on S_γ whose edges are given by taking homotopy classes of edges of $\lambda \cap S_\gamma$.

An essential step is to relate the quantum Teichmüller spaces of S and S_γ .

Proposition 1 For every ideal triangulation λ and every multicurve γ on S , there exists an algebra homomorphism

$$\Theta_{\gamma,\lambda}^q: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \longrightarrow \mathcal{T}_\lambda^q(S)$$

described explicitly by sending each generator of $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$ to certain monomials in $\mathcal{T}_\lambda^q(S)$.

The existence of this homomorphism is the algebraic translation of the fact that the Weil–Petersson Poisson structure extends naturally to the completion $\overline{\mathcal{T}(S)}$ of the Teichmüller space for the Weil–Petersson metric (see H Masur [15] and S Wolpert [23]). This completion is called the *augmented Teichmüller space* and was introduced by W Abikoff [2] and L Bers [5]. As a set, it is the union of $\mathcal{T}(S)$ and of the $\mathcal{T}(S_\gamma)$ for every multicurve γ . The action of $\text{MCG}(S)$ on $\mathcal{T}(S)$ extends to $\overline{\mathcal{T}(S)}$ and the quotient $\overline{\mathcal{T}(S)}/\text{MCG}(S)$ can be identified topologically with $\overline{\mathcal{M}(S)}$. The key geometric ingredient is given by an extension of the shear parameters on $\mathcal{T}(S)$ to the strata $\mathcal{T}(S_\gamma)$ of this augmentation.

When q is a primitive N^{th} root of unity with N odd, Bonahon and Liu associate an irreducible representation $\rho_m: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ of the Chekhov–Fock algebra to each metric $m \in \mathcal{T}(S)$ and weights $p_1, \dots, p_s \in \{0, \dots, N-1\}$ labelling the punctures v_1, \dots, v_s of S . The representation ρ_m is defined uniquely up to isomorphism and varies continuously with the metric m . We proceed with studying the behavior of ρ_m when m approaches a lower-dimensional stratum $\mathcal{T}(S_\gamma)$ in $\overline{\mathcal{T}(S)}$.

For simplicity, let us restrict attention in the introduction to the case when γ consists of a single curve.

Theorem 2 Let $m_t \in \mathcal{T}(S)$ be a continuous family of hyperbolic metrics such that, as $t \rightarrow 0$, m_t converges to $m_\gamma \in \mathcal{T}(S_\gamma)$ in $\overline{\mathcal{T}(S)}$. Let $\rho_t: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ be a continuous family of irreducible representations classified by m_t and weights $p_1, \dots, p_s \in \{0, \dots, N-1\}$ labelling the punctures v_1, \dots, v_s of S .

Then, as $t \rightarrow 0$, the representation

$$\rho_t \circ \Theta_{\gamma,\lambda}^q: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}(V)$$

approaches

$$\bigoplus_{i=0}^{N-1} \rho_\gamma^i: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}\left(\bigoplus_{i=0}^{N-1} V_i\right)$$

where, for each i , $\rho_\gamma^i: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}(V_i)$ is the irreducible representation classified by m_γ , the weights p_1, \dots, p_s labelling the old punctures and the weight i labelling the two new punctures v' and v'' of S_γ .

The next step is to show that this decomposition is well-behaved under changes of triangulations. More precisely, for any two triangulations λ and λ' of S , Chekhov and Fock introduce *quantum coordinate change isomorphisms* $\Phi_{\lambda\lambda'}^q: \widehat{\mathcal{T}}_{\lambda'}^q(S) \rightarrow \widehat{\mathcal{T}}_{\lambda}^q(S)$ between the fraction division algebras associated to the Chekhov–Fock algebras $\mathcal{T}_{\lambda'}^q(S)$ and $\mathcal{T}_{\lambda}^q(S)$. The *quantum Teichmüller space* $\mathcal{T}^q(S)$ of S is then defined to be the union of the $\widehat{\mathcal{T}}_{\lambda}^q(S)$ for every triangulation λ , modulo the relation which identifies $X' \in \widehat{\mathcal{T}}_{\lambda'}^q(S)$ to $\Phi_{\lambda\lambda'}^q(X') \in \widehat{\mathcal{T}}_{\lambda}^q(S)$.

The coordinate change isomorphisms are only defined between fraction algebras. For certain representations ρ_{λ} of $\mathcal{T}_{\lambda}^q(S)$ the composition with $\Phi_{\lambda,\lambda'}^q$ still makes sense and we obtain a representation

$$\rho_{\lambda} \circ \Phi_{\lambda,\lambda'}^q: \mathcal{T}_{\lambda'}^q(S) \rightarrow \text{End}(V).$$

A representation ρ of $\mathcal{T}^q(S)$ is then a family of representations $\{\rho_{\lambda}\}_{\lambda}$ of $\mathcal{T}_{\lambda}^q(S)$ such that $\rho_{\lambda} \circ \Phi_{\lambda\lambda'}^q = \rho_{\lambda'}$ for every pair of triangulations λ and λ' . In particular, given $m \in \mathcal{T}(S)$ and weights p_1, \dots, p_s labelling the punctures of S , one obtains a representation $\rho_m = \{\rho_{m,\lambda}\}_{\lambda}$ where $\rho_{m,\lambda}: \mathcal{T}_{\lambda}^q(S) \rightarrow \text{End}(V)$ is the irreducible representation of $\mathcal{T}_{\lambda}^q(S)$ classified by these data.

Theorem 3 *Let $\rho_t = \{\rho_{t,\lambda}\}_{\lambda}$ be a continuous family of irreducible representations of $\mathcal{T}^q(S)$ classified by weights p_1, \dots, p_s and a continuous family $m_t \in \mathcal{T}(S)$ such that m_t approaches $m_{\gamma} \in \mathcal{T}(S_{\gamma})$ as $t \rightarrow 0$. For each triangulation λ of S , we let the limit*

$$\lim_{t \rightarrow 0} \rho_{t,\lambda} \circ \Theta_{\gamma,\lambda}^q = \bigoplus_i \rho_{\gamma,\lambda}^i$$

be given as in Theorem 2.

Then, for any triangulations λ and λ' and weight i , we have

$$\rho_{\gamma,\lambda'}^i = \rho_{\gamma,\lambda}^i \circ \Phi_{\lambda\lambda'}^q.$$

Hence the family of representations $\{\rho_{\gamma,\lambda}^i\}_{\lambda}$ determines an irreducible representation $\rho_{\gamma}^i: \mathcal{T}^q(S_{\gamma}) \rightarrow \text{End}(V)$, which is classified by m_{γ} , the weights p_1, \dots, p_s labelling the old punctures, and the weight i labelling the new punctures v' and v'' of S_{γ} .

Interpretation and perspectives

In his seminal work [22], E Witten described a correspondence between Chern–Simons theory and conformal field theory for compact gauge groups. In this context, the theory is rational, implying in particular that the spaces of conformal blocks are finite-dimensional. The dimensions of the subspaces in the decompositions for the

corresponding factorization rules must satisfy so-called *fusion rules* leading to the celebrated Verlinde formula.

In analogy with Witten's work, H Verlinde proposed in [20] to relate Chern–Simons theory for the non-compact group $\mathrm{SL}_2(\mathbb{R})$ to non-rational conformal field theory in the context of quantum Liouville theory (see more recently Teschner [17]). The moduli space of flat $\mathrm{SL}_2(\mathbb{R})$ -connections being non-compact, its quantization gives rise to an infinite-dimensional Hilbert space \mathcal{H} , which Verlinde proposes to identify with the space of conformal blocks of a conformal field theory. Via this identification, and restricting to the Teichmüller space, Verlinde argues that the factorization rule for \mathcal{H} corresponds to its decomposition into the eigenspaces of the action of the quantum length operators L_γ associated to simple closed curves γ . Unlike the rational case, the fusion rules are trivial: the spectrum of these operators is simple and equal to $(0, \infty)$, corresponding geometrically to the range of the length function ℓ_γ (see Kashaev [13]). Following the same approach as in the rational case, one can try to construct a family of projective representations of the mapping class groups by studying the “change of bases” corresponding to decompositions of \mathcal{H} for different pants decompositions of S , that is, maximal collections of essential simple closed curves [18].

In our context, given an ideal triangulation λ , the relevant operator associated to a curve γ is the one introduced at the beginning of the proof of Theorem 17, called X_γ . It is the quantum analogue of the exponential graph length function x_γ (see Lemma 7). In particular, given a finite-dimensional irreducible representation of the Chekhov–Fock algebra $\rho_m: \mathcal{T}_\lambda^q(S) \rightarrow \mathrm{End}(V)$ associated to a point $m \in \mathcal{T}(S)$, the decomposition of V exhibited in Theorem 2 is given by the eigenspaces of the operator X_γ . Its spectrum is simple and of the form $\{c_\gamma(m)q^i \mid i = 0, \dots, N-1\}$, for some parameter $c_\gamma(m) \in (1, \infty)$ determined by m . As such, the fusion rules for the theory are once again trivial.

The specificity of the finite-dimensional approach to quantum Teichmüller theory, however, is its geometrical flavor. In particular, when letting m approach a metric m_γ pinched along γ , one can see that the length function ℓ_γ is asymptotically equivalent to $\log x_\gamma$. The parameter in the spectrum of X_γ is in fact $c_\gamma(m) = \sqrt[N]{x_\gamma(m)}$, possibly after a shift by an N^{th} root of unity. As such, although the decomposition of V can be obtained at any point $m \in \mathcal{T}(S)$, it becomes geometrically meaningful when approaching a lower-dimensional stratum $\mathcal{T}(S_\gamma)$.

The correct framework for studying this behavior further should be in terms of the projective vector bundle $\widetilde{\mathcal{K}}^q$ over the moduli space $\mathcal{M}(S)$ discussed in the introduction. The first question to investigate will be whether the decomposition discussed above provides a natural extension of this bundle to the Deligne–Mumford compactification

$\overline{\mathcal{M}(S)}$ of the moduli space. The independence of the factorization rule under changes of triangulations (see Proposition 25) should be the key ingredient in this extension. A tantalizing question at this point is if this extension can be used to construct a family of finite-dimensional representations of mapping class groups. We plan to study this question further; however, we do not conjecture at this point if such a construction exists or not.

Plan of the paper

In the first section we introduce the necessary background on Teichmüller space and its augmentation, shear coordinates and the Weil–Petersson Poisson structure. Section 2 is dedicated to the study of a natural extension of shear coordinates to the augmented Teichmüller space and to the proof that the Weil–Petersson Poisson bracket extends to the augmentation for this extension. In Section 3, we apply this construction to the study of the representation theory of the Chekhov–Fock algebras, which are versions of the quantum Teichmüller space for a fixed triangulation. The last section is dedicated to the study of the independence of the result under changes of triangulations, thus extending it to the quantum Teichmüller space itself.

Acknowledgments

Most of the content of this paper was written as a graduate student at the University of Southern California under the supervision of Francis Bonahon. It is with great pleasure that I thank him for his support and continued interest in my work. His healthy skepticism during our frequent conversations made my modest victories all the more rewarding. I would also like to thank Ko Honda and Bob Penner for their support, as well as my fellow students at the time, Roman Golovko and Dmytro Chebotarov, for our frequent mathematical discussions. I would also like to thank Stéphane Baseilhac and Feng Luo for several enlightening conversations regarding this work.

This research was partially supported by the grant DMS-0604866 from the National Science Foundation.

1 Geometric background

Throughout this paper, S will be an oriented surface of genus g obtained from a closed surface \overline{S} without boundary by removing s punctures v_1, \dots, v_s . We will assume that S has at least one puncture and has Euler characteristic $\chi(S) = 2 - 2g - s < 0$. The simplest such surfaces are the spheres with 3 or 4 punctures and the once-punctured torus.

1.1 Teichmüller spaces

For the purpose of this paper we will need two variants of Teichmüller space. The first and most classical one, denoted simply as $\mathcal{T}(S)$, will be the set of isotopy classes of complete hyperbolic metrics on S with finite area. We will call this space simply the *Teichmüller space* of S . However, we will need to drop the finite area condition to describe the exponential shear coordinates, which are essential to the definition of the quantum Teichmüller space. Let $\text{Conv}(S, m)$ denote the *convex core* of S , that is, the smallest non-empty closed convex subset of (S, m) . It is a surface with cusps and geodesic boundaries whose interior is homeomorphic to S . If (S, m) has finite area then its convex core consists of the whole surface. Otherwise, some of the punctures of (S, m) will have a neighborhood isometric to an infinite area funnel bounded by one of the geodesic boundaries of $\text{Conv}(S, m)$. We let $\widetilde{\mathcal{T}}(S)$ be the space of isotopy classes of complete hyperbolic metrics on S , possibly with infinite volume, together with an orientation of each of the boundary components of $\text{Conv}(S, m)$. The set $\widetilde{\mathcal{T}}(S)$ is called the *enhanced Teichmüller space* of S . In particular, since, for a complete hyperbolic metric, $\text{Conv}(S, m)$ has no boundary component, there is a natural embedding of $\mathcal{T}(S)$ into $\widetilde{\mathcal{T}}(S)$.

1.2 The augmented Teichmüller space

We will also need to consider the *augmented Teichmüller space* $\overline{\mathcal{T}}(S)$, which was introduced by Abikoff [1; 2] and Bers [5], and was further studied by Masur [15] (see more recently Wolpert [24] and references therein). We will briefly recall its construction and some of its properties.

Let $\gamma = \gamma_1 \cup \cdots \cup \gamma_k$ be the union of k disjoint, non-homotopic, essential simple closed curves. Such a γ will be called a *multicurve*. Alternatively, γ corresponds to a $(k - 1)$ -simplex in $C(S)$, the *complex of curves* of S . We will denote by S_γ the surface obtained from S by removing the multicurve γ . It is a possibly disconnected surface with two new punctures for each curve removed. As a set, we define

$$\overline{\mathcal{T}}(S) = \mathcal{T}(S) \cup \bigcup_{\gamma \in C(S)} \mathcal{T}(S_\gamma),$$

where $\mathcal{T}(S_\gamma)$ is the product of Teichmüller spaces associated to the connected components of S_γ . The $\mathcal{T}(S_\gamma)$ are called the *strata* of $\overline{\mathcal{T}}(S)$.

A topology on $\overline{\mathcal{T}}(S)$ can be defined as follows: a sequence of metrics $(m_n)_n$ in $\mathcal{T}(S)$ converges to $m_\gamma \in \mathcal{T}(S_\gamma)$ if, as $n \rightarrow \infty$, the length $\ell_{m_n}(\gamma_i)$ of (the geodesic representative of) γ_i for m_n tends to 0 for every i , and m_n converges uniformly to m_γ on every compact subset of S_γ .

The action of the mapping class group $\text{MCG}(S)$ of S on the Teichmüller space extends to $\overline{\mathcal{T}(S)}$ and the quotient $\overline{\mathcal{T}(S)}/\text{MCG}(S)$ can be identified, as a topological space, with the Deligne–Mumford compactification $\overline{\mathcal{M}(S)}$ of the moduli space $\mathcal{M}(S) = \mathcal{T}(S)/\text{MCG}(S)$ [2].

1.3 Exponential shear coordinates

One of the main ingredients for the quantization of Teichmüller space as described first in [10] is the notion of shear coordinates introduced by W Thurston [19] (see also Bonahon [6]). We will describe here their exponential version, following for example [14]. We use the notation $\mathbb{R}_+ = (0, \infty)$.

Since $\chi(S) < 0$ and S has at least one puncture, it admits an *ideal triangulation* $\lambda = \{\lambda_1, \dots, \lambda_n\}$, that is, a triangulation of \overline{S} with vertices v_1, \dots, v_s and edges $\lambda_1, \dots, \lambda_n$, where the edges are considered up to isotopy. The number of edges of an ideal triangulation depends only on the Euler characteristic of S and is given by $n = -3\chi(S) = 6g + 3s - 6$. If we endow S with a hyperbolic metric m , each edge λ_i is isotopic to a unique geodesic g_i for this metric. One can then associate to λ_i a number $x_i \in \mathbb{R}_+$, called the *exponential shear parameter* of m along λ_i , obtained as follows: let \tilde{g}_i be a lift of g_i to the universal cover of (S, m) , which we identify with the upper half-space \mathbb{H}^2 . \tilde{g}_i separates two triangles \tilde{T}_i^1 and \tilde{T}_i^2 , bounded by lifts of edges of λ so that the union $\tilde{Q}_i = \tilde{g}_i \cup \tilde{T}_i^1 \cup \tilde{T}_i^2$ forms a square in \mathbb{H}^2 with vertices on the projective real line bounding \mathbb{H}^2 . For a given orientation of \tilde{g}_i , we name the vertices of \tilde{Q}_i by z_-, z_+, z_r, z_l , such that \tilde{g}_i goes from z_- to z_+ , and z_r and z_l are respectively to the right and to the left of \tilde{g}_i . The exponential shear parameter of m along λ_i is defined as

$$x_i = -\text{cross-ratio}(z_r, z_l, z_-, z_+) = -\frac{(z_r - z_-)(z_l - z_+)}{(z_r - z_+)(z_l - z_-)} \in \mathbb{R}_+.$$

Geometrically, $\log x_i$ corresponds to the (signed) distance between the orthogonal projections of z_r and z_l onto \tilde{g}_i .

Conversely, one can construct a (possibly incomplete) hyperbolic metric m from any choice of parameters $x_1, \dots, x_n \in \mathbb{R}_+$ associated to the edges of λ , obtained by gluing ideal hyperbolic triangles into squares whose vertices have the prescribed cross-ratio. Its completion is a hyperbolic surface S' with geodesic boundaries and cusps, for which each end of the edges of λ either converges to a cusp or spirals around a geodesic boundary. The direction of the spiraling provides an orientation of the geodesic boundary. This metric on S' admits a unique extension to a complete metric on S whose convex core is S' . In this way, one obtains a homeomorphism $\phi_\lambda: \overline{\mathcal{T}(S)} \rightarrow \mathbb{R}_+^n$ for every ideal triangulation λ of S .

Given x_1, \dots, x_n the shear parameters of $m \in \widetilde{\mathcal{T}}(S)$ associated to a triangulation λ , one can read the geometry of (S, m) around each puncture v_j of S as follows: let $p_j = x_1^{k_{1j}} \dots x_n^{k_{nj}}$ where $k_{ij} \in \{0, 1, 2\}$ is the number of ends of the edge λ_i that converge to v_j . Then, if $p_j = 1$, v_j is a cusp. Otherwise, $|\log p_j|$ is the length of the boundary component of $\text{Conv}(S, m)$ facing v_j . Its sign corresponds to the orientation of the boundary component with respect to the orientation of $\text{Conv}(S, m)$. As a consequence, the homeomorphism ϕ_λ restricts to an embedding of $\mathcal{T}(S)$ into \mathbb{R}_+^n , corresponding to setting all the p_j equal to 1.

The shear parameters are defined along each edge of λ or equivalently for pairs of adjacent triangles. This definition generalizes to any pair of triangles in the universal cover of S as follows: let $\widetilde{\lambda}$ be the lift of λ to the universal cover \widetilde{S} of (S, m) . Let P and Q be two ideal triangles in \widetilde{S} delimited by $\widetilde{\lambda}$. Let $\widetilde{\lambda}_{i_1}, \dots, \widetilde{\lambda}_{i_l}$, lifts of $\lambda_{i_1}, \dots, \lambda_{i_l}$ respectively, be the set of edges of $\widetilde{\lambda}$ separating P and Q . We include in this set the edges of P and Q that are closest to each other. The shearing cocycle σ of $m \in \mathcal{T}(S)$ associated to λ is defined for such triangles by

$$\sigma(P, Q) = \sum_{j=1}^l \log x_{i_j},$$

where x_{i_j} is the shearing parameter of m for the edge λ_{i_j} . Some properties of the shearing cocycles will be needed later on and we refer to [6] for more details.

1.4 The Weil–Petersson Poisson structure

The enhanced Teichmüller space can be endowed with a Poisson structure that admits a simple expression in the logarithmic shear coordinates associated to a triangulation λ of S (see for example [9]). The collection of triangles $S \setminus \lambda$ has $2n$ spikes converging toward the punctures, each of them delimited by edges λ_i and λ_j not necessarily distinct. For $i, j \in \{1, \dots, n\}$, let $a_{ij} \in \{0, 1, 2\}$ be the number of spikes of $S \setminus \lambda$ that are delimited to the left by λ_i and to the right by λ_j , when looking toward the end of the spikes. The Weil–Petersson Poisson structure is given in coordinates by the following bi-vector

$$\Pi_{WP} = \sum_{i,j} \sigma_{ij} \frac{\partial}{\partial \log x_i} \wedge \frac{\partial}{\partial \log x_j},$$

where $\sigma_{ij} = a_{ij} - a_{ji} \in \{-2, -1, 0, 1, 2\}$. Its degeneracy is well understood: The lengths $\log p_j$ associated to the punctures v_j are Casimir functions for Π_{WP} and the cusped Teichmüller space

$$\mathcal{T}(S) = \{\log p_1 = \dots = \log p_s = 0\}$$

is a symplectic leaf for this structure. The induced symplectic form on $\mathcal{T}(S)$ can then be identified with the Kähler form associated to the usual Weil–Petersson metric on Teichmüller space.

2 Pinching along curves: geometric aspects

We suppose once again that S is an oriented surface of genus g with $s \geq 1$ punctures v_1, \dots, v_s and such that $\chi(S) < 0$. Let $\gamma = \gamma_1 \cup \dots \cup \gamma_k$ be a multicurve and $S_\gamma = S \setminus \gamma$. It is homeomorphic to a surface with $s + 2k$ punctures: the “old” ones v_1, \dots, v_s and two new punctures v'_i and v''_i corresponding to the removal of γ_i for $i = 1, \dots, k$. Alternatively one can think of S_γ as being obtained from S by pinching the multicurve γ to k nodes and removing them. Note that S_γ may be disconnected.

The goal of this section is to describe explicitly the behavior of the shear coordinates and the Weil–Petersson Poisson structure when going from $\mathcal{T}(S)$ to $\mathcal{T}(S_\gamma)$ in the topology of the augmented Teichmüller space $\overline{\mathcal{T}(S)}$.

2.1 Induced ideal triangulations

Given an ideal triangulation $\lambda = \{\lambda_1, \dots, \lambda_n\}$ of S , we want to define an induced ideal triangulation λ_γ of S_γ . We can choose $\gamma = \bigcup_i \gamma_i$ so that the γ_i never cross the same edge twice in a row. Let $\lambda \setminus \gamma$ denote the family of arcs obtained from the edges of λ intersected with S_γ . We group these arcs into distinct isotopy classes μ_1, \dots, μ_l in S_γ , each consisting of a certain number of segments of $\lambda_1, \dots, \lambda_n$. Hence $\{\mu_1, \dots, \mu_l\}$ is a family of non-intersecting and non-homotopic arcs in S_γ .

Lemma 4 *The family $\lambda_\gamma = \{\mu_1, \dots, \mu_n\}$ is an ideal triangulation of S_γ consisting of n distinct isotopy classes of arcs.*

Proof Since the γ_i do not backtrack, $\lambda \setminus \gamma$ decomposes S_γ into pieces of the form given in Figure 1, where the dashed lines represent γ and the first piece can have 0, 1, 2 or 3 dashed sides. The first piece is an ideal triangle in S_γ and the other two are bigons. One can collapse these bigons successively to arcs with one or two vertices among the new punctures of S_γ . This can be done for all the bigons successively unless one of the situations described in Figure 2 occurs. This cannot happen, however, since the γ_i are essential and non-homotopic to each other. The remaining arcs correspond to the homotopy classes μ_1, \dots, μ_l and decompose S_γ into ideal triangles. Since $\chi(S_\gamma) = \chi(S)$ and the number of edges of an ideal triangulation depends only on the Euler characteristic, the ideal triangulation $\lambda_\gamma = \{\mu_1, \dots, \mu_n\}$ of S_γ has the same number of edges as λ . \square




Figure 1




Figure 2

We call λ_γ the *ideal triangulation of S_γ induced by λ* .

Remark 5 In practice each edge μ_i of λ_γ is obtained from λ by considering a maximal sequence of adjacent bigons in the decomposition of S_γ by $\lambda \setminus \gamma$ and collapsing it to an edge. Via this process, ideal triangles for λ_γ on S_γ are identified naturally with ideal triangles for λ on S (see Figure 3).




Figure 3

2.2 Extension of the shear coordinates

Let $\lambda = \{\lambda_1, \dots, \lambda_n\}$ be an ideal triangulation of S and $\lambda_\gamma = \{\mu_1, \dots, \mu_n\}$ be the induced triangulation of S_γ . We suppose that, for $i = 1, \dots, n$, μ_i corresponds to the

homotopy class of k_{ij} segments from λ_j for $j = 1, \dots, n$. The following proposition relates the shear coordinates on $\mathcal{T}(S)$ associated to λ to the ones on $\mathcal{T}(S_\gamma)$ associated to λ_γ .

Proposition 6 *Let $m_t \in \mathcal{T}(S)$ be a continuous family of hyperbolic metrics on S , $(x_1(t), \dots, x_n(t)) \in \mathbb{R}_+^n$ their shear parameters for λ , and $m_\gamma \in \mathcal{T}(S_\gamma)$, $(y_1, \dots, y_n) \in \mathbb{R}_+^n$ its shear parameters for λ_γ . Then*

$$m_t \xrightarrow[t \rightarrow 0]{} m_\gamma \text{ in } \overline{\mathcal{T}(S)} \implies \lim_{t \rightarrow 0} x_1^{k_{i1}}(t) \cdots x_n^{k_{in}}(t) = y_i \text{ for } i = 1, \dots, n.$$

Proof Let μ_i be an edge of λ_γ in S_γ and $\tilde{\mu}_i$ be one of its lifts to the universal cover of (S_γ, m_γ) . Then $\tilde{\mu}_i$ is the diagonal of a square consisting of two ideal triangles P_0^i and Q_0^i . Note that the universal cover of (S_γ, m_γ) is isometric to several copies of \mathbb{H}^2 , one for each connected component of S_γ . We denote by \tilde{S}_γ the one containing $\tilde{\mu}_i$. By Remark 5, μ_i corresponds to a rectangle composed of a succession of bigons in the decomposition of S_γ by $\lambda \smallfrown \gamma$, ending at the sides of two (non-necessarily distinct) triangles. We consider a lift of this rectangle to the universal cover \tilde{S}_t of (S, m_t) . It ends at the sides of two triangles P_t^i and Q_t^i , which are separated by k_{ij} lifts of the edge λ_j for $j = 1, \dots, n$. Hence the shearing cocycle σ_t associated to m_t satisfies

$$\sigma_t(P_t^i, Q_t^i) = \sum_j k_{ij} \log x_j(t).$$

In addition, since $m_t \rightarrow m_\gamma$, these lifts can be chosen so that, with the right identification of \tilde{S}_γ with \tilde{S}_t , P_t^i and Q_t^i approach P_0^i and Q_0^i , respectively, as $t \rightarrow 0$.

We can then use the following inequality, derived from [6, Lemma 8]: if a_t (resp. b_t) is the projection of the third vertex of P_t^i (resp. Q_t^i) onto g_t (resp. h_t), where g_t and h_t are the edges of P_t^i and Q_t^i that are the closest to each other, and if b'_t is the projection of b_t onto g_t , then

$$|\sigma_t(P_t^i, Q_t^i) - d(a_t, b'_t)| \leq \ell_{m_t}(\gamma).$$

We refer to Figure 4 for an example with notation.

Since P_t^i and Q_t^i approach P_0^i and Q_0^i respectively, we have $a_t \rightarrow a_0$ and $b_t, b'_t \rightarrow b_0$ where a_0 and b_0 are the projections of the third vertex of P_0^i and Q_0^i , respectively, onto $g_0 = h_0 = \tilde{\mu}_i$. Then

$$\begin{aligned} \lim_{t \rightarrow 0} \sum_j k_{ij} \log x_j(t) &= \lim_{t \rightarrow 0} \sigma_t(P_t^i, Q_t^i) = \lim_{t \rightarrow 0} d(a_t, b'_t) \\ &= d(a_0, b_0) \\ &= \sigma_0(P_0^i, Q_0^i) = \log y_i. \quad \square \end{aligned}$$




Figure 4

In other words, the shear parameter on $\mathcal{T}(S_\gamma)$ associated to an edge μ_i of λ_γ is the limit of a monomial in the shear parameters on $\mathcal{T}(S)$ associated to λ . This monomial is given by the product of parameters associated to the (segments of) edges of λ constituting the homotopy class of μ_i .

Another important monomial can be associated to γ itself, when it consists of one simple closed curve. We denote by c_i the number of times γ intersects the edge λ_i and let

$$x_\gamma = x_1^{c_1} \cdots x_n^{c_n}.$$

We call this monomial the *exponential graph length* of γ .

If $m_t \in \mathcal{T}(S)$, $t \in (0, 1]$, is a family of hyperbolic metrics with shear parameters $x_1(t), \dots, x_n(t)$ and $x_\gamma(t)$ is the associated exponential graph length of γ , we have the following lemma.

Lemma 7 *If, as $t \rightarrow 0$, $\ell_{m_t}(\gamma)$ approaches 0, then the exponential graph length $x_\gamma(t)$ of γ approaches 1.*

Proof This is a consequence of the formula for the length of γ in shear coordinates as described for example in [10]. In particular, we have

$$(1) \quad 2 \cosh(\ell_{m_t}(\gamma)/2) = x_\gamma^{1/2}(t) + x_\gamma^{-1/2}(t) + \dots,$$

where all the terms in the sum are positive.

Since the left hand side approaches 2, we see that $x_\gamma(t)$ must approach 1 as $t \rightarrow 0$. \square

It also follows from this proof that the other terms on the right hand side of (1) approach 0 as $t \rightarrow 0$. This means that, near the stratum $\mathcal{T}(S_\gamma)$, the length function $\ell_\cdot(\gamma)$ is asymptotically equivalent to the graph length function $\log x_\gamma$. This explains the essential rôle the quantum analogue X_γ of x_γ will play later on.

2.3 Extension of the Weil–Petersson Poisson structure

Masur [15] (see also Wolpert [23]) proved that the Weil–Petersson Kähler metric on $\mathcal{T}(S)$ extends in an appropriate sense to its augmentation $\overline{\mathcal{T}(S)}$ and can be identified with the Weil–Petersson metric on the lower-dimensional strata $\mathcal{T}(S_\gamma)$. We would like to know how this fact together with Proposition 6 translate in terms of the expression of the Weil–Petersson Poisson structures on $\mathcal{T}(S)$ and $\mathcal{T}(S_\gamma)$ in the shear coordinates associated to λ and λ_γ respectively. To do so we use Lemma 8 given below, which is a homological interpretation of the Weil–Petersson structure as described for example in [7]. Proposition 9 can then be interpreted as a topological translation of the result of Masur.

Let $\sigma = (\sigma_{ij})_{ij}$ be the matrix of coefficients of the Weil–Petersson Poisson structure on $\widetilde{\mathcal{T}(S)}$ in the coordinates $(\log x_1, \dots, \log x_n)$ associated to an ideal triangulation $\lambda = \{\lambda_1, \dots, \lambda_n\}$, as was described in Section 1.4. Setting $\sigma(\lambda_i, \lambda_j) = \sigma_{ij}$, σ can be identified with an antisymmetric bilinear form on $\mathcal{H}(\lambda, \mathbb{Z}) \cong \mathbb{Z}^n$, the free abelian group generated over the set of edges of λ . We are going to use a homological interpretation of σ as given for example in [7]. This formulation follows [6] where it is used to describe the Thurston symplectic form.

Let G be the dual graph of λ and \widehat{G} be the oriented graph obtained from G by keeping the same vertex set and replacing each edge of G by two oriented edges that have the same endpoints as the original edge but with opposite orientations.

There is a unique way to thicken \widehat{G} into a surface \widehat{S} such that:

- (1) \widehat{S} deformation retracts to \widehat{G} .
- (2) As one goes around a vertex \widehat{v} of \widehat{G} in \widehat{S} , the orientation of the edges of \widehat{G} ending at \widehat{v} points alternately toward and away from \widehat{v} .
- (3) The natural projection $p: \widehat{G} \rightarrow G$ extends to a 2-fold cover $\widehat{S} \rightarrow S$, branched along the vertex set of \widehat{G} .

Let $\eta: \widehat{S} \rightarrow \widehat{S}$ be the covering involution of the branched cover $p: \widehat{S} \rightarrow S$.

Lemma 8 *The group $\mathcal{H}(\lambda, \mathbb{Z})$ can be identified with the subgroup of $H_1(\widehat{S})$ consisting of those $\widehat{\alpha}$ such that $\eta_*(\widehat{\alpha}) = -\widehat{\alpha}$. In addition, if $\alpha, \beta \in \mathcal{H}(\lambda, \mathbb{Z})$ correspond to $\widehat{\alpha}, \widehat{\beta} \in H_1(\widehat{S})$, then $\sigma(\alpha, \beta) = \widehat{\alpha} \cdot \widehat{\beta}$; their algebraic intersection number.*

Proof See [7, Lemmas 6 and 7]. □

The identification of Lemma 8 is given as follows: if $e_i \in \mathcal{H}(\lambda, \mathbb{Z})$ is the element associating weight 1 to the edge λ_i and weight 0 to the other edges, \hat{e}_i is the lift in \hat{G} of the edge of G dual to λ_i . The closed curve \hat{e}_i comes with a natural orientation given by the one on \hat{G} , and we identify \hat{e}_i with its homology class in $H_1(\hat{S})$. More generally, to $\alpha = \sum \alpha_i e_i$, we can then associate the homology class $\hat{\alpha} = \sum \alpha_i \hat{e}_i$.

Suppose now that $\tau = (\tau_{ij})_{ij}$ is the matrix of coefficients of the Weil–Petersson Poisson structure on $\tilde{\mathcal{T}}(S_\gamma)$ for the induced ideal triangulation $\lambda_\gamma = \{\mu_1, \dots, \mu_n\}$. We suppose that, for $i = 1, \dots, n$, μ_i corresponds to the homotopy class of k_{ij} segments from λ_j for $j = 1, \dots, n$. We let $\mathbf{k}_i = (k_{i1}, \dots, k_{in}) \in \mathbb{Z}^n$ and identify σ with a bilinear form on $\mathbb{Z}^n \cong \mathcal{H}(\lambda, \mathbb{Z})$. The following proposition relates the entries of τ and σ .

Proposition 9 *With the notation above, the coefficients of the Weil–Petersson Poisson structure on $\mathcal{T}(S)$ and $\mathcal{T}(S_\gamma)$ for λ and λ_γ , respectively, are related via the following formula: for $i, j = 1 \dots n$,*

$$\tau_{ij} = \sigma(\mathbf{k}_i, \mathbf{k}_j) = \sum_{s,t} k_{is} k_{jt} \sigma_{st}.$$

Proof Let G be the graph dual to λ in S , G_γ be the graph dual to λ_γ and G'_γ the graph dual to $\lambda \setminus \gamma$ in S_γ . We recall that $\lambda \setminus \gamma$ denotes the family of arcs obtained from the edges of λ intersected with S_γ , where we *do not* consider the arcs up to homotopy. It decomposes S_γ into triangles and bigons, hence the vertices of G'_γ are either bivalent or trivalent. We will call a *maximal chain* in G'_γ any chain of edges connected via bi-valent vertices and with endpoints at trivalent vertices. In particular, edges connecting trivalent vertices *are* maximal chains. Following Remark 5, the maximal chains of G'_γ are in one-to-one correspondence with the edges of G_γ , and this defines a natural homeomorphism $G'_\gamma \cong G_\gamma$. In addition, the identification of ideal triangles for λ and λ_γ gives an identification of G and G'_γ in a neighborhood of each of their trivalent vertices.

We also consider the oriented graphs \hat{G} and \hat{G}_γ (see beginning of section), as well as the oriented graph \hat{G}'_γ obtained from G'_γ by keeping the same set of trivalent vertices and replacing each maximal chain by two such chains connected to the same (6–valent) endpoints, endowed with opposite orientations. This graph is naturally homeomorphic to \hat{G}_γ . As described above, \hat{G} thickens into \hat{S} , and both \hat{G}_γ and \hat{G}'_γ thicken into \hat{S}_γ . We have covering maps $p: \hat{S} \rightarrow S$ and $p_\gamma: \hat{S}_\gamma \rightarrow S_\gamma$, which restrict to the corresponding graphs.

Recall that, by definition, $S_\gamma = S \setminus \gamma \subset S$. Similarly one can identify \hat{S}_γ with $\hat{S} \setminus \hat{\gamma}$, where $\hat{\gamma} = p^{-1}(\gamma)$. Indeed, let $U \subset S_\gamma \subset S$ be a union of small discs around

the trivalent vertices of G'_γ where it is identified with G . By construction, we have $p_\gamma^{-1}(U) \cong p^{-1}(U) \subset \widehat{S}$. Outside of U both coverings are trivial, so we also have a natural identification $p_\gamma^{-1}(S_\gamma \setminus U) \cong p^{-1}(S_\gamma \setminus U) \subset \widehat{S}$. Hence we obtain that $\widehat{S}_\gamma \cong p^{-1}(S_\gamma) \subset \widehat{S}$. Note, in addition, that γ can be chosen so that it does not pass through any of the ramification points of p (that is, the vertices of G). With this assumption, $p^{-1}(S_\gamma) = \widehat{S} \setminus \widehat{\gamma}$, where $p^{-1}(\gamma) = \widehat{\gamma}$ consists of two non-intersecting multicurves and we can identify \widehat{S}_γ with $\widehat{S} \setminus \widehat{\gamma}$ sitting in \widehat{S} . Accordingly, p_γ is identified with the restriction of p to $S \setminus \gamma$.

The inclusion $\widehat{\iota}: \widehat{S}_\gamma \hookrightarrow \widehat{S}$ induces a map $\widehat{\iota}_*: H_1(\widehat{S}_\gamma) \rightarrow H_1(\widehat{S})$ at the level of homology. By construction, if we denote by η_γ the covering involution associated to p_γ , we have that $\widehat{\iota}_* \circ \eta_{\gamma*} = \eta_* \circ \widehat{\iota}_*$.

On the other hand, there is a natural map $\pi: G'_\gamma \rightarrow G$ defined by sending each edge of G'_γ onto the edge of G dual to the same edge of the triangulation λ , which lifts to a η -invariant map $\widehat{\pi}: \widehat{G}'_\gamma \rightarrow \widehat{G}$. One can then consider retractions $\widehat{r}: \widehat{S} \rightarrow \widehat{G}$ and $\widehat{r}_\gamma: \widehat{S}_\gamma \rightarrow \widehat{G}'_\gamma$ such that $\widehat{\pi} \circ \widehat{r}_\gamma$ is equal to $\widehat{r} \circ \widehat{\iota}$, giving the following commutative diagram:

$$(2) \quad \begin{array}{ccc} H_1(\widehat{S}_\gamma) & \xrightarrow{\widehat{\iota}_*} & H_1(\widehat{S}) \\ \downarrow \widehat{r}_{\gamma*} & & \downarrow \widehat{r}_* \\ H_1(\widehat{G}'_\gamma) & \xrightarrow{\widehat{\pi}_*} & H_1(\widehat{G}) \end{array}$$

Let e_i be the generator of $\mathcal{H}(\lambda, \mathbb{Z})$ assigning weight 1 to λ_i and 0 to the other edges of λ , and f_i be the generator of $\mathcal{H}(\lambda_\gamma, \mathbb{Z})$ assigning weight 1 to μ_i and 0 to the other edges of λ_γ . Following Lemma 8, we associate to e_i the homology class $\widehat{e}_i \in H_1(\widehat{S})$ corresponding to the lift in \widehat{G} of the edge of G dual to λ_i . To f_i on the other hand, we associate $\widehat{f}_i \in H_1(\widehat{S}_\gamma)$, which corresponds to the lift in \widehat{G}'_γ of the maximal chain of G'_γ dual to μ_i . By construction, this chain consists of the lift of k_{is} edges dual to λ_s for $s = 1, \dots, n$. Hence, as curves, $\widehat{\pi}(\widehat{f}_i)$ covers k_{is} times each \widehat{e}_s . Homologically, using the commutativity of diagram (2), we obtain:

$$(3) \quad \widehat{\iota}_*(\widehat{f}_i) = \widehat{r}_*^{-1} \circ \widehat{\pi}_* \circ \widehat{r}_{\gamma*}(\widehat{f}_i) = \sum_{s=1}^n k_{is} \widehat{e}_s \text{ for } i = 1, \dots, n$$

Hence we have the following equalities:

$$\begin{aligned} \tau_{ij} &= \widehat{f}_i \cdot \widehat{f}_j = \widehat{\iota}_*(\widehat{f}_i) \cdot \widehat{\iota}_*(\widehat{f}_j) \\ &= \sum_s k_{is} \widehat{e}_s \cdot \sum_t k_{jt} \widehat{e}_t = \sum_{s,t} k_{is} k_{jt} \widehat{e}_s \cdot \widehat{e}_t = \sum_{s,t} k_{is} k_{jt} \sigma_{st}. \quad \square \end{aligned}$$

In terms of shear coordinates and with the notation of Section 2.2, Proposition 9 implies that the Poisson brackets associated to the Weil–Petersson structure on $\mathcal{T}(S)$ and $\mathcal{T}(S_\gamma)$ are related via the formula

$$\{\log y_i, \log y_j\}_{\mathcal{T}(S_\gamma)} = \{\log(x_1^{k_{i1}} \cdots x_n^{k_{in}}), \log(x_1^{k_{j1}} \cdots x_n^{k_{jn}})\}_{\mathcal{T}(S)}$$

which is consistent with Proposition 6 and Masur [15] and Wolpert [23].

3 Pinching along curves: quantum aspect

3.1 The Chekhov–Fock algebra

Let $\lambda = \{\lambda_1, \dots, \lambda_n\}$ be an ideal triangulation of S and fix a non-zero complex number $q \in \mathbb{C}^*$. Following [14], we define the *Chekhov–Fock algebra* $\mathcal{T}_\lambda^q(S)$ of S associated to λ to be the algebra over \mathbb{C} with generators $X_i^{\pm 1}$ associated to the edges λ_i of λ and subject to the relations

$$X_i X_j = q^{2\sigma_{ij}} X_j X_i$$

for every i, j , where the $\sigma_{ij} \in \{-2, -1, 0, 1, 2\}$ are the coefficients of the Weil–Petersson Poisson structure on $\widetilde{\mathcal{T}}(S)$ in the shear coordinates associated to λ . We will sometimes use the notation $\mathcal{T}_\lambda^q(S) = \mathbb{C}[X_1, \dots, X_n]_\lambda^q$ to specify the generators of the algebra.

If A and B are two monomials in the variables X_1, \dots, X_n , then they satisfy a relation of the form $AB = q^{2\alpha} BA$ for some integer α , and we will use the notation $\sigma(A, B) = \alpha$. This coefficient is independent of the order of the generators inside each monomial.

For $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n$, if A is a monomial consisting of k_i times the generator X_i for $i=1, \dots, n$, in any given order, we define the following element in $\mathcal{T}_\lambda^q(S)$:

$$[A] = X_{\mathbf{k}} = q^{-\sum_{i < j} k_i k_j \sigma_{ij}} X_1^{k_1} \cdots X_n^{k_n}.$$

This is known as the *Weyl quantum ordering*. These monomials satisfy the following relations:

$$X_{\mathbf{k}} X_{\mathbf{l}} = q^{\sigma(\mathbf{k}, \mathbf{l})} X_{\mathbf{k} + \mathbf{l}} = q^{2\sigma(\mathbf{k}, \mathbf{l})} X_{\mathbf{l}} X_{\mathbf{k}},$$

where we once again identify σ with a bilinear form on \mathbb{Z}^n . The different notation for σ coincides in the sense that $\sigma(X_{\mathbf{k}}, X_{\mathbf{l}}) = \sigma(\mathbf{k}, \mathbf{l})$.

In particular, if α is a path between two vertices in the dual graph G of λ (which does not backtrack), we can identify it with the element $\alpha = (\alpha_1, \dots, \alpha_n)$ of $\mathcal{H}(\lambda, \mathbb{Z})$, where α_i is the number of times the path α passes through the edge of G dual to λ_i .

Then we associate to α the monomial X_α defined as above. If β is another path in G , we have

$$X_\alpha X_\beta = q^{2\hat{\alpha} \cdot \hat{\beta}} X_\beta X_\alpha$$

by Lemma 8, where $\hat{\alpha}$ and $\hat{\beta}$ are the associated elements in $H_1(\hat{S})$. Of particular interest will be the element X_γ associated to a simple closed curve γ in S , which we identify with its retraction to a cycle in G .

If Σ is another surface with ideal triangulation μ , a homomorphism between $\mathcal{T}_\lambda^q(S)$ and $\mathcal{T}_\mu^q(\Sigma)$ does not in general preserve the quantum ordering. However, we have the following elementary lemma, which will be useful later on.

Lemma 10 *Let A_1, \dots, A_s be monomials in $\mathcal{T}_\lambda^q(S)$, B_1, \dots, B_s be monomials in $\mathcal{T}_\mu^q(\Sigma)$. If $\Psi: \mathcal{T}_\lambda^q(S) \rightarrow \mathcal{T}_\mu^q(\Sigma)$ is an algebra homomorphism such that $\Psi([A_i]) = [B_i]$ for all i , then $\Psi([A_1 \cdots A_s]) = [B_1 \cdots B_s]$.*

3.2 A homomorphism between Chekhov–Fock algebras

As a direct consequence of Propositions 6 and 9, we construct a natural homomorphism between the Chekhov–Fock algebras associated to S and S_γ . We recall that, by Lemma 4, λ induces an ideal triangulation $\lambda_\gamma = \{\mu_1, \dots, \mu_n\}$ of S_γ , where μ_i is the homotopy class in S_γ of k_{ij} segments from λ_j , for $j = 1, \dots, n$. We let $\mathbf{k}_i = (k_{i1}, \dots, k_{in})$ for $i = 1, \dots, n$.

Proposition 11 *The map*

$$\Theta_{\gamma, \lambda}^q: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) = \mathbb{C}[Y_1, \dots, Y_n]_{\lambda_\gamma}^q \longrightarrow \mathbb{C}[X_1, \dots, X_n]_{\lambda}^q = \mathcal{T}_\lambda^q(S)$$

defined on the generators by $\Theta_{\gamma, \lambda}^q(Y_i) = X_{\mathbf{k}_i}$ extends to an algebra homomorphism.

Proof We check it on the generators of $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$:

$$\Theta_{\gamma, \lambda}^q(Y_i Y_j) = \Theta_{\gamma, \lambda}^q(Y_i) \Theta_{\gamma, \lambda}^q(Y_j) = X_{\mathbf{k}_i} X_{\mathbf{k}_j} = q^{2\sigma(\mathbf{k}_i, \mathbf{k}_j)} X_{\mathbf{k}_j} X_{\mathbf{k}_i} = q^{2\tau_{ij}} \Theta_{\gamma, \lambda}^q(Y_j Y_i),$$

where the last equality is given by Proposition 9. □

The following lemma states that, if we pinch the curves constituting γ in different orders, the resulting homomorphisms given by Proposition 11 are the same.

Lemma 12 *Consider any sequence of integers i_1, \dots, i_k such that $\{i_1, \dots, i_k\} = \{1, \dots, k\}$ and let $\gamma^l = \bigcup_{j=1}^l \gamma_{i_j}$ for $l \leq k$. Then*

$$\Theta_{\gamma, \lambda}^q = \Theta_{\gamma_{i_k}, \lambda_{\gamma^{k-1}}}^q \circ \Theta_{\gamma_{i_{k-1}}, \lambda_{\gamma^{k-2}}}^q \circ \cdots \circ \Theta_{\gamma_{i_2}, \lambda_{\gamma^1}}^q \circ \Theta_{\gamma_{i_1}, \lambda}^q.$$

Proof The definition of the edges μ_i of λ_γ as homotopy classes of segments from $\lambda_1, \dots, \lambda_n$ does not depend on the order in which the curves $\gamma_1, \dots, \gamma_k$ are pinched. Hence the generators Y_i of $\mathcal{T}_\lambda^q(S_\gamma)$ are sent to the same monomials by the two maps, *up to ordering*. Since all the maps considered send generators to quantum ordered monomials, Lemma 10 implies that the quantum orders are respected on each side and hence the maps coincide. \square

Remark 13 For future reference, we want to interpret $\Theta_{\gamma,\lambda}^q$ in terms of dual graphs. As in the proof of Proposition 9, we let G be the graph dual to λ in S and G'_γ be the graph dual to $\lambda \setminus \gamma$ in S_γ . There is a natural map $\pi: G'_\gamma \rightarrow G$ defined by sending each edge of G'_γ onto the edge of G dual to the same edge of λ . If α is any path between trivalent vertices in G'_γ , we denote by $Y_\alpha \in \mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$ the quantum ordered product of generators associated to the edges crossed by α (after making the identification $G'_\gamma \cong G_\gamma$ with the graph dual to λ_γ), and we define similarly $X_\beta \in \mathcal{T}_\lambda^q(S)$ for β any path between vertices in G . Then, by definition of $\Theta_{\gamma,\lambda}^q$, we have

$$\Theta_{\gamma,\lambda}^q(Y_\alpha) = X_{\pi(\alpha)}.$$

4 Application to the representation theory of Chekhov–Fock algebras

4.1 Representations of the Chekhov–Fock algebras

The irreducible finite-dimensional representations of $\mathcal{T}_\lambda^q(S)$ for q a root of unity have been studied in detail in [7]. We will recall the main results that will be needed for our purpose.

An important step is to describe the center of these algebras. For $i = 1, \dots, s$, consider the element

$$P_i = X_{\mathbf{p}_i} = [X_1^{p_{i1}} X_2^{p_{i2}} \dots X_n^{p_{in}}]$$

of $\mathcal{T}_\lambda^q(S)$ associated to the puncture v_i of S , where $p_{ij} \in \{0, 1, 2\}$ is the number of ends of the edge λ_j that converge to v_i and $\mathbf{p}_i = (p_{i1}, \dots, p_{in})$. Namely, P_i is the quantum ordered product of generators associated to the edges ending at v_i . If α_i is a small loop around v_i and we identify it with its retraction to a cycle in the dual graph G , we also have $P_i = X_{\alpha_i}$, with the notation introduced previously.

In addition, let $\mathbf{h} = (1, \dots, 1)$ and $H = X_{\mathbf{h}} = [X_1 \cdots X_n]$. The following proposition describes the center of the Chekhov–Fock algebras for certain non-generic values of the parameter q .

Proposition 14 (Bonahon–Liu) *If q^2 is a primitive N^{th} root of unity with N odd, the center \mathcal{Z}_λ^q of $\mathcal{T}_\lambda^q(S)$ is generated by the elements X_i^N for $i = 1, \dots, n$, P_j for $j = 1, \dots, s$ and H .*

A complete classification of the finite-dimensional irreducible representations of $\mathcal{T}_\lambda^q(S)$ when q^2 is a root of unity was obtained in [7]. A particular case of this classification can be summarized by the following theorem. Note the further restriction to a condition on q instead of q^2 .

Theorem 15 (Bonahon–Liu) *Suppose that q is a primitive N^{th} root of unity with N odd. Every irreducible finite-dimensional representation $\rho: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ has dimension N^{3g+s-3} and is determined completely by its restriction to the center \mathcal{Z}_λ^q of $\mathcal{T}_\lambda^q(S)$.*

In particular, given $p_j \in \{0, \dots, N - 1\}$ integers labelling the punctures v_j of S and $m \in \mathcal{T}(S)$ with shear parameters (x_1, \dots, x_n) associated to λ , there is a finite-dimensional irreducible representation $\rho: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ such that:

- $\rho(X_i^N) = x_i \text{Id}_V$ for $i = 1, \dots, n$.
- $\rho(P_j) = q^{p_j} \text{Id}_V$ for $j = 1, \dots, s$.

These conditions determine ρ uniquely up to isomorphism.

Proof The first part is given by [7, Theorem 20]. The second part is a specialization of [7, Theorem 21] to the case where the x_i are positive real numbers and $x^{p_{j1}} \dots x^{p_{jn}} = 1$ for $j = 1, \dots, s$, corresponding to the shear parameters of a complete hyperbolic metric on S . In this case $\rho(H)$ is completely determined by the fact that $\rho(H^N) = \text{Id}_V$ and $H^2 = P_1 \dots P_s$. □

We call $\mathbf{p} = (p_1, \dots, p_s)$ the *weights* of the representation ρ associated to the s -tuple of punctures (v_1, \dots, v_s) . Depending on the context we will use the notation ρ_m or $\rho_m^{\mathbf{p}}$ to emphasize the dependence of an irreducible representation on its associated weights and metric.

Remark 16 Theorem 15 generalizes directly to the case of a disconnected surface $S = S_1 \sqcup S_2$, except for the statement about the dimensions. More precisely, if $\lambda = \lambda_1 \cup \lambda_2$ is a triangulation of S , then there is a natural isomorphism

$$\mathcal{T}_\lambda^q(S) \cong \mathcal{T}_{\lambda_1}^q(S_1) \otimes \mathcal{T}_{\lambda_2}^q(S_2)$$

since the generators associated to λ_1 and λ_2 commute with each other. Hence an irreducible representation of $\mathcal{T}_\lambda^q(S)$ is a tensor product of irreducible representations

$\rho_1 \otimes \rho_2$ of each factor and has dimension $N^{3g_1+s_1-3} \times N^{3g_2+s_2-3} = N^{3g+s-6}$. In particular, given a multicurve $\gamma = \gamma_1 \cup \dots \cup \gamma_k$, the dimension of an irreducible representation of $\mathcal{T}_\lambda^q(S)$ is N^k times the dimension of an irreducible representation of $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$, regardless of the connectivity of S and S_γ .

4.2 Convergence of representations in the augmented Teichmüller space

We suppose once again that q is a primitive N^{th} root of unity with N odd. Theorem 15 implies that, to $\mathbf{p} = (p_1, \dots, p_s) \in \{0, \dots, N - 1\}^s$ and a continuous family of hyperbolic metrics $m_t \in \mathcal{T}(S)$, $t \in (0, 1]$, one can associate a continuous family of irreducible representations $\rho_t: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ as follows: let $x_1(t), \dots, x_n(t)$ be the shearing parameters associated to m_t for the triangulation λ , and let ρ_1 be an irreducible representation classified by m_1 and weights p_1, \dots, p_n . Then, by Theorem 15, there are elements A_1, \dots, A_n of $\text{End}(V)$ such that $A_i^N = \text{Id}_V$ and $\rho_1(X_i) = \sqrt[N]{x_i(1)} A_i$, for $i = 1, \dots, n$. One can then construct a family of representations ρ_t , defined on the generators by $\rho_t(X_i) = \sqrt[N]{x_i(t)} A_i$. In this way, we obtain a family of representations ρ_t classified by m_t , $t \in (0, 1]$, and weights \mathbf{p} . It is continuous in the sense that, for any element $X \in \mathcal{T}_\lambda^q(S)$, $\rho_t(X)$ is a continuous family in $\text{End}(V)$.

By composing with the homomorphism $\Theta_{\gamma,\lambda}^q$ from Proposition 11, any representation ρ of $\mathcal{T}_\lambda^q(S)$ gives a representation

$$\rho \circ \Theta_{\gamma,\lambda}^q \text{ of } \mathcal{T}_{\lambda_\gamma}^q(S_\gamma).$$

If ρ_m is an irreducible representation classified by $m \in \mathcal{T}(S)$ and weights \mathbf{p} , then, by the dimension count done in Remark 16, $\rho_m \circ \Theta_{\gamma,\lambda}^q$ is a *reducible* representation. We would like to know how this representation decomposes into irreducible subrepresentations when m approaches $m_\gamma \in \mathcal{T}(S_\gamma)$.

We recall that the punctures of S_γ are v_1, \dots, v_s corresponding to the same punctures in S together with the new punctures $v'_1, v''_1, \dots, v'_k, v''_k$ corresponding to the removal of the curves $\gamma_1, \dots, \gamma_k$. Given $\mathbf{p} = (p_1, \dots, p_s)$ weights labelling the punctures of S , we say that $\mathbf{p}_\gamma \in \{0, \dots, N - 1\}^{s+2k}$ are *compatible weights* labelling $(v_1, \dots, v_s, v'_1, v''_1, \dots, v'_k, v''_k)$ if they are of the form

$$\mathbf{p}_\gamma = (p_1, \dots, p_s, i_1, i_1, \dots, i_k, i_k).$$

Theorem 17 *Let $m_t \in \mathcal{T}(S)$ be a continuous family of hyperbolic metrics such that m_t converges to $m_\gamma \in \mathcal{T}(S_\gamma)$ in $\overline{\mathcal{T}(S)}$ as $t \rightarrow 0$. Let $\rho_t: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ be a continuous family of irreducible representations classified by m_t and weights $\mathbf{p} \in \{0, \dots, N - 1\}^s$ labelling the punctures v_1, \dots, v_s of S .*

Then, as $t \rightarrow 0$, the representation

$$\rho_t \circ \Theta_{\gamma, \lambda}^q: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}(V)$$

approaches

$$\bigoplus_{\mathbf{p}_\gamma} \rho_\gamma^{\mathbf{p}_\gamma}: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}\left(\bigoplus_{\mathbf{p}_\gamma} V_{\mathbf{p}_\gamma}\right)$$

where the direct sum is over all possible compatible weights \mathbf{p}_γ on S_γ and

$$\rho_\gamma^{\mathbf{p}_\gamma}: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}(V_{\mathbf{p}_\gamma})$$

is an irreducible representation classified by the metric m_γ and the weights \mathbf{p}_γ .

Proof We first suppose that $k = 1$, that is, γ consists of a single curve.

Let P_1, \dots, P_s be the central elements of $\mathcal{T}_\lambda^q(S)$ associated to the punctures v_1, \dots, v_s of S and P'_1, \dots, P'_s be the central elements of $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$ associated to the same punctures in S_γ . We also have the two central elements P' and P'' associated to the two new punctures v' and v'' . Consider also the monomial $X_\gamma \in \mathcal{T}_\lambda^q(S)$ associated to the retraction of γ to a cycle in G . In practice, if γ crosses c_i times the edge λ_i of λ for $i = 1, \dots, n$, we have

$$X_\gamma = [X_1^{c_1} \dots X_n^{c_n}].$$

This element is the quantum analogue of the exponential graph length x_γ discussed at the end of Section 2.2.

Lemma 18 *The elements defined above satisfy:*

- (1) $\Theta_{\gamma, \lambda}^q(P'_i) = P_i$ for $i = 1, \dots, s$.
- (2) $\Theta_{\gamma, \lambda}^q(P') = \Theta_{\gamma, \lambda}^q(P'') = X_\gamma$.

Proof This is a consequence of the interpretation of $\Theta_{\gamma, \lambda}^q$ given in Remark 13.

For (1), let α_i be a small curve going around v_i once in $S_\gamma \subset S$. If we identify α_i with its retraction to a cycle in G'_γ , we have that $P'_i = Y_{\alpha_i}$. Since $\pi(\alpha_i)$ corresponds to the retraction of α_i onto G , we see that

$$\Theta_{\gamma, \lambda}^q(P'_i) = X_{\pi(\alpha_i)} = P_i.$$

For (2), let γ' be a curve parallel to γ such that γ' is homotopic to v' in S_γ . Then γ' retracts to a cycle in G'_γ and $P' = Y_{\gamma'}$. In addition $\pi(\gamma')$ corresponds to the retraction of γ' , and hence of γ , onto G , so

$$\Theta_{\gamma, \lambda}^q(P') = X_{\pi(\gamma')} = X_\gamma.$$

The same argument holds for P'' if one considers a curve γ'' parallel to γ and homotopic to v'' in S_γ . □

Lemma 19 *There exists $\Delta \in \mathcal{T}_\lambda^q(S)$ such that $X_\gamma \Delta = q^4 \Delta X_\gamma$.*

Proof By Lemma 8, it suffices to find a path δ in G such that $\hat{\gamma} \cdot \hat{\delta} = \pm 2$, where $\hat{\gamma}$ and $\hat{\delta}$ are (the classes in $H_1(\hat{S})$ of) oriented η -anti-invariant lifts of γ and δ to \hat{S} . Then $\Delta = X_\delta^{\pm 1}$ will satisfy the lemma, since $X_\gamma X_\delta = q^{2\hat{\gamma} \cdot \hat{\delta}} X_\delta X_\gamma$.

If γ is non-separating, and since γ is an essential simple closed curve, there exists δ another essential simple closed curve in S that intersects γ exactly once. In addition, one can choose representatives of γ and δ not passing through the ramification points of the covering $p: \hat{S} \rightarrow S$, that is, not passing through the vertices of G . Then, by construction, $\hat{\delta} = p^{-1}(\delta)$ is such that $\hat{\gamma} \cdot \hat{\delta} = \pm 2|\gamma \cap \delta| = \pm 2$.

If γ is separating then it divides the set of vertices of G into two non-empty subsets. Let δ be an arc in S with endpoints at vertices of G on each side of γ and intersecting γ exactly once. Then $\hat{\delta} = p^{-1}(\delta)$ is a closed curve in \hat{S} with a natural orientation and by construction $\hat{\gamma} \cdot \hat{\delta} = 2$. □

End of proof of Theorem 17 Since N and 4 are coprime, Lemma 19 implies that, under the action of $\rho_t(X_\gamma)$, V decomposes into eigenspaces V_i of dimension N^{3g+p-4} with associated eigenvalues $c(t)q^i$, where $i \in \{0, \dots, N-1\}$ and $c(t) \in \mathbb{C}^*$. In addition

$$\rho_t(X_\gamma^N) = \rho_t((X_1^N)^{c_1} \cdots (X_n^N)^{c_n}) = x_1^{c_1}(t) \cdots x_n^{c_n}(t) \text{Id}_V = x_\gamma(t) \text{Id}_V,$$

where $x_\gamma(t)$ is the exponential graph length of γ . Hence, after a shift by some N^{th} root of 1, we can consider that $c(t) = \sqrt[N]{x_\gamma(t)} \in \mathbb{R}_+$.

P' and P'' are central in $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$, so the eigenspaces of

$$\rho_t \circ \Theta_{\gamma,\lambda}^q(P') = \rho_t \circ \Theta_{\gamma,\lambda}^q(P'') = \rho_t(X_\gamma)$$

are invariant under the action of $\rho_t \circ \Theta_{\gamma,\lambda}^q$. In other words

$$\rho_t \circ \Theta_{\gamma,\lambda}^q = \bigoplus_i \rho_{t,\gamma}^i,$$

where $\rho_{t,\gamma}^i: \mathcal{T}_{\lambda_\gamma}^q(S_\gamma) \rightarrow \text{End}(V_i)$ is such that

$$\rho_{t,\gamma}^i(P') = \rho_{t,\gamma}^i(P'') = c(t)q^i \text{Id}_{V_i}.$$

For dimensional reasons these representations are irreducible by Theorem 15.

By Lemma 18, we have that $\rho_t \circ \Theta_{\gamma,\lambda}^q(P_j^\gamma) = \rho_t(P_j) = q^{p_j} \text{Id}_V$, hence

$$\rho_{t,\gamma}^i(P_j^\gamma) = q^{p_j} \text{Id}_{V_i},$$

and by definition of $\Theta_{\gamma,\lambda}^q$,

$$\rho_t \circ \Theta_{\gamma,\lambda}^q(Y_j^N) = \rho_t((X_1^N)^{k_1} \cdots (X_n^N)^{k_n}) = x_1^{k_{j1}}(t) \cdots x_n^{k_{jn}}(t) \text{Id}_V,$$

hence

$$\rho_{t,\gamma}^i(Y_j^N) = x_1^{k_{j1}}(t) \cdots x_n^{k_{jn}}(t) \text{Id}_{V_i}.$$

Then, by Proposition 6, as $t \rightarrow 0$, $x_1^{k_{j1}}(t) \cdots x_n^{k_{jn}}(t) \rightarrow y_j$, the shear parameters of m_γ , and by Lemma 7, $x_\gamma(t) \rightarrow 1$. This implies that, as $t \rightarrow 0$, $\rho_{t,\gamma}^i$ approaches the irreducible representation ρ_γ^i of $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$ that is classified by the weights p_1, \dots, p_s associated to the punctures v_1, \dots, v_s , the weight i associated to v' and v'' , and the hyperbolic metric $m_\gamma \in \mathcal{T}(S_\gamma)$.

Using Lemma 12, the case of a multicurve $\gamma = \{\gamma_1, \dots, \gamma_k\}$ follows by induction on k . □

5 Behavior under changes of coordinates: the quantum Teichmüller space

5.1 The quantum Teichmüller space

We want to apply the results of the preceding section to the representations of the quantum Teichmüller space $\mathcal{T}^q(S)$. Let us first recall its construction as given in [14]. If λ is an ideal triangulation of S , we denote by $\widehat{\mathcal{T}}_\lambda^q(S)$ the fraction division algebra of the Chekhov–Fock algebra $\mathcal{T}_\lambda^q(S)$. Chekhov and Fock constructed a family of isomorphisms

$$\Phi_{\lambda\lambda'}^q: \widehat{\mathcal{T}}_{\lambda'}^q(S) \rightarrow \widehat{\mathcal{T}}_\lambda^q(S)$$

called (*quantum*) *coordinate change isomorphisms*, defined for any two triangulations λ, λ' of S . In particular, if λ'' is another triangulation, they satisfy the composition relation $\Phi_{\lambda\lambda''}^q = \Phi_{\lambda\lambda'}^q \circ \Phi_{\lambda'\lambda''}^q$. The main example is given by the case when λ and λ' differ by a diagonal exchange in an embedded square in S as in Figure 5. Then $\Phi_{\lambda\lambda'}^q(X_n') = X_n$ if $n \neq 1, \dots, 5$, $\Phi_{\lambda\lambda'}^q(X_0') = X_0^{-1}$ and:

$$\begin{aligned} \Phi_{\lambda\lambda'}^q(X_1') &= (1 + qX_0)X_1, & \Phi_{\lambda\lambda'}^q(X_2') &= (1 + qX_0^{-1})^{-1}X_2, \\ \Phi_{\lambda\lambda'}^q(X_3') &= (1 + qX_0)X_3, & \Phi_{\lambda\lambda'}^q(X_4') &= (1 + qX_0^{-1})^{-1}X_4. \end{aligned}$$

We refer to [14] or [7] for similar formulas when some of the edges of the square are identified. One can then construct $\Phi_{\lambda\lambda'}^q$ for any triangulations λ, λ' , using the composition relation and the fact that one can get from any triangulation λ to another triangulation λ' by a succession of diagonal exchanges and re-indexings (see Penner [16] for a proof). The fact that the maps so-obtained do not depend on the choice of a sequence of triangulations from λ to λ' is one of the achievements of [10].




Figure 5

Using these isomorphisms we can construct the *quantum Teichmüller space* $\mathcal{T}^q(S)$ as the quotient

$$\mathcal{T}^q(S) = \bigsqcup_{\lambda} \widehat{\mathcal{T}}_{\lambda}^q(S) / \sim,$$

where the disjoint union is over all triangulations λ of S , and the equivalence relation \sim identifies $X' \in \widehat{\mathcal{T}}_{\lambda'}^q(S)$ with $\Phi_{\lambda\lambda'}^q(X') \in \widehat{\mathcal{T}}_{\lambda}^q(S)$.

5.2 Representations of the quantum Teichmüller space

A first attempt at defining a representation of $\mathcal{T}^q(S)$ would be to consider a family of representations ρ_{λ} of $\widehat{\mathcal{T}}_{\lambda}^q(S)$ for every triangulation λ of S such that $\rho_{\lambda} \circ \Phi_{\lambda\lambda'}^q = \rho_{\lambda'}$ for every λ, λ' . However, one can easily check that such representations cannot be finite-dimensional. On the other hand, when q^2 is an N^{th} root of unity, the Chekhov–Fock algebras $\mathcal{T}_{\lambda}^q(S)$ admit many finite-dimensional representations. Hence, for our purpose, a representation of the quantum Teichmüller space $\mathcal{T}^q(S)$ will be a family of representations of $\mathcal{T}_{\lambda}^q(S)$, for every triangulation λ of S , satisfying certain compatibility relations when changing triangulations.

More precisely, let $\rho_{\lambda}: \mathcal{T}_{\lambda}^q(S) \rightarrow \text{End}(V)$ be an algebra homomorphism satisfying the following condition: for every Laurent polynomial $X' \in \mathcal{T}_{\lambda'}^q(S)$, the rational fraction $\Phi_{\lambda\lambda'}^q(X') \in \widehat{\mathcal{T}}_{\lambda'}^q(S)$ can be written as

$$\Phi_{\lambda\lambda'}^q(X') = P_1 Q_1^{-1} = Q_2^{-1} P_2,$$

where $P_1, Q_1, P_2, Q_2 \in \mathcal{T}_\lambda^q(S)$ are Laurent polynomials for which $\rho_\lambda(Q_1)$ and $\rho_\lambda(Q_2)$ are invertible in $\text{End}(V)$. If ρ_λ satisfies such a condition, we say that the composition

$$\rho_\lambda \circ \Phi_{\lambda\lambda'}^q: \mathcal{T}_{\lambda'}^q(S) \rightarrow \text{End}(V)$$

makes sense and is defined naturally as

$$\rho_\lambda \circ \Phi_{\lambda\lambda'}^q(X') = \rho_\lambda(P_1)\rho_\lambda(Q_1)^{-1} = \rho_\lambda(Q_2)^{-1}\rho_\lambda(P_2) \in \text{End}(V).$$

One can check that this definition does not depend on the decomposition of $\Phi_{\lambda\lambda'}^q(X')$ as a quotient of polynomials, and that this indeed defines an algebra homomorphism.

Definition 20 A representation $\rho = \{\rho_\lambda\}_\lambda$ of the quantum Teichmüller space $\mathcal{T}^q(S)$ over the vector space V consists of the data of an algebra homomorphism $\rho_\lambda: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ for every triangulation λ such that, for every λ, λ' , the representation $\rho_\lambda \circ \Phi_{\lambda\lambda'}^q: \mathcal{T}_{\lambda'}^q(S) \rightarrow \text{End}(V)$ makes sense and is equal to $\rho_{\lambda'}$.

We will sometimes use the notation $\rho: \mathcal{T}^q(S) \rightarrow \text{End}(V)$ for such a representation, keeping in mind that ρ consists in fact of a family of homomorphisms:

$$\{\rho_\lambda: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)\}_\lambda.$$

Such representations were called representations of the polynomial core of $\mathcal{T}^q(S)$ in [7].

To prove that a family of representations $\{\rho_\lambda\}_\lambda$ of the Chekhov–Fock algebras is in fact a representation of $\mathcal{T}^q(S)$, one can use the following lemma (Lemma 25 in [7]).

Lemma 21 Let an algebra homomorphism $\rho_\lambda: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ be given for every ideal triangulation λ . Suppose that $\rho_\lambda \circ \Phi_{\lambda\lambda'}^q: \mathcal{T}_{\lambda'}^q(S) \rightarrow \text{End}(V)$ makes sense and is equal to $\rho_{\lambda'}$ whenever λ and λ' differ by a diagonal exchange or a re-indexing. Then $\rho = \{\rho_\lambda\}_\lambda$ is a representation of $\mathcal{T}^q(S)$.

Given an irreducible representation $\rho_\lambda: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ for some ideal triangulation λ , one can show that the composition $\rho_\lambda \circ \Phi_{\lambda\lambda'}^q: \mathcal{T}_{\lambda'}^q(S) \rightarrow \text{End}(V)$ makes sense for any other ideal triangulation λ' and defines an irreducible representation $\rho_{\lambda'}$ of $\mathcal{T}_{\lambda'}^q(S)$. Such a family $\rho = \{\rho_\lambda\}_\lambda$ is thus called an *irreducible representation* of $\mathcal{T}^q(S)$.

In addition, if ρ_λ is classified by the weights $\mathbf{p} = (p_1, \dots, p_s)$ and the metric $m \in \mathcal{T}(S)$ expressed in the shear coordinates for λ , then the representation $\rho_{\lambda'}$ is also classified by the same weights and the metric m expressed in the shear coordinates for λ' . In

this case we say that $\rho = \{\rho_\lambda\}_\lambda$ is the irreducible representation of $\mathcal{T}^q(S)$ classified by m and \mathbf{p} (see [7, Lemma 29 and Theorem 30]).

Finally, if $m_t \in \mathcal{T}(S)$ is a continuous family of hyperbolic metrics and $\rho_{t,\lambda}$ is a continuous family of representations of $\mathcal{T}_\lambda^q(S)$ classified by m_t and weights \mathbf{p} , we say that the representations $\rho_t = \{\rho_{t,\lambda}\}_\lambda$ obtained in this way form a *continuous family of representations of $\mathcal{T}^q(S)$* .

5.3 Main theorem

In this section we restrict once again to the case when q is an N^{th} root of unity with N odd. The next step is to study how the decomposition obtained in Theorem 17 is affected by changing the triangulation. Given two triangulations λ, λ' , we can consider the following diagram:

$$(4) \quad \begin{array}{ccc} \widehat{\mathcal{T}}_{\lambda'}^q(S) & \xrightarrow{\Phi_{\lambda\lambda'}^q} & \widehat{\mathcal{T}}_\lambda^q(S) \\ \Theta_{\gamma,\lambda'}^q \uparrow & & \uparrow \Theta_{\gamma,\lambda}^q \\ \widehat{\mathcal{T}}_{\lambda'_\gamma}^q(S_\gamma) & \xrightarrow{\Phi_{\lambda_\gamma\lambda'_\gamma}^q} & \widehat{\mathcal{T}}_{\lambda_\gamma}^q(S_\gamma) \end{array}$$

which is in general *non-commutative*.




Figure 6

We focus on the case when λ and λ' differ only by a diagonal exchange in a square Q as in Figure 5.

If the curve γ never crosses Q vertically or horizontally, that is, never crosses successively $\lambda_1, \lambda_0, \lambda_3$ or $\lambda_2, \lambda_0, \lambda_4$, the triangulations λ_γ and λ'_γ also differ by a diagonal exchange and can be identified outside of a square Q_γ (see Figure 6).




Figure 7

If γ does cross Q vertically or horizontally, the triangulations λ_γ and λ'_γ can be identified (see Figure 7). In this case we introduce two maps $\Psi^q_{\lambda\lambda',v}$ and $\Psi^q_{\lambda\lambda',h}$, defined for each case of identifications of the boundary of Q as follows: in each case $\Psi^q_{\lambda\lambda',v}(X'_i) = \Psi^q_{\lambda\lambda',h}(X'_i) = X_i$ for $i \geq 5$, $\Psi^q_{\lambda\lambda',v}(X'_0) = \Psi^q_{\lambda\lambda',h}(X'_0) = X_0^{-1}$ and:

- If Q is embedded,

$$\begin{aligned} \Psi^q_{\lambda\lambda',v}(X'_1) &= qX_0X_1, & \Psi^q_{\lambda\lambda',v}(X'_2) &= X_2, \\ \Psi^q_{\lambda\lambda',v}(X'_3) &= qX_0X_3, & \Psi^q_{\lambda\lambda',v}(X'_4) &= X_4, \end{aligned}$$

and

$$\begin{aligned} \Psi^q_{\lambda\lambda',h}(X'_1) &= X_1, & \Psi^q_{\lambda\lambda',h}(X'_2) &= q^{-1}X_0X_2, \\ \Psi^q_{\lambda\lambda',h}(X'_3) &= X_3, & \Psi^q_{\lambda\lambda',h}(X'_4) &= q^{-1}X_0X_4. \end{aligned}$$

- If $\lambda_1 = \lambda_3$ and $\lambda_2 \neq \lambda_4$,

$$\begin{aligned} \Psi_{\lambda\lambda',v}^q(X'_1) &= q^4 X_0^2 X_1, & \Psi_{\lambda\lambda',v}^q(X'_2) &= X_2, \\ \Psi_{\lambda\lambda',v}^q(X'_4) &= X_4, \end{aligned}$$

and

$$\begin{aligned} \Psi_{\lambda\lambda',h}^q(X'_1) &= X_1, & \Psi_{\lambda\lambda',h}^q(X'_2) &= q^{-1} X_0 X_2, \\ \Psi_{\lambda\lambda',h}^q(X'_4) &= q^{-1} X_0 X_4. \end{aligned}$$

- If $\lambda_1 = \lambda_2$ and $\lambda_3 \neq \lambda_4$,

$$\begin{aligned} \Psi_{\lambda\lambda',v}^q(X'_1) &= X_0 X_1, \\ \Psi_{\lambda\lambda',v}^q(X'_3) &= q X_0 X_3, & \Psi_{\lambda\lambda',v}^q(X'_4) &= X_4, \end{aligned}$$

and

$$\begin{aligned} \Psi_{\lambda\lambda',h}^q(X'_1) &= X_0 X_1, \\ \Psi_{\lambda\lambda',h}^q(X'_3) &= X_3, & \Psi_{\lambda\lambda',h}^q(X'_4) &= q^{-1} X_0 X_4. \end{aligned}$$

- If $\lambda_1 = \lambda_3$ and $\lambda_2 = \lambda_4$, that is, S is a once-punctured torus:

$$\begin{aligned} \Psi_{\lambda\lambda',v}^q(X'_1) &= q^4 X_0^2 X_1, & \Psi_{\lambda\lambda',h}^q(X'_1) &= X_1, \\ \Psi_{\lambda\lambda',v}^q(X'_2) &= X_2, & \Psi_{\lambda\lambda',h}^q(X'_2) &= q^{-4} X_0^2 X_2. \end{aligned}$$

The other cases are inverses of the ones above. Note that we exclude the case when $\lambda_1 = \lambda_2$ and $\lambda_3 = \lambda_4$ corresponding to a sphere with three holes since there are no essential simple closed curves on S in this case.

One can easily check that $\Psi_{\lambda\lambda',v}^q$ and $\Psi_{\lambda\lambda',h}^q$ are algebra homomorphisms.

Proposition 22 *If λ and λ' differ by a diagonal exchange in a square Q as in Figure 5 then one of the following is true:*

- (1) *The multicurve γ does not cross Q horizontally or vertically and*

$$\Phi_{\lambda\lambda'}^q \circ \Theta_{\gamma,\lambda'}^q = \Theta_{\gamma,\lambda}^q \circ \Phi_{\lambda,\lambda'}^q.$$

- (2) *γ crosses Q vertically at least once and*

$$\Psi_{\lambda\lambda',v}^q \circ \Theta_{\gamma,\lambda'}^q = \Theta_{\gamma,\lambda}^q.$$

- (3) *γ crosses Q horizontally at least once and*

$$\Psi_{\lambda\lambda',h}^q \circ \Theta_{\gamma,\lambda'}^q = \Theta_{\gamma,\lambda}^q.$$

Proof We use the notation of Figures 5, 6 and 7. The following lemma is a simple computation.

Lemma 23 Let $A_i, i = 1, 2, 3, 4$, monomials in $\mathcal{T}_\lambda^q(S)$, be the products of generators associated to the edges of λ converging to the corner c_i of Q , A_5 be the product of generators when one crosses Q vertically, and A_6 the product of generators when one crosses Q horizontally. Define similarly B_1, \dots, B_6 monomials in $\mathcal{T}_{\lambda'}^q(S)$ for the triangulation λ' . Then, for $i, j = 1, \dots, 4$

$$\Phi_{\lambda\lambda'}^q([B_i]) = \Psi_{\lambda\lambda',v}^q([B_i]) = \Psi_{\lambda\lambda',h}^q([B_i]) = [A_i]$$

and

$$\Psi_{\lambda\lambda',v}^q([B_5]) = [A_5] \quad \text{and} \quad \Psi_{\lambda\lambda',h}^q([B_6]) = [A_6].$$

Lemma 23 says that $\Phi_{\lambda\lambda'}^q, \Psi_{\lambda\lambda',v}^q$ and $\Psi_{\lambda\lambda',h}^q$ send the products of generators at a corner of Q for λ' to the respective products for λ , respecting the quantum orderings. In addition $\Psi_{\lambda\lambda',v}^q$ and $\Psi_{\lambda\lambda',h}^q$ respect the quantum ordered products of generators when crossing Q vertically and horizontally respectively.

We first consider the case where both Q and Q_γ are embedded in S and S_γ , respectively, and we look at the generators Y_1 and Y'_1 associated to the edges μ_1 and μ'_1 . We recall that, by Remark 5, the edges of λ_γ can be identified with rectangles in the decomposition of S_γ by $\lambda \setminus \gamma$, formed by maximal chains of bigons.

For (1), we notice that μ_1 corresponds to a rectangle for $\lambda \setminus \gamma$ that starts along λ_1 , and then crosses Q l_i times around the corner c_i for $i = 1, 2, 3, 4$. It also crosses λ_i k_i times for $i = 5, \dots, n$. The same is true for μ'_1 . We let $Z = X_5^{k_5} \dots X_n^{k_n}$ and $Z' = X_5^{k_5} \dots X_n^{k_n}$. Then

$$\Theta_{\gamma,\lambda}^q(Y_1) = [X_1 A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} Z] = q^\alpha X_1 [A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} Z],$$

where α is some integer. We note that $\sigma(X_1, A_i) = \sigma(X'_1, B_i)$ for $i = 1, 2, 3, 4$, hence

$$\Theta_{\gamma,\lambda}^q(Y'_1) = [X'_1 B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} Z'] = q^\alpha X'_1 [B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} Z']$$

for the same integer α . We also note that μ_0 corresponds to the edge λ_0 and hence $\Theta_{\gamma,\lambda}^q(Y_0) = X_0$. Using Lemma 23 together with Lemma 10 we obtain

$$\begin{aligned} \Phi_{\lambda\lambda'}^q \circ \Theta_{\gamma,\lambda'}^q(Y'_1) &= \Phi_{\lambda\lambda'}^q(q^\alpha X'_1 [B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} Z']) \\ &= q^\alpha (1 + qX_0) X_1 [A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} Z] \\ &= \Theta_{\gamma,\lambda}^q((1 + qY_0)Y_1) \\ &= \Theta_{\gamma,\lambda}^q \circ \Phi_{\lambda_\gamma\lambda'_\gamma}^q(Y'_1). \end{aligned}$$

A similar computation works for Y'_2, Y'_3 and Y'_4 . For $i > 4$, μ_i corresponds to a rectangle with neither side ending along Q but which may still cross it at the corners, and one shows in the same way that the equality holds for Y'_5, \dots, Y'_n . Hence (1) is true in the case of embedded squares.

For (2), μ_1 corresponds to a rectangle for $\lambda \setminus \gamma$ that starts along λ_1 , crosses the square Q l_i times around the corner c_i for $i = 1, 2, 3, 4$, and crosses Q vertically l_5 times. For $\lambda' \setminus \gamma$, it corresponds to a rectangle that starts along λ'_0 , crosses λ'_1 , then crosses Q in the same way. It also crosses λ_i (resp. λ'_i) k_i times, for $i = 5, \dots, n$. Then

$$\Theta_{\gamma,\lambda}^q(Y_1) = [X_1 A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} A_5^{l_5} Z] = q^\alpha X_1 [A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} A_5^{l_5} Z],$$

where α is some integer. We note that $\sigma(X_1, A_i) = \sigma(X'_0 X'_1, B_i)$ for $i = 1, 2, 3, 4, 5$, hence

$$\begin{aligned} \Theta_{\gamma,\lambda}^q(Y'_1) &= [X'_0 X'_1 B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} B_5^{l_5} Z'] = q^\alpha [X'_0 X'_1] [B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} B_5^{l_5} Z'] \\ &= q^{\alpha-1} X'_0 X'_1 [B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} B_5^{l_5} Z']. \end{aligned}$$

Using the definition of $\Psi_{\lambda\lambda',v}^q$ in case 1, together with Lemmas 23 and 10, we obtain

$$\begin{aligned} \Psi_{\lambda\lambda',v}^q \circ \Theta_{\gamma,\lambda}^q(Y'_1) &= \Psi_{\lambda\lambda',v}^q (q^{\alpha-1} X'_0 X'_1 [B_1^{l_1} B_2^{l_2} B_3^{l_3} B_4^{l_4} B_5^{l_5} Z']) \\ &= q^{\alpha-1} X_0^{-1} q X_0 X_1 [A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} A_5^{l_5} Z] \\ &= q^\alpha X_1 [A_1^{l_1} A_2^{l_2} A_3^{l_3} A_4^{l_4} A_5^{l_5} Z] \\ &= \Theta_{\gamma,\lambda}^q(Y_1). \end{aligned}$$

A similar argument works for the other generators Y'_2, \dots, Y'_n .

The case (3) is similar to (2), where μ_1 corresponds to a rectangle that does not cross Q vertically but crosses it horizontally l_6 times. The argument is the same replacing B_5, l_5 and $\Psi_{\lambda\lambda',v}^q$ with B_6, l_6 and $\Psi_{\lambda\lambda',h}^q$, respectively.

If some of the edges of Q or Q_γ are identified, the same method works using the corresponding formulae for $\Phi_{\lambda\lambda'}^q, \Psi_{\lambda\lambda',v}^q$ and $\Psi_{\lambda\lambda',h}^q$. One also needs to take into account that, in this case, an edge μ_i of λ_γ may correspond to a rectangle in $S \setminus \gamma$ with both ends along edges of Q . The formulae for $\Theta_{\gamma,\lambda}^q(Y_i)$ and $\Theta_{\gamma,\lambda}^q(Y'_i)$ have to be changed accordingly. □

The maps $\Psi_{\lambda\lambda',v}^q$ and $\Psi_{\lambda\lambda',h}^q$ can be interpreted as the “limit” of the coordinate change $\Phi_{\lambda\lambda'}^q$ when the length of γ approaches 0, depending on whether γ crosses Q vertically or horizontally. This is made clearer by the following lemma.

Lemma 24 *Let ρ_t be a continuous family of irreducible representations of $\mathcal{T}_\lambda^q(S)$ classified by a continuous family $m_t \in \mathcal{T}(S)$ such that, as $t \rightarrow 0$, m_t approaches $m_\gamma \in \mathcal{T}(S_\gamma)$ in $\overline{\mathcal{T}(S)}$. Then, if λ' differs from λ by a diagonal exchange as in Figure 5 and γ crosses Q vertically ($i = v$) or horizontally ($i = h$), we have*

$$\rho_{t,\lambda} \circ \Phi_{\lambda\lambda'}^q \sim \rho_{t,\lambda} \circ \Psi_{\lambda\lambda',i}^q \quad \text{as } t \rightarrow 0.$$

Proof We suppose that $(x_1(t), \dots, x_n(t))$ are the shear parameters associated to m_t for the triangulation λ . Then, as in Section 4.2, there are matrices A_1, \dots, A_n such that $\rho(X_i) = \sqrt[n]{x_i(t)} A_i$ for every t .

For simplicity, we consider the case when Q is embedded in S . The computations are similar in the non-embedded cases. If γ crosses Q vertically then $x_0(t) \rightarrow \infty$ as $t \rightarrow 0$ and, considering for example the generator X_1 , we have

$$\begin{aligned} \rho_{t,\lambda} \circ \Phi_{\lambda\lambda'}^q(X_1) &= (I + q \sqrt[n]{x_0(t)} A_0)^{-1} \sqrt[n]{x_1(t)} A_1 \\ &\sim q \sqrt[n]{x_0(t)} \sqrt[n]{x_1(t)} A_0 A_1 = \rho_{t,\lambda} \circ \Psi_{\lambda\lambda',v}^q(X_1) \end{aligned}$$

and similarly for the other generators.

If γ crosses Q horizontally then $x_0(t) \rightarrow 0$ as $t \rightarrow 0$ and, considering for example the generator X_2 , we have

$$\begin{aligned} \rho_{t,\lambda} \circ \Phi_{\lambda\lambda'}^q(X_2) &= (I + q(\sqrt[n]{x_0(t)} A_0)^{-1})^{-1} \sqrt[n]{x_2(t)} A_2 \\ &\sim q^{-1} \sqrt[n]{x_0(t)} \sqrt[n]{x_2(t)} A_0 A_2 = \rho_{t,\lambda} \circ \Psi_{\lambda\lambda',h}^q(X_2) \end{aligned}$$

and similarly for the other generators. □

Proposition 25 *Suppose that $\rho_{t,\lambda}: \mathcal{T}_\lambda^q(S) \rightarrow \text{End}(V)$ and $\rho_{t,\lambda'}: \mathcal{T}_{\lambda'}^q(S) \rightarrow \text{End}(V)$ are two continuous families of irreducible representations classified by the same weights and by a continuous family $m_t \in \mathcal{T}(S)$ such that, as $t \rightarrow 0$, m_t approaches $m_\gamma \in \mathcal{T}(S_\gamma)$ in $\overline{\mathcal{T}(S)}$. By Theorem 17 and using the same notation, we have*

$$\lim_{t \rightarrow 0} \rho_{t,\lambda} \circ \Theta_{\gamma,\lambda}^q = \bigoplus_{\mathbf{p}_\gamma} \rho_{\gamma,\lambda}^{\mathbf{p}_\gamma} \quad \text{and} \quad \lim_{t \rightarrow 0} \rho_{t,\lambda'} \circ \Theta_{\gamma,\lambda'}^q = \bigoplus_{\mathbf{p}_\gamma} \rho_{\gamma,\lambda'}^{\mathbf{p}_\gamma}.$$

Suppose in addition that, for all t , we have $\rho_{t,\lambda} \circ \Phi_{\lambda\lambda'}^q = \rho_{t,\lambda'}$. Then

$$\rho_{\gamma,\lambda'}^{\mathbf{p}_\gamma} = \rho_{\gamma,\lambda}^{\mathbf{p}_\gamma} \circ \Phi_{\lambda_\gamma,\lambda'_\gamma}^q \quad \text{for all compatible } \mathbf{p}_\gamma.$$

Proof We show the result for the case when $\gamma = \gamma_1$ is a simple closed curve. The general case follows by induction on k using Lemma 12.

By Lemma 21, it suffices to show the result for a diagonal exchange in a square Q as in Figure 5. We let P_γ be the quantum ordered product of generators associated to the edges of λ_γ ending at one of the new punctures of S_γ . If λ'_γ is different from λ_γ , we define similarly P'_γ . Following the proof of Theorem 17, we have that

$$\rho_{t,\lambda} \circ \Theta_{\gamma,\lambda}^q = \bigoplus_i \rho_{t,\lambda}^i,$$

where $\rho_{t,\lambda}^i$ are representations of $\mathcal{T}_{\lambda_\gamma}^q(S_\gamma)$ onto V_i , the eigenspace of $\rho_{t,\lambda} \circ \Theta_{\gamma,\lambda}^q(P_\gamma)$ with eigenvalue q^i . Then, by definition, $\rho_{\gamma,\lambda}^i$ is the limit as t approaches 0 of $\rho_{t,\lambda}^i$.

We have

$$\rho_{t,\lambda} \circ \Theta_{\gamma,\lambda}^q \circ \Phi_{\lambda_\gamma\lambda'_\gamma}^q \rightarrow \left(\bigoplus_i \rho_{\gamma,\lambda}^i\right) \circ \Phi_{\lambda_\gamma\lambda'_\gamma}^q$$

and by hypothesis

$$\rho_{t,\lambda} \circ \Phi_{\lambda\lambda'}^q \circ \Theta_{\gamma,\lambda'}^q = \rho_{t,\lambda'} \circ \Theta_{\gamma,\lambda'}^q \rightarrow \bigoplus_i \rho_{\gamma,\lambda'}^i.$$

If γ does not cross Q vertically or horizontally, then $\Phi_{\lambda\lambda'}^q \circ \Theta_{\gamma,\lambda'}^q = \Theta_{\gamma,\lambda}^q \circ \Phi_{\lambda_\gamma\lambda'_\gamma}^q$ by Proposition 22. Composing on both sides of this equation by $\rho_{t,\lambda}$ on the left and taking the limit as t approaches 0, we get $\bigoplus_i \rho_{\gamma,\lambda'}^i = \left(\bigoplus_i \rho_{\gamma,\lambda}^i\right) \circ \Phi_{\lambda_\gamma\lambda'_\gamma}^q$. Note that $\Phi_{\lambda_\gamma\lambda'_\gamma}^q(P'_\gamma) = P_\gamma$ so it sends the eigenspaces of P'_γ onto those of P_γ with the same eigenvalues. Hence $\rho_{\gamma,\lambda'}^i = \rho_{\gamma,\lambda}^i \circ \Phi_{\lambda_\gamma\lambda'_\gamma}^q$ for every i .

If γ crosses Q , say vertically, then $\Phi_{\lambda_\gamma\lambda'_\gamma}^q$ is the identity. By Proposition 22,

$$\Psi_{\lambda\lambda',v}^q \circ \Theta_{\gamma,\lambda'}^q = \Theta_{\gamma,\lambda}^q$$

and, by Lemma 24, $\rho_{t,\lambda} \circ \Phi_{\lambda\lambda'}^q \sim \rho_{t,\lambda} \circ \Psi_{\lambda\lambda',v}^q$. Composing on both sides of this equivalence by $\Theta_{\gamma,\lambda'}^q$ on the right and taking the limit as t approaches 0 we get that

$$\bigoplus_i \rho_{\gamma,\lambda'}^i = \bigoplus_i \rho_{\gamma,\lambda}^i.$$

The decomposition $V = \bigoplus V_i$ is the same on each side, given by the eigenspaces of $P_\gamma = P'_\gamma$, hence we have $\rho_{\gamma,\lambda'}^i = \rho_{\gamma,\lambda}^i$ for all i in this case. □

In the notation of Proposition 25, we see that $\rho_{\gamma,\lambda}^{P_\gamma}$ may differ from $\rho_{\gamma,\lambda'}^{P'_\gamma}$ only if λ_γ is different from λ'_γ . Hence we can rename these representations $\rho_{\gamma,\lambda_\gamma}^{P_\gamma}$.

If Ω is a subset of the set $\Lambda(S)$ of ideal triangulations of S and if $\{\rho_\mu\}_{\mu \in \Omega}$ is a family of compatible representations of the Chekhov–Fock algebras $\mathcal{T}_\mu^q(S)$, we can extend this family to a representation $\rho = \{\rho_\lambda\}_{\lambda \in \Lambda(S)}$ of $\mathcal{T}^q(S)$ by setting $\rho_\lambda = \rho_\mu \circ \Phi_{\mu\lambda}^q$ for any $\lambda \in \Lambda(S)$, for some fixed $\mu \in \Omega$. The composition rule for the coordinate

change isomorphisms implies that this definition does not depend on μ . Of particular interest here is the subset $\Lambda_\gamma = \{\lambda_\gamma \text{ induced by } \lambda \mid \lambda \in \Lambda(S)\}$ of the set $\Lambda(S_\gamma)$ of ideal triangulations of S_γ .

By Proposition 25, if $\rho_t = \{\rho_{t,\lambda}\}_\lambda$ is such a continuous family of representations of $\mathcal{T}^q(S)$, the limiting irreducible factors $\rho_{\gamma,\lambda_\gamma}^{\mathbf{p}_\gamma}$ for each $\rho_{t,\lambda}$, as given by Theorem 17, satisfy the compatibility relations

$$\rho_{\gamma,\lambda'_\gamma}^{\mathbf{p}_\gamma} = \rho_{\gamma,\lambda_\gamma}^{\mathbf{p}_\gamma} \circ \Phi_{\lambda_\gamma,\lambda'_\gamma}^q.$$

Hence, taken together, they can be extended to form irreducible representations of $\mathcal{T}^q(S_\gamma)$, proving the following theorem.

Theorem 26 *Let $\rho_t = \{\rho_{t,\lambda}\}_\lambda$ be a continuous family of irreducible representations of $\mathcal{T}^q(S)$ classified by weights $\mathbf{p} \in \{0, \dots, N-1\}^s$ and a continuous family of metrics $m_t \in \mathcal{T}(S)$ such that m_t approaches $m_\gamma \in \mathcal{T}(S_\gamma)$ in $\overline{\mathcal{T}(S)}$ as $t \rightarrow 0$. For each triangulation λ , we let the limit*

$$\lim_{t \rightarrow 0} \rho_{t,\lambda} \circ \Theta_{\gamma,\lambda}^q = \bigoplus_{\mathbf{p}_\gamma} \rho_{\gamma,\lambda_\gamma}^{\mathbf{p}_\gamma}$$

be given as in Theorem 17.

Then, for compatible weights \mathbf{p}_γ on S_γ , the family of representations $\{\rho_{\gamma,\lambda_\gamma}^{\mathbf{p}_\gamma}\}_{\lambda_\gamma \in \Lambda_\gamma}$ extends to an irreducible representation $\rho_{\gamma}^{\mathbf{p}_\gamma}$ of $\mathcal{T}^q(S_\gamma)$ classified by the metric m_γ and the weights \mathbf{p}_γ .

References

- [1] **W Abikoff**, *Augmented Teichmüller spaces*, Bull. Amer. Math. Soc. 82 (1976) 333–334 MR0432919
- [2] **W Abikoff**, *Degenerating families of Riemann surfaces*, Ann. of Math. 105 (1977) 29–44 MR0442293
- [3] **H Bai**, *A uniqueness property for the quantization of Teichmüller spaces*, Geom. Dedicata 128 (2007) 1–16 MR2350143
- [4] **H Bai, F Bonahon, X Liu**, *Local representations of the quantum Teichmüller space* arXiv:0707.2151
- [5] **L Bers**, *Spaces of degenerating Riemann surfaces*, from: “Discontinuous groups and Riemann surfaces”, (L Greenberg, editor), Ann. of Math. Studies 79, Princeton Univ. Press (1974) 43–55 MR0361051
- [6] **F Bonahon**, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston’s symplectic form*, Ann. Fac. Sci. Toulouse Math. 5 (1996) 233–297 MR1413855

- [7] **F Bonahon, X Liu**, *Representations of the quantum Teichmüller space and invariants of surface diffeomorphisms*, Geom. Topol. 11 (2007) 889–937 MR2326938
- [8] **L D Faddeev, R M Kashaev**, *Quantum dilogarithm*, Modern Phys. Lett. A 9 (1994) 427–434 MR1264393
- [9] **V V Fock**, *Dual Teichmüller space* arXiv:dg-ga/9702018
- [10] **V V Fock, L O Chekhov**, *Quantum Teichmüller spaces*, Teoret. Mat. Fiz. 120 (1999) 511–528 MR1737362 In Russian; translated in Theoret. Math. Phys. 120 (1999) 1245–1259
- [11] **V V Fock, A B Goncharov**, *The quantum dilogarithm and representations of quantum cluster varieties*, Invent. Math. 175 (2009) 223–286 MR2470108
- [12] **R M Kashaev**, *Quantization of Teichmüller spaces and the quantum dilogarithm*, Lett. Math. Phys. 43 (1998) 105–115 MR1607296
- [13] **R M Kashaev**, *On the spectrum of Dehn twists in quantum Teichmüller theory*, from: “Physics and combinatorics”, (A N Kirillov, N Liskova, editors), World Sci. Publ. (2001) 63–81 MR1872252
- [14] **X Liu**, *The quantum Teichmüller space as a noncommutative algebraic object*, J. Knot Theory Ramifications 18 (2009) 705–726 MR2527682
- [15] **H Masur**, *Extension of the Weil–Petersson metric to the boundary of Teichmüller space*, Duke Math. J. 43 (1976) 623–635 MR0417456
- [16] **R C Penner**, *The decorated Teichmüller space of punctured surfaces*, Comm. Math. Phys. 113 (1987) 299–339 MR919235
- [17] **J Teschner**, *On the relation between quantum Liouville theory and the quantized Teichmüller spaces*, from: “Proceedings of 6th International Workshop on Conformal Field Theory and Integrable Models”, Internat. J. Modern Phys. A 19 (2004) 459–477 MR2087126
- [18] **J Teschner**, *An analog of a modular functor from quantized Teichmüller theory*, from: “Handbook of Teichmüller theory. Vol. I”, (A Papadopoulos, editor), IRMA Lect. Math. Theor. Phys. 11, Eur. Math. Soc., Zürich (2007) 685–760 MR2349683
- [19] **W P Thurston**, *Minimal stretch maps between hyperbolic surfaces* arXiv math/9801039
- [20] **H Verlinde**, *Conformal field theory, two-dimensional quantum gravity and quantization of Teichmüller space*, Nuclear Phys. B 337 (1990) 652–680 MR1057726
- [21] **H Verlinde, E Verlinde**, *Conformal field theory and geometric quantization*, from: “Superstrings ’89”, (M Green, R Iengo, S Randjbar-Daemi, E Sezgin, A Strominger, editors), World Sci. Publ. (1990) 422–449 MR1159975
- [22] **E Witten**, *Quantum field theory and the Jones polynomial*, Comm. Math. Phys. 121 (1989) 351–399 MR990772

- [23] **S Wolpert**, *On the Weil–Petersson geometry of the moduli space of curves*, Amer. J. Math. 107 (1985) 969–997 MR796909
- [24] **S Wolpert**, *The Weil–Petersson metric geometry*, from: “Handbook of Teichmüller theory. Vol. II”, (A Papadopoulos, editor), IRMA Lect. Math. Theor. Phys. 13, Eur. Math. Soc., Zürich (2009) 47–64 MR2497791

*Department of Mathematics, Rutgers University
110 Frelinghuysen Rd, Piscataway, NJ 08854-8019, USA*

`juroger@math.rutgers.edu`

`http://math.rutgers.edu/~juroger`

Received: 8 February 2013