

AG
T

*Algebraic & Geometric
Topology*

Volume 15 (2015)

**Lagrangian cobordisms via generating families:
Construction and geography**

FRÉDÉRIC BOURGEOIS

JOSHUA M SABLOFF

LISA TRAYNOR



Lagrangian cobordisms via generating families: Construction and geography

FRÉDÉRIC BOURGEOIS
JOSHUA M SABLOFF
LISA TRAYNOR

Embedded Lagrangian cobordisms between Legendrian submanifolds are produced by isotopy, spinning, and handle-attachment constructions that employ the technique of generating families. Moreover, any Legendrian with a generating family has an immersed Lagrangian filling with a compatible generating family. These constructions are applied in several directions, in particular to a nonclassical geography question: any graded group satisfying a duality condition can be realized as the generating family homology of a connected Legendrian submanifold in \mathbb{R}^{2n+1} or in the 1–jet space of any compact n –manifold with $n \geq 2$.

Corrections were submitted on 7 Oct 2023 and posted online on 15 Oct 2023.

57R17; 53D12, 53D42

1 Introduction

Lagrangian cobordisms between Legendrian submanifolds have recently enjoyed increasing interest, partly due to their centrality in symplectic field theory (see Eliashberg, Givental and Hofer [19], and see also Bourgeois, Ekholm and Eliashberg [7] and Ekholm [13; 14] for recent examples) and wrapped Fukaya category frameworks (see Abbondandolo and Schwarz [1], Abouzaid and Seidel [2], and Fukaya, Seidel and Smith [24]), and partly due to their connections to smooth topology, including the computation of the slice genus of a knot (see Chantraine [9]).

This paper extends the study of Lagrangian cobordisms in two ways: First, we develop tools to construct Lagrangian cobordisms between Legendrian submanifolds in 1–jet bundles using the Morse-theoretic framework of generating families. We call the resulting Lagrangian cobordism \bar{L} between the Legendrians Λ_- and Λ_+ a *gf-compatible* cobordism and write $(\Lambda_-, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+)$; see below for a precise definition. An important special case is a *gf-compatible* null-cobordism $(\emptyset, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+)$, which we call a *gf-filling*. Second, we investigate applications of our constructions, with an emphasis on solutions to geographic questions for Legendrian submanifolds.

1.1 Constructions

We study four constructions of Lagrangian cobordisms:

Immersed fillings (see [Proposition 3.1](#)) Every Legendrian submanifold with a generating family has an immersed gf-filling.

Spinning (see [Proposition 3.2](#)) An embedded, gf-compatible, n -dimensional cobordism may be “spun” to an $(n + 1)$ -dimensional embedded gf-compatible cobordism.

Legendrian isotopy (see [Proposition 3.6](#)) A Legendrian isotopy starting at a Legendrian with a generating family induces a gf-compatible embedded cobordism.

Handle attachment/embedded surgery (see [Theorem 4.2](#)) In an appropriate setup, it is possible to create a gf-compatible cobordism by attaching a Lagrangian handle, thereby realizing an embedded Legendrian surgery. This construction may also be performed locally to create a Lagrangian cobordism even without the presence of a global generating family.

The construction of an immersed filling is novel, but others have studied the spinning, isotopy, and surgery constructions. The fact that Legendrian isotopy induces a Lagrangian cobordism (though without the gf-compatibility) has previously appeared in work of Chantraine [9] and Eliashberg and Gromov [20, Lemma 4.2.5]. The surgery construction and its relation to generating families is closely related to work of Entov [21], though his work applies to a somewhat different setting. During the preparation of this paper, alternative approaches, without reference to generating families, to spinning (see Golovko [25; 26]) and surgery (see Dimitroglou Rizell [12] and Ekholm, Honda and Kálmán [17]) were developed.

1.2 Applications

With these constructions in hand, we turn to their applications. The first type of application involves finding Lagrangian fillings of Legendrian submanifolds in $J^1\mathbb{R}^n$ using the surgery and isotopy constructions. The 3-dimensional case is particularly interesting, as Chantraine proved that a Lagrangian filling for a Legendrian knot realizes the smooth 4-ball genus of the underlying smooth knot [9]. In [Section 5](#), we demonstrate the existence of Lagrangian fillings for several families of links: Legendrian links that topologically are twisted Whitehead doubles or 0-closures of positive braids (see [Propositions 5.1](#) and [5.4](#)). The techniques we discuss here also have interesting applications that have appeared elsewhere:

- Any positive knot has a Legendrian representative with a Lagrangian filling (see Hayden and Sabloff [27]).

- There exist Legendrian links with nonhomeomorphic fillings, answering a question of Boileau and Fourrier [5] about the uniqueness of fillings of links by complex curves (see Cao, Gallup, Hayden and Sabloff [8]).
- For any Legendrian knot, it is possible to construct a Lagrangian cobordism so that Λ_- is a Legendrian unknot; for twist, torus, and low crossing knots this can be done so that the cobordism has minimal smooth genus (see Boranda, Traynor and Yan [6]).

We note that Ekholm, Honda, and Kálmán's construction of Lagrangian cobordisms in [17] (see also Dimitriglou Rizell [12]) would work equally well for these applications. In contrast, there are also several applications of the fact that the cobordisms we construct in this paper are gf-compatible:

- There exist Legendrian spheres in $J^1\mathbb{R}^n$ that have arbitrarily many different generating family homologies (see [8] again).
- Denote the space of all Legendrian spheres in $J^1\mathbb{R}^n$ by \mathcal{L}^n . For any $n > 1$, there exists a Legendrian sphere Λ^n such that $\pi_1(\mathcal{L}^n; \Lambda^n)$ is nontrivial (see Sabloff and Sullivan [38]).

The second type of application is to geography questions for Legendrian submanifolds:

Which collections of invariants can be realized by a Legendrian submanifold?

We first restrict the geography question to the generating family cohomology invariant, which is defined in Section 2 below. Denote the Poincaré polynomial of the generating family cohomology $GH^*(f)$ by $\Gamma_f(t)$.

Question (Nonclassical generating family geography) Given a Laurent polynomial $P(t) \in \mathbb{N}[t, t^{-1}]$, is there a connected Legendrian submanifold $\Lambda \subset J^1M$ (where M is compact or equal to \mathbb{R}^n) with a linear-at-infinity generating family f such that $\Gamma_f(t) = P(t)$?

Any investigation of geography must begin by understanding obstructions. The first obstruction to a polynomial being the Poincaré polynomial for the generating family homology is the duality result of Traynor and Sabloff [39], which we strengthen below in Theorem 6.1; see also Ekholm, Etnyre and Sabloff [15], Fuchs and Rutherford [23], and Sabloff [37]. In particular, duality implies the following:

Theorem 1.1 *Given a Legendrian $\Lambda \subset J^1M$ with a linear-at-infinity generating family f , the generating family Poincaré polynomial is of the form*

$$(1-1) \quad \Gamma_f(t) = (q_0 + q_1t + \cdots + q_n t^n) + p(t) + t^{n-1} p(t^{-1}),$$

where $p(t) = \sum_{i \in \mathbb{Z}, i \geq \lfloor (n-1)/2 \rfloor} p_i t^i$, $q_k + q_{n-k}$ is the k^{th} Betti number of Λ , and $q_n \neq 0$.

For connected Legendrians, the polynomial in (1-1) has $q_n = 1$ and $q_0 = 0$. A polynomial of the form of the right hand side of Equation (1-1) with $q_n \neq 0$ is said to be *compatible with duality*; if, in addition, the polynomial has $q_n = 1$ and $q_0 = 0$, then the polynomial is in *connected form*. The spinning, isotopy, and embedded surgery constructions, coupled with the cobordism exact sequence of [39] (see below), allow us to prove the following complete answer to the nonclassical generating family geography question.

Theorem 1.2 *If the Laurent polynomial $P(t)$ is compatible with duality and is in connected form, then for any $n \geq 2$, there exists a connected Legendrian submanifold of J^1M with a generating family f so that $\Gamma_f(t) = P(t)$.*

The $n = 1$ case was proven by Melvin and Shrestha for Legendrian contact homology [31], and the work of Fuchs and Rutherford [23] implies that the results of Melvin and Shrestha also hold for generating family homology.

In fact, the Legendrians constructed to prove the theorem above are all gf-compatibly Lagrangian cobordant to higher-dimensional analogues of the Hopf link, and (hence) are all Lagrangian null-cobordant (though not necessarily gf-compatibly). Our techniques are suitably functorial so that, using the results of [15] in place of Theorem 1.1 to restrict the admissible polynomials, and [11; 25] for Legendrian contact homology versions of the cobordism exact sequence, Theorem 1.2 also holds for Legendrian contact homology.

Our investigations into the nonclassical geography question also yield results for the classical Thurston–Bennequin and rotation number invariants in higher dimensions:

Question (Classical fillable geography) *Given a pair of integers (τ, ρ) , does there exist a fillable Legendrian n -sphere $\Lambda \subset J^1\mathbb{R}^n$ whose Thurston–Bennequin and rotation numbers are τ and ρ , respectively?*

It is well-known that if Λ is fillable, then its rotation number vanishes; see the discussion in [9, Section 2.2], for example. Further, for even-dimensional Legendrians, the

Thurston–Bennequin number is determined by the Euler characteristic; see [16] or [33, Appendix A], for example. Thus, the only interesting question for classical fillable geography involves the Thurston–Bennequin number in odd dimensions, where tb must be odd [33, Appendix A]. We provide a complete answer to the classical fillable geography question in Section 6.3:

Theorem 1.3 *For any odd n and τ , there exists an odd-dimensional fillable Legendrian sphere $\Lambda \subset J^1\mathbb{R}^n$ with $tb(\Lambda) = \tau$.*

Plan of the paper We set notation and sketch the necessary background for Lagrangian cobordisms and generating families in Section 2. In Section 3, we work in the setting of gf-compatible Legendrians and Lagrangians to construct immersed Lagrangian fillings, spun Legendrian submanifolds and Lagrangian cobordisms, and Lagrangian cobordisms arising from Legendrian isotopy. The construction of the attachment of a Lagrangian handle is described in Section 4. Applications of the constructions begin in Section 5 with several examples of gf-fillings for Legendrian knots in $J^1\mathbb{R}$. Finally, in Section 6, we use constructions developed in this paper, together with the cobordism exact sequence of [39], to investigate the geography questions mentioned above.

Acknowledgements The authors thank the American Institute of Mathematics, the Université de Nantes, the Royal Academies for Science and the Arts of Belgium, and the Banff International Research Station for hosting conferences at which the authors initiated and completed the work discussed in this paper. The authors also thank Matt Hedden for his help in straightening out some references. The present work is part of Bourgeois’ activities within CAST, a Research Network Program of the European Science Foundation. Bourgeois is partially supported by ERC Starting Grant StG-239781-ContactMath. Sabloff is partially supported by NSF grants DMS-0909273 and DMS-1406093. Traynor is partially supported by NSF grant DMS-0909021.

2 Background notions

In this section, we briefly review the language of generating families for Legendrian submanifolds of J^1M and for Lagrangian cobordisms between them. See [23; 28; 39; 40] for the original definitions and for more details.

2.1 Generating families for Legendrians

The technique of generating families may be used to construct Legendrian submanifolds of the 1-jet bundle of a smooth manifold M . For future reference, we denote the *front projection* by $\pi_{xz}: J^1M \rightarrow J^0M$ and the *base projection* by $\pi_x: J^1M \rightarrow M$.

Given a smooth manifold M , let $f: M^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ be a smooth function, where $M \times \mathbb{R}^N$ has coordinates (x, η) . Unless otherwise noted, we assume that M is either compact or \mathbb{R}^n . Suppose that $\mathbf{0}$ is a regular value of the map $\partial_\eta f: M \times \mathbb{R}^N \rightarrow \mathbb{R}^N$. We define the *fiber critical set* of f to be the n -dimensional submanifold $\Sigma_f = (\partial_\eta f)^{-1}(\mathbf{0})$. Define immersions $i_f: \Sigma_f \rightarrow T^*M$ and $j_f: \Sigma_f \rightarrow J^1M$ in local coordinates by

$$i_f(x, \eta) = (x, \partial_x f(x, \eta)), \quad j_f(x, \eta) = (x, \partial_x f(x, \eta), f(x, \eta)).$$

The image L of i_f is an immersed Lagrangian submanifold; the image Λ of j_f is an immersed Legendrian submanifold. We say that f *generates* L and Λ , or that f is a *generating family (of functions)*.

Not every Legendrian submanifold has a generating family — see [34], for example — but for those that do, the Morse theory of the set of generating families gives rise to interesting nonclassical invariants. Since the domain of a generating family $f: M \times \mathbb{R}^N \rightarrow \mathbb{R}$ is not compact, we need to control the behavior of f at infinity. We define f to be *linear-at-infinity*, or *tame*, if there exists a nonzero linear function $A: \mathbb{R}^N \rightarrow \mathbb{R}$ such that $f(x, \eta) = A(\eta)$ outside a compact subset of $M \times \mathbb{R}^N$.

The Morse-theoretic invariant that a generating family attaches to a Legendrian submanifold is defined using the *difference function* $\delta: M \times \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$:

$$(2-1) \quad \delta(x, \eta, \tilde{\eta}) = f(x, \tilde{\eta}) - f(x, \eta).$$

It is not hard to show that there is a one-to-one correspondence between the Reeb chords of Λ (ie line segments parallel to the z axis that begin and end on Λ) and the critical points of δ with positive critical value. Choose $\omega > \epsilon > 0$ so that all positive critical values of δ_f lie between ω and ϵ , and define the *relative (resp. total) generating family cohomology* of f to be the relative cohomology of the ω and ϵ (resp. $-\epsilon$) sublevel sets of δ :

$$GH^k(f) = H^{k+N+1}(\delta^\omega, \delta^\epsilon), \quad \widetilde{GH}^k(f) = H^{k+N+1}(\delta^\omega, \delta^{-\epsilon}).$$

The *generating family homology* is defined analogously with the same index shift. In most other sources, the relative generating family cohomology is simply called the generating family cohomology, and we will adopt this convention throughout this paper except in the [appendix](#). The long exact sequence of the triple $(\delta^\omega, \delta^\epsilon, \delta^{-\epsilon})$, together with the Thom isomorphism, relates the relative and total generating family cohomologies:

$$(2-2) \quad \dots \rightarrow GH^k(f) \rightarrow \widetilde{GH}^k(f) \rightarrow H^{k+1}(\Lambda) \rightarrow \dots$$

The set of all generating family cohomology groups taken over all possible generating families for Λ forms an invariant of Λ up to Legendrian isotopy (see [23; 40]).

2.2 Generating families for Lagrangian cobordisms

Let (X, α) be a contact manifold and denote its symplectization by $(\mathbb{R} \times X, d(e^t \alpha))$. A Lagrangian submanifold \bar{L} of the symplectization is said to be a *Lagrangian cobordism* between the Legendrian submanifolds $\Lambda_{\pm} \subset X$ if there exist $t_{\pm} > 0$ such that

$$\begin{aligned} \bar{L} \cap ((-\infty, t_-] \times X) &= (-\infty, t_-] \times \Lambda_-, \\ \bar{L} \cap ([t_+, \infty) \times X) &= [t_+, \infty) \times \Lambda_+. \end{aligned}$$

We denote such a Lagrangian cobordism by $\Lambda_- \prec_{\bar{L}} \Lambda_+$, and we denote the compact manifold $\bar{L} \cap [t_-, t_+] \times X$ by L .

If the contact manifold is a 1-jet bundle $J^1 M$, then there is a symplectomorphism between its symplectization and $T^*(\mathbb{R}_+ \times M)$ with its canonical symplectic structure. As in [39, Section 4], we shift our perspective to $T^*(\mathbb{R}_+ \times M)$ so that we may use generating families to describe Lagrangian cobordisms. Lagrangians constructed through generating families will be exact and these will map to exact Lagrangians in the symplectization of $J^1 M$. For ease of notation, we denote a Lagrangian cobordism \bar{L} and its image in $T^*(\mathbb{R}_+ \times M)$ by the same symbol.

If the functions $f_{\pm}: M \times \mathbb{R}^N \rightarrow \mathbb{R}$ and $F: \mathbb{R}_+ \times M \times \mathbb{R}^N \rightarrow \mathbb{R}$ satisfy the following relation for some $t_- < t_+$:

$$(2-3) \quad F(t, x, \eta) = \begin{cases} t f_-(x, \eta) & t \leq t_-, \\ t f_+(x, \eta) & t \geq t_+, \end{cases}$$

then we say that the triple (F, f_-, f_+) is *compatible*. A *gf-compatible* Lagrangian cobordism consists of a Lagrangian cobordism $\Lambda_- \prec_{\bar{L}} \Lambda_+$ together with a compatible triple of generating families for the three objects involved. We denote a gf-compatible cobordism by

$$(\Lambda_-, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+);$$

for a filling, ie when $\Lambda_- = \emptyset$, we will frequently use the shorthand notation $\emptyset \prec_{(\bar{L}, F)} (\Lambda, f)$, with the understanding that f_- is a linear function. We require the triple of compatible functions to be *tame* in the sense that f_{\pm} are linear-at-infinity and F is *slicewise linear at infinity*, ie each $F(t, \cdot, \cdot)$ is equal to a linear function $A_t(\eta)$ outside a compact set of $\{t\} \times M \times \mathbb{R}^N$.

A key finding of [39] is the following:

Theorem 2.1 (Cobordism exact sequence) *A gf-compatible Lagrangian cobordism $(\Lambda_-, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+)$ induces a linear map $\Psi_F: GH^k(f_-) \rightarrow GH^k(f_+)$ that fits into the long exact sequence*

$$(2-4) \quad \dots \longrightarrow GH^k(f_-) \xrightarrow{\Psi_F} GH^k(f_+) \longrightarrow H^{k+1}(L, \Lambda_+) \longrightarrow \dots$$

3 Basic constructions

In this section, we discuss three global constructions of gf-compatible Lagrangian cobordisms. The first is an explicit proof of the Gromov–Lees theorem in this setting, namely that a Legendrian submanifold with a linear-at-infinity generating family has a gf-compatible, immersed Lagrangian filling. The second is a translation of the Ekhholm–Etnyre–Sullivan spinning construction [16] to the generating family setting, together with a generalization to properly embedded Lagrangian submanifolds in half-spaces; see also [26]. The final construction is a translation to the generating family setting of Chantraine’s proof that Legendrian isotopy induces a Lagrangian cobordism [9]; see also [20] and [25].

3.1 Immersed Lagrangian fillings

If a Legendrian knot $\Lambda \subset \mathbb{R}^3$ has a generating family, then $r(\Lambda) = 0$ (see [23], for example) and thus has an immersed Lagrangian filling by work of Chantraine [9, Remark 4.2]. We strengthen this statement by extending it to Legendrian submanifolds in arbitrary dimensions and by showing that the filling can be constructed via a compatible generating family. The idea is to smoothly deform a generating family to a linear function.

Proposition 3.1 *If $\Lambda \subset J^1M$ is a Legendrian submanifold with a linear-at-infinity generating family f , then there exists an immersed gf-compatible cobordism $\emptyset \prec_{(\bar{L}, F)} (\Lambda, f)$.*

Proof Let $f: M \times \mathbb{R}^N \rightarrow \mathbb{R}$ be a linear-at-infinity generating family for Λ ; assume that f agrees with the nonzero linear function $A(\eta)$ outside a compact set. To show the existence of a gf-compatible immersed filling, it suffices to construct $F: \mathbb{R}_+ \times M \times \mathbb{R}^N \rightarrow \mathbb{R}$ so that:

- (1) For all t outside a compact set of $\{t\} \times M \times \mathbb{R}^N$, $F(t, x, \eta)$ is a nonzero linear function $B_t(\eta)$.
- (2) There exists a t_- such that, for $t \leq t_-$, $F(t, x, \eta) = B_t(\eta)$.

- (3) There exists a t_+ such that, for $t \geq t_+$, $F(t, x, \eta) = tf(x, \eta)$.
- (4) $\mathbf{0} \in \mathbb{R}^N$ is a regular value of $\partial_\eta F$.

To begin the process, choose a smooth, increasing function $\sigma: \mathbb{R}_+ \rightarrow \mathbb{R}$ that is 0 on $(0, 1]$ and 1 on $[2, \infty)$. Define a function $G: \mathbb{R}_+ \times M \times \mathbb{R}^N \rightarrow \mathbb{R}$ by

$$G(t, x, \eta) = t \cdot (\sigma(t)f(x, \eta) + (1 - \sigma(t))A(\eta)).$$

Then, for fixed $t_- < 1$ and $t_+ > 2$, G satisfies conditions (1–3) desired for F . For later purposes, note that outside a compact set of $\{t\} \times M \times \mathbb{R}^N$ and for $t \leq t_-$, $G(t, x, \eta) = tA(\eta)$.

We now modify G to guarantee that condition (4) is satisfied. Since f is a generating family, $\mathbf{0} \in \mathbb{R}^N$ is a regular value of $\partial_\eta f$. Since f is linear-at-infinity, we may find an open, convex ball $U \subset \mathbb{R}^N$ around $\mathbf{0}$ that consists of regular values of $\partial_\eta f$. By Sard’s theorem, there exists $\varepsilon_G \in U$ that is a regular value of $\partial_\eta G$ and satisfies for all $t \in [1, 2]$ that, as functions, $\varepsilon_G \cdot \eta \neq tA(\eta)$. Choose a smooth path $\varepsilon: \mathbb{R}_+ \rightarrow U$ so that $\varepsilon(t) \equiv \varepsilon_G$ when $t \leq 2$, $\varepsilon(t) \cdot \eta \neq tA(\eta)$ when $t \in [2, t_+]$, and $\varepsilon(t) \equiv 0$ when $t \geq t_+$. Now consider the function

$$F(t, x, \eta) = G(t, x, \eta) - \varepsilon(t) \cdot \eta = \begin{cases} G(t, x, \eta) - \varepsilon_G \cdot \eta & t \leq 2, \\ tf(x, \eta) - \varepsilon(t) \cdot \eta & t \in [2, t_+], \\ tf(x, \eta), & t \geq t_+. \end{cases}$$

Conditions (1–3) are still satisfied by F . Further, since ε_G is a regular value of $\partial_\eta G$, it follows that $\mathbf{0}$ is a regular value of $\partial_\eta F$ when $t \leq 2$. It remains to show that $\mathbf{0}$ is a regular value of $\partial_\eta F$ when $t \geq 2$. When $t \geq 2$, $\partial_\eta F(t, x, \eta)$ vanishes if and only if $\partial_\eta f(x, \eta) = \varepsilon(t)/t$. Since $\varepsilon(t)$ lies in the convex set of regular values U and $t \geq 2$, $\varepsilon(t)/t \in U$ is a regular value of $\partial_\eta f$ for all $t \geq 2$. Thus when $t \geq 2$, at a point (t, x, η) where $\partial_\eta F = 0$, we see that the final $n + N$ columns of the $N \times (1 + n + N)$ matrix $D(\partial_\eta F)$ form a submatrix of rank N , as desired. □

3.2 Spinning constructions

Let H^n denote the closed upper half-space of \mathbb{R}^n , ie

$$H^n = \{\mathbf{x} \in \mathbb{R}^n : x_n \geq 0\}.$$

Ekholm, Etnyre, and Sullivan [16] described a method of producing an $(n + 1)$ -dimensional Legendrian submanifold of $J^1\mathbb{R}^{n+1}$ from an n -dimensional Legendrian in J^1H^n by spinning a front diagram about the z axis. Ekholm and Kálmán generalized the construction to twist-spun Legendrians [18]. In this section, we generalize these constructions to certain properly embedded Legendrian submanifolds of J^1H^n , show

how these constructions can be performed using generating families, and discuss spinning Lagrangian cobordisms in the spirit of Golovko [25].

A smooth, properly embedded Legendrian submanifold Λ of $J^1 H^n$ is a *spinnable Legendrian* if for all $p \in \partial\Lambda \subset \partial J^1 H^n$, the local parametrization of Λ near p , $\phi: H^n \rightarrow J^1(H^n)$, extends to a smooth map $\phi: \mathbb{R}^n \rightarrow J^1(\mathbb{R}^n)$ by $\phi(x_1, \dots, x_n) = \phi(x_1, \dots, -x_n)$ when $x_n < 0$. A spinnable Legendrian Λ gives rise to a *spun Legendrian* $\Lambda^s \subset J^1\mathbb{R}^{n+1}$ whose front projection is obtained by rotating the front projection of Λ about the subspace $\{x_n = x_{n+1} = 0\}$. More generally, a *spinnable Legendrian loop* consists of a smooth isotopy Λ_θ , $\theta \in S^1 = [0, 2\pi]/\sim$, of spinnable Legendrians such that for all θ , a neighborhood of $\partial\Lambda_\theta$ does not vary with respect to θ . A spinnable isotopy Λ_θ , in turn, gives rise to a *twist-spun Legendrian* $\Lambda^{ts} \subset J^1\mathbb{R}^{n+1}$ by following the isotopy during a rotation about $\{x_n = x_{n+1} = 0\}$. More specifically, if $\phi_\theta: U \rightarrow H^n \times \mathbb{R}$ are smooth, local parametrizations of the fronts Λ_θ given by

$$\phi_\theta(q) = (x_1^\theta(q), \dots, x_n^\theta(q), z^\theta(q)),$$

then a parametrization of the front of the twist-spun submanifold Λ^{ts} is given by

$$\phi^{ts}(q, \theta) = (x_1^\theta(q), \dots, x_{n-1}^\theta(q), x_n^\theta(q) \cos \theta, x_n^\theta(q) \sin \theta, z^\theta(q)).$$

We say that a generating family $f: H^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ is a *spinnable generating family* if f has a smooth extension to $f: \mathbb{R}^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ with $f(x_1, \dots, x_n, \eta) = f(x_1, \dots, -x_n, \eta)$, for $x_n < 0$. A *spinnable loop of generating families* consists of a smooth path of spinnable generating families f_θ , $\theta \in S^1 = [0, 2\pi]/\sim$, such that in a neighborhood of $\partial H^n \times \mathbb{R}^N$, f_θ does not depend on θ . By construction, spinnable (loops of) generating families generate spinnable (loops of) Legendrians. A *spinnable loop of linear-at-infinity generating families* consists of a spinnable loop of generating families f_θ such that outside a compact set of $H^n \times \mathbb{R}^n$, $f_\theta(x, \eta)$ agrees with a nonzero linear function $A(\eta)$ for all θ .

Proposition 3.2 *If a Legendrian submanifold $\Lambda \subset J^1 H^n$ has a spinnable linear-at-infinity generating family then the spun Legendrian submanifold Λ^s has a linear-at-infinity generating family. More generally, if Λ_θ is generated by a spinnable loop of tame generating families then the twist-spun Legendrian submanifold Λ^{ts} has a linear-at-infinity generating family.*

Proof The first part of the claim follows from the second via the constant isotopy, so we only prove the twist-spun claim. If $f_\theta: H^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ is a spinnable loop of linear-at-infinity generating families for Λ_θ , then we claim that the function $f^{ts}: \mathbb{R}^{n-1} \times \mathbb{R}^2 \times \mathbb{R}^N \rightarrow \mathbb{R}$ given by

$$f^{ts}(x_1, \dots, x_{n-1}, x_n, \theta, \eta) = f_\theta(x_1, \dots, x_{n-1}, x_n, \eta),$$

where (x_n, θ) denote polar coordinates on \mathbb{R}^2 , is linear-at-infinity and generates Λ^{ts} . Since $f_\theta|_{\partial H^n \times \mathbb{R}^N}$ does not depend on θ , $f_\theta(x_1, \dots, x_{n-1}, 0, \eta)$ is independent of θ and so f^{ts} is well-defined. The condition that all f_θ coincide on a neighborhood of $\partial H^n \times \mathbb{R}^N$ guarantee that f^{ts} is smooth. Since each f_θ agrees with the same linear function $A(\eta)$ outside a compact set, f^{ts} is linear-at-infinity. A straightforward calculation shows that since $\mathbf{0}$ is a regular value of $\partial_\eta f_\theta$ for all θ , $\mathbf{0}$ is a regular value of $\partial_\eta f^{ts}$, and thus f^{ts} is a generating family for Λ^{ts} . \square

Example 3.3 One way to create higher-dimensional analogues of the Legendrian unknot is to cut the unknot in half along the z axis and spin the result; see Figure 1. Iterating this construction yields “flying saucers” with generating families in all dimensions.




Figure 1: Spinning half of a Legendrian unknot yields a “flying saucer”

Remark 3.4 In [29], Kálmán constructed a loop of Legendrian trefoil knots — ie a spinnable isotopy — that is not contractible in the space of Legendrian trefoils; this isotopy was viewed in the context of spinning in [18]. Even though the trefoil has a generating family, and hence the isotopy gives rise to a path of generating families for the trefoils along the isotopy, the fact that the twist-spun Legendrian torus does not have an augmentation shows that this path of generating families is not a loop.

The spinning construction for generating families may be extended to Lagrangian cobordisms; see [25] for a relative of this construction that does not consider generating families. As above, a *spinnable Lagrangian cobordism* is a properly embedded Lagrangian cobordism $\bar{L} \subset T^*(\mathbb{R}_+ \times H^n)$ such that for all $p \in \partial \bar{L} \subset \partial T^*(\mathbb{R}_+ \times H^n)$, the local parametrization of \bar{L} near p , $\phi: \mathbb{R} \times H^n \rightarrow T^*(\mathbb{R}_+ \times H^n)$, extends to a smooth map $\phi: \mathbb{R} \times \mathbb{R}^n \rightarrow T^*(\mathbb{R}_+ \times \mathbb{R}^n)$ by $\phi(t, x_1, \dots, x_n) = \phi(t, x_1, \dots, -x_n)$. We say that a generating family $F: \mathbb{R}_+ \times H^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ is a *spinnable generating family* if F has a smooth extension to $F: \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ by $F(t, x_1, \dots, x_n, \eta) = F(t, x_1, \dots, -x_n, \eta)$. By construction, a spinnable generating family generates a spinnable Lagrangian cobordism.

The construction of a spun Lagrangian cobordism with a spun generating family then follows exactly the same steps.

Proposition 3.5 *Given a spinnable Lagrangian cobordism \bar{L} with tame, spinnable generating family F of the form $(\Lambda_-, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+)$ in $T^*(\mathbb{R}_+ \times H^n)$, the spun Lagrangian cobordism $\Lambda_-^s \prec_{\bar{L}^s} \Lambda_+^s$ has a compatible, tame generating family.*

3.3 Legendrian isotopy

In this subsection, we translate the fact that a Legendrian isotopy induces a Lagrangian cobordism [9; 20; 25] to the generating family setting. Namely, we prove:

Proposition 3.6 *Suppose that Λ_- is a Legendrian submanifold of $J^1 M$ with a tame generating family f_- , and that Λ_- is Legendrian isotopic to Λ_+ . Then there exists an embedded gf-compatible Lagrangian cobordism $(\Lambda_-, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+)$. Further, if f_- is tame, then F is tame.*

The proof of Proposition 3.6 will use the notion of a difference function $\delta(x, \eta, \tilde{\eta})$, as introduced in Section 2.1. The following lemma is our key technical tool:

Lemma 3.7 (Family of functions construction) *Given a 1-parameter family of functions $f_t: M \times \mathbb{R}^N \rightarrow \mathbb{R}$, $t \in \mathbb{R}_+$, define the function $F: \mathbb{R}_+ \times M \times \mathbb{R}^N \rightarrow \mathbb{R}$ by*

$$F(t, x, \eta) = t f_t(x, \eta).$$

If

- (1) $\mathbf{0}$ is a regular value of $\partial_\eta F$, and
- (2) for all $(x, \eta, \tilde{\eta})$ with $\eta \neq \tilde{\eta}$ in the fiber critical set of the difference function $\delta_t(x, \eta, \tilde{\eta}) = f_t(x, \tilde{\eta}) - f_t(x, \eta)$, we have

$$(3-1) \quad \frac{\partial}{\partial x} \delta_t(x, \eta, \tilde{\eta}) = 0 \implies \delta_t(x, \eta, \tilde{\eta}) \neq -t \frac{\partial}{\partial t} \delta_t(x, \eta, \tilde{\eta}),$$

then F generates an embedded Lagrangian submanifold of $T^*(\mathbb{R}_+ \times M)$.

Proof If $\mathbf{0}$ is a regular value of $\partial_\eta F$, then F generates an immersed Lagrangian $\bar{L} \subset T^*(\mathbb{R}_+ \times M)$ given by

$$(3-2) \quad \bar{L} = \{(t, x, f_t(x, \eta) + t \partial_t f_t(x, \eta), t \partial_x f_t(x, \eta)) : \partial_\eta f_t(x, \eta) = 0\}.$$

A direct calculation shows that the double points of \bar{L} are in bijective correspondence with points $(t, x, \eta, \tilde{\eta})$ with $\eta \neq \tilde{\eta}$ satisfying:

- (1) $(x, \eta, \tilde{\eta})$ is in the fiber critical set of δ_t ,
- (2) $\frac{\partial}{\partial x} \delta_{f_t}(x, \eta, \tilde{\eta}) = 0$, and
- (3) $\delta_{f_t}(x, \eta, \tilde{\eta}) = -t \frac{\partial}{\partial t} \delta_{f_t}(x, \eta, \tilde{\eta})$.

By hypothesis, it is impossible to simultaneously satisfy these three conditions, and thus the Lagrangian generated by F is embedded. □

We are now ready to prove [Proposition 3.6](#).

Proof of Proposition 3.6 Let Λ_t be a 1-parameter family of Legendrian submanifolds of J^1M such that $\Lambda_t = \Lambda_+$ for $t \geq t_+$ and $\Lambda_t = \Lambda_-$ for $t \leq t_-$. By the persistence of linear-at-infinity generating families under Legendrian isotopy (see, for example, [10; 28]), we obtain a 1-parameter family of linear-at-infinity generating families $f_t: M \times \mathbb{R}^N \rightarrow \mathbb{R}$ that generate Λ_t for $t \in [t_-, t_+]$. Extend f_t to a smooth family for all $t \in \mathbb{R}_+$ so that outside of a compact interval, we have $f_t = f_{\pm}$.

We will now check that for an appropriate parametrization of the family f_t , the function $F(t, x, \eta) = tf_t(x, \eta)$ satisfies the hypotheses of [Lemma 3.7](#), and hence generates the desired Lagrangian cobordism. First, it is straightforward to verify that since $\mathbf{0}$ is a regular value of $\partial_{\eta} f_t$ for all t , $\mathbf{0}$ is also a regular value of $\partial_{\eta} F$. Second, notice that $\frac{\partial}{\partial x} \delta_t(x, \eta, \tilde{\eta}) = 0$ when (x, η) and $(x, \tilde{\eta})$ correspond to the endpoints of a Reeb chord of Λ_t ; the “length” of the Reeb chord is precisely $\delta_t(x, \eta, \tilde{\eta})$, which may be positive or negative. Let $h > 0$ denote the minimum absolute value of the lengths of Reeb chords of all of the Legendrians in the isotopy Λ_t . It then suffices to show that for points $(x, \eta, \tilde{\eta})$ in the fiber critical set of δ_t , we have

$$(3-3) \quad \frac{h}{t} > |\partial_t \delta_t(x, \eta, \tilde{\eta})|.$$

Since, for each t , the fiber critical set of δ_t is compact, and $\partial_t \delta_t = 0$ for t outside of a compact interval, it follows that $\partial_t \delta_t$ is bounded on the domain of interest. Thus, after an orientation-preserving diffeomorphism ρ of \mathbb{R}_+ , we may assume that $\tilde{F}(t, x, \eta) = tf_{\rho(t)}(x, \eta)$ will satisfy [Equation \(3-3\)](#). □

We end this section with several remarks arising from the proof of [Proposition 3.6](#) and its consequences.

Remark 3.8 Since the Lagrangian cobordism generated in the proof of [Proposition 3.6](#) is, in fact, a concordance (ie is diffeomorphic to $\Lambda \times \mathbb{R}$), the cobordism exact sequence in [Theorem 2.1](#) tells us that the cobordism map Ψ_F is an isomorphism.

Remark 3.9 It is important to point out that if a Lagrangian is constructed with the generating family $F(t, x, \eta) = tf_t(x, \eta)$, where f_t generates a Legendrian Λ_t , the “slices” of the Lagrangian are not, in general, Λ_t . An examination of Equation (3-2) shows that if $f_t(x, \eta)$ generates the Legendrian Λ_t , the corresponding s -slice, $s \in \mathbb{R}$, of the Lagrangian in the symplectization $\mathbb{R} \times J^1M$ generated by $F(t, x, \eta, \tilde{\eta})$ will agree with Λ_t if and only if for some neighborhood of t , f_t is constant with respect to t . When f_t is not constant, the corresponding slice of the Lagrangian differs from Λ_t by a contribution of $t\partial_t f_t(x, \eta)$ to the z -coordinate of Λ_t .

4 Attaching Lagrangian handles

In this section, we explain how an embedded $(q-1)$ -surgery on a Legendrian $\Lambda_- \subset J^1M$ with a generating family induces a gf-compatible Lagrangian cobordism given by attaching a q -handle. As noted in the introduction, this construction is closely related to those of Entov [21] and Dimitroglou Rizell [12].

4.1 Attaching regions and handle attachment

We begin by specifying the data necessary to attach a Lagrangian q -handle along a Legendrian submanifold Λ . We denote the cusps of the front of a Legendrian Λ by Λ^\succ , and if Λ has a generating family, then the cusps that represent births/deaths between critical points of indices j and $j + 1$ are denoted $\Lambda^{\succ j}$. Finally, denote a k -dimensional disk of radius r by $D^k(r)$.

Definition 4.1 For $1 \leq q \leq n$, a smooth embedding

$$\sigma: D^q(1 + \lambda) \times D^{n-q}(\lambda) \times D^1(\lambda) \rightarrow M^n \times \mathbb{R},$$

for some small λ , is a q -attaching region for a Lagrangian q -handle along the front of a Legendrian submanifold Λ if

- (1) $\sigma^{-1}(\Lambda) = \{(u, v, w) : w^2 = (\|u\|^2 - \|v\|^2 - 1)^3\}$,
- (2) $\sigma^{-1}(\Lambda^\succ) = \{(u, v, 0) : \|u\|^2 - \|v\|^2 = 1\}$ with the image of this set called the *surgery domain*, and
- (3) for each fixed $(u_0, v_0) \in D^q(1 + \lambda) \times D^{n-q}(\lambda)$, $\sigma(\{u_0\} \times \{v_0\} \times D^1(\lambda))$ is parallel to the z direction in J^1M .

The embedding σ is a *gf-attaching region* if, in addition, property (2) is modified so that the surgery domain lies in $\Lambda^{\succ j}$ for some fixed $j \geq 0$. The *core disk* of the attaching region is the image of $D^q(1) \times \{0\} \times \{0\}$.




Figure 2: A schematic picture of the domain of an attaching region $\sigma: D^q(1 + \lambda) \times D^{n-q}(\lambda) \times D^1(\lambda) \rightarrow M^n \times \mathbb{R}$, with the preimage of the Legendrian Λ and the cusps Λ^\succ shown in solid and dotted curves respectively.

See Figure 2 for a schematic picture of an attaching region. After an isotopy of Λ , we may assume that the image of the attaching region lies in the vertical slice defined by $c - \lambda \leq z \leq c + \lambda$ and that the core disk lies in the hypersurface defined by $z = c$.

We are now ready to formally state the surgery construction.

Theorem 4.2 *Let Λ_- be a Legendrian submanifold of J^1M with a tame generating family $f_-: M^n \times \mathbb{R}^N \rightarrow \mathbb{R}$. Given a q -attaching region σ for Λ_- , there exists a smooth 1-parameter family of functions $f_t: M \times \mathbb{R}^N \rightarrow \mathbb{R}$ such that $F(t, x, \eta) = t f_t(x, \eta)$ is tame and generates an embedded Lagrangian cobordism $(\Lambda_-, f_-) \prec_{(\bar{L}, F)} (\Lambda_+, f_+)$ satisfying:*

- (1) *The cobordism \bar{L} has the homotopy type of a cylinder over Λ_- with a q -cell attached.*
- (2) *The Legendrian Λ_+ is obtained from Λ_- by an embedded $(q-1)$ -surgery along the boundary of the core disk.*

See Figure 3 for an example of performing 0-surgery on a two-dimensional Legendrian in \mathbb{R}^5 . Note that, as a special case, the theorem allows the attachment of a gf-compatible Lagrangian 0-handle, which results in a null-cobordism for the standard n -dimensional flying saucer.

The remainder of this section is devoted to a proof of the surgery construction theorem. The proof begins by transferring the attaching region to the domain of the generating family f_- . We then attach a handle in $\mathbb{R}_+ \times M \times \mathbb{R}^N$ to the fiber critical set of f_- . With this scaffolding in place, we construct F itself. The proof ends with a verification that F is, indeed, a generating family, and the Lagrangian cobordism that it generates is embedded.




Figure 3: (a) The core disk for a 0 surgery and (b) the result of embedded surgery along the core disk for a Legendrian surface in $\mathbb{R}^5 = J^1\mathbb{R}^2$

4.2 Transferring the attaching region to $M \times \mathbb{R}^N$

Condition (3) of Definition 4.1 guarantees that we may think of $D^q(1 + \lambda) \times D^{n-q}(\lambda)$ as being embedded in M . The region of the domain of f_- involved in the attaching of the handle is then

$$\tilde{E} = (D^q(1 + \lambda) \times D^{n-q}(\lambda) \times \mathbb{R}^N) \cap f_-^{-1}(c - \lambda, c + \lambda).$$

We let $\tilde{E}_{(u,v)}$ denote points in \tilde{E} whose first two coordinates are (u, v) .

The structure of the fiber critical set Σ_- of f_- in \tilde{E} is quite simple to describe using conditions (1) and (2) of Definition 4.1. In terms of coordinates (u, v) on $D^q(1 + \lambda) \times D^{n-q}(\lambda)$, we have:

- (1) If $\|u\|^2 - \|v\|^2 < 1$, then $\Sigma_- \cap \tilde{E}_{(u,v)} = \emptyset$.
- (2) If $\|u\|^2 - \|v\|^2 = 1$, then $\Sigma_- \cap \tilde{E}_{(u,v)}$ consists of a single point.
- (3) If $\|u\|^2 - \|v\|^2 > 1$, then $\Sigma_- \cap \tilde{E}_{(u,v)}$ consists of two points.

Thus, after a fiber-preserving diffeomorphism of \tilde{E} that preserves each $\tilde{E}_{(u,v)}$, we may assume that $\Sigma_- \cap \tilde{E}$ is the set

$$\{(u, v, \eta) : \|u\|^2 - \|v\|^2 - \eta_1^2 = 1, \eta_2 = \dots = \eta_N = 0\}.$$

For future use, we let $E = \tilde{E} \cap \{\eta_2 = \dots = \eta_N = 0\}$ and we let $Q(u, v, \eta_1) = -\|u\|^2 + \|v\|^2 + \eta_1^2$.

4.3 Attaching a handle to Σ_-

We next build the fiber critical set Σ of F by attaching a q -handle to Σ_- . The set $\Sigma \subset \mathbb{R}_+ \times M \times \mathbb{R}^N$ will consist of:

- (1) A cylindrical extension $(0, t_-] \times \Sigma_-$.

- (2) A q -handle attached to this cylinder for $t \in [t_-, t_+]$, whose boundary is $\Sigma_- \cup \Sigma_+ = \Sigma \cap \{t_{\pm}\}$.
- (3) A cylindrical extension $[t_+, \infty) \times \Sigma_+$.




Figure 4: The handle H in the domain of f_- , with $\pi_x(H)$ shown in grey

To construct the q -handle, we first form its projection H to $E \subset M \times \mathbb{R}^N$. As in the standard construction of a handle (see [32, Section 3]), we let H be the deformation retract of the sublevel set $Q^{\leq 1}$ to the region diffeomorphic to $D^q \times D^{n-q+1}$ depicted in Figure 4. We use the identification of H with $D^q \times D^{n-q+1}$ to split the boundary of H into three pieces:

- (1) $S = S^q(1 + \lambda) \times S^{n-q+1}(\lambda)$.
- (2) $\partial_- H = S^{q-1} \times D^{n-q} \setminus S$, ie the portion of the boundary of H that lies in Σ_- (not including S).
- (3) $\partial_+ H = D^q \times S^{n-q-1} \setminus S$. Note that the closure of $\partial_+ H$ is meant to be tangent to Σ_- (to all orders).

To place the handle H into the domain of $F: \mathbb{R}_+ \times M \times \mathbb{R}^N \rightarrow \mathbb{R}$, we consider a smooth function $h: H \setminus S \rightarrow \mathbb{R}$ such that:

- (1) h has a single critical point at the origin of critical value $t_c \in (t_-, t_+)$.
- (2) $h(u, v, \eta_1) = Q(u, v, \eta_1) + t_c$ near the origin.
- (3) $h^{-1}(t_{\pm}) = \partial_{\pm} H$.
- (4) Over $\partial_{\pm} H$, the graph of h is tangent (to all orders) to the vertical cylinder $\partial_{\pm} H \times \mathbb{R}$.

We construct Σ by taking the union

$$\Sigma = (0, t_-] \times \Sigma_- \cup (\Sigma_- \setminus \partial_- H) \times [t_-, t_+] \cup \text{graph}(h) \cup [t_+, \infty) \times \Sigma_+.$$

That Σ is smooth follows from the last two conditions in the definition of h .

A key feature of the Σ constructed above is that for all $x \in \pi_x(\text{Int } H)$, the cardinality of the set

$$\Sigma_{(t,x)} = \Sigma \cap h^{-1}(\{t\}) \cap \pi_x^{-1}(\{x\}) \cap f^{-1}(c - \lambda, c + \lambda)$$

is an increasing function of t , passing from 0 to 1 (for at most one value of t) to 2.

The following lemma is obvious from the construction above, as we essentially use the classical Morse-theoretic picture of [32].

Lemma 4.3 Σ_+ is obtained from Σ_- by a $(q-1)$ -surgery.

4.4 Constructing F using Σ_-

The next step in the proof of Theorem 4.2 will be to construct a 1-parameter family $f_t: M \times \mathbb{R}^N \rightarrow \mathbb{R}$, with $t \in \mathbb{R}_+$, so that Σ is the fiber critical set of $F(t, x, \eta) = t f_t(x, \eta)$. Constructing the family f_t is equivalent to constructing a smooth family of functions $f_{(t,x)}: \mathbb{R}^N \rightarrow \mathbb{R}$ for $(t, x) \in \mathbb{R}_+ \times M$ so that the critical points of $f_{(t,x)}$ are precisely the points of $\Sigma_{(t,x)}$.

For $x \notin \pi_x(H)$, simply let $f_{(t,x)} = (f_-)_x$. Note that in this case, $f_{(t,x)}$ is clearly linear-at-infinity.

Now suppose that $x \in \pi_x(H)$. As noted above, the cardinality of $\Sigma_{(t,x)}$ increases from 0 to at most 2, and contains a single point for at most one value t_0 of t . Outside of a neighborhood of $\Sigma_{(t_0,x)}$, let $f_{(t,x)} = (f_-)_x$. If there exists $t_0 \in [t_-, t_+]$ so that $\Sigma_{(t_0,x)}$ consists of a single point (t_0, x, η_x) , then the construction of $f_{(t,x)}$ proceeds as follows: working one x slice at a time, we modify $(f_-)_x$ in a neighborhood of (x, η_x) so that there are no new critical points for $t_- \leq t < t_0$, there is one birth-death critical point for $t = t_0$, and for $t_+ \geq t > t_0$ there is a pair of nondegenerate critical points of indices $j + 1$ and j at positions dictated by $\Sigma_{(t,x)}$. The local nature of this modification shows that $f_{(t,x)}$ has exactly the same behavior at infinity as f_- . This finishes the construction of $f_{(t,x)}$, and hence the construction of the generating family F .

The proof of Theorem 4.2 will be completed by the proofs of the following two claims:

Claim 4.4 After a small perturbation, F is a generating family for a Lagrangian cobordism between Λ_- and Λ_+ .

This claim essentially follows from the fact that we constructed $\Sigma = (\partial_\eta F)^{-1}(0)$ to be a submanifold of $\mathbb{R}_+ \times M \times \mathbb{R}^N$; if necessary, a slight perturbation of F will guarantee that $\mathbf{0}$ is a regular value of $\partial_\eta F$.

Claim 4.5 *The Lagrangian L generated by F is embedded.*

To prove this claim, let $\delta_t(x, \eta, \tilde{\eta})$ be the difference function of f_t . From Lemma 3.7, we know the Lagrangian generated by F will be embedded as long as for all $(x, \eta, \tilde{\eta})$ with $\eta \neq \tilde{\eta}$ in the fiber-critical set of δ_t , we have

$$\partial_x \delta_t(x, \eta, \tilde{\eta}) = 0 \implies \delta_t(x, \eta, \tilde{\eta}) \neq -t \partial_t \delta_t(x, \eta, \tilde{\eta}).$$

First, consider the case where $(\pi_x^{-1}(\pi_x(H)) \cap \Sigma_- \subset H$; that is, in the front projection, there are no additional portions of $\pi_{xz}(\Lambda_-)$ above or below the image of the core disk. In this case, the only potential immersion points arise from pairs of points inside H . By modifying f_t , we can guarantee that the only potential immersion points arise from a pair at the ‘‘center’’ of the handle: namely, for $(x, \eta, \tilde{\eta})$ with $\eta \neq \tilde{\eta}$ in the fiber-critical set of δ_t ,

$$\partial_x \delta_t(x, \eta, \tilde{\eta}) = 0 \iff t > t_c \text{ and } x = 0.$$

If η and $\tilde{\eta}$ are labeled so that $\delta_t(x, \eta, \tilde{\eta}) > 0$, then by modifying the movement of the critical values of f_t , we can guarantee that $\partial_t \delta_t(x, \eta, \tilde{\eta}) > 0$, and embeddedness of L follows.

Second, consider the case where potential immersion points arise from pairs of points (t, x, η) and $(t, x, \tilde{\eta})$ where one point is inside H and the other point lies in the portion of Σ that is cylindrical over Σ_- . In this case, by perturbing f_t we can ensure that there is a compact set of potential immersion points where $\partial_x \delta_t(x, \eta, \tilde{\eta}) = 0$, and that $|\delta_t(x, \eta, \tilde{\eta})| \geq h > 0$ on the domain of interest. Then, using an argument similar to the proof of Proposition 3.6, by increasing t_+ and reparametrizing f_t , we can guarantee that $|\delta_t(x, \eta, \tilde{\eta})|/t \geq h/t > |\partial_t \delta_t(x, \eta, \tilde{\eta})|$ at all the potential immersion points. This completes the proof of Claim 4.5, and hence the proof of Theorem 4.2.

A brief examination of the proof shows that the hypothesis of a global generating family for the Legendrian Λ_- is unnecessary so long as we do not expect either the Lagrangian cobordism \bar{L} or the Legendrian Λ_+ to have a generating family. All we need is a generating family for Λ_- in a neighborhood of the attaching sphere of the core disk, and such a family is easily constructed in a local model. Thus, we have the following corollary:

Corollary 4.6 *Let Λ_- be a Legendrian submanifold of $J^1 M$. Given a q -attaching region for Λ_- , there exists an embedded exact Lagrangian cobordism $\Lambda_- \prec_{\bar{L}} \Lambda_+$ such that:*

- (1) The cobordism \bar{L} has the homotopy type of a cylinder over Λ_- with a q -cell attached.
- (2) The Legendrian Λ_+ is obtained from Λ_- by an embedded $(q - 1)$ -surgery along the attaching sphere of the core disk.

Remark 4.7 Note that the surgery construction used in [Corollary 4.6](#) is equivalent to the ambient surgery construction by Dimitroglou Rizell in [\[12\]](#). There, the Lagrangian handle is defined (see [\[12, Section 4.2.2\]](#)) in the front projection as the union of two graphs, instead of our generating family description.

5 Constructions in dimension 3

In the special case of Legendrian links in \mathbb{R}^3 or $J^1 S^1$, the handle attachment construction in the previous section reduces to the existence of a Lagrangian cobordism between Legendrians whose front diagrams are depicted in [Figure 5](#). These cobordisms are related to the work of Ekholm, Honda, and Kálmán [\[17\]](#), but with the added benefit that if the Legendrian knot at the bottom of the cobordism has a linear-at-infinity generating family and the cusps have corresponding indices, then the link at the top also has a generating family and the cobordism has a compatible generating family.

In this section, we illustrate the possibilities of the construction in two families of examples. As noted in the introduction, deeper applications of these constructions appear in [\[6; 8; 27; 38\]](#).

While the description of these examples follows the “bottom up” construction specified by the handle attachment of the previous section, we note that these examples were discovered by working “top down” from the resulting links, using graded normal rulings to keep track of which pairs of critical points could be canceled (see [\[34\]](#) and [\[22\]](#), for example).

5.1 Example: Whitehead doubles

Recall that for any Legendrian knot $\Lambda \subset \mathbb{R}^3$, one can form the Legendrian Whitehead double, denoted $Wh_{tb}(\Lambda)$, as follows. First form the 2-copy of Λ , which is a link consisting of Λ and a small push-off of Λ in the z direction; then make this Legendrian link into a Legendrian knot by replacing a 0-tangle with a cusped ∞ -tangle. The top of [Figure 6](#) illustrates $Wh_{tb}(\Lambda)$ where Λ is the Legendrian unknot with $tb = -3$ and $r = 0$. Topologically, $Wh_{tb}(\Lambda)$ is the $tb(\Lambda)$ -twisted, positively clasped Whitehead double of the underlying knot type K of Λ .




Figure 5: The modifications to the front of a 1–dimensional Legendrian that arise from attaching a 0–handle and a 1–handle

It is not difficult to show that no matter the original knot Λ , its Legendrian Whitehead double has $tb(Wh_{tb}(\Lambda)) = 1$ and $r(Wh_{tb}(\Lambda)) = 0$. Further, it has at least one graded normal ruling, and hence a generating family by [23]. Moreover, we can use the techniques of the previous two sections to prove:

Proposition 5.1 *If Λ is a Legendrian knot in \mathbb{R}^3 , then $Wh_{tb}(\Lambda)$ has a Lagrangian filling of genus 1. Further, if $r(\Lambda) = 0$, then the filling is gf-compatible.*

Proof We will construct a Lagrangian filling of $Wh_{tb}(\Lambda)$ as follows: first, attach a 0–handle to obtain an unknot. Using the trace of Λ , use Reidemeister type I moves at the cusps and type II moves at the crossings to drag one cusp of the unknot along Λ until it lies next to the other cusp of the original unknot; see the left side of Figure 6 for an illustration when Λ is a Legendrian unknot with no crossings. Perform two more Reidemeister type I moves, one each on the top and bottom strands of the original unknot. Finally, attach two 1–handles as indicated in the center of Figure 6 to obtain $Wh_{tb}(\Lambda)$; if $r(\Lambda) = 0$, then the indices of the outer cusps match. The composition of the 0–handle attachment, the isotopy, and the two 1–handle attachments yields a genus 1 Lagrangian filling for $Wh_{tb}(\Lambda)$ which is gf-compatible if $r(\Lambda) = 0$. \square

By combining Proposition 5.1 with [39, Theorem 1.5] we obtain:

Corollary 5.2 *The set linearized contact homology or generating family polynomials for the Whitehead double of any Legendrian knot with rotation number 0 must contain the polynomial $2 + t$.*

Denote by $\overline{tb}(K)$ the topological invariant given by the maximal Thurston–Bennequin invariant of any Legendrian representative of a smooth knot K . By adding s stabilizations to a maximal tb representative of K , we obtain a Legendrian representative




Figure 6: The construction of a Lagrangian null-cobordism for a Whitehead double

Λ_s of K with $tb(\Lambda_s) = \overline{tb}(K) - s$. Applying the construction above, for all $s \geq 0$ we find a genus 1 Lagrangian null-cobordism for $Wh_{tb}(\Lambda_s)$, which is topologically the $(\overline{tb}(K) - s)$ -twisted Whitehead double of the underlying knot type K of Λ . Combining this construction with Chantraine’s result that a Lagrangian filling realizes the smooth 4–ball genus of a Legendrian knot [9], we obtain a result of Rudolph:

Corollary 5.3 (Rudolph [36]) *For any $r \leq \overline{tb}(K)$, the r -twisted Whitehead double of K has smooth 4–ball genus equal to 1.*

5.2 Example: Positive braids

Given a positive braid $B = \sigma_{i_1} \cdots \sigma_{i_k}$ on s strands, form a Legendrian link Λ_B as in the top of Figure 7.

Proposition 5.4 *For any positive braid B with k crossings on s strands, the Legendrian link Λ_B with c components has a gf-filling of genus $\frac{1}{2}(2 - c + k - s)$.*




Figure 7: The construction of a Lagrangian null-cobordism of genus 0 for the zero closure of the positive braid $\sigma_2\sigma_1$. The 1–handles are attached in succession from the outermost to innermost cusps.

Proof We construct a gf-filling of Λ_B as follows: first, for each generator σ_{ij} , attach $(s - 1)$ 0–handles and perform an isotopy to obtain a nested unlink U_j of $(s - 1)$ components with a single crossing mimicking the σ_{ij} crossing in Λ ; for an example, see the left side of Figure 7. Position these unlinks next to each other, ordered from U_1 to U_k . Finally, successively attach s 1–handles between each pair of adjacent unlinks, starting at the outermost cusps, as in Figure 7 to obtain Λ_B . If Λ_B has c components, then this filling has genus $\frac{1}{2}(2 - c + k - s)$. \square

In particular, when Λ_B is a knot, we see that the smooth 4–ball genus of Λ_B is $\frac{1}{2}(1 + k - s)$, as originally proved by Rudolph [35, Section 3] as a corollary of Kronheimer and Mrowka’s work on embedded surfaces in 4–manifolds [30].

Remark 5.5 In a similar spirit, Lagrangian fillings of positive, Legendrian rational links are constructed in [6]. In particular, it is shown that the smooth 4–ball genus of these positive rational knots can be calculated from its rational notation; see [6, Remark 1.8].

Remark 5.6 The discussion in this subsection is the beginning of a more interesting story about the relationship between various notions of positivity (braid positivity, positivity, and (strong) quasi-positivity) and the existence of a Lagrangian filling; see [27] for a deeper exploration.

6 Legendrian geography

With the isotopy, spinning and, most importantly, handle attachment constructions in hand, we proceed to apply them to two questions about the geography of Legendrian submanifolds: a nonclassical geography question (which generating family polynomials can be realized by Legendrian submanifolds?) and the classical fillable geography question (what Thurston–Bennequin numbers can be realized by fillable Legendrian submanifolds?). Throughout this section, we work with coefficients in a field \mathbb{F} .

6.1 Duality and compatible polynomials

After strengthening the duality exact sequence for generating family cohomology of [39] to better take into account the algebraic topology of the underlying Legendrian, we prove [Theorem 1.1](#), which restricts the possible Poincaré polynomials for the generating family cohomology. Analogous versions of these results for linearized contact homology appear in [15]. The strengthening of the duality exact sequence takes two forms. First, we relate the maps in the duality exact sequence to the Poincaré duality of the Legendrian; second, we prove the existence of a “fundamental class” for the generating family homology.

Theorem 6.1 (Duality) *If Λ is a Legendrian submanifold of J^1M with linear-at-infinity generating family f , then there is a long exact sequence*

$$(6-1) \quad \dots \longrightarrow GH^{k-1}(f) \xrightarrow{\rho_k} GH_{n-k}(f) \xrightarrow{\sigma_k} H^k(\Lambda) \xrightarrow{\delta_k} \dots .$$

The maps δ_k satisfy two further properties:

- (1) *If $\gamma: H^k(\Lambda) \rightarrow H_{n-k}(\Lambda)$ is the Poincaré duality isomorphism, then, when using coefficients in a field, $\gamma \circ \sigma_k$ is the adjoint of the map δ_{n-k} .*
- (2) *The map δ_n does not vanish. In particular, over a field, it is an isomorphism when Λ is connected.*

We will delay the proof of this theorem until the [appendix](#), as the somewhat technical proof uses methods that are quite different than those in the rest of the paper. We call an element $\alpha \in \text{Im } \delta_k$ a *manifold class*. If Λ is connected, then the image under δ_n of the top class of Λ in $GH^n(f)$ is called the *fundamental class*.

Remark 6.2 If we were to use the total generating family cohomology instead of the relative version, then part (2) of the duality theorem above and long exact sequence (2-2) would imply that there is no fundamental class in $\widetilde{GH}^n(f)$, but that there is a manifold

class in degree 0. This indicates that the total generating family cohomology would be more convenient for the study of the cohomology ring, as the degree-0 manifold class would constitute a unit.

The strengthened duality theorem is the key ingredient in the proof of [Theorem 1.1](#), which states that every generating family polynomial is compatible with duality.

Proof of Theorem 1.1 To set notation, let $d_k = \dim GH^k(f)$, let $q_k = \dim \text{Im } \delta_k$, let $b_k = \dim H^k(\Lambda)$, and let $p_k = d_k - q_k$.

We begin by proving that under the hypotheses of [Theorem 6.1](#), when using field coefficients, we have

$$(6-2) \quad b_k = q_k + q_{n-k}.$$

First note that since δ_{n-k} and σ_k are adjoints up to the isomorphism γ , the dimensions of their kernels are the same. The relation (6-2) now follows from the rank–nullity theorem and the exactness of the sequence (6-1) at $H^k(\Lambda)$.

We next claim that for all $k \in \mathbb{Z}$, we have

$$(6-3) \quad p_k = p_{n-1-k}.$$

The rank–nullity theorem for ρ_k and the exactness of the sequence (6-1) at $GH^k(f)$ imply that

$$(6-4) \quad d_k = q_k + \dim \ker \sigma_{k+1}.$$

[Equation \(6-2\)](#) and the rank–nullity theorem for δ_k then imply that $q_{n-1-k} = b_{k+1} - q_{k+1} = \dim \ker \delta_{k+1}$. Combining this fact with the rank–nullity theorem for σ_{k+1} and the exactness of the sequence (6-1) at $GH_{n-1-k}(f)$, we obtain

$$(6-5) \quad d_{n-k-1} = \dim \ker \sigma_{k+1} + q_{n-1-k}.$$

Combining [Equations \(6-4\)](#) and (6-5) then yields

$$(6-6) \quad d_k = q_k + p_{n-1-k},$$

and [Equation \(6-3\)](#) now follows from the definition of p_k .

The theorem is now a consequence of [Equations \(6-2\)](#), (6-3), and (6-6) with q_k and p_k forming the coefficients of $q(t)$ and $p(t)$, respectively. The fact that $q_n \neq 0$ is a consequence of [Theorem 6.1](#). \square

6.2 Nonclassical geography

In the previous section we proved that every generating family polynomial of a connected Legendrian submanifold of $J^1 M$ is compatible with duality; in this section, we use the constructions of Sections 3 and 4 to prove [Theorem 1.2](#), namely that every Laurent polynomial in connected form that is compatible with duality is the generating family polynomial for *some* connected Legendrian submanifold of dimension $n \geq 2$. We will construct the Legendrian in a $J^1 \mathbb{R}^n$ coordinate chart inside $J^1 M$.

Before beginning the constructions proper, we set down a useful computation for the generating family cohomology of a gf-compatible 0–surgery connecting two connected Legendrians.

Lemma 6.3 (0–surgery lemma) *Suppose Λ_- has a tame generating family f_- such that $\Gamma_{f_-}(t)$ is of the form*

$$\Gamma_{f_-}(t) = (2t^n + q_{n-1}t^{n-1} + \dots + q_1t + q_0) + p(t) + t^{n-1}p(t^{-1}),$$

where $p(t) = \sum_{i \in \mathbb{Z}, i \geq \lfloor (n-1)/2 \rfloor} p_i t^i$. If Λ_+ is a connected Legendrian obtained from Λ_- by a gf-compatible 0–surgery, then Λ_+ has a tame generating family f_+ with

$$\Gamma_{f_+}(t) = \Gamma_{f_-}(t) - t^n.$$

Proof Let L denote the $(n + 1)$ –dimensional Lagrangian cobordism between the Legendrians Λ_- and Λ_+ described in the hypotheses of the lemma. Since L is obtained from Λ_- by attaching a 1–handle, L is obtained from Λ_+ by attaching an n –handle. When k is neither n nor $n - 1$, the cobordism exact sequence of [Theorem 2.1](#) implies that $GH^k(f_-) \simeq GH^k(f_+)$. The remaining terms of the cobordism exact sequence are

$$(6-7) \quad 0 \rightarrow GH^{n-1}(f_-) \rightarrow GH^{n-1}(f_+) \rightarrow \mathbb{F} \rightarrow GH^n(f_-) \rightarrow GH^n(f_+) \rightarrow 0.$$

Since $\dim GH^n(f_-) = 2 + p_n$ by hypothesis, the exactness of the sequence above implies that either $\dim GH^n(f_+) = 1 + p_n$ or $\dim GH^n(f_+) = 2 + p_n$. To see which of these is correct, we turn to the duality exact sequence [\(6-1\)](#), which tells us that

$$0 \rightarrow GH^{-1}(f_+) \rightarrow GH_n(f_+) \rightarrow H^0(\Lambda_+) \rightarrow \dots$$

We already know the isomorphism $GH^{-1}(f_+) \simeq GH^{-1}(f_-)$, and thus we have that $\dim GH^{-1}(f_+) = p_{-1} = p_n$. Since Λ_+ is connected, we use [Theorem 6.1](#) (2) to conclude that $\dim GH_n(f_+) = \dim GH^n(f_+)$ must be $1 + p_n$. It then follows from the exactness of the sequence [\(6-7\)](#) that $\dim GH^{n-1}(f_+) = q_{n-1} + p_{n-1}$, thus proving the lemma. □

One situation in which we can use this lemma is in the connected sum of two connected Legendrians in $J^1\mathbb{R}^n$. Given two Legendrians with tame generating families (Λ_i, f_i) , $i = 1, 2$, sufficiently separated in the horizontal direction, there is a tame generating family f_- for their disjoint union with

$$\Gamma_{f_-}(t) = \Gamma_{f_1}(t) + \Gamma_{f_2}(t);$$

see [39, Proposition 3.19] for more details. Let (Λ_+, f_+) be the result of a gf-compatible 0–surgery on (Λ_-, f_-) that connects the two components of Λ_- .

Corollary 6.4 (Connected sums) *The generating family cohomology of the connect sum (Λ_+, f_+) may be computed as*

$$\Gamma_{f_+}(t) = \Gamma_{f_1}(t) + \Gamma_{f_2}(t) - t^n.$$

The first important step in the proof of Theorem 1.2 is to build up Legendrians with interesting manifold classes, which are represented by the polynomial $q_0 + q_1t + \dots + q_nt^n$ in Equation (1-1).

Lemma 6.5 (Manifold lemma) *For all integers $a = 1, \dots, n - 1$, there exists a connected, n –dimensional Legendrian submanifold $\Lambda_a \subset J^1M$ with a tame generating family f_a such that*

$$\Gamma_{f_a}(t) = t^n + t^a.$$

In addition, there exists a 2–component n –dimensional Legendrian link Λ_0 with a tame generating family f_0 such that

$$\Gamma_{f_0}(t) = t^n + t^0.$$

Proof In this proof and in the proofs below, we will construct the Legendrian Λ_a in a $J^1\mathbb{R}^n$ coordinate chart inside J^1M . Let Λ_- denote the n –dimensional Legendrian sphere obtained from spinning the front of the Legendrian unknot shown in Figure 8. Since Legendrian isotopy commutes with spinning, we know that Λ_- is isotopic to the standard n –dimensional flying saucer. This implies that Λ_- is gf-fillable (by attaching a 0–handle as in Theorem 4.2, and by Proposition 3.6). By construction, Λ_- contains an $(n - 1)$ –dimensional sphere of cusps that bounds a horizontal disk D^n . For $a = 0, 1, \dots, n - 1$, perform an $(n - a - 1)$ –surgery on Λ_- , which when combined with the Lagrangian filling of Λ_- also yields a gf-filling (L_a, F_a) of (Λ_a, f_a) . For $a > 0$, the resulting Legendrian Λ_a is connected.

We now proceed to compute the generating family cohomology of the links Λ_a . Since the Lagrangian L_a was constructed by attaching an $(n - a)$ –handle to a 0–handle, its




Figure 8: Spinning this unknot results in an n -dimensional Legendrian submanifold isotopic to the standard flying saucer, but having an S^{n-1} family of cusps that bounds a disk D^n on the outside.

homology is supported in dimensions 0 and $n - a$, both with dimension 1. Poincaré duality then implies that

$$H^k(L_a, \Lambda_a) = \begin{cases} \mathbb{F} & k = a + 1, n + 1, \\ 0 & \text{otherwise.} \end{cases}$$

The cobordism exact sequence implies that $GH^k(f_a) \simeq H^{k+1}(L_a, \Lambda_a)$, and the lemma follows. □

Remark 6.6 The proof of Lemma 6.5 shows that the Legendrians Λ_a are all gf-fillable.

Taking connect sums of the Legendrians constructed in Lemma 6.5 and applying Corollary 6.4 allows us to build up Legendrians with (almost) arbitrary manifold classes.

Corollary 6.7 (Manifold class building block) *For arbitrary nonnegative integers q_1, q_2, \dots, q_{n-1} , there exists a connected, n -dimensional Legendrian submanifold that has a tame generating family f with*

$$\Gamma_f(t) = t^n + q_{n-1}t^{n-1} + \dots + q_1t^1.$$

We are now ready to build up the “duality classes” recorded by the polynomials $p(t)$ and $t^{n-1}p(t^{-1})$. We begin by constructing Hopf links with the desired classes.

Lemma 6.8 (Hopf link lemma) *For $n \geq 2$ and $a \neq 0, n - 1$, there exists an n -dimensional Hopf link Λ_a with a tame generating family f such that*

$$\Gamma_f(t) = 2t^n + t^a + t^{n-1}t^{-a}.$$

Proof By the symmetry of these polynomials, it suffices to show it is possible to realize the polynomial $2t^n + t^a + t^{n-1-a}$, for $a \geq \lfloor (n-1)/2 \rfloor$ and $a \neq n-1$.




Figure 9: Two Legendrian Hopf links used in the proof of Lemma 6.8. Both have six Reeb chords along the central axis, with the length of the Reeb chord from the bottom of the upper component to the top of the lower component shorter than then Reeb chord between the bottoms of the two components.

We begin by considering the 1-dimensional Legendrian Hopf link on the left side of Figure 9. We claim that this link has a generating family with the indicated (relative) fiber indices. Start with two linear-at-infinity generating families f_0 and f_1 for the lower and upper unknots, respectively, on the left side of Figure 9; suppose that they have the same fiber dimensions and that their fiber critical sets have the same fiber indices. Stabilize f_1 with a nondegenerate quadratic function of index $a + 1$ and stabilize f_0 with a nondegenerate quadratic function of the same dimension, but of index 0. Finally, translate the domain of f_1 in the fiber so that its support is disjoint from that of f_0 and let f_L be the sum of f_0 and f_1 in the sense of [39, Definition 3.18]; the resulting f_L is a generating family for the full Hopf link with the desired indices on the fiber critical sets.

Spinning the two fronts in Figure 9 yields isotopic n -dimensional Legendrian Hopf links Λ_L and Λ_R ; note that Λ_L has a generating family (again called f_L) by Proposition 3.2.

Since Λ_L and Λ_R are isotopic and Λ_L has a generating family, the persistence of generating families under isotopy implies that Λ_R has a generating family f_R that

has the same generating family homology as f_L . The difference functions for both f_L and f_R have six critical points with positive critical values. The indices of these critical points, listed in order of decreasing critical value, are:

Λ_L	$n + 1 + a$	n	$n + a$	n	$n - 1 - a$	a
Λ_R	$a + 1$	n	a	n	$n - 1 - a$	a

It follows that the critical points of f_L with indices $n + 1 + a$ and $n + a$ cannot contribute to the homology; similarly, for f_R , the critical points with indices $a + 1$ and a cannot contribute to the homology, and thus the total homology is at most 4-dimensional.

Except in the cases $a = n - a, n + 1, n - 1$, index arguments imply that the generating family homology must be 4-dimensional, ie that

$$\Gamma_f(t) = 2t^n + t^a + t^{n-1-a},$$

as desired.

By assumption, $a \neq n - 1$, so it remains to show that the generating family homology is 4-dimensional when $a = n - a$ and when $a = n + 1$.

When $a = n - a$, then the critical points in degrees a and $n - 1 - a = a - 1$ survive in homology since the critical value of the critical point of index $a - 1$ is larger than that of the critical point of index a . Thus, this case also yields the desired 4-dimensional homology.

Finally, when $a = n + 1$, we can apply a duality argument to show that the total homology is 4-dimensional. In this case, $n - 1 - a = -2$, and if the homology were not 4-dimensional, then we would have $\Gamma_f(t) = t^n + t^{-2}$. This is a contradiction to the duality long exact sequence, as the part of the sequence

$$GH^{n+1}(f) \rightarrow GH_{-2}(f) \rightarrow H^{n+1}(\Lambda)$$

would become

$$0 \rightarrow \mathbb{F} \rightarrow 0.$$

This completes the proof of the lemma. □

Remark 6.9 As all of the Hopf links constructed in the lemma above are Legendrian isotopic to the Hopf link Λ_0 of [Lemma 6.5](#), they are all fillable, though not necessarily gf-fillable.

We next use the Hopf links constructed above to produce Legendrian spheres whose generating family homologies have a given pair of dual classes.

Lemma 6.10 (Sphere lemma) *For all integers a , there exists a Legendrian n -sphere Λ with a tame generating family f such that*

$$\Gamma_f(t) = t^n + t^a + t^{n-1}t^{-a}.$$

Proof By symmetry, it suffices to restrict to the case where $a \geq \lfloor (n-1)/2 \rfloor$.

First, we generalize the first Reidemeister move to higher dimensions as follows: for any Legendrian submanifold $\Lambda \subset J^1\mathbb{R}^n$, the Legendrian submanifold Λ' obtained by replacing a graph-like portion of the front projection of Λ with one of the fronts depicted in Figure 10 is Legendrian isotopic to Λ . To see why, let $\Lambda_0 \subset \Lambda$ be an open subset obtained as the 1-jet of a function and contained in a Darboux chart of $J^1\mathbb{R}^n$. We claim that the 1-parameter family of generating families $f_t: [-1 \times 1] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$f_t(x, \eta) = \pm(\|\eta\|^4 + t\|\eta\|^2) + x \cdot \eta,$$

where $x = (x_1, \dots, x_n)$ and $\eta = (\eta_1, \dots, \eta_n)$, describes a Legendrian isotopy between Λ_0 (for $t = +1$) and the Legendrian submanifolds whose fronts are depicted in Figure 10 (for $t = -1$). The \pm sign can be chosen in order to obtain the front on the bottom or top of the figure.




Figure 10: The higher-dimensional first Reidemeister moves

With this technique in hand, take a Hopf link constructed in Lemma 6.8 and apply $a + 1$ Reidemeister moves; in the case $a = 0$, use the link from Lemma 6.5. The result

has a generating family g_a with the fiber indices indicated in Figure 11. Applying a 0-surgery along a horizontal line indicated in Figure 11 produces a gf-compatible Lagrangian cobordism from the Hopf link to a sphere Λ_a with generating family f_a .




Figure 11: A scheme to construct a Legendrian sphere with $\Gamma_{f_a}(t) = t^n + t^a + t^{n-1}t^{-a}$ with $a \geq \lfloor (n-1)/2 \rfloor$.

To finish the proof of the lemma, we verify that the generating family cohomology of (Λ_a, f_a) is given by $\Gamma_{f_a} = t^n + t^a + t^{n-1}t^{-a}$. If $(a, n-1-a) \neq (n-1, 0)$, then the 0-surgery lemma, Lemma 6.3, implies the desired result. If, on the other hand, we are in the case $(a, n-1-a) = (n-1, 0)$, the cobordism exact sequence implies that $\dim GH^0(f_0) = 1$ and that $\dim GH^k(f_a) = 0$ for all other $k \neq 0, n-1, n$. The remaining part of the cobordism exact sequence is

$$0 \rightarrow GH^{n-1}(f_0) \rightarrow \mathbb{F} \rightarrow \mathbb{F} \rightarrow GH^n(f_0).$$

Thus, $GH^{n-1}(f_0)$ and $GH^n(f_0)$ either simultaneously vanish or have dimension 1. Theorem 6.1 (2) implies that the latter is, indeed, the case, which completes the proof. □

Remark 6.11 Continuing Remark 6.9, we note that the spheres constructed in the lemma above are all fillable since they arise from attaching Lagrangian handles to fillable Legendrians.

From this lemma, we obtain spherical building blocks for duality classes:

Corollary 6.12 (Sphere building block) *For any $n \geq 2$ and any polynomial $p(t) = \sum_{i \in \mathbb{Z}, i \geq \lfloor (n-1)/2 \rfloor} p_i t^i$, there exists an n -dimensional Legendrian sphere with tame generating family f with polynomial*

$$\Gamma_f(t) = t^n + p(t) + t^{n-1}p(t^{-1}).$$

The corollary follows from [Lemma 6.10](#) in an analogous fashion to how [Corollary 6.7](#) followed from [Lemma 6.5](#) and [Corollary 6.4](#).

Finally, we have all the building blocks to answer the nonclassical generating family geography question.

Proof of Theorem 1.2 This proof follows from taking a connect sum of a manifold constructed in [Corollary 6.7](#) and a sphere constructed in [Corollary 6.12](#); once again, the generating family polynomial is computed using [Corollary 6.4](#). \square

6.3 Classical fillable geography

The constructions built up in the previous section, especially the sphere building block [Corollary 6.12](#), also suffice for the proof of [Theorem 1.3](#). We will also need the fact that the Thurston–Bennequin number of a Legendrian submanifold $\Lambda \subset J^1\mathbb{R}^n$ with generating family f may be computed by taking the Euler characteristic of the generating family homology:

$$(6-8) \quad tb(\Lambda) = (-1)^{(n-2)(n-1)/2} \Gamma_f(-1).$$

This fact was proven in [\[16\]](#) for linearized Legendrian contact homology, and can be translated to generating family homology using [\[39, Proposition 3.2\]](#).

Proof of Theorem 1.3 Fix an odd $n \geq 2$ and any odd integer $2k + 1$. Suppose that the quantity $(n - 2)(n - 1)/2$ is even; the proof in the other case is entirely similar. If $2k + 1 > 0$, it is straightforward to compute that the Legendrian sphere Λ with generating family polynomial

$$\Gamma_f(t) = t^n + (k + 1)t^{n-1} + (k + 1)t^0$$

has $tb(\Lambda) = 2k + 1$. If $2k + 1 < 0$, use the Legendrian sphere with generating family polynomial

$$\Gamma_f(t) = (|k| + 1)t^n + |k|t^{-1}.$$

By [Remark 6.11](#), all of the spheres used above are fillable, though only the first set is gf-fillable. This completes the proof of [Theorem 1.3](#). \square

Appendix: Proof of the duality theorem

We conclude this paper by proving [Theorem 6.1](#), repeating some of the proof of the original statement so as to properly set notation for the strengthening. The key ingredient in the proof of the duality exact sequence is [\[39, Lemma 7.1\]](#), which we restate here for the reader’s convenience:

Lemma A.1 [39] Assume $f: M^n \times \mathbb{R}^N \rightarrow \mathbb{R}$ is linear-at-infinity; let $\delta: M^n \times \mathbb{R}^{2N} \rightarrow \mathbb{R}$ be its associated difference function. For sufficiently large ω and for all $a \in \mathbb{R}$, there is an isomorphism

$$\beta: H^j(\delta^\omega, \delta^{-a}) \xrightarrow{\sim} H_{2N+n-j}(\delta^a, \delta^{-\omega}).$$

Proof of Theorem 6.1(1) The duality exact sequence is modeled on the cohomology long exact sequence of the triple $(\delta^\omega, \delta^\epsilon, \delta^{-\epsilon})$ for sufficiently large ω and sufficiently small $\epsilon > 0$; we have already encountered this triple in the long exact sequence (2-2). In fact, using this long exact sequence and the duality isomorphism from Lemma A.1, we obtain the following commutative diagram:

$$\begin{array}{ccccccc} \longrightarrow & H^{k+N}(\delta^\omega, \delta^\epsilon) & \xrightarrow{p_k} & H^{k+N}(\delta^\omega, \delta^{-\epsilon}) & \xrightarrow{s_k} & H^{k+N}(\delta^\epsilon, \delta^{-\epsilon}) & \xrightarrow{\delta_k} \longrightarrow \\ & \simeq \downarrow \alpha & & \simeq \downarrow \beta & & \simeq \downarrow \gamma & \\ \longrightarrow & H_{n-k+N}(\delta^{-\epsilon}, \delta^{-\omega}) & \longrightarrow & H_{n-k+N}(\delta^\epsilon, \delta^{-\omega}) & \xrightarrow{s'_{n-k}} & H_{n-k+N}(\delta^\epsilon, \delta^{-\epsilon}) & \longrightarrow \end{array}$$

We now identify several terms in the diagram above. By definition, $H^{k+N}(\delta^\omega, \delta^\epsilon) = GH^{k-1}(f)$; by the Thom isomorphism, $H^{k+N}(\delta^\epsilon, \delta^{-\epsilon}) \simeq H^k(\Lambda)$. To identify the middle term on the bottom line, observe that the term $H_{n-k+N}(\delta^\epsilon, \delta^{-\omega})$ is part of the homology long exact sequence of $(\delta^\omega, \delta^\epsilon, \delta^{-\omega})$. Owing to the fact that $H_*(\delta^\omega, \delta^{-\omega})$ vanishes, the connecting homomorphism $\partial_{n-k}: H_{n-k+N+1}(\delta^\omega, \delta^\epsilon) \rightarrow H_{n-k+N}(\delta^\epsilon, \delta^{-\omega})$ is an isomorphism for all k . Since, by definition, $GH_{n-k}(f) = H_{n-k+N+1}(\delta^\omega, \delta^\epsilon)$, we see that the middle bottom term is isomorphic, via ∂_{n-k} , to $GH_{n-k}(f)$.

The map $\rho_k: GH^{k-1}(f) \rightarrow GH_{n-k}(f)$ is defined as $\rho_k = \partial_{n-k}^{-1} \circ \beta \circ p_k$, and the map $\sigma_k: GH_{n-k}(f) \rightarrow H^k(\Lambda)$ is defined as $\sigma_k = s_k \circ \beta^{-1} \circ \partial_{n-k}$. Since field coefficients are used, the adjoint δ_{n-k}^* of ∂_{n-k} fits into the following commutative diagram, obtained from the long exact sequences of the triples $(\delta^\omega, \delta^\epsilon, \delta^{-\epsilon})$ and $(\delta^\omega, \delta^\epsilon, \delta^{-\omega})$:

$$\begin{array}{ccc} \longrightarrow & H_{n-k+N+1}(\delta^\omega, \delta^\epsilon) & \xrightarrow{\delta_{n-k}^*} & H_{n-k+N}(\delta^\epsilon, \delta^{-\epsilon}) & \longrightarrow \\ & \uparrow \text{id} & & \uparrow i_* = s'_{n-k} & \\ \longrightarrow & H_{n-k+N+1}(\delta^\omega, \delta^\epsilon) & \xrightarrow{\partial_{n-k}} & H_{n-k+N}(\delta^\epsilon, \delta^{-\omega}) & \longrightarrow \end{array}$$

Thus $\delta_{n-k}^* = (s'_{n-k}) \circ \partial_{n-k} = (\gamma \circ s_k \circ \beta^{-1}) \circ \partial_{n-k} = \gamma \circ \sigma_k$, as claimed. □

In order to prove the second part of Theorem 6.1, namely the existence of the fundamental class, we need to use a slight extension of the standard Morse–Bott machinery

described in [3]. We begin by recalling some of the basic objects in Morse–Bott theory to set notation. Fix a generating family f for Λ and consider its difference function δ and a Morse–Bott–Smale metric on the domain $M \times \mathbb{R}^{2N}$ of δ . Given nondegenerate critical points q and r of δ , we denote by $\mathcal{M}(q; r)$ the moduli space of gradient trajectories of δ from r to q modulo reparametrization by a constant shift. It is a smooth manifold of dimension $\text{Ind}_\delta q - \text{Ind}_\delta r - 1$. Let $\Sigma \subset M \times \Delta$ be the critical submanifold of δ in its zero level set. Similarly, we denote by $\mathcal{M}(q; \Sigma)$ the moduli space of gradient trajectories from Σ to q modulo reparametrization by a constant shift. It is a smooth manifold of dimension $\text{Ind}_\delta q - N - 1$. It is equipped with a smooth evaluation map $\text{ev}: \mathcal{M}(q; \Sigma) \rightarrow \Sigma$ at $-\infty$.

The key construction for the proof of [Theorem 6.1](#) (2) is a new moduli space of gradient flow lines that begin on the “diagonal” $M \times \Delta \subset M \times \mathbb{R}^N \times \mathbb{R}^N$. Given a nondegenerate critical point q of δ , we define $\mathcal{M}(q; M \times \Delta)$ to be the space of gradient flow lines $\gamma: [0, \infty) \rightarrow M \times \mathbb{R}^{2N}$ of δ with $\gamma(0) \in M \times \Delta$ and $\lim_{t \rightarrow \infty} \gamma(t) = q$. Finally, we define $\mathcal{M}(\Sigma; M \times \Delta)$ to be the space of gradient flow lines $\gamma: [0, \infty) \rightarrow M \times \mathbb{R}^{2N}$ of δ with $\gamma(0) \in M \times \Delta$ and $\lim_{t \rightarrow \infty} \gamma(t) \in \Sigma$. The moduli space $\mathcal{M}(\Sigma; M \times \Delta)$ consists entirely of constant trajectories, so we may identify the moduli space with Σ and its evaluation map at $+\infty$ with the identity.

Lemma A.2 *The moduli space $\mathcal{M}(q; M \times \Delta)$ is a smooth manifold of dimension $\text{Ind}_\delta q - N$. If it has dimension 1, it is compactified by the 0–dimensional moduli spaces*

$$\mathcal{M}(\Sigma; M \times \Delta) \times_\Sigma \mathcal{M}(q; \Sigma) \quad \text{and} \quad \bigcup_{\text{Ind}_\delta r = N} \mathcal{M}(r; M \times \Delta) \times \mathcal{M}(q; r)$$

so that each element in these moduli spaces is in the closure of a single end of $\mathcal{M}(q; M \times \Delta)$.

Proof The moduli space $\mathcal{M}(q; M \times \Delta)$ is the intersection of the stable manifold of q with $M \times \Delta$. After a small perturbation of the metric on $M \times \mathbb{R}^{2N}$, this intersection is transverse, so that $\mathcal{M}(q; M \times \Delta)$ is a smooth manifold of dimension $\text{Ind}_\delta q + (n + N) - (n + 2N) = \text{Ind}_\delta q - N$.

The compactification of the space of gradient trajectories $\gamma: \mathbb{R} \rightarrow M \times \mathbb{R}^N \times \mathbb{R}^N$ whose restriction to $[0, \infty)$ is in $\mathcal{M}(q; M \times \Delta)$ consists of broken gradient trajectories [3, Lemma 3.3]. Any such broken gradient trajectory contains a gradient trajectory passing through $M \times \Delta$ at $t = 0$, so that its restriction to $[0, \infty)$ is either an element of $\mathcal{M}(\Sigma; M \times \Delta)$ or an element of $\mathcal{M}(r; M \times \Delta)$ for some nondegenerate critical point r of δ . Note that the part of the broken trajectory passing through $M \times \Delta$ cannot be

a piece of a trajectory from a critical point with negative critical value to q since the image of the whole broken trajectory is symmetric under the exchange of the two \mathbb{R}^N factors. The compactification of $\mathcal{M}(q; M \times \Delta)$ therefore consists of broken trajectories with such an element followed by an ordinary broken gradient trajectory from Σ or r to q .

Let us first assume that the gradient trajectory passing through $M \times \Delta$ at $t = 0$ is an element of $\mathcal{M}(\Sigma; M \times \Delta)$. The next gradient trajectory in the broken gradient trajectory runs from Σ to a critical point s of δ with $\delta(s) > 0$, an element of the moduli space $\mathcal{M}(s; \Sigma)$ with dimension $\text{Ind}_\delta s - N - 1 \geq 0$. If $\text{Ind}_\delta q = N + 1$, this forces $s = q$ so that the broken trajectory is an element of $\mathcal{M}(\Sigma; M \times \Delta) \times_\Sigma \mathcal{M}(q; \Sigma)$. Conversely, the Morse–Bott gluing theorem [3, Theorem A.11] shows that any element of this moduli space is the limit of a unique end of $\mathcal{M}(q; M \times \Delta)$.

We now assume that the gradient trajectory passing through $M \times \Delta$ at $t = 0$ is an element of $\mathcal{M}(r; M \times \Delta)$. If $\text{Ind}_\delta q = N + 1$, there can only be one more gradient trajectory in the above broken gradient trajectory: an element of $\mathcal{M}(q; r)$ with $\text{Ind}_\delta r = N$. Conversely, the usual gluing theorem in Morse theory shows that each element of $\mathcal{M}(r; M \times \Delta) \times \mathcal{M}(q; r)$ is the limit of a unique end of $\mathcal{M}(q; M \times \Delta)$. □

With this technical lemma in hand, we are ready for the proof of the existence of a fundamental class.

Proof of Theorem 6.1(2) Let $\Lambda_1, \dots, \Lambda_k$ be the connected components of Λ and let p_i be the generator of $H^0(\Lambda_i)$ for $i = 1, \dots, k$. We claim that $p = \sum_{i=1}^k p_i \in \ker \delta_0$. In view of the property (1) proved above, this means that δ_n has rank at least 1.

Note that $\delta_0(p)$ results from the count of gradient trajectories with cascades [4] from a minimum of an auxiliary Morse function on Σ to a critical point q of the difference function with $\text{Ind} q = N + 1$. Such cascade gradient trajectories are in bijective correspondence with elements of $\mathcal{M}(q; \Sigma)$.

In order to prove our claim, define the cochain $c \in C^N(\delta^\omega, \delta^\epsilon)$ by

$$c = \sum_{\text{ind } r=N} \#\mathcal{M}(r; M \times \Delta) r.$$

Denoting the Morse codifferential of c by dc , the claim above follows from the assertion that $\delta_0(p) + dc = 0$. Lemma A.2 implies that this relation encodes a count of the boundary of the one-dimensional moduli space $\bigcup_{\text{Ind} q=N+1} \mathcal{M}(q; M \times \Delta)$: here $\delta_0(p)$ accounts for the first type of broken flow line in the lemma, while dc accounts for the second type. □

References

- [1] **A Abbondandolo, M Schwarz**, *On the Floer homology of cotangent bundles*, Comm. Pure Appl. Math. 59 (2006) 254–316 [MR2190223](#)
- [2] **M Abouzaid, P Seidel**, *An open string analogue of Viterbo functoriality*, Geom. Topol. 14 (2010) 627–718 [MR2602848](#)
- [3] **DM Austin, P J Braam**, *Morse–Bott theory and equivariant cohomology*, from “The Floer memorial volume” (H Hofer, C H Taubes, A Weinstein, E Zehnder, editors), Progr. Math. 133, Birkhäuser, Basel (1995) 123–183 [MR1362827](#)
- [4] **A Banyaga, D E Hurtubise**, *Cascades and perturbed Morse–Bott functions*, Algebr. Geom. Topol. 13 (2013) 237–275 [MR3031642](#)
- [5] **M Boileau, L Fourier**, *Knot theory and plane algebraic curves*, Chaos Solitons Fractals 9 (1998) 779–792 [MR1628756](#)
- [6] **B Boranda, L Traynor, S Yan**, *The surgery unknotting number of Legendrian links*, Involve 6 (2013) 273–299 [MR3101761](#)
- [7] **F Bourgeois, T Ekholm, Y Eliashberg**, *Effect of Legendrian surgery*, Geom. Topol. 16 (2012) 301–389 [MR2916289](#)
- [8] **C Cao, N Gallup, K Hayden, J M Sabloff**, *Topologically distinct Lagrangian and symplectic fillings*, Math. Res. Lett. 21 (2014) 85–99 [MR3247041](#)
- [9] **B Chantraine**, *Lagrangian concordance of Legendrian knots*, Algebr. Geom. Topol. 10 (2010) 63–85 [MR2580429](#)
- [10] **Y V Chekanov**, *Critical points of quasifunctions, and generating families of Legendrian manifolds*, Funktsional. Anal. i Prilozhen. 30 (1996) 56–69 [MR1402081](#)
- [11] **G Dimitroglou Rizell**, *Lifting pseudo-holomorphic polygons to the symplectisation of $P \times R$ and applications*, preprint [arXiv:1305.1572](#)
- [12] **G Dimitroglou Rizell**, *Legendrian ambient surgery and Legendrian contact homology*, preprint (2012) [arXiv:1205.5544v1](#)
- [13] **T Ekholm**, *Rational symplectic field theory over \mathbb{Z}_2 for exact Lagrangian cobordisms*, J. Eur. Math. Soc. (JEMS) 10 (2008) 641–704 [MR2421157](#)
- [14] **T Ekholm**, *A version of rational SFT for exact Lagrangian cobordisms in 1–jet spaces*, from “New perspectives and challenges in symplectic field theory” (M Abreu, F Lalonde, L Polterovich, editors), CRM Proc. Lecture Notes 49, Amer. Math. Soc. (2009) 173–199 [MR2555937](#)
- [15] **T Ekholm, J B Etnyre, J M Sabloff**, *A duality exact sequence for Legendrian contact homology*, Duke Math. J. 150 (2009) 1–75 [MR2560107](#)
- [16] **T Ekholm, J Etnyre, M Sullivan**, *Non-isotopic Legendrian submanifolds in \mathbb{R}^{2n+1}* , J. Differential Geom. 71 (2005) 85–128 [MR2191769](#)

- [17] **T Ekholm, K Honda, T Kálmán**, *Legendrian knots and exact Lagrangian cobordisms*, preprint [arXiv:1212.1519](https://arxiv.org/abs/1212.1519)
- [18] **T Ekholm, T Kálmán**, *Isotopies of Legendrian 1–knots and Legendrian 2–tori*, *J. Symplectic Geom.* 6 (2008) 407–460 [MR2471099](https://doi.org/10.1112/S1074748008000199)
- [19] **Y Eliashberg, A Givental, H Hofer**, *Introduction to symplectic field theory*, *Geom. Funct. Anal.* (2000) 560–673 [MR1826267](https://doi.org/10.1007/s00036-000-0026-7)
- [20] **Y Eliashberg, M Gromov**, *Lagrangian intersection theory: Finite-dimensional approach*, from “Geometry of differential equations” (A Khovanskii, A Varchenko, V Vassiliev, editors), *Amer. Math. Soc. Transl. Ser. 2* 186, Amer. Math. Soc. (1998) 27–118 [MR1732407](https://doi.org/10.1090/ps/1998/00186)
- [21] **M Entov**, *Surgery on Lagrangian and Legendrian singularities*, *Geom. Funct. Anal.* 9 (1999) 298–352 [MR1692478](https://doi.org/10.1007/s00036-000-0026-7)
- [22] **D Fuchs**, *Chekanov–Eliashberg invariant of Legendrian knots: existence of augmentations*, *J. Geom. Phys.* 47 (2003) 43–65 [MR1985483](https://doi.org/10.1016/S0167-8954(03)00048-3)
- [23] **D Fuchs, D Rutherford**, *Generating families and Legendrian contact homology in the standard contact space*, *J. Topol.* 4 (2011) 190–226 [MR2783382](https://doi.org/10.1017/S1446788711000072)
- [24] **K Fukaya, P Seidel, I Smith**, *The symplectic geometry of cotangent bundles from a categorical viewpoint*, from “Homological mirror symmetry” (A Kapustin, M Kreuzer, K-G Schlesinger, editors), *Lecture Notes in Phys.* 757, Springer, Berlin (2009) 1–26 [MR2596633](https://doi.org/10.1007/978-3-540-78663-3_1)
- [25] **R Golovko**, *A note on Lagrangian cobordisms between Legendrian submanifolds of \mathbb{R}^{2n+1}* , *Pacific J. Math.* 261 (2013) 101–116 [MR3037561](https://doi.org/10.1007/s12015-013-0101-1)
- [26] **R Golovko**, *A note on the front spinning construction*, *Bull. Lond. Math. Soc.* 46 (2014) 258–268
- [27] **K Hayden, J M Sabloff**, *Positive knots and Lagrangian fillability*, *Proc. Amer. Math. Soc.* 143 (2015) 1813–1821 [MR3314092](https://doi.org/10.1090/S0002-9947-2015-0154092-0)
- [28] **J Jordan, L Traynor**, *Generating family invariants for Legendrian links of unknots*, *Algebr. Geom. Topol.* 6 (2006) 895–933 [MR2240920](https://doi.org/10.1016/S1472-7480(06)00020-0)
- [29] **T Kálmán**, *Contact homology and one parameter families of Legendrian knots*, *Geom. Topol.* 9 (2005) 2013–2078 [MR2209366](https://doi.org/10.1017/S1446788705000366)
- [30] **P B Kronheimer, T S Mrowka**, *Gauge theory for embedded surfaces, I*, *Topology* 32 (1993) 773–826 [MR1241873](https://doi.org/10.1016/0165-1486(93)90036-3)
- [31] **P Melvin, S Shrestha**, *The nonuniqueness of Chekanov polynomials of Legendrian knots*, *Geom. Topol.* 9 (2005) 1221–1252 [MR2174265](https://doi.org/10.1017/S144678870500265)
- [32] **J Milnor**, *Morse theory*, *Annals of Mathematics Studies* 51, Princeton Univ. Press (1963) [MR0163331](https://doi.org/10.2307/2372331)

- [33] **E Murphy**, *Loose Legendrian embeddings in high dimensional contact manifolds*, preprint [arXiv:1201.2245](https://arxiv.org/abs/1201.2245)
- [34] **P E Pushkar', Y V Chekanov**, *Combinatorics of fronts of Legendrian links, and Arnol'd's 4-conjectures*, Uspekhi Mat. Nauk 60 (2005) 99–154 [MR2145660](https://arxiv.org/abs/1201.2245) In Russian; translated in Russian Math. Surveys 60 (2005) 95–149
- [35] **L Rudolph**, *Quasipositivity as an obstruction to sliceness*, Bull. Amer. Math. Soc. 29 (1993) 51–59 [MR1193540](https://arxiv.org/abs/1201.2245)
- [36] **L Rudolph**, *An obstruction to sliceness via contact geometry and "classical" gauge theory*, Invent. Math. 119 (1995) 155–163 [MR1309974](https://arxiv.org/abs/1201.2245)
- [37] **J M Sabloff**, *Duality for Legendrian contact homology*, Geom. Topol. 10 (2006) 2351–2381 [MR2284060](https://arxiv.org/abs/1201.2245)
- [38] **J Sabloff, M Sullivan**, *Families of Legendrian submanifolds via generating families* [arXiv:1311.0528](https://arxiv.org/abs/1311.0528) to appear in Quantum Topol.
- [39] **J M Sabloff, L Traynor**, *Obstructions to Lagrangian cobordisms between Legendrians via generating families*, Algebr. Geom. Topol. 13 (2013) 2733–2797 [MR3116302](https://arxiv.org/abs/1201.2245)
- [40] **L Traynor**, *Generating function polynomials for Legendrian links*, Geom. Topol. 5 (2001) 719–760 [MR1871403](https://arxiv.org/abs/1201.2245)

Laboratoire de Mathématiques d'Orsay, Université Paris-Sud,
Bâtiment 425, F-91405 Orsay, France

Department of Mathematics and Statistics, Haverford College
370 Lancaster Ave, Haverford, PA 19041, United States

Department of Mathematics, Bryn Mawr College
Bryn Mawr, PA 19010, United States

bourgeois@math.u-psud.fr, jsabloff@haverford.edu,
ltraynor@brynmawr.edu

<http://www.math.u-psud.fr/~bourgeois/>, <http://www.haverford.edu/math/jsabloff/>, <http://www.brynmawr.edu/math/Math/people/traynor/intro.html>

Received: 18 September 2014

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
School of Mathematics
Georgia Institute of Technology
Atlanta GA 30332-0160
USA

Kathryn Hess
kathryn.hess@epfl.ch
Département de mathématiques
École Polytechnique Fédérale de Lausanne
CH-1015 Lausanne
Switzerland

MANAGING EDITOR

Colin Rourke
agt@msp.warwick.ac.uk
Mathematics Institute
University of Warwick
Coventry CV4 7AL
UK

BOARD OF EDITORS

Matthew Ando	University of Illinois mando@math.uiuc.edu	Tsuyoshi Kobayashi	Nara Women's University tsuyoshi09@gmail.com
Joan Birman	Columbia University jb@math.columbia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Fred Cohen	University of Rochester cohf@math.rochester.edu	Andrés Navas Flores	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Tomotada Ohtsuki	Kyoto University tomotada@kurims.kyoto-u.ac.jp
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Robert Oliver	Université Paris-Nord bobol@math.univ-paris13.fr
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Luis Paris	Université de Bourgogne lparis@u-bourgogne.fr
Soren Galatius	Stanford University galatius@math.stanford.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Sheffield j.greenlees@sheffield.ac.uk	Peter Scott	University of Michigan pscott@umich.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Karen Vogtmann	Cornell University vogtmann@math.cornell.edu
Jesse Johnson	Oklahoma State University jjohnson@math.okstate.edu	Daniel T. Wise	McGill University, Canada daniel.wise@mcgill.ca
Mikhail Khovanov	Columbia University khovanov@math.columbia.edu		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2015 is US \$350/year for the electronic version, and \$575/year (+ \$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP. Algebraic & Geometric Topology is indexed by *Mathematical Reviews*, *Zentralblatt MATH*, *Current Mathematical Publications* and the *Science Citation Index*.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) at Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published 6 times per year and continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 15

Issue 4 (pages 1863–2477)

2015

The Chillingworth class is a signed stable length INGRID IRMER	1863
Stable immersions in orbifolds ALDEN WALKER	1877
Moments of a length function on the boundary of a hyperbolic manifold NICHOLAS G VLAMIS	1909
Rectification of enriched ∞ -categories RUNE HAUGSENG	1931
Analytic families of quantum hyperbolic invariants STÉPHANE BASEILHAC and RICCARDO BENEDETTI	1983
Completed power operations for Morava E -theory TOBIAS BARTHEL and MARTIN FRANKLAND	2065
On the slice-ribbon conjecture for pretzel knots ANA G LECUONA	2133
Braids, complex volume and cluster algebras KAZUHIRO HIKAMI and REI INOUE	2175
Ellis enveloping semigroup for almost canonical model sets of an Euclidean space JEAN-BAPTISTE AUJOGUE	2195
A relative Lubin–Tate theorem via higher formal geometry AARON MAZEL-GEE, ERIC PETERSON and NATHANIEL STAPLETON	2239
Positive curvature and rational ellipticity MANUEL AMANN and LEE KENNARD	2269
Quasi-unital ∞ -categories YONATAN HARPAZ	2303
An alternative approach to extending pseudo-Anosovs over compression bodies ROBERT ACKERMANN	2383
The fundamental group and Betti numbers of toric origami manifolds TARA S HOLM and ANA RITA PIRES	2393
Resolving rational cohomological dimension via a Cantor group action MICHAEL LEVIN	2427
Lagrangian cobordisms via generating families: Construction and geography FRÉDÉRIC BOURGEOIS, JOSHUA M SABLOFF and LISA TRAYNOR	2439