

Semistability and simple connectivity at ∞ of finitely generated groups with a finite series of commensurated subgroups

MICHAEL L MIHALIK

A subgroup H of a group G is *commensurated* in G if for each $g \in G$, $gHg^{-1} \cap H$ has finite index in both H and gHg^{-1} . If there is a sequence of subgroups $H = Q_0 < Q_1 < \cdots < Q_k < Q_{k+1} = G$ where Q_i is commensurated in Q_{i+1} for all i , then Q_0 is *subcommensurated* in G . In this paper we introduce the notion of the simple connectivity at ∞ of a finitely generated group (in analogy with that for finitely presented groups). Our main result is this: if a finitely generated group G contains an infinite finitely generated subcommensurated subgroup H of infinite index in G , then G is one-ended and semistable at ∞ . If, additionally, G is recursively presented and H is finitely presented and one-ended, then G is simply connected at ∞ . A normal subgroup of a group is commensurated, so this result is a strict generalization of a number of results, including the main theorems in works of G Conner and M Mihalik, B Jackson, V M Lew, M Mihalik, and J Profio. We also show that Grigorchuk's group (a finitely generated infinite torsion group) and a finitely presented ascending HNN extension of this group are simply connected at ∞ , generalizing the main result of a paper of L Funar and DE Otera.

[20F65](#); [20F69](#), [57M10](#)

1 Introduction and background

In 1962, J Stallings defined what it means for a space to be n -connected at ∞ , and proved the following:

Theorem 1.1 (Stallings [23]) *If V^n , $n \geq 5$, is a contractible PL n -manifold without boundary, then V is PL homeomorphic to R^n if and only if V is simply connected at ∞ .*

In 1974, R Lee and F Raymond first considered the fundamental group of an end of a group. In particular, they considered groups that are simply connected at ∞ .

Theorem 1.2 (Lee and Raymond [14]) *Let G be a finitely presented group with normal subgroup N isomorphic to \mathbb{Z}^k and quotient $K = G/N$. Assume when $k = 1$ that K is one-ended and that when $k = 2$ that K is not finite, and no restrictions when $k > 2$. Then G is simply connected at ∞ .*

For a reasonable space X (or finitely presented group G), one needs to know that X (respectively G) is semistable at ∞ in order to have the fundamental group of an end of X (respectively G) defined independent of base ray. In 1982, B Jackson generalized [Theorem 1.2](#) and in 1983, M Mihalik proved the first semistability at ∞ theorem for a class of finitely presented groups. These two results serve as a starting point for this paper.

Theorem 1.3 (Jackson [11]) *If H is an infinite finitely presented normal subgroup of infinite index in the finitely presented group G , and either H or G/H is one-ended, then G is simply connected at ∞ .*

Theorem 1.4 (Mihalik [16]) *If H is an infinite finitely generated normal subgroup of infinite index in the finitely presented group G , then G is semistable at ∞ .*

In 1985, the following connections were drawn between semistability and simple connectivity at ∞ , and group cohomology.

Theorem 1.5 (Geoghegan and Mihalik [6]) *If G is a finitely presented and semistable at ∞ group then $H^2(G, \mathbb{Z}G)$ is free abelian. If G is simply connected at ∞ then $H^2(G, \mathbb{Z}G) = 0$.*

It is unknown whether or not all finitely presented groups are semistable at ∞ . It is also unknown whether or not for all finitely presented groups G , $H^2(G, \mathbb{Z}G)$ is free abelian. The main theorem in the unpublished 1993 PhD dissertation of V Ming Lew generalized [Theorem 1.4](#) and the main theorem of the 1990 PhD dissertation of J Profio generalized [Theorem 1.3](#).

Theorem 1.6 (Lew [15]) *Suppose H is an infinite finitely generated subnormal subgroup of the finitely generated group G , then*

$$H = N_0 \triangleleft N_1 \triangleleft N_2 \triangleleft \cdots \triangleleft N_k = G$$

for $k \geq 1$, and H has infinite index in G . Then G is one-ended and semistable at ∞ .

Theorem 1.7 (Profio [21]) *Suppose $H \triangleleft N \triangleleft G$ is a normal series with H and G finitely presented, and H one-ended and of infinite index in G . Then G is simply connected at ∞ .*

Given a subgroup H of a group G , the element $g \in G$ is in the *commensurator* of H in G (denoted $\text{Comm}(H, G)$) if $gHg^{-1} \cap H$ has finite index in both H and gHg^{-1} . The subgroup H is *commensurated* in G if $\text{Comm}(H, G) = G$, so normal subgroups are commensurated. The main result of G Conner and Mihalik [3] generalizes Theorems 1.4 and 1.3 in a direction different than these last two results.

Theorem 1.8 (Conner and Mihalik [3]) *If a finitely generated group G has an infinite finitely generated commensurated subgroup Q , and Q has infinite index in G , then G is one-ended and semistable at ∞ . Furthermore, if G and Q are finitely presented and either Q is one-ended or the pair (G, Q) has one filtered end, then G is simply connected at ∞ .*

Example For p a prime, the group $\text{SL}_n(\mathbb{Z}[\frac{1}{p}])$ is finitely presented. When $n > 2$, the only normal subgroups of this group are either finite or of finite index; see [22]. For $n > 2$, the finitely presented one-ended subgroup $\text{SL}_n(\mathbb{Z})$ is commensurated in $\text{SL}_n(\mathbb{Z}[\frac{1}{p}])$ and so by Theorem 1.8, $\text{SL}_n(\mathbb{Z}[\frac{1}{p}])$ is one-ended and simply connected at ∞ .

While Lew's theorem improves Theorem 1.4 by replacing normality by subnormality, Profio's result is the best attempt in the last 30 years to improve the normality hypothesis of Theorem 1.3 to subnormality. As a corollary of our main theorem, we obtain the subnormal version of Jackson's Theorem 1.3. The semistability part of Theorem 1.9 is proved first and then used in an essential way in the proof of the simply connected at ∞ part of Theorem 1.9. A new idea, the simple connectivity at ∞ of a finitely generated group, is introduced and used in a fundamental way to prove the second part of Theorem 1.9. We point out that we cannot prove this part of Theorem 1.9, even in the finitely presented case, without this new concept.

If Q is a commensurated subgroup of G we use the notation $Q < G$. The main theorem of this article is the following.

Theorem 1.9 (main theorem) *Suppose H is a finitely generated infinite subgroup of infinite index in the finitely generated group G , and H is subcommensurated in G :*

$$H = Q_0 < Q_1 < \cdots < Q_k < G.$$

Then G is one-ended and semistable at ∞ . If, additionally, H is one-ended and finitely presented and G is finitely generated and recursively presented, then G is simply connected at ∞ .

In the next section we define what it means for a finitely generated (and recursively presented) group to be simply connected at ∞ (a strict generalization of simple connectivity at ∞ for finitely presented groups).

Example In Mihalik [20], short exact sequences are produced for each $n > 0$ of the form

$$1 \rightarrow H \rightarrow (\mathbb{Z}^n * \mathbb{Z}) \times (\mathbb{Z}^n * \mathbb{Z}) \rightarrow \mathbb{Z}^n \rightarrow 1,$$

where H is one-ended and finitely generated. The group \mathbb{Z}^n is $(n-2)$ -connected at ∞ , yet $(\mathbb{Z}^n * \mathbb{Z}) \times (\mathbb{Z}^n * \mathbb{Z})$ is not simply connected at ∞ . These examples show that the finitely presented hypothesis on H in Theorems 1.3 and 1.9, cannot be easily relaxed.

R Grigorchuk constructed a finitely generated infinite torsion group G and a finitely presented HNN extension H of G ; see Grigorchuk [7; 8]. The group G contains a subgroup of finite index T , and T is isomorphic to $T \times T$; see de la Harpe [9, Chapter VIII, Theorem 28]. We show that a recursively presented direct product of a one-ended finitely generated group with an infinite finitely generated group is simply connected at ∞ , and so Grigorchuk's group G is simply connected at ∞ . We also show that an ascending HNN extension of a one-ended finitely generated recursively presented semistable at ∞ and simply connected at ∞ group is simply connected at ∞ . This implies that the HNN extension of Grigorchuk's group is simply connected at ∞ .

The remainder of the paper is organized as follows. In Section 2, the working definitions and notation are established. We introduce our definition of a subgroup being simply connected at ∞ inside an overgroup. This definition is then used to define the simple connectivity at ∞ of a finitely generated group. We end Section 2 with a few important technical lemmas.

In Section 3, we prove the semistability part of our main theorem. This is an induction argument that starts with base case given by Theorem 1.8.

In Section 4, we prove the simple connectivity at ∞ part of our main theorem. This is also an induction argument that starts with the base case given by the simple connectivity part of Theorem 1.8. The semistability result of Section 3 is used in conjunction with Lemma 2.9 to set up the proof of the simple connectivity part of Theorem 1.9.

Finally, in Section 5 we prove two simply connected at ∞ results, and apply the results to show Grigorchuk's group and its HNN extension are simply connected at ∞ .

2 Definitions and a technical lemma

Geoghegan's book [5] is a general reference to all that is in this section. A continuous function $f: X \rightarrow Y$ is *proper* if for each compact subset C of Y , $f^{-1}(C)$ is compact in X . A proper map $r: [0, \infty) \rightarrow X$ is called a *ray* in X . If K is a locally finite, connected CW-complex, then one can define an equivalence relation \sim on the set A of all rays in K by setting $r \sim s$ if and only if for each compact set $C \subset K$, there exists an integer $N(C)$ such that $r([N(C), \infty))$ and $s([N(C), \infty))$ are contained in the same unbounded path component of $K - C$ (a path component of $K - C$ is *unbounded* if it is not contained in any compact subset of K). An equivalence class of A/\sim is called *an end of K* , the set of equivalence classes of A/\sim is called *the set of ends of K* , and two rays in K in the same equivalence class are said to *converge to the same end*. The cardinality of A/\sim , denoted by $e(K)$, is the *number of ends of K* .

If G is a finitely generated group with generating set S , then the *Cayley graph of G with respect to S* , denoted $\Gamma_{(G,S)}$, has vertex set G and an edge between vertices v and w if $vs = w$ for some $s \in S$. We define the *number of ends of G* , denoted by $e(G)$, to be the number of ends of the Cayley graph of G with respect to a finite generating set. (In particular, $e(G) = e(\Gamma_{(G,S)})$.) This definition is independent of the choice of finite generating set for G . If G is finitely generated, then $e(G)$ is either 0, 1, 2, or is infinite (in which case it has the cardinality of the real numbers). We let $*$ denote the base point of $\Gamma_{(G,S)}$, which corresponds to the identity of G .

If f and g are rays in K , then one says that f and g are *properly homotopic* if there is a proper map $H: [0, 1] \times [0, \infty) \rightarrow K$ such that $H|_{\{0\} \times [0, \infty)} = f$ and $H|_{\{1\} \times [0, \infty)} = g$. If $f(0) = g(0) = v$ and $H|_{[0, 1] \times \{0\}} = v$, one says f and g are *properly homotopic relative to v* (or $\text{rel}\{v\}$).

Definition 2.1 A locally finite connected CW-complex K is *semistable at ∞* if any two rays in K converging to the same end are properly homotopic. The space K is *simply connected at ∞* if for any compact set $C \subset K$ there is a compact $D \subset K$ such that loops in $K - D$ are homotopically trivial in $K - C$.

In a locally finite CW-complex, any ray is properly homotopic to an edge path ray. So in order to show semistability in such a complex, it is enough to prove edge path rays converging to the same end are properly homotopic.

[16, Theorem 2.1] and [18, Lemma 9] provide several equivalent notions of semistability. The space considered in [16] is simply connected, but simple connectivity is not important in that argument. A slight modification of proofs gives the following result; see [3].

Theorem 2.2 *Suppose K is a locally finite connected one-ended CW-complex. Then the following are equivalent:*

- (1) K is semistable at ∞ .
- (2) For any ray $r: [0, \infty) \rightarrow K$ and compact set C , there is a compact set D such that for any third compact set E and loop α based on r and with image in $K - D$, α is homotopic rel $\{r\}$ to a loop in $K - E$, by a homotopy with image in $K - C$.
- (3) For some (equivalently any) ray r in K and any collection of compact sets C_i such that $\bigcup_{i=1}^{\infty} C_i = K$ and C_{i-1} is a subset of the interior of C_i , the inverse system

$$\pi_1(X - C_1, r) \leftarrow \pi_1(X - C_2, r) \leftarrow \dots$$

with bonding maps induced by inclusion along r , is proisomorphic to an inverse system of groups with epimorphic bonding maps.

- (4) For any compact set C there is a compact set D such that if r and s are rays based at v and with image in $K - D$, then r and s are properly homotopic rel $\{v\}$ by a proper homotopy in $K - C$.

If K is simply connected (or if a group acting by homeomorphisms on K , acts transitively on the vertices of K) then a fifth equivalent condition can be added to this list:

- (5) If r and s are rays based at v , then r and s are properly homotopic rel $\{v\}$.

If finite connected CW-complexes X and Y have isomorphic fundamental groups, then the universal cover of X is semistable (simply connected) at ∞ if and only if the universal cover of Y is semistable (simply connected) at ∞ . This result can be seen from the early work of F E A Johnson [12; 13], or the proof of [14, Theorem 3]. For a complete argument see the first three sections of [5, Chapter 5].

Definition 2.3 If G is a one-ended, finitely presented group and X is some (equivalently any) finite CW-complex with fundamental group G , then we say G is *semistable at ∞* if the universal cover of X is semistable at ∞ . We say G is *simply connected at ∞* if the universal cover of X is simply connected at ∞ .

The notion of semistability for a finitely generated group was first defined in [19]. We give the definition for one-ended groups since this is the case that concerns us. Suppose G is a one-ended finitely generated group with generating set $\mathcal{S} := \{g_1, g_2, \dots, g_n\}$ and let $\Gamma_{(G, \mathcal{S})}$ be the Cayley graph of G with respect to this generating set. Suppose $\{\alpha_1, \alpha_2, \dots, \alpha_m\}$ is a finite set of relations in G written in the letters $\{g_1^{\pm}, g_2^{\pm}, \dots, g_n^{\pm}\}$.

For any vertex $v \in \Gamma_{(G,S)}$, there is an edge path cycle labeled α_i at v . The 2-dimensional CW-complex $\Gamma_{(G,S)}(\alpha_1, \dots, \alpha_m)$ is obtained by attaching, at each vertex of $\Gamma_{(G,S)}$, 2-cells corresponding to the relations $\alpha_1, \dots, \alpha_n$.

We show in [19] that if \mathcal{S} and \mathcal{T} are finite generating sets for the group G and there are finitely many \mathcal{S} -relations P such that $\Gamma_{(G,S)}(P)$ is semistable at ∞ , then there are finitely many \mathcal{T} -relations Q such that $\Gamma_{(G,T)}(Q)$ is semistable at ∞ ; hence the following definition.

Definition 2.4 A finitely generated group G is *semistable at ∞* if for some (equivalently any) finite generating set \mathcal{S} for G and finite set of \mathcal{S} -relations P the complex $\Gamma_{(G,S)}(P)$ is semistable at ∞ .

Note that if G has finite presentation $\langle \mathcal{S} : P \rangle$, then G is semistable at ∞ with respect to Definition 2.3 if and only if G is semistable at ∞ with respect to Definition 2.4 if and only if $\Gamma_{(G,S)}(P)$ is semistable at ∞ .

The following definition defines what it means for a finitely generated subgroup of a finitely presented group to be simply connected at ∞ relative to the finitely presented overgroup.

Definition 2.5 A finitely generated subgroup A of a finitely presented group G is *simply connected at ∞ in G* (or *relative to G*) if for some (equivalently any by Lemma 2.9 with $N = 0$) finite presentation $\langle \mathcal{A}, \mathcal{B} : R \rangle$ of the group G (where \mathcal{A} generates A and $\mathcal{A} \cup \mathcal{B}$ generates G), the 2-complex $\Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R)$ has the following property: given any compact set $C \subset \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R)$, there is a compact set $D \subset \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R)$ such that any edge path loop in $\Gamma_{(A, \mathcal{A})} - D$ is homotopically trivial in $\Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) - C$.

In order to define what it means for a finitely generated group G to be simply connected at ∞ , we must know that G embeds in some finitely presented group. In 1961, G Higman proved the following.

Theorem 2.6 (Higman [10]) *A finitely generated infinite group G can be embedded in a finitely presented group if and only if G is recursively presented.*

Definition 2.7 A finitely generated and recursively presented group A is *simply connected at ∞* if for any finitely presented group G and subgroup A' isomorphic to A , A' is simply connected at ∞ in G .

Suppose that A is a finitely presented group and A satisfies the simply connected at ∞ condition of Definition 2.3, then A satisfies Definition 2.7. If A is a finitely presented group satisfying Definition 2.7, then if we let $A = G$ in Definition 2.7, we see that A satisfies Definition 2.3, and there is no ambiguity.

We conclude this section with Lemmas 2.9 and 2.10, but first some terminology. Suppose $\langle S : R \rangle$ is a finite presentation for a group G . If A is a subcomplex of $\Gamma_{(G,S)}(R)$, then $\text{St}(A)$ is the subcomplex of $\Gamma_{(G,S)}(R)$ whose vertices $V(\text{St}(A))$ are the vertices of A along with each vertex of $\Gamma_{(G,S)}(R)$ that is connected to a vertex of A by an edge. The edges $E(\text{St}(A))$ of $\text{St}(A)$ are all edges of A union all edges of $\Gamma_{(G,S)}(R)$, both of whose vertices are contained in $V(\text{St}(A))$. The 2-cells $F(\text{St}(A))$ of $\text{St}(A)$ are all 2-cells of A union all 2-cells F such that all vertices of F belong to $V(\text{St}(A))$. If A is an arbitrary subset of $\Gamma_{(G,S)}(R)$ then let \hat{A} be the smallest subcomplex of $\Gamma_{(G,S)}(R)$ containing A and define $\text{St}(A)$ to be $\text{St}(\hat{A})$. Inductively, $\text{St}^L(A) = \text{St}(\text{St}^{L-1}(A))$ for $L > 1$.

- Lemma 2.8** (1) Suppose A and B are subcomplexes of $\Gamma_{(G,S)}(R)$ and $\text{St}(A) \cap B \neq \emptyset$. Then $A \cap \text{St}(B) \neq \emptyset$.
- (2) Suppose A is a subcomplex of $\Gamma_{(G,S)}(R)$ and B is an arbitrary subset of $\Gamma_{(G,S)}(R)$ and $\text{St}(B) \cap A \neq \emptyset$ then $\text{St}^{L+1}(A) \cap B \neq \emptyset$, where L is the length of the longest relation in R .

Proof Case (1) If $\text{St}(A) \cap B \neq \emptyset$ then there is a vertex $v \in \text{St}(A) \cap B$. If v is in A then we are finished. Otherwise, there is a vertex $w \in A$ and an edge from v to w . Then $w \in A \cap \text{St}(B)$.

Case (2) Let v be a vertex in $\text{St}(B) \cap A = \text{St}(\hat{B}) \cap A$. If v is in B we are finished. Otherwise, v is in \hat{B} or v is adjacent to a vertex $w \in \hat{B}$. If v is in \hat{B} then there is an edge e containing a point $b \in B$ and v is a vertex of e , or there is a 2-cell F containing a point $b \in B$ and v is a vertex of F . In either case, b is in $\text{St}^L(v)$, so b is in $B \cap \text{St}^L(A)$. If v is adjacent to a vertex $w \in \hat{B}$ then as above, there is $b \in B \cap \text{St}^L(w) \subset B \cap \text{St}^{L+1}(A)$. □

The following technical lemma has a somewhat standard proof.

Lemma 2.9 Suppose A is a finitely generated subgroup of the finitely presented group G . Then A is simply connected at ∞ in G if and only if the following holds:

- (†) For $\langle S : R \rangle$ an arbitrary finite presentation for G , $N \geq 0$ an integer and C a compact subset of $\Gamma_{(G,S)}(R)$, there is a compact set $D(C, N) \subset \Gamma$ such that if α is an edge path loop in $\Gamma - D$ and each vertex of α is within N of some vertex of $A (\subset \Gamma)$, then α is homotopically trivial in $\Gamma - C$.

Proof If condition (\dagger) holds with $N = 0$ then clearly A is simply connected at ∞ . For the converse, assume A is simply connected at ∞ and $\langle \mathcal{A}, \mathcal{B} : T \rangle$ is a presentation for G satisfying the conditions of Definition 2.5. Define $\Gamma_1 := \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(T)$ and $\Gamma_2 := \Gamma_{(G, \mathcal{S})}(R)$. Recall that the vertices of Γ_1 and of Γ_2 are both the elements of G . In order to avoid confusion if v is a vertex of Γ_1 we denote by v' the corresponding vertex of Γ_2 . We define proper maps respecting the action of G , $f_1: \Gamma_1 \rightarrow \Gamma_2$ and $f_2: \Gamma_2 \rightarrow \Gamma_1$ such that for each vertex $g \in G$ of Γ_1 , $f_1(g) = g'$ and $f_2(g') = g$. If e is an edge of Γ_1 with initial vertex v , terminal vertex w and label $s \in \mathcal{A} \cup \mathcal{B}$, then choose an edge path τ_s in Γ_2 from $v' := f_1(v)$ to $w' := f_1(w)$. Define $f_1(e)$ to be τ_s . If $g \in G$ define f_1 on ge to be $g\tau_s$. Similarly define f_2 from the 1-skeleton of Γ_2 to the 1-skeleton of Γ_1 . Let M_1 be the length of the longest path τ_s for $s \in \mathcal{A} \cup \mathcal{B}$ and M_2 be the length of the longest path $\tau_{s'}$ for $s' \in \mathcal{S}$. Note that if e is an edge of Γ_1 , with initial vertex v and terminal vertex w , then $f_2 f_1(e)$ is an edge path of length at most $M_1 M_2$ from v to w , and similarly if e is an edge of Γ_2 .

In particular, if x is a point of an edge of Γ_1 then $f_2(f_1(x)) \in \text{St}^{M_1 M_2}(x)$, and likewise if x belongs to an edge of Γ_2 .

If F is a 2-cell of Γ_1 , then the boundary of F is an edge path β_F with edge labels the same as an element of T . Then $f_1(\beta_F)$ is an edge path loop in Γ_2 . Choose $P_1 > 0$ so that if F is any 2-cell of Γ_1 , then the edge path loop $f_1(\beta_F)$ is homotopically trivial in $\text{St}^{P_1}(v')$ for any vertex v' of $f_1(\beta_F)$. The map f_1 is defined so that $f_1|_F$ (the restriction of f_1 to any 2-cell F) realizes this homotopy and respects the action of G on Γ_1 and Γ_2 . Similarly map the 2-cells of Γ_2 to Γ_1 and choose P_2 for f_2 . Let L be the length of the longest relator of $T \cup R$.

If x is a point of a 2-cell F of Γ_1 and v is a vertex of F , then $f_1(x) \in \text{St}^{P_1}(f_1(v)) = \text{St}^{P_1}(v')$. This means there is an edge path τ in Γ_2 of length at most P_1 from v' to a vertex w' and w' belongs to an edge b or 2-cell B containing $f_1(x)$.

If w' belongs to an edge b then $f_2(f_1(x))$ belongs to $f_2((\tau, b))$ an edge path of length at most $M_2(P_1 + 1)$ that begins at v . In this case $f_2(f_1(x)) \in \text{St}^{M_2(P_1+1)}(x)$.

Otherwise, w' belongs to a 2-cell B containing $f_1(x)$ and $f_2(f_1(x))$ belongs to $\text{St}^{P_2}(f_2(w')) = \text{St}^{P_2}(w)$. Then $f_2(\tau)$ is an edge path of length at most $P_1 M_2$ from v to w , and $f_2 f_1(x) \in \text{St}^{P_1 M_1 + P_2}(v)$. As $x \in \text{St}^L(v)$, we have $f_2(f_1(x)) \in \text{St}^{P_1 M_1 + P_2 + L}(x)$. Combining, we make the following claim.

Claim 1 *There is an integer M such that if x is a point of Γ_1 (respectively Γ_2), then $f_2(f_1(x)) \in \text{St}^M(x)$ (respectively $f_1(f_2(x)) \in \text{St}^M(x)$).*

Let $\Gamma_3 := \Gamma_{(\mathcal{A}, \mathcal{A})}$ be the corresponding subgraph of Γ_1 . Then for any compact set C in Γ_1 there is a compact set D in Γ_1 so that any edge path loop in $\Gamma_3 - D$, is

homotopically trivial in $\Gamma_1 - C$. Let $\Gamma_4 := f_1(\Gamma_3)$. Then A is a subset of the vertices of Γ_4 and we call these vertices the pseudo vertices of Γ_4 . For each edge e of Γ_3 , $f_1(e)$ is an edge path of Γ_4 (connecting two pseudo vertices) that we call a pseudo edge of Γ_4 .

Claim 2 *Given a compact set C in Γ_2 , there is a compact set $D_1(C)$ in Γ_2 such that any pseudo edge path loop β in $\Gamma_4 - D_1$ is homotopically trivial in $\Gamma_2 - C$.*

Proof Assume C is a compact subcomplex of Γ_2 . Then $\text{St}^{M+L}(f_2(C))$ is a compact subcomplex of Γ_1 . As Γ_1 satisfies [Definition 2.5](#), there is a compact subcomplex E of Γ_1 such that any edge path loop in $\Gamma_3 - E$ is homotopically trivial in $\Gamma_1 - \text{St}^{M+L}(f_2(C))$.

Choose D_1 a compact subcomplex of Γ_2 such that if $w \in G$ is a vertex of E then $f_2(w) := w' \in D_1$. If β' is a pseudo edge path loop in $\Gamma_4 - D_1$, let β be an edge path loop in Γ_3 such that $f_1(\beta) = \beta'$. Note that no vertex of β belongs to E and so β avoids E . Then there is a homotopy H that kills β in $\Gamma_1 - \text{St}^{M+L}(f_2(C))$ and f_1H kills β' in Γ_2 . It remains to show that the image of f_1H avoids C . If $\text{im}(f_1H) \cap C \neq \emptyset$, then $\text{im}(f_2 f_1 H) \cap f_2(C) \neq \emptyset$. By [Claim 1](#), $\text{im}(f_2 f_1 H) \subset \text{St}^M(\text{im}(H))$ and so $\text{St}^M(\text{im}(H)) \cap f_2(C) \neq \emptyset$. By [Lemma 2.8\(1\)](#), $\text{St}(\text{im}(H)) \cap \text{St}^{M-1}(f_2(C)) \neq \emptyset$ and by [Lemma 2.8\(2\)](#) $\text{im}(H) \cap \text{St}^{M+L}(f_2(C)) \neq \emptyset$. But $\text{im}(H) \cap \text{St}^{M+L}(f_2(C)) = \emptyset$. □

Now we complete the proof of [Lemma 2.9](#). Recall, $N \geq 0$ is an arbitrary fixed integer. Choose N_1 such that if two pseudo vertices of Γ_4 are within $2N + 1$ of one another in Γ_2 then there is a pseudo edge path of Γ_2 -length at most N_1 connecting them. Let C be compact in Γ_2 . Choose N_2 so that if τ is an edge path loop in Γ_2 of length at most $N_1 + 2N + 1$, then τ is homotopically trivial in $\text{St}^{N_2}(w')$ for any vertex w' of τ . Now suppose α is an edge path loop of $\Gamma_2 - \text{St}^{N_2}(D_1(C))$ and each vertex of α is within N of A (the pseudo vertices of Γ_4). By the definition of N_2 , α is homotopic to a pseudo edge path loop α' in $\Gamma_4 - D_1$ by a homotopy in $\Gamma_2 - D_1$. Since α' is homotopically trivial in $\Gamma_2 - C$, α is as well. □

Remark For spaces (and finitely presented groups), simple connectivity at ∞ is stronger than semistability at ∞ . In fact [Theorem 2.2\(3\)](#) states that a space K is semistable at ∞ if and only if $\text{pro} - \pi_1^{\text{end}}(K)$ is proisomorphic to an inverse system of groups with epimorphic bonding maps. The space K is simply connected at ∞ if and only if $\text{pro} - \pi_1^{\text{end}}(K)$ is protrivial. It is not clear whether or not our definition of simple connectivity at ∞ for a finitely generated group implies the group is semistable at ∞ .

Interestingly, these two notions can be combined in effective ways (see [Lemma 2.10](#) and [Theorem 5.2](#)).

[Lemma 2.9](#) implies the following. Suppose the finitely generated group A is simply connected at ∞ in the finitely presented group G , (S, R) is a finite presentation for G , and v_1, \dots, v_n are vertices of $\Gamma_{(G,S)}(R)$. Then for any compact $C \subset \Gamma$ and integer $N \geq 0$ there is a compact set $D(C, N, \{v_1, \dots, v_n\})$ such that any loop in $\Gamma - D$, each of whose vertices is within N of $v_i A$ for some $i \in \{1, \dots, n\}$, is homotopically trivial in $\Gamma - C$. What is not guaranteed is a compact set $D(C, N)$ satisfying the following: for all $v \in G$ and any edge path loop α in $\Gamma - D$ with each vertex of α within N of vA , the loop α is homotopically trivial in $\Gamma - C$. We do gain this enhanced condition (see [Lemma 2.10](#)) if A is both simply connected at ∞ in G and semistable at ∞ .

Lemma 2.10 *Suppose A is a finitely generated semistable at ∞ group and A is simply connected at ∞ in the finitely presented group G , then the following holds:*

- (\ddagger) *For $\langle S : R \rangle$ an arbitrary finite presentation for G , $N \geq 0$ an integer and C a compact subcomplex of $\Gamma_{(G,S)}(R)$, there is a compact set $D(C, N) \subset \Gamma$ such that if α is an edge path loop in $\Gamma - D$, v is an element of G , and each vertex of α is within N of the coset $vA (\subset \Gamma)$, then α is homotopically trivial in $\Gamma - C$.*

Proof By [Lemma 2.9](#), condition (\ddagger) holds for any finite collection of vertices v . For the sake of simplicity, we assume that S contains a set of generators \mathcal{A} for A , and R contains a finite set of \mathcal{A} -relations $R_{\mathcal{A}}$ so that $\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}})$ is semistable at ∞ . Choose an integer M so that if α is an edge path loop in $\Gamma_{(G,S)}(R)$ and each vertex of α is within N of $A \subset \Gamma_{(G,S)}(R)$ then α is homotopic to an edge path loop α' in $\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}})$ by a homotopy in $\text{St}^M(\text{image}(\alpha)) \subset \Gamma_{(G,S)}(R)$. Choose a finite set of vertices v_1, \dots, v_n such that if $v\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}}) \cap \text{St}^M(C) \neq \emptyset$ then $vA = v_i A$ for some $i \in \{1, \dots, n\}$. Choose $D(\text{St}^M, N, \{v_1, \dots, v_n\})$ compact (as in the remark on page 3624) so that any loop in $\Gamma_{(G,S)}(R) - D$, each of whose vertices is within N of $v_i A$ for some $i \in \{1, \dots, n\}$, is homotopically trivial in $\Gamma_{(G,S)}(R) - C$. If $vA \neq v_i A$ for any $i \in \{1, \dots, n\}$, and α is a loop in $\Gamma_{(G,S)}(R) - D$ such that each vertex of α is within N of vA , then by the definition of M , α is homotopic to a loop α' in $v\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}})$ by a homotopy avoiding C . Since A is simply connected in G there is a compact set E such that any loop in $v\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}}) - E$ is homotopically trivial in $\Gamma_{(G,S)}(R) - C$. Since A is semistable at ∞ , α' is homotopic in $v\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}})$ to a loop α'' in $v\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}}) - E$. Since $C \cap v\Gamma_{(A,\mathcal{A})}(R_{\mathcal{A}}) = \emptyset$ this homotopy avoids C . Now α'' is homotopically trivial in $\Gamma_{(G,S)}(R) - C$ and so α is also. \square

3 Semistability

For the remainder of the paper, we assume that G is a finitely generated group, H is an infinite finitely generated subgroup of infinite index in G and (as in the statement of [Theorem 1.9](#)) H is subcommensurated in G :

$$H = Q_0 < Q_1 < \dots < Q_k < Q_{k+1} = G.$$

Let $\mathcal{H} := \{h_1, \dots, h_n\}$ be a finite generating set for H , and suppose the group G has generating set $\mathcal{G} := \{h_1, \dots, h_n, s_1, \dots, s_m\}$. Let $\mathcal{S} := \{s_1, \dots, s_m\}$.

Lemma 3.1 [[2](#), Lemma 3.1] *Suppose Q and B are subgroups of the group G and $Q < G$. Then $Q \cap B < B$.*

Lemma 3.2 *If H is a subgroup of A and A is a subgroup of G ($H < A < G$), then*

$$H = Q_0 < Q_1 \cap A < \dots < Q_k \cap A < A.$$

Proof Recall $Q_{k+1} := G$. For $i = 1, \dots, k + 1$, define $B_i := A \cap Q_i$. As $Q_{i-1} < Q_i$ and $B_i < Q_i$, [Lemma 3.1](#) implies that $Q_{i-1} \cap B_i < B_i$. Equivalently, $Q_{i-1} \cap A < Q_i \cap A$. □

Lemma 3.3 *Suppose $i \in \{1, 2, \dots, k + 1\}$, $g \in Q_i$ and Y is a subgroup of Q_{i-1} . Then $g^{-1}Q_{i-1}g \cap Y$ has finite index in Y and so $gYg^{-1} \cap Q_{i-1}$ has finite index in gYg^{-1} . Note that if Y is finitely generated, then $g^{-1}Q_{i-1}g \cap Y$ and $Q_{i-1} \cap gYg^{-1}$ are as well.*

Proof The group $g^{-1}Q_{i-1}g \cap Q_{i-1}$ has finite index in Q_{i-1} . Therefore, the group $g^{-1}Q_{i-1}g \cap Q_{i-1} \cap Y = g^{-1}Q_{i-1}g \cap Y$ has finite index in Y . Conjugating, we have that the group $Q_{i-1} \cap gYg^{-1}$ has finite index in gYg^{-1} . □

For $s \in S^{\pm 1}$ let \mathcal{A}_s be a finite generating set for $Q_k \cap s^{-1}Hs$, and define

$$\mathcal{A}_1 := \bigcup_{s \in S^{\pm 1}} \mathcal{A}_s \quad \text{and} \quad \mathcal{A} := \mathcal{H} \cup \mathcal{A}_1.$$

Then $A := \langle \mathcal{A} \rangle$ is a finitely generated subgroup of Q_k .

The following two lemmas imply the semistability part of [Theorem 1.9](#).

Lemma 3.4 *If H has finite index in A , then H is commensurated in G (and so G is one-ended and semistable at ∞ by [Theorem 1.8](#)).*

Lemma 3.5 *If H has infinite index in A , then H is subcommensurated in A :*

$$H = Q_0 < Q_1 \cap A < \dots < Q_k \cap A = A,$$

and both A and G are one-ended and semistable at ∞ .

Proof of Lemma 3.4 It suffices to show that for $s \in S^{\pm 1}$, $s^{-1}Hs \cap H$ has finite index in both H and $s^{-1}Hs$. Since H has finite index in A , and $\langle A_s \rangle = Q_k \cap s^{-1}Hs < A$, the group $H \cap (Q_k \cap s^{-1}Hs) = H \cap s^{-1}Hs$ has finite index in $Q_k \cap s^{-1}Hs$. By Lemma 3.3 (with $Y = H$ and $g = s^{-1}$), the group $Q_k \cap s^{-1}Hs$ has finite index in $s^{-1}Hs$ and so $H \cap s^{-1}Hs$ has finite index in $s^{-1}Hs$ for all $s \in S^{\pm 1}$. Conjugating we have $sHs^{-1} \cap H$ has finite index in H for all $s \in S^{\pm 1}$. Combining we have $s^{-1}Hs \cap H$ has finite index in both H and $s^{-1}Hs$ for all $s \in S^{\pm 1}$. \square

Proof of Lemma 3.5 Now suppose H has infinite index in A . The subcommensurated sequence $H = Q_0 < Q_1 < \dots < Q_k < G$ has length $k + 1$. Theorem 1.8 shows that if $k = 0$, then G is one-ended and semistable at ∞ . Inductively, we assume that if G' is finitely generated and there is a subcommensurated sequence $H' = Q'_0 < Q'_1 < \dots < Q'_{k-1} < G'$ of length k such that H' is finitely generated and has infinite index in G' , then G' is one-ended and semistable at ∞ .

In our case, H has infinite index in A , and the length k subcommensurated series $H = Q_0 < Q_1 \cap A < \dots < Q_{k-1} \cap A < A$ implies that A is one-ended and semistable at ∞ . Hence we may choose a finite set P of \mathcal{A} -relations so that $\Gamma_{(A,A)}(P)$ is one-ended and semistable at ∞ .

If $s \in S^{\pm 1}$ and $a \in A_s$, then there is a G -relation of the form $a = s^{-1}a_s s$ for some \mathcal{H} -word a_s . Let R be the (finite) collection of all such relations. Define

$$\tilde{\Gamma} := \Gamma_{(G, A \cup S)}(P \cup R).$$

We simultaneously show $\tilde{\Gamma}$ is one-ended and semistable at ∞ by showing that all proper edge path rays in $\tilde{\Gamma}$ are properly homotopic (completing the proof of the semistability part of Theorem 1.9).

Claim 3 *Let K be the length of the longest R -relation. If $v \in G$ (so v is a vertex of $\tilde{\Gamma}$), $s \in S^{\pm 1}$ and r is an A_s -proper ray at v , then r is properly homotopic $\text{rel}\{v\}$ to a ray of the form $(s^{-1}, h'_1, h'_2, \dots)$, where $h'_i \in \mathcal{H}$. Furthermore, this proper homotopy has image in $\text{St}^K(\text{im}(r))$.*

Proof Suppose $r = (a_1, a_2, \dots)$ with $a_i \in A_s$. Then r is properly homotopic $\text{rel}\{v\}$ to $(s^{-1}, (a_1)_s, s, s^{-1}(a_2)_s^{-1}, s, \dots)$, simply by using the 2-cells for the R -relation $a_i = s^{-1}(a_i)_s s$. Then r is properly homotopic $\text{rel}\{v\}$ to $(s^{-1}, (a_1)_s, (a_2)_s, \dots)$ by a proper homotopy in $\text{St}^K(\text{im}(r))$. \square

If $v \in G$ and C_v is a compact subcomplex of $v\Gamma_{(\mathcal{A},\mathcal{A})}(P) \subset \tilde{\Gamma}$ then there is a compact subcomplex D_v of $v\Gamma_{(\mathcal{A},\mathcal{A})}(P)$ such that if r and s are edge path rays at $w \in v\Gamma_{(\mathcal{A},\mathcal{A})}(P) - D_v$, then r and s are properly homotopic $\text{rel}\{w\}$ by a proper homotopy in $v\Gamma_{(\mathcal{A},\mathcal{A})} - C_v$. Hence, if C is a compact subcomplex of $\tilde{\Gamma}$ and we let $C_v := C \cap v\Gamma_{(\mathcal{A},\mathcal{A})}(P)$ (for the finite set of vertices v such $C \cap v\Gamma_{(\mathcal{A},\mathcal{A})}(P) \neq \emptyset$) and let $D := \bigcup D_v$, then any two \mathcal{A} -rays r and s at $w \in v\Gamma_{(\mathcal{A},\mathcal{A})}(P) - D$ are properly homotopic $\text{rel}\{w\}$ in $\tilde{\Gamma} - C$.

[18, Lemma 2] is an elementary graph theory result that states: for each $v \in G$, there is an \mathcal{H} -ray r_v at v such that for any compact set $C \subset \tilde{\Gamma}$ there are only finitely many $v \in G$ such that r_v intersects C . Also, for each $s \in \mathcal{S}^{\pm 1}$, there is an \mathcal{A}_s -ray $r_{(s,v)}$ at v such that for any compact set $C \subset \tilde{\Gamma}$, only finitely many v are such that $r_{(s,v)}$ intersects C .

Choose a sequence of compact subcomplexes $\{C_i\}_{i=1}^\infty$ of $\tilde{\Gamma}$ satisfying the following conditions:

- (1) $\bigcup_{i=1}^\infty C_i = \tilde{\Gamma}$.
- (2) $\text{St}^K(C_i)$ (see Claim 3) is contained in the interior of C_{i+1} , and the finite set of vertices v such that r_v or $r_{(s,v)}$ ($s \in \mathcal{S}^{\pm 1}$) intersects C_i , is a subset of C_{i+1} .
- (3) If r and s are \mathcal{A} -rays in $\tilde{\Gamma} - C_i$ both based at a vertex v , then r and s are properly homotopic $\text{rel}\{v\}$ by a proper homotopy in $v\Gamma_{(\mathcal{A},\mathcal{A})}(P) - C_{i-1}$; see Theorem 2.2(4).

For convenience, define $C_i := \emptyset$ for $i < 1$ and observe that conditions (1)–(3) (see Theorem 2.2(5)) remain valid for all C_i . The next lemma implies Lemma 3.5 and concludes the proof of the semistability part of Theorem 1.9.

Lemma 3.6 *If v is a vertex of $\tilde{\Gamma}$, and $t = (e_1, e_2, \dots)$ is an arbitrary ray at v , then t is properly homotopic to r_v , $\text{rel}\{v\}$.*

Proof Assume that t has consecutive vertices $v = v_0, v_1, \dots$. By construction, if $v_j \in C_{i+1} - C_i$, then r_{v_j} avoids C_{i-1} . Assume j is the largest integer such that C_j avoids e_i . Then $r_{v_{i-1}}$ and r_{v_i} avoid C_{j-1} . We will show $r_{v_{i-1}}$ is properly homotopic to $e_i * r_{v_i}$ $\text{rel}\{v_{i-1}\}$ by a proper homotopy H_i with image avoiding C_{j-3} .

If $e_i \in \mathcal{A}^{\pm 1}$, this is clear by condition (3) with H_i avoiding C_{j-2} . If $e_i \in \mathcal{S}^{\pm 1}$, then $r_{v_{i-1}}$ and $r_{(e_i, v_{i-1})}$ are \mathcal{A} -rays avoiding C_{j-1} , and so by (3) are properly homotopic $\text{rel}\{v_{i-1}\}$ by a homotopy avoiding C_{j-2} . By Claim 3 and condition (2), $r_{(e_i, v_{i-1})}$ is properly homotopic $\text{rel}\{v_{i-1}\}$ to a ray (e_i, h'_1, h'_2, \dots) , where $h'_i \in \mathcal{H}^{\pm 1}$ and the




Figure 1: Inductive construction of H

homotopy avoids C_{j-2} . By condition (3), (h'_1, h'_2, \dots) is properly homotopic $\text{rel}\{v_i\}$ to r_{v_i} by a proper homotopy that avoids C_{j-3} . Patch these three proper homotopies together to obtain H_i ; see Figure 1.

Let H be the homotopy $\text{rel}\{v\}$ of t to r_v , obtained by patching together the homotopies H_i . We need to check that H is proper. Let $C \subset \tilde{\Gamma}$ be compact. Choose an index j such that $C \subseteq C_j$. Since t is a proper edge path to ∞ , choose an index N such that all edges after the N^{th} edge of t avoid C_{j+3} . Then for all $i > N$, H_i avoids C_j , so H is proper. \square

This concludes the proof of Lemma 3.5 and the first part of Theorem 1.9. \square

4 Simple connectivity at ∞

It is straightforward to check that the proof of the simply connected at ∞ part of Theorem 1.8 given in [3] extends to the finitely generated case (as follows): if $\langle S : R \rangle$ is a finite presentation of the group G then $\Gamma_{(G,S)}(R)$ is simply connected. The only time the simple connectivity of Γ is used in the proof of Theorem 1.8 is via this fact:

- (*) If C is a compact subset of Γ and N is a fixed positive integer, then there is an integer $M(N, C)$ such that any edge path loop α of length at most N in $\Gamma - \text{St}^M(C)$ is homotopically trivial in $\Gamma - C$.

Suppose G is a finitely generated subgroup of a finitely presented group W and W has presentation $\langle \mathcal{W} : R \rangle$, where \mathcal{W} contains a set of generators \mathcal{G} for G . When proving a finitely generated version of the simply connected at ∞ part of Theorem 1.8, all work is done in the simply connected space $\Gamma_{(W, \mathcal{W})}(R)$, and one only needs (*) for edge path loops α with edge labels in $\mathcal{G}^{\pm 1}$. Hence, the proof of the simply connected at ∞ part of Theorem 1.8 directly extends to the stronger finitely generated version.

Theorem 4.1 (Conner and Mihalik improved) *Suppose H is a one-ended, finitely presented infinite subgroup of infinite index in the finitely generated and recursively presented group G . If H is commensurated in G , then G is one-ended and simply connected at ∞ .*

In order to finish the proof of our main theorem it remains to prove the following.

Theorem 4.2 *Suppose that H is a one-ended finitely presented subcommensurated subgroup of infinite index in the finitely generated and recursively presented group G :*

$$H = Q_0 < Q_1 < \dots < Q_k < Q_{k+1} = G.$$

Then G is simply connected at ∞ .

Proof We say H is $(k+1)$ -subcommensurated in G . When $k = 0$, [Theorem 4.1](#) implies that G is simply connected at ∞ . Assume (inductively) the statement of [Theorem 4.2](#) is valid when H is $(n+1)$ -subcommensurated for $n < k$. Let

$$\begin{aligned} \mathcal{H} &:= \{h_1, \dots, h_n\} \text{ generate } H, \\ \mathcal{G} &:= \{h_1, \dots, h_n, s_1, \dots, s_m\} \text{ generate } G, \text{ and} \\ \mathcal{S} &:= \{s_1, \dots, s_m\}. \end{aligned}$$

For p an element of a group P with generating set \mathcal{P} , let $|g|_{\mathcal{P}}$ be word length in \mathcal{P} : the smallest integer ℓ such that $g = p_1 \cdots p_{\ell}$ where $p_i \in \mathcal{P}^{\pm 1}$. We use the notation $|g| := |g|_{\mathcal{P}}$ for all $g \in P$.

For each $s \in \mathcal{S}^{\pm 1}$ let \mathcal{A}_s be a finite generating set for $sHs^{-1} \cap Q_k$ (see [Lemma 3.3](#) with $Y = H$) and let $\mathcal{A}'_s := s^{-1}\mathcal{A}_s s \subset H$. Choose an integer L_1 such that

$$L_1 \geq |a|_{\mathcal{H}} \quad \text{for all } a \in \bigcup_{s \in \mathcal{S}^{\pm 1}} \mathcal{A}'_s.$$

We have that

$$A_s := \langle \mathcal{A}_s \rangle < Q_k$$

has finite index in sHs^{-1} , and

$$A'_s := \langle \mathcal{A}'_s \rangle = s^{-1}A_s s < H$$

has finite index in H . As in [Section 3](#), define

$$A_1 := \bigcup_{s \in \mathcal{S}^{\pm 1}} A_s, \quad \mathcal{A} := \mathcal{H} \cup A_1 \quad \text{and} \quad A := \langle \mathcal{A} \rangle < Q_k.$$

For each $s \in \mathcal{S}^{\pm 1}$ and $a \in A_s$ there is an \mathcal{H} -word $w(s, a)$ of length at most L_1 , such that $s^{-1}asw^{-1}(a, s)$ is an (A, A) -relator, which we denote by $r(a, s)$. Let

$$R_1 := \{r(a, s) \mid s \in \mathcal{S}^{\pm 1}, a \in A_s\}.$$

For each $g \in G$ let B_g be a finite generating set for the group $gAg^{-1} \cap Q_k$; see Lemma 3.3 with $Y = A$. Let $B'_g := g^{-1}B_g g \subset A \cap g^{-1}Q_k g$. If g is in A , then $gAg^{-1} = A$ and so we define $B_g := A := B'_g$. Then

$$B_g := \langle B_g \rangle = gAg^{-1} \cap Q_k$$

has finite index in gAg^{-1} , and

$$B'_g := \langle B'_g \rangle = g^{-1}B_g g = A \cap g^{-1}Q_k g$$

has finite index in A . For each $g \in G$, let N_g be an integer so that in the Cayley graph $\Gamma_{(G, \mathcal{G})}$, each vertex of A is within N_g of a vertex of B'_g . Let

$$B_j := A \cup \left(\bigcup_{\{g \in G: |g| \leq j\}} B_g \right) \subset Q_k \quad \text{and} \quad N_j := \max\{N_g : g \in G \text{ and } |g| \leq j\}.$$

Lemma 4.3 *Suppose g is in G and y in gA . Then in $\Gamma_{(G, \mathcal{G})}$, y is within $N_g + |g|$ of a point of B_g .*

Proof Let $y = ga$ for some $a \in A$. There is $b' \in B'_g = g^{-1}Q_k g \cap A$ within N_g of a . Then $y' := gb'$ is within N_g of $y = ga$. As $y'g^{-1} = gb'g^{-1} \in Q_k \cap gAg^{-1} = B_g$, y' is within $|g|$ of B_g and so y is within $N_g + |g|$ of B_g . \square

If H has finite index in A , then by Lemma 3.4, H is commensurated in G and so G is simply connected at ∞ by Theorem 4.1. So we may assume that H has infinite index in A . Our induction hypothesis, Lemma 3.2 and the results of Section 3 imply the following.

Lemma 4.4 *The finitely generated subgroups A and $B_j := \langle B_j \rangle$ of Q_k are one-ended, semistable at ∞ and simply connected at ∞ for all $j \geq 1$.*

Next assume that G is a subgroup of a finitely presented (over) group W . Then for all $j \geq 1$, A and B_j are simply connected at ∞ in W . Let \mathcal{W} be a finite generating set for W containing \mathcal{A} and \mathcal{G} , and let $\langle \mathcal{W} : R \rangle$ be a finite presentation for W . Assume that R contains a set R' of \mathcal{A} -relations so that $\Gamma_{(A, \mathcal{A})}(R')$ is semistable at ∞ . We also assume that R' contains the set of conjugation relations R_1 . If v is a G -vertex of $\Gamma_{(W, \mathcal{W})}(R)$, let $v\Gamma_{(A, \mathcal{A})}(R')$ be the copy of $\Gamma_{(A, \mathcal{A})}(R')$ at v . To ease notation, if p is a $G^{\pm 1}$ -word and \bar{p} is the corresponding element of G , define $B_p := B_{\bar{p}}$ and $B'_p := B'_{\bar{p}}$.

As a direct consequence of Lemma 4.3, we have another lemma.

Lemma 4.5 Suppose v is a G -vertex of $\Gamma_{(W, \mathcal{W})}(R)$ and (e_1, \dots, e_i) labels a \mathcal{G} -edge path with consecutive vertices $v = v_0, v_1, \dots, v_i$. If $j \in \{1, 2, \dots, i\}$ and w is a vertex of the Cayley graph $v_j \Gamma_{(A, A)}$, then w is within $j + N_j$ of a vertex of $v \Gamma_{(B_j, B_j)}$.

Note that [Theorem 4.2](#) does not follow directly from [Lemma 2.9](#) and the fact that B_ℓ is simply connected at ∞ in W for all ℓ , but it does follow from the next lemma.

Lemma 4.6 Given any compact subcomplex C of $\Gamma_{(W, \mathcal{W})}(R)$ there is a compact subcomplex D of $\Gamma_{(W, \mathcal{W})}(R)$ such that any \mathcal{G} -loop α at a G -vertex of $\Gamma_{(W, \mathcal{W})}(R) - D$ is homotopically trivial in $\Gamma_{(W, \mathcal{W})}(R) - C$.

Proof If v is a vertex of $\Gamma_{(W, \mathcal{W})}(R)$ and $v \Gamma_{(A, A)}(R') \cap C$ is empty, then by [Theorem 2.2\(5\)](#), any two A -rays at v are properly homotopic relative to v by a proper homotopy in $v \Gamma_{(A, A)}(R')$ (and so the homotopy avoids C). There are only finitely many $v \Gamma_{(A, A)}(R')$ that intersect C . Since $\Gamma_{(A, A)}(R')$ is semistable at ∞ there is a compact subcomplex D_1 of $\Gamma_{(W, \mathcal{W})}(R)$ such that any two A -edge path rays based at a G -vertex v and with image in $\Gamma_{(W, \mathcal{W})}(R) - D_1$, are properly homotopic relative to v by a homotopy in $v \Gamma_{(A, A)}(R') - C$. There are only finitely many Cayley graphs of the form $v \Gamma_{(H, \mathcal{H})}$ or $v \Gamma_{(A_s, A_s)}$ (for $v \in G$ and $s \in S^{\pm 1}$) that intersect D_1 . Choose a finite subcomplex D of $\Gamma_{(W, \mathcal{W})}(R)$ such that D contains D_1 and the bounded components of both $v \Gamma_{(A_s, A_s)} - \text{St}^{L_1}(D_1)$ and $v \Gamma_{(H, \mathcal{H})} - D_1$, for all $v \Gamma_{(A_s, A_s)}$ and $v \Gamma_{(H, \mathcal{H})}$ that intersect D_1 . (Recall, if $s \in S^{\pm 1}$ and $a \in A_s$, then there is an \mathcal{H} -word $w(a, s)$ of length at most L_1 , such that $s^{-1} a s w(a, s)^{-1} \in R_1$.)

Therefore,

- (*) if e is an edge of $\Gamma_{(W, \mathcal{W})}(R) - D$ with initial vertex $v \in G$, terminal vertex w and label $s \in S^{\pm 1}$, then there is a proper A_s -ray q_v at v avoiding $\text{St}^{L_1}(D_1)$ and hence an \mathcal{H} -ray s_w at w avoiding D_1 , such that q_v and (e, s_w) are homotopic relative to v by a homotopy (using only 2-cells arising from R_1 -conjugation relations) in $\Gamma_{(W, \mathcal{W})}(R) - C$ (see [Figure 2](#)); and
- (**) if v is a G -vertex of $\Gamma_{(W, \mathcal{W})}(R) - D$, there is an \mathcal{H} -ray at v in $\Gamma_{(W, \mathcal{W})}(R) - D_1$.

Assume α is a \mathcal{G} -loop based at the G -vertex v in $\Gamma_{(W, \mathcal{W})}(R) - D$. We wish to show that α is homotopically trivial in $\Gamma_{(W, \mathcal{W})} - C$. Since G is one-ended, we may assume that $\Gamma_{(G, \mathcal{G})} - D$ is connected, and so there is an edge path in $\Gamma_{(G, \mathcal{G})} - D$ from v to a vertex of H . Hence we assume, without loss, that $v \in H$. Let ℓ be the length of α . By [Lemma 4.4](#), B_ℓ is simply connected at ∞ and [Lemma 2.9](#) implies that there is a compact subcomplex $E(C, \ell + N_\ell)$ of $\Gamma_{(W, \mathcal{W})}(R)$ with the following property: if β is an edge path loop with image in $\Gamma_{(W, \mathcal{W})}(R) - E$ and each vertex of β is within $\ell + N_\ell$ of B_ℓ , then β is homotopically trivial in $\Gamma_{(W, \mathcal{W})}(R) - C$.




Figure 2: For e an edge of $\Gamma_{(W, \mathcal{W})}(R) - D$, there is a proper A_s -ray q_v avoiding $\text{St}^{L_1}(D_1)$ and hence an \mathcal{H} -ray s_w avoiding D_1 , such that q_v and (e, s_w) are homotopic in $\Gamma_{(W, \mathcal{W})}(R) - C$.




Figure 3: A \mathcal{G} -loop based at a G -vertex in $\Gamma_{(W, \mathcal{W})}(R) - D$ is homotopically trivial in $\Gamma_{(W, \mathcal{W})}(R) - C$.

It is enough to show that α is homotopic to such a β in $\Gamma_{(W, \mathcal{W})}(R) - C$ (and this is where the semistability of A comes in). Let s_v be an \mathcal{H} -proper edge path ray at v in $\Gamma_{(W, \mathcal{W})}(R) - D_1$; see (**). If all edges of α are \mathcal{H} -edges, then by the definition of D_1 , the rays s_{v_0} and (α, s_{v_0}) are properly homotopic relative to v , by \hat{H} a proper homotopy with image in $v\Gamma_{(A, A)}(R') - C$. Otherwise, write α as $(\tau_0, e_0, \tau_1, e_1, \dots, \tau_j, e_j, \tau_{j+1})$, where τ_i is a (possibly trivial) \mathcal{H} -path and e_i is an S -edge. Let the initial vertex of e_i be v'_i and the terminal vertex of e_i be v_{i+1} ; see Figure 3.

If e_i is labeled by $s \in \mathcal{S}^{\pm 1}$ then let q_i be a proper A_s -ray at v'_i , avoiding $\text{St}^{L_1}(D_1)$. For each edge a of q_i , there is a 2-cell with boundary label $(s^{-1}, a, s, w^{-1}(a, s))$, where $w(a, s)$ is an \mathcal{H} -word of length at most L_1 (see the definition of R_1). So if q_i is labeled (a_1, a_2, \dots) then $s_{v_{i+1}}$, the \mathcal{H} -ray at v_{i+1} , with labeling $(w(a_1, s), w(a_2, s), \dots)$ is such that q_i is properly homotopic to $(e_i, s_{v_{i+1}})$ relative to v'_i by a homotopy (only using R_1 -cells) with image in $\Gamma_{(W, \mathcal{W})}(R) - C$. The ray s_{v_i} has image in $\Gamma_{(W, \mathcal{W})}(R) - D_1$ and so by the semistability of $\Gamma_{(A, A)}(R')$ we have for $i \in \{0, 1, \dots, j\}$,

q_i is properly homotopic to (τ_i^{-1}, s_{v_i}) relative to v'_i by a proper homotopy \hat{H}_i in $v'_i\Gamma_{(A,A)} - C$. Finally, define \hat{H}_{j+1} to be a proper homotopy in $v_0\Gamma_{(A,A)} - C$ of s_{v_0} to $(\tau_{j+1}^{-1}, s_{v_{j+1}})$.

Assume that v'_i is the $j(i)^{\text{th}}$ vertex of α . By Lemma 4.5, every vertex of $v'_i\Gamma_{(A,A)}(R')$ is within $N_{j(i)} + j(i) (\leq N_\ell + \ell)$ of $B_{j(i)} \subset B_\ell$. Hence for $i \in \{0, \dots, j\}$, each vertex of (the image of) \hat{H}_i is within $N_\ell + \ell$ of a vertex of B_ℓ .

Combining the homotopies $\hat{H}_0, \dots, \hat{H}_{j+1}$ along with those of q_i to $(e_i, s_{v_{i+1}})$, we have a proper cellular homotopy \hat{H} (relative to v) of s_{v_0} to (α, s_{v_0}) with image in $\Gamma_{(W,W)}(R) - C$, and each vertex of the image of the homotopy \hat{H} is within $N_\ell + \ell$ of a vertex of B_ℓ . Then

$$\hat{H}: [0, 1] \times [0, \infty) \rightarrow \Gamma_{(W,W)}(R) - C$$

such that $\hat{H}|_{[0,1] \times \{0\}}$ is α , $\hat{H}|_{\{0\} \times [0,\infty)}$ and $\hat{H}|_{\{1\} \times [0,\infty)}$ both agree with s_{v_0} , and each vertex of the image of \hat{H} is within $N_\ell + \ell$ of B_ℓ . Choose N such that $\hat{H}^{-1}(E) \subset [0, 1] \times [0, N]$. The loop $\hat{H}|_{[0,1] \times \{N+1\}}$ provides a \mathcal{G} -loop β such that α is homotopic to β in $\Gamma_{(W,W)}(R) - C$, and each vertex of β is within $N_\ell + \ell$ of B_ℓ . By the definition of E , the loop β (and hence α) is homotopically trivial in $\Gamma_{(W,W)}(R) - C$. □

This completes the proof of Theorem 4.2. □

5 Grigorchuk’s group is simply connected at ∞

In 1985, Mihalik [17] proved that if H is an ascending HNN-extension of a finitely presented group G , then H is semistable at ∞ . If, additionally, G is one-ended, then H is simply connected at ∞ . Finitely presented ascending HNN extensions of one-ended finitely generated groups are not even known to be semistable at ∞ .

Grigorchuk [7; 8] constructed an infinite finitely generated torsion group G . The group G has a subgroup of finite index T , and T has a subgroup of finite index that is isomorphic to $T \times T$. There is a finitely presented ascending HNN-extension of G (and so G is recursively presented). It is easy to see that if G is finitely generated and H has finite index in G , then G is semistable at ∞ if and only if H is semistable at ∞ ; see [5, Proposition 16.5.3]. [16, Theorem 2.2] implies that $T \times T$ (and hence G) is semistable at ∞ . Theorem 5.1 implies that Grigorchuk’s group G is simply connected at ∞ and Theorem 5.2 implies that the finitely presented HNN-extension of G is also simply connected at ∞ . This last result generalizes the main result of [4] which states that G is quasimply filtered (qsf).

Theorem 5.1 *Suppose the recursively presented group G is finitely generated and isomorphic to $A \times B$ where A and B are finitely generated infinite groups and A is one-ended. Then G is simply connected at ∞ .*

Proof Suppose G is a subgroup of a finitely presented group Q . Let $\mathcal{P} := \langle Q : R_Q \rangle$ be a finite presentation for Q such that \mathcal{Q} contains generators \mathcal{A} for A and \mathcal{B} for B , and R_Q contains the commutation relations R between these generators. Define $\Gamma(\mathcal{P}) := \Gamma_{(\mathcal{Q}, \mathcal{Q})}(R_Q)$ (the Cayley 2-complex for \mathcal{P}). The subspace $\Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) \subset \Gamma(\mathcal{P})$ is G -equivariantly homeomorphic to $\Gamma_{(A, \mathcal{A})} \times \Gamma_{(B, \mathcal{B})}$. Let $\phi_A: \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) \rightarrow \Gamma_{(A, \mathcal{A})}$ and $\phi_B: \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) \rightarrow \Gamma_{(B, \mathcal{B})}$ be projections.

Claim 5.1.1 *Suppose C is a compact subcomplex of $\Gamma(\mathcal{P})$. Then there is a compact subcomplex D of $\Gamma(\mathcal{P})$ such that if γ is an edge path loop in $\Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) - D$ with edge labels all in $\mathcal{A}^{\pm 1}$ or all in $\mathcal{B}^{\pm 1}$, then γ is homotopically trivial in $\Gamma(\mathcal{P}) - C$.*

Proof The set $\phi_B(C)$ union the bounded components of $\Gamma_{(B, \mathcal{B})} - \phi_B(C)$ is compact and so contains finitely many vertices. If v is one of these vertices, let β_v be an edge path from v to a vertex v' of an unbounded path component of $\Gamma_{(B, \mathcal{B})} - \phi_B(C)$. If v is a vertex of an unbounded component of $\Gamma_{(B, \mathcal{B})} - \phi_B(C)$, then β_v is trivial and $v' := v$. In any case, let τ_v be a proper edge path ray at v' with image in $\Gamma_{(B, \mathcal{B})} - \phi_B(C)$. Let L be the length of the longest β_v . Let $D := \text{St}^L(C)$. Suppose each edge of γ has label in $\mathcal{A}^{\pm 1}$. Then $\phi_B(\gamma)$ is a single vertex v of $\Gamma_{(B, \mathcal{B})}$. Let w be the initial vertex of γ , and lift the ray (β_v, τ_v) to w (ie consider the \mathcal{B} -ray at w with the same edge labeling as that of (β_v, τ_v)). Call the lift $(\tilde{\beta}_v, \tilde{\tau}_v)$. Since each edge of $(\tilde{\beta}_v, \tilde{\tau}_v)$ has label in $\mathcal{B}^{\pm 1}$ and each edge of γ has label in $\mathcal{A}^{\pm 1}$, there is a proper homotopy of $H: [0, 1] \times [0, \infty) \rightarrow \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R)$ that uses the product structure of $\Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R)$ to basically slide the loop γ along the ray $(\tilde{\beta}_v, \tilde{\tau}_v)$. In particular, $H|_{\{0\} \times [0, \infty)} = H|_{\{1\} \times [0, \infty)} = (\tilde{\beta}_v, \tilde{\tau}_v)$ and $H|_{[0, 1] \times \{n\}}$ is an edge path loop with the same labeling as γ for each integer $n \geq 0$. The image of $\phi_B(H)$ is simply the image of (β_v, τ_v) . Say β_v has length $K (\leq L)$. Then $H|_{[0, 1] \times [0, K]}$ has image in $\text{St}^K(\text{image}(\gamma)) \subset \Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) - C$. As $\phi_B(H|_{[0, 1] \times [K, \infty)})$ has the same image as τ_v (which avoids $\phi_B(C)$), the image of H avoids C . Note that $H|_{[0, 1] \times \{n\}}$ is the translate of γ to the n^{th} vertex of $(\tilde{\beta}_v, \tilde{\tau}_v)$. Since this ray is proper, there is an integer N such that the translate of γ to the N^{th} vertex of $(\tilde{\beta}_v, \tilde{\tau}_v)$ is homotopically trivial in $\Gamma(\mathcal{P}) - C$. Since $H|_{[0, 1] \times [0, N]}$ provides a homotopy (avoiding C) of γ to a loop that bounds in $\Gamma(\mathcal{P}) - C$, we have that γ is homotopically trivial in $\Gamma(\mathcal{P}) - C$.

A completely analogous argument in the case all edges of γ have labels in $\mathcal{B}^{\pm 1}$ finishes the proof of the claim. Note that the one-endedness of A is not important yet. \square

Let C be a compact subcomplex of $\Gamma(\mathcal{P})$ and D the associated compact set of Claim 5.1.1. Since A is one-ended, $\Gamma_{(A,A)} - \phi_A(D)$ has one unbounded component K and $\Gamma_{(A,A)} - K$ is compact. Let z_0 be a vertex of K . For each vertex v of $\Gamma_{(A,A)}$ choose an edge path τ_v from v to z_0 so that if $v \in K$ then τ_v has image in K . There are only finitely many vertices of $\Gamma_{(A,A)}$ not in K . Of these, let M be the length of the longest τ_v . Let $E := \text{St}^M(D)$. We show that edge path loops in $\Gamma_{(G,A \cup B)}(R) - E$ are homotopically trivial in $\Gamma(\mathcal{P}) - C$.

Let γ be a loop in $\Gamma_{(G,A \cup B)}(R) - E$. Write γ as $(\alpha_1, \delta_1, \dots, \alpha_n, \delta_n)$, where edge labels in the edge paths α_i (respectively δ_i) are in $\mathcal{A}^{\pm 1}$ (respectively $\mathcal{B}^{\pm 1}$). Let a_i be the initial vertex of α_i and d_i the initial vertex of δ_i . Let $a'_i := \phi_A(a_i)$ and $d'_i := \phi_A(d_i)$. Then $d'_i = a'_{i+1}$ since δ_i is a \mathcal{B} -path. Let $\tilde{\tau}_i$ be the lift of the path $\tau_{a'_i}$ to a_i and $\hat{\tau}_i$ be the lift of $\tau_{d'_i}$ to d_i ; see Figure 4. Note that

- (1) $\tau_{d'_i} = \tau_{a'_{i+1}}$, so $\hat{\tau}_i$ and $\tilde{\tau}_{i+1}$ have the same edge labeling,
- (2) $\tilde{\tau}_i$ and $\hat{\tau}_i$ share the same end point v_i , where $\phi_A(v_i) = z_0 \in K$, and
- (3) the translate of δ_i to v_i (call it δ'_i) ends at v_{i+1} .

By (1) there is a product homotopy H_i of $(\delta_i, \tilde{\tau}_{i+1})$ to $(\hat{\tau}_i, \delta'_i)$. We show that the image of H_i avoids D .

If $d'_i (= \phi_A(d_i)) \in K$ then the image of $\phi_A(H_i)$ is equal to the image of $\tau_{d'_i}$ which is a subset of $K \subset \Gamma_{(A,A)} - \phi_A(D)$. In this case, H_i has image in $\Gamma_{(G,A \cup B)}(R) - D$. If $\phi_A(d_i) \notin K$, then the length of $\hat{\tau}_i$ is $\leq M$ and the image of H_i is in $\text{St}^M(\text{image}(\delta_i)) \subset \Gamma_{(G,A \cup B)}(R) - D$ (since δ_i has image in $\Gamma_{(G,A \cup B)}(R) - E$ and $E = \text{St}^M(D)$).

The \mathcal{A} -loops $(\alpha_i, \hat{\tau}_i, \tilde{\tau}_i^{-1})$ have image in $\Gamma_{(G,A \cup B)}(R) - D$ and so, by Claim 5.1.1 are homotopically trivial by a homotopy H'_i in $\Gamma(\mathcal{P}) - C$. Combining all homotopies H_i and H'_i we have that γ is homotopic to $(\delta'_1, \dots, \delta'_n)$ by a homotopy in $\Gamma_{(G,A \cup B)}(R) - D$. Since all edge labels of this last loop are in $\mathcal{B}^{\pm 1}$, Claim 5.1.1 implies $(\delta'_1, \dots, \delta'_n)$ (and hence γ) is homotopically trivial in $\Gamma(\mathcal{P}) - C$. □

If $\phi: G \rightarrow G$ is a monomorphism then the HNN extension

$$H := \langle G, t : t^{-1}gt = \phi(g) \text{ for all } g \in G \rangle$$

is called an *ascending HNN extension* of G with *stable letter* t , and denoted $G *_{\phi,t}$. When ϕ is not an epimorphism, then H is a *strictly ascending HNN extension*. There is an ascending HNN extension H of Grigorchuk’s group G that is finitely presented. The main result of [4] shows that H has the quasimply filtered (qsf) property. Simple connectivity at ∞ implies qsf [1]. Our next result implies that H is simply connected at ∞ and so generalizes [4]. We need not assume that H is finitely presented.




Figure 4: Loops in $\Gamma_{(G, \mathcal{A} \cup \mathcal{B})}(R) - E$ are homotopically trivial in $\Gamma(\mathcal{P}) - C$.

Theorem 5.2 Suppose H is an ascending HNN extension of a one-ended finitely generated semistable at ∞ and simply connected at ∞ group G . Then H is simply connected at ∞ .

Proof Say $H := G *_{\phi, t}$. Assume that Q is finitely presented and H is a subgroup of Q . We must show that H is simply connected at ∞ in Q . Let \mathcal{G} be a finite set of generators for G , and R_1 a finite set of relations so that $\Gamma_{(G, \mathcal{G})}(R_1)$ is semistable at ∞ . Since t is the stable letter in $G *_{\phi, t}$, the finite set $\mathcal{H} := \mathcal{G} \cup \{t\}$ generates H . For each generator $g \in \mathcal{G}$ there is a \mathcal{G} -word (usually denoted $\phi(g)$) giving rise to the relator $t^{-1}gt = \phi(g)$. These (finitely many) relators are called the *conjugation relations* of H with respect to \mathcal{G} . Let R_2 be this set of conjugation relations and let $R := R_1 \cup R_2$. Let \mathcal{P} be a finite presentation for Q with \mathcal{H} a subset of the generators and R a subset of the relations of \mathcal{P} . We assume $\Gamma_{(G, \mathcal{G})}(R_1)$ is a subset of $\Gamma_{(H, \mathcal{H})}(R)$ which is a subset of $\Gamma(\mathcal{P})$.

There is a homomorphism from $\mu: H \rightarrow \mathbb{Z}$ with kernel equal to the normal closure of G . We say an element $h \in H$ (ie a vertex of $\Gamma_{(H, \mathcal{H})}$) is in level n if $\mu(h) = n$. If C is a compact subcomplex of Γ we say C is in levels $L(C)$ through $M(C)$ if L is maximal and M is minimal such that for each vertex v of C , $L \leq \mu(v) \leq M$. If e is a \mathcal{G} -edge of $\Gamma_{(H, \mathcal{H})}(R)$ in level K and e has label $g \in \mathcal{G}^{\pm 1}$ then the relation $t^{-1}gt = \phi(g)$ defines a 2-cell of $\Gamma_{(H, \mathcal{H})}(R)$ that slides e to a path in level $K + 1$. The path in level $K + 1$ can be slid to level $K + 2$ and so e can be slid to any level above level K .




Figure 5: Sliding \mathcal{G} -edges of γ up in any level $L - 1$ through M to level $M + 1$ by a homotopy avoiding C .

The basic idea is to show that there is a compact set D in $\Gamma_{(H, \mathcal{H})}(R)$ such that any loop in $\Gamma_{(H, \mathcal{H})}(R) - D$ is homotopic to a loop that can be slid to a single level (usually above the levels of C), by a homotopy avoiding C . Any path in a single level belongs to a translate of $\Gamma_{(G, \mathcal{G})}(R_1)$. The semistability of G is used to move this loop (by a homotopy in that copy of $\Gamma_{(G, \mathcal{G})}(R_1)$) to a loop far from C . The simple connectivity at ∞ of G shows this last loop is homotopically trivial in $\Gamma(P) - C$ and so H is simply connected at ∞ in Q .

Let $D(C)$ be a compact subcomplex of $\Gamma_{(H, \mathcal{H})}(R)$ such that if e is a \mathcal{G} -edge of $\Gamma_{(H, \mathcal{H})}(R)$ between levels $L(C) - 1$ and $M(C)$ and the homotopy that slides e to level $M + 1$ intersects C , then e is an edge of D . Also assume that if X is a translate of $\Gamma_{(G, \mathcal{G})}$ in level $L - 1$, then D contains all bounded components of $X - D$ (there are only finitely many such X that intersect D). Now let γ be an edge path loop in $\Gamma_{(H, \mathcal{H})}(R) - D$. If no vertex of γ is in level L through M , then γ can be slid up to be entirely in level $L - 1$ or into some level above level M by a homotopy avoiding C .

Otherwise, each \mathcal{G} -edge of γ in a level $L - 1$ through M can be slid to level $M + 1$ by a homotopy missing C . Call the resulting path γ_1 (so any edge of γ_1 between levels $L - 1$ and M is labeled t). Consider a maximal segment α of γ_1 with vertices all in levels $L - 1$ and below. Then α is preceded (respectively followed) by a segment of γ_1 of the form t^{-n} (respectively t^n) that begins (respectively ends) in level $M + 1$. Let a be the initial point of α , and b be the end point of α ; see Figure 5.

Note that a and b are points of γ and so do not belong to D . Hence there are proper \mathcal{G} -edge path rays r_a and r_b at a and b respectively that avoid D . Slide α (relative

to a and b) to level $L - 1$ and call the resulting path α_1 . This homotopy is below the levels of C and so avoids C . Now since $\Gamma_{(G, \mathcal{G})}(R_1)$ is semistable at ∞ , there is a proper homotopy A (in $a\Gamma_{(G, \mathcal{G})}(R_1)$) of r_a to (α_1, r_b) relative to a . This homotopy gives an edge path of the form $\delta := (\delta_1, \delta_2, \delta_3)$ that is homotopic to α_1 , where δ_1 is an initial segment of r_a (and so avoids D), δ_3^{-1} is an initial segment of r_b (and so avoids D) and δ_2 is a “far out” path in the image of A (and so it also avoids D). By the definition of D each edge of δ can be slid to level $M + 1$ by a homotopy avoiding C .

This shows that γ_1 (and hence γ) is homotopic to a loop γ_2 in levels all above level M . Slide this loop up to a loop γ_3 in a single level (by a homotopy avoiding C , since C is below level $M + 1$). Note that γ_3 is a \mathcal{G} -path (since it has no t -edges). In any case, γ is homotopic (in $\Gamma(\mathcal{P}) - C$) to a \mathcal{G} -loop γ_3 in a translate, $v\Gamma_{(G, \mathcal{G})}(R_1)$, of $\Gamma_{(G, \mathcal{G})}(R_1)$ that avoids C . The simple connectivity at ∞ of G implies that there is a compact set E such that any loop in $v\Gamma_{(G, \mathcal{G})}(R_1) - E$, is homotopically trivial in $\Gamma(\mathcal{P}) - C$. The semistability of G implies γ_3 is homotopic in $v\Gamma_{(G, \mathcal{G})}(R_1)$ to a loop in $\Gamma_{(G, \mathcal{G})}(R_1) - E$ and so γ_3 and hence γ , is homotopically trivial in $\Gamma(\mathcal{P}) - C$. \square

References

- [1] **S G Brick, M L Mihalik**, *The QSF property for groups and spaces*, Math. Z. 220 (1995) 207–217 [MR1355026](#)
- [2] **G Conner, M Mihalik**, *Commensurated subgroups and ends of groups*, J. Group Theory 16 (2013) 107–139 [MR3008315](#)
- [3] **G R Conner, M L Mihalik**, *Commensurated subgroups, semistability and simple connectivity at infinity*, Algebr. Geom. Topol. 14 (2014) 3509–3532 [MR3302969](#)
- [4] **L Funar, D E Otera**, *On the wgsc and qsf tameness conditions for finitely presented groups*, Groups Geom. Dyn. 4 (2010) 549–596 [MR2653975](#)
- [5] **R Geoghegan**, *Topological methods in group theory*, Graduate Texts in Mathematics 243, Springer (2008) [MR2365352](#)
- [6] **R Geoghegan, M L Mihalik**, *Free abelian cohomology of groups and ends of universal covers*, J. Pure Appl. Algebra 36 (1985) 123–137 [MR787167](#)
- [7] **RI Grigorchuk**, *On a problem of Day on nonelementary amenable groups in the class of finitely presented groups*, Mat. Zametki 60 (1996) 774–775 [MR1619929](#) In Russian; translation in Math. Notes 60 (1996) 580–582
- [8] **RI Grigorchuk**, *An example of a finitely presented amenable group that does not belong to the class EG*, Mat. Sb. 189 (1998) 79–100 [MR1616436](#) In Russian; translation in Sb. Math. 189 (1998) 75–95
- [9] **P de la Harpe**, *Topics in geometric group theory*, University of Chicago Press (2000) [MR1786869](#)

- [10] **G Higman**, *Subgroups of finitely presented groups*, Proc. Roy. Soc. Ser. A 262 (1961) 455–475 [MR0130286](#)
- [11] **B Jackson**, *End invariants of group extensions*, Topology 21 (1982) 71–81 [MR630881](#)
- [12] **F E A Johnson**, *Manifolds of homotopy type $K(\pi, 1)$, I*, Proc. Cambridge Philos. Soc. 70 (1971) 387–393 [MR0290358](#)
- [13] **F E A Johnson**, *On the first end invariant of an exact sequence*, Mathematika 22 (1975) 60–70 [MR0427483](#)
- [14] **R Lee, F Raymond**, *Manifolds covered by Euclidean space*, Topology 14 (1975) 49–57 [MR0365581](#)
- [15] **V M Lew**, *The semistability at infinity for multiple extension groups*, PhD thesis, Cornell University, Ithaca, NY (1993) [MR2689313](#) Available at <http://search.proquest.com/docview/304035762>
- [16] **M L Mihalik**, *Semistability at the end of a group extension*, Trans. Amer. Math. Soc. 277 (1983) 307–321 [MR690054](#)
- [17] **M L Mihalik**, *Ends of groups with the integers as quotient*, J. Pure Appl. Algebra 35 (1985) 305–320 [MR777262](#)
- [18] **M L Mihalik**, *Ends of double extension groups*, Topology 25 (1986) 45–53 [MR836723](#)
- [19] **M Mihalik**, *Semistability at ∞ of finitely generated groups, and solvable groups*, Topology Appl. 24 (1986) 259–269 [MR872498](#)
- [20] **M L Mihalik**, *Semistability at infinity, simple connectivity at infinity and normal subgroups*, Topology Appl. 72 (1996) 273–281 [MR1406313](#)
- [21] **J S Profio**, *Using subnormality to show the simple connectivity at infinity of a finitely presented group*, Trans. Amer. Math. Soc. 320 (1990) 281–292 [MR961627](#)
- [22] **Y Shalom, G A Willis**, *Commensurated subgroups of arithmetic groups, totally disconnected groups and adelic rigidity*, Geom. Funct. Anal. 23 (2013) 1631–1683 [MR3102914](#)
- [23] **J Stallings**, *The piecewise-linear structure of Euclidean space*, Proc. Cambridge Philos. Soc. 58 (1962) 481–488 [MR0149457](#)

Department of Mathematics, Vanderbilt University
Nashville, TN 37240, United States

michael.l.mihalik@vanderbilt.edu

Received: 6 January 2016 Revised: 15 March 2016