

AG  
T

*Algebraic & Geometric  
Topology*

Volume 23 (2023)

**Discrete real specializations of sesquilinear representations  
of the braid groups**

NANCY SCHERICH



# Discrete real specializations of sesquilinear representations of the braid groups

NANCY SCHERICH

Using Salem numbers, this paper gives real specializations of sesquilinear representations of the braid groups that make the images discrete groups. This method is applied to the Burau, Jones and Lawrence–Krammer representations, and some details on the commensurability of the target groups are given.

20C08, 57M99

## 1 Introduction

Representations of the braid groups have attracted attention because of their wide variety of applications from discrete geometry to quantum computing. This paper takes the point of view that one should ask structural questions about the image of a braid group representation, in particular whether the image is discrete for specializations of the parameter. Venkataramana in [15] also followed this pursuit for discrete specializations of the Burau representation but with a different approach toward arithmeticity.

Since the Jones representations are used in modeling quantum computations, much work has been done to understand specializations at roots of unity, as explored by Funar and Kohno in [7], Freedman, Larsen and Wang in [6], and many others. However, there seems to be a lack of exploration of the *real* specializations of these representations. This paper takes a more general approach to find discrete real specializations of any sesquilinear group representation, and show how this can be applied to representations of the braid groups. The main theorem follows.

**Theorem 1.1** *Let  $\rho_t : G \rightarrow \mathrm{GL}_m(\mathbb{Z}[t, t^{-1}])$  be a group representation with parameter  $t$ . Suppose there exists a matrix  $J_t$  such that:*

- (1) *For all  $M$  in the image of  $\rho_t$ ,  $M^* J_t M = J_t$ , where  $M^*(t) = M^\top(1/t)$ .*

- (2)  $J_t = (J_{1/t})^\top$ .
- (3)  $J_t \in \mathrm{GL}_m(\mathbb{Q}(t))$ , where no entry of  $J_t$  has denominator with 1 as a root.
- (4)  $J_t$  is positive definite for  $t$  in a neighborhood  $\eta$  of 1 in  $\mathbb{C}$ .

Then there exist infinitely many Salem numbers  $s$  such that the specialization representation  $\rho_s$  at  $t = s$  is discrete.

Further applying a classification theorem of hermitian forms from Scharlau [13] proves the following commensurability result of the target groups.

**Corollary 1.2** For  $\rho_t: G \rightarrow \mathrm{SL}_{2m+1}(\mathbb{Z}[t, t^{-1}])$  as in [Theorem 1.1](#), there exist infinitely many Salem numbers  $s$  such that for infinitely many integers  $n$  and  $k$  the specializations  $\rho_{s^k}$  at  $t = s^k$  and  $\rho_{s^n}$  at  $t = s^n$  map into commensurable lattices.

Squier showed in [14] that the reduced Burau representation is sesquilinear and satisfies the criteria for [Theorem 1.1](#). [Example 3.2](#) gives explicit Salem numbers such that specializing the reduced Burau representation to these numbers is discrete. Using the Burau representation as motivation, [Section 2](#) introduces the main tools of discrete generalized unitary groups and Salem numbers. In [Section 3](#) we apply [Theorem 1.1](#) to the Jones and Lawrence–Krammer representations of the braid group, and we suspect it also applies to all of the BMW representations. Lastly, [Section 4](#) discusses the lattice structure and commensurability of the target groups for the Salem number specializations.

## Acknowledgements

The results of this paper formed part of the author’s dissertation. The author would like to thank her advisor, Darren Long, for endless insight, guidance and support on this project, Vaughan Jones for helpful conversations, and Warren Scherich for moral support. This paper was funded in part by the NSF grant DMS-1463740.

## 2 Discrete representations using Salem numbers

### 2.1 Motivation from Squier and the Burau representation

The (reduced) Burau representation  $\rho_{n,t}: B_{n+1} \rightarrow \mathrm{GL}_n(\mathbb{Z}[t, t^{-1}])$  is an irreducible representation of the braid group. These representations depend on  $n$ , where  $n + 1$

is the number of braid strands, and are parametrized by a variable  $t$ . Squier showed in [14] that there is a nondegenerate  $n$ -dimensional matrix  $J_{n,t}$  satisfying the equation

$$(2-1) \quad M^* J_{n,t} M = J_{n,t},$$

for all  $M$  in the image of  $\rho_{n,t}$ . Here  $M^*$  is the transpose of  $M$  after replacing  $t$  with  $1/t$  in the entries of  $M$ ,  $M^*(t) := M(1/t)^\top$ .  $J_{n,t}$  is sesquilinear with respect to  $*$ ,  $J_{n,t}^* = J_{n,t}$ , and letting  $t = x^2$ ,  $J_{n,t}$  is given by the tridiagonal matrix

$$J_{n,t} = \begin{bmatrix} x + x^{-1} & -1 & & & \\ & -1 & \ddots & \ddots & \\ & & \ddots & \ddots & \\ & & & -1 & \\ & & & -1 & x + x^{-1} \end{bmatrix}.$$

If  $t$  is a unit complex number, (2-1) agrees with the usual unitary relation  $(\overline{M})^\top M = \text{Id}$ . Representations that satisfy (2-1) are called *sesquilinear*, and are said to map into a *generalized unitary group*. This terminology will be made precise in the next section.

These generalized unitary groups are the key to finding discrete specializations. The method described here is to show that carefully chosen specializations of the parameter  $t$  make the entire generalized unitary group discrete, thus making the image of the representation discrete.

## 2.2 Unitary groups

In general, unitary groups are matrix groups which respect a form, or inner product. These notions heavily rely on the ring of coefficients and an involution of that ring. Let  $R$  be a ring and  $\phi$  an order 2 automorphism of  $R$ . For a matrix  $M$  defined over  $R$ , let  $M^* = (M^\phi)^\top$ , where  $M^\phi$  is the matrix obtained by applying  $\phi$  to the entries of  $M$ . For the Burau representation in Section 2.1,  $\phi$  is the map given by  $t \mapsto 1/t$ , and  $R = \mathbb{Z}[t, t^{-1}]$ .

**Definition 2.1** For a matrix  $J$  such that  $J^* = J$ , the *generalized unitary group* is

$$U_m(J, \phi, R) := \{M \in \text{GL}_m(R) \mid M^* J M = J\}.$$

Here,  $J$  is called a *sesquilinear form* with respect to  $\phi$ . For example, in this notation the familiar unitary group  $U_m$  can be written as  $U_m(\text{Id}, -, \mathbb{C})$ , where “ $-$ ” is complex conjugation. A representation is called *sesquilinear* if its image is contained in a generalized unitary group.

**2.2.1 Creating discrete unitary groups** The Burau representation can be written as  $\rho_{n,t}: B_{n+1} \rightarrow U_n(J_{n,t}, \phi, \mathbb{Z}[t, t^{-1}])$ . With the goal of parameter specialization in mind, the relevant choice for the coefficient ring is a number ring. Discreteness of the unitary group is a delicate relationship between the form  $J$  and the algebraic structure of the number ring. More precisely, let  $L$  be a totally real algebraic field extension of  $\mathbb{Q}$  and let  $K$  be a degree 2 field extension of  $L$ , with  $L, K \subseteq \mathbb{C}$ . Let  $\phi$  be the order 2 generator of  $\text{Gal}(K/L)$ , and let  $\mathcal{O}_K$  and  $\mathcal{O}_L$  denote the rings of integers of  $K$  and  $L$ , respectively:

$$\begin{array}{ccc}
 \begin{array}{c} \curvearrowright^\phi \\ \mathcal{O}_K \\ \downarrow \\ \mathcal{O}_L \end{array} &
 \begin{array}{c} \curvearrowright^\phi \\ K \\ \downarrow 2 \\ L \end{array} &
 \begin{array}{c} \curvearrowright^{\phi^\sigma} \\ K^\sigma \subseteq \mathbb{C} \\ \\ L^\sigma \subseteq \mathbb{R} \end{array}
 \end{array}$$

Let  $\sigma$  be a complex place of  $K$ , which in this setting is a field homomorphism  $\sigma: K \rightarrow \mathbb{C}$  different from  $\phi$  and the identity map. We write  $X^\sigma = \sigma(X)$  for any  $X$  in  $K$ . The algebraic structure is passed along by  $\sigma$ , meaning  $\mathcal{O}_{K^\sigma} = (\mathcal{O}_K)^\sigma$  is the ring of integers for  $K^\sigma$  and  $\phi^\sigma = \sigma\phi\sigma^{-1}$  is an involution on  $K^\sigma$ .

Let  $J$  be a matrix over  $\mathcal{O}_K$  that is sesquilinear with respect to  $\phi$ . Then  $J^\sigma$  is sesquilinear with respect to  $\phi^\sigma$ . So, in particular,

$$U_m(J^\sigma, \phi^\sigma, \mathcal{O}_{K^\sigma}) = \{M \in \text{GL}_m(\mathcal{O}_{K^\sigma}) \mid (M^{\phi^\sigma})^\top J^\sigma M = J^\sigma\}.$$

Since  $\sigma$  is a homomorphism, we can see that  $(U_m(J, \phi, \mathcal{O}_K))^\sigma = U_m(J^\sigma, \phi^\sigma, \mathcal{O}_{K^\sigma})$  by applying  $\sigma$  to the equation  $J = M^*JM$ .

The following results outline compatibility requirements between  $J$  and  $\mathcal{O}_K$ , which show that  $U_m(J, \phi, \mathcal{O}_K)$  is a discrete subgroup of  $\text{GL}_m(\mathbb{R})$  under the standard euclidean topology.

**Proposition 2.2**  $U_m(J^\sigma, \phi^\sigma, \mathcal{O}_{K^\sigma})$  is a bounded group when  $J^\sigma$  is positive definite and  $\phi^\sigma$  is complex conjugation.

**Proof** Because  $J^\sigma$  is positive definite, by Sylvester’s law of inertia and the Gram-Schmidt process, there exists a matrix  $Q \in \text{GL}_m(\mathbb{C})$  such that  $J^\sigma = Q^* \text{Id } Q$ . This implies that  $QU_m(J^\sigma, \phi^\sigma, \mathcal{O}_{K^\sigma})Q^{-1} \subseteq U_m(\text{Id}, \phi^\sigma, \mathbb{C})$ , which is a subgroup of the compact group  $U_m$ . □

**Theorem 2.3**  $U_m(J, \phi, \mathcal{O}_K)$  is discrete if, for every complex place  $\sigma$  of  $K$ ,  $J^\sigma$  is positive definite and  $\phi^\sigma$  is complex conjugacy.

**Proof** Assume that  $\{M_n\}$  converges to the identity in  $U_m(J, \phi, \mathcal{O}_K)$ . To show  $\{M_n\}$  is eventually constant, we will show that for  $n$  large there are only finitely many possibilities for the entries  $(M_n)_{ij}$ .

By assumption, for each  $\sigma$  the group  $U_m(J^\sigma, \phi^\sigma, \mathcal{O}_{K^\sigma})$  is bounded by [Proposition 2.2](#). Also, for every  $M_n$ ,  $M_n^\sigma \in U_m(J^\sigma, \phi^\sigma, \mathcal{O}_{K^\sigma})$ . Thus, there exists a  $B$  such that for large  $n$ , for all  $i, j$ , and for all  $\sigma$ , we have that  $|(M_n^\sigma)_{ij}| < B$ .

For every  $M \in U_m(J, \phi, \mathcal{O}_K)$ , the equation  $M^*JM = J$  can be rearranged to  $JMJ^{-1} = ((M^\phi)^\top)^{-1}$ , showing that  $M$  and  $((M^\phi)^\top)^{-1}$  are simultaneously conjugate. Thus  $\{M_n^\phi\}$  also converges to the identity. Convergent sequences are bounded, so for large enough  $n$ ,  $|(M_n)_{ij}| < B$  and  $|(M_n)_{ij}^\phi| < B$  for every  $ij$ -entry.

$L$  is a totally real degree 2 subfield of  $K$ , and  $\phi$  generates  $\text{Gal}(K/L)$ . So  $K$  has one nonidentity real embedding  $\phi$ , and all other embeddings are complex. Thus we have shown above that for large  $n$  there is a uniform bound  $B$  for each entry  $(M_n)_{ij}$  and each Galois conjugate of  $(M_n)_{ij}$ . There are only finitely many algebraic integers  $\alpha$  such that  $\deg(\alpha) \leq \deg(K/\mathbb{Q})$ , and with the property that  $\alpha$  and all of the Galois conjugates of  $\alpha$  have absolute values bounded above by  $B$ . So there are only finitely many possible entries for  $(M_n)_{ij}$ , which implies the sequence  $\{M_n\}$  is eventually constant.  $\square$

**Corollary 2.4** If  $\rho: G \rightarrow U_m(J, \phi, \mathcal{O}_K)$  is a representation of a group  $G$  such that for every nonidentity place  $\sigma$  of  $K$ ,  $J^\sigma$  is positive definite and  $\phi^\sigma$  is complex conjugacy, then  $\rho$  is a discrete representation.

At first glance, the requirements for [Corollary 2.4](#) seem very specific and perhaps it is doubtful that any such a representation could exist. However, as described in [Section 2.1](#), Squier showed that the Burau representation maps into a generalized unitary group over  $\mathbb{Z}[t, t^{-1}]$ , so the next task is to find values of  $t$  such that the form and coefficient ring satisfy the specific hypothesis of [Corollary 2.4](#).

### 2.3 Salem numbers

Salem numbers are the key ingredient to the application of [Corollary 2.4](#), which requires a real algebraic number field with tight control and understanding of each of its complex embeddings.




Figure 1: A schematic picture of an order 6 Salem number.

**Definition 2.5** A Salem number  $s$  is a real algebraic unit greater than 1, with one real Galois conjugate  $1/s$ , and all of the complex Galois conjugates of  $s$  have absolute value equal to 1.

For example, the largest real root of Lehmer’s polynomial, called Lehmer’s number,

$$x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1,$$

is a Salem number. Trivial Salem numbers of degree 2 are solutions to  $s^2 - ns + 1$  for  $n \in \mathbb{N}$  and  $n > 2$ . It is well known that there are infinitely many Salem numbers of arbitrarily large absolute value and degree. In particular, if  $s$  is a Salem number, then  $s^m$  is also a Salem number for every positive integer  $m$ . One geometric consequence of the property that powers of Salem numbers are Salem numbers is that by taking powers, one can control the spatial configuration of the complex Galois conjugates of a Salem number, as described in [Lemma 2.6](#).

**Lemma 2.6** For any Salem number  $s$ , and for any interval containing 1 on the complex unit circle, there exist infinitely many integers  $m$  such that every complex Galois conjugate of  $s^m$  lies in the interval.

**Proof** Let  $e^{i\theta_1}, \dots, e^{i\theta_k}$  be all the Galois conjugates of the Salem number  $s$  with positive imaginary part. Suppose that  $\prod_{j=1}^k (e^{i\theta_j})^{m_j} = 1$ . Let  $\phi$  be the automorphism of the Galois closure of  $s$  with the property that  $\phi(e^{i\theta_1}) = s$ . Since  $\phi$  must permute the




Figure 2: A schematic picture of [Lemma 2.6](#).

Galois conjugates of  $s$ , for  $j \neq 1$ ,  $\phi(e^{i\theta_j})$  is again on the complex unit circle. Thus,

$$1 = \phi\left(\prod_{j=1}^k (e^{i\theta_j})^{m_j}\right) = s^{m_1} \prod_{j=2}^k \phi(e^{i\theta_j})^{m_j}, \quad \text{which implies} \quad \prod_{j=2}^k \phi(e^{i\theta_j})^{m_j} = 1/s^{m_1}.$$

Since each  $\phi(e^{i\theta_j})$  is a unit complex number, it must be the case that each  $m_j = 0$ . This shows that the point  $p = (e^{i\theta_1}, \dots, e^{i\theta_k})$  satisfies the criteria for Kronecker's theorem. In particular, the set  $\{p^m \mid m \in \mathbb{Z}\}$  is dense in the torus  $T^k$ . □

Fixing an arbitrary Salem number  $s$ , let  $K = \mathbb{Q}(s)$ ,  $L = \mathbb{Q}(s + 1/s)$  and  $\mathcal{O}_K$  be the ring of integers of  $K$ :

$$\begin{array}{c} \mathbb{Q}(s) = K \\ \quad \quad \quad \downarrow 2 \\ \mathbb{Q}(s + 1/s) = L \\ \quad \quad \quad \downarrow \\ \mathbb{Q} \end{array}$$

Since  $s$  and  $1/s$  are real and all other Galois conjugates of  $s$  are complex,  $K$  has exactly two real embeddings. For a complex embedding  $\sigma$  of  $K$ ,  $(s + 1/s)^\sigma = 2 \operatorname{Re}(s^\sigma)$ , which is real. This shows that all embeddings of  $L$  are real, and that  $L$  is a totally real subfield of  $K$ . Since  $s$  is a root of  $X^2 - (s + 1/s)X + 1$ ,  $K$  is degree 2 over  $L$ .

The Galois group of  $K/L$  is generated by  $\phi$ , which maps  $s$  to  $1/s$ . (This exactly matches the involution  $t \mapsto 1/t$  needed in the sesquilinear condition for the Burau representation.) On the complex unit circle, inversion is the same as complex conjugation. So for the complex embeddings  $\sigma$  of  $K$ ,  $\phi^\sigma$  is complex conjugacy. Notice for a sesquilinear matrix  $J_t$  over  $\mathcal{O}_K$  with a parameter  $t$ , specializing  $t = s$  leaves  $J_s^\sigma$  hermitian.

**Theorem 1.1** *Let  $\rho_t: G \rightarrow \operatorname{GL}_m(\mathbb{Z}[t, t^{-1}])$  be a representation of a group  $G$ . Suppose there exists a matrix  $J_t$  such that:*

- (1)  $M^* J_t M = J_t$  for all  $M$  in the image of  $\rho_t$ .
- (2)  $J_t = (J_{1/t})^\top$ .
- (3)  $J_t \in \operatorname{GL}_m(\mathbb{Q}(t))$ , where no entry of  $J_t$  has denominator with 1 as a root.
- (4)  $J_t$  is positive definite for  $t$  in a neighborhood  $\eta$  of 1 in  $\mathbb{C}$ .

*Then there exist infinitely many Salem numbers  $s$  such that the specialization  $\rho_s$  at  $t = s$  is discrete.*



**Proof** The neighborhood  $\eta$  can be chosen so that no entry of  $J_t$  has a denominator with a root in  $\eta$ . By [Lemma 2.6](#), there are infinitely many Salem numbers with the property that all the complex Galois conjugates lie in  $\eta$ . Let  $s$  be one such Salem number. Specializing  $t$  to  $s$  gives  $\rho_s: G \rightarrow U_m(J_s, \phi, O_{\mathbb{Q}(s)})$ , where  $\phi$  is the usual map given by  $s \mapsto 1/s$ .

Let  $\sigma$  be a complex place of  $\mathbb{Q}(s)$  which is given by  $s \mapsto z$  for  $z$  a complex Galois conjugate of  $s$ . Then  $J_s^\sigma = J_z$ , and since  $z \in \eta$ ,  $J_z$  is positive definite. By [Corollary 2.4](#), the specialization  $\rho_s$  at  $t = s$  is discrete. □

**Remark 2.7** If the representations in [Theorem 1.1](#) all have determinant 1, then the image is more than just discrete, and in fact is a subgroup of a lattice. See [Section 4](#) for more details.

### 3 Applications to braid group representations

#### 3.1 The Burau representation

**Proposition 3.1** *There are infinitely many Salem numbers  $s$  such that the Burau representation specialized to  $t = s$  is discrete.*

**Proof** The specialization of  $\rho_{n,1}$  at  $t = 1$  collapses to an irreducible representation of the symmetric group. As a representation of a finite group,  $\rho_{n,1}$  fixes a positive definite form which is unique up to scaling, by [Lemma 3.7](#), which is proved later. At  $t = 1$ ,  $J_{n,1}$  is positive definite, and the signature of  $J_{n,t}$  can only change at zeroes of its determinant.

An inductive computation shows that  $\det(J_{n,t}) = (t^{2n+2} - 1)/(t^n(t^2 - 1))$  for  $t \neq 1$ , and the zeroes of  $\det(J_{n,t})$  occur at  $(n+1)^{\text{th}}$  roots of unity. Thus,  $J_{n,t}$  remains positive definite for unit complex values of  $t$  with argument less than  $2\pi/(n+1)$ . This shows the Burau representation satisfies the criteria of [Theorem 1.1](#). □

**Example 3.2** The Burau representation  $\rho_{4,t}$  of  $B_4$  is discrete when specializing  $t$  to the following Salem numbers:

- Lehmer’s number raised to the powers 16, 32 and 47,
- the largest real root of  $1 - x^4 - x^5 - x^6 + x^{10}$  raised to the powers 17, 23, and 43.

## 3.2 The Jones representations

The *Hecke algebra* (of type  $A_n$ ), denoted by  $H_n(q)$ , is the complex algebra generated by invertible elements  $g_1, \dots, g_{n-1}$  with relations

$$(3-1) \quad \begin{aligned} g_i g_{i+1} g_i &= g_{i+1} g_i g_{i+1} && \text{for all } i < n, \\ g_i g_j &= g_j g_i && \text{for } |i - j| > 1, \\ g_i^2 &= (1 - q)g_i + q && \text{for all } i < n. \end{aligned}$$

Here,  $q$  is viewed as a complex parameter.  $H_n(q)$  is a quotient of  $\mathbb{C}[B_n]$  by the relation (3-1). The *Jones representations* of  $B_n$  are defined by precomposing a representation of  $H_n(q)$  by the quotient map from  $\mathbb{C}[B_n]$ . The Jones representations have matrix entries in  $\mathbb{Z}[t, t^{-1}]$ .

**3.2.1 The Jones representations are sesquilinear** The Hecke algebras have a natural automorphism, denoted here by  $\phi$ , which sends  $q$  to  $1/q$ . Taking  $q$  to be a unit complex number, this automorphism becomes complex conjugacy. The Jones representations are known to be sesquilinear with respect to  $\phi$  for various complex specializations of  $q$  and with many different types of proofs, as in [2; 3; 9; 17]. To be overtly clear that all of the criteria of [Theorem 1.1](#) are satisfied by the Jones representations, we provide a simple proof of sesquilinearity here that is very similar to [3, Proposition 3.7] by Brunat and Marin.

The irreducible representations of  $H_n(q)$  are parametrized by the Young diagrams. (For a more detailed discussion of Young diagrams see [18], and for a construction of the Jones representations see [16].) Every Young diagram contains sub-Young diagrams, obtained by removing boxes in a way that retains the weakly decreasing row length condition. If  $\lambda$  is a Young diagram with  $n$  boxes, then we will call the sub-Young diagrams found by removing one box from  $\lambda$  the  $(n-1)$ -subdiagrams of  $\lambda$ . A Young diagram is completely determined by any two of its  $(n-1)$ -subdiagrams. These  $(n-1)$ -subdiagrams also determine representations of the Hecke algebras in a powerful way. The following theorem, originally due to Curtis, Iwahori and Kilmoyer in [5] and popularized by Jones in [8], states concretely the relationship between Young diagrams and the representations of the Hecke algebras.

**Theorem 3.3** *Up to equivalence, the finite-dimensional irreducible representations of  $H_n(q)$ , for generic  $q$ , are in one to one correspondence with the Young diagrams of  $n$  boxes. Moreover, if  $\rho$  is a representation corresponding to Young diagram  $\lambda$ , then  $\rho$*

restricted to  $H_{n-1}(q)$  is equivalent to the representation  $\bigoplus_{i=1}^k \rho_{\lambda_i}$ , where  $\lambda_1, \dots, \lambda_k$  are all of the  $(n-1)$ -subdiagrams of  $\lambda$  and each  $\rho_{\lambda_i}$  is an irreducible representation of  $H_{n-1}(q)$  corresponding to  $\lambda_i$ .

Here equivalence means the existence of an intertwining isomorphism, made precise by the following definition.

**Definition 3.4** The representations  $\varphi : G \rightarrow GL(V)$  and  $\psi : G \rightarrow GL(W)$  are *equivalent* if there exists a linear isomorphism  $T : V \rightarrow W$  such that  $T\varphi(g)(v) = \psi(g)T(v)$  for all  $g \in G$  and  $v \in V$ , or that the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\varphi(g)} & V \\ \downarrow T & & \downarrow T \\ W & \xrightarrow{\psi(g)} & W \end{array}$$

Choosing bases for  $V$  and  $W$ , the equivalence  $T$  gives the matrix equation

$$[T][\varphi(g)][T]^{-1} = [\psi(g)].$$

At the level of matrices, representations are equivalent exactly when they are simultaneously conjugate. In the context of [Theorem 3.3](#), the restriction of  $\rho$  to  $H_{n-1}(q)$  is equivalent to the representation  $\bigoplus_{i=1}^k \rho_{\lambda_i}$ , which means there is a change of basis such that the restriction of  $\rho$  is block diagonal. These restriction rules are combinatorially depicted in the Young lattice of Young diagrams; see [\[18\]](#).

A representation is sesquilinear if there exists an invertible matrix  $J$  such that for every  $M$  in the image of the representation the following equation is satisfied:

$$(3-2) \quad M^*JM = J.$$

Rearranging this equation, we see that  $M = J^{-1}((M\phi)^\tau)^{-1}J$ , showing that  $M$  and  $((M\phi)^\tau)^{-1}$  are simultaneously conjugate and the conjugating matrix  $J$  is the sesquilinear form. Changing views slightly, consider the following definition.

**Definition 3.5** For  $\varphi : G \rightarrow GL(V)$  a complex linear representation,  $\tilde{\varphi} : G \rightarrow GL(V^*)$  is called the  $\phi$ -twisted contragredient representation of  $\varphi$  and is given by


$$\tilde{\varphi}(g)f(v) = f(\varphi(g^{-1})^\phi v)$$

for every  $g \in G$ ,  $v \in V$  and  $f \in V^*$ .

If a basis for  $V$  is chosen, then as matrices,  $[\tilde{\varphi}(g)] = ([\varphi(g)^\phi]^\top)^{-1}$ . So another way to view a sesquilinear representation is one that is *equivalent to its  $\phi$ -twisted contragredient*.

**Lemma 3.6** *Every finite-dimensional irreducible representation of the Hecke algebra is equivalent to its  $\phi$ -twisted contragredient representation, when  $q$  is a generic complex number.*

**Proof** We can establish this result for  $n = 3$ . There are three nonequivalent irreducible representations of  $H_3(q)$  corresponding to the following Young diagrams:



Up to equivalence, the first two representations are 1-dimensional, given by  $g_i \mapsto q$  and  $g_i \mapsto -1$ , and are in fact equal to their  $\phi$ -twisted contragredient representations. The third representation is known to be the Burau representation for  $B_3$ . As described earlier, Squier showed that the Burau representations are sesquilinear and are therefore equivalent to their  $\phi$ -twisted contragredient.

Inductively moving forward, let  $\rho: H_n(q) \rightarrow GL(V)$  be a finite-dimensional irreducible representation and  $\tilde{\rho}$  be the  $\phi$ -twisted contragredient representation of  $\rho$ . Up to equivalence,  $\rho$  corresponds to a Young diagram  $\lambda$ . To show that  $\rho$  and  $\tilde{\rho}$  are equivalent, it suffices to show that both representations correspond to the same  $\lambda$ . A Young diagram is completely characterized by its list of  $(n-1)$ -subdiagrams, which correspond to the restriction of the representation to  $H_{n-1}(q)$ . So it is enough to show that the restrictions of  $\rho$  and  $\tilde{\rho}$  correspond to the same list of  $(n-1)$ -subdiagrams.

Writing  $\rho| = \rho|_{H_{n-1}(q)}$ , by [Theorem 3.3](#) there is an equivalence  $T$  such that

$$T\rho|(h)T^{-1} = \bigoplus_{i=1}^k \rho_{\lambda_i}(h) \quad \text{for every } h \in H_{n-1}(q),$$

where each  $\lambda_i$  is an  $(n-1)$ -subdiagram of  $\lambda$ ,  $k$  is the number of  $(n-1)$ -subdiagrams of  $\lambda$ , and  $\rho_{\lambda_i}$  is an irreducible representation of  $H_{n-1}(q)$  corresponding to  $\lambda_i$ . Choosing a basis for  $V$ , the matrix for  $[T\rho|(h)T^{-1}]$  is block diagonal. Taking the  $\phi$ -twisted contragredient of a block diagonal matrix preserves the block decomposition, which gives

$$([T^\phi]^\top)^{-1}[\tilde{\rho}|(h)][T^\phi]^\top = \bigoplus_{i=1}^k [\tilde{\rho}_{\lambda_i}(h)] \quad \text{for every } h \in H_{n-1}(q).$$

This equation shows that  $|\tilde{\rho}|$  is equivalent to  $\bigoplus \tilde{\rho}_{\lambda_i}$ . Since each  $\rho_{\lambda_i}$  is an irreducible representation of  $H_{n-1}(q)$ , we can inductively assume that  $\rho_{\lambda_i}$  is equivalent to  $\tilde{\rho}_{\lambda_i}$ , for all  $i \leq k$ . Therefore,  $\rho_{\lambda_i}$  and  $\tilde{\rho}_{\lambda_i}$  correspond to the same Young diagram  $\lambda_i$ . Thus the restrictions of  $\rho$  and  $\tilde{\rho}$  correspond to the same list of  $(n-1)$ -subdiagrams.  $\square$

**Lemma 3.7** *If an absolutely irreducible matrix representation has an invertible matrix  $J$  satisfying  $M^*JM = J$  for all  $M$  in the representation, then  $J$  is unique up to scaling.*

**Proof** Suppose there were two such matrices  $J_1$  and  $J_2$ . Then (3-2) gives, for all matrices  $M$  in the representation,

$$J_1MJ_1^{-1} = ((M^\phi)^\top)^{-1} = J_2MJ_2^{-1} \implies (J_1^{-1}J_2)^{-1}M(J_1^{-1}J_2) = M.$$

This shows that  $J_1^{-1}J_2$  is in the centralizer of the entire irreducible representation. Schur’s lemma gives that  $J_1^{-1}J_2 = \alpha \cdot \text{Id}$  for some scalar  $\alpha$ , and finally  $J_2 = \alpha J_1$ .  $\square$

**Proposition 3.8** *If  $\rho$  is an irreducible Jones representation of  $B_n$  and  $q$  is a generic unit complex number close to 1, then there exists a nondegenerate, positive definite, sesquilinear matrix  $J$  with entries in  $\mathbb{Q}(q)$  such that for all  $M$  in the image of  $\rho$ ,  $(M^\phi)^\top JM = J$ .*

**Proof** Let  $\rho$  be a finite-dimensional irreducible representation of  $H_n(q)$  over  $V$ . By Lemma 3.6,  $\rho$  is equivalent to its  $\phi$ -twisted contragredient representation  $\tilde{\rho}$  by an equivalence  $T$ . Choose a basis for  $V$  and its dual basis for  $V^*$ , and let  $\mathcal{T}$  be the matrix for  $T$  with respect to these bases. We will use this matrix  $\mathcal{T}$  to find the desired matrix  $J$ . Let the superscript  $*$  denote the  $\phi$ -twisted transpose of a matrix to ease computation. For all  $g \in H_n(q)$ , we get the matrix equations

$$\begin{aligned} (3-3) \quad \mathcal{T}[\rho(g)]\mathcal{T}^{-1} = [\tilde{\rho}(g)] = ([\rho(g)]^{-1})^* &\implies (\mathcal{T}^{-1})^*[\rho(g)]^*\mathcal{T}^* = [\rho(g)]^{-1} \\ &\implies \mathcal{T}^*[\rho(g)](\mathcal{T}^*)^{-1} = ([\rho(g)]^{-1})^*. \end{aligned}$$

This shows that  $\mathcal{T}$  and  $\mathcal{T}^*$  are two possible forms for  $\rho$ . By Lemma 3.7,  $\mathcal{T} = \alpha\mathcal{T}^*$  for some  $\alpha \in \mathbb{C}$ . Applying  $*$  again gives  $\mathcal{T} = \alpha\alpha^*\mathcal{T}$  and  $\alpha\alpha^* = 1$ .

Define  $J = \beta\mathcal{T} + \beta^*\mathcal{T}^* = (\alpha\beta + \beta^*)\mathcal{T}^*$  where  $\beta$  is as follows. (Here  $\beta$  is needed to ensure that  $J$  is invertible.) If  $\alpha \neq -1$ , let  $\beta = 1$ , which gives that  $\det J = \det((\alpha + 1)\mathcal{T})$ , which is nonzero. If  $\alpha = -1$ , let  $\beta \in \mathbb{C}$  be such that  $\beta^* \neq \beta$ . Then

$$\det J = \det[(\alpha\beta + \beta^*)\mathcal{T}^*] = \det[(-\beta + \beta^*)\mathcal{T}]$$

is nonzero. So, in both cases,  $J$  is invertible.

Next note that  $J$  is sesquilinear, that is,  $J^* = (\beta\mathcal{T} + \beta^*\mathcal{T}^*)^* = \beta^*\mathcal{T}^* + \beta\mathcal{T} = J$ . If  $M$  is a matrix in the image of  $\rho$ , rearranging the right-hand equation of (3-3) gives  $M^*\mathcal{T}^*M = \mathcal{T}$ . So inserting  $J$  gives

$$M^*JM = M^*(\alpha\beta + \beta^*)\mathcal{T}^*M = (\alpha\beta + \beta^*)M^*\mathcal{T}^*M = (\alpha\beta + \beta^*)\mathcal{T} = J.$$

To show that the entries of  $J$  are in  $\mathbb{Q}(q)$  we will proceed by induction on  $n$ , as in the proof of Lemma 3.6. As a base case with  $n = 3$ , Squier's form for the Burau representation has entries in  $\mathbb{Q}(q)$ . Let  $\rho$  be an irreducible Jones representation of  $B_n$  with  $\rho|_{B_{n-1}} = \bigoplus_{i=1}^k \rho_i$ , where each  $\rho_i$  is an irreducible Jones representation of  $B_{n-1}$ . We can inductively assume each  $\rho_i$  is sesquilinear with form  $J_i$  whose coefficients are in  $\mathbb{Q}(q)$ . Thus, there exist some scalars  $\alpha_i$  such that  $J = [\alpha_i J_i]$ , the block diagonal matrix, and  $J$  is the sesquilinear form for  $\rho$ .

It remains to show that  $J$  is positive definite. Taking  $q = 1$ ,  $\rho$  is an irreducible representation of the symmetric group  $\Sigma_n$ . As a linear representation of a finite group,  $V$  admits an inner product that is invariant under the action of  $\Sigma_n$ , given by a positive definite nondegenerate matrix  $\hat{J}$ . Lemma 3.7 guarantees that  $\hat{J}$  is unique up to scaling. Since  $J|_{q=1}$  is also a form for this representation, it must be that  $\hat{J}$  is a multiple of  $J|_{q=1}$ , which gives that  $J$  is positive definite for  $q = 1$ . Since  $J$  is Hermitian for unit complex  $q$  it has real eigenvalues, and continuity of the determinant map finally gives that either  $J$  or  $-J$  is positive definite for  $q$  close to 1.  $\square$

**Corollary 3.9** *For each irreducible Jones representation, there are infinitely many Salem numbers  $s$  such that specializing  $q = s^m$ , for some  $m$ , is a discrete representation.*

### 3.3 The Lawrence–Krammer and BMW representations

The BMW algebras are a 2-parameter family of algebras, denoted by  $C_n(l, m)$ , with  $n - 1$  generators and parameters  $l$  and  $m$ . The BMW representations of the braid group come from representations of the BMW algebras [1; 12]. Similar to how the Burau representation is one irreducible summand of the Jones representations, Zinno proved in [19] that the Lawrence–Krammer representation is one summand of the BMW representations. To make sense of the  $*$  operation, the relevant involution for the BMW algebra is given by  $l \mapsto 1/l$ ,  $m \mapsto m$  and  $\alpha \mapsto 1/\alpha$ , where  $m = \alpha + 1/\alpha$ . Budney proved that the Lawrence–Krammer representation is sesquilinear [4], and Brunat and Marin give a more general proof that all the BMW representations are sesquilinear [3]; see also [10]. It is also known that the sesquilinear forms  $J_{l,m}$  are

positive definite for a neighborhood of  $(1, 1)$  in the unit complex sphere in  $\mathbb{C}^2$  [17]; see [2, Theorem 1.2] for a concise restatement. It is suspected that the forms  $J_{l,m}$  have coefficients in  $\mathbb{Q}(m, l)$ , and this is known to be true for the Lawrence–Krammer representation.

**Corollary 3.10** *For the Lawrence–Krammer representation there are infinitely many Salem numbers  $s$  such that specializing  $l = s^{k_1}$  and  $\alpha = s^{k_2}$ , for some  $k_1$  and  $k_2$ , is a discrete representation.*

**Example 3.11** Let  $\rho$  be the Lawrence–Krammer representation of  $B_4$  given on page 272 of [1]. Taking the Salem number  $S = 1/2 + 1/\sqrt{2} + 1/(2\sqrt{-1 + 2\sqrt{2}})$ , specializing  $\alpha = S^{15}$  and  $l = S^3$  makes  $\rho$  a discrete representation.

## 4 Commensurability

Ideally, we would like to find *real* specializations so that the Jones representations have images that are not just discrete, but are arithmetic groups or lattices in  $\mathrm{GL}_n(\mathbb{R})$ . A first step in this direction is to further study the unitary groups coming from Salem number specializations, and consider when the images are subsets of lattices. Specializing to two different powers of the same Salem number can give *commensurable* unitary groups, but the defining sesquilinear forms might be very different.

Recall the notation of  $K$ ,  $L$ ,  $\mathcal{O}_K$  and  $\phi$  from Section 2.3. In general, fixing a number ring  $\mathcal{O}_K$  and dimension  $m$ , the group  $U_m(J, \phi, \mathcal{O}_K)$  is determined by the form  $J$ . Notice that  $U_m(J, \phi, \mathcal{O}_K) = U_m(\lambda J, \phi, \mathcal{O}_K)$  for every  $\lambda \in L$ , and that the form  $J$  is not completely unique. This motivates the following definition.

**Definition 4.1** Matrices  $J$  and  $H$  are equivalent over  $K$  if  $Q^* J Q = \lambda H$  for some  $Q \in \mathrm{GL}_m(K)$  and  $\lambda \in \mathrm{Fix}(\phi)$ .

It would be nice if equivalent forms gave rise to *equal* unitary groups, but this is not true in general. However, in the careful scenario that the unitary group is a lattice in  $\mathrm{SL}_m(\mathbb{R})$ , changing the form by equivalence yields “the same” lattice, up to commensurability in the following sense.

**Definition 4.2** Two groups  $G_1$  and  $G_2$  are *commensurable* if there are finite-index subgroups  $H_1 \subseteq G_1$  and  $H_2 \subseteq G_2$  such that  $H_1$  is isomorphic to  $H_2$ .

**Definition 4.3** A lattice in a semisimple Lie group is a discrete subgroup with finite covolume.

For our purposes, we will take  $SL_m(\mathbb{R})$  or  $PSL_m(\mathbb{R})$  as the semisimple Lie group.

**Proposition 4.4** Assume that  $SU_m(J_1, \phi, \mathcal{O}_K)$  and  $SU_m(J_2, \phi, \mathcal{O}_K)$  are lattices in  $SL_m(\mathbb{R})$ . If  $J_1$  and  $J_2$  are equivalent over  $K$ , then  $SU_m(J_1, \phi, \mathcal{O}_K)$  is commensurable to  $SU_m(J_2, \phi, \mathcal{O}_K)$

**Proof** Let  $\lambda J_1 = Q^* J_2 Q$  for some  $Q \in GL_m(K)$  and  $\lambda \in \text{Fix}(\phi)$ . For notational clarity, write  $SU(J_i, \mathcal{O}_K) = SU_m(J_i, \phi, \mathcal{O}_K)$ .

Since scalar multiplication commutes with matrix multiplication,  $M^* J M = J$  if and only if  $M^* \lambda J M = \lambda J$ . So scaling the form preserves the unitary group, and without loss of generality we may assume  $\lambda = 1$ .

It is clear that  $M^* J M = J$  if and only if  $(Q^* M^* Q^{*-1})(Q^* J Q)(Q^{-1} M Q) = Q^* J Q$ , which seems like it implies that  $SU(Q^* J_1 Q, \mathcal{O}_K) = Q^{-1} SU(J_1, \mathcal{O}_K) Q$ . However, since  $Q$  has coefficients in  $K$ ,  $Q^{-1} M Q$  may not have coefficients in  $\mathcal{O}_K$ , so we can only conclude that  $Q^{-1} SU(J, \mathcal{O}_K) Q \subseteq SU(Q^* J Q, K)$ . To avoid this, we need to pass to a finite-index subgroup.

Since  $K$  is the ring of fractions of  $\mathcal{O}_K$ , there exists  $\gamma \in \mathcal{O}_K$  such that  $\gamma Q \in M_m(\mathcal{O}_K)$ . As a ring of integers of an algebraic extension,  $\mathcal{O}_K$  is a Dedekind domain and every quotient is finite. So  $\mathcal{O}_K / \langle \gamma^2 \rangle$  is finite and  $SU(J_1, \mathcal{O}_K / \langle \gamma^2 \rangle)$  is finite. The kernel  $N$  of the quotient map  $SU(J_1, \mathcal{O}_K) \rightarrow SU(J_1, \mathcal{O}_K / \langle \gamma^2 \rangle)$  has finite index in  $SU(J_1, \mathcal{O}_K)$ .

Any element  $B$  in the kernel has the form  $B = \text{Id} + \gamma^2 A$  for some matrix  $A \in M_m(\mathcal{O}_K)$ . Substituting  $Q^* J_2 Q$  for  $J_1$  in the equation  $B^* J_1 B = J_1$  gives that  $Q B Q^{-1}$  fixes the form  $J_2$ . Because  $Q$  has coefficients over  $K$ ,  $Q B Q^{-1}$  has coefficients in  $K$  and not necessarily in  $\mathcal{O}_K$ . However, since  $Q B Q^{-1} = \text{Id} + (\gamma Q) A (\gamma Q^{-1})$ , and both  $A$  and  $\gamma Q$  are integral,  $Q B Q^{-1}$  is also integral. Thus  $Q B Q^{-1} \in SU(J_2, \mathcal{O}_K)$ .

Since  $SU(J_1, \mathcal{O}_K)$  is a lattice and  $N$  is a finite-index subgroup,  $N$  is also a lattice in  $SL(\mathbb{R})$  with finite covolume. Thus  $Q N Q^{-1}$  has finite covolume in  $SL(\mathbb{R})$  and is therefore a lattice. So  $Q N Q^{-1}$  is a sublattice of  $SU(J_2, \mathcal{O}_K)$  and must have finite index by Margulis's theorem for lattices.

This shows that  $N$  is a finite-index subgroup of  $SU(J_1, \mathcal{O}_K)$  and  $Q N Q^{-1}$  has finite index in  $SU(J_2, \mathcal{O}_K)$ . □



So how does this lattice information apply to the Jones representations? Firstly, after rescaling and reparametrization the Jones representations can be made to have determinant  $\pm 1$ , allowing the image to land in  $\text{PSU}(J, \phi, \mathcal{O}_K)$  instead of just  $U(J, \phi, \mathcal{O}_K)$ . Secondly, an arithmetic group theory result of Harish and Chandra that is formalized in our setting in Chapter 6 of Morris [11], states that  $\text{SU}_m(J, \phi, \mathcal{O}_K)$  is a lattice in  $\text{SL}_m(\mathbb{R})$  under the exact Salem number circumstances as required by Theorem 1.1. So Corollary 3.9 can be restated using this new vocabulary.

**Corollary 4.5** *For each irreducible Jones representation, after a change of parameter, there are infinitely many Salem numbers  $s$  such that specializing  $q$  to a power of  $s$  maps into a lattice in  $\text{PSL}_m(\mathbb{R})$ .*

**Proof** Let  $\rho_q$  be an irreducible Jones representation of dimension  $m$ . The images of the braid generators under  $\rho_q$  have determinant  $\pm q^k$  for some  $k \in \mathbb{N}$ . After a change of variable  $q = y^m$  and scaling the generators by  $1/y^{m-k}$ , this adjusted representation  $\tilde{\rho}_y$  maps into  $\text{PSU}_m(J^y, \mathbb{Z}[y^{\pm 1}])$ .

The subgroup  $B_n^{\text{even}}$  of even braids (the preimage of the alternating group under the standard projection to  $S_n$ ) is a noncentral normal subgroup of  $B_n$  of finite index. The restriction  $\tilde{\rho}_y|_{B_n^{\text{even}}}$  maps  $B_n^{\text{even}}$  into  $\text{SU}_m(J_y, \mathbb{Z}[y^{\pm 1}])$ , and by Theorem 1.1 there exist infinitely many Salem numbers  $s$  such that the specialization  $\rho_s|_{B_n^{\text{even}}}$  at  $y = s$  is discrete. Further, by the results in Chapter 6 of [11] described above, these specializations make  $\text{SU}_m(J_s, \mathcal{O}_K)$  lattices in  $\text{SL}_m(\mathbb{R})$ . Finite-index arguments imply  $\text{PSU}_m(J_s, \mathcal{O}_K)$  is a lattice in  $\text{PSL}_m(\mathbb{R})$ .  $\square$

Since the goal is to obtain commensurable lattices as images of our Jones representations, and it is more natural to think of lattices in  $\text{SL}_m(\mathbb{R})$  instead of in  $\text{PSL}_m(\mathbb{R})$ , one may simply pass to the finite-index subgroup  $B_n^{\text{even}}$  and continue to think only about lattices in  $\text{SL}_m(\mathbb{R})$ . To apply Proposition 4.4 requires equivalent defining forms. In general, it is difficult to determine when two forms are equivalent. The following theorem gives a complete classification of the sesquilinear forms in a very specific algebraic setting that applies to the Salem number field scenario.

**Theorem 4.6** (Scharlau [13, Chapter 10]) *If  $L$  is a global field and  $K = L(\sqrt{\delta})$ , sesquilinear forms over  $K/L$  are classified by dimension, determinant class and the signatures for those orderings of  $L$  for which  $\delta$  is negative.*

This classification relies on the *determinant class*, which is defined here. Recall for a Salem number  $s$  the following tower of fields:

$$\begin{array}{ccc}
 \mathbb{Q}(s) = K & & K^\times \\
 \downarrow 2 & & \downarrow \text{Norm} \\
 \mathbb{Q}(s + 1/s) = L & & L^\times \quad (L^\times)^2 \subseteq \text{Norm}(K^\times) \\
 \downarrow & & \\
 \mathbb{Q} & & 
 \end{array}$$

The Galois group of  $K/L$  is generated by  $\phi$ , which maps  $s$  to  $1/s$ . There is a multiplicative group homomorphism  $\text{Norm}: K^\times \rightarrow L^\times$  given by  $\text{Norm}(\alpha) = \alpha\alpha^\phi$ , where  $K^\times = K - \{0\}$ . Notice for  $\beta \in L$  we have  $\text{Norm}(\beta) = \beta\beta^\phi = \beta^2$ , and so  $(L^\times)^2 \subseteq \text{Norm}(K)$ .

**Definition 4.7** The *determinant class* of a sesquilinear form  $H$  over  $K/L$  is the coset of  $\det(H)$  in  $K^\times / \text{Norm}(K^\times)$ :

$$[\det(H)] = \det(H) \text{Norm}(K).$$

Taking  $\delta = (s - 1/s)^2$ ,  $K$  can be rewritten as  $K = L(\sqrt{\delta})$ . Thus we can restate Scharlau’s classification in the specific context of Salem numbers.

**Theorem 4.8** (Scharlau restated) *Sesquilinear forms over  $K/L$  are classified by dimension, determinant class and the signatures for those orderings of  $L$  for which  $(s - 1/s)^2$  is negative.*

In odd dimensions it is very simple to show that *all* sesquilinear forms have the same determinant class, up to scaling. However, for even dimensions, the situation is very unclear.

**Proposition 4.9** *For every odd-dimensional invertible sesquilinear matrices  $H$  and  $J$  over  $K$ ,  $[\det(H)] = [\det(\lambda J)]$  for  $\lambda \in L$ .*

**Proof** Let  $H$  and  $J$  be sesquilinear matrices over  $K$  of dimension  $2k + 1$ . Since  $H$  and  $J$  are Hermitian, they are both diagonalizable with diagonal entries fixed by  $\phi$ . So, the determinants of both  $H$  and  $J$  are elements in  $L$ . Let  $d_H$  and  $d_J$  denote the nonzero determinants of  $H$  and  $J$ . Thus,

$$d_H = \frac{d_H}{d_J} d_J \stackrel{\text{mod}(L^\times)^2}{\equiv} \left(\frac{d_H}{d_J}\right)^{2k+1} d_J = \det\left(\frac{d_H}{d_J} J\right).$$

Since  $(L^\times)^2 \subseteq \text{Norm}(K)$ , we have that  $H$  and  $\lambda J$  have the same determinant class for  $\lambda = d_H/d_J \in L$ . □

As a result, to determine whether two forms of the same odd dimension are equivalent, it suffices to only check that they have the same signatures.

**Theorem 4.10** *For  $J_t$  a sesquilinear form that is positive definite for  $t$  in a neighborhood  $\eta$  of 1, there are infinitely many Salem numbers  $s$  and integers  $n$  and  $m$  such that, in all odd dimensions,  $\text{SU}_{2k+1}(J_{s^n}, \phi, \mathcal{O}_K)$  and  $\text{SU}_{2k+1}(J_{s^m}, \phi, \mathcal{O}_K)$  are commensurable, discrete groups.*

**Proof** By Lemma 2.6 there are infinitely many Salem numbers  $s$  and integers  $n$  and  $m$  such that every complex Galois conjugate of  $s^m$  and  $s^n$  lies in  $\eta$ . Fix one such Salem number  $s$ , and  $K, L$  and  $\delta$  as above.

By Theorem 4.6, sesquilinear forms are completely classified by dimension, determinant class, and the signatures for the places of  $L$  for which  $(s - 1/s)^2$  is negative. By Proposition 4.9,  $J_{s^n}$  and  $\lambda J_{s^m}$  have the same determinant class for  $\lambda$  in  $L$ , namely  $\lambda = \det J_{s^n} / \det J_{s^m}$ .

Let  $\sigma$  be a complex placement of  $L$ . Then  $\sigma(s^m)$  is a complex Galois conjugates of  $s^m$ , and similarly for  $\sigma(s^n)$  and  $s^n$ . Since  $n$  and  $m$  were chosen so that all of the complex Galois conjugates of  $s^m$  and  $s^n$  have arguments in  $\eta$ ,  $J_{\sigma(s^m)}$  and  $J_{\sigma(s^n)}$  are positive definite. Moreover,  $\det J_{s^n} / \det J_{s^m}$  and  $\sigma(\det J_{s^n} / \det J_{s^m})$  are both positive, making  $\lambda > 0$ . So regardless of whether  $\sigma((s - 1/s)^2)$  is positive or negative, the forms  $J_{\sigma(s^i)}$  have the same signature.

Therefore  $J_{s^n}$  is equivalent to  $\lambda J_{s^m}$ , and so  $\text{SU}(J_{s^n}, \phi, \mathcal{O}_K)$  is commensurable to  $\text{SU}(J_{s^m}, \phi, \mathcal{O}_K)$ . The groups are discrete by Theorem 2.3. □

**Corollary 4.11** *Let  $\rho_t: G \rightarrow \text{SL}_{2k+1}(\mathbb{Z}[t, t^{-1}])$  be a group representation with a parameter  $t$ . Suppose there exists a matrix  $J_t$  such that:*

- (1) *For all  $M$  in the image of  $\rho_t$ ,  $M^* J_t M = J_t$ , where  $M^*(t) = M^\top(1/t)$ .*
- (2)  *$J_t = (J_{1/t})^\top$ .*
- (3)  *$J_t \in \text{GL}_m(\mathbb{Q}(t))$ , where no entry of  $J_t$  has denominator with 1 as a root.*
- (4)  *$J_t$  is positive definite for  $t$  in an neighborhood  $\eta$  of 1.*

Then there exist infinitely many Salem numbers  $s$  such that for infinitely many integers  $n$  and  $m$  the specializations  $\rho_{s^m}$  at  $t = s^m$  and  $\rho_{s^n}$  at  $t = s^n$  map into commensurable lattices of  $\mathrm{SL}_{2k+1}(\mathbb{R})$ .

**Example 4.12** The reduced Burau representation of  $B_4$  is 3–dimensional and, after the appropriate rescaling to have determinant 1, satisfies [Corollary 4.11](#). So certain powers of the specializations in [Example 3.2](#) map into commensurable lattices in  $\mathrm{SL}_3(\mathbb{R})$ .

## References

- [1] **J S Birman, H Wenzl**, *Braids, link polynomials and a new algebra*, Trans. Amer. Math. Soc. 313 (1989) 249–273 [MR](#) [Zbl](#)
- [2] **O Brunat, K Magaard, I Marin**, *Image of the braid groups inside the finite Iwahori–Hecke algebras*, J. Reine Angew. Math. 733 (2017) 161–182 [MR](#) [Zbl](#)
- [3] **O Brunat, I Marin**, *Image of the braid groups inside the finite Temperley–Lieb algebras*, Math. Z. 277 (2014) 651–664 [MR](#) [Zbl](#)
- [4] **RD Budney**, *On the image of the Lawrence–Krammer representation*, J. Knot Theory Ramifications 14 (2005) 773–789 [MR](#) [Zbl](#)
- [5] **C W Curtis, N Iwahori, R Kilmoyer**, *Hecke algebras and characters of parabolic type of finite groups with  $(B, N)$ –pairs*, Inst. Hautes Études Sci. Publ. Math. 40 (1971) 81–116 [MR](#) [Zbl](#)
- [6] **MH Freedman, MJ Larsen, Z Wang**, *The two-eigenvalue problem and density of Jones representation of braid groups*, Comm. Math. Phys. 228 (2002) 177–199 [MR](#) [Zbl](#)
- [7] **L Funar, T Kohno**, *On Burau’s representations at roots of unity*, Geom. Dedicata 169 (2014) 145–163 [MR](#) [Zbl](#)
- [8] **V FR Jones**, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (2) 126 (1987) 335–388 [MR](#) [Zbl](#)
- [9] **I Marin**, *Caractères de rigidité du groupe de Grothendieck–Teichmüller*, Compos. Math. 142 (2006) 657–678 [MR](#) [Zbl](#)
- [10] **I Marin**, *Braids inside the Birman–Wenzl–Murakami algebra*, Algebr. Geom. Topol. 10 (2010) 1865–1886 [MR](#) [Zbl](#)
- [11] **D W Morris**, *Introduction to arithmetic groups*, Deductive, Lethbridge, AB (2015) [MR](#) [Zbl](#)
- [12] **J Murakami**, *The Kauffman polynomial of links and representation theory*, Osaka J. Math. 24 (1987) 745–758 [MR](#) [Zbl](#)

- [13] **W Scharlau**, *Quadratic and Hermitian forms*, Grundle Math. Wissen. 270, Springer (1985) [MR](#) [Zbl](#)
- [14] **C C Squier**, *The Burau representation is unitary*, Proc. Amer. Math. Soc. 90 (1984) 199–202 [MR](#) [Zbl](#)
- [15] **T N Venkataramana**, *Image of the Burau representation at  $d^{\text{th}}$  roots of unity*, Ann. of Math. (2) 179 (2014) 1041–1083 [MR](#) [Zbl](#)
- [16] **H Wenzl**, *Hecke algebras of type  $A_n$  and subfactors*, Invent. Math. 92 (1988) 349–383 [MR](#) [Zbl](#)
- [17] **H Wenzl**, *Quantum groups and subfactors of type  $B$ ,  $C$ , and  $D$* , Comm. Math. Phys. 133 (1990) 383–432 [MR](#) [Zbl](#)
- [18] **Y Zhao**, *Young tableaux and the representations of the symmetric group*, Harvard College Math. Rev. 2 (2008) 33–45
- [19] **M G Zinno**, *On Krammer's representation of the braid group*, Math. Ann. 321 (2001) 197–211 [MR](#) [Zbl](#)

ICERM, Brown University  
Providence, RI, United States

[nancy.scherich@gmail.com](mailto:nancy.scherich@gmail.com)

Received: 13 October 2019      Revised: 3 December 2021

# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](http://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
[etnyre@math.gatech.edu](mailto:etnyre@math.gatech.edu)  
Georgia Institute of Technology

Kathryn Hess  
[kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

Julie Bergner	University of Virginia <a href="mailto:jeb2md@eservices.virginia.edu">jeb2md@eservices.virginia.edu</a>	Robert Lipshitz	University of Oregon <a href="mailto:lipshitz@uoregon.edu">lipshitz@uoregon.edu</a>
Steven Boyer	Université du Québec à Montréal <a href="mailto:cohf@math.rochester.edu">cohf@math.rochester.edu</a>	Norihiko Minami	Nagoya Institute of Technology <a href="mailto:nori@nitech.ac.jp">nori@nitech.ac.jp</a>
Tara E. Brendle	University of Glasgow <a href="mailto:tara.brendle@glasgow.ac.uk">tara.brendle@glasgow.ac.uk</a>	Andrés Navas	Universidad de Santiago de Chile <a href="mailto:andres.navas@usach.cl">andres.navas@usach.cl</a>
Indira Chatterji	CNRS & Université Côte d'Azur (Nice) <a href="mailto:indira.chatterji@math.cnrs.fr">indira.chatterji@math.cnrs.fr</a>	Thomas Nikolaus	University of Münster <a href="mailto:nikolaus@uni-muenster.de">nikolaus@uni-muenster.de</a>
Alexander Dranishnikov	University of Florida <a href="mailto:dranish@math.ufl.edu">dranish@math.ufl.edu</a>	Robert Oliver	Université Paris 13 <a href="mailto:bobol@math.univ-paris13.fr">bobol@math.univ-paris13.fr</a>
Corneli Druţu	University of Oxford <a href="mailto:cornelia.drutu@maths.ox.ac.uk">cornelia.drutu@maths.ox.ac.uk</a>	Birgit Richter	Universität Hamburg <a href="mailto:birgit.richter@uni-hamburg.de">birgit.richter@uni-hamburg.de</a>
Tobias Ekholm	Uppsala University, Sweden <a href="mailto:tobias.ekholm@math.uu.se">tobias.ekholm@math.uu.se</a>	Jérôme Scherer	École Polytech. Féd. de Lausanne <a href="mailto:jerome.scherer@epfl.ch">jerome.scherer@epfl.ch</a>
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México <a href="mailto:mario@matem.unam.mx">mario@matem.unam.mx</a>	Zoltán Szabó	Princeton University <a href="mailto:szabo@math.princeton.edu">szabo@math.princeton.edu</a>
David Futер	Temple University <a href="mailto:dfuter@temple.edu">dfuter@temple.edu</a>	Ulrike Tillmann	Oxford University <a href="mailto:tillmann@maths.ox.ac.uk">tillmann@maths.ox.ac.uk</a>
John Greenlees	University of Warwick <a href="mailto:john.greenlees@warwick.ac.uk">john.greenlees@warwick.ac.uk</a>	Maggy Tomova	University of Iowa <a href="mailto:maggy-tomova@uiowa.edu">maggy-tomova@uiowa.edu</a>
Ian Hambleton	McMaster University <a href="mailto:ian@math.mcmaster.ca">ian@math.mcmaster.ca</a>	Nathalie Wahl	University of Copenhagen <a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>
Hans-Werner Henn	Université Louis Pasteur <a href="mailto:henn@math.u-strasbg.fr">henn@math.u-strasbg.fr</a>	Chris Wendl	Humboldt-Universität zu Berlin <a href="mailto:wendl@math.hu-berlin.de">wendl@math.hu-berlin.de</a>
Daniel Isaksen	Wayne State University <a href="mailto:isaksen@math.wayne.edu">isaksen@math.wayne.edu</a>	Daniel T. Wise	McGill University, Canada <a href="mailto:daniel.wise@mcgill.ca">daniel.wise@mcgill.ca</a>
Christine Lescop	Université Joseph Fourier <a href="mailto:lescop@ujf-grenoble.fr">lescop@ujf-grenoble.fr</a>		

---

See inside back cover or [msp.org/agt](http://msp.org/agt) for submission instructions.


The subscription price for 2023 is US \$650/year for the electronic version, and \$940/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 23      Issue 5 (pages 1935–2414)      2023

---

Splitting Madsen–Tillmann spectra, II: The Steinberg idempotents and Whitehead conjecture	1935
TAKUJI KASHIWABARA and HADI ZARE	
Free and based path groupoids	1959
ANDRÉS ÁNGEL and HELLEN COLMAN	
Discrete real specializations of sesquilinear representations of the braid groups	2009
NANCY SCHERICH	
A model for configuration spaces of points	2029
RICARDO CAMPOS and THOMAS WILLWACHER	
The Hurewicz theorem in homotopy type theory	2107
J DANIEL CHRISTENSEN and LUIS SCOCCOLA	
A concave holomorphic filling of an overtwisted contact 3–sphere	2141
NAOHIKO KASUYA and DANIELE ZUDDAS	
Modifications preserving hyperbolicity of link complements	2157
COLIN ADAMS, WILLIAM H MEEKS III and ÁLVARO K RAMOS	
Golod and tight 3–manifolds	2191
KOUYEMON IRIYE and DAISUKE KISHIMOTO	
A remark on the finiteness of purely cosmetic surgeries	2213
TETSUYA ITO	
Geodesic complexity of homogeneous Riemannian manifolds	2221
STEPHAN MESCHER and MAXIMILIAN STEGEMEYER	
Adequate links in thickened surfaces and the generalized Tait conjectures	2271
HANS U BODEN, HOMAYUN KARIMI and ADAM S SIKORA	
Homotopy types of gauge groups over Riemann surfaces	2309
MASAKI KAMEKO, DAISUKE KISHIMOTO and MASAHIRO TAKEDA	
Diffeomorphisms of odd-dimensional discs, glued into a manifold	2329
JOHANNES EBERT	
Intrinsic symmetry groups of links	2347
CHARLES LIVINGSTON	
Loop homotopy of 6–manifolds over 4–manifolds	2369
RUIZHI HUANG	
Infinite families of higher torsion in the homotopy groups of Moore spaces	2389
STEVEN AMELOTTE, FREDERICK R COHEN and YUXIN LUO	