

Algebra & Number Theory

Volume 3

2009

No. 7

Cox rings of degree one del Pezzo surfaces

Damiano Testa, Anthony Várilly-Alvarado and Mauricio Velasco



mathematical sciences publishers

Cox rings of degree one del Pezzo surfaces

Damiano Testa, Anthony Várilly-Alvarado and Mauricio Velasco

Let X be a del Pezzo surface of degree one over an algebraically closed field, and let $\text{Cox}(X)$ be its total coordinate ring. We prove the missing case of a conjecture of Batyrev and Popov, which states that $\text{Cox}(X)$ is a quadratic algebra. We use a complex of vector spaces whose homology determines part of the structure of the minimal free $\text{Pic}(X)$ -graded resolution of $\text{Cox}(X)$ over a polynomial ring. We show that sufficiently many Betti numbers of this minimal free resolution vanish to establish the conjecture.

1. Introduction

Let k be an algebraically closed field and let X be a smooth projective integral scheme over k . Assume that the Picard group $\text{Pic}(X)$ is freely generated by the classes of divisors D_0, D_1, \dots, D_r . The *total coordinate ring*, or *Cox ring* of X with respect to this basis is given by

$$\text{Cox}(X) := \bigoplus_{(m_0, \dots, m_r) \in \mathbb{Z}^{r+1}} \mathrm{H}^0(X, \mathcal{O}_X(m_0 D_0 + \dots + m_r D_r)),$$

with multiplication induced by the multiplication of functions in $k(X)$. Different choices of bases yield (noncanonically) isomorphic Cox rings.

The first appearance of Cox rings was in the context of toric varieties in [Cox 1995]. In that paper Cox proves that if X is a toric variety then its total coordinate ring is a finitely generated multigraded polynomial ring, and that X can be recovered as a quotient of an open subset of $\text{Spec}(\text{Cox}(X))$ by the action of a torus.

Cox rings are finitely generated k -algebras in several other cases, including del Pezzo surfaces [Batyrev and Popov 2004], rational surfaces with big anticanonical divisor [Testa et al. 2009], blow-ups of \mathbb{P}^n at points lying on a rational normal curve [Castravet and Tevelev 2006] and wonderful varieties [Brion 2007]. All these varieties are examples of *Mori dream spaces* [Hu and Keel 2000], and for this

MSC2000: 14J26.

Keywords: Cox rings, total coordinate rings, del Pezzo surfaces.

The first author was partially supported by Jacobs University Bremen, DFG grant STO-299/4-1 and EPSRC grant number EP/F060661/1; the third author is partially supported by NSF grant DMS-0802851.

class the Cox ring of X captures much of the birational geometry of the variety. For example, the effective and nef cones of X are finitely generated polyhedral cones and there are only finitely many varieties isomorphic to X in codimension one, satisfying certain mild restrictions. To add to the long list of consequences of the landmark paper [Birkar et al. 2008], we note that log Fano varieties are also Mori dream spaces.

Colliot-Thélène and Sansuc introduced universal torsors to aid the study of the Hasse principle and weak approximation on an algebraic variety X over a number field [Colliot-Thélène and Sansuc 1980; 1987]; see also [Colliot-Thélène et al. 1984]. If the Cox ring of X is finitely generated, then the universal torsor of X is an open subset of $\text{Spec}(\text{Cox}(X))$, an affine variety for which explicit presentations have been calculated in many cases [Hassett and Tschinkel 2004; Hassett 2004]. Starting with Salberger [1998], universal torsors have been successfully applied to the problem of counting points of bounded height on many classes of varieties. The expository article [Peyre 2004] has a very readable account of the ideas involved. The explicit descriptions of universal torsors via Cox rings have led to explicit examples of generalized del Pezzo surfaces that satisfy Manin's conjectures for points of bounded height [de la Bretèche and Browning 2007; de la Bretèche et al. 2007].

Batyrev and Popov [2004] systematically study the Cox rings of del Pezzo surfaces X/k . They show that $\text{Cox}(X)$ is a finitely generated k -algebra; moreover, if $\deg(X) \leq 6$ then $\text{Cox}(X)$ is generated by sections whose classes have anticanonical degree one [Batyrev and Popov 2004, Theorem 3.2]. Let \mathcal{G} denote a minimal set of homogeneous generators of $\text{Cox}(X)$, and denote by $k[\mathcal{G}]$ the polynomial ring whose variables are indexed by the elements of \mathcal{G} . As a result, $\text{Cox}(X)$ is a quotient of $k[\mathcal{G}]$:

$$\text{Cox}(X) \cong k[\mathcal{G}]/I_X.$$

Batyrev and Popov provided a conjectural description of the generators of I_X .

Conjecture 1.1 [Batyrev and Popov 2004]. Let X be a del Pezzo surface. The ideal I_X is generated by quadrics.

Quadratic relations have a clear geometric meaning: linear systems associated to degree two nef divisors on X have many more reducible elements than their dimension; thus there are linear dependence relations among degree two monomials in $k[\mathcal{G}]$. All quadratic relations arise in this way (see page 758).

Remark 1.2. Del Pezzo surfaces of degree at least six are toric varieties and hence their Cox rings are polynomial rings. The Cox ring of a del Pezzo surface of degree five is the homogeneous coordinate ring of the Grassmannian $Gr(2, 5)$ [Batyrev and Popov 2004].

Several partial results are known about this conjecture. Stillman, Testa, and Velasco established it for del Pezzo surfaces of degree four, and general del Pezzo surfaces of degree three [Stillman et al. 2007]. The conjecture was proved in [Laface and Velasco 2009] for surfaces of degree at least two. Recently, Serganova and Skorobogatov [2007; 2008] established Conjecture 1.1 up to radical for surfaces of degree at least two, using representation theoretic methods, (see also [Derenthal 2007] for related work). In all cases, the conjecture for degree one surfaces eluded proof. The purpose of this paper is to fill in this gap for surfaces defined over a field of characteristic different from two, as well as for certain general surfaces in characteristic two that we call *sweeping* (see Definition 4.3).

Theorem 1.3. *Let X be a del Pezzo surface of degree one; if the characteristic of k is two, then assume that X is sweeping. The ideal I_X is generated by quadrics.*

Remark 1.4. Our method of proof is cohomological in nature and relies mainly on the Kawamata–Viehweg vanishing theorem. By [Terakawa 1998] and [Xie 2008], this theorem is independent of the characteristic of k for rational surfaces.

The argument we give works for del Pezzo surfaces of every degree. We include a short proof of the conjecture for del Pezzo surfaces of degree at least two in Section 10. An alternative proof of Conjecture 1.1 for general del Pezzo surfaces in characteristic zero, obtained independently by computational means, appears in a paper by Sturmfels and Xu [2008]. They apply the theory of sagbi bases to construct an initial toric ideal of I_X which is generated by quadrics. Geometrically this corresponds to a degeneration of the universal torsor on X to a suitable toric variety.

In order to prove Theorem 1.3, we modify the approach taken in [Laface and Velasco 2009]. We use a complex of vector spaces whose homology determines part of the structure of the minimal free $\text{Pic}(X)$ -graded resolution of $\text{Cox}(X)$ over a polynomial ring. We show that sufficiently many Betti numbers of this minimal free resolution vanish to establish the theorem. We hope that similar techniques can be applied to obtain presentations of Cox rings of other classes of varieties, for example singular del Pezzo surfaces, blow-ups of \mathbb{P}^n at points lying on the rational normal curve of degree n and $\overline{M}_{0,n}$.

The paper is organized as follows. In Section 2 we define del Pezzo surfaces, fix presentations for their Cox rings and establish notation for the rest of the paper. In Section 3 we study the nef cone of del Pezzo surfaces and prove some geometric results. In Section 4 we analyze low-degree linear systems on del Pezzo surfaces of degree at most two. Our proof of Theorem 1.3 uses [Batyrev and Popov 2004, Proposition 3.4]. However, the proof given by those authors for their proposition has a gap: it applies only to general del Pezzo surfaces of degree one in characteristic not two [Popov 2004]. Thus we prove Proposition 4.4 in order to establish

Theorem 1.3 for all del Pezzo surfaces of degree one in characteristic not two. In Section 5 we review the approach of [Laface and Velasco 2009] to study the ideal of relations of finitely generated Cox rings, adapting it to the case of degree one del Pezzo surfaces. In Section 6 we define the notions of *capturability* of a divisor and *stopping criterion*, and prove Theorem 6.7, the main ingredients in our proof of the Batyrev–Popov conjecture. In Section 7 we show the capturability of most divisors on del Pezzo surfaces of degree one. We then handle the remaining cases for del Pezzo surfaces of degree one (Sections 8, 9) and del Pezzo surfaces of higher degree (Section 10). In Section 11 we finish the proof of the Batyrev–Popov conjecture and give the first multigraded Betti numbers of the Cox rings of del Pezzo surfaces.

2. Notation and background on del Pezzo surfaces

We briefly review some facts about del Pezzo surfaces and establish much of the paper’s notation along the way.

Definition 2.1. A *del Pezzo surface* X is a smooth, projective surface over k whose anticanonical divisor $-K_X$ is ample. The *degree* of X is the integer $(K_X)^2$.

Picard groups and Cox rings. A del Pezzo surface X not isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$ is isomorphic to a blow-up of \mathbb{P}^2 centered at $r \leq 8$ points *in general position*: this means no three points on a line, no six on a conic and no eight on a singular cubic with a singularity at one of the points. Let L be the inverse image of a line in \mathbb{P}^2 and let E_1, \dots, E_r be the exceptional divisors corresponding to the blown-up points. Then (L, E_1, \dots, E_r) is a basis for $\text{Pic}(X)$, and

$$\text{Cox}(X) := \bigoplus_{(m_0, \dots, m_r) \in \mathbb{Z}^{r+1}} H^0(X, \mathcal{O}_X(m_0L + m_1E_1 + \dots + m_rE_r)).$$

If X has degree one, then with the above choice of basis, the classes in $\text{Pic}(X)$ of the 240 exceptional curves are given in Table 1 [Manin 1986].

# of curves	Picard degree (up to a permutation of E_1, \dots, E_8)
8	E_1
28	$L - E_1 - E_2$
56	$2L - E_1 - E_2 - E_3 - E_4 - E_5$
56	$3L - 2E_1 - E_2 - E_3 - E_4 - E_5 - E_6 - E_7$
56	$4L - 2E_1 - 2E_2 - 2E_3 - E_4 - E_5 - E_6 - E_7 - E_8$
28	$5L - 2E_1 - 2E_2 - 2E_3 - 2E_4 - 2E_5 - 2E_6 - E_7 - E_8$
8	$6L - 3E_1 - 2E_2 - 2E_3 - 2E_4 - 2E_5 - 2E_6 - 2E_7 - 2E_8$

Table 1. Exceptional curves on X .

For a del Pezzo surface X of any degree, we let \mathcal{C} denote the set of exceptional curves on X . If X has degree one, we let

$$\mathcal{G} := \mathcal{C} \cup \{K_1, K_2\},$$

where $K_1, K_2 \in |-K_X|$; if X has degree at least two, then we let

$$\mathcal{G} := \mathcal{C}.$$

In all cases, we write $k[\mathcal{G}]$ for the polynomial ring whose variables are indexed by the elements of \mathcal{G} . By [Batyrev and Popov 2004, Theorem 3.2], we know that $\text{Cox}(X)$ is a finitely generated k -algebra, generated by sections whose divisor classes have anticanonical degree one. For all $G \in \mathcal{G}$, we let $g \in H^0(X, \mathcal{O}_X(G))$ be a nonzero element. We typically use uppercase letters for divisors on X and we denote the generator of $k[\mathcal{G}]$ associated to an exceptional curve E by the corresponding lowercase letter e . If X has degree one, then we denote the generators of $k[\mathcal{G}]$ corresponding to K_1 and K_2 by k_1 and k_2 , respectively. Furthermore, given any exceptional curve $E \in \mathcal{C}$, we let E' denote the unique exceptional curve whose divisor class is $-2K_X - E$, and we denote accordingly the generator corresponding to E' by e' .

There is a surjective morphism

$$k[\mathcal{G}] \rightarrow \text{Cox}(X)$$

that maps g to g . We denote the kernel of this map by I_X .

For any divisor D on X , any integer n and any $\text{Pic}(X)$ -graded ideal $J \subset k[\mathcal{G}]$, denote by J_D the vector space of homogeneous elements of $\text{Pic}(X)$ -degree D , by J_n the vector space of the homogeneous elements of anticanonical degree n and call them, respectively, the degree D part of J and the degree n part of J .

Finally, let J_X be the ideal generated by $(I_X)_2$; since $J_X \subset I_X$, there is a surjection

$$(k[\mathcal{G}]/J_X)_D \rightarrow (k[\mathcal{G}]/I_X)_D.$$

This map plays a role in Section 8.

3. Remarks on the nef cone of del Pezzo surfaces

In this section we collect basic results on del Pezzo surfaces that we use in the paper. The following result is well known [Debarre 2001, page 148, 6.5].

Proposition 3.1. *Let X be a del Pezzo surface of degree $d \leq 7$. A divisor class $C \in \text{Pic}(X)$ is nef (respectively ample) if and only if $C \cdot E \geq 0$ (respectively $C \cdot E > 0$) for all (-1) -curves $E \subset X$.*

Definition 3.2. Let X be a del Pezzo surface. The *minimal ample divisor on X* is the ample divisor A_X defined in the following table:

X	A_X	$(K_X)^2$
\mathbb{P}^2	$-\frac{1}{3}K_X$	9
$\mathbb{P}^1 \times \mathbb{P}^1$	$-\frac{1}{2}K_X$	8
$Bl_p(\mathbb{P}^2)$	$2L - E$	8
$Bl_{p_1, \dots, p_r}(\mathbb{P}^2)$	$-K_X$	$9 - r$

We also extend the list to include $X = \mathbb{P}^1$ and we define the minimal ample divisor on \mathbb{P}^1 to be the class of a point. To simplify the notation, if $b: X \rightarrow Y$ is a morphism, then we may denote $b^*(A_Y)$ by A_X . Minimal ample divisors allow us to give a geometric description of nef divisors.

Corollary 3.3. *Let X_r be a del Pezzo surface of degree $9 - r$. Let $D \in \text{Pic}(X_r)$ be a nef divisor. Then we can find nonnegative integers n_0, n_1, \dots, n_r and a sequence of morphisms $X_r \rightarrow X_{r-1} \rightarrow \dots \rightarrow X_1 \rightarrow X_0$ such that*

- *each morphism is the contraction of a (-1) -curve, except for $X_1 \rightarrow X_0$, which is allowed to be a conic bundle; and*
- *D can be expressed as*

$$D = n_r A_{X_r} + n_{r-1} A_{X_{r-1}} + \dots + n_0 A_{X_0}.$$

Proof. We proceed by induction on r , the cases $r \leq 1$ being immediate. Suppose that $r \geq 2$ and let $n := \min\{L \cdot D \mid L \subset X \text{ is a } (-1)\text{-curve}\}$. By assumption we have $n \geq 0$. Let $\bar{D} := D + nK_{X_r}$; note that \bar{D} is nef by Proposition 3.1. Choose a (-1) -curve $L_0 \subset X$ such that $\bar{D} \cdot L_0 = 0$. Thus \bar{D} is the pull-back of a nef divisor on the del Pezzo surface X_{r-1} obtained by contracting L_0 . The result follows by the inductive hypothesis. □

Observe that the integer n_r in the statement of the corollary is the nef threshold ([Reid 1997, page 126]) of the divisor D with respect to the minimal ample divisor of X_r . The minimal ample divisor is minimal in the sense that for every ample divisor A on X , the divisor $A - A_X$ is nef: this follows from Corollary 3.3.

By [Kollár 1996, Proposition III.3.4] and Corollary 3.3 we deduce that every nef divisor N on a del Pezzo surface is effective and that $|2N|$ is base-point free. Moreover, if N is a nef nonbig divisor, then N is a multiple of a conic bundle.

Let $X \rightarrow Y$ be a morphism with connected fibers and let A be the pull-back to X of the minimal ample divisor on Y . If $Y \simeq \mathbb{P}^1$, then we call A a *conic*. If $Y \simeq \mathbb{P}^2$, then we call A a *twisted cubic*, by analogy with the case of cubic surfaces. Finally, if Y is a del Pezzo surface of degree d , then we call A an *anticanonical divisor of degree d in X* .

We summarize and systematize the previous discussion in the following lemma.

Lemma 3.4. *Let X be a del Pezzo surface. The cone of nef divisors on X is generated by the following divisors:*

- (1) *the conics;*
- (2) *the twisted cubics;*
- (3) *the anticanonical divisors of degree $d \leq 3$ in X .*

In particular, if N is any nonzero nef divisor on X , then we may find $r := 8 - \deg(X)$ distinct exceptional curves E_1, E_2, \dots, E_r on X such that $N - E_1, \dots, N - E_r$ are either nef or the sum of a nef divisor and an exceptional curve, unless $\deg(X) = 1$ and $N = -K_X$.

Proof. The divisors in the list are clearly nef. Conversely, if D is any nef divisor, then either D is a multiple of a conic, and we are done, or a positive multiple of D induces a morphism with connected fibers $X \rightarrow Y$. It is clear that if A is the pull-back to X of the minimal ample divisor on Y , then $D - A$ is nef. By induction on $n := -K_X \cdot D$, we therefore reduce to showing that the anticanonical divisors of degree $d \geq 4$ in X are nonnegative linear combinations of the divisors listed. This is immediate.

For the second statement, note that it suffices to check it for the divisors in the list, and for $-2K_X$ if $\deg(X) = 1$, where the result is easy to verify. \square

The following lemma will be used in the proof of Lemma 7.7.

Lemma 3.5. *Let X be a del Pezzo surface, let $b: X \rightarrow \mathbb{P}^1$ be a conic bundle with fiber class Q and let C be an exceptional curve such that $C \cdot Q = 2$. There are exactly five reducible fibers of b such that C intersects both components. In particular, if $\deg(X) = 1$, then there are two reducible fibers of b such that C is disjoint from one of the two components in each fiber.*

Remark 3.6. If $\deg(X) \geq 4$, then there are no exceptional curves C such that $C \cdot Q = 2$, and hence the lemma applies nontrivially only to the cases $\deg(X) \leq 3$.

Proof. Let S and T be the components of a reducible fiber of b . Since $C \cdot Q = 2$, there are only two possibilities for the intersection numbers $C \cdot S$ and $C \cdot T$: they are either both equal to one, or one of them is zero and the other is two. Suppose that there are k reducible fibers of b such that the curve E intersects both components. Thus contracting all the components in fibers of b disjoint from E and one component in each of the remaining reducible fibers, we obtain a relatively minimal ruled surface $b': X' \rightarrow \mathbb{P}^1$, together with a smooth rational curve C' in X' , the image of C , having anticanonical degree $k + 1$, square $k - 1$ and intersection number two with a fiber of b' . Since X' is isomorphic to either $\mathbb{P}^1 \times \mathbb{P}^1$ or $Bl_p(\mathbb{P}^2)$, a direct calculation shows that this is only possible if $X' \simeq \mathbb{P}^1 \times \mathbb{P}^1$ and $k = 5$.

The last statement follows from the fact that any conic bundle structure on X contains exactly $8 - \deg(X)$ reducible fibers. \square

4. Low-degree linear systems

The lemmas in this section determine subsets of monomials that span $\text{Cox}(X)_D$ when $(\deg(X), D) \in \{(2, -K_X), (1, -2K_X)\}$.

Lemma 4.1. *Let X be a del Pezzo surface of degree two. The linear system $|-K_X|$ is spanned by any five of its reducible elements.*

Proof. The linear system $|-K_X|$ defines a separable morphism of degree two $\varphi: X \rightarrow \mathbb{P}^2$ such that every pair of exceptional curves E, E' on X with $E + E' = -K_X$ maps under φ to the same line in \mathbb{P}^2 . To prove the result it suffices to show that any five of these lines have no base-point. Suppose that $p \in \mathbb{P}^2$ is contained in $k_p \geq 4$ such lines; let $q \in X$ be a point in the inverse image of p . By construction, the point q is contained in k_p exceptional curves. Let E be an exceptional curve containing q and let E' be the exceptional curve such that $E + E' = -K_X$. The curve E' is disjoint from the exceptional curves F such that $F \cdot E = 1$, since $F \cdot E' = 0$. Thus contracting E' we obtain a del Pezzo surface Y of degree three with a point contained in at least $k_p - 1$ exceptional curves. The anticanonical linear system embeds Y as a smooth cubic surface in \mathbb{P}^3 , and exceptional curves through a point $y \in Y$ are lines contained in the tangent plane to Y at y . This implies that $k_p - 1 \leq 3$, and the result follows. \square

To study the case of degree one del Pezzo surfaces we begin with a lemma.

Lemma 4.2. *Let $R \subset \mathbb{P}^3$ be a curve, let $r \in R$ be a smooth point and let $p \in \mathbb{P}^3$ be a closed point different from r . Let H be a plane through p intersecting R at r with order of contact $m \geq 2$. Let $\pi_R: R \rightarrow \mathbb{P}^2$ be the projection away from p , let $\bar{R} = \pi_R(R)$, and let $\gamma: R \dashrightarrow (\mathbb{P}^2)^\vee$ be the composition of π_R and the Gauss map of \bar{R} , i.e., the map sending a general point $q \in R$ to the tangent line to \bar{R} at $\pi_R(q)$. Then the map γ is defined at r and one of the following occurs:*

- (1) *the tangent line to R at r contains p , or*
- (2) *the length of the localization at r of the fiber of γ at $\gamma(r)$ equals $m - 1$ if $\text{char}(k) \nmid m$ and it equals m if $\text{char}(k) \mid m$.*

Proof. The rational map γ is defined at r since $r \in R$ is a smooth point and the range of γ is projective. Choose homogeneous coordinates X_0, X_1, X_2, X_3 on \mathbb{P}^3 so that $p = [1, 0, 0, 1]$ and $r = [0, 0, 0, 1]$. In these coordinates H is defined by the linear form $AX_1 + BX_2$, for some $A, B \in k$. In the affine coordinates $(x_0 = X_0/X_3, x_1 = X_1/X_3, x_2 = X_2/X_3)$ the curve R is defined near the origin r by a complete intersection (f_1, f_2) and thus its embedded tangent space is the kernel of the matrix $DF(r) = ((\partial f_i / \partial x_j)(r))$ with $i \in \{1, 2\}$ and $j \in \{0, 1, 2\}$. Therefore either the first column of $DF(r)$ vanishes — that is, $\partial f_1 / \partial x_0$ and $\partial f_2 / \partial x_0$ vanish at r — and the tangent line to R through r contains p , or some 2×2 minor of

$DF(r)$ containing the first column is nonzero, since otherwise all 2×2 minors of $DF(r)$ vanish, contradicting the assumption that R is nonsingular at r . In the latter case, the completion of the local ring of R at r is isomorphic to $k[[t]]$ via a parametrization in formal power series of the form $(x_0(t), x_1(t), t)$. Since H has order of contact m at r , the power series $Ax_1(t) + Bt$ vanishes to order m so $A \neq 0$ and we have

$$x_1(t) = -\frac{B}{A}t + c_m t^m + (\text{higher order terms})$$

with $c_m \neq 0$. The equation of the tangent line to \bar{R} at $\pi_R(x_0(t), x_1(t), t, 1)$ is

$$-(X_1 - x_1(t)) + (X_2 - t)x'_1(t) = 0,$$

so the morphism γ is given by

$$(x_0(t), x_1(t), t, 1) \mapsto [-1, x'_1(t), x_1(t) - tx'_1(t)].$$

The localization at r of the fiber of γ at $\gamma(r)$ is thus given by the ideal

$$I = (Ax'_1(t) + B, x_1(t) - tx'_1(t)).$$

Since we have

$$Ax'_1(t) + B = Ac_m m t^{m-1} + (\text{higher order terms}),$$

$$x_1(t) - tx'_1(t) = c_m(1 - m)t^m + (\text{higher order terms}),$$

it follows that either $\text{char}(k) \nmid m$ and $I = (t^{m-1})$, or $\text{char}(k) \mid m$ and $I = (t^m)$ as we wanted to show. □

Let X be a del Pezzo surface of degree one. The linear system $|-2K_X|$ is base-point free and the image of the associated morphism $\kappa : X \rightarrow |-2K_X|^* \simeq \mathbb{P}^3$ is a cone W over a smooth conic; the morphism κ is a separable double cover of W branched at the cone vertex $w \in W$ and along a divisor $R \subset W$. We call *tritangent planes of X* the elements of $|-2K_X|$ supported on exceptional curves, and think of them as planes in $|-2K_X|^* = \mathbb{P}^3$. There are 120 such planes and they do not contain the vertex w of the cone, since the planes containing w correspond to elements of $|-2K_X|$ supported on the sum of two effective anticanonical divisors (see [Kollár 1996, Section III.3] and [Manin 1986, Chapter IV] for details).

Definition 4.3. A del Pezzo surface X of degree one over a field k is *sweeping* if any 119 tritangent planes of X span $|-2K_X|$.

Our goal in the remainder of this section is to show that every del Pezzo surface of degree one in characteristic not two is sweeping; if the characteristic is two, then we only show that a general del Pezzo surface of degree one is sweeping. In the following sections we prove that if X is a sweeping del Pezzo surface of degree one, then Conjecture 1.1 holds for X .

From here through page 740 we assume that k does not have characteristic 2. Then the curve R is smooth and it is the complete intersection of W with a cubic surface. Hence R is a canonical curve of genus four and degree six admitting a unique morphism of degree three to \mathbb{P}^1 (up to changes of coordinates on \mathbb{P}^1) obtained by projecting away from w . The tritangent planes to R are planes in \mathbb{P}^3 not containing w whose intersection with R is twice an effective divisor.

Proposition 4.4. *If X is a del Pezzo surface of degree one over a field k of characteristic different from two, then X is sweeping. More precisely, any 113 of the tritangent planes of X span $|-2K_X|$.*

Proof. Let $p \in \mathbb{P}^3$ be a closed point. We say that a tritangent plane H containing p is p -regular (resp. p -singular) if p does not belong to (resp. p belongs to) a tangent line to R at some point in $H \cap R$; denote by k_p the number of tritangent planes containing p and by k_p^r (resp. k_p^s) the number of p -regular (resp. p -singular) tritangent planes. Suppose that the point p belongs to R ; then there are no p -regular tritangent planes, since every tritangent plane containing p contains also the tangent line to R at p . Thus if q is any point on the tangent line to R at p , we have $k_p \leq k_q^s \leq k_q$, so it suffices to prove the proposition assuming that $p \in \mathbb{P}^3$ is a closed point not in R . Therefore let $p \in \mathbb{P}^3 \setminus R$ be a closed point; projecting away from p we obtain a morphism $\pi_R : R \rightarrow \mathbb{P}^2$; let $\bar{R} = \pi_R(R)$. Let $\bar{\gamma} : \bar{R} \dashrightarrow (\mathbb{P}^2)^\vee$ be the Gauss map, let $\gamma : R \rightarrow (\mathbb{P}^2)^\vee$ be the unique morphism that extends the composition $\bar{\gamma} \circ (\pi_R)$, and let $\check{R} = \gamma(R)$. The morphism γ factors as $R \xrightarrow{g} N \xrightarrow{\nu} \check{R}$, where $\nu : N \rightarrow \check{R}$ is the normalization of \check{R} . We summarize these definitions in a commutative diagram:

$$\begin{array}{ccc}
 \bar{R} & \xrightarrow{\bar{\gamma}} & \check{R} \\
 \pi_R \uparrow & \nearrow \gamma & \uparrow \nu \\
 R & \xrightarrow{g} & N
 \end{array} \tag{4-1}$$

The argument consists in identifying the contributions of the tritangent planes through p in terms of combinatorial invariants of Diagram 1. More precisely, the p -singular tritangent planes correspond (at most 7 : 1) to the points of \bar{R} where the morphism π_R is ramified; the p -regular tritangent planes correspond to points of \check{R} where (the separable part of) γ is ramified.

We now estimate the number k_p^s of p -singular tritangent planes.

Lemma 4.5. *As above, let R be the ramification divisor of κ .*

- (1) *If $\ell \subset \mathbb{P}^3$ is a tangent line to R , then there are at most 7 tritangent planes containing ℓ ;*
- (2) *Let $p \in \mathbb{P}^3$ be a closed point not in R . There are at most 42 p -singular tritangent planes.*

Proof of Lemma 4.5. (1) Let $r \in R$ be a point such that ℓ is tangent to R at r and let $\alpha: R \rightarrow \mathbb{P}^1$ be the morphism obtained by considering the pencil of planes in \mathbb{P}^3 containing ℓ . The morphism α has degree at most four since every plane containing ℓ is tangent to R at r and hence intersects R at r with multiplicity at least two. If α were not separable, then the characteristic of k would be three (recall that we assume $\text{char}(k) \neq 2$), α would be the Frobenius morphism, and the curve R would be rational, which is not the case. We deduce that α is separable; if α has degree three, then ℓ contains the vertex w of the cone W and in this case no tritangent plane contains ℓ . Moreover, α cannot have degree two since R is not hyperelliptic. Thus α is separable and we reduce to the case in which the degree of α is four. If H is a tritangent plane to R containing ℓ , then we have $R \cap H = 2((r) + (p) + (q))$, and hence the contribution of H to the ramification divisor of α is at least two. From the Hurwitz formula we deduce that the ramification divisor has degree 14 and r is contained in at most $14/2 = 7$ tritangent planes.

(2) First, we reduce to the case that the morphism π_R has degree at most two (and in particular it is separable). Indeed, the degree of π_R divides six, and since R is not contained in a plane, the image \bar{R} cannot be a line; therefore it suffices to analyze the cases in which the degree of π_R equals three. If π_R were not separable of degree three, then it would be purely inseparable and \bar{R} would have degree two but geometric genus four, which is not possible. If π_R is separable of degree three, then p is the vertex w of the cone W and hence $k_p^s = 0$, since there are no tritangent planes through w . This completes the reduction. Note that if the degree of π_R equals two, then the image of π_R is a smooth plane cubic, since R is not hyperelliptic.

Second, by [Hartshorne 1977, Proposition IV.3.4], the morphism π_R ramifies at every tangent line to R through p .

Finally, the image of a ramification point of π_R is either a singular point of \bar{R} or a ramification point of the morphism induced by π_R to the normalization of \bar{R} . Using the Hurwitz formula, the formula for the arithmetic genus of a plane curve, and the equality $\text{deg}(\bar{R}) \cdot \text{deg}(\pi_R) = 6$, we deduce that the number of points in \mathbb{P}^2 corresponding to ramification points of π_R is at most six. Thus there are at most six lines in \mathbb{P}^3 containing p such that every p -singular tritangent plane contains one of these lines. We conclude using Lemma 4.5(1). □

By Lemma 4.5(2) we have $k_p^s \leq 42$. Thus to prove the proposition, it suffices to show that $k_p^r \leq 70$.

Define the torsion sheaf Δ on \check{R} by the exact sequence

$$0 \rightarrow \mathcal{O}_{\check{R}} \rightarrow \nu_* \mathcal{O}_N \rightarrow \Delta \rightarrow 0. \tag{4-2}$$

From the long exact cohomology sequence, we deduce that

$$h^1(N, \mathcal{O}_N) + h^0(\Delta) = h^1(\check{R}, \mathcal{O}_{\check{R}}). \tag{4-3}$$

Applying the Hurwitz formula to the composition of π_R with a projection of \bar{R} from a general point of \mathbb{P}^2 , we obtain $\text{deg}(\check{R}) \leq 18$, and hence $h^1(\check{R}, \mathbb{C}_{\check{R}}) \leq 136$.

For each tritangent plane H through p denote by $\ell_H \in \check{R} \subset (\mathbb{P}^2)^\vee$ the point corresponding to the image in \mathbb{P}^2 of H under the projection away from p . We want to estimate the contribution of the point $\ell_H \in \check{R}$ to $h^0(\Delta)$. Note that this contribution is at least $\text{length}(\mathbb{C}_{v^{-1}(\ell_H)}) - 1$, and that

$$\text{length}(\mathbb{C}_{v^{-1}(\ell_H)}) \cdot \text{deg}(g) = \text{length}(\mathbb{C}_{\gamma^{-1}(\ell_H)}). \tag{4-4}$$

Factor g as the composition of a power of the Frobenius morphism, followed by a separable morphism g_s . Let d_i be the inseparable degree of g and $d_s = \text{deg}(g_s)$ be the separable degree of g , let also \check{d} denote the degree of \check{R} ; recall that we have $d_i d_s \check{d} \leq 18$.

Suppose first that $d_i d_s = 1$. By Lemma 4.2 and (4-4) each p -regular tritangent plane contributes at least 2 to $h^0(\Delta)$; hence we have $h^0(\Delta) \geq 2k_p^r$, and we conclude that $k_p^r \leq 68$.

Suppose now that $d_i d_s > 1$ and hence $\check{d} \leq 9$. By the Hurwitz formula, the degree of the ramification divisor of the morphism g_s is at most $6 + 2 \text{deg}(g_s) \leq 6 + 36 = 42$. The curve \check{R} has degree at most nine, and hence it has at most 28 singular points.

Let $U \subset N \times \mathbb{P}^3$ be the universal family of planes through p tangent to R and let $Z := (U \cap N \times R)$. Thus Z defines a family of closed subschemes of $R \subset \mathbb{P}^3$ of dimension zero and degree six. The geometric generic fiber of $Z \rightarrow N$ therefore determines a partition of 6 that we call the generic splitting type. Alternatively, if H is a plane through p tangent to R , then $H \cap R$ determines an effective divisor of degree six on R ; if H is general with the required properties, then the multiplicities of the geometric points of $H \cap R$ are the generic splitting type.

By [Hefez and Kleiman 1985, Theorem 3.5] and [Kaji 1992, page 529], the generic splitting type of γ is of the form $(a, \dots, a, 1, \dots, 1)$, where either $a = 2$ and the morphism is separable or a is the inseparable degree of the rational map $\bar{R} \rightarrow \check{R}$. The possibilities are

$$(5, 1), (4, 1, 1), (3, 3), (3, 1, 1, 1), (2, 2, 2), (2, 2, 1, 1), (2, 1, 1, 1, 1).$$

If the generic splitting type is $(2, 2, 2)$, then the point p is the cone vertex w and in this case $k_p = 0$. The partition $(2, 1, 1, 1, 1)$ corresponds to the case $d_i d_s = 1$ and we already analyzed it. The partitions $(5, 1)$ and $(4, 1, 1)$ can also be excluded, since they imply that R is birational to a plane curve of degree at most $\lfloor 18/4 \rfloor = 4$ and hence cannot have arithmetic genus four.

Examining the remaining generic splitting types and using Lemma 4.2, we deduce that the p -regular tritangent planes correspond to ramification points of g_s . In particular, at most $42 + 28 = 70$ such points exist, and the proposition follows. \square

Remark 4.6. In characteristic two the curve R above has degree 3, contains the cone vertex w , and is not necessarily smooth [Cragolini and Oliverio 2000]. While we are not able to prove that every del Pezzo surface of degree one in characteristic two is sweeping, we prove it for a general such surface. We write an explicit example of a del Pezzo surface S of degree two defined over $\overline{\mathbb{F}}_2$ and five tritangent planes to S no four of which share a common point. Thus S is sweeping; since the property of being sweeping is open, we conclude that a general del Pezzo surface of degree one in characteristic two is sweeping. We found this example using Magma [Bosma et al. 1997].

Let $\mathbb{P} := \mathbb{P}(1, 1, 2, 3)$ be the weighted projective space over $\overline{\mathbb{F}}_2$ with coordinates x, y, z, w and respective weights 1, 1, 2, 3. Let $S \subset \mathbb{P}$ be the surface defined by

$$S: \{w^2 + z^3 + wxz + wy^3 + x^6 = 0\}.$$

It is immediate to check that S is smooth. Let $t \in \overline{\mathbb{F}}_2$ be an element satisfying $t^2 + t + 1 = 0$; any four of the homogeneous forms

$$z + x^2, \quad z + y^2, \quad z + tx^2, \quad z + tx^2 + t^2xy + y^2, \quad z + tx^2 + xy + t^2y^2$$

are linearly independent. Moreover, they represent tritangent planes since we have

$$\begin{aligned} F(x, y, x^2, w) &= w(w + x^3 + y^3), \\ F(x, y, y^2, w) &= (w + (x^3 + txy^2 + x^2y + ty^3))(w + (x^3 + t^2xy^2 + x^2y + t^2y^3)), \\ F(x, y, tx^2, w) &= w(w + tx^3 + y^3), \\ F(x, y, tx^2 + t^2xy + y^2, w) &= (w + y(x^2 + xy + ty^2))(w + (tx^3 + t^2y^3 + tx^2y)), \\ F(x, y, tx^2 + xy + t^2y^2, w) &= (w + ty(x^2 + txy + y^2))(w + t(x^3 + tx^2y + ty^3)), \end{aligned}$$

and we deduce that S is sweeping.

The proof of Corollary 4.7 below given in [Batyrev and Popov 2004, Proposition 3.4] contains a gap pointed out in [Popov 2004]: the original argument implicitly assumes that the characteristic of the base field is not two, and reduces the proof to the fact that $H^0(X, \mathcal{O}_X(-2K_X))$ is spanned by the sections supported on exceptional curves. Proposition 4.4 fixes this gap.

Corollary 4.7. *Let X be a del Pezzo surface of degree one in characteristic not two. If $D \neq -K_X$ is an effective divisor, then $H^0(X, \mathcal{O}_X(D))$ is spanned by global sections supported on exceptional curves. \square*

Remark 4.8. A general del Pezzo surface of degree one in characteristic two is sweeping; therefore Corollary 4.7 also holds for such surfaces.

5. Betti numbers and the Batyrev–Popov conjecture

We review the approach of [Laface and Velasco 2009] to study the ideal of relations of Cox rings. As usual, let X be a del Pezzo surface, and let $R = k[\mathcal{G}]$. Throughout this section D denotes a divisor on X . Since R is positively graded (by anticanonical degree), every finitely generated $\text{Pic}(X)$ -graded R -module has a unique minimal $\text{Pic}(X)$ -graded free resolution. For the module $\text{Cox}(X) = R/I_X$ this resolution is of the form

$$\dots \rightarrow \bigoplus_{D \in \text{Pic}(X)} R(-D)^{b_{2,D}} \rightarrow \bigoplus_{D \in \text{Pic}(X)} R(-D)^{b_{1,D}} \rightarrow R \rightarrow 0,$$

where the rightmost nonzero map is given by a row matrix whose nonzero entries are a set of minimal generators of the ideal I_X . Since the differential of the resolution has degree 0, it follows that I_X has exactly $b_{1,D}(\text{Cox}(X))$ minimal generators of Picard degree D .

Let \mathbb{K} be the Koszul complex on \mathcal{G} . Consider the degree D part of the complex $\text{Cox}(X) \otimes_R \mathbb{K}$. Then

$$\begin{aligned} H_i((\text{Cox}(X) \otimes_R \mathbb{K})_D) &= (H_i(\text{Cox}(X) \otimes_R \mathbb{K}))_D \\ &= (\text{Tor}_i^R(\text{Cox}(X), k))_D = k^{b_{i,D}(\text{Cox}(X))}, \end{aligned}$$

where the equalities on the last line follow since $\text{Tor}_i^R(A, B)$ is symmetric in A and B and the Koszul complex is the minimal free resolution of k over R . Hence we have the equality

$$\dim_k(H_i((\text{Cox}(X) \otimes_R \mathbb{K})_D)) = b_{i,D}(\text{Cox}(X)).$$

Thus, Conjecture 1.1 is equivalent to the statement that $b_{1,D}(\text{Cox}(X)) = 0$ for all $D \in \text{Pic}(X)$ with $-K_X \cdot D \geq 3$. This is the form of the conjecture that we prove.

Let X be a del Pezzo surface of degree one. To compute the Betti numbers $b_{1,D}(\text{Cox}(X))$, denote by C_1, \dots, C_{240} the exceptional curves of X , and let

$$C_{240+i} := K_i$$

for $i \in \{1, 2\}$. With this notation, the part of the complex relevant to our task is

$$\bigoplus_{1 \leq i < j \leq 242} H^0(X, \mathcal{O}_X(D - C_i - C_j)) \xrightarrow{d_2} \bigoplus_{i=1}^{242} H^0(X, \mathcal{O}_X(D - C_i)) \xrightarrow{d_1} H^0(X, \mathcal{O}_X(D)),$$

where d_2 sends $\sigma_{ij} \in H^0(X, \mathcal{O}_X(D - C_i - C_j))$ to

$$(0, \dots, 0, \sigma_{ij}c_j, 0, \dots, 0, -\sigma_{ij}c_i, 0, \dots, 0)$$

and d_1 sends $\sigma_i \in H^0(X, \mathcal{O}_X(D - C_i))$ to $\sigma_i c_i$.

A cycle is an element of $\ker d_1$; a boundary is an element of $\text{im } d_2$. The support of a cycle $\sigma = (\dots, \sigma_i, \dots)$ is

$$\|\sigma\| = \{C_i : \sigma_i \neq 0\}$$

and the size of the support is the cardinality of $\|\sigma\|$, denoted by $|\sigma|$.

Strategy. In order to show that $b_{1,D}(\text{Cox}(X)) = 0$ whenever $-K_X \cdot D \geq 3$, we split the divisor classes D as follows:

- (1) D is ample and has anticanonical degree at least four (Section 7);
- (2) D is ample and has anticanonical degree three (Section 8);
- (3) D is not ample (Section 9).

The case when D is not ample follows by induction on the degree of the del Pezzo surface, and the ample divisors of anticanonical degree three are dealt with algebraically. In order to show that $b_{1,D}(\text{Cox}(X)) = 0$ whenever D is ample and $-K_X \cdot D \geq 4$ we show that every cycle is a boundary by arguing as follows.

- We describe constructions through which a divisor may be removed from the support of a cycle using boundaries, at the cost of possibly introducing new divisors in the support of the cycle.
- We apply these constructions to all cycles in a systematic way to reduce their support to at most two elements.

We conclude using the following lemma.

Lemma 5.1. *Any cycle σ with $|\sigma| \leq 2$ is a boundary.*

Proof. We deduce from [Elizondo et al. 2004, Theorem 1.1] that the Cox rings of del Pezzo surfaces are unique factorization domains. The statement follows immediately. □

6. Capturability and stopping criteria

We introduce the following terminology to aide us in our search of ways to reduce the support of a cycle. We keep the notation of the previous section.

Definition 6.1. A capture move for a divisor D is a pair (\mathcal{S}, C) , where $\mathcal{S} \subset \mathcal{G}$, $C \in \mathcal{G} \setminus \mathcal{S}$ and the map

$$\bigoplus_{S \in \mathcal{S}} H^0(X, \mathcal{O}_X(D - S - C)) \otimes H^0(X, \mathcal{O}_X(S)) \longrightarrow H^0(X, \mathcal{O}_X(D - C))$$

induced by tensor product of sections is surjective. We say that C is the captured curve, and that C is capturable for D by \mathcal{S} .

Let T be a set; for a direct sum $\bigoplus_{t \in T} V_t$ of vector spaces indexed by T and $T' \subset T$, we write $\sum_{t \in T'} a_t \hat{e}_t$ for the element $(b_t)_{t \in T}$ of $\bigoplus_{t \in T} V_t$ where $b_t = a_t$ if $t \in T'$ and $b_t = 0$ otherwise. As usual, if $S \in \mathcal{G}$, we denote by s the chosen global section of $\mathcal{O}_X(S)$. Let (\mathcal{S}, C) be a capture move for D , let $\sigma_C \in H^0(X, \mathcal{O}_X(D - C))$, and let $\sigma \in \bigoplus H^0(X, \mathcal{O}_X(D - C_i))$ be the element corresponding to σ_C . Then we have

$$\sigma_C = \sum_{S \in \mathcal{S}} p_S s, \quad p_S \in H^0(X, \mathcal{O}_X(D - S - C)),$$

and thus we obtain

$$\sigma = \sum_{S \in \mathcal{S}} p_S c \hat{e}_S + d_2 \left(\sum_{S \in \mathcal{S}} \varepsilon_{CS} p_S \hat{e}_{C+S} \right) \in \bigoplus_i H^0(X, \mathcal{O}_X(D - C_i)) \tag{6-1}$$

where $\varepsilon_{CS} \in \{\pm 1\}$.

Hence, if σ is a cycle and there is a capture move (\mathcal{S}, C) , then we can modify σ by a boundary so that $C \notin \|\sigma\|$. In this sense we have *captured* the curve C from the support of σ . Observe, however, that this modification adds a subset of \mathcal{S} to $\|\sigma\|$, so a priori the size of the support may not have decreased. We need to find and apply capture moves in an organized way to ensure that we are genuinely decreasing the size of the support of a cycle.

Finally note that if (\mathcal{S}, C) is a capture move and $\mathcal{S}' \supset \mathcal{S}$ with $C \notin \mathcal{S}'$, then (\mathcal{S}', C) is also a capture move. We frequently use this observation without explicitly mentioning it.

Lemma 6.2. *Let A, B and C be distinct exceptional curves on X , with A and B disjoint, and let D be any divisor. If $H^1(X, \mathcal{O}_X(D - A - B - C)) = 0$, then $(\{A, B\}, C)$ is a capture move for D . In particular, if $(-K_X + D - A - B - C)$ is nef, and if either it is big or it has anticanonical degree two, then $(\{A, B\}, C)$ is a capture move for D .*

Proof. Since A and B are disjoint, there is an exact sequence of sheaves

$$0 \rightarrow \mathcal{O}_X(-A - B) \rightarrow \mathcal{O}_X(-A) \oplus \mathcal{O}_X(-B) \rightarrow \mathcal{O}_X \rightarrow 0.$$

Tensoring with $\mathcal{O}_X(D - C)$ we obtain the short exact sequence

$$0 \rightarrow \mathcal{O}_X(D - A - B - C) \rightarrow \mathcal{O}_X(D - A - C) \oplus \mathcal{O}_X(D - B - C) \rightarrow \mathcal{O}_X(D - C) \rightarrow 0$$

and the desired surjectivity follows from the associated long exact sequence in cohomology and the assumption that

$$H^1(X, \mathcal{O}_X(D - A - B - C)) = 0.$$

The last statement follows from the Kawamata–Viehweg vanishing theorem when $-K_X + D - A - B - C$ is nef and big. Otherwise $-K_X + D - A - B - C = Q$ for

some conic Q and the statement follows from the exact sequence

$$0 \rightarrow \mathcal{O}_X(K_X) \rightarrow \mathcal{O}_X(K_X + Q) \rightarrow \mathcal{O}_Q(K_X + Q) \rightarrow 0,$$

by considering the associated long exact sequence in cohomology. □

Lemma 6.3. *If X is sweeping, then the pairs (\mathcal{C}, K_1) and (\mathcal{C}, K_2) are capture moves for all $D \neq -2K_X$.*

Proof. Since $D + K_X \neq -K_X$, it follows from Corollary 4.7 and Remark 4.8 that $H^0(X, \mathcal{O}_X(D + K_X))$ is spanned by global sections supported on exceptional curves. Thus the following map is surjective

$$\bigoplus_{S \in \mathcal{C}} H^0(X, \mathcal{O}_X(D - S + K_X)) \otimes H^0(X, \mathcal{O}_X(S)) \longrightarrow H^0(X, \mathcal{O}_X(D + K_X))$$

and (\mathcal{C}, K_1) and (\mathcal{C}, K_2) are capture moves for D . □

Definition 6.4. A *stopping criterion* for a divisor D is a set $\mathcal{S} \subset \mathcal{G}$ such that the following complex is exact:

$$\bigoplus_{\substack{C_i, C_j \in \mathcal{S} \\ i < j}} H^0(X, \mathcal{O}_X(D - C_i - C_j)) \xrightarrow{d_2} \bigoplus_{C_i \in \mathcal{S}} H^0(X, \mathcal{O}_X(D - C_i)) \xrightarrow{d_1} H^0(X, \mathcal{O}_X(D))$$

Remark 6.5. By Lemma 5.1, a subset of \mathcal{G} of cardinality two is a stopping criterion for any divisor D .

The name stopping criterion is motivated by Theorem 6.7: whenever we can capture all curves in a given degree D by curves contained in a stopping criterion, then there are no relations in degree D . In this case, we may *stop* looking for capture moves.

Definition 6.6. Let $D \in \text{Pic}(X)$ be a divisor class and let $\mathcal{M} := (E_1, \dots, E_n)$ be a sequence of elements of \mathcal{G} ; define $\mathcal{S}_i := \mathcal{G} \setminus \{E_1, E_2, \dots, E_i\}$ for all $i \in \{1, \dots, n\}$. We say that D is *capturable* (by \mathcal{M}) if

- (1) (\mathcal{S}_i, E_i) is a capture move for D for all $i \in \{1, \dots, n\}$,
- (2) \mathcal{S}_n is a stopping criterion for D .

Theorem 6.7. *Fix a divisor class $D \in \text{Pic}(X)$. If D is capturable then*

$$b_{1,D}(\text{Cox}(X)) = 0$$

and hence there are no minimal generators of the ideal I_X in degree D .

Proof. Let $\mathcal{M} := (E_1, \dots, E_n)$ be such that D is capturable by \mathcal{M} . Let $\mathcal{S}_0 = \mathcal{G}$ and, for $i \in \{1, \dots, n\}$ define $\mathcal{S}_i := \mathcal{G} \setminus \{E_1, E_2, \dots, E_i\}$. We want to show that every cycle σ is a boundary. Let $j(\sigma) := \max\{i \in \{0, \dots, n\} \mid \|\sigma\| \subset \mathcal{S}_i\}$. By definition if $\|\sigma\| \subset \mathcal{S}_n$, or equivalently if $j(\sigma) = n$, then σ is a boundary.

Suppose that $j = j(\sigma) < n$; since $(\mathcal{S}_{j+1}, E_{j+1})$ is a capture move for D , we deduce as in (6-1) that σ is homologous to a cycle τ such that $j(\tau) \geq j + 1$. Repeating this argument we conclude that σ is homologous to a cycle with support contained in \mathcal{S}_n and thus that σ is itself a boundary. \square

7. Ample divisors of anticanonical degree at least four

Let X be a del Pezzo surface of degree one. Assume throughout this section that if X is defined over a field of characteristic two, then X is sweeping. In this section we prove that there are no minimal generators of the ideal I_X in all sufficiently large ample degrees. More precisely, if D is an ample divisor such that $-K_X \cdot D \geq 4$ then we show that D is capturable.

The general strategy is the following. First we capture K_1 and K_2 using \mathcal{C} via Lemma 6.3. Next, assume C is an exceptional curve and we want to capture it using $\mathcal{S} \subset \mathcal{C}$. We prove that there exist disjoint exceptional curves S and T in \mathcal{S} such that $H^1(X, \mathcal{O}_X(K_X + L_{ST})) = 0$, where $L_{ST} := -K_X + D - C - S - T$. Then $(\{S, T\}, C)$ is a capture move for D by Lemma 6.2. Often we show that L_{ST} is nef and either big or of anticanonical degree two, omitting any reference to Lemma 6.2. In all cases we capture enough curves to conclude that D is a capturable divisor, using Lemma 5.1 as stopping criterion.

Lemma 7.1. *Let X be a del Pezzo surface of degree one and let $b: X \rightarrow Y$ be a birational morphism. Denote by K_Y the pull-back to X of a canonical divisor on Y and let N be a nef divisor on X . Assume that $\deg(Y) \geq 3$; then the divisor $D = -K_X - K_Y + N$ is capturable.*

Proof. Let \mathcal{S} be the set of exceptional curves contracted by b and note that \mathcal{S} consists of at least two disjoint exceptional curves and that $-K_Y - S - T$ is big and nef for all $S, T \in \mathcal{S}$, $S \neq T$. Let $C \in \mathcal{C} \setminus \mathcal{S}$ be any exceptional curve and let $C' := -2K_X - C$. First we capture the curves C such that $C' \in \mathcal{S}$ using \mathcal{S} : $L_{C'T} := -K_Y + N - T$ is big and nef for any choice of $T \in \mathcal{S} \setminus \{C'\}$.

Second we choose any two distinct $S, T \in \mathcal{S}$, we let $\mathcal{S}' := \{S, T\}$ and we capture all curves $C \in \mathcal{C} \setminus \mathcal{S}'$ such that $C' \notin \mathcal{S}'$: the divisor $L_{ST} = C' + (-K_Y - S - T) + N$ is big and nef provided $C' \cdot (-K_Y - S - T) > 0$; since the only curves orthogonal to $(-K_Y - S - T)$ are the curves in $\mathcal{S} \setminus \mathcal{S}'$, we conclude. \square

Lemma 7.2. *Let X be a del Pezzo surface of degree one and let $b: X \rightarrow Y$ be a birational morphism. Denote by A the pull-back to X of the minimal ample divisor on Y and let N be a nef divisor on X . Assume that $\deg(Y) \geq 8$; then the divisor $D = -K_X + A + N$ is capturable.*

Proof. Note that Y is isomorphic to \mathbb{P}^2 , $Bl_p(\mathbb{P}^2)$ or $\mathbb{P}^1 \times \mathbb{P}^1$.

Case 1: $Y \not\cong \mathbb{P}^1 \times \mathbb{P}^1$. In this case $A = L + Q$, where L is a twisted cubic and Q is either zero or a conic. Thus it suffices to treat the case $D = -K_X + L + N$.

Let \mathcal{S} be the set of curves contracted by L ; thus \mathcal{S} consists of eight disjoint exceptional curves. Note that $\sum_{S \neq T \in \mathcal{S}} L - S - T = 7(-K_X + L)$ has intersection number at least 7 with every exceptional curve on X . Since the intersection number of two exceptional curves on X is at most 3, it follows that every exceptional curve on X intersects positively at least two of the curves $\{L - S - T \mid S \neq T \in \mathcal{S}\}$. For any $C \in \mathcal{C} \setminus \mathcal{S}$ let $S, T \in \mathcal{S}$ be such that $(-2K_X - C) \cdot (L - S - T) > 0$ and such that no (rational) multiple of N equals $(-2K_X - C) + (L - S - T)$. Note that the second condition ensures that N does not contract both $-2K_X - C$ and $(L - S - T)$ when $-2K_X - C + (L - S - T)$ is a conic. We have that $L_{ST} := -2K_X + L - C - S - T + N = (-2K_X - C) + (L - S - T) + N$ is nef and either it has anticanonical degree two or it is also big. Thus (\mathcal{S}, C) is a capture move for all $C \in \mathcal{C} \setminus \mathcal{S}$. Let $S, T \in \mathcal{S}$ be distinct elements and let $\mathcal{S}' := \{S, T\}$. For any $C \in \mathcal{S} \setminus \mathcal{S}'$ the divisor $L_{ST} := (-2K_X - C) + (L - S - T) + N$ is big and nef since $(-2K_X - C) \cdot (L - S - T) = 2$.

Case 2: $Y \cong \mathbb{P}^1 \times \mathbb{P}^1$. Let \mathcal{S} be the set of curves contracted by A ; thus \mathcal{S} consists of seven disjoint exceptional curves. Note that $\sum_{S \neq T \in \mathcal{S}} A - S - T = -6K_X + 9A$ has intersection number at least six with every exceptional curve on X and that the summands are conics. Since the intersection number of an exceptional curve and a conic on X is at most four, it follows that every exceptional curve on X intersects positively at least two of the conics $\{A - S - T \mid S \neq T \in \mathcal{S}\}$. For any $C \in \mathcal{C} \setminus \mathcal{S}$ let $S, T \in \mathcal{S}$ be such that $(-2K_X - C) \cdot (A - S - T) > 0$. We have that $L_{ST} := -2K_X + A - C - S - T + N = (-2K_X - C) + (A - S - T) + N$ is big and nef. Thus (\mathcal{S}, C) is a capture move for all $C \in \mathcal{C} \setminus \mathcal{S}$. Let $S, T \in \mathcal{S}$ be distinct elements and let $\mathcal{S}' := \{S, T\}$. For any $C \in \mathcal{S} \setminus \mathcal{S}'$ the divisor

$$L_{ST} := (-2K_X - C) + (A - S - T) + N$$

is big and nef since $(-2K_X - C) \cdot (A - S - T) = 4$. □

Lemma 7.3. *Let X be a del Pezzo surface of degree one, E an exceptional curve on X and $N \neq 0$ a nef divisor on X such that $N \cdot E = 0$. The divisor $-(n+2)K_X + E + N$ is capturable for $n \geq 0$.*

Proof. By Lemma 3.4, for $i \in \{1, 2\}$, there is an exceptional curve F_i on X such that $F_i \cdot E = 0$ and $N - F_i$ is either nef or the sum of a nef divisor and an exceptional curve, with $F_1 \neq F_2$. Let $\mathcal{S} := \{E, F_1, F_2\}$. For any $C \in \mathcal{C} \setminus \mathcal{S}$, let $F \in \{F_1, F_2\}$ be such that $(-2K_X - C)$ is not a fixed component of $(N - F)$; we have that $L_{EF} = -(n+3)K_X + E - C - E - F + N = -K_X + (-2K_X - C) + (N - F) - nK_X$ is big and nef and we conclude that we can capture all curves in $\mathcal{C} \setminus \mathcal{S}$, using only

the three curves in \mathcal{S} . Finally, by the same argument, we find $(\{E, F_1\}, F_2)$ is a capture move and we are done. \square

Lemma 7.4. *Let X be a del Pezzo surface of degree one, E an exceptional curve on X and N a nef divisor on X . The divisor $D := -(n+3)K_X + E + N$ is capturable for $n \geq 0$.*

Proof. Let E, E_2, \dots, E_8 be eight disjoint exceptional curves on X and let $\tilde{E}_i := -K_X + E - E_i$ for $2 \leq i \leq 8$. Let $\mathcal{S} := \{E, E_2, \dots, E_8, \tilde{E}_2, \dots, \tilde{E}_8\}$. It is clear from Table 1 that every exceptional curve $C \in \mathcal{C} \setminus \mathcal{S}$ intersects positively at least two of the curves in \mathcal{S} ; in particular for every $C \in \mathcal{C} \setminus \mathcal{S}$ there is $S \in \mathcal{S}$ such that $S \cdot E = 0, S \cdot C > 0$ and N is not a (rational) multiple of $-4K_X - (S + C)$. Thus $L_{ES} = -K_X + D - E - S - C = (-4K_X - C - S) + N - nK_X$ is nef and it either has anticanonical degree two or it is big. In either case $H^1(X, \mathcal{O}_X(K_X + L_{ES})) = 0$ and we may capture all the curves in $\mathcal{C} \setminus \mathcal{S}$ using the curves in \mathcal{S} .

Note that for all integers $i, j, k, l \in \{2, 3, \dots, 8\}, i \neq j, k \neq l$, we have

$$(-4K_X - E_i - \tilde{E}_j) \cdot (-4K_X - E_k - \tilde{E}_l) = 0$$

if and only if $i = k$ and $j = l$. In particular, if N is a multiple of a conic, it is proportional to at most one of the divisors $(-4K_X - E_i - \tilde{E}_j)$; relabeling the indices if necessary, we may assume that N is not proportional to $-4K_X - (E_i + \tilde{E}_j)$ for all $i, j \in \{2, 3, \dots, 8\}$ and $i \neq j, (i, j) \neq (7, 8)$. Thus for all $i \in \{3, 4, \dots, 8\}$ the divisor $L_{E\tilde{E}_2} = (-4K_X - E_i - \tilde{E}_2) + N - nK_X$ is either a conic or big and nef and we may capture E_3, E_4, \dots, E_8 using E, \tilde{E}_2 . Similarly we can capture $\tilde{E}_2, \tilde{E}_3, \dots, \tilde{E}_8$ using E, E_2 and we are done. \square

Lemma 7.5. *Let X be a del Pezzo surface of degree one. Let $D = -(n+4)K_X + N$ where $n \geq 0$ and N is a nef divisor. Let C, S and T be exceptional curves such that $S \cdot T = 0$ and that $C \cdot S, C \cdot T$ are at least 2. Then $(\{S, T\}, C)$ is a capture move for D if either*

- (1) $C \cdot (S + T) = 5$, or
- (2) $C \cdot (S + T) = 4$ and the divisor N is either 0 or not a multiple of the conic $T' + S' + C' + K_X$.

Proof. Consider $L_{ST} := -K_X + D - S - T - C = S' + T' + C' - (n-1)K_X + N$. Since the inequalities $C' \cdot S' \geq 2$ and $C' \cdot T' \geq 2$ hold, the curves S', C' and T' , taken together, are not all pullbacks of exceptional curves on a del Pezzo surface Y of degree at least two. It follows that every exceptional curve $V \notin \{S', T', C'\}$ intersects at least one curve in this set and thus L_{ST} is nef. If $C \cdot (S + T) = 5$ then either $C' + S' = -2K_X$ or $C' + T' = -2K_X$ and L_{ST} is big. If $N = 0$ then L_{ST} has anticanonical degree 2 and capturability follows from Lemma 6.2. Finally,

assume that $C \cdot S = C \cdot T = 2$. In this case $S' + T' + C' + K_X = Q$ is a conic and $L_{ST} = Q - nK_X + N$ is big unless $N = mQ$ for some $m \geq 1$. \square

Lemma 7.6. *Let X be a del Pezzo surface of degree one and let N be a nef divisor on X . The divisor $-(n + 4)K_X + N$ is capturable for $n \geq 0$.*

Proof. Let L be a twisted cubic and let E_1, \dots, E_8 be exceptional curves on X contracted by L . First we capture all curves $C \in \mathcal{C}$ such that $L \cdot C = 3$. From Table 1 it is clear that $C = -K_X + E_i - E_j$ for some $i \neq j$. Set

$$S = E_j, \quad T_1 = 2L - E_i - E_{i_1} - E_{i_2} - E_{i_3} - E_{i_4}, \quad T_2 = 2L - E_i - E_{j_1} - E_{j_2} - E_{j_3} - E_{j_4},$$

with $\{i, j\} \cap \{i_1, \dots, i_4\} = \emptyset$, $\{i, j\} \cap \{j_1, \dots, j_4\} = \emptyset$, $\{i_1, \dots, i_4\} \neq \{j_1, \dots, j_4\}$. Note that $L \cdot S, L \cdot T_i \neq 3$, $C \cdot S = C \cdot T_i = 2$ and $S \cdot T_i = 0$ for $i = 1, 2$. Let T be an exceptional curve in $\{T_1, T_2\}$ such that $N \neq m(T' + S' + C' + K_X)$. By Lemma 7.5(2), $(\{S, T\}, C)$ is a capture move.

Next, we capture curves C with $L \cdot C \in \{4, 5\}$. It is clear from Table 1 that for any such curve there exists a pair of distinct exceptional curves $S, T \in \{E_1, \dots, E_8\}$ such that $(\{S, T\}, C)$ is a capture move via Lemma 7.5(2). Similarly we capture curves C with $L \cdot C \in \{1, 2\}$ using some pair $S, T \in \{E'_1, \dots, E'_8\}$. Next we capture E_3, \dots, E_8 using E'_1 and E'_2 via Lemma 7.5(1) and finally, we capture E'_1, \dots, E'_8 using E_1 and E_2 . This concludes the proof. \square

Lemma 7.7. *Let X be a del Pezzo surface of degree one and let Q be a conic on X . The divisor $D = -(n + 1)K_X + (m + 1)Q$ is capturable for $m, n \geq 0$ and $m + n \geq 1$.*

Proof. Suppose first that $n \geq 1$. Let \mathcal{S} be the set of curves contracted by Q and for any $S \in \mathcal{S}$, let \tilde{S} be the unique curve such that $S + \tilde{S} = Q$. If S, T, C are exceptional curves let $B_{ST} = -K_X + C' + Q - S - T$ and note that if B_{ST} is nef and big then so is $L_{ST} := -K_X + D - C - S - T = B_{ST} + mQ - (n - 1)K_X$ because $n \geq 1$. Moreover B_{ST} has anticanonical degree 2 and $h^2(X, \mathcal{O}_X(B_{ST})) = 0$ so by Riemann–Roch B_{ST} is nef if and only if $B_{ST}^2 = 2Q \cdot C' - 2C' \cdot S - 2C' \cdot T \geq 0$.

We first show that the curves C in $\mathcal{C} \setminus \mathcal{S}$ can be captured with the curves in \mathcal{S} . To do so, we split them into cases according to the value of $Q \cdot C'$.

If $Q \cdot C' = 0$ choose $T \in \mathcal{S}$ disjoint from C' . In this case $B_{C'T} = -K_X + \tilde{T}$ is nef and big, and $(\{C', T\}, C)$ is a capture move. If $Q \cdot C' = 1$ choose disjoint curves S and $T \in \mathcal{S}$ which are also disjoint from C' and note that B_{ST} has anticanonical degree 2 and square 2 so it is the pullback to X of the anticanonical divisor of a del Pezzo surface of degree 2 and thus it is nef and big; whence $(\{S, T\}, C)$ is a capture move. If $Q \cdot C' = 2$, by Lemma 3.5 there exist disjoint curves S, T in \mathcal{S} such that $S \cdot C' = T \cdot C' = 0$ so B_{ST} has square 4 and hence $B_{ST} = -2K$ is nef and big. Thus $(\{S, T\}, C)$ is a capture move in this case. Finally if $Q \cdot C' = 3$, then let S and T be disjoint curves in \mathcal{S} such that $S \cdot C' = T \cdot C' = 1$; then B_{ST} has square 2 and it is a nef and big divisor since it is the pullback to X of the anticanonical

divisor of a del Pezzo surface of degree 2. Thus $(\{S, T\}, C)$ is a capture move in this case.

Now choose S, T disjoint exceptional curves in \mathcal{S} . We show that all curves C in $\mathcal{S} \setminus \{S, T\}$ can be captured from $\{S, T\}$. In this case $Q \cdot C' = 4$ and $S \cdot C', T \cdot C' \leq 2$, so $B_{ST}^2 \geq 0$. If B_{ST} is big the statement follows since so is L_{ST} . If $B_{ST}^2 = 0$ there are two cases to consider, either $n = 1$ and $m = 0$ and the statement follows from Lemma 6.2, or at least one of n, m is greater than one. In this case $L_{ST} = B_{ST} - (n - 1)K_X + mQ$ is big because $B_{ST} \cdot Q = 6 \neq 0$. It follows that we can capture every exceptional curve in $\mathcal{S} \setminus \{S, T\}$ from $\{S, T\}$.

Suppose now that $n = 0$ and $m \geq 1$. Note that if $Q \cdot C' \leq 1$ then

$$Q \cdot (D - C) = Q \cdot (K_X + C' + (m + 1)Q) = -2 + Q \cdot C' < 0,$$

so $D - C$ is not an effective divisor. In this case C is captured vacuously and thus we restrict our attention to curves C with $Q \cdot C' \geq 2$. Let \mathcal{S} be the set of exceptional curves contracted by Q and for any $S \in \mathcal{S}$ let \tilde{S} be the unique curve such that $S + \tilde{S} = Q$.

For any $C \in \mathcal{C} \setminus \mathcal{S}$ which satisfies $Q \cdot C' \geq 2$ there exist disjoint curves S, T in \mathcal{S} such that \tilde{S} and \tilde{T} intersect C' . By construction $\tilde{S} + \tilde{T} + C'$ is nef and big and so is $L_{ST} := -K_X + D - S - T - C = \tilde{S} + \tilde{T} + C' + (m - 1)Q$. It follows that we can capture $\mathcal{C} \setminus \mathcal{S}$ with \mathcal{S} .

Finally, fix two disjoint curves $S, T \in \mathcal{S}$ and let C be any curve in $\mathcal{S} \setminus \{S, T\}$. Since $Q \cdot C = 0$ we have $Q \cdot C' = 4$ so C' intersects both \tilde{S} and \tilde{T} . It follows that L_{ST} is nef and big and that \mathcal{S} can be captured from $\{S, T\}$. □

Theorem 7.8. *Let X be a sweeping del Pezzo surface of degree one and let D be an ample divisor. If $-K_X \cdot D \geq 4$, then D is capturable.*

Proof. Write $D = -nK_X + N$, where $n \geq 1$, the divisor N is nef and not ample and $n - K_X \cdot N \geq 4$. If $N = 0$, then $n \geq 4$ and we conclude using Lemma 7.6. If $N \neq 0$, write $N = mA + N'$ where $X \rightarrow Y$ is a morphism with connected fibers, A is the pull-back to X of the minimal ample divisor on Y , N' is the pull-back of a nef divisor on Y and $m \geq 1$. If Y is a surface, then we conclude using Lemmas 7.1–7.4. If $Y \simeq \mathbb{P}^1$, then we conclude using Lemma 7.7. □

8. Ample divisors of anticanonical degree three

The only ample divisors D that seem to elude the strategy of Section 7 are those of anticanonical degree three. These divisors are $-3K_X, -2K_X + E, -K_X + Q$, where E is any exceptional curve and Q is any conic. Recall that there is a surjection

$$(k[\mathcal{C}]/J_X)_D \rightarrow (k[\mathcal{G}]/I_X)_D;$$

see Section 2. We show the dimension of $(k[\mathcal{G}]/J_X)_D$ is at most $h^0(X, \mathcal{O}_X(D)) = \dim(k[\mathcal{G}]/I_X)$ and conclude that $(I_X)_D = (J_X)_D$.

Lemma 8.1. *Let X be a del Pezzo surface of degree one and let A, B, G be exceptional curves with $A + B = -K_X + G$. Then k_1g, k_2g and ab form a basis for $H^0(X, \mathcal{O}_X(-K_X + G))$. In particular, if C, D are exceptional curves with $C + D = -K_X + G$ then $cd = \alpha_1k_1g + \alpha_2k_2g + \alpha_3ab$ and the coefficient α_3 is nonzero.*

Proof. The morphism $X \rightarrow \mathbb{P}^2$ associated to $|-K_X + G|$ contracts G and is ramified along a smooth plane quartic R . The image of G is a point not lying on any bitangent to R since otherwise X would have had a (-2) -curve. The images of K_1 and K_2 in \mathbb{P}^2 are distinct lines through the image of G and therefore the three lines in \mathbb{P}^2 corresponding to $A + B, K_1 + G, K_2 + G$ have no common point and are independent. □

Lemma 8.2. *Let X be a del Pezzo surface of degree one; then the ideal I_X has no minimal generators in degree $D = -3K_X$.*

Proof. The only ways of writing $-3K_X$ as a sum of three effective divisors are given in the following table.

Monomial	Description	
$h_1h_2h_3$	$h_1, h_2, h_3 \in \{k_1, k_2\}$	
haa'	$h \in \{k_1, k_2\}, A \in \mathcal{C}$	(8-1)
abc	$A, B, C \in \mathcal{C}$ $A \cdot B = A \cdot C = B \cdot C = 2$	

Indeed, let $-3K_X = A + B + C$ be any expression of $-3K_X$ as a sum of three effective divisors. If one among A, B, C is in $|-K_X|$, then we are in one of the first two cases above. If $A, B, C \in \mathcal{C}$, then intersecting with A, B, C successively both sides of the equation $-3K_X = A + B + C$ we obtain the system

$$\begin{cases} A \cdot B + A \cdot C = 4, \\ A \cdot B + B \cdot C = 4, \\ A \cdot C + B \cdot C = 4, \end{cases}$$

whose only solution is $A \cdot B = A \cdot C = B \cdot C = 2$.

By definition $(I_X)_2 = (J_X)_2$; we deduce that the span of the monomials of the first two forms in (8-1) in $k[\mathcal{G}]/J_X$ has dimension at most six, being the image of $H^0(X, \mathcal{O}_X(-K_X)) \otimes H^0(X, \mathcal{O}_X(-2K_X))$ under the multiplication map (note that $\dim H^0(X, \mathcal{O}_X(-K_X)) = 2, \dim H^0(X, \mathcal{O}_X(-2K_X)) = 4$ and the linearly independent elements $k_1 \otimes k_2^2e - k_2 \otimes k_1k_2$ and $k_2 \otimes k_1^2e - k_1 \otimes k_1k_2$ are in the kernel).

Fix any monomial $p = abc$ of the third form in (8-1) and denote the span of p together with the monomials of the first two forms in (8-1) by V . Let $q = def$ be any monomial of the third form in (8-1); we prove that q can be written as the sum of an element of V and an element of J_X .

The result is clear if $c = f$, since in this case we can use the relation coming from degree $A + B$ involving the monomials k_1gc, k_2gc, abc and dec to conclude (see Lemma 8.1). By the same reasoning we reduce to the case in which $\{a, b, c\}$ and $\{d, e, f\}$ are disjoint.

Suppose that $A \cdot D = 2$; then the divisor $G := -3K_X - A - D$ satisfies $G^2 = K_X \cdot G = -1$, and hence G is an exceptional curve. Moreover from the fact that $D \cdot G = 2$, we deduce that G is an exceptional curve on the del Pezzo surface obtained by contracting D' and whose anticanonical divisor pulled-back to X is $E + F$; let $H := E + F - G$ and note that H is also an exceptional curve. In particular we can use a relation coming from degree $E + F$ involving the monomials dk_1d', dk_2d', dgh and def to reduce to the case $\{a, b, c\} \cap \{d, e, f\} \neq \emptyset$.

Finally, if $A \cdot D = A \cdot E = A \cdot F = 1$, then A is a conic in the del Pezzo surface obtained by contracting F' . From Table 1 it follows that there are exceptional curves intersecting any given conic twice; denote the strict transform in X of one such curve of X by H and the exceptional curve $E + F - H$ by J . Thus we can use a relation coming from degree $E + F$ involving the monomials dk_1d', dk_2d', dhj and def to reduce to the case in which $A \cdot D = 2$. □

Lemma 8.3. *Let E be an exceptional curve on X ; then the ideal I_X has no minimal generators in degree $D = -2K_X + E$.*

Proof. The monomials of degree D are of the following forms:

Monomial	Description
ks	$k \in \{k_1, k_2\}$, $s \in k[\mathcal{G}]_{-K_X+E}$
$aa'e$	$A \cdot A' = 3$
abc	$A \cdot E = 1$ $B \cdot E = C \cdot E = 0$ $A \cdot B = A \cdot C = 2$ $B \cdot C = 1$

Indeed let $m \in k[\mathcal{G}]_D$ be a monomial. If $k \in \{k_1, k_2\}$, or e divides m , then m is of one of the first two forms. Otherwise let $m = abc$, with $\{a, b, c\} \cap \{k_1, k_2, e\} = \emptyset$; if two of the curves A, B, C have intersection number three, then the remaining one is E , which we are excluding. If the intersection numbers among the curves A, B, C are all at most two, then the required conditions follow multiplying successively the equality $A + B + C = -2K_X + E$ by A, B, C .

We show that the monomials of degree D span a subspace of dimension at most six of $(k[\mathcal{G}]/J_X)_D$; since $\dim \text{Cox}(X)_D = 6$, the result follows.

First, the image of $H^0(X, \mathcal{O}_X(-K_X)) \otimes H^0(X, \mathcal{O}_X(-K_X + E))$ in $k[\mathcal{G}]_D$ has dimension at most five: $\dim H^0(X, \mathcal{O}_X(-K_X)) = 2$, $\dim H^0(X, \mathcal{O}_X(-K_X + E)) = 3$ and the element $k_1 \otimes k_2 e - k_2 \otimes k_1 e$ is in the kernel.

Second, the span of the monomials of the first two forms has dimension at most six. Let $aa'e, bb'e$ be any two monomials of the second form. There is a quadratic relation q involving $aa', bb', k_1^2, k_1k_2, k_2^2$ since these five vectors correspond to five elements of the four dimensional space $H^0(X, \mathcal{O}_X(-2K_X))$. Moreover aa' and bb' are independent from k_1^2, k_1k_2, k_2^2 , since the monomials in k_1, k_2 correspond to sections having a base-point, and neither of the remaining elements aa' and bb' vanishes at the base-point. Thus in the relation q the coefficients of aa' and bb' are both nonzero. We deduce that the span of the elements of the first two forms has dimension at most six in $(k[\mathcal{G}]/J_X)_D$.

Third, let abc be a monomial of the third form. The divisor $A + B$ contracts the unique exceptional curve $F := K_X + A + B$, and the divisor $G := A + B - E$ is an exceptional curve on the del Pezzo surface Y obtained from X by contracting F . Therefore there is a quadratic relation involving the monomials ab, ge, k_1f, k_2f and the monomials ab and ge are independent from k_1f, k_2f , by Lemma 8.1. Thus we may use this relation to write the image of abc in $k[\mathcal{G}]/J_X$ as a combination of divisors of the first two forms and the proof is complete. \square

Lemma 8.4. *Let Q be a conic on X ; then the ideal I_X has no minimal generators in degree $D = -K_X + Q$.*

Proof. The monomials of degree D are of the following forms:

Monomial	Description
ks	$k \in \{k_1, k_2\}, s \in k[\mathcal{G}]_Q$
efe_1	$E_1 \cdot Q = 0$ $E \cdot F = 2$ $E_1 \cdot (E + F) = 2$ $Q \cdot E = Q \cdot F = 1$
ee_1e_2	$E_1 \cdot Q = E_2 \cdot Q = E_1 \cdot E_2 = 0$ $Q \cdot E = E_1 \cdot E = E_2 \cdot E = 2$

Note that E_1, E_2 are disjoint components of reducible fibers of the conic bundle associated to Q . Indeed let

$$-K_X + Q \sim A + B + C. \tag{8-2}$$

Multiplying both sides of (8-2) by Q we find

$$2 = A \cdot Q + B \cdot Q + C \cdot Q;$$

thus at least one of the intersection products is zero. Suppose that $C \cdot Q = 0$, and let $\tilde{C} = Q - C$; then $A + B = -K_X + \tilde{C}$ is a reducible divisor in the linear system associated to the pull-back of the anticanonical divisor on the del Pezzo surface of degree two obtained from X by contracting \tilde{E} . The first case in (8-2) corresponds to the sections containing a divisor in the linear system $|-K_X|$; the second to one not containing it and containing two curves not in the linear system $|Q|$; the last one to one containing two elements of $|Q|$.

Since $(k[\mathcal{G}]/J_X)_2 = \text{Cox}(X)_2$, it follows that the dimension of the span of the monomials of the first form in (8-2) modulo $(J_X)_{-K_X+Q}$ is at most four. Let $p = ee_1e_2$ be a monomial in $k[\mathcal{G}]_{-K_X+Q}$ of the third form in (8-2) and let V denote the span of the monomials of the first form in (8-2) together with p in $(k[\mathcal{G}]/J_X)_{-K_X+Q}$. Since $\dim(\text{Cox}(X)_{-K_X+Q}) = 5$, the result follows if we show that

$$V = (k[\mathcal{G}]/J_X)_{-K_X+Q}.$$

Let $q = \bar{e}\bar{e}_1\bar{e}_2$ be a monomial in $k[\mathcal{G}]_{-K_X+Q}$ of the third form in (8-2).

If $\bar{e}_1 = e_1$, then let $E_3 = K_X + E + E_2 = K_X + \bar{E} + \bar{E}_2$ be the exceptional curve contracted by $E + E_2$. By Lemma 8.1, applied to the monomials $\bar{e}\bar{e}_2, ee_2, k_1e_3, k_2e_3$, we conclude that the image of q in $(k[\mathcal{G}]/J_X)_{-K_X+Q}$ belongs to V .

If $\{\bar{E}_1, \bar{E}_2\} \cap \{E_1, E_2\} = \emptyset$, then at least one curve in $\{\bar{E}_1, \bar{E}_2\}$ is disjoint from one curve in $\{E_1, E_2\}$ and relabeling the indices if necessary we may assume that $\bar{E}_1 \cdot E_1 = 0$. By the same reasoning above $q = \bar{e}\bar{e}_1\bar{e}_2$ is in the span of V and $\tilde{q} = \bar{e}\bar{e}_1e_1$ (note that $\tilde{E} := -K_X + Q - \bar{E}_1 - E_1$ is an exceptional curve, since it has anticanonical degree one and square negative one), and we conclude since \tilde{q} is in the span of V .

Finally, if $q = \bar{e}\bar{f}\bar{e}_1$, then at least one curve in E_1, E_2 is disjoint from \bar{E}_1 and relabeling the indices if necessary we may assume that $\bar{E}_1 \cdot E_1 = 0$. Reasoning as above, q is in the span of V and $\bar{e}e_1\bar{e}_1$, and we are done. □

Corollary 8.5. *Let X be a sweeping del Pezzo surface of degree one and let D be an ample divisor on X such that $-K_X \cdot D \geq 3$. Then the ideal I_X has no minimal generators in degree D .*

Proof. By Theorem 7.8, if $-K_X \cdot D \geq 4$ then D is capturable and the result follows from Theorem 6.7. Since the only ample divisors of anticanonical three are $-3K_X, -2K_X + E$ and $-K_X + Q$, where E is an exceptional curve and Q is a conic, the result is true if $-K_X \cdot D = 3$, by Lemmas 8.2–8.4. □

9. Nonample divisors

In this section we prove that the relations in degree D , where D is a nonample divisor on X of anticanonical degree at least three, coming from J_X are sufficient to show that the quotient $(k[\mathcal{G}]/J_X)_D$ is in fact spanned by monomials coming from a del Pezzo surface of smaller degree. This is the basis for the inductive procedure of Section 11. Throughout this section we assume that del Pezzo surfaces X of degree one defined over fields of characteristic two are sweeping.

Lemma 9.1. *Let X be a del Pezzo surface of degree at most five and let D be a divisor on X . Suppose that E is an exceptional curve on X such that $D \cdot E = 0$. Then $(k[\mathcal{G}]/J_X)_D$ is spanned by products of variables corresponding to exceptional curves disjoint from E .*

Proof. Let $m \in k[\mathcal{G}]_D$ be a monomial and write $m = e^p \cdot s$, where $p \geq 0$ and s is a product of variables different from E . Note that if $p = 0$, then s is a product of variables corresponding to divisors disjoint from E , since by assumption $D \cdot E = 0$. We shall show that if $p \geq 1$, then using the quadratic relations we may decrease p ; the result then follows by induction on p .

Suppose that $p \geq 1$. Since $D \cdot E = 0$, there is a variable $c \in \mathcal{G}$ such that $E \cdot C > 0$ and c divides s . The monomial ce is a monomial of anticanonical degree two corresponding to a nef divisor. By definition of J_X , the vector space $(k[\mathcal{G}]/J_X)_{C+E}$ coincides with $\text{Cox}(X)_{C+E}$; thus it suffices to show that there is a basis of $\text{Cox}(X)_{C+E}$ consisting of sections vanishing along exceptional curves different from E . To conclude, we analyze all the possibilities for the divisor $C+E$.

Case 1. The divisor $C+E$ is a conic. The linear system $|C+E|$ contains $8 - \deg(X)$, i.e., at least three, distinct reducible elements and any two of these span it.

Case 2. The divisor $C+E$ is $-b^*K_Y$, where $b: X \rightarrow Y$ is a birational morphism, Y has degree two. The linear system $|C+E|$ contains 28 distinct reducible elements and any five of these span it (Lemma 4.1).

Case 3. The divisor $C+E$ is $-2K_X$ and the degree of X is one. The linear system $|-2K_X|$ contains 120 distinct reducible elements whose irreducible components are (-1) -curves and any 113 of these span it (Proposition 4.4).

In all these cases the sections supported on (-1) -curves distinct from E span $\text{Cox}(X)_{C+E}$, and the lemma follows. \square

Lemma 9.2. *Let X be a del Pezzo surface of degree at most five and let D be a divisor on X . Suppose that E is an exceptional curve on X such that $D \cdot E < 0$. Then $\dim(k[\mathcal{G}]/J_X)_D = \dim(k[\mathcal{G}]/J_X)_{D-E}$ and $\dim \text{Cox}(X)_D = \dim \text{Cox}(X)_{D-E}$.*

Proof. If D is not effective, then $D-E$ is not effective and $k[\mathcal{G}]_D = k[\mathcal{G}]_{D-E} = 0$. If D is effective, then $E \cdot D < 0$ implies that E is a component of D and thus every

monomial in $k[\mathcal{G}]_D$ is divisible by e ; the same argument also proves the statement about the Cox ring. \square

10. Del Pezzo surfaces of degree at least two

Lemma 10.1. *Let X be a del Pezzo surface of degree two. Any ample divisor is capturable except $-K_X$.*

Proof. Let $D \neq -K_X$ be an ample divisor and write $D = -nK_X + N$, with $n \geq 1$ and N nef and not ample; by assumption either $n \geq 2$ or $N \neq 0$.

Suppose first that $N \neq 0$; if N contracts more than one exceptional curve, then we let \mathcal{S} be the set of exceptional curves contracted by N . If N contracts exactly one exceptional curve E , then we let F be any exceptional curve disjoint from E and let $\mathcal{S} := \{E, F\}$.

Let $C \in \mathcal{C} \setminus \mathcal{S}$ and let $C' := -K_X - C$. If $C' \in \mathcal{S}$, then the divisor $L_{C'T} = N + (-K_X - T) - (n - 1)K_X$ is big and nef for all $T \in \mathcal{S}$, $T \neq C'$ since

$$(-K_X - T) \cdot N = -K_X \cdot N > 0;$$

thus we can capture all such curves using \mathcal{S} . If $C' \notin \mathcal{S}$, then let $S, T \in \mathcal{S}$ be disjoint and note that $N - T$ is either nef or N is a multiple of a conic and it is the sum of a multiple of a conic and a single exceptional curve. Moreover

- $S' \cdot (N - T) = -K_X \cdot N - 1 > 0$;
- if N is a multiple of a conic, then S' also intersects the fixed component of $N - T$;
- $C' \cdot (N - T) = -K_X \cdot N - 1 - C \cdot T$ is zero if and only if N is a conic and $C \cdot T = 1$.

In this last case, note that $C \cdot N \neq 0$, and C intersects at least one component in every reducible fiber of the conic bundle determined by N ; hence we may choose T to be the other component if $C' \cdot (N - T) = 0$. In all cases we may choose $S, T \in \mathcal{S}$ disjoint such that L_{ST} is big and nef, and hence capture all $C \in \mathcal{C} \setminus \mathcal{S}$ using only curves in \mathcal{S} . In particular, we are done if N contracts at most two exceptional curves.

To conclude it suffices to treat the two cases $N = L + N'$ and $N = Q + N'$, where L is a twisted cubic, Q is a conic and N' is nef; note also that $S \cdot N' = 0$, for all $S \in \mathcal{S}$.

Choose any two disjoint $S, T \in \mathcal{S}$ and let $C \in \mathcal{S}$, $C \neq S, T$. If $N = L + N'$, we have that the divisor $L_{ST} = -K_X + (-K_X - C) + (L - S - T) + N'$ is big and nef since $(-K_X - C) \cdot (L - S - T) = 1$ and we conclude. If $N = Q + N'$, then the divisor $L_{ST} = (-K_X - S) + (-K_X - T) + (Q - C) + N'$ is big and nef since $(-K_X - S) \cdot (Q - C) = (-K_X - T) \cdot (Q - C) = 1$ and again we conclude.

Suppose now that $N = 0$ and hence $D = -nK_X$, where $n \geq 2$. Let \mathcal{S} be any set of seven disjoint exceptional curves. For all $C \in \mathcal{C} \setminus \mathcal{S}$, there are distinct $S, T \in \mathcal{S}$ such that $C \cdot S, C \cdot T > 0$: this is clear from the list of exceptional curves. Thus for such a choice of S, T , we have that the divisor $L_{ST} = -3K_X - C - S - T = C' + S' + T'$ is big and nef and we may capture all the curves in $\mathcal{C} \setminus \mathcal{S}$ using \mathcal{S} . Finally let $S, T \in \mathcal{S}$ be distinct elements and let $S' := -K_X - S, T' := -K_X - T$; for all $C \in \mathcal{S}$ the divisor $L_{S'T'} = -3K_X - C - S' - T' = C' + S + T$ is big and nef and we conclude. \square

Lemma 10.2. *Let X be a del Pezzo surface of degree three. Any ample divisor is capturable.*

Proof. Let D be an ample divisor and write $D = -K_X + N$, with N a nef divisor.

Let \mathcal{S} be a set of six disjoint exceptional curves. For all $C \in \mathcal{C} \setminus \mathcal{S}$ there are two distinct $S, T \in \mathcal{S}$ such that $C \cdot S = C \cdot T = 1$. With these choices of S, T we have that the divisor $L_{ST} = -2K_X + N - C - S - T = (-K_X - T) + (-K_X - C - S) + N$ is big and nef since $(-K_X - T) \cdot (-K_X - C - S) = 1$. Thus we may capture all the exceptional curves $C \in \mathcal{C} \setminus \mathcal{S}$ using the curves in \mathcal{S} . Let \mathcal{S}' be the set of six disjoint exceptional curves each intersecting all the curves in \mathcal{S} except for one (thus $\mathcal{S} \cup \mathcal{S}'$ is a *Schläfli double-six*). Choose $S, T \in \mathcal{S}$ and let $S', T' \in \mathcal{S}'$ be two corresponding curves, that is $S \cdot S' = T \cdot T' = 0$. Using the moves above we can capture $\{S', T'\}$ with \mathcal{S} and we can capture $\mathcal{S} \setminus \{S, T\}$ using $\mathcal{S}' \setminus \{S', T'\}$, and finally we capture $\mathcal{S}' \setminus \{S', T'\}$ using $\{S, T\}$. \square

Lemma 10.3. *Let X be a del Pezzo surface of degree four. Any ample divisor is capturable.*

Proof. Let D be an ample divisor and write $D = -K_X + N$, with N a nef divisor. The criterion to capture curves is that $(\{E, F\}, C)$ is a capture move if $C \notin \{E, F\}$, $E \cdot F = 0$ and $C \cdot E + C \cdot F > 0$: if $C \cdot E > 0$, then the divisor

$$L_{EF} = (-K_X - C - E) + (-K_X - F) + N$$

is the sum of a conic, a twisted cubic and a nef divisor and hence it is big and nef.

We refer to Table 1. Use E_3, E_4 to capture $L - E_i - E_j$ for all $3 \leq i < j < 5$. Use $L - E_1 - E_3, L - E_1 - E_4, L - E_1 - E_5$ to capture E_3, E_4, E_5 and finally use E_1, E_2 to capture all the remaining curves. \square

Lemma 10.4. *Let X be a del Pezzo surface of degree five. Any ample divisor is capturable.*

Proof. Same criterion and strategy as in Lemma 10.3, ignoring any reference to the index 5. \square

Lemma 10.5. *Let X be a del Pezzo surface of degree six. Any ample divisor is capturable.*

Proof. Same criterion as in Lemma 10.3. Capture $L - E_1 - E_2$ using E_1, E_2 and then capture the remaining curves using $L - E_1 - E_3, L - E_2 - E_3$. \square

11. Quadratic generation

Main result. We collect all the information gathered in the previous sections to prove the main result of this paper: the ideal I_X is generated by its degree two part.

Proof of the Batyrev–Popov Conjecture. Let n be an integer; by induction on $r_X := 9 - \deg(X)$ and by induction on n we show that $(J_X)_n = (I_X)_n$. By definition, the statement is true if $n \leq 2$, for all del Pezzo surfaces X ; if $r_X \leq 4$, then the result is well known for all n .

Suppose that $r_X > 4$, $n \geq 3$, that for all del Pezzo surfaces Y such that $r_Y < r_X$ we have that $J_Y = I_Y$, and that $(J_X)_{n-1} = (I_X)_{n-1}$. Let D be a divisor on X of anticanonical degree n ; if there is an exceptional curve E such that $D \cdot E \leq 0$, then the result follows by Lemma 9.2 or by Lemma 9.1. Otherwise D is ample and the result follows from Corollary 8.5 if $\deg(X) = 1$, or from the lemmas of Section 10 if $\deg(X) \geq 2$. \square

Quadratic generators. Let X be a del Pezzo surface of degree d . We briefly explain how all generators of I_X arise. The nef divisors D with anticanonical degree two on X are:

- (1) the conics Q ,
- (2) $-K_X$ if $d = 2$,
- (3) $-K_X + E$ if $d = 1$, where E is an exceptional curve on X ,
- (4) $-2K_X$ if $d = 1$.

These divisors are precisely sums of pairs of intersecting exceptional curves on X .

We count the relations coming from conics as follows. Every conic Q has $8 - d$ reducible sections and $h^0(X, \mathcal{O}_X(Q)) = 2$; thus each conic gives rise to $6 - d$ quadratic relations. If there are B conics on X then there are $(6 - d)B$ generators of I_X induced by conic bundles. For example, when $d = 1$ we obtain $5 \times 2,160 = 10,800$ relations.

When $d = 2$, we also have relations in degree $-K_X$. There are 28 monomials in $k[\mathcal{G}]_{-K_X}$, and $h^0(X, \mathcal{O}_X(-K_X)) = 3$, giving 25 linear dependence relations among these quadratic monomials. These relations yield the remaining generators of I_X .

When $d = 1$, there are 30 monomials in $k[\mathcal{G}]_{-K_X+E}$: k_1e, k_2e and the 28 monomials coming from the anticanonical divisor of the del Pezzo surface of degree two obtained by contracting E . This gives $(30 - 3) \times 240 = 6,480$ generators in I_X . Finally, we also have relations coming from $-2K_X$. There are 123 monomials in $k[\mathcal{G}]_{-2K_X}$: k_1^2, k_1k_2, k_2^2 and the 120 monomials of the form ee' , where $E, E' \in \mathcal{C}$ and

$\deg(X)$	D	Number of monomials in $k[\mathcal{G}]_D$	$h^0(X, \mathbb{O}_X(D))$	$b_{1,D}(\text{Cox}(X))$	Number of divisors of type D
1	Q	7	2	5	2160
	$-K_X + E$	$28 + 2$	3	27	240
	$-2K_X$	$120 + 3$	4	119	1
	(Total)	22443		17399	2401
2	Q	6	2	4	126
	$-K_X$	28	3	25	1
	(Total)	784		529	127
3	Q	5	2	3	27
	(Total)	135		81	27
4	Q	4	2	2	10
	(Total)	40		20	10
5	Q	3	2	1	5
	(Total)	15		5	5

Table 2. First Betti numbers $b_{1,D}(\text{Cox}(X))$ for del Pezzo surfaces.

$E + E' = -2K_X$. Since $h^0(X, \mathbb{O}_X(-2K_X)) = 4$, we obtain 119 linear dependence relations among these quadratic monomials. These relations yield the remaining generators of I_X . This information is summarized in Table 2.

Acknowledgments

We are grateful to Bernd Sturmfels and David Eisenbud for many profitable discussions. We thank Ragni Piene for valuable conversations on the Gauss map for curves in positive characteristic during the Spring 2009 Algebraic Geometry semester at MSRI. We would also like to thank the anonymous referee for carefully reading the manuscript and for suggesting useful improvements.

References

- [Batyrev and Popov 2004] V. V. Batyrev and O. N. Popov, “The Cox ring of a del Pezzo surface”, pp. 85–103 in *Arithmetic of higher-dimensional algebraic varieties* (Palo Alto, CA, 2002), edited by B. Poonen and Y. Tschinkel, Progr. Math. **226**, Birkhäuser, Boston, MA, 2004. MR 2005h:14091 Zbl 1075.14035
- [Birkar et al. 2008] C. Birkar, P. Cascini, C. D. Hacon, and J. McKernan, “Existence of minimal models for varieties of log general type”, preprint, 2008. arXiv 0610203v2
- [Bosma et al. 1997] W. Bosma, J. Cannon, and C. Playoust, “The Magma algebra system, I: The user language”, *J. Symbolic Comput.* **24**:3-4 (1997), 235–265. MR 1484478 Zbl 0898.68039

- [de la Bretèche and Browning 2007] R. de la Bretèche and T. D. Browning, “On Manin’s conjecture for singular del Pezzo surfaces of degree 4, I”, *Michigan Math. J.* **55**:1 (2007), 51–80. MR 2008b:14028 Zbl 1132.14019
- [de la Bretèche et al. 2007] R. de la Bretèche, T. D. Browning, and U. Derenthal, “On Manin’s conjecture for a certain singular cubic surface”, *Ann. Sci. École Norm. Sup. (4)* **40**:1 (2007), 1–50. MR 2008e:11038 Zbl 1125.14008
- [Brion 2007] M. Brion, “The total coordinate ring of a wonderful variety”, *J. Algebra* **313**:1 (2007), 61–99. MR 2008d:14067 Zbl 1123.14024
- [Castravet and Tevelev 2006] A.-M. Castravet and J. Tevelev, “Hilbert’s 14th problem and Cox rings”, *Compos. Math.* **142**:6 (2006), 1479–1498. MR 2007i:14044 Zbl 1117.14048
- [Colliot-Thélène and Sansuc 1980] J.-L. Colliot-Thélène and J.-J. Sansuc, “La descente sur les variétés rationnelles”, pp. 223–237 in *Journées de Géométrie Algébrique d’Angers, Juillet 1979* (Alphen aan den Rijn, 1979), Sijthoff & Noordhoff, 1980. Zbl 0451.14018
- [Colliot-Thélène and Sansuc 1987] J.-L. Colliot-Thélène and J.-J. Sansuc, “La descente sur les variétés rationnelles, II”, *Duke Math. J.* **54**:2 (1987), 375–492. MR 89f:11082 Zbl 0659.14028
- [Colliot-Thélène et al. 1984] J.-L. Colliot-Thélène, J.-J. Sansuc, and P. Swinnerton-Dyer, “Intersections de deux quadriques et surfaces de Châtelet”, *C. R. Acad. Sci. Paris Sér. I Math.* **298**:16 (1984), 377–380. MR 85j:11031 Zbl 0575.14044
- [Cox 1995] D. A. Cox, “The homogeneous coordinate ring of a toric variety”, *J. Algebraic Geom.* **4**:1 (1995), 17–50. MR 95i:14046 Zbl 0846.14032
- [Cragolini and Oliverio 2000] P. Cragolini and P. A. Oliverio, “Lines on del Pezzo surfaces with $K_S^2 = 1$ in characteristic 2 in the smooth case”, *Portugal. Math.* **57**:1 (2000), 59–95. MR 2001c:14057 Zbl 0989.14012
- [Debarre 2001] O. Debarre, *Higher-dimensional algebraic geometry*, Universitext **13**, Springer, New York, 2001. MR 2002g:14001 Zbl 0978.14001
- [Derenthal 2007] U. Derenthal, “Universal torsors of del Pezzo surfaces and homogeneous spaces”, *Adv. Math.* **213**:2 (2007), 849–864. MR 2008i:14052 Zbl 1125.14020
- [Elizondo et al. 2004] E. J. Elizondo, K. Kurano, and K.-i. Watanabe, “The total coordinate ring of a normal projective variety”, *J. Algebra* **276**:2 (2004), 625–637. MR 2005b:14013 Zbl 1074.14006
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Grad. Texts in Math. **52**, Springer, New York, 1977. MR 57 #3116 Zbl 0367.14001
- [Hassett 2004] B. Hassett, “Equations of universal torsors and Cox rings”, pp. 135–143 in *Mathematisches Institut, Georg-August-Universität Göttingen: Seminars Summer Term 2004*, edited by Y. Tschinkel, Universitätsdrucke Göttingen, Göttingen, 2004. MR 2007a:14046 Zbl 1108.14304
- [Hassett and Tschinkel 2004] B. Hassett and Y. Tschinkel, “Universal torsors and Cox rings”, pp. 149–173 in *Arithmetic of higher-dimensional algebraic varieties* (Palo Alto, CA, 2002), edited by B. Poonen and Y. Tschinkel, Progr. Math. **226**, Birkhäuser, Boston, MA, 2004. MR 2005a:14049 Zbl 1077.14046
- [Hefez and Kleiman 1985] A. Hefez and S. L. Kleiman, “Notes on the duality of projective varieties”, pp. 143–183 in *Geometry today* (Rome, 1984), edited by E. Arbarello et al., Progr. Math. **60**, Birkhäuser, Boston, MA, 1985. MR 88f:14046 Zbl 0579.14047
- [Hu and Keel 2000] Y. Hu and S. Keel, “Mori dream spaces and GIT”, *Michigan Math. J.* **48** (2000), 331–348. MR 2001i:14059 Zbl 1077.14554
- [Kaji 1992] H. Kaji, “On the inseparable degrees of the Gauss map and the projection of the conormal variety to the dual of higher order for space curves”, *Math. Ann.* **292**:3 (1992), 529–532. MR 93j:14038 Zbl 0734.14017

- [Kollár 1996] J. Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Math. (3) **32**, Springer, Berlin, 1996. MR 98c:14001 Zbl 0877.14012
- [Laface and Velasco 2009] A. Laface and M. Velasco, “Picard-graded Betti numbers and the defining ideals of Cox rings”, *J. Algebra* **322**:2 (2009), 353–372. MR MR2529093
- [Manin 1986] Y. I. Manin, *Cubic forms*, 2nd ed., North-Holland Mathematical Library **4**, North-Holland Publishing Co., Amsterdam, 1986. Algebra, geometry, arithmetic, Translated from the Russian by M. Hazewinkel. MR 87d:11037 Zbl 0582.14010
- [Peyre 2004] E. Peyre, “Counting points on varieties using universal torsors”, pp. 61–81 in *Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002)*, edited by B. Poonen and Y. Tschinkel, Progr. Math. **226**, Birkhäuser, Boston, MA, 2004. MR 2004m:11103
- [Popov 2004] O. N. Popov, “The Cox ring of a Del Pezzo surface has rational singularities”, preprint, 2004. arXiv 0402154v1
- [Reid 1997] M. Reid, “Chapters on algebraic surfaces”, pp. 3–159 in *Complex algebraic geometry* (Park City, UT, 1993), edited by J. Kollár, IAS/Park City Math. Ser. **3**, Amer. Math. Soc., Providence, RI, 1997. MR 98d:14049 Zbl 0910.14016
- [Salberger 1998] P. Salberger, “Tamagawa measures on universal torsors and points of bounded height on Fano varieties”, pp. 91–258 in *Nombre et répartition de points de hauteur bornée* (Paris, 1996), vol. 251, edited by E. Peyre, 1998. MR 2000d:11091 Zbl 0959.14007
- [Serganova and Skorobogatov 2007] V. V. Serganova and A. N. Skorobogatov, “Del Pezzo surfaces and representation theory”, *Algebra Number Theory* **1**:4 (2007), 393–419. MR 2009b:14070 Zbl 1170.14026
- [Serganova and Skorobogatov 2008] V. Serganova and A. Skorobogatov, “On the equations for universal torsors over del Pezzo surfaces”, preprint, 2008. arXiv 0806.0089
- [Stillman et al. 2007] M. Stillman, D. Testa, and M. Velasco, “Gröbner bases, monomial group actions, and the Cox rings of del Pezzo surfaces”, *J. Algebra* **316**:2 (2007), 777–801. MR 2008i:14054
- [Sturmfels and Xu 2008] B. Sturmfels and Z. Xu, “Sagbi Bases of Cox–Nagata rings”, preprint, 2008. to appear in *J. European Math. Soc.* arXiv 0803.0892
- [Terakawa 1998] H. Terakawa, “On the Kawamata–Viehweg vanishing theorem for a surface in positive characteristic”, *Arch. Math. (Basel)* **71**:5 (1998), 370–375. MR 99k:14031 Zbl 0940.14010
- [Testa et al. 2009] D. Testa, A. Várilly-Alvarado, and M. Velasco, “Big rational surfaces”, preprint, 2009. arXiv 0901.1094v2
- [Xie 2008] Q. Xie, “Kawamata–Viehweg vanishing on rational surfaces in positive characteristic”, 2008. arXiv 0710.2706v5

Communicated by Jean-Louis Colliot-Thélène

Received 2008-03-08

Revised 2009-06-05

Accepted 2009-09-14

adomani@gmail.com

*Mathematical Institute, 24-29 St Giles', Oxford OX1 3LB,
United Kingdom*
<http://www.maths.ox.ac.uk/node/7850>

varilly@rice.edu

*Department of Mathematics, Rice University, MS 136,
Houston, TX 77005, United States*
<http://math.rice.edu/~av15>

velasco@math.berkeley.edu

*Department of Mathematics, University of California,
Berkeley, CA 94720, United States*
<http://math.berkeley.edu/~velasco>

