

Algebra & Number Theory

Volume 3

2009

No. 7

The half-twist for $U_q(\mathfrak{g})$ representations

Noah Snyder and Peter Tingley



mathematical sciences publishers

The half-twist for $U_q(\mathfrak{g})$ representations

Noah Snyder and Peter Tingley

We introduce the notion of a half-ribbon Hopf algebra, which is a Hopf algebra \mathcal{H} along with a distinguished element $t \in \mathcal{H}$ such that (\mathcal{H}, R, C) is a ribbon Hopf algebra, where $R = (t^{-1} \otimes t^{-1})\Delta(t)$ and $C = t^{-2}$. The element t is closely related to the topological “half-twist”, which twists a ribbon by 180 degrees. We construct a functor from a topological category of ribbons with half-twists to the category of representations of any half-ribbon Hopf algebra. We show that $U_q(\mathfrak{g})$ is a (topological) half-ribbon Hopf algebra, but that t^{-2} is not the standard ribbon element. For $U_q(\mathfrak{sl}_2)$, we show that there is no half-ribbon element t such that t^{-2} is the standard ribbon element. We then discuss how ribbon elements can be modified, and some consequences of these modifications.

1. Introduction	809
2. Conventions	811
3. Background	812
4. Half-ribbon Hopf algebras	816
5. Ribbon and half-ribbon elements for $U_q(\mathfrak{g})$	824
6. Questions	832
Acknowledgments	833
References	833

1. Introduction


Let $\mathcal{RIBBON}(S)$ be the category whose morphisms consist of tangles of oriented directed ribbons up to isotopy, each labeled with an element of some set S . There is a notion of a ribbon Hopf algebra \mathcal{H} (see [Chari and Pressley 1994], for example), which is related to this topological category by the fact that there is a monoidal functor \mathcal{F}' from $\mathcal{RIBBON}(\mathcal{H}\text{-rep})$ to the category of representations $\mathcal{H}\text{-rep}$. This allows one to construct invariants of oriented framed links, and from there invariants of ordinary links.

MSC2000: primary 17B37; secondary 57T05, 57M05.

Keywords: quantum group, Hopf algebra, ribbon category.

Both authors were supported in part by RTG grant DMS-0354321. Tingley was also partly supported by the Australian Research Council grant DP0879951 and NSF grant DMS-0902649.

There is a morphism in $\mathcal{RIBBON}(S)$ which twists a ribbon by 360 degrees, but not one that twists a ribbon by 180 degrees (since negatively oriented objects are not allowed). We propose that one should consider a slightly larger category, denoted $\mathcal{HTRIB}(S)$, where this 180 degree twist, the “half-twist,” is allowed. The half-twist can be applied to several ribbons at once, and all morphisms in $\mathcal{HTRIB}(S)$ can be constructed out of half-twists, along with various caps and cups. The following isotopy shows how the crossing in $\mathcal{HTRIB}(S)$ is constructed out of the half-twist:



Recall that a ribbon Hopf algebra \mathcal{H} is a Hopf algebra along with two extra features:

- A universal R -matrix $R \in \mathcal{H} \otimes \mathcal{H}$. The functor \mathcal{F}' takes a simple crossing of ribbons labeled V and W to $\text{Flip} \circ R$ acting on $V \otimes W$.
- A central “ribbon” element $C \in \mathcal{H}$. The functor \mathcal{F}' takes a 360 degree twist of a ribbon labeled V to C acting on V .

The elements R and C must satisfy various compatibility conditions (see [Section 4A](#)).

In the present work we define a half-ribbon Hopf algebra to be a ribbon Hopf algebra along with a distinguished element t such that

- (i) $R = (t^{-1} \otimes t^{-1})\Delta(t)$,
- (ii) $C = t^{-2}$.

We show that, if \mathcal{H} is a half-ribbon Hopf algebra, then \mathcal{F}' can be extended to a functor \mathcal{F} from $\mathcal{HTRIB}(S)$ to \mathcal{H} -rep.

Our main interest is the case where \mathcal{H} is the quantized universal enveloping algebra $U_q(\mathfrak{g})$ of a finite-dimensional complex simple Lie algebra \mathfrak{g} . In this case $U_q(\mathfrak{g})$ is actually a topological ribbon Hopf algebra, meaning that R and C only lie in some completion, not in $U_q(\mathfrak{g})$ itself. We define a topological half-ribbon Hopf algebra by allowing t to lie in a completion of \mathcal{H} , and show that $U_q(\mathfrak{g})$ has this structure. The main ingredient is a formula for the R matrix of $U_q(\mathfrak{g})$, due to Kirilov and Reshetikhin [1990] and Levendorskiĭ and Soibelman [1991], of the form

$$R = (X^{-1} \otimes X^{-1})\Delta(X). \tag{2}$$

The correspondence to condition (i) above should be clear, and we show that X is in fact a half-ribbon element for $U_q(\mathfrak{g})$. Interestingly, X^{-2} is not the standard ribbon element, and in fact we show that in certain cases it is not possible to find a half-ribbon element t for $U_q(\mathfrak{g})$ such that t^{-2} is the standard ribbon element.

In Section 5 we discuss the different ribbon elements for $U_q(\mathfrak{g})$, and in some small cases we describe exactly which ones arise from half-ribbon elements. We then discuss some consequences of using X^{-2} in place of the usual ribbon element. In particular, it simplifies the correspondence between certain skein theoretic constructions of link invariants and the quantum group constructions of the same objects, essentially by explaining some annoying negative signs that appear in, for example, [Ohtsuki 2002] or [Kuperberg 1996].

We feel it would be of considerable interest to study which ribbon Hopf algebras can be given the structure of half-ribbon Hopf algebras. It would also be nice to give straightforward conditions on an element t in a general Hopf algebra \mathcal{H} , such that \mathcal{H} along with t is a half-ribbon Hopf algebra (that is, \mathcal{H} is a ribbon Hopf algebra, with $R = (t^{-1} \otimes t^{-1})\Delta(t)$ and $C = t^{-2}$). At the end of the paper we discuss these and other possible future directions.

2. Conventions

We first fix some notation. For the most part we follow conventions from [Chari and Pressley 1994].

- \mathfrak{g} is a complex simple Lie algebra with Cartan algebra \mathfrak{h} , and $A = (a_{ij})_{i,j \in I}$ is its Cartan matrix.

- $\langle \cdot, \cdot \rangle$ denotes the pairing between \mathfrak{h} and \mathfrak{h}^* and (\cdot, \cdot) denotes the usual symmetric bilinear form on either \mathfrak{h} or \mathfrak{h}^* . Fix the usual bases α_i for \mathfrak{h}^* and H_i for \mathfrak{h} , and recall that $\langle H_i, \alpha_j \rangle = a_{ij}$.

- $d_i = (\alpha_i, \alpha_i)/2$, so that $(H_i, H_j) = d_j^{-1}a_{ij}$. Let B denote the matrix $(d_j^{-1}a_{ij})$.
- $q_i = q^{d_i}$.
- ρ is the element of \mathfrak{h}^* such that $(\alpha_i, \rho) = d_i$ for all i .
- ρ^\vee is the element of \mathfrak{h} such that $\langle \alpha_i, \rho^\vee \rangle = 1$ for all i .
- s_i is the element of the Weyl group which is defined by

$$s_i(\alpha_j) = \alpha_j - \langle \alpha_i, \alpha_j^\vee \rangle \alpha_i. \tag{3}$$

- θ is the diagram automorphism such that $w_0(\alpha_i) = -\alpha_{\theta(i)}$, where w_0 is the longest element in the Weyl group.

- $U_q(\mathfrak{g})$ is the quantized universal enveloping algebra associated to \mathfrak{g} , generated over $\mathbb{C}(q)$ by E_i, F_i for all $i \in I$, and K_w for w in the coweight lattice of \mathfrak{g} . As usual, let $K_i = K_{H_i}$. We actually must adjoin a fixed k th root of q to the base field, for some k depending on \mathfrak{g} . This causes no real difficulty, and for the most part

we ignore it. For convenience, we recall the exact formula for the coproduct and antipode:

$$\begin{cases} \Delta E_i = E_i \otimes K_i + 1 \otimes E_i, \\ \Delta F_i = F_i \otimes 1 + K_i^{-1} \otimes F_i, \\ \Delta K_i = K_i \otimes K_i, \end{cases} \quad \begin{cases} S(E_i) = -E_i K_i^{-1}, \\ S(F_i) = -K_i F_i, \\ S(K_i) = K_i^{-1}. \end{cases} \quad (4)$$

- $[n] = (q^n - q^{-n}) / (q - q^{-1})$, and $X^{(n)} = (X^n) / ([n][n-1] \cdots [2])$.
- V_λ is the irreducible representation of $U_q(\mathfrak{g})$ with highest weight λ , and v_λ is a highest weight vector.
- P is the weight lattice for \mathfrak{g} and Q is the root lattice.
- The standard ribbon element C for $U_q(\mathfrak{g})$ acts on V_λ as the constant

$$q^{-(\lambda, \lambda) - 2(\lambda, \rho)}.$$

- $U_q(\mathfrak{g})$ -rep is the category of finite-dimensional Type 1 representations.
- (\mathcal{H}, R) is a quasitriangular Hopf algebra over a field F , where
 - (i) $\mu : \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H}$ is multiplication;
 - (ii) $\iota : F \rightarrow \mathcal{H}$ is the unit;
 - (iii) $\Delta : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$ is the comultiplication;
 - (iv) $\varepsilon : \mathcal{H} \rightarrow F$ is the counit;
 - (v) $S : \mathcal{H} \rightarrow \mathcal{H}$ is the antipode.

• Flip : $V \otimes W \rightarrow W \otimes V$ is defined by $\text{Flip}(v \otimes w) = w \otimes v$. For longer tensor products, define $\text{rev} : V_1 \otimes \cdots \otimes V_k \rightarrow V_k \otimes \cdots \otimes V_1$ by $v_1 \otimes \cdots \otimes v_k \mapsto v_k \otimes \cdots \otimes v_1$.

• If V is a representation of a Hopf algebra H , then we define the left dual V^* to be the dual vector space with the action $x \circ f(v) = f(S(x)v)$, and the right dual *V to be the dual vector space with the action $x \circ f(v) = f(S^{-1}(x)v)$. Note that $({}^*V)^* = V = {}^*(V^*)$.

3. Background

Much of the motivation for this paper comes from studying an expression for the R -matrix

$$R = (X^{-1} \otimes X^{-1}) \Delta(X), \quad (5)$$

where X is an element in some completion of $U_q(\mathfrak{g})$. This was first introduced by Kirilov and Reshetikhin [1990] and Levendorskiĭ and Soibelman [1991], and has recently proven useful in studying the relationship between the braiding and crystal bases [Kamnitzer and Tingley 2009]. In this section we review this formula, and also recall the definition and basic properties of a ribbon Hopf algebra.

3A. A completion of $U_q(\mathfrak{g})$. The element X from (5) is not actually in $U_q(\mathfrak{g})$, but only in a completion. In order to be precise, we briefly review the completion of $U_q(\mathfrak{g})$ that we use.

Definition 3.1. $\widetilde{U_q(\mathfrak{g})}$ is the completion of $U_q(\mathfrak{g})$ in the weak topology generated by all matrix elements of all (finite-dimensional Type 1) representations. Similarly, $\widetilde{U_q(\mathfrak{g})} \otimes \widetilde{U_q(\mathfrak{g})}$ is the completion of $U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g})$ in the weak topology defined by all matrix elements of representations $V_\lambda \otimes V_\mu$, for all ordered pairs (λ, μ) .

Theorem 3.2 [Kamnitzer and Tingley 2009]. $\widetilde{U_q(\mathfrak{g})}$ is the direct product of the endomorphism rings of all irreducible representations of $U_q(\mathfrak{g})$. That is,

$$\widetilde{U_q(\mathfrak{g})} = \prod_{\lambda \in \Lambda_+} \text{End}(V_\lambda). \tag{6}$$

Comment 3.3. It is straightforward to see that $\widetilde{U_q(\mathfrak{g})}$ is a topological Hopf algebra.

By Theorem 3.2, specifying an element of $\widetilde{U_q(\mathfrak{g})}$ is exactly the same as specifying how it acts on each isomorphism class of irreducible representation. Similarly, if we want to specify an element of $\widetilde{U_q(\mathfrak{g})}^{\otimes 2}$ we just need to say how it acts on every tensor product of any two irreducible representations.

3B. A method of constructing commutativity constraints. The next result is a development of an idea introduced in [Henriques and Kamnitzer 2006] to study the crystal commutor; the proof is a straightforward exercise.

Proposition 3.4 [Kamnitzer and Tingley 2009, Proposition 3.11]. *Let Y be an invertible element in $\widetilde{U_q(\mathfrak{g})}$ such that the map $C_Y : X \rightarrow YXY^{-1}$ restricts to an algebra automorphism of $U_q(\mathfrak{g})$. Then C_Y is a coalgebra antiautomorphism if and only if, for every pair of representations V and W , the map*

$$\begin{aligned} \sigma_{V,W}^Y : V \otimes W &\rightarrow W \otimes V \\ v \otimes w &\rightarrow \text{Flip} \circ (Y^{-1} \otimes Y^{-1}) \Delta(Y)v \otimes w \end{aligned} \tag{7}$$

is an isomorphism.

Given an element Y satisfying the conditions of Proposition 3.4, we use the notation σ^Y to denote the system of isomorphisms $\{\sigma_{V,W}^Y\}$.

3C. The element X . We now explicitly describe the element X from (5). We will need a way to specify a lowest weight vector of V_λ , depending linearly on a chosen highest weight vector. We do this using the action of the braid group on V_λ . We very briefly review this theory, and refer the reader to, for example, [Chari and Pressley 1994, Chapter 8.1.2] or [Lusztig 1993] for more details.

Definition 3.5 [Lusztig 1993, 5.2.1]. T_i is the element of $\widetilde{U_q(\mathfrak{g})}$ that acts on a weight vector v by:

$$T_i(v) = \sum_{\substack{a,b,c \geq 0 \\ a-b+c = (\text{wt}(v), \alpha_i)}} (-1)^b q_i^{ac-b} E_i^{(a)} F_i^{(b)} E_i^{(c)} v.$$

By [Lusztig 1993, Theorem 39.4.3], these T_i generate an action of the braid group on each V_λ , and thus a map from the braid group to $\widetilde{U_q(\mathfrak{g})}$. This realization of the braid group is often referred to as the quantum Weyl group. It is related to the classical Weyl group by the fact that, for any weight vector $v \in V$, $\text{wt}(T_i(v)) = s_i(\text{wt}(v))$.

Definition 3.6. Let $w_0 = s_{i_1} s_{i_2} \cdots s_{i_N}$ be a reduced expression for the longest word w_0 in the Weyl group. Then $T_{w_0} = T_{i_1} T_{i_2} \cdots T_{i_N}$.

It is clear that T_{w_0} is well defined because the elements T_i satisfy the braid relations. Note that T_{w_0} interchanges the highest and lowest weight spaces of V_λ .

Definition 3.7. v_λ^{low} is the element in the lowest weight space of V_λ defined by

$$T_{w_0}(v_\lambda^{\text{low}}) = v_\lambda,$$

where v_λ is the chosen highest weight vector.

Definition 3.8. J is the element of $\widetilde{U_q(\mathfrak{g})}$ defined by, for any weight vector $v \in V_\lambda$,

$$J(v) = q^{(\text{wt}(v), \text{wt}(v))/2 + (\text{wt}(v), \rho)} v.$$

Definition 3.9. $X = JT_{w_0}$.

Lemma 3.10 [Kamnitzer and Tingley 2009, Section 5.2]. *The element X has the following properties:*

- (i) $X(v_\lambda^{\text{low}}) = q^{(\lambda, \lambda)/2 + (\lambda, \rho)} v_\lambda$.
- (ii) $X(v_\lambda) = (-1)^{\langle 2\lambda, \rho^\vee \rangle} q^{(\lambda, \lambda)/2 + (\lambda, \rho)} v_\lambda^{\text{low}}$.
- (iii) X^2 is central, and acts on V_λ as multiplication by the scalar

$$(-1)^{\langle 2\lambda, \rho^\vee \rangle} q^{(\lambda, \lambda) + 2(\lambda, \rho)}.$$

- (iv) C_X is given by

$$\begin{cases} C_X(E_i) = -F_{\theta(i)}, \\ C_X(F_i) = -E_{\theta(i)}, \\ C_X(K_i) = K_{\theta(i)}^{-1}. \end{cases} \quad (8)$$

We next present a key result:

Theorem 3.11 ([Kirillov and Reshetikhin 1990, Theorem 3; Levendorskiĭ and Soĭbel'man 1991, Theorem 1]; see [Kamnitzer and Tingley 2009] for this exact statement.). σ^X is the standard braiding. Equivalently, the standard R -matrix for $U_q(\mathfrak{g})$ can be realized as

$$R = (X^{-1} \otimes X^{-1})\Delta(X). \tag{9}$$

In fact, we will take (9) as the definition of the universal R -matrix.

3D. Ribbon Hopf algebras.

Definition 3.12 [Chari and Pressley 1994, Definition 4.2.8]. A ribbon Hopf algebra (\mathcal{H}, R, v) is a quasitriangular Hopf algebra (\mathcal{H}, R) equipped with an invertible central element v such that

- (i) $v^2 = uS(u)$, where $u = \mu(S \otimes id)R_{21}$,
- (ii) $S(v) = v$,
- (iii) $\varepsilon(v) = 1$,
- (iv) $\Delta(v) = (v \otimes v)(R_{21}R_{12})^{-1}$.

Proposition 3.13 [Chari and Pressley 1994]. The element $g := v^{-1}u$ is grouplike (where u is as in Definition 3.12(i)). □

The following four maps are crucial to studying ribbon Hopf algebras:

Definition 3.14 [Chari and Pressley 1994, page 163]. Let (\mathcal{H}, R, v) be a ribbon Hopf algebra, and (V, π) a representation of \mathcal{H} . Let $f \in V^*$ and $v \in V$. Let e_i and e^i be dual bases of V and V^* respectively. Then

- (i) $ev(f \otimes v) = f(v)$;
- (ii) $\widetilde{ev}(v \otimes f) = f(gv)$;
- (iii) $\text{coev}(1) = \sum e_i \otimes e^i$;
- (iv) $\widetilde{\text{coev}}(1) = \sum e^i \otimes g^{-1}e_i$.

Recall that $V \otimes V^*$ can be identified with $\text{End}(V)$. Under this identification \widetilde{ev} is the quantum trace of [Reshetikhin and Turaev 1991].

We will be working with $U_q(\mathfrak{g})$, which is not a ribbon Hopf algebra according to the above definition, since the elements v and R actually lie in the completions $\widetilde{U_q(\mathfrak{g})}$ and $U_q(\mathfrak{g}) \otimes \widetilde{U_q(\mathfrak{g})}$ discussed in Section 3A. This is known as a topological ribbon Hopf algebra. The theory goes through just as well in the topological case.

Definition 3.15. Let C be the element of $\widetilde{U_q(\mathfrak{g})}$ which acts on the representation V_λ as multiplication by $q^{-(\lambda, \lambda) - 2(\lambda, \rho)}$.

The following is well known (see [Chari and Pressley 1994, Corollary 8.3.16], for example).

Theorem 3.16. $(U_q(\mathfrak{g}), R, C)$ is a topological ribbon Hopf algebra. □

3E. The Temperley–Lieb category. The unoriented Temperley–Lieb category TL has as objects collections of points on the real line up to isotopy, and a morphism from A to B is a formal linear combination of “planar arc diagrams” each of which is a collection of nonintersecting segments in $\mathbb{R} \times [0, 1]$ whose boundary consists of $A \times 0$ and $B \times 1$ modulo isotopy in the plane and the following relation:

$$\bigcirc \longrightarrow -q - q^{-1}. \tag{10}$$

As discussed in, for example, [Goodman and Wenzl 2002], TL is a rigid monoidal category, where composition is given by vertical stacking, tensor product is given by disjoint union, and the dual is given by 180-degree rotation. Since the double dual acts trivially, this is a pivotal category with the trivial pivotal structure.

4. Half-ribbon Hopf algebras

This section contains the definition of a half-ribbon Hopf algebra. It also explains the relationship with the topological category $\mathcal{HRLB}(S)$.

4A. Definition and basic properties.

Definition 4.1. A half-ribbon Hopf algebra is a Hopf algebra, together with a distinguished element $t \in \mathcal{H}$, such that (\mathcal{H}, R, v) is a ribbon Hopf algebra, where $R = (t^{-1} \otimes t^{-1})\Delta(t)$ and $v = t^{-2}$.

In the case where t only exists in a completion of \mathcal{H} , we say \mathcal{H} is a topological half-ribbon Hopf algebra.

Proposition 4.2. *A half-ribbon element t in a half-ribbon Hopf algebra has the following properties:*

- (i) *The algebra automorphism $C_t : \mathcal{H} \rightarrow \mathcal{H}$ defined by $x \rightarrow txt^{-1}$ is also a coalgebra antiautomorphism.*
- (ii) *$\varepsilon(t) = 1$. Equivalently, t acts as the identity on the trivial representation.*
- (iii) *$S(t)^2 = t^2$.*

Proof. (i) follows from $R\Delta(x) = \Delta^{op}(x)R$.

(ii) follows from the fact that R acts as the identity on $\mathbf{1} \otimes V$, where $\mathbf{1}$ is the trivial representation.

(iii) follows because t^{-2} is a ribbon element so $S(t^{-2}) = t^{-2}$. □

Proposition 4.3. *The various important elements in a ribbon Hopf algebra can be written in terms of a half-ribbon elements t as follows:*

R	v	u	g	(11)
$(t^{-1} \otimes t^{-1})\Delta(t) = \Delta^{op}(t)(t^{-1} \otimes t^{-1})$	t^{-2}	$S(t^{-1})t^{-1}$	$S(t)t^{-1} = S(t^{-1})t$	

Proof. (R): The first formula is part of the definition of a half-ribbon element. The second formula follows by [Proposition 4.2\(i\)](#) since this implies that, for all $x \in \mathcal{H}$,

$$(t^{-1} \otimes t^{-1})\Delta(x)(t \otimes t) = \Delta^{op}(t^{-1}xt).$$

(v): This is part of the definition of a half-ribbon element.

(u): By definition, $u = \mu \circ (S \otimes 1)R_{21}$, where μ is multiplication. Thus:

$$u = \mu \circ (S \otimes 1)[(\Delta^{op}(t)(t^{-1} \otimes t^{-1})]_{21}) \tag{12}$$

$$= \mu[((S \otimes 1)(\Delta(t))(t^{-1} \otimes t^{-1})] \tag{13}$$

$$= \sum S(t^{-1})S(t_1)t_2t^{-1} \tag{14}$$

$$= S(t^{-1})[t \circ \varepsilon(t)]t^{-1} \tag{15}$$

$$= S(t^{-1})t^{-1}. \tag{16}$$

In (14) we have used Sweedler’s notation $\Delta(x) = \sum x_1 \otimes x_2$. The equality (15) follows from the antipode axiom of a Hopf algebra, and (16) follows by [Proposition 4.2\(ii\)](#).

(g): By definition, $g = v^{-1}u$. Recall that v is central. To get the second equation, apply [Proposition 4.2\(iii\)](#). □

Comment 4.4. For historical reasons the ribbon element v represents a negative twist. We use t for the positive half-twist, which explains the fact that $v = t^{-2}$.

4B. A topological category of ribbons with half-twists. There is a functor from a certain topological category to the category of representations of any ribbon Hopf algebra, which allows one to construct topological invariants. For a half-ribbon Hopf algebra, this functor can be extended to a larger topological category. In this section we define the two relevant categories, beginning with the large one.

Definition 4.5. The category $\mathcal{HRI\mathcal{B}}(S)$ of topological half-ribbons with labels in some set S is defined by:

- Objects in $\mathcal{HRI\mathcal{B}}(S)$ consist of a finite number of disjoint closed intervals on the real line, each labeled with an element of S , each with a choice of shading (shaded or unshaded), and each directed (up or down). These objects are considered up to isotopy of the real line. For example:

$$\begin{array}{cccccc} \uparrow & \uparrow & \downarrow & \downarrow & \downarrow & \uparrow \\ \boxed{A} & \boxed{B} & \boxed{B} & \boxed{C} & \boxed{A} & \boxed{D} \end{array} . \tag{17}$$

- A morphism between two objects $A, B \in \mathcal{HRI\mathcal{B}}(S)$ consist of a “tangle of orientable, directed ribbons” in $\mathbb{R}^2 \times I$, whose loose ends are exactly

$$(A, 0, 0) \cup (B, 0, 1) \subset \mathbb{R} \times \mathbb{R} \times I,$$

along with a choice of direction and shading of each ribbon. These must be such that the direction (up or down) of each interval in $A \cup B$ agrees with the direction of the ribbon whose end lies at that interval, and the shading (light or dark) of each interval agrees with the shading on the visible side of the ribbon near that interval. These are considered up to isotopy.

- Composition of two morphisms is given by stacking them on top of each other, and then shrinking the vertical axis by a factor of two. Note that we read our diagrams bottom to top. For example:

Definition 4.6. $RIBBON(S)$ is the full subcategory of $\mathcal{HRIB}(S)$ consisting of those objects all of whose intervals are unshaded.

The definition of $\mathcal{HRIB}(S)$ can easily be made rigorous, in the same way as is done for the subcategory $RIBBON(S)$; see for example [Chari and Pressley 1994]. In fact, as with $RIBBON(S)$, $\mathcal{HRIB}(S)$ is a rigid monoidal category, where tensor products and duals are shown below:

$$\begin{aligned}
 (\uparrow A \downarrow \uparrow B \downarrow) \otimes (\uparrow A \downarrow \uparrow C \downarrow \uparrow C \downarrow \uparrow B \downarrow) &= \uparrow A \downarrow \uparrow B \downarrow \uparrow A \downarrow \uparrow C \downarrow \uparrow C \downarrow \uparrow B \downarrow, \\
 (\uparrow A \downarrow \uparrow B \downarrow \uparrow C \downarrow \uparrow B \downarrow)^* &= \uparrow B \downarrow \uparrow C \downarrow \uparrow B \downarrow \uparrow A \downarrow.
 \end{aligned}$$

4C. A functor from $\mathcal{HRIB}(S)$ to $\mathcal{H}\text{-rep}$. We will now show that the category of representations of a half-ribbon Hopf algebra admits a natural functor from $\mathcal{HRIB}(\mathcal{H}\text{-rep})$. This theorem is an extension of the corresponding result relating ribbon Hopf algebras and $RIBBON(\mathcal{H}\text{-rep})$, so we begin by stating this known result.

Theorem 4.7 (see [Reshetikhin 1989], [Shum 1994], or [Chari and Pressley 1994, Theorem 5.3.2]). *Let (\mathcal{H}, R, C) be a ribbon Hopf algebra. There is a unique monoidal functor \mathcal{F}' from $RIBBON(\mathcal{H}\text{-rep})$ to $\mathcal{H}\text{-rep}$ satisfying the following conditions:*

(i) $\mathcal{F}'(\uparrow V \downarrow) = V$ and $\mathcal{F}'(\uparrow V \downarrow) = V^*$,

$$(ii) \quad \begin{aligned} \mathcal{F}' \left(\begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} \right) &= \text{ev}, & \mathcal{F}' \left(\begin{array}{c} \text{---} \downarrow \text{---} \\ \text{---} \uparrow \text{---} \end{array} \right) &= \text{coev}, \\ \mathcal{F}' \left(\begin{array}{c} \text{---} \downarrow \text{---} \\ \text{---} \uparrow \text{---} \end{array} \right) &= \tilde{\text{ev}}, & \mathcal{F}' \left(\begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} \right) &= \widetilde{\text{coev}}. \end{aligned}$$

$$(iii) \quad \mathcal{F}' \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) = C,$$

thought of as a morphism from V to V or from V^* to V^* , depending on the orientation.

$$(iv) \quad \mathcal{F}' \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) = \text{Flip} \circ R$$

as a morphism from the tensor product of the bottom two objects to the tensor product of the top two objects, regardless of labeling and orientation.

Comment 4.8. Warning: this functor is not a strict rigid functor. Notice that $\mathcal{F}'(\text{---} \downarrow \text{---}^*) = \mathcal{F}'(\text{---} \uparrow \text{---}) = V$, while $\mathcal{F}'(\text{---} \downarrow \text{---})^* = (V^*)^* = V^{**}$.

Our main result for this section is that \mathcal{F}' can be extended to a functor $\mathcal{F} : \mathcal{HRIB}(\mathcal{H}\text{-rep}) \rightarrow \mathcal{H}\text{-rep}$ as follows.

Definition 4.9. Let V be a representation of a half-ribbon Hopf algebra \mathcal{H} and for each $x \in \mathcal{H}$ let $\pi_V(x)$ be the element of $\text{End}(V)$ defined by x . Define a new representation V^{C_t} which is equal to V as a vector space, but with the action of $x \in \mathcal{H}$ defined by $\pi_{V^{C_t}}(x) = \pi_V(xt^{-1})$.

Definition 4.10. The *topological half-twist* is the morphism I_n in \mathcal{HRIB} which takes n ribbons of any shading and orientation, and twists them all together by 180 degrees:



Theorem 4.11. There is a unique monoidal functor \mathcal{F} from $\mathcal{HRIB}(\mathcal{H}\text{-rep})$ to $\mathcal{H}\text{-rep}$ satisfying:

- (i) $\mathcal{F}(\text{---} \uparrow \text{---}) = V$, $\mathcal{F}(\text{---} \downarrow \text{---}) = V^*$, $\mathcal{F}(\text{---} \uparrow \text{---}) = V^{C_t}$ and $\mathcal{F}(\text{---} \downarrow \text{---}) = (V^*)^{C_t}$.
- (ii) On unshaded caps and cups \mathcal{F} agrees with \mathcal{F}' .
- (iii) The topological half-twist on n ribbons is sent to $\text{rev} \circ \Delta^n(t)$ acting on \mathcal{F} of the bottom object, regardless of shading and orientation.

Furthermore, the restriction of \mathcal{F} to $\mathcal{RIBBON}(\mathcal{H}\text{-rep})$ agrees with \mathcal{F}' .

The proof of [Theorem 4.11](#) requires several lemmas, and will occupy the rest of this section.

Comment 4.12. Like \mathcal{F}' , \mathcal{F} is not a strict rigid functor. In particular, we have

$$\mathcal{F}(\text{HV}) = (V^*)^{C_t},$$

which is not equal to $(V^{C_t})^*$ (although these two representations are isomorphic). In fact, $(V^*)^{C_t}$ is equal to the right dual $*(V^{C_t})$. To see why, notice that, as vector spaces, both $(V^*)^{C_t}$ and $*(V^{C_t})$ are equal to $\text{Hom}_F(V, F)$. Each comes with a chosen action of \mathcal{H} , and, using the fact that C_t is a coalgebra antiautomorphism, one can show that these two actions are identical.

Comment 4.13. A good mnemonic for remembering what \mathcal{F} does to objects is to think of the ribbon as always being labeled with a representation on its light side, so that when you look at the dark side you see that label rotated 180 degrees around the y axis. For example, instead of $V_i^{C_t}$ you would see ${}_iV$. This works for remembering how duals interact with half twists since when you rotate V_i^* by 180 degrees about the y axis you see ${}_iV$.

Comment 4.14. There is more than one extension of \mathcal{F}' from $\mathcal{RIBBON}(\mathcal{H}\text{-rep})$ to the larger category $\mathcal{HRIB}(\mathcal{H}\text{-rep})$, but only one satisfies [Theorem 4.11\(iii\)](#).

We first prove the following different characterization of \mathcal{F} . This is simpler to prove, and is also more general than [Theorem 4.11](#). However, this result is less satisfying in other ways. For instance, [Proposition 4.15](#) does not imply that all half twists are sent to t acting on \mathcal{F} of the bottom object.

Proposition 4.15. *Let (\mathcal{H}, R, v) be a ribbon Hopf algebra, and let $t \in \mathcal{H}$ be an invertible element. There is a unique monoidal functor \mathcal{F} from $\mathcal{HRIB}(\mathcal{H}\text{-rep})$ to $\mathcal{H}\text{-rep}$ such that:*

- (i) \mathcal{F} agrees with \mathcal{F}' on the full subcategory $\mathcal{RIBBON}(\mathcal{H}\text{-rep})$.
- (ii) $\mathcal{F}(\text{HV}) = V^{C_t}$ and $\mathcal{F}(\text{VH}) = (V^*)^{C_t}$.
- (iii) $\mathcal{F}(\text{X}) = t$ as a morphism from V to V^{C_t} or from V^* to $(V^*)^{C_t}$ depending on orientation.
 $\mathcal{F}(\text{X}) = t^{-1}$ as a morphism from V^{C_t} to V or from $(V^*)^{C_t}$ to V^* depending on orientation (notice that this is a negative half-twist, hence the use of t^{-1}).

Proof. This is the simplest case of Reshetikhin and Turaev’s [1990] more general extension of \mathcal{F}' to ribbon graphs with coupons, where we think of the half-twists defined above as coupons on a single ribbon. So we just sketch the proof. Every morphism in $\mathcal{HRIB}(\mathcal{H}\text{-rep})$ is isotopic to ribbon in the following standard form: every shaded object at the bottom has a positive half-twist next to it, then in the

middle of the diagram there's a ribbon tangle, then at the top of the diagram every shaded object has a negative half-twist next to it. Furthermore two such diagrams are isotopic if and only if the ribbon tangles in the middle of their standard diagrams are isotopic. Hence there is at most one possible functor \mathcal{F} . That \mathcal{F} is in fact a functor follows from the fact that t and t^{-1} are inverses of each other, so that composition of diagrams is just given by composing their middle parts. \square

The following lemma shows how \mathcal{F} acts on those elementary morphisms not already specified in the statement of [Proposition 4.15](#).

Lemma 4.16. *Let (\mathcal{H}, t) be a half-ribbon Hopf algebra, and define \mathcal{F} as in [Proposition 4.15](#).*

- (i) \mathcal{F} sends any positive half-twist to t acting on \mathcal{F} of the bottom object regardless of orientation or shading. \mathcal{F} sends any negative half-twist to t^{-1} regardless of orientation or shading.
- (ii) \mathcal{F} takes any simple crossing to $\text{Flip} \circ R$ acting on \mathcal{F} of the bottom object, regardless of orientations and shadings.
- (iii) \mathcal{F} sends shaded caps and cups to the following maps:

$$\mathcal{F} \left(\begin{array}{c} \text{shaded cap} \\ \uparrow \text{ on left, } \downarrow \text{ on right} \end{array} \right) = \text{ev}_{*(V^{C_t})} : v \otimes f \mapsto f(v), \tag{19}$$

$$\mathcal{F} \left(\begin{array}{c} \text{shaded cup} \\ \downarrow \text{ on left, } \uparrow \text{ on right} \end{array} \right) = \tilde{\text{ev}}_{*(V^{C_t})} : f \otimes v \mapsto f(\pi_V(g)v), \tag{20}$$

$$\mathcal{F} \left(\begin{array}{c} \text{unshaded cap} \\ \uparrow \text{ on left, } \downarrow \text{ on right} \end{array} \right) = \widetilde{\text{coev}}_{*(V^{C_t})} : 1 \mapsto \sum_i \pi_V(g)^{-1} v_i \otimes v^i, \tag{21}$$

$$\mathcal{F} \left(\begin{array}{c} \text{unshaded cup} \\ \downarrow \text{ on left, } \uparrow \text{ on right} \end{array} \right) = \text{coev}_{*(V^{C_t})} : 1 \mapsto \sum_i v^i \otimes v_i. \tag{22}$$


For the explicit formulas, we have used the fact that V^{C_t} and $(V^)^{C_t}$ are identical to V and V^* , respectively, as vector spaces, and chosen dual basis v_i and v^i for V and V^* .*

Comment 4.17. In order to apply the maps $\text{ev}_{*(V^{C_t})}$, etc. above, one must use the canonical isomorphism $(*(V^{C_t}))^* \cong V^{C_t}$, and the equality $(V^*)^{C_t} = *(V^{C_t})$ from [Comment 4.12](#). It is a straightforward exercise to show that these maps are given in coordinates by the above formulas.

Proof of Lemma 4.16. We prove each of these formulas by a direct calculation.

For the half-twists and ribbon full-twists we only use that $t^2 = v^{-1}$. For example, the positive half-twist going from unshaded to shaded is a composition of the negative half-twist with the full twist. So this result follows from $t^{-1}v^{-1} = t$.

The formulas for \mathcal{F} applied to a crossing with one or both of the ribbons shaded all follow from the naturality of the braiding. For example, to find \mathcal{F} of a crossing where one strand is shaded, consider the following isotopy:



Thus \mathcal{F} of the left side must be equal to $(t \otimes 1) \circ \text{Flip} \circ R \circ (1 \otimes t^{-1})$. By the naturality of the braiding,

$$(t \otimes 1) \circ \text{Flip} \circ R \circ (1 \otimes t^{-1}) = \text{Flip} \circ R \circ (1 \otimes t) \circ (1 \otimes t^{-1}) = \text{Flip} \circ R. \tag{23}$$

This holds independently of the orientations of the ribbons. A similar argument gives the same formula for crossings where the other ribbon is shaded, or where both are shaded.

It remains to compute the formula for shaded cups and caps. These equations use $g = S(t)t^{-1} = S(t^{-1})t$. We explicitly show two of the four cases. The other two are similar.

By the definition of \mathcal{F} , one has

$$\mathcal{F} \left(\begin{array}{c} \text{shaded cup} \\ \uparrow \quad \downarrow \\ \text{v} \end{array} \right) (v \otimes f) = \mathcal{F} \left(\begin{array}{c} \text{unshaded cup} \\ \uparrow \quad \downarrow \\ \text{v} \end{array} \right) (v \otimes f) \tag{24}$$

$$= \tilde{e}v(tv \otimes t^{-1}f) \tag{25}$$

$$= t^{-1}f(gtv) \tag{26}$$

$$= f(S(t^{-1})gtv) \tag{27}$$

$$= f(v). \tag{28}$$

Here (24) is an isotopy, (25) follows from functoriality and our computation of \mathcal{F} on all half-twists, (26) is the definition of $\tilde{e}v$, (27) is the definition of the action on the dual space, and (28) uses the formula $g = S(t)t^{-1}$.

Now for a cup. Notice that, if $e^i \in V^*$ and $e_j \in V$ are dual bases and x is an invertible element in \mathcal{H} , then $v^i = xe^i$ and $v_i = S(x)^{-1}e_i$ are also dual bases. By the definition of \mathcal{F} , we have:

$$\mathcal{F} \left(\text{cup diagram with } v^i \text{ and } v_i \text{ labels} \right) (1) = \mathcal{F} \left(\text{cup diagram with } e^i \text{ and } e_j \text{ labels} \right) (1) \tag{29}$$

$$= \sum t e_i \otimes t^{-1} e^i \tag{30}$$

$$= \sum t S(t^{-1}) v_i \otimes v^i \tag{31}$$


$$= \sum g^{-1} v_i \otimes v^i. \tag{32}$$

Here (29) is an isotopy, (30) follows from functoriality and our computation of \mathcal{F} on all half-twists, (31) is the change of dual bases described above with $x = t$, and (32) uses the formula $g = S(t)t^{-1}$. \square

Note that above we used the formulas relating t to g (which was derived using the formula relating R and t), but we did not use the formula for R directly. That formula is used in the following proposition which computes how \mathcal{F} acts on the half-twist applied to many strands.

Proposition 4.18. *Suppose that (\mathcal{H}, t) is a half-ribbon Hopf algebra, let \mathcal{F} be the unique functor guaranteed by Proposition 4.15. Then, $\mathcal{F}(I_n) = \text{rev} \circ \Delta^n(t)$.*

Proof. We proceed by induction on n , the case $n = 1$ being trivial. Consider the isotopy



Thus, by the definition of \mathcal{F} ,

$$\mathcal{F}(I_n) = (t \otimes t \otimes \cdots \otimes t) \mathcal{F}(T_{w_0}^{(n)}), \tag{33}$$

where $T_{w_0}^{(n)}$ is the braid group element corresponding to the longest element of S_n . Let $\sigma^{(n)}$ be the braid group element



Let $s^{(n)}$ be the image of $\sigma^{(n)}$ in S_n , and let $\text{rev}^{(n)}$ denote the longest element of S_n . Clearly $T_{w_0}^{(n)} = (T_{w_0}^{(n-1)} \otimes 1) \circ \sigma^{(n)}$. Hence

$$\mathcal{F}(I_n) = (t \otimes t \otimes \cdots \otimes t) \circ \mathcal{F}(T_{w_0}^{(n)}) \tag{34}$$

$$= (t \otimes t \otimes \cdots \otimes t) \circ (\mathcal{F}(T_{w_0}^{(n-1)}) \otimes 1) \circ \mathcal{F}(\sigma^{(n)}) \tag{35}$$

$$= (\mathcal{F}(I_{n-1}) \otimes t) \circ \mathcal{F}(\sigma^{(n)}) \tag{36}$$

$$= (\text{rev}^{(n-1)} \Delta^{n-1}(t) \otimes t) \circ s^{(n)} \circ (1 \otimes \Delta^{n-1})(R) \tag{37}$$

$$= (\text{rev}^{(n-1)} \Delta^{n-1}(t) \otimes t) \circ s^{(n)} \circ (1 \otimes \Delta^{n-1})((t^{-1} \otimes t^{-1})\Delta(t)) \tag{38}$$

$$= \text{rev}^{(n)} \Delta^n(t). \tag{39}$$

Here (36) follows from (33) in the case $n - 1$, (37) follows from the inductive assumption and the quasitriangularity of R , (38) holds because t is a half-ribbon element, and (39) is a straightforward calculation.

This proof works regardless of the orientations and shadings of the ribbons. \square

Proof of Theorem 4.11. Using the isotopy from (1), the crossing is a composition of the two strand topological half-twist and two copies of the inverse of the one strand topological half-twist. Thus the conditions in the statement uniquely determine \mathcal{F} on all the elementary morphisms listed in the condition of Proposition 4.15. So Proposition 4.15 shows that there is a unique candidate for \mathcal{F} . Proposition 4.18 shows that this \mathcal{F} satisfies the remaining condition of the theorem. \square

5. Ribbon and half-ribbon elements for $U_q(\mathfrak{g})$

We show that $U_q(\mathfrak{g})$ is always a topological half-ribbon Hopf algebra. That is, there exists an element X in a completion of $U_q(\mathfrak{g})$ such that $(X^{-1} \otimes X^{-1})\Delta(X)$ is the standard R -matrix, and X^{-2} is a ribbon element. Interestingly, X^{-2} is not the standard ribbon element C . We also classify the different ribbon elements for $U_q(\mathfrak{g})$, and discuss how one might decide which of these arise from half-ribbon elements. In particular, we show that the standard ribbon element for $U_q(\mathfrak{sl}_2)$ does not arise from a half-ribbon element. We then explain some consequences of varying the ribbon element. It turns out the ribbon element X^{-2} is in some ways particularly nice (see Lemma 5.7 and Section 5E).

5A. A half-ribbon Hopf algebra structure on $U_q(\mathfrak{g})$. We first need the following result relating different ribbon elements for the same quasitriangular Hopf algebra.

Lemma 5.1. *Let (\mathcal{H}, R, v) be a ribbon Hopf algebra, and $s \in \mathcal{H}$ be a central grouplike element that squares to 1. Then (\mathcal{H}, R, vs) is also a ribbon Hopf algebra. Furthermore all ribbon elements for (\mathcal{H}, R) are of the form vs for some such s .*

Proof. This is a straightforward application of the definition of a ribbon element [Barrett and Westbury 1999, Remark 3.4]. \square

Theorem 5.2. *Let X be the element from Theorem 3.11. Then $(U_q(\mathfrak{g}), X)$ is a topological half-ribbon Hopf algebra. Furthermore $(X^{-1} \otimes X^{-1})\Delta(X)$ is the standard R -matrix.*

Proof. By Theorem 3.11, $(U_q(\mathfrak{g}), R)$ is a quasitriangular Hopf algebra, where $R = (X^{-1} \otimes X^{-1})\Delta(X)$ is the standard R matrix. Thus it suffices to show that X^{-2} is a ribbon element for this quasitriangular Hopf algebra. We already know that $(U_q(\mathfrak{g}), R, C)$ is a ribbon Hopf algebra. Thus by Lemma 5.1 it suffices to show that (i) X^2C is central; (ii) X^2C is grouplike; and (iii) $(X^2C)^2 = 1$.

(i) By Lemma 3.10(iii) and Definition 3.15, X^2C acts on V_λ as multiplication by the scalar $(-1)^{\langle 2\lambda, \rho^\vee \rangle}$. Thus it is clearly central.

(ii) Any highest weight ν for $V_\lambda \otimes V_\mu$ has weight $\lambda + \mu - \gamma$ for some γ in the root lattice. Since $\langle 2\gamma, \rho^\vee \rangle$ is an even integer for any γ in the root lattice, it follows that X^2C is grouplike.

(iii) $(X^2C)^2$ acts on V_λ by $(-1)^{2 \cdot \langle 2\lambda, \rho^\vee \rangle} = 1$. □

5B. The Frobenius–Schur indicator. We must now discuss a tool for comparing ribbon elements. The following definition of the Frobenius–Schur indicator for a pivotal category was given by [Ng and Schauenburg 2007] (see also [Linchenko and Montgomery 2000] for the case of Hopf algebras and [Fuchs et al. 1999] for C^* sovereign categories).

Recall that a pivotal structure is a natural collection of maps $p_V : V \rightarrow V^{**}$ which defines a natural isomorphism $\text{Id} \rightarrow **$ of monoidal functors [Barrett and Westbury 1999]. The category of representations of a ribbon Hopf algebra is always pivotal with the pivotal structure being given by the grouplike element $g = v^{-1}u$ as follows: fix $v \in V$. Then $p_V(v)$ is the element of V^{**} defined by, for all $f \in V^*$, $p_V(v)(f) = f(gv)$ (see for example [Chari and Pressley 1994]).

Definition 5.3. Given an F -linear category with a chosen pivotal structure p , define the Frobenius–Schur indicator of an absolutely simple object V as follows: if $V \not\cong V^*$, then $FS_p(V) = 0$. Otherwise, choose an isomorphism $f : V \rightarrow V^*$. By Schur’s lemma there exists some constant, which we define to be $FS_p(V)$, such that $f = FS_p(V)f^* \circ p$.

Comment 5.4. As shown in [Ng and Schauenburg 2007], $FS_p(V)$ does not depend on the choice of f , and we have $FS_p(V) = \pm 1$ (or 0).

Comment 5.5. Notice that the Frobenius–Schur indicator depends on the pivotal structure, which in turn depends on the choice of ribbon element. As described in [Ng and Schauenburg 2007], we use $FS_v(V)$ to denote the Frobenius–Schur indicator for V , calculated with the ribbon element v .

Comment 5.6. The reason for the term Frobenius–Schur indicator is that in the case of groups (and indeed involutory Hopf algebras [Linchenko and Montgomery 2000]) with the trivial pivotal structure, it agrees with the usual Frobenius–Schur indicator, which is 1 when there is an invariant orthogonal form on $V \otimes V$, -1 when there is an invariant symplectic form on $V \otimes V$, and 0 otherwise.

Lemma 5.7. *Let V_λ be an irreducible representation of $U_q(\mathfrak{g})$ and assume V_λ is self-dual. Then $FS_{X^{-2}}(V_\lambda) = 1$.*

Proof. Let $f : V_\lambda \rightarrow V_\lambda^*$ be an isomorphism. It suffices to show that $f = f^* \circ p_{V_\lambda}$. Since both f and $f^* \circ p_{V_\lambda}$ are isomorphisms, it suffices to show that their actions on v_λ^{low} agree. But $f(v_\lambda^{\text{low}})$ is in the lowest weight space of V_λ^* , which is one-dimensional and pairs nondegenerately with v_λ . Thus, it in fact suffices to show that

$$f(v_\lambda^{\text{low}})(v_\lambda) = f^* \circ p_{V_\lambda}(v_\lambda^{\text{low}})(v_\lambda). \quad (40)$$

Using the formula for g given in Proposition 4.3,

$$S(X) = Xg^{-1} = gX. \quad (41)$$

Let $k = (-1)^{\langle 2\lambda, \rho^\vee \rangle} q^{(\lambda, \lambda)/2 + (\lambda, \rho)}$. By Lemma 3.10(ii), $X(v_\lambda) = k(v_\lambda^{\text{low}})$. Thus the left side of (40) can be simplified as follows

$$f(v_\lambda^{\text{low}})(v_\lambda) = f(k^{-1}X(v_\lambda))(v_\lambda) \quad (42)$$

$$= k^{-1}(Xf(v_\lambda))(v_\lambda) \quad (43)$$

$$= k^{-1}f(v_\lambda)(S(X)v_\lambda) \quad (44)$$

$$= k^{-1}f(v_\lambda)(Xg^{-1}v_\lambda) \quad (45)$$

$$= k^{-1}f(v_\lambda)(gXv_\lambda) \quad (46)$$

$$= k^{-1}f(v_\lambda)(kgv_\lambda^{\text{low}}) \quad (47)$$

$$= f(v_\lambda)(gv_\lambda^{\text{low}}). \quad (48)$$

Here (43) follows because f is an isomorphism, (45) and (46) follow by (41) and (47) by the above formula for the action of X on v_λ .

Now consider the right side of (40). By definition, $p_{V_\lambda}(v_\lambda^{\text{low}})$ is the element of V_λ^{**} which takes $\phi \in V_\lambda^*$ to $\phi(gv_\lambda^{\text{low}})$. Thus

$$f^* \circ p_{V_\lambda}(v_\lambda^{\text{low}})(v_\lambda) = f(v_\lambda)(gv_\lambda^{\text{low}}). \quad (49)$$

We have shown that the two sides of (40) agree, so the result follows. \square

Turaev [1994] calls the property that Lemma 5.7 holds unimodality and requires it when defining geometric 3j and 6j symbols.

Lemma 5.8. $FS_C(V)$ is the usual Frobenius–Schur indicator of the corresponding classical representation.

Proof. Since the Frobenius–Schur indicator is a discrete invariant, it doesn't change under continuous deformation. \square

5C. Which ribbon elements for $U_q(\mathfrak{g})$ arise from half-ribbon elements? Here we classify ribbon elements for $U_q(\mathfrak{g})$. We then prove that the standard ribbon element for $U_q(\mathfrak{sl}_2)$ is not equal to t^{-2} for any half-ribbon element t .

Definition 5.9. Suppose that ϕ is a character of P/Q , where P is the weight lattice of \mathfrak{g} , and Q is the root lattice. Define $s(\phi) \in \widetilde{U_q(\mathfrak{g})}$ to act on V_λ as multiplication by the scalar $\phi(\lambda)$.

Theorem 5.10. The central grouplike elements of $\widetilde{U_q(\mathfrak{g})}$ are precisely the $s(\phi)$.

Proof. Suppose that s is a central grouplike element of $\widetilde{U_q(\mathfrak{g})}$. Since it is central, s acts by a scalar on any irreducible representation. Furthermore, since it is grouplike, s acts by the same scalar on any two irreducible representations that appear in the same arbitrary tensor product of irreducible representations. Thus such an s gives a function of P/\sim , where \sim is the equivalence relation on weights generated by $\lambda \sim \mu$ if V_λ and V_μ appear in the same tensor product of irreducible representations. It suffices to show that $\lambda \sim \mu$ if and only if $\lambda + Q = \mu + Q \in P/Q$. One can easily see that \sim respects the additive structure of the weight lattice, so it is enough to prove that V_λ appears in some tensor product also containing the trivial representation V_0 precisely when λ is in the root lattice.

Since the Littlewood–Richardson coefficients are the same in the quantum and classical cases, this reduces to the same question in the classical case. So let G be the simply connected Lie group attached to $U(\mathfrak{g})$, and let $Z(G)$ be the center of G . The central character map gives a pairing between $Z(G)$ and the weight lattice mod the root lattice. Since any tensor product of irreducible representations has a well-defined central character, we see that V_λ is equivalent to V_0 only if λ is in the root lattice.

To see the other direction, let V be a faithful representation of the compact adjoint form K of G .

Claim: For any λ in the root lattice, there exists N so that, for all $n > N$, V_λ occurs as a subrepresentation of $V^{\otimes n}$.

Proof of claim: since λ is in the root lattice, V_λ descends to a representation of K . On this compact form we can use character theory. Because K has no center, $|\chi_V(g)| < \dim V$ for any $g \neq 1$. Let

$$k_n = \left| \int_K \chi_V^n d\mu \right|, \quad (50)$$

where $d\mu$ is the normalized Haar measure. Since K is compact and $|\chi_V(g)| < \chi_V(1)$ for all $g \neq 1$, we see that

$$\lim_{n \rightarrow \infty} \frac{\langle \chi_{V^{\otimes n}}, \chi_{V_\lambda} \rangle}{k_n} = \lim_{n \rightarrow \infty} \frac{1}{k_n} \int_K \chi_V(g)^n \chi_{V_\lambda}(g^{-1}) d\mu = \chi_{V_\lambda}(1) = \dim V_\lambda. \quad (51)$$

In particular, for $n \geq N$ we see that $\langle \chi_{V^{\otimes n}}, \chi_{V_\lambda} \rangle$ is nonzero, so V_λ occurs in $V^{\otimes n}$.

Applying the above fact to 0 and an arbitrary root vector λ , we see that for a sufficiently high power $N' = \max(N_0, N_\lambda)$ the tensor power $V^{\otimes N'}$ contains both V_0 and V_λ , and hence $0 \sim \lambda$. □

Theorem 5.11. *Ribbon elements of $U_q(\widetilde{\mathfrak{g}})$ are exactly elements of the form*

$$s(\phi)X^{-2},$$

where ϕ is a character of the weight lattice mod the root lattice of order ≤ 2 .

Proof. This follows from [Lemma 5.1](#) and [Theorem 5.10](#). □

We have the following relationship between ribbon elements and the Frobenius–Schur indicator.

Proposition 5.12. *If V_λ is self dual, then $FS_{s(\phi)X^{-2}}(V_\lambda) = \phi(\lambda)$.*

Proof. The only part of the definition of the Frobenius–Schur indicator that changes when you change the ribbon element from X^{-2} to $s(\phi)X^{-2}$ is p_{V_λ} , and this is multiplied by $\phi(\lambda)$. Hence the proposition follows from [Lemma 5.7](#). □

We wish to understand which of these ribbon element extend to a half-ribbon elements on $U_q(\widetilde{\mathfrak{g}})$. As we shall see, the ratio of two half-ribbon elements is grouplike, so to classify all of them we need to understand the grouplike elements in $U_q(\widetilde{\mathfrak{g}})$. In general this seems to pose some technical challenges, but we can do the case of $U_q(\mathfrak{sl}_2)$.

Proposition 5.13. *Let (\mathcal{H}, R) be a quasitriangular Hopf algebra, and assume that t_1 and t_2 are two half-ribbon elements. Then $t_1 t_2^{-1}$ is grouplike. (Note that we do not assume here that $t_1^{-2} = t_2^{-2}$.)*

Proof. This follows from the equality $(t_1^{-1} \otimes t_1^{-1})\Delta(t_1) = (t_2^{-1} \otimes t_2^{-1})\Delta(t_2)$. □

Lemma 5.14. *Let V be the standard representation of $U_q(\mathfrak{sl}_2)$, with basis $\{v_+, v_-\}$ such that $E(v_-) = v_+$, $Kv_+ = qv_+$, and $Kv_- = q^{-1}v_-$. Then $V \otimes V$ contains a copy of the trivial representation V_0 . Furthermore, this is spanned by the element*

$$v_+ \otimes v_- - q^{-1}v_- \otimes v_+.$$

Proof. It suffices to check that $\Delta(E)(v_+ \otimes v_- - q^{-1}v_- \otimes v_+) = 0$. This follows from the definition of Δ . □

Definition 5.15. Let $a \in \mathbb{C}(q)$. Define K_a to be the element of $\widetilde{U_q(\mathfrak{sl}_2)}$ that acts on any weight vector $v \in V_\lambda$ by $K_a(v) = a^{\text{wt}(v)}$. Note that K_a is grouplike.

Lemma 5.16. All grouplike elements in $\widetilde{U_q(\mathfrak{sl}_2)}$ are of the form K_a , for $a \in \mathbb{C}(q)$.

Proof. A grouplike element is determined by its action on the fundamental representation V . Suppose that σ is grouplike. Define constants a, b, c and d by $\sigma v_+ = av_+ + bv_-$ and $\sigma v_- = cv_+ + dv_-$. Since σ is grouplike it must act trivially on the trivial subrepresentation of $V \otimes V$. Thus by Lemma 5.14,

$$v_+ \otimes v_- - q^{-1}v_- \otimes v_+ = \sigma \otimes \sigma(v_+ \otimes v_- - q^{-1}v_- \otimes v_+) \tag{52}$$

$$= ac(1 - q^{-1})v_+ \otimes v_+ + (ad - q^{-1}bc)v_+ \otimes v_- \tag{53}$$

$$- q^{-1}(ad - qbc)v_- \otimes v_+ + (1 - q^{-1})bdv_- \otimes v_-. \tag{54}$$

Comparing coefficients we see that $ac = bd = bc = 0$ and $ad = 1$. Hence σ acts on V in exactly the same way that K_a does. Since they are both grouplike and V is a tensor generator for the category of representations of $U_q(\mathfrak{sl}_2)$, we see that $\sigma = K_a$. □

Theorem 5.17. There is no topological half-ribbon element t for $U_q(\mathfrak{sl}_2)$ such that t^{-2} is the usual ribbon element C .

Proof. By Theorem 3.16, X is a half-ribbon element for $(U_q(\mathfrak{sl}_2), R)$. By Proposition 5.13, any other half-ribbon element is of the form $X\alpha$ for some grouplike element α . By Lemma 5.16, $\alpha = K_a$ for some $a \in \mathbb{C}(q)$. Since X sends the λ weight space to the $-\lambda$ weight, it follows that $K_a X K_a v = Xv$ for any weight space v . Therefore, we have that $(XK_a)^{-2} = X^{-2}$. Hence every half-ribbon element t for $U_q(\mathfrak{sl}_2)$ has $t^{-2} = X^{-2}$. But X^{-2} acts as $-C$ on the standard representation, so is not equal to C . □

5D. A summary of ribbon and half-ribbon elements for $U_q(\mathfrak{sl}_2)$, $U_q(\mathfrak{sl}_3)$ and $U_q(\mathfrak{sl}_4)$. We now describe the possible ribbon elements of these three quantum groups, and describe which are the squares of half-ribbon elements. Notice that the three cases all behave differently.

- $U_q(\mathfrak{sl}_2)$. There are two ribbon elements C , and X^{-2} . Only X^{-2} can be realized using a half-ribbon element (see Theorem 5.17).
- $U_q(\mathfrak{sl}_3)$. In this case $P/Q \cong \mathbb{Z}/3$. Thus, there are no central grouplikes of order two, so there can be only one ribbon element. This unique ribbon element can be realized using the half-ribbon element X .
- $U_q(\mathfrak{sl}_4)$. In this case $P/Q \cong \mathbb{Z}/4$. This has two characters of order ≤ 2 , so there are two ribbon elements. In this case both come from half-ribbon elements. A straightforward calculation on the standard representation of $U_q(\mathfrak{sl}_4)$ shows that X^{-2} is not the standard ribbon element. To see the other half-ribbon element,

consider the central element $s \in \widetilde{U_q(\mathfrak{sl}_4)}$ which acts on V_λ as multiplication by $i^{4(\lambda, \lambda)}$, where i is the complex fourth root of unity. Recall that $(\lambda, \alpha) \in \mathbb{Z}$ for any root α , which implies that s is grouplike. Clearly we also have $s^4 = 1$, and from these two facts it follows that $X' := sX$ is a half-ribbon element for $(U_q(\mathfrak{sl}_4), R)$. One can check that s acts as multiplication by $-i$ on the standard representation, so $X'^{-2} = s^2 X^{-2} \neq X^{-2}$. Since X^{-2} is the nonstandard ribbon element, X'^{-2} must be the standard ribbon element.

5E. Varying the ribbon element and diagrammatic categories. Often you'll find a result in the literature saying that a certain diagram category "is the same as" as certain category coming from quantum groups (for example, [Ohtsuki 2002, Appendix H] and [Kuperberg 1996, page 11]). However, in the details of this claim there's an annoying sign difference between the diagram category and the quantum group category. The reason for this is that, although the two categories match up as braided tensor categories, they are different as pivotal categories. This can be fixed by changing the ribbon element on the quantum group side, thus changing the pivotal structure.

We illustrate this here by considering the case of $U_q(\mathfrak{sl}_2)$ -rep and the Temperley–Lieb category (see Section 3E). In the correspondence discussed in [Ohtsuki 2002, Appendix H], the standard representation of $U_q(\mathfrak{sl}_2)$ corresponds to the elementary object (i.e. a single \bullet) in the Temperley–Lieb category. We now use the Frobenius–Schur indicator to show that, for any such statement to hold on the level of pivotal categories, one must use the ribbon element X^{-2} .

Proposition 5.18. *Let \bullet denote a single point in Temperley–Lieb (with the trivial pivotal structure discussed in Section 3E). Then $FS(\bullet) = 1$.*

Proof. The single strand is an isomorphism between \bullet and $\bullet^* = \bullet$. The dual of the single strand (given by rotating 180 degrees about the z axis) is again the single strand. □

Proposition 5.19. *Let V be the standard representation of $U_q(\mathfrak{sl}_2)$. Then*

$$FS_C(V) = -1 \quad \text{and} \quad FS_{X^{-2}}(V) = 1.$$

Proof. This is a straightforward calculation using the definitions of C (Definition 3.15) and Lemma 3.10(iii). It also follows from the discussion in Section 5C. □

If \bullet in Temperley–Lieb is going to correspond to the standard representation, the Frobenius–Schur indicators should agree. Clearly this can only happen if we use the ribbon element X^{-2} .

Comment 5.20. A similar difficulty arises in relating the type C quantum group knot invariants with (a specialization of) the Kauffman polynomial. Once again, one can fix the problem by switching to the ribbon element X^{-2} .

Comment 5.21. This difficulty can also be addressed by changing the diagrammatic category. The Temperley–Lieb category has another pivotal structure p such that $p_\bullet = -\text{id}_\bullet$. With this pivotal structure $FS(\bullet) = -1$. This other pivotal structure is given diagrammatically by the disoriented Temperley–Lieb category of [Clark, Morrison and Walker 2009].

5F. The effect of varying the ribbon element on knot invariants. Suppose that (\mathcal{H}, R, v) is a ribbon Hopf algebra over a base field F . Fix a simple \mathcal{H} -module V . The functor from Theorem 4.7 sends a link with every component labeled with V to an element of F , denoted by $\mathcal{F}^V(L)$, which is an invariant of framed oriented links. Note that $\mathcal{F}^V(L)$ actually depends on the choice of ribbon element v , so when we need to be clear about which ribbon element we are using we will denote it by $\mathcal{F}_v^V(L)$. Since V is simple the ribbon element acts by a scalar $\theta_v(V)$ on V . It follows as in, for example, [Ohtsuki 2002, Section 3.3] that $\theta_v(V)^{w(L)}\mathcal{F}_v^V(L)$ is an invariant of oriented but unframed links (where $w(L)$ is the writhe of the link diagram, as described in, for example, [Ohtsuki 2002, page 11] and the current Wikipedia entry for “Writhe”).

Proposition 5.22. *Suppose that \mathcal{H} is a quasitriangular Hopf algebra with*

$$\text{End}(\mathbf{1}) = F,$$

and that V is a simple \mathcal{H} -module. Suppose that v_1 and v_2 are two different ribbon elements for \mathcal{H} . Let $\theta_{v_1}(V)$ and $\theta_{v_2}(V)$ be the scalars by which v_1 and v_2 act on V . Let $\mathcal{F}_{v_1}^V$ and $\mathcal{F}_{v_2}^V$ be the functors attached to these two ribbon Hopf algebras. Then for any link L ,

$$\theta_{v_1}^{w(L)}\mathcal{F}_{v_1}^V(L) = \left(\frac{FS_{v_1}(V)}{FS_{v_2}(V)}\right)^{\#L} \theta_{v_2}^{w(L)}\mathcal{F}_{v_2}^V(L),$$

where $\#L$ is the number of components of L .

Proof. By Theorem 5.11, $\alpha := v_1/v_2$ is central grouplike element of order 2. Let α_V be the scalar by which α acts on V . By Proposition 5.12,

$$\alpha_V = FS_{v_1}(V)/FS_{v_2}(V).$$

Let $g_i = v_i^{-1}u$ where u is the Drinfeld element. It follows immediately from definitions that $\widetilde{e}v_2 = \alpha_V^{-1}\widetilde{e}v_1$ and $\widetilde{\text{coe}}v_2 = \alpha_V^{-1}\widetilde{\text{coe}}v_1$.

The only elementary morphisms for which $\mathcal{F}_{v_1}^V$ and $\mathcal{F}_{v_2}^V$ disagree are left-going cups (corresponding to $\widetilde{\text{coe}}v$), left-going caps (corresponding to $\widetilde{e}v$), and full-twists. Thus,

$$\theta_{v_1}^{w(L)}\mathcal{F}_{v_1}^V(L) = (\alpha_V)^{-N_L}\theta_{v_2}^{w(L)}\mathcal{F}_{v_2}^V(L), \tag{55}$$

where

$$N_L := \#\{\text{left going caps and cups}\} + w(L). \tag{56}$$

It is easy to see that $N_L \pmod{2}$ is an invariant of oriented links (for example, check all the Turaev moves [Ohtsuki 2002, Section 3.2]). Furthermore, $N_L \pmod{2}$ doesn't change when you replace a positive crossing by a negative crossing. Since every link can be unknotted by replacing positive crossings with negative crossings, we see that $N_L \pmod{2}$ depends only on the number of components of L . By looking at the k -component unlink we see that $N_L \equiv \#L \pmod{2}$. \square

6. Questions

Question 1. Which ribbon Hopf algebras can be endowed with a half-ribbon element?

There are several aspects to this question. One could look for examples of half-ribbon Hopf algebras other than $U_q(\mathfrak{g})$. One could also try to find nonexamples, in the sense of finding ribbon Hopf algebras that do not contain the required element t . Such examples exist (at least for topological ribbon Hopf algebras). For instance we showed in Section 5C that $U_q(\mathfrak{sl}_2)$, with the standard ribbon element, cannot be made into a half-ribbon Hopf algebra. However, it can be modified by multiplying the ribbon element by a central grouplike element, and then it does have the required t . One could also ask if there are examples of ribbon Hopf algebras that cannot be made into half-ribbon Hopf algebras, even allowing this sort of modification. More ambitiously, one could look for a general method of determining when a ribbon Hopf algebra \mathcal{H} contains an element t such that (\mathcal{H}, t) is a half-twist Hopf algebra.

In the current work we have mainly considered $U_q(\mathfrak{g})$, which is infinite-dimensional, and only has a topological half-ribbon element in the sense that t only belongs to a completion of $U_q(\mathfrak{g})$. We feel it would be interesting to look at the case of finite-dimensional Hopf algebras as well.

Question 2. Fix a Hopf algebra \mathcal{H} . Is there a natural set of conditions one can impose on an element $t \in \mathcal{H}$ which guarantees that (\mathcal{H}, t) is a half-ribbon Hopf algebra?

We would like to be able to start with a Hopf algebra, which is not a priori quasitriangular, and endow it with a ribbon (and half-ribbon) structure by finding a certain $t \in \mathcal{H}$. One can of course write down the conditions t needs to satisfy by insisting that $(t^{-1} \otimes t^{-1})\Delta(t)$ is a quasitriangular structure, and t^{-2} is a ribbon element. However, these are very difficult to deal with, so the real question is to find nicer conditions on t . This would give a new method of constructing quasitriangular Hopf algebras.

Question 3. What happens if you weaken the conditions on t ?

Checking that $(t^{-1} \otimes t^{-1})\Delta t$ is a quasitriangular structure is difficult, but for some applications one can weaken this condition by insisting only that

(i) for any representation V and W of \mathcal{H} , the map

$$\text{Flip} \circ (t^{-1} \otimes t^{-1})\Delta(t) : V \otimes W \rightarrow W \otimes V \quad (57)$$

is an isomorphism; and

(ii) t^{-2} acts as a ribbon element, in the sense that

$$t^{-2} \circ \text{mult} \circ (S \otimes 1) \left((t^{-1} \otimes t^{-1})\Delta(t) \right) \quad (58)$$

is grouplike, and so can be used to develop a theory of quantum trace.

These conditions seem easier to check, and a Hopf algebra \mathcal{H} with such a t already has some nice structure. This could be used, for instance, in studying coboundary categories [Drinfel'd 1989].

Acknowledgments

We thank Joel Kamnitzer, Scott Morrison, and Nicolai Reshetikhin for helpful conversations. We also thank the anonymous referee for very thorough and helpful comments.

References

- [Barrett and Westbury 1999] J. W. Barrett and B. W. Westbury, “Spherical categories”, *Adv. Math.* **143**:2 (1999), 357–375. [MR 2000c:18007](#) [Zbl 0930.18004](#)
- [Chari and Pressley 1994] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge University Press, New York, 1994. [MR 95j:17010](#) [Zbl 0839.17009](#)
- [Clark, Morrison and Walker 2009] D. Clark, S. Morrison, and K. Walker, “Fixing the functoriality of Khovanov homology”, *Geom. Topol.* **13**:3 (2009), 1499–1582. [MR 2496052](#) [Zbl 1169.57012](#)
- [Drinfel'd 1989] V. G. Drinfel'd, “Quasi-Hopf algebras”, *Algebra i Analiz* **1**:6 (1989), 114–148. In Russian; translated in *Leningrad Math. J.* **1**:6 (1990), 1419–1457. [MR 91b:17016](#)
- [Fuchs et al. 1999] J. Fuchs, A. C. Ganchev, K. Szlachányi, and P. Vecsernyés, “ S_4 symmetry of $6j$ symbols and Frobenius–Schur indicators in rigid monoidal C^* categories”, *J. Math. Phys.* **40**:1 (1999), 408–426. [MR 99k:81111](#) [Zbl 0986.81044](#)
- [Goodman and Wenzl 2002] F. Goodman and H. Wenzl, “Ideals in the Temperley Lieb category”, preprint, 2002. [arXiv 0206301](#)
- [Henriques and Kamnitzer 2006] A. Henriques and J. Kamnitzer, “Crystals and coboundary categories”, *Duke Math. J.* **132**:2 (2006), 191–216. [MR 2007m:17020](#) [Zbl 1123.22007](#)
- [Kamnitzer and Tingley 2009] J. Kamnitzer and P. Tingley, “The crystal commutator and Drinfel'd's unitarized R -matrix”, *J. Algebraic Combin.* **29**:3 (2009), 315–335. [MR 2496310](#)
- [Kirillov and Reshetikhin 1990] A. N. Kirillov and N. Reshetikhin, “ q -Weyl group and a multiplicative formula for universal R -matrices”, *Comm. Math. Phys.* **134**:2 (1990), 421–431. [MR 92c:17023](#) [Zbl 0723.17014](#)

- [Kuperberg 1996] G. Kuperberg, “Spiders for rank 2 Lie algebras”, *Comm. Math. Phys.* **180**:1 (1996), 109–151. [MR 97f:17005](#) [Zbl 0870.17005](#)
- [Levendorskii and Soĭbel’man 1991] S. Z. Levendorskii and Y. S. Soĭbel’man, “The quantum Weyl group and a multiplicative formula for the R -matrix of a simple Lie algebra”, *Funktsional. Anal. i Prilozhen.* **25**:2 (1991), 73–76. In Russian; translated in *Funct. Anal. Appl.* **25**:2 (1991), 143–145. [MR 93a:17017](#)
- [Linchenko and Montgomery 2000] V. Linchenko and S. Montgomery, “A Frobenius–Schur theorem for Hopf algebras”, *Algebr. Represent. Theory* **3**:4 (2000), 347–355. [MR 2001k:16073](#) [Zbl 0971.16018](#)
- [Lusztig 1993] G. Lusztig, *Introduction to quantum groups*, Progress in Math. **110**, Birkhäuser, Boston, MA, 1993. [MR 94m:17016](#) [Zbl 0788.17010](#)
- [Ng and Schauenburg 2007] S.-H. Ng and P. Schauenburg, “Higher Frobenius–Schur indicators for pivotal categories”, pp. 63–90 in *Hopf algebras and generalizations*, edited by L. H. Kauffman et al., Contemp. Math. **441**, Amer. Math. Soc., Providence, RI, 2007. [MR 2008m:18015](#) [Zbl 1153.18008](#)
- [Ohtsuki 2002] T. Ohtsuki, *Quantum invariants: a study of knots, 3-manifolds, and their sets*, Series on Knots and Everything **29**, World Scientific, River Edge, NJ, 2002. [MR 2003f:57027](#) [Zbl 0991.57001](#)
- [Reshetikhin 1989] N. Y. Reshetikhin, “Quasitriangular Hopf algebras and invariants of links”, *Algebra i Analiz* **1**:2 (1989), 169–188. In Russian; translated in *Leningrad Math. J.*, **1**:2 (1990), 491–513. [MR 90k:17033](#)
- [Reshetikhin and Turaev 1990] N. Y. Reshetikhin and V. G. Turaev, “Ribbon graphs and their invariants derived from quantum groups”, *Comm. Math. Phys.* **127**:1 (1990), 1–26. [MR 91c:57016](#) [Zbl 0768.57003](#)
- [Reshetikhin and Turaev 1991] N. Reshetikhin and V. G. Turaev, “Invariants of 3-manifolds via link polynomials and quantum groups”, *Invent. Math.* **103**:3 (1991), 547–597. [MR 92b:57024](#) [Zbl 0725.57007](#)
- [Shum 1994] M. C. Shum, “Tortile tensor categories”, *J. Pure Appl. Algebra* **93**:1 (1994), 57–110. [MR 95a:18008](#) [Zbl 0803.18004](#)
- [Turaev 1994] V. G. Turaev, *Quantum invariants of knots and 3-manifolds*, de Gruyter Studies in Math. **18**, Walter de Gruyter & Co., Berlin, 1994. [MR 95k:57014](#) [Zbl 0812.57003](#)

Communicated by Susan Montgomery

Received 2008-10-06

Revised 2009-09-11

Accepted 2009-10-17

nsnyder@math.columbia.edu

*Department of Mathematics, Columbia University,
Rm 626, MC 4403, 2990 Broadway, New York, NY 10027,
United States*

<http://math.columbia.edu/~nsnyder/>

ptingley@math.mit.edu

*Department of Mathematics,
Massachusetts Institute of Technology,
77 Massachusetts Ave, Cambridge, MA 02139, United States*

<http://www-math.mit.edu/~ptingley/>