

# *Algebra & Number Theory*

Volume 8

2014

No. 6

**On moduli spaces for quasitilted algebras**

Grzegorz Bobiński



# On moduli spaces for quasitilted algebras

Grzegorz Bobiński

We prove that if a quasitilted algebra is tame, then the associated moduli spaces are products of projective spaces. Together with an earlier result of Chindris this gives a geometric characterization of the tame quasitilted algebras. In our proof we use knowledge of the representation theory of the tame quasitilted algebras and a construction of semi-invariants as determinants.

Throughout the article  $\mathbb{k}$  is an algebraically closed field of characteristic 0. By  $\mathbb{Z}$ ,  $\mathbb{N}$  and  $\mathbb{N}_+$  we denote the sets of integers, nonnegative integers and positive integers, respectively. Finally, if  $i, j \in \mathbb{Z}$ , then  $[i, j] := \{k \in \mathbb{Z} \mid i \leq k \leq j\}$  (in particular,  $[i, j] = \emptyset$  if  $i > j$ ).

There is a well-known dichotomy for finite-dimensional algebras due to Drozd [1980]: every algebra is either tame or wild, but not both. Here a finite-dimensional algebra is called tame if, for each dimension  $d$ , the indecomposable  $d$ -dimensional modules form finitely many one-parameter families. On the other hand, an algebra  $\Lambda$  is wild if the classification of  $\Lambda$ -modules is as difficult as the classification of the pairs of two (noncommuting) endomorphisms of a finite-dimensional vector space (the latter problem is considered to be hopeless).

The above definitions of tame and wild algebras are of a geometric nature. This encourages people to look for characterizations of representation type, which use properties of geometric objects associated with them, for example, module varieties (some results of this type can be found in [Bobiński and Skowroński 1999; Skowroński and Zvara 1998]). In particular, Skowroński and Weyman [2000] have proved that a hereditary algebra  $\Lambda$  is tame if and only if all the corresponding rings of semi-invariants are complete intersections. Inspired by this result Chindris [2009; 2011; 2013] has initiated a programme, whose aim is to characterize representation type in terms of (rational) invariant theory (see also [Domokos 2011] for an earlier result in this direction). As a result of his studies he and Carroll published the following conjecture, which they attribute to Weyman:

**Conjecture** [Carroll and Chindris 2012]. *Let  $\Lambda$  be an algebra. Then the following conditions are equivalent:*

---

MSC2010: primary 16G10; secondary 16G60, 13A50.

Keywords: quasitilted algebra, moduli space, semi-invariant.

- (1)  $\Lambda$  is of tame representation type.
- (2) For any dimension vector  $\mathbf{d}$ , for any irreducible component  $\mathcal{C}$  of the variety  $\text{mod}_\Lambda(\mathbf{d})$  of  $\Lambda$ -modules of dimension vector  $\mathbf{d}$ , and for any weight  $\theta$  such that  $\mathcal{C}_\theta^{\text{ss}} \neq \emptyset$ ,  $\mathcal{M}(\mathcal{C})_\theta^{\text{ss}}$  is a product of projective spaces.

Here  $\mathcal{C}_\theta^{\text{ss}}$  denotes the open subset of  $\theta$ -semistable modules in  $\mathcal{C}$  and  $\mathcal{M}(\mathcal{C})_\theta^{\text{ss}}$  denotes the associated moduli space (see [Section 5](#)).

This conjecture has no chance of holding in such generality. Obvious counterexamples are local wild algebras. There is also a counterexample due to Ringel of a triangular (no cycles in the Gabriel quiver) wild algebra, such that all the associated moduli spaces are points.

The aim of this paper is to verify this conjecture for the quasitilted algebras. The quasitilted algebras form an important class of finite-dimensional algebras. Using covering techniques, the study of some classes of algebras can be reduced to the study of quasitilted algebras. In particular, every self-injective algebra of polynomial growth is a socle deformation of an orbit algebra of the repetitive algebra of a tame quasitilted algebra with positive semidefinite Euler form (for more in this direction see the survey article [\[Skowroński 2006\]](#)).

The main result of the paper is the following:

**Theorem 1.** *Let  $\Lambda$  be a quasitilted algebra. Then the following conditions are equivalent:*

- (1)  $\Lambda$  is of tame representation type.
- (2) For any dimension vector  $\mathbf{d}$ , for any irreducible component  $\mathcal{C}$  of  $\text{mod}_\Lambda(\mathbf{d})$ , and for any weight  $\theta$  such that  $\mathcal{C}_\theta^{\text{ss}} \neq \emptyset$ ,  $\mathcal{M}(\mathcal{C})_\theta^{\text{ss}}$  is a product of projective spaces.

The implication (2)  $\implies$  (1) has been proved for the tilted algebras by Chindris [\[2013, Proposition 4.1\]](#). In fact, as has been explained to me by Chindris, his proof of the implication (2)  $\implies$  (1) generalizes to the quasitilted algebras. More precisely, Chindris [\[2013\]](#) used a result of Kerner [\[1989\]](#) stating that every wild tilted algebra has a convex subalgebra which is wild concealed. In the case of the quasitilted algebras one has to use results of Lenzing and Skowroński [\[1996\]](#) (every wild quasitilted algebra has a convex subalgebra which is wild almost concealed-canonical) and Meltzer [\[1996\]](#) (every wild almost concealed-canonical algebra has a convex subcategory which is wild concealed). Thus in the paper we concentrate on the proof of the following theorem:

**Theorem 2.** *Let  $\Lambda$  be a tame quasitilted algebra. Then for any dimension vector  $\mathbf{d}$ , for any irreducible component  $\mathcal{C}$  of  $\text{mod}_\Lambda(\mathbf{d})$ , and for any weight  $\theta$  such that  $\mathcal{C}_\theta^{\text{ss}} \neq \emptyset$ ,  $\mathcal{M}(\mathcal{C})_\theta^{\text{ss}}$  is a product of projective spaces.*

The paper is organized as follows. In [Section 1](#) we recall basic facts about quivers and their representations. [Section 2](#) is devoted to a short introduction of quasitilted algebras. Next, in [Sections 3](#) and [4](#) we introduce module varieties and semi-invariants, respectively. Moreover, in [Section 4](#) some reduction results for semi-invariants are obtained. Finally, in [Section 5](#) we recall King's construction of moduli spaces and in [Section 6](#) we prove that the moduli spaces for the tame quasitilted algebras are products of projective spaces.

## 1. Quivers and their representations

In this section we present facts about quivers and their representations, which we use in the paper. As a general background we suggest [[Assem et al. 2006](#); [Auslander et al. 1997](#); [Ringel 1984](#)].

By a quiver  $Q$  we mean a finite set  $Q_0$  (called the set of vertices of  $Q$ ) together with a finite set  $Q_1$  (called the set of arrows of  $Q$ ) and two maps  $s, t : Q_1 \rightarrow Q_0$ , which assign to each arrow  $\alpha$  its starting vertex  $s\alpha$  and its terminating vertex  $t\alpha$ , respectively. By a path of length  $n \in \mathbb{N}_+$  in a quiver  $Q$  we mean a sequence  $\sigma = (\alpha_1, \dots, \alpha_n)$  of arrows such that  $s\alpha_i = t\alpha_{i+1}$  for each  $i \in [1, n-1]$ . In the above situation we put  $\ell\sigma := n$ ,  $s\sigma := s\alpha_n$  and  $t\sigma := t\alpha_1$ . We treat every arrow in  $Q$  as a path of length 1. Moreover, for each vertex  $x$  we have a trivial path  $\mathbf{1}_x$  at  $x$  such that  $\ell\mathbf{1}_x := 0$  and  $s\mathbf{1}_x := x =: t\mathbf{1}_x$ . For the rest of the paper we assume that the considered quivers do not have oriented cycles, where by an oriented cycle we mean a path  $\sigma$  of positive length such that  $s\sigma = t\sigma$ .

Let  $Q$  be a quiver. By the path algebra  $\mathbb{k}Q$  of  $Q$ , we mean the vector space with a basis formed by the paths in  $Q$  and multiplication induced by the concatenation of paths. If  $x$  and  $y$  are vertices of  $Q$ , we put  $\mathbb{k}Q(x, y) := \mathbf{1}_y \mathbb{k}Q \mathbf{1}_x$ ; i.e.,  $\mathbb{k}Q(x, y)$  is the space spanned by the paths with the starting vertex  $x$  and the terminating vertex  $y$ . The (finite-dimensional)  $\mathbb{k}Q$ -modules may be identified with the  $\mathbb{k}$ -representations of  $Q$ , where by a  $\mathbb{k}$ -representation of  $Q$  we mean  $V$  consisting of finite-dimensional  $\mathbb{k}$ -vector spaces  $V(x)$ ,  $x \in Q_0$ , and  $\mathbb{k}$ -linear maps  $V(\alpha) : V(s\alpha) \rightarrow V(t\alpha)$ ,  $\alpha \in Q_1$ . In particular, if  $M$  is a  $\mathbb{k}Q$ -module and  $V$  is the corresponding representation, then  $V(x) := \mathbf{1}_x M$  for each  $x \in Q_0$ . We will usually identify  $\mathbb{k}Q$ -modules with the corresponding representations of  $\mathbb{k}$ . If  $V$  and  $W$  are representations of a quiver  $Q$ , then a morphism  $\varphi : V \rightarrow W$  is given by linear maps  $\varphi(x) : V(x) \rightarrow W(x)$ ,  $x \in Q_0$ , such that  $W(\alpha)\varphi(s\alpha) = \varphi(t\alpha)V(\alpha)$  for each  $\alpha \in Q_1$ . We denote the category of  $\mathbb{k}$ -representations of  $Q$  by  $\text{rep } Q$ . If  $V$  is a representation,  $x, y \in Q_0$ , and  $\omega \in \mathbb{k}Q(x, y)$ , then one defines  $V(\omega) : V(x) \rightarrow V(y)$  in an obvious way. Given a representation  $V$  of  $Q$  we denote by  $\mathbf{dim } V$  its dimension vector, defined by the formula  $(\mathbf{dim } V)(x) := \dim_{\mathbb{k}} V(x)$ , for  $x \in Q_0$ . Observe that  $\mathbf{dim } V \in \mathbb{N}^{Q_0}$  for each representation  $V$  of  $Q$ . We call the elements of  $\mathbb{N}^{Q_0}$  dimension vectors. If  $d$

is a dimension vector, then we denote by  $\text{supp } \mathbf{d}$  the subquiver of  $Q$  induced by the vertices  $x$  such that  $\mathbf{d}(x) \neq 0$ . A dimension vector  $\mathbf{d}$  is called connected if the quiver  $\text{supp } \mathbf{d}$  is connected. A dimension vector  $\mathbf{d}$  is called sincere if  $\text{supp } \mathbf{d} = Q$ .

By a bound quiver  $(Q, I)$  we mean a quiver  $Q$  together with an ideal  $I$  of  $\mathbb{k}Q$  such that  $I \subseteq \langle Q_1 \rangle^2$ , where by  $\langle Q_1 \rangle$  we denote the ideal of  $\mathbb{k}Q$  generated by the arrows. Given a bound quiver  $(Q, I)$ , we call the algebra  $\mathbb{k}Q/I$  the path algebra of  $(Q, I)$ . Note that if  $(Q, I)$  is a bound quiver, then the  $\mathbb{k}Q/I$ -modules may be identified with the representations  $V$  of  $Q$  such that  $V(\omega) = 0$  for each  $\omega \in I \cap (\bigcup_{x,y \in Q_0} \mathbb{k}Q(x, y))$ . If  $\Lambda$  is the path algebra of a bound quiver  $(Q, I)$ , then we call  $Q$  the Gabriel quiver of  $\Lambda$ . Gabriel proved that (up to isomorphism)  $Q$  is uniquely determined by  $\Lambda$ . Moreover, Gabriel's theorem implies that each quasitilted algebra is Morita equivalent to the path algebra of a bound quiver (since we only consider quivers without oriented cycles, we also need [Happel et al. 1996, Proposition III.1.1(b)] for this result). Thus from now on all algebras considered are the path algebras of bound quivers. Observe that if  $J$  is an ideal in an algebra  $\Lambda$ , then the Gabriel quiver of  $\Lambda/J$  is a subquiver of the Gabriel quiver of  $\Lambda$  (here this is important that there are no oriented cycles in the considered quivers). If  $(Q, I)$  is a bound quiver, then an algebra  $\Lambda'$  is called a convex subalgebra of  $\mathbb{k}Q/I$  if there exists a convex subquiver  $Q'$  of  $Q$  such that  $\Lambda' = \mathbb{k}Q'/(I \cap \mathbb{k}Q')$ . Recall that a subquiver  $Q'$  of  $Q$  is called convex if for every path  $(\alpha_1, \dots, \alpha_n)$  in  $Q$  with  $s\alpha_n, t\alpha_1 \in Q'_0$  we have  $\alpha_i \in Q'_1$  for each  $i \in [1, n]$  (and, consequently,  $s\alpha_i \in Q'_0$  for each  $i \in [1, n-1]$ ).

Let  $\Lambda$  be an algebra with Gabriel quiver  $Q$ . For a vertex  $x$  of  $Q$  we put  $P_\Lambda(x) := \Lambda \mathbf{1}_x$ . Then  $P_\Lambda(x)$  is an indecomposable projective  $\Lambda$ -module and every indecomposable projective  $\Lambda$ -module is (up to isomorphism) of this form. If  $V$  is a  $\Lambda$ -module, then  $\text{Hom}_\Lambda(P_\Lambda(x), V) = V(x)$  for each  $x \in Q_0$ . In particular,  $\text{Hom}_\Lambda(P_\Lambda(x), P_\Lambda(y)) = \mathbb{k}Q(y, x)$  for any  $x, y \in Q_0$ . Moreover, if  $x, y \in Q_0$ ,  $\omega \in \mathbb{k}(y, x)$  and  $V$  is a  $\Lambda$ -module, then

$$\text{Hom}_\Lambda(\omega, V) : \text{Hom}_\Lambda(P_\Lambda(y), V) \rightarrow \text{Hom}_\Lambda(P_\Lambda(x), V)$$

is just  $V(\omega) : V(y) \rightarrow V(x)$ .

For an algebra  $\Lambda$  we denote by  $\text{mod } \Lambda$  the category of  $\Lambda$ -modules. Next, if  $\Lambda$  is an algebra and  $\Lambda^{\text{op}}$  is the opposite algebra of  $\Lambda$ , then we denote by  $D_\Lambda$  the duality  $\text{mod } \Lambda \rightarrow \text{mod } \Lambda^{\text{op}}$  given by

$$D_\Lambda(M) := \text{Hom}_k(M, k) \quad (M \in \text{mod } \Lambda).$$

Finally, for an algebra  $\Lambda$  we denote by  $\tau_\Lambda$  the corresponding Auslander–Reiten translation, which assigns to each  $\Lambda$ -module  $M$  another  $\Lambda$ -module  $\tau_\Lambda M$  (see [Assem et al. 2006, Section IV.2] for a definition). We will need the following consequence of the Auslander–Reiten formula [Assem et al. 2006, Theorem IV.2.13]: if  $M$  and  $N$

are  $\Lambda$ -modules and  $\text{pdim}_\Lambda M \leq 1$ , then

$$\dim_{\mathbb{k}} \text{Ext}_\Lambda^1(M, N) = \dim_{\mathbb{k}} \text{Hom}_\Lambda(N, \tau_\Lambda M). \tag{1}$$

Let  $\Lambda$  be an algebra with Gabriel quiver  $Q$ . Since there are no cycles in  $Q$ ,  $\text{gldim } \Lambda < \infty$ . Consequently, we may define the bilinear form  $\langle -, - \rangle_\Lambda : \mathbb{Z}^{Q_0} \times \mathbb{Z}^{Q_0} \rightarrow \mathbb{Z}$  by the condition


$$\langle \mathbf{dim } M, \mathbf{dim } N \rangle_\Lambda = \sum_{i \in \mathbb{N}} \dim_{\mathbb{k}} \text{Ext}_\Lambda^i(M, N)$$

for all  $\Lambda$ -modules  $M$  and  $N$ . We denote the corresponding quadratic form, called the Euler form, by  $\chi_\Lambda$ .

### 2. Quasitilted algebras

A module  $T$  over an algebra  $\Lambda$  is called tilting if  $\text{pdim}_\Lambda T \leq 1$ ,  $\text{Ext}_\Lambda^1(T, T) = 0$ , and  $T$  is a direct sum of  $n$  pairwise nonisomorphic indecomposable  $\Lambda$ -modules, where  $n$  is the number of vertices of the Gabriel quiver of  $\Lambda$ . By a tilted algebra we mean the opposite algebra of the endomorphism algebra of a tilting module over the path algebra of a quiver.

An algebra  $\Lambda$  is called quasitilted if  $\Lambda$  is the opposite algebra of the endomorphism algebra of a tilting object in a connected hereditary abelian  $\mathbb{k}$ -category with finite-dimensional homomorphism and extension spaces. Equivalently,  $\text{gldim } \Lambda \leq 2$  and either  $\text{pdim}_\Lambda X \leq 1$  or  $\text{idim}_\Lambda X \leq 1$  for each indecomposable  $\Lambda$ -module  $X$  (see [Happel et al. 1996, Theorem 2.3]). Two prominent examples of quasitilted algebras are the tilted algebras introduced above and the Ringel canonical algebras  $\Lambda(\mathbf{m}, \boldsymbol{\lambda})$ , where  $\mathbf{m} = (m_1, \dots, m_n)$ ,  $n \geq 3$ , is a sequence of integers greater than 1 and  $\boldsymbol{\lambda} = (\lambda_3, \dots, \lambda_n)$ . In the above situation  $\Lambda(\mathbf{m}, \boldsymbol{\lambda})$  is the path algebra of the quiver



modulo the ideal generated by the relations

$$\alpha_{1,1} \cdots \alpha_{1,m_1} + \lambda_i \alpha_{2,1} \cdots \alpha_{2,m_2} - \alpha_{i,1} \cdots \alpha_{i,m_i}, \quad i \in [3, n].$$

Due to [Happel 2001, Theorem 3.1] every quasitilted algebra is either a tilted algebra or is of canonical type (i.e., is derived equivalent to a canonical algebra).

The structure of the module categories over tilted algebras has been investigated in [Kerner 1989], while the structure of the module categories over quasitilted algebras of canonical type has been studied in [Lenzing and Skowroński 1996]. We also refer to [Skowroński 1998] for a characterization of the tame quasitilted algebras and to [Ringel 1984] for a description of the module categories over so-called tubular algebras, which form an important subclass of the tame quasitilted algebras. We list some consequences of these investigations.

Let  $\Lambda$  be a tame quasitilted algebra with Gabriel quiver  $Q$ . If  $\mathbf{d}$  is a dimension vector, then there exists an indecomposable  $\Lambda$ -module with dimension vector  $\mathbf{d}$  if and only if  $\mathbf{d}$  is a root of  $\chi_\Lambda$ , i.e.,  $\mathbf{d}$  is a connected nonzero dimension vector such that  $\chi_\Lambda(\mathbf{d}) \in \{0, 1\}$ . We call a root  $\mathbf{d}$  isotropic if  $\chi_\Lambda(\mathbf{d}) = 0$ . We call a root  $\mathbf{d}$  a Schur root if there exists a  $\Lambda$ -module  $X$  (necessarily indecomposable) with dimension vector  $\mathbf{d}$  and trivial endomorphism algebra. Let  $\mathbf{d}_1$  and  $\mathbf{d}_2$  be two isotropic roots with  $\text{supp } \mathbf{d}_1 \cap \text{supp } \mathbf{d}_2 \neq \emptyset$ . Then  $\text{Hom}_\Lambda(X_1, X_2) \neq 0$  for all indecomposable  $\Lambda$ -modules  $X_1$  and  $X_2$  with  $\mathbf{dim } X_1 = \mathbf{d}_1$  and  $\mathbf{dim } X_2 = \mathbf{d}_2$ , or  $\text{Hom}_\Lambda(X_2, X_1) \neq \emptyset$  for all indecomposable  $\Lambda$ -modules  $X_1$  and  $X_2$  with  $\mathbf{dim } X_1 = \mathbf{d}_1$  and  $\mathbf{dim } X_2 = \mathbf{d}_2$ , or  $\mathbf{d}_1$  and  $\mathbf{d}_2$  are multiplicities of the same isotropic Schur root.

Now assume that  $\Lambda$  is a canonical algebra. The indecomposable  $\Lambda$ -modules can be divided into three classes,  $\mathcal{L}_\Lambda$ ,  $\mathcal{T}_\Lambda$  and  $\mathcal{R}_\Lambda$ : the class  $\mathcal{L}_\Lambda$  is formed by the indecomposable  $\Lambda$ -modules  $X$  such that  $\dim X(0) > \dim X(\infty)$ , the class  $\mathcal{T}_\Lambda$  is formed by the indecomposable  $\Lambda$ -modules  $X$  such that  $\dim X(0) = \dim X(\infty)$ , and the class  $\mathcal{R}_\Lambda$  is formed by the indecomposable  $\Lambda$ -modules  $X$  such that  $\dim X(0) < \dim X(\infty)$ . An algebra  $\Lambda$  is called (almost) concealed-canonical if  $\Lambda$  is the opposite algebra of the endomorphism algebra of a tilting module  $T$ , which is a direct sum of indecomposable modules from  $\mathcal{L}_\Lambda$  ( $\mathcal{L}_\Lambda \cup \mathcal{T}_\Lambda$ , respectively).

### 3. Module varieties

Let  $\Lambda$  be the path algebra of a bound quiver  $(Q, I)$ . For a dimension vector  $\mathbf{d}$  we denote by  $\text{mod}_\Lambda(\mathbf{d})$  the set of representations  $M$  of  $(Q, I)$  (recall that we identify the  $\Lambda$ -modules with the representations of  $(Q, I)$ ) such that  $M(x) = \mathbb{k}^{\mathbf{d}(x)}$  for each  $x \in Q_0$ . This set can be naturally identified with a closed subset of the affine space  $\text{rep}_Q(\mathbf{d}) := \prod_{\alpha \in Q_1} \mathbb{M}(\mathbf{d}(t\alpha), \mathbf{d}(s\alpha))$ ; thus it has the structure of an affine variety (note that under this identification  $\text{rep}_Q(\mathbf{d}) = \text{mod}_{\mathbb{k}Q}(\mathbf{d})$ ). The reductive group  $\text{GL}(\mathbf{d}) := \prod_{x \in Q_0} \text{GL}(\mathbf{d}(x))$  acts on  $\text{mod}_\Lambda(\mathbf{d})$  via

$$(g * M)(\alpha) := g(t\alpha) \cdot M(\alpha) \cdot g(s\alpha)^{-1} \quad (g \in \text{GL}(\mathbf{d}), \alpha \in Q_1).$$

If  $M \in \text{mod}_\Lambda(\mathbf{d})$ , then we denote its orbit with respect to this action by  $\mathcal{O}(M)$ . One has  $\mathcal{O}(M) = \mathcal{O}(N)$  if and only if  $M \simeq N$ .

Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be closed irreducible subsets of varieties  $\text{mod}_\Lambda(\mathbf{d}_1)$  and  $\text{mod}_\Lambda(\mathbf{d}_2)$ , respectively. By  $\mathcal{C}_1 \oplus \mathcal{C}_2$  we denote the closure of the set consisting of all  $M \in \text{mod}_\Lambda(\mathbf{d}_1 + \mathbf{d}_2)$  such that  $M \simeq M_1 \oplus M_2$  for some  $M_1 \in \mathcal{C}_1$  and  $M_2 \in \mathcal{C}_2$ . In the above situation we call  $\mathcal{C}_1$  and  $\mathcal{C}_2$  summands of  $\mathcal{C}$ .

An irreducible component  $\mathcal{C}$  of  $\text{mod}_\Lambda(\mathbf{d})$  is called indecomposable if the indecomposable modules in  $\mathcal{C}$  form a dense subset of  $\mathcal{C}$ . If  $\mathcal{C}$  is an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , then there exist uniquely determined (up to ordering) indecomposable irreducible components  $\mathcal{C}_1, \dots, \mathcal{C}_n$  of  $\mathcal{C}$  such that

$$\mathcal{C} = \mathcal{C}_1 \oplus \dots \oplus \mathcal{C}_n$$

[Crawley-Boevey and Schröer 2002, Theorem 1.1] (see also [de la Peña 1991]). We call the above presentation the generic decomposition of  $\mathcal{C}$ . Moreover, if, for  $i \in [1, n]$ ,  $\mathcal{C}_i \subseteq \text{mod}_\Lambda(\mathbf{d}_i)$ , then we call  $\mathbf{d}_1, \dots, \mathbf{d}_n$  the generic summands of  $\mathbf{d}$  at  $\mathcal{C}$ .

Now we present a description of the indecomposable irreducible components in the case of the tame quasitilted algebras, which follows from [Bobiński and Skowroński 1999]. First, if  $\mathbf{d}$  is a dimension vector, then there is at most one indecomposable irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ . Thus if it exists we denote it by  $\mathcal{C}(\mathbf{d})$ . Moreover, there exists an indecomposable irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$  if and only if  $\mathbf{d}$  is a Schur root. Moreover, if  $\mathbf{d}$  is not isotropic, then  $\mathcal{C}(\mathbf{d})$  is an orbit closure, i.e., there exists a  $\Lambda$ -module  $M$  such that  $\mathcal{C}(\mathbf{d}) = \overline{\mathcal{O}(M)}$ .

#### 4. Semi-invariants

Let  $Q$  be a quiver,  $\mathbf{d}$  a dimension vector, and  $\mathcal{C}$  a  $\text{GL}(\mathbf{d})$ -invariant closed subset of  $\text{rep}_Q(\mathbf{d})$ . The action of  $\text{GL}(\mathbf{d})$  on  $\mathcal{C}$  induces an action on the coordinate ring  $\mathbb{k}[\mathcal{C}]$  via

$$(g * f)(M) := f(g^{-1} * M) \quad (g \in \text{GL}(\mathbf{d}), f \in \mathbb{k}[\mathcal{C}], M \in \mathcal{C}).$$

If  $\mathcal{C}$  is irreducible, then there is a unique closed orbit in  $\mathcal{C}$ , that of the semisimple module with dimension vector  $\mathbf{d}$ , hence there are only trivial  $\text{GL}(\mathbf{d})$ -invariant regular functions on  $\mathcal{C}$ , i.e.,  $\mathbb{k}[\mathcal{C}]^{\text{GL}(\mathbf{d})} = \mathbb{k}$ . However, one may still have nontrivial semi-invariants. A regular function  $f \in \mathbb{k}[\mathcal{C}]$  is called a semi-invariant of weight  $\theta \in \mathbb{Z}^{Q_0}$  if

$$g * f = \chi^\theta(g) \cdot f$$

for each  $g \in \text{GL}(\mathbf{d})$ . Here  $\chi^\theta : \text{GL}(\mathbf{d}) \rightarrow \mathbb{k}^\times$  is given by

$$\chi^\theta(g) := \prod_{x \in Q_0} \det^{\theta(e_x)}(g(x)) \quad (g \in \text{GL}(\mathbf{d})),$$



where  $\mathbf{e}_x, x \in Q_0$ , are the standard basis vectors of  $\mathbb{Z}^{Q_0}$ . We denote the space of semi-invariants of weight  $\theta$  by  $\text{SI}[\mathcal{C}]_\theta$ . One easily observes that  $\theta(\mathbf{d}) = 0$  provided  $\text{SI}[\mathcal{C}]_\theta \neq 0$ .

We present a method of constructing semi-invariants, which in the case of quivers is due to Schofield [1991], and has been generalized to the case of bound quivers independently by Derksen and Weyman [2002] and Domokos [2002] (we also refer to the latter two articles for proofs). Let  $Q$  be a quiver and  $\mathbf{d}$  a dimension vector. Fix sequences  $\mathbf{x} = (x_1, \dots, x_n)$  and  $\mathbf{y} = (y_1, \dots, y_m)$  of vertices of  $Q$ . Put

$$\mathbb{k}Q(\mathbf{x}, \mathbf{y}) := \prod_{\substack{i \in [1, n] \\ j \in [1, m]}} \mathbb{k}Q(x_i, y_j).$$

If  $\phi = (\phi_{i,j})_{i \in [1, n], j \in [1, m]} \in \mathbb{k}Q(\mathbf{x}, \mathbf{y})$  and  $M \in \text{rep}_Q(\mathbf{d})$  for a dimension vector  $\mathbf{d}$ , then we obtain a map

$$M(\phi) := [M(\phi_{i,j})]_{\substack{i \in [1, n] \\ j \in [1, m]}} : M(\mathbf{x}) := \bigoplus_{i \in [1, n]} M(x_i) \rightarrow M(\mathbf{y}) := \bigoplus_{j \in [1, m]} M(y_j).$$

If, in addition,  $\sum_{i \in [1, n]} \mathbf{d}(x_i) = \sum_{j \in [1, m]} \mathbf{d}(y_j)$ , then we may define a regular function  $c_{\mathbf{d}}^\phi : \text{rep}_Q(\mathbf{d}) \rightarrow \mathbb{k}$  by

$$c_{\mathbf{d}}^\phi(M) := \det M(\phi) \quad (M \in \text{rep}_Q(\mathbf{d})).$$

Then  $c_{\mathbf{d}}^\phi$  is a semi-invariant of weight  $\theta^\phi$ , where

$$\theta^\phi(\mathbf{c}) := \sum_{i \in [1, n]} \mathbf{c}(x_i) - \sum_{j \in [1, m]} \mathbf{c}(y_j) \quad (\mathbf{c} \in \mathbb{Z}^{Q_0})$$

(note that, in particular,  $\theta^\phi(\mathbf{d}) = 0$ ). If  $\mathcal{C}$  is a  $\text{GL}(\mathbf{d})$ -invariant closed subset of  $\text{rep}_Q(\mathbf{d})$ , then we denote the restriction  $c_{\mathbf{d}}^\phi|_{\mathcal{C}}$  of  $c_{\mathbf{d}}^\phi$  to  $\mathcal{C}$  by  $c_{\mathcal{C}}^\phi$ .

We list some obvious consequences (cf. [Derksen and Weyman 2000, Lemma 1]).

**Lemma 4.1.** *Let  $\mathbf{x}$  and  $\mathbf{y}$  be sequences of vertices of a quiver  $Q$ ,  $\phi \in \mathbb{k}Q(\mathbf{x}, \mathbf{y})$ ,  $\mathbf{d}$  a dimension vector such that  $\theta^\phi(\mathbf{d}) = 0$ , and  $M = M_1 \oplus M_2 \in \text{mod}_\Lambda(\mathbf{d})$ .*

- (1) *If  $\theta^\phi(\mathbf{dim} M_1) \neq 0$  (hence, equivalently,  $\theta^\phi(\mathbf{dim} M_2) \neq 0$ ), then  $c_{\mathbf{d}}^\phi(M) = 0$ .*
- (2) *If  $\theta^\phi(\mathbf{dim} M_1) = 0$  (hence, equivalently,  $\theta^\phi(\mathbf{dim} M_2) = 0$ ), then  $c_{\mathbf{d}}^\phi(M) = c_{\mathbf{dim} M_1}^\phi(M_1) \cdot c_{\mathbf{dim} M_2}^\phi(M_2)$ . □*

If we have sequences  $\mathbf{x}, \mathbf{x}', \mathbf{y}$  and  $\mathbf{y}'$  of vertices of a quiver  $Q$ ,  $\phi \in \mathbb{k}(\mathbf{x}, \mathbf{y})$  and  $\phi' \in \mathbb{k}(\mathbf{x}', \mathbf{y}')$ , then we may define an element  $\phi \oplus \phi' \in \mathbb{k}Q(\mathbf{x} \cdot \mathbf{x}', \mathbf{y} \cdot \mathbf{y}')$  in the obvious way, where  $\mathbf{x} \cdot \mathbf{x}'$  and  $\mathbf{y} \cdot \mathbf{y}'$  are the concatenations of the respective sequences. Observe that

$$M(\phi \oplus \phi') = \begin{bmatrix} M(\phi) & 0 \\ 0 & M(\phi') \end{bmatrix} : M(\mathbf{x}) \oplus M(\mathbf{x}') \rightarrow M(\mathbf{y}) \oplus M(\mathbf{y}')$$

for each  $M \in \text{rep}_Q(\mathbf{d})$ . Consequently, we get the following.

**Lemma 4.2.** *Let  $\mathbf{x}, \mathbf{x}', \mathbf{y}$  and  $\mathbf{y}'$  be sequences of vertices of a quiver  $Q$ ,  $\phi \in \mathbb{k}(\mathbf{x}, \mathbf{y})$  and  $\phi' \in \mathbb{k}(\mathbf{x}', \mathbf{y}')$ . If  $\mathbf{d}$  is a dimension vector and  $\theta^\phi(\mathbf{d}) = 0 = \theta^{\phi'}(\mathbf{d})$ , then*

$$c_{\mathbf{d}}^{\phi \oplus \phi'} = c_{\mathbf{d}}^\phi \cdot c_{\mathbf{d}}^{\phi'}.$$

In particular,  $c_{\mathbf{d}}^{\phi \oplus \phi'}$  is a semi-invariant of weight  $\theta^\phi + \theta^{\phi'}$ . □

We can interpret the above construction using projective presentations. Let  $\Lambda$  be a factor algebra of  $\mathbb{k}Q$  for a quiver  $Q$ . As above, let  $\mathbf{x} = (x_1, \dots, x_n)$  and  $\mathbf{y} = (y_1, \dots, y_m)$  be sequences of vertices of a quiver  $Q$ , and  $\phi \in \mathbb{k}Q(\mathbf{x}, \mathbf{y})$ . If we put

$$P_\Lambda(\mathbf{x}) := \bigoplus_{i \in [1, n]} P_\Lambda(x_i) \quad \text{and} \quad P_\Lambda(\mathbf{y}) := \bigoplus_{j \in [1, m]} P_\Lambda(y_j),$$

then we may view  $\phi$  as a map  $P_\Lambda(\mathbf{y}) \rightarrow P_\Lambda(\mathbf{x})$  (note that every map between projective  $\Lambda$ -modules is of this form, for some  $\mathbf{x}, \mathbf{y}$  and  $\phi$ ). Observe that

$$\theta^\phi(\mathbf{dim} M) = \dim_{\mathbb{k}} \text{Hom}_\Lambda(P_\Lambda(\mathbf{x}), M) - \dim_{\mathbb{k}} \text{Hom}_\Lambda(P_\Lambda(\mathbf{y}), M)$$

for each  $\Lambda$ -module  $M$ . If  $M \in \text{mod}_\Lambda(\mathbf{d})$  for a dimension vector  $\mathbf{d}$ , then  $M(\phi)$  may be identified with the induced map

$$\text{Hom}_\Lambda(\phi, M) : \text{Hom}_\Lambda(P_\Lambda(\mathbf{x}), M) \rightarrow \text{Hom}_\Lambda(P_\Lambda(\mathbf{y}), M).$$

This implies in particular that if  $\theta^\phi(\mathbf{d}) = 0$ , then  $c_{\mathbf{d}}^\phi(M) \neq 0$  if and only if  $\text{Hom}_\Lambda(\text{Coker } \phi, M) = 0$ . We associate a semi-invariant with  $\text{Coker } \phi$  (independently of  $\phi$ ). In order to simplify the presentation, we make some additional assumptions.

Let  $\Lambda$  be an algebra with Gabriel quiver  $Q$ . Moreover, let  $V$  be a  $\Lambda$ -module with projective dimension at most 1. If  $\phi : Q \rightarrow P$  is a projective presentation of  $V$  such that  $\phi$  is a monomorphism, then  $\theta^\phi = \langle \mathbf{dim} V, - \rangle_\Lambda$ , hence is independent of  $\phi$ . We denote this weight by  $\theta^V$ . Moreover, if  $\theta^V(\mathbf{d}) = 0$ , and  $\phi$  and  $\phi'$  are projective presentations of  $V$  such that  $\phi$  and  $\phi'$  are monomorphisms, then  $c_{\text{mod}_\Lambda(\mathbf{d})}^\phi$  and  $c_{\text{mod}_\Lambda(\mathbf{d})}^{\phi'}$  coincide up to a nonzero scalar. Thus we may define  $c_{\Lambda, \mathbf{d}}^V \in \mathbb{k}[\text{mod}_\Lambda(\mathbf{d})]$  by  $c_{\Lambda, \mathbf{d}}^V := c_{\text{mod}_\Lambda(\mathbf{d})}^\phi$ , where  $\phi$  is a chosen projective presentation of  $V$ . Then  $c_{\Lambda, \mathbf{d}}^V$  is a semi-invariant of weight  $\theta^V$  and  $c_{\Lambda, \mathbf{d}}^V(M) \neq 0$  if and only if  $\text{Hom}_\Lambda(V, M) = 0$  or, equivalently,  $\text{Hom}_\Lambda(M, \tau_\Lambda V) = 0$  (for the latter statement we need (1)). If  $\mathcal{C}$  is a closed  $\text{GL}(\mathbf{d})$ -invariant subset of  $\text{mod}_\Lambda(\mathbf{d})$ , then we denote by  $c_{\Lambda, \mathcal{C}}^V$  the restriction of  $c_{\Lambda, \mathbf{d}}^V$  to  $\mathcal{C}$ .

It will be often useful to associate semi-invariants to modules of projective dimension at most 1 in a “regular” way. Thus assume in addition that  $\mathbf{c}$  is a dimension vector such that  $\mathcal{P}_\Lambda(\mathbf{c}) \neq \emptyset$ , where  $\mathcal{P}_\Lambda(\mathbf{c})$  is the subset of  $\text{mod}_\Lambda(\mathbf{c})$  consisting of the modules of projective dimension at most 1. Let  $\mathbf{x}$  be a sequence of vertices

of  $Q$  such that  $P_\Lambda(\mathbf{x}) = \bigoplus_{x \in Q_0} P_\Lambda(x)^{c(x)}$ . Since there exists an epimorphism  $P_\Lambda(\mathbf{x}) \twoheadrightarrow V$  for each  $\Lambda$ -module  $V$  with dimension vector  $\mathbf{c}$  and  $\mathcal{P}_\Lambda(\mathbf{c}) \neq \emptyset$ , there exists a sequence  $\mathbf{y}$  of vertices of  $Q$  such that  $\mathbf{dim} P_\Lambda(\mathbf{x}) - \mathbf{dim} P_\Lambda(\mathbf{y}) = \mathbf{c}$ . Let  $\mathcal{X}_\Lambda(\mathbf{c}) \subseteq \mathbb{k}Q(\mathbf{x}, \mathbf{y})$  be the set of monomorphisms  $P_\Lambda(\mathbf{y}) \hookrightarrow P_\Lambda(\mathbf{x})$ . Obviously, if  $\phi \in \mathcal{X}_\Lambda(\mathbf{c})$ , then  $\text{Coker } \phi \in \mathcal{P}_\Lambda(\mathbf{c})$ . On the other hand, if  $V \in \mathcal{P}_\Lambda(\mathbf{c})$ , then there exists  $\phi \in \mathcal{X}_\Lambda(\mathbf{c})$  such that  $\text{Coker } \phi \simeq V$ . In fact we have even more:

**Lemma 4.3.** *Let  $\Lambda$  be an algebra and  $\mathbf{c}$  a dimension vector. If  $\mathcal{V}$  is a nonempty open subset of  $\mathcal{P}_\Lambda(\mathbf{c})$  and*

$$\mathcal{U} := \{\phi \in \mathcal{X}_\Lambda(\mathbf{c}) \mid \text{Coker } \phi \in \mathcal{V}\},$$

*then  $\mathcal{U}$  is a nonempty open subset of  $\mathcal{X}_\Lambda(\mathbf{c})$ .*

*Proof.* Let  $P := \bigoplus_{x \in Q_0} P_\Lambda(x)^{c(x)}$ . Let  $\mathcal{Y}_\Lambda(\mathbf{c})$  be the set of  $\psi = (\psi(x))_{x \in Q_0}$  such that, for each  $x \in Q_0$ ,  $\psi(x) : P(x) \rightarrow \mathbb{k}^{c(x)}$  is a linear map. We denote by  $\mathcal{Z}_\Lambda(\mathbf{c})$  the set of pairs  $(\phi, \psi)$  such that  $\phi \in \mathcal{X}_\Lambda(\mathbf{c})$ ,  $\psi \in \mathcal{Y}_\Lambda(\mathbf{c})$  and  $\psi \circ \phi = 0$ . If  $\pi : \mathcal{Z}_\Lambda(\mathbf{c}) \rightarrow \mathcal{X}_\Lambda(\mathbf{c})$  is the canonical projection, then  $\pi$  is a vector bundle. Consequently, if  $\mathcal{Z}'_\Lambda(\mathbf{c})$  is the set of pairs  $(\phi, \psi) \in \mathcal{Z}_\Lambda(\mathbf{c})$  such that  $\psi$  is a surjection and  $\pi'$  is the restriction of  $\pi$  to  $\mathcal{Z}'_\Lambda(\mathbf{c})$ , then  $\pi'$  is locally trivial (with fiber isomorphic to  $\text{GL}(\mathbf{c})$ ). In particular, if  $\mathcal{W}$  is an open subset of  $\mathcal{Z}'_\Lambda(\mathbf{c})$ , then  $\pi'(\mathcal{W})$  is an open subset of  $\mathcal{X}_\Lambda(\mathbf{c})$ .

There exists a regular map  $\Theta : \mathcal{Z}'_\Lambda(\mathbf{c}) \rightarrow \text{mod}_\Lambda(\mathbf{c})$  such that  $\Theta(\phi, \psi) \simeq \text{Coker } \phi$  for all  $(\phi, \psi) \in \mathcal{Z}'_\Lambda(\mathbf{c})$  (the proof is analogous to the proof of [Richmond 2001, Lemma 9], hence we omit it). Since  $\mathcal{U} = \pi(\Theta^{-1}(\mathcal{V}))$ , the claim follows.  $\square$

We will also need the following:

**Lemma 4.4.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  a  $\text{GL}(\mathbf{d})$ -invariant irreducible closed subset of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\mathbf{c}$  a dimension vector such that  $\mathcal{P}_\Lambda(\mathbf{c}) \neq \emptyset$ .*

(1) *If  $\mathcal{U}$  is a nonempty open subset of  $\mathcal{X}_\Lambda(\mathbf{c})$ , then*

$$\text{span}\{c_{\mathcal{C}}^\phi \mid \phi \in \mathcal{U}\} = \text{span}\{c_{\mathcal{C}}^\phi \mid \phi \in \mathcal{X}_\Lambda(\mathbf{c})\}.$$

(2) *If  $\mathcal{V}$  is a nonempty open subset of  $\mathcal{P}_\Lambda(\mathbf{c})$ , then*

$$\text{span}\{c_{\Lambda, \mathcal{C}}^V \mid V \in \mathcal{V}\} = \text{span}\{c_{\Lambda, \mathcal{C}}^V \mid V \in \mathcal{P}_\Lambda(\mathbf{c})\}.$$

*Proof.* Using Lemma 4.3, it is sufficient to prove the first assertion. Let  $\phi_1, \dots, \phi_m \in \mathcal{X}_\Lambda(\mathbf{c})$  be such that  $c_{\mathcal{C}}^{\phi_1}, \dots, c_{\mathcal{C}}^{\phi_m}$  form a basis of  $\text{span}\{c_{\mathcal{C}}^\phi \mid \phi \in \mathcal{X}_\Lambda(\mathbf{c})\}$ . There exist  $M_1, \dots, M_m \in \mathcal{C}$  such that

$$\det[c_{\mathcal{C}}^{\phi_i}(M_j)]_{1 \leq i, j \leq m} \neq 0.$$

It suffices to show there exist  $\psi_1, \dots, \psi_m \in \mathcal{U}$  such that

$$\det[c_{\mathcal{C}}^{\psi_i}(M_j)]_{1 \leq i, j \leq m} \neq 0.$$

However, the regular function

$$\phi : \mathcal{X}_\Lambda(\mathbf{c})^m \rightarrow \mathbb{k}, \quad (\psi_1, \dots, \psi_m) \mapsto \det[c_\mathcal{C}^{\psi_i}(M_j)]_{1 \leq i, j \leq m},$$

is not a zero function, hence the claim follows.  $\square$

Now we use the above construction to describe generating sets of semi-invariants. We present two such sets. Depending on the situation, it will be more convenient to use one of them.

Let  $\Lambda$  be an algebra with Gabriel quiver  $Q$ ,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . There exists unique  $\mathbf{c}_\theta \in \mathbb{Z}^{Q_0}$  such that  $\theta = \langle \mathbf{c}_\theta, - \rangle_{\mathbb{k}Q}$ . Since  $\text{SI}[\text{rep}_Q(\mathbf{d})]_\theta \neq 0$ , we may assume that  $\mathbf{c}_\theta$  is a dimension vector. We explain this more precisely.

If  $\theta'$  and  $\theta''$  are weights, then  $\text{SI}[\text{rep}_Q(\mathbf{d})]_{\theta'}$  and  $\text{SI}[\text{rep}_Q(\mathbf{d})]_{\theta''}$  are equal and both nonzero if and only if  $\theta'$  and  $\theta''$  are  $\mathbf{d}$ -equivalent, i.e.,  $\theta'(x) = \theta''(x)$  for all  $x \in (\text{supp } \mathbf{d})_0$ . Now [Derksen and Weyman 2000, Theorem 1] (see also [Schofield and van den Bergh 2001, Theorem 2.3]) implies that there exists a dimension vector  $\mathbf{c}$  such that the weights  $\theta$  and  $\langle \mathbf{c}, - \rangle_{\mathbb{k}Q}$  are  $\mathbf{d}$ -equivalent. Consequently, we may assume that we only consider weights of this form.

It is clear that  $\mathcal{P}_{\mathbb{k}Q}(\mathbf{c}_\theta) \neq \emptyset$  (the category  $\text{rep } Q$  is hereditary); hence also  $\mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta) \neq \emptyset$ . It follows from [Chindris 2009, Corollary 2.5] that the semi-invariants  $c_\mathbf{d}^\phi$ , for  $\phi \in \mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta)$ , span  $\text{SI}[\text{rep}_Q(\mathbf{d})]_\theta$ . Since  $\mathcal{C}$  is a closed  $\text{GL}(\mathbf{d})$ -invariant subset of  $\text{rep}_Q(\mathbf{d})$  and  $\text{char } \mathbb{k} = 0$ , it follows that the semi-invariants  $c_\mathcal{C}^\phi$ ,  $\phi \in \mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta)$ , span  $\text{SI}[\mathcal{C}]_\theta$ .

We list some consequences:

**Lemma 4.5.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . If  $\mathcal{C}'$  is a summand of  $\mathcal{C}$ , then  $\text{SI}[\mathcal{C}']_\theta \neq 0$ .*

*Proof.* Write  $\mathcal{C} = \mathcal{C}' \oplus \mathcal{C}''$ . By assumption there exist  $\phi \in \mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta)$  and  $M \in \mathcal{C}$  such that  $c_\mathcal{C}^\phi(M) \neq 0$ . Without loss of generality we may assume that  $M = M' \oplus M''$  for  $M' \in \mathcal{C}'$  and  $M'' \in \mathcal{C}''$ . Lemma 4.1(1) implies that  $\theta(\dim M_1) = 0 = \theta(\dim M_2)$ . Consequently, Lemma 4.1(2) implies that  $c_\mathcal{C}^\phi(M) = c_{\mathcal{C}'}^\phi(M') \cdot c_{\mathcal{C}''}^\phi(M'')$ . In particular,  $c_{\mathcal{C}'}^\phi(M') \neq 0$ .  $\square$

**Lemma 4.6.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . Then there exists an open subset  $\mathcal{U}$  of  $\mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta)$  such that  $c_\mathcal{C}^\phi \neq 0$  for all  $\phi \in \mathcal{U}$ .*

*Proof.* There exist  $\phi_0 \in \mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta)$  and  $M \in \mathcal{C}$  such that  $c_\mathcal{C}^{\phi_0}(M) \neq 0$ . We define a function  $\Phi : \mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta) \rightarrow \mathbb{k}$  by

$$\Phi(\phi) := c_\mathcal{C}^\phi(M) \quad (\phi \in \mathcal{X}_{\mathbb{k}Q}(\mathbf{c}_\theta)).$$

This is a regular function and we take  $\mathcal{U} := \Phi^{-1}(\mathbb{k}^\times)$ . □

Now we present the second construction. As above let  $\Lambda$  be an algebra with Gabriel quiver  $\mathcal{Q}$ ,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . Next, let  $\Lambda_{\mathcal{C}} := \Lambda / \text{Ann } \mathcal{C}$ , where  $\text{Ann } \mathcal{C} := \bigcap_{M \in \mathcal{C}} \text{Ann } M$ . Consequently,  $\mathcal{C}$  is a faithful irreducible component of  $\text{mod}_{\Lambda_{\mathcal{C}}}(\mathbf{d})$ . Again there exists  $\mathbf{c}_{\theta, \mathcal{C}} \in \mathbb{Z}^{\mathcal{Q}_0}$  such that  $\theta = \langle \mathbf{c}_{\theta, \mathcal{C}}, - \rangle_{\Lambda_{\mathcal{C}}}$ . Since  $\text{SI}[\mathcal{C}]_\theta \neq 0$ ,  $\mathbf{c}_{\theta, \mathcal{C}}$  is a dimension vector and  $\mathcal{P}_{\mathcal{C}}(\theta) := \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_{\theta, \mathcal{C}}) \neq \emptyset$  (see [Derksen and Weyman 2002, Theorem 1]). Moreover, [Derksen and Weyman 2002, Theorem 1] also says that  $\text{SI}[\mathcal{C}]_\theta$  is spanned by the semi-invariants  $c_{\Lambda_{\mathcal{C}}, \mathcal{C}}^V$ ,  $V \in \mathcal{P}_{\mathcal{C}}(\theta)$ .

It is known that  $\overline{\mathcal{P}}_{\mathcal{C}}(\theta)$  is an irreducible component of  $\text{mod}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_{\theta, \mathcal{C}})$  [Barot and Schröer 2001, Proposition 3.1]. It is quite easy to observe that the generic decomposition of  $\overline{\mathcal{P}}_{\mathcal{C}}(\theta)$  is of the form

$$\overline{\mathcal{P}}_{\mathcal{C}}(\theta) = \overline{\mathcal{P}}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_1) \oplus \cdots \oplus \overline{\mathcal{P}}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_n)$$

for some dimension vectors  $\mathbf{c}_1, \dots, \mathbf{c}_n$  such that  $\mathbf{c}_{\theta, \mathcal{C}} = \mathbf{c}_1 + \cdots + \mathbf{c}_n$ . Obviously  $\mathbf{c}_1, \dots, \mathbf{c}_n$  are the generic summands of  $\mathbf{c}_{\theta, \mathcal{C}}$  (at  $\overline{\mathcal{P}}_{\mathcal{C}}(\theta)$ ). If we put  $\theta_i := \langle \mathbf{c}_i, - \rangle_{\Lambda_{\mathcal{C}}}$ , then we call the presentation

$$\theta = \theta_1 + \cdots + \theta_n$$

the generic decomposition of  $\theta$  at  $\mathcal{C}$ .

As a first consequence we get the following:

**Lemma 4.7.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . If  $\theta = \theta_1 + \cdots + \theta_n$  is the generic decomposition of  $\theta$  at  $\mathcal{C}$ , then the image of the map*

$$\text{SI}[\mathcal{C}]_{\theta_1} \times \cdots \times \text{SI}[\mathcal{C}]_{\theta_n} \rightarrow \text{SI}[\mathcal{C}]_\theta, \quad (f_1, \dots, f_n) \mapsto f_1 \cdots f_n,$$

spans  $\text{SI}[\mathcal{C}]_\theta$ . In particular,  $\text{SI}[\mathcal{C}]_{\theta_i} \neq 0$  for each  $i \in [1, n]$ .

*Proof.* Let  $\mathbf{c}_1, \dots, \mathbf{c}_n$  be the dimension vectors corresponding to the weights  $\theta_1, \dots, \theta_n$ , respectively, in the sense explained above. The set  $\mathcal{V}$  of  $V \in \mathcal{P}_{\mathcal{C}}(\theta)$  such that  $V \simeq V_1 \oplus \cdots \oplus V_n$  for  $V_1 \in \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_1), \dots, V_n \in \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_n)$  contains an open subset of  $\mathcal{P}_{\mathcal{C}}(\theta)$ . Lemma 4.4(2) implies that  $\text{SI}[\mathcal{C}]_\theta$  is spanned by the semi-invariants  $c_{\Lambda_{\mathcal{C}}, \mathcal{C}}^{V_1 \oplus \cdots \oplus V_n}$ ,  $V_1 \in \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_1), \dots, V_n \in \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_n)$ . Moreover, Lemma 4.2 implies that

$$c_{\Lambda_{\mathcal{C}}, \mathcal{C}}^{V_1 \oplus \cdots \oplus V_n} = c_{\Lambda_{\mathcal{C}}, \mathcal{C}}^{V_1} \cdots c_{\Lambda_{\mathcal{C}}, \mathcal{C}}^{V_n}$$

for all  $V_1 \in \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_1), \dots, V_n \in \mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}_n)$ , hence the claim follows. □

As a next consequence we obtain the following useful fact:

**Lemma 4.8.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . If  $\mathcal{P}_{\mathcal{C}}(\theta)$  contains a dense orbit, then  $\dim_{\mathbb{k}} \text{SI}[\mathcal{C}]_\theta = 1$ .*

*Proof.* If  $\mathcal{O}(V)$  is a dense orbit in  $\mathcal{P}_{\mathcal{C}}(\theta)$ , then Lemma 4.4(2) implies that  $\text{SI}[\mathcal{C}]_\theta$  is spanned by the semi-invariant  $c_{\Lambda_{\mathcal{C}}, \mathcal{C}}^V$ , hence the claim follows.  $\square$

Consequently, we get the following:

**Corollary 4.9.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . If  $\mathbf{c}'$  is a generic summand of  $\mathbf{c}_{\theta, \mathcal{C}}$  at  $\overline{\mathcal{P}}_{\mathcal{C}}(\theta)$  such that  $\mathcal{P}_{\Lambda_{\mathcal{C}}}(\mathbf{c}')$  contains a dense orbit, then*

$$\text{SI}[\mathcal{C}]_\theta \simeq \text{SI}[\mathcal{C}]_{\theta - \theta'},$$

where  $\theta' := \langle \mathbf{c}', - \rangle_{\Lambda_{\mathcal{C}}}$ .

*Proof.* Lemma 4.7 implies that  $\text{SI}[\mathcal{C}]_{\theta'} \neq 0$ . Together with Lemma 4.8 this implies that  $\dim_{\mathbb{k}} \text{SI}[\mathcal{C}]_{\theta'} = 1$ . Fix a nonzero semi-invariant  $f \in \text{SI}[\mathcal{C}]_{\theta'}$ . Lemma 4.7 implies that the map

$$\text{SI}[\mathcal{C}]_{\theta - \theta'} \rightarrow \text{SI}[\mathcal{C}]_\theta, \quad c \mapsto f \cdot c,$$

is surjective. Since  $\mathcal{C}$  is irreducible, this is also injective, and the claim follows.  $\square$

### 5. Moduli spaces

Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector, and  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ . If  $\theta$  is a weight, then a  $\Lambda$ -module  $M \in \mathcal{C}$  is called  $\theta$ -semistable if there exists  $f \in \text{SI}[\mathcal{C}]_{p\theta}$ , for some  $p \in \mathbb{N}_+$ , such that  $f(M) \neq 0$ . King [1994] has proved that  $M$  is  $\theta$ -semistable if and only if  $\theta(\dim M) = 0$  and  $\theta(\dim N) \leq 0$  for each submodule  $N$  of  $M$ . We denote by  $\mathcal{C}_\theta^{\text{ss}}$  the set of  $\theta$ -semistable  $\Lambda$ -modules in  $\mathcal{C}$ . King has also constructed a coarse moduli  $\mathcal{M}(\mathcal{C})_\theta^{\text{ss}}$  for the  $\theta$ -semistable modules in  $\mathcal{C}$  (up to an equivalence, which identifies modules which have the same simple composition factors within the category of  $\theta$ -semistable modules). By definition,

$$\mathcal{M}(\mathcal{C})_\theta^{\text{ss}} = \text{Proj} \left( \bigoplus_{p \in \mathbb{N}} \text{SI}[\mathcal{C}]_{p\theta} \right).$$

**Lemma 5.1.** *Let  $\Lambda$  be an algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\mathcal{C}_\theta^{\text{ss}} \neq \emptyset$ . If  $\mathcal{C} = \mathcal{C}_1 \oplus \mathcal{C}_2$  for irreducible components  $\mathcal{C}_1$  and  $\mathcal{C}_2$  of  $\text{mod}_\Lambda(\mathbf{d}_1)$  and  $\text{mod}_\Lambda(\mathbf{d}_2)$ , respectively, and  $\mathcal{C}_2$  is an orbit closure, then*

$$\mathcal{M}(\mathcal{C})_\theta^{\text{ss}} \simeq \mathcal{M}(\mathcal{C}_1)_\theta^{\text{ss}}.$$

*Proof.* Without loss of generality we may assume that  $\text{SI}[\mathcal{C}]_\theta \neq 0$ . Then we will show that

$$\text{SI}[\mathcal{C}]_{p\theta} \simeq \text{SI}[\mathcal{C}_1]_{p\theta}$$

for each  $p \in \mathbb{N}$ .

Let  $\mathcal{C}_2 = \overline{\mathcal{O}(M)}$ . Consider the map  $\Phi : \mathcal{C}_1 \rightarrow \mathcal{C}$  given by

$$\Phi(N) := N \oplus M \quad (N \in \mathcal{C}_1).$$

We will show that  $\Phi^* : \mathbb{k}[\mathcal{C}] \rightarrow \mathbb{k}[\mathcal{C}_1]$  induces an isomorphism  $\Phi_p^* : \text{SI}[\mathcal{C}]_{p\theta} \simeq \text{SI}[\mathcal{C}_1]_{p\theta}$  for each  $p \in \mathbb{N}$ .

Fix  $p \in \mathbb{N}$ . Since  $\text{GL}(\mathbf{d}) \times (\mathcal{C}_1 \oplus \{M\})$  is a dense subset of  $\mathcal{C}$ , it is clear that  $\Phi_p^*$  is a monomorphism. Thus it remains to show that  $\Phi_p^*$  is an epimorphism. Let  $\mathcal{X} := \mathcal{X}_{\mathbb{k}Q}(p\theta)$ , where  $Q$  is the Gabriel quiver of  $\Lambda$ . Using [Lemma 4.4\(1\)](#), it suffices to show that there exists an open subset  $\mathcal{U}$  of  $\mathcal{X}$  such that  $c_{\mathcal{C}_1}^\phi$  is in the image of  $\Phi_p^*$  for each  $\phi \in \mathcal{U}$ .

It follows from [Lemma 4.5](#) that  $\text{SI}[\mathcal{C}_2]_{p\theta} \neq 0$ . Using [Lemma 4.6](#), we obtain that there exists an open subset  $\mathcal{U}$  of  $\mathcal{X}$  such that  $c_{\mathcal{C}_2}^\phi \neq 0$  for each  $\phi \in \mathcal{U}$ . In particular,  $c_{\mathcal{C}_2}^\phi(M) \neq 0$  for each  $\phi \in \mathcal{U}$ . Now it follows from [Lemma 4.1](#) that

$$c_{\mathcal{C}_1}^\phi = \Phi_p^* \left( \frac{1}{c_{\mathcal{C}_2}^\phi(M)} c_{\mathcal{C}}^\phi \right)$$

for each  $\phi \in \mathcal{U}$ . □

### 6. Moduli spaces for the tame quasitilted algebras

The aim of this section is to prove [Theorem 2](#). Let  $\Lambda$  be a tame quasitilted algebra,  $\mathbf{d}$  a dimension vector,  $\mathcal{C}$  an irreducible component of  $\text{mod}_\Lambda(\mathbf{d})$ , and  $\theta$  a weight such that  $\mathcal{C}_\theta^{\text{ss}} \neq \emptyset$ . We show that  $\mathcal{M}(\mathcal{C})_\theta^{\text{ss}}$  is a product of projective spaces. Let  $\Lambda' := \Lambda / \text{Ann } \mathcal{C}$ .

We know that there exist Schur roots  $\mathbf{d}_1, \dots, \mathbf{d}_n$  such that

$$\mathcal{C} = \mathcal{C}_1 \oplus \dots \oplus \mathcal{C}_n,$$

where  $\mathcal{C}_i := \mathcal{C}(\mathbf{d}_i)$ ,  $i \in [1, n]$ . Using [Lemma 5.1](#), we may assume that  $\mathbf{d}_1, \dots, \mathbf{d}_n$  are isotropic.

Now let  $\mathbf{c}_1, \dots, \mathbf{c}_m$  be the generic summands of  $\mathbf{c}_{\theta, \mathcal{C}}$  at  $\overline{\mathcal{P}}_{\Lambda'}(\mathbf{c}_{\theta, \mathcal{C}})$ . Using [Corollary 4.9](#), we may assume that for each  $j \in [1, m]$ ,  $\mathcal{P}_{\Lambda'}(\mathbf{c}_j)$  does not contain a dense orbit. Since  $\overline{\mathcal{P}}_{\Lambda'}(\mathbf{c}_j)$  is an indecomposable irreducible component, this implies that there exist infinitely many indecomposable  $\Lambda'$ -modules of dimension vector  $\mathbf{c}_j$ , for each  $j \in [1, m]$ . This also means that, for each  $j \in [1, m]$ , there exist infinitely many indecomposable  $\Lambda$ -modules of dimension vector  $\mathbf{c}_j$ , hence

$\mathbf{c}_1, \dots, \mathbf{c}_m$  are isotropic roots of  $\chi_\Lambda$  (using more detailed knowledge of  $\text{mod } \Lambda$  one could also show that they are Schur roots, but we will not use this).

Before we formulate the next lemma let us recall that if  $i \in [1, n]$ ,  $j \in [1, m]$ ,  $M \in \mathcal{C}_i$  and  $V \in \mathcal{P}_{\Lambda'}(\mathbf{c}_j)$ , then  $c_{\mathcal{C}_i}^V(M) = 0$  if and only if  $\text{Hom}_\Lambda(V, M) \neq 0$  or  $\text{Hom}_\Lambda(M, \tau_{\Lambda'} V) \neq 0$ .

**Lemma 6.1.** *In the above notation, either  $\mathbf{c}_j$  is a multiplicity of  $\mathbf{d}_i$  or the intersection  $\text{supp } \mathbf{d}_i \cap \text{supp } \mathbf{c}_j$  is empty for all  $i \in [1, n]$  and  $j \in [1, m]$ .*

*Proof.* Fix  $i \in [1, n]$  and  $j \in [1, m]$ . Note that Lemmas 4.5 and 4.7 imply that  $\text{SI}[\mathcal{C}_i]_{\theta'} \neq 0$ , where  $\theta' := \langle \mathbf{c}_j, - \rangle_{\Lambda'}$ . Assume that neither  $\mathbf{c}_j$  is a multiplicity of  $\mathbf{d}_i$  nor  $\text{supp } \mathbf{d}_i \cap \text{supp } \mathbf{c}_j = \emptyset$ . Then Section 2 implies that one of the following holds:

- (1)  $\text{Hom}_\Lambda(V, M) \neq 0$  for each indecomposable  $\Lambda$ -module  $M$  with dimension vector  $\mathbf{d}_i$  and each indecomposable  $\Lambda$ -module  $V$  with dimension vector  $\mathbf{c}_j$ .
- (2)  $\text{Hom}_\Lambda(M, V) \neq 0$  for each indecomposable  $\Lambda$ -module  $M$  with dimension vector  $\mathbf{d}_i$  and each indecomposable  $\Lambda$ -module  $V$  with dimension vector  $\mathbf{c}_j$ .

In the first case we immediately obtain that  $\text{SI}[\mathcal{C}_i]_{\theta'} = 0$ , a contradiction. We show that we get the same conclusion in the second case.

Since  $\Lambda$ , hence also  $\Lambda'$ , are tame and there are infinitely many indecomposable  $\Lambda'$ -modules in  $\mathcal{P}_{\Lambda'}(\mathbf{c}_j)$ , [Crawley-Boevey 1988, Theorem D] implies that there is a nonempty open subset  $\mathcal{V}$  of  $\mathcal{P}_{\Lambda'}(\mathbf{c}_j)$  such that  $\tau_{\Lambda'} V \simeq V$  for each  $V \in \mathcal{V}$ . In particular,  $\mathbf{dim } \tau_{\Lambda'} V = \mathbf{c}_j$  for each  $V \in \mathcal{V}$ . Thus (2) together with Lemma 4.4(2) implies that  $\text{SI}[\mathcal{C}_i]_{\theta'} = 0$ , and this finishes the proof.  $\square$

Let  $I'$  be the set of  $i \in [1, n]$  such that there exists  $j \in [1, m]$  with  $\mathbf{c}_j$  a multiplicity of  $\mathbf{d}_i$ . Let  $I'' := [1, n] \setminus I'$ . Lemma 6.1 implies that  $\text{supp } \mathbf{d}_p \cap \text{supp } \mathbf{d}_q = \emptyset$  if  $p \in I'$  and  $q \in I''$  (since  $\text{supp } \mathbf{d}_p = \text{supp } \mathbf{c}_j$  for some  $j \in J$ ). Thus  $\mathcal{C} = \mathcal{C}_1 \times \mathcal{C}_2$ , where

$$\mathcal{C}_1 := \bigoplus_{i \in I'} \mathcal{C}(\mathbf{d}_i) \quad \text{and} \quad \mathcal{C}_2 := \bigoplus_{i \in I''} \mathcal{C}(\mathbf{d}_i).$$

Consequently,

$$\mathcal{M}(\mathcal{C})_{\theta}^{\text{ss}} = \mathcal{M}(\mathcal{C}_1)_{\theta}^{\text{ss}} \times \mathcal{M}(\mathcal{C}_2)_{\theta}^{\text{ss}}.$$

If  $I' \neq [1, n] \neq I''$ , then we get Theorem 2 by induction. Hence we have only two cases to consider: either  $I' = [1, n]$  or  $I'' = [1, n]$ .

First assume that  $I'' = [1, n]$ . Since  $\mathcal{C}$  is a faithful component over  $\Lambda'$ ,  $\mathbf{d}$  is a sincere dimension vector over  $\Lambda'$ . On the other hand,  $\text{supp } \mathbf{c}_j \cap \text{supp } \mathbf{d} = \emptyset$ , hence  $\mathbf{c}_j = 0$ . Consequently,  $\theta = 0$ . Thus

$$\text{SI}[\mathcal{C}]_{p\theta} = \mathbb{k}[\mathcal{C}]^{\text{GL}(\mathbf{d})} = \mathbb{k}$$

for each  $p \in \mathbb{N}$ , and

$$\mathcal{M}(\mathcal{C})_{\theta}^{\text{ss}} = \text{Proj}(\mathbb{k}[T]) = \{*\}.$$



Now assume that  $I' = [1, n]$ . We can make another reduction in this case. Let  $\mathbf{d}'_1, \dots, \mathbf{d}'_l$  be the pairwise different vectors among  $\mathbf{d}_1, \dots, \mathbf{d}_n$ . For each  $p \in [1, l]$ , let  $I_p$  be the set of  $i \in [1, n]$  such that  $\mathbf{d}_i = \mathbf{d}'_p$ . Let  $\mathcal{C}'_p := \bigoplus_{i \in I_p} \mathcal{C}(\mathbf{d}_i)$  for  $p \in [1, l]$ . [Lemma 6.1](#) again implies that  $\text{supp } \mathbf{d}'_p \cap \text{supp } \mathbf{d}'_q = \emptyset$  if  $p, q \in [1, l]$  and  $p \neq q$ . Consequently,

$$\mathcal{C} = \mathcal{C}'_1 \times \dots \times \mathcal{C}'_l$$

and

$$\mathcal{M}(\mathcal{C})_{\theta}^{\text{ss}} = \mathcal{M}(\mathcal{C}'_1)_{\theta}^{\text{ss}} \times \dots \times \mathcal{M}(\mathcal{C}'_l)_{\theta}^{\text{ss}}.$$

If  $l > 1$ , then [Theorem 2](#) follows by induction again, thus we may assume  $l = 1$ . In this case [\[Bobiński and Skowroński 1999, Theorem 2\]](#) implies that  $\text{mod}_{\Lambda}(\mathbf{d})$  is irreducible, hence  $\mathcal{C} = \text{mod}_{\Lambda}(\mathbf{d})$ . Thus [Theorem 2](#) is a result of the following:

**Proposition 6.2.** *Let  $\Lambda$  be a tame quasitilted algebra and let  $\mathbf{h}$  be an isotropic Schur root of  $\chi_{\Lambda}$ . If  $n, p \in \mathbb{N}_+$ , then*

$$\mathcal{M}(\text{mod}_{\Lambda}(n\mathbf{h}))_{p\langle \mathbf{h}, - \rangle_{\Lambda}}^{\text{ss}} \simeq \mathbb{P}_{\mathbb{k}}^n.$$

*Proof.* This is a part of [\[Domokos and Lenzing 2002, Theorem 7.1\]](#). One may also give a more direct proof, using a description of the semi-invariants for concealed-canonical algebras (the support of  $\mathbf{h}$  is a concealed-canonical algebra) which implies that  $\bigoplus_{q \in \mathbb{N}} \text{SI}[\text{mod}_{\Lambda}(n\mathbf{h})]_{q\langle \mathbf{h}, - \rangle_{\Lambda}}$  is the polynomial ring in  $n + 1$  variables (see [\[Bobiński 2015, Proposition 6.2\]](#)).  $\square$

## Acknowledgements

The paper was written during the author's stay at the Bielefeld University, which was supported by CRC 701. The author also acknowledges the support of National Science Center grant no. 2011/03/B/ST1/00847.

## References

- [Assem et al. 2006] I. Assem, D. Simson, and A. Skowroński, *Elements of the representation theory of associative algebras, Vol. 1: Techniques of representation theory*, London Mathematical Society Student Texts **65**, Cambridge University Press, Cambridge, 2006. [MR 2006j:16020](#) [Zbl 1092.16001](#)
- [Auslander et al. 1997] M. Auslander, I. Reiten, and S. O., *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics **36**, Cambridge University Press, Cambridge, 1997. [MR 98e:16011](#)
- [Barot and Schröer 2001] M. Barot and J. Schröer, "Module varieties over canonical algebras", *J. Algebra* **246**:1 (2001), 175–192. [MR 2003e:16013](#) [Zbl 1036.16010](#)
- [Bobiński 2015] G. Bobiński, "Semi-invariants for concealed-canonical algebras", *J. Pure Appl. Algebra* **219**:1 (2015), 59–76. [MR 3240823](#) [Zbl 06334325](#)
- [Bobiński and Skowroński 1999] G. Bobiński and A. Skowroński, "Geometry of modules over tame quasi-tilted algebras", *Colloq. Math.* **79**:1 (1999), 85–118. [MR 2000i:14067](#) [Zbl 0994.16009](#)

- [Carroll and Chindris 2012] A. T. Carroll and C. Chindris, “On the invariant theory for acyclic gentle algebras”, preprint, 2012. [arXiv 1210.3579](#)
- [Chindris 2009] C. Chindris, “Orbit semigroups and the representation type of quivers”, *J. Pure Appl. Algebra* **213**:7 (2009), 1418–1429. [MR 2010a:16024](#) [Zbl 1207.16012](#)
- [Chindris 2011] C. Chindris, “Geometric characterizations of the representation type of hereditary algebras and of canonical algebras”, *Adv. Math.* **228**:3 (2011), 1405–1434. [MR 2012h:16033](#) [Zbl 1252.16014](#)
- [Chindris 2013] C. Chindris, “On the invariant theory for tame tilted algebras”, *Algebra Number Theory* **7**:1 (2013), 193–214. [MR 3037894](#) [Zbl 06167117](#)
- [Crawley-Boevey 1988] W. W. Crawley-Boevey, “On tame algebras and bocses”, *Proc. London Math. Soc.* (3) **56**:3 (1988), 451–483. [MR 89c:16028](#) [Zbl 0661.16026](#)
- [Crawley-Boevey and Schröer 2002] W. Crawley-Boevey and J. Schröer, “Irreducible components of varieties of modules”, *J. Reine Angew. Math.* **553** (2002), 201–220. [MR 2004a:16020](#) [Zbl 1062.16019](#)
- [Derksen and Weyman 2000] H. Derksen and J. Weyman, “Semi-invariants of quivers and saturation for Littlewood–Richardson coefficients”, *J. Amer. Math. Soc.* **13**:3 (2000), 467–479. [MR 2001g:16031](#) [Zbl 0993.16011](#)
- [Derksen and Weyman 2002] H. Derksen and J. Weyman, “Semi-invariants for quivers with relations”, *J. Algebra* **258**:1 (2002), 216–227. [MR 2003m:16018](#) [Zbl 1048.16005](#)
- [Domokos 2002] M. Domokos, “Relative invariants for representations of finite dimensional algebras”, *Manuscripta Math.* **108**:1 (2002), 123–133. [MR 2003d:16017](#) [Zbl 1031.16014](#)
- [Domokos 2011] M. Domokos, “On singularities of quiver moduli”, *Glasg. Math. J.* **53**:1 (2011), 131–139. [MR 2012a:16029](#) [Zbl 1241.16010](#)
- [Domokos and Lenzing 2002] M. Domokos and H. Lenzing, “Moduli spaces for representations of concealed-canonical algebras”, *J. Algebra* **251**:1 (2002), 371–394. [MR 2003d:16016](#) [Zbl 1013.16006](#)
- [Drozd 1980] J. A. Drozd, “Tame and wild matrix problems”, pp. 242–258 in *Representation theory, II: Proceedings of the Second International Conference on Representations of Algebras* (Ottawa, 1979), edited by V. Dlab and P. Gabriel, Lecture Notes in Math. **832**, Springer, Berlin, 1980. [MR 83b:16024](#) [Zbl 0457.16018](#)
- [Happel 2001] D. Happel, “A characterization of hereditary categories with tilting object”, *Invent. Math.* **144**:2 (2001), 381–398. [MR 2002a:18014](#) [Zbl 1015.18006](#)
- [Happel et al. 1996] D. Happel, I. Reiten, and S. O., *Tilting in abelian categories and quasitilted algebras*, Memoirs of the American Mathematical Society **575**, Amer. Math. Soc., Providence, RI, 1996. [MR 97j:16009](#) [Zbl 0849.16011](#)
- [Kerner 1989] O. Kerner, “Tilting wild algebras”, *J. London Math. Soc.* (2) **39**:1 (1989), 29–47. [MR 90d:16025](#) [Zbl 0675.16013](#)
- [King 1994] A. D. King, “Moduli of representations of finite-dimensional algebras”, *Quart. J. Math. Oxford Ser.* (2) **45**:180 (1994), 515–530. [MR 96a:16009](#) [Zbl 0837.16005](#)
- [Lenzing and Skowroński 1996] H. Lenzing and A. Skowroński, “Quasi-tilted algebras of canonical type”, *Colloq. Math.* **71**:2 (1996), 161–181. [MR 97j:16019](#) [Zbl 0870.16007](#)
- [Meltzer 1996] H. Meltzer, “Auslander–Reiten components for concealed-canonical algebras”, *Colloq. Math.* **71**:2 (1996), 183–202. [MR 98b:16014](#) [Zbl 0923.16016](#)
- [de la Peña 1991] J. A. de la Peña, “On the dimension of the module-varieties of tame and wild algebras”, *Comm. Algebra* **19**:6 (1991), 1795–1807. [MR 92i:16016](#) [Zbl 0818.16013](#)

- [Richmond 2001] N. J. Richmond, “A stratification for varieties of modules”, *Bull. London Math. Soc.* **33**:5 (2001), 565–577. [MR 2002d:16017](#) [Zbl 1054.16007](#)
- [Ringel 1984] C. M. Ringel, *Tame algebras and integral quadratic forms*, Lecture Notes in Mathematics **1099**, Springer, Berlin, 1984. [MR 87f:16027](#) [Zbl 0546.16013](#)
- [Schofield 1991] A. Schofield, “Semi-invariants of quivers”, *J. London Math. Soc.* (2) **43**:3 (1991), 385–395. [MR 92g:16019](#) [Zbl 0779.16005](#)
- [Schofield and van den Bergh 2001] A. Schofield and M. van den Bergh, “Semi-invariants of quivers for arbitrary dimension vectors”, *Indag. Math. (N.S.)* **12**:1 (2001), 125–138. [MR 2003e:16016](#) [Zbl 1004.16012](#)
- [Skowroński 1998] A. Skowroński, “Tame quasi-tilted algebras”, *J. Algebra* **203**:2 (1998), 470–490. [MR 99b:16019](#) [Zbl 0908.16013](#)
- [Skowroński 2006] A. Skowroński, “Selfinjective algebras: finite and tame type”, pp. 169–238 in *Trends in representation theory of algebras and related topics* (Querétaro, 2004), edited by J. A. de la Peña and R. Bautista, Contemp. Math. **406**, Amer. Math. Soc., Providence, RI, 2006. [MR 2007f:16045](#) [Zbl 1129.16013](#)
- [Skowroński and Weyman 2000] A. Skowroński and J. Weyman, “The algebras of semi-invariants of quivers”, *Transform. Groups* **5**:4 (2000), 361–402. [MR 2001m:16017](#) [Zbl 0986.16004](#)
- [Skowroński and Zwara 1998] A. Skowroński and G. Zwara, “Degenerations for indecomposable modules and tame algebras”, *Ann. Sci. École Norm. Sup.* (4) **31**:2 (1998), 153–180. [MR 99k:16032](#) [Zbl 0915.16011](#)

Communicated by David Benson

Received 2013-12-19

Revised 2014-02-14

Accepted 2014-06-14

[gregbob@mat.umk.pl](mailto:gregbob@mat.umk.pl)

*Faculty of Mathematics and Computer Science,  
Nicolaus Copernicus University, Ulica Chopina 12/18,  
87-100 Toruń, Poland*

# Algebra & Number Theory

[msp.org/ant](http://msp.org/ant)

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
Dave Benson	University of Aberdeen, Scotland	Raman Parimala	Emory University, USA
Richard E. Borcherds	University of California, Berkeley, USA	Jonathan Pila	University of Oxford, UK
John H. Coates	University of Cambridge, UK	Anand Pillay	University of Notre Dame, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Victor Reiner	University of Minnesota, USA
Brian D. Conrad	University of Michigan, USA	Peter Sarnak	Princeton University, USA
Hélène Esnault	Freie Universität Berlin, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Bernd Sturmfels	University of California, Berkeley, USA
Roger Heath-Brown	Oxford University, UK	Richard Taylor	Harvard University, USA
Craig Huneke	University of Virginia, USA	Ravi Vakil	Stanford University, USA
Yujiro Kawamata	University of Tokyo, Japan	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Yuri Manin	Northwestern University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Barry Mazur	Harvard University, USA	Efim Zelmanov	University of California, San Diego, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Shou-Wu Zhang	Princeton University, USA
Susan Montgomery	University of Southern California, USA		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2014 is US \$225/year for the electronic version, and \$400/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 8    No. 6    2014

---

Decompositions of commutative monoid congruences and binomial ideals THOMAS KAHLE and EZRA MILLER	1297
Locally analytic representations and sheaves on the Bruhat–Tits building DEEPAM PATEL, TOBIAS SCHMIDT and MATTHIAS STRAUCH	1365
Complétés universels de représentations de $GL_2(\mathbb{Q}_p)$ PIERRE COLMEZ and GABRIEL DOSPINESCU	1447
On moduli spaces for quasitilted algebras GRZEGORZ BOBIŃSKI	1521