

# *Algebra & Number Theory*

Volume 10

2016

No. 7

**Local deformation rings for  $GL_2$  and a  
Breuil–Mézard conjecture when  $\ell \neq p$**

Jack Shotton



# Local deformation rings for $\mathrm{GL}_2$ and a Breuil–Mézard conjecture when $\ell \neq p$

Jack Shotton

We compute the deformation rings of two dimensional mod  $l$  representations of  $\mathrm{Gal}(\bar{F}/F)$  with fixed inertial type for  $l$  an odd prime,  $p$  a prime distinct from  $l$ , and  $F/\mathbb{Q}_p$  a finite extension. We show that in this setting an analogue of the Breuil–Mézard conjecture holds, relating the special fibres of these deformation rings to the mod  $l$  reduction of certain irreducible representations of  $\mathrm{GL}_2(\mathcal{O}_F)$ .

## 1. Introduction

Let  $p$  be a prime and let  $F$  be a finite extension of  $\mathbb{Q}_p$  with absolute Galois group  $G_F$ . We study the (framed) deformation rings for two-dimensional mod  $l$  representations of  $G_F$ , where  $l$  is an odd prime distinct from  $p$ . More specifically, let  $E$  be a finite extension of  $\mathbb{Q}_l$  with ring of integers  $\mathcal{O}$ , uniformiser  $\lambda$ , and residue field  $\mathbb{F}$ . Let

$$\bar{\rho} : G_F \rightarrow \mathrm{GL}_2(\mathbb{F})$$

be a continuous representation. Then there is a universal lifting (or framed deformation) ring  $R^\square(\bar{\rho})$  parametrising lifts of  $\bar{\rho}$ . Our main result relates congruences between irreducible components of  $\mathrm{Spec} R^\square(\bar{\rho})$  to congruences between certain representations of  $\mathrm{GL}_2(\mathcal{O}_F)$ , where  $\mathcal{O}_F$  is the ring of integers of  $F$ . Our method is to give explicit equations for the components of  $\mathrm{Spec} R^\square(\bar{\rho})$ , which may be of independent use.

If  $\tau : I_F \rightarrow \mathrm{GL}_2(E)$  is a continuous representation that extends to a representation of  $G_F$  (an *inertial type*), then we say that a representation  $\rho : G_F \rightarrow \mathrm{GL}_2(\bar{E})$  has type  $\tau$  if its restriction to  $I_F$  is isomorphic to  $\tau$ . Say that an irreducible component of  $\mathrm{Spec} R^\square(\bar{\rho})$  has type  $\tau$  if a Zariski dense subset of its  $\bar{E}$ -points correspond to representations of type  $\tau$ . We define (Definition 4.1) a formal sum  $\mathcal{C}(\bar{\rho}, \tau)$  of irreducible components of the special fibre  $\mathrm{Spec} R^\square(\bar{\rho}) \otimes_{\mathcal{O}} \mathbb{F}$ . For semisimple  $\tau$ , this is obtained as the intersection with the special fibre of those components of  $\mathrm{Spec} R^\square(\bar{\rho})$  having type  $\tau$ ; for nonsemisimple  $\tau$  this must be slightly modified.

MSC2010: 11S37.

Keywords: Galois representations, deformation rings, local Langlands, Breuil–Mézard.

To an inertial type  $\tau$  we also associate an irreducible  $E$ -representation  $\sigma(\tau)$  of  $\mathrm{GL}_2(\mathcal{O}_F)$ , by a slight variant on the definition of [Henniart 2002] (see Section 3C). For an irreducible  $\mathbb{F}$ -representation  $\theta$  of  $\mathrm{GL}_2(\mathcal{O}_F)$ , define  $m(\theta, \overline{\sigma(\tau)})$  to be the multiplicity of  $\theta$  as a Jordan–Hölder factor of the mod  $\lambda$  reduction of  $\sigma(\tau)$ . Then we can state our main theorem:

**Theorem 4.2.** *Let  $\bar{\rho} : G_F \rightarrow \mathrm{GL}_2(\mathbb{F})$  be a continuous representation. For each irreducible  $\mathbb{F}$ -representation  $\theta$  of  $\mathrm{GL}_2(\mathcal{O}_F)$ , there is a formal sum  $\mathcal{C}(\bar{\rho}, \theta)$  of irreducible components of  $\mathrm{Spec} R^\square(\bar{\rho}) \otimes \mathbb{F}$  such that, for each inertial type  $\tau$ , we have the equality*

$$\mathcal{C}(\bar{\rho}, \tau) = \sum_{\theta} m(\theta, \overline{\sigma(\tau)}) \mathcal{C}(\bar{\rho}, \theta).$$

In fact the  $\mathcal{C}(\bar{\rho}, \theta)$  are uniquely determined (at least for those  $\theta$  which actually occur in some  $\overline{\sigma(\tau)}$ ).

This theorem is an analogue for mod  $l$  representations of  $G_F$  of the Breuil–Mézard conjecture [2002], which pertains to mod  $p$  representations of  $G_{\mathbb{Q}_p}$ . Our statement is not in the language of Hilbert–Samuel multiplicities used in [Breuil and Mézard 2002], but rather in the geometric language of [Emerton and Gee 2014]. The original conjecture of Breuil and Mézard was proved in most cases by Kisin [2009a]; further cases were proved by Paškūnas [2015] by local methods, and the full conjecture was proved when  $p > 3$  in [Hu and Tan 2013]. The conjecture was generalised to  $n$ -dimensional representations of  $G_F$  in [Emerton and Gee 2014]; the only case known, outside of those just mentioned, is that of two-dimensional potentially Barsotti–Tate representations (see [Gee and Kisin 2014]).

In the  $l \neq p$  setting, a comparison of special fibres of (very particular) local deformation rings was used by Taylor [2008] to prove the change of level results needed to obtain nonminimal automorphy lifting theorems. This is another motivation for our result.

Our method of proof is to explicitly determine equations for deformation rings of fixed type and, indeed, obtaining these explicit descriptions is another goal of this paper. We reduce to the tamely ramified case, in which we use the relation

$$\phi \sigma \phi^{-1} = \sigma^q,$$

for  $\phi \in G_F$  a lift of Frobenius and  $\sigma \in I_F$  a generator of tame inertia. Since we are considering lifts  $\rho$  of fixed type, and so with fixed characteristic polynomial of  $\rho(\sigma)$ , we may use the Cayley–Hamilton theorem to reduce this equation to one of degree at most two in the entries of  $\rho(\phi)$  and  $\rho(\sigma)$ . These explicit descriptions show that the irreducible components of  $\mathrm{Spec} R^\square(\bar{\rho}) \otimes \bar{E}$  are always smooth (which is also proved in [Pilloni 2008]) and that the reduced deformation rings in which the semisimplification of the restriction to inertia is fixed are always Cohen–Macaulay

(see Section 5E). It is natural to ask whether these properties persist beyond the case of two-dimensional representations. We note that the generic fibres of our local deformation rings have been studied in [Pilloni 2008; Reduzzi 2013], but their methods say little about the integral structure.

In a forthcoming paper, we will extend Theorem 4.2 to the case of  $n$ -dimensional representations using global methods.

The structure of this paper is as follows. In Section 2 we define the universal deformation rings and show how to reduce their study to the case when  $\bar{\rho}$  is tamely ramified. We also prove some lemmas that will be useful in the calculations that follow. In Section 3 we define the deformation rings with fixed inertial type that we will need and discuss the construction of the representations  $\sigma(\tau)$ . In Section 4 we state and prove the main theorem, modulo the calculations of Section 5 and results of Section 6. Section 5 contains the calculations of explicit equations for local deformation rings, divided into cases according to the value of  $q \bmod l$ . Finally, in Section 6 we prove the results on the mod  $l$  reduction of the  $\sigma(\tau)$  that are stated in Section 3D (and used in the proof of Theorem 4.2).

## 2. Preliminaries

**2A. Fields and Galois groups.** Suppose that  $l \neq p$  are primes with  $l > 2$ .

Let  $F/\mathbb{Q}_p$  be a finite extension with ring of integers  $\mathcal{O}_F$ , maximal ideal  $\mathfrak{p}_F$ , uniformiser  $\varpi_F$ , and residue field  $k_F$  of order  $q$ . Let  $F$  have absolute Galois group  $G_F$ , inertia group  $I_F$ , and wild inertia group  $P_F$ . Let  $I_F \rightarrow I_F/\tilde{P}_F \cong \mathbb{Z}_l$  be the maximal pro- $l$  quotient of  $I_F$ , so that  $\tilde{P}_F/P_F \cong \prod_{l' \neq l, p} \mathbb{Z}_{l'}$ . Note that  $\tilde{P}_F$  is normal in  $G_F$  and write  $T_F = G_F/\tilde{P}_F$ . The short exact sequence  $1 \rightarrow I_F/\tilde{P}_F \rightarrow T_F \rightarrow G_F/I_F \rightarrow 1$  splits, so that  $T_F \cong \mathbb{Z}_l \rtimes \hat{\mathbb{Z}}$ . We fix topological generators  $\sigma$  of this  $\mathbb{Z}_l$  and  $\phi$  of this  $\hat{\mathbb{Z}}$  such that  $\phi$  is a lift of arithmetic Frobenius. Then the action of  $\hat{\mathbb{Z}}$  on  $\mathbb{Z}_l$  is given by

$$\phi \sigma \phi^{-1} = \sigma^q. \quad (1)$$

Let  $L/F$  be an unramified quadratic extension, with residue field  $k_L$ .

Now let  $E/\mathbb{Q}_l$  be a finite extension with ring of integers  $\mathcal{O}$ , residue field  $\mathbb{F}$  and uniformiser  $\lambda$ . Let  $\epsilon : G_F \rightarrow \mathbb{Z}_l^\times$  be the  $l$ -adic cyclotomic character, and let  $\mathbb{1} : G_F \rightarrow \mathbb{Z}_l^\times$  be the trivial character. If  $A$  is any  $\mathcal{O}$ -algebra then we will regard these as maps to  $A^\times$  via the structure maps  $\mathbb{Z}_l \rightarrow \mathcal{O} \rightarrow A$ .

Define two integers  $a$  and  $b$  by  $a = v_l(q - 1)$  and  $b = v_l(q + 1)$ , where  $v_l$  is the  $l$ -adic valuation; at most one of  $a$  and  $b$  is nonzero since  $l$  is odd.

**2B. Deformation rings.** Suppose that  $\bar{M}$  is an  $n$ -dimensional  $\mathbb{F}$ -vector space and that  $\bar{\rho} : G_F \rightarrow GL(\bar{M})$  is a continuous representation. Let  $(\bar{e}_i)_{i=1}^n$  be a basis for  $\bar{M}$ , so that  $\bar{\rho}$  gives a map  $\bar{\rho} : G_F \rightarrow GL_n(\mathbb{F})$ .

Let  $\mathcal{C}_{\mathcal{O}}$  denote the category of artinian local  $\mathcal{O}$ -algebras with residue field  $\mathbb{F}$ , and  $\mathcal{C}_{\mathcal{O}}^{\wedge}$  the category of complete noetherian local  $\mathcal{O}$ -algebras with residue field  $\mathbb{F}$ . If  $A$  is an object of  $\mathcal{C}_{\mathcal{O}}$  or  $\mathcal{C}_{\mathcal{O}}^{\wedge}$ , let  $\mathfrak{m}_A$  be its maximal ideal. Define two functors

$$D(\bar{\rho}), D^{\square}(\bar{\rho}) : \mathcal{C}_{\mathcal{O}} \rightarrow \mathbf{Set}$$

as follows:

- $D(\bar{\rho})(A)$  is the set of equivalence classes of  $(M, \iota)$  where  $M$  is a free rank- $n$   $A$ -module,  $\rho : G_F \rightarrow \text{Aut}_A(M)$  a continuous homomorphism, and  $\iota : M \otimes_A \mathbb{F} \xrightarrow{\sim} \bar{M}$  an isomorphism commuting with the actions of  $G_F$ .
- $D^{\square}(\bar{\rho})(A)$  is the set of equivalence classes of  $(M, \rho, (e_i)_{i=1}^n)$  where  $M$  is a free  $A$ -module of rank  $n$ ,  $\rho : G_F \rightarrow \text{Aut}_A(M)$  is a continuous homomorphism, and  $(e_i)_{i=1}^n$  is a basis of  $M$  as an  $A$ -module, such that the isomorphism  $\iota : M \otimes_A \mathbb{F} \xrightarrow{\sim} \bar{M}$  defined by  $\iota : e_i \otimes 1 \mapsto \bar{e}_i$  commutes with the actions of  $G_F$ .

In the first case,  $(M, \rho, \iota)$  and  $(M', \rho', \iota')$  are equivalent if there is an isomorphism  $\alpha : M \rightarrow M'$ , commuting with the actions of  $G_F$ , such that  $\iota = \iota' \circ \alpha$ ; in the second case,  $(M, \rho, (e_i)_i)$  and  $(M', \rho', (e'_i)_i)$  are isomorphic if the map  $M \rightarrow M'$  defined by  $e_i \mapsto e'_i$  commutes with the actions of  $G_F$ . There is a natural transformation of functors  $D^{\square}(\bar{\rho}) \rightarrow D(\bar{\rho})$  given by forgetting the basis.

Alternatively, when  $\bar{\rho}$  is regarded as a homomorphism to  $\text{GL}_n(\mathbb{F})$ , we have the equivalent definitions

$$D^{\square}(\bar{\rho})(A) = \{\rho : G_F \rightarrow \text{GL}_n(A) \mid \rho \text{ is continuous and lifts } \bar{\rho}\}$$

and

$$D(\bar{\rho})(A) = D^{\square}(\bar{\rho})(A) / \text{conjugacy by } 1 + M_n(\mathfrak{m}_A).$$

The functor  $D(\bar{\rho})$  is not usually prorepresentable, but the functor  $D^{\square}(\bar{\rho})$  always is (see, for example, [Kisin 2009b, 2.3.4]).

**Definition 2.1.** The *universal lifting ring* (or universal framed deformation ring) of  $\bar{\rho}$  is the object  $R^{\square}(\bar{\rho})$  of  $\mathcal{C}_{\mathcal{O}}^{\wedge}$  that prorepresents the functor  $D^{\square}(\bar{\rho})$ . The universal lift is denoted  $\rho^{\square} : G_F \rightarrow \text{GL}_n(R^{\square}(\bar{\rho}))$ .

We recall a useful calculation (see, e.g., [Barnet-Lamb et al. 2014, Section 1.2]):

**Lemma 2.2.** *The ring  $R^{\square}(\bar{\rho})[1/l]$  is generically formally smooth of dimension  $n^2$ .*

The next lemma enables us to reduce to the case where the residual representation is trivial on  $\tilde{P}_F$ . Suppose that  $\theta$  is an irreducible  $\mathbb{F}$ -representation of  $\tilde{P}_F$ . Then by [Clozel et al. 2008, Lemma 2.4.11] there is a lift of  $\theta$  to an  $\mathcal{O}$ -representation of  $\tilde{P}_F$ , which may be extended to an  $\mathcal{O}$ -representation  $\tilde{\theta}$  of  $G_{\theta}$ , where  $G_{\theta}$  is the group  $\{g \in G_F \mid g\theta g^{-1} \cong \theta\}$ . For each irreducible representation  $\theta$  of  $\tilde{P}_F$  we pick such

a  $\tilde{\theta}$  and a finite free  $\mathcal{O}$ -module  $N(\theta)$  on which  $\tilde{P}_F$  acts as  $\tilde{\theta}$ . If  $M$  is a set-finite  $\mathcal{O}$ -module with a continuous action  $\rho$  of  $G_F$ , then define

$$M_\theta = \text{Hom}_{\tilde{P}_F}(\tilde{\theta}, M).$$

The module  $M_\theta$  has a natural continuous action  $\rho_\theta$  of  $G_\theta$  given by  $(gf)(v) = gf(g^{-1}v)$ ; the subgroup  $\tilde{P}_F$  of  $G_\theta$  acts trivially.

**Lemma 2.3** (tame reduction). (1) *Let  $M$  be a set-finite  $\mathcal{O}$ -module with a continuous action of  $G_F$ . Then there is a natural isomorphism*

$$M = \bigoplus_{[\theta]} \text{Ind}_{G_\theta}^{G_F}(N(\theta) \otimes_{\mathcal{O}} M_\theta),$$

where  $\theta$  runs through a set of representatives for the  $G_F$ -conjugacy classes of irreducible representations of  $\tilde{P}_F$ .

(2) *The isomorphism of part (1) induces a natural isomorphism of functors:*

$$D(\bar{\rho}) \xrightarrow{\sim} \prod_{[\theta]} D(\bar{\rho}_\theta),$$

where  $\theta$  runs through a set of representatives for the  $G_F$ -conjugacy classes of irreducible representations of  $\tilde{P}_F$ .

(3) *If  $R^\square(\bar{\rho}_\theta)$  is the universal framed deformation ring for the representation  $\bar{\rho}_\theta$  of  $G_\theta/\tilde{P}_F$ , then*

$$R^\square(\bar{\rho}) \cong \left( \widehat{\bigotimes}_{[\theta]} R^\square(\bar{\rho}_\theta) \right) \llbracket X_1, \dots, X_{n^2 - \sum n_\theta^2} \rrbracket,$$

where  $n_\theta = \dim \rho_\theta$ . This isomorphism lies above  $D(\bar{\rho}) \xrightarrow{\sim} \prod_{[\theta]} D(\bar{\rho}_\theta)$ , the isomorphism of part (2).

*Proof.* The first two parts are in [Clozel et al. 2008]: part (1) is Lemma 2.4.12 and part (2) is Corollary 2.4.13. Part (3) is the refinement to framed deformations obtained by keeping track of a basis in the construction of part (1) of the proposition, as in [Choi 2009, Proposition 2.0.5].

As [Choi 2009] is not easily available, we sketch the argument for part (3). Let  $[\theta_1], [\theta_2], \dots$  be the  $G_F$ -conjugacy classes of irreducible  $\tilde{P}_F$ -representations. Pick left coset representatives  $(g_{ij})_j$  for  $G_{\theta_i}$  in  $G_F$ . Write  $N_i$  for  $N(\theta_i)$ , and choose an  $\mathcal{O}$ -basis  $(f_{ik})_k$  of  $N_i$ .

Let  $A$  be an object of  $\mathcal{C}_{\mathcal{O}}$ ,  $M$  be a free rank  $n$   $A$ -module with a continuous action of  $G_F$ , and  $M_{\theta_i}$  be as above. Given (for each  $i$ ) a basis  $(e_{il})_{l=1}^{n_{\theta_i}}$  of  $M_{\theta_i}$ , we can produce a basis  $(e_{ijkl})_{j,k,l}$  of

$$M_{\theta_i} = A[G_F] \otimes_{A[G_\theta]} (N_i \otimes_{\mathcal{O}} M_{\theta_i})$$

defined by

$$e_{ijkl} = g_{ij} \otimes f_{ik} \otimes e_{il}.$$

Then  $(e_{ijkl})_{i,j,k,l}$  is a basis of  $M$ .

Let  $\mathcal{F}(A)$  be the set of  $\mathbf{Y} = (Y_{ijkl,i'j'k'l'})$  which are  $n \times n$  matrices of elements of  $\mathfrak{m}_A$  such that

$$Y_{ijkl,i'j'k'l'} = 0 \quad \text{if } i = i' \text{ and } j = j' = k = k' = l$$

(so that  $n^2 - \sum n_{\theta_i}^2$  “free” entries of  $\mathbf{Y}$  remain). Then  $\mathcal{F}$  defines a functor on  $\mathcal{C}_{\mathcal{O}}$  prorepresented by  $\mathcal{O}[[X_1, \dots, X_{n^2 - \sum n_{\theta_i}^2}]]$  (the variables  $X$  being simply an enumeration of those  $Y_{ijkl,i'j'k'l'}$  which can be nonzero).

We then have a natural transformation of functors

$$\mathcal{F} \times \prod_{[\theta]} D^{\square}(\bar{\rho}_{\theta}) \rightarrow D^{\square}(\bar{\rho})$$

taking the tuple  $(\mathbf{Y}, (M_{\theta_i}, \rho_{\theta_i}, e_{il})_i)$  to the tuple

$$\left( \bigoplus_i \text{Ind}_{G_{\theta_i}}^{G_F}(N_i \otimes_{\mathcal{O}} M_{\theta_i}), \bigoplus_i \text{Ind}_{G_{\theta_i}}^{G_F}(\tilde{\theta}_i \otimes_{\mathcal{O}} \rho_{\theta_i}), (I_n + \mathbf{Y})(e_{ijkl})_{i,j,k,l} \right).$$

Then one can check (and this is what is done in [Choi 2009, Proposition 2.0.5]) that this is in fact an isomorphism, and so we get the claimed isomorphism of prorepresenting objects.  $\square$

### 2C. Twisting.

**Lemma 2.4.** *Suppose that  $\chi : G_F \rightarrow \mathcal{O}^{\times}$  is any character. Then there is a natural isomorphism*

$$R^{\square}(\bar{\rho}) \xrightarrow{\sim} R^{\square}(\bar{\rho} \otimes \bar{\chi}).$$

Moreover, if  $\chi_1$  and  $\chi_2$  satisfy  $\bar{\chi}_1 = \bar{\chi}_2$  then they induce the same maps

$$R^{\square}(\bar{\rho}) \otimes \mathbb{F} \xrightarrow{\sim} R^{\square}(\bar{\rho} \otimes \bar{\chi}_i) \otimes \mathbb{F}.$$

*Proof.* This follows easily from the isomorphism of functors

$$D^{\square}(\bar{\rho}) \rightarrow D^{\square}(\bar{\rho} \otimes \bar{\chi})$$

given by tensoring with  $\chi$  (remembering that we are considering  $\mathcal{O}$ -algebras). For the last statement, observe that if the functors are restricted to  $\mathbb{F}$ -algebras then the isomorphism only depends on  $\bar{\chi}$ .  $\square$

Since every  $\mathbb{F}$ -valued character lifts to  $\mathcal{O}$  (using the Teichmüller lift) this shows that  $R^{\square}(\bar{\rho}) \cong R^{\square}(\bar{\rho} \otimes \bar{\chi})$  for every  $\bar{\chi} : G_F \rightarrow \mathbb{F}^{\times}$ .

We also need the calculation of the universal deformation ring of a character, to which some of our calculations reduce. This is completely standard, but we include it as a simple illustration of the method.



**Lemma 2.5.** *Let  $\bar{\chi} : G_F \rightarrow \mathbb{F}^\times$  be a continuous character. Then*

$$R^\square(\bar{\chi}) = \frac{\mathcal{O}[[X, Y]]}{((1 + X)^{l^a} - 1)}$$

*has  $l^a$  irreducible components, indexed by the  $l^a$ -th roots of unity. They are formally smooth of relative dimension one over  $\mathcal{O}$ .*

*Proof.* By Lemma 2.4, we may take  $\bar{\chi}$  to be trivial. If  $\chi$  is any lift of  $\bar{\chi}$  to an object  $A$  of  $\mathcal{C}_{\mathcal{O}}$ , then for  $g \in \tilde{P}_F$  we must have  $\chi(g)^n = 1$  for some  $n$  coprime to  $l$ , and therefore  $\chi(g) = 1$ , so that we are reduced to considering characters of  $T_F$ . We must have that  $\chi(\sigma)^q = \chi(\sigma)$  and  $\chi(\sigma) \equiv 1 \pmod{\mathfrak{m}_A}$ , and therefore that  $\chi(\sigma)^{l^a} = 1$ . We are then free to choose  $\chi(\phi)$ . Writing  $\chi(\sigma) = 1 + X$  and  $\chi(\phi) = 1 + Y$ , we have shown that

$$D^\square(\bar{\chi})(A) = \text{Hom}_{\mathcal{C}_{\mathcal{O}}} \left( \frac{\mathcal{O}[[X, Y]]}{((1 + X)^{l^a} - 1)}, A \right)$$

functorially, and so the universal framed deformation ring is as claimed. □

**2D. Multiplicities and cycles.** Suppose that  $X$  is a noetherian scheme and that  $\mathcal{F}$  is a coherent sheaf on  $X$ . Let  $Y$  be the scheme-theoretic support of  $\mathcal{F}$ , and let  $d \geq \dim Y$ . Let  $\mathcal{Z}^d(X)$  be the free abelian group on the  $d$ -dimensional points of  $X$ ; elements of  $\mathcal{Z}^d(X)$  are called  $d$ -dimensional cycles. If  $\mathfrak{a} \in X$  is a point of dimension  $d$  write  $[\mathfrak{a}]$  for the corresponding element of  $\mathcal{Z}^d(X)$  and define the multiplicity  $e(\mathcal{F}, \mathfrak{a})$  to be the length of  $\mathcal{F}_{\mathfrak{a}}$  as an  $\mathcal{O}_{Y, \mathfrak{a}}$ -module (this is zero if  $\mathfrak{a} \notin Y$ ).

**Definition 2.6.** The cycle  $Z^d(\mathcal{F})$  associated to  $\mathcal{F}$  is the element

$$\sum_{\mathfrak{a}} e(\mathcal{F}, \mathfrak{a})[\mathfrak{a}] \in \mathcal{Z}^d(X).$$

If  $X = \text{Spec } A$  is affine and  $\mathcal{F} = \tilde{M}$  for a finitely generated  $A$ -module  $M$ , then we will write  $Z^d(M)$  for  $Z^d(\mathcal{F})$ .

If  $i : X \rightarrow X'$  is a closed immersion of  $X$  in a noetherian scheme  $X'$ , then there is a natural inclusion  $i_* : \mathcal{Z}^d(X) \rightarrow \mathcal{Z}^d(X')$  for each  $d$ . For a coherent sheaf  $\mathcal{F}$  on  $X$  whose support has dimension at most  $d$ , we then have

$$i_*(Z^d(\mathcal{F})) = Z^d(i_*(\mathcal{F})).$$

We will often use this compatibility without comment.

A cycle is *effective* if it is of the form  $\sum n_{\mathfrak{a}}[\mathfrak{a}]$  for  $n_{\mathfrak{a}} \geq 0$ . We say that an effective cycle  $C_1$  is a *subcycle* of an effective cycle  $C_2$  if  $C_2 - C_1$  is also effective.



**2E. A determinantal ring.** For  $a, b,$  and  $c$  natural numbers, if  $I$  is the ideal generated by the  $a \times a$  minors of a  $b \times c$  matrix with independent indeterminate entries over a Cohen–Macaulay ring  $A,$  then  $A/I$  is always Cohen–Macaulay (see [Eisenbud 1995, Theorem 18.18]). We include a simple proof in the very special case that we need below.

**Proposition 2.7.** *Let  $k \geq 2$  be an integer and let  $A$  be either a field or a discrete valuation ring. Let  $R = A[X_1, \dots, X_k, Y_1, \dots, Y_k]$  and let  $I \triangleleft R$  be the ideal generated by the  $2 \times 2$  minors of*

$$\begin{pmatrix} X_1 & X_2 & \cdots & X_k \\ Y_1 & Y_2 & \cdots & Y_k \end{pmatrix}.$$

*Let  $S = R/I.$  Then  $S$  is a Cohen–Macaulay domain and is flat over  $A.$  It is Gorenstein if and only if  $k = 2.$*

*The same is true if we replace  $S$  by its completion  $S^\wedge$  at the “irrelevant” ideal  $(X_1, \dots, X_k, Y_1, \dots, Y_k).$*

*Proof.* Note that  $R$  and  $S$  are naturally graded  $A$ -algebras.

Suppose that  $A$  is a field. It is easy to see that  $\text{Proj}(S)$  is a smooth irreducible projective variety over  $A$  of dimension  $k + 1$  — it is covered by the open sets  $\{X_i \neq 0\}$  and  $\{Y_i \neq 0\},$  each of which is isomorphic to  $(\mathbb{A}_A^1 \setminus \{0\}) \times \mathbb{A}_A^k.$  Thus  $S$  is a domain. We may extend  $A$  so that its cardinality is at least  $k + 1,$  and choose pairwise distinct  $\alpha_1, \dots, \alpha_k \in A^\times.$

I claim that  $(X_1 - \alpha_1 Y_1, \dots, X_k - \alpha_k Y_k, Y_1 + \dots + Y_k)$  is a regular sequence in  $S.$  To see this, observe that  $\text{Proj}(S/(X_1 - \alpha_1 Y_1, \dots, X_i - \alpha_i Y_i))$  is reduced (we may check this on the affine pieces) and that its irreducible components are all of the form

$$\text{Proj}\left(\frac{R}{(X_j - \alpha_{i_0} Y_j)_{1 \leq j \leq k} + (X_j, Y_j)_{1 \leq j \leq i, j \neq i_0}}\right),$$

for  $1 \leq i_0 \leq i$  or of the form

$$\text{Proj}(S/(X_1, \dots, X_i, Y_1, \dots, Y_i)).$$

Now it is easy to check that  $X_{i+1} - \alpha_{i+1} Y_{i+1}$  (if  $i < k$ ) or  $Y_1 + \dots + Y_k$  (if  $i = k$ ) is a nonzerodivisor on each of these components, and so is a nonzerodivisor on  $S/(X_1 - \alpha_1 Y_1, \dots, X_i - \alpha_i Y_i)$  as required.

Now

$$S/((X_i - \alpha_i Y_i)_i, Y_1 + \dots + Y_k) \cong A[Y_2, \dots, Y_k]/(Y_2, \dots, Y_k)^2$$

is Gorenstein if and only if  $k = 2,$  as required.

If  $A$  is a DVR then the following easy lemma (a specialisation of [Snowden 2011, Proposition 2.2.1]) gives the result.

**Lemma 2.8.** *If  $A$  is a DVR and  $S$  is a finitely generated  $A$ -algebra such that  $S \otimes A/\mathfrak{m}_A$  and  $S \otimes \text{Frac } A$  are domains of the same dimension, then  $S$  is flat over  $A$  (that is, a uniformiser of  $A$  is a regular parameter in  $S$ ).*

The final statement of the proposition follows from the facts that both localisation and completion preserve the properties of being Gorenstein, Cohen–Macaulay, or  $A$ -flat;  $S^\wedge$  is a domain because its associated graded ring is  $S$ , which is a domain.  $\square$

### 3. Types

#### 3A. Inertial types.

**Definition 3.1.** An *inertial type*  $\tau$  (of dimension  $n$ ) is an equivalence class of pairs  $(r_\tau, N_\tau)$  such that:

- $r_\tau : I_F \rightarrow GL_n(\bar{E})$  is a representation with open kernel.
- $N_\tau$  is a nilpotent  $n \times n$  matrix over  $\bar{E}$ .
- $(r_\tau, N_\tau)$  extends to a Weil–Deligne representation of  $G_F$ .

In particular,  $N_\tau$  commutes with the image of  $r_\tau$ . Two such pairs are equivalent if they are conjugate by an element of  $GL_n(\bar{E})$ .

We say that a continuous representation  $\rho : G_F \rightarrow GL_n(\bar{E})$  has inertial type  $\tau$  if the restriction to inertia of the associated Weil–Deligne representation is equivalent to  $\tau$ .

We define some particular two-dimensional types which will often arise. They will all be of the form  $(r, N)$  with  $r|_{\tilde{P}_F}$  trivial, and are therefore determined by  $r(\sigma)$  and  $N$ . Define:

- $\tau_{\zeta, s}$  by  $r(\sigma) = \begin{pmatrix} \zeta & 0 \\ 0 & \zeta \end{pmatrix}$  and  $N = 0$ , where  $\zeta$  is an  $l^a$ -th root of unity ( $s$  is for “split”).
- $\tau_{\zeta, ns}$  by  $r(\sigma) = \begin{pmatrix} \zeta & 0 \\ 0 & \zeta \end{pmatrix}$  and  $N = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$  where  $\zeta$  is an  $l^a$ -th root of unity ( $ns$  is for “nonsplit”).
- $\tau_{\zeta_1, \zeta_2}$  by  $r(\sigma) = \begin{pmatrix} \zeta_1 & 0 \\ 0 & \zeta_2 \end{pmatrix}$  and  $N = 0$  where,  $\zeta_1$  and  $\zeta_2$  are distinct  $l^a$ -th roots of unity.
- $\tau_\xi$  by  $r(\sigma) = \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix}$  and  $N = 0$  where,  $\xi$  is a *nontrivial*  $l^b$ -th root of unity.

To see that  $\tau_\xi$  is a type, note that if  $L/F$  is the unramified quadratic extension, then there is a character of  $G_L/\tilde{P}_F$  mapping  $\sigma$  to  $\xi$ , which when induced to  $G_F$  gives a representation of type  $\tau_\xi$ .

#### 3B. Deformation rings with fixed type.

**Definition 3.2.** Let  $\tau$  be an inertial type. Then  $R^\square(\bar{\rho}, \tau)$  is the maximal reduced,  $l$ -torsion free quotient of  $R^\square(\bar{\rho})$  with the following property: if  $x : R^\square(\bar{\rho}) \rightarrow GL_n(\bar{E})$  is a continuous homomorphism such that the associated representation  $\rho_x : G_F \rightarrow GL_n(\bar{E})$  has type  $\tau$ , then  $x$  factors through  $R^\square(\bar{\rho}, \tau)$ .

The rings  $R^\square(\bar{\rho}) \otimes \mathbb{F}$  and  $R^\square(\bar{\rho}, \tau) \otimes \mathbb{F}$  will occur very often, and so we denote them respectively by  $\bar{R}^\square(\bar{\rho})$  and  $\bar{R}^\square(\bar{\rho}, \tau)$ .

From now on suppose that  $n = 2$ . Write  $\tau = (r_\tau, N_\tau)$  and assume that  $E$  is large enough that all of the roots of the characteristic polynomial of  $r_\tau$  lie in  $E$ . Let  $R^\square(\bar{\rho}, \tau)^\circ$  be the maximal quotient of  $R^\square(\bar{\rho})$  on which:

- If  $r_\tau$  is not scalar then, for all  $g \in I_F$ , the characteristic polynomial of  $\rho^\square(g)$  agrees with that of  $r_\tau$ .
- If  $r_\tau$  is scalar and  $N_\tau = 0$  then, for all  $g \in I_F$ ,  $\rho^\square(g)$  is scalar and agrees with  $r_\tau$ .
- If  $r_\tau$  is scalar and  $N_\tau \neq 0$  then, for all  $g \in I_F$ , the characteristic polynomial of  $\rho^\square(g)$  agrees with that of  $r_\tau$ . Moreover, we have

$$q(\text{tr } \rho^\square(\phi))^2 = (q + 1)^2 \det(\rho^\square(\phi)). \tag{2}$$

It is clear that these quotients exist and that the conditions imposed are deformation problems for  $\bar{\rho}$ .

**Lemma 3.3.** *The ring  $R^\square(\bar{\rho}, \tau)$  is a reduced  $l$ -torsion free quotient of  $R^\square(\bar{\rho}, \tau)^\circ$ .*

*If  $N_\tau = 0$ , then we have that  $R^\square(\bar{\rho}, \tau)$  is equal to the maximal reduced  $l$ -torsion free quotient of  $R^\square(\bar{\rho}, \tau)^\circ$ .*

*Proof.* The first part is clear unless  $r_\tau$  is scalar and  $N_\tau \neq 0$ . In this case, we must show that any representation  $\rho : G_F \rightarrow \text{GL}_2(\bar{E})$  of type  $\tau$  satisfies equation (2). The Weil–Deligne representation  $(r, N)$  corresponding to such a  $\rho$  satisfies  $r|_{I_F} = r_\tau$  and  $N \neq 0$ . Then  $r(\phi)N = qNr(\phi)$  implies that  $r(\phi)$  preserves the line  $\ker N$  and the quotient  $\bar{E}^2 / \ker N$ . If it acts as  $\alpha$  on the former and  $\beta$  on the latter then we must have  $\alpha = q\beta$ ; as  $\alpha$  and  $\beta$  are the eigenvalues of  $\rho(\phi)$  equation (2) is easily verified.

The final claim follows from the simple observation that any  $\bar{E}$ -point of  $R^\square(\bar{\rho}, \tau)^\circ$  has associated Galois representation of type  $\tau$ , except perhaps if  $r_\tau$  is scalar and  $N_\tau \neq 0$ . □

**Remark 3.4.** If  $R$  is a reduced,  $l$ -torsion free quotient of  $R^\square(\bar{\rho})$  such that  $R^\square(\bar{\rho}, \tau)$  is a quotient of  $R$ , then  $R = R^\square(\bar{\rho}, \tau)$  if and only if the closed points of type  $\tau$  are Zariski dense in  $\text{Spec } R[1/l]$ . In our calculations, when this is true it will always be clear by inspection.

**3C.  $K$ -types.** Let  $G = \text{GL}_2(F)$ ,  $K = \text{GL}_2(\mathcal{O}_F)$ , and for  $N \geq 1$  let  $K(N) = 1 + M_2(\mathfrak{p}_F^N)$  and  $K_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : c \in \mathfrak{p}_F^N \right\}$ . Let  $U_0 = \mathcal{O}_F^\times$  and for  $N \geq 1$  let  $U_N = 1 + \mathfrak{p}_F^N$ . The *exponent* of a character  $\chi$  of  $\mathcal{O}_F^\times$  is the smallest  $N \geq 0$  such that  $\chi$  is trivial on  $U_N$ . If  $\pi$  is an irreducible admissible representation of  $\text{GL}_m(F)$  (we only need  $m = 1$  and  $m = 2$ ) over  $\bar{E}$ , let  $\text{rec}(\pi)$  be the continuous representation of  $W_F$  over  $\bar{E}$  associated to  $\pi$  under the local Langlands correspondence (normalised so as to be preserved by automorphisms of  $\bar{E}$ ).

For each two-dimensional inertial type  $\tau = (r_\tau, N_\tau)$ , we define an irreducible representation  $\sigma(\tau)$  by the following recipe:

- If  $\tau = \tau_{1,s}$ , then  $\sigma(\tau)$  is the trivial representation of  $K$ .
- If  $\tau = \tau_{1,ns}$ , then  $\sigma(\tau)$  is the inflation to  $K$  of the Steinberg representation  $\text{St}$  of  $GL_2(k_F)$ .
- If  $\tau = (\mathbb{1} \oplus \text{rec}(\epsilon)|_{I_F}, 0)$  for a nontrivial character  $\epsilon$  of  $F^\times$  of exponent  $N$ , then

$$\sigma(\tau) = \text{Ind}_{K_0(N)}^K \epsilon,$$

where  $\epsilon\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \epsilon(a)$ .

- If  $\tau = (\text{rec}(\pi)|_{I_F}, 0)$  for a cuspidal representation  $\pi$  of  $GL_2(F)$ , then by [Bushnell and Henniart 2006, Theorem 15.5] there is a certain subgroup  $J \subset G$ , containing the centre of  $G$  and compact modulo centre, and a representation  $\Lambda$  of  $J$  such that

$$\pi = \text{c-Ind}_J^G \Lambda.$$

By conjugating, we may suppose that the maximal compact subgroup  $J^0$  of  $J$  is contained in  $K$ . We then have

$$\sigma(\tau) = \text{Ind}_{J^0}^K (\Lambda|_{J^0}).$$

- If  $\tau = \tau' \otimes \text{rec}(\chi)|_{I_F}$ , then  $\sigma(\tau) = \sigma(\tau') \otimes (\chi|_{U_0} \circ \det)$ .

This is a slightly modified version of the construction in [Henniart 2002] — the construction there only depends on  $r_\tau$ , and agrees with ours whenever  $r_\tau$  is not scalar. The following is an easy consequence of [Henniart 2002]:

**Proposition 3.5.** *If  $\sigma(\tau)$  is contained in an irreducible admissible representation  $\pi$  of  $GL_2(F)$  and  $\text{rec}(\pi) = (r, N)$ , then  $r|_{I_F} \cong r_\tau$  and either  $N \cong N_\tau$  or  $N_\tau \neq 0$  and  $N = 0$ .*

*If  $\pi$  is infinite-dimensional, then the converse is true.*

**3D. Reduction of types.** Suppose that  $\bar{r} : I_F \rightarrow GL_2(\bar{\mathbb{F}})$  is such that  $\bar{r}$  extends to  $G_F$ .

**Definition 3.6.** The set  $L(\bar{r})$  is the set of types  $\tau$  such that there exists a representation  $\rho : G_F \rightarrow GL_2(\mathcal{O}_{\bar{E}})$  of type  $\tau$  satisfying

$$\bar{\rho}|_{I_F} \cong \bar{r}.$$

If  $\bar{r}|_{\bar{P}_F}$  is nonscalar then we abuse notation and also write  $L(\bar{r})$  for the set of  $r$  such that  $(r, 0) \in L(\bar{r})$ , as in this case every element of  $L(\bar{r})$  is of this form.

**Lemma 3.7.** *Suppose that  $\bar{r}$  is trivial on  $\tilde{P}_F$ . Then each element of  $L(\bar{r})$  is one of the types  $\tau_{\zeta,s}$ ,  $\tau_{\zeta,ns}$ ,  $\tau_{\zeta_1,\zeta_2}$ ,  $\tau_\xi$  defined in Section 3A.*

*Proof.* Suppose that  $\rho : G_F \rightarrow \mathrm{GL}_2(\mathcal{O}_{\bar{E}})$  is of type  $\tau$  and is such that  $\bar{\rho}|_{I_F} \cong \bar{r}$ . As  $\bar{r}|_{\tilde{P}_F}$  is trivial,  $\rho$  must also be trivial on  $\tilde{P}_F$  and its type is determined by the eigenvalues of  $\rho(\sigma)$  and by a nilpotent matrix  $N$  commuting with  $\rho(\sigma)$ . Now, the fundamental relation  $\phi\sigma\phi^{-1} = \sigma^q$  shows that the eigenvalues of  $\rho(\sigma)$  are the same (but perhaps in a different order) as those of  $\rho(\sigma)^q$ , and this implies that they are  $(q^2 - 1)$ -th roots of unity. Moreover, they are congruent to 1 modulo the maximal ideal of  $\mathcal{O}_{\bar{E}}$ , and so must in fact be either  $l^a$ -th or  $l^b$ -th roots of unity (recall that at most one of  $a$  and  $b$  is nonzero, since  $l \neq 2$ ). If they are distinct  $l^a$ -th roots of unity, then  $N$  must be zero and  $\tau = \tau_{\zeta_1, \zeta_2}$ ; if they are equal  $l^a$ -th roots of unity then  $\tau = \tau_{\zeta, s}$  or  $\tau_{\zeta, ns}$ ; if they are  $l^b$ -th roots of unity then they must be  $\xi$  and  $\xi^q = \xi^{-1}$  for an  $l^b$ -th root of unity  $\xi$ . Moreover the case  $\xi = 1$  has already been dealt with and so we may assume that  $\xi \neq 1$ , in which case  $N = 0$  and  $\tau = \tau_{\xi}$ .  $\square$

- Lemma 3.8.** (1) *Suppose that  $\bar{r}|_{\tilde{P}_F}$  is irreducible. There is a lift  $r$  of  $\bar{r}$  to  $\mathrm{GL}_2(\bar{E})$ , which we fix. Then  $L(\bar{r}) = \{r \otimes \chi\}_{\chi}$  as  $\chi$  runs over the set of characters  $\chi : I_F \rightarrow \bar{E}^{\times}$  which extend to  $G_F$  and reduce to the trivial character.*
- (2) *Suppose that  $\bar{r}|_{\tilde{P}_F} \cong (\bar{r}_1 \oplus \bar{r}_2)|_{\tilde{P}_F}$  where  $\bar{r}_1$  and  $\bar{r}_2$  are distinct characters of  $G_F$ . There are lifts  $r_1$  and  $r_2$  of  $\bar{r}_1$  and  $\bar{r}_2$  to  $\bar{E}^{\times}$ , which we fix. Then  $L(\bar{r}) = \{(r_1|_{I_F} \otimes \chi_1) \oplus (r_2|_{I_F} \otimes \chi_2)\}_{\chi_1, \chi_2}$  where  $\chi_1, \chi_2$  run over all pairs of characters  $I_F \rightarrow \bar{E}^{\times}$  which extend to  $G_F$  and reduce to the trivial character.*
- (3) *Suppose that  $\bar{r}|_{\tilde{P}_F} \cong (\bar{r}_1 \oplus \bar{r}_1^c)|_{\tilde{P}_F}$  where  $\bar{r}_1$  and  $\bar{r}_1^c$  are distinct characters of  $G_L$  which are conjugate by an element of  $G_F$  (recall that  $L/F$  is the unramified quadratic extension). There is a lift  $r_1$  of  $\bar{r}_1$  to  $\bar{E}^{\times}$ . Then  $L(\bar{r}) = \{(r_1|_{I_F} \otimes \chi) \oplus (r_1^c|_{I_F} \otimes \chi^c)\}_{\chi}$  as  $\chi$  runs over all characters  $I_F \rightarrow \bar{E}^{\times}$  which extend to  $G_L$  and reduce to the trivial character.*

*Proof.* This follows from Proposition 5.1 below; the ingredients in the proof of that proposition are Lemma 2.3 (reduction to the tame case) and Lemma 2.4 (lifting ring of a character).  $\square$

**Lemma 3.9.** *If  $\tau = (r, 0)$  is an inertial type with  $r|_{\tilde{P}_F}$  nonscalar, then  $\overline{\sigma(\tau)}$  is irreducible. If  $\tau'$  is any other inertial type, then  $\overline{\sigma(\tau')}$  contains  $\overline{\sigma(\tau)}$  if and only if  $\tau' \in L(\bar{r})$  (in which case  $\overline{\sigma(\tau)} \cong \overline{\sigma(\tau')}$ ).*

*Proof.* These are the results of Propositions 6.4 and 6.5.  $\square$

If  $\tau = (r, N)$  with  $r|_{\tilde{P}_F}$  scalar, then  $\overline{\sigma(\tau)}$  need not be irreducible. We give the (well-known) analysis of these  $\overline{\sigma(\tau)}$  in Section 6A. For now, we just give names to the following representations of  $\mathrm{GL}_2(k_F)$  (and hence, by inflation, of  $K$ ) over  $\mathbb{F}$

- the trivial representation,  $\mathbb{1}$ ,
- the Steinberg representation,  $\mathrm{St}$  (irreducible if  $q \not\equiv -1 \pmod{l}$ ),
- if  $q \equiv -1 \pmod{l}$ , the cuspidal (but not supercuspidal) subrepresentation  $\pi_1$  of  $\mathrm{St}$ .

### 4. The “Breuil–Mézard conjecture”

Let  $\bar{\rho} : G_F \rightarrow \mathrm{GL}_2(\mathbb{F})$  be a continuous representation, and suppose that  $E$  is sufficiently large that

- every subrepresentation of  $\bar{\rho} \otimes \bar{\mathbb{F}}$  is already defined over  $\mathbb{F}$ ,
- $E$  contains all of the  $(q^2 - 1)$ -th roots of unity,
- for every  $\tau \in L(\bar{\rho}|_{I_F})$ ,  $\sigma(\tau)$  is defined over  $E$ .

We state our analogue of the Breuil–Mézard conjecture when  $l \neq p$ . By [Lemma 2.2](#) and the fact that  $R^\square(\bar{\rho}, \tau)$  is defined to be  $\mathcal{O}$ -flat, we have

$$\dim \bar{R}^\square(\bar{\rho}, \tau) \leq 4.$$

**Definition 4.1.** We associate to each type  $\tau = (r, N)$  a cycle  $\mathcal{C}(\bar{\rho}, \tau) \in \mathcal{Z}^4(\bar{R}^\square(\bar{\rho}))$  as follows:

- If  $N = 0$ , set

$$\mathcal{C}(\bar{\rho}, \tau) = Z^4(\bar{R}^\square(\bar{\rho}, \tau)).$$

- If  $N \neq 0$  (in which case  $r$  must be scalar) let  $\tau' = (r, 0)$  and set

$$\mathcal{C}(\bar{\rho}, \tau) = Z^4(\bar{R}^\square(\bar{\rho}, \tau)) + Z^4(\bar{R}^\square(\bar{\rho}, \tau')).$$

**Theorem 4.2.** For each irreducible  $\bar{\mathbb{F}}$ -representation  $\theta$  of  $\mathrm{GL}_2(\mathcal{O}_F)$ , there is an effective cycle  $\mathcal{C}(\bar{\rho}, \theta) \in \mathcal{Z}^4(\bar{R}^\square(\bar{\rho}))$  such that, for any inertial type  $\tau$ , we have an equality of cycles

$$\mathcal{C}(\bar{\rho}, \tau) = \sum_{\theta} m(\theta, \overline{\sigma(\tau)}) \mathcal{C}(\bar{\rho}, \theta), \tag{3}$$

where  $m(\theta, \overline{\sigma(\tau)})$  is the multiplicity of  $\theta$  as a Jordan–Hölder factor of  $\overline{\sigma(\tau)}$  and the sum runs over all  $\theta$ .

*Proof.* We proceed case by case, using the results of [Section 3D](#) and of [Sections 5](#) and [6A](#) below.

Suppose that  $\bar{\rho}|_{\tilde{p}_F}$  is nonscalar. Then by [Lemma 3.9](#), the representations  $\overline{\sigma(\tau)}$  for  $\tau \in L(\bar{\rho}|_{I_F})$  are all irreducible and isomorphic to a common irreducible representation, which we call  $\theta_0$ . By [Corollary 5.2](#),  $\bar{R}^\square(\bar{\rho})$  has a unique minimal prime, denoted  $\mathfrak{a}$ , which has dimension 4. So we have

$$\mathcal{Z}^4(\mathrm{Spec}(\bar{R}^\square(\bar{\rho}))) = \mathbb{Z} \cdot [\mathfrak{a}].$$

Define  $\mathcal{C}(\bar{\rho}, \theta_0) = [\mathfrak{a}]$  and  $\mathcal{C}(\bar{\rho}, \theta) = 0$  for  $\theta \neq \theta_0$ . By [Corollary 5.2](#),

$$\mathcal{C}(\bar{\rho}, \tau) = [\mathfrak{a}] = \mathcal{C}(\bar{\rho}, \theta_0)$$

if  $\tau \in L(\bar{\rho}|_{\tilde{p}_F})$ , otherwise

$$\mathcal{C}(\bar{\rho}, \tau) = 0.$$

In other words, for all  $\tau$  we have

$$\mathcal{C}(\bar{\rho}, \tau) = \sum_{\theta} m(\theta, \overline{\sigma(\tau)}) \mathcal{C}(\bar{\rho}, \theta),$$

as required.

If  $\bar{\rho}|_{\bar{F}}$  is scalar, then we may twist  $\bar{\rho}$  by a character of  $G_F$  and apply [Lemma 2.4](#) and so *suppose for the rest of the proof that  $\bar{\rho}|_{\bar{F}}$  is trivial.*

If  $q \not\equiv \pm 1 \pmod{l}$ , then  $L(\bar{\rho}|_{I_F}) \subset \{\tau_{1,s}, \tau_{1,ns}\}$ . By the discussion of [Section 6A](#), we have that

$$\overline{\sigma(\tau_{1,s})} = \mathbb{1} \quad \text{and} \quad \overline{\sigma(\tau_{1,ns})} = \text{St}$$

are irreducible and nonisomorphic, and that neither is a Jordan–Hölder factor of any other  $\overline{\sigma(\tau)}$ . So the fact that we *can* define the  $\mathcal{C}(\bar{\rho}, \theta)$  so as to satisfy [\(3\)](#) is a trivality, as there are no relations amongst the  $\overline{\sigma(\tau)}$  for different  $\tau$ . We work out what the  $\mathcal{C}(\bar{\rho}, \theta)$  are explicitly: for  $\theta \neq \mathbb{1}$  or St we define  $\mathcal{C}(\bar{\rho}, \theta) = 0$ . Otherwise, there are four cases to consider:

- If  $\bar{\rho}(\phi)$  has eigenvalues with ratio not in  $\{1, \pm q\}$  then by [Proposition 5.3](#) there is a unique minimal prime  $\mathfrak{a}_{nr}$  of  $\bar{R}^{\square}(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_{nr}].$$

- If  $\bar{\rho}$  is an extension of the trivial character by itself then by [Proposition 5.5](#) part 1 there is a unique minimal prime  $\mathfrak{a}_{nr}$  of  $\bar{R}^{\square}(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_{nr}].$$

- If  $\bar{\rho}$  is a nonsplit extension of the trivial character by the cyclotomic character then by [Proposition 5.5](#) part 2 there is a unique minimal prime  $\mathfrak{a}_N$  of  $\bar{R}^{\square}(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = 0, \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_N].$$

- If  $\bar{\rho}$  is the direct sum of the trivial character and the cyclotomic character then by [Proposition 5.5](#) part 2 there are two minimal primes of  $\bar{R}^{\square}(\bar{\rho})$ , denoted there by  $\mathfrak{a}_{nr}$  and  $\mathfrak{a}_N$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_{nr}] + [\mathfrak{a}_N].$$

It is then easy to verify that equation [\(3\)](#) holds; we just do the last case. We see from [Proposition 5.5](#) part 2 that

$$\mathcal{C}(\bar{\rho}, \tau_{1,s}) = [\mathfrak{a}_{nr}] = \mathcal{C}(\bar{\rho}, \mathbb{1}), \quad \mathcal{C}(\bar{\rho}, \tau_{1,ns}) = [\mathfrak{a}_{nr}] + [\mathfrak{a}_N] = \mathcal{C}(\bar{\rho}, \text{St}),$$

and  $\mathcal{C}(\bar{\rho}, \tau) = 0$  for all other  $\tau$ , exactly as required by [\(3\)](#).



If  $q \equiv -1 \pmod{l}$ , then  $L(\bar{\rho}|_{I_F}) \subset \bigcup_{\xi} \{\tau_{1,s}, \tau_{1,ns}, \tau_{\xi}\}$  for  $\xi$  a nontrivial  $l^b$ -th root of unity. By the discussion of [Section 6A](#), we have that

$$\overline{\sigma(\tau_{1,s})} = \mathbb{1}, \quad \overline{\sigma(\tau_{\xi})} = \pi_1, \quad \overline{\sigma(\tau_{1,ns})}^{ss} = \mathbb{1} \oplus \pi_1,$$

where  $\mathbb{1}$  and  $\pi_1$  are irreducible and nonisomorphic, and are not Jordan–Hölder factors of any other  $\overline{\sigma(\tau)}$ . For  $\theta \neq \mathbb{1}$  or  $\pi_1$  we define  $\mathcal{C}(\bar{\rho}, \theta) = 0$ . Otherwise, there are four cases to consider:

- If  $\bar{\rho}(\phi)$  has eigenvalues with ratio not in  $\{\pm 1\}$  then by [Proposition 5.3](#) there is a unique minimal prime  $\mathfrak{a}_{nr}$  of  $\bar{R}^{\square}(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \pi_1) = 0.$$

- If  $\bar{\rho}$  is an extension of the trivial character by itself then by [Proposition 5.6](#) part 1 there is a unique minimal prime  $\mathfrak{a}_{nr}$  of  $\bar{R}^{\square}(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \pi_1) = 0.$$

- If  $\bar{\rho}$  is a nonsplit extension of the trivial character by the cyclotomic character then by [Proposition 5.6](#) part 2a there is a unique minimal prime, denoted  $\mathfrak{a}_N$  in that proposition, of  $\bar{R}^{\square}(\bar{\rho}, \tau_{1,ns})$ , which we regard as a prime of  $\bar{R}^{\square}(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = 0, \quad \mathcal{C}(\bar{\rho}, \pi_1) = [\mathfrak{a}_N].$$

- If  $\bar{\rho}$  is the direct sum of the trivial character by the cyclotomic character then in [Proposition 5.6](#) part 2b three four-dimensional primes of  $\bar{R}^{\square}(\bar{\rho})$  are defined, denoted there  $\mathfrak{a}_{nr}$ ,  $\mathfrak{a}_N$  and  $\mathfrak{a}_{N'}$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \pi_1) = [\mathfrak{a}_N] + [\mathfrak{a}_{N'}].$$

It is then easy to verify that [\(3\)](#) holds using [Proposition 5.3](#) in the first case and [Proposition 5.6](#) parts 1, 2a, and 2b in the second, third, and fourth cases; again we just do the fourth case, which is the most complicated. [Equation \(3\)](#) is equivalent to the equations

$$\begin{aligned} \mathcal{C}(\bar{\rho}, \tau_{1,s}) &= \mathcal{C}(\bar{\rho}, \mathbb{1}) &&= [\mathfrak{a}_{nr}], \\ \mathcal{C}(\bar{\rho}, \tau_{1,ns}) &= \mathcal{C}(\bar{\rho}, \mathbb{1}) + \mathcal{C}(\bar{\rho}, \pi_1) &&= [\mathfrak{a}_{nr}] + [\mathfrak{a}_N] + [\mathfrak{a}_{N'}], \\ \mathcal{C}(\bar{\rho}, \tau_{\xi}) &= \mathcal{C}(\bar{\rho}, \pi_1) &&= [\mathfrak{a}_N] + [\mathfrak{a}_{N'}], \end{aligned}$$

and

$$\mathcal{C}(\bar{\rho}, \tau) = 0 \quad \text{if } \tau \notin \bigcup_{\xi} \{\tau_{1,s}, \tau_{1,ns}, \tau_{\xi}\}.$$

But by [Proposition 5.6](#) part 2b we have

$$\begin{aligned} \mathcal{C}(\bar{\rho}, \tau_{1,s}) &= Z^4(\bar{R}(\bar{\rho}, \tau_{1,s})) &&= [\mathfrak{a}_{nr}], \\ \mathcal{C}(\bar{\rho}, \tau_{1,ns}) &= Z^4(\bar{R}(\bar{\rho}, \tau_{1,s})) + Z^4(\bar{R}(\bar{\rho}, \tau_{1,ns})) &&= [\mathfrak{a}_{nr}] + [\mathfrak{a}_N] + [\mathfrak{a}_{N'}], \\ \mathcal{C}(\bar{\rho}, \tau_\xi) &= Z^4(\bar{R}(\bar{\rho}, \tau_\xi)) &&= [\mathfrak{a}_N] + [\mathfrak{a}_{N'}], \end{aligned}$$

and

$$\mathcal{C}(\bar{\rho}, \tau) = 0 \quad \text{if } \tau \notin \bigcup_\xi \{\tau_{1,s}, \tau_{1,ns}, \tau_\xi\},$$

as required.

If  $q \equiv 1 \pmod{l}$ , then  $L(\bar{\rho}|_{I_F}) \subset \bigcup_{\zeta, \zeta_1, \zeta_2} \{\tau_{\zeta,s}, \tau_{\zeta,ns}, \tau_{\zeta_1, \zeta_2}\}$  for  $\zeta, \zeta_1$  and  $\zeta_2$  (possibly trivial)  $l^a$ -th roots of unity with  $\zeta_1 \neq \zeta_2$ . By the discussion of [Section 6A](#), we have that

$$\overline{\sigma(\tau_{\zeta,s})} = \mathbb{1}, \quad \overline{\sigma(\tau_{\zeta,ns})} = \text{St}, \quad \overline{\sigma(\tau_{\zeta_1, \zeta_2})} = \mathbb{1} \oplus \text{St},$$

where  $\mathbb{1}$  and  $\text{St}$  are irreducible and nonisomorphic, and are not Jordan–Hölder factors of any other  $\overline{\sigma(\tau)}$ . For  $\theta \neq \mathbb{1}$  or  $\text{St}$  we define  $\mathcal{C}(\bar{\rho}, \theta) = 0$ . Otherwise, there are four cases to consider:

- If  $\bar{\rho}(\phi)$  has eigenvalues with ratio not in  $\{\pm 1\}$  then by [Proposition 5.3](#) there is a unique minimal prime  $\mathfrak{a}_{nr}$  of  $\bar{R}^\square(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_{nr}].$$

- If  $\bar{\rho}$  is a ramified extension of the trivial character by itself then by [Proposition 5.8](#) part 1 there is a unique minimal prime  $\mathfrak{a}_N$  of  $\bar{R}^\square(\bar{\rho}, \tau_{1,ns})$  which we regard as a four-dimensional prime of  $\bar{R}^\square(\bar{\rho})$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = 0, \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_N].$$

- If  $\bar{\rho}$  is a unramified extension of the trivial character by itself then by [Proposition 5.8](#) parts 2 and 3 there are four-dimensional primes of  $\bar{R}^\square(\bar{\rho})$  which are denoted there by  $[\mathfrak{a}_{nr}]$  and  $[\mathfrak{a}_N]$ . In this case, define

$$\mathcal{C}(\bar{\rho}, \mathbb{1}) = [\mathfrak{a}_{nr}], \quad \mathcal{C}(\bar{\rho}, \text{St}) = [\mathfrak{a}_{nr}] + [\mathfrak{a}_N].$$

It is then easy to verify that [\(3\)](#) holds using [Proposition 5.3](#) in the first case, [Proposition 5.8](#) part 1 in the second case, and [Proposition 5.8](#) parts 2 and 3 in the third case (according as  $\bar{\rho}$  is split or not); again we just do the third case, which is the most complicated. [Equation \(3\)](#) is equivalent to the equations

$$\begin{aligned} \mathcal{C}(\bar{\rho}, \tau_{\zeta,s}) &= \mathcal{C}(\bar{\rho}, \mathbb{1}) &&= [\mathfrak{a}_{nr}], \\ \mathcal{C}(\bar{\rho}, \tau_{\zeta,ns}) &= \mathcal{C}(\bar{\rho}, \text{St}) &&= [\mathfrak{a}_{nr}] + [\mathfrak{a}_N], \\ \mathcal{C}(\bar{\rho}, \tau_{\zeta_1, \zeta_2}) &= \mathcal{C}(\bar{\rho}, \mathbb{1}) + \mathcal{C}(\bar{\rho}, \text{St}) &&= [\mathfrak{a}_{nr}] + [\mathfrak{a}_{nr}] + [\mathfrak{a}_N], \end{aligned}$$

and

$$\mathcal{C}(\bar{\rho}, \tau) = 0 \quad \text{if } \tau \notin \bigcup_{\zeta, \zeta_1, \zeta_2} \{\tau_{\zeta,s}, \tau_{\zeta,ns}, \tau_{\zeta_1, \zeta_2}\}.$$

But by [Proposition 5.8](#) parts 2 and 3 we have:

$$\begin{aligned} \mathcal{C}(\bar{\rho}, \tau_{\zeta,s}) &= Z^4(\bar{R}(\bar{\rho}, \tau_{\zeta,s})) &&= [\mathfrak{a}_{nr}], \\ \mathcal{C}(\bar{\rho}, \tau_{\zeta,ns}) &= Z^4(\bar{R}(\bar{\rho}, \tau_{\zeta,s})) + Z^4(\bar{R}(\bar{\rho}, \tau_{1,ns})) &&= [\mathfrak{a}_{nr}] + [\mathfrak{a}_N], \\ \mathcal{C}(\bar{\rho}, \tau_{\zeta_1,\zeta_2}) &= Z^4(\bar{R}(\bar{\rho}, \tau_{\zeta_1,\zeta_2})) &&= 2[\mathfrak{a}_{nr}] + [\mathfrak{a}_N], \end{aligned}$$

and

$$\mathcal{C}(\bar{\rho}, \tau) = 0 \quad \text{if } \tau \notin \bigcup_{\zeta, \zeta_1, \zeta_2} \{\tau_{\zeta,s}, \tau_{\zeta,ns}, \tau_{\zeta_1,\zeta_2}\},$$

as required. □

**Remark 4.3.** Although the definition of  $\mathcal{C}(\bar{\rho}, \tau)$  may seem ad-hoc, it in fact has the following natural interpretation: it is the reduction modulo  $\lambda$  of the cycle in  $\mathcal{Z}^4(R^\square(\bar{\rho}))$  obtained by taking the Zariski closure of the closed points  $x \in \text{Spec } R^\square(\bar{\rho})[1/l]$  such that  $\text{rec}^{-1}(\rho_x)|_K$  contains  $\sigma(\tau)$ .

**Remark 4.4.** We conjecture that [Theorem 4.2](#) remains true when  $l = 2$ .

### 5. Calculations

Let  $\bar{\rho} : G_F \rightarrow GL_2(\mathbb{F})$  be a continuous representation. The aims of this section are to give explicit presentations for the rings  $R^\square(\bar{\rho}, \tau)$  and to compute the cycles  $Z(\bar{R}^\square(\bar{\rho}, \tau)) \in \mathcal{Z}^4(\text{Spec } \bar{R}^\square(\bar{\rho}))$ . We continue to assume that  $E$  is sufficiently large, as defined at the start of the previous section.

**5A. Simple cases.** When  $\bar{\rho}|_{\bar{p}_F}$  is not scalar, then [Lemma 2.3](#) allows us to determine the universal framed deformation rings. Recall that if  $\bar{r} : I_F \rightarrow GL_2(\mathbb{F})$  is a representation that extends to  $G_F$  then we have defined the set  $L(\bar{r})$  of types that lift  $\bar{r}$ .

**Proposition 5.1.** *If  $\bar{\rho}|_{\bar{p}_F}$  is irreducible, then*

$$R^\square(\bar{\rho}) \cong \mathcal{O}[[X, Y, Z_1, Z_2, Z_3]] / ((1 + X)^{l^a} - 1).$$

*The  $l^a$  irreducible components of  $\text{Spec } R^\square(\bar{\rho})$  are precisely the  $\text{Spec } R^\square(\bar{\rho}, \tau)$  for  $\tau \in L(\bar{\rho}|_{I_F})$ .*

*If  $\bar{\rho}|_{\bar{p}_F}$  is a sum of distinct characters which extend to  $G_F$ , then*

$$R^\square(\bar{\rho}) \cong \mathcal{O}[[X_1, X_2, Y_1, Y_2, Z_1, Z_2]] / ((1 + X_1)^{l^a} - 1, (1 + X_2)^{l^a} - 1).$$

*The  $l^{2a}$  irreducible components of  $\text{Spec } R^\square(\bar{\rho})$  are precisely the  $\text{Spec } R^\square(\bar{\rho}, \tau)$  for  $\tau \in L(\bar{\rho}|_{I_F})$ .*

*If  $\bar{\rho}|_{\bar{p}_F}$  is a sum of distinct characters which are conjugate by the nontrivial element of  $G_L \setminus G_F$ , then*

$$R^\square(\bar{\rho}) \cong \mathcal{O}[[X, Y, Z_1, Z_2, Z_3]] / ((1 + X)^{l^b} - 1).$$

The  $l^b$  irreducible components of  $\text{Spec } R^\square(\bar{\rho})$  are precisely the  $\text{Spec } R^\square(\bar{\rho}, \tau)$  for  $\tau \in L(\bar{\rho}|_{I_F})$ .

*Proof.* This follows straightforwardly from [Lemma 2.3](#). Suppose first that  $\bar{\rho}|_{\tilde{P}_F}$  is irreducible. Then there is a unique irreducible representation  $\theta$  of  $\tilde{P}_F$  such that  $\bar{\rho}_\theta$  (in the notation of [Lemma 2.3](#)) is nonzero. For that  $\theta$ ,  $\bar{\rho}_\theta$  is an unramified one-dimensional representation of  $G_F$ . So by [Lemmas 2.3](#) and [2.5](#)

$$R^\square(\bar{\rho}) \cong R^\square(\bar{\rho}_\theta)[[Z_1, Z_2, Z_3]] \cong \mathcal{O}[[X, Y, Z_1, Z_2, Z_3]]/((1 + X)^{l^a} - 1).$$

We have  $\rho^\square \cong \tilde{\theta} \otimes \chi^\square$  where  $\chi^\square$  is the universal character  $G_F \rightarrow R^\square(\bar{\rho}_\theta)^\times$ .

Suppose now that  $\bar{\rho}|_{\tilde{P}_F} = \theta_1 \oplus \theta_2$  for distinct characters  $\theta_1$  and  $\theta_2$ . Suppose first that the  $\theta_i$  are not  $G_F$ -conjugate. As in [Lemma 2.3](#), we pick  $\mathcal{O}$ -characters  $\tilde{\theta}_1$  and  $\tilde{\theta}_2$  of  $G_F$  lifting and extending  $\theta_1$  and  $\theta_2$ . Then (in the notation of [Lemma 2.3](#))  $\bar{\rho}_{\theta_1}$  and  $\bar{\rho}_{\theta_2}$  are both unramified characters. By [Lemmas 2.3](#) and [2.5](#)

$$\begin{aligned} R^\square(\bar{\rho}) &\cong (R^\square(\bar{\rho}_{\theta_1}) \hat{\otimes} R^\square(\bar{\rho}_{\theta_2}))[[Z_1, Z_2]] \\ &\cong \mathcal{O}[[X_1, X_2, Y_1, Y_2, Z_1, Z_2]]/((1 + X_1)^{l^a} - 1, (1 + X_2)^{l^a} - 1). \end{aligned}$$

We have

$$\rho^\square \cong \tilde{\theta}_1 \otimes \chi_1^\square \oplus \tilde{\theta}_2 \otimes \chi_2^\square,$$

where each  $\chi_i^\square$  is the universal character over  $R^\square(\bar{\rho}_{\theta_i})$ .

Suppose finally that  $\theta_1$  and  $\theta_2$  are  $G_F$ -conjugate. We take  $\theta = \theta_1$ ; then  $G_\theta = G_L$  where  $L$  is a quadratic extension of  $F$ . In fact, since  $\tilde{P}_F \subset G_L$  and  $l$  is odd, we must have that  $G_L$  is the unramified quadratic extension of  $F$ . As in [Lemma 2.3](#), pick an  $\mathcal{O}$ -character  $\tilde{\theta}$  of  $G_L$  lifting and extending  $\theta$ . Then (in the notation of [Lemma 2.3](#))  $\bar{\rho}_\theta$  is an unramified character of  $G_L$ . By [Lemmas 2.3](#) and [2.5](#)

$$\begin{aligned} R^\square(\bar{\rho}) &\cong R^\square(\bar{\rho}_\theta)[[Z_1, Z_2, Z_3]] \\ &\cong \mathcal{O}[[X, Y, Z_1, Z_2, Z_3]]/((1 + X)^{l^b} - 1), \end{aligned}$$

since  $v_l(q^2 - 1) = l^b$ . We have

$$\rho^\square \cong \text{Ind}_{G_L}^{G_F}(\tilde{\theta} \otimes \chi^\square),$$

where  $\chi^\square$  is the universal character over  $R^\square(\bar{\rho}_\theta)$ .

We show that  $f : \text{Spec}(R^\square(\bar{\rho}, \tau)) \mapsto \tau$  is a bijection from the set of irreducible components of  $\text{Spec}(R^\square(\bar{\rho}))$  to  $L(\bar{\rho}|_{I_F})$ . It is easy to see that  $f$  is an injection (from our explicit expressions for  $\rho^\square$ ). The type of the  $\bar{E}$ -points of  $\text{Spec}(R^\square(\bar{\rho}, \tau))$  is constant on irreducible components, so to show that a particular  $\tau$  is in the image of  $f$  it suffices to produce a lift of  $\bar{\rho}$  to  $\bar{E}$  of type  $\tau$ . Each  $\tau \in L(\bar{\rho}|_{I_F})$  is, by definition, the type of a lift of some  $\bar{\rho}'$  with  $\bar{\rho}'|_{I_F} \cong \bar{\rho}|_{I_F}$ . But it is clear from the

calculations above that the image of  $f$  only depends on  $\bar{\rho}|_{I_F}$ , and so  $f$  is surjective as required.  $\square$

**Corollary 5.2.** *If  $\bar{\rho}|_{\tilde{P}_F}$  is not scalar, then  $\bar{R}^\square(\bar{\rho})$  has a unique minimal prime  $\mathfrak{a}$ , which has dimension 4. For  $\tau$  an inertial type we have that*

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = [\mathfrak{a}]$$

if  $\tau \in L(\bar{\rho}|_{\tilde{P}_F})$  and  $Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = 0$  otherwise.

We may now assume that  $\bar{\rho}|_{\tilde{P}_F}$  is scalar; after a twist (invoking [Clozel et al. 2008, Lemma 2.4.11] to extend the character occurring in  $\bar{\rho}|_{\tilde{P}_F}$  to the whole Galois group), we may assume that  $\bar{\rho}|_{\tilde{P}_F}$  is trivial, so that any lift of  $\bar{\rho}|_{\tilde{P}_F}$  is also trivial. In this case, then,  $\bar{\rho}|_{I_F}$  is inflated from a representation of the (pro)cyclic pro- $l$  group  $I_F/\tilde{P}_F$  over a field of characteristic  $l$ . Any irreducible representation in characteristic  $l$  of an  $l$ -group is trivial, and so  $\bar{\rho}|_{I_F}$  must be an extension of the trivial representation by the trivial representation. Now, because  $\phi\sigma\phi^{-1} = \sigma^q$ ,  $\bar{\rho}(\phi)$  maps the subspace of fixed vectors of  $\bar{\rho}(\sigma)$  to itself; therefore,  $\bar{\rho}$  must be an extension of unramified characters. That is, there is a short exact sequence

$$0 \rightarrow \chi_1 \rightarrow \bar{\rho} \rightarrow \chi_2 \rightarrow 0$$

for unramified characters  $\chi_1$  and  $\chi_2$ . Such an extension corresponds to an element of  $H^1(G_F, \chi_1\chi_2^{-1})$ ; by a simple calculation with the local Euler characteristic formula and local Tate duality, this cohomology group is nonzero if and only if  $\chi_1 = \chi_2$  or  $\chi_1 = \chi_2\epsilon$ . So we can easily deal with the case where neither of these two possibilities can occur.

**Proposition 5.3.** *Suppose that  $\bar{\rho}|_{\tilde{P}_F}$  is trivial and that  $\bar{\rho}(\phi)$  has eigenvalues  $\bar{\alpha}, \bar{\beta} \in \mathbb{F}$  with  $\bar{\alpha}/\bar{\beta} \notin \{1, q, q^{-1}\}$ . Then*

$$R^\square(\bar{\rho}) \cong \frac{\mathcal{O}[[A, B, P, Q, X, Y]]}{((1 + P)^{l^a} - 1, (1 + Q)^{l^a} - 1)},$$

and  $\rho^\square(\sigma)$  is diagonalizable with eigenvalues  $1 + P$  and  $1 + Q$ .

For  $\zeta$  an  $l^a$ -th root of unity (possibly equal to 1), we have that

$$\begin{aligned} R^\square(\bar{\rho}, \tau_{\zeta, s}) &= \mathcal{O}[[A, B, P, Q, X, Y]] / (1 + P - \zeta, 1 + Q - \zeta) \\ &\cong \mathcal{O}[[A, B, X, Y]] \end{aligned}$$

is formally smooth of relative dimension 4 over  $\mathcal{O}$  and that  $R^\square(\bar{\rho}, \tau_{\zeta, ns}) = 0$ . If  $q \equiv 1 \pmod{l}$  and  $\zeta_1, \zeta_2$  are distinct  $l^a$ -th roots of unity, then

$$\begin{aligned} R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2}) &= \frac{\mathcal{O}[[A, B, P, Q, X, Y]]}{(2 + P + Q - \zeta_1 - \zeta_2, PQ - (\zeta_1 - 1)(\zeta_2 - 1))} \\ &\cong \mathcal{O}[[A, B, P, X, Y]] / (1 + P - \zeta_1)(1 + P - \zeta_2). \end{aligned}$$

For all other  $\tau$ ,  $R^\square(\bar{\rho}, \tau) = 0$ .

The ideal  $\mathfrak{a}_{nr}$  defining  $\bar{R}^\square(\bar{\rho}, \tau_{1,s})$  is the unique minimal prime of  $\bar{R}^\square(\bar{\rho})$ . We have

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} [\mathfrak{a}_{nr}] & \text{if } \tau = \tau_{\zeta,s}, \\ 2[\mathfrak{a}_{nr}] & \text{if } \tau = \tau_{\zeta_1,\zeta_2}, \\ 0 & \text{if } \tau = \tau_{\zeta,ns}. \end{cases}$$

*Proof.* First note that, by the above cohomology calculation,  $\bar{\rho}(\sigma)$  must be trivial.

Let  $\alpha$  and  $\beta$  be lifts of  $\bar{\alpha}$  and  $\bar{\beta}$  to  $\mathcal{O}$ . Suppose that  $\mathcal{A}$  is an object of  $\mathcal{C}_{\mathcal{O}}$  and that  $M$  is a free  $\mathcal{A}$ -module of rank 2 with a continuous action of  $G_F$  given by  $\rho : G_F \rightarrow \text{Aut}_{\mathcal{A}}(M)$ , reducing to  $\bar{\rho}$  modulo  $\mathfrak{m}_{\mathcal{A}}$ . Suppose that the characteristic polynomial of  $\rho(\phi)$  is  $(X - \alpha - A)(X - \beta - B)$ , where  $A, B \in \mathfrak{m}_{\mathcal{A}}$ —note that by Hensel’s lemma the characteristic polynomial *does* have roots in  $\mathcal{A}$  reducing to  $\bar{\alpha}$  and  $\bar{\beta}$ . Then there is a decomposition

$$M = (\rho(\phi) - \alpha - A)M \oplus (\rho(\phi) - \beta - B)M.$$

Here it is crucial that  $\alpha + A, \beta + B$  and,  $\alpha - \beta + A - B$  are all invertible in  $\mathcal{A}$ . If  $\bar{v}_\alpha, \bar{v}_\beta$  is a basis of eigenvectors of  $\bar{\rho}(\phi)$  in  $M \otimes \mathbb{F}$  and  $v_\alpha, v_\beta$  is a basis of  $M$  lifting  $\bar{v}_\alpha, \bar{v}_\beta$  then there are unique  $X, Y \in \mathfrak{m}_{\mathcal{A}}$  such that  $v_\alpha + Xv_\beta, v_\beta + Yv_\alpha$  are eigenvectors of  $\rho(\phi)$ . Moreover, replacing  $(v_\alpha, v_\beta)$  by  $(\mu v_\alpha, \mu v_\beta)$  for  $\mu \in 1 + \mathfrak{m}_{\mathcal{A}}$  does not change  $X$  and  $Y$ .

Therefore we may assume that  $\bar{\rho}(\phi) = \begin{pmatrix} \bar{\alpha} & 0 \\ 0 & \bar{\beta} \end{pmatrix}$  and that

$$\begin{aligned} \rho(\phi) &= \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}^{-1} \begin{pmatrix} \alpha + A & 0 \\ 0 & \beta + B \end{pmatrix} \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}, \\ \rho(\sigma) &= \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 + P & R \\ S & 1 + Q \end{pmatrix} \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}, \end{aligned}$$

where  $\rho$  determines  $X, Y, P, R, S, Q \in \mathfrak{m}_{\mathcal{A}}$  uniquely. The equation  $\phi\sigma\phi^{-1} = \sigma^q$  implies that

$$\begin{pmatrix} \alpha + A & 0 \\ 0 & \beta + B \end{pmatrix} \begin{pmatrix} 1 + P & R \\ S & 1 + Q \end{pmatrix} \begin{pmatrix} \alpha + A & 0 \\ 0 & \beta + B \end{pmatrix}^{-1} = \begin{pmatrix} 1 + P & R \\ S & 1 + Q \end{pmatrix}^q.$$

Looking at the top right and bottom left entries gives that  $R = S = 0$ . Then looking at the diagonal entries gives that  $(1 + P)^{q-1} = (1 + Q)^{q-1} = 1$ , which is equivalent to  $(1 + P)^{l^a} = (1 + Q)^{l^a} = 1$ . Thus

$$R^\square(\bar{\rho}) \cong \frac{\mathcal{O}[[A, B, P, Q, X, Y]]}{((1 + P)^{l^a} - 1, (1 + Q)^{l^a} - 1)}.$$

The possible inertial types are  $\tau_{\zeta,s}$  and  $\tau_{\zeta_1,\zeta_2}$  ( $\tau_{\zeta,ns}$  cannot occur since all lifts are diagonalisable). Clearly  $R^\square(\bar{\rho}, \tau_{\zeta,s})$  is defined by the equations  $1 + P =$

$1 + Q = \zeta$ . The ring  $R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})^\circ$  is cut out by the equations  $2 + P + Q = \zeta_1 + \zeta_2$ ,  $(1 + P)(1 + Q) = \zeta_1 \zeta_2$  and the redundant equations  $(1 + P)^{l^a} = (1 + Q)^{l^a} = 1$ . But

$$R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})^\circ \cong \mathcal{O}[[A, B, P, X, Y]] / ((1 + P - \zeta_1)(1 + P - \zeta_2))$$

is reduced and  $\lambda$ -torsion free and so is equal to  $R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})$ .

For the reduction modulo  $\lambda$ , simply note that

$$\begin{aligned} \bar{R}^\square(\bar{\rho}) &= \mathbb{F}[[A, B, P, Q, X, Y]] / (P^{l^a}, Q^{l^a}), \\ \bar{R}^\square(\bar{\rho}, \tau_{\zeta, s}) &= \mathbb{F}[[A, B, P, Q, X, Y]] / (P, Q), \end{aligned}$$

and

$$\bar{R}^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2}) = \mathbb{F}[[A, B, P, Q, X, Y]] / (P^2, Q^2, P + Q).$$

So  $\mathfrak{a}_{nr} = (P, Q)$  is the unique minimal prime of  $\bar{R}^\square(\bar{\rho})$  and the multiplicities are as claimed.  $\square$

We extract the first half of the proof of this proposition for future use:

**Lemma 5.4.** *If  $\bar{\rho}(\phi)$  has distinct eigenvalues, we may assume that it is diagonal. In that case, there exists a unique matrix  $\begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix} \in GL_2(R^\square(\bar{\rho}))$ , reducing to the identity modulo the maximal ideal, such that  $\rho^\square(\phi) = \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}^{-1} \Phi \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}$  for a diagonal matrix  $\Phi$ .*  $\square$

**5B.  $q \not\equiv \pm 1 \pmod{l}$ .** Suppose that  $q \not\equiv \pm 1 \pmod{l}$ . By [Proposition 5.3](#), we have already dealt with the cases in which the eigenvalues of  $\bar{\rho}(\phi)$  are not in the ratio 1 or  $q^{\pm 1}$ . All other cases are dealt with by the following (after twisting and conjugating  $\bar{\rho}$ ). Note that, by [Lemma 3.7](#), the only possible types when  $\bar{\rho}|_{\tilde{F}_F}$  is trivial are  $\tau_{1,s}$  and  $\tau_{1,ns}$ .

**Proposition 5.5.** *Suppose that  $q \not\equiv \pm 1 \pmod{l}$ , and that  $\bar{\rho}|_{\tilde{F}_F}$  is trivial.*

(1) *Suppose  $\bar{\rho}(\sigma)$  is trivial, and  $\bar{\rho}(\phi) = \begin{pmatrix} 1 & \bar{y} \\ 0 & 1 \end{pmatrix}$  for  $\bar{y} \in \mathbb{F}$ . Then  $R^\square(\bar{\rho}, \tau_{1,s}) = R^\square(\bar{\rho})$  is formally smooth of relative dimension 4 over  $\mathcal{O}$ , while  $R^\square(\bar{\rho}, \tau_{1,ns}) = 0$ .*

(2) *Suppose that  $\bar{\rho}(\sigma) = \begin{pmatrix} 1 & \bar{x} \\ 0 & 1 \end{pmatrix}$  and  $\bar{\rho}(\phi) = \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix}$ .*

*If  $\bar{x} \neq 0$ , then  $R^\square(\bar{\rho}, \tau_{1,ns}) = R^\square(\bar{\rho})$  is formally smooth of relative dimension 4 over  $\mathcal{O}$ , while  $R^\square(\bar{\rho}, \tau_{1,s}) = 0$ .*

*If  $\bar{x} = 0$  then*

$$R^\square(\bar{\rho}) \cong \mathcal{O}[[X_1, \dots, X_5]] / (X_1 X_2).$$

*The quotients by the two minimal primes are  $R^\square(\bar{\rho}, \tau_{1,s})$  and  $R^\square(\bar{\rho}, \tau_{1,ns})$ , so that both are formally smooth of relative dimension 4 over  $\mathcal{O}$ . The minimal primes  $\mathfrak{a}_{nr}$  and  $\mathfrak{a}_N$  of  $\bar{R}^\square(\bar{\rho})$  which respectively define  $\bar{R}^\square(\bar{\rho}, \tau_{1,s})$  and  $\bar{R}^\square(\bar{\rho}, \tau_{1,ns})$  are distinct.*



*Proof.* For the first part, write

$$\rho^\square(\sigma) = \begin{pmatrix} 1+A & B \\ C & 1+D \end{pmatrix}, \quad \rho^\square(\phi) = \begin{pmatrix} 1+P & y+R \\ S & 1+Q \end{pmatrix},$$

where  $y$  is a lift of  $\bar{y}$  (taken to be zero if  $\bar{y} = 0$ ) and  $A, B, C, D, P, Q, R, S \in \mathfrak{m}$ .

Let  $I = (A, B, C, D)$ . Considering the equation  $\rho^\square(\phi)\rho^\square(\sigma) = \rho^\square(\sigma)^q\rho^\square(\phi)$  modulo the ideal  $I\mathfrak{m}$  gives equations  $Cy \equiv (q - 1)A, B + Dy \equiv qAy + qB, C \equiv qC,$  and  $(q - 1)D + qCy \equiv 0,$  all modulo  $I\mathfrak{m}$ . As  $q \not\equiv 1 \pmod{l}$  we find that  $I = I\mathfrak{m}$ . Therefore, by Nakayama’s lemma,  $I = 0$  and  $\rho^\square$  is unramified. So  $R^\square(\bar{\rho}) = R^\square(\bar{\rho}, \tau_{1,s}) \cong \mathcal{O}[[P, Q, R, S]]$  as claimed. Note that this proof is still valid if  $q \equiv -1 \pmod{l}$ .

The proof of the second part is similar. By Lemma 5.4, we may write

$$\rho^\square(\sigma) = \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1+A & x+B \\ C & 1+D \end{pmatrix} \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix},$$

$$\rho^\square(\phi) = \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}^{-1} \begin{pmatrix} q(1+P) & 0 \\ 0 & 1+Q \end{pmatrix} \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix},$$

with  $x$  a lift of  $\bar{x}$  (taken to be zero if  $\bar{x} = 0$ ) and  $A, B, C, D, X, Y, P, Q \in \mathfrak{m}$ .

Let  $I = (A, C, D)$ . Considering the relation  $\phi\sigma\phi^{-1} = \sigma^q$  modulo  $I\mathfrak{m}$  and applying Nakayama’s lemma as before now yields  $A = C = D = 0$  (using that  $q^2 \not\equiv 1 \pmod{l}$ ). The relation (not modulo any ideal) gives that  $(x + B)(P - Q) = 0,$  and it is easy to see if this equality holds then the given formulae for  $\rho^\square$  do indeed define a representation so that

$$R^\square(\bar{\rho}) = \frac{\mathcal{O}[[B, P, Q, X, Y]]}{((x + B)(P - Q))}.$$

If  $\bar{x} \neq 0$  then this implies that  $P = Q$ . Then  $R^\square(\bar{\rho}) = \mathcal{O}[[B, P, X, Y]]$ . It is clear that  $R^\square(\bar{\rho}) = R^\square(\bar{\rho}, \tau_{1,ns})$ , and the proposition follows.

If  $\bar{x} = 0$  then, writing  $U = P - Q$ , we have  $R^\square(\bar{\rho}) = \mathcal{O}[[B, P, U, X, Y]]/(BU)$ . In these coordinates, it is clear from the description of  $\rho^\square$  that

$$R^\square(\bar{\rho}, \tau_{1,s}) = R^\square(\bar{\rho})/(B) \quad \text{and} \quad R^\square(\bar{\rho}, \tau_{1,ns}) = R^\square(\bar{\rho})/(U).$$

The proposition follows. □

**5C.  $q \equiv -1 \pmod{l}$ .** Suppose that  $q \equiv -1 \pmod{l}$ . By Proposition 5.3 we have already dealt with the cases in which the eigenvalues of  $\bar{\rho}(\phi)$  are not in the ratio 1 or  $-1$ . All other cases are dealt with by the following result (after twisting and conjugating  $\bar{\rho}$ ). By Lemma 3.7, the only possible types when  $\bar{\rho}|_{\tilde{\rho}_F}$  is trivial are  $\tau_{1,s}, \tau_{1,ns},$  and  $\tau_\xi$  for  $\xi$  a nontrivial  $l^b$ -th root of unity.

**Proposition 5.6.** *Suppose that  $q \equiv -1 \pmod{l}$  and that  $\bar{\rho}|_{\tilde{\rho}_F}$  is trivial.*

(1) Suppose that  $\bar{\rho}(\sigma) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $\bar{\rho}(\phi) = \begin{pmatrix} 1 & \bar{y} \\ 0 & 1 \end{pmatrix}$  for  $\bar{y} \in \mathbb{F}$ . Then

$$R^\square(\bar{\rho}, \tau_{1,s}) = R^\square(\bar{\rho})$$

is formally smooth of relative dimension 4 over  $\mathcal{O}$ , while

$$R^\square(\bar{\rho}, \tau_{1,ns}) = R^\square(\bar{\rho}, \tau_\xi) = 0.$$

If  $\mathfrak{a}_{nr}$  is the unique minimal prime of  $\bar{R}^\square(\bar{\rho})$ , then we have

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} [\mathfrak{a}_{nr}] & \text{if } \tau = \tau_{1,s}, \\ 0 & \text{if } \tau = \tau_{1,ns}, \\ 0 & \text{if } \tau = \tau_\xi. \end{cases}$$

(2) Suppose that  $\bar{\rho}(\sigma) = \begin{pmatrix} 1 & \bar{x} \\ 0 & 1 \end{pmatrix}$  and  $\bar{\rho}(\phi) = \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix}$  for  $\bar{x} \in \mathbb{F}$ .

(a) If  $\bar{x} \neq 0$ , then  $R^\square(\bar{\rho}, \tau_{1,ns})$  and  $R^\square(\bar{\rho}, \tau_\xi)$  are formally smooth of relative dimension 4 over  $\mathcal{O}$ , while  $R^\square(\bar{\rho}, \tau_{1,s}) = 0$ . If  $\mathfrak{a}_N$  is the prime ideal of  $\bar{R}^\square(\bar{\rho})$  cutting out  $\bar{R}^\square(\bar{\rho}, \tau_{1,ns})$  then we have

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} 0 & \text{if } \tau = \tau_{1,s}, \\ [\mathfrak{a}_N] & \text{if } \tau = \tau_{1,ns}, \\ [\mathfrak{a}_N] & \text{if } \tau = \tau_\xi. \end{cases} \tag{4}$$

(b) If  $\bar{x} = 0$ , then  $R^\square(\bar{\rho}, \tau_{1,s})$  is formally smooth of relative dimension 4 over  $\mathcal{O}$  and

$$R^\square(\bar{\rho}, \tau_{1,ns}) \cong \frac{\mathcal{O}[[X_1, \dots, X_6]]}{((X_1, X_3) \cap (X_2, X_3 - (q+1)))}$$

is a non-Cohen–Macaulay ring of relative dimension 4 over  $\mathcal{O}$ . Its spectrum is the scheme theoretic union of two formally smooth components that do not intersect in the generic fibre. Lastly,

$$R^\square(\bar{\rho}, \tau_\xi) \cong \frac{\mathcal{O}[[X_1, \dots, X_5]]}{(X_1 X_2 - (\xi - \xi^{-1})^2)}$$

is a complete intersection domain of relative dimension 4 over  $\mathcal{O}$  with formally smooth generic fibre. If  $\mathfrak{a}_{nr}$  is the prime of  $\bar{R}^\square(\bar{\rho})$  corresponding to  $\bar{R}^\square(\bar{\rho}, \tau_{1,s})$  and  $\mathfrak{a}_N, \mathfrak{a}'_N$  are the prime ideals of  $\bar{R}^\square(\bar{\rho})$  corresponding to the two minimal primes of  $\bar{R}^\square(\bar{\rho}, \tau_{1,ns})$ , then we have

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} [\mathfrak{a}_{nr}] & \text{if } \tau = \tau_{1,s}. \\ [\mathfrak{a}_N] + [\mathfrak{a}'_N] & \text{if } \tau = \tau_{1,ns}, \\ [\mathfrak{a}_N] + [\mathfrak{a}'_N] & \text{if } \tau = \tau_\xi. \end{cases} \tag{5}$$

*Proof.* The proof of the first part is identical to that of [Proposition 5.5](#), part 1.

For the second part, by [Lemma 5.4](#) we may write

$$\rho^\square(\sigma) = \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix} \begin{pmatrix} 1+A & x+B \\ C & 1+D \end{pmatrix} \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix},$$

$$\rho^\square(\phi) = \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix} \begin{pmatrix} -(1+P) & 0 \\ 0 & 1+Q \end{pmatrix} \begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix},$$

with  $x$  a lift of  $\bar{x}$  (taken to be zero if  $\bar{x} = 0$ ) and  $A, B, C, D, X, Y, P, Q \in \mathfrak{m}$ .

Firstly, it is clear that  $R^\square(\bar{\rho}, \tau_{1,s}) = 0$  if  $\bar{x} \neq 0$  and, if  $\bar{x} = 0$ , that

$$R^\square(\bar{\rho}, \tau_{1,s}) \cong \mathcal{O}[[P, Q, X, Y]].$$

Next we deal with  $\tau_{1,ns}$ . On  $R^\square(\bar{\rho}, \tau_{1,ns})$  we have the equations

$$\begin{aligned} \text{tr}(\rho^\square(\sigma)) &= 2, \\ \det(\rho^\square(\sigma)) &= 1, \\ q \text{tr}(\rho(\phi))^2 &= (q + 1)^2 \det(\rho(\phi)), \end{aligned}$$

and

$$\rho^\square(\phi)\rho^\square(\sigma)\rho^\square(\phi)^{-1} = \rho^\square(\sigma)^q.$$

The first two of these may be rewritten as

$$A = -D \quad \text{and} \quad A^2 + (x + B)(C) = 0$$

and the third can be written as

$$(q + 1 + P + qQ)(q + 1 + Q + qP) = 0.$$

By the Cayley–Hamilton theorem,  $(\rho^\square(\sigma) - 1)^2 = 0$  on  $R^\square(\bar{\rho}, \tau_{1,ns})^\circ$ ; it follows that  $\rho^\square(\sigma)^q - 1 = q(\rho^\square(\sigma) - 1)$  on  $R^\square(\bar{\rho}, \tau_{1,ns})^\circ$  and so the relation  $\phi\sigma\phi^{-1} = \sigma^q$  together with  $D = -A$  yields the equation

$$\begin{pmatrix} A & -(x+B)\frac{1+P}{1+Q} \\ -C\frac{1+Q}{1+P} & -A \end{pmatrix} = \begin{pmatrix} qA & q(x+B) \\ qC & -qA \end{pmatrix}.$$

Equating coefficients and using that 2 and  $q - 1$  are invertible we obtain that  $A = D = 0$  and that

$$(x + B)(q + 1 + qQ + P) = 0, \tag{6}$$

$$C(q + 1 + Q + qP) = 0, \tag{7}$$

$$(x + B)C = 0, \tag{8}$$

$$(q + 1 + Q + qP)(q + 1 + qQ + P) = 0 \tag{9}$$

is a complete set of equations cutting out  $R^\square(\bar{\rho}, \tau_{1,ns})^\circ$  (the last two equations being, respectively, the conditions on  $\det(\rho^\square(\sigma))$  and on  $\rho^\square(\phi)$ ).

If  $\bar{x} \neq 0$  then these equations are equivalent to  $q + 1 + qQ + P = 0$  and  $C = 0$  and so we see that

$$R^\square(\bar{\rho}, \tau_{1,ns}) \cong \mathcal{O}[[B, P, X, Y]].$$

If  $\bar{x} = 0$  then the left hand sides of the four equations given generate the ideal

$$I = (B, q + 1 + Q + qP) \cap (C, q + 1 + qQ + P)$$

in  $\mathcal{O}[[B, C, P, Q, X, Y]]$ . Since  $\mathcal{O}[[B, C, P, Q, X, Y]]/I$  is reduced and  $\lambda$ -torsion free and a Zariski dense set of its  $\bar{E}$ -points have type  $\tau_{1,ns}$ , it is equal to  $R^\square(\bar{\rho}, \tau_{1,ns})$ . After the change of variables

$$X_3 = \frac{q(q + 1 + Q + qP)}{(q - 1)(1 + P)}, \quad (X_1, X_2, X_4, X_5, X_6) = (B, C, P, X, Y)$$

we get the presentation given in the proposition.

Let

$$\mathcal{S} = \frac{\mathcal{O}[[X_1, X_2, X_3]]}{(X_1, X_3) \cap (X_2, X_3 - (q + 1))}.$$

Then  $\mathcal{S}$  has dimension two. We show that  $\mathcal{S}$  is not Cohen–Macaulay; the same is then true for  $R^\square(\bar{\rho}, \tau_{1,ns})$ . Now  $\lambda$  is a nonzerodivisor in  $\mathcal{S}$ , and

$$\mathcal{S}/\lambda = \frac{\mathbb{F}[[X_1, X_2, X_3]]}{(X_1X_2, X_1X_3, X_2X_3, X_3^2)}.$$

The maximal ideal of  $\mathcal{S}/\lambda$  is annihilated by  $X_3$ , and  $X_3 \neq 0$  in  $\mathcal{S}/\lambda$ . So  $\mathcal{S}/\lambda$ , and hence  $\mathcal{S}$ , is not Cohen–Macaulay. The remaining statements about  $R^\square(\bar{\rho}, \tau_{1,ns})$  are clear.

Now suppose that  $\tau = \tau_\xi$ . On  $R^\square(\bar{\rho}, \tau_\xi)$  we have

$$\begin{aligned} \text{tr}(\rho^\square(\sigma)) &= \xi + \xi^{-1}, \\ \det(\rho^\square(\sigma)) &= 1, \end{aligned}$$

and

$$\rho^\square(\phi)\rho^\square(\sigma)\rho^\square(\phi)^{-1} = \rho^\square(\sigma)^q.$$

The first two of these may be rewritten as

$$A + D = \xi + \xi^{-1} - 2 \quad \text{and} \quad AD - (x + B)C = 2 - \xi - \xi^{-1}.$$

By the Cayley–Hamilton theorem,  $(\rho^\square(\sigma) - \xi)(\rho^\square(\sigma) - \xi^{-1}) = 0$ . As

$$T^q \equiv \xi + \xi^{-1} - T \pmod{(T - \xi)(T - \xi^{-1})}$$

in  $\mathbb{Z}[T]$ , the relation  $\phi\sigma\phi^{-1} = \sigma^q$  yields

$$\begin{pmatrix} 1+A & -(x+B)\frac{1+P}{1+Q} \\ -C\frac{1+Q}{1+P} & 1+D \end{pmatrix} = \begin{pmatrix} \xi + \xi^{-1} - 1 - A & -(x+B) \\ -C & \xi + \xi^{-1} - 1 - D \end{pmatrix}.$$

Equating coefficients and combining with the equation  $\det(\rho^\square(\sigma)) = 1$  we get

$$A = D = \frac{\xi + \xi^{-1}}{2} - 1, \tag{10}$$

$$(x + B)(P - Q) = 0, \tag{11}$$

$$C(P - Q) = 0, \tag{12}$$

$$4(x + B)C = (\xi - \xi^{-1})^2. \tag{13}$$

If  $\bar{x} \neq 0$  then these equations are equivalent to  $P = Q$  and  $C = (\xi - \xi^{-1})^2 / (4(x + B))$ , so that

$$R^\square(\bar{\rho}, \tau_\xi) \cong \mathcal{O}[[X, Y, B, P]].$$

If  $\bar{x} = 0$ , then the equations imply that

$$0 = BC(P - Q) = \left(\frac{\xi - \xi^{-1}}{2}\right)^2 (P - Q)$$

and hence that  $P = Q$ , as  $R^\square(\bar{\rho}, \tau_\xi)$  is  $\lambda$ -torsion free by definition. Thus

$$R^\square(\bar{\rho}, \tau_\xi) \cong \frac{\mathcal{O}[[X, Y, B, C, P]]}{(4BC - (\xi - \xi^{-1})^2)}.$$

The remaining statements about  $R^\square(\bar{\rho}, \tau_\xi)$  are clear.

Now we calculate the various  $Z^4(\bar{R}^\square(\bar{\rho}, \tau))$ . For part 1, this is trivial. For part 2, we have computed each  $\bar{R}^\square(\bar{\rho}, \tau)$  as a quotient of the ring  $\mathbb{F}[[A, B, C, D, P, Q, X, Y]]$  by an ideal which we call  $I(\tau)$ . We see that if  $\bar{x} \neq 0$  then  $I(\tau_{1,ns}) = I(\tau_\xi)$ , and  $\bar{R}^\square(\bar{\rho}, \tau_{1,s}) = 0$ , from which (4) follows. If  $\bar{x} = 0$  then

$$I(\tau_{1,s}) = (A, B, C, D),$$

$$I(\tau_{1,ns}) = (A, D, BC, B(Q - P), C(Q - P), (Q - P)^2),$$

and

$$I(\tau_\xi) = (A, D, BC, Q - P).$$

The minimal primes above these  $I(\tau)$  in  $\mathbb{F}[[A, \dots, Y]]$  are  $\mathfrak{a}_{nr} = (A, B, C, D)$ ,  $\mathfrak{a}_N = (A, C, D, Q - P)$  and  $\mathfrak{a}_{N'} = (A, B, D, Q - P)$ ; the multiplicities in (5) are then easily verified. □

**Remark 5.7.** When  $\bar{\rho}$  is unramified and  $\bar{\rho}(\phi) = \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix}$ , the ring  $R^\square(\bar{\rho}, \tau_{1,ns})$  is not Cohen–Macaulay. However the ring  $R^\square(\bar{\rho}, \text{unip})$ , which is defined to be the maximal reduced quotient of  $R^\square(\bar{\rho})$  on which  $\rho^\square(\sigma)$  is unipotent (so that  $\text{Spec } R^\square(\bar{\rho}, \text{unip})$  is the scheme-theoretic union in  $\text{Spec } R^\square(\bar{\rho})$  of  $\text{Spec } R^\square(\bar{\rho}, \tau_{1,s})$  with  $\text{Spec } R^\square(\bar{\rho}, \tau_{1,ns})$ ), is Cohen–Macaulay. Indeed it is easy to see from the proof that

$$R^\square(\bar{\rho}, \text{unip}) \cong \frac{\mathcal{O}[[X_1, \dots, X_6]]}{(X_1 X_2, X_1(X_3 - (q + 1)), X_2 X_3)},$$

which is Cohen–Macaulay ( $(\lambda, X_1 + X_2 + X_3, X_4, X_5, X_6)$  is a regular sequence).

**5D.  $q \equiv 1 \pmod{l}$ .** Suppose that  $q \equiv 1 \pmod{l}$ . By Proposition 5.3 we have already dealt with the cases in which the eigenvalues of  $\bar{\rho}(\phi)$  are distinct. All other cases are dealt with by the following (after twisting and conjugating  $\bar{\rho}$ ). Note that by Lemma 3.7, the only possible types when  $\bar{\rho}|_{\bar{\rho}_F}$  is trivial are  $\tau_{\zeta,s}$ ,  $\tau_{\zeta,ns}$  and  $\tau_{\zeta_1,\zeta_2}$  for  $\zeta$  any  $l^a$ -th root of unity and  $\zeta_1, \zeta_2$  any distinct  $l^a$ -th roots of unity.

**Proposition 5.8.** *Suppose that  $q \equiv 1 \pmod{l}$  and that  $\bar{\rho}|_{\bar{\rho}_F}$  is trivial. Suppose that  $\bar{\rho}(\sigma) = \begin{pmatrix} 1 & \bar{x} \\ 0 & 1 \end{pmatrix}$  and  $\bar{\rho}(\phi) = \begin{pmatrix} 1 & \bar{y} \\ 0 & 1 \end{pmatrix}$  for  $\bar{x}, \bar{y} \in \mathbb{F}$ .*

(1) *If  $\bar{x} \neq 0$  then  $R^\square(\bar{\rho}, \tau_{\zeta,s}) = 0$ , while  $R^\square(\bar{\rho}, \tau_{\zeta,ns})$  and  $R^\square(\bar{\rho}, \tau_{\zeta_1,\zeta_2})$  are formally smooth over  $\mathcal{O}$  of relative dimension 4.*

*If  $\mathfrak{a}_N$  is the four-dimensional prime of  $\bar{R}^\square(\bar{\rho})$  corresponding to  $\bar{R}^\square(\bar{\rho}, \tau_{1,ns})$  then we have*

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} 0 & \text{if } \tau = \tau_{\zeta,s}, \\ [\mathfrak{a}_N] & \text{if } \tau = \tau_{\zeta,ns}, \\ [\mathfrak{a}_N] & \text{if } \tau = \tau_{\zeta_1,\zeta_2}. \end{cases} \tag{14}$$

(2) *If  $\bar{x} = 0$  and  $\bar{y} \neq 0$ , then  $R^\square(\bar{\rho}, \tau_{\zeta,s})$  and  $R^\square(\bar{\rho}, \tau_{\zeta,ns})$  are formally smooth over  $\mathcal{O}$  of relative dimension 4 while*

$$R^\square(\bar{\rho}, \tau_{\zeta_1,\zeta_2}) \cong \mathcal{O}[[X_1, \dots, X_5]]/(X_1^2 X_2 - (\zeta_1 - \zeta_2)^2)$$

*is a complete intersection domain of relative dimension 4 over  $\mathcal{O}$ .*

*If  $\mathfrak{a}_{nr}$  and  $\mathfrak{a}_N$  are the prime ideals of  $\bar{R}^\square(\bar{\rho})$  corresponding to  $\bar{R}^\square(\bar{\rho}, \tau_{1,s})$  and  $\bar{R}^\square(\bar{\rho}, \tau_{1,ns})$  respectively, then*

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} [\mathfrak{a}_{nr}] & \text{if } \tau = \tau_{\zeta,s}, \\ [\mathfrak{a}_N] & \text{if } \tau = \tau_{\zeta,ns}, \\ 2[\mathfrak{a}_{nr}] + [\mathfrak{a}_N] & \text{if } \tau = \tau_{\zeta_1,\zeta_2}. \end{cases} \tag{15}$$

(3) *If  $\bar{x} = \bar{y} = 0$ , then  $R^\square(\bar{\rho}, \tau_{\zeta,s})$  is formally smooth over  $\mathcal{O}$  of relative dimension 4 while  $R^\square(\bar{\rho}, \tau_{\zeta,ns})$  and  $R^\square(\bar{\rho}, \tau_{\zeta_1,\zeta_2})$  are non-Gorenstein Cohen–Macaulay domains of relative dimension 4 over  $\mathcal{O}$ .*

*Both  $\bar{R}^\square(\bar{\rho}, \tau_{\zeta,s})$  and  $\bar{R}^\square(\bar{\rho}, \tau_{\zeta,ns})$  are domains; let the corresponding primes of  $\bar{R}^\square(\bar{\rho})$  be  $\mathfrak{a}_{nr}$  and  $\mathfrak{a}_N$  respectively. Then*

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau)) = \begin{cases} [\mathfrak{a}_{nr}] & \text{if } \tau = \tau_{\zeta,s}, \\ [\mathfrak{a}_N] & \text{if } \tau = \tau_{\zeta,ns}, \\ 2[\mathfrak{a}_{nr}] + [\mathfrak{a}_N] & \text{if } \tau = \tau_{\zeta_1,\zeta_2}. \end{cases} \tag{16}$$

*Proof.* Write

$$\rho^\square(\sigma) = \begin{pmatrix} 1+A & x+B \\ C & 1+D \end{pmatrix}, \quad \rho^\square(\phi) = \begin{pmatrix} 1+P & y+R \\ S & 1+Q \end{pmatrix}$$

with  $A, B, C, D, P, Q, R,$  and  $S \in \mathfrak{m}$  and  $x, y$  lifts of  $\bar{x}, \bar{y}$  (taken to be zero if  $\bar{x}$  or  $\bar{y} = 0$ ).

First, we have that  $R^\square(\bar{\rho}, \tau_{\zeta,s}) = 0$  if  $\bar{x} \neq 0$  and

$$R^\square(\bar{\rho}, \tau_{\zeta,s}) \cong \mathcal{O}[[P, Q, R, S]]$$

otherwise.

Next, we look at  $R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})$  for  $\zeta_1$  and  $\zeta_2$  distinct  $l^a$ -th roots of unity. The condition that  $\rho^\square(\sigma)$  has characteristic polynomial  $(t - \zeta_1)(t - \zeta_2)$  is equivalent to the equations

$$A + D = \zeta_1 + \zeta_2 - 2 \quad \text{and} \quad AD - (x + B)C = (\zeta_1 - 1)(\zeta_2 - 1).$$

Since  $(t - \zeta_1)(t - \zeta_2) \mid t^{q-1} - 1$ , by the Cayley–Hamilton theorem we have

$$\rho^\square(\sigma)^q = \rho^\square(\sigma)$$

on  $R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})^\circ$ . So the relation  $\phi\sigma\phi^{-1} = \sigma^q$  yields

$$\begin{pmatrix} 1+A & x+B \\ C & 1+D \end{pmatrix} \begin{pmatrix} 1+P & y+R \\ S & 1+Q \end{pmatrix} = \begin{pmatrix} 1+P & y+R \\ S & 1+Q \end{pmatrix} \begin{pmatrix} 1+A & x+B \\ C & 1+D \end{pmatrix}.$$

Equating coefficients, eliminating  $D$  and writing  $U = P - Q$  and  $F = A - D = 2A - (\zeta_1 + \zeta_2 - 2)$  we see that  $R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})$  is the reduced,  $l$ -torsion-free quotient of  $\mathcal{O}[[B, C, F, P, R, S, U]]$  by the relations

$$(x + B)S = (y + R)C, \tag{17}$$

$$F(y + R) = U(x + B), \tag{18}$$

$$FS = UC, \tag{19}$$

$$(\zeta_1 - \zeta_2)^2 = F^2 + 4(x + B)C. \tag{20}$$

If  $\bar{x} \neq 0$  then these equations are equivalent to  $U = F(y + R)(x + B)^{-1}$ ,  $C = \frac{1}{4}((\zeta_1 - \zeta_2)^2 - F^2)(x + B)^{-1}$ , and  $S = C(y + R)(x + B)^{-1}$  so that

$$R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2}) \cong \mathcal{O}[[B, F, P, R]].$$

If  $\bar{x} = 0$  and  $\bar{y} \neq 0$ , then  $F = BU(y + R)^{-1}$  and  $C = BS(y + R)^{-1}$  will be a solution to equations (17)–(20) provided that

$$(\zeta_1 - \zeta_2)^2 = \left(\frac{B}{y + R}\right)^2 (U^2 + 4(y + R)S).$$

Writing  $(X_1, \dots, X_5) = (B(y + R)^{-1}, U^2 + 4(y + R)S, P, R, U)$  we get

$$R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2}) \cong \frac{\mathcal{O}[[X_1, \dots, X_5]]}{X_1^2 X_2 - (\zeta_1 - \zeta_2)^2}$$



as claimed. The other statements about  $R^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2})$  follow easily.

If  $\bar{x} = \bar{y} = 0$ , then let  $\mathcal{A} = \mathcal{O}[[B, C, F, P, R, S, U]]$  and  $I \triangleleft \mathcal{A}$  be the ideal

$$I = ((\zeta_1 - \zeta_2)^2 - F^2 - 4BC, BS - CR, FR - BU, FS - CU).$$

Note that the ideal

$$J = (BS - CR, FR - BU, FS - CU)$$

is generated by the  $2 \times 2$  minors of  $\begin{pmatrix} B & C & F \\ R & S & U \end{pmatrix}$ . So, by [Proposition 2.7](#),  $\mathcal{A}/J$  is a Cohen–Macaulay, non-Gorenstein domain. Since  $F^2 - 4BC$  is not zero in the domain  $\mathcal{A}/J \otimes \mathbb{F}$ ,  $(\lambda, F^2 - 4BC)$  is a regular sequence in  $\mathcal{A}/J$ . Hence  $(F^2 - 4BC - (\zeta_1 - \zeta_2)^2, \lambda)$  is a regular sequence in  $\mathcal{A}/J$ , and therefore  $\mathcal{A}/I$  is  $\mathcal{O}$ -flat, Cohen–Macaulay and non-Gorenstein. It is reduced because it is Cohen–Macaulay and, as we shall show in the next paragraph, generically reduced.

To show that  $\mathcal{A}/I$  is irreducible, it suffices to show that  $\mathcal{X} = \text{Spec}(\mathcal{A}/I \otimes E)$  is irreducible. This follows if we can show that  $\mathcal{X}$  is formally smooth and connected. As  $F^2 - 4BC \neq 0$  on  $\mathcal{X}$ , it is covered by the affine open subsets  $\mathcal{U}_B = \{B \neq 0\}$  and  $\mathcal{U}_F = \{F \neq 0\}$ . By the argument used in the  $\bar{x} \neq 0$  case,  $\mathcal{U}_B$  is formally smooth. A similar argument works for  $\mathcal{U}_F$ , the projection map

$$p : \mathcal{X} \rightarrow \text{Spec}\left(\frac{\mathcal{O}[[F, B, C, U, P]]}{(F^2 + 4BC - (\zeta_1 - \zeta_2)^2)} \otimes E\right)$$

is an isomorphism from  $\mathcal{U}_F$  onto an open subscheme, but the right hand side is easily seen to be formally smooth. Hence  $\mathcal{X}$  is formally smooth. Note that the composition of the map  $p$  with the projection away from  $U$  is a continuous map with connected fibres and connected image, which admits a continuous section (obtained by taking  $R = S = U = 0$ ); it follows that  $\mathcal{X}$  is connected, as required. Since  $\mathcal{X}$  is formally smooth it is certainly reduced; therefore  $\mathcal{A}/I$  is generically reduced (as it is  $\mathcal{O}$ -flat), just as we claimed above.

Now we turn to  $R^\square(\bar{\rho}, \tau_{\zeta, ns})$ . By [Lemma 2.4](#) we may assume that  $\zeta = 1$ . The condition that the characteristic polynomial of  $\rho^\square(\sigma)$  be  $(t - 1)^2$  is equivalent to the equations

$$A + D = 0 \quad \text{and} \quad AD - (x + B)C = 0.$$

Writing  $T = P + Q$  and  $U = P - Q$  the condition that

$$q \text{tr}(\rho^\square(\phi))^2 = (q + 1)^2 \det(\rho^\square(\phi))$$

becomes

$$(q - 1)^2(T + 2)^2 = (q + 1)^2(U^2 + 4(y + R)S).$$

Since  $t^q - 1 \equiv q(t - 1) \pmod{(t - 1)^2}$ , the Cayley–Hamilton theorem shows that

$$\rho^\square(\sigma)^q - 1 = q(\rho^\square(\sigma) - 1)$$

on  $R^\square(\bar{\rho}, \tau_{1,ns})$ . From  $\phi\sigma\phi^{-1} = \sigma^q$  we therefore get the equation

$$(\phi - 1)(\sigma - 1) - (\sigma - 1)(\phi - 1) = (q - 1)(\sigma - 1)\phi$$

on  $R^\square(\bar{\rho}, \tau_{1,ns})$ . Equating coefficients and substituting  $D = -A$  we get the equations

$$A^2 + (x + B)C = 0, \tag{21}$$

$$(q - 1)^2(T + 2)^2 = (q + 1)^2(U^2 + 4(y + R)S), \tag{22}$$

$$C(y + R) - S(x + B) = (q - 1)(A(1 + P) + (x + B)S), \tag{23}$$

$$U(x + B) - 2A(y + R) = (q - 1)(A(y + R) + (x + B)(1 + Q)), \tag{24}$$

$$2AS - CU = (q - 1)(C(1 + P) - AS), \tag{25}$$

$$S(x + B) - C(y + R) = (q - 1)(C(y + R) - A(1 + Q)). \tag{26}$$

After replacing  $P$  with  $(T + U)/2$  and  $Q$  with  $(T - U)/2$ , this is a complete set of equations for  $R^\square(\bar{\rho}, \tau_{1,ns})$  in  $\mathcal{O}[[A, B, C, R, S, T, U]]$ .

We replace equations (23) and (26) by their sum and difference

$$(q - 1)(AU + (x + B)S + C(y + R)) = 0, \tag{27}$$

$$(q + 1)(C(y + R) - (x + B)S) = (q - 1)A(2 + T). \tag{28}$$

As  $R^\square(\bar{\rho}, \tau_{1,ns})$  is  $\lambda$ -torsion free, (27) implies that

$$AU + (x + B)S + C(y + R) = 0. \tag{29}$$

We could also write this equation as  $\text{tr}((\sigma - 1)\phi) = 0$ .

Putting  $\alpha(T) = ((q - 1)(2 + T))/(q + 1)$ , we find that (21), (22), (24), (25) and (28)+(29) may respectively be rewritten as

$$\begin{aligned} A^2 + (x + B)C &= 0, \\ 4(y + R)S + (U - \alpha(T))(U + \alpha(T)) &= 0, \\ 2A(y + R) - (x + B)(U - \alpha(T)) &= 0, \\ 2AS - C(U + \alpha(T)) &= 0, \\ 2C(y + R) + A(U - \alpha(T)) &= 0, \\ 2(x + B)S + A(U + \alpha(T)) &= 0. \end{aligned}$$

Let  $I$  be the ideal of  $\mathcal{O}[[A, B, C, R, S, T, U]]$  generated by these equations and let  $R' = \mathcal{O}[[A, B, C, R, S, T, U]]/I$ , so that  $R^\square(\bar{\rho}, \tau_{1,ns})$  is the maximal reduced  $l$ -torsion free quotient of  $R'$ .

If  $\bar{x} \neq 0$  then  $C$ ,  $U$ , and  $S$  are uniquely determined by  $A$ ,  $B$ ,  $R$ , and  $T$  so that

$$R^\square(\bar{\rho}, \tau_{1,ns}) \cong \mathcal{O}[[A, B, R, T]].$$

If  $\bar{y} \neq 0$ , then  $S$ ,  $C$ , and  $A$  are uniquely determined by  $B$ ,  $R$ ,  $T$ , and  $U$  so that

$$R^\square(\bar{\rho}, \tau_{1,ns}) \cong \mathcal{O}[[B, R, T, U]].$$

If  $\bar{x} = \bar{y} = 0$ , so that  $x = y = 0$ , observe that

$$R' \cong \frac{\mathcal{B}}{J_0 + J_1},$$

where

$$\mathcal{B} = \mathcal{O}[[X_1, \dots, X_4, Y_1, \dots, Y_4, T]],$$

the ideal  $J_0$  is generated by the  $2 \times 2$  minors of

$$\begin{pmatrix} X_1 & X_2 & X_3 & X_4 \\ Y_1 & Y_2 & Y_3 & Y_4 \end{pmatrix}$$

and  $J_1 = (X_1 + Y_2, X_3 - Y_4 + 2(q-1)/(q+1))$ . (The change of variables is  $X_1 = A$ ,  $X_2 = B$ ,  $Y_1 = C$ ,  $Y_2 = -A$ ,  $X_3 = -2R/(2+T)$ ,  $Y_4 = 2S(2+T)$ ,  $Y_3 = (U - \alpha(T))/(2+T)$ , and  $X_4 = (U + \alpha(T))/(2+T)$ .) Then, by [Proposition 2.7](#),  $\mathcal{B}/J_0$  is a Cohen–Macaulay non-Gorenstein domain. Moreover,  $(\lambda, X_1 + Y_2, X_3 - Y_4)$  may be checked to be a regular sequence on  $\mathcal{B}/J_0$ . Therefore  $(X_1 + Y_2, X_3 + Y_4 + 2(q-1)/(q+1), \lambda)$  is also regular, and so  $\mathcal{B}/(J_0 + J_1)$  is Cohen–Macaulay,  $\mathcal{O}$ -flat, and not Gorenstein. The same is then true for  $R'$ .

We show that  $R' \otimes \mathbb{F}$  is a domain, which implies that  $R'$  is a domain. Let  $\bar{I}$  be the image of  $I$  in  $\mathbb{F}[[A, B, C, R, S, T, U]]$ . Then  $\bar{I}$  is homogeneous so  $\text{gr}(R' \otimes \mathbb{F}) = \mathbb{F}[A, B, C, R, S, T, U]/\bar{I}$  and it suffices to check that this is a domain (by [Eisenbud 1995](#), Corollary 5.5). It is therefore sufficient to check that  $\text{Proj}(\text{gr}(R' \otimes \mathbb{F}))$  is reduced and irreducible. But it is easy to check this on the usual seven affine pieces. This argument is from [Taylor 2009](#).

Next we show that  $R^\square(\bar{\rho}, \tau_{1,ns})$  is reduced. In fact, we show that

$$\mathcal{Y} = \text{Spec}(R^\square(\bar{\rho}, \tau_{1,ns}) \otimes E)$$

is formally smooth, which implies that  $R^\square(\bar{\rho}, \tau_{1,ns})$  is reduced because it is Cohen–Macaulay and  $\mathcal{O}$ -flat. For  $\star = B, C, R, S, U - \alpha(T)$ , or  $U + \alpha(T)$  let  $\mathcal{U}_\star = \{\star \neq 0\} \subset \mathcal{Y}$  be the corresponding affine open subscheme. Then the  $\mathcal{U}_\star$  are an affine open cover of  $\mathcal{Y}$ . For  $\star = B, C, R$  or  $S$  we see that  $\mathcal{U}_\star$  is formally smooth by the same argument as for the cases  $\bar{x} \neq 0$  and  $\bar{y} \neq 0$  above. For  $\mathcal{U}_{U \pm \alpha(T)}$ , the projection morphism

$$p : \mathcal{U}_{U - \alpha(T)} \rightarrow \text{Spec}\left(\frac{\mathcal{O}[[C, R, S, T]]}{4RS - (U + \alpha(T))(U - \alpha(T))} \otimes E\right)$$

is an isomorphism onto an open subscheme. But the right hand scheme is easily seen to be formally smooth as required.

Finally we calculate the  $Z^4(\bar{R}(\bar{\rho}, \tau))$ . We do this when  $\bar{x} = \bar{y} = 0$ , as the other cases are similar but easier. We have written each  $\bar{R}^\square(\bar{\rho}, \tau)$  as the quotient of  $\mathbb{F}\llbracket A, B, C, R, S, T, U \rrbracket$  by an ideal which we call  $I(\tau)$ . Recall the presentations

$$\begin{aligned} I(\tau_{\zeta,s}) &= (A, B, C), \\ I(\tau_{\zeta,ns}) &= (A^2 + BC, 4RS + U^2, 2CR + AU, 2BS + AU, 2AR - BU, 2AS - CU), \\ I(\tau_{\zeta_1, \zeta_2}) &= (A^2 + BC, BS - CR, 2AR - BU, 2AS - CU) \end{aligned}$$

(using that  $A + D = 0$  in  $\bar{R}^\square(\bar{\rho}, \tau)$  for each  $\tau$ , we have eliminated  $D$  and written  $F = A - D = 2A$ ). We have already shown that  $I(\tau_{\zeta,s})$  and  $I(\tau_{\zeta,ns})$  are prime — they are the ideals denoted  $\mathfrak{a}_{nr}$  and  $\mathfrak{a}_N$  in the statement of the theorem. It is clear that

$$Z^4(\bar{R}^\square(\bar{\rho}, \tau_{\zeta,s})) = [\mathfrak{a}_{nr}], \quad Z^4(\bar{R}^\square(\bar{\rho}, \tau_{\zeta,ns})) = [\mathfrak{a}_N].$$

Suppose that  $\mathfrak{p}$  is a prime ideal of  $\mathbb{F}\llbracket A, B, C, R, S, T, U \rrbracket$  containing  $I(\tau_{\zeta_1, \zeta_2})$ . We show that  $\mathfrak{p}$  contains  $\mathfrak{a}_{nr}$  or  $\mathfrak{a}_N$ . If  $B, C \in \mathfrak{p}$  then  $A \in \mathfrak{p}$  as  $A^2 + BC \in I(\tau_{\zeta_1, \zeta_2})$  and we have  $\mathfrak{a}_{nr} \subset \mathfrak{p}$ . Otherwise, suppose that  $B \notin \mathfrak{p}$ . As  $A^2 + BC \in \mathfrak{p}$ , either both  $A$  and  $C$  are in  $\mathfrak{p}$  or neither is. If  $A, C \in \mathfrak{p}$  then from  $2AR - BU \in \mathfrak{p}$  we deduce that  $U \in \mathfrak{p}$ , while from  $BS - CR \in \mathfrak{p}$  we deduce that  $S \in \mathfrak{p}$ . It is then easy to see that  $\mathfrak{a}_N \subset \mathfrak{p}$ . If  $A, B, C \notin \mathfrak{p}$  then because  $B(2CR + AU)$  and  $C(2BS + AU)$  are in  $I(\tau_{\zeta_1, \zeta_2})$  we see that  $2CR + AU, 2BS + AU \in \mathfrak{p}$ . This implies that  $A(4RS + U^2) \in \mathfrak{p}$ , and so  $4RS + U^2 \in \mathfrak{p}$  and hence  $\mathfrak{a}_N \subset \mathfrak{p}$  as required.

To finish, it is easy to check that

$$e(\bar{R}^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2}), \mathfrak{a}_{nr}) = 2 \quad \text{and} \quad e(\bar{R}^\square(\bar{\rho}, \tau_{\zeta_1, \zeta_2}), \mathfrak{a}_N) = 1,$$

and so we get [Equation \(16\)](#). □

**5E. Cohen–Macaulayness.** If  $\tau_0$  is a semisimple representation of  $I_F$  over  $E$ , let  $R(\bar{\rho}, \tau_0)'$  be the maximal reduced and  $l$ -torsion-free quotient of  $R(\bar{\rho})$  all of whose  $\bar{E}$ -points give rise to representations  $\rho$  of  $G_F$  with  $\rho|_{I_F}^{ss} \cong \tau_0$ . Then I claim that  $R(\bar{\rho}, \tau_0)'$  is always Cohen–Macaulay. Indeed, if  $\tau_0$  is nonscalar then this is proved above. If  $\tau_0$  is scalar, then we may twist and assume that it is trivial. If  $q \not\equiv \pm 1 \pmod{l}$ , this follows from [Proposition 5.5](#). If  $q \equiv 1 \pmod{l}$  then we can deduce the claim from [Proposition 5.8](#) together with [[Eisenbud 1995](#), Exercise 18.13], which says that if  $R/I$  and  $R/J$  are  $d$ -dimensional Cohen–Macaulay quotients of a noetherian local ring  $R$ , and  $\dim R/(I + J) = d - 1$ , then  $R/(I \cap J)$  is Cohen–Macaulay if and only if  $R/(I + J)$  is. We take  $R = R^\square(\bar{\rho})$ , and  $I$  and  $J$  to be the ideals cutting out  $R^\square(\bar{\rho}, \tau_s)$  and  $R^\square(\bar{\rho}, \tau_{ns})$  respectively. Then  $R/I$  and  $R/J$  are Cohen–Macaulay, and  $R/(I + J)$  is a quotient of the formally smooth ring  $R/I$  by the single equation  $q \operatorname{tr}(\rho^\square(\phi))^2 = (q + 1)^2 \det(\rho^\square(\phi))$ , and so is Cohen–Macaulay.

Therefore  $R/(I \cap J)$  is Cohen–Macaulay as required. When  $q \equiv -1 \pmod{l}$  the claim follows from [Proposition 5.6](#) unless  $\bar{\rho}$  is the direct sum of the trivial and cyclotomic characters, in which case we use [Remark 5.7](#).

For  $n$ -dimensional representations the unrestricted framed deformation ring  $R^\square(\bar{\rho})$  is always Cohen–Macaulay (in fact, a complete intersection; this is due to David Helm, building on work of Choi [\[2009\]](#)). It is natural to wonder whether the rings obtained by fixing the semisimplified restriction to inertia are always Cohen–Macaulay. Note that they are not always Gorenstein.

For a discussion of how the Cohen–Macaulay property of local deformation rings can be used to show that certain global Galois deformation rings are flat over  $\mathcal{O}$ , see [\[Snowden 2011, Section 5\]](#).

## 6. Reduction of types — proofs.

The aim of this section is to analyse the reduction modulo  $l$  of the  $K$ -types  $\sigma(\tau)$  defined in [Section 3](#), and in particular to prove [Lemma 3.9](#).

**6A. The essentially tame case.** Suppose that  $\tau = (r_\tau, N_\tau)$  where  $r_\tau$  is a tamely ramified, semisimple representation of  $I_F$ . Then  $\sigma(\tau)$  is inflated from a representation of  $\mathrm{GL}_2(k_F)$ . We will always use the same notation for a representation of  $\mathrm{GL}_2(k_F)$  and its inflation to  $\mathrm{GL}_2(\mathcal{O}_F)$ . For this subsection let  $G = \mathrm{GL}_2(k_F)$ , let  $B$  be the subgroup of upper-triangular matrices, let  $U$  be the subgroup of unipotent elements of  $B$ , let  $Z$  be the centre of  $G$ , and fix an embedding  $\alpha : k_L^\times \hookrightarrow G$ . Fix a nontrivial additive character  $\psi$  of  $U$ . Then we have (see, e.g., [\[Bushnell and Henniart 2006, Chapter 6\]](#)):

- If  $r_\tau = (\mathrm{rec}(\tilde{\chi}) \oplus \mathrm{rec}(\tilde{\chi}))|_{I_F}$  and  $N_\tau \neq 0$ , where  $\tilde{\chi}|_{\mathcal{O}_F^\times}$  is inflated from a character  $\chi$  of  $k_F^\times$ , then  $\sigma(\tau) = (\chi \circ \det) \otimes \mathrm{St}$ , where  $\mathrm{St}$  is the Steinberg representation of  $G$ .
- If  $r_\tau = (\mathrm{rec}(\tilde{\chi}) \oplus \mathrm{rec}(\tilde{\chi}))|_{I_F}$  and  $N_\tau = 0$ , where  $\tilde{\chi}|_{\mathcal{O}_F^\times}$  is inflated from a character  $\chi$  of  $k_F^\times$ , then  $\sigma(\tau) = \chi \circ \det$ .
- If  $r_\tau = (\mathrm{rec}(\tilde{\chi}_1) \oplus \mathrm{rec}(\tilde{\chi}_2))|_{I_F}$ , where  $\tilde{\chi}_1|_{\mathcal{O}_F^\times}$  and  $\tilde{\chi}_2|_{\mathcal{O}_F^\times}$  are inflated from *distinct* characters  $\chi_1$  and  $\chi_2$  of  $k_F^\times$ , then

$$\sigma(\tau) = \mu(\chi_1, \chi_2) := \mathrm{Ind}_B^G(\chi_1 \otimes \chi_2).$$

- If  $r_\tau = (\mathrm{Ind}_{G_L}^{G_F} \mathrm{rec}(\tilde{\theta}))|_{I_F}$  where  $\tilde{\theta}|_{\mathcal{O}_L^\times}$  is inflated from a character  $\theta$  of  $k_L^\times$  which is not equal to its  $\mathrm{Gal}(k_L/k_F)$  conjugate  $\theta^c$ , then

$$\sigma(\tau) = \pi_\theta := \mathrm{Ind}_{ZU}^G(\theta|_Z \psi) - \mathrm{Ind}_{\alpha(k_L^\times)}^G \theta$$

(this virtual representation is a genuine irreducible representation that is independent of the choice of  $\psi$ ).

The only isomorphisms between these representations are of the form  $\mu(\chi_1, \chi_2) \cong \mu(\chi_2, \chi_1)$  and  $\pi_\theta \cong \pi_{\theta^c}$ .

We want to understand the reductions of these representations modulo  $l$ , for this see [Helm 2010]. We will use analogous notation for representations of  $G$  in characteristic zero and in characteristic  $l$ ; hopefully this will not cause confusion.

If  $q \not\equiv \pm 1 \pmod{l}$ , then reduction modulo  $l$  is a bijection between irreducible  $\overline{\mathbb{F}}_l$ -representations of  $G$  and irreducible  $\overline{E}$ -representations of  $G$ , as  $G$  has order  $q(q+1)(q-1)^2$  which is coprime to  $l$ .

If  $q \equiv 1 \pmod{l}$ , then the distinct irreducible representations of  $\mathrm{GL}_2(k_F)$  over  $\overline{\mathbb{F}}$  are  $\chi \circ \det$  and  $\mathrm{St} \otimes (\chi \circ \det)$  for  $\chi : k_F^\times \rightarrow \overline{\mathbb{F}}^\times$ ,  $\mu(\chi_1, \chi_2)$  for  $\chi_1, \chi_2 : k_F^\times \rightarrow \overline{\mathbb{F}}^\times$  a pair of distinct characters, and  $\pi_\theta$  for  $\theta : k_L^\times \rightarrow \overline{\mathbb{F}}^\times$  a character which is not isomorphic to its conjugate. The notation is all entirely analogous to the characteristic zero case. Once again, the only isomorphisms are  $\mu(\chi_1, \chi_2) \cong \mu(\chi_2, \chi_1)$  and  $\pi_\theta \cong \pi_{\theta^c}$ . The reductions of the characteristic zero representations are:

- $\overline{\chi \circ \det} = \overline{\chi} \circ \det$ .
- $\overline{\mathrm{St} \otimes \chi \circ \det} = \mathrm{St} \otimes (\overline{\chi} \circ \det)$ .
- $\overline{\mu(\chi_1, \chi_2)} = \mu(\overline{\chi}_1, \overline{\chi}_2)$  if  $\overline{\chi}_1 \neq \overline{\chi}_2$ .
- $\overline{\mu(\chi_1, \chi_2)} = (\overline{\chi} \circ \det) \oplus \mathrm{St} \otimes (\overline{\chi} \circ \det)$  if  $\overline{\chi}_1 = \overline{\chi}_2 = \overline{\chi}$ .
- $\overline{\pi_\theta} = \pi_{\overline{\theta}}$ .

For the last of these, we must observe that  $\theta/\theta^c$  is a character of  $k_L^\times/k_F^\times$ , a group which has order  $q+1$  and so coprime to  $l$  (as  $l > 2$ ). Therefore if  $\theta \neq \theta^c$  then  $\overline{\theta} \neq \overline{\theta^c}$ .

If  $q \equiv -1 \pmod{l}$ , then the distinct irreducible representations are  $\chi \circ \det$  for  $\chi : k_F^\times \rightarrow \overline{\mathbb{F}}^\times$ ,  $\mu(\chi_1, \chi_2)$  for  $\chi_1, \chi_2 : k_F^\times \rightarrow \overline{\mathbb{F}}^\times$  unordered pair of distinct characters,  $\pi_\theta$  for  $\theta : k_L^\times \rightarrow \overline{\mathbb{F}}^\times$  a character which is not isomorphic to its conjugate, and  $(\chi \circ \det) \otimes \pi_1$  for  $\chi : k_F^\times \rightarrow \overline{\mathbb{F}}^\times$  a character. This last needs some explanation,  $\pi_1$  is the reduction modulo  $l$  of  $\pi_\theta$  for any character  $\theta : k_L^\times/k_F^\times \rightarrow \overline{E}^\times$  which is not equal to  $\theta^c$  but whose reduction modulo  $l$  is trivial. Once again, the only isomorphisms are  $\mu(\chi_1, \chi_2) \cong \mu(\chi_2, \chi_1)$  and  $\pi_\theta \cong \pi_{\theta^c}$ . The reductions of the characteristic 0 representations are:

- $\overline{\chi \circ \det} = \overline{\chi} \circ \det$ .
- $\overline{\mu(\chi_1, \chi_2)} = \mu(\overline{\chi}_1, \overline{\chi}_2)$ .
- $\overline{\pi_\theta} = \pi_{\overline{\theta}}$  if  $\overline{\theta} \neq \overline{\theta^c}$ .
- $\overline{\pi_\theta} = \pi_1 \otimes (\overline{\theta}|_{k_F^\times} \circ \det)$  if  $\overline{\theta} = \overline{\theta^c}$ .
- $\overline{\mathrm{St} \otimes \chi \circ \det}$  has  $\pi_1 \otimes (\overline{\chi} \circ \det)$  as a submodule with quotient  $\overline{\chi} \circ \det$ .

In particular, comparing this analysis with Lemma 3.8 shows:

**Lemma 6.1.** *If  $\tau = (r, 0)$  and  $\tau' = (r', 0)$  are scalar on  $P_F$  but not on  $\bar{P}_F$ , then  $\bar{\sigma}(\tau)$  and  $\bar{\sigma}(\tau')$  are irreducible and are isomorphic if and only if  $r \equiv r' \pmod{l}$ .*

**6B. The wild case.** If  $\tau = (r, 0)$  and all twists of  $r$  are wildly ramified (we say that  $\tau$  is “essentially wildly ramified”), then the following lemma will allow us to show that  $\bar{\sigma}(\tau)$  is irreducible. If  $\rho$  is a  $\bar{\mathbb{Z}}_l$ -representation of a group  $H$ , we write  $\bar{\rho}$  for  $\rho \otimes \bar{\mathbb{F}}_l$ .

**Lemma 6.2.** *Suppose that  $H \triangleleft J \subset K$  are profinite groups such that  $H$  is open in  $K$ ,  $H$  has pro-order coprime to  $l$ , and  $J/H$  is an abelian  $l$ -group. Suppose that  $\lambda$  is a  $\bar{\mathbb{Z}}_l$ -representation of  $J$ , and write  $\eta$  for the restriction of  $\lambda$  to  $H$ . Suppose that  $\eta$  (and hence  $\lambda$ ) is irreducible. Suppose that if  $g \in K$  intertwines  $\eta$ , then  $g \in J$ . Then*

- (1) *The representations of  $J$  extending  $\eta$  are precisely  $\lambda_i = \lambda \otimes v_i$  as  $v_i$  run through the characters of  $J/H$ . There is an isomorphism  $\text{Ind}_H^J \eta \otimes \bar{E} \cong \bigoplus_i \lambda_i$ . The unique  $\bar{\mathbb{F}}_l$ -representation extending  $\bar{\eta}$  is  $\bar{\lambda}$ , and all of the Jordan–Hölder factors of  $\text{Ind}_H^J \bar{\eta}$  are isomorphic to  $\bar{\lambda}$ .*
- (2) *A  $\bar{\mathbb{F}}_l$ -representation  $\rho$  of  $J$  contains  $\bar{\lambda}$  as a subrepresentation if and only if it contains  $\bar{\lambda}$  as a quotient.*
- (3) *The representations  $\text{Ind}_J^K \lambda_i$  and  $\text{Ind}_J^K \bar{\lambda}$  are irreducible.*

*Proof.*

- (1) In characteristic 0 we argue as follows. First note that the representations  $\lambda_i$  are distinct, otherwise  $\lambda|_H$  would have a nonscalar endomorphism, contradicting Schur’s lemma. By Frobenius reciprocity, the  $\lambda_i$  are distinct irreducible constituents of  $\text{Ind}_H^J \eta$ . Since the sum of their dimensions is  $\dim \text{Ind}_H^J \eta$ , they are the only irreducible constituents. By Frobenius reciprocity, any representation extending  $\eta$  must occur in  $\text{Ind}_H^J \eta$  and so must be one of the  $\lambda_i$ , as required. In characteristic  $l$ , first note that  $\bar{\lambda}$  is irreducible since the pro-order of  $H$  is coprime to  $l$ . It follows from this and the fact that  $\bar{v}_i$  is trivial for all  $i$  that the Jordan–Hölder factors of  $\text{Ind}_H^J \bar{\eta}$  are isomorphic to  $\bar{\lambda}$ . Frobenius reciprocity then implies that  $\bar{\lambda}$  is the unique irreducible representation of  $J$  extending  $H$ .
- (2) It follows from part 1 that  $\text{Hom}_J(\bar{\lambda}, \rho) \neq 0$  if and only if  $\text{Hom}_J(\text{Ind}_H^J \bar{\eta}, \rho) \neq 0$ . By Frobenius reciprocity, this is equivalent to  $\text{Hom}_H(\bar{\eta}, \rho) \neq 0$ . But by the assumption on the pro-order of  $H$ ,  $\bar{\mathbb{F}}_l$ -representations of  $H$  are semisimple, and so this is equivalent to  $\text{Hom}_H(\rho, \bar{\eta}) \neq 0$ , which by the same argument is equivalent to  $\text{Hom}_J(\rho, \text{Ind}_H^J \bar{\eta}) \neq 0$ .
- (3) First, note that  $\dim \text{Hom}_K(\text{Ind}_J^K \bar{\lambda}, \text{Ind}_J^K \bar{\lambda}) = 1$ , by Mackey’s decomposition formula and the assumption that elements of  $K \setminus J$  do not intertwine  $\eta$ . Now suppose that  $\rho$  is an irreducible subrepresentation of  $\text{Ind}_J^K \bar{\lambda}$ . By Frobenius reciprocity and part 2 we may deduce that  $\rho$  is also an irreducible quotient of  $\text{Ind}_J^K \bar{\lambda}$ . The composition  $\text{Ind}_J^K \bar{\lambda} \twoheadrightarrow \rho \hookrightarrow \text{Ind}_J^K \bar{\lambda}$  is then a nonzero element

of  $\text{Hom}_K(\text{Ind}_J^K \bar{\lambda}, \text{Ind}_J^K \bar{\lambda})$ , and is therefore scalar. But this is only possible if  $\rho = \text{Ind}_J^K \bar{\lambda}$ , as required. The statement about  $\text{Ind}_J^K \lambda_i$  follows.  $\square$

**Proposition 6.3.** *Let  $\tau = (r, 0)$  be an essentially wildly ramified inertial type. Then there exists a subgroup  $J \subset K$ , an irreducible representation  $\lambda$  of  $J$ , and a subgroup  $\tilde{J} \triangleleft J$ , such that  $(\tilde{J}, J, K, \lambda)$  satisfy the hypotheses on  $(H, J, K, \lambda)$  in Lemma 6.2 and such that  $\sigma(\tau) = \text{Ind}_{\tilde{J}}^K \lambda$ .*

*In particular,  $\overline{\sigma(\tau)}$  is irreducible.*

*Proof.* Suppose first that  $r$  is the restriction to  $I_F$  of a reducible representation of  $G_F$ . Then  $\sigma(\tau) = \text{Ind}_{K_0(N)}^K \epsilon \otimes (\chi \circ \det)$  for a character  $\epsilon$  of  $\mathcal{O}_F^\times$  of exponent  $N \geq 2$  and a character  $\chi$  of  $\mathcal{O}_F^\times$ . Let  $J = K_0(N)$ , and let

$$\tilde{J} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in J : a \text{ has order coprime to } l \text{ modulo } \mathfrak{p}_F \right\}.$$

Then  $\tilde{J}$ ,  $J$ , and  $\epsilon$  satisfy all the required hypotheses — the only one to check is that  $\epsilon|_{\tilde{J}}$  is not intertwined by any element of  $K \setminus J$ . We deduce this (in somewhat circular fashion) from the irreducibility of  $\text{Ind}_J^K(\epsilon)$ , since this is shorter than a direct proof. If  $g \in K$  intertwines  $\epsilon|_{\tilde{J}}$ , then  $\text{Hom}_{\tilde{J} \cap g \tilde{J} g^{-1}}(\epsilon, \epsilon^g) \neq 0$ . By Mackey’s formula

$$\dim(\text{Hom}_{\tilde{J}}(\epsilon, \text{Ind}_J^K \epsilon)) = \sum_{g \in \tilde{J} \backslash K / \tilde{J}} \dim \text{Hom}_{\tilde{J} \cap g \tilde{J} g^{-1}}(\epsilon, \epsilon^g).$$

The left hand side is in turn equal to  $\dim \text{Hom}_K(\text{Ind}_J^K \epsilon, \text{Ind}_J^K \epsilon)$ . But  $\text{Ind}_J^K \epsilon = \bigoplus_i \text{Ind}_J^K \epsilon_i$  where  $\epsilon_i$  are the characters of  $J$  extending  $\epsilon|_{\tilde{J}}$ , and by the appendix to [Breuil and Mézard 2002], these  $\text{Ind}_J^K \epsilon_i$  are irreducible and distinct. Therefore the left hand side is equal to  $(J : \tilde{J})$ . The right hand side has a contribution of 1 from each  $g \in J / \tilde{J}$ , and therefore from no other  $g$ , as required.

Now suppose that  $r$  is the restriction to  $I_F$  of an irreducible representation of  $G_F$ . Then  $\sigma(\tau) = \text{Ind}_J^K \lambda$  for an irreducible representation  $\lambda$  of  $J$  extending an irreducible representation  $\eta$  of a pro- $p$  normal subgroup  $J^1$  of  $J$  (see [Bushnell and Henniart 2006, Sections 15.5, 15.6 and 15.7] — note that our  $J$  is the maximal compact subgroup of their  $J_\alpha$ , but our  $J^1$  agrees with their  $J_\alpha^1$ ). We have  $J/J^1 = k^\times$ , where  $k$  is the residue field of a quadratic extension of  $F$ , and so  $J$  has a normal subgroup  $\tilde{J}$  of pro-order coprime to  $l$  such that  $J/\tilde{J}$  is an  $l$ -group. Then  $(\tilde{J}, J, K, \lambda)$  satisfy all the required hypotheses — the intertwining statement follows from [Bushnell and Henniart 2006, 15.6 Proposition 2].  $\square$

**Proposition 6.4.** *Let  $\tau = (r, 0)$  and  $\tau' = (r', 0)$  be inertial types that are not scalar on  $\tilde{P}_F$ . If  $r \equiv r' \pmod{l}$ , then  $\overline{\sigma(\tau)}$  and  $\overline{\sigma(\tau')}$  are isomorphic.*

*Proof.* If either of  $r$  and  $r'$  is (after a twist) tamely ramified, then so is the other and this is contained in Lemma 6.1. Otherwise, by Lemma 3.8, we are in one of the following cases:



- (1)  $r = (\chi_1 \oplus \chi_2)|_{I_F}$  for characters  $\chi_1$  and  $\chi_2$  of  $G_F$  that are distinct on  $P_F$ , and  $r' = (\chi'_1 \oplus \chi'_2)|_{I_F}$  for characters  $\chi'_1$  and  $\chi'_2$  of  $G_F$  with  $\chi_i \equiv \chi'_i$  for  $i = 1, 2$ .
- (2)  $r = (\text{Ind}_{G_L}^{G_F} \xi)_{I_F}$  and  $r' = (\text{Ind}_{G_L}^{G_F} \xi')_{I_F}$  for wildly ramified characters  $\xi$  and  $\xi'$  of  $G_L$  such that  $\xi \equiv \xi'$ , and such that  $\xi|_{\tilde{P}_F}$  does not extend to  $G_F$ .
- (3)  $r|_{\tilde{P}_F}$  is irreducible and  $r' = r \otimes \chi$  for a character  $\chi$  of  $I_F$  that extends to  $G_F$  and such that  $\chi \equiv 1 \pmod{l}$ .

In the first case, we may write  $\chi_i = \text{rec}(\epsilon_i)$  and  $\chi'_i = \text{rec}(\epsilon'_i)$  with  $\epsilon_i$  and  $\epsilon'_i$  characters of  $F^\times$  such that  $\epsilon_i \equiv \epsilon'_i \pmod{l}$  and such that  $\epsilon = \epsilon_1/\epsilon_2$  has exponent  $N \geq 1$ . Since  $\epsilon' = \epsilon'_1/\epsilon'_2$  also has exponent  $N$ , we have

$$\begin{aligned} \sigma(\tau) &= \epsilon_2 \otimes \text{Ind}_{K_0(N)}^K \epsilon \\ &\equiv \epsilon'_2 \otimes \text{Ind}_{K_0(N)}^K \epsilon' \pmod{l} \\ &= \sigma(\tau'). \end{aligned}$$

In the second case, by twisting we may reduce to the case where  $(L/F, \text{rec}^{-1}(\xi))$  is an unramified minimal admissible pair [Bushnell and Henniart 2006, § 19.6]. Then, following through the explicit construction of [Bushnell and Henniart 2006, SS 19.3 and 19.4] we see that there are

- (1) a simple stratum  $(\mathfrak{A}, n, \alpha)$  with associated compact open subgroups  $J_1 \subset J \subset K$ , with  $J_1$  pro- $p$  and  $J/J_1 \cong k_L^\times$ ,
- (2) a representation  $\eta$  of  $J^1$  and extensions  $\lambda$  and  $\lambda'$  of  $\eta$  to  $J$  such that  $\text{Ind}_J^K(\lambda) = \sigma(\tau)$  and  $\text{Ind}_J^K(\lambda') = \sigma(\tau')$ .

Indeed, up to conjugacy,  $(\mathfrak{A}, n, \alpha)$ ,  $J_1$ , and  $\eta$  are determined by  $\text{rec}^{-1}(\xi)|_{U_L^1} = \text{rec}^{-1}(\xi')|_{U_L^1}$ . The representations  $\lambda$  and  $\lambda'$  are defined in terms of  $\text{rec}^{-1}(\xi)$  and  $\text{rec}^{-1}(\xi')$  by the formulae of [Bushnell and Henniart 2006, 19.3.1 and Corollary 19.4] (together with the correction factor of paragraph 34.4, an unramified twist  $\Delta_\xi$ , that makes no difference to the argument). It is clear from these that if  $\xi \equiv \xi'$  then  $\lambda \equiv \lambda'$  as required.

In the final case,  $r' = r \otimes \chi$  for a character  $\chi$  of  $I_F$  that extends to  $G_F$ . By compatibility of  $\tau \mapsto \sigma(\tau)$  with twisting,

$$\begin{aligned} \sigma(\tau') &= \sigma(\tau) \otimes \text{rec}^{-1}(\chi) \circ \det \\ &\equiv \sigma(\tau) \pmod{l} \end{aligned}$$

as required. □

**Proposition 6.5.** *Let  $\tau = (r, 0)$  and  $\tau' = (r', 0)$  be inertial types that are not scalar on  $\tilde{P}_F$ . If  $\overline{\sigma(\tau)}$  and  $\overline{\sigma(\tau')}$  are isomorphic, then  $r \equiv r' \pmod{l}$ .*

*Proof.* If one of  $r$  and  $r'$  has a twist which is trivial on  $P_F$ , then so does the other and in this case the proposition follows from Lemma 6.1.

Otherwise we may, by twisting, assume that  $\sigma(\tau)$  and  $\sigma(\tau')$  satisfy  $l(\sigma) \leq l(\sigma \otimes \chi)$  for all characters  $\chi$  of  $\mathcal{O}_F^\times$  (the definition of  $l(\sigma)$  is as in [Bushnell and Henniart 2006, § 12.6]). In this case  $\sigma(\tau)$  and  $\sigma(\tau')$  contain the same, nonempty, sets of fundamental strata (because this only depends on the restriction to pro- $p$  subgroups).

If one of  $\sigma(\tau)$  and  $\sigma(\tau')$  contains a split fundamental stratum [Bushnell and Henniart 2006, 13.2] then so does the other. In this case, [Bushnell and Henniart 2006, Corollary 13.3] implies that they cannot be cuspidal types and so we have

$$\sigma(\tau) = \text{Ind}_{K_0(N)}^K(\epsilon) \quad \text{and} \quad \sigma(\tau') = \text{Ind}_{K_0(N')}^K(\epsilon')$$

for some  $\epsilon$  and  $\epsilon'$  of exponents  $N$  and  $N'$ . It is easy to see that in fact we must have  $N = N'$ . From Lemma 6.2 we deduce that  $\epsilon \equiv \epsilon' \pmod{l}$ , and so  $\tau \equiv \tau' \pmod{l}$  as required.

Otherwise,  $\sigma(\tau) = \text{Ind}_J^K \lambda$  and  $\sigma(\tau') = \text{Ind}_J^K \lambda'$  for a simple stratum  $(\mathfrak{A}, n, \alpha)$  with associated groups  $J^1 \subset J$  and representations  $\lambda$  and  $\lambda'$  extending the representation  $\eta$  of  $J$ . From Lemma 6.2 we deduce that  $\lambda' = \lambda \otimes \eta$  for a character  $\eta$  of  $J/J^1$  with  $\eta \equiv 1 \pmod{l}$ .

If  $\mathfrak{A}$  is unramified, then by the reverse of the argument in the second case of the previous proposition we see that  $\tau = (\text{Ind}_{G_L}^{G_F} \xi)|_{I_F}$  and  $\tau' = (\text{Ind}_{G_L}^{G_F} \xi')|_{I_F}$  for  $\xi$  and  $\xi'$  characters of  $G_L$  with  $\xi|_{I_L} \equiv \xi'|_{I_L}$ , whence the result.

If  $\mathfrak{A}$  is ramified, then  $\eta$  can be regarded as a character of  $J/J^1 \cong k_M^\times = k_F^\times$  with  $\eta \equiv 1 \pmod{l}$  for some ramified quadratic extension  $M/F$ . I claim that there is a character  $\chi$  of  $\mathcal{O}_F^\times$  with  $\eta = \chi \circ \det$  and  $\chi \equiv 1 \pmod{l}$ . Indeed, as  $l > 2$  we can take the inflation to  $\mathcal{O}_F^\times$  of the character  $\chi$  of  $k_F^\times$  satisfying  $\chi \equiv 1 \pmod{l}$  and  $\chi^2 = \eta$ . Then  $\sigma(\tau) = \sigma(\tau') \otimes (\chi \circ \det)$  and so

$$\begin{aligned} \tau &= \tau' \otimes \text{rec}(\chi) \\ &\equiv \tau' \pmod{l}, \end{aligned}$$

as required. □

### Acknowledgements

This work forms part of my Imperial College, London PhD thesis, and I am grateful to my supervisor Toby Gee for suggesting this topic, for comments on drafts of this paper, and for answering many questions. I also thank Kevin Buzzard, David Helm, Yongquan Hu, Tristan Kalloniatis, Vytautas Paškūnas for helpful comments and corrections, and Jack Thorne for encouraging me to investigate the Cohen–Macaulay property of these deformation rings.

This research was supported by the Engineering and Physical Sciences Research Council and the Philip Leverhulme Trust, and part of this work was done during a visit to the University of Chicago supported by the London Mathematical Society and the Cecil King Foundation.

## References

- [Barnet-Lamb et al. 2014] T. Barnet-Lamb, T. Gee, D. Geraghty, and R. Taylor, “Potential automorphy and change of weight”, *Ann. of Math. (2)* **179**:2 (2014), 501–609. MR 3152941
- [Breuil and Mézard 2002] C. Breuil and A. Mézard, “Multiplicités modulaires et représentations de  $GL_2(\mathbb{Z}_p)$  et de  $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$  en  $l = p$ ”, *Duke Math. J.* **115**:2 (2002), 205–310. MR 1944572
- [Bushnell and Henniart 2006] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for  $GL_2$* , Grundlehren der Mathematischen Wissenschaften **335**, Springer, Berlin, 2006. MR 2234120
- [Choi 2009] S. H. Choi, *Local deformation lifting spaces of mod  $l$  Galois representations*, Ph.D. thesis, Harvard University, 2009.
- [Clozel et al. 2008] L. Clozel, M. Harris, and R. Taylor, “Automorphy for some  $l$ -adic lifts of automorphic mod  $l$  Galois representations”, *Publ. Math. Inst. Hautes Études Sci.* 108 (2008), 1–181. MR 2470687
- [Eisenbud 1995] D. Eisenbud, *Commutative algebra*, Graduate Texts in Mathematics **150**, Springer, New York, 1995. MR 1322960
- [Emerton and Gee 2014] M. Emerton and T. Gee, “A geometric perspective on the Breuil–Mézar­d conjecture”, *J. Inst. Math. Jussieu* **13**:1 (2014), 183–223. MR 3134019
- [Gee and Kisin 2014] T. Gee and M. Kisin, “The Breuil–Mézar­d conjecture for potentially Barsotti–Tate representations”, *Forum Math. Pi* **2** (2014), e1, 56. MR 3292675
- [Helm 2010] D. Helm, “On  $l$ -adic families of cuspidal representations of  $GL_2(\mathbb{Q}_p)$ ”, *Math. Res. Lett.* **17**:5 (2010), 805–822. MR 2727610
- [Henniart 2002] G. Henniart, appendix to [Breuil and Mézard 2002], 2002.
- [Hu and Tan 2013] Y. Hu and F. Tan, “The Breuil–Mézar­d conjecture for non-scalar split residual representations”, preprint, 2013. arXiv 1309.1658
- [Kisin 2009a] M. Kisin, “The Fontaine–Mazur conjecture for  $GL_2$ ”, *J. Amer. Math. Soc.* **22**:3 (2009), 641–690. MR 2505297
- [Kisin 2009b] M. Kisin, “Moduli of finite flat group schemes, and modularity”, *Ann. of Math. (2)* **170**:3 (2009), 1085–1180. MR 2600871
- [Paškūnas 2015] V. Paškūnas, “On the Breuil–Mézar­d conjecture”, *Duke Math. J.* **164**:2 (2015), 297–359. MR 3306557
- [Pilloni 2008] V. Pilloni, “The study of 2-dimensional  $p$ -adic Galois deformations in the  $l$  not  $p$  case”, 2008, available at <http://perso.ens-lyon.fr/vincent.pilloni/Defo.pdf>.
- [Reduzzi 2013] D. Reduzzi, “On the number of irreducible components of local deformation rings in the unequal characteristic case”, 2013, available at [http://msp.org/msp\\_org/media/files/D.A.Reduzzi-Irreducible\\_Components\\_Local\\_Deformations.pdf](http://msp.org/msp_org/media/files/D.A.Reduzzi-Irreducible_Components_Local_Deformations.pdf).
- [Snowden 2011] A. Snowden, “Singularities of ordinary deformation rings”, 2011. arXiv 1111.3654
- [Taylor 2008] R. Taylor, “Automorphy for some  $l$ -adic lifts of automorphic mod  $l$  Galois representations II”, *Publ. Math. Inst. Hautes Études Sci.* 108 (2008), 183–239. MR 2470688
- [Taylor 2009] R. Taylor, “Modularity lifting theorems”, Lecture notes, Harvard University, 2009.

Communicated by Brian Conrad

Received 2015-04-28

Revised 2016-04-14

Accepted 2016-07-18

[jack.shotton@gmail.com](mailto:jack.shotton@gmail.com)

University of Chicago, 5442 S Ellis Ave, Chicago, IL 60615,  
United States

# Algebra & Number Theory

[msp.org/ant](http://msp.org/ant)

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Susan Montgomery	University of Southern California, USA
Richard E. Borcherds	University of California, Berkeley, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
John H. Coates	University of Cambridge, UK	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Roger Heath-Brown	Oxford University, UK	Ravi Vakil	Stanford University, USA
Craig Huneke	University of Virginia, USA	Michel van den Bergh	Hasselt University, Belgium
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Yuri Manin	Northwestern University, USA	Efim Zelmanov	University of California, San Diego, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Shou-Wu Zhang	Princeton University, USA

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2016 is US \$290/year for the electronic version, and \$485/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 10    No. 7    2016

---

Arithmetic invariant theory and 2-descent for plane quartic curves	1373
JACK A. THORNE	
Furstenberg sets and Furstenberg schemes over finite fields	1415
JORDAN S. ELLENBERG and DANIEL ERMAN	
Local deformation rings for $GL_2$ and a Breuil–Mézard conjecture when $l \neq p$	1437
JACK SHOTTON	
Generalized Kuga–Satake theory and rigid local systems, II: rigid Hecke eigensheaves	1477
STEFAN PATRIKIS	
Lifting preprojective algebras to orders and categorifying partial flag varieties	1527
LAURENT DEMONET and OSAMU IYAMA	
A fibered power theorem for pairs of log general type	1581
KENNETH ASCHER and AMOS TURCHET	