

Algebra & Number Theory

Volume 10

2016

No. 7

**Generalized Kuga–Satake theory
and rigid local systems, II:
rigid Hecke eigensheaves**

Stefan Patrikis



Generalized Kuga–Satake theory and rigid local systems, II: rigid Hecke eigensheaves

Stefan Patrikis

We use rigid Hecke eigensheaves, building on Yun’s work on the construction of motives with exceptional Galois groups, to produce the first robust examples of “generalized Kuga–Satake theory” outside the Tannakian category of motives generated by abelian varieties. To strengthen our description of the “motivic” nature of Kuga–Satake lifts, we digress to establish a result that should be of independent interest: for any quasiprojective variety over a (finitely generated) characteristic-zero field, the associated graded of the weight filtration on its intersection cohomology arises from a motivated motive in the sense of André, and in particular from a classical homological motive if one assumes the standard conjectures. This extends work of de Cataldo and Migliorini.

1. Background: generalized Kuga–Satake theory

The aim of this paper is to produce nontrivial examples of the generalized Kuga–Satake theory proposed in [Patrikis 2014b]. The classical Kuga–Satake construction is a miracle of Hodge theory that associates to any complex K3 surface X a complex abelian variety $\text{KS}(X)$ and an inclusion of \mathbb{Q} -Hodge structures

$$H^2(X, \mathbb{Q}) \subset H^1(\text{KS}(X), \mathbb{Q})^{\otimes 2}.$$

This construction takes its clearest conceptual form within the motivic Galois formalism. Let $\mathcal{M}_{\mathbb{C}}^{\text{hom}}$ denote the category of pure motives over \mathbb{C} for homological equivalence. Assuming the standard conjectures, this is a neutral Tannakian category over \mathbb{Q} with fiber functor given by Betti cohomology:

$$H_B : \mathcal{M}_{\mathbb{C}}^{\text{hom}} \rightarrow \text{Vect}_{\mathbb{Q}}.$$

This work was supported by NSF grant DMS-1303928.

MSC2010: primary 14C15; secondary 11F80, 14D24.

Keywords: Galois representations, rigid local systems, Kuga–Satake construction, geometric Langlands.

Let $\mathcal{G}_{\mathbb{C}}^{\text{hom}} = \text{Aut}^{\otimes}(H_B)$ denote the corresponding Tannakian group. Then we can phrase the Kuga–Satake construction as follows: the motive $H^2(X)$ admits a (symmetric) polarization, hence (normalizing by a Tate-twist to weight zero) corresponds to a motivic Galois representation $\rho : \mathcal{G}_{\mathbb{C}}^{\text{hom}} \rightarrow \text{SO}(H_B^2(X)(1))$.¹ The motive $H^1(\text{KS}(X))$ then *is* the motivic Galois representation corresponding to the composite $r \circ \tilde{\rho}$ in the diagram

$$\begin{array}{ccc}
 \text{GSpin}(H_B^2(X)(1)) & \xrightarrow{r} & \text{GL}(C^+(H_B^2(X)(1))) \\
 \tilde{\rho} \nearrow \text{---} & & \downarrow \\
 \mathcal{G}_{\mathbb{C}}^{\text{hom}} & \xrightarrow{\rho} & \text{SO}(H_B^2(X)(1))
 \end{array} \tag{1}$$

in which $\tilde{\rho}$ is a suitable lift of ρ , and r is the natural representation of GSpin on the even Clifford algebra. The strongest possible version of the Kuga–Satake construction is the statement that such a lift $\tilde{\rho}$ exists; this is far from known at present, as it implicitly includes deep cases of the Lefschetz standard conjecture. A weaker, but still highly nontrivial, analogue is known when $\mathcal{G}_{\mathbb{C}}^{\text{hom}}$ is replaced by the motivic Galois group of André’s category of motives for motivated cycles; see [André 1996a].

But the formulation itself is highly suggestive, pointing towards deep and largely unexplored generalizations, some of whose essential difficulties are orthogonal to the usual impenetrable conjectures of algebraic and arithmetic geometry — Lefschetz, Hodge, Tate, etc. In what follows we will work with motives over number fields and their ℓ -adic realizations, rather than motives over \mathbb{C} and their Hodge–Betti realizations, but there are analogues of the results of this paper in the latter setting. We now state a conjecture that captures the most refined form of a “generalized Kuga–Satake theory” for motives over number fields. For two number fields F and E , we let $\mathcal{M}_{F,E}$ denote the category of motives for motivated cycles over F with coefficients in E ; it is (unconditionally) neutral Tannakian over E , and by choosing an embedding $F \hookrightarrow \mathbb{C}$, the (E -linear) Betti fiber functor gives us its motivic Galois group $\mathcal{G}_{F,E}$ (see [André 1996b] for background).

Conjecture 1.1 (see Section 4.3 of [Patrikis 2014b]). *Let $\tilde{H} \rightarrow H$ be a surjection of linear algebraic E -groups whose kernel is equal to a central torus in \tilde{H} , and let*

$$\rho : \mathcal{G}_{F,E} \rightarrow H$$

be a motivic Galois representation. Then if either F is totally imaginary, or the “Hodge numbers” of ρ satisfy the (necessary) parity condition of [Patrikis 2015,

¹That this motivic Galois representation should be special orthogonal, rather than merely orthogonal, is nontrivial; it follows from work of André [1996a].

Proposition 5.5], then there exists a finite extension E'/E and a lifting

$$\begin{array}{ccc}
 & & \tilde{H}_{E'} \\
 & \nearrow \tilde{\rho} & \downarrow \\
 \mathcal{G}_{F, E'} & \xrightarrow{\rho \otimes_E E'} & H_{E'}
 \end{array}$$

of motivic Galois representations.

For a leisurely overview of this conjecture, see the introduction to [Patrikis 2014a]; for a detailed discussion of the arithmetic evidence, see [Patrikis 2014b]. Even working with motivated rather than homological motives, this conjecture is highly refined: in the classical setting of diagram (1), the existence of such a $\tilde{\rho}$ requires not only the existence of $\text{KS}(X)$, but also the full force of the theorem of Deligne–André that Hodge cycles on abelian varieties are motivated.² At first approximation, though, we can replace Conjecture 1.1 with the following variant:

Definition 1.2. Setting $\Gamma_F = \text{Gal}(\bar{F}/F)$ for an algebraic closure \bar{F} of F , we let $\rho : \Gamma_F \rightarrow H(\bar{\mathbb{Q}}_\ell)$ be a geometric Galois representation valued in an arbitrary linear algebraic group H over $\bar{\mathbb{Q}}_\ell$.

- We say that ρ is weakly motivic if there exists a faithful representation $r : H \hookrightarrow \text{GL}(V_r)$ such that $r \circ \rho$ is isomorphic to the $(\iota : E \hookrightarrow \bar{\mathbb{Q}}_\ell)$ -realization $H_\iota(M)$ of some object M of $\mathcal{M}_{F, E}$.
- Suppose that we are given such a weakly motivic $\rho : \Gamma_F \rightarrow H(\bar{\mathbb{Q}}_\ell)$, and let $\tilde{\rho}$ be a geometric lift to \tilde{H} :

$$\begin{array}{ccc}
 & & \tilde{H}(\bar{\mathbb{Q}}_\ell) \\
 & \nearrow \tilde{\rho} & \downarrow \\
 \Gamma_F & \xrightarrow{\rho} & H(\bar{\mathbb{Q}}_\ell)
 \end{array} \tag{2}$$

(That such geometric lifts typically exist is [Patrikis 2014b, Theorem 3.2.10] and [Patrikis 2015, Proposition 5.5].)³ We say that $\tilde{\rho}$ satisfies the generalized Kuga–Satake property if it is weakly motivic as an \tilde{H} -representation.

²For $F = \mathbb{C}$, that is. When F is a number field, the existence of $\tilde{\rho}$ requires the Tate conjecture for abelian varieties, and rather delicate descent arguments — see [Patrikis 2014b, §4.2].

³This should be contrasted with the situation in which the kernel of $\tilde{H} \rightarrow H$ is finite, where geometric lifts, even after allowing a finite base change on F , need not exist: for a simple example, consider the case $\text{SL}_2 \rightarrow \text{PGL}_2$ in which ρ is the projective representation associated to the Tate module of an elliptic curve (or even more simply, consider multiplication by $N > 1$ on \mathbb{G}_m , and let ρ be the cyclotomic character). For the full story, see [Wintenberger 1995].

In sum, our aim in establishing certain cases of this “generalized Kuga–Satake property” is to verify (motivated refinements of) certain cases of the Fontaine–Mazur conjecture.

With this framework in place, we can introduce the particular setting of this paper. Our aim is to study certain families of weakly motivic $\rho : \Gamma_F \rightarrow H(\bar{\mathbb{Q}}_\ell)$ for which it is possible to find lifts $\tilde{\rho} : \Gamma_F \rightarrow \tilde{H}(\bar{\mathbb{Q}}_\ell)$ satisfying the generalized Kuga–Satake property. Outside of the context of the classical Kuga–Satake construction, where ρ is the representation on $H^2(X_{\bar{F}}, \bar{\mathbb{Q}}_\ell)$, for X/F a K3 surface — or closely related examples in which the motives in question are still generated by motives of abelian varieties⁴ — there were no nontrivial examples of such a lifting until [Patrikis 2014a]. But that paper is restricted to low-dimensional examples in which $\tilde{H} = \mathrm{GSpin}_5 \rightarrow H = \mathrm{SO}_5$, and relies heavily on low-dimensional coincidences in the Dynkin classification. Thus the primary desiderata for our examples are that:

- (D.1) the motives in question not lie in the Tannakian subcategory of \mathcal{M}_F generated by abelian varieties and Artin motives;
- (D.2) the examples exist in arbitrary rank, or at least for “interesting” groups H ;
- (D.3) the lift $\tilde{\rho}$ should not be realizable within the Tannakian category of geometric representations generated by ρ , characters, and Artin representations.

We make explicit this last desideratum just to point out that for some choices of \tilde{H} , for instance $\tilde{H} = H \times \mathbb{G}_m$, the existence of a weakly motivic lift $\tilde{\rho}$ is completely trivial. Condition (D.3) is a way to ensure the results we prove have nontrivial content.

The examples of this paper meet all three criteria of interest. For our ρ we take the remarkable weakly motivic Galois representations constructed in [Yun 2014a, Theorem 4.2, Proposition 4.6]. Let us recall a somewhat simplified version of the main result of [Yun 2014a]. Let G be a split, simple, simply connected group of type A_1 , D_n with n even, G_2 , E_7 , or E_8 , and let G^\vee denote the split \mathbb{Q} -form of its dual group. We have to say a word about the coefficients of the Galois representations and motives. For definiteness, fix an embedding $\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_\ell$, implicit whenever we take “the” ℓ -adic realization of a motive with coefficients in $\bar{\mathbb{Q}}$, and let i be a square-root of -1 in $\bar{\mathbb{Q}}$. All the local systems considered can be arranged to have coefficients in the (possibly trivial) extension $\mathbb{Q}'_\ell = \mathbb{Q}_\ell(i)$. The motives will have coefficients in the subfield $\mathbb{Q}' \subset \bar{\mathbb{Q}}$ given by

$$\mathbb{Q}' = \begin{cases} \mathbb{Q} & \text{in types } D_{4m}, G_2, E_8, \\ \mathbb{Q}(i) & \text{in types } A_1, D_{4m+2}, E_7. \end{cases} \tag{3}$$

There is a certain two-fold cover $(^2)Z_G \twoheadrightarrow Z_G$ (see Definition 4.1 and Lemma 4.2) of the center Z_G of G — regard $(^2)Z_G$ as a group scheme over \mathbb{Q} — and we call a

⁴For an “axiomatic” generalization of this context, see [André 1996a], which, for instance, further allows X to be a hyperkähler variety, or a cubic four-fold.

character

$$\chi : {}^{(2)}Z_G(\overline{\mathbb{Q}}) \rightarrow \overline{\mathbb{Q}}^\times$$

odd if it is nontrivial on the kernel of ${}^{(2)}Z_G \rightarrow Z_G$.

Theorem 1.3 [Yun 2014a]. *For any odd character $\chi : {}^{(2)}Z_G(\overline{\mathbb{Q}}) \rightarrow \overline{\mathbb{Q}}^\times$, there exists a local system*

$$\rho_\chi : \pi_1(\mathbb{P}_{\mathbb{Q}}^1 - \{0, 1, \infty\}) \rightarrow G^\vee(\mathbb{Q}'_\ell)$$

whose geometric monodromy group is G^\vee , except in type D_{2m} , in which case the geometric monodromy group is SO_{4m-1} . For all number fields F such that

$$F \supseteq \begin{cases} \mathbb{Q} & \text{if } G \text{ is of type } D_{4m}, G_2, \text{ or } E_8, \\ \mathbb{Q}(\sqrt{-1}) & \text{if } G \text{ is of type } A_1, D_{4m+2}, \text{ or } E_7, \end{cases}$$

and all specializations $t : \mathrm{Spec} F \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$, the pullback $\rho_{\chi,t} : \Gamma_F \rightarrow G^\vee(\mathbb{Q}'_\ell)$ is weakly motivic. To be precise, the composition of $\rho_{\chi,t}$ with the quasiminuscule representation of G^\vee is isomorphic to the \mathbb{Q}'_ℓ -realization of an object of $\mathcal{M}_{F,\mathbb{Q}'}$.

We can now state the first main result of this paper. There is a minor technicality in the phrasing of this theorem that results very naturally from the way the geometric Satake isomorphism descends to number fields—see Section 4B for a careful explanation. Namely, for any connected reductive group H , let ρ^\vee denote the usual half-sum of the positive coroots (for any choice of based root datum), and set $H_1 = (H \times \mathbb{G}_m) / \langle (2\rho^\vee(-1) \times -1) \rangle$. In the case $H = G^\vee$, to avoid cluttered notation we write G_1^\vee for $(G^\vee)_1$; this should not cause any confusion. Yun’s construction is most naturally viewed as the construction of a local system

$$\rho_\chi : \pi_1(\mathbb{P}_{\mathbb{Q}}^1 - \{0, 1, \infty\}) \rightarrow G_1^\vee(\mathbb{Q}'_\ell) = (G^\vee \times \mathbb{G}_m)(\mathbb{Q}'_\ell)$$

in which the G^\vee component is as in Theorem 1.3, and the \mathbb{G}_m component is the cyclotomic character; the equality here uses the fact that G is simply connected.

Theorem 1.4. *Let $\tilde{H} \twoheadrightarrow G^\vee$ be any surjection of split connected reductive groups with kernel equal to a central torus in \tilde{H} . Then:*

- (1) *There exists a local system $\tilde{\rho}_\chi : \pi_1(\mathbb{P}_{\mathbb{Q}(\sqrt{-1})}^1 - \{0, 1, \infty\}) \rightarrow \tilde{H}_1(\mathbb{Q}'_\ell)$ lifting ρ_χ , i.e., such that the diagram*

$$\begin{array}{ccc} & & \tilde{H}_1(\mathbb{Q}'_\ell) \\ & \nearrow \tilde{\rho}_\chi & \downarrow \\ \pi_1(\mathbb{P}_{\mathbb{Q}(\sqrt{-1})}^1 - \{0, 1, \infty\}) & \xrightarrow{\rho_\chi} & G_1^\vee(\mathbb{Q}'_\ell) \end{array}$$

commutes. When G is of type D_{4m} , we may replace $\mathbb{Q}(\sqrt{-1})$ by \mathbb{Q} in this assertion.

- (2) For all number field specializations $t : \text{Spec } F \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$ (assuming $F \supset \mathbb{Q}(\sqrt{-1})$ in types A_1 , D_{4m+2} , and E_7), $\tilde{\rho}_{\chi,t}$ is weakly motivic, i.e., satisfies the generalized Kuga–Satake property.

The real content of this result is for G of types D_{2m} and E_7 . When $\pi_1(G) = \{1\}$ (types G_2 , E_8), there can never be any generalized Kuga–Satake lift satisfying criterion (D.3). In type A_1 , the construction is not completely trivial, but the motives in question are generated by abelian varieties and Artin motives, so fail to satisfy our criterion (D.1).⁵ But in the essential cases of types D_{2m} and E_7 , all of our desiderata are met, the key point being that, for suitable choice of \tilde{H} , the group \tilde{H}_1 has irreducible representations restricting to each of the minuscule representations of the simply connected cover G_{sc}^\vee of G^\vee ; these are representations not possessed by the original (adjoint) group G^\vee . See Section 6 for details.

We now briefly summarize the approach to constructing the lifted local systems $\tilde{\rho}_\chi$ (see the beginning of Section 2 for more orientation). Yun’s ρ_χ is constructed as the eigen-local system associated to a Hecke eigensheaf on a certain moduli space Bun of G -bundles on \mathbb{P}^1 with level structure at the points $\{0, 1, \infty\}$. Simply put, we enlarge the center of the semisimple group G to form a reductive group \tilde{G} (whose dual group \tilde{G}^\vee plays the role of \tilde{H} above); then we study an analogous moduli space $\widetilde{\text{Bun}}$ of \tilde{G} -bundles with level structure, and show that Yun’s eigensheaves can be extended to eigensheaves on $\widetilde{\text{Bun}}$. The weakly motivic nature of the lifts $\tilde{\rho}_{\chi,t}$ is realized in the (restricting to the interesting cases in type A_1 , D_{2m} , E_7) minuscule representations of \tilde{G}^\vee (or rather, of \tilde{G}_1^\vee); as in [Yun 2014a], the motives themselves are closely related to the (intersection) cohomology of certain open subvarieties of affine Schubert varieties.

To put this approach in perspective, let us note that it is a geometric analogue of the classical automorphic construction parallel to the lifting problem (2). Namely, extending an automorphic representation of G to \tilde{G} heuristically corresponds to lifting a representation $\mathcal{L}_F \rightarrow G^\vee(\mathbb{C})$ of the “automorphic Langlands group” \mathcal{L}_F to $\tilde{G}^\vee(\mathbb{C})$. We are carrying out an analogue for certain Hecke eigensheaves, being careful to retain hold of the explicit “motivic” nature of the corresponding eigen-local systems.

In fact, we prove something considerably stronger than Theorem 1.4, strengthening the “motivic” result even in Yun’s original context. Rather than showing (as in Theorem 1.4(2)) that the $\tilde{\rho}_{\chi,t}$ (or $\rho_{\chi,t}$) are weakly motivic, we show (Theorem 6.1) that, for any finite-dimensional representation r of \tilde{H}_1 , $r \circ \tilde{\rho}_{\chi,t}$ is motivated. The content of this assertion is the following: the arguments showing that $\rho_{\chi,t}$ and $\tilde{\rho}_{\chi,t}$ are weakly motivic rest on the fact that quasiminuscule and minuscule affine

⁵Also, in this case, a more elementary construction of the lift can be achieved using Katz’s theory [1996] of rigid local systems; this is a simple case of the strategy of [Patrikis 2014a].

Schubert varieties have very mild singularities (punctual in the quasiminuscule case; none at all in the minuscule case). For such varieties (and their close cousins that appear in the proof), we can in quite elementary terms describe their intersection cohomology groups as the ℓ -adic realizations of motivated motives. The claim that *all* $r \circ \tilde{\rho}_{\chi,t}$ are motivated depends on a similar description, but for varieties with singularities as bad as those of any affine Schubert variety. This essentially means we need a “motivated” description of the intersection cohomology $\mathrm{IH}^*(Y_{\bar{k}}, \mathbb{Q}_\ell)$ of an arbitrarily singular, and not necessarily projective, variety Y over a characteristic-zero field k ; to be precise, since motivated motives do not reflect “mixed” behavior, we prove such an assertion for the associated weight graded $\mathrm{Gr}_i^W \mathrm{IH}^*(Y_{\bar{k}}, \mathbb{Q}_\ell)$. This is deduced as a consequence of a stronger “motivated” variant of the decomposition theorem, and especially from a “motivated support decomposition” — see [Theorem 8.13](#) and [Corollary 8.14](#). Here is the specialized statement for intersection cohomology:

Theorem 1.5 (compare [Corollary 8.15](#)). *Let k be a finitely generated field of characteristic zero, and let Y/k be any quasiprojective variety. Then there is an object $M \in \mathcal{M}_k$ whose ℓ -adic realization is isomorphic as a Γ_k -representation to $\mathrm{Gr}_i^W \mathrm{IH}^m(Y_{\bar{k}}, \mathbb{Q}_\ell)$. If Γ is a finite group scheme over k acting on Y , and $e \in \bar{\mathbb{Q}}[\Gamma(\bar{k})]^{\Gamma_k}$ is an idempotent, then for any embedding $\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_\ell$ there is an object of $\mathcal{M}_{k, \bar{\mathbb{Q}}}$ whose $(\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_\ell)$ -realization is isomorphic as a Γ_k -representation to $\mathrm{Gr}_i^W e(\mathrm{IH}^m(Y_{\bar{k}}, \bar{\mathbb{Q}}_\ell))$.*

The same holds for intersection cohomology with compact supports.

When Y is projective, in which case $\mathrm{IH}^m(Y_{\bar{k}}, \mathbb{Q}_\ell)$ is pure of weight m , and k is algebraically closed, this⁶ is a recent result of de Cataldo and Migliorini [[2014](#), [Theorem 3.2.2](#)], part of their beautiful series of papers (see for instance [[de Cataldo and Migliorini 2005](#); [2010](#); [de Cataldo 2012](#)]) reestablishing the decomposition theorem and its associated mixed Hodge-theoretic package by “geometric”, rather than “sheaf-theoretic”, methods. These papers chart a fundamental advance in our understanding of the geometry of perverse sheaves, and I expect there will be many more, and far deeper, motivic applications than the one here. Since the arguments establishing [Theorem 1.5](#) are independent of those of the rest of this paper, I refer the reader to [Section 7A](#) for a fuller introduction, and for an overview of the approach to [Theorem 1.5](#). Also see [Remark 8.16](#) for additional applications, such as a p -adic de Rham comparison isomorphism for intersection cohomology.

⁶Not exactly, of course, since as we have phrased the result the theorem is vacuous for k algebraically closed; in that case substitute for the ℓ -adic cohomology the collection of Betti, de Rham, and ℓ -adic realizations.

2. Bundles with level structure

Before plunging into the technical details, we give a little more orientation for the reader not familiar with Yun’s argument. The principle underlying Yun’s strategy is that, for a suitable reductive group G and a careful choice of level structure (i.e., ramification), there will be an essentially unique automorphic form on G over the function field $F = \mathbb{F}_q((t))$. The usual double coset space

$$G(F) \backslash G(\mathcal{A}_F) / \prod_x G(\mathcal{O}_x),$$

the product taken over closed points of X , and \mathcal{O}_x the complete local ring at x , on which automorphic forms are defined admits an interpretation as the \mathbb{F}_q -points of the moduli stack Bun_G of G -bundles on \mathbb{P}^1 . Appropriate moduli spaces of bundles with level structure then have \mathbb{F}_q -points corresponding to taking other level structures in the above double quotient. The general aspiration of (“classical”) geometric Langlands is to upgrade these automorphic functions to perverse sheaves (via Grothendieck’s function-sheaf correspondence) on the appropriate moduli stack of bundles. The automorphic interpretation plays no direct role in Yun’s work, but it serves as motivation. Setting the motivation aside, the problem becomes one of finding level structures corresponding to some moduli space Bun whose simple perverse sheaves can be explicitly described (in fact, what is described in Yun’s case is a subset of “odd” perverse sheaves). To construct eigen-local systems associated to any of these perverse sheaves, one needs to know that they are Hecke eigensheaves. It is here that the uniqueness properties of the construction are absolutely essential: the classical analogue to keep in mind is the statement that in a one-dimensional space of classical modular forms, every element must be a Hecke eigenform! The motivic nature of the resulting local systems is only revealed by carefully tracing through the construction.

The present section reviews facts about spaces of bundles with level structure. The next section then lays out carefully the construction of Yun’s moduli space Bun , and of the enlarged moduli space $\widetilde{\text{Bun}}$ essential to our generalization; this latter space will be chosen to make as easy as possible a comparison of perverse sheaves on the two spaces.

We now proceed to the formal exposition. In this section only, we allow G to be any connected reductive group over a field k , and X to be any smooth projective geometrically connected curve over k . Our aim is to review a construction from [Yun 2011, §4.2] of moduli spaces of G -bundles on X with level structure at a finite set $S = \{x_1, \dots, x_n\} \subset X(k)$ of k -points. Here and throughout, we denote by LG and L^+G the “abstract” loop group and positive loop group of G , i.e., the functors of k -algebras given by $R \mapsto G(R((t)))$ and $R \mapsto G(R[[t]])$ (a group ind-scheme and pro-algebraic group, respectively, over k), where t is a formal parameter. Now

let x be a closed point of X , and denote by \mathcal{O}_x the complete local ring of X at x , with residue field $\kappa(x)$ and fraction field \mathcal{K}_x . Then we denote by $L_x G$ and $L_x^+ G$ the functors $R \mapsto G(R \widehat{\otimes}_{\kappa(x)} \mathcal{K}_x)$ and $R \mapsto G(R \widehat{\otimes}_{\kappa(x)} \mathcal{O}_x)$.

Definition 2.1. Let $\text{Bun}_{G,S,\infty} \rightarrow \text{Ring}_k$ be the stack associated to the following prestack $\text{Bun}_{G,S,\infty}^{\text{pre}}$ over k : for any k -algebra R , $\text{Bun}_{G,S,\infty}^{\text{pre}}(R)$ is the groupoid of triples $(\alpha, \mathcal{P}, \tau)$ where:

- $\alpha = (\alpha_{x_i})_{i=1,\dots,n}$ is a collection of local coordinates $\alpha_{x_i} : R[[t]] \xrightarrow{\sim} \mathcal{O}_{x_i}$ (here we regard x_i as an R -point $x_i : \text{Spec } R \rightarrow X_R$ and take the formal completion of X_R along the graph $\Gamma(x_i)$).
- \mathcal{P} is a G -torsor on X_R .
- $\tau = (\tau_{x_i})_{i=1,\dots,n}$ is a collection of full level structures

$$\tau_{x_i} : G \times \mathcal{D}_{x_i} \xrightarrow{\sim} \mathcal{P}|_{\mathcal{D}_{x_i}},$$

where $\mathcal{D}_{x_i} = \text{Spec}(\mathcal{O}_{x_i})$.

Let $\text{Aut}_{\mathcal{O}}$ denote the pro-algebraic group of continuous automorphisms of $k[[t]]$. The semidirect product

$$(LG \rtimes \text{Aut}_{\mathcal{O}})^n$$

acts on the right on $\text{Bun}_{G,S,\infty}$ as follows.

Definition 2.2. For $g = (g_i)_{i=1,\dots,n} \in G(R((t)))^n$ and $\sigma = (\sigma_i)_{i=1,\dots,n} \in \text{Aut}(R[[t]])^n$, and $(\alpha, \mathcal{P}, \tau) \in \text{Bun}_{G,S,\infty}^{\text{pre}}(R)$, let (g, σ) act on $(\alpha, \mathcal{P}, \tau)$ by

$$R_{g,\sigma}(\alpha, \mathcal{P}, \tau) = (\alpha \circ \sigma, \mathcal{P}^g, \tau^g),$$

where:

- $\alpha \circ \sigma = (\alpha_{x_i} \circ \sigma_i)_i$.
- \mathcal{P}^g is the G -bundle on X_R obtained by gluing $\mathcal{P}|_{X_R - \bigcup_i \Gamma(x_i)}$ to the trivial G -bundles on the completions $\mathcal{D}_{x_i} = \mathcal{O}_{x_i}$ along the punctured discs $\mathcal{D}_{x_i}^\times$ via the isomorphisms

$$G \times \mathcal{D}_{x_i}^\times \xrightarrow{\alpha_{x_i} \circ g_i \circ \alpha_{x_i}^{-1}} G \times \mathcal{D}_{x_i}^\times \xrightarrow{\tau_{x_i}} \mathcal{P}|_{\mathcal{D}_{x_i}^\times}.$$

- $\tau^g = (\tau_{x_i}^g)_{i=1,\dots,n}$ consists of the *tautological* trivializations of \mathcal{P}^g over each \mathcal{D}_{x_i} coming from the definition of \mathcal{P}^g .

At each of the points x_i , we now fix a pro-algebraic subgroup P_i of LG that is stable under the action of $\text{Aut}_{\mathcal{O}}$; we additionally require that, for some integer m , P_i should contain the subgroup

$$I(m) = \{g \in L^+ G : g \equiv 1 \pmod{t^m}\}$$

in finite codimension.

Definition 2.3. Having fixed $S = \{x_1, \dots, x_n\}$ and $\mathbf{P}_1, \dots, \mathbf{P}_n$ as above, we define $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)$ to be the stack associated to the quotient prestack

$$R \mapsto \text{Bun}_{G,S,\infty}(R) / \prod_{i=1}^n (\mathbf{P}_i \rtimes \text{Aut}_{\mathcal{O}}(R)).$$

When there is no risk of confusion, we omit the subscript S from the notation and simply write $\text{Bun}_G(\mathbf{P}_1, \dots, \mathbf{P}_n)$.

Note that since the action of $(LG \rtimes \text{Aut}_{\mathcal{O}})^n$ does not necessarily preserve the isomorphism class of the G -torsor \mathcal{P} on X_R , the moduli space $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)$ need not have a projection to Bun_G . The action does not alter $\mathcal{P}|_{X_R - \bigcup_i \Gamma(x_i)}$, however, so an object of $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)(R)$ does yield a well-defined G -torsor on this complement. Also, the category $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)$ has a tautological object given by taking the image of an object of $\text{Bun}_{G,S,\infty}^{\text{pre}}(k)$ given by the trivial bundle with its tautological level structures and any fixed choice of local coordinates α_{x_i} . (For any two such choices, the resulting objects of $\text{Bun}_{G,S,\infty}(k)$ become uniquely isomorphic modulo the $\text{Aut}_{\mathcal{O}}^n$ -action.)

Lemma 2.4. $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)$ is an algebraic stack locally of finite type.

Proof. This follows exactly as in [Yun 2014a, Corollary 4.2.6], by first deducing the result for

$$\text{Bun}_{G,S}(\mathbf{I}_1(m), \dots, \mathbf{I}_n(m))$$

from the (well-known) result for Bun_G , and from there deducing the case of $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)$. □

Just as in [Yun 2014a, Lemma 4.2.5], we also have:

Lemma 2.5. For each $i = 1, \dots, n$, let

$$\Omega_{x_i} = N_{LG}(\mathbf{P}_i) / \mathbf{P}_i.$$

Then there is a right-action of Ω_{x_i} on $\text{Bun}_{G,S}(\mathbf{P}_1, \dots, \mathbf{P}_n)$.

Finally, we can replace any \mathbf{P}_i by some finite cover, still acting on $\text{Bun}_{G,S,\infty}$ on the right through \mathbf{P}_i ; Lemmas 2.4 and 2.5 continue to hold.

Remark 2.6. For the reader’s convenience, we put this statement in its classical context: Let

$$\Theta := G(F) \backslash G(\mathbb{A}_F) / \left(\prod_{x \neq x_i} G(\mathcal{O}_x) \times \prod_i \mathbf{P}_i \right).$$

Then on automorphic forms $f : \Theta \rightarrow \bar{\mathbb{Q}}_{\ell}$ over a function field F , we have the usual action by Hecke correspondences arising from decomposing the double coset $\mathbf{P}_i w \mathbf{P}_i$ into single cosets. But when w normalizes \mathbf{P}_i , the Hecke action comes from an actual automorphism (right-translation) of the moduli space Θ .

3. Our setting

Now we describe in detail the moduli spaces of G -bundles studied in this paper, taking [Yun 2014a] as our starting point. Let G be a split (almost-)simple simply connected group over k , satisfying the following two hypotheses:

- G is oddly laced;
- -1 belongs to the Weyl group W_G of G .

Explicitly, we take G to be a split simple simply connected group of type A_1 , D_{2n} , G_2 , E_7 , or E_8 in the Dynkin classification. In fact, as we will see, the results of this paper are only nontrivial when the simply connected and adjoint forms of G differ — so for all practical purposes, we are working with types A_1 , D_{2n} , and E_7 .

Let \tilde{G} be a split connected reductive group over k with derived group equal to G , so that the quotient $\tilde{G}/G = S$ is a torus; call the quotient map $\nu : \tilde{G} \rightarrow S$. Fix a maximal torus \tilde{T} of \tilde{G} and a Borel subgroup \tilde{B} containing \tilde{T} , likewise giving $T = \tilde{T} \cap G$, $B = \tilde{B} \cap G$, and determining based root data for \tilde{G} and G , and an explicit Weyl group W_G defined in terms of T . We denote by \tilde{Z} and Z_G the centers of \tilde{G} and G , and we let \tilde{Z}^0 be the identity component of \tilde{Z} . Note that in all cases under consideration $Z_G = Z_G[2]$. The cases of particular interest for us — in which there is a nontrivial Kuga–Satake lifting problem — are those in which $Z_G \neq \{1\}$, namely types A_1 , D_{2n} , and E_7 . From now on we

assume the characteristic of k is not 2. (4)

In particular, Z_G is a discrete group scheme over k , and the order of the kernel of the isogeny $\tilde{Z} \rightarrow S$ is invertible in k . Our first task is to define the moduli spaces of \tilde{G} -bundles on $X = \mathbb{P}^1$ with level structure that will supply us with Hecke eigensheaves. We first recall the construction in [Yun 2014a]. Yun works with the following conjugacy class of parahoric subgroups in LG (see [Yun 2014a, §§2.2–2.3]). In the apartment $\mathcal{A}(T)$ associated to T of the building of LG , we can choose as origin the point corresponding to the subgroup L^+G , with the resulting identification $\mathcal{A}(T) \cong X_\bullet(T) \otimes \mathbb{R}$. Then under this identification $\frac{1}{2}\rho^\vee$ lies in a unique facet, and we let $P_{\frac{1}{2}\rho^\vee}$ be the parahoric subgroup associated to this facet. More precisely, Bruhat–Tits theory provides, for any facet a in the building of LG , a smooth group scheme \mathcal{P}_a over $k[[t]]$ with connected fibers whose generic fiber is $G \times_{\text{Spec } k} \text{Spec } k((t))$. We define P_a to be the pro-algebraic subgroup of LG representing the functor (of k -algebras)

$$R \mapsto \mathcal{P}_a(R[[t]]).$$

We then apply this construction to the case where a is the facet containing $\frac{1}{2}\rho^\vee$. Let K denote the maximal reductive quotient of $P_{\frac{1}{2}\rho^\vee}$; since G is simply connected, K is

connected. Moreover, Yun shows [2014a, §2.5] that K has a canonical connected double cover:

Definition 3.1. Let ${}^{(2)}K$ denote the connected double cover of K , so there is an exact sequence

$$1 \rightarrow \mu_2^{\ker} \rightarrow {}^{(2)}K \rightarrow K \rightarrow 1.$$

Note that our notation differs from that of [Yun 2014a, §2.5], where this group is denoted \tilde{K} ; we reserve $(\tilde{*})$ for groups associated with the enlargement \tilde{G} of G .

We now define the particular moduli stacks of interest, beginning with the ones used in [Yun 2014a]. Let $P_0 \subset L_0G$ be the parahoric subgroup in the conjugacy class of $P_{\frac{1}{2}\rho^\vee}$ that contains the Iwahori $I_0 \subset L_0^+G$, defined in terms of B . Moreover, let

$${}^{(2)}P_0 = P_0 \times_{K_0} {}^{(2)}K_0,$$

and let P_0^+ denote the pro-unipotent radical of P_0 . Next let P_∞ be the parahoric in the conjugacy class of $P_{\frac{1}{2}\rho^\vee}$ that contains the Iwahori $I_\infty^{\text{op}} \subset L_\infty^+G$ defined in terms of B^{op} . Finally, let $I_1 \subset L_1^+G$ denote the Iwahori subgroup defined again in terms of B . In the notation of Section 2, we now let $S = \{0, 1, \infty\} \subset \mathbb{P}^1(k)$; for later reference, we let X^0 be the variety $\mathbb{P}^1 - S$ over k . The primary object of study in [Yun 2014a] is the moduli space (see Definition 2.3)

$$\text{Bun} = \text{Bun}_G({}^{(2)}P_0, I_1, P_\infty).$$

This sits in the diagram

$$\begin{array}{ccccc} \text{Bun}^+ & \longrightarrow & \text{Bun}_G(P_0^+, P_\infty) & & \\ \downarrow & & \downarrow & \searrow & \\ \text{Bun} & \longrightarrow & \text{Bun}_G({}^{(2)}P_0, P_\infty) & \longrightarrow & \text{Bun}_G(P_0, P_\infty) \end{array} \tag{5}$$

in which $\text{Bun}^+ = \text{Bun}_G(P_0^+, I_1, P_\infty)$. The vertical maps are ${}^{(2)}K_0$ -torsors, and the square is 2-cartesian. Note too that the fibers of horizontal arrows in the square are isomorphic to the flag variety fl_G of G .

Next we modify these constructions to define the corresponding moduli stacks of \tilde{G} -bundles on X . There are various ways of doing this; we take care to choose the new level structures so that the moduli spaces in the G and \tilde{G} cases are most easily compared.

Definition 3.2. Let \tilde{P}_∞ be the subgroup scheme of $L_\infty\tilde{G}$ generated by P_∞ and $L_\infty^+(\tilde{Z}^0)$. Let $\tilde{P}_0(1)$ be the subgroup scheme of $L_0\tilde{G}$ generated by P_0 and the pro-algebraic group $\tilde{Z}^0(1)$ defined as the kernel of reduction modulo t (a local coordinate at zero),

$$\tilde{Z}^0(1) = \ker(L_0^+(\tilde{Z}^0) \rightarrow \tilde{Z}^0).$$

Note that $\tilde{\mathbf{P}}_0(1)$ is isomorphic to the direct product $\mathbf{P}_0 \times \tilde{\mathbf{Z}}^0(1)$: the restriction of ν to $\nu : \tilde{\mathbf{Z}}^0 \rightarrow S$ can be identified with a product of maps $\mathbb{G}_m \rightarrow \mathbb{G}_m$, each given by multiplication by some $n \in \{\pm 1, \pm 2\}$, so (working one coordinate at a time), for any $x \in R[[t]]$, the equation

$$1 = \nu(1 + tx) = 1 + ntx + (\text{higher order terms})$$

forces $x = 0$, since we have assumed (see (4)) that 2 is invertible in k . Moreover, the pro-unipotent radical of $\tilde{\mathbf{P}}_0(1)$ is $\mathbf{P}_0^+ \cdot \tilde{\mathbf{Z}}^0(1)$, so the maximal reductive quotient of $\tilde{\mathbf{P}}_0(1)$ is also K_0 . In particular, we can form the analogous group ${}^{(2)}\tilde{\mathbf{P}}_0(1)$ by pullback.

Finally, let $\tilde{\mathbf{I}}_1$ denote the Iwahori subgroup associated to $\tilde{\mathbf{B}}$ in $L_1^+ \tilde{\mathbf{G}}$. With this notation in place, we introduce our main object of study:

Definition 3.3. Let $\widetilde{\text{Bun}}$ denote the algebraic stack $\text{Bun}_{\tilde{\mathbf{G}}}({}^{(2)}\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{I}}_1, \tilde{\mathbf{P}}_\infty)$.

Similarly setting $\widetilde{\text{Bun}}^+ = \text{Bun}_{\tilde{\mathbf{G}}}(\mathbf{P}_0^+ \cdot \tilde{\mathbf{Z}}^0(1), \tilde{\mathbf{I}}_1, \tilde{\mathbf{P}}_\infty)$, we then have the $\tilde{\mathbf{G}}$ -analogue of the basic diagram (5):

$$\begin{array}{ccccc}
 \widetilde{\text{Bun}}^+ & \longrightarrow & \text{Bun}_{\tilde{\mathbf{G}}}(\mathbf{P}_0^+ \cdot \tilde{\mathbf{Z}}^0(1), \tilde{\mathbf{P}}_\infty) & & \\
 \downarrow & & \downarrow & \searrow & \\
 \widetilde{\text{Bun}} & \longrightarrow & \text{Bun}_{\tilde{\mathbf{G}}}({}^{(2)}\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{P}}_\infty) & \longrightarrow & \text{Bun}_{\tilde{\mathbf{G}}}(\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{P}}_\infty)
 \end{array} \tag{6}$$

Here the vertical maps are still ${}^{(2)}K_0$ -torsors, the diagram is 2-cartesian, and again the fibers of the horizontal arrows in the square are copies of $\mathfrak{fl}_{\tilde{\mathbf{G}}}$.

We now must recall the Birkhoff decomposition and uniformization results for G - (or $\tilde{\mathbf{G}}$ -) bundles on $X = \mathbb{P}^1$. Consider the “trivial G -bundle on \mathbb{A}^1 with tautological \mathbf{P}_0 -level structure” $\mathcal{P}_{\mathbb{A}^1}^0$; to be precise, $\mathcal{P}_{\mathbb{A}^1}^0$ is defined as in Section 2, and is not literally a G -bundle on \mathbb{A}^1 . Likewise, let $\tilde{\mathcal{P}}_{\mathbb{A}^1}^0$ be the trivial $\tilde{\mathbf{G}}$ -bundle on \mathbb{A}^1 with tautological $\tilde{\mathbf{P}}_0(1)$ level structure at zero. Let Γ_0 and $\tilde{\Gamma}_0$ denote the group ind-schemes of automorphisms of $\mathcal{P}_{\mathbb{A}^1}^0$ and $\tilde{\mathcal{P}}_{\mathbb{A}^1}^0$, respectively. Also let W^{aff} denote the affine Weyl group $X_\bullet(T) \rtimes W_G$, and let $\tilde{W} = X_\bullet(\tilde{T}) \rtimes W_G$ denote the Iwahori–Weyl group of $\tilde{\mathbf{G}}$. The Weyl group of the reductive quotient K_∞ of \mathbf{P}_∞ can be identified with a subgroup of W^{aff} : take the subgroup generated by simple reflections that fix the alcove of \mathbf{P}_∞ . The same holds for the reductive quotient K_0 of \mathbf{P}_0 and its Weyl group, and in both cases we write the resulting subgroup of W^{aff} as W_K .

Lemma 3.4. *There are isomorphisms of stacks*

$$[\Gamma_0 \backslash L_\infty G / \mathbf{P}_\infty] \xrightarrow{\sim} \text{Bun}_G(\mathbf{P}_0, \mathbf{P}_\infty), \tag{7}$$

$$[\tilde{\Gamma}_0 \backslash L_\infty \tilde{\mathbf{G}} / \tilde{\mathbf{P}}_\infty] \xrightarrow{\sim} \text{Bun}_{\tilde{\mathbf{G}}}(\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{P}}_\infty), \tag{8}$$

and Birkhoff decompositions

$$L_\infty G(\bar{k}) = \coprod_{W_K \setminus W^{\text{aff}}/W_K} \Gamma_0(\bar{k})w\mathbf{P}_\infty(\bar{k}), \tag{9}$$

$$L_\infty \tilde{G}(\bar{k}) = \coprod_{W_K \setminus \tilde{W}/W_K} \tilde{\Gamma}_0(\bar{k})w\tilde{\mathbf{P}}_\infty(\bar{k}). \tag{10}$$

Proof. See [Yun 2014a, §3.2.2] and [Heinloth et al. 2013, Proposition 1.1]. \square

It follows easily from diagram (6) that $\pi_0(\widetilde{\text{Bun}})$ is naturally in bijection with $\pi_0(\text{Bun}_{\tilde{G}}(\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{P}}_\infty))$; this is in turn in bijection (since G is simply connected) with

$$\pi_0(\text{Bun}_{\tilde{G}}) \xrightarrow{\nu} \pi_0(\text{Bun}_S) \xleftarrow{\sim} \pi_0(L_\infty S/L_\infty^+ S) \xleftarrow{\sim} X_\bullet(S).$$

We can describe the connected components of $\widetilde{\text{Bun}}$ in terms of this uniformization. First note that replacing \mathbf{P}_0 and Γ_0 with ${}^{(2)}\mathbf{P}_0$ and ${}^{(2)}\Gamma_0$, and $\tilde{\mathbf{P}}_0(1)$ and $\tilde{\Gamma}_0$ with ${}^{(2)}\tilde{\mathbf{P}}_0(1)$ and ${}^{(2)}\tilde{\Gamma}_0$, we get obvious analogues of Lemma 3.4. Then, for each $w \in W_K \setminus \tilde{W}/W_K$ we obtain an object $\tilde{\mathcal{P}}_w$ of $\text{Bun}_{\tilde{G}}({}^{(2)}\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{P}}_\infty)(k)$ by gluing $\tilde{\mathcal{P}}_{\mathbb{A}^1}^0$ with $\text{Ad}(w)\tilde{\mathbf{P}}_\infty$; and we can make the corresponding construction of $\mathcal{P}_w \in \text{Bun}_G({}^{(2)}\mathbf{P}_0, \mathbf{P}_\infty)$ for $w \in W^{\text{aff}}$. The stabilizers of \mathcal{P}_w and $\tilde{\mathcal{P}}_w$ are, respectively,

$$\text{Stab}_w^G = (\Gamma_0 \cap w\mathbf{P}_\infty w^{-1}) \times_{K_0} {}^{(2)}K_0, \tag{11}$$

$$\text{Stab}_w^{\tilde{G}} = (\tilde{\Gamma}_0 \cap w\tilde{\mathbf{P}}_\infty w^{-1}) \times_{K_0} {}^{(2)}K_0. \tag{12}$$

In other words, $\text{Bun}_G({}^{(2)}\mathbf{P}_0, \mathbf{P}_\infty)$ has a stratification by substacks $[\{\mathcal{P}_w\}/\text{Stab}_w^G]$; likewise, $\text{Bun}_{\tilde{G}}({}^{(2)}\tilde{\mathbf{P}}_0(1), \tilde{\mathbf{P}}_\infty)$ has a stratification by substacks $[\{\tilde{\mathcal{P}}_w\}/\text{Stab}_w^{\tilde{G}}]$. By taking the preimages in Bun and $\widetilde{\text{Bun}}$, we obtain stratifications by substacks that we denote Bun_w (for $w \in W_K \setminus W^{\text{aff}}/W_K$) and $\widetilde{\text{Bun}}_w$ (for $w \in W_K \setminus \tilde{W}/W_K$), respectively. For $w = \lambda \rtimes w_G \in \tilde{W} = X_\bullet(\tilde{T}) \rtimes W_G$, the substack $\widetilde{\text{Bun}}_w$ lies in the component corresponding to $\nu \circ \lambda \in X_\bullet(S)$. In particular, we can identify the connected component of $\widetilde{\text{Bun}}$ containing the tautological object \mathcal{P}_1 as

$$\widetilde{\text{Bun}}^0 = \coprod_{\substack{w=\lambda \rtimes w_G \in W_K \setminus \tilde{W}/W_K \\ \lambda \in X_\bullet(\tilde{T})}} \widetilde{\text{Bun}}_w = \coprod_{w \in W_K \setminus W^{\text{aff}}/W_K} \widetilde{\text{Bun}}_w.$$

Taking the associated \tilde{G} -bundle defines a map $\text{Bun} \rightarrow \widetilde{\text{Bun}}$, and for w in $W_K \setminus W^{\text{aff}}/W_K$ it respects the above stratifications, yielding a map $\text{Bun}_w \rightarrow \widetilde{\text{Bun}}_w$. The crucial point is the following:

Proposition 3.5. *The map $\text{Bun} \rightarrow \widetilde{\text{Bun}}^0$ is an equivalence, i.e., an isomorphism of stacks.*

Proof. We check this stratum by stratum. It suffices to show that, for all w in $X_*(T) \rtimes W_G = W^{\text{aff}} \subset \tilde{W}$, we have $\text{Stab}_w^G = \text{Stab}_w^{\tilde{G}}$, i.e., that the natural map

$$\Gamma_0 \cap w \mathbf{P}_\infty w^{-1} \rightarrow \tilde{\Gamma}_0 \cap w \tilde{\mathbf{P}}_\infty w^{-1}$$

is an isomorphism. For a k -algebra R , an element of $(\tilde{\Gamma}_0 \cap w \tilde{\mathbf{P}}_\infty w^{-1})(R)$ gives rise fppf-locally on R to an equation of the form $p_0 z_0 = w z_\infty p_\infty w^{-1}$ with $p_0 \in \mathbf{P}_0(R)$, $z_0 \in \tilde{Z}^0(1)(R)$, $p_\infty \in \mathbf{P}_\infty(R)$, and $z_\infty \in L_\infty^+(\tilde{Z}^0)$. Applying ν , we find $\nu(z_0) = \nu(z_\infty)$; but since $1 + tR[[t]] \cap R[[t^{-1}]]^\times = \{1\}$, we see that $\nu(z_0) = \nu(z_\infty) = 1$. This forces (as in the argument following Definition 3.2, by our assumption on $\text{char}(k)$) $z_0 = 1$, and $z_\infty \in \mathbf{P}_\infty(R)$. We may as well then assume $z_\infty = 1$ (incorporating z_∞ into p_∞), and so we actually have an equality $p_0 = w p_\infty w^{-1}$ bearing witness to an element of $(\Gamma_0 \cap w \mathbf{P}_\infty w^{-1})(R)$. This implies that

$$\Gamma_0 \cap w \mathbf{P}_\infty w^{-1} \rightarrow \tilde{\Gamma}_0 \cap w \tilde{\mathbf{P}}_\infty w^{-1}$$

is an epimorphism, and, as it is obviously injective, we are done. □

4. The eigensheaves

4A. Construction of the eigensheaves. In this section, we combine the equivalence $\text{Bun} \xrightarrow{\sim} \widetilde{\text{Bun}}^0$ of Proposition 3.5 with the analysis of the sheaf theory of Bun carried out in [Yun 2014a, Theorem 3.2] to produce our desired Hecke eigensheaves on $\widetilde{\text{Bun}}$. The key simplification arises from applying Lemma 2.5 at the point $x = 1$, where we have taken \tilde{I}_1 level structure. In this case we identify the group Ω_1 with the stabilizer in \tilde{W} of the alcove corresponding to the standard Iwahori \tilde{I}_1 , and $\nu : \Omega_1 \xrightarrow{\sim} X_*(S)$ also identifies Ω_1 with $\pi_0(\widetilde{\text{Bun}})$. For $\gamma \in \Omega_1$, we denote by

$$\mathbb{T}_\gamma : \widetilde{\text{Bun}} \rightarrow \widetilde{\text{Bun}}$$

the action given by Lemma 2.5. Writing $\widetilde{\text{Bun}}^\gamma$ for the connected component corresponding to γ , we see that \mathbb{T}_γ induces isomorphisms

$$\mathbb{T}_\gamma : \widetilde{\text{Bun}}^0 \xrightarrow{\sim} \widetilde{\text{Bun}}^\gamma.$$

In particular, all connected components of $\widetilde{\text{Bun}}$ are isomorphic (compare [Heinloth et al. 2013, Corollary 1.2]).⁷ The idea is to take Yun’s construction of a perverse Hecke eigensheaf on $\text{Bun} \xrightarrow{\sim} \widetilde{\text{Bun}}^0$, and then use the (“ramified Hecke operators”) \mathbb{T}_γ to propagate the eigensheaf to the other connected components of $\widetilde{\text{Bun}}$. We begin by reviewing Yun’s construction [2014a, §3]. The tautological object in $\text{Bun}_G(\mathbf{P}_0, \mathbf{P}_\infty)$ (with automorphism group K_0) has preimage in Bun equivalent to

⁷Note that this is a special, and highly simplifying, feature of our particular context; for contrast, observe that the degree-0 and degree-1 connected components of Bun_{GL_2} ($X = \mathbb{P}^1$ still) are not isomorphic, since no degree-1 vector bundle has GL_2 as its automorphism group.

a quotient $[(^{(2)}K_0 \setminus \text{fl}_G)]$, for a suitable action of K_0 on fl_G (see [Yun 2014a, §3.2.4]). The group K_0 acts on fl_G with finitely many orbits, so there is a unique open orbit $U \subset \text{fl}_G$, giving open embeddings

$$[(^{(2)}K_0 \setminus U)] \subset [(^{(2)}K_0 \setminus \text{fl}_G)] \subset \text{Bun}.$$

As in [Yun 2014a, §3.2.5], we fix a point $u_0 \in U(\mathbb{Z}[1/N])$ (for some N sufficiently large, and for an integral model of U arising from extending K_0 and G to split reductive group schemes over some $\mathbb{Z}[1/M]$), and

denote by $u_0 \in U(k)$ the induced k -point

$$\text{for all } k \text{ of sufficiently large characteristic.} \tag{13}$$

This choice is in effect from now on. As an element of $U(k) \subset \text{fl}_G(k)$, u_0 corresponds to a Borel subgroup $B_0 \subset G$ over k which is in general position with respect to K_0 :

Definition 4.1. Let A denote the finite group scheme $B_0 \cap K_0$ over k . Let $(^{(2)}A)$ denote the double cover of A given by pullback along $(^{(2)}K_0 \twoheadrightarrow K_0)$.⁸ Finally, let $Z(^{(2)}A)$ denote the center of $(^{(2)}A)$.

Recall the following results [Yun 2014a, §2.6] on the structure and representation theory of the finite 2-group $(^{(2)}A(\bar{k}))$. Recall that we have set

$$\mathbb{Q}' = \begin{cases} \mathbb{Q} & \text{if } G \text{ is of type } D_{4n}, G_2, \text{ or } E_8, \\ \mathbb{Q}(\mathbf{i}) & \text{if } G \text{ is of type } A_1, D_{4n+2}, \text{ or } E_7, \end{cases} \tag{14}$$

and have also set $\mathbb{Q}'_\ell = \mathbb{Q}_\ell(\mathbf{i})$. All sheaves considered will be \mathbb{Q}'_ℓ -sheaves. In parallel to this condition on the coefficients, we impose the following restriction on the field of definition k , in effect for the rest of this paper:

$$\sqrt{-1} \in k \text{ for } G \text{ of type } A_1, D_{4m+2}, \text{ or } E_7. \tag{15}$$

Lemma 4.2. Assume k satisfies condition (15), so that Γ_k acts trivially on $Z(^{(2)}A)(\bar{k})$.

- (1) $(^{(2)}Z_G) = Z(^{(2)}A)$.
- (2) Restriction to $(^{(2)}Z_G)(\bar{k})$ gives a bijection between irreducible odd representations of $(^{(2)}A)(\bar{k})$ and odd characters of $Z(^{(2)}A)(\bar{k})$:

$$\text{Irr}_{\mathbb{Q}}(^{(2)}A(\bar{k}))_{\text{odd}} \xrightarrow{\sim} \text{Hom}(Z(^{(2)}A)(\bar{k}), \bar{\mathbb{Q}}^\times)_{\text{odd}} = \text{Hom}(Z(^{(2)}A)(\bar{k}), \mathbb{Q}'^\times)_{\text{odd}}. \tag{16}$$

- (3) If k is a finite field, local field, or number field, then for each odd $\chi : Z(^{(2)}A)(\bar{k}) \rightarrow \mathbb{Q}'^\times$ the corresponding irreducible representation V_χ of $(^{(2)}A)(\bar{k})$ descends to an irreducible representation of $(^{(2)}A)(\bar{k}) \rtimes \Gamma_k$, whose coefficients can be taken to be $\mathbb{Q}(\mathbf{i})$.

⁸Note that this is what Yun denotes \tilde{A} .

Proof. The first claim is [Yun 2014a, Lemma 2.6(2)]. The second claim is elementary: the inverse of the isomorphism (16) is given by inducing the central character, up to some multiplicity. The third claim is a variant of [Yun 2014a, Lemma 2.7], whose proof is not complete.⁹ The obstruction to descending V_χ to a representation of $({}^2\mathbf{A}(\bar{k}) \rtimes \Gamma_k)$ is a class in $H^2(\Gamma_k, \bar{\mathbb{Q}}^\times)$. This Galois cohomology group vanishes for the claimed k ; this is elementary for k finite, and for local and especially number fields it is a beautiful theorem of Tate [Serre 1977, Theorem 4]. The argument showing the descended V_χ can be defined with $\mathbb{Q}(i)$ coefficients as in [Yun 2014a, Lemma 2.7]. \square

For clarity, we collect in one place the various conditions in effect on the field of definition k :

Definition 4.3. Consider any odd central character $\chi : Z({}^2\mathbf{A})(\bar{k}) \rightarrow \mathbb{Q}^\times$, with associated irreducible representation V_χ of $({}^2\mathbf{A}(\bar{k}))$. Let k be any field satisfying conditions (13) and (15), and *moreover* for which V_χ satisfies the conclusion of Lemma 4.2(3). Then, from now on, let V_χ denote a fixed choice of descent to an irreducible representation of $({}^2\mathbf{A}(\bar{k}) \rtimes \Gamma_k)$, with $\mathbb{Q}(i)$ coefficients.

We now recall the crucial result analyzing the sheaf theory of Bun , or, in our case, $\widetilde{\text{Bun}}^0$. Throughout, for an algebraic stack \mathfrak{X} over a field k , we will write $D^b(\mathfrak{X})$ for the derived category of bounded complexes of \mathbb{Q}'_ℓ -sheaves with constructible cohomology, as in [Laszlo and Olsson 2008] (if we need to specify another field of coefficients, \mathbb{Q}_ℓ for instance, we will write $D^b(\mathfrak{X}, \mathbb{Q}_\ell)$). Recall [Yun 2014a, §3.3.1] the subcategory

$$D^b(\text{Bun})_{\text{odd}} \subset D^b(\text{Bun})$$

of odd sheaves, on which $\mu_2^{\text{ker}} = \ker({}^2K_0 \rightarrow K_0)$ acts by the sign character. We can similarly define $D^b(\widetilde{\text{Bun}})_{\text{odd}}$, since μ_2^{ker} is also contained in the automorphism group of every object of $\widetilde{\text{Bun}}$. For future reference, let us also note a refinement of this observation: the automorphism group of every object of $\text{Bun}_{\widetilde{G}}(\widetilde{P}_0(1), \widetilde{I}_1, \widetilde{P}_\infty)$ contains the center Z_G of G , and likewise the automorphism group of every object of $\widetilde{\text{Bun}}$ contains the double cover (pullback under $({}^2K_0 \rightarrow K_0)$) $({}^2Z_G)$ of Z_G . We can therefore decompose $D^b(\widetilde{\text{Bun}})$ into a direct sum of categories $D^b(\widetilde{\text{Bun}})_\psi$, indexed over characters $\psi : ({}^2Z_G \rightarrow \bar{\mathbb{Q}}_\ell^\times)$. We, of course, will be interested in the corresponding decomposition of $D^b(\widetilde{\text{Bun}})_{\text{odd}}$ into a direct sum over the odd characters ψ .

We recall the main result analyzing odd sheaves on Bun . Let $j : [({}^2K_0 \setminus U)] \hookrightarrow \text{Bun}$ denote the open inclusion.

⁹Namely, that argument uses the incorrect assertion that $H^2(\bar{\Gamma}, \bar{\mathbb{Q}}^\times) = 0$ for $\bar{\Gamma}$ a finite group isomorphic to a direct sum of $\mathbb{Z}/2\mathbb{Z}$ s.

Theorem 4.4 [Yun 2014a, Theorem 3.2]. *Assume G is the split simple simply connected group of type $A_1, D_{2n}, E_7, E_8,$ or G_2 . Then the restriction*

$$j^* : D^b(\text{Bun})_{\text{odd}} \rightarrow D^b([{}^{(2)}K_0 \setminus U]_{\text{odd}})$$

is an equivalence of categories with quasi-inverse given by $j_! = j_$.*

The analysis of connected components of $\widetilde{\text{Bun}}$ then implies:

Corollary 4.5. *For all $\gamma \in \Omega_1$, consider the composite*

$$j_\gamma = \mathbb{T}_\gamma \circ j : [{}^{(2)}K_0 \setminus U] \hookrightarrow \widetilde{\text{Bun}}^\gamma.$$

Then the restriction

$$j_\gamma^* : D^b(\widetilde{\text{Bun}}^\gamma)_{\text{odd}} \rightarrow D^b([{}^{(2)}K_0 \setminus U]_{\text{odd}})$$

is an equivalence with inverse $j_{\gamma,!} = j_{\gamma,}$.*

Assume k is as in Definition 4.3. We can now define the hoped-for eigensheaves on $\widetilde{\text{Bun}}$ over k , starting from Yun’s construction on Bun . Fix an odd character (recall equation (16))

$$\chi : Z({}^{(2)}A)(\bar{k}) \rightarrow \mathbb{Q}^{\times}, \tag{17}$$

to which we have associated (Lemma 4.2 and Definition 4.3) an irreducible representation V_χ of $({}^{(2)}A)(\bar{k}) \rtimes \Gamma_k$ having χ as central character. By [Yun 2014a, Lemma 3.3], $V_\chi \otimes_{\mathbb{Q}(i)} \mathbb{Q}'_\ell$ is isomorphic to the pullback under u_0 of a geometrically irreducible local system

$$\mathcal{F}_\chi \in \text{Loc}_{({}^{(2)}K_0)}(U, \mathbb{Q}'_\ell)_{\text{odd}},$$

which we view as an object of $D^b([{}^{(2)}K_0 \setminus U]_{\text{odd}})$. Yun’s eigensheaf is then, by [Yun 2014a, Theorem 4.2],

$$j_!(\mathcal{F}_\chi) = j_*(\mathcal{F}_\chi) \in D^b(\text{Bun})_{\text{odd}}.$$

Definition 4.6. Assume k is as in Definition 4.3. Let $\chi : Z({}^{(2)}A)(\bar{k}) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ be any odd character. We let $A_\chi \in D^b(\widetilde{\text{Bun}})_{\text{odd}}$ be the perverse sheaf on $\widetilde{\text{Bun}}$ whose restriction A_χ^γ , for all $\gamma \in \Omega_1$, to $\widetilde{\text{Bun}}^\gamma$ is given by

$$A_\chi^\gamma = A_\chi|_{\widetilde{\text{Bun}}^\gamma} = j_{\gamma,!}\mathcal{F}_\chi = j_{\gamma,*}\mathcal{F}_\chi.$$

That is, we make the only definition compatible with the requirement that A_χ^0 be Yun’s eigensheaf, and that A_χ be eigen for the ramified Hecke operators \mathbb{T}_γ at $1 \in X$.

4B. Geometric Satake equivalence. We recall a convenient form of the geometric Satake equivalence. See [Mirković and Vilonen 2007] and [Yun 2014a, §4.1] for more background. Let \mathcal{G} be any split connected reductive group over k (\mathcal{G} will of course eventually be either G or \tilde{G}). Let $\text{Gr}_{\mathcal{G}} = L\mathcal{G}/L^+\mathcal{G}$ as usual denote the affine Grassmannian of \mathcal{G} . The main result of [Mirković and Vilonen 2007] describes the category $\text{Sat}_{\mathcal{G}}^{\text{geom}}$ of $(L^+\mathcal{G})_{\bar{k}}$ -equivariant perverse sheaves on $\text{Gr}_{\mathcal{G},\bar{k}}$ as follows: $\text{Sat}_{\mathcal{G}}^{\text{geom}}$ admits a convolution product making it a neutral Tannakian category over \mathbb{Q}_{ℓ} with fiber functor

$$H^* : \text{Sat}_{\mathcal{G}}^{\text{geom}} \rightarrow \text{Vect}_{\mathbb{Q}_{\ell}}, \quad \mathcal{K} \mapsto H^*(\text{Gr}_{\mathcal{G},\bar{k}}, \mathcal{K}). \tag{18}$$

This fiber functor induces an equivalence

$$\text{Sat}_{\mathcal{G}}^{\text{geom}} \xrightarrow{\sim} \text{Rep}(\mathcal{G}^{\vee}), \tag{19}$$

where we write \mathcal{G}^{\vee} for the (split form over \mathbb{Q}_{ℓ} of the) dual group of \mathcal{G} . We need a version of $\text{Sat}_{\mathcal{G}}^{\text{geom}}$ over k rather than \bar{k} . It is natural for us to deviate from [Yun 2014a, §4.1] and instead follow the suggestion of [Heinloth et al. 2013, Remark 2.9] and [Frenkel and Gross 2009, §2]. Recall that the simple objects of $\text{Sat}_{\mathcal{G}}^{\text{geom}}$ are given by the intersection cohomology sheaves of the affine Schubert varieties $\text{Gr}_{\mathcal{G},\leq\lambda}$. For all dominant $\lambda \in X_{\bullet}(T)$, we write

$$j_{\lambda} : \text{Gr}_{\mathcal{G},\lambda} \hookrightarrow \text{Gr}_{\mathcal{G}}$$

for the inclusion of the $L^+\mathcal{G}$ -orbit containing t^{λ} . Then by definition the intersection cohomology sheaf of the closure $\text{Gr}_{\mathcal{G},\leq\lambda}$ of $\text{Gr}_{\mathcal{G},\lambda}$ is

$$\text{IC}_{\lambda} = j_{\lambda,*} \mathbb{Q}_{\ell}[\langle 2\rho, \lambda \rangle],$$

the shift reflecting that the dimension of $\text{Gr}_{\mathcal{G},\lambda}$ is $\langle 2\rho, \lambda \rangle$. We will define $\text{Sat}_{\mathcal{G}}$ to be the full subcategory of perverse sheaves on $\text{Gr}_{\mathcal{G}}$ consisting of finite direct sums of arbitrary Tate twists $\text{IC}_{\lambda}(m)$, for all $\lambda \in X_{\bullet}(T)^+$ and $m \in \mathbb{Z}$. Note that, in contrast to [Yun 2014a, §4.1], we do not normalize the weights of the IC_{λ} to be zero; this bookkeeping device frees us from having to choose a square-root of the cyclotomic character,¹⁰ and it ensures that the local systems we eventually construct will specialize (at points of $X^0(K)$, for K/\mathbb{Q}_{ℓ} finite) to de Rham Galois representations. Adapting the argument of [Yun 2014a, §4.1] to our normalization, a result of Arkhipov and Bezrukavnikov [2009, §3] implies that $\text{Sat}_{\mathcal{G}}$ is closed under convolution: to be precise, we have

$$\text{IC}_{\lambda} * \text{IC}_{\mu} \cong \bigoplus_{\nu} \text{IC}_{\nu}(\langle \nu - \lambda - \mu, \rho \rangle)^{\oplus m_{\lambda,\mu}^{\nu}}$$

¹⁰Which of course cannot be done over $k = \mathbb{Q}$, although it is possible over many quadratic extensions of \mathbb{Q} .

for some multiplicities $m_{\lambda,\mu}^v$. Note that $\langle v - \lambda - \mu, \rho \rangle$ is an integer, since only v for which there is an inclusion of highest-weight representations $V_v \hookrightarrow V_\mu \otimes V_\lambda$, and in particular for which $\lambda + \mu - v$ lies in the root lattice, will appear on the right-hand side. We would like to combine the tensor functor

$$H_k^* : \text{Sat}_{\mathcal{G}} \rightarrow \text{Sat}_{\mathcal{G}}^{\text{geom}} \xrightarrow{H^*} \text{Rep}(\mathcal{G}^\vee) \tag{20}$$

with a mechanism for keeping track of the weight and Tate twist. Thus we define a fully faithful tensor functor

$$H_w^* : \text{Sat}_{\mathcal{G}} \rightarrow \text{Rep}(\mathcal{G}^\vee \times \mathbb{G}_m)$$

by additively extending the assignment on simple objects

$$\text{IC}_\lambda(n) \mapsto H_k^*(\text{IC}_\lambda(n)) \boxtimes (z \mapsto z^{(2\rho,\lambda)-2n}).$$

Composing with the canonical fiber functor ω of $\text{Rep}(\mathcal{G}^\vee \times \mathbb{G}_m)$, this yields a surjective homomorphism $\mathcal{G}^\vee \times \mathbb{G}_m \rightarrow \text{Aut}^\otimes(\omega \circ H_w^*)$ whose kernel

$$\left\{ (g, z) \in \mathcal{G}^\vee \times \mathbb{G}_m : \text{for all dominant } \lambda \in X_\bullet(T) \text{ and all } n \in \mathbb{Z}, \right. \\ \left. g \text{ acts on } V_\lambda \text{ by } z^{2n - \langle 2\rho, \lambda \rangle} \right\}$$

is clearly equal to the subgroup $\langle (2\rho(-1), -1) \rangle \subset \mathcal{G}^\vee \times \mathbb{G}_m$. That is, we have a tensor equivalence $\text{Sat}_{\mathcal{G}} \xrightarrow{\sim} \text{Rep}(\mathcal{G}_1^\vee)$, where (following [Frenkel and Gross 2009])


$$\mathcal{G}_1^\vee = (\mathcal{G}^\vee \times \mathbb{G}_m) / \langle (2\rho(-1), -1) \rangle. \tag{21}$$

Note that if \mathcal{G} is simply connected, then \mathcal{G}_1^\vee is isomorphic to $G^\vee \times \mathbb{G}_m$, since $2\rho(-1) = 1$.

4C. Geometric Hecke operators. We briefly recall the definition of geometric Hecke operators in our context, as well as the notion of a Hecke eigensheaf. Recall that the Hecke stack $\widetilde{\text{Hk}}$ associated to $\widetilde{\text{Bun}}$ is the category of tuples $(R, x, \mathcal{P}, \mathcal{P}', \iota)$ where:

- R is a k -algebra;
- $x \in X^0(R)$;
- \mathcal{P} and \mathcal{P}' are objects of $\widetilde{\text{Bun}}(R)$;
- ι is an isomorphism of \mathcal{P} and \mathcal{P}' away from the graph of x .

Projecting such data to (R, x, \mathcal{P}) (the map \overleftarrow{h}) or (R, x, \mathcal{P}') (the map \overrightarrow{h}) gives a correspondence diagram



As explained in [Yun 2014a, §4.1.3] (using the fact that \overrightarrow{h} and \overleftarrow{h} are locally trivial fibrations in the smooth topology, with fibers isomorphic to $\mathrm{Gr}_{\tilde{G}}$ — see [Heinloth et al. 2013, Remark 4.1]), or slightly differently in [Yun 2013, §4.3.1], for each $\mathcal{K} \in \mathrm{Sat}_{\tilde{G}}$ there is an object $\mathcal{K}_{\mathrm{HK}} \in D^b(\mathrm{HK}, \mathbb{Q}_\ell)$ whose restriction to each geometric fiber of \overrightarrow{h} is isomorphic to \mathcal{K} . As usual, the (universal) geometric Hecke operator is the functor

$$\begin{aligned} \mathbb{T} : \mathrm{Sat}_{\tilde{G}} \times D^b(\widetilde{\mathrm{Bun}} \times X^0) &\rightarrow D^b(\widetilde{\mathrm{Bun}} \times X^0), \\ (\mathcal{K}, \mathcal{F}) &\mapsto \overrightarrow{h}_! (\overleftarrow{h}^*(\mathcal{F}) \otimes_{\mathbb{Q}_\ell} \mathcal{K}_{\mathrm{HK}}). \end{aligned} \tag{22}$$

The induced functor

$$\mathrm{Sat}_{\tilde{G}} \rightarrow \mathrm{End}(D^b(\widetilde{\mathrm{Bun}} \times X^0)) \tag{23}$$

is monoidal. When the input from $D^b(\widetilde{\mathrm{Bun}} \times X^0)$ is of the form $\mathcal{F} \boxtimes \overline{\mathbb{Q}}_\ell$ for some $\mathcal{F} \in D^b(\widetilde{\mathrm{Bun}})$, we write

$$\mathbb{T}_{\mathcal{K}}(\mathcal{F}) = \mathbb{T}(\mathcal{K}, \mathcal{F} \boxtimes \overline{\mathbb{Q}}_\ell).$$

Finally, recall the definition of a Hecke eigensheaf:

Definition 4.7. Let \mathcal{F} be an object of $D^b(\widetilde{\mathrm{Bun}})$. We say that \mathcal{F} is a Hecke eigensheaf if there exists

- a tensor functor $\tilde{\mathcal{E}} : \mathrm{Sat}_{\tilde{G}} \rightarrow \mathrm{Loc}(X^0)$;
- a system of isomorphisms, for all $\mathcal{K} \in \mathrm{Sat}_{\tilde{G}}$,

$$\epsilon_{\mathcal{K}} : \mathbb{T}_{\mathcal{K}}(\mathcal{F}) \xrightarrow{\sim} \mathcal{F} \boxtimes \tilde{\mathcal{E}}(\mathcal{K}),$$

satisfying compatibility conditions that will not concern us (see [Gaitsgory 2007, following Proposition 2.8]).

In this case we call $\tilde{\mathcal{E}}$ the eigen-local system of \mathcal{F} .

4D. Proof of the eigensheaf property. Recall that we have fixed a point

$$u_0 : \mathrm{Spec} k \rightarrow U.$$

We also write u_0 for the induced maps $\mathrm{Spec} k \rightarrow [{}^{(2)}K_0 \setminus U] \subset \widetilde{\mathrm{Bun}}^0 \subset \widetilde{\mathrm{Bun}}$. For all $\gamma \in \Omega_1$, we can compose with \mathbb{T}_γ to obtain

$$u_\gamma : \mathrm{Spec} k \rightarrow \widetilde{\mathrm{Bun}}^\gamma.$$

From Corollary 4.5, we obtain equivalences

$$(u_\gamma \times \mathrm{id})^* : D^b(\widetilde{\mathrm{Bun}}^\gamma \times X^0)_{\mathrm{odd}} \xrightarrow{\sim} D_{(2)A}(X^0)_{\mathrm{odd}}, \tag{24}$$

where $(2)A$ acts trivially on X^0 . The strategy for proving that A_χ is an eigensheaf (χ as in (17)) is to show that, for all $\gamma \in \Omega_1$ and all $\mathcal{K} \in \mathrm{Sat}_{\tilde{G}}$, $(u_\gamma \times \mathrm{id})^* \mathbb{T}_{\mathcal{K}}(A_\chi)$

is concentrated in a single perverse degree. Such sheaves A_χ can then be explicitly described via [Corollary 4.5](#) and an analogue of [[Yun 2014a](#), Lemma 3.4]. In preparation for this computation, note that the $\mathbb{T}_{\gamma,!}$ and \mathbb{T}_γ^* commute with the $\mathbb{T}_\mathcal{K}$. Informally, this is the statement that ‘‘Hecke operators at different places commute’’; more formally, the stack $\widetilde{\text{Hk}}$ carries an Ω_1 -action compatible with its two projections \overleftarrow{h} and \overrightarrow{h} . Furthermore, the spread-out sheaves $\mathcal{K}_{\widetilde{\text{Hk}}}$ (for all $\mathcal{K} \in \text{Sat}_{\widetilde{\mathcal{G}}}$) are Ω_1 -equivariant, so we find that

$$\begin{aligned} (u_\gamma \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi) &\cong (u_0 \times \text{id})^* (\mathbb{T}_\gamma \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi) \\ &\cong (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(\mathbb{T}_\gamma^* A_\chi) \cong (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi). \end{aligned} \tag{25}$$

Now consider the following diagram, where declaring the squares cartesian defines the new objects $\widetilde{\text{GR}}$ and $\widetilde{\text{GR}}_\gamma^U$:

$$\begin{array}{ccccc} & & \widetilde{\text{GR}}_\gamma^U & \xrightarrow{j_\gamma} & \widetilde{\text{GR}} & \longrightarrow & \widetilde{\text{Hk}} \\ & \swarrow \omega_\gamma^U & & \searrow \omega & \downarrow \pi & & \downarrow \overrightarrow{h} \\ [^{(2)}K_0 \setminus U] & \xrightarrow{j_\gamma} & \widetilde{\text{Bun}} & & X^0 & \xrightarrow{u_0 \times \text{id}} & \widetilde{\text{Bun}} \times X^0 \end{array} \tag{26}$$

Here ω is the remaining projection corresponding to \overleftarrow{h} on $\widetilde{\text{Hk}}$. Note that $\widetilde{\text{GR}}$ is the analogue of the Beilinson–Drinfeld Grassmannian in this context.¹¹ Let us also denote by

$$\pi_\gamma^U : \widetilde{\text{GR}}_\gamma^U \rightarrow X^0$$

the composite $\pi \circ j_\gamma$. Repeated application of proper base change yields

$$\begin{aligned} (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(j_{\gamma,!} \mathcal{F}_\chi) &= (u_0 \times \text{id})^* \overrightarrow{h}_! (\overleftarrow{h}^* (j_{\gamma,!} \mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{Hk}}}) \\ &\cong \pi_{\gamma,!}^U (\omega_\gamma^{U,*} (\mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}}), \end{aligned} \tag{27}$$

where $\mathcal{K}_{\widetilde{\text{GR}}}$ denotes the pullback of $\mathcal{K}_{\widetilde{\text{Hk}}}$ to $\widetilde{\text{GR}}$. The sheaf $\mathcal{K}_{\widetilde{\text{GR}}}[1]$ is perverse (recall that the fibers of $\mathcal{K}_{\widetilde{\text{GR}}}$ at $x \in X^0$ are copies of \mathcal{K}), and \mathcal{F}_χ is a local system (in cohomological degree zero), so $\omega_\gamma^{U,*} (\mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}}[1]$ is perverse. Our immediate aim is to show that each $(u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(j_{\gamma,!} \mathcal{F}_\chi)[1]$ is a perverse sheaf on X^0 . Any object \mathcal{K} of $\text{Sat}_{\widetilde{\mathcal{G}}}$ is a direct sum of simple objects, so we may assume \mathcal{K} is simple and therefore supported on some $\text{Gr}_{\widetilde{\mathcal{G}}, \leq \lambda}$, $\lambda \in X_*(\widetilde{T})$. The corresponding $\mathcal{K}_{\widetilde{\text{GR}}}$ is then supported on a corresponding substack $\widetilde{\text{GR}}_{\leq \lambda}$, which pulls back in diagram (26) to a substack $\widetilde{\text{GR}}_{\gamma, \leq \lambda}^U$ of $\widetilde{\text{GR}}_\gamma^U$.

¹¹Note that we continue to adhere to the notational pattern of using $\widetilde{(*)}$ to denote the \widetilde{G} -version of an object that could similarly be defined for G . Our notation is, as a result, not always consistent with that of [[Yun 2014a](#)]: for instance, $\widetilde{\text{GR}}^U$ denotes there (the version for G of) what we will call \mathfrak{G}^U below (see diagram (36)).

We now come to the crucial geometric lemma. We note that Yun has found [2014b, Lemma 4.4.7] an argument that applies much more generally; the following, an elaboration of [Yun 2014a, Lemma 4.8] will suffice for us.

Lemma 4.8. *For all $\gamma \in \Omega_1$, the map $\pi_\gamma^U : \widetilde{\text{Gr}}_{\gamma, \leq \lambda}^U \rightarrow X^0$ is affine.*

Proof. Since $[(^{(2)}K_0 \setminus U) \subset [^{(2)}K_0 \setminus \text{fl}_G]$ is affine, we may replace $\widetilde{\text{Gr}}_{\gamma, \leq \lambda}^U$ with the preimage of

$$[^{(2)}K_0 \setminus \text{fl}_G] \xrightarrow{j_\gamma} \widetilde{\text{Bun}}^\gamma.$$

Let us call this preimage $\widetilde{\text{Gr}}_{\gamma, \leq \lambda}^{\text{fl}}$. By construction as the preimage of $\mathbb{B}(K_0) \subset \text{Bun}_{\widetilde{G}}(\widetilde{P}_0(1), \widetilde{P}_\infty)$ (under the morphism (28) below), and using Lemma 3.1 of [Yun 2014a], $\widetilde{\text{Gr}}_{\gamma, \leq \lambda}^{\text{fl}}$ (resp. $\widetilde{\text{Gr}}_{\gamma, \leq \lambda}^{\text{fl}}$) is the nonvanishing locus of a nonzero section s of a line bundle \mathcal{L} on $\widetilde{\text{Gr}}_\gamma$ (resp. $\widetilde{\text{Gr}}_{\gamma, \leq \lambda}$). It suffices to show the line bundle in question is ample. By [Lazarsfeld 2004, Proposition 1.7.8], this can be checked on geometric fibers, since the morphism $\widetilde{\text{Gr}}_{\gamma, \leq \lambda} \rightarrow X^0$ is proper. Thus, let $x : \text{Spec } K \rightarrow X^0$ be a geometric point of X^0 , and consider the section x^*s of $x^*\mathcal{L}$. The fiber $\widetilde{\text{Gr}}_{\gamma, \leq \lambda, x}$ is isomorphic to the γ component, truncated by λ of the affine Grassmannian $\text{Gr}_{\widetilde{G}}$; we denote this by $\text{Gr}_{\widetilde{G}, \leq \lambda}^\gamma$. We claim that $x^*\mathcal{L}$ is ample on $\text{Gr}_{\widetilde{G}, \leq \lambda}^\gamma$, so in particular its restriction to the closed subscheme $\text{Gr}_{\widetilde{G}, \leq \lambda}^\gamma$ is ample. This claim results from the following two assertions:

- $\text{Pic}(\text{Gr}_{\widetilde{G}}^\gamma) \cong \mathbb{Z}$;
- x^*s is a nonzero global section of $x^*\mathcal{L}$ (which by the previous item must then be ample).

The first item follows from [Faltings 2003, Corollary 12]. That result shows that $\text{Pic}(\text{Gr}_G) \cong \mathbb{Z}$ (for G our simply connected group), but the same then follows for each connected component of $\text{Gr}_{\widetilde{G}}$. To be absolutely precise: consider, along with the affine Grassmannian, the affine flag variety $\text{Fl}_{\widetilde{G}} = L\widetilde{G}/\widetilde{I}$, where \widetilde{I} denotes the Iwahori. The connected components $\text{Fl}_{\widetilde{G}}^0$ and $\text{Gr}_{\widetilde{G}}^0$ are, up to taking reduced subschemes, isomorphic to their semisimple counterparts Fl_G and Gr_G (see, e.g., [Pappas and Rapoport 2008, Proposition 6.6]). As in Section 4A, the different components of $\text{Fl}_{\widetilde{G}}$ are isomorphic via ramified Hecke operators $\text{Fl}_{\widetilde{G}}^0 \xrightarrow{\mathbb{T}_\gamma} \text{Fl}_{\widetilde{G}}^\gamma$. In addition, $\text{Pic}(\text{Gr}_{\widetilde{G}}^0)$ is isomorphic to the subgroup of $\text{Pic}(\text{Fl}_{\widetilde{G}}^0)$ corresponding to the unique minimal parahoric P properly containing \widetilde{I} but not contained in $L^+\widetilde{G}$ (see the proof of [Faltings 2003, Corollary 12]); by the same argument, $\text{Pic}(\text{Gr}_{\widetilde{G}}^\gamma)$ can be described inside of $\text{Pic}(\text{Fl}_{\widetilde{G}}^\gamma)$ as the subspace spanned by the natural $\mathcal{O}(1)$ on $\mathbb{T}_\gamma(P)/\widetilde{I}$. For the second item, recall that the pair (\mathcal{L}, s) is the pullback along the composite

$$\widetilde{\text{Gr}}_\gamma \rightarrow \widetilde{\text{Bun}}^\gamma \xrightarrow{\mathbb{T}_\gamma^{-1}} \widetilde{\text{Bun}}^0 \rightarrow \text{Bun}_{\widetilde{G}}(\widetilde{P}_0(1), \widetilde{P}_\infty)^0 \xleftarrow{\sim} \text{Bun}_G(P_0, P_\infty), \quad (28)$$

where the original section is nonvanishing on the locus $\mathbb{B}K_0 \subset \text{Bun}_G(\mathbf{P}_0, \mathbf{P}_\infty)$ corresponding to the tautological object. It suffices then to show that the geometric fibers of $\widetilde{\text{GR}}_\gamma$ over $\mathbb{B}K_0 \times X^0$ are nonempty. To see this, note that $\widetilde{\text{Hk}} \rightarrow \widetilde{\text{Bun}} \times X^0$ has geometric fibers isomorphic to $\text{Gr}_{\widetilde{G}}$. Choosing an element \mathcal{P} of the fiber over (\mathcal{P}_{u_0}, x) that lies in the γ component of $\text{Gr}_{\widetilde{G}}$, we are done: the isomorphism $\iota : \mathcal{P}|_{X-\{x\}} \xrightarrow{\sim} \mathcal{P}_{u_0}|_{X-\{x\}}$ automatically implies that \mathcal{P} projects to an object isomorphic to the tautological object of $\text{Bun}_G(\mathbf{P}_0, \mathbf{P}_\infty)$. \square

With Lemma 4.8 in hand, we can prove the main result of this section:

Theorem 4.9. *For all odd characters $\chi : Z^{(2)\text{A}} \rightarrow \overline{\mathbb{Q}}_\ell^\times$, A_χ is a Hecke eigensheaf.*

Proof. Since $\omega_\gamma^{U,*}(\mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}}[1]$ is perverse, and π_γ^U is affine,

$$(u_0 \times \text{id})^* \mathbb{T}_{\mathcal{K}}(j_{\gamma,!} \mathcal{F}_\chi) \cong \pi_{\gamma,!}^U (\omega_\gamma^{U,*}(\mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}}) \in {}^p D^{\geq 1}(X^0).$$

But by Corollary 4.5, this is also

$$(u_0 \times \text{id})^* \mathbb{T}_{\mathcal{K}}(j_{\gamma,*} \mathcal{F}_\chi) \cong \pi_!(\omega^* j_{\gamma,*}(\mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}}). \tag{29}$$

There is a natural isomorphism $\omega^* \circ j_{\gamma,*} \xrightarrow{\sim} j_{\gamma,*} \circ \omega_\gamma^{U,*}$; as in the proof of [Yun 2014a, Proposition 4.7], this follows from the fact that \overleftarrow{h} is a locally trivial fibration in the smooth topology. Thus, identifying $\pi_! = \pi_*$ on the support of $\mathcal{K}_{\widetilde{\text{GR}}}$ ($\pi : \widetilde{\text{GR}}_{\leq \lambda} \rightarrow X^0$ is proper), and using the projection formula and the Leray spectral sequence, we can carry on the identification (29) as

$$(u_0 \times \text{id})^* \mathbb{T}_{\mathcal{K}}(j_{\gamma,!} \mathcal{F}_\chi) \cong \pi_*((j_{\gamma,*} \omega_\gamma^{U,*} \mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}}) \cong \pi_{\gamma,*}^U (\omega_\gamma^{U,*} \mathcal{F}_\chi \otimes \mathcal{K}_{\widetilde{\text{GR}}}). \tag{30}$$

(This is just the obvious variant of [Yun 2014a, (4.19)].) Since π_γ^U is affine, we can dually conclude that

$$(u_0 \times \text{id})^* \mathbb{T}_{\mathcal{K}}(j_{\gamma,!} \mathcal{F}_\chi) \in {}^p D^{\leq 1}(X^0), \tag{31}$$

hence that $(u_0 \times \text{id})^* \mathbb{T}_{\mathcal{K}}(j_{\gamma,!} \mathcal{F}_\chi)[1]$ is perverse. Consequently, $(u_0 \times \text{id})^* \mathbb{T}_{\mathcal{K}}(A_\chi)[1]$ is perverse.

Now, for each component $\widetilde{\text{Bun}}^\gamma$ of $\widetilde{\text{Bun}}$, we apply [Yun 2014a, Lemma 3.4] to $(u_\gamma \times \text{id})^* \mathbb{T}_{\mathcal{K}}(A_\chi)$ to conclude

$$\begin{aligned} \mathbb{T}_{\mathcal{K}}(A_\chi)|_{\widetilde{\text{Bun}}^\gamma \times X^0} &\cong \bigoplus_{\substack{\psi : Z^{(2)\text{A}} \rightarrow \overline{\mathbb{Q}}_\ell^\times \\ \psi \text{ is odd}}} (j_{\gamma,!} \mathcal{F}_\psi) \boxtimes (V_\psi^* \otimes (u_\gamma \times \text{id})^* \mathbb{T}_{\mathcal{K}}(A_\chi)_\psi)^{(2)\text{A}} \\ &= (j_{\gamma,!} \mathcal{F}_\chi) \boxtimes (V_\chi^* \otimes (u_\gamma \times \text{id})^* \mathbb{T}_{\mathcal{K}}(A_\chi))^{(2)\text{A}}, \end{aligned} \tag{32}$$

where for the second equality we use the fact that the Hecke operators $\mathbb{T}_{\mathcal{K}}$ carries the subcategory $D^b(\widetilde{\text{Bun}})_\psi$ to $D^b(\widetilde{\text{Bun}} \times X^0)_\psi$ for any $\psi : {}^{(2)}Z_G \rightarrow \overline{\mathbb{Q}}_\ell^\times$ (recall from Lemma 4.2 that $Z^{(2)\text{A}}$ is equal to the double cover ${}^{(2)}Z_G \rightarrow Z_G$ of $Z_G = Z_G[2]$).

We have already observed that $(u_\gamma \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi) \cong (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi)$ is independent of γ ; we conclude that

$$\mathbb{T}_\mathcal{K}(A_\chi) \cong A_\chi \boxtimes (V_\chi^* \otimes (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi))^{(2)\mathbb{A}}, \tag{33}$$

and we claim that A_χ is a Hecke eigensheaf with “eigenvalue”

$$\tilde{\mathcal{E}}_\chi : \text{Sat}_{\tilde{G}} \rightarrow \text{Loc}(X^0), \quad \mathcal{K} \mapsto (V_\chi^* \otimes (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi))^{(2)\mathbb{A}}. \tag{34}$$

That is, what remains to show is that $\tilde{\mathcal{E}}_\chi(\mathcal{K})$ is in fact a local system, and that $\tilde{\mathcal{E}}_\chi$ is a tensor functor satisfying the conditions of [Definition 4.7](#). This follows (by the monoidal property of the Hecke operators) by the same argument as [\[Heinloth et al. 2013, §4.2\]](#), since we have seen that $(V_\chi^* \otimes (u_0 \times \text{id})^* \mathbb{T}_\mathcal{K}(A_\chi))^{(2)\mathbb{A}}$ lies in perverse degree one. □

To summarize:

Corollary 4.10. *Assume k is as in [Definition 4.3](#). For every odd character $\chi : Z^{(2)\mathbb{A}}(\bar{k}) \rightarrow \bar{\mathbb{Q}}_\ell^\times$, the object A_χ of $D^b(\widetilde{\text{Bun}})_{\text{odd}}$ given by $A_\chi|_{\widetilde{\text{Bun}}^\gamma} = j_{\gamma,!}(\mathcal{F}_\chi)$ is a Hecke eigensheaf with eigen-local system*

$$\tilde{\mathcal{E}}_\chi : \text{Sat}_{\tilde{G}} \rightarrow \text{Loc}(X^0, \mathbb{Q}'_\ell),$$

giving rise by the Tannakian formalism to a monodromy representation (recall the notation from equation (21))

$$\tilde{\rho}_\chi : \pi_1(X^0) \rightarrow \tilde{G}_1^\vee(\mathbb{Q}'_\ell).$$

The restriction of $\tilde{\mathcal{E}}_\chi$ to the full subcategory $\text{Sat}_G \subset \text{Sat}_{\tilde{G}}$ is naturally isomorphic to the eigen-local system (there denoted \mathcal{E}'_χ) of [\[Yun 2014a, Theorem 4.2\]](#).

Moreover, if $\mathcal{K} = \text{IC}_\chi(m)$ is simple, then $\tilde{\mathcal{E}}_\chi(\mathcal{K})$ is pure of weight $\langle 2\rho, \lambda \rangle - 2m$.

Proof. We have established everything except the purity claim, which follows from the argument of [Theorem 4.9](#). Namely, equations (29) and (30) imply that $\tilde{\mathcal{E}}_\chi(\mathcal{K})$ is mixed of weights \leq and $\geq \langle 2\rho, \lambda, \cdot \rangle - 2m$ (by [\[Deligne 1980\]](#)). □

Consequently, we have a commutative diagram

$$\begin{array}{ccc} & & \tilde{G}_1^\vee(\mathbb{Q}'_\ell) \\ & \nearrow \tilde{\rho}_\chi & \downarrow \\ \pi_1(X^0) & \xrightarrow{\rho_\chi} & G_1^\vee(\mathbb{Q}'_\ell) \end{array}$$


in which ρ_χ (of course, these monodromy representations are only well-defined up to \tilde{G}^\vee or G^\vee conjugation) is Yun’s local system.

5. The motives

Having established the Hecke eigensheaf property, we can now describe the local systems $\tilde{\mathcal{E}}_\chi(\mathcal{K})$ for all $\mathcal{K} \in \text{Sat}_{\tilde{\mathcal{G}}}$. We continue to assume k is as in Definition 4.3; in particular, the k -group scheme $Z^{(2)}(\mathbf{A})$ is discrete. Let us fix a dominant weight $\lambda \in X^\bullet(\tilde{T}^\vee) = X_\bullet(\tilde{T})$, and restrict to the case of $\mathcal{K} = \text{IC}_\lambda$. In this case the sheaf $\mathcal{K}_{\tilde{\text{Hk}}}$ is supported on a substack $\tilde{\text{Hk}}_{\leq \lambda}$, and the sheaf

$$\overleftarrow{h}^*(A_\chi^0) \otimes \mathcal{K}_{\tilde{\text{Hk}}}$$


is supported on the locus of $(\mathcal{P}, \mathcal{P}', x, \iota)$, where $\mathcal{P} \in \widetilde{\text{Bun}}^0$ and \mathcal{P} and \mathcal{P}' are in relative position $\leq \lambda$, i.e., $\text{ev}(\mathcal{P}, \mathcal{P}', \iota, x)$ lies in the $\leq \lambda$ strata of $[(L^+\tilde{G} \backslash L\tilde{G} / L^+\tilde{G}) / \text{Aut}_{\mathcal{O}}]$. This forces \mathcal{P}' to lie in the component $\widetilde{\text{Bun}}^{\nu\lambda}$, where, recall, $\nu : \tilde{G} \rightarrow S$ is the multiplier character. It follows that to compute $\mathbb{T}_{\mathcal{K}}(A_\chi^0)$ we can restrict $\overleftarrow{h} : \tilde{\text{Hk}}_{\leq \lambda} \rightarrow \widetilde{\text{Bun}}$ to the preimage of $\widetilde{\text{Bun}}^0$, and thus consider instead the correspondence diagram



In terms of this diagram, we find that

$$\mathbb{T}_{\mathcal{K}}(A_\chi^0) \simeq A_\chi^{\nu\lambda} \boxtimes \tilde{\mathcal{E}}_\chi(\mathcal{K}). \tag{35}$$

Recall that we are trying to describe $\tilde{\mathcal{E}}_\chi(\mathcal{K})$. The argument is that of [Yun 2014a, Lemma 4.3], except we have to keep track of the different connected components. Pulling back (35) by $(u_{\nu\lambda} \times \text{id})$, we obtain, just as in (26) and (27), a diagram with cartesian squares



and, letting $\pi_{u_{\nu\lambda}}^U$ denote the composite map $\widetilde{\text{GR}}_{u_{\nu\lambda}, \leq \lambda}^U \rightarrow X^0$, we obtain an identification

$$V_\chi \otimes \tilde{\mathcal{E}}_\chi(\mathcal{K}) \cong (u_{\nu\lambda} \times \text{id})^* \mathbb{T}_{\mathcal{K}}(A_\chi^0) \cong \pi_{u_{\nu\lambda}}^U \left(\omega_{u_{\nu\lambda}}^{U,*}(\mathcal{F}_\chi) \otimes \mathcal{K}_{\widetilde{\text{GR}}} \right). \tag{37}$$

(We will write $\mathcal{K}_{\widetilde{\text{GR}}}$ for the pullback of $\mathcal{K}_{\tilde{\text{Hk}}}$ to either of $\widetilde{\text{GR}}_{u_{\nu\lambda}, \leq \lambda}$ or $\widetilde{\text{GR}}_{u_{\nu\lambda}, \leq \lambda}^U$.) Also let

$$\pi_{\tilde{\mathfrak{G}}_{\leq \lambda}}^U : \tilde{\mathfrak{G}}_{\leq \lambda}^U \rightarrow X^0$$

denote the corresponding projection. We now exploit the fact that $\tilde{\mathfrak{S}}_{\leq \lambda}^U$ carries a $({}^{(2)}\mathbf{A} \times ({}^2)\mathbf{A})$ -action; for clarity, the first copy, acting via the pullback on the \overleftarrow{h} (or as here, $\omega_{u_{\nu\circ\lambda}}$) projection, will be denoted $({}^{(2)}\mathbf{A}(1)$, and the second copy, acting via pullback on the \overrightarrow{h} projection, will be denoted $({}^{(2)}\mathbf{A}(2)$. Decomposing the regular representation of $({}^{(2)}\mathbf{A})$, we obtain a $({}^{(2)}\mathbf{A}(1)$ -equivariant isomorphism

$$(\nu_{0,*}\mathbb{Q}'_{\ell})_{\text{odd}} \cong \bigoplus_{\substack{\chi: Z({}^{(2)}\mathbf{A}) \rightarrow \overline{\mathbb{Q}}_{\ell}^{\times} \\ \chi \text{ is odd}}} V_{\chi}^* \otimes \omega_{u_{\nu\circ\lambda}}^{U,*} \mathcal{F}_{\chi}.$$

Here $({}^{(2)}\mathbf{A}(1)$ acts on V_{χ}^* . Since the isomorphism (37) is $({}^{(2)}\mathbf{A}(2)$ -equivariant (acting on V_{χ} on the left-hand side, and on the right-hand side since $\mathcal{K}_{\overline{\text{GR}}}$ is the pullback of $\mathcal{K}_{\overline{\text{HK}}}$), we obtain a $({}^{(2)}\mathbf{A} \times ({}^2)\mathbf{A})$ -equivariant isomorphism

$$\begin{aligned} (\pi_{\tilde{\mathfrak{S}}_{\leq \lambda}^U, !} \nu_0^* \mathcal{K}_{\overline{\text{GR}}})_{\text{odd}} &\cong (\pi_{u_{\nu\circ\lambda}, !}^U \nu_0^* \mathcal{K}_{\overline{\text{GR}}})_{\text{odd}} \\ &\cong \bigoplus_{\substack{\chi: Z({}^{(2)}\mathbf{A}) \rightarrow \overline{\mathbb{Q}}_{\ell}^{\times} \\ \chi \text{ is odd}}} V_{\chi}^* \otimes \pi_{u_{\nu\circ\lambda}, !}^U (\omega_{u_{\nu\circ\lambda}}^{U,*} \mathcal{F}_{\chi} \otimes \mathcal{K}_{\overline{\text{GR}}}) \\ &\cong \bigoplus_{\substack{\chi: Z({}^{(2)}\mathbf{A}) \rightarrow \overline{\mathbb{Q}}_{\ell}^{\times} \\ \chi \text{ is odd}}} V_{\chi}^* \otimes V_{\chi} \otimes \tilde{\mathcal{E}}_{\chi}(\mathcal{K}). \end{aligned} \tag{38}$$

Writing $\mathbb{Q}'_{\ell}[({}^{(2)}\mathbf{A})_{\chi}]$ for the $({}^{(2)}\mathbf{A} \times ({}^2)\mathbf{A})$ -equivariant local system on $\text{Spec } k$ corresponding to the representation $V_{\chi}^* \otimes V_{\chi}$ of the group

$$({}^{(2)}\mathbf{A}(\bar{k}) \times ({}^2)\mathbf{A}(\bar{k})) \rtimes \Gamma_{\bar{k}},^{12}$$

we summarize what we have shown (compare [Yun 2014a, Lemma 4.3]):

Lemma 5.1. *There is a canonical isomorphism of $({}^{(2)}\mathbf{A} \times ({}^2)\mathbf{A})$ -equivariant local systems on X^0*

$$(\pi_{\tilde{\mathfrak{S}}_{\leq \lambda}^U, !} \nu_0^* \mathcal{K}_{\overline{\text{GR}}})_{\text{odd}} \cong \bigoplus_{\substack{\chi: Z({}^{(2)}\mathbf{A}) \rightarrow \overline{\mathbb{Q}}_{\ell}^{\times} \\ \chi \text{ is odd}}} \mathbb{Q}'_{\ell}[({}^{(2)}\mathbf{A})_{\chi}] \otimes \tilde{\mathcal{E}}_{\chi}(\mathcal{K}). \tag{39}$$

In particular, the left-hand side is a local system.

It is explained in [Yun 2014a, §3.3.4] how to take the invariants of an equivariant perverse sheaf under a (not necessarily discrete) finite group scheme. Applying this we have:

¹²Equivalently, regarding $\mathbb{Q}'_{\ell}[({}^{(2)}\mathbf{A}(\bar{k}))]$ as a $({}^{(2)}\mathbf{A}(\bar{k}) \times ({}^2)\mathbf{A}(\bar{k}))$ -module via $(a_1, a_2) \cdot a = a_1 a a_2^{-1}$, and extracting the constituent where $Z({}^{(2)}\mathbf{A}(2))$ acts by χ .

Corollary 5.2. *For all $\mathcal{K} \in \text{Sat}_{\tilde{G}}$ and all odd $\chi : Z({}^{(2)}\mathbf{A}) \rightarrow \overline{\mathbb{Q}}_\ell^\times$, there is an isomorphism of local systems on X^0*

$$\tilde{\mathcal{E}}_\chi(\mathcal{K}) \cong (\mathbb{Q}'_\ell[{}^{(2)}\mathbf{A}]_\chi^* \otimes (\pi_{\tilde{\mathfrak{G}}_{\leq \lambda}, !} \nu_0^* \mathcal{K}_{\text{GR}})_{\text{odd}})^{{}^{(2)}\mathbf{A} \times {}^{(2)}\mathbf{A}}.$$

6. The case of minuscule weights

We now want to make this description of $\tilde{\mathcal{E}}_\chi(\mathcal{K})$ explicit. Our ultimate goal is the following:

Theorem 6.1. *Let k be \mathbb{Q} or $\mathbb{Q}(\sqrt{-1})$ according to whether G is of type D_{4n} , G_2 , E_8 or A_1 , D_{4n+2} , E_7 . Consider any odd $\chi : Z({}^{(2)}\mathbf{A}) \rightarrow \overline{\mathbb{Q}}_\ell^\times$ and any $\mathcal{K} \in \text{Sat}_{\tilde{G}}$. Let F be any number field containing k . Then, for any point $t \in X^0(F)$, the specialization*

$$\tilde{\rho}_{\chi, \mathcal{K}, t} : \pi_1(\text{Spec } F) \xrightarrow{t} \pi_1(X^0) \xrightarrow{\tilde{\rho}_\chi} \tilde{G}_1^\vee(\mathbb{Q}'_\ell) \rightarrow \text{GL}(H_w^*(\mathcal{K}))$$

(where the representation of \tilde{G}_1^\vee is that induced by \mathcal{K} under the Satake isomorphism, as in Section 4B) is, as an Γ_F -representation, isomorphic to the \mathbb{Q}'_ℓ -realization of an object of $\mathcal{M}_{F, \mathbb{Q}'}$.

The case of \mathcal{K} corresponding to a quasiminuscule weight is considered in [Yun 2014a, §4.3]. Although our discussion is valid for any \tilde{G} as in Section 3, there are certain cases in which it is uninteresting: for instance, if $G = \text{SL}_2$, we gain nothing by taking $\tilde{G} = \text{SL}_2 \times \mathbb{G}_m$; however, by taking $\tilde{G} = \text{GL}_2$, we gain the representations of $\text{SL}_2 = G_{\text{sc}}^\vee$ (the simply connected cover of G^\vee), and it is these new representations that will be of interest. Just as in the classical setting, the Kuga–Satake abelian variety is found via the spin representation of Spin_{21} , while the motive of the K3 arises from the standard 21-dimensional representation.

To show that $\tilde{\rho}_{\chi, \mathcal{K}, t}$ arises from an object of $\mathcal{M}_{F, \mathbb{Q}'}$ demands a significant digression into understanding intersection cohomology of varieties with arbitrarily bad singularities. A good first approximation to understanding the motivic nature of $\tilde{\rho}_{\chi, t} : \Gamma_F \rightarrow \tilde{G}_1^\vee(\mathbb{Q}'_\ell)$ is to verify this after composition with a single faithful finite-dimensional representation of \tilde{G}_1^\vee (i.e., to show $\tilde{\rho}_{\chi, t}$ is weakly motivic in the sense of Definition 1.2). That is what we will do in this section.

First, we make a robust choice of \tilde{G} , such that \tilde{G}^\vee has representations restricting to each of the minuscule representations of G_{sc}^\vee . For instance, we can take:

- (A_1) $\tilde{G} = \text{GL}_2$.
- (E_7) Let c denote the nontrivial element of $Z_G = \mu_2$. Then take

$$\tilde{G} = (G \times \mathbb{G}_m) / \langle (c, -1) \rangle.$$

- (D_n , n even) Let c and z be generators of $Z_G \cong \mu_2 \times \mu_2$. Then take

$$\tilde{G} = (G \times \mathbb{G}_m \times \mathbb{G}_m) / \langle (c, -1, 1), (z, 1, -1) \rangle.$$

Each minuscule representation of $G_{\text{sc}}^{\vee 13}$ extends to an irreducible representation of \tilde{G}^{\vee} , and then to an irreducible representation of \tilde{G}_1^{\vee} . Taking a direct sum, we obtain a faithful family of representations

$$r_{\min} : \tilde{G}_1^{\vee} \rightarrow \text{GL}(V_{\min}), \tag{40}$$

and set ourselves the goal of showing that each $r_{\min} \circ \tilde{\rho}_{\chi,t}$ is motivated. The full force of [Theorem 6.1](#) is considerably deeper (it is new even in Yun’s original setting), so in the present section we will only treat the case of these minuscule weights, which has the added advantage that the relevant geometry — of the corresponding affine Schubert varieties — is especially simple.

We begin, however, with some generalities: continue to let $\mathcal{K} \in \text{Sat}_{\tilde{G}}$ be any irreducible object of the form $\mathcal{K} = \text{IC}_{\lambda}$, $\lambda \in X^*(\tilde{T}^{\vee})$ (the discussion will apply equally well to \mathcal{K} of the form $\text{IC}_{\lambda}(m)$, but we take $m = 0$ to simplify the notation). What we denoted above by $\mathcal{K}_{\tilde{\text{GR}}[1]}$ is the intersection complex of $\tilde{\text{GR}}_{u_{v\circ\lambda}, \leq \lambda}^U$ (or the same before restricting to U). Since the map

$$v_0 : \tilde{\mathfrak{S}}_{\leq \lambda}^U \rightarrow \tilde{\text{GR}}_{u_{v\circ\lambda}, \leq \lambda}$$

is étale, $v_0^* \mathcal{K}_{\tilde{\text{GR}}[1]}$ is again the intersection complex of $\tilde{\mathfrak{S}}_{\leq \lambda}^U$. Recall that the stratification of the affine Grassmannian induces one for the Beilinson–Drinfeld Grassmannian:

$$\tilde{\text{GR}}_{u_{v\circ\lambda}, \leq \lambda} = \coprod_{\substack{\mu \leq \lambda \\ \mu \text{ dominant}}} \tilde{\text{GR}}_{u_{v\circ\lambda}, \mu}.$$

The terms on the right-hand side are defined by replacing $\tilde{\text{Hk}}_{\leq \lambda}$ by $\tilde{\text{Hk}}_{\mu}$ in [\(36\)](#). Note that $v \circ \mu = v \circ \lambda$ since $\lambda - \mu \in X_*(T)$ lies in the coroot lattice of G . The dense open locus $\tilde{\text{GR}}_{u_{v\circ\lambda}, \lambda}$ is smooth over X^0 : fiberwise it is the smooth stratum $\text{Gr}_{\tilde{G}, \lambda}$ of $\text{Gr}_{\tilde{G}, \leq \lambda}$. We write $\tilde{\mathfrak{S}}_{\lambda}^U$ and $\tilde{\mathfrak{S}}_{< \lambda}^U$ for the preimages in $\tilde{\mathfrak{S}}_{\leq \lambda}^U$ of $\tilde{\text{GR}}_{u_{v\circ\lambda}, \lambda}$ and $\coprod_{\mu < \lambda} \tilde{\text{GR}}_{u_{v\circ\lambda}, \mu}$.

Taking the t -fiber ($t \in X^0(F)$) of the isomorphism in [Lemma 5.1](#), we obtain a (quasi-)isomorphism

$$\text{IH}_c(\tilde{\mathfrak{S}}_{\leq \lambda, t}^U)_{\text{odd}} \cong \bigoplus_{\chi \text{ odd}} \mathbb{Q}'_{\ell}[(2)\mathbf{A}]_{\chi} \otimes \tilde{\rho}_{\chi, \mathcal{K}, t}. \tag{41}$$

Let us explain the notation. For any irreducible variety Y over a field F , the intersection complex IC_Y is a perverse sheaf in cohomological degrees $[-\dim Y, 0]$. It is pure of weight $\dim Y$. We denote by $\text{IH}_c(Y)$ the complex $R\Gamma_c(\text{IC}_Y)$ on $\text{Spec } F$; it lies in cohomological degrees $[-\dim Y, \dim Y]$, and is pure of weights $\leq \dim Y$. As usual, we then define the compactly supported intersection cohomology $\text{IH}_c^i(Y_{\bar{F}})$

¹³These are, in the three cases: the standard representation of SL_2 , the 56-dimensional representation of E_7 , and the standard ($2n$ -dimensional) and two half-spin representations of Spin_{2n} .

(a Γ_F -representation) as $H^{i-\dim Y}(\mathrm{IH}_c(Y))$ (note the degree shift). We also observe that while (compactly supported) intersection cohomology is not in general functorial for (proper) morphisms of varieties, it is for (proper) étale morphisms: $\mathrm{IC}_{\tilde{\mathfrak{G}}_{\leq \lambda, t}^U}$ is still $({}^{(2)}\mathbb{A} \times {}^{(2)}\mathbb{A})$ -equivariant, as is the isomorphism (41). Now as in [Yun 2014a, §4.3.2], we let $e_\chi \in \mathbb{Q}'[{}^{(2)}\mathbb{A}(\bar{k}) \times {}^{(2)}\mathbb{A}(\bar{k})]^{\Gamma_k}$ be the idempotent whose action on the $({}^{(2)}\mathbb{A}(\bar{k}) \times {}^{(2)}\mathbb{A}(\bar{k}))$ -module $\mathbb{Q}'[{}^{(2)}\mathbb{A}(\bar{k})]$ projects to the component $\mathbb{Q}'[{}^{(2)}\mathbb{A}]_\chi$ and then onto the line spanned by $\mathrm{id} \in \mathrm{End}(V_\chi)$ (a direct factor of the representation $\mathbb{Q}'[{}^{(2)}\mathbb{A}]_\chi$ after restricting to the diagonal copy $({}^{(2)}\mathbb{A}(\bar{k}) \hookrightarrow {}^{(2)}\mathbb{A}(\bar{k}) \times {}^{(2)}\mathbb{A}(\bar{k}))$). Explicitly,

$$e_\chi = \frac{1}{|{}^{(2)}\mathbb{A}(\bar{k}) \times {}^{(2)}\mathbb{A}(\bar{k})|} \sum_{(a_1, a_2)} \theta_\chi(a_1 a_2^{-1})(a_1, a_2),$$

where θ_χ denotes the character of the $({}^{(2)}\mathbb{A}(\bar{k}))$ -representation V_χ .

Proposition 6.2. *Let $\mathcal{K} = \mathrm{IC}_\lambda \in \mathrm{Sat}_{\tilde{\mathfrak{G}}}$, let $\chi : Z({}^{(2)}\mathbb{A}) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ be odd, and let $t \in X^0(F)$ for any number field F containing k . Then*

$$\tilde{\rho}_{\chi, \mathcal{K}, t} \cong \mathrm{Gr}_{(2\rho, \lambda)}^W(e_\chi \mathrm{IH}_c^{(2\rho, \lambda)}(\tilde{\mathfrak{G}}_{\leq \lambda, t}^U)).$$

Proof. Apply e_χ to equation (41), noting that the right-hand side is concentrated in degree zero. Since we have seen that $\tilde{\mathcal{E}}_\chi(\mathrm{IC}_\lambda)$ is pure of weight $(2\rho, \lambda)$, the claim is immediate. \square

Proposition 6.2 reduces Theorem 6.1 to a special case of the following general theorem:

Theorem 6.3. *Let k be a finitely generated field of characteristic zero, and let Y/k be a quasiprojective variety. Then, for all $i, r \in \mathbb{Z}$, $\mathrm{Gr}_i^W(\mathrm{IH}_c^r(Y))$ is as a Γ_k -representation isomorphic to the ℓ -adic realization of an object of \mathcal{M}_k .*

Next consider the case in which Y is acted on by a finite k -group scheme Γ . Let $e \in \bar{\mathbb{Q}}[\Gamma(\bar{k})]^{\Gamma_k}$ be an idempotent. Fix an embedding $\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_\ell$. Then, for all $i, r \in \mathbb{Z}$, $\mathrm{Gr}_i^W(e \mathrm{IH}_c^r(Y, \bar{\mathbb{Q}}_\ell))$ is as a Γ_k -representation isomorphic to the $(\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_\ell)$ -realization of an object of $\mathcal{M}_{k, \bar{\mathbb{Q}}}$.

Remark 6.4. See Section 7A for what is meant by the weight gradings Gr_\bullet^W . Note that in the application we only need the case $i = r$.

Theorem 6.3 will be proven in Corollary 8.15. For the remainder of this section, we content ourselves with showing that $\tilde{\rho}_{\chi, t}$ is weakly motivic. Thus, it suffices to assume that λ restricts to a minuscule weight of G_{sc}^\vee . In this case, $\mathrm{Gr}_{\tilde{\mathfrak{G}}, \leq \lambda} = \mathrm{Gr}_{\tilde{\mathfrak{G}}, \lambda}$ has nonsingular reduced part, so that

$$\tilde{\rho}_{\chi, \mathrm{IC}_\lambda, t} \cong \mathrm{Gr}_{(2\rho, \lambda)}^W(e_\chi H_c^{(2\rho, \lambda)}(\tilde{\mathfrak{G}}_{\lambda, t}^U)).$$

That the right-hand side is isomorphic to the ℓ -adic realization of an object of $\mathcal{M}_{F, \mathbb{Q}'}$ follows from the standard description (originating in [Deligne 1971a]) of

the weight filtration on the cohomology of a smooth variety, via the Leray spectral sequence for its inclusion into a smooth compactification with boundary given by a smooth normal crossings divisor. See [Yun 2014a, §4.3.1] or [Patrikis and Taylor 2015, discussion between Remark 2.6 and Lemma 2.7] for this equivariant version. We conclude:

Corollary 6.5. *For all choices of \tilde{G} as in Section 3, there exists a faithful finite-dimensional representation $r : \tilde{G}_1^\vee \hookrightarrow \mathrm{GL}(V_r)$ such that, for all number field specializations $\mathrm{Spec} F \xrightarrow{L} X^0$ with F satisfying condition (15),*

$$r \circ \tilde{\rho}_{\chi,t} : \Gamma_F \rightarrow \mathrm{GL}(V_r \otimes \mathbb{Q}'_\ell)$$

is isomorphic to the \mathbb{Q}'_ℓ -realization of an object of $\mathcal{M}_{F,\mathbb{Q}'}$. For all G , we may choose \tilde{G} and r such that $r|_{G_{\mathrm{sc}}^\vee}$ is isomorphic to the direct sum of all the minuscule representations of G_{sc}^\vee .

In particular, the lifts $\tilde{\rho}_{\chi,t}$ of Yun’s $\rho_{\chi,t}$ satisfy the generalized Kuga–Satake property of Definition 1.2.

7. Intersection cohomology is motivated

7A. Overview. In the remaining sections, which are logically independent of the rest of the paper, we prove Theorem 6.3. Let k be a field of characteristic zero, and fix an algebraic closure \bar{k} of k . As usual, let $\Gamma_k = \mathrm{Gal}(\bar{k}/k)$. Let Y/k be any quasiprojective variety. If Y is irreducible of dimension d_Y , we can form the ℓ -adic intersection cohomology groups

$$\mathrm{IH}^{r+d_Y}(Y) = H^r(Y_{\bar{k}}, \mathrm{IC}_Y|_{Y_{\bar{k}}}),$$

as well as their analogues with compact supports, $\mathrm{IH}_c^{r+d_Y}(Y)$. If Y is reducible, the definitions need a little more care, working component by component; see [de Cataldo 2012, §4.6] for an explanation. The intersection complex $\mathrm{IC}_{Y_{\bar{k}}}$ is Γ_k -equivariant, so Γ_k acts on $\mathrm{IH}^*(Y)$ and $\mathrm{IH}_c^*(Y)$. Since we do not assume Y is projective, these Γ_k -representations are not pure; in particular, Theorem 6.3 cannot hold for the groups $\mathrm{IH}_c^*(Y)$ themselves. Thus we first need to make sense of the weight filtration on $\mathrm{IH}_c^*(Y)$, in order even to speak of the Γ_k -representations $\mathrm{Gr}_\bullet^W \mathrm{IH}_c^*(Y)$.

There are two basic templates, one “sheaf-theoretic” and one “geometric”, for endowing the cohomology of a variety with a weight filtration. The models for the former approach are [Deligne 1980; Beilinson et al. 1982]; the models for the latter are [Deligne 1971b; 1974]. The latter approach typically depends on having resolution of singularities over the field k , and is consequently restricted to characteristic zero; but when available, it yields more robust, because more “motivic”, results. Thus we will explain, at least for k finitely generated over \mathbb{Q} , how to give

an *a priori* “sheaf-theoretic” sense to $\mathrm{Gr}_\bullet^W \mathrm{IH}^*(Y)$, but then our main aim will be to give a “geometric” construction, as part of the proof of [Theorem 6.3](#), that recovers the sheaf-theoretic definition of the Γ_k -representations $\mathrm{Gr}_\bullet^W \mathrm{IH}_c^*(Y)$. Let us begin then by recalling the sheaf-theoretic construction of a weight filtration on $\mathrm{IH}_c^*(Y)$.

Since we work with k of characteristic zero, the basic case of positive characteristic addressed in [[Deligne 1980](#); [Beilinson et al. 1982](#)] is not sufficient. But the results of those papers have been extended in a form suitable for our purposes, and indeed much more generally than we require, in [[Huber 1997](#); [Morel 2012](#)].¹⁴ Namely, the intersection complex IC_Y is a horizontal, pure perverse sheaf in the sense of [[Morel 2012](#), §2], and [[Morel 2012](#), Théorème 3.2, Proposition 6.1] implies that $\mathrm{IH}_c^*(Y)$ (likewise $\mathrm{IH}^*(Y)$) carries a unique weight filtration W_\bullet . In particular, this means that each $\mathrm{Gr}_r^W \mathrm{IH}_c^*(Y)$ is pure of weight r in the following sense: the underlying lisse sheaf on $\mathrm{Spec} k$ arises by base change from a lisse sheaf \mathcal{G} on some smooth subalgebra $A \subset k$, of finite-type over \mathbb{Z} , and with $\mathrm{Frac}(A) = k$; and for all specializations at closed points x of $\mathrm{Spec} A$, $x^* \mathcal{G}$ is pure of weight r in the usual finite field sense. This characterizing property will hold for the output of our geometric construction; this is verified step-by-step as the construction proceeds.

We now outline the approach to [Theorem 6.3](#). By Poincaré duality for intersection cohomology (which is Γ_k -equivariant), we may restrict to the case of $\mathrm{IH}^*(Y)$. First, we remark that the basic difficulty, and interest, of this problem is that both intersection cohomology and weight filtrations are *a priori* “sheaf-theoretically” defined. The theorem shows that these sheafy constructions can in fact be realized just by playing with the cohomology of smooth projective varieties. There are two, essentially orthogonal, special cases of this problem:

- Y may be smooth but nonprojective. In this case, $\mathrm{IH}^r(Y) = H^r(Y_{\bar{k}}, \mathbb{Q}_\ell)$, and the result follows from the geometric approach of [[Deligne 1971b](#)]; namely, if \bar{Y} is a smooth compactification of Y with $\bar{Y} \setminus Y$ equal to a union of smooth divisors D_α with normal crossings, then the (E_3 -degenerate) Leray spectral sequence for the inclusion $Y \subset \bar{Y}$ yields a description of $\mathrm{Gr}_\bullet^W H^r(Y_{\bar{k}}, \mathbb{Q}_\ell)$ in terms of the divisors D_α and their various (smooth, projective) intersections; see [Theorem 7.2\(3\)](#) below, for a slight rephrasing.
- Y may be projective but singular. In this case, the result, when k is algebraically closed,¹⁵ has been proven by de Cataldo and Migliorini. We briefly describe the two

¹⁴The basic notions of horizontal sheaf, perverse t-structure on the “derived” category of horizontal sheaves, and weights for horizontal sheaves are developed in, respectively, Sections 1, 2, and 3 of [[Huber 1997](#)]. Morel’s paper builds on these foundations, generalizing the results of [[Huber 1997](#)] to any finitely generated k , and establishing a sort of six operations functoriality for complexes having weight filtrations.

¹⁵In this case one should work not just with ℓ -adic cohomology but also with (compatible) Betti and de Rham realizations, in order for the assertion to have any content.

crucial geometric inputs (assume for this informal description that k is algebraically closed). Let $f : X \rightarrow Y$ be a resolution of singularities. Roughly speaking, $\mathrm{IH}^*(Y)$ occurs as a “main term” in $H^*(X, \mathbb{Q}_\ell) = H^*(Y, f_*\mathbb{Q}_\ell)$ corresponding (via the decomposition theorem) to the summand of $f_*\mathbb{Q}_\ell$ in perverse degree $\dim X$ and supported along the open dense stratum (the nonsingular locus) Y^0 of Y . The first key result is that the perverse (Leray) filtration on $H^*(Y, f_*\mathbb{Q}_\ell)$ admits [de Cataldo 2012, Theorem 3.3.5] a remarkable geometric description in terms of a suitably generic “flag filtration”. The second is that the factor of $f_*\mathbb{Q}_\ell$ supported along Y^0 can, at least in cohomology, also be extracted “geometrically” — this follows from the novel approach to the decomposition theorem pioneered by de Cataldo and Migliorini in a series of papers (see [de Cataldo and Migliorini 2014, §1.3.3] for a precise statement).

Our task is to fuse these two approaches, and to get everything to work over an arbitrary (not algebraically closed) field k of characteristic zero. The chief obstruction to getting the relevant arguments of [de Cataldo and Migliorini 2014] to work over any k is that the “generic flags” mentioned above would need to be defined k -rationally. This it turns out is not so hard to achieve, using Bertini’s theorem over k and, crucially, the fact that flag varieties are rational, so that any Zariski open set over k necessarily has k -points.

Rather more complicated is integrating the approaches of [Deligne 1971b] and [de Cataldo and Migliorini 2014] in order to prove Theorem 6.3 for any quasiprojective Y . The basic difficulty is that, since motivated motives are only defined in the pure case, the argument (resting on [Deligne 1971b]) in the smooth case is not obviously “functorial in Y ”. Fortunately, it can be upgraded to one that is, using the results of [Guillén and Navarro Aznar 2002] on the existence of “weight complexes” of motivated motives whose cohomology computes $\mathrm{Gr}_\bullet^W H^*(Y)$ for any k -variety Y . We will also use a version for cohomology with compact supports — due independently to Gillet and Soulé [1996] and Guillén and Navarro, it is somewhat simpler, but not suited for describing the perverse Leray filtration as in [de Cataldo 2012], even for cohomology with compact supports. It is crucial, however, that we exploit both theories: the inductive construction of the support decomposition as in [de Cataldo and Migliorini 2014, Proposition 2.2.1] requires having motivated versions both of pullback in H^* and pullback for proper morphisms in H_c^* (note that these two kinds of pullbacks are not related by Poincaré duality; one cannot be formally reduced to the other). Once this setup is in place, however, the arguments of [de Cataldo and Migliorini 2014] go through *mutatis mutandis*. We consequently establish stronger results on finding “motivated” splittings of the perverse Leray filtration, and a motivated support decomposition, closely in parallel to the main results of [de Cataldo and Migliorini 2014] — see Theorem 8.13 and Corollary 8.14, which should be regarded as the main results of this half of the paper.

Notation 7.1. Except where we explicitly allow more general fields, from now on k will be a finitely generated field extension of \mathbb{Q} . Whenever we speak of the weight grading Gr_\bullet^W on various cohomology groups of a variety over k , the grading is unique, and can be shown to exist by [Morel 2012, Théorème 3.2, Proposition 6.1]. As before, \mathcal{M}_k denotes André’s category of motives for motivated cycles over k (with \mathbb{Q} -coefficients). For a smooth projective variety X over k , we write $H(X)$ for the canonical object of \mathcal{M}_k associated to X . Finally, given a map of varieties $f : X \rightarrow Y$, we always mean the derived functors when we write f_* , $f_!$, etc.

7B. Weight-graded motivated motives associated to smooth varieties. Here is the theorem of Guillén and Navarro Aznar, specialized to the precise statement we require:¹⁶

Theorem 7.2 (see Théorème 5.10 of [Guillén and Navarro Aznar 2002]). *Let k be a field of characteristic zero, and let Sch/k denote the category of finite-type separated k -schemes. Then there exists a contravariant functor*

$$h : \text{Sch}/k \rightarrow K^b(\mathcal{M}_k), \tag{42}$$

valued in the homotopy category of bounded complexes in \mathcal{M}_k , such that:

- (1) *If X is a smooth projective k -scheme, then $h(X)$ is naturally isomorphic to the canonical motivated motive $H(X)$ associated to X .*
- (2) *If X is a smooth projective k -scheme, and $D = \bigcup_{\alpha=1}^t D_\alpha$ is a normal crossings divisor equal to the union of smooth divisors D_α , we can form a cubical diagram of smooth projective varieties*

$$S_\bullet(D) \rightarrow X,$$

*where, for every nonempty subset $\Sigma \subset \{1, \dots, t\}$, $S_\Sigma(D)$ is the (smooth) intersection $D_\Sigma = \bigcap_{\alpha \in \Sigma} D_\alpha$, with the obvious inclusion maps $S_\Sigma(D) \rightarrow S_{\Sigma'}(D)$ whenever $\Sigma' \subset \Sigma$. Using the **covariant** functoriality arising from Gysin maps, we can then associate a cubical diagram $h_*(S_\bullet(D) \rightarrow X)$ in \mathcal{M}_k ; to be precise, $h_*(S_\Sigma(D))$ is the object of \mathcal{M}_k*

$$h(D_\Sigma)(\dim D_\Sigma),$$

with the Gysin maps $h_(S_\Sigma(D)) \rightarrow h_*(S_{\Sigma'}(D))$ whenever $\Sigma' \subset \Sigma$. Then $h(X \setminus D)$ is isomorphic to the simple complex associated to this cubical diagram (see the proof for what this means):*

$$h(X \setminus D) \cong s(h_*(S_\bullet(D) \rightarrow X))(-\dim X). \tag{43}$$

¹⁶They prove something stronger, with Chow motives in place of motivated motives.

- (3) In particular, $h(X \setminus D)$ is a complex whose degree- r homology¹⁷ $H_r(h(X \setminus D))$ is an object of \mathcal{M}_k whose ℓ -adic realization is given by (for k finitely generated over \mathbb{Q})

$$H_\ell(H_r(h(X \setminus D))) \cong \bigoplus_q \mathrm{Gr}_{q+r}^W H^q((X \setminus D)_{\bar{k}}, \mathbb{Q}_\ell).$$

Proof. Except for the third assertion, this is all explicitly in [Guillén and Navarro Aznar 2002, Théorème 5.10]. The remaining claim follows from the usual description [Deligne 1971b] of the weight gradeds for $H^*((X \setminus D)_{\bar{k}}, \mathbb{Q}_\ell)$: ignoring for notational convenience the Tate twists, the degree- r term $h(X \setminus D)_r$ (to be precise, after the identification of equation (43)) is $\bigoplus_{|\Sigma|=r} h(D_\Sigma)$, with the boundary map $h(X \setminus D)_r \rightarrow h(X \setminus D)_{r-1}$ given by an alternating sum of Gysin maps. The ℓ -adic realization of this complex can be identified (up to a sign in the boundary maps, at least — see [Guillén and Navarro Aznar 1990, (1.8) Proposition]) with the complex

$$\dots \rightarrow K_r = \bigoplus_q E_1^{-r, q+r} \xrightarrow{\bigoplus_q d_1^{-r, q+r}} K_{r-1} = \bigoplus_q E_1^{-r+1, q+r} \rightarrow \dots$$

built out of the E_1 terms of the (weight) spectral sequence of the filtered complex (bête filtration)

$$E_1^{-r, q+r} = H^q(X_{\bar{k}}, \mathrm{Gr}_r^W j_* \mathbb{Q}_\ell) \implies H^q(X_{\bar{k}}, j_* \mathbb{Q}_\ell) = H^q((X \setminus D)_{\bar{k}}, \mathbb{Q}_\ell).$$

This spectral sequence degenerates at the E_2 page (by the yoga of weights), and its E_2 terms then give the weight gradeds of $H^q((X \setminus D)_{\bar{k}}, \mathbb{Q}_\ell)$; part (3) of the theorem follows. □

This is not a full description of the result of Guillén and Navarro Aznar, but it contains the two points of interest for us: the explicit description of the objects $H_r(h(X \setminus D))$, and in particular their connection with the weight filtration on $H^*((X \setminus D)_{\bar{k}}, \mathbb{Q}_\ell)$; and, crucially, the fact that h is functorial. In particular, for any morphism $\phi: U \rightarrow V$ in Sch/k , we get, for all r , morphisms $H_r(h(V)) \rightarrow H_r(h(U))$ in \mathcal{M}_k .

Here is the compact-supports version:

Theorem 7.3 [Gillet and Soulé 1996, Theorem 2; Guillén and Navarro Aznar 2002, Théorème 5.2]. *Let k be a field of characteristic zero, and let Sch_c/k denote the category of separated finite-type k -schemes with morphisms given by proper maps. Then there exists a contravariant functor*

$$W : \mathrm{Sch}_c/k \rightarrow K^b(\mathcal{M}_k) \tag{44}$$

such that:

¹⁷We use homological conventions here.

- (1) If X is a smooth projective k -scheme, then $W(X)$ is naturally isomorphic to the canonical motivated motive $H(X)$ associated to X .
- (2) If X is a smooth projective k -scheme, and $D = \bigcup_{\alpha=1}^t D_\alpha$ is a normal crossings divisor equal to the union of smooth divisors D_α , then $W(X \setminus D)$ is isomorphic to the simple complex (we now use cohomological conventions and normalize $W(X \setminus D)$ to live in cohomological degrees $[0, t]$)

$$H(X) \rightarrow \bigoplus_{\alpha} H(D_\alpha) \rightarrow \cdots \rightarrow \bigoplus_{|\Sigma|=s} H(D_\Sigma) \rightarrow \cdots \tag{45}$$

with coboundaries given by an alternating sum of restriction maps $H(D_{\Sigma'}) \rightarrow H(D_\Sigma)$ whenever $\Sigma' \subset \Sigma$. (See [Gillet and Soulé 1996, Proposition 3].)

- (3) In particular, $W(X \setminus D)$ is a complex whose degree- s cohomology $H^s(W(X \setminus D))$ is an object of \mathcal{M}_k whose ℓ -adic realization is given by (for k finitely generated over \mathbb{Q})

$$H_\ell(H^s(W(X \setminus D))) \cong \bigoplus_p \mathrm{Gr}_p^W H_c^{p+s}((X \setminus D)_{\bar{k}}, \mathbb{Q}_\ell).$$

In the setting of parts 2 and 3 of Theorems 7.2 and 7.3, let $U = X \setminus D$. Poincaré duality for U descends to a duality relation in \mathcal{M}_k between the cohomologies of the complexes $h(U)$ and $W(U)$. Before stating it, we introduce a little more notation:

Definition 7.4. Let $H_r^q(h(U))$ be the canonical summand of $H_r(h(U))$ in \mathcal{M}_k of weight $q + r$. Let $W^p(U)$ be the canonical complex of weight- p summands of the terms of $W(U)$, and let $H^s(W^p(U))$ be the degree- s cohomology.

Remark 7.5. The object $H_r^q(h(U))$ of \mathcal{M}_k has ℓ -adic realization $\mathrm{Gr}_{q+r}^W H^q(U_{\bar{k}}, \mathbb{Q}_\ell)$. The object $H^s(W^p(U))$ of \mathcal{M}_k has ℓ -adic realization $\mathrm{Gr}_p^W H_c^{p+s}(U_{\bar{k}}, \mathbb{Q}_\ell)$.

Lemma 7.6. Let $U = X \setminus D$ as above, and assume U is equidimensional of dimension d . Then there is a canonical isomorphism in \mathcal{M}_k

$$H_r^q(h(U))^\vee \cong H^r(W^{2d-q-r}(U))(d). \tag{46}$$

Proof. Poincaré duality for each D_Σ induces a perfect duality between $h(U)_s$ and $W(U)^s$ (the degree- s terms of each complex) for all s . The Gysin maps $H(D_{\Sigma'})(d - |\Sigma'|) \rightarrow H(D_\Sigma)(d - |\Sigma|)$ are Poincaré dual to the pullback maps $H(D_\Sigma) \rightarrow H(D_{\Sigma'})$ for all $\Sigma \subset \Sigma'$, and we can deduce perfect dualities (in \mathcal{M}_k)

$$H_s(h(U))^\vee \cong H^s(W(U))(d).$$

The result follows from decomposing these dualities into each of their graded components. □

8. The perverse Leray filtration

8A. Relation to flag filtrations. In this section we recall the beautiful and fundamental result of de Cataldo and Migliorini that describes the perverse Leray filtration for a map of varieties in terms of a certain flag filtration — see [de Cataldo and Migliorini 2010] and, for the specific result we use, Theorem 3.3.5 of [de Cataldo 2012]. These results are worked out over an algebraically closed field of characteristic zero, and so our first aim in this section is to check the analogue for any $k \supset \mathbb{Q}$.

We first recall the Jouanolou trick.

Definition 8.1. Let Y be a variety over k . An *affinement* of Y is a map $\mathcal{Y} \xrightarrow{p} Y$ in Sch/k with \mathcal{Y} an affine k -scheme, such that p is a torsor for some vector bundle on Y .

Proposition 8.2 [Jouanolou 1973, Lemme 1.5]. *Suppose $Y \in \text{Sch}/k$ is quasiprojective. Then an affinement of Y exists.*

Jouanolou’s result in fact holds for arbitrary quasiprojective schemes, but we are only interested in the case of varieties over k .

Now let $f : X \rightarrow Y$ be a morphism of k -varieties. It induces the (increasing) perverse Leray filtration on $H^*(X_{\bar{k}}, \mathbb{Q}_\ell)$ via

$$\begin{aligned} \mathcal{P}_j^f(H^*(X_{\bar{k}}, \mathbb{Q}_\ell)) &= \text{im}(H^*(Y_{\bar{k}}, {}^p\tau_{\leq j} f_* \mathbb{Q}_\ell) \rightarrow H^*(Y_{\bar{k}}, f_* \mathbb{Q}_\ell)) \\ &\subseteq H^*(Y_{\bar{k}}, f_* \mathbb{Q}_\ell) = H^*(X_{\bar{k}}, \mathbb{Q}_\ell). \end{aligned} \tag{47}$$

Here ${}^p\tau_{\leq j}$ denotes perverse truncation.¹⁸ We make the analogous definition of the perverse Leray filtration on $H_c^*(X_{\bar{k}}, \mathbb{Q}_\ell)$, replacing $f_* \mathbb{Q}_\ell$ by $f_! \mathbb{Q}_\ell$ (the only case of interest to us will be when f is proper, so $f_* = f_!$).

Theorem 8.3 [de Cataldo 2012, Theorem 3.3.5]. *Assume $k = \bar{k}$ is an algebraically closed field of characteristic zero. Let $f : X \rightarrow Y$ be a morphism in Sch/k with Y quasiprojective. Let $p : \mathcal{Y} \rightarrow Y$ be an affinement of Y of relative dimension $d(p)$,¹⁹ and choose a closed embedding $\mathcal{Y} \hookrightarrow \mathbb{A}^N$ of \mathcal{Y} into some affine space. Let*

$$\mathbb{A}_\bullet = \{\emptyset = \mathbb{A}_{-N-1} \subset \mathbb{A}_{-N} \subset \cdots \subset \mathbb{A}_0 = \mathbb{A}^N\}$$

¹⁸It is defined for the complex $f_* \mathbb{Q}_\ell$ on Y itself, and we omit the base change to \bar{k} in the notation of these cohomology groups.

¹⁹If Y is not connected, $d(p)$ is a function $\pi_0(Y) \rightarrow \mathbb{Z}$; we can always reduce to the case of connected Y , so do not dwell on this.

be a full flag of affine linear sections of \mathbb{A}^N , and form the cartesian diagram

$$\begin{array}{ccccc}
 & & r_{\bullet} & & \\
 & \curvearrowright & & \searrow & \\
 \mathcal{X}_{\bullet} & \xrightarrow{i_{\bullet}} & \mathcal{X} & \xrightarrow{p} & X \\
 \downarrow & & \downarrow & & \downarrow f \\
 \mathcal{Y}_{\bullet} & \longrightarrow & \mathcal{Y} & \xrightarrow{p} & Y \\
 \downarrow & & \downarrow & & \\
 \mathbb{A}_{\bullet} & \longrightarrow & \mathbb{A}^N & &
 \end{array} \tag{48}$$

We define the associated (increasing) flag filtrations

$$F_j^{\mathbb{A}_{\bullet}} H^*(X_{\bar{k}}, \mathbb{Q}_{\ell}) = \ker(r_{-j}^* : H^*(X_{\bar{k}}, \mathbb{Q}_{\ell}) \rightarrow H^*((\mathcal{X}_{-j})_{\bar{k}}, \mathbb{Q}_{\ell})) \tag{49}$$

and (see Remark 8.4)

$$F_j^{\mathbb{A}_{\bullet}} H_c^*(X_{\bar{k}}, \mathbb{Q}_{\ell}) = \text{im}(r_{!,j} : H_c^*((\mathcal{X}_j)_{\bar{k}}, \mathbb{Q}_{\ell}) \rightarrow H_c^*(X_{\bar{k}}, \mathbb{Q}_{\ell})). \tag{50}$$

Then, for a general flag \mathbb{A}_{\bullet} ,

$$\mathcal{P}_j^f H^q(X_{\bar{k}}, \mathbb{Q}_{\ell}) = F_{1+d(p)-q+j}^{\mathbb{A}_{\bullet}} H^q(X_{\bar{k}}, \mathbb{Q}_{\ell}) \tag{51}$$

and

$$\mathcal{P}_j^f H_c^q(X_{\bar{k}}, \mathbb{Q}_{\ell}) = F_{j-q-d(p)}^{\mathbb{A}_{\bullet}} H_c^q(X_{\bar{k}}, \mathbb{Q}_{\ell}). \tag{52}$$

Remark 8.4. (1) Let us spell out the construction of the maps $r_{!,j}$. There is a canonical identification

$$H_c^k(X_{\bar{k}}, \mathbb{Q}_{\ell}) \cong H_c^k(\mathcal{X}_{\bar{k}}, p^! \mathbb{Q}_{\ell}) \cong H_c^{k+2d(p)}(\mathcal{X}_{\bar{k}}, \mathbb{Q}_{\ell})(d(p)),$$

and then adjunction gives maps

$$H_c^k((\mathcal{X}_{-j})_{\bar{k}}, i_{-j}^! \mathbb{Q}_{\ell}) \rightarrow H_c^k(\mathcal{X}_{\bar{k}}, \mathbb{Q}_{\ell}).$$

As part of the definition of “general position”, we may assume the \mathcal{X}_{-j} are smooth, so by cohomological purity these adjunction maps are identified with (Gysin) maps

$$H_c^{k-2j}((\mathcal{X}_{-j})_{\bar{k}}, \mathbb{Q}_{\ell})(-j) \rightarrow H_c^k(\mathcal{X}_{\bar{k}}, \mathbb{Q}_{\ell}).$$

The “corestriction” maps $r_{!,-j}$ are then given by the composites

$$H_c^{k+2d(p)-2j}((\mathcal{X}_{-j})_{\bar{k}}, \mathbb{Q}_{\ell})(d(p) - j) \rightarrow H_c^{k+2d(p)}(\mathcal{X}_{\bar{k}}, \mathbb{Q}_{\ell})(d(p)) \rightarrow H_c^k(X_{\bar{k}}, \mathbb{Q}_{\ell}).$$

Important for our purposes is that these are precisely the maps Poincaré dual to the pullback maps arising from the maps $\mathcal{X}_{-j} \rightarrow X$.

(2) We need to specify what is meant by a *general* flag; this will be done in [Section 8B](#). What matters for our purposes is that there exists a Zariski open, dense subspace Flag^{gen} inside the variety Flag of full (affine linear) flags in \mathbb{A}^N such that all points $\mathbb{A}_\bullet \in \text{Flag}^{\text{gen}}(k)$ are “general”.

(3) Note that the degree shift between the perverse and flag filtrations depends on the degree (q above) of cohomology. We will ultimately work one degree of cohomology at a time, and all that matters for us is that *some* shift of the flag filtration agrees with the perverse filtration. To extract the exact degree shift $j \mapsto 1 + d(p) - q + j$, use [[de Cataldo 2012](#), Theorem 3.3.5, (3.8), Example 3.1.6, and (3.16)], and similarly for cohomology with compact supports.

We now explain why this result can be refined k -rationally, so that the diagram (48) for which the conclusion of [Theorem 8.3](#) holds can be taken to be a diagram in Sch/k . In the process, we will say more explicitly what is meant by a “general” flag in the case of *proper* $f : X \rightarrow Y$ (this case is somewhat simpler — see [[de Cataldo 2012](#), Remark 3.2.13] — and it is all we need).

8B. Stratifications. To define “general” flags, we need to say something about stratifications. From now on we will consider a proper map $f : X \rightarrow Y$ of varieties over k with Y quasiprojective. For the purposes of [Theorem 8.3](#), we need only find a stratification Σ of Y such that $f_*\mathbb{Q}_\ell$ is Σ -constructible. That is, we require a decomposition $Y = \bigsqcup_{\sigma \in \Sigma} Y_\sigma$ of Y into locally closed, irreducible, nonsingular varieties such that $f_*\mathbb{Q}_\ell|_{Y_\sigma}$ is lisse for all σ . This is easily arranged; note that the strata Y_σ may be irreducible but not geometrically connected. Then we can deduce:

Corollary 8.5. *Let k be a field of characteristic zero, and let $f : X \rightarrow Y$ be a proper morphism in Sch/k with Y quasiprojective. Then there exists a diagram (48) defined over k for which the conclusions (51) and (52) of [Theorem 8.3](#) hold.*

Proof. Choose as before an affinement $p : \mathcal{Y} \rightarrow Y$ and an embedding $\mathcal{Y} \hookrightarrow \mathbb{A}^N$, and let Flag denote the variety over k of full affine linear flags in \mathbb{A}^N . Fix a stratification Σ of Y such that $f_*\mathbb{Q}_\ell$ is Σ -constructible, and pull it back to a stratification $p^{-1}\Sigma$ of \mathcal{Y} . We then consider full flags

$$\{\mathbb{A}_{-N} \subset \cdots \subset \mathbb{A}_{-1} \subset \mathbb{A}^N\}$$

such that \mathbb{A}_{-1} intersects every stratum \mathcal{Y}_σ transversally; and, refining each $\mathbb{A}_{-1} \cap \mathcal{Y}_\sigma$ to the disjoint union of its connected components, \mathbb{A}_{-2} intersects the induced stratification of $\mathcal{Y} \cap \mathbb{A}_{-1}$ transversally; and so on, inductively. By Bertini’s theorem in exactly the form [[Jouanolou 1979](#), Théorème 6.3(2)], applied inductively to each of the (smooth) strata in each $\mathcal{Y} \cap \mathbb{A}_{-i}$, the collection of such flags defines a Zariski open (over k) dense subset $\text{Flag}^{\text{gen}} \subset \text{Flag}$. Since Flag is a rational variety (for instance, by Bruhat decomposition), and k has characteristic zero,

$\text{Flag}^{\text{gen}}(k)$ is nonempty. The corollary then follows by the proof of [de Cataldo 2012, Theorems 3.3.1 and 3.3.5]. \square

So that we can directly invoke the results of [de Cataldo and Migliorini 2014], in what follows we will make further demands on the stratification, as explained in Section 1.3.2 of that paper. For a fixed proper map $f : X \rightarrow Y$, we consider stratifications of X and Y as the disjoint unions of smooth, locally closed, irreducible (over k) subvarieties, such that every stratum of X maps smoothly and surjectively onto a stratum of Y . Organizing the strata of Y by dimension, we write $Y = \bigsqcup_{l=0}^{\dim Y} S_l$, where S_l has pure dimension l . Each S_l is a disjoint union of smooth and irreducible components of dimension l ; these irreducible components need not be geometrically irreducible, but that does not affect our arguments. We then have Zariski open (dense) subsets $U_l = \bigsqcup_{m \geq l} S_m$, and we get associated closed and open immersions $\alpha_l : S_l \hookrightarrow U_l$ (closed) and $\beta_l : U_{l+1} \hookrightarrow U_l$ (open), with $U_l = S_l \sqcup U_{l+1}$. For more background on these stratifications, see [de Cataldo and Migliorini 2005, §3.2].

8C. Motivated perverse Leray filtration. By Corollary 8.5, the perverse Leray filtrations on $H^*(X_{\bar{k}}, \mathbb{Q}_\ell)$ and $H_c^*(X_{\bar{k}}, \mathbb{Q}_\ell)$ have been Γ_k -equivariantly identified with certain flag filtrations, given in terms of maps of k -varieties $X \rightarrow \mathcal{X}_{-j}$. With an eye toward our final application, in which case $f : X \rightarrow Y$ will be a resolution of singularities of Y , we continue to assume f is proper, but also require that X is nonsingular and irreducible,²⁰ and we use Theorem 7.2, Lemma 7.6 and Corollary 8.5 to define the “perverse Leray filtration” on the motivated motives $H_r(h(X))$ and $H^s(W(X))$. Consider a diagram (48) over k for which the conclusion of Theorem 8.3 holds. Since $h : \text{Sch}/k \rightarrow K^b(\mathcal{M}_k)$ is a functor, we obtain a commutative diagram

$$\begin{array}{ccc} h(X) & \xrightarrow{r_{-i}^*} & h(\mathcal{X}_{-i}) \\ & \searrow^{r_{-i-1}^*} & \downarrow \\ & & h(\mathcal{X}_{-i-1}) \end{array}$$

in $K^b(\mathcal{M}_k)$. Recall that since \mathcal{M}_k is canonically weight-graded (it has Künneth projectors), we can apply the composite functor H_r^q given by taking cohomology H_r of this diagram and projecting to the weight- $(q + r)$ component for any $q \in \mathbb{Z}$, obtaining a commutative diagram in \mathcal{M}_k

$$\begin{array}{ccc} H_r^q(h(X)) & \xrightarrow{r_{-i}^*} & H_r^q(h(\mathcal{X}_{-i})) \\ & \searrow^{r_{-i-1}^*} & \downarrow \\ & & H_r^q(h(\mathcal{X}_{-i-1})) \end{array}$$

²⁰The irreducibility assumption is only for convenience in certain intermediate results, in which, for instance, we wish to invoke Poincaré duality without complicating the notation. Eventually, we extend component by component to the reducible case.

Recall that the ℓ -adic realization of $H_r^q(h(X))$ is (by [Theorem 7.2](#)) isomorphic to $\mathrm{Gr}_{q+r}^W H^q(X_{\bar{k}}, \mathbb{Q}_\ell)$; this accounts for the notation.

Definition 8.6. The perverse Leray filtration of $H_r^q(h(X))$ is defined to be

$$\mathcal{P}_j^f H_r^q(h(X)) = \ker \{r_{-(1+d(p)-q+j)}^* : H_r^q(h(X)) \rightarrow H_r^q(h(\mathcal{X}_{-(1+d(p)-q+j)}))\}. \quad (53)$$

The gradeds for the perverse filtration, still objects of \mathcal{M}_k , are then denoted

$$\mathrm{Gr}_j^{\mathcal{P}^f} H_r^q(h(X)) = \mathcal{P}_j^f H_r^q(h(X)) / \mathcal{P}_{j-1}^f H_r^q(h(X)).$$

Remark 8.7. Our indexing convention is somewhat different from that of de Cataldo and Migliorini (compare [\[de Cataldo and Migliorini 2005, Definition 2.2.2\]](#)).

Now, we already have a definition (equation (47)) of \mathcal{P}^f on $H^q(X_{\bar{k}}, \mathbb{Q}_\ell)$ before passing to the weight gradeds; the two versions of \mathcal{P}^f are compatible in the following sense:

Lemma 8.8. *The ℓ -adic realization $H_\ell(\mathcal{P}_j^f H_r^q(h(X)))$ is isomorphic to*

$$\mathrm{Gr}_{q+r}^W \mathcal{P}_j^f H^q(X_{\bar{k}}, \mathbb{Q}_\ell),$$

and likewise with $\mathrm{Gr}_j^{\mathcal{P}^f}$ in place of \mathcal{P}_j^f .

Proof. The ℓ -adic realization functor is exact, and the maps on cohomology induced by the morphisms $\mathcal{X}_\bullet \rightarrow X$ are strict for the associated weight filtrations, so this follows from the choice of \mathcal{X}_\bullet as in [Corollary 8.5](#). \square

We also need a “motivated” description of the perverse Leray filtration in compactly supported cohomology, i.e., a filtration by submotives on each $H^s(W^p(X))$. Taking our cue from [Remark 8.4](#), we formally define a filtration on $H_r^q(h(X))^\vee$ by

$$(H_r^q(h(X)) / \mathcal{P}_j^f H_r^q(h(X)))^\vee \subset H_r^q(h(X))^\vee, \quad (54)$$

and then invoke duality to define:

Definition 8.9. The perverse Leray filtration of $H^s(W^p(X))$ is defined to be

$$\begin{aligned} \mathcal{P}_j^f H^r(W^{2 \dim X - q - r}(X)) &= (H_r^q(h(X)) / \mathcal{P}_{-j+2 \dim X - 1}^f H_r^q(h(X)))^\vee(-\dim X) \\ &\subseteq H_r^q(h(X))^\vee(-\dim X) \xrightarrow{\sim} H^r(W^{2 \dim X - q - r}(X)). \end{aligned} \quad (55)$$

We check that this definition is compatible with the usual one in cohomology:

Lemma 8.10. *The ℓ -adic realization $H_\ell(\mathcal{P}_j^f H^r(W^{2 \dim X - q - r}(X)))$ is canonically isomorphic to*

$$\mathrm{Gr}_{2 \dim X - q - r}^W \mathcal{P}_j^f H_c^{2 \dim X - q}(X_{\bar{k}}, \mathbb{Q}_\ell),$$

and likewise with $\mathrm{Gr}_j^{\mathcal{P}^f}$ in place of \mathcal{P}_j^f .

Proof. By the description (Remark 8.4) of $r_{l,-j}$ as the map Poincaré dual to r_{-j}^* , we see that Poincaré duality for X induces a duality (here F_\bullet denotes the flag filtrations for general flags)

$$(F_j H^q(X_{\bar{k}}, \mathbb{Q}_\ell))^\vee \cong (H_c^{2 \dim X - q}(X_{\bar{k}}, \mathbb{Q}_\ell) / F_{-j} H_c^{2 \dim X - q}(X_{\bar{k}}, \mathbb{Q}_\ell))(\dim X),$$

i.e.,

$$(\mathcal{P}_{-l+2 \dim X - 1}^f H^q(X_{\bar{k}}, \mathbb{Q}_\ell))^\vee \cong (H_c^{2 \dim X - q}(X_{\bar{k}}, \mathbb{Q}_\ell) / \mathcal{P}_l^f H_c^{2 \dim X - q}(X_{\bar{k}}, \mathbb{Q}_\ell))(\dim X).$$


The lemma follows by passing to Gr_\bullet^W . □

By definition, we obtain the following duality in \mathcal{M}_k , a motivated analogue of [de Cataldo and Migliorini 2014, §1.3.3(12)]:

$$\text{Gr}_j^{\mathcal{P}^f} H^r(W^{2 \dim X - q - r}(X)) \times \text{Gr}_{-j+2 \dim X}^{\mathcal{P}^f} H_r^q(h(X)) \rightarrow \mathbb{Q}(-\dim X). \quad (56)$$

We next check a functoriality property of these motivated perverse Leray filtrations.

Lemma 8.11. *Suppose*



is a commutative diagram in Sch/k . Then the pullback maps $H_r^q(h(X)) \rightarrow H_r^q(h(\mathcal{T}))$ induce morphisms (in \mathcal{M}_k)

$$\mathcal{P}_j^f H_r^q(h(X)) \rightarrow \mathcal{P}_j^g H_r^q(h(\mathcal{T})).$$

If g factors as $\mathcal{T} \xrightarrow{\gamma} Z \xrightarrow{\iota} Y$ with ι a closed immersion, then the filtrations $\mathcal{P}_\bullet^\gamma$ and \mathcal{P}_\bullet^g on $H^*(\mathcal{T}_{\bar{k}}, \mathbb{Q}_\ell)$, or on $H_r^q(h(\mathcal{T}))$, coincide.

If $\mathcal{T} \rightarrow X$ is proper, then the proper pullback $H^s(W^p(X)) \rightarrow H^s(W^p(\mathcal{T}))$ also preserves the perverse Leray filtrations (55).

Proof. Since the ℓ -adic realization functor on \mathcal{M}_k is exact, it suffices to check the statement in cohomology. Here it is elementary — see for instance [de Cataldo and Migliorini 2005, Remark 4.2.3]. For the second statement, use the fact that ι_* is exact for the perverse t-structure (so commutes with perverse truncation). □

8D. Motivated support decomposition. Now we proceed as in [de Cataldo and Migliorini 2014, Proposition 2.2.1] to establish a “motivated support decomposition” of the $\text{Gr}_j^{\mathcal{P}^f} H_r^q(h(X))$, corresponding to the support decomposition of the perverse sheaf ${}^p H^j(f_* \mathbb{Q}_\ell)$. We begin by checking that the desired support decomposition exists k -rationally. Continue to let $f : X \rightarrow Y$ be our proper map of quasiprojective

varieties over k with X nonsingular. Let $Y = \bigsqcup_{l=0}^{\dim Y} S_l$ be a stratification for f as in Section 8B, with the collection of closed and open immersions

$$S_l \xrightarrow{\alpha_l} U_l \xleftarrow{\beta_l} U_{l+1}. \tag{57}$$

Lemma 8.12. *In the above setting, there is a canonical isomorphism in $\text{Perv}(Y)$*

$${}^p H^j(f_* \mathbb{Q}_\ell[\dim X]) \xrightarrow{\sim} \bigoplus_{l=0}^{\dim Y} \text{IC}_{\bar{S}_l}(\alpha_l^* H^{-l}({}^p H^j(f_* \mathbb{Q}_\ell[\dim X]))), \tag{58}$$

where the $\alpha_l^* H^{-l}({}^p H^j(f_* \mathbb{Q}_\ell[\dim X]))$ are (geometrically semisimple) local systems on S_l . Replacing α_l by the inclusion $S \xrightarrow{\iota_S} S_l \xrightarrow{\alpha_l} U_l$ of an irreducible (= connected) component S of S_l , we obtain the refined k -rational support decomposition

$${}^p H^j(f_* \mathbb{Q}_\ell[\dim X]) \xrightarrow{\sim} \bigoplus_{l=0}^{\dim Y} \bigoplus_{S \in \pi_0(S_l)} \text{IC}_{\bar{S}}((\alpha_l \circ \iota_S)^* H^{-l}({}^p H^j(f_* \mathbb{Q}_\ell[\dim X]))). \tag{59}$$

Proof. The second claim follows from the first, so we focus on establishing (58). This statement in $\text{Perv}(Y_{\bar{k}})$ is a precise form — see [de Cataldo and Migliorini 2005, Theorem 2.1.1(c)] — of the semisimplicity assertion of the decomposition theorem, so it suffices to check that the map in equation (58) can be defined in $\text{Perv}(Y)$. For notational simplicity, denote the perverse sheaf ${}^p H^j(f_* \mathbb{Q}_\ell[\dim X])$ on Y simply by K . We follow closely the argument of [de Cataldo and Migliorini 2005, Lemma 4.1.3], and, as there, the claim will follow from the following assertion: for all $l = 0, \dots, \dim Y$, there is a canonical isomorphism

$$K|_{U_l} \xrightarrow{\sim} \beta_{l,*}(K|_{U_{l+1}}) \oplus H^{-l}(K|_{U_l})[l].$$

We now explain this isomorphism, which itself follows from the corresponding geometric statement in [de Cataldo and Migliorini 2005, Lemma 4.1.3, §6]. The second projection comes from the truncation triangle

$$\tau_{\leq -l-1} K|_{U_l} \rightarrow \tau_{\leq -l} K|_{U_l} \rightarrow H^{-l} K|_{U_l}[l] \xrightarrow{\pm 1},$$

whose middle term is canonically $K|_{U_l}$, and whose right-hand term is perverse (by the support conditions in the definition of perverse sheaves; see [de Cataldo and Migliorini 2005, §4.1]).

To define the first projection, recall the successive truncation description of intermediate extension as

$$\tau_{\leq -l-1} \beta_{l,*} \beta_l^* K|_{U_l} \cong \beta_{l,*}(K|_{U_{l+1}}).$$

This suggests applying $\text{Hom}(K|_{U_l}, \cdot)$ to the truncation triangle

$$\tau_{\leq -l-1} \beta_{l*} \beta_l^* K|_{U_l} \rightarrow \tau_{\leq -l} \beta_{l*} \beta_l^* K|_{U_l} \rightarrow H^{-l}(\beta_{l*} \beta_l^* K|_{U_l})[l] \xrightarrow{\pm 1} .$$

There is a canonical map

$$K|_{U_l} \xleftarrow{\sim} \tau_{\leq -l} K|_{U_l} \longrightarrow \tau_{\leq -l} \beta_{l*} \beta_l^* K|_{U_l}$$

\curvearrowright
 a

and to construct the projection $K|_{U_l} \rightarrow \beta_{l!*}(K|_{U_{l+1}})$, it suffices to check that the image of a in

$$\text{Hom}(H^{-l}(K|_{U_l}), H^{-l}(\beta_{l*} \beta_l^* K|_{U_l})) \xrightarrow{\sim} \text{Hom}(K|_{U_l}, H^{-l}(\beta_{l*} \beta_l^* K|_{U_l})[l])$$

(recall $\tau_{\leq -l} K|_{U_l} \xrightarrow{\sim} K|_{U_l}$) is zero. But we can check whether a map of constructible sheaves on Y is zero by passing to $Y_{\bar{k}}$, so the geometric assertion [de Cataldo and Migliorini 2005, §6] implies our corresponding arithmetic assertion. \square

We have maneuvered into a position to invoke the argument of [de Cataldo and Migliorini 2014, Proposition 2.2.1] to prove:

Theorem 8.13. *Let $f : X \rightarrow Y$ be a proper map of quasiprojective varieties over k with X nonsingular. Then, for each triple of integers j, q, r , there exists a decomposition in \mathcal{M}_k*

$$\text{Gr}_j^{\mathcal{P}^f} H_r^q(h(X)) \xrightarrow{\sim} \bigoplus_{l=0}^{\dim Y} \bigoplus_{S \in \pi_0(S_l)} \text{Gr}_{j,S}^{\mathcal{P}^f} H_r^q(h(X)) \tag{60}$$

whose ℓ -adic realization is the output of applying $\text{Gr}_{q+r}^W H^q(Y_{\bar{k}}, \bullet)$ to the splitting of equation (59).²¹

The same holds for cohomology with compact supports, i.e., for the motives $\text{Gr}_j^{\mathcal{P}^f} H^s(W^p(X))$.

Proof. When X is projective, this is established in [de Cataldo and Migliorini 2014, Theorem 3.2.2], via the argument of Proposition 2.1.1 of the same paper; the impediment to that argument going through for nonprojective X is dealt with by our systematic use of the motivated motives $H_r^q(h(X))$ and $H^s(W^p(X))$. We do not repeat the proof, but we will remark on the key points. The argument proceeds by induction on $\dim X$; the inductive step is achieved by using [de Cataldo and Migliorini 2014, equations (13) and (14)] to define the summands $\text{Gr}_{j,S}^{\mathcal{P}^f} H_r^q(h(X))$ in terms of already-defined terms for lower-dimensional X . Note that it is essential that we have at our disposal motives corresponding both to cohomology without supports (the $H_r^q(h(X))$) and to cohomology with compact supports (the $H^s(W^p(X))$), with

²¹Rather, the slight relabeling of this splitting that results from replacing $f_* \mathbb{Q}_\ell[\dim X]$ in (59) with $f_* \mathbb{Q}_\ell$.

their respective pullback functorialities (Theorems 7.2 and 7.3) and the duality (Lemma 7.6 and equation (56)) relating them. \square

The corresponding result in [de Cataldo and Migliorini 2014] uses the relative hard Lefschetz theorem to obtain an absolute Hodge splitting (in the case $k = \bar{k}$) of $H^*(X_{\bar{k}}, \mathbb{Q}_\ell)$ corresponding to the full splitting of $f_*\mathbb{Q}_\ell$ given by the decomposition theorem, rather than as here merely the support decomposition in a particular perverse degree; a similar strengthening can be established in our context, which we now briefly sketch, although it is not needed for our primary goal, Theorem 6.3. For X as in the theorem, consider as usual a smooth compactification \bar{X} with $\bar{X} \setminus X = \bigcup_\alpha D_\alpha$ equal to a union of smooth divisors with normal crossings. We may assume \bar{X} is projective, and then take η to be a hyperplane line bundle arising from some projective embedding. The required motivated version of the relative hard Lefschetz theorem is that there are isomorphisms

$$\bigcup \eta^j : \mathrm{Gr}_{-j+\dim X}^{\mathcal{P}^f} H_r^q(h(X)) \xrightarrow{\sim} \mathrm{Gr}_{j+\dim X}^{\mathcal{P}^f} H_r^{q+2j}(h(X))(j). \quad (61)$$

To construct this isomorphism, note first that we can pull η back to any of the intersections $D_\Sigma = \bigcap_{\alpha \in \Sigma} D_\alpha$ and obtain a morphism of complexes $h(X) \rightarrow h(X)(1)$. This comes from the projection formula: writing η_Σ for the pullback of η to D_Σ , we have, for any inclusion $\iota : D_\Sigma \hookrightarrow D_{\Sigma'}$,

$$\iota_*(a \cup \eta_\Sigma) = \iota_*(a) \cup \eta_{\Sigma'},$$

i.e., cup-product with η commutes with the boundary (Gysin) maps of the complex $h(X)$. Passing to cohomology, η induces maps

$$\eta : H_r^q(h(X)) \rightarrow H_r^{q+2}(h(X))(1).$$

The required compatibility

$$\eta : \mathcal{P}_j^f H_r^q(h(X)) \rightarrow \mathcal{P}_{j+2}^f H_r^{q+2}(h(X))(1)$$

with the perverse filtrations follows directly from Definition 8.6. We therefore have constructed the maps appearing in (61); that they are isomorphisms, as are the corresponding maps for each term of the support decomposition, then follows as usual from the corresponding statement in cohomology. The formalism of “hard Lefschetz triples” [de Cataldo and Migliorini 2014, §1.3.4] in the abelian category \mathcal{M}_k allows us to enhance Theorem 8.13 with the following:

Corollary 8.14. *The choice of η gives rise to a distinguished splitting*

$$H_r^q(h(X)) \xrightarrow{\sim} \bigoplus_j \mathrm{Gr}_j^{\mathcal{P}^f} H_r^q(h(X))$$

of the motivated perverse Leray filtration.

Of course, \mathcal{M}_k is semisimple, so we already knew that some splitting exists. The combination of [Theorem 8.13](#) and [Corollary 8.14](#) can be regarded as a “motivated decomposition theorem”. Finally, we reach the motivating application:

Corollary 8.15 (includes [Theorem 6.3](#) above). *Let Y/k be any quasiprojective variety. Then there is an object $M \in \mathcal{M}_k$ whose ℓ -adic realization is isomorphic as a Γ_k -representation to $\mathrm{Gr}_i^W \mathrm{IH}^q(Y_{\bar{k}}, \mathbb{Q}_\ell)$. If A is a finite group scheme over k acting on Y , and $e \in \overline{\mathbb{Q}}[A(\bar{k})]^{\Gamma_k}$, then for any embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_\ell$ there is an object of $\mathcal{M}_{k, \overline{\mathbb{Q}}}$ whose $(\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_\ell)$ -realization is isomorphic as a Γ_k -representation to $\mathrm{Gr}_i^W e(\mathrm{IH}^q(Y_{\bar{k}}, \overline{\mathbb{Q}}_\ell))$.*

The same holds for intersection cohomology with compact supports.

Proof. We can assume Y is irreducible. Let $f : X \rightarrow Y$ be a resolution of singularities; X is then irreducible of dimension $\dim X$. For the motive M having ℓ -adic realization $\mathrm{Gr}_{q+r}^W \mathrm{IH}^q(Y_{\bar{k}}, \mathbb{Q}_\ell)$, we can take

$$M = \mathrm{Gr}_{\dim X, Y^{\mathrm{sm}}}^{\mathcal{P}^f} H_r^q(h(X)),$$

where Y^{sm} denotes the smooth locus of Y . (Compare [\[de Cataldo and Migliorini 2014, Remark 1.4.2\]](#), noting that we have normalized the perverse filtration differently than they do.)

For the equivariant statement, take $f : X \rightarrow Y$ to be an A -equivariant resolution of singularities; for the existence of these resolutions, including our case in which A is not necessarily a discrete group scheme, see, e.g., [\[Kollár 2005, Proposition 9.1\]](#). By [Lemma 8.11](#), each $\gamma \in \Gamma(\bar{k})$ induces an automorphism of $\mathrm{Gr}_{\dim X}^{\mathcal{P}^f} H_r^q(h(X_{\bar{k}}))$. For $e \in \overline{\mathbb{Q}}[A(\bar{k})]^{\Gamma_k}$, we obtain (after extending scalars to $\overline{\mathbb{Q}}$) an endomorphism of $\mathrm{Gr}_{\dim X}^{\mathcal{P}^f} H_r^q(h(X))$. That this endomorphism preserves the canonical submotive ([Theorem 8.13](#))

$$\mathrm{Gr}_{\dim X, Y^{\mathrm{sm}}}^{\mathcal{P}^f} H_r^q(h(X)) \subset \mathrm{Gr}_{\dim X}^{\mathcal{P}^f} H_r^q(h(X))$$

is then verified by checking the corresponding statement for ℓ -adic realizations.

The statement for compact supports follows similarly, or by now invoking Poincaré duality. □

Remark 8.16. (1) The motive underlying $\mathrm{Gr}_i^W \mathrm{IH}^k(Y_{\bar{k}}, \mathbb{Q}_\ell)$ is canonical in the following sense. The only ambiguity in its construction is that we may take a second resolution $f' : X' \rightarrow Y$ before applying the argument of [Corollary 8.15](#). But any two resolutions of singularities can be dominated by a third, and so the functoriality property of [Lemma 8.11](#) implies that by passing through this third resolution we can deduce an isomorphism in \mathcal{M}_k

$$\mathrm{Gr}_{\dim X, Y^{\mathrm{sm}}}^{\mathcal{P}^f} H_r^q(h(X)) \cong \mathrm{Gr}_{\dim X', Y^{\mathrm{sm}}}^{\mathcal{P}^f} H_r^q(h(X')).$$

To be precise, if we take a resolution $f'' : X'' \rightarrow Y$ dominating the resolutions f and f' , we compare the intersection cohomology motives coming from f and f'' , f' and f'' , and thus f and f' . As just noted, [Lemma 8.11](#) provides a canonical (pullback) map

$$\mathrm{Gr}_j^{\mathcal{P}^f} H_r^q(h(X)) \rightarrow \mathrm{Gr}_j^{\mathcal{P}^{f''}} H_r^q(h(X'')),$$

for all j (likewise for f'). We take $j = \dim Y$, and decompose the source and target according to [Theorem 8.13](#); our task is to show that the respective Y^{sm} -summands map isomorphically to one another. But this may be checked after taking (exact) ℓ -adic realizations, where it follows from the fact that f and f'' are isomorphisms over Y^{sm} (hence so is the map $X'' \rightarrow X$), and that IC_Y is the unique summand (in the decomposition theorem) of ${}^p H^0(f_* \mathbb{Q}_\ell[\dim Y])$ and ${}^p H^0(f''_* \mathbb{Q}_\ell[\dim Y])$ supported on Y^{sm} .

(2) In particular, [Corollary 8.15](#) completes the proof of [Theorem 6.1](#).

(3) Let us now take k to be a finite extension of \mathbb{Q}_p , with $\ell = p$. Let Y/k be a projective variety — this way we avoid discussing weight filtrations, and in particular do not have to be concerned that k is not finitely generated over \mathbb{Q} — so that $\mathrm{IH}^q(Y_{\bar{k}}, \mathbb{Q}_p)$ has underlying motive $\mathrm{Gr}_{\dim X, Y^{\mathrm{sm}}}^{\mathcal{P}^f}(H^q(X))$, where $f : X \rightarrow Y$ is any resolution of singularities. By [Remark 8.16](#)(1), we can then canonically define the intersection de Rham cohomology of Y/k to be the de Rham realization (a filtered k -vector space) of the motive $\mathrm{Gr}_{\dim X, Y^{\mathrm{sm}}}^{\mathcal{P}^f}(H^q(X))$, and by general properties of \mathcal{M}_k we obtain a p -adic de Rham comparison isomorphism, compatible with morphisms of motivated motives.

(4) Finally, taking k to be a totally real field, [[Patrikis and Taylor 2015](#), Corollary B] extends from smooth projective varieties over k with Hodge-regular cohomology in some degree to arbitrary projective varieties over k with Hodge-regular intersection cohomology in some degree. Here we use the theorems of Gabber that $\{\mathrm{IH}^q(Y_{\bar{k}}, \mathbb{Q}_\ell)\}_\ell$ forms a weakly compatible system of pure Γ_k -representations. Consequently, these compatible systems (in the regular case) are strongly compatible, and the corresponding L-functions admit meromorphic continuation to the whole complex plane, with the expected functional equation. Is it possible to construct examples of such singular varieties Y ? Note that Yun’s construction in type G_2 and D_{2n} (the latter regarded as SO_{4n-1} -valued) do give families of examples of potentially automorphic motives — this is a special case of the examples arising from Katz’s theory, as discussed in [[Patrikis and Taylor 2015](#), §2]. The lifts of Yun’s examples constructed in [Corollary 4.10](#) are no longer Hodge-Tate regular, so no further examples of potentially automorphic motives result from the constructions of the present paper.

Acknowledgements

This paper could not exist without the beautiful work of Zhiwei Yun [2014a]. I am, moreover, very grateful to Yun for generously sharing his understanding of that paper with me. I am grateful to the anonymous referee for a careful reading and helpful comments. I thank Bhargav Bhatt for answering a question about stratifications, and Mark de Cataldo for his feedback on a draft of this paper. I thank the Institut Mathématique de Jussieu for its hospitality during part of the preparation of this paper.

References

- [André 1996a] Y. André, “On the Shafarevich and Tate conjectures for hyper-Kähler varieties”, *Math. Ann.* **305**:2 (1996), 205–248. [MR](#) [Zbl](#)
- [André 1996b] Y. André, “Pour une théorie inconditionnelle des motifs”, *Inst. Hautes Études Sci. Publ. Math.* **83** (1996), 5–49. [MR](#) [Zbl](#)
- [Arkhipov and Bezrukavnikov 2009] S. Arkhipov and R. Bezrukavnikov, “Perverse sheaves on affine flags and Langlands dual group”, *Israel J. Math.* **170** (2009), 135–183. [MR](#) [Zbl](#)
- [Beilinson et al. 1982] A. A. Beilinson, J. Bernstein, and P. Deligne, “Faisceaux pervers”, pp. 5–171 in *Analysis and topology on singular spaces, I* (Luminy, 1981), Astérisque **100**, Soc. Math. France, Paris, 1982. [MR](#) [Zbl](#)
- [de Cataldo 2012] M. A. A. de Cataldo, “The perverse filtration and the Lefschetz hyperplane theorem, II”, *J. Algebraic Geom.* **21**:2 (2012), 305–345. [MR](#) [Zbl](#)
- [de Cataldo and Migliorini 2005] M. A. A. de Cataldo and L. Migliorini, “The Hodge theory of algebraic maps”, *Ann. Sci. École Norm. Sup. (4)* **38**:5 (2005), 693–750. [MR](#) [Zbl](#)
- [de Cataldo and Migliorini 2010] M. A. A. de Cataldo and L. Migliorini, “The perverse filtration and the Lefschetz hyperplane theorem”, *Ann. of Math. (2)* **171**:3 (2010), 2089–2113. [MR](#) [Zbl](#)
- [de Cataldo and Migliorini 2014] M. A. A. de Cataldo and L. Migliorini, “The projectors of the decomposition theorem are motivic”, preprint, 2014. [arXiv](#)
- [Deligne 1971a] P. Deligne, “Théorie de Hodge, I”, pp. 425–430 in *Actes du Congrès International des Mathématiciens, Tome I* (Nice, 1970), Gauthier-Villars, Paris, 1971. [MR](#) [Zbl](#)
- [Deligne 1971b] P. Deligne, “Théorie de Hodge, II”, *Inst. Hautes Études Sci. Publ. Math.* **40** (1971), 5–57. [MR](#) [Zbl](#)
- [Deligne 1974] P. Deligne, “Théorie de Hodge, III”, *Inst. Hautes Études Sci. Publ. Math.* **44** (1974), 5–77. [MR](#) [Zbl](#)
- [Deligne 1980] P. Deligne, “La conjecture de Weil, II”, *Inst. Hautes Études Sci. Publ. Math.* **52** (1980), 137–252. [MR](#) [Zbl](#)
- [Faltings 2003] G. Faltings, “Algebraic loop groups and moduli spaces of bundles”, *J. Eur. Math. Soc. (JEMS)* **5**:1 (2003), 41–68. [MR](#) [Zbl](#)
- [Frenkel and Gross 2009] E. Frenkel and B. Gross, “A rigid irregular connection on the projective line”, *Ann. of Math. (2)* **170**:3 (2009), 1469–1512. [MR](#) [Zbl](#)
- [Gaitsgory 2007] D. Gaitsgory, “On de Jong’s conjecture”, *Israel J. Math.* **157** (2007), 155–191. [MR](#) [Zbl](#)
- [Gillet and Soulé 1996] H. Gillet and C. Soulé, “Descent, motives and K -theory”, *J. Reine Angew. Math.* **478** (1996), 127–176. [MR](#) [Zbl](#)

- [Guillén and Navarro Aznar 1990] F. Guillén and V. Navarro Aznar, “Sur le théorème local des cycles invariants”, *Duke Math. J.* **61**:1 (1990), 133–155. [MR](#) [Zbl](#)
- [Guillén and Navarro Aznar 2002] F. Guillén and V. Navarro Aznar, “Un critère d’extension des foncteurs définis sur les schémas lisses”, *Publ. Math. Inst. Hautes Études Sci.* **95** (2002), 1–91. [MR](#) [Zbl](#)
- [Heinloth et al. 2013] J. Heinloth, B.-C. Ngô, and Z. Yun, “Kloosterman sheaves for reductive groups”, *Ann. of Math. (2)* **177**:1 (2013), 241–310. [MR](#) [Zbl](#)
- [Huber 1997] A. Huber, “Mixed perverse sheaves for schemes over number fields”, *Compositio Math.* **108**:1 (1997), 107–121. [MR](#) [Zbl](#)
- [Jouanolou 1973] J. P. Jouanolou, “Une suite exacte de Mayer–Vietoris en K -théorie algébrique”, pp. 293–316. *Lecture Notes in Math.*, Vol. 341 in *Algebraic K-theory, I: Higher K-theories* (Battelle Memorial Institute, Seattle, WA, 1972), edited by H. Bass, Springer, Berlin, 1973. [MR](#) [Zbl](#)
- [Jouanolou 1979] J.-P. Jouanolou, *Théorèmes de Bertini et applications*, Université Louis Pasteur, Département de Mathématique, Institut de Recherche Mathématique Avancée, Strasbourg, 1979. [MR](#) [Zbl](#)
- [Katz 1996] N. M. Katz, *Rigid local systems*, *Annals of Mathematics Studies* **139**, Princeton University Press, 1996. [MR](#) [Zbl](#)
- [Kollár 2005] J. Kollár, “Resolution of singularities – Seattle lecture”, preprint, 2005. [arXiv](#)
- [Laszlo and Olsson 2008] Y. Laszlo and M. Olsson, “The six operations for sheaves on Artin stacks, II: Adic coefficients”, *Publ. Math. Inst. Hautes Études Sci.* **107** (2008), 169–210. [MR](#) [Zbl](#)
- [Lazarsfeld 2004] R. Lazarsfeld, *Positivity in algebraic geometry, I*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* **48**, Springer, Berlin, 2004. [MR](#) [Zbl](#)
- [Mirković and Vilonen 2007] I. Mirković and K. Vilonen, “Geometric Langlands duality and representations of algebraic groups over commutative rings”, *Ann. of Math. (2)* **166**:1 (2007), 95–143. [MR](#) [Zbl](#)
- [Morel 2012] S. Morel, “Complexes mixtes sur un schéma de type fini sur \mathbb{Q} ”, preprint, 2012, available at https://web.math.princeton.edu/~smorel/sur_Q.pdf.
- [Pappas and Rapoport 2008] G. Pappas and M. Rapoport, “Twisted loop groups and their affine flag varieties”, *Adv. Math.* **219**:1 (2008), 118–198. [MR](#) [Zbl](#)
- [Patrikis 2014a] S. Patrikis, “Generalized Kuga–Satake theory and rigid local systems, I: the middle convolution”, preprint, 2014. [arXiv](#)
- [Patrikis 2014b] S. Patrikis, “Variations on a theorem of Tate”, preprint, 2014. To appear in *Memoirs of the AMS*. [arXiv](#)
- [Patrikis 2015] S. Patrikis, “On the sign of regular algebraic polarizable automorphic representations”, *Math. Ann.* **362**:1-2 (2015), 147–171. [MR](#) [Zbl](#)
- [Patrikis and Taylor 2015] S. Patrikis and R. Taylor, “Automorphy and irreducibility of some l -adic representations”, *Compos. Math.* **151**:2 (2015), 207–229. [MR](#) [Zbl](#)
- [Serre 1977] J.-P. Serre, “Modular forms of weight one and Galois representations”, pp. 193–268 in *Algebraic number fields: L -functions and Galois properties* (University of Durham, 1975), edited by A. Fröhlich, Academic Press, London, 1977. [MR](#) [Zbl](#)
- [Wintenberger 1995] J.-P. Wintenberger, “Relèvement selon une isogénie de systèmes l -adiques de représentations galoisiennes associés aux motifs”, *Invent. Math.* **120**:2 (1995), 215–240. [MR](#) [Zbl](#)
- [Yun 2011] Z. Yun, “Global Springer theory”, *Adv. Math.* **228**:1 (2011), 266–328. [MR](#) [Zbl](#)

[Yun 2013] Z. Yun, “Rigidity in the Langlands correspondence and applications”, preprint, 2013, available at <http://stanford.edu/~zwyun/Rigid-ICCM.pdf>. To appear in *Sixth International Congress of Chinese Mathematicians*.

[Yun 2014a] Z. Yun, “Motives with exceptional Galois groups and the inverse Galois problem”, *Invent. Math.* **196**:2 (2014), 267–337. [MR](#) [Zbl](#)

[Yun 2014b] Z. Yun, “Rigidity in automorphic representations and local systems”, pp. 73–168 in *Current developments in mathematics 2013* (Harvard University, Cambridge, MA), edited by D. Jerison et al., International Press, Somerville, MA, 2014. [MR](#) [Zbl](#)

Communicated by Hélène Esnault

Received 2015-06-06

Revised 2016-01-06

Accepted 2016-06-11

patrikis@math.utah.edu

*Department of Mathematics, University of Utah,
155 S 1400 E, Salt Lake City, UT 84112, United States*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Susan Montgomery	University of Southern California, USA
Richard E. Borcherds	University of California, Berkeley, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
John H. Coates	University of Cambridge, UK	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Roger Heath-Brown	Oxford University, UK	Ravi Vakil	Stanford University, USA
Craig Huneke	University of Virginia, USA	Michel van den Bergh	Hasselt University, Belgium
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Yuri Manin	Northwestern University, USA	Efim Zelmanov	University of California, San Diego, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2016 is US \$290/year for the electronic version, and \$485/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 10 No. 7 2016

Arithmetic invariant theory and 2-descent for plane quartic curves	1373
JACK A. THORNE	
Furstenberg sets and Furstenberg schemes over finite fields	1415
JORDAN S. ELLENBERG and DANIEL ERMAN	
Local deformation rings for GL_2 and a Breuil–Mézard conjecture when $l \neq p$	1437
JACK SHOTTON	
Generalized Kuga–Satake theory and rigid local systems, II: rigid Hecke eigensheaves	1477
STEFAN PATRIKIS	
Lifting preprojective algebras to orders and categorifying partial flag varieties	1527
LAURENT DEMONET and OSAMU IYAMA	
A fibered power theorem for pairs of log general type	1581
KENNETH ASCHER and AMOS TURCHET	