

Algebra & Number Theory

Volume 10

2016

No. 9

**Cluster algebras and category \mathcal{O}
for representations of Borel subalgebras
of quantum affine algebras**

David Hernandez and Bernard Leclerc



Cluster algebras and category \mathcal{O} for representations of Borel subalgebras of quantum affine algebras

David Hernandez and Bernard Leclerc

Let \mathcal{O} be the category of representations of the Borel subalgebra of a quantum affine algebra introduced by Jimbo and the first author. We show that the Grothendieck ring of a certain monoidal subcategory of \mathcal{O} has the structure of a cluster algebra of infinite rank, with an initial seed consisting of prefundamental representations. In particular, the celebrated Baxter relations for the 6-vertex model get interpreted as Fomin–Zelevinsky mutation relations.

1. Introduction	2015
2. Quantum loop algebra and Borel algebras	2018
3. Representations of Borel algebras	2020
4. Cluster algebras	2024
5. Properties of the category \mathcal{O}^+	2026
6. Proof of the main theorem	2033
7. Conjectures and evidence	2043
References	2050

1. Introduction

Let $U_q(\mathfrak{g})$ be an untwisted quantum affine algebra (we assume throughout this paper that $q \in \mathbb{C}^*$ is not a root of unity). M. Jimbo and Hernandez [2012] introduced a category \mathcal{O} of representations of a Borel subalgebra $U_q(\mathfrak{b})$ of $U_q(\mathfrak{g})$. Finite-dimensional representations of $U_q(\mathfrak{g})$ are objects in this category as well as the infinite-dimensional prefundamental representations of $U_q(\mathfrak{b})$ constructed in

Hernandez is supported in part by the European Research Council under the European Union's Framework Programme H2020 with ERC Grant Agreement number 647353 QAffine.

MSC2010: primary 17B37; secondary 13F60, 17B10, 82B23.

Keywords: quantum affine algebras, cluster algebras, category \mathcal{O} , monoidal categorification, Baxter relations.

[Hernandez and Jimbo 2012]. They are obtained as asymptotic limits of Kirillov–Reshetikhin modules, which form a family of simple finite-dimensional representations of $U_q(\mathfrak{g})$. These prefundamental representations, denoted by $L_{i,a}^+$ and $L_{i,a}^-$, are simple $U_q(\mathfrak{b})$ -modules parametrized by a complex number $a \in \mathbb{C}^*$ and $1 \leq i \leq n$, where n is the rank of the underlying finite-dimensional simple Lie algebra. Such prefundamental representations were first constructed explicitly for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ by Bazhanov, Lukyanov, and Zamolodchikov [Bazhanov et al. 1997], for $\widehat{\mathfrak{sl}}_3$ by Bazhanov, Hibberd, and Khoroshkin [Bazhanov et al. 2002] and for $\widehat{\mathfrak{sl}}_n$ with $i = 1$ by Kojima [2008].

The category \mathcal{O} and the prefundamental representations were used by E. Frenkel and Hernandez [2015] to prove a conjecture of Frenkel–Reshetikhin on the spectra of quantum integrable systems. Let us recall that the partition function Z of a quantum integrable system is crucial to understanding its physical properties. It may be written in terms of the eigenvalues λ_j of the transfer matrix T . Therefore, to compute Z one needs to find the spectrum of T . In his seminal paper, Baxter [1972] tackled this question for the 6-vertex and 8-vertex models. He observed moreover that the eigenvalues λ_j of T have the very remarkable form

$$\lambda_j = A(z) \frac{Q_j(zq^2)}{Q_j(z)} + D(z) \frac{Q_j(zq^{-2})}{Q_j(z)}, \tag{1-1}$$

where q, z are parameters of the model (quantum and spectral), the functions $A(z), D(z)$ are universal (in the sense that they are the same for all eigenvalues), and Q_j is a polynomial. The above relation is now called *Baxter’s relation* (or *Baxter’s TQ relation*). Frenkel and Reshetikhin [1999] conjectured that the spectra of more general quantum integrable systems constructed from a representation V of a quantum affine algebra $U_q(\mathfrak{g})$ have a similar form. (In this framework, the 6-vertex model is the particular case when $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ and V is irreducible of dimension 2.) One of the main steps in the proof of this conjecture, given in [Frenkel and Hernandez 2015], is to interpret the expected generalized Baxter relations as algebraic identities in the Grothendieck ring of the category \mathcal{O} for $U_q(\mathfrak{b})$ (see [Hernandez 2015] for a short overview). For example, if $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ and V is the 2-dimensional simple representation of $U_q(\mathfrak{g})$ with q -character $\chi_q(V) = Y_{1,a} + Y_{1,aq^2}^{-1}$, one gets the following categorical version of Baxter’s relation (1-1):

$$[V \otimes L_{1,aq}^+] = [\omega_1][L_{1,aq^{-1}}^+] + [-\omega_1][L_{1,aq^3}^+]. \tag{1-2}$$

(Here, $[\omega_1]$ and $[-\omega_1]$ denote the classes of certain one-dimensional representations of $U_q(\mathfrak{b})$; see Definition 3.4 below.)

In another direction, the notion of monoidal categorification of cluster algebras was introduced in [Hernandez and Leclerc 2010]. The cluster algebra $\mathcal{A}(Q)$ attached to a quiver Q is a commutative \mathbb{Z} -algebra with a distinguished set of generators called

cluster variables and obtained inductively via the Fomin–Zelevinsky procedure of mutation [2002]. By definition, the rank of $\mathcal{A}(Q)$ is the number of vertices of Q (finite or infinite). A monoidal category \mathcal{C} is said to be a *monoidal categorification* of $\mathcal{A}(Q)$ if there exists a ring isomorphism

$$\mathcal{A}(Q) \xrightarrow{\sim} K_0(\mathcal{C})$$

which induces a bijection between cluster variables and classes of simple modules which are *prime* (without nontrivial tensor factorization) and *real* (with simple tensor square). Various examples of monoidal categorifications have been established in terms of quantum affine algebras [Hernandez and Leclerc 2010; 2016], perverse sheaves on quiver varieties [Nakajima 2011; Kimura and Qin 2014; Qin 2015], and Khovanov–Lauda–Rouquier algebras [Kang et al. 2014; 2015].

In this paper, we propose new monoidal categorifications of cluster algebras in terms of the category \mathcal{O} of a Borel subalgebra $U_q(\mathfrak{b})$ of an untwisted quantum affine algebra $U_q(\mathfrak{g})$. More precisely, in [Hernandez and Leclerc 2016] we attached to \mathfrak{g} a semi-infinite quiver G^- and we proved that the cluster algebra $\mathcal{A}(G^-)$ is isomorphic to the Grothendieck ring of a monoidal category $\mathcal{C}_{\mathbb{Z}}^-$ of finite-dimensional representations of $U_q(\mathfrak{g})$. Moreover, the classes of the Kirillov–Reshetikhin modules in $\mathcal{C}_{\mathbb{Z}}^-$ are the images under this isomorphism of a subset of the cluster variables. Let Γ be the doubly infinite quiver corresponding to G^- , as defined in [Hernandez and Leclerc 2016, Section 2.1.2]. The main result of this paper (Theorem 4.2) is that the completed cluster algebra $\mathcal{A}(\Gamma)$ attached to this doubly infinite quiver is isomorphic to the Grothendieck ring of a certain monoidal subcategory $\mathcal{O}_{2\mathbb{Z}}^+$ of \mathcal{O} . This subcategory $\mathcal{O}_{2\mathbb{Z}}^+$ is generated by finite-dimensional representations and positive prefundamental representations whose spectral parameters satisfy an integrality condition (see Definitions 3.8 and 4.1 and Proposition 5.16). Moreover, the classes of the positive prefundamental representations form the cluster variables of an initial seed of $\mathcal{A}(\Gamma)$. In particular, when $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ the counterparts (1-2) of Baxter’s relations (1-1) get interpreted as instances of Fomin–Zelevinsky mutation relations.¹ For general types, the one-step mutation relations are interpreted as other remarkable relations in the Grothendieck ring $K_0(\mathcal{O}^+)$ (Formula (6-14)).

Along the way we get interesting additional results, for instance, (i) the construction of new asymptotic representations beyond the case of prefundamental representations (Theorem 7.6), and (ii) the tensor factorization of arbitrary simple modules of \mathcal{O}^+ into prime modules when $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ (Theorem 7.9). We conjecture that the category $\mathcal{O}_{2\mathbb{Z}}^+$ is a monoidal categorification of the cluster algebra $\mathcal{A}(\Gamma)$ (Conjecture 7.2). We prove this conjecture for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ (Theorem 7.11). An essential

¹The generalized Baxter relations of [Frenkel and Hernandez 2015] can also be regarded as relations in the cluster algebra, but not as mutation relations since they involve more than 2 terms in general.

tool in several of our proofs is a duality D between the Grothendieck rings of certain subcategories \mathcal{O}^+ and \mathcal{O}^- of \mathcal{O} (Theorem 5.17), which maps classes of simple objects to classes of simple objects (Theorem 7.7).

The paper is organized as follows. In Section 2 we give some background on quantum affine (or loop) algebras and their Borel subalgebras. In Section 3 we review the main properties of the category \mathcal{O} introduced in [Hernandez and Jimbo 2012] and we introduce the subcategories \mathcal{O}^+ and \mathcal{O}^- of interest for this paper (Definition 3.9). In Section 4 we state the main result on the isomorphism between $\mathcal{A}(\Gamma)$ and the Grothendieck ring of $\mathcal{O}_{2\mathbb{Z}}^+$. In Section 5 we establish relevant properties of \mathcal{O}^+ ; in particular, we introduce and study the duality D between \mathcal{O}^+ and \mathcal{O}^- . The proof of Theorem 4.2 is given in Section 6. In Section 7 we present the conjecture on monoidal categorifications and we give various evidence supporting it, in particular the existence of asymptotic representations (Section 7B). To conclude we present additional conjectural relations in $K_0(\mathcal{O}^+)$, extending the generalized Baxter relations of [Frenkel and Hernandez 2015] (Conjecture 7.15).

The main results of this paper were presented in several conferences (the Oberwolfach workshop “Enveloping algebras and geometric representation theory” in May 2015, “Categorical representation theory and combinatorics” in Seoul in December 2015, and “A bridge between representation theory and physics” in Canterbury in January 2016). An announcement was also published in the Oberwolfach Report [Hernandez 2015].

2. Quantum loop algebra and Borel algebras

2A. Quantum loop algebra. Let $C = (C_{i,j})_{0 \leq i,j \leq n}$ be an indecomposable Cartan matrix of nontwisted affine type. We denote by \mathfrak{g} the Kac–Moody Lie algebra associated with C . Set $I = \{1, \dots, n\}$, and denote by $\dot{\mathfrak{g}}$ the finite-dimensional simple Lie algebra associated with the Cartan matrix $(C_{i,j})_{i,j \in I}$. Let $\{\alpha_i\}_{i \in I}$, $\{\alpha_i^\vee\}_{i \in I}$, $\{\omega_i\}_{i \in I}$, $\{\omega_i^\vee\}_{i \in I}$, $\dot{\mathfrak{h}}$ be the simple roots, the simple coroots, the fundamental weights, the fundamental coweights and the Cartan subalgebra of $\dot{\mathfrak{g}}$, respectively. We set

$$Q = \bigoplus_{i \in I} \mathbb{Z}\alpha_i, \quad Q^+ = \bigoplus_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i, \quad P = \bigoplus_{i \in I} \mathbb{Z}\omega_i.$$

We will also use $P_{\mathbb{Q}} = P \otimes \mathbb{Q}$ with its partial ordering defined by $\omega \leq \omega'$ if and only if $\omega' - \omega \in Q^+$. Let $D = \text{diag}(d_0, \dots, d_n)$ be the unique diagonal matrix such that $B = DC$ is symmetric and the d_i are relatively prime positive integers. We denote by $(\cdot, \cdot) : Q \times Q \rightarrow \mathbb{Z}$ the invariant symmetric bilinear form such that $(\alpha_i, \alpha_i) = 2d_i$. We use the numbering of the Dynkin diagram as in [Kac 1990]. Let a_0, \dots, a_n be the Kac labels as on pages 55–56 of the same work. We have $a_0 = 1$ and we set $\alpha_0 = -(a_1\alpha_1 + a_2\alpha_2 + \dots + a_n\alpha_n)$.

Throughout this paper, we fix a nonzero complex number q which is not a root of unity. We set $q_i = q^{d_i}$. We fix once and for all $h \in \mathbb{C}$ such that $q = e^h$, and we define $q^r = e^{rh}$ for any $r \in \mathbb{Q}$. Since q is not a root of unity, for $r, s \in \mathbb{Q}$ we have that $q^r = q^s$ if and only if $r = s$.

We will use the standard symbols for q -integers:

$$[m]_z = \frac{z^m - z^{-m}}{z - z^{-1}}, \quad [m]_z! = \prod_{j=1}^m [j]_z, \quad \begin{bmatrix} s \\ r \end{bmatrix}_z = \frac{[s]_z!}{[r]_z! [s-r]_z!}.$$

The quantum loop algebra $U_q(\mathfrak{g})$ is the \mathbb{C} -algebra defined by generators $e_i, f_i, k_i^{\pm 1}$ ($0 \leq i \leq n$) and the following relations for $0 \leq i, j \leq n$:

$$\begin{aligned} k_i k_j &= k_j k_i, & [e_i, f_j] &= \delta_{i,j} \frac{k_i - k_i^{-1}}{q_i - q_i^{-1}}, \\ k_0^{a_0} k_1^{a_1} \cdots k_n^{a_n} &= 1, & 0 &= \sum_{r=0}^{1-C_{i,j}} (-1)^r e_i^{(1-C_{i,j}-r)} e_j e_i^{(r)} \quad (i \neq j), \\ k_i e_j k_i^{-1} &= q_i^{C_{i,j}} e_j, & 0 &= \sum_{r=0}^{1-C_{i,j}} (-1)^r f_i^{(1-C_{i,j}-r)} f_j f_i^{(r)} \quad (i \neq j), \\ k_i f_j k_i^{-1} &= q_i^{-C_{i,j}} f_j, \end{aligned}$$

Here we have set $x_i^{(r)} = x_i^r / [r]_{q_i}!$ ($x_i = e_i, f_i$). The algebra $U_q(\mathfrak{g})$ has a Hopf algebra structure given by

$$\begin{aligned} \Delta(e_i) &= e_i \otimes 1 + k_i \otimes e_i, & S(e_i) &= -k_i^{-1} e_i, \\ \Delta(f_i) &= f_i \otimes k_i^{-1} + 1 \otimes f_i, & S(f_i) &= -f_i k_i, \\ \Delta(k_i) &= k_i \otimes k_i, & S(k_i) &= k_i^{-1}, \end{aligned}$$

where $i = 0, \dots, n$.

The algebra $U_q(\mathfrak{g})$ can also be presented in terms of the Drinfeld generators [Drinfeld 1987; Beck 1994]

$$x_{i,r}^{\pm} \quad (i \in I, r \in \mathbb{Z}), \quad \phi_{i,\pm m}^{\pm} \quad (i \in I, m \geq 0), \quad k_i^{\pm 1} \quad (i \in I).$$

Example 2.1. For $\mathfrak{g} = \mathfrak{sl}_2$, we have $e_1 = x_{1,0}^+$, $e_0 = k_1^{-1} x_{1,1}^-$, $f_1 = x_{1,0}^-$, $f_0 = x_{1,-1}^+ k_1$.

We shall use the generating series ($i \in I$)

$$\phi_i^{\pm}(z) = \sum_{m \geq 0} \phi_{i,\pm m}^{\pm} z^{\pm m} = k_i^{\pm 1} \exp\left(\pm (q_i - q_i^{-1}) \sum_{m > 0} h_{i,\pm m} z^{\pm m}\right).$$

We also set $\phi_{i,\pm m}^{\pm} = 0$ for $m < 0$, $i \in I$.

The algebra $U_q(\mathfrak{g})$ has a \mathbb{Z} -grading defined by $\deg(e_i) = \deg(f_i) = \deg(k_i^{\pm 1}) = 0$ for $i \in I$ and $\deg(e_0) = -\deg(f_0) = 1$. It satisfies $\deg(x_{i,m}^{\pm}) = \deg(\phi_{i,m}^{\pm}) = m$ for

$i \in I, m \in \mathbb{Z}$. For $a \in \mathbb{C}^*$, there is a corresponding automorphism $\tau_a : U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g})$ such that $\tau_a(x) = a^m x$ for every $x \in U_q(\mathfrak{g})$ of degree $m \in \mathbb{Z}$.

By Proposition 1.6 of [Chari 1995], there exists an involutive automorphism $\hat{\omega} : U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g})$ defined by ($i \in I, m, r \in \mathbb{Z}, r \neq 0$)

$$\hat{\omega}(x_{i,m}^\pm) = -x_{i,-m}^\mp, \quad \hat{\omega}(\phi_{i,\pm m}^\pm) = \phi_{i,\mp m}^\mp, \quad \hat{\omega}(h_{i,r}) = -h_{i,-r}.$$

Let $U_q(\mathfrak{g})^\pm$ (resp. $U_q(\mathfrak{g})^0$) be the subalgebra of $U_q(\mathfrak{g})$ generated by the $x_{i,r}^\pm$, where $i \in I, r \in \mathbb{Z}$ (resp. by the $\phi_{i,\pm r}^\pm$, where $i \in I, r \geq 0$). We have a triangular decomposition [Beck 1994]

$$U_q(\mathfrak{g}) \simeq U_q(\mathfrak{g})^- \otimes U_q(\mathfrak{g})^0 \otimes U_q(\mathfrak{g})^+. \quad (2-3)$$

2B. Borel algebra.

Definition 2.2. The Borel algebra $U_q(\mathfrak{b})$ is the subalgebra of $U_q(\mathfrak{g})$ generated by e_i and $k_i^{\pm 1}$ with $0 \leq i \leq n$.

This is a Hopf subalgebra of $U_q(\mathfrak{g})$. The algebra $U_q(\mathfrak{b})$ contains the Drinfeld generators $x_{i,m}^+, x_{i,r}^-, k_i^{\pm 1}, \phi_{i,r}^+$, where $i \in I, m \geq 0$ and $r > 0$.

Let $U_q(\mathfrak{b})^\pm = U_q(\mathfrak{g})^\pm \cap U_q(\mathfrak{b})$ and $U_q(\mathfrak{b})^0 = U_q(\mathfrak{g})^0 \cap U_q(\mathfrak{b})$. Then we have

$$U_q(\mathfrak{b})^+ = \langle x_{i,m}^+ \rangle_{i \in I, m \geq 0}, \quad U_q(\mathfrak{b})^0 = \langle \phi_{i,r}^+, k_i^{\pm 1} \rangle_{i \in I, r > 0}.$$

It follows from [Beck 1994; Damiani 1998] that we have a triangular decomposition

$$U_q(\mathfrak{b}) \simeq U_q(\mathfrak{b})^- \otimes U_q(\mathfrak{b})^0 \otimes U_q(\mathfrak{b})^+. \quad (2-4)$$

3. Representations of Borel algebras

In this section we review results on representations of the Borel algebra $U_q(\mathfrak{b})$, in particular on the category \mathcal{O} defined in [Hernandez and Jimbo 2012] and on finite-dimensional representations of $U_q(\mathfrak{g})$. We also introduce the subcategories \mathcal{O}^+ and \mathcal{O}^- of particular interest for this paper.

3A. Highest ℓ -weight modules. For a $U_q(\mathfrak{b})$ -module V and $\omega \in P_{\mathbb{Q}}$, we have the weight space

$$V_\omega = \{v \in V \mid k_i v = q_i^{\omega(\alpha_i^\vee)} v \text{ for all } i \in I\}. \quad (3-5)$$

We say that V is *Cartan-diagonalizable* if $V = \bigoplus_{\omega \in P_{\mathbb{Q}}} V_\omega$.

For V a Cartan-diagonalizable $U_q(\mathfrak{b})$ -module, we define the structure of a $U_q(\mathfrak{b})$ -module on its graded dual $V^* = \bigoplus_{\beta \in P_{\mathbb{Q}}} V_\beta^*$ by

$$(xu)(v) = u(S^{-1}(x)v) \quad (u \in V^*, v \in V, x \in U_q(\mathfrak{b})).$$

Definition 3.1. A series $\Psi = (\Psi_{i,m})_{i \in I, m \geq 0}$ of complex numbers such that $\Psi_{i,0} \in q_i^{\mathbb{Q}}$ for all $i \in I$ is called an ℓ -weight.

We denote by P_ℓ the set of ℓ -weights. Identifying $(\Psi_{i,m})_{m \geq 0}$ with its generating series, we shall write

$$\Psi = (\Psi_i(z))_{i \in I}, \quad \Psi_i(z) = \sum_{m \geq 0} \Psi_{i,m} z^m.$$

We will often use the involution $\Psi \mapsto \bar{\Psi}$ on P_ℓ , where $\bar{\Psi}$ is obtained from Ψ by replacing every pole and zero of Ψ by its inverse.

Since each $\Psi_i(z)$ is an invertible formal power series, P_ℓ has a natural group structure by componentwise multiplication. We have a surjective morphism of groups $\varpi : P_\ell \rightarrow P_{\mathbb{Q}}$ given by $\Psi_i(0) = q_i^{\varpi(\Psi)(\alpha_i^\vee)}$. For a $U_q(\mathfrak{b})$ -module V and $\Psi \in P_\ell$, the linear subspace

$$V_\Psi = \{v \in V \mid \text{there exists a } p \geq 0 \text{ such that,} \\ \text{for all } i \in I \text{ and } m \geq 0, (\phi_{i,m}^+ - \Psi_{i,m})^p v = 0\} \quad (3-6)$$

is called the ℓ -weight space of V of ℓ -weight Ψ . Note that since $\phi_{i,0}^+ = k_i$, we have $V_\Psi \subset V_\omega$, where $\omega = \varpi(\Psi)$.

Definition 3.2. A $U_q(\mathfrak{b})$ -module V is said to be of highest ℓ -weight $\Psi \in P_\ell$ if there is $v \in V$ such that $V = U_q(\mathfrak{b})v$ and the following hold:

$$e_i v = 0 \quad (i \in I), \quad \phi_{i,m}^+ v = \Psi_{i,m} v \quad (i \in I, m \geq 0).$$

The ℓ -weight $\Psi \in P_\ell$ is uniquely determined by V . It is called the highest ℓ -weight of V . The vector v is said to be a highest ℓ -weight vector of V . For any $\Psi \in P_\ell$, there exists a simple highest ℓ -weight module $L(\Psi)$ of highest ℓ -weight Ψ . This module is unique up to isomorphism.

For Ψ an ℓ -weight, we set $\tilde{\Psi} = (\varpi(\Psi))^{-1}\Psi$ and we introduce the simple $U_q(\mathfrak{b})$ -module

$$\tilde{L}(\Psi) := L(\tilde{\Psi}).$$

This is the simple $U_q(\mathfrak{b})$ -module obtained from $L(\Psi)$ by shifting all ℓ -weights by $\varpi(\Psi)^{-1}$ (see [Hernandez and Jimbo 2012, Remark 2.5]).

The submodule of $L(\Psi) \otimes L(\Psi')$ generated by the tensor product of the highest ℓ -weight vectors is of highest ℓ -weight $\Psi\Psi'$. In particular, $L(\Psi\Psi')$ is a subquotient of $L(\Psi) \otimes L(\Psi')$.

Definition 3.3 [Hernandez and Jimbo 2012]. For $i \in I$ and $a \in \mathbb{C}^\times$, let

$$L_{i,a}^\pm = L(\Psi_{i,a}^{\pm 1}), \quad \text{where } (\Psi_{i,a}^{\pm 1})_j(z) = \begin{cases} (1 - za)^{\pm 1}, & j = i, \\ 1, & j \neq i. \end{cases} \quad (3-7)$$

We call $L_{i,a}^+$ (resp. $L_{i,a}^-$) a positive (resp. negative) *prefundamental representation*.

Definition 3.4. For $\omega \in P_{\mathbb{Q}}$, let $[\omega] = L(\Psi_\omega)$, where $(\Psi_\omega)_i(z) = q_i^{\omega(\alpha_i^\vee)}$ ($i \in I$).

Note that the representation $[\omega]$ is 1-dimensional with a trivial action of e_0, \dots, e_n .

For $a \in \mathbb{C}^\times$, the subalgebra $U_q(\mathfrak{b})$ is stable by τ_a . Denote its restriction to $U_q(\mathfrak{b})$ by the same letter. Then the pullbacks of the $U_q(\mathfrak{b})$ -modules $L_{i,b}^\pm$ by τ_a is $L_{i,ab}^\pm$.

3B. Category \mathcal{O} . For $\lambda \in P_{\mathbb{Q}}$, we set $D(\lambda) = \{\omega \in P_{\mathbb{Q}} \mid \omega \leq \lambda\}$.

Definition 3.5 [Hernandez and Jimbo 2012]. A $U_q(\mathfrak{b})$ -module V is said to be in category \mathcal{O} if:

- (i) V is Cartan-diagonalizable;
- (ii) for all $\omega \in P_{\mathbb{Q}}$ we have $\dim(V_\omega) < \infty$;
- (iii) there exist a finite number of elements $\lambda_1, \dots, \lambda_s \in P_{\mathbb{Q}}$ such that the weights of V are in $\bigcup_{j=1, \dots, s} D(\lambda_j)$.

The category \mathcal{O} is a monoidal category.

Remark 3.6. The definition of \mathcal{O} is slightly different from that in [Hernandez and Jimbo 2012] as we allow only rational powers of q for the eigenvalues of k_i .

Let P_ℓ^r be the subgroup of P_ℓ consisting of Ψ such that $\Psi_i(z)$ is a rational function of z for any $i \in I$.

Theorem 3.7 [Hernandez and Jimbo 2012]. *Let $\Psi \in P_\ell$. A simple object in the category \mathcal{O} is of highest ℓ -weight and the simple module $L(\Psi)$ is in category \mathcal{O} if and only if $\Psi \in P_\ell^r$. Moreover, for V in category \mathcal{O} , $V_\Psi \neq 0$ implies $\Psi \in P_\ell^r$.*

Given a map $c : P_\ell^r \rightarrow \mathbb{Z}$, consider its support

$$\text{supp}(c) = \{\Psi \in P_\ell^r \mid c(\Psi) \neq 0\}.$$

Let \mathcal{E}_ℓ be the additive group of maps $c : P_\ell^r \rightarrow \mathbb{Z}$ such that $\varpi(\text{supp}(c))$ is contained in a finite union of sets of the form $D(\mu)$, and such that for every $\omega \in P_{\mathbb{Q}}$ the set $\text{supp}(c) \cap \varpi^{-1}(\{\omega\})$ is finite. Similarly, let \mathcal{E} be the additive group of maps $c : P_{\mathbb{Q}} \rightarrow \mathbb{Z}$ whose support is contained in a finite union of sets of the form $D(\mu)$. The map ϖ is naturally extended to a surjective homomorphism $\varpi : \mathcal{E}_\ell \rightarrow \mathcal{E}$.

As for the category \mathcal{O} of a classical Kac–Moody Lie algebra, the multiplicity of a simple module in a module of our category \mathcal{O} is well-defined (see [Kac 1990, Section 9.6]) and we have the Grothendieck ring $K_0(\mathcal{O})$. Its elements are the formal sums

$$\chi = \sum_{\Psi \in P_\ell^r} \lambda_\Psi [L(\Psi)],$$

where the $\lambda_\Psi \in \mathbb{Z}$ are set so that

$$\sum_{\Psi \in P_\ell^r, \omega \in P_{\mathbb{Q}}} |\lambda_\Psi| \dim((L(\Psi))_\omega)[\omega]$$

is in \mathcal{E} . For V, V' representations in \mathcal{O} , the product $[V].[V']$ in $K_0(\mathcal{O})$ is naturally obtained by considering the multiplicities of simple modules in $V \otimes V'$ in the sense of [Kac 1990, Section 9.6].

We naturally identify \mathcal{E} with the Grothendieck ring of the category of representations of \mathcal{O} with constant ℓ -weights, the simple objects of which are the $[\omega]$, $\omega \in P_{\mathbb{Q}}$. Thus as in [Kac 1990, Section 9.7] we will regard elements of \mathcal{E} as formal sums

$$c = \sum_{\omega \in \text{supp}(c)} c(\omega)[\omega].$$

The multiplication is given by $[\omega][\omega'] = [\omega + \omega']$ and \mathcal{E} is regarded as a subring of $K_0(\mathcal{O})$. If $(c_i)_{i \in \mathbb{N}}$ is a countable family of elements of \mathcal{E} such that, for any $\omega \in P_{\mathbb{Q}}$, we have $c_i(\omega) \neq 0$ for finitely many $i \in \mathbb{N}$, then the sum $\sum_{i \in \mathbb{N}} c_i$ is well-defined as a map $P_{\mathbb{Q}} \rightarrow \mathbb{Z}$. When this map is in \mathcal{E} we say that $\sum_{i \in \mathbb{N}} c_i$ is a countable sum of elements in \mathcal{E} . Note that we have the analogous notion of a countable sum in $K_0(\mathcal{O})$, compatible with countable sums of characters in \mathcal{E} .

3C. Finite-dimensional representations. Let \mathcal{C} be the category of (type-1) finite-dimensional representations of $U_q(\mathfrak{g})$.

For $i \in I$, let $P_i(z) \in \mathbb{C}[z]$ be a polynomial with constant term 1. Set

$$\Psi = (\Psi_i(z))_{i \in I}, \quad \Psi_i(z) = q_i^{\deg(P_i)} \frac{P_i(zq_i^{-1})}{P_i(zq_i)}.$$

Then $L(\Psi)$ is finite-dimensional. Moreover the action of $U_q(\mathfrak{b})$ can be uniquely extended to an action of $U_q(\mathfrak{g})$, and any simple object in the category \mathcal{C} is of this form. Hence \mathcal{C} is a subcategory of \mathcal{O} and the inclusion functor preserves simple objects.

For $i \in I$ and $a \in \mathbb{C}^*$, we denote by $V_{i,a}$ the simple finite-dimensional representation associated with the polynomials

$$P_i(z) = 1 - za, \quad P_j(z) = 1 \quad (j \neq i).$$

The modules $V_{i,a}$ are called the *fundamental representations*.

3D. The categories \mathcal{O}^+ and \mathcal{O}^- . We introduce two new subcategories \mathcal{O}^+ and \mathcal{O}^- of the category \mathcal{O} .

Definition 3.8. An ℓ -weight is said to be positive (resp. negative) if it is a monomial in the following ℓ -weights:

- $Y_{i,a} = q_i \Psi_{i,aq_i}^{-1} \Psi_{i,aq_i^{-1}}$, where $i \in I$, $a \in \mathbb{C}^*$,
- $\Psi_{i,a}$ (resp. $\Psi_{i,a}^{-1}$), where $i \in I$, $a \in \mathbb{C}^*$,
- $[\omega]$, where $\omega \in P_{\mathbb{Q}}$.

Definition 3.9. \mathcal{O}^+ (resp. \mathcal{O}^-) is the category of representations in \mathcal{O} whose simple constituents have a positive (resp. negative) highest ℓ -weight.

Remark 3.10. (i) By construction, $\mathcal{O}^+, \mathcal{O}^-$ are stable by extensions. We will prove they are also stable by tensor products (Theorem 5.17).

(ii) There are other remarkable subcategories of \mathcal{O} , for example, the category $\widehat{\mathcal{O}}$ of representations of $U_q(\mathfrak{g})$ which belong to \mathcal{O} as representations of $U_q(\mathfrak{b})$. This category $\widehat{\mathcal{O}}$ was introduced in [Hernandez 2005] and further studied in [Mukhin and Young 2014].

(iii) One motivation of Definition 3.9 is that \mathcal{O}^\pm contains \mathcal{C} as well as the prefundamental representations $L_{i,a}^\pm$. We have the following inclusion diagram:

$$\begin{array}{ccc} \mathcal{O} & \supset & \mathcal{O}^+, \mathcal{O}^- \\ \cup & & \cup \\ \widehat{\mathcal{O}} & \supset & \mathcal{C} \end{array}$$

Note that $\widehat{\mathcal{O}}$ is not contained in \mathcal{O}^+ or \mathcal{O}^- , and conversely neither \mathcal{O}^+ nor \mathcal{O}^- is contained in $\widehat{\mathcal{O}}$. For instance, for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, the representation $L((1 - zq)/(q - z))$ is in the category $\widehat{\mathcal{O}}$ by [Mukhin and Young 2014, Theorem 3.6], but not in \mathcal{O}^+ or \mathcal{O}^- because its highest ℓ -weight has no factorization as in equation (5-12) below. On the other hand, the prefundamental representations $L_{1,a}^\pm$ are in the category \mathcal{O}^\pm but not in $\widehat{\mathcal{O}}$ (see [Hernandez and Jimbo 2012, Section 4.1] or [Mukhin and Young 2014, Theorem 3.6]).

(iv) All generalized Baxter’s relations established in [Frenkel and Hernandez 2015] hold in the Grothendieck rings $K_0(\mathcal{O}^+)$ or $K_0(\mathcal{O}^-)$ (see Theorem 5.5 below).

(v) The factorization of real simple modules in \mathcal{O} into prime representations is not unique, so the full category \mathcal{O} is not a good candidate for the notion of monoidal categorification discussed in the introduction. For example, for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, it follows from [Mukhin and Young 2014, Remark 4.3, Theorem 4.6] that

$$L\left(q^{-5} \frac{1 - q^4 z}{1 - q^{-6} z}\right) \otimes L\left(q^{-9} \frac{1 - q^8 z}{1 - q^{-10} z}\right) \simeq L\left(q^{-7} \frac{1 - q^4 z}{1 - q^{-10} z}\right) \otimes L\left(q^{-7} \frac{1 - q^8 z}{1 - q^{-6} z}\right).$$

Moreover the tensor product is simple real and each simple factor has the character of a Verma module. Consequently each factor is not isomorphic to a tensor product of prefundamental representations and so is prime.

4. Cluster algebras

We state the main results of this paper. We refer the reader to [Fomin and Zelevinsky 2003] and [Gekhtman et al. 2010] for an introduction to cluster algebras, and for any undefined terminology.

4A. An infinite-rank cluster algebra. Let us recall the infinite quiver G introduced in [Hernandez and Leclerc 2016, Section 2.1.3]. Put $\tilde{V} = I \times \mathbb{Z}$. Let $\tilde{\Gamma}$ be the quiver with vertex set \tilde{V} . The arrows of $\tilde{\Gamma}$ are given by

$$((i, r) \rightarrow (j, s)) \iff (C_{i,j} \neq 0 \text{ and } s = r + d_i C_{i,j}).$$

By [Hernandez and Leclerc 2016], the quiver $\tilde{\Gamma}$ has two isomorphic connected components. We pick one of the two isomorphic connected components of $\tilde{\Gamma}$ and call it Γ . The vertex set of Γ is denoted by V . A second labeling of the vertices of Γ is deduced from the first one by means of the function ψ defined by

$$\psi(i, r) = (i, r + d_i) \quad ((i, r) \in V). \tag{4-8}$$

Let $W \subset I \times \mathbb{Z}$ be the image of V under ψ . We shall denote by G the same quiver as Γ but with vertices labeled by W .

By analogy with [Hernandez and Leclerc 2016, Section 2.2.1], consider an infinite set of indeterminates

$$z = \{z_{i,r} \mid (i, r) \in V\}$$

over \mathbb{Q} . Let $\mathcal{A}(\Gamma)$ be the cluster algebra defined by the initial seed (z, Γ) . Thus, $\mathcal{A}(\Gamma)$ is the subring of the field of rational functions $\mathbb{Q}(z)$ generated by all the cluster variables, that is, the elements obtained from some element of z via a finite sequence of seed mutations. Each element of $\mathcal{A}(\Gamma)$ is a linear combination of finite monomials in some cluster variables. By the Laurent phenomenon [Fomin and Zelevinsky 2002], $\mathcal{A}(\Gamma)$ is contained in $\mathbb{Z}[z_{i,r}^{\pm 1}]_{(i,r) \in V}$.

For our purposes, it is always possible to work with sufficiently large finite subseeds of the seed (z, Γ) , and replace $\mathcal{A}(\Gamma)$ by the genuine cluster subalgebras attached to them. On the other hand, statements become nicer if we allow ourselves to formulate them in terms of the infinite-rank cluster algebra $\mathcal{A}(\Gamma)$.

Define an \mathcal{E} -algebra homomorphism $\chi : \mathbb{Z}[z_{i,r}^{\pm 1}] \otimes_{\mathbb{Z}} \mathcal{E} \rightarrow \mathcal{E}$ by setting

$$\chi(z_{i,r}^{\pm 1}) = \left[\left(\frac{\mp r}{2d_i} \right) \omega_i \right] \quad ((i, r) \in V).$$

For $A \in \mathcal{A}(\Gamma) \otimes_{\mathbb{Z}} \mathcal{E}$, we write $\chi(A) = \sum_{\omega} A_{\omega}[\omega]$ and $|\chi|(A) = \sum_{\omega} |A_{\omega}|[\omega]$. We will consider the completed tensor product

$$\mathcal{A}(\Gamma) \hat{\otimes}_{\mathbb{Z}} \mathcal{E},$$

that is, the algebra of countable sums $\sum_{i \in \mathbb{N}} A_i$ of elements $A_i \in \mathcal{A}(\Gamma) \otimes_{\mathbb{Z}} \mathcal{E}$ such that $\sum_{i \in \mathbb{N}} |\chi|(A_i)$ is in \mathcal{E} as a countable sum (as defined in Section 3B). Note that in particular we have the analogous notion of a countable sum in $\mathcal{A}(\Gamma) \hat{\otimes}_{\mathbb{Z}} \mathcal{E}$.

4B. Main theorem.

Definition 4.1. Define the category $\mathcal{O}_{2\mathbb{Z}}^+$ as the subcategory of representations in \mathcal{O}^+ whose simple constituents have a highest ℓ -weight Ψ such that the roots and the poles of $\Psi_i(z)$ are of the form q^r with $(i, r) \in V$.

We will write for short $\ell_{i,a}^+ = [L_{i,a}^+]$.

Theorem 4.2. *The category $\mathcal{O}_{2\mathbb{Z}}^+$ is monoidal and the identification*

$$z_{i,r} \otimes \left[\frac{r}{2d_i} \omega_i \right] \equiv \ell_{i,q^r}^+ \quad ((i, r) \in V) \tag{4-9}$$

defines an isomorphism of \mathcal{E} -algebras

$$\mathcal{A}(\Gamma) \hat{\otimes}_{\mathbb{Z}} \mathcal{E} \simeq K_0(\mathcal{O}_{2\mathbb{Z}}^+)$$

compatible with countable sums.

Remark 4.3. (i) The identification (4-9) gives an isomorphism of \mathcal{E} -algebras

$$\mathcal{E}[z_{i,r}^{\pm 1}]_{(i,r) \in V} \xrightarrow{\sim} \mathcal{E}[(\ell_{i,q^r}^+)^{\pm 1}]_{(i,r) \in V},$$

which can be extended to countable sums as above. So the main point of the proof of Theorem 4.2 will be to show that the subalgebra $\mathcal{A}(G) \hat{\otimes}_{\mathbb{Z}} \mathcal{E}$ is mapped to $K_0(\mathcal{O}_{2\mathbb{Z}}^+)$ by this isomorphism.

(ii) As in the case of finite-dimensional representations, the description of the simple objects of \mathcal{O}^+ essentially reduces to the description of the simple objects of $\mathcal{O}_{2\mathbb{Z}}^+$ (the decomposition explained in [Hernandez and Leclerc 2010, Section 3.7] can be extended to our more general situation by using the asymptotic approach of Section 7B below). Hence the Grothendieck ring $K_0(\mathcal{O}_{2\mathbb{Z}}^+)$ contains all the interesting information on $K_0(\mathcal{O}^+)$.

The proof of Theorem 4.2 will be given in Section 6, using material presented in Section 5.

5. Properties of the category \mathcal{O}^+

5A. q -characters. The q -character morphism was first considered in [Frenkel and Reshetikhin 1999] and is a very useful tool for our proofs.

Recall from Section 3B the notation \mathcal{E} and \mathcal{E}_ℓ . Because of the support condition, we can endow \mathcal{E} with a ring structure defined by

$$(c \cdot d)(\omega) = \sum_{\omega' + \omega'' = \omega} c(\omega')d(\omega'') \quad (c, d \in \mathcal{E}, \omega \in P_{\mathbb{Q}}).$$

Similarly, \mathcal{E}_ℓ also has a ring structure given by

$$(c \cdot d)(\Psi) = \sum_{\Psi' \Psi'' = \Psi} c(\Psi')d(\Psi'') \quad (c, d \in \mathcal{E}_\ell, \Psi \in P_\ell^{\text{r}}),$$

and such that ϖ becomes a ring homomorphism.

For $\Psi \in P_\ell^{\text{r}}$ and $\omega \in P_{\mathbb{Q}}$, we define the delta functions $[\Psi] = \delta_{\Psi, \cdot} \in \mathcal{E}_\ell$ and $[\omega] = \delta_{\omega, \cdot} \in \mathcal{E}$, where as usual δ denotes the Kronecker symbol. Note that the above multiplications give

$$[\Psi'] \cdot [\Psi''] = [\Psi' \Psi''], \quad [\omega'] \cdot [\omega''] = [\omega' + \omega''].$$

Let V be a $U_q(\mathfrak{b})$ -module in category \mathcal{O} . We define [Frenkel and Reshetikhin 1999; Hernandez and Jimbo 2012] the q -character and the character of V :

$$\chi_q(V) := \sum_{\Psi \in P_\ell^{\text{r}}} \dim(V_\Psi)[\Psi] \in \mathcal{E}_\ell, \quad \chi(V) := \varpi(\chi_q(V)) = \sum_{\omega \in P_{\mathbb{Q}}} \dim(V_\omega)[\omega] \in \mathcal{E}.$$

If $V \in \mathcal{O}$ has a unique ℓ -weight Ψ whose weight $\varpi(\Psi)$ is maximal, we also consider its normalized q -character $\tilde{\chi}_q(V)$ and normalized character $\tilde{\chi}(V)$ defined by

$$\tilde{\chi}_q(V) := [\Psi^{-1}] \cdot \chi_q(V), \quad \tilde{\chi}(V) := \varpi(\tilde{\chi}_q(V)).$$

Note that

$$\chi_q(\tilde{L}(\Psi)) = [\tilde{\Psi}] \cdot \tilde{\chi}_q(L(\Psi)) \neq \tilde{\chi}_q(L(\Psi)).$$

Proposition 5.1 [Hernandez and Jimbo 2012]. *The q -character morphism*

$$\chi_q : K_0(\mathcal{O}) \rightarrow \mathcal{E}_\ell, \quad [V] \mapsto \chi_q(V),$$

is an injective ring morphism.

Following [Frenkel and Reshetikhin 1999], consider the ring of Laurent polynomials $\mathcal{Y} = \mathbb{Z}[Y_{i,a}^{\pm 1}]_{i \in I, a \in \mathbb{C}^*}$ in the indeterminates $\{Y_{i,a}\}_{i \in I, a \in \mathbb{C}^*}$. Let \mathcal{M} be the multiplicative group of Laurent monomials in \mathcal{Y} . For example, for $i \in I$ and $a \in \mathbb{C}^*$ define $A_{i,a} \in \mathcal{M}$ by

$$A_{i,a} = Y_{i,aq_i^{-1}} Y_{i,aq_i} \left(\prod_{j:C_{j,i}=-1} Y_{j,a} \prod_{j:C_{j,i}=-2} Y_{j,aq^{-1}} Y_{j,aq} \prod_{j:C_{j,i}=-3} Y_{j,aq^{-2}} Y_{j,a} Y_{j,aq^2} \right)^{-1}.$$

For a monomial $m = \prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{u_{i,a}} \in \mathcal{M}$, we consider its “evaluation at $\phi^+(z)$ ”. By definition it is the element $m(\phi(z)) \in P_\ell^{\text{r}}$ given by

$$m(\phi(z)) = \prod_{i \in I, a \in \mathbb{C}^*} (Y_{i,a}(\phi(z)))^{u_{i,a}}, \quad \text{where } (Y_{i,a}(\phi(z)))_j = \begin{cases} q_i \frac{1-aq_i^{-1}z}{1-aq_i z}, & j = i, \\ 1, & j \neq i. \end{cases}$$

This defines an injective group morphism $\mathcal{M} \rightarrow P_\ell^\mathbb{F}$. We identify a monomial $m \in \mathcal{M}$ with its image in $P_\ell^\mathbb{F}$. This is compatible with the notation $Y_{i,a}$ used in Definition 3.8. Note that $\varpi(Y_{i,a}) = \omega_i$ and $\varpi(A_{i,a}) = \alpha_i$.

It is proved in [Frenkel and Reshetikhin 1999] that a finite-dimensional $U_q(\mathfrak{g})$ -module V satisfies $V = \bigoplus_{m \in \mathcal{M}} V_{m(\phi(z))}$. In particular, $\chi_q(V)$ can be viewed as an element of \mathcal{Y} .

A monomial $M \in \mathcal{M}$ is said to be dominant if $M \in \mathbb{Z}[Y_{i,a}]_{i \in I, a \in \mathbb{C}^*}$. Given a finite-dimensional simple $U_q(\mathfrak{g})$ -module $L(\Psi)$, there exists a dominant monomial $M \in \mathcal{M}$ such that $\Psi = M(\phi(z))$. We will also set $L(\Psi) = L(M)$.

For example, for $i \in I$, $a \in \mathbb{C}^*$, $k \geq 0$, we have the Kirillov–Reshetikhin module

$$W_{k,a}^{(i)} = L(Y_{i,a} Y_{i,aq^2} \cdots Y_{i,aq_i^{2(k-1)}}). \quad (5-10)$$

Example 5.2. If $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, we have ($k \geq 0$, $a \in \mathbb{C}^*$) [Frenkel and Reshetikhin 1999]

$$\begin{aligned} \chi_q(W_{k,aq^{1-2k}}^{(1)}) \\ = Y_{aq^{-1}} Y_{aq^{-3}} \cdots Y_{aq^{-2k+1}} (1 + A_{1,a}^{-1} + A_{1,a}^{-1} A_{1,aq^{-2}}^{-1} + \cdots + A_{1,a}^{-1} \cdots A_{1,aq^{-2(k-1)}}^{-1}). \end{aligned}$$

Theorem 5.3. (i) [Hernandez and Jimbo 2012; Frenkel and Hernandez 2015]. For any $a \in \mathbb{C}^*$, $i \in I$ we have

$$\chi_q(L_{i,a}^+) = [\Psi_{i,a}] \chi(L_{i,a}^+) = [\Psi_{i,a}] \chi(L_{i,a}^-),$$

where $\chi(L_{i,a}^+) = \chi(L_{i,a}^-)$ does not depend on a .

(ii) [Hernandez and Jimbo 2012]. For any $a \in \mathbb{C}^*$, $i \in I$ we have

$$\chi_q(L_{i,a}^-) \in [\Psi_{i,a}^{-1}] (1 + A_{i,a}^{-1} \mathbb{Z} \llbracket A_{j,b}^{-1} \rrbracket_{j \in I, b \in \mathbb{C}^*}).$$

Example 5.4. In the case $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, we have

$$\begin{aligned} \chi_q(L_{1,a}^+) &= [(1 - za)] \sum_{r \geq 0} [-2r\omega_1], \\ \chi_q(L_{1,a}^-) &= \left[\frac{1}{(1 - za)} \right] \sum_{r \geq 0} A_{1,a}^{-1} A_{1,aq^{-2}}^{-1} \cdots A_{1,aq^{-2(r-1)}}^{-1}. \end{aligned}$$

Thus, although positive and negative prefundamental representations have the same character,

$$\chi(L_{1,a}^+) = \chi(L_{1,a}^-) = \sum_{r \geq 0} [-2r\omega_1],$$

their q -characters are very different: the normalized q -character $\tilde{\chi}_q(L_{1,a}^+)$ is independent of the spectral parameter a , whereas $\tilde{\chi}_q(L_{1,a}^-)$ does depend on a .

5B. Baxter relations. We can now state the *generalized Baxter relations*:

Theorem 5.5 [Frenkel and Hernandez 2015, Theorem 4.8, Remark 4.10].² *Let V be a finite-dimensional representation of $U_q(\mathfrak{g})$. Replace in $\chi_q(V)$ each variable $Y_{i,a}$ by $[\omega_i][L_{i,aq_i}^+]/[L_{i,aq_i}^+]$ (resp. $[-\omega_i][L_{i,aq_i}^-]/[L_{i,aq_i}^-]$) and $\chi_q(V)$ by $[V]$ (resp. $[V^*]$). Then multiplying by a common denominator we get a relation in the Grothendieck ring $K_0(\mathcal{O})$.*

Example 5.6. Taking $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ and $V = L(Y_{1,a})$, we obtain relation (1-2), namely,

$$\begin{aligned} [L(Y_{1,a})][L_{1,aq}^+] &= [L_{1,aq^{-1}}^+][\omega_1] + [L_{1,aq^3}^+][-\omega_1], \\ [L(Y_{1,aq^2})][L_{1,aq}^-] &= [L_{1,aq^{-1}}^-][-\omega_1] + [L_{1,aq^3}^-][\omega_1]. \end{aligned}$$

Remark 5.7. By our main result, Theorem 4.2, (and its dual version; see Remark 5.18), these generalized Baxter relations are interpreted as relations in the cluster algebra $\mathcal{A}(G)$. Moreover, when $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, the original Baxter relation (1-2) gets interpreted as a Fomin–Zelevinsky mutation relation.

The right-hand side of a generalized Baxter relation is an \mathcal{E} -linear combination of classes of tensor products of prefundamental representations. As shown in [Frenkel and Hernandez 2015], this can be seen as a tensor product decomposition in $K(\mathcal{O})$. Indeed, we have:

Theorem 5.8 [Frenkel and Hernandez 2015]. *Any tensor product of positive (resp. negative) prefundamental representations $L_{i,a}^+$ (resp. $L_{i,a}^-$) is simple.*

5C. Prefundamental characters of representations and duality. Let K_0^\pm be the \mathcal{E} -subalgebra of $K_0(\mathcal{O}^\pm)$ generated by the $[V_{i,a}], [L_{i,a}^\pm]$ ($i \in I, a \in \mathbb{C}^*$).

It follows from Theorem 5.5 that the fraction field of $\mathcal{E}[\ell_{i,a}^\pm]_{i \in I, a \in \mathbb{C}^*}$ contains K_0^\pm . More precisely, each element of K_0^\pm is a Laurent polynomial in $\mathcal{E}[(\ell_{i,a}^\pm)^{\pm 1}]_{i \in I, a \in \mathbb{C}^*}$. Note that the $\ell_{i,a}^\pm$ are algebraically independent (this is for example a consequence of Theorem 5.8, which implies that the monomials in the $\ell_{i,a}^\pm$ are linearly independent in $K_0(\mathcal{O}^+)$). In particular the expansion in $\mathcal{E}[(\ell_{i,a}^\pm)^{\pm 1}]_{i \in I, a \in \mathbb{C}^*}$ is unique and we can define the injective ring morphism

$$\chi^+ : K_0^+ \rightarrow \mathcal{E}[(\ell_{i,a}^+)^{\pm 1}]_{i \in I, a \in \mathbb{C}^*},$$

which is called the positive prefundamental character morphism.

In the same way, by using the relations in Theorem 5.5 in terms of the prefundamental representations $[L_{i,a}^-]$ and by setting $\ell_{i,a}^- = [L_{i,a^{-1}}^-]$, we get the negative

²The result in [Frenkel and Hernandez 2015] is stated in terms of the $L_{i,a}^+$, and the $R_{i,a}^+$ such that $(R_{i,a}^+)^* \simeq L_{i,a}^-$. In the last case, the variable $Y_{i,a}$ has to be replaced by $[\omega_i][R_{i,aq_i^{-1}}^+]/[R_{i,aq_i}^+]$ to get $[V]$. That is why in the statement of Theorem 5.5 in terms of the $L_{i,a}^-$ the representations $[-\omega_i] \simeq [\omega_i]^*$ and V^* appear.

prefundamental character morphism, which is an injective ring morphism

$$\chi^- : K_0^- \rightarrow \mathcal{E}[(\ell_{i,a}^-)^{\pm 1}]_{i \in I, a \in \mathbb{C}^*}.$$

Example 5.9. We can reformulate Theorem 5.5 as follows. For a finite-dimensional representation V of $U_q(\mathfrak{g})$, the positive (resp. negative) prefundamental character $\chi^+(V)$ (resp. $\chi^-(V^*)$) is obtained from the q -character $\chi_q(V)$ by replacing each variable $Y_{i,a}$ by $[\omega_i] \ell_{i,aq_i}^+ (\ell_{i,aq_i}^+)^{-1}$ (resp. $[-\omega_i] \ell_{i,a^{-1}q_i}^- (\ell_{i,a^{-1}q_i}^-)^{-1}$). In fact, this can also be seen as a change of variables analogous to that of [Hernandez and Leclerc 2016, Section 5.2.2]. For example, for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, using Example 5.6 we have

$$\begin{aligned} \chi^+(L(Y_{1,a})) &= \frac{[\omega_1] \ell_{1,aq^{-1}}^+ + [-\omega_1] \ell_{1,aq^3}^+}{\ell_{1,aq}^+}, \\ \chi^-(L(Y_{1,a})) &= \frac{[-\omega_1] \ell_{1,a^{-1}q^3}^- + [\omega_1] \ell_{1,a^{-1}q^{-1}}^-}{\ell_{i,a^{-1}q}^-}. \end{aligned} \tag{5-11}$$

Example 5.10. Set $\mathfrak{g} = \widehat{\mathfrak{sl}}_3$. Let

$$\Psi = ((1 - zq^{-2})/(1 - z), (1 - qz)) = [-\omega_1] Y_{1,q^{-1}} \Psi_{2,q}.$$

Let us compute $\chi^+(L(\Psi))$. For $k \geq 0$, let $W_k = \widetilde{L}(M_k)$, where

$$M_k = (Y_{2,q^{2k}} \cdots Y_{2,q^4} Y_{2,q^2}) Y_{1,q^{-1}}.$$

We can prove as in [Hernandez and Jimbo 2012, Section 7.2] that $(W_k)_{k \geq 0}$ gives rise to a limiting $U_q(\mathfrak{b})$ -module W_∞ whose q -character is

$$\chi_q(W_\infty) = \Psi(1 + A_{1,1}^{-1}) \chi(L_{2,q}^+) = \chi_q(L_{2,q}^+) [-\omega_1] Y_{1,q^{-1}} + [-\omega_1 + \omega_2] Y_{1,q}^{-1} \chi_q(L_{2,q^{-1}}^+).$$

But it follows from Theorem 7.6(ii) that $L(\Psi)$ and W_∞ have the same character, $\lim_{k \rightarrow +\infty} \chi(\widetilde{L}(M_k)) = \lim_{k \rightarrow +\infty} \chi(\widetilde{L}(\overline{M}_k^{-1}))$. As $L(\Psi)$ is a subquotient of W_∞ , they are isomorphic. Consequently

$$\chi^+(L(\Psi)) = \frac{\ell_{2,q}^+ \ell_{1,q^{-2}}^+ + [-\alpha_1] \ell_{1,q^2}^+ \ell_{2,q^{-1}}^+}{\ell_{1,1}^+}.$$

Proposition 5.11. A representation V in \mathcal{O}^\pm satisfying $[V] \in K_0^\pm$ has finite length.

Remark 5.12. An object in the category \mathcal{O} does not necessarily have finite length. The subcategory of objects of finite length is not stable by tensor product [Boos et al. 2009, Lemma C.1].

Proof. Suppose $[V] \in K_0^+$. Then there is a monomial M in the $\ell_{i,a}^+$ such that $M\chi^+(V)$ is a polynomial in the $\ell_{i,a}^+$. There is a tensor product L of positive prefundamental representations such that $\chi^+(L) = M$. Then $L \otimes V$ has finite length, hence the result. \square

Proposition 5.13. *The identification of the variables $\ell_{i,a}^+$ and $\ell_{i,a}^-$ induces a unique isomorphism of \mathcal{E} -algebras*

$$D : K_0^+ \rightarrow K_0^-.$$

Proof. The identification gives a well-defined injective ring morphism

$$D' : K_0^+ \rightarrow \mathcal{E}[(\ell_{i,a}^-)^{\pm 1}]_{i \in I, a \in \mathbb{C}^*}.$$

It suffices to prove that its image is $\chi^-(K_0^-)$. For $V = L_{i,b}^+$ a prefundamental, its image by D' is $\ell_{i,b}^- = \chi^-([L_{i,b-1}^-])$. For $V = L(Y_{i,b})$ a fundamental, its image by D' is obtained from its q -character $\chi_q(V)$ by replacing each variable $Y_{i,a}$ by $[\omega_i][L_{i,aq_i}^-]/[L_{i,aq_i}^-]$, that is, by $[\omega_i]\ell_{i,a-1q_i}^-/\ell_{i,a-1q_i}^-$. In the construction of χ^- , this corresponds to $Y_{i,a-1}^{-1}$ (see the formula in Theorem 5.5). By [Hernandez 2007, Lemma 4.10], there is a finite-dimensional representation V' whose q -character is obtained from $\chi_q(V)$ by replacing each $Y_{i,a}$ by $Y_{i,a-1}^{-1}$. Hence $D'(V) = \chi^-((V')^*) \in \chi^-(K_0^-)$. We have proved $\text{Im}(D') \subset \chi^-(K_0(\widehat{\mathcal{O}}_{f,\mathbb{Z}}^-))$. Similarly, for W such that $V = W^*$, we have $D'(W') = \chi^-(V)$ and we get the other inclusion. \square

Example 5.14. (i) For any $(i, a) \in I \times \mathbb{C}^*$, we have

$$D([L_{i,a}^+]) = [L_{i,a-1}^-].$$

(ii) For any dominant monomial m , we have

$$D([L(m)]) = [L(m_1)],$$

where m_1 is obtained from m by replacing each $Y_{i,a}$ by $Y_{i,a-1}$. Indeed $L(m_1)$ is isomorphic to $((L(m))')^*$, whose highest weight monomial is the inverse of the lowest weight monomial $(m_1)^{-1}$ of $L(m)'$. For example, for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, we have $D([L(Y_{1,a})]) = [L(Y_{1,a-1})]$, as this can be observed by comparing the two formulas in (5-11).

Remark 5.15. The duality D is compatible with characters by (i) in Theorem 5.3. However it is not compatible with q -characters (for example, negative and positive prefundamental representations have very different q -characters, as explained above).

Proposition 5.16. *An element in the Grothendieck group $K_0(\mathcal{O}^\pm)$ is a (possibly countable) sum of elements in K_0^\pm .*

Proof. Let us prove it for $K_0(\mathcal{O}^+)$ (the proof is analogous for $K_0(\mathcal{O}^-)$). By definition, the positive ℓ -weights label the simple modules in \mathcal{O}^+ . Moreover, an ℓ -weight is positive if and only if it is a product of highest ℓ -weights of representations $L_{i,a}^+$, $V_i(a)$ and $[\omega]$. This implies that, for each positive ℓ -weight Ψ , we can choose (and fix) a monomial in the $[L_{i,a}^+]$, $[V_i(a)]$, $[\omega]$ such that the corresponding representation has highest ℓ -weight equal to Ψ . Hence the positive ℓ -weights also

label the linearly independent family of these monomials in K_0^+ . Expanding these monomials we get finite sums of classes of simple modules by Proposition 5.11. We get an (infinite) transition matrix from the classes of simple objects in \mathcal{O}^+ to such products in K_0^+ , and this matrix is unitriangular (for the standard partial ordering with respect to weight). Hence the result. \square

This implies immediately the following.

Theorem 5.17. \mathcal{O}^+ and \mathcal{O}^- are monoidal and the morphism D extends uniquely to an isomorphism of \mathcal{E} -algebras

$$D : K_0(\mathcal{O}^+) \rightarrow K_0(\mathcal{O}^-).$$

Remark 5.18. Consequently, our main result in Theorem 4.2 may also be written in terms of the subcategory $\mathcal{O}_{\mathbb{Z}}^-$ of $\mathcal{O}_{\mathbb{Z}}^-$ whose Grothendieck ring is $D(K_0(\mathcal{O}_{\mathbb{Z}}^+))$.

Proof. For L, L' simple in \mathcal{O}^+ , we may consider a decomposition of $[L], [L']$ as a countable sum of elements in K_0^+ as in Proposition 5.16. Then $[L][L']$ is also such a countable sum and is in $K_0(\mathcal{O}^+)$. Hence \mathcal{O}^+ is monoidal. This is analogous for \mathcal{O}^- .

The isomorphism of Proposition 5.13 is extended by linearity to $K_0(\mathcal{O}^+)$ by using Proposition 5.16. This map $D : K_0(\mathcal{O}^+) \rightarrow K_0(\mathcal{O}^-)$ is an injective ring morphism. The ring morphism $D^{-1} : K_0(\mathcal{O}^-) \rightarrow K_0(\mathcal{O}^+)$ is constructed in the same way and so D is a ring isomorphism. \square

Proposition 5.19. A simple object in \mathcal{O}^+ (resp. in \mathcal{O}^-) is a subquotient of a tensor product of two simple representations $V \otimes L$, where V is finite-dimensional and L is a tensor product of positive (resp. negative) prefundamental representations.

Proof. Let $L(\Psi)$ be simple in \mathcal{O}^\pm . By definition, its highest ℓ -weight is a product of highest ℓ -weights of representations $[\omega], L_{i,a}^\pm, V_i(a)$, where $\omega \in P_{\mathbb{Q}}, i \in I, a \in \mathbb{C}^*$. So Ψ can be factorized as

$$\Psi = [\omega] \times m \times \prod_{i \in I, a \in \mathbb{C}^*} \Psi_{i,a}^{u_{i,a}}, \tag{5-12}$$

where $\omega \in P_{\mathbb{Q}}, \pm u_{i,a} \geq 0$ and $m \in \mathcal{M}$ is a dominant monomial. The result follows by taking $V = [\omega] \otimes L(m)$ and $L = \bigotimes_{i \in I, a \in \mathbb{C}^*} (L_{i,a}^\pm)^{\otimes |u_{i,a}|}$, which is simple by Theorem 5.8. \square

Proposition 5.20. The normalized q -character of a simple object in \mathcal{O}^- belongs to the ring $\mathbb{Z}[[A_{i,a}^{-1}]]_{i \in I, a \in \mathbb{C}^*}$.


Proof. The result is known for the category \mathcal{C} by [Frenkel and Mukhin 2001]. For negative prefundamental representations, the result is known by [Hernandez and Jimbo 2012, Theorem 6.1]. The general result follows from Proposition 5.19. \square

Note that this property is not satisfied in \mathcal{O}^+ ; see Example 5.4.


6. Proof of the main theorem

6A. Examples of mutations.

6A1. Let $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$. We display a sequence of 3 mutations starting from the initial seed of $\mathcal{A}(G)$. The mutated cluster variables are indicated by a framebox.



6A2. Let $\mathfrak{g} = \widehat{\mathfrak{sl}}_3$. Example 5.10 can be reformulated as a one-step mutation from the initial seed, as follows:



Recall that here $\Psi = [-\omega_1]Y_{1,q-1}\Psi_{2,q}$, as in Example 5.10.

6A3. For an arbitrary \mathfrak{g} , let us calculate the first mutation relation for each cluster variable $z_{i,r}$ of the initial seed, generalizing Section 6A2. We denote by $z_{i,r}^*$ the new cluster variable obtained by mutating $z_{i,r}$. Then we have

$$z_{i,r}^* z_{i,r} = \prod_{j:C_{j,i} \neq 0} z_{j,r-d_j C_{j,i}} + \prod_{j:C_{j,i} \neq 0} z_{j,r+d_j C_{j,i}}$$

We claim that $z_{i,r}^* = [\lambda][L(\Psi)]$, where

$$\Psi = [-\omega_i]Y_{i,q^{r-d_i}} \prod_{j:C_{j,i} < 0} \Psi_{j,q^{r-d_j}C_{j,i}} \quad \text{and} \quad \lambda = \frac{\alpha_i}{2} - r \sum_{j:C_{j,i} < 0} \frac{\omega_j}{2d_j}.$$

As in Section 6A2, this is derived from the explicit q -character formula

$$\chi_q(L(\Psi)) = [\Psi](1 + A_{i,q^r}^{-1}) \prod_{j:C_{j,i} < 0} \chi_j, \tag{6-13}$$

where $\chi_j = \chi(L_{j,a}^+)$ does not depend on a . (By considering $L(\Psi) \otimes L(Y_{i,q^{r-d_i}} \Psi^{-1})$ and $L(\Psi) \otimes L_{i,q^r}^+$ we prove that the multiplicities in $\chi_q(L(\Psi))$ are larger than in the right-hand side of (6-13). The reverse inequality is established by considering $L(\overline{M}_R^{-1}) \otimes L(\Psi \overline{M}_R)$, where the monomials M_R are defined for $\overline{\Psi}^{-1}$ as in Theorem 7.6.) The mutation relation thus becomes the following relation in the Grothendieck ring $K_0(\mathcal{O}^+)$:

$$[L(\Psi) \otimes L_{i,q^r}^+] = \left[\bigotimes_{j:C_{j,i} \neq 0} L_{j,q^{r-d_j}C_{j,i}}^+ \right] + [-\alpha_i] \left[\bigotimes_{j:C_{j,i} \neq 0} L_{j,q^{r+d_j}C_{j,i}}^+ \right]. \tag{6-14}$$

By Theorem 5.8, the two terms on the right-hand side are simple. Hence this is the decomposition of the class of the tensor product into simple modules.

6B. Proof of Theorem 4.2. We identify the \mathcal{E} -algebras

$$\mathcal{E}[z_{i,r}^{\pm 1}]_{(i,r) \in V} \quad \text{and} \quad \mathcal{E}[(\ell_{i,q^r}^+)^{\pm 1}]_{(i,r) \in V}$$

as in Remark 4.3.

Proposition 6.1. *We have $K_0^+ \cap K_0(\mathcal{O}_{2\mathbb{Z}}^+) \subseteq \mathcal{A}(G) \otimes_{\mathbb{Z}} \mathcal{E}$.*

Note that by Proposition 5.16 this implies $K_0(\mathcal{O}_{2\mathbb{Z}}^+) \subseteq \mathcal{A}(G) \hat{\otimes}_{\mathbb{Z}} \mathcal{E}$.

Proof. Clearly, $\ell_{i,q^s}^+ = z_{i,s}[(s/2d_i)\omega_i]$ belongs to $\mathcal{A}(G)_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{E}$ for any $(i, s) \in V$. By Proposition 5.16, it remains to show that $[V_{i,q^r}]$ belongs to $\mathcal{A}(G)_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{E}$ for any $(i, r) \in W$.

Remember from Section 4A that we denote by G the same quiver as Γ with vertices labeled by W instead of V . In the next discussion, we divide the vertices of G and Γ into *columns*, as in [Hernandez and Leclerc 2016, Example 2.3], and we denote by k the number of columns. As in [Hernandez and Leclerc 2016, Section 2.1.3], consider the full subquiver G^- of G whose vertex set is $W^- = \{(i, r) \in W \mid r \leq 0\}$. The definition of G shows that there is only one vertex of $G \setminus G^-$ in each column which is connected to G^- by some arrow. Let H^- denote the ice quiver obtained from G^- by adding these k vertices together with their connecting arrows, and by declaring the new vertices frozen.

Consider the cluster algebras $\mathcal{A}(H^-)$ (with k frozen variables) and $\mathcal{A}(G^-)$ (with no frozen variable). It follows from the definitions that $\mathcal{A}(H^-)$ can be regarded as a subalgebra of $\mathcal{A}(G)$, and $\mathcal{A}(G^-)$ is the coefficient-free counterpart of $\mathcal{A}(H^-)$, studied in [Hernandez and Leclerc 2016]. Let f_j ($1 \leq j \leq k$) be the frozen variable of $\mathcal{A}(H^-)$ sitting in column j . The cluster of the initial seed of $\mathcal{A}(H^-)$ thus consists of the frozen variables f_j ($1 \leq j \leq k$) and the ordinary cluster variables $z_{i,s}$ ($(i, s) \in V^-$), where $V^- = \{(i, s) \in V \mid s + d_i \leq 0\}$. Let us denote by $u_{i,s}$ ($(i, s) \in V^-$) the cluster variables of the initial seed of $\mathcal{A}(G^-)$. We can use a similar change of variables as in [Hernandez and Leclerc 2016, Section 2.2.2]:

$$y_{i,r} = u_{i,r-d_i} \quad \text{if } r + d_i > 0, \quad y_{i,r} = \frac{u_{i,r-d_i}}{u_{i,r+d_i}} \quad \text{otherwise.}$$

Let

$$F : \mathbb{Z}[u_{i,s}^{\pm 1} \mid (i, s) \in V^-] \rightarrow \mathbb{Z}[f_j^{\pm 1}, z_{i,s}^{\pm 1} \mid 1 \leq j \leq k, (i, s) \in V^-]$$

be the ring homomorphism defined by

$$F(u_{i,s}) = \frac{z_{i,s}}{f_j} \quad \text{if } (i, s) \text{ sits in column } j.$$

Thus

$$F(y_{i,r}) = \begin{cases} \frac{z_{i,r-d_i}}{f_j} & \text{if } r + d_i > 0 \text{ and } (i, r) \text{ sits in column } j, \\ f_j & \\ \frac{z_{i,r-d_i}}{z_{i,r+d_i}} & \text{otherwise.} \end{cases}$$

We introduce a \mathbb{Z}^k -grading on $\mathbb{Z}[f_j^{\pm 1}, z_{i,s}^{\pm 1} \mid 1 \leq j \leq k, (i, s) \in V^-]$ by declaring that

$$\deg(f_j) = e_j, \quad \deg(z_{i,s}) = e_j \quad \text{if } (i, s) \text{ sits in column } j,$$

where e_j ($1 \leq j \leq k$) denotes the canonical basis of \mathbb{Z}^k .

Let x be the cluster variable of $\mathcal{A}(G^-)$ obtained from the initial seed $(\{u_{i,s}\}, G^-)$ via a sequence of mutations σ , and let y be the cluster variable of $\mathcal{A}(H^-)$ obtained from the initial seed $(\{z_{i,s}, f_j\}, H^-)$ via the *same* sequence of mutations σ . We want to compare the Laurent polynomials y and $F(x)$. Since $\deg(F(u_{i,s})) = (0, \dots, 0)$ for every (i, s) , we see that $F(x)$ is multihomogeneous of degree $(0, \dots, 0)$ for the above grading. On the other hand, it is easy to check that for every nonfrozen vertex (i, s) of the ice quiver H^- the sum of the multidegrees of the initial cluster variables and frozen variables sitting at the targets of the arrows going out of (i, s) is equal to the sum of the multidegrees of the initial cluster variables and frozen variables sitting at the sources of the arrows going into (i, s) . Therefore, $\mathcal{A}(H^-)$ is a multigraded cluster algebra, in the sense of [Grabowski and Launois 2014]. It follows that y is

also multihomogeneous, of degree (a_1, \dots, a_k) . Now, by construction, we have

$$F(x)|_{f_1=1, \dots, f_k=1} = y|_{f_1=1, \dots, f_k=1}.$$

Therefore,

$$y = F(x) \prod_{j=1}^k f_j^{a_j}.$$

Taking a cluster expansion with respect to the initial cluster of $\mathcal{A}(H^-)$, we write $y = N/D$ where D is a monomial in the nonfrozen cluster variables and N is a polynomial in the nonfrozen and frozen variables. Moreover N is not divisible by any of the f_j . It follows that $\prod_{j=1}^k f_j^{a_j}$ is the smallest monomial such that $F(x) \prod_{j=1}^k f_j^{a_j}$ contains only nonnegative powers of the variables f_j .

Now we can conclude using [Hernandez and Leclerc 2016, Theorem 3.1], which implies that, for all $(i, r) \in W^-$ with $r \ll 0$, the q -character of V_{i,q^r} (expressed in terms of the variables $y_{i,s} \equiv Y_{i,q^s}$) is a cluster variable x of $\mathcal{A}(G^-)$. By [Frenkel and Mukhin 2001, Corollary 6.14], for $r \ll 0$ this cluster variable does not contain any variable $y_{i,s}$ with $s + 2d_i > 0$, hence $F(x)$ does not contain any frozen variable f_j . Therefore $y = F(x)$, and the q -character of V_{i,q^r} (expressed in terms of the variables $z_{i,s}$) is a cluster variable of $\mathcal{A}(H^-)$, that is, a cluster variable of $\mathcal{A}(\Gamma)$. This proves the claim for every fundamental module V_{i,q^r} with $r \ll 0$. But by definition of the cluster algebra $\mathcal{A}(\Gamma)$, the set of cluster variables is invariant under the change of variables $z_{i,s} \mapsto z_{i,s+2d_i}$. Thus we are done. \square

Proposition 6.2. *We have $\mathcal{A}(G) \hat{\otimes}_{\mathbb{Z}} \mathcal{E} \subseteq K_0(\mathcal{O}_{2\mathbb{Z}}^+)$.*

To prove this proposition, we need to establish some preliminary results. We first prove in Lemma 6.4 that, for $\chi \in \mathcal{A}(G)$, at least one negative ℓ -weight occurs in $\chi_q(D(\chi))$. Then in Lemma 6.5 we construct a family of distinguished elements

$$F(\Psi) \in \chi_q(D(K_0(\mathcal{O}^+))).$$

The first result allows us to write each element of $\chi_q(D(\mathcal{A}(G)))$ as a linear combination of the $F(\Psi)$. This implies the inclusion in Proposition 6.2.

So consider an element χ in $\mathcal{A}(G)$. By the Laurent phenomenon [Fomin and Zelevinsky 2002], χ is a Laurent polynomial in the initial cluster variables:

$$\chi = P(\{z_{i,r}\}_{(i,r) \in V}).$$

Hence $\mathcal{A}(G)$ is a subalgebra of the fraction field of $K_0(\mathcal{O}_{2\mathbb{Z}}^+)$, and the duality D of Proposition 5.13 can be algebraically extended to $\mathcal{A}(G)$. In particular we have

$$D(\chi) = P(\{D(z_{i,r})\}_{(i,r) \in V}) \in \text{Frac}(K_0(\mathcal{O}_{2\mathbb{Z}}^-))$$

in the fraction field of $K_0(\mathcal{O}_{2\mathbb{Z}}^-)$. The q -character morphism can also be algebraically extended to $\text{Frac}(K_0(\mathcal{O}_{2\mathbb{Z}}^-))$. Then $\chi_q(D(\chi))$ is obtained by replacing each $z_{i,r}$ by

the corresponding q -character

$$\chi_q(D(z_{i,r})) = \left[\left(\frac{-r}{2d_i} \right) \omega_i \right] \chi_q(L_{i,q^{-r}}^-) = \left[\left(\frac{-r}{2d_i} \right) \omega_i \right] \Psi_{i,q^{-r}}^{-1} (1 + \mathcal{A}_{i,r}), \quad (6-15)$$

where, by Theorem 7.6, $\mathcal{A}_{i,r}$ is a formal power series in the $A_{j,a}^{-1}$ without constant term. In particular, we have an analogous formula for the inverse,

$$(\chi_q(D(z_{i,r})))^{-1} = \left[\left(\frac{r}{2d_i} \right) \omega_i \right] \Psi_{i,q^{-r}} (1 + \mathcal{B}_{i,r}),$$

where

$$\mathcal{B}_{i,r} = \sum_{k \geq 1} (-\mathcal{A}_{i,r})^k$$

is a formal power series in the $A_{j,a}^{-1}$ without constant term. In particular $\chi_q(D(\chi))$ is in \mathcal{E}_ℓ and we get a sum of the form

$$\chi_q(D(\chi)) = \sum_{1 \leq \alpha \leq R} \lambda_\alpha [\omega_\alpha] m_\alpha (1 + \mathcal{A}_\alpha) \in \mathcal{E}_\ell, \quad (6-16)$$

where ω_α is a weight, m_α is a Laurent monomial in the $\Psi_{i,q^{-r}}$, \mathcal{A}_α is a formal power series in the $A_{j,a}^{-1}$ without constant term and $\lambda_\alpha \in \mathbb{Z}$.

We recall the notion of negative ℓ -weight introduced in Definition 3.8.

Remark 6.3. We say that a sequence $(\Psi^{(m)})_{m \geq 0}$ of ℓ -weights converges pointwise as a rational fraction to an ℓ -weight Ψ if, for every $i \in I$ and $z \in \mathbb{C}$, the ratio $\Psi_i^{(m)}(z)/\Psi_i(z)$ converges to 1 when $N \rightarrow +\infty$ and $|q| > 1$. For example, defining the monomials $M_{i,r,N}$ as in equation (6-17) below, the sequence $(\widetilde{M}_{i,r,N})_{N \geq 0}$ converges pointwise as a rational fraction to $\Psi_{i,q^{-r}}^{-1}$.

Lemma 6.4. Let $\chi \in \mathcal{A}(G)$ be nonzero. Then at least one negative ℓ -weight occurs in $\chi_q(D(\chi))$.

Proof. We will use the following partial ordering \leq on the set of ℓ -weights Ψ satisfying $\varpi(\Psi) = 1$: for such ℓ -weights Ψ, Ψ' , we set $\Psi \geq \Psi'$ if

$$\Psi'(\Psi)^{-1} = \prod_{i,r \geq -M} \widetilde{A_{i,q^r}}^{-v_{i,r}}$$

is a possibly infinite product (that is, pointwise the limit of the partial products) with the $v_{i,r} \geq 0$. If $\Psi = \widetilde{m}$ and $\Psi' = \widetilde{m'}$ with m, m' monomials in \mathcal{M} , then $\Psi \leq \Psi'$ is equivalent to $m \leq m'$ for the partial ordering considered in [Nakajima 2004].

As the sum (6-16) is finite, there is α_0 such that m_{α_0} is maximal for \leq . We prove that m_{α_0} is a negative ℓ -weight.

Let N be such that all the cluster variables $z_{i,r}$ of the initial seed occurring in the Laurent monomials of equation (6-16) satisfy $r > -2d(N + 2)$, where $d = \text{Max}_{i \in I}(d_i)$ is the lacing number of \mathfrak{g} . We consider as above the semi-infinite

cluster algebra $\mathcal{A}(H_N^+)$ obtained from $\mathcal{A}(G)$, where the cluster variables sitting at $(i, r) \in V$, $r \leq -2d(N + 2)$ have been removed. As explained in the proof of Proposition 6.1, $\mathcal{A}(H_N^+)$ can be regarded as a subalgebra of $\mathcal{A}(G)$. We replace every cluster variable $z_{i,r}$ of the initial seed by the class of the Kirillov–Reshetikhin module $W_{i,r,N}$ of highest monomial

$$M_{i,r,N} = \prod_{\substack{k \geq 0 \\ r+2kd_i \leq 2dN}} Y_{i,q^{-r-d_i-2kd_i}}. \tag{6-17}$$

(Here $M_{i,r,N}$ is set to be 1 if $r > 2dN$.) We obtain

$$\phi_N(\chi) \in \text{Frac}(K_0(\mathbb{C})),$$

the image of χ in the fraction field of $K_0(\mathbb{C})$. By using the duality D , we reverse all spectral parameters (by (ii) in Example 5.14 illustrating Proposition 5.13, or by [Hernandez and Leclerc 2010, Section 3.4]). We obtain the same³ as in [Hernandez and Leclerc 2016, Section 2.2.2]. Then by [Hernandez and Leclerc 2016, Theorem 5.1], $D(\phi_N(\chi))$ belongs to the Grothendieck ring of \mathbb{C} . So by applying D again, $\phi_N(\chi)$ is in $K_0(\mathbb{C})$.

Now we get, as for equation (6-16),

$$\chi_q(\phi_N(\chi)) = \sum_{1 \leq \alpha \leq R} \lambda_\alpha m_\alpha^{(N)} (1 + \mathcal{A}_\alpha^{(N)}), \tag{6-18}$$

where $m_\alpha^{(N)}$ is a monomial and $\mathcal{A}_\alpha^{(N)}$ is a formal power series in the $A_{i,a}^{-1}$. Note that, when $N \rightarrow +\infty$, $\mathcal{A}_\alpha^{(N)}$ converges to \mathcal{A}_α as a formal power series in the $A_{i,a}^{-1}$. Let $\omega_{i,r,N}$ be the highest weight of $W_{i,r,N}$. Now if the initial cluster variable $z_{i,r}$ is replaced by

$$\left[\left(\frac{r}{2d_i} \right) \omega_i - \omega_{i,r,N} \right] [W_{i,r,N}] \tag{6-19}$$

instead of $[W_{i,r,N}]$, we just have to replace in (6-18) each $m_\alpha^{(N)}$ by $[\omega_\alpha](m_\alpha^{(N)})^\sim$ (the $\mathcal{A}_\alpha^{(N)}$ are unchanged). Then $(m_\alpha^{(N)})^\sim$ converges to m_α when $N \rightarrow +\infty$ pointwise as a rational fraction.

Let us show that there are infinitely many N such that $(m_{\alpha_0}^{(N)})^\sim$ is maximal among the $(m_\alpha^{(N)})^\sim$ for \leq . Otherwise, since equation (6-16) has finitely many summands, there is α such that $(m_{\alpha_0}^{(N)})^\sim < (m_\alpha^{(N)})^\sim$ for infinitely many N . In the limit, we get that $m_{\alpha_0} < m_\alpha$, contradiction.

For N such that $(m_{\alpha_0}^{(N)})^\sim$ is maximal for \leq , $m_{\alpha_0}^{(N)}$ is necessarily dominant, as $\phi_N(\chi) \in K_0(\mathbb{C})$ (see [Frenkel and Mukhin 2001, Section 5.4]). Then the limit m_{α_0} of the $(m_{\alpha_0}^{(N)})^\sim$ is negative, as it is easy to check that a limit of dominant monomials

³We have a term $-d_i$ which does not occur in [Hernandez and Leclerc 2016, Section 2.2.2], as we use the labeling by V and not by W .

is a negative ℓ -weight. Finally, since m_{α_0} is maximal for \preceq , it necessarily occurs with a nonzero coefficient in the expansion of $\chi_q(D(\chi))$. \square

Lemma 6.5. *Let Ψ be a negative ℓ -weight such that the roots and the poles of $\Psi_i(z)$ are of the form q^r with $(i, r) \in V$. Then there is a unique $F(\Psi) \in \chi_q(D(K_0(\mathcal{O}_{2\mathbb{Z}}^+)))$ such that Ψ is the unique negative ℓ -weight occurring in $F(\Psi)$ and its coefficient is 1.*

Moreover $F(\Psi)$ is of the form

$$F(\Psi) = [\Psi] + \sum_{\Psi': \varpi(\Psi') < \varpi(\Psi)} \lambda_{\Psi'} [\Psi'], \tag{6-20}$$

for the usual partial ordering on weights and with the $\lambda_{\Psi'} \in \mathbb{Z}$.

Proof. The uniqueness follows from Proposition 6.1 and Lemma 6.4. For each negative ℓ -weight Ψ as in the lemma, there is a representation $M(\Psi)$ in $\mathcal{O}_{2\mathbb{Z}}^+$ such that $\chi_q(D([M(\Psi)]))$ is of the form

$$\chi_q(D([M(\Psi)])) = [\Psi] + \sum_{\Psi': \varpi(\Psi') < \varpi(\Psi)} \mu_{\Psi', \Psi} [\Psi'].$$

Indeed it suffices to consider a tensor product of fundamental and positive pre-fundamental representations. Now if the $F(\Psi)$ do exist, we have an infinite triangular transition matrix from the $(F(\Psi))$ to the $(\chi_q(D([M(\Psi)]))$ with 1 on the diagonal and whose off-diagonal coefficients are the $\mu_{\Psi', \Psi}$ for Ψ, Ψ' negative. So to prove the existence, it suffices to consider the inverse of this matrix (which is well-defined, as for given Ψ, Ψ' there is a finite number of Ψ'' satisfying $\varpi(\Psi') \preceq \varpi(\Psi'')$ and $\mu_{\Psi'', \Psi} \neq 0$). \square

We can now finish the proof of Proposition 6.2:


Proof. Let χ be in $\mathcal{A}(G)$. For Ψ a negative ℓ -weight, we denote by λ_{Ψ} the coefficient of Ψ in $\chi_q(D(\chi))$. Then by Lemma 6.4 we have

$$\chi_q(D(\chi)) = \sum_{\Psi \text{ negative}} \lambda_{\Psi} F(\Psi).$$

As the $F(\Psi)$ are of the form (6-20), this sum is well-defined in $\chi_q(D(K_0(\mathcal{O}_{2\mathbb{Z}}^+)))$ and we get $\chi \in K_0(\mathcal{O}_{2\mathbb{Z}}^+)$. \square

Example 6.6. Let us illustrate the proof of Lemma 6.4, which is the crucial technical point for the proof of Proposition 6.2. Consider the sequence of mutations of 6A1.


Let us write the cluster variables $\phi_N(\chi)$:




The cluster variable corresponding to $[L(Y_q)]$ has its q -character, which can be written in the form of equation (6-16):

$$\begin{aligned} \chi_q(L(Y_q)) &= Y_q + Y_{q^3}^{-1} = \frac{\chi_q(W_{N-1, q^{1-2N}}) + \chi_q(W_{N+1, q^{1-2N}})}{\chi_q(W_{N, q^{1-2N}})} \\ &= Y_q^{-1} \frac{1 + A_{q^{-2}}^{-1} (1 + A_{q^{-4}}^{-1} (1 + \dots (1 + A_{q^{2(1-N)}}^{-1})) \dots)}{1 + A_1^{-1} (1 + A_{q^{-2}}^{-1} (1 + \dots (1 + A_{q^{2(1-N)}}^{-1})) \dots)} \\ &\quad + Y_q \frac{1 + A_{q^2}^{-1} (1 + A_1^{-1} (1 + \dots (1 + A_{q^{2(1-N)}}^{-1})) \dots)}{1 + A_1^{-1} (1 + A_{q^{-2}}^{-1} (1 + \dots (1 + A_{q^{2(1-N)}}^{-1})) \dots)}. \end{aligned}$$


We see as in the statement of Lemma 6.4 that the monomial Y_q maximal for \leq is dominant and so negative in the sense of Definition 3.8. In the limit $N \rightarrow +\infty$ (with the renormalized weights) we get the representations in $K_0(\mathcal{O}^-)$:



The duality again gives the representations in $K_0(\mathcal{O}^+)$:



Example 6.7. We now illustrate the proof of Lemma 6.4 by means of the mutation of Section 6A2. Let us write the cluster variables $\phi_N(\chi)$:




Here $m^{(N)} = Y_{1,q}(Y_{2,q^{2(1-N)}}Y_{2,q^{4-2N}} \cdots Y_{2,q^{-2}})$. The q -character corresponding to the cluster variable $[L(m^{(N)})]$ can be written in the form of equation (6-16):


$$\begin{aligned} \chi_q(L(m^{(N)})) &= \frac{\chi_q(W_{N-1,q}^{(1)})\chi_q(W_{N,q}^{(2)})}{\chi_q(W_{N,q}^{(1)})} + \frac{\chi_q(W_{N+1,q}^{(1)})\chi_q(W_{N-1,q}^{(2)})}{\chi_q(W_{N,q}^{(1)})} \\ &= Y_{1,q}^{-1}(Y_{2,q^{2(1-N)}}Y_{2,q^{4-2N}} \cdots Y_{2,1})(1 + \mathcal{A}_1^{(N)}) \\ &\quad + Y_{1,q}(Y_{2,q^{2(1-N)}}Y_{2,q^{4-2N}} \cdots Y_{2,q^{-2}})(1 + \mathcal{A}_2^{(N)}). \end{aligned}$$

Here $\mathcal{A}_1^{(N)}$ and $\mathcal{A}_2^{(N)}$ are formal power series in the A_{1,q^a}^{-1} , A_{2,q^a}^{-1} ($a \in \mathbb{C}^*$) without constant term. The monomial $(m^{(N)})^\sim$ is maximal for \preceq . Its limit for $N \rightarrow +\infty$ is $[-\omega_1]Y_{1,q}\Psi_{2,q^{-1}}^{-1}$, which is negative in the sense of Definition 3.8.

Example 6.8. In this example we check that the images of the initial cluster variables considered in the proof of Lemma 6.4 do match. Let us consider type B_2 with the following initial seed and the initial cluster variables replaced by the $W_{i,r}$:



In the limit $N \rightarrow +\infty$ (with the renormalized weights) we get the images of the initial cluster variables:



7. Conjectures and evidence

7A. A conjecture. The concept of a monoidal categorification of a cluster algebra was introduced in [Hernandez and Leclerc 2016, Definition 2.1]. We say that a simple object S of a monoidal category is *real* if $S \otimes S$ is simple. Let us recall that a cluster monomial is a monomial in the cluster variables of a single cluster.

Definition 7.1. Let \mathcal{A} be a cluster algebra and let \mathcal{M} be an abelian monoidal category. We say that \mathcal{A} is a monoidal categorification of \mathcal{A} if there is an isomorphism between \mathcal{A} and the Grothendieck ring of \mathcal{M} such that the cluster monomials of \mathcal{A} are the classes of all the real simple objects of \mathcal{M} (up to invertibles).

See [Hernandez and Leclerc 2013, Section 2] for a discussion on applications of monoidal categorifications. In view of Theorem 4.2, it is natural to formulate the following conjecture.

Conjecture 7.2. *The isomorphism of Theorem 4.2 defines a monoidal categorification; that is, the cluster monomials in $\mathcal{A}(\Gamma)$ get identified with real simple objects in $\mathcal{O}_{2\mathbb{Z}}^+$ up to invertible representations.*

Remark 7.3. By using the duality in Proposition 5.13, the statements of Theorem 4.2 and Conjecture 7.2 can also be formulated in terms of the category $\mathcal{O}_{2\mathbb{Z}}^-$.

Note that Theorem 5.8 implies that all cluster monomials of the initial seed are identified with real simple objects, more precisely, with simple tensor products of positive prefundamental representations, in agreement with Conjecture 7.2. To give other evidence supporting Conjecture 7.2, we will use the results in the next subsection.

7B. Limiting characters. We will be using the dual category \mathcal{O}^* considered in [Hernandez and Jimbo 2012], whose definition we now recall.

Definition 7.4. Let \mathcal{O}^* be the category of Cartan-diagonalizable $U_q(\mathfrak{b})$ -modules V such that V^* is in category \mathcal{O} .

A $U_q(\mathfrak{b})$ -module V is said to be of lowest ℓ -weight $\Psi \in P_\ell$ if there is $v \in V$ such that $V = U_q(\mathfrak{b})v$ and the following hold:

$$U_q(\mathfrak{b})^- v = \mathbb{C}v, \quad \phi_{i,m}^+ v = \Psi_{i,m} v \quad (i \in I, m \geq 0).$$

For $\Psi \in P_\ell$, there exists up to isomorphism a unique simple $U_q(\mathfrak{b})$ -module $L'(\Psi)$ of lowest ℓ -weight Ψ . This module belongs to \mathcal{O}^* . More precisely, we have:

Proposition 7.5 [Hernandez and Jimbo 2012]. *For $\Psi \in P_\ell$, $(L'(\Psi))^* \simeq L(\Psi^{-1})$.*

We can also define as in Section 5A notions of characters and q -characters for \mathcal{O}^* .

We now explain that characters (resp. q -characters) of certain simple objects in the category \mathcal{O}^- can be obtained as limits of characters (resp. q -characters)

of finite-dimensional representations. This is known for negative prefundamental representations [Hernandez and Jimbo 2012].

Let $L(\Psi)$ be a simple module whose highest ℓ -weight can be written as a finite product

$$\Psi = [\omega] \times m \times \prod_{i \in I} \left(\prod_{r \geq -R_0} \Psi_{i,q^r}^{u_{i,q^r}} \right),$$

where $\omega \in P_{\mathbb{Q}}$, $R_0 \geq 0$, $u_{i,q^r} \leq 0$ and $m \in \mathcal{M}$ is a dominant monomial. For $R \geq R_0$, set

$$M_R = m \prod_{i \in I} \left(\prod_{\substack{r \geq -R_0 \\ r' \geq 0 \\ r-2d_i r' \geq -R}} Y_{i,q^{r-2d_i r'}}^{-u_{i,q^r}} \right). \tag{7-21}$$

Theorem 7.6. (1) *We have the limit as formal power series*

$$\tilde{\chi}_q(L(M_R)) \xrightarrow{R \rightarrow +\infty} \tilde{\chi}_q(L(\Psi)) \in \mathbb{Z} \llbracket A_{i,a}^{-1} \rrbracket_{i \in I, a \in \mathbb{C}^*}.$$

(2) *We have $\tilde{\chi}(L(\bar{\Psi}^{-1})) = \tilde{\chi}(L(\Psi))$ and so we have the limit as formal power series*

$$\tilde{\chi}(L(M_R)) \xrightarrow{R \rightarrow +\infty} \tilde{\chi}(L(\bar{\Psi}^{-1})) \in \mathbb{Z} \llbracket -\alpha_i \rrbracket_{i \in I}.$$

The proof of Theorem 7.6 is essentially the same as that of [Hernandez and Jimbo 2012, Theorem 6.1], so we just give an outline.

Proof. First let us prove that the dimensions of weight spaces of $\tilde{L}(\Psi)$ are larger than those of $\tilde{L}(M_R)$. Consider the tensor product

$$T = \tilde{L}(\Psi) \otimes \tilde{L}(M_R \Psi^{-1}).$$

By definition of M_R , the ℓ -weight $M_R \Psi^{-1}$ is a product of Ψ_{i,q^r}^+ times $[\lambda]$ for some $\lambda \in P_{\mathbb{Q}}$, so by Theorem 5.8 the module $\tilde{L}(M_R \Psi^{-1})$ is a tensor product of positive fundamental representations. Moreover T and $\tilde{L}(M_R)$ have the same highest ℓ -weight, so $\tilde{L}(M_R)$ is a subquotient of T . By [Frenkel and Mukhin 2001, Theorem 4.1], each ℓ -weight of $\tilde{L}(M_R)$ is the product of the highest ℓ -weight $M_R(\varpi(M_R))^{-1}$ by a product of $A_{j,b}^{-1}$, $j \in i$, $b \in \mathbb{C}^*$. Hence, by Theorem 5.3, an ℓ -weight of T is an ℓ -weight of $\tilde{L}(M_R)$ only if it is of the form

$$\Psi' \varpi(\Psi)^{-1} (\tilde{M}_R \tilde{\Psi}^{-1}),$$

where Ψ' is an ℓ -weight of $L(\Psi)$ and $\tilde{M}_R \tilde{\Psi}^{-1}$ is the highest ℓ -weight of $\tilde{L}(M_R \Psi^{-1})$. We get the result for the dimensions.

Then we prove as in [Hernandez and Jimbo 2012, Section 4.2] that we can define an inductive linear system

$$L(M_0) \rightarrow L(M_1) \rightarrow \dots \rightarrow L(M_R) \rightarrow L(M_{R+1}) \rightarrow \dots$$

from the $L(M_R)$ so that we have the convergence of the action of the subalgebra $\tilde{U}_q(\mathfrak{g})$ of $U_q(\mathfrak{g})$ generated by the $x_{i,r}^+$ and the $k_i^{-1}x_{i,r}^-$. We get a limiting representation of $\tilde{U}_q(\mathfrak{g})$ from which one can construct a representation L_1 of $U_q(\mathfrak{b})$ in the category \mathcal{O} and a representation L_2 of $U_q(\mathfrak{b})$ in the category \mathcal{O}^* [Hernandez and Jimbo 2012, Proposition 2.4]. Moreover, L_1 (resp. L_2) is of highest (resp. lowest) ℓ -weight Ψ (resp. $\bar{\Psi}$).

By construction, the normalized q -character of L_1 is the limit of the normalized q -characters $\tilde{\chi}_q(L(M_R))$ as formal power series. Combining with the result of the first paragraph of this proof, the representation L_1 is necessarily simple isomorphic to $L(\Psi)$. We have proved the first statement in the theorem.

Now, by construction L_2^* is in the category \mathcal{O} with highest ℓ -weight $\bar{\Psi}^{-1}$ and satisfies $\tilde{\chi}(L_2^*) = \tilde{\chi}(L_1)$. To conclude, it suffices to prove that L_2^* is irreducible. This is proved as in [Hernandez and Jimbo 2012, Theorem 6.3]. \square

We have the following application:

Theorem 7.7. *Let $L(\Psi)$ be a simple module in the category \mathcal{O}^- such that $\tilde{\Psi} = \Psi$. Then its image by D^{-1} in $K_0(\mathcal{O}^+)$ is simple equal to $D^{-1}([L(\Psi)]) = [L(\bar{\Psi}^{-1})]$.*

Proof. From Example 5.14(i), the property is satisfied by negative prefundamental representations. Since these representations generate the fraction field of $K_0(\mathcal{O}^-)$, it suffices to show that the assignment $[L(\Psi)] \mapsto [L(\bar{\Psi}^{-1})]$ for ℓ -weights Ψ satisfying $\Psi = \tilde{\Psi}$ is multiplicative (recall that D is a morphism of \mathcal{E} -algebras). Let us use the same notation as in the proof of Theorem 7.6 above. For $L(\Psi) \simeq L_1$ and $L(\Psi') \simeq L'_1$ simple modules in \mathcal{O}^- with $\tilde{\Psi} = \Psi$ and $\tilde{\Psi}' = \Psi'$, we have the corresponding modules L_2, L'_2 in \mathcal{O}^* . We consider the decomposition

$$[L(\Psi) \otimes L(\Psi')] = \sum_{\Psi'', \tilde{\Psi}'' = \Psi''} m_{\Psi''} [L(\Psi'')]$$

in $K_0(\mathcal{O}^-)$ with $m_{\Psi''} \in \mathcal{E}$. Each $\chi_q(L(\Psi''))$ is obtained as a limit as in Theorem 7.6, and by construction the corresponding modules $L'(\Psi'')$ in the category \mathcal{O}^* satisfy

$$[L_2 \otimes L'_2] = \sum_{\Psi'', \tilde{\Psi}'' = \Psi''} m'_{\Psi''} [L'(\bar{\Psi}'')],$$

where each $m'_{\Psi''} \in \mathcal{E}$ is obtained from $m_{\Psi''}$ via the substitution $[\omega] \mapsto [-\omega]$. This implies

$$[L_2^* \otimes (L'_2)^*] = \sum_{\Psi'', \tilde{\Psi}'' = \Psi''} m_{\Psi''} [(L'(\bar{\Psi}''))^*],$$

that is, in view of Proposition 7.5,

$$[L(\bar{\Psi}^{-1})] \otimes [L(\bar{\Psi}'^{-1})] = \sum_{\Psi'', \tilde{\Psi}'' = \Psi''} m_{\Psi''} [L(\bar{\Psi}''^{-1})]. \quad \square$$

Example 7.8. Applying D to (6-14), we get the relation in $K_0(\mathcal{O}^-)$

$$[L(\bar{\Psi}^{-1}) \otimes L_{i,q^{-r}}^-] = \left[\bigotimes_{j:C_{j,i} \neq 0} L_{j,q}^{-r+d_j C_{j,i}} \right] + [-\alpha_i] \left[\bigotimes_{j:C_{j,i} \neq 0} L_{j,q}^{-r-d_j C_{j,i}} \right],$$

where

$$\bar{\Psi}^{-1} = [-\omega_i] Y_{i,q^{-r+d_i}} \prod_{j:C_{j,i} < 0} \Psi_{j,q}^{-r+d_j C_{j,i}}.$$

7C. Proof of Conjecture 7.2 for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$. In this section we give an explicit description of all simple modules in \mathcal{O}^+ and in \mathcal{O}^- for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$.

A q -set is a subset of \mathbb{C}^* of the form $\{aq^{2r} \mid R_1 \leq r \leq R_2\}$ for some $a \in \mathbb{C}^*$ and $R_1 \leq R_2 \in \mathbb{Z} \cup \{-\infty, +\infty\}$. The KR-modules $W_{k,a}, W_{k',b}$ are said to be in *special position* if the union of $\{a, aq^2, \dots, aq^{2(k-1)}\}$ and $\{b, bq^2, \dots, bq^{2(k'-1)}\}$ is a q -set which contains both properly. The KR-module $W_{k,a}$ and the prefundamental representation L_b^+ are said to be in special position if the union of $\{a, aq^2, aq^4, \dots, aq^{2(k-1)}\}$ and $\{bq, bq^3, bq^5, \dots\}$ is a q -set which contains both properly. Two positive prefundamental representations are never in special position. Two representations are in *general position* if they are not in special position.

The invertible elements in the category \mathcal{O}^+ are the 1-dimensional representations $[\omega]$.

Theorem 7.9. *Suppose that $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$. The prime simple objects in the category \mathcal{O}^+ are the positive prefundamental representations and the KR-modules (up to invertibles). Any simple object in \mathcal{O}^+ can be factorized in a unique way as a tensor product of prefundamental representations and KR-modules (up to permutation of the factors and to invertibles). Moreover, such a tensor product is simple if and only if all its factors are pairwise in general position.*

Proof. As in the classical case of finite-dimensional representations, it is easy to check that every positive ℓ -weight has a unique factorization as a product of highest ℓ -weights of KR-modules and positive prefundamental representations in pairwise general position. Hence it suffices to prove the equivalence in the last sentence. By [Chari and Pressley 1994], the result is known for finite-dimensional representations. Now, by using Section 7B, this result implies that a tensor product with factors which are in general position is simple. Conversely, it is known that a tensor product of KR-modules which are in special position is not simple. Also, it is easy to see that the tensor product of a KR-module and a positive prefundamental representation which are in special position is not simple. □

Remark 7.10. (i) This is a generalization of the factorization of simple representations in \mathcal{C} when $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ [Chari and Pressley 1994].

- (ii) This result for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$ implies that all simple objects in \mathcal{O}^+ are real and that their factorization into prime representations is unique.
- (iii) In [Mukhin and Young 2014], a factorization is proved for simple modules in $\hat{\mathcal{O}}$ when $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$. But the factorization is not unique in the category $\hat{\mathcal{O}}$; see Remark 3.10.
- (iv) By Proposition 5.13, our result implies a similar factorization in the category \mathcal{O}^- .
- (v) The combinatorics of q -sets in pairwise general position is very similar to the combinatorics of triangulations of the ∞ -gon studied in [Grabowski and Gratz 2014], in relation with certain cluster structures of infinite rank. However, in that paper only arcs (m, n) joining two integers m and n are considered, whereas we also allow arcs of the form $(m, +\infty)$ corresponding to positive prefundamental representations. Also, we are only interested in one mutation class, namely the mutation class of the initial triangulation $\{(m, +\infty) \mid m \in \mathbb{Z}\}$.

Theorem 7.11. *Conjecture 7.2 is true in the \mathfrak{sl}_2 -case.*

Proof. Theorem 7.9 provides an explicit factorization of simple objects in $\mathcal{O}_{2\mathbb{Z}}^+$ into positive prefundamental representations and finite-dimensional KR-modules. In particular, we get an explicit q -character formula for such a simple object and so a complete explicit description of the Grothendieck ring $K_0(\mathcal{O}_{2\mathbb{Z}}^+)$. The cluster algebra $\mathcal{A}(\Gamma)$ can also be explicitly described by using triangulations of the ∞ -gon (see Remark 7.10). Hence we can argue as in [Hernandez and Leclerc 2010, Section 13.4]. □

7D. Equivalence of conjectures. In general, we have the following:

Theorem 7.12. *Conjecture 7.2 is equivalent to Conjecture 5.2 of [Hernandez and Leclerc 2016].*

Combining with the recent results in [Qin 2015], this would imply a part of Conjecture 7.2 for ADE types, namely that all cluster monomials are classes of real simple objects.

As recalled in the introduction, [Hernandez and Leclerc 2016, Conjecture 5.2] states that $\mathcal{C}_{\mathbb{Z}}^-$ is the monoidal categorification of a cluster algebra $\mathcal{A}(G^-)$. Note that $\mathcal{C}_{\mathbb{Z}}^-$ is a subcategory of $\mathcal{O}_{2\mathbb{Z}}^-$ and $\mathcal{O}_{2\mathbb{Z}}^+$.

Proof. For $N > 0$, let \mathcal{C}_N be the category of finite-dimensional $U_q(\mathfrak{g})$ -modules V satisfying

$$[V] \in \mathbb{Z} \llbracket V_{i,q^m} \rrbracket_{i \in I, -2dN - d_i \leq m < d(N+2) - d_i} \subset K_0(\mathcal{C}).$$

It is a monoidal category similar to the categories considered in [Hernandez and Leclerc 2010]. It contains the KR-module $W_{i,r,N}$ with highest monomial $M_{i,r,N}$ given in equation (6-17), where $i \in I$ and $-d(N+2) < r \leq 2dN$. The Grothendieck

ring $K_0(\mathcal{C}_N)$ has a cluster algebra structure with an initial seed consisting of these KR-modules $W_{i,r,N}$ (here we use the initial seed as in [Hernandez and Leclerc 2016]). We have established in the proof of Theorem 4.2 that $K_0(\mathcal{C}_N) \otimes \mathcal{E}$ may be seen as a subalgebra of $K_0(\mathcal{O}_{2\mathbb{Z}}^+)$ by using the identification of $z_{i,r}$ with the element defined in equation (6-19). This induces embeddings $K_0(\mathcal{C}_N) \subset K_0(\mathcal{C}_{N+1})$,

$$K_0(\mathcal{C}_1) \subset K_0(\mathcal{C}_2) \subset K_0(\mathcal{C}_3) \subset \cdots \subset K_0(\mathcal{O}_{2\mathbb{Z}}^+),$$

which are not the naive embeddings obtained from the inclusion of categories $\mathcal{C}_N \subset \mathcal{C}_{N+1}$. The cluster monomials in $K_0(\mathcal{C}_N)$ corresponds now to cluster monomials in $K_0(\mathcal{O}_{2\mathbb{Z}}^+)$.

Note that by Theorem 7.7 we may consider simultaneously the statement of Conjecture 7.2 for $\mathcal{O}_{2\mathbb{Z}}^+$ or for $\mathcal{O}_{2\mathbb{Z}}^-$.

Suppose that [Hernandez and Leclerc 2016, Conjecture 5.2] is true. This implies that the cluster monomials in $K_0(\mathcal{C}_N)$ are the real simple modules for any $N > 0$. Consider a cluster monomial in $\mathcal{O}_{2\mathbb{Z}}^+$. Then for N large enough, we have a corresponding real representation V_N in $K_0(\mathcal{C}_N)$. The highest monomial of V_N is a Laurent monomial in the $m_{i,r,N}$ of the form considered in Theorem 7.6. By Theorem 7.6, $\chi_q(V_N)$ converges to the q -character of a simple module V in $\mathcal{O}_{2\mathbb{Z}}^-$ when $N \rightarrow +\infty$. Moreover V is real, as the q -character of $V \otimes V$ is obtained as a limit of simple q -character $\chi_q(V_N \otimes V_N)$ by Theorem 7.6. Conversely, every real simple module V in $\mathcal{O}_{2\mathbb{Z}}^-$ is obtained as such a limit of simple modules V_N . Moreover since V is real, V_N is real (since $\chi_q(V_N \otimes V_N)$ is an upper q -character of $V \otimes V$ in the sense of [Hernandez 2010, Corollary 5.8]). For N large enough the modules V_N correspond to the same cluster monomial, which is therefore identified with V .

Conversely, suppose that Conjecture 7.2 is true. Consider a cluster monomial χ in $K_0(\mathcal{C}_{\mathbb{Z}}^-)$. The cluster variables occurring in χ are produced via sequences of mutations from a finite number of KR-modules in the initial seed. By the proof of Proposition 6.1, there is a seed in $K_0(\mathcal{O}_{2\mathbb{Z}}^-)$ containing these KR-modules (and the quiver of this seed has the same arrows joining the corresponding vertices). By our hypothesis, χ is the class of a simple real module as an element of $K_0(\mathcal{O}_{2\mathbb{Z}}^-) \supset K_0(\mathcal{C}_{\mathbb{Z}}^-)$. Hence it is also real simple in $K_0(\mathcal{C}_{\mathbb{Z}}^-)$. Now consider a real simple module $[V]$ in $K_0(\mathcal{C}_{\mathbb{Z}}^-)$. It corresponds to a cluster monomial in $K_0(\mathcal{O}_{2\mathbb{Z}}^-)$ which is a cluster monomial in $K_0(\mathcal{C}_{\mathbb{Z}}^-)$ by the same arguments. \square

7E. Web property theorem. Let us prove the following generalization of the main result of [Hernandez 2010]. If Conjecture 7.2 holds, then the statement of the next theorem is a necessary condition for simple modules in the same seed.

Theorem 7.13. *Let S_1, \dots, S_N be simple objects in \mathcal{O}^+ (resp. in \mathcal{O}^-). Then $S_1 \otimes \cdots \otimes S_N$ is simple if and only if the tensor products $S_i \otimes S_j$ are simple for $i \leq j$.*

Proof. By Proposition 5.13, this is equivalent to proving the statement in the category \mathcal{O}^- . Note that the “only if” part is clear. For the “if” part of the statement, we may assume without loss of generality that the zeros and poles of the highest ℓ -weights of the S_i are in $q^{\mathbb{Z}}$ (see (ii) in Remark 4.3). For each simple module S_i , consider a corresponding simple finite-dimensional module $L(M_{R,i})$ as in Section 7B. Since $S_i \otimes S_j$ is simple, there exists R_1 such that for $R \geq R_1$ the tensor product $L(M_{R,i}) \otimes L(M_{R,j})$ is simple. Indeed, by Theorem 7.6, $\tilde{\chi}_q(S_i \otimes S_j)$ is the limit of the $\tilde{\chi}_q(L(M_{R,i}M_{R,j}))$. More precisely, there is R_1 such that for $R \geq R_1$ the image of $\tilde{\chi}_q(S_i \otimes S_j)$ in $\mathbb{Z}[A_{i,q^r}^{-1}]_{i \in I, r \geq -R_1+r_i}$ is equal to $\tilde{\chi}_q(L(M_{R,i}M_{R,j}))$. This implies that $\tilde{\chi}_q(L(M_{R,i}M_{R,j})) = \tilde{\chi}_q(L(M_{R,i}))\tilde{\chi}_q(L(M_{R,j}))$.

Now, by [Hernandez 2010], $L(M_{R,1}) \otimes \cdots \otimes L(M_{R,N})$ is simple isomorphic to $L(M_{R,1} \cdots M_{R,N})$. This implies that the character of $S_1 \otimes \cdots \otimes S_N$ is the same as the character of the simple module with the same highest ℓ -weight. Hence they are isomorphic. \square

Remark 7.14. This provides an alternative proof of Theorem 7.9.

7F. Another conjecture. To conclude, let us state another general conjecture. Although the cluster algebra structure presented in this paper does not appear in the statement, this conjecture arises naturally if we compare Theorem 4.2 with the results of [Hernandez and Leclerc 2016].

We consider a simple finite-dimensional representation $L(m)$ whose dominant monomial m satisfies $m \in \mathbb{Z}[Y_{i,aq^r}]_{(i,r) \in W, r \leq R}$ for a given $R \in \mathbb{Z}$. We have the corresponding truncated q -character [Hernandez and Leclerc 2010]

$$\chi_q^{\leq R}(L(m)) \in m\mathbb{Z}[A_{i,q^r}^{-1}]_{i \in I, r \leq R-d_i},$$

which is the sum (with multiplicity) of the monomials m' occurring in $\chi_q(L(m))$ and satisfying $m(m')^{-1} \in \mathbb{Z}[A_{i,q^r}]_{i \in I, r \leq R-d_i}$. As in the statement of Theorem 5.5, we consider

$$(\chi^+)^{\leq R}(L(m)) \in \text{Frac}(K_0(\mathcal{O}^+)),$$

obtained from $\chi_q^{\leq R}(L(m))$ by replacing each variable $Y_{i,a}$ by $[\omega_i]_{i,aq_i}^+ (\ell_{i,aq_i}^+)^{-1}$.

Let us set

$$W_R = \{(i, r) \in W \mid R \geq r > R - 2d_i\}.$$

For $(i, r) \in W_R$, we set $u_{i,r}$ to be the maximum of 0 and the powers $u_{i,q^r}(m')$ of Y_{i,q^r} in all monomials m' occurring in $\chi_q^{\leq R}(L(m))$. Let

$$\Psi_R = \prod_{(i,r) \in W_R} \Psi_{i,q^{r+d_i}}^{u_{i,r}} \quad \text{and} \quad \Psi = m\Psi_R.$$

The representations $L(\Psi)$, $L(\Psi_R)$ are in the category \mathcal{O}^+ . By Theorem 5.8, $L(\Psi_R)$ is a simple tensor product of positive prefundamental representations.

Conjecture 7.15. *We have the relation in $\text{Frac}(K_0(\mathcal{O}^+))$*

$$\chi^+(L(\Psi)) = (\chi^+)^{\leq R}(L(m)) \prod_{(i,r) \in W_R} (\ell_{i,q^{r+d_i}}^+)^{u_{i,r}} = (\chi^+)^{\leq R}(L(m)) \chi^+(L(\Psi_R)).$$

Remark 7.16. By taking the q -character, the statement is equivalent to the following q -character formula:

$$\chi_q(L(\Psi)) = \chi_q^{\leq R}(L(m)) \chi_q(L(\Psi_R)).$$

In some cases the conjecture is already proved:

Example 7.17. (i) In the case $\chi_q^{\leq R}(L(m)) = \chi_q(L(m))$, we have $u_{i,r} = 0$ for $(i,r) \in W_R$. Here the conjecture reduces to the generalized Baxter relations of Theorem 5.5.

(ii) In the case $R = r + d_i$ and $m = Y_{i,q^{r-d_i}}$, we have $\chi_q(L(m))^{\geq R} = m(1 + A_{i,q^r}^{-1})$. The conjecture reduces to the relations we have established in formula (6-14).

(iii) As discussed above, it follows from Theorem 7.12 and from the main result of [Qin 2015] that, for ADE -types, all cluster monomials are classes of real simple objects. In particular, for ADE -types, Conjecture 7.15 holds for all simple modules $L(m)$ which are cluster monomials (for the cluster algebra structure defined in [Hernandez and Leclerc 2016]). Indeed we may assume that $R = 0$. It is proved in the same paper that, for any dominant monomial m , the truncated q -character $\chi_q^{\leq 0}(L(m))$ is an element of the cluster algebra $\mathcal{A}(G^-)$ defined in the proof of Proposition 6.1. In $\chi_q^{\leq 0}(L(m))$ we perform the same substitution as in Theorem 5.5 above (that is, we apply the ring homomorphism F of the proof of Proposition 6.1). If we assume that $\chi_q^{\leq 0}(L(m))$ is a cluster variable of $\mathcal{A}(G^-)$, then it follows from the proof of Proposition 6.1 that

$$y := F(\chi_q^{\leq 0}(L(m))) \prod_{(i,r) \in W_0} z_{i,q^{r+d_i}}^{u_{i,r}}$$

is a cluster variable in $\mathcal{A}(H^-)$. Using Theorem 7.12, we deduce that y is the positive prefundamental character of a simple module. Since F is multiplicative, the argument readily extends to simple modules $L(m)$ such that $\chi_q^{\leq 0}(L(m))$ is a cluster monomial.

References

[Baxter 1972] R. J. Baxter, “Partition function of the eight-vertex lattice model”, *Ann. Physics* **70** (1972), 193–228. MR 0290733 Zbl 0236.60070

[Bazhanov et al. 1997] V. V. Bazhanov, S. L. Lukyanov, and A. B. Zamolodchikov, “Integrable structure of conformal field theory, II: Q-operator and DDV equation”, *Comm. Math. Phys.* **190**:2 (1997), 247–278. MR 1489571 Zbl 0908.35114

- [Bazhanov et al. 2002] V. V. Bazhanov, A. N. Hibberd, and S. M. Khoroshkin, “Integrable structure of \mathcal{W}_3 conformal field theory, quantum Boussinesq theory and boundary affine Toda theory”, *Nuclear Phys. B* **622**:3 (2002), 475–547. MR 1880098 Zbl 0983.81088
- [Beck 1994] J. Beck, “Braid group action and quantum affine algebras”, *Comm. Math. Phys.* **165**:3 (1994), 555–568. MR 1301623 Zbl 0807.17013
- [Boos et al. 2009] H. Boos, M. Jimbo, T. Miwa, F. Smirnov, and Y. Takeyama, “Hidden Grassmann structure in the XXZ model, II: Creation operators”, *Comm. Math. Phys.* **286**:3 (2009), 875–932. MR 2472021 Zbl 1173.82003
- [Chari 1995] V. Chari, “Minimal affinizations of representations of quantum groups: the rank 2 case”, *Publ. Res. Inst. Math. Sci.* **31**:5 (1995), 873–911. MR 1367675 Zbl 0855.17010
- [Chari and Pressley 1994] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge University Press, 1994. MR 1300632 Zbl 0839.17009
- [Damiani 1998] I. Damiani, “La R -matrice pour les algèbres quantiques de type affine non tordu”, *Ann. Sci. École Norm. Sup. (4)* **31**:4 (1998), 493–523. MR 1634087 Zbl 0911.17005
- [Drinfeld 1987] V. G. Drinfeld, “A new realization of Yangians and of quantum affine algebras”, *Dokl. Akad. Nauk SSSR* **296**:1 (1987), 13–17. In Russian; translated in *Soviet Math. Dokl.* **36**:2 (1988), 212–216. MR 914215
- [Fomin and Zelevinsky 2002] S. Fomin and A. Zelevinsky, “Cluster algebras, I: Foundations”, *J. Amer. Math. Soc.* **15**:2 (2002), 497–529. MR 1887642 Zbl 1021.16017
- [Fomin and Zelevinsky 2003] S. Fomin and A. Zelevinsky, “Cluster algebras: notes for the CDM-03 conference”, pp. 1–34 in *Current developments in mathematics, 2003*, edited by B. Mazur et al., International Press, Somerville, MA, 2003. MR 2132323 Zbl 1119.05108
- [Frenkel and Hernandez 2015] E. Frenkel and D. Hernandez, “Baxter’s relations and spectra of quantum integrable models”, *Duke Math. J.* **164**:12 (2015), 2407–2460. MR 3397389 Zbl 1332.82022
- [Frenkel and Mukhin 2001] E. Frenkel and E. Mukhin, “Combinatorics of q -characters of finite-dimensional representations of quantum affine algebras”, *Comm. Math. Phys.* **216**:1 (2001), 23–57. MR 1810773 Zbl 1051.17013
- [Frenkel and Reshetikhin 1999] E. Frenkel and N. Reshetikhin, “The q -characters of representations of quantum affine algebras and deformations of \mathcal{W} -algebras”, pp. 163–205 in *Recent developments in quantum affine algebras and related topics* (Raleigh, NC, 1998), edited by N. Jing and K. C. Misra, Contemp. Math. **248**, Amer. Math. Soc., Providence, RI, 1999. Corrected version available at <https://arxiv.org/abs/math/9810055>. MR 1745260 Zbl 0973.17015
- [Gekhtman et al. 2010] M. Gekhtman, M. Shapiro, and A. Vainshtein, *Cluster algebras and Poisson geometry*, Mathematical Surveys and Monographs **167**, American Mathematical Society, Providence, RI, 2010. MR 2683456 Zbl 1217.13001
- [Grabowski and Gratz 2014] J. E. Grabowski and S. Gratz, “Cluster algebras of infinite rank”, *J. Lond. Math. Soc. (2)* **89**:2 (2014), 337–363. MR 3188622 Zbl 1308.13034
- [Grabowski and Launois 2014] J. E. Grabowski and S. Launois, “Graded quantum cluster algebras and an application to quantum Grassmannians”, *Proc. Lond. Math. Soc. (3)* **109**:3 (2014), 697–732. MR 3260291 Zbl 1315.13036
- [Hernandez 2005] D. Hernandez, “Representations of quantum affinizations and fusion product”, *Transform. Groups* **10**:2 (2005), 163–200. MR 2195598 Zbl 1102.17009
- [Hernandez 2007] D. Hernandez, “On minimal affinizations of representations of quantum groups”, *Comm. Math. Phys.* **276**:1 (2007), 221–259. MR 2342293 Zbl 1141.17011
- [Hernandez 2010] D. Hernandez, “Simple tensor products”, *Invent. Math.* **181**:3 (2010), 649–675. MR 2660455 Zbl 1221.17015

- [Hernandez 2015] D. Hernandez, “Asymptotical representations, spectra of quantum integrable systems and cluster algebras”, pp. 1399–1402 in *Enveloping Algebras and Geometric Representation Theory*, 12 2, 2015. Abstracts from the workshop held May 10–16, 2015, Organized by Iain Gordon, Bernard Leclerc and Wolfgang Soergel.
- [Hernandez and Jimbo 2012] D. Hernandez and M. Jimbo, “Asymptotic representations and Drinfeld rational fractions”, *Compos. Math.* **148**:5 (2012), 1593–1623. MR 2982441 Zbl 1266.17010
- [Hernandez and Leclerc 2010] D. Hernandez and B. Leclerc, “Cluster algebras and quantum affine algebras”, *Duke Math. J.* **154**:2 (2010), 265–341. MR 2682185 Zbl 1284.17010
- [Hernandez and Leclerc 2013] D. Hernandez and B. Leclerc, “Monoidal categorifications of cluster algebras of type A and D ”, pp. 175–193 in *Symmetries, integrable systems and representations*, edited by K. Iohara et al., Springer Proc. Math. Stat. **40**, Springer, 2013. MR 3077685 Zbl 1317.13052
- [Hernandez and Leclerc 2016] D. Hernandez and B. Leclerc, “A cluster algebra approach to q -characters of Kirillov–Reshetikhin modules”, *J. Eur. Math. Soc. (JEMS)* **18**:5 (2016), 1113–1159. MR 3500832 Zbl 06578174
- [Kac 1990] V. G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, 1990. MR 1104219 Zbl 0716.17022
- [Kang et al. 2014] S.-J. Kang, M. Kashiwara, M. Kim, and S.-j. Oh, “Monoidal categorification of cluster algebras”, preprint, 2014. arXiv 1412.8106
- [Kang et al. 2015] S.-J. Kang, M. Kashiwara, M. Kim, and S.-j. Oh, “Monoidal categorification of cluster algebras, II”, preprint, 2015. arXiv 1502.06714
- [Kimura and Qin 2014] Y. Kimura and F. Qin, “Graded quiver varieties, quantum cluster algebras and dual canonical basis”, *Adv. Math.* **262** (2014), 261–312. MR 3228430 Zbl 1331.13016
- [Kojima 2008] T. Kojima, “Baxter’s Q -operator for the W -algebra W_N ”, *J. Phys. A* **41**:35 (2008), 355206, 16. MR 2426018 Zbl 1146.81032
- [Mukhin and Young 2014] E. Mukhin and C. A. S. Young, “Affinization of category \mathcal{O} for quantum groups”, *Trans. Amer. Math. Soc.* **366**:9 (2014), 4815–4847. MR 3217701 Zbl 1306.17005
- [Nakajima 2004] H. Nakajima, “Quiver varieties and t -analogs of q -characters of quantum affine algebras”, *Ann. of Math. (2)* **160**:3 (2004), 1057–1097. MR 2144973 Zbl 1140.17015
- [Nakajima 2011] H. Nakajima, “Quiver varieties and cluster algebras”, *Kyoto J. Math.* **51**:1 (2011), 71–126. MR 2784748 Zbl 1223.13013
- [Qin 2015] F. Qin, “Triangular bases in quantum cluster algebras and monoidal categorification conjectures”, preprint, 2015. arXiv 1501.04085

Communicated by Edward Frenkel

Received 2016-04-19 Accepted 2016-08-09

david.hernandez@imj-prg.fr

*Institut de Mathématiques de Jussieu-Paris Rive Gauche,
Université Paris-Diderot Paris 7, Bâtiment Sophie Germain,
Case 7012, 75205 Paris Cedex 13, France*

bernard.leclerc@unicaen.fr

*Laboratoire de Mathématiques Nicolas Oresme,
Université de Caen Basse-Normandie, 14032 Caen, France*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Susan Montgomery	University of Southern California, USA
Richard E. Borcherds	University of California, Berkeley, USA	Shigefumi Mori	RIMS, Kyoto University, Japan
John H. Coates	University of Cambridge, UK	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Roger Heath-Brown	Oxford University, UK	Ravi Vakil	Stanford University, USA
Craig Huneke	University of Virginia, USA	Michel van den Bergh	Hasselt University, Belgium
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Yuri Manin	Northwestern University, USA	Efim Zelmanov	University of California, San Diego, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2016 is US \$290/year for the electronic version, and \$485/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 10 No. 9 2016

Torsion pour les variétés abéliennes de type I et II MARC HINDRY and NICOLAS RATAZZI	1845
Galois-generic points on Shimura varieties ANNA CADORET and ARNO KRET	1893
Extremality of loci of hyperelliptic curves with marked Weierstrass points DAWEI CHEN and NICOLA TARASCA	1935
$\overline{\mathcal{R}}_{15}$ is of general type GREGOR BRUNS	1949
A vanishing theorem for weight-one syzygies LAWRENCE EIN, ROBERT LAZARSFELD and DAVID YANG	1965
Effective cones of cycles on blowups of projective space IZZET COSKUN, JOHN LESIEUTRE and JOHN CHRISTIAN OTTEM	1983
Cluster algebras and category \mathcal{O} for representations of Borel subalgebras of quantum affine algebras DAVID HERNANDEZ and BERNARD LECLERC	2015



1937-0652(2016)10:9;1-G