

Algebra & Number Theory

Volume 11

2017

No. 5

Some sums over irreducible polynomials

David E. Speyer



Some sums over irreducible polynomials

David E. Speyer

We prove a number of conjectures due to Dinesh Thakur concerning sums of the form $\sum_P h(P)$ where the sum is over monic irreducible polynomials P in $\mathbb{F}_q[T]$, the function h is a rational function and the sum is considered in the T^{-1} -adic topology. As an example of our results, in $\mathbb{F}_2[T]$, the sum $\sum_P 1/(P^k - 1)$ always converges to a rational function, and is 0 for $k = 1$.

1. Introduction

Our goal is to explain some identities experimentally discovered by Dinesh Thakur, involving sums over irreducible polynomials in finite fields. We begin by stating the simplest of these identities: Let \mathcal{P} be the set of irreducible polynomials in $\mathbb{F}_2[T]$. Then

$$\sum_{P \in \mathcal{P}} \frac{1}{P - 1} = 0.$$

Here the sum must be interpreted as a sum of power series in T^{-1} . For example, the first five summands are

$$\begin{aligned} \frac{1}{T - 1} &= T^{-1} + T^{-2} + T^{-3} + \dots \\ \frac{1}{(T + 1) - 1} &= T^{-1} \\ \frac{1}{(T^2 + T + 1) - 1} &= T^{-2} + T^{-3} + \dots \\ \frac{1}{(T^3 + T + 1) - 1} &= T^{-3} + \dots \\ \frac{1}{(T^3 + T^2 + 1) - 1} &= T^{-3} + \dots \end{aligned}$$

As the reader can see, only finitely many terms contribute to the coefficient of each power of T^{-1} , and the coefficient of T^{-j} is 0 for each j .

MSC2010: primary 11M38; secondary 05E05, 11M32.

Keywords: zeta function, special value, function field.

We now introduce the notation necessary to state our general results. To aid the reader’s comprehension, we adopt the following conventions: Integers will always be denoted by lower case Roman letters (k, p, q, \dots); polynomials over finite fields will always be denoted by capital Roman letters (A, F, P, \dots), sets of such polynomials will always be denoted by calligraphic letters ($\mathcal{A}, \mathcal{P}, \mathcal{R}, \dots$), symmetric polynomials will be denoted by bold letters ($\mathbf{e}_k, \mathbf{p}_k, \dots$). Of course, there will be other sorts of mathematical objects as well, which we trust the reader to accommodate as they occur.

Let p be a prime and q a power of p . Let \mathbb{F}_q be the field with q elements. Let \mathcal{R} be the polynomial ring $\mathbb{F}_q[T]$. Let \mathcal{K} be the fraction field $\mathbb{F}_q(T)$ and let $\widehat{\mathcal{K}}$ be the T^{-1} -adic completion of \mathcal{K} . All infinite sums will be understood in the T^{-1} -adic topology.

Let \mathcal{P} be the set of irreducible polynomials in \mathcal{R} ; let \mathcal{P}_1 be the set of monic irreducible polynomials. Here is our main result for the case $p = 2$.

Theorem 1.1. *If $p = 2$ then, for any positive integer $k \equiv 0 \pmod{q - 1}$, the sum*

$$\sum_{P \in \mathcal{P}_1} \frac{1}{P^k - 1}$$

is in \mathcal{K} .

The reader may wonder what happens if we sum over all irreducible polynomials rather than monic ones; that is an easy corollary:

Corollary 1.2. *Let $p = 2$. For any positive integer k , the sum*

$$\sum_{P \in \mathcal{P}} \frac{1}{P^k - 1}$$

is in \mathcal{K} .

Proof. We rewrite the sum as $\sum_{P \in \mathcal{P}_1} \sum_{a \in \mathbb{F}_q^\times} 1/((aP)^k - 1)$. The corollary then follows from the identity

$$\sum_{a \in \mathbb{F}_q^\times} \frac{1}{(aX)^k - 1} = \frac{1}{X^{\text{LCM}(k, q-1)} - 1}$$

in $\mathbb{F}_q(U)$. To prove this identity, write

$$\frac{1}{(aX)^k - 1} = \sum_{j=1}^{\infty} 1/(aX)^{kj}$$

and recall that

$$\sum_{a \in \mathbb{F}_q^\times} a^m = \begin{cases} 1, & m \equiv 0 \pmod{q - 1}, \\ 0, & \text{otherwise.} \end{cases}$$

□

We now discuss the case of a general prime. Define the rational function $G_p(U)$ by

$$G_p(U) = \frac{(1 - U^p) - (1 - U)^p}{p(1 - U)^p}.$$

When $p = 2$, we have $G_2(U) = (2U - 2U^2)/(2(1 - U)^2) = U/(1 - U)$, so $G_2(1/P) = 1/(P - 1)$. When p is odd, we have the following alternate expressions for G_p :

Proposition 1.3. *If p is odd, then, as rational functions in $\mathbb{F}_p(U)$, we have*

$$G_p(U) = \frac{\sum_{j=1}^{p-1} U^j/j}{(1 - U)^p} = \sum_{\substack{0 \leq j < \infty \\ j \not\equiv 0 \pmod p}} \frac{U^j}{j}.$$

Proof. If p is odd, then $(1 - U^p) - (1 - U)^p = \sum_{j=1}^{p-1} (-1)^{j-1} \binom{p}{j} U^j$. We have

$$\frac{(-1)^{j-1} \binom{p}{j}}{p} = \frac{(-1)^{j-1} (p-1)(p-2) \cdots (p-j+1)}{1 \cdots 2 \cdots (j-1)j} \equiv \frac{1}{j} \pmod p.$$

This proves the first equality, and the second is immediate. □

Theorem 1.4. *For any positive integer $k \equiv 0 \pmod{q-1}$, the sum*

$$\sum_{P \in \mathcal{P}_1} G_p(1/P^k)$$

is in \mathcal{K} .

As we noted, $G_2(1/X) = 1/(X - 1)$, so [Theorem 1.4](#) implies [Theorem 1.1](#).

Remark 1.5. When $p = 2$, we do *not* have $G_2(U) = \sum_{j \not\equiv 0 \pmod p} U^j/j$; the latter sum is $H(U) := U/(1 - U^2)$. However, it is true that $\sum_{P \in \mathcal{P}_1} H(1/P^k)$ is in \mathcal{K} , because $H(U) = G(U) - G(U^2)$.

Once again, we have a trivial variant where we sum over \mathcal{P} :

Corollary 1.6. *For any positive integer k , the sum*

$$\sum_{P \in \mathcal{P}} G_p(1/P^k)$$

is in \mathcal{K} .

Proof. If $p = 2$, we proved this in [Corollary 1.2](#), so we may (and do) assume p is odd. As in the proof of [Corollary 1.2](#), we rewrite the sum as $\sum_{P \in \mathcal{P}_1} \sum_{a \in \mathbb{F}_q^\times} G_p(1/(aP)^k)$. We now need the identity

$$\sum_{a \in \mathbb{F}_q^\times} G_p((aU)^k) = \text{GCD}(q-1, k) G_p(U^{\text{LCM}(q-1, k)})$$

in $\mathbb{F}_q(U)$. To prove this identity, we use the formula $G_p(U) = \sum_{j \not\equiv 0 \pmod p} U^j/j$ and the identity

$$\sum_{a \in \mathbb{F}_q^\times} a^m = \begin{cases} q-1, & m \equiv 0 \pmod{q-1}, \\ 0, & \text{otherwise.} \end{cases}$$

So

$$\sum_{a \in \mathbb{F}_q^\times} G_p(1/(aU)^k) = \sum_{j \not\equiv 0 \pmod p} \sum_{a \in \mathbb{F}_q^\times} \frac{1}{j(aU)^{kj}} = (q-1) \sum_{\substack{j \not\equiv 0 \pmod p \\ kj \equiv 0 \pmod{q-1}}} \frac{1}{jU^{kj}}.$$

Putting $kj = \text{LCM}(q-1, k)\ell$, this is

$$\begin{aligned} (q-1) \sum_{\ell \not\equiv 0 \pmod p} \frac{k}{\text{LCM}(q-1, k)\ell U^{\text{LCM}(q-1, k)\ell}} &= \frac{k(q-1)}{\text{LCM}(q-1, k)} G_p(U^{\text{LCM}(q-1, k)}) \\ &= \text{GCD}(q-1, k) G_p(U^{\text{LCM}(q-1, k)}), \end{aligned}$$

as required. □

We also compute explicit values for the sum when k is not too large.

Theorem 1.7. *Let $k = (q-1)\ell$. If $1 \leq \ell \leq q/p$, then $\sum_{P \in \mathcal{P}_1} G_p(1/P^k) = 0$. If $q/p + 1 < \ell \leq 2q/p$, then*

$$\sum_{P \in \mathcal{P}_1} G_p(1/P^k) = \ell \frac{(T^q - T)^{q+1}}{(T^{q^2} - T^q)(T^{q^2} - T)}.$$

In principle, our methods are capable of computing $\sum_{P \in \mathcal{P}_1} G_p(1/P^k)$ for any $k \equiv 0 \pmod{q-1}$, but they become impractical beyond $\ell = 2q/p$.

History of the problem. Dinesh Thakur suspected such relations should exist, based on heuristics concerning ζ deformation. He experimentally discovered most of the relations described above in characteristic two, and suspected there should be similar results in odd characteristic. Thakur [2015] published these computations in a preprint entitled ‘‘Surprising symmetries in distribution of prime polynomials’’. At Thakur’s suggestion, Terence Tao [2015] promoted the problem in posts on his blog and on the Polymath blog. I am grateful to Thakur for finding such an elegant problem and to Tao for bringing it to my attention. My thanks also to all who participated in the discussion on the Polymath blog: Noam Elkies, Ian Finn, Ofir Gorodetsky, Jesse, Gil Kalai, David Lowry-Duda, Dustin G. Mixon, John Nicol, Partha Solapurkar, John Voight, Victor Wang, Qiaochu Yuan, Joshua Zelinsky, and additional thanks to Ofir Gorodetsky for suggesting several improvements to the manuscript.

The author is supported by NSF grant DMS-1600223.

2. The Carlitz exponential, and symmetric polynomials

The main tool in our proofs is the theory of the Carlitz exponential. Put

$$D_i = (T^{q^i} - T)(T^{q^i} - T^q)(T^{q^i} - T^{q^2}) \dots (T^{q^i} - T^{q^{i-1}}).$$

Define

$$e_C(Z) = \sum_{j=0}^{\infty} \frac{Z^{q^j}}{D_j},$$

this sum is T^{-1} -adically convergent for any $Z \in \widehat{\mathcal{K}}$. We will make use of the product identity

$$\frac{e_C(\bar{\pi} Z)}{\bar{\pi} Z} = \prod_{A \in \mathcal{R} \setminus \{0\}} \left(1 + \frac{Z}{A}\right),$$

where $\bar{\pi} \in \widehat{\mathcal{K}}(\sqrt[q-1]{-T})$ is given by

$$\bar{\pi} = \frac{T^{\sqrt[q-1]{-T}}}{\prod_{A \in \mathcal{R} \setminus \{0\}} (1 - (TA)^{-1})}.$$

See, for example, [Goss 1996, Theorem 3.2.8]. This identity should be thought of as similar to Euler's identity,

$$\frac{\sin(\pi z)}{\pi z} = \prod_{a \in \mathbb{Z} \setminus \{0\}} \left(1 + \frac{z}{a}\right).$$

We introduce the notations \mathcal{A} for the nonzero polynomials of \mathcal{R} , and \mathcal{A}_1 for the monic polynomials.

Writing e_k for the elementary symmetric function of degree k , this implies

$$e_k(1/A)_{A \in \mathcal{A}} = \begin{cases} \bar{\pi}^k / D_j, & k = q^j - 1, \\ 0, & \text{otherwise.} \end{cases}$$

Since the ring of symmetric polynomials is generated by the e_k , we deduce:

Proposition 2.1. *If f is a homogenous symmetric polynomial of degree k , then $f(1/A)_{A \in \mathcal{A}}$ is in $\bar{\pi}^k \mathcal{K}$.*

Here we note that $f(1/A)_{A \in \mathcal{A}}$ is always defined, since only finitely many terms contribute to the coefficient of any particular power of T^{-1} .

The above considers symmetric polynomials in $\{1/A\}_{A \in \mathcal{A}}$, but we would rather restrict to the case of A monic. To this end, we have

Proposition 2.2.

$$e_\ell(1/A^{q-1})_{A \in \mathcal{A}_1} = \begin{cases} (-1)^\ell \bar{\pi}^{\ell(q-1)} / D_j, & \ell = (q^j - 1) / (q - 1), \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Grouping together scalar multiples of the same polynomial in the Carlitz product identity, we have

$$\frac{e_C(\bar{\pi} Z)}{\bar{\pi} Z} = \prod_{A \in \mathcal{A}_1} \left(1 - \frac{Z^{q-1}}{A^{q-1}}\right).$$

Equate coefficients of $Z^{\ell(q-1)}$ on both sides. □

Corollary 2.3. *If f is a homogenous symmetric polynomial of degree ℓ , then $f(1/A^{q-1})_{A \in \mathcal{A}_1}$ is in $\bar{\pi}^{\ell(q-1)}\mathcal{K}$.*

3. Proofs of rationality

We now have enough background to prove [Theorem 1.4](#) and, hence, [Theorem 1.1](#). Throughout, let $k \equiv 0 \pmod{q-1}$.

Consider the symmetric polynomial

$$g_p(X_1, X_2, \dots) := \frac{1}{p} \left(\left(\sum X_i \right)^p - \sum X_i^p \right).$$

The polynomial g_p has integer coefficients, so we may discuss plugging elements of \mathcal{K} into it.

Let C be the cyclic group of order p , and let C act on \mathcal{A}_1^p by rotating coordinates. Let Δ denote the diagonal: $\Delta := \{(A, A, \dots, A)\} \subset \mathcal{A}_1^p$. Then

$$g_p(1/A^k)_{A \in \mathcal{A}_1} = \sum_{(A_1, \dots, A_p) \in (\mathcal{A}_1^p \setminus \Delta)/C} \frac{1}{A_1^k A_2^k \cdots A_p^k}.$$

The sum is over cosets for the free action of C on $\mathcal{A}^p \setminus \Delta$.

Let

$$\Phi = \{(A_1, \dots, A_p) \in \mathcal{A}_1^p : \text{GCD}(A_1, \dots, A_p) = 1\}.$$

Any $(A_1, \dots, A_p) \in \mathcal{A}_1^p$ can be uniquely factored as $A_i = DB_i$ for some $D \in \mathcal{A}_1$ and $(B_1, \dots, B_p) \in \Phi$. So we can factor the above sum as

$$g_p(1/A^k)_{A \in \mathcal{A}_1} = \left(\sum_{D \in \mathcal{A}_1} \frac{1}{D^{kp}} \right) \left(\sum_{(B_1, \dots, B_p) \in (\Phi \setminus \{(1, \dots, 1)\})/C} \frac{1}{B_1^k B_2^k \cdots B_p^k} \right).$$

Now, from [Corollary 2.3](#), $g_p(1/A^k)_{A \in \mathcal{A}_1}$ is in $\bar{\pi}^{pk}\mathcal{K}$. Also from [Corollary 2.3](#), $\sum_{D \in \mathcal{A}_1} 1/D^{kp}$ is in $\bar{\pi}^{pk}\mathcal{K}$, and a quick computation shows that this sum is 1 plus terms in $T^{-1}\mathbb{F}_q[[T^{-1}]]$, so it is not zero. We deduce that

$$\sum_{(B_1, \dots, B_p) \in (\Phi \setminus \{(1, \dots, 1)\})/C} \frac{1}{B_1^k B_2^k \cdots B_p^k} \in \mathcal{K}.$$

For $B \in \mathcal{A}_1$, let $\Psi(B)$ be the set of p -tuples (B_1, B_2, \dots, B_p) for which $\prod B_i = B$ and $\text{GCD}(B_1, \dots, B_p) = 1$. Let $\psi(B) = \#\Psi(B)$. So we have shown that

$$\sum_{B \in \mathcal{A}_1 \setminus \{1\}} \frac{\psi(B)/p}{B^k} \in \mathcal{K}.$$

Here, to interpret the numerator, we must divide $\psi(B)$ by p as integers and only then consider the quotient in \mathbb{F}_p .

If $B = P_1^{k_1} P_2^{k_2} \dots P_r^{k_r}$ then there is an easy bijection between $\Psi(B)$ and $\Psi(P_1^{k_1}) \times \Psi(P_2^{k_2}) \times \dots \times \Psi(P_r^{k_r})$, so $\psi(B) = \prod \psi(P_i^{k_i})$. If P is irreducible then $\psi(P^r)$ is divisible by p for any $r > 0$, since C acts freely on $\Psi(P^r)$. So, if B is divisible by two different irreducible polynomials, then $\psi(B)$ is divisible by p^2 . So we can rewrite the sum as

$$\sum_{P \in \mathcal{P}_1} \sum_{r=1}^{\infty} \frac{\psi(P^r)/p}{P^{rk}}.$$

We now compute $\psi(P^r)$; which is the number of p -tuples $(P^{r_1}, \dots, P^{r_p})$ with $\prod P^{r_i} = P^r$ and $\text{GCD}(P^{r_1}, \dots, P^{r_p}) = 1$. In other words, we must count $(r_1, \dots, r_p) \in \mathbb{Z}_{\geq 0}^p$ with $\sum r_i = r$ and $\min(r_1, \dots, r_p) = 0$. The number of $(r_1, \dots, r_p) \in \mathbb{Z}_{\geq 0}^p$ with $\sum r_i = r$ is the coefficient of U^r in $1/(1-U)^p$. In order to impose $\min(r_1, \dots, r_p) = 0$, we subtract off the terms with $\min(r_1, \dots, r_p) > 0$. These are in bijection with $(s_1, \dots, s_p) \in \mathbb{Z}_{\geq 0}^p$ with $p + \sum s_i = r$. So $\psi(P^r)$ is the coefficient of U^r in $1/(1-U)^p - U^p/(1-U)^p$. In other words, $\sum_{r=0}^{\infty} \psi(P^r)U^r = (1-U^p)/(1-U)^p$. So

$$\sum_{r=1}^{\infty} \frac{\psi(P^r)}{p} U^r = \frac{1}{p} \left(\frac{1-U^p}{(1-U)^p} - 1 \right) = G_p(U).$$

We deduce that $\sum_{r=1}^{\infty} (\psi(P^r)/p)/P^{rk} = G_p(1/P^k)$. We have now shown that $\sum_{P \in \mathcal{P}_1} G_p(1/P^k) \in \mathcal{K}$, as claimed. □

We record the specific formula we have proved:

Proposition 3.1. *Let k be a positive integer. Then*

$$\sum_{P \in \mathcal{P}_1} G_p(1/P^k) = \frac{\mathbf{g}_p(1/A^k)_{A \in \mathcal{A}_1}}{\sum_{A \in \mathcal{A}_1} 1/A^{pk}}.$$

We will rewrite this formula in various ways in [Section 5](#). We remark that this formula is correct even if k is not divisible by $q - 1$, although we have only shown the ratio is in \mathcal{K} when $k \equiv 0 \pmod{q - 1}$. The denominator of this formula is $\zeta(pk) = \zeta(k)^p$, where ζ is the Goss ζ -function [\[1979\]](#).

4. Vanishing

We will now prove the claim in [Theorem 1.7](#) that the sum vanishes when $k = (q - 1)\ell$ for $1 \leq \ell \leq q/p$. From [Proposition 3.1](#), it is equivalent to show that $\mathbf{g}_p(1/A^{\ell(q-1)})_{A \in \mathcal{A}_1} = 0$. To this end, we must explicitly write $\mathbf{g}_p(1/A^{\ell(q-1)})$ as a polynomial in the $\mathbf{e}_k(1/A^{q-1})$.

The variables λ or μ will always denote partitions, meaning weakly decreasing sequences $(\lambda_1, \lambda_2, \dots, \lambda_r)$ of positive integers; sums over λ or μ implicitly contain the condition that the summation variable is a partition.

We define $\mathbf{e}_\lambda = \prod_s \mathbf{e}_{\lambda_s}$. The symmetric polynomials \mathbf{e}_λ form an integer basis for the symmetric polynomials with integer coefficients.

Lemma 4.1. *Write*

$$\mathbf{g}_p(X_1^\ell, X_2^\ell, \dots) = \sum_{|\lambda|=p\ell} c_\lambda \mathbf{e}_\lambda(X_1, X_2, \dots)$$

for some integers c_λ . Then $c_{11\dots 1} = 0$.

Proof. Note that $\mathbf{e}_{11\dots 1}$ is the only \mathbf{e}_λ with a nonzero coefficient of $X_1^{p\ell}$. The coefficient of $X_1^{p\ell}$ in $\mathbf{g}_p(X_1^\ell, X_2^\ell, \dots)$ is clearly 0. □

Now, suppose that $\ell \leq q/p$, so we have $p\ell < q + 1$. So any partition $(\lambda_1, \dots, \lambda_r)$ of $p\ell$ other than $(1, 1, \dots, 1)$ contains a λ_i between 2 and q . By [Proposition 2.2](#), $\mathbf{e}_m(1/A^{q-1})_{A \in \mathcal{A}_1} = 0$ for $2 \leq m \leq q$, so $\mathbf{e}_\lambda(1/A^{q-1})_{A \in \mathcal{A}_1} = 0$ whenever λ is a partition of $p\ell$ other than $(1, 1, \dots, 1)$. We deduce that $\mathbf{g}_p(1/A^{q-1})_{A \in \mathcal{A}_1} = 0$ as desired. □

5. Computations for small k

In this section, we will discuss the computation of $\sum_{P \in \mathcal{P}_1} G_p(1/P^k)$ for $k \equiv 0 \pmod{q-1}$ and, in particular, prove the remaining half of [Theorem 1.7](#). Our strategy is to combine [Propositions 3.1](#) and [2.2](#). We must compute $\mathbf{g}_p(1/A^{\ell(q-1)})_{A \in \mathcal{A}_1}$ and $\sum_{A \in \mathcal{A}_1} 1/A^{pk}$. Note the latter is $(\mathbf{p}_\ell(1/A^{q-1})_{A \in \mathcal{A}_1})^p$, where $\mathbf{p}_d(X_1, X_2, \dots)$ is the power sum symmetric function $\sum X_i^d$. We write $k = (q - 1)\ell$.

Put

$$\begin{aligned} \mathbf{g}_p(X_1^\ell, X_2^\ell, \dots) &= \sum_{|\lambda|=\ell p} c_\lambda \mathbf{e}_\lambda(X_1, X_2, \dots), \\ \mathbf{p}_\ell(X_1, X_2, \dots) &= \sum_{|\mu|=\ell} d_\mu \mathbf{e}_\mu(X_1, X_2, \dots). \end{aligned}$$

Note that $\mathbf{e}_m(1/A^{q-1})_{A \in \mathcal{A}_1} = 0$ unless m is of the form $(q^j - 1)/(q - 1)$. So we only need to sum over partitions where all the parts of λ are of the form $(q^j - 1)/(q - 1)$.

From now on, we now impose that $q/p + 1 \leq \ell \leq 2q/p$. So $\ell < q + 1$. Any partition of ℓ cannot contain any parts of size $(q^j - 1)/(q - 1)$, for $j > 1$. Similarly,

$p\ell < 2q + 2$, so a partition of $p\ell$ can contain at most one part of size $q + 1 = (q^2 - 1)/(q - 1)$ and no parts of size $(q^j - 1)/(q - 1)$ for $j > 2$. We deduce that the only terms which contribute to our final answer come from $\lambda = (1, 1, \dots, 1)$ or $\lambda = (q + 1, 1, 1, \dots, 1)$ when computing $\mathbf{g}_p(1/A^{\ell(q-1)})_{A \in \mathcal{A}_1}$, and from $\mu = (1, 1, \dots, 1)$ in computing $(p_\ell(1/A^{q-1}))_{A \in \mathcal{A}_1}^p$. Moreover, from Lemma 4.1, the coefficient $c_{(1,1,\dots,1)}$ is zero.

We deduce that

$$\begin{aligned} \sum_{P \in \mathcal{P}_1} G_p(1/P^k) &= \frac{c_{(q+1,1^{p\ell-q-1})} \mathbf{e}_{(q+1,1^{p\ell-q-1})}(1/A^{q-1})_{A \in \mathcal{A}_1}}{(d_{1^\ell} \mathbf{e}_1(1/A^{q-1})_{A \in \mathcal{A}_1})^p} \\ &= \frac{c_{(q+1,1^{p\ell-q-1})} \mathbf{e}_{q+1}(1/A^{q-1})_{A \in \mathcal{A}_1} (\mathbf{e}_1(1/A^{q-1})_{A \in \mathcal{A}_1})^{p\ell-q-1}}{d_{1^\ell} (\mathbf{e}_1(1/A^{q-1})_{A \in \mathcal{A}_1})^{p\ell}} \\ &= \frac{c_{(q+1,1^{p\ell-q-1})} \mathbf{e}_{q+1}(1/A^{q-1})_{A \in \mathcal{A}_1}}{d_{1^\ell} (\mathbf{e}_1(1/A^{q-1})_{A \in \mathcal{A}_1})^{q+1}}. \end{aligned}$$

Here 1^r is shorthand for the partition with r parts equal to 1.

We now use Proposition 2.2. The powers of $\bar{\pi}$ and (-1) cancel to give

$$\sum_{P \in \mathcal{P}_1} G_p(1/P^k) = \frac{c_{(q+1,1^{p\ell-q-1})} D_1^{q+1}}{d_{1^\ell} D_2} = \frac{c_{(q+1,1^{p\ell-q-1})}}{d_{1^\ell}} \frac{(T^q - T)^{q+1}}{(T^{q^2} - T^q)(T^{q^2} - T)}.$$

To finish the computation, we must find $c_{q+1,1^{ps-1}}$ and d_{1^ℓ} . The latter is easy: Comparing coefficients of X_1^ℓ on both sides of

$$p_\ell(X_1, X_2, \dots) = \sum_{|\mu|=\ell} d_\mu \mathbf{e}_\mu(X_1, X_2, \dots),$$

we deduce that $d_{1^\ell} = 1$.

To compute $c_{(q+1,1^{p\ell-q-1})}$, we begin with the formula

$$\mathbf{g}_p(X_1^\ell, X_2^\ell, \dots) = \frac{1}{p} (p_\ell(X_1, X_2, \dots)^p - p_{p\ell}(X_1, X_2, \dots)).$$

For brevity, we write $\mathbf{f}(X)$ to indicate that the inputs to a symmetric polynomial are (X_1, X_2, \dots) . Note that we are working with symmetric polynomials with integer coefficients, so it makes sense to divide by p .

We rewrite the right hand side of the previous equation as

$$\frac{1}{p} \left((\mathbf{e}_1(X)^\ell + \dots)^p - (\mathbf{e}_1(X)^{p\ell} + d_{q+1,1^{p\ell-q-1}} \mathbf{e}_{q+1}(X) \mathbf{e}_1(X)^{p\ell-q-1} + \dots) \right).$$

Here the ellipses denote terms \mathbf{e}_λ where λ has some part that is not of the form $(q^j - 1)/(q - 1)$. We deduce that

$$c_{q+1,1^{p\ell-q-1}} = -\frac{1}{p} d_{q+1,1^{p\ell-q-1}}.$$

Now, observe the identity

$$\begin{aligned} \sum_j \frac{(-1)^{j-1} \mathbf{p}_j(X) U^j}{j} &= \sum_i \log(1 + X_i U) \\ &= \log \prod_i (1 + X_i U) = \log \left(1 + \sum_{m=1}^{\infty} \mathbf{e}_m(X) U^m \right). \end{aligned}$$

The coefficient of $U^{p\ell}$ on the left is $((-1)^{p\ell}/p\ell) \mathbf{p}_{p\ell}$. Expanding the log on the right hand side as a Taylor series, only one term contributes to $U^{p\ell} \mathbf{e}_{q+1} \mathbf{e}_1^{p\ell-q-1}$.

So we obtain

$$\frac{(-1)^{p\ell-1}}{p\ell} \mathbf{p}_{p\ell}(X) = \frac{(-1)^{p\ell-q-1}}{p\ell-q} \binom{p\ell-q}{1} \mathbf{e}_{q+1}(X) \mathbf{e}_1^{p\ell-q-1}(X) + \dots,$$

where the ellipses denote a sum of \mathbf{e}_λ other than $\mathbf{e}_{q+1}(X) \mathbf{e}_1^{p\ell-q-1}(X)$. So

$$d_{q+1, 1^{p\ell-q-1}} = (-1)^q p\ell \quad \text{and} \quad c_{q+1, 1^{ps-1}} = (-1)^{q-1} \ell.$$

Plugging into our previous formula, and using that $(-1)^{q-1} \equiv 1 \pmod p$,

$$\sum_{P \in \mathcal{P}_1} G_p(1/P^k) = \ell \frac{(T^q - T)^{q+1}}{(T^{q^2} - T^q)(T^{q^2} - T)}.$$

This concludes the proof of [Theorem 1.7](#). □

We conclude by verifying one of Thakur’s conjectures which goes beyond the range $\ell \leq 2q/p$. Let $p = q = 2$. Thakur conjectures

$$\sum_{P \in \mathcal{P}_1} \frac{1}{P^3 - 1} = \frac{1}{T^4 + T^2}.$$

We begin by computing

$$\begin{aligned} \mathbf{p}_3(X) &= \mathbf{e}_1(X)^3 + 3\mathbf{e}_3(X) - 3\mathbf{e}_2(X)\mathbf{e}_1(X), \\ \mathbf{p}_3(X)^2 &= \mathbf{e}_1(X)^6 + 6\mathbf{e}_1(X)^3\mathbf{e}_3(X) + 9\mathbf{e}_3(X)^2 + \dots \end{aligned}$$

Here and in the following equations, the ellipses denote \mathbf{e}_λ terms where λ contains a part other than 1 and 3. (Note that $(2^3 - 1)/(2 - 1) = 7$, too large to contribute to a symmetric polynomial of degree 6.) Similarly,

$$\mathbf{p}_6(X) = \mathbf{e}_1(X)^6 + 6\mathbf{e}_1(X)^3\mathbf{e}_3(X) + 3\mathbf{e}_3(X)^2 + \dots$$

So

$$\mathbf{g}_2(X_1^3, X_2^3, \dots) = \frac{1}{2}(\mathbf{p}_3(X)^2 - \mathbf{p}_6(X)) = 3\mathbf{e}_3(X)^2 + \dots$$

and (recall that we are working modulo 2)

$$\mathbf{g}_2(1/A^3)_{A \in \mathcal{A}_1} = (\mathbf{e}_3(1/A)_{A \in \mathcal{A}_1})^2 = \frac{\bar{\pi}^6}{D^2} = \frac{\bar{\pi}^6}{(T^4 - T^2)^2(T^4 - T)^2}.$$

Similarly,

$$\begin{aligned} p_6(1/A)_{A \in \mathcal{A}_1} &= (e_1(1/A)_{A \in \mathcal{A}_1})^6 + (e_3(1/A)_{A \in \mathcal{A}_1})^2 \\ &= \left(\frac{\bar{\pi}}{D_1}\right)^6 + \left(\frac{\bar{\pi}^3}{D_2}\right)^2 \\ &= \bar{\pi}^6 \left(\left(\frac{1}{T^2 - T}\right)^6 + \left(\frac{1}{(T^4 - T^2)(T^4 - T)}\right)^2 \right). \end{aligned}$$

We verify Thakur's claim:

$$\sum_{P \in \mathcal{P}_1} \frac{1}{P^3 - 1} = \frac{1/((T^4 - T^2)^2(T^4 - T)^2)}{1/(T^2 - T)^6 + 1/((T^4 - T^2)^2(T^4 - T)^2)} = \frac{1}{T^4 + T^2}.$$

References

- [Goss 1979] D. Goss, “*v*-adic zeta functions, *L*-series and measures for function fields”, *Invent. Math.* **55**:2 (1979), 107–119. [MR](#) [Zbl](#)
- [Goss 1996] D. Goss, *Basic structures of function field arithmetic*, Ergebnisse der Mathematik (3) **35**, Springer, 1996. [MR](#) [Zbl](#)
- [Tao 2015] T. Tao, “Polymath proposal: explaining identities for irreducible polynomials”, 2015, Available at <https://polymathprojects.org/2015/12/28/>.
- [Thakur 2015] D. S. Thakur, “Surprising symmetries in distribution of prime polynomials”, preprint, 2015. [arXiv](#)

Communicated by Kiran S. Kedlaya

Received 2016-10-17 Accepted 2017-04-03

speyer@umich.edu

*Department of Mathematics, University of Michigan,
2844 East Hall, 530 Church Street,
Ann Arbor, MI 48109-1043, United States*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Raman Parimala	Emory University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	University of California, Santa Cruz, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2017 is US \$325/year for the electronic version, and \$520/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 11 No. 5 2017

Hybrid sup-norm bounds for Maass newforms of powerful level ABHISHEK SAHA	1009
Collinear CM-points YURI BILU, FLORIAN LUCA and DAVID MASSER	1047
A uniform classification of discrete series representations of affine Hecke algebras DAN CIUBOTARU and ERIC OPDAM	1089
An explicit bound for the least prime ideal in the Chebotarev density theorem JESSE THORNER and ASIF ZAMAN	1135
Modular curves of prime-power level with infinitely many rational points ANDREW V. SUTHERLAND and DAVID ZYWINA	1199
Some sums over irreducible polynomials DAVID E. SPEYER	1231