

Algebra & Number Theory

Volume 17
2023
No. 11

Differential operators, retracts, and toric face rings

Christine Berkesch, C-Y. Jean Chan, Patricia Klein,
Laura Felicia Matusevich, Janet Page and Janet Vassilev



Differential operators, retracts, and toric face rings

Christine Berkesch, C-Y. Jean Chan, Patricia Klein,
Laura Felicia Matusevich, Janet Page and Janet Vassilev.

We give explicit descriptions of rings of differential operators of toric face rings in characteristic 0. For quotients of normal affine semigroup rings by radical monomial ideals, we also identify which of their differential operators are induced by differential operators on the ambient ring. Lastly, we provide a criterion for the Gorenstein property of a normal affine semigroup ring in terms of its differential operators.

Our main technique is to realize the \mathbb{k} -algebras we study in terms of a suitable family of their algebra retracts in a way that is compatible with the characterization of differential operators. This strategy allows us to describe differential operators of any \mathbb{k} -algebra realized by retracts in terms of the differential operators on these retracts, without restriction on $\text{char}(\mathbb{k})$.

Introduction	1959
1. \mathbb{k} -algebras realized by retracts	1961
2. Rings of differential operators on \mathbb{k} -algebras realized by retracts	1963
3. Applications	1968
4. Rings of differential operators of quotient rings	1972
5. Characterizing Gorenstein rings via differential operators	1975
Acknowledgements	1981
References	1982

Introduction

Differential operators play a notable role in branches of mathematics as seemingly disparate as partial differential equations and local cohomology, dynamical systems and invariant theory. Recently, there have been many exciting new developments concerning differential operators and their applications in commutative algebra, including to topics such as Bernstein–Sato polynomials, connections between singularities and local cohomology, equivariant D -modules, and Hodge ideals; see, for example, [1; 2; 9; 16; 20; 21; 23; 24; 25; 31; 36; 37; 38; 39; 40; 41]. One obstruction to the even greater use of rings of differential operators is the notorious difficulty of computing them explicitly. In fact, there are very few classes of rings whose differential operators are systematically computed, namely polynomial rings, Stanley–Reisner rings, affine semigroup rings, and coordinate rings of curves; see, for example, [11; 12;

Berkesch was partially supported by NSF Grants DMS 1661962 and 2001101. Matusevich was partially supported by the Simons Foundation Collaboration Grant for Mathematicians.

MSC2020: primary 16S32; secondary 13F55, 13N05.

Keywords: differential operators, toric face rings, algebra retracts, affine semigroup rings.

22; 26; 27; 28; 29; 30; 32; 42; 43; 44; 47; 48]. The goal of this article is to give a new class of explicitly computed rings of differential operators.

A *retract* of a \mathbb{k} -algebra R is a subring of R which is isomorphic to a quotient of R . We call R a \mathbb{k} -algebra realized by retracts if it can be embedded into a finite direct sum of domains, each of which is a retract of R . Algebra retracts are of considerable interest; it is common in the literature to prove desirable properties of retracts using information from the ambient ring; see, for example, [3; 6; 7; 10; 14; 15; 46]. In this work, we do the opposite; we use knowledge about the differential operators on algebra retracts to compute differential operators of an ambient ring (see Theorems 2.6 and 2.8). Naturally, this approach is most productive when the differential operators on the retracts are known. This is the case for a combinatorially interesting class of \mathbb{k} -algebras realized by retracts known as *toric face rings*.

Toric face rings were first introduced by Stanley [45], and further developed in [3; 17; 18; 33; 34; 35; 49], among others. They include both Stanley–Reisner rings and affine semigroup rings as special cases, thus bringing under a single umbrella two of the mainstays of combinatorial commutative algebra. In this setting, the retracts we are interested in are affine semigroup rings, whose rings of differential operators are known in characteristic zero; here we use the presentation given by Saito and Traves in [42; 43]. Thus we can directly apply Theorem 2.8 to compute the ring of differential operators of a toric face ring in Proposition 3.3. As a consequence, we recover results on differential operators on Stanley–Reisner rings given by Tripp, Eriksson, and Traves in [12; 47; 48] over arbitrary fields.

Theorem 2.6 finds the differential operators of a \mathbb{k} -algebra realized by retracts as a subring of the direct sum of the rings of differential operators of the retracts. This direct sum of the retracts is in general much larger than the original ring. On the other hand, the differential operators on Stanley–Reisner rings and affine semigroup rings are induced from differential operators over their natural ambient rings. In general, however, the richness of the direct sum is really needed. To illustrate this, we provide a description of which differential operators on a quotient of a normal affine semigroup ring by a radical monomial ideal are induced from the differential operators on the ambient semigroup ring (see Theorem 4.3) and show that this does not necessarily give the whole ring of differential operators.

Outline. In Section 2, we describe the ring of differential operators of a \mathbb{k} -algebra realized by retracts in terms of differential operators on these retracts. In Section 3, we apply the results of Section 2 in characteristic zero to compute the rings of differential operators of Stanley–Reisner rings and of toric face rings more generally. In Section 4, we consider the quotient of a normal affine semigroup ring R by a radical monomial ideal J in characteristic zero and provide an explicit formula for the differential operators on R/J that are induced by operators on R . Finally, in Section 5, we use differential operators to provide a new condition for the Gorenstein property of an affine normal semigroup ring.

Throughout this paper, let \mathbb{k} denote an algebraically closed field; in Sections 3, 4, and 5, assume that the characteristic of \mathbb{k} is zero.

1. \mathbb{k} -algebras realized by retracts

In this section we introduce \mathbb{k} -algebras realized by retracts and provide examples. Let S and R be \mathbb{k} -algebras. An injective \mathbb{k} -algebra homomorphism $\iota: S \rightarrow R$ is called an *algebra retract* of R if there exists a surjective homomorphism of \mathbb{k} -algebras $\pi: R \rightarrow S$ such that $\pi \circ \iota = \text{id}_S$. When S and R are graded, we also assume that the homomorphisms ι and π are also graded. See [15] for a local version of this definition.

Definition 1.1. Let R be a \mathbb{k} -algebra, and let $\{S_\ell\}_{\ell \in \Lambda}$ be a finite collection of domains that are algebra retracts of R with defining \mathbb{k} -algebra homomorphisms $\iota_\ell: S_\ell \hookrightarrow R$ and $\pi_\ell: R \twoheadrightarrow S_\ell$ satisfying $\pi_\ell \circ \iota_\ell = \text{id}_{S_\ell}$. For each ℓ , let $P_\ell := \ker \pi_\ell$. We call R a \mathbb{k} -algebra realized by retracts and say that R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$ when the following two conditions hold:

(i) The following map is injective:

$$\phi: R \rightarrow \bigoplus_{\ell \in \Lambda} S_\ell \quad \text{given by } f \mapsto (\pi_\ell(f) \mid \ell \in \Lambda); \tag{1.2}$$

in other words, $\bigcap_{\ell \in \Lambda} P_\ell = 0$.

(ii) The S_ℓ are irredundant for (1.2); more precisely, for each $i \in \Lambda$, we have $\bigcap_{\substack{\ell \neq i \\ \ell \in \Lambda}} P_\ell \neq 0$.

Since ι_ℓ is injective, if $f \in S_\ell$, we write $f = \iota_\ell(f)$ in R . For $f \in R$, we use $\phi(f)_i$ to denote the i -th coordinate of $\phi(f)$ in $\bigoplus_{\ell \in \Lambda} S_\ell$.

For example, suppose that R is a quotient of a \mathbb{k} -algebra T by a radical ideal I with associated primes P_1, \dots, P_r . Now if T/P_i are algebra retracts of R for all $1 \leq i \leq r$, then R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\} = \{T/P_1, T/P_2, \dots, T/P_r\}$. In the remainder of this section, we show that Stanley–Reisner rings and toric face rings are \mathbb{k} -algebras realized by retracts.

1A. Stanley–Reisner rings as \mathbb{k} -algebras realized by retracts. Let Δ be a simplicial complex on a finite vertex set $V = \{1, 2, \dots, d\}$. The *Stanley–Reisner ring of Δ* is the \mathbb{k} -algebra given by

$$\mathbb{k}[\Delta] := \frac{\mathbb{k}[t_1, t_2, \dots, t_d]}{\langle t^{\mathbf{a}} \mid \mathbf{a} \in \mathbb{N}^d, \text{supp}(\mathbf{a}) \notin \Delta \rangle},$$

where $t^{\mathbf{a}} = t_1^{a_1} t_2^{a_2} \cdots t_d^{a_d}$ and $\text{supp}(\mathbf{a}) := \{i \in V \mid a_i \neq 0\}$. Retracts have been previously studied in this context. For example, in [10], Epstein and Nguyen show that every graded algebra retract of a Stanley–Reisner ring $\mathbb{k}[\Delta]$ is a Stanley–Reisner ring. Further, all such retracts are isomorphic to $\mathbb{k}[\Delta|_W]$, where $\Delta|_W$ is the restriction of Δ to a subset W of V .

One way to view $\mathbb{k}[\Delta]$ as a \mathbb{k} -algebra realized by retracts is via the facets of Δ . If $\{F_\ell\}_{\ell \in \Lambda}$ denotes the collection of facets of Δ , then $\mathbb{k}[F_\ell] \cong \mathbb{k}[t_i \mid i \in F_\ell]$, and

$$\begin{aligned} \iota_\ell: \mathbb{k}[F_\ell] \hookrightarrow \mathbb{k}[\Delta] \quad \text{and} \quad \pi_\ell: \mathbb{k}[\Delta] \twoheadrightarrow \mathbb{k}[F_\ell] \\ t_i \mapsto \begin{cases} t_i & \text{if } i \in F_\ell, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

are the maps needed to see that

$$\phi: \mathbb{k}[\Delta] \rightarrow \bigoplus_{\ell \in \Lambda} \mathbb{k}[F_\ell] \quad \text{given by } t_i \mapsto (\pi_\ell(t_i) \mid \ell \in \Lambda)$$

is an injective map. Since the retracts are domains, this implies that $\mathbb{k}[\Delta]$ is a \mathbb{k} -algebra realized by retracts.

1B. Toric face rings as \mathbb{k} -algebras realized by retracts. The building blocks of toric face rings are affine semigroup rings, so we begin with those.

Notation 1.3. Let M be a finitely generated submonoid of \mathbb{Z}^d . The *affine semigroup ring defined by M* is

$$\mathbb{k}[M] := \bigoplus_{a \in M} \mathbb{k} \cdot t^a,$$

where $t^a = t_1^{a_1} t_2^{a_2} \cdots t_d^{a_d}$ for $\mathbf{a} = (a_1, a_2, \dots, a_d)$.

Finitely generated submonoids of \mathbb{Z}^d are usually called affine semigroups. It is convenient to view the generators of an affine semigroup M as the columns of a $d \times n$ integer matrix A ; in this case, we use the notation $M = \mathbb{N}A$. Throughout this article, we assume that the group generated by the columns of A is the full ambient lattice, so $\mathbb{Z}A = \mathbb{Z}^d$ and also that the real positive cone over A , $\mathbb{R}_{\geq 0}A$, is pointed, meaning that it contains no lines.

A semigroup $\mathbb{N}A$ is *normal* if $\mathbb{N}A = \mathbb{R}_{\geq 0}A \cap \mathbb{Z}A$. In this case, the semigroup ring $\mathbb{k}[\mathbb{N}A]$ is normal in the sense of commutative algebra.

A hyperplane H in \mathbb{R}^d is a supporting hyperplane of $\mathbb{R}_{\geq 0}A$ if this cone lies entirely in one of the closed half spaces defined by H . A *face* σ of $\mathbb{R}_{\geq 0}A$ (or A or $\mathbb{N}A$) is the intersection of $\mathbb{N}A$ with a supporting hyperplane of $\mathbb{R}_{\geq 0}A$. Such a face is called a *facet* if its linear span has dimension $d - 1$. This is somewhat nonstandard terminology, as our faces and facets are submonoids of $\mathbb{N}A$ instead of cones.

Recall that the \mathbb{Z}^d -graded prime ideals in $\mathbb{k}[\mathbb{N}A]$ are in one-to-one correspondence with the faces of A (or $\mathbb{R}_{\geq 0}A$) [19, Proposition 1.5], as a face τ of A corresponds to the multigraded prime $\mathbb{k}[\mathbb{N}A]$ -ideal $P_\tau = \langle t^a \mid \mathbf{a} \in \mathbb{N}A \setminus \tau \rangle$.

Next, let $\Sigma \subset \mathbb{R}^d$ be a rational polyhedral fan consisting of strongly convex (or pointed) cones. A *monoidal complex* \mathcal{M} supported on Σ is a collection of monoids $\{M_\sigma \mid \sigma \in \Sigma\}$ such that:

- (i) $M_\tau \subseteq \tau \cap \mathbb{Z}^d$ and $\mathbb{R}_{\geq 0}M_\tau = \tau$.
- (ii) if $\sigma, \tau \in \Sigma$ and $\sigma \subseteq \tau$, then $M_\sigma = \sigma \cap M_\tau$.

Denote $|\mathcal{M}| = \bigcup_{\tau \in \Sigma} M_\tau$.

Definition 1.4. The *toric face ring* of \mathcal{M} over \mathbb{k} is given as a graded vector space by

$$\mathbb{k}[\mathcal{M}] = \bigoplus_{a \in |\mathcal{M}|} \mathbb{k} \cdot t^a$$

with multiplication defined by

$$t^{\mathbf{a}} \cdot t^{\mathbf{b}} = \begin{cases} t^{\mathbf{a}+\mathbf{b}} & \text{if there is } \tau \in \Sigma \text{ such that } \mathbf{a}, \mathbf{b} \in \tau, \\ 0 & \text{otherwise.} \end{cases}$$

Given $\tau \in \Sigma$, the semigroup ring $\mathbb{k}[M_\tau]$ is a subring of $\mathbb{k}[\mathcal{M}]$; we denote by

$$\iota_\tau: \mathbb{k}[M_\tau] \hookrightarrow \mathbb{k}[\mathcal{M}]$$

the natural inclusion. For $\tau \in \Sigma$, let

$$P_\tau := \langle t^{\mathbf{a}} \mid \mathbf{a} \in \mathcal{M} \setminus M_\tau \rangle.$$

Then $\mathbb{k}[\mathcal{M}]/P_\tau$ is isomorphic to the semigroup ring $\mathbb{k}[M_\tau]$, so that P_τ is a monomial prime ideal of $\mathbb{k}[\mathcal{M}]$. We identify $\mathbb{k}[\mathcal{M}]/P_\tau$ with $\mathbb{k}[M_\tau]$, and denote by

$$\pi_\tau: \mathbb{k}[\mathcal{M}] \rightarrow \mathbb{k}[M_\tau]$$

the natural projection onto the quotient. Clearly, $\mathbb{k}[M_\tau]$ is a retract of $\mathbb{k}[\mathcal{M}]$.

A *facet* of Σ is a cone in Σ which is maximal with respect to inclusion among all elements of Σ . Let $\mathcal{F}(\Sigma)$ denote the collection of all facets of Σ . Since $\bigcap_{\tau \in \mathcal{F}(\Sigma)} P_\tau = 0$, the ring homomorphism

$$\phi: \mathbb{k}[\mathcal{M}] \rightarrow \bigoplus_{\tau \in \mathcal{F}(\Sigma)} \mathbb{k}[M_\tau] \quad \text{given by } f \mapsto (\pi_\tau(f) \mid \tau \in \mathcal{F}(\Sigma)) \tag{1.5}$$

is injective. It follows that $\mathbb{k}[\mathcal{M}]$ is a \mathbb{k} -algebra realized by retracts.

In general, the algebra retracts of toric face rings of $\mathbb{k}[\mathcal{M}]$ are given by restricting \mathcal{M} to a subfan Γ of Σ [10, Proposition 4.5]. The retracts that we consider here are those given by the restriction of Σ to one of its maximal cones Γ , yielding an affine semigroup ring.

Example 1.6. Stanley–Reisner rings of simplicial complexes are toric face rings. To see this, let Δ be a simplicial complex on the vertex set $V = \{1, 2, \dots, d-1\}$. For each subset F of V , associate the pointed cone C_F generated by the set of elements of the form $\mathbf{e}_i + \mathbf{e}_d$ for $i \in F$, where \mathbf{e}_i denotes the i -th standard basis vector in \mathbb{R}^d . If Σ denotes the fan in \mathbb{R}^d consisting of the cones C_F for $F \in \Delta$, then the toric face ring $\mathbb{k}[\mathcal{M}]$ is isomorphic to the Stanley–Reisner ring $\mathbb{k}[\Delta]$. \square

Example 1.7. On the other hand, when Σ has a unique maximal cone, then $\mathbb{k}[\mathcal{M}]$ is simply an affine semigroup ring. An affine semigroup ring $\mathbb{k}[\mathbb{N}A]$ modulo a radical monomial ideal J is also a toric face ring. In this case, if the ideal $J = \bigcap_{i=1}^r P_{\tau_i}$, then the fan Σ consists of faces of the cone $\mathbb{R}_{\geq 0}A$ that are contained in τ_i for some $1 \leq i \leq r$. We will examine this case more closely in Sections 4 and 5. \square

2. Rings of differential operators on \mathbb{k} -algebras realized by retracts

Fix a \mathbb{k} -algebra R and R -module M , and note that in this section we have no requirements on the characteristic of \mathbb{k} unless explicitly mentioned. The \mathbb{k} -linear differential operators $D(R, M)$ are defined

inductively by the degree i of the operator. The degree 0 differential operators are

$$D^0(R, M) := \text{Hom}_R(R, M),$$

and, for $i > 0$, the degree i differential operators are

$$D^i(R, M) := \{\delta \in \text{Hom}_{\mathbb{k}}(R, M) \mid [\delta, r] \in D^{i-1}(R, M) \text{ for all } r \in R\},$$

where $[\delta, r] := \delta \circ r - r \circ \delta$ is the commutator. The module of differential operators on M , denoted $D(R, M)$, is $\bigcup_{i \geq 0} D^i(R, M)$. We write $D(R)$ for $D(R, R)$. Note that $D(R, -)$ is a left-exact functor, and that, in particular, for any R -ideal J , $D(R, J) = \{\delta \in D(R) \mid \delta(R) \subset J\}$.

Example 2.1. If $\text{char}(\mathbb{k}) = 0$, the ring of differential operators of the polynomial ring in d variables with coefficients in \mathbb{k} is the Weyl algebra

$$W = \mathbb{k}[t_1, \dots, t_d] \langle \partial_1, \dots, \partial_d \rangle,$$

where the relations defined on the generators are $t_i t_j - t_j t_i = 0 = \partial_i \partial_j - \partial_j \partial_i$, and $\partial_i t_j - t_j \partial_i = \delta_{ij}$, the Kronecker- δ function. The ring of differential operators of the ring of Laurent polynomials in d variables is the extended Weyl algebra $\mathbb{k}[t_1^{\pm 1}, \dots, t_d^{\pm 1}] \langle \partial_1, \dots, \partial_d \rangle$ with the relations as in the ordinary Weyl algebra together with the additional relation $t_i^{-1} \partial_j - \partial_j t_i^{-1} = t_i^{-2} \delta_{ij}$. □

Let R be a \mathbb{k} -algebra realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$, as in Definition 1.1. Given $\ell \in \Lambda$, and $\delta \in D(R)$, set $\delta_\ell = \pi_\ell \circ \delta \circ \iota_\ell$. In this section, we show that the map:

$$D(R) \rightarrow \bigoplus_{\ell \in \Lambda} D(S_\ell) \quad \text{given by } \delta \mapsto (\delta_\ell := \pi_\ell \circ \delta \circ \iota_\ell \mid \ell \in \Lambda), \tag{2.2}$$

is injective and compute the ring of differential operators $D(R)$ in terms of the rings of differential operators $D(S_\ell)$ for $\ell \in \Lambda$. We do this in two ways (see Theorems 2.6 and 2.8). First, we include two lemmas.

Lemma 2.3. *Assume that the \mathbb{k} -algebra R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$. Let $\delta \in D^i(R)$. If $\ell \in \Lambda$, then $\delta_\ell \in D^i(S_\ell)$.*

Proof. We use induction on i . If $\delta \in D^0(R) = \text{Hom}_R(R, R)$, then δ is given by multiplication by a fixed element of R , say r . But then δ_ℓ is given by multiplication by $\pi_\ell(r)$, and so $\delta_\ell \in D^0(S_\ell)$.

Now assume the result is true for operators of order $i - 1$, and let $\delta \in D^i(R)$. The \mathbb{k} -linearity of δ_ℓ follows from \mathbb{k} -linearity of δ , since ι_ℓ and π_ℓ are \mathbb{k} -linear. To show that $\delta_\ell \in D^i(S_\ell)$, it is enough to verify that $[\delta_\ell, f] \in D^{i-1}(S_\ell)$ for all $f \in S_\ell$. This follows by induction since, for $f \in S_\ell$, $[\delta_\ell, f] = [\delta, \iota_\ell(f)]_\ell$.

To see this, note that $\iota_\ell \circ f$ is the map $\iota_\ell(f)\iota_\ell$ and for any $g \in R$, $\pi_\ell \circ g$ is the map $\pi_\ell(g)\pi_\ell$. Then we have

$$\begin{aligned} [\delta_\ell, f] &= \pi_\ell \circ \delta \circ \iota_\ell \circ f - f \circ \pi_\ell \circ \delta \circ \iota_\ell \\ &= \pi_\ell \circ \delta \circ \iota_\ell(f)\iota_\ell - \pi_\ell(\iota_\ell(f))\pi_\ell \circ \delta \circ \iota_\ell \\ &= \pi_\ell \circ \delta \circ \iota_\ell(f)\iota_\ell - \pi_\ell \circ \iota_\ell(f)\delta \circ \iota_\ell \\ &= \pi_\ell \circ (\delta \circ \iota_\ell(f) - \iota_\ell(f)\delta) \circ \iota_\ell \\ &= [\delta, \iota_\ell(f)]_\ell. \end{aligned} \quad \square$$

Lemma 2.4. *Assume that the \mathbb{k} -algebra R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$. Let $\delta \in D^i(R)$, $\ell \in \Lambda$, and $f \in R$. If $\pi_\ell(f) = 0$, then $\pi_\ell(\delta(f)) = 0$.*

Example 2.5. Lemma 2.4 is a key ingredient for the main results of this section and provides an easily checked necessary condition for a linear operator on R to be a differential operator on R . To see this more concretely, consider $R = \mathbb{k}[x, y]/\langle xy \rangle$. This Stanley–Reisner ring is realized by the retracts $\mathbb{k}[x] \cong R/\langle y \rangle$ and $\mathbb{k}[y] \cong R/\langle x \rangle$, where the canonical projections are denoted π_x and π_y , respectively. The ring of differential operators on R is generated as a \mathbb{k} -algebra by the operators $x^i \partial_x^j$ and $y^k \partial_y^\ell$ such that $i \geq j$ and $k \geq \ell$. To see why ∂_x is not a differential operator on R , note that $\pi_y(x) = 0$, but $\pi_y(\partial_x(x)) = \pi_y(1) \neq 0$, contradicting the conclusion of Lemma 2.4. \square

Proof of Lemma 2.4. By induction on i as before, if $i = 0$, then $\delta(f) = r \cdot f$ for a fixed $r \in R$. If $\pi_\ell(f) = 0$, it follows that $\pi_\ell(r \cdot f) = 0$, since π_ℓ is a ring homomorphism.

Now assume the result is true for operators of order $i - 1$, and let $\delta \in D^i(R)$. Suppose $\pi_\ell(f) = 0$ and let $P = \bigcap_{f \notin P_j} P_j$ where $P_j = \ker \pi_j$. Let g be a nonzero element in P such that $\pi_\ell(g) \neq 0$. Such an element exists since R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$. Then since $f \cdot g \in (\bigcap_{f \in P_i} P_i) (\bigcap_{f \notin P_j} P_j) \subseteq \bigcap_{\ell \in \Lambda} P_\ell = 0$, we have $f \cdot g = 0$ in R . By the inductive hypothesis, $\pi_\ell([\delta, g](f)) = 0$. Then

$$[\delta, g](f) = \delta(g \cdot f) - g\delta(f) = -g\delta(f),$$

where the last equality holds because $\delta(g \cdot f) = \delta(0) = 0$. Hence

$$0 = \pi_\ell([\delta, g](f)) = \pi_\ell(-g\delta(f)) = -\pi_\ell(g)\pi_\ell(\delta(f)).$$

As g was chosen so that $\pi_\ell(g) \neq 0$, it follows that $\pi_\ell(\delta(f)) = 0$ since S_ℓ is a domain. \square

We are now ready to give a first description of $D(R)$.

Theorem 2.6. *Assume that the \mathbb{k} -algebra R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$ with the injective map $\phi: R \rightarrow \bigoplus_{\ell \in \Lambda} S_\ell$. Let $\delta: R \rightarrow R$ be \mathbb{k} -linear. Then $\delta \in D^i(R)$ if and only if the following two conditions hold:*

- (1) $\delta_\ell \in D^i(S_\ell)$ for all $\ell \in \Lambda$.
- (2) If $\ell \in \Lambda$ and $\pi_\ell(f) = 0$, then $\pi_\ell(\delta(f)) = 0$.

Proof. If $\delta \in D^i(R)$, then δ satisfies Conditions (1) and (2) by Lemmas 2.3 and 2.4.

Now assume that δ is \mathbb{k} -linear and satisfies Conditions (1) and (2). Fix $f \in R$. For each $\ell \in \Lambda$, set $f_\ell = \iota_\ell(\pi_\ell(f))$. Then $\pi_\ell(f) = \pi_\ell(f_\ell)$ for all $\ell \in \Lambda$, and so $\pi_\ell(f - f_\ell) = 0$ for all $\ell \in \Lambda$. By Condition (2), we have $\pi_\ell(\delta(f - f_\ell)) = 0$. Hence,

$$\pi_\ell(\delta(f)) = \pi_\ell(\delta(f_\ell)) \quad \text{for each } \ell \in \Lambda. \tag{2.7}$$

We now show that $\delta \in D^i(R)$ by induction on i . Set $r = \delta(1)$, and assume that, for $\ell \in \Lambda$, $\delta_\ell \in D^0(S_\ell)$ by Condition (1). Note that $\delta_\ell = \pi_\ell \circ \delta \circ \iota_\ell$. As an S_ℓ -module homomorphism on $D^0(S_\ell)$, δ_ℓ is determined by the image of the identity element 1_{S_ℓ} in S_ℓ under δ_ℓ . That is, for any $g \in S_\ell$,

$$\delta_\ell(g) = \delta_\ell(1_{S_\ell} \cdot g) = \delta_\ell(1_{S_\ell})g.$$

Applying (2.7) to $f = 1$ in R , we have $\pi_\ell(\delta(1_\ell)) = \pi_\ell(\delta(1)) = \pi_\ell(r)$. On the other hand,

$$\pi_\ell(\delta(1_\ell)) = (\pi_\ell \circ \delta)(\iota_\ell(\pi_\ell(1))) = (\pi_\ell \circ \delta \circ \iota_\ell)(\pi_\ell(1)) = \delta_\ell(\pi_\ell(1)) = \delta_\ell(1_{S_\ell}).$$

It follows that δ_ℓ is given by multiplication by $\pi_\ell(r)$ for each $\ell \in \Lambda$. Next note that

$$\pi_\ell(\delta(f_\ell)) = \pi_\ell \circ \delta \circ \iota_\ell(\pi_\ell(f)) = \delta_\ell(\pi_\ell(f)) = \pi_\ell(r)\pi_\ell(f) = \pi_\ell(rf).$$

Combining these two observations, we have, by definition of ϕ ,

$$\phi(\delta(f)) = (\pi_\ell(\delta(f)))_{\ell \in \Lambda} = (\pi_\ell(rf))_{\ell \in \Lambda} = \phi(rf).$$

Since ϕ is injective, $\delta(f) = rf$, which implies that $\delta \in D^0(R)$.

Assume now that the result is true for operators of order $i - 1$, and let δ satisfy the required conditions for order i : (1) $\delta_\ell \in D^i(S_\ell)$ for all $\ell \in \Lambda$. (2) If $\ell \in \Lambda$ and $\pi_\ell(g) = 0$ then $\pi_\ell(\delta(g)) = 0$.

We need to show that $[\delta, f] \in D^{i-1}(R)$ for $f \in R$. By induction, it suffices to show that Conditions (1) and (2) hold for $[\delta, f]$. To address Condition (1), fix $\ell \in \Lambda$. Since $[\delta, f]_\ell = \pi_\ell \circ [\delta, f] \circ \iota_\ell$, we will show that $\pi_\ell \circ [\delta, f] \circ \iota_\ell \in D^{i-1}(S_\ell)$ for all $\ell \in \Lambda$. Let us now expand:

$$\begin{aligned} \pi_\ell \circ [\delta, f] \circ \iota_\ell &= \pi_\ell \circ (\delta \circ f - f \delta) \circ \iota_\ell \\ &= \pi_\ell \circ \delta \circ f \circ \iota_\ell - \pi_\ell(f \delta) \circ \iota_\ell \\ &= \pi_\ell \circ \delta \circ f_\ell \circ \iota_\ell - \pi_\ell(f)\pi_\ell \circ \delta \circ \iota_\ell && \text{by (2.7)} \\ &= \pi_\ell \circ \delta \circ \iota_\ell(\pi_\ell(f)) \circ \iota_\ell - \pi_\ell(f)\pi_\ell \circ \delta \circ \iota_\ell && \text{by the definition of } f_\ell \\ &= \pi_\ell \circ \delta \circ \iota_\ell \circ \pi_\ell(f) - \pi_\ell(f)\pi_\ell \circ \delta \circ \iota_\ell && \text{as } \iota_\ell \text{ is a homomorphism} \\ &= \delta_\ell \circ \pi_\ell(f) - \pi_\ell(f)\delta_\ell && \text{by the definition of } \delta_\ell \\ &= [\delta_\ell, \pi_\ell(f)]. \end{aligned}$$

Since $\delta_\ell \in D^i(S_\ell)$ by assumption, $[\delta_\ell, \pi_\ell(f)] \in D^{i-1}(S_\ell)$ by the definition of an order i differential operator.

To address Condition (2), assume $\ell \in \Lambda$ and $\pi_\ell(g) = 0$ for some $g \in R$. Because π_ℓ is a homomorphism and δ satisfies Condition (2), $\pi_\ell(fg) = \pi_\ell(f)\pi_\ell(g) = 0$, and so $\pi_\ell(\delta(fg)) = 0$. Then

$$\pi_\ell \circ [\delta, f](g) = \pi_\ell(\delta(fg)) - \pi_\ell(f\delta(g)) = 0 - \pi_\ell(f)\pi_\ell(\delta(g)) = 0.$$

Hence $[\delta, f]$ satisfied Condition (2) as well as Condition (1), and so, by induction, $[\delta, f] \in D^{i-1}(R)$, and then $\delta \in D^i(R)$. \square

To state our second characterization of $D(R)$, if $\lambda \subset \Lambda$, we need some additional notation. Set $S_\lambda = \bigcap_{\ell \in \lambda} S_\ell$. Since S_ℓ is an algebra retract of R for all $\ell \in \lambda$, S_λ is also an algebra retract of R in a natural way (and indeed S_λ is also an algebra retract of S_ℓ for all $\ell \in \lambda$). We define $\iota_{\lambda,\ell}$ to be the natural inclusion that identifies S_λ as a subring of S_ℓ , and $\pi_{\ell,\lambda}$ is the natural projection from S_ℓ to S_λ , where the latter is considered as a quotient of the former.

Theorem 2.8. *Assume that the \mathbb{k} -algebra R is realized by the retracts $\{S_\ell \mid \ell \in \Lambda\}$. The map*

$$D(R) \rightarrow \bigoplus_{\ell \in \Lambda} D(S_\ell) \quad \text{given by } \delta \mapsto (\delta_\ell = \pi_\ell \circ \delta \circ \iota_\ell \mid \ell \in \Lambda)$$

from (2.2) is an injective ring homomorphism. A tuple $(\rho_\ell \mid \ell \in \Lambda) \in \bigoplus_{\ell \in \Lambda} D(S_\ell)$ is in the image of (2.2) if and only if it satisfies the following two conditions:

- (a) If $\lambda \subset \Lambda$ and $j, k \in \lambda$, then $\pi_{j,\lambda} \circ \rho_j \circ \iota_{\lambda,j} = \pi_{k,\lambda} \circ \rho_k \circ \iota_{\lambda,k}$.
- (b) If $\lambda = \{j, k\}$, then $\pi_{j,\lambda}(\rho_j(f)) = 0$ if $\pi_k(f) = 0$.

Proof. Since (2.2) is given by composition and direct sums of ring homomorphisms, it is itself a ring homomorphism.

First, for $\delta \in D(R)$, its image satisfies Conditions (a) (by construction) and (b) by Condition (2) of Theorem 2.6. To prove injectivity, as well as verify the description of the image of (2.2), we construct an inverse.

For $\lambda \subset \Lambda$ and $\rho \in D(S_\lambda)$, set $\bar{\rho} = \iota_\lambda \circ \rho \circ \pi_\lambda$. When ρ satisfies Condition (b), then, by an argument similar to that at the end of the proof of Theorem 2.6, $\bar{\rho} \in D(R)$.

If $(\rho_\ell \mid \ell \in \Lambda) \in \bigoplus_{\ell \in \Lambda} D(S_\ell)$ satisfies Condition (a) and $\emptyset \neq \lambda \subset \Lambda$, choose $\ell \in \lambda$, and set $\rho_\lambda = \pi_{\ell,\lambda} \circ \rho_\ell \circ \iota_{\lambda,\ell}$. By Condition (a), ρ_λ is independent of the choice of $\ell \in \lambda$. Now let

$$\rho = \sum_{\emptyset \neq \lambda \subset \Lambda} (-1)^{|\lambda|-1} \bar{\rho}_\lambda.$$

We claim that if $(\rho_\ell \mid \ell \in \Lambda) \in \bigoplus_{\ell \in \Lambda} D(S_\ell)$ satisfies Conditions (a) and (b), then $\rho \in D(R)$, and the image of ρ under the map (2.2) is $(\rho_\ell \mid \ell \in \Lambda)$. This gives the desired inverse and finishes the proof. \square

3. Applications

In this section, we apply the results of Section 2 to compute rings of differential operators for examples of \mathbb{k} -algebras realized by retracts discussed in Section 1. Throughout this section, we assume that the characteristic of \mathbb{k} is zero.

3A. Differential operators of Stanley–Reisner rings. Traves [47] gave a nice classification of the ring of differential operators for Stanley–Reisner ring. In particular, he proved that $D(\mathbb{k}[\Delta])$ is generated as a \mathbb{k} -algebra by

$$\{t^a \partial^b \mid t^a \in P \text{ or } t^b \notin P \text{ for each minimal prime } P \text{ of } R\}.$$

where $t^a \partial^b = t_1^{a_1} \dots t_d^{a_d} \partial_1^{b_1} \dots \partial_d^{b_d}$ and $\partial_i = \frac{\partial}{\partial t_i}$.

One can see directly that this description matches our description of the ring of differential operators in terms of algebra retracts. In particular, for a Stanley–Reisner ring $\mathbb{k}[\Delta]$, recall that $\mathbb{k}[\Delta]$ is realized by the retracts $\{\mathbb{k}[F_\ell] \mid \ell \in \Lambda\}$, where $\{F_\ell \mid \ell \in \Lambda\}$ are the facets of Δ . Further, we know that the minimal primes of $\mathbb{k}[\Delta]$ are exactly those corresponding to the facets, namely $P_{F_\ell} = \langle t^a \mid a \in \mathbb{N}^d, \text{supp}(a) \not\subseteq F_\ell \rangle$; see for example [13].

For any \mathbb{k} -linear map $\delta : \mathbb{k}[\Delta] \rightarrow \mathbb{k}[\Delta]$, Theorem 2.6 tells us that $\delta \in D^i(\mathbb{k}[\Delta])$ if and only if

- (1) $\delta_\ell \in D^i(\mathbb{k}[F_\ell])$ for all $\ell \in \Lambda$ and
- (2) $\pi_\ell(\delta(f)) = 0$ for all $\ell \in \Lambda$ and $f \notin \mathbb{k}[F_\ell]$.

Now for any facet F_ℓ of Δ , notice that $\mathbb{k}[F_\ell] \cong \mathbb{k}[t_i \mid i \in F_\ell]$, so that $D(\mathbb{k}[F_\ell])$ is the standard Weyl algebra on the variables $\{t_i \mid i \in F_\ell\}$. By Condition (1), $D(\mathbb{k}[\Delta])$ must be generated by elements of the form $x^a \partial^b$.

Further, we have that $\pi_\ell(t^a \partial^b(f)) = 0$ for all $f \notin \mathbb{k}[F_\ell]$ if and only if $t^a \notin \mathbb{k}[F_\ell]$ or $\partial^b(f) = 0$ for all $f \notin \mathbb{k}[F_\ell]$. This happens if and only if $\text{supp}(a) \not\subseteq F_\ell$ or $\text{supp}(b) \subseteq F_\ell$. In other words, we have Condition (2) if and only if $t^a \in P_{F_\ell}$ or $t^b \notin P_{F_\ell}$ for every $\ell \in \Lambda$.

3B. Differential operators of toric face rings. The algebra retracts of toric face rings that we will consider are affine semigroup rings. Saito and Traves [42] described the ring of differential operators for an affine semigroup ring $\mathbb{k}[\mathbb{N}A]$ over the complex numbers, when viewed as a subring of the ring of differential operators of the Laurent polynomials, i.e.,

$$D(\mathbb{k}[\mathbb{Z}^d]) = \mathbb{k}[t_1^{\pm 1}, \dots, t_d^{\pm 1}](\partial_1, \dots, \partial_d),$$

where ∂_i denotes the differential operator $\frac{\partial}{\partial t_i}$. Setting $\theta_j = t_j \partial_j$ for $1 \leq j \leq d$ and noting that $\theta_i \theta_j = \theta_j \theta_i$ for all i and j , the ring $\mathbb{k}[\theta] = \mathbb{k}[\theta_1, \theta_2, \dots, \theta_d]$ is a polynomial ring. Set

$$\Omega(\mathbf{m}) := \{\mathbf{a} \in \mathbb{N}A \mid \mathbf{a} + \mathbf{m} \notin \mathbb{N}A\} = \mathbb{N}A \setminus (-\mathbf{m} + \mathbb{N}A).$$

The idealizer of $\Omega(\mathbf{m})$ is defined to be the $\mathbb{k}[\theta]$ -ideal

$$\mathbb{I}(\Omega(\mathbf{m})) := \langle f(\theta) \in \mathbb{k}[\theta] \mid f(\mathbf{a}) = 0 \text{ for all } \mathbf{a} \in \Omega(\mathbf{m}) \rangle,$$

with the θ_i of degree $\mathbf{0}$. In fact, $\mathbb{1}(\Omega(\mathbf{m}))$ consists of $f(\theta)$ such that $t^{\mathbf{m}} f(\theta) \in D(\mathbb{k}[\mathbb{N}A])$. This is a consequence of the work of Saito and Traves [42, Theorem 2.1], where they show that

$$D(\mathbb{k}[\mathbb{N}A]) = \bigoplus_{\mathbf{m} \in \mathbb{Z}^d} t^{\mathbf{m}} \cdot \mathbb{1}(\Omega(\mathbf{m})).$$

To compute $\mathbb{1}(\Omega(\mathbf{m}))$ for a normal semigroup ring, consider a facet σ of A , recalling that by this we mean a submonoid of $\mathbb{N}A$ whose linear span has dimension $d - 1$. The primitive integral support function (or simply support function) F_σ is the unique linear form on \mathbb{R}^d such that

$$(1) F_\sigma(\mathbb{R}_{\geq 0}A) \geq 0, \quad (2) F_\sigma(\sigma) = 0, \quad \text{and} \quad (3) F_\sigma(\mathbb{Z}^d) = \mathbb{Z}.$$

Saito and Traves used the $F_\sigma(\mathbf{m})$ to determine the precise form of $\mathbb{1}(\Omega(\mathbf{m}))$ for normal $\mathbb{k}[\mathbb{N}A]$. The ideas of the computation in [42] can be traced back to [22] and [28]. Set

$$G_{\mathbf{m}}(\theta) := \prod_{F_\sigma(\mathbf{m}) < 0} \prod_{i=0}^{-F_\sigma(\mathbf{m})-1} (F_\sigma(\theta) - i). \tag{3.1}$$

Theorem 3.2 [42, Theorem 3.2.2]. *Let R be a normal affine semigroup ring of dimension d . Let $F_{\sigma_1}, \dots, F_{\sigma_r}$ be the support functions of the facets σ_i of the semigroup defining R . Let \mathbf{m} be a multidegree in \mathbb{Z}^d . Then*

$$D(R)_{\mathbf{m}} = t^{\mathbf{m}} \cdot \langle G_{\mathbf{m}}(\theta) \rangle.$$

We now provide a graded description of $D^i(R)$ when $R = \mathbb{k}[\mathcal{M}]$ is a toric face ring. Let $\sigma \in \Sigma$, and let $\rho \in D^i(\mathbb{k}[M_\sigma])$. Then ρ is a sum of operators of order i each of which is homogeneous with respect to the natural $\mathbb{Z}\sigma$ -grading (where $\mathbb{Z}\sigma$ is the abelian group generated by M_σ). We claim that if ρ satisfies Condition (b) of Theorem 2.8, then each homogeneous component does as well. This holds since applying an operator of multidegree $\mathbf{b} \in \mathbb{Z}\sigma$ to a monomial $t^{\mathbf{a}}$ yields a (possibly zero) scalar multiple of $t^{\mathbf{a}+\mathbf{b}}$. As ρ is a finite sum of operators of multidegrees $\mathbf{b}_1, \dots, \mathbf{b}_n$, $\rho(t^{\mathbf{a}}) = 0$ if and only if the constant multiple of $t^{\mathbf{a}+\mathbf{b}_i}$ is 0 for all $1 \leq i \leq n$.

The following result then provides the final key to describe differential operators on toric face rings, as we illustrate in examples later.

Proposition 3.3. *Let $\sigma \in \mathcal{F}(\Sigma)$, and let $\rho \in D^i(\mathbb{k}[M_\sigma])$ be homogeneous of degree $\mathbf{b} \in \mathbb{Z}\sigma$ satisfying Condition (b) from Theorem 2.8. Then $\rho = t^{\mathbf{b}}q(\theta)$ for some $q \in \mathbb{k}[\theta]$ such that*

- (i) $q(\mathbf{a}) = 0$ if $\mathbf{a} + \mathbf{b} \notin M_\sigma$ when $\mathbf{a} \in M_\sigma$, and
- (ii) for $\tau \in \mathcal{F}(\Sigma) \setminus \{\sigma\}$ and $\mathbf{a} \in M_\sigma \setminus M_{\sigma \cap \tau}$ such that $\mathbf{a} + \mathbf{b} \in M_{\sigma \cap \tau}$, we have $q(\mathbf{a}) = 0$.

Proof. Condition (i) follows from the description of $D(\mathbb{k}[M_\tau])$. Condition (ii) follows from the assumption that Condition (b) of Theorem 2.8 is satisfied for ρ . □

Remark 3.4. The two conditions in the previous result arise from the multigraded nature of our descriptions of rings of differential operators. For a given multidegree \mathbf{b} , operators of this degree are spanned

by operators of the form $\delta = t^b q(\theta)$, where q is a polynomial. Then $\delta(t^a) = q(a)t^{a+b}$. If a belongs to our semigroup but $a + b$ does not, then we must have $q(a) = 0$ if δ is to be a differential operator on the semigroup ring. This is the first condition in Proposition 3.3.

Similarly, if we are working with a semigroup coming from a monoidal complex, it may happen that a belongs to a semigroup, and $a + b$ belongs to both the semigroup and some other semigroup in the complex. In this case the projection of $\delta(t^a) = q(a)t^{a+b}$ onto this second semigroup will not vanish unless $q(a) = 0$. This explains the second condition in Proposition 3.3.

For each b , the set of a at which the polynomials q are forced to vanish lie on a finite collection of translates of the linear spans of the faces of the maximal cones in the fan Σ . This explains why their vanishing ideal is generated by products of shifts of linear forms corresponding to supporting hyperplanes of those faces. □

We now include concrete examples of differential operators in the toric face ring setting.

Example 3.5. Let

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}, \quad B = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad C = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 2 \end{pmatrix}.$$

Two of the facets of the integral cone $\mathbb{N}A$ are $\sigma = \mathbb{N}B$ and $\tau = \mathbb{N}C$. Define

$$R := \frac{\mathbb{k}[\mathbb{N}A]}{P_\sigma \cap P_\tau} = \frac{\mathbb{k}[x, xy, xy^2, xz, xz^2]}{\langle x^2yz, x^2y^2z, x^2yz^2, x^2y^2z^2 \rangle}.$$

Note that there is a natural map

$$\phi: R \rightarrow \frac{R}{P_\sigma} \oplus \frac{R}{P_\tau} \cong \mathbb{k}[\mathbb{N}B] \oplus \mathbb{k}[\mathbb{N}C] \cong \mathbb{k}[x, xy, xy^2] \oplus \mathbb{k}[x, xz, xz^2].$$

The differential operators defined on the face $\sigma \cap \tau$ in each multidegree $(u, 0, 0)$ can be realized by operators in $\mathbb{k}[\mathbb{N}B] \oplus \mathbb{k}[\mathbb{N}C]$ in multidegree $((u, 0), (u, 0))$ generated by

$$(\rho_u, \delta_u) := \left(x^u \cdot \prod_{i=0}^{-2u-1} (2\theta_x - \theta_y - i), x^u \cdot \prod_{i=0}^{-2u-1} (2\theta_x - \theta_z - i) \right) \tag{3.6}$$

This is the case by Theorem 2.8 since

$$\pi_{\sigma, \sigma \cap \tau} \circ \rho_u \circ \iota_{\sigma \cap \tau, \sigma} = x^u \cdot \prod_{i=0}^{-2u-1} (2\theta_x - i) = \pi_{\tau, \sigma \cap \tau} \circ \delta_u \circ \iota_{\sigma \cap \tau, \tau}$$

and

$$\pi_{\sigma, \sigma \cap \tau}(\rho_u(x^u y^v)) = 0 \quad \text{for } (u, v) \in \sigma \setminus \tau,$$

as well as

$$\pi_{\tau, \sigma \cap \tau}(\rho_u(x^u z^v)) = 0 \quad \text{for } (u, v) \in \tau \setminus \sigma.$$

The operators on $\sigma \setminus \sigma \cap \tau$ in each multidegree $(u, v, 0)$ (for $v \neq 0$) can be realized by operators in $\mathbb{k}[\mathbb{N}B] \oplus \mathbb{k}[\mathbb{N}C]$ in multidegree $((u, v), (0, 0))$ generated by

$$(\rho_{u,v}, 0) := \left(x^u y^v \cdot \left(\prod_{i=0}^{-2u+v-1} (2\theta_x - \theta_y - i) \right) \left(\prod_{i=0}^{-v} (\theta_y - i) \right), 0 \right). \tag{3.7}$$

Since

$$\pi_{\sigma, \sigma \cap \tau}(\rho_{u,v}(x^u y^v)) = 0 \quad \text{for } (u, v) \in \sigma,$$

both Conditions (a) and (b) of Theorem 2.8 are clearly satisfied.

A similar argument shows that the operators on $\tau \setminus \sigma \cap \tau$ in each multidegree $(u, 0, v)$ (for $v \neq 0$) can be realized by operators in $\mathbb{k}[\mathbb{N}B] \oplus \mathbb{k}[\mathbb{N}C]$ in multidegree $((0, 0), (u, v))$ generated by

$$(0, \delta_{u,v}) := \left(0, x^u z^v \cdot \left(\prod_{i=0}^{-2u+v-1} (2\theta_x - \theta_z - i) \right) \left(\prod_{i=0}^{-v} (\theta_z - i) \right) \right). \tag{3.8}$$

Combinations of operators of the forms represented by (3.6), (3.7) and (3.8) also produce operators in our ring of differential operators for all $v \neq 0$. □

Example 3.9. Set

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Recall $\mathbb{N}A$ gives the integral cone of the ring $S = \mathbb{k}[\mathbb{N}A] = \mathbb{k}[x, xy, xz, xyz]$. The facets of $\mathbb{N}A$ are

$$\begin{aligned} \sigma_1 &= \mathbb{N}\langle \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2 \rangle, & \sigma_3 &= \mathbb{N}\langle \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3 \rangle, \\ \sigma_2 &= \mathbb{N}\langle \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_3 \rangle, & \sigma_4 &= \mathbb{N}\langle \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3 \rangle. \end{aligned}$$

The support functions of the facets of $\mathbb{N}A$ are $\theta_z, \theta_y, \theta_x - \theta_y, \theta_x - \theta_z$, respectively. By Theorem 3.2, the differential operators on R in multidegree \mathbf{m} are given by

$$x^{m_1} y^{m_2} z^{m_3} \cdot \left\langle \left(\prod_{i=1}^{-m_3-1} (\theta_z - i) \right) \left(\prod_{i=1}^{-m_2-1} (\theta_y - i) \right) \left(\prod_{i=1}^{-m_1+m_3-1} (\theta_x - \theta_z - i) \right) \left(\prod_{i=1}^{-m_1+m_2-1} (\theta_x - \theta_y - i) \right) \right\rangle.$$

Set

$$R = \frac{S}{P_{\sigma_1} \cap P_{\sigma_2} \cap P_{\sigma_3} \cap P_{\sigma_4}} = \frac{\mathbb{k}[x, xy, xz, xyz]}{\langle x^2 yz \rangle}.$$

Note that $T := \mathbb{k}[a, b, c, d]/\langle ad, bc \rangle \cong R$. We know the differential operators of T by [47] since T is a Stanley–Reisner ring. The generators of $D(T)$ in multidegree $-\mathbf{e}_1$ are

$$a^{-1} \theta_a (\theta_a - 1), \quad a^{-1} \theta_a \theta_b, \quad a^{-1} \theta_a \theta_c.$$

There are not enough operators in the extended Weyl algebra in three variable to express these three operators. Given the mappings

$$\begin{array}{ccc}
 D(T) & \longrightarrow & D(\mathbb{k}[a, b]) \oplus D(\mathbb{k}[a, c]) \oplus D(\mathbb{k}[b, d]) \oplus D(\mathbb{k}[c, d]) \\
 \downarrow \cong & & \downarrow \cong \\
 D(R) & \longrightarrow & D(\mathbb{k}[x, xy]) \oplus D(\mathbb{k}[x, xz]) \oplus D(\mathbb{k}[xy, xyz]) \oplus D(\mathbb{k}[xz, xyz])
 \end{array}$$

with horizontal maps as in Theorem 2.8, we see that

$$\begin{aligned}
 a^{-1}\theta_a(\theta_a-1) &\mapsto (a^{-1}\theta_a(\theta_a-1), a^{-1}\theta_a(\theta_a-1), 0, 0) \cong \left(x^{-1} \prod_{i=0}^1 (\theta_x - \theta_y - i), x^{-1} \prod_{i=0}^1 (\theta_x - \theta_z - i), 0, 0 \right) \\
 a^{-1}\theta_a\theta_b &\mapsto (a^{-1}\theta_a\theta_b, 0, 0, 0) \cong (x^{-1}(\theta_x - \theta_y)\theta_y, 0, 0, 0) \\
 a^{-1}\theta_a\theta_c &\mapsto (0, a^{-1}\theta_a\theta_c, 0, 0) \cong (0, x^{-1}(\theta_x - \theta_z)\theta_z, 0, 0).
 \end{aligned}$$

To get an operator acting as $a^{-1}\theta_a(\theta_a - 1)$, $a^{-1}\theta_a\theta_b$, or $a^{-1}\theta_a\theta_c$ in terms of the linear support functions on S , we would need fractional expressions of the forms

$$x^{-1} \cdot \frac{\prod_{i=0}^1 (\theta_x - \theta_y - i) \prod_{i=0}^1 (\theta_x - \theta_z - i)}{\prod_{i=0}^1 (\theta_x - i)}, \quad x^{-1} \cdot \frac{(\theta_x - \theta_y)\theta_y(\theta_x - \theta_z)}{\theta_x}, \quad x^{-1} \cdot \frac{(\theta_x - \theta_y)(\theta_x - \theta_z)\theta_z}{\theta_x},$$

which do not come from the extended Weyl algebra. □

4. Rings of differential operators of quotient rings

In this section, we consider a special class of toric face rings, namely quotients of normal affine semigroup rings by radical monomial ideals. Our main goal is to characterize which differential operators on the quotient arise from operators on the ambient affine semigroup ring. We build off the techniques of [44, Proposition 1.6] and provide a careful inductive argument on the number of facets of the Newton polytope in order to generalize beyond the case where the ambient ring S is regular. In Section 5, we show that, if S is regular (or even Gorenstein) and J is the interior ideal of S , then $JD(S) = D(S, J)$, which gives a direct argument that our result agrees with [44, Proposition 1.6] in that case.

For any $\delta \in D(R)$, we will use δJ to denote the set consisting of products $\delta \circ f$ of differential operators for any $f \in J$ and $\delta * f$ to denote the action $\delta(f)$ to avoid any possible confusion.

Proposition 4.1. *Let R be commutative \mathbb{k} -algebra and J be an ideal of R . Let*

$$\mathbb{I}(J) := \{ \delta \in D(R) \mid \delta * J \subset J \}.$$

The differential operators on R that induce maps on R/J are precisely those in $\mathbb{I}(J)$. Further, there is an embedding of rings

$$\frac{\mathbb{I}(J)}{D(R, J)} \hookrightarrow D(R/J).$$

Proof. The first statement follows from the universal property of quotients. In fact, for any $\delta \in D(R)$, if δ induces an operator in $D(R/J)$, i.e., a map from R/J to R/J , then we must have $\delta * J \subset J$. So, δ belongs to $\mathbb{L}(J)$ by definition.

Every operator in $\mathbb{L}(J)$ induces a differential operator from R/J to itself. Now consider the map

$$\rho: \mathbb{L}(J) \rightarrow D(R/J) \quad \text{given by } \rho(\delta) = \delta'.$$

The kernel is $\ker(\rho) = \{\delta \mid \delta * R \subseteq J\} = D(R, J)$ by [28, 1.2]. Hence ρ induces an injective map $\bar{\rho}: \mathbb{L}(J)/D(R, J) \hookrightarrow D(R/J)$ given by $\bar{\rho}(\bar{\delta}) = \delta'$, as desired. \square

Before stating the main theorem, we will need some notation. Let R be the normal affine semigroup ring defined by a matrix A . When referring to an arbitrary face or facet of A , we will use τ or σ , respectively. Every face of A can be expressed as an intersection of facets. Recall the correspondence between the \mathbb{Z}^d -graded primes of R and the faces of A . A face τ of A corresponds to the prime ideal $P_\tau := \langle t^{\mathbf{m}} \mid \mathbf{m} \in \mathbb{N}A \setminus \mathbb{N}\tau \rangle$.

Consider the \mathbb{Z}^d -graded radical monomial ideal $J := \bigcap_{i=1}^r P_{\tau_i}$, with the face $\tau_i := \bigcap_{j=1}^{k_i} \sigma_{i,j}$ for facets $\sigma_{i,j}$ of A . For a fixed facet σ of A , set

$$H_{\sigma, \mathbf{m}}(\theta) := F_\sigma(\theta) + F_\sigma(\mathbf{m}). \tag{4.2}$$

When referring to a facet $\sigma_{i,j}$ from the definition of J , we will replace σ in F_σ and $H_{\sigma, \mathbf{m}}(\theta)$ with i, j , writing instead $F_{i,j}$ and $H_{i,j, \mathbf{m}}$. Lastly, for an integer k , let $[k]$ denote the set $\{1, 2, \dots, k\}$, and let $\mathbf{j} = (j_1, j_2, \dots, j_r) \in K_J := [k_1] \times [k_2] \times \dots \times [k_r]$ denote an r -tuple of integers in the allowable range with respect to the ideal J .

Finally whenever we have a product L of linear factors, let $\text{rad}(L)$ denote the (monic) generator of the radical of the ideal generated by L ; in other words, $\text{rad}(L)$ is a product of distinct linear polynomials in θ .

Theorem 4.3. *Let R be a normal affine semigroup ring defined by the $d \times n$ matrix A , and let $J = \bigcap_{i=1}^r P_{\tau_i}$ be the radical monomial ideal corresponding to the faces $\tau_i = \bigcap_{j=1}^{k_i} \sigma_{i,j}$. Let $G_{\mathbf{m}}(\theta)$ and $H_{i,j, \mathbf{m}}(\theta)$ be as in (3.1) and (4.2), respectively. Then for $\mathbf{m} \in \mathbb{Z}^d$,*

$$\left[\frac{\mathbb{L}(J)}{D(R, J)} \right]_{\mathbf{m}} = t^{\mathbf{m}} \cdot \frac{\langle G_{\mathbf{m}}(\theta) \cdot \text{rad}(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) < 0} H_{i,j, \mathbf{m}}(\theta)) \mid \mathbf{j} \in K_J \rangle}{\langle G_{\mathbf{m}}(\theta) \cdot \text{rad}(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) \leq 0} H_{i,j, \mathbf{m}}(\theta)) \mid \mathbf{j} \in K_J \rangle},$$

which is precisely the contribution from $D(R)$ within the \mathbf{m} -th graded piece of $D(R/J)$ induced by the operators in $D(R)$.

To clarify notation, we will consider an example before proceeding with the proof of Theorem 4.3.

Example 4.4. Let $R = \mathbb{k}[x_1, x_2, x_3]$ and $J = \langle x_1x_2, x_1x_3 \rangle$, in which case $r = 2$, $k_1 = 1$, $k_2 = 2$, and $K_J = \{1\} \times \{1, 2\}$. The prime decomposition of J is $J = \langle x_1 \rangle \cap \langle x_2, x_3 \rangle = P_{\sigma_{1,1}} \cap P_{\sigma_{2,1} \cap \sigma_{2,2}}$, and an element of K_J corresponds to a choice of one prime from the set $\{\langle x_1 \rangle\}$ and one prime from the set $\{\langle x_2 \rangle, \langle x_3 \rangle\}$.

In the notation of Theorem 4.3,

$$\mathbb{L}(J)_m = x^m \cdot \left\langle G_m(\theta) \prod_{\substack{F_{1,1}(m) < 0, \\ F_{2,1}(m) < 0}} H_{1,1,m}(\theta) H_{2,1,m}(\theta), G_m(\theta) \prod_{\substack{F_{1,1}(m) < 0, \\ F_{2,2}(m) < 0}} H_{1,1,m}(\theta) H_{2,2,m}(\theta) \right\rangle.$$

It is the presence of $H_{1,1,m}(\theta)$ as a factor of each generator f of $\mathbb{L}(J)_m$ that guarantees that whenever $x^{m'} \in J$, we have $f * x^{m'} \in \langle x_1 \rangle$. Similarly, it is the presence of either $H_{2,1,m}(\theta)$ or $H_{2,2,m}(\theta)$ that guarantees that $f * x^{m'} \in \langle x_2, x_3 \rangle$. Similarly, we have

$$D(R, J)_m = x^m \cdot \left\langle G_m(\theta) \prod_{\substack{F_{1,1}(m) \leq 0, \\ F_{2,1}(m) \leq 0}} H_{1,1,m}(\theta) H_{2,1,m}(\theta), G_m(\theta) \prod_{\substack{F_{1,1}(m) \leq 0, \\ F_{2,2}(m) \leq 0}} H_{1,1,m}(\theta) H_{2,2,m}(\theta) \right\rangle.$$

As in the general formula, the difference between the computations of $\mathbb{L}(J)_m$ and $D(R, J)_m$ is seen in the difference between the strict inequalities $F_{i,j}(m) < 0$ of $\mathbb{L}(J)_m$ and the weak inequalities $F_{i,j}(m) \leq 0$ of $D(R, J)_m$. Finally, the differential operators in $D(R)_m$ which induce maps on R/J are precisely those in $\mathbb{L}(J)_m$ making $\mathbb{L}(J)_m/D(R, J)_m$ a submodule of $D(R/J)_m$, which respects the grading on numerator and denominator. □

Proof of Theorem 4.3. Observe that

$$t^m \cdot \mathbb{L}(\Omega(m)) = t^m \cdot \left\langle \prod_{F_\sigma(m) < 0} \prod_{i=0}^{-F_\sigma(m)-1} (F_\sigma(\theta) - i) \right\rangle.$$

Now for each $m' \in \mathbb{Z}^d$, $t^{m'} \in J$ if and only if $F_\sigma(m') > 0$ for all facets σ of A . Because $\theta_i * t^{m'} = m'_i t^{m'}$ for each i ,

$$\left[t^m \cdot \prod_{F_\sigma(m) < 0} \prod_{i=0}^{-F_\sigma(m)-1} (F_\sigma(\theta) - i) \right] * t^{m'} = t^{m+m'} \cdot \prod_{F_\sigma(m) < 0} \prod_{i=0}^{-F_\sigma(m)-1} (F_\sigma(m') - i).$$

Hence,

$$[t^m \cdot G_m(\theta)] * t^{m'} = t^{m+m'} \cdot G_m(m') \in J$$

exactly when at least one of the following two conditions is satisfied:

- (1) $t^{m+m'} \in J$.
- (2) $G_m(m') = 0$.

First, for each m' such that $t^{m'} \in J$, we must have that $F_\sigma(m') + F_\sigma(m) = F_\sigma(m + m') \geq 0$ for all facets σ and that, for each $i \in [r]$, there exists some $j \in [k_i]$ so that $F_{i,j}(m + m') > 0$. We consider two conditions:

- (i) m satisfies $F_{\sigma'}(m) < 0$ for some facet σ' .
- (ii) $F_{\sigma'}(m) \geq 0$ for all facets σ' and there exists some $i \in [r]$ for which $F_{i,j}(m + m') = 0$ for all $j \in [k_i]$.

We claim that if either Condition (i) or Condition (ii) holds, then there exists some \mathbf{m}' with $t^{\mathbf{m}'} \in J$ and $G_{\mathbf{m}}(\mathbf{m}') \neq 0$. If Condition (i) holds, then $G_{\mathbf{m}}(\mathbf{m}') \neq 0$ for exactly the \mathbf{m}' that satisfy $F_{\sigma'}(\mathbf{m}) = -F_{\sigma'}(\mathbf{m}')$ and $F_{\sigma}(\mathbf{m}') \gg F_{\sigma}(\mathbf{m})$ for all facets $\sigma \neq \sigma'$. However, for all such \mathbf{m} , $H_{\sigma',\mathbf{m}}(\mathbf{m}')G_{\mathbf{m}}(\mathbf{m}') = 0$ for all \mathbf{m}' that satisfy $F_{\sigma'}(\mathbf{m}) = -F_{\sigma'}(\mathbf{m}')$ and $F_{\sigma}(\mathbf{m}') \gg -F_{\sigma}(\mathbf{m})$ for all facets $\sigma \neq \sigma'$.

If Condition (ii) holds, then $G_{\mathbf{m}}(\mathbf{m}')$ will fail to vanish exactly for the \mathbf{m}' for which there exists an i with $F_{i,j}(\mathbf{m}) = -F_{i,j}(\mathbf{m}')$ for all $j \in [k_i]$ and $F_{i',j}(\mathbf{m}') \gg F_{i',j}(\mathbf{m})$ for all facets $\sigma_{i',j}$ with $i \neq i'$. However, for all such \mathbf{m} , $H_{i,j,\mathbf{m}}(\mathbf{m}')G_{\mathbf{m}}(\mathbf{m}') = 0$ for all $i \in [r]$, $j \in [k_i]$, and \mathbf{m}' satisfying the hypotheses that $F_{i,j}(\mathbf{m}) = -F_{i,j}(\mathbf{m}')$ for all $j \in [k_i]$ and $F_{i',j}(\mathbf{m}') \gg F_{i',j}(\mathbf{m})$ for all facets $\sigma_{i',j}$ for $i \neq i'$.

Combining these calculations,

$$\mathbb{1}(J)_{\mathbf{m}} = t^{\mathbf{m}} \cdot \left\langle G_{\mathbf{m}}(\theta) \cdot \text{rad} \left(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) < 0} H_{i,j,\mathbf{m}}(\theta) \right) \mid \mathbf{j} \in K_J \right\rangle.$$

We compute $D(R, J)$ similarly. Now we begin with an arbitrary $\mathbf{m} \in \mathbb{Z}^d$ and $t^{\mathbf{m}'} \in R$. The only distinction between this calculation and the previous calculations for $\mathbb{1}(J)$ is that \mathbf{m}' is now taken from a larger set. Namely, $F_{\sigma}(\mathbf{m}')$ can now be 0 as well. Then, Condition (1) holds when $F_{\sigma}(\mathbf{m} + \mathbf{m}') > 0$ whenever $t^{\mathbf{m}'} \in R$, a condition automatically satisfied when $F_{\sigma}(\mathbf{m}) > 0$. Again, in order to address Condition (2), we consider the two Conditions (i) and (ii), above. We note that, by the same argument used to compute $\mathbb{1}(J)$, if either Condition (i) or Condition (ii) holds, then there exists some vector $\mathbf{j} \in K_J$ for which

$$G_{\mathbf{m}}(\mathbf{m}') \cdot \text{rad} \left(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) < 0} H_{i,j,\mathbf{m}}(\mathbf{m}') \right)$$

is nonzero for some \mathbf{m}' with $t^{\mathbf{m}'} \in R$. Hence

$$D(R, J)_{\mathbf{m}} = t^{\mathbf{m}} \cdot \left\langle G_{\mathbf{m}}(\theta) \cdot \text{rad} \left(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) \leq 0} H_{i,j,\mathbf{m}}(\theta) \right) \mid \mathbf{j} \in K_J \right\rangle.$$

It now follows from Proposition 4.1 that

$$\frac{\mathbb{1}(J)_{\mathbf{m}}}{D(R, J)_{\mathbf{m}}} = t^{\mathbf{m}} \cdot \frac{\left\langle G_{\mathbf{m}}(\theta) \cdot \text{rad} \left(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) < 0} H_{i,j,\mathbf{m}}(\theta) \right) \mid \mathbf{j} \in K_J \right\rangle}{\left\langle G_{\mathbf{m}}(\theta) \cdot \text{rad} \left(\prod_{i=1}^r \prod_{F_{i,j}(\mathbf{m}) \leq 0} H_{i,j,\mathbf{m}}(\theta) \right) \mid \mathbf{j} \in K_J \right\rangle},$$

as desired. □

5. Characterizing Gorenstein rings via differential operators

In this section, we compare $D(R, J)$ and $JD(R)$ when R is a normal affine semigroup ring. We find that the equality $D(R, J) = JD(R)$ holds if J is a principal monomial ideal (see Proposition 5.5). We then restrict to the special case of $J = \omega_R$, the intersection of all graded height one prime ideals of R . That is, ω_R is the defining ideal of the union of all facets. We choose the notation ω_R for this ideal, sometimes called the interior ideal, because it is a canonical module for R ; see, for example, [8, Proposition 8.2.9].

Recall that R is Gorenstein if and only if ω_R is principal; see, for example, [5, Theorem 3.3.7]. The main result of the section is as follows:

Theorem 5.1. *Let R be a normal affine semigroup ring, and let ω_R be the intersection of all graded height one prime ideals of R . Then R is Gorenstein if and only if $\omega_R D(R) = D(R, \omega_R)$.*

Before giving the proof of Theorem 5.1, we include two examples.

Example 5.2. Let

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \end{bmatrix}, \quad R = \mathbb{k}[\mathbb{N}A] = \mathbb{k}[s, st, st^2] \quad \text{and} \quad J = \omega_R = \langle st \rangle.$$

Note that R is Gorenstein since ω_R is principal. We will see in this case that $\omega_R D(R) = D(R, \omega_R)$.

We denote the faces of A by $\sigma_{1,1}$ and $\sigma_{2,1}$, and so that the primitive integral support functions are

$$F_{1,1}(\theta) = \theta_2 \quad \text{and} \quad F_{2,1}(\theta) = 2\theta_1 - \theta_2.$$

Then recall that

$$H_{1,1,m}(\theta) := F_{1,1}(\theta) + F_{1,1}(\mathbf{m}) \quad \text{and} \quad H_{2,1,m}(\theta) := F_{2,1}(\theta) + F_{2,1}(\mathbf{m})$$

and

$$G_{\mathbf{m}}(\theta) := \prod_{i=0}^{-F_{1,1}(\mathbf{m})-1} (F_{1,1}(\theta) - i) \prod_{j=0}^{-F_{2,1}(\mathbf{m})-1} (F_{2,1}(\theta) - j).$$

By (the proof of) Theorem 4.3,

$$\begin{aligned} D(R, \omega_R)_{\mathbf{m}} &= s^{m_1} t^{m_2} \left\langle G_{\mathbf{m}}(\theta) \cdot \left(\prod_{i=1}^2 \prod_{F_{i,1}(\mathbf{m}) \leq 0} H_{i,1,m}(\theta) \right) \right\rangle \\ &= s^{m_1} t^{m_2} \left\langle \prod_{i=0}^{-F_{1,1}(\mathbf{m})} (F_{1,1}(\theta) - i) \prod_{j=0}^{-F_{2,1}(\mathbf{m})} (F_{2,1}(\theta) - j) \right\rangle \\ &= s^{m_1} t^{m_2} \left\langle \prod_{i=0}^{-m_2} (\theta_2 - i) \prod_{j=0}^{-2m_1+m_2} (2\theta_1 - \theta_2 - j) \right\rangle, \end{aligned}$$

where, by convention, an empty product is 1 and $\langle - \rangle$ denotes an ideal in $\mathbb{k}[\theta_1, \theta_2]$. Multiplying the expression for $D(R)$ given in Theorem 3.2 by ω_R , we obtain

$$\omega_R D(R) = \langle st \rangle \cdot \bigoplus_{m \in \mathbb{Z}^2} s^{m_1} t^{m_2} \cdot \langle G_{\mathbf{m}}(\theta) \rangle.$$

We define $\mathbf{1} := (1, 1)$. Then we have

$$\begin{aligned}
 (\omega_R D(R))_{\mathbf{m}} &= s^{m_1} t^{m_2} \cdot \langle G_{\mathbf{m}-\mathbf{1}}(\theta) \rangle \\
 &= s^{m_1} t^{m_2} \cdot \left\langle \prod_{i=0}^{-F_{1,1}(\mathbf{m}-\mathbf{1})-1} (F_{1,1}(\theta) - i) \prod_{i=0}^{-F_{2,1}(\mathbf{m}-\mathbf{1})-1} (F_{2,1}(\theta) - i) \right\rangle \\
 &= s^{m_1} t^{m_2} \cdot \left\langle \prod_{i=0}^{-F_{1,1}(\mathbf{m})+F_{1,1}(\mathbf{1})-1} (F_{1,1}(\theta) - i) \prod_{i=0}^{-F_{2,1}(\mathbf{m})+F_{2,1}(\mathbf{1})-1} (F_{2,1}(\theta) - i) \right\rangle \\
 &= s^{m_1} t^{m_2} \cdot \left\langle \prod_{i=0}^{-F_{1,1}(\mathbf{m})} (F_{1,1}(\theta) - i) \prod_{i=0}^{-F_{2,1}(\mathbf{m})} (F_{2,1}(\theta) - i) \right\rangle \\
 &= s^{m_1} t^{m_2} \left\langle \prod_{i=0}^{-m_2} (\theta_2 - i) \prod_{j=0}^{-2m_1+m_2} (2\theta_1 - \theta_2 - j) \right\rangle.
 \end{aligned}$$

It follows that $\omega_R D(R) = D(R, \omega_R)$. □

Example 5.3. Let

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \end{bmatrix}, \quad R = \mathbb{k}[\mathbb{N}A] = \mathbb{k}[s, st, st^2, st^3] \quad \text{and} \quad J = \omega_R = \langle st, st^2 \rangle.$$

Note that R is not Gorenstein. We denote the faces of A by $\sigma_{1,1}$ and $\sigma_{2,1}$, and so that the primitive integral support functions are

$$F_{1,1}(\theta) = \theta_2 \quad \text{and} \quad F_{2,1}(\theta) = 3\theta_1 - \theta_2.$$

Then for $\mathbf{m} = (m_1, m_2)$, we have

$$H_{1,1,\mathbf{m}}(\theta) = F_{1,1}(\theta) + F_{1,1}(\mathbf{m}) \quad \text{and} \quad H_{2,1,\mathbf{m}}(\theta) = F_{2,1}(\theta) + F_{2,1}(\mathbf{m})$$

and

$$G_{\mathbf{m}}(\theta) = \prod_{i=0}^{-F_{1,1}(\mathbf{m})-1} (F_{1,1}(\theta) - i) \prod_{i=0}^{-F_{2,1}(\mathbf{m})-1} (F_{2,1}(\theta) - i).$$

Multiplying the expression for $D(R)$ given in Theorem 3.2 by ω_R , we obtain

$$\omega_R D(R) = \langle st, st^2 \rangle \cdot \bigoplus_{\mathbf{m} \in \mathbb{Z}^2} s^{m_1} t^{m_2} \cdot \langle G_{\mathbf{m}}(\theta) \rangle,$$

so that

$$(\omega_R D(R))_{\mathbf{m}} = s^{m_1} t^{m_2} \cdot \langle G_{\mathbf{m}-(1,1)}(\theta), G_{\mathbf{m}-(1,2)}(\theta) \rangle,$$

where

$$\begin{aligned}
 G_{\mathbf{m}-(1,1)}(\theta) &= \prod_{j=0}^{-F_{1,1}(\mathbf{m}-(1,1))-1} (F_{1,1}(\theta) - j) \prod_{j=0}^{-F_{2,1}(\mathbf{m}-(1,1))-1} (F_{2,1}(\theta) - j) \\
 &= \prod_{j=0}^{-F_{1,1}(\mathbf{m})} (F_{1,1}(\theta) - j) \prod_{j=0}^{-F_{2,1}(\mathbf{m})+1} (F_{2,1}(\theta) - j) \\
 &= \prod_{j=0}^{-m_2} (\theta_2 - j) \prod_{j=0}^{-3m_1+m_2+1} (3\theta_2 - \theta_1 - j)
 \end{aligned}$$

when $F_{1,1}(\mathbf{m})=m_2 \leq 0$ and $F_{2,1}(\mathbf{m})=3m_1 - m_2 \leq 1$ and

$$\begin{aligned}
 G_{\mathbf{m}-(1,2)}(\theta) &= \prod_{j=0}^{-F_{1,1}(\mathbf{m}-(1,2))-1} (F_{1,1}(\theta) - j) \prod_{j=0}^{-F_{2,1}(\mathbf{m}-(1,2))-1} (F_{2,1}(\theta) - j) \\
 &= \prod_{j=0}^{-F_{1,1}(\mathbf{m})+1} (F_{1,1}(\theta) - j) \prod_{j=0}^{-F_{2,1}(\mathbf{m})} (F_{2,1}(\theta) - j) \\
 &= \prod_{j=0}^{-m_2+1} (\theta_2 - j) \prod_{j=0}^{-3m_1+m_2} (3\theta_1 - \theta_2 - j)
 \end{aligned}$$

when $F_{1,1}(\mathbf{m})=m_2 \leq 1$ and $F_{2,1}(\mathbf{m})=3m_1 - m_2 \leq 0$.

On the other hand, by Theorem 4.3 and a computation similar to the one in Example 5.2,

$$\begin{aligned}
 D(R, \omega_R)_{\mathbf{m}} &= s^{m_1} t^{m_2} \cdot \left\langle \prod_{i=0}^{-F_{1,1}(\mathbf{m})} (F_{1,1}(\theta) - i) \prod_{i=0}^{-F_{2,1}(\mathbf{m})} (F_{2,1}(\theta) - i) \right\rangle \\
 &= s^{m_1} t^{m_2} \cdot \left\langle \prod_{i=0}^{-m_2} (\theta_2 - i) \prod_{i=0}^{-3m_1+m_2} (3\theta_1 - \theta_2 - i) \right\rangle
 \end{aligned}$$

when $F_{1,1}(\mathbf{m})=m_2 \leq 0$ and $F_{2,1}(\mathbf{m})=3m_1 - m_2 \leq 0$.

To see $\omega_R D(R)$ and $D(R, \omega_R)$ are different, consider any degree, say $\mathbf{m} = (-1, -1)$. Note that the degree \mathbf{m} piece of $D(R, \omega_R)$ is generated by one element, namely

$$s^{-1} t^{-1} \prod_{i=0}^1 (F_{1,1}(\theta) - i) \prod_{i=0}^2 (F_{2,1}(\theta) - i)$$

On the other hand, the degree \mathbf{m} piece of $\omega_R D(R)$ is generated by two elements; it is generated in $\mathbb{k}[\theta_1, \theta_2]$ by

$$s^{-1} t^{-1} \prod_{i=0}^1 (F_{1,1}(\theta) - i) \prod_{i=0}^3 (F_{2,1}(\theta) - i) \quad \text{and} \quad s^{-1} t^{-1} \prod_{i=0}^2 (F_{1,1}(\theta) - i) \prod_{i=0}^2 (F_{2,1}(\theta) - i)$$

In particular,

$$s^{-1}t^{-1} \prod_{i=0}^1 (F_{1,1}(\theta) - i) \prod_{i=0}^2 (F_{2,1}(\theta) - i) \in D(R, \omega_R)$$

but

$$s^{-1}t^{-1} \prod_{i=0}^1 (F_{1,1}(\theta) - i) \prod_{i=0}^2 (F_{2,1}(\theta) - i) \notin \omega_R D(R). \quad \square$$

In order to prove Theorem 5.1, we will need some preliminary results.

Lemma 5.4 [4, Theorem 6.33]. *Let R be a normal affine semigroup ring of dimension d , and let $F_{\sigma_1}, \dots, F_{\sigma_r}$ be the support functions of the facets σ_i of the semigroup defining R . Let ω_R be the intersection of all graded height one prime ideals of R . Then ω_R is a principal ideal if and only if there exists $t^c \in \omega$ for some $c \in \mathbb{Z}^d$ such that $F_{\sigma_i}(c) = 1$ for all $i = 1, \dots, k$.*

We first show that principal monomial ideals J behave well (i.e., $JD(R) = D(R, J)$) for affine normal semigroup rings.

Proposition 5.5. *Let R be a normal affine semigroup ring and let $J = \langle t^c \rangle$, then*

$$JD(R) = D(R, J).$$

Proof. Since $JD(R) \subseteq D(R, J)$, it suffices to prove that $JD(R)_m = D(R, J)_m$ for all multidegrees m . We do so by showing that they are both cyclic and are generated by the same differential operator.

Since $J = \langle t^c \rangle$, by the expression obtained in Theorem 3.2, we have

$$JD(R)_m = t^m D(R)_{m-c} = t^m \langle G_{m-c}(\theta) \rangle.$$

Since F_i is linear, we have

$$F_i(-(\mathbf{m} - \mathbf{c})) - F_i(\mathbf{c}) = -F_i(\mathbf{m}) + F_i(\mathbf{c}) - F_i(\mathbf{c}) = -F_i(\mathbf{m}).$$

Hence the single generator for the graded piece $JD(R)_m$ can be viewed as

$$t^m \cdot \left\langle G_m(\theta) \prod_{i=1}^r H_{i,m}(\theta) \right\rangle,$$

which is exactly the generator of $D(R, J)_m$, completing the proof. □

Now we are ready to prove the main theorem of the section: Theorem 5.1. As above, we will write F_i and $H_{i,m}$ for F_{σ_i} and $H_{\sigma_i,m}$, respectively.

Proof of Theorem 5.1. Suppose first that R is Gorenstein. Then ω_R is principal, say $\omega_R = \langle t^c \rangle$. Now by Proposition 5.5, we obtain $\omega_R D(R) = D(R, \omega_R)$.

Conversely assume that R is not Gorenstein, that is, ω_R is not principal. Since $D(R, \omega_R)$ and $\omega_R D(R)$ are both multigraded and it is always true that $\omega_R D(R) \subseteq D(R, \omega_R)$, we may prove that they are not equal by identifying their difference at $m = \mathbf{0}$.

As mentioned above, $D(R, \omega_R)_m$ is generated by a single element $t^m \langle G_m(\theta) \prod_{i=1}^r H_{i,m}(\theta) \rangle$. Thus

$$D(R, \omega_R)_0 = \left\langle \prod_{i=1}^r H_{i,0}(\theta) \right\rangle = \langle F_1(\theta) \cdots F_r(\theta) \rangle.$$

Let ω_R be minimally generated by $t^{c_1}, t^{c_2}, \dots, t^{c_h}$. Then

$$(\omega D(R))_m = \sum_{j=1}^h t^{c_j} t^{m-c_j} \cdot D(R)_{m-c_j} = \sum_{j=1}^h t^m \cdot \langle G_{m-c_j}(\theta) \rangle = t^m \cdot \langle G_{m-c_j}(\theta) \mid j = 1, 2, \dots, h \rangle.$$

In particular, when $m = \mathbf{0}$,

$$(\omega D(R))_0 = \langle G_{-c_j}(\theta) \mid j = 1, 2, \dots, h \rangle.$$

We know from Lemma 5.4 that some $F_{\sigma_i}(\mathbf{c}_j) \neq 1$. Therefore

$$(\omega D(R))_0 = \sum_{j=1}^h \left\langle \prod_{i=1}^r F_i(\theta) (F_i(\theta) - 1) \cdots (F_i(\theta) - (F_i(\mathbf{c}_j) - 1)) \right\rangle \tag{5.6}$$

$$= \langle F_1(\theta) F_2(\theta) \cdots F_r(\theta) \rangle \cap \sum_{j=1}^h \left\langle \prod_{i=1}^r (F_i(\theta) - 1) \cdots (F_i(\theta) - (F_i(\mathbf{c}_j) - 1)) \right\rangle \tag{5.7}$$

We claim that the ideal

$$\sum_{j=1}^h \left\langle \prod_{i=1}^r (F_i(\theta) - 1) \cdots (F_i(\theta) - (F_i(\mathbf{c}_j) - 1)) \right\rangle \tag{5.8}$$

from (5.6) is a proper ideal of $\mathbb{k}[\theta]$. Once this is established, it will follow from (5.6) that the inclusion

$$(\omega_R D(R))_0 \subseteq D(R, \omega_R)_0 = \langle F_1(\theta) \cdots F_r(\theta) \rangle,$$

is proper, which will conclude the proof.

To see that (5.8) is a proper ideal of $\mathbb{k}[\theta]$, we may assume after reordering that \mathbf{c}_1 generates a ray of the cone

$$\mathcal{C} := \mathbb{R}_{\geq 0}\{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_h\} = \{\mathbf{m} \in \mathbb{N}A \mid F_1(\mathbf{m}) \geq 1, \dots, F_r(\mathbf{m}) \geq 1\} \subseteq \mathbb{R}^d$$

over the exponents of the minimal generators of ω_R and, in light of Lemma 5.4, that there are $d - 1$ linearly independent primitive integral support functions, which after relabeling can be taken as F_1, \dots, F_{d-1} , such that

$$F_1(\mathbf{c}_1) = F_2(\mathbf{c}_1) = \cdots = F_{d-1}(\mathbf{c}_1) = 1 \quad \text{and} \quad F_d(\mathbf{c}_1) > 1.$$

(There may be more i such that $F_i(\mathbf{c}_1) = 1$, but we need to make use of only a linearly independent set of them, which is necessarily of size $d - 1$, since \mathbf{c}_1 generates a ray of \mathcal{C} .)




Figure 1. Hypothetical generators of ω_R in dimension 3.

Example 5.9. We pause our proof to give an example illustrating the newly introduced notation. Consider the case $d = 3$ and refer to Figure 1 as a visualization. Fix \mathbf{c}_1 to be a vertex of the convex hull of the \mathbf{c}_i , or, rather, a generator of a ray of the Newton polyhedron of ω_R . Without loss of generality,

$$F_1(\mathbf{c}_1) = 1 = F_2(\mathbf{c}_1) \quad \text{and} \quad F_3(\mathbf{c}_1) > 1.$$

Further, there exists an r such that

$$F_1(\mathbf{c}_i) > 1 \quad \text{for all } r + 1 \leq i \leq h \quad \text{and} \quad F_1(\mathbf{c}_i) = 1 \quad \text{for all } 1 \leq i \leq r.$$

Note that in Figure 1, we have chosen $r = 3$. Now, since \mathbf{c}_1 is a vertex, it must be that

$$F_2(\mathbf{c}_i) > 1 \quad \text{for all } 2 \leq i \leq r.$$

Hence, $\langle F_1(\theta) - 1, F_2(\theta) - 1, F_3(\theta) - 1 \rangle$ is a primary component of (5.8). □

We resume our proof in full generality. Given that \mathbf{c}_1 is a ray of the cone \mathcal{C} with $F_i(\mathbf{c}_1) = 1$ for every i with $1 \leq i \leq d - 1$, it follows that, for every $j > 1$, there is an i_j with $1 \leq i_j \leq d - 1$ such that $F_{i_j}(\mathbf{c}_j) > 1$. Now set $i_1 = d$ because $F_d(\mathbf{c}_1) > 1$. Then since $1 \leq i_j \leq d$ for all j and the hyperplanes defined by $\{F_i(\theta) = 0\}_{i=1}^d$ are necessarily nonparallel, the ideal

$$\langle F_{i_1}(\theta) - 1, F_{i_2}(\theta) - 1, \dots, F_{i_d}(\theta) - 1 \rangle$$

is a proper, primary (in fact, prime) component of (5.8). Thus (5.8) is a proper ideal of $\mathbb{k}[\theta]$, as desired to establish the final needed claim. □

Acknowledgements

We would like to thank the organizers, Karen Smith, Sandra Spiroff, Irena Swanson, and Emily Witt, of *Workshop: Women in Commutative Algebra (WICA)*, where this work commenced; additionally, we appreciate the hospitality of BIRS in hosting this meeting. We thank AIM for supporting our collaboration, which greatly helped the advancement of this project. We are grateful to Jack Jeffries for valuable

discussions that clarified our results; we also thank Monica Lewis, Pasha Pylyavskyy, and Vic Reiner for helpful conversations. Finally, we thank the anonymous referee for feedback on an earlier version of our manuscript.

References

- [1] J. Àlvarez Montaner, D. J. Hernández, J. Jeffries, L. Núñez Betancourt, P. Teixeira, and E. E. Witt, “Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands”, *Commun. Contemp. Math.* **24**:10 (2022), art. id. 2150083. MR Zbl
- [2] H. Brenner, J. Jeffries, and L. Núñez Betancourt, “Quantifying singularities with differential operators”, *Adv. Math.* **358** (2019), art. id. 106843. MR Zbl
- [3] W. Bruns and J. Gubeladze, “Polytopal linear retractions”, *Trans. Amer. Math. Soc.* **354**:1 (2002), 179–203. MR Zbl
- [4] W. Bruns and J. Gubeladze, *Polytopes, rings, and K -theory*, Springer, 2009. MR Zbl
- [5] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, 1993. MR
- [6] A. Conca and J. Herzog, “Ladder determinantal rings have rational singularities”, *Adv. Math.* **132**:1 (1997), 120–147. MR Zbl
- [7] D. L. Costa, “Retracts of polynomial rings”, *J. Algebra* **44**:2 (1977), 492–502. MR Zbl
- [8] D. A. Cox, J. B. Little, and H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics **124**, American Mathematical Society, Providence, RI, 2011. MR Zbl
- [9] A. De Stefani, E. Grifo, and J. Jeffries, “A Zariski–Nagata theorem for smooth \mathbb{Z} -algebras”, *J. Reine Angew. Math.* **761** (2020), 123–140. MR
- [10] N. Epstein and H. D. Nguyen, “Algebra retracts and Stanley–Reisner rings”, *J. Pure Appl. Algebra* **218**:9 (2014), 1665–1682. MR Zbl
- [11] E. Eriksen, “Differential operators on monomial curves”, *J. Algebra* **264**:1 (2003), 186–198. MR Zbl
- [12] A. Eriksson, “The ring of differential operators of a Stanley–Reisner ring”, *Comm. Algebra* **26**:12 (1998), 4007–4013. MR Zbl
- [13] C. A. Francisco, J. Mermin, and J. Schweig, “A survey of Stanley–Reisner theory”, pp. 209–234 in *Connections between algebra, combinatorics, and geometry*, edited by S. M. Cooper and S. Sather-Wagstaff, Springer Proc. Math. Stat. **76**, Springer, 2014. MR Zbl
- [14] N. Gupta, “On the cancellation problem for the affine space \mathbb{A}^3 , in characteristic p ”, *Invent. Math.* **195**:1 (2014), 279–288. MR Zbl
- [15] J. Herzog, “Algebra retracts and Poincaré-series”, *Manuscripta Math.* **21**:4 (1977), 307–314. MR Zbl
- [16] M. Hochster and J. Jeffries, “Extensions of primes, flatness, and intersection flatness”, pp. 63–81 in *Commutative algebra—150 years with Roger and Sylvia Wiegand*, edited by N. R. Baeth et al., Contemp. Math. **773**, Amer. Math. Soc., Providence, RI, 2021. MR Zbl
- [17] B. Ichim and T. Römer, “On toric face rings”, *J. Pure Appl. Algebra* **210**:1 (2007), 249–266. MR Zbl
- [18] B. Ichim and T. Römer, “On canonical modules of toric face rings”, *Nagoya Math. J.* **194** (2009), 69–90. MR Zbl
- [19] M.-N. Ishida, “The local cohomology groups of an affine semigroup ring”, pp. 141–153 in *Algebraic geometry and commutative algebra*, vol. 1, edited by H. Hijikata et al., Kinokuniya, Tokyo, 1988. MR Zbl
- [20] J. Jeffries and A. K. Singh, “Differential operators on classical invariant rings do not lift modulo p ”, *Adv. Math.* **432** (2023), art. id. 109276. MR
- [21] J. Jeffries, L. Núñez Betancourt, and E. Quinlan-Gallego, “Bernstein–Sato theory for singular rings in positive characteristic”, *Trans. Amer. Math. Soc.* **376**:7 (2023), 5123–5180. MR Zbl
- [22] A. G. Jones, “Rings of differential operators on toric varieties”, *Proc. Edinburgh Math. Soc.* (2) **37**:1 (1994), 143–160. MR Zbl

- [23] A. C. Lőrincz and M. Perlman, “Equivariant \mathcal{D} -modules on alternating senary 3-tensors”, *Nagoya Math. J.* **243** (2021), 61–82. MR
- [24] A. C. Lőrincz and C. Raicu, “Iterated local cohomology groups and Lyubeznik numbers for determinantal rings”, *Algebra Number Theory* **14**:9 (2020), 2533–2569. MR Zbl
- [25] A. C. Lőrincz and U. Walther, “On categories of equivariant \mathcal{D} -modules”, *Adv. Math.* **351** (2019), 429–478. MR Zbl
- [26] J. L. Muhasky, “The differential operator ring of an affine curve”, *Trans. Amer. Math. Soc.* **307**:2 (1988), 705–723. MR Zbl
- [27] I. M. Musson, “Some rings of differential operators which are Morita equivalent to the Weyl algebra A_1 ”, *Proc. Amer. Math. Soc.* **98**:1 (1986), 29–30. MR Zbl
- [28] I. M. Musson, “Rings of differential operators on invariant rings of tori”, *Trans. Amer. Math. Soc.* **303**:2 (1987), 805–827. MR Zbl
- [29] I. M. Musson, “Differential operators on toric varieties”, *J. Pure Appl. Algebra* **95**:3 (1994), 303–315. MR Zbl
- [30] I. M. Musson and M. Van den Bergh, “Invariants under tori of rings of differential operators and related topics”, pp. viii+85 in *Mem. Amer. Math. Soc.* **650**, Amer. Math. Soc., Providence, RI, 1998. MR Zbl
- [31] M. Mustață and M. Popa, “Hodge ideals”, pp. v+80 in *Mem. Amer. Math. Soc.* **1268**, Amer. Math. Soc., Providence, RI, 2019. MR Zbl
- [32] M. Mustață, G. G. Smith, H. Tsai, and U. Walther, “ \mathcal{D} -modules on smooth toric varieties”, *J. Algebra* **240**:2 (2001), 744–770. MR Zbl
- [33] D. H. Nguyen, “Seminormality and local cohomology of toric face rings”, *J. Algebra* **371** (2012), 536–553. MR Zbl
- [34] D. H. Nguyen, “On the Koszul property of toric face rings”, *J. Commut. Algebra* **6**:2 (2014), 233–259. MR Zbl
- [35] R. Okazaki and K. Yanagawa, “Dualizing complex of a toric face ring”, *Nagoya Math. J.* **196** (2009), 87–116. MR Zbl
- [36] M. Perlman, “Equivariant \mathcal{D} -modules on $2 \times 2 \times 2$ hypermatrices”, *J. Algebra* **544** (2020), 391–416. MR Zbl
- [37] M. Perlman, “Lyubeznik numbers for Pfaffian rings”, *J. Pure Appl. Algebra* **224**:5 (2020), art. id. 106247. MR Zbl
- [38] M. Perlman and C. Raicu, “Hodge ideals for the determinant hypersurface”, *Selecta Math. (N.S.)* **27**:1 (2021), art. id. 1. MR Zbl
- [39] E. Quinlan-Gallego, “Bernstein–Sato roots for monomial ideals in positive characteristic”, *Nagoya Math. J.* **244** (2021), 25–34. MR Zbl
- [40] E. Quinlan-Gallego, “Bernstein–Sato theory for arbitrary ideals in positive characteristic”, *Trans. Amer. Math. Soc.* **374**:3 (2021), 1623–1660. MR Zbl
- [41] C. Raicu, “Characters of equivariant \mathcal{D} -modules on spaces of matrices”, *Compos. Math.* **152**:9 (2016), 1935–1965. MR Zbl
- [42] M. Saito and W. N. Traves, “Differential algebras on semigroup algebras”, pp. 207–226 in *Symbolic computation: solving equations in algebra, geometry, and engineering* (South Hadley, MA, 2000), edited by E. L. Green et al., *Contemp. Math.* **286**, Amer. Math. Soc., Providence, RI, 2001. MR Zbl
- [43] M. Saito and W. N. Traves, “Finite generation of rings of differential operators of semigroup algebras”, *J. Algebra* **278**:1 (2004), 76–103. MR Zbl
- [44] S. P. Smith and J. T. Stafford, “Differential operators on an affine curve”, *Proc. London Math. Soc.* (3) **56**:2 (1988), 229–259. MR Zbl
- [45] R. Stanley, “Generalized H -vectors, intersection cohomology of toric varieties, and related results”, pp. 187–213 in *Commutative algebra and combinatorics* (Kyoto, 1985), edited by M. Nagata and H. Matsumura, *Adv. Stud. Pure Math.* **11**, North-Holland, Amsterdam, 1987. MR Zbl
- [46] H. Terao, “Algebras generated by reciprocals of linear forms”, *J. Algebra* **250**:2 (2002), 549–558. MR Zbl
- [47] W. N. Traves, “Differential operators on monomial rings”, *J. Pure Appl. Algebra* **136**:2 (1999), 183–197. MR Zbl
- [48] J. R. Tripp, “Differential operators on Stanley–Reisner rings”, *Trans. Amer. Math. Soc.* **349**:6 (1997), 2507–2523. MR Zbl
- [49] K. Yanagawa, “Dualizing complexes of seminormal affine semigroup rings and toric face rings”, *J. Algebra* **425** (2015), 367–391. MR Zbl

Communicated by Irena Peeva

Received 2022-01-31 Revised 2023-01-18 Accepted 2023-02-23

cberkes@umn.edu

*School of Mathematics, University of Minnesota, Minneapolis, MN,
United States*

chan1cj@cmich.edu

*Department of Mathematics, Central Michigan University, Mount Pleasant,
MI, United States*

pjklein@tamu.edu

*Department of Mathematics, Texas A&M University, College Station,
TX, United States*

matusevich@tamu.edu

*Department of Mathematics, Texas A&M University, College Station,
TX, United States*

janet.page@ndsu.edu

*Mathematics Department, North Dakota State University, Fargo, ND,
United States*

jvassil@unm.edu

*Department of Mathematics and Statistics, University of New Mexico,
Albuquerque, NM, United States*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR
Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR
David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2023 is US \$485/year for the electronic version, and \$705/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 17 No. 11 2023

On self-correspondences on curves JOËL BELLAÏCHE	1867
Fitting ideals of class groups for CM abelian extensions MAHIRO ATSUTA and TAKENORI KATAOKA	1901
The behavior of essential dimension under specialization, II ZINOVY REICHSTEIN and FEDERICO SCAVIA	1925
Differential operators, retracts, and toric face rings CHRISTINE BERKESCH, C-Y. JEAN CHAN, PATRICIA KLEIN, LAURA FELICIA MATUSEVICH, JANET PAGE and JANET VASSILEV	1959
Bézoutians and the \mathbb{A}^1 -degree THOMAS BRAZELTON, STEPHEN MCKEAN and SABRINA PAULI	1985
Axiomatizing the existential theory of $\mathbb{F}_q((t))$ SYLVY ANSCOMBE, PHILIP DITTMANN and ARNO FEHM	2013
The diagonal coinvariant ring of a complex reflection group STEPHEN GRIFFETH	2033