

# ANALYSIS & PDE

Volume 5

No. 1

2012

TRISTAN ROY

A REMARK ON BARELY  $\dot{H}^{s_p}$ -SUPERCRITICAL WAVE  
EQUATIONS



mathematical sciences publishers

# A REMARK ON BARELY $\dot{H}^{s_p}$ -SUPERCRITICAL WAVE EQUATIONS

TRISTAN ROY

We prove that a good  $\dot{H}^{s_p}$  critical theory for the 3D wave equation  $\partial_{tt}u - \Delta u = -|u|^{p-1}u$  can be extended to prove global well-posedness of smooth solutions of at least one 3D barely  $\dot{H}^{s_p}$ -supercritical wave equation  $\partial_{tt}u - \Delta u = -|u|^{p-1}ug(|u|)$ , with  $g$  growing slowly to infinity, provided that a Kenig-Merle type condition is satisfied. This result is related to those obtained by Tao and the author for the particular case  $s_p = 1$ , showing global regularity for  $g$  growing logarithmically with radial data and for  $g$  growing doubly logarithmically with general data.

## 1. Introduction

For fixed  $p > 3$ , let  $\tilde{H}^2 := \dot{H}^2(\mathbb{R}^3) \cap \dot{H}^{s_p}(\mathbb{R}^3)$  and  $\tilde{H}^1 := \dot{H}^1(\mathbb{R}^3) \cap \dot{H}^{s_p-1}(\mathbb{R}^3)$ , where  $s_p := \frac{3}{2} - \frac{2}{p-1}$ . We consider the wave equation

$$\begin{cases} \partial_{tt}u - \Delta u = -|u|^{p-1}ug(|u|), \\ u(0) := u_0 \in \tilde{H}^2, \\ \partial_t u(0) := u_1 \in \tilde{H}^1, \end{cases} \quad (1-1)$$

where  $u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$  is a complex-valued scalar field and  $g$  is a smooth, real-valued positive function defined on the set of nonnegative numbers and satisfying

$$0 \leq g'(x) \lesssim \frac{1}{x}. \quad (1-2)$$

This condition says that  $g$  grows more slowly than any positive power of  $u$ .

We shall see that (1-1) has many connections with the defocusing power-type wave equation

$$\begin{cases} \partial_{tt}u - \Delta u = -|u|^{p-1}u, \\ u(0) := u_0 \in \dot{H}^{s_p}(\mathbb{R}^3), \\ \partial_t u(0) := u_1 \in \dot{H}^{s_p-1}(\mathbb{R}^3). \end{cases} \quad (1-3)$$

It is known that if  $u$  satisfies (1-3), then  $u_\lambda$  defined by

$$u_\lambda(t, x) := \frac{1}{\lambda^{2/(p-1)}} u\left(\frac{t}{\lambda}, \frac{x}{\lambda}\right), \quad (1-4)$$

satisfies the same equation, but with data

$$u_\lambda(0, x) = \frac{1}{\lambda^{2/(p-1)}} u_0\left(\frac{x}{\lambda}\right) \quad \text{and} \quad \partial_t u_\lambda(0, x) = \frac{1}{\lambda^{2/(p-1)+1}} u_1\left(\frac{x}{\lambda}\right).$$

MSC2000: 35Q55.

Keywords: wave equation, global existence, barely supercritical.



Notice that (1-3) is  $\dot{H}^{s_p}(\mathbb{R}^3)$  critical, which means that the  $\dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)$ -norm of  $(u(0), \partial_t u(0))$  is invariant under the scaling defined above.

We recall the local existence theory. From [Ginibre and Velo 1989; Lindblad and Sogge 1995], we know that there exists a positive constant  $\delta := \delta(\|(u_0, u_1)\|_{\dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)}) > 0$  and a time of local existence  $T_l > 0$  such that if

$$\left\| \cos(tD)u_0 + \frac{\sin(tD)}{D} \right\|_{L_t^{2(p-1)} L_x^{2(p-1)}([0, T_l] \times \mathbb{R}^3)} \leq \delta \quad (1-5)$$

then there exists a unique solution  $(u, \partial_t u)$  in

$$\mathcal{C}([0, T_l], \dot{H}^{s_p}(\mathbb{R}^3)) \cap L_t^{2(p-1)} L_x^{2(p-1)}([0, T_l] \times \mathbb{R}^3) \cap D^{\frac{1}{2}-s_p} L_t^4 L_x^4([0, T_l] \times \mathbb{R}^3) \times \mathcal{C}([0, T_l], \dot{H}^{s_p-1}(\mathbb{R}^3))$$

of (1-3)<sup>1</sup> in the integral equation sense, i.e.,  $u$  satisfies the Duhamel formula

$$u(t) := \cos(tD)u_0 + \frac{\sin(tD)}{D}u_1 - \int_0^t \frac{\sin(t-t')D}{D}(|u|^{p-1}u)(t') dt'. \quad (1-6)$$

It follows that we can define a maximal time interval of existence  $I_{\max} = (-T_-, T_+)$ . Moreover,

$$\|u\|_{L_t^{2(p-1)} L_x^{2(p-1)}(J)} < \infty, \quad \|D^{s_p-\frac{1}{2}}u\|_{L_t^4 L_x^4(J)} < \infty, \quad \text{and} \quad \|(u, \partial_t u)\|_{L_t^\infty \dot{H}^{s_p} \times L_t^\infty \dot{H}^{s_p-1}(J)} < \infty$$

for any compact subinterval  $J \subset I_{\max}$ . See [Kenig and Merle 2006] or [Tao 2006a] for more explanations.

Now we turn to the global well-posedness theory of “(1-3)”. In view of the local well-posedness theory, one can prove (see [Kenig and Merle 2011] and references), after some effort, that it is enough to find a finite upper bound of  $\|u\|_{L_t^{2(p-1)} L_x^{2(p-1)}(I \times \mathbb{R}^3)}$  on arbitrary long time intervals  $I$ , and, if this is the case, then the solution scatters to a solution of the linear wave equation. No blow-up has been observed for (1-3). Therefore it is believed that the following scattering conjecture is true:

**Conjecture 1.1** (scattering conjecture). *Assume that  $u$  is the solution of (1-3) with data  $(u_0, u_1) \in \dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)$ . Then  $u$  exists for all time  $t$  and there exists  $C_1 := C_1(\|(u_0, u_1)\|_{\dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)})$  such that*

$$\|u\|_{L_t^{2(p-1)} L_x^{2(p-1)}(\mathbb{R} \times \mathbb{R}^3)} \leq C_1. \quad (1-7)$$

The case  $s_p = 1$  (equivalently,  $p = 5$ ) is particular. Indeed the solution

$$(u, \partial_t u) \in \mathcal{C}([0, T_l], \dot{H}^1(\mathbb{R}^3)) \times \mathcal{C}([0, T_l], L^2(\mathbb{R}^3))$$

satisfies the conservation of the energy  $E(t)$  defined by

$$E(t) := \frac{1}{2} \int_{\mathbb{R}^3} |\partial_t u(t, x)|^2 dx + \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u(t, x)|^2 dx + \frac{1}{6} \int_{\mathbb{R}^3} |u|^6(t, x) dx. \quad (1-8)$$

<sup>1</sup>The  $L_t^{2(p-1)} L_x^{2(p-1)}(\mathbb{R} \times \mathbb{R}^3)$ -norm of  $u$  is invariant under the scaling (1-4). The choice of the space  $L_t^{2(p-1)} L_x^{2(p-1)}$  in which we place the solution  $u$  is not unique. There exists an infinite number of spaces of the form  $L_t^q L_x^r$  scale invariant in which we can establish a local well-posedness theory.

In other words,  $E(t) = E(0)$ . This is why this equation is often called energy-critical: the exponent  $s_p = 1$  corresponds precisely to the minimal regularity required for (1-8) to be defined. The global well-posedness of (1-8) in the energy class and in higher regularity spaces is now understood. Rauch [1981] proved the global existence of smooth solutions of this equation with small data. Struwe [1988] showed that the result still holds for large data but with the additional assumption of spherical symmetry of the data. The general case (large data, no symmetry assumption) was finally settled by Grillakis [1990; 1992]. Shatah and Struwe [1994] and independently Kapitanski [1994] proved global existence of solutions in the energy class. Bahouri and Gérard [1999] reproved this result by using a compactness method and results from Bahouri and Shatah [1998]. In particular, they showed that the  $L_t^{2(5-1)} L_x^{2(5-1)}(\mathbb{R} \times \mathbb{R}^3)$ -norm of the solution is bounded by an unspecified finite quantity. Lately Tao [2006b] found an exponential tower type bound of this norm. All these proofs of global existence of solutions of the energy-critical wave equation have as a common key point the conservation of energy, which leads, in particular, to the control of the  $\dot{H}^1 \times L^2$ -norm of the solution  $(\partial_t u(t), u(t))$ .

If  $s_p < 1$ , or equivalently,  $p < 5$ , we are in the energy-subcritical equation. The scattering conjecture is an open problem. Nevertheless, some partial results are known if we consider the same problem (1-3), but with data  $(u_0, u_1) \in H^s \times H^{s-1}$ ,  $s_p < s$ . More precisely, it is proved in [Kenig et al. 2000; Gallagher and Planchon 2003; Bahouri and Chemin 2006; Roy 2007; Roy 2009a] that there exists  $s_0 := s_0(p)$  such that  $s_p < s_0 < 1$  and such that (1-3) is globally well-posed in  $H^s \times H^{s-1}$ , for  $s > s_0$ .

If  $s_p > 1$ , or, equivalently,  $p > 5$ , we are in the energy-supercritical regime. The global behavior of the solution is, in this regime, very poorly understood. Indeed, following the theory of the energy-critical wave equation, the first step would be to prove that the  $\dot{H}^{s_p} \times \dot{H}^{s_p-1}$ -norm of the solution is bounded for all time by a finite quantity depending only on the  $\dot{H}^{s_p} \times \dot{H}^{s_p-1}$ -norm of the initial data. Unfortunately, the control of this norm is a very challenging problem, since there are no known conservation laws in high regularity Sobolev spaces. Kenig and Merle [2011] recently proved, at least for radial data, that this step would be the last, by using their concentration compactness/rigidity theorem method [Kenig and Merle 2006]. More precisely, they showed that if  $\sup_{t \in I_{\max}} \|(u(t), \partial_t u(t))\|_{\dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)} < \infty$ , then Conjecture 1.1 is true.

As mentioned before, the energy supercritical regime is almost *terra incognita*. Nevertheless, Tao [2007] observed that the technology used to prove global well-posedness of smooth solutions of (1-3) can be extended, after some effort, to some equations of the type (1-1), with  $p = 5$  and radial data. More precisely, he proved global regularity of (1-1) with  $g(x) := \log(2 + x^2)$ . This phenomenon, in fact, does not depend on the symmetry of the data: it was proved in [Roy 2009b] that there exists a unique global smooth solution of (1-1) with  $g(x) := \log^c \log(10 + x^2)$  and  $0 < c < \frac{8}{225}$ .

Equations of the type (1-1) are called *barely  $\dot{H}^{s_p}$ -supercritical* wave equations. Indeed, the condition (1-2) basically says that for every  $\epsilon > 0$ , there exist two constants  $c_1 := c_1(p)$  and  $c_2 := c_2(p, \epsilon)$  such that

$$c_1(p) \leq g(|u|) \leq c_2(p, \epsilon)|u|^\epsilon \quad \text{for } |u| \text{ large.} \quad (1-9)$$

Since the critical exponent of the equation  $\partial_{tt} u - \Delta u = -|u|^{p-1+\epsilon} u$  is  $s_{p+\epsilon} = s_p + O(\epsilon)$ , the nonlinearity of (1-1) is barely  $\dot{H}^{s_p}$ -supercritical.

The goal of this paper is to check that this phenomenon, observed for  $s_p = 1$ , still holds for other values of  $s_p$ . The standard local well-posedness theory shows us that it is enough to control the pointwise-in-time  $\tilde{H}^2 \times \tilde{H}^1$ -norm of the solution. In this paper, we will use an alternative local well-posedness theory. We shall prove:

**Proposition 1.2** (local existence for barely  $\dot{H}^{s_p}$ -supercritical wave equation). *Assume that  $g$  satisfies (1-2) and*

$$g''(x) = O\left(\frac{1}{x^2}\right). \quad (1-10)$$

*Let  $M$  be such that  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq M$ . Then there exists  $\delta := \delta(M) > 0$  small such that, if  $T_l$  satisfies*

$$\left\| \cos(tD)u_0 + \frac{\sin tD}{D}u_1 \right\|_{L_t^{2(p-1)}L_x^{2(p-1)}([0, T_l] \times \mathbb{R}^3)} \leq \delta, \quad (1-11)$$

*then there exists a unique  $(u, \partial_t u)$  in*

$$\mathcal{C}([0, T_l], \tilde{H}^2) \cap L_t^{2(p-1)}L_x^{2(p-1)}([0, T_l]) \cap D^{\frac{1}{2}-s_p}L_t^4L_x^4([0, T_l]) \cap D^{\frac{1}{2}-2}L_t^4L_x^4([0, T_l]) \times \mathcal{C}([0, T_l], \tilde{H}^1)$$

*that solves (1-1) in the integral equation sense; i.e.,  $u$  satisfies the Duhamel formula*

$$u(t) := \cos(tD)u_0 + \frac{\sin tD}{D}u_1 - \int_0^t \frac{\sin(t-t')D}{D}(|u(t')|^{p-1}u(t')g(|u(t')|)) dt'. \quad (1-12)$$

Notice the many similarities between Proposition 1.2 and the local well-posedness theory for (1-3).

This allows us to define a maximum time interval of existence  $I_{\max, g} = [-T_{-, g}, T_{+, g}]$  such that, for any compact subinterval  $J \subset I_{\max, g}$ , the quantities

$$\|u\|_{L_t^{2(p-1)}L_x^{2(p-1)}(J)}, \quad \|D^{s_p-\frac{1}{2}}u\|_{L_t^4L_x^4(J)}, \quad \|D^{2-\frac{1}{2}}u\|_{L_t^4L_x^4(J)}, \quad \|(u, \partial_t u)\|_{L_t^\infty\tilde{H}^2(J) \times L_t^\infty\tilde{H}^1(J)}$$

are all finite. Again, see [Kenig and Merle 2006] or [Tao 2006a] for more explanations.

Now we set up the problem. In view of the comments above for  $s_p = 1$ , we need to make two assumptions. First we will work with a “good”  $\dot{H}^{s_p}(\mathbb{R}^3)$  theory: therefore we will assume that Conjecture 1.1 is true. Then, we also would like to work with  $\dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)$  bounded solutions  $(u(t), \partial_t u(t))$ ; more precisely, we will assume this:

**Condition 1.3** (of Kenig–Merle type). *Let  $g$  be a function that satisfies (1-2) and that is constant for  $x$  large. Then there exists  $C_2 := C_2(\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}, g)$  such that*

$$\sup_{t \in I_{\max, g}} \|(u(t), \partial_t u(t))\|_{\dot{H}^{s_p}(\mathbb{R}^3) \times \dot{H}^{s_p-1}(\mathbb{R}^3)} \leq C_2. \quad (1-13)$$

**Remark 1.4.** In the particular case  $s_p = 1$ , it is not difficult to see that Condition 1.3 is satisfied. Indeed,  $u$  satisfies the energy conservation law

$$E_b(t) := \frac{1}{2} \int_{\mathbb{R}^3} (\partial_t u(t, x))^2 dx + \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u(t, x)|^2 dx + \int_{\mathbb{R}^3} F(u(t, x), \bar{u}(t, x)) dx, \quad (1-14)$$

with

$$F(z, \bar{z}) = |z|^{5+1} \int_0^1 t^5 \operatorname{Re}(g(t|z|)) dt = |z|^{5+1} \int_0^1 t^5 g(t|z|) dt. \quad (1-15)$$

Since  $g$  is bounded, we have  $|F(z, \bar{z})| \lesssim |z|^6$ . By using the Sobolev embeddings  $\|u_0\|_{L_x^6} \lesssim \|u_0\|_{\dot{H}^2}$  and  $\|u(t)\|_{L_x^6} \lesssim \|u(t)\|_{\dot{H}^2}$ , we easily conclude that [Condition 1.3](#) holds. The energy conservation law was often in [[Tao 2007](#); [Roy 2009b](#)].

Here is the main result of this paper:

**Theorem 1.5.** *Let  $p$  be fixed.*

(1) *There exists a function  $\tilde{g}$  satisfying (1-2) and*

$$\lim_{x \rightarrow \infty} \tilde{g}(x) = \infty \quad (1-16)$$

*and such that the solution of (1-1) (with  $g := \tilde{g}$ ) exists for all time, provided that the scattering conjecture and [Condition 1.3](#) are satisfied.*

(2) *There exists a function  $f$  depending on  $T$  and  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}$  such that*

$$\|u\|_{L_t^\infty \tilde{H}^2([-T, T])} + \|\partial_t u\|_{L_t^\infty \tilde{H}^1([-T, T])} \leq f(T, \|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}). \quad (1-17)$$

[Theorem 1.5](#) shows that a “good”  $\dot{H}^{sp}(\mathbb{R}^3)$  theory for (1-3) can be extended, at least, to one barely  $\dot{H}^{sp}(\mathbb{R}^3)$ -supercritical equation, with  $\tilde{g}$  going to infinity.

**Remark 1.6.** Apart from its dependence on  $p$ , the function  $\tilde{g}$  is universal: it does not depend on an upper bound of the initial data. Moreover,  $\tilde{g}$  is unbounded: it goes to infinity with as  $x$ .

**Remark 1.7.** In fact, [Theorem 1.5](#) holds for a weaker version of [Condition 1.3](#): there exists a function  $C_2$  such that for all subinterval  $I \subset I_{\max, g}$

$$\sup_{t \in I} \|(u(t), \partial_t u(t))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} \leq C_2, \quad (1-18)$$

with  $C_2 := C_2(\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}, g, |I|)$ . See the proof of [Theorem 1.5](#) and, in particular, (5-21), (5-33) and (5-48).

We recall some basic properties and estimates. If  $t_0 \in [t_1, t_2]$ , if  $F \in L_t^{\tilde{q}} L_x^{\tilde{r}}([t_1, t_2])$  and if  $(u, \partial_t u) \in C([t_1, t_2], \dot{H}^m(\mathbb{R}^3)) \times C([t_1, t_2], \dot{H}^{m-1}(\mathbb{R}^3))$  satisfy

$$u(t) : \cos(tD)u_0 + \frac{\sin tD}{D}u_1 - \int_{t_0}^t \frac{\sin(t-t')D}{D}F(t')dt', \quad (1-19)$$

with data  $(u(t_0), \partial_t u(t_0)) \in \dot{H}^m(\mathbb{R}^3) \times \dot{H}^{m-1}(\mathbb{R}^3)$ , then we have the Strichartz estimates [[Ginibre and Velo 1995](#); [Lindblad and Sogge 1995](#)]

$$\begin{aligned} \|u\|_{L_t^q L_x^r([t_1, t_2])} + \|u\|_{L_t^\infty \dot{H}^m(\mathbb{R}^3)([t_1, t_2])} + \|\partial_t u\|_{L_t^\infty \dot{H}^{m-1}(\mathbb{R}^3)([t_1, t_2])} \\ \lesssim \|(u(t_0), \partial_t u(t_0))\|_{\dot{H}^m(\mathbb{R}^3) \times \dot{H}^{m-1}(\mathbb{R}^3)} + \|F\|_{L_t^{\tilde{q}} L_x^{\tilde{r}}([t_1, t_2])}. \end{aligned} \quad (1-20)$$

Here  $(q, r)$  is  $m$ -wave admissible, i.e.,

$$(q, r) \in (2, \infty] \times [2, \infty] \quad \text{and} \quad \frac{1}{q} + \frac{3}{r} = \frac{3}{2} - m; \quad (1-21)$$

moreover,

$$\frac{1}{q} + \frac{3}{r} = \frac{1}{\tilde{q}} + \frac{3}{\tilde{r}} - 2. \quad (1-22)$$

We set some notation that will appear throughout the paper.

We write  $A \lesssim B$  if there exists a universal nonnegative constant  $C' > 0$  such that  $A \leq C'B$ . The notation  $A = O(B)$  means  $A \lesssim B$ . More generally, we write  $A \lesssim_{a_1, \dots, a_n} B$  if there exists a nonnegative constant  $C' = C(a_1, \dots, a_n)$  such that  $A \leq C'B$ . We say that  $C''$  is the constant determined by  $\lesssim$  in  $A \lesssim_{a_1, \dots, a_n} B$  if  $C''$  is the smallest possible  $C'$  such that  $A \leq C'B$ . We write  $A \ll_{a_1, \dots, a_n} B$  if there exists a universal small nonnegative constant  $c = c(a_1, \dots, a_n)$  such that  $A \leq cB$ . Following [Kenig and Merle 2011], we define, on an interval  $I$ ,

$$\|u\|_{S(I)} := \|u\|_{L_t^{2(p-1)} L_x^{2(p-1)}(I)}, \quad \|u\|_{W(I)} := \|u\|_{L_t^4 L_x^4(I)}, \quad \|u\|_{\bar{W}(I)} := \|u\|_{L_t^{\frac{4}{3}} L_x^{\frac{4}{3}}(I)}. \quad (1-23)$$

We also define the quantity

$$Q(I, u) := \|D^{s_p - \frac{1}{2}} u\|_{W(I)} + \|D^{2 - \frac{1}{2}} u\|_{W(I)} + \|u\|_{L_t^\infty \tilde{H}^2(I)} + \|\partial_t u\|_{L_t^\infty \tilde{H}^1(I)} \quad (1-24)$$

Let  $X$  be a Banach space and  $r \geq 0$ . Then

$$B(X, r) := \{f \in X : \|f\|_X \leq r\} \quad (1-25)$$

We recall also the well-known Sobolev embeddings. We have

$$\|h\|_{L^\infty(\mathbb{R}^3)} \lesssim \|h\|_{\tilde{H}^2}, \quad (1-26)$$

$$\|h\|_{S(I)} \lesssim \|D^{s_p - \frac{1}{2}} h\|_{L_t^{2(p-1)} L_x^{\frac{6(p-1)}{2p-3}}(I)}. \quad (1-27)$$

We shall combine (1-27) with the Strichartz estimates, since  $(2(p-1), \frac{6(p-1)}{2p-3})$  is  $\frac{1}{2}$ -wave admissible.

We also recall some Leibnitz rules [Christ and Weinstein 1991; Kenig et al. 1993]. We have

$$\|D^\alpha F(u)\|_{L_t^q L_x^r(I)} \lesssim \|F'(u)\|_{L_t^{q_1} L_x^{r_1}(I)} \|D^\alpha u\|_{L_t^{q_2} L_x^{r_2}(I)}, \quad (1-28)$$

with  $\alpha > 0$ ,  $r, r_1, r_2$  lying in  $[1, \infty]$ ,  $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$ , and  $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$ .

The Leibnitz rule for products is

$$\|D^\alpha(uv)\|_{L_t^q L_x^r(I)} \lesssim \|D^\alpha u\|_{L_t^{q_1} L_x^{r_1}(I)} \|v\|_{L_t^{q_2} L_x^{r_2}(I)} + \|D^\alpha u\|_{L_t^{q_3} L_x^{r_3}(I)} \|v\|_{L_t^{q_4} L_x^{r_4}(I)}, \quad (1-29)$$

with  $\alpha > 0$ ,  $r, r_1, r_2$  lying in  $[1, \infty]$ ,  $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$ ,  $\frac{1}{q} = \frac{1}{q_3} + \frac{1}{q_4}$ ,  $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$ , and  $\frac{1}{r} = \frac{1}{r_3} + \frac{1}{r_4}$ .

If  $F \in C^2$ , we can write

$$F(x) - F(y) = \int_0^1 F'(tx + (1-t)y)(x-y) dt. \quad (1-30)$$

By using (1-28) and (1-29) the Leibnitz rule for differences can be formulated as

$$\begin{aligned} \|D^\alpha(F(u) - F(v))\|_{L_t^q L_x^{r_1}(I)} &\lesssim \sup_{t \in [0,1]} \|F'(tu + (1-t)v)\|_{L_t^{q_1} L_x^{r_1}(I)} \|D^\alpha(u - v)\|_{L_t^{q_2} L_x^{r_2}(I)} \\ &+ \sup_{t \in [0,1]} \|F''(tu + (1-t)v)\|_{L_t^{q'_1} L_x^{r'_1}(I)} \left( \|D^\alpha u\|_{L_t^{q'_2} L_x^{r'_2}(I)} + \|D^\alpha v\|_{L_t^{q'_2} L_x^{r'_2}(I)} \right) \|u - v\|_{L_t^{q'_3} L_x^{r'_3}(I)}, \end{aligned} \quad (1-31)$$

with  $\alpha > 0$ ,  $r_1, r_2, r'_1, r'_2, r'_3$  lying in  $[1, \infty]$ ,  $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$ ,  $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$ ,  $\frac{1}{q} = \frac{1}{q'_1} + \frac{1}{q'_2} + \frac{1}{q'_3}$ , and  $\frac{1}{r} = \frac{1}{r'_1} + \frac{1}{r'_2} + \frac{1}{r'_3}$ .

We shall apply these formulas to several formulas of  $F(u)$ , and, in particular, to  $F(u) := |u|^{p-1}ug(|u|)$ . Notice that, by (1-2) and (1-10), we have  $F'(x) \sim |x|^{p-1}g(|x|)$  and  $F''(x) \sim |x|^{p-2}g(|x|)$ . Notice also that, by (1-2) again, we have, for  $t \in [0, 1]$ ,

$$g(|tx + (1-t)y|) \leq g(2 \max(|x|, |y|)) \leq g(\max(|x|, |y|) + \log 2) \lesssim g(|x|) + g(|y|). \quad (1-32)$$

This will allow us to estimate easily

$$\sup_{t \in [0,1]} \|F'(tu + (1-t)v)\|_{L_t^{q_1} L_x^{r_1}(I)} \quad \text{and} \quad \sup_{t \in [0,1]} \|F''(tu + (1-t)v)\|_{L_t^{q'_1} L_x^{r'_1}(I)}.$$

Now we explain the main ideas of this paper. We shall prove, in Section 3, that very many values functions  $g$ , a special property for the solution of (1-1) holds.

**Proposition 1.8** (control of  $S(I)$ -norm and of norm of initial data imply control of  $L_t^\infty \tilde{H}^2(I) \times L_t^\infty \tilde{H}^1(I)$  norm). *Let  $I$  be a compact subinterval of  $I_{\max, g}$  (so  $\|u\|_{S(I)} < \infty$ ) and assume that  $0 \in I$ . Assume that  $g$  satisfies (1-2), (1-10) and<sup>2</sup>*

$$\int_1^\infty \frac{1}{yg^2(y)} dy = \infty. \quad (1-33)$$

*Let  $A \geq 0$  such that  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq A$ . Let  $u$  be the solution of (1-1). There exists a constant  $C > 0$  such that*

$$\|(u, \partial_t u)\|_{L_t^\infty \tilde{H}^2(I) \times L_t^\infty \tilde{H}^1(I)} \leq (2C)^N A, \quad (1-34)$$

*with  $N := N(I)$ , such that*

$$\int_{2CA}^{(2C)^N A} \frac{1}{yg^2(y)} dy \gg \|u\|_{S(I)}^{2(p-1)}. \quad (1-35)$$

Moreover we shall give a criterion of global well-posedness (proved in Section 4):

**Proposition 1.9** (criterion of global well-posedness). *Assume that  $|I_{\max, g}| < \infty$ . Assume that  $g$  satisfies (1-2), (1-10) and (1-33). Then*

$$\|u\|_{S(I_{\max, g})} = \infty. \quad (1-36)$$

The first step is to prove global well-posedness of (1-1), with  $g := g_1$  a nondecreasing function that is constant for  $x$  large (say  $x \geq C'_1$ , with  $C'_1$  to be determined). By Proposition 1.9, it is enough to find an upper bound of the  $S([-T, T])$ -norm of the solution  $u_{[1]}$  for  $T$  arbitrarily large. This can indeed be done, by proving that  $g_1$  can be considered as a subcritical perturbation of the nonlinearity. In other words,  $g_1(|u|)|u|^{p-1}u$  will play the same role as that of  $|u|^{p-1}u(1 - |u|^{-\alpha})$  for some  $\alpha > 0$ . Once we have noticed

<sup>2</sup>Condition (1-33) basically says that  $g$  grows slowly on average.



that this comparison is possible, we shall estimate the relevant norms (in particular,  $\|u_{[1]}\|_{S([-T, T])}$ ) using perturbation theory, [Conjecture 1.1](#) and [Condition 1.3](#), in the spirit of [\[Zhang 2006\]](#). We expect to find a bound of the form

$$\|u_{[1]}\|_{S([-T, T])} \leq C_3 \left( \|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}, T \right), \quad (1-37)$$

with  $C_3$  increasing as  $T$  or  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}$  grows. Notice that if we restrict  $[-T, T]$  to the interval  $[-1, 1]$  and if the  $\tilde{H}^2 \times \tilde{H}^1$ -norm of the initial data  $(u_0, u_1)$  is bounded by 1, then we can prove, using (1-37), (1-26) and [Proposition 1.8](#), that the  $L_t^\infty L_x^\infty([-T, T])$ -norm of the solution  $u_{[1]}$  is bounded by a constant (denoted by  $C_1$ ) on  $[-1, 1]$ . Therefore, if  $h$  is a smooth extension of  $g_1$  outside  $[0, C_1]$ , and if  $u$  is the solution of (1-1) (with  $g := h$ ), we expect to prove that  $u = u_{[1]}$  on  $[-1, 1]$  and for data  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq 1$ . This implies in particular, by (1-37), that we have a finite upper bound  $\|u\|_{S([-1, 1])}$ .

We are not done yet. There are two problems. First,  $g_1$  does not go to infinity. Second, we only control  $\|u\|_{S([-1, 1])}$  for data  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq 1$ : we would like to control  $\|u\|_{S(\mathbb{R})}$  for arbitrary data. In order to overcome these difficulties we iterate the procedure described above. More precisely, given a function  $g_{i-1}$  that is constant for  $x \geq C_{i-1}$  and such that  $u_{[i-1]}$ , a solution of (1-1) with  $g = g_{i-1}$ , satisfies  $\|u_{[i-1]}\|_{S([-i-1, i-1])} < \infty$ , we construct a function  $g_i$  that

- is an extension of  $g_{i-1}$  outside  $[0, C_{i-1}]$ , and
- is increasing and constant (say equal to  $i + 1$ ) for  $x \geq C'_i$ , with  $C'_i$  to be determined.

Again, we shall prove that the  $g_i$  may be regarded as a subcritical perturbation of the nonlinearity  $(i+1)|u|^{p-1}u$ . This allow us to control  $\|u_{[i]}\|_{S([-i, i])}$ , by using perturbation theory, [Conjecture 1.1](#), and [Condition 1.3](#). Using [Proposition 1.8](#) and (1-26), we can find a finite upper bound for  $\|u_{[i]}\|_{L_t^\infty L_x^\infty([-i, i])}$ . We assign the value of this upper bound to  $C_i$ . To conclude the argument we let  $\tilde{g} = \lim_{i \rightarrow \infty} g_i$ . Given  $T > 0$ , we can find a  $j$  such that  $[-T, T] \subset [-j, j]$  and  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq j$ . We prove that  $u = u_{[j]}$  on  $[-j, j]$ , where  $u$  is a solution of (1-1) with  $g := \tilde{g}$ . Since we have a finite upper bound of  $\|u_{[j]}\|_{S([-j, j])}$ , we also control  $\|u\|_{S([-j, j])}$  and  $\|u\|_{S([-T, T])}$ . [Theorem 1.5](#) follows from [Proposition 1.9](#).

## 2. Proof of [Proposition 1.2](#)

In this section we prove [Proposition 1.2](#) for barely  $\dot{H}^{s_p}(\mathbb{R}^3)$ -supercritical wave equations (1-1). The proof is based upon standard arguments. Here we have chosen to modify an argument in [\[Kenig and Merle 2011\]](#).

For  $\delta, T_l, C, M$  to be chosen and such that (1-11) holds we define

$$\begin{aligned} B_1 &:= \mathbf{B}(\mathcal{C}([0, T_l], \tilde{H}^2) \cap D^{\frac{1}{2}-s_p} W([0, T_l]) \cap D^{\frac{1}{2}-2} W([0, T_l]), 2CM), \\ B_2 &:= \mathbf{B}(S([0, T_l]), 2\delta), \\ B' &:= \mathbf{B}(\mathcal{C}([0, T_l], \tilde{H}^1), 2CM), \end{aligned} \quad (2-1)$$

and

$$X := \left\{ (u, \partial_t u) : u \in B_1 \cap B_2, \partial_t u \in B' \right\}. \quad (2-2)$$

Let

$$\Psi(u, \partial_t u) := \begin{pmatrix} \cos(tD)u_0 + \frac{\sin(tD)}{D}u_1 - \int_0^t \frac{\sin(t-t')D}{D} (|u(t')|^{p-1}u(t')g(|u(t')|)) dt' \\ -D \sin(tD)u_0 + \cos(tD)u_1 - \int_0^t \cos(t-t')D (|u(t')|^{p-1}u(t')g(|u(t')|)) dt' \end{pmatrix}. \quad (2-3)$$

$\Psi$  maps  $X$  to  $X$ . Indeed, in view of (1-11), (1-20), and the fractional Leibnitz rule (1-28) applied to  $\alpha \in \{s_p - \frac{1}{2}, 2 - \frac{1}{2}\}$  and

$$F(u) := |u|^{p-1}ug(|u|)$$

and by applying the multipliers  $D^{2-\frac{1}{2}}$  and  $D^{s_p-\frac{1}{2}}$  to the Strichartz estimates with  $m = \frac{1}{2}$ , we have

$$\begin{aligned} & Q([0, T_l]) \\ & \lesssim \|(u_0, u_1)\|_{\tilde{H}^2(\mathbb{R}^3) \times \tilde{H}^1(\mathbb{R}^3)} + \|D^{s_p-\frac{1}{2}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}([0, T_l])} + \|D^{2-\frac{1}{2}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}([0, T_l])} \\ & \leq CM + C(\|D^{s_p-\frac{1}{2}}u\|_{W([0, T_l])} + \|D^{2-\frac{1}{2}}u\|_{W([0, T_l])})\|u\|_{S([0, T_l])}^{p-1}g(\|u\|_{L_t^\infty L_x^\infty([0, T_l])}) \\ & \leq CM + (2\delta)^{p-1}C(2CM)g(2CM) \end{aligned} \quad (2-4)$$

for some  $C > 0$  and

$$\begin{aligned} \|u\|_{S([0, T_l])} - \delta & \lesssim \|D^{s_p-\frac{1}{2}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}([0, T_l])} \\ & \lesssim \|u\|_{S([0, T_l])}^{p-1} \|D^{s_p-\frac{1}{2}}u\|_{W([0, T_l])} g(\|u\|_{L_t^\infty L_x^\infty([0, T_l])}) \lesssim (2\delta)^{p-1}(2CM)g(2CM). \end{aligned} \quad (2-5)$$

Choosing  $\delta = \delta(M) > 0$  small enough we see that  $\Psi(X) \subset X$ .

$\Psi$  is a contraction. Indeed we have

$$\begin{aligned} & \|\Psi(u) - \Psi(v)\|_X \\ & \lesssim \|D^{s_p-\frac{1}{2}}(|u|^{p-1}ug(|u|) - |v|^{p-1}vg(|v|))\|_{\tilde{W}([0, T_l])} + \|D^{2-\frac{1}{2}}(|u|^{p-1}ug(|u|) - |v|^{p-1}vg(|v|))\|_{\tilde{W}([0, T_l])} \\ & \lesssim (g(\|u\|_{L_t^\infty L_x^\infty([0, T_l])}) + g(\|v\|_{L_t^\infty L_x^\infty([0, T_l])})) \\ & \quad \times \left( (\|u\|_{S([0, T_l])}^{p-1} + \|v\|_{S([0, T_l])}^{p-1}) (\|D^{s_p-\frac{1}{2}}(u-v)\|_{W([0, T_l])} + \|D^{2-\frac{1}{2}}(u-v)\|_{W([0, T_l])}) \right. \\ & \quad \left. + (\|u\|_{S([0, T_l])}^{p-2} + \|v\|_{S([0, T_l])}^{p-2}) \|u-v\|_{S([0, T_l])} \right. \\ & \quad \left. \times (\|D^{s_p-\frac{1}{2}}u\|_{W([0, T_l])} + \|D^{2-\frac{1}{2}}u\|_{W([0, T_l])} + \|D^{s_p-\frac{1}{2}}v\|_{W([0, T_l])} + \|D^{2-\frac{1}{2}}v\|_{W([0, T_l])}) \right) \\ & \lesssim (g(2CM)(2\delta)^{p-1} + (2\delta)^{p-2}(2CM))\|u-v\|_X. \end{aligned} \quad (2-6)$$

In these computations, we applied the Leibnitz rule for differences to  $\alpha \in \{s_p - \frac{1}{2}, 2 - \frac{1}{2}\}$  and

$$F(u) := |u|^{p-1}ug(|u|).$$

Therefore, if  $\delta = \delta(M) > 0$  is small enough,  $\Psi$  is a contraction.

### 3. Proof of Proposition 1.8

To show Proposition 1.8, it is enough to prove that  $Q(I) < \infty$ . Without loss of generality we can assume that  $A \gg 1$ . Then we divide  $I$  into subintervals  $(I_i)_{1 \leq i \leq N}$  such that

$$\|u\|_{S(I_i)} = \frac{\eta}{g^{1/(p-1)}((2C)^i A)} \quad (3-1)$$

for some  $C \gtrsim 1$  and  $\eta > 0$  constants to be chosen later, except maybe the last one. Notice that such a partition always exists since by (1-33) we get, for  $N := N(I)$  large enough,

$$\sum_{i=1}^N \frac{1}{g^2((2C)^i A)} \geq \int_1^N \frac{1}{g^2((2C)^x A)} dx \gtrsim \int_{2CA}^{(2C)^N A} \frac{1}{yg^2(y)} dy \gg \|u\|_{S(I)}^{2(p-1)}. \quad (3-2)$$

We get, by a similar reasoning as used in Section 2

$$\begin{aligned} Q(I_1, u) &\lesssim \|(u_0, u_1)\|_{\tilde{H}^2(\mathbb{R}^3) \times \tilde{H}^1(\mathbb{R}^3)} + \|D^{s_{p-\frac{1}{2}}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}(I_1)} + \|D^{2-\frac{1}{2}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}(I_1)} \\ &\lesssim A + (\|D^{s_{p-\frac{1}{2}}}u\|_{W(I_1)} + \|D^{2-\frac{1}{2}}u\|_{W(I_1)})\|u\|_{S(I_1)}^{p-1}g(\|u\|_{L_t^\infty L_x^\infty(I_1)}) \\ &\lesssim A + \|u\|_{S(I_1)}^{p-1}Q(I_1, u)g(Q(I_1, u)). \end{aligned} \quad (3-3)$$

We choose  $C$  to be equal to the constant determined by  $\lesssim$  in (3-3). Without loss of generality we can assume that  $C > 1$ . By a continuity argument, iteration on  $i$ , we get, for  $\eta \ll 1$ , (1-34).

### 4. Proof of Proposition 1.9

To prove Proposition 1.9, we argue as follows: by time reversal symmetry it is enough to prove that  $T_{+,g} < \infty$ . If  $\|u\|_{S(I_{\max,g})} < \infty$  then we have  $Q([0, T_{+,g}], u) < \infty$ : this follows by slightly adapting the proof of Proposition 1.8. Consequently, by the dominated convergence theorem, there would exist a sequence  $t_n \rightarrow T_{+,g}$  such that  $\|u\|_{S([t_n, T_{+,g}])} \ll \delta$  and  $\|D^{s_{p-\frac{1}{2}}}u\|_{W([t_n, T_{+,g}])} \ll \delta$  if  $n$  is large enough, with  $\delta$  defined in Proposition 1.2. But, by (1-19) and (1-20),

$$\begin{aligned} &\|\cos((t - t_n)D)u(t_n) + \frac{\sin(t - t_n)D}{D}u_1\|_{S([t_n, T_{+,g}])} \\ &\lesssim \|u\|_{S([t_n, T_{+,g}])} + \|u\|_{S([t_n, T_{+,g}])}^{p-1}\|D^{s_{p-\frac{1}{2}}}u\|_{W([t_n, T_{+,g}])}g(Q([0, T_{+,g}], u)) \ll \delta, \end{aligned} \quad (4-1)$$

and consequently, by continuity, there would exist  $\tilde{T} > T_{+,g}$  such that

$$\left\| \cos((t - t_n)D)u(t_n) + \frac{\sin(t - t_n)D}{D}\partial_t u(t_n) \right\|_{S([t_n, \tilde{T}])} \leq \delta, \quad (4-2)$$

which would contradict the definition of  $T_{+,g}$ .

**Remark 4.1.** Notice that if we have the stronger bound  $\|u\|_{S(I_{\max,g})} \leq C$  with  $C := C(\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}) < \infty$ , then not only  $I_{\max,g} = (-\infty, +\infty)$  but also  $u$  scatters as  $t \rightarrow \pm\infty$ . Indeed, by Proposition 1.9,  $I_{\max,g} = \mathbb{R}$ . Then by time reversal symmetry it is enough to assume that  $t \rightarrow \infty$ . Let  $v(t) := (u(t), \partial_t u(t))$ .

We are looking for  $v_{+,0} := (u_{+,0}, u_{+,1})$  such that

$$\|v(t) - K(t)v_{+,0}\|_{\tilde{H}^2 \times \tilde{H}^1} \rightarrow 0 \quad (4-3)$$

as  $t \rightarrow \infty$ . Here

$$K(t) := \begin{pmatrix} \cos t D & (\sin t D)/D \\ -D \sin t D & \cos t D \end{pmatrix} \quad (4-4)$$

We have

$$K^{-1}(t) = \begin{pmatrix} \cos t D & -(\sin t D)/D \\ D \sin t D & \cos t D \end{pmatrix}. \quad (4-5)$$

Notice that  $K^{-1}(t)$  and  $K(t)$  are bounded in  $\tilde{H}^2 \times \tilde{H}^1$ . Therefore it is enough to prove that  $K^{-1}(t)v(t)$  has a limit as  $t \rightarrow \infty$ . But since  $K^{-1}(t)v(t) = (u_0, u_1) - K^{-1}(t)(u_{\text{nl}}(t), \partial_t u_{\text{nl}}(t))$  — where

$$u_{\text{nl}}(t) := - \int_0^t \frac{\sin(t-t')D}{D} (|u(t')|^{p-1}u(t')g(|u(t')|)) dt'$$

denotes the nonlinear part of the solution (1-12) — it suffices to prove that  $K^{-1}(t)(u_{\text{nl}}(t), \partial_t u_{\text{nl}}(t))$  has a limit. But

$$\begin{aligned} & \|K^{-1}(t_1)u_{\text{nl}}(t_1) - K^{-1}(t_2)u_{\text{nl}}(t_2)\|_{\tilde{H}^2 \times \tilde{H}^1} \\ & \lesssim \|(u_{\text{nl}}, \partial_t u_{\text{nl}})\|_{L_t^\infty \tilde{H}^2([t_1, t_2]) \times L_t^\infty \tilde{H}^1([t_1, t_2])} \\ & \lesssim (\|D^{sp-\frac{1}{2}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}([t_1, t_2])} + \|D^{2-\frac{1}{2}}(|u|^{p-1}ug(|u|))\|_{\tilde{W}([t_1, t_2])}) \\ & \lesssim (\|D^{sp-\frac{1}{2}}u\|_{W([t_1, t_2])} + \|D^{2-\frac{1}{2}}u\|_{W([t_1, t_2])})\|u\|_{S([t_1, t_2])}^{p-1}g(\|u\|_{L_t^\infty L_x^\infty(\mathbb{R})}). \end{aligned} \quad (4-6)$$

It remains to prove that  $Q(\mathbb{R}) < \infty$  in order to conclude that the Cauchy criterion is satisfied, which would imply scattering. This follows from  $\|u\|_{S(\mathbb{R})} < \infty$  and a slight modification of the proof of Proposition 1.8.

## 5. Construction of the function $g$

In this section we prove Theorem 1.5. Let

$$\text{Up}(i) := \{(T, (u_0, u_1)) : 0 \leq T \leq i, \|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq i\} \quad (5-1)$$

As  $i$  ranges over  $\{1, 2, \dots\}$  we construct, for each set  $\text{Up}(i)$ , a function  $g_i$  satisfying (1-2) and (1-10). Moreover it is constant for large values of  $|x|$ . The function  $g_{i+1}$  depends on  $g_i$ ; the construction of  $g_i$  is made by induction on  $i$ . More precisely:

**Lemma 5.1.** *Let  $A \gg 1$ . There exist two sequences of numbers  $\{C_i\}_{i \geq 0}$ ,  $\{C'_i\}_{i \geq 0}$  and a sequence of functions  $\{g_i\}_{i \geq 0}$  such that, for all  $(T, (u_0, u_1)) \in \text{Up}(i)$ , we have*

- $g_0 := 1, C_0 := 0, C'_0 = 0$ ;
- $\{C_i\}_{i \geq 0}$  and  $\{C'_i\}_{i \geq 0}$  are positive, nondecreasing, and satisfy

$$AC_{i-1} < C'_i < AC_i \quad (5-2)$$

for  $i \geq 1$  and

$$C_i \geq i; \quad (5-3)$$

- $g_i$  is smooth, nondecreasing, and satisfies (1-2), (1-10),

$$\int_1^{C'_i} \frac{1}{tg_i^2(t)} dt \rightarrow \infty \quad \text{as } i \rightarrow \infty, \quad (5-4)$$

and

$$g_i(|x|) = \begin{cases} g_{i-1}(|x|) & \text{if } |x| \leq AC_{i-1}, \\ i+1 & \text{if } |x| \geq C'_i; \end{cases} \quad (5-5)$$

- the solution  $u_{[i]}$  of the wave equation

$$\begin{cases} \partial_{tt}u_{[i]} - \Delta u_{[i]} = -|u_{[i]}|^{p-1}u_{[i]}g_i(|u_{[i]}|), \\ u_{[i]}(0) = u_0 \in \tilde{H}^2, \\ \partial_t u_{[i]}(0) = u_1 \in \tilde{H}^1 \end{cases} \quad (5-6)$$

satisfies

$$\max \left( \|u_{[i]}\|_{S([-i,i])}, \|(u_{[i]}, \partial_t u_{[i]})\|_{L_t^\infty \tilde{H}^2([-T,T]) \times L_t^\infty \tilde{H}^1([-T,T])} \right) \leq C_i. \quad (5-7)$$

We postpone the proof until page 212. Assume the lemma is true and let  $\tilde{g} = \lim_{i \rightarrow \infty} g_i$ . Clearly  $\tilde{g}$  is smooth; it satisfies (1-2) and (1-10). It also goes to infinity. Moreover let  $u$  be the solution of (1-1) with  $g := \tilde{g}$ . We want to prove that the solution  $u$  exists for all time. Let  $T_0 \geq 0$  be a fixed time. Let  $j := j(T_0, \|u_0\|_{\tilde{H}^2}, \|u_1\|_{\tilde{H}^1}) > 0$  be the smallest positive integer such that  $[-T_0, T_0] \subset [-j, j]$  and  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq j$ . We claim that

$$\|(u, \partial_t u)\|_{L_t^\infty \tilde{H}^2([-T_0, T_0]) \times L_t^\infty \tilde{H}^1([-T_0, T_0])} \leq C_j \quad \text{and} \quad \|u\|_{S([-T_0, T_0])} \leq C_j.$$

Indeed, let

$$F_j := \{t \in [0, j] : \|(u, \partial_t u)\|_{L_t^\infty \tilde{H}^2([-t, t]) \times L_t^\infty \tilde{H}^1([-t, t])} \leq C_j \text{ and } \|u\|_{S([-t, t])} \leq C_j\}. \quad (5-8)$$

We must show that  $F_j$  coincides with  $[0, j]$ . Certainly  $F_j$  is nonempty, since it contains 0; see (5-3).

$F_j$  is closed. Indeed, let  $\tilde{t} \in \bar{F}_j$ . There exists a sequence  $(t_n)_{n \geq 1}$  in  $[0, j]$  such that  $t_n \rightarrow \tilde{t}$ ,  $\|u\|_{S([-t_n, t_n])} \leq C_j$ , and  $\|(u, \partial_t u)\|_{L_t^\infty \tilde{H}^2([-t_n, t_n]) \times L_t^\infty \tilde{H}^1([-t_n, t_n])} \leq C_j$ . It is enough to prove that  $\|u\|_{S([- \tilde{t}, \tilde{t}])}$  is finite and then apply dominated convergence. There are two cases:

- If  $\text{card}\{t_n : t_n \leq \tilde{t}\} < \infty$ , there exists  $n_0$  large enough such that  $t_n \geq \tilde{t}$  for  $n \geq n_0$  and

$$\|u\|_{S([- \tilde{t}, \tilde{t}])} \leq \|u\|_{S([-t_n, t_n])} < \infty. \quad (5-9)$$

- If  $\text{card}\{t_n : t_n \leq \tilde{t}\} = \infty$ , we can assume by passing to a subsequence that  $t_n \leq \tilde{t}$ . Let  $n_0 \geq 1$  be fixed. Since

$$\left\| \cos(t - t_{n_0}) Du(t_{n_0}) + \frac{\sin(t - t_{n_0})D}{D} \partial_t u(t_{n_0}) \right\|_{S([t_{n_0}, \tilde{t}])} \lesssim \|(u(t_{n_0}), \partial_t u(t_{n_0}))\|_{\tilde{H}^2 \times \tilde{H}^1} \lesssim C_j, \quad (5-10)$$

we conclude from the dominated convergence theorem that there is  $n_1 := n_1(n_0)$  large enough that

$$\left\| \cos(t - t_{n_0}) Du(t_{n_0}) + \frac{\sin(t - t_{n_0})D}{D} \partial_t u(t_{n_0}) \right\|_{S([t_{n_1}, \tilde{t}])} \leq \delta, \quad (5-11)$$



with  $\delta := \delta(C_j)$  defined in [Proposition 1.2](#). Therefore, by [Proposition 1.2](#), we have  $\|u\|_{S([t_{n_1}, \tilde{t}])} < \infty$ . Similarly,  $\|u\|_{S([- \tilde{t}, -t_{n_1}])} < \infty$ . Combining these inequalities with  $\|u\|_{S([-t_{n_1}, t_{n_1}])} \leq C_j$ , we eventually get  $\|u\|_{S([- \tilde{t}, \tilde{t}])} < \infty$ , as desired.

*$F_j$  is open.* Indeed, let  $\tilde{t} \in F_j$ . By [Proposition 1.2](#) there exists  $\alpha > 0$  such that if  $t \in (\tilde{t} - \alpha, \tilde{t} + \alpha) \cap [0, j]$  then  $[-t, t] \subset I_{\max, \tilde{g}}$  and  $\|u\|_{L_t^\infty L_x^\infty([-t, t])} \lesssim \|u\|_{L_t^\infty \tilde{H}^2([-t, t])} \lesssim C_j$ . Also, by (5-7),  $[-t, t] \subset I_{\max, g_j}$ . In view of these remarks, we conclude, after slightly adapting the proof of [Proposition 1.8](#), that  $Q([-t, t], u) \lesssim_j 1$  and  $Q([-t, t], u_{[j]}) \lesssim_j 1$ . We divide  $[-t, t]$  into a finite number of subintervals  $(I_i)_{i \leq k} = ([a_i, b_i])_{1 \leq i \leq k}$  that satisfy, for  $\eta \ll 1$  to be defined later, the following properties:

- (1)  $1 \leq i \leq k$ :  $\|u_{[j]}\|_{S(I_i)} \leq \eta$ ,  $\|u\|_{S(I_i)} \leq \eta$ ,  $\|D^{s_p - \frac{1}{2}}u_{[j]}\|_{W(I_i)} \leq \eta$ ,  $\|D^{s_p - \frac{1}{2}}u\|_{W(I_i)} \leq \eta$ ,  $\|D^{2 - \frac{1}{2}}u\|_{W(I_i)} \leq \eta$ , and  $\|D^{2 - \frac{1}{2}}u_{[j]}\|_{W(I_i)} \leq \eta$ .
- (2)  $1 \leq i < k$ :  $\|u_{[j]}\|_{S(I_i)} = \eta$  or  $\|u\|_{S(I_i)} = \eta$  or  $\|D^{s_p - \frac{1}{2}}u_{[j]}\|_{W(I_i)} = \eta$  or  $\|D^{s_p - \frac{1}{2}}u\|_{W(I_i)} = \eta$  or  $\|D^{2 - \frac{1}{2}}u\|_{W(I_i)} = \eta$ , or  $\|D^{2 - \frac{1}{2}}u_{[j]}\|_{W(I_i)} = \eta$ .

Notice that, by (1-2), we have

$$\|g_j(|u|) - g_j(|u_{[j]}|)\|_{L_t^\infty L_x^\infty(I_i)} \lesssim \|u - u_{[j]}\|_{L_t^\infty L_x^\infty(I_i)} \lesssim \|u - u_{[j]}\|_{L_t^\infty \tilde{H}^2(I_i)}. \quad (5-12)$$

Consider  $w = u - u_{[j]}$ . Applying the Leibnitz rules (1-28), (1-31), and (1-29), together with (5-12), we have

$Q(I_1, w)$

$$\begin{aligned} &\lesssim \|D^{s_p - \frac{1}{2}}(|u|^{p-1}u(\tilde{g} - g_j)(|u|))\|_{\tilde{W}(I_1)} + \|D^{2 - \frac{1}{2}}(|u|^{p-1}u(\tilde{g} - g_j)(|u|))\|_{\tilde{W}(I_1)} \\ &\quad + \|D^{s_p - \frac{1}{2}}(|u|^{p-1}u - |u_{[j]}|^{p-1}u_{[j]})g_j(|u|)\|_{\tilde{W}(I_1)} + \|D^{2 - \frac{1}{2}}(|u|^{p-1}u - |u_{[j]}|^{p-1}u_{[j]})g_j(|u|)\|_{\tilde{W}(I_1)} \\ &\quad + \|D^{s_p - \frac{1}{2}}(|u_{[j]}|^{p-1}u_{[j]}(g_j(|u|) - g_j(|u_{[j]}|)))\|_{\tilde{W}(I_1)} + \|D^{2 - \frac{1}{2}}(|u_{[j]}|^{p-1}u_{[j]}(g_j(|u|) - g_j(|u_{[j]}|)))\|_{\tilde{W}(I_1)} \\ &\lesssim (\tilde{g} - g_j)(\|u\|_{L_t^\infty \tilde{H}^2(I_1)}) (\|D^{s_p - \frac{1}{2}}u\|_{W(I_1)} + \|D^{2 - \frac{1}{2}}u\|_{W(I_1)}) \|u\|_{S(I_1)}^{p-1} \\ &\quad + g_j(\|u\|_{L_t^\infty \tilde{H}^2(I_1)}) \left( (\|u_{[j]}\|_{S(I_1)}^{p-1} + \|u\|_{S(I_1)}^{p-1}) (\|D^{s_p - \frac{1}{2}}w\|_{W(I_1)} + \|D^{2 - \frac{1}{2}}w\|_{W(I_1)}) \right. \\ &\quad \left. + (\|u_{[j]}\|_{S(I_1)}^{p-2} + \|u\|_{S(I_1)}^{p-2}) \|w\|_{S(I_1)} \right. \\ &\quad \left. \times (\|D^{s_p - \frac{1}{2}}u\|_{W(I_1)} + \|D^{2 - \frac{1}{2}}u\|_{W(I_1)} + \|D^{s_p - \frac{1}{2}}u_{[j]}\|_{W(I_1)} + \|D^{2 - \frac{1}{2}}u_{[j]}\|_{W(I_1)}) \right) \\ &\quad + \|g'_j(|u|)\|_{L_t^\infty L_x^\infty(I_1)} (\|D^{s_p - \frac{1}{2}}u\|_{W(I_1)} + \|D^{2 - \frac{1}{2}}u\|_{W(I_1)}) (\|u\|_{S(I_1)}^{p-2} + \|u_{[j]}\|_{S(I_1)}^{p-2}) \|w\|_{S(I_1)} \\ &\quad + \|g_j(|u|) - g_j(|u_{[j]}|)\|_{L_t^\infty L_x^\infty(I_1)} \|D^{s_p - \frac{1}{2}}u_{[j]}\|_{W(I_1)} \|u_{[j]}\|_{S(I_1)}^{p-1} \\ &\quad + \|u_{[j]}\|_{S(I_1)}^{p-1} \|u_{[j]}\|_{L_t^\infty \tilde{H}^2(I_1)} (\|w\|_{L_t^\infty \tilde{H}^2(I_1)} (\|D^{s_p - \frac{1}{2}}u\|_{W(I_1)} + \|D^{s_p - \frac{1}{2}}u_{[j]}\|_{W(I_1)}) + \|D^{s_p - \frac{1}{2}}w\|_{W(I_1)}) \\ &\lesssim g_j(C_j) \eta^{p-1} Q(I_1, w) + \eta^{p-1} Q(I_1, w) + \eta^p Q(I_1, w) + C_j \eta^{p-1} (\eta Q(I_1, w) + Q(I_1, w)), \end{aligned} \quad (5-13)$$

since, by choosing  $A$  large enough and by the construction of  $\tilde{g}$ , we have

$$(\tilde{g} - g_j)(\|u\|_{L_t^\infty \tilde{H}^2(I_1)}) = 0. \quad (5-14)$$

We conclude via a continuity argument that  $Q(I_1, w) = 0$ , so  $u = u_{[j]}$  on  $I_1$ . In particular,  $u(b_1) = u_{[j]}(b_1)$ . By iteration on  $i$ , it is not difficult to see that  $u = u_{[j]}$  on  $[-t, t]$ . Hence  $(\bar{t} - \alpha, \bar{t} + \alpha) \cap [0, j] \subset F_j$ , by (5-7). Thus  $F_j$  is open.

The upshot is that  $F_j = [0, j]$ , so  $\|u\|_{S([-T_0, T_0])} \leq C_j$ . This proves global well-posedness. Moreover, since  $j$  depends on  $T_0$  and  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}$ , we get (1-17).

*Proof of Lemma 5.1.* The proof extends to the end of the paper. We must establish *a priori* bounds.

Step 1: Construction of  $g_1$ .

Basically,  $g_1$  is a nonnegative function that increases and is equal to 2 for  $x$  large. Recall that  $[-T, T] \subset [-1, 1]$  and  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq 1$ . Let  $I \subset [-T, T]$ .

Observe that the point  $(\infty-, 3+) := (\frac{3+\epsilon}{\epsilon}, 3+\epsilon)$  with  $\epsilon \ll 1$  is  $\frac{1}{2}$ -wave admissible.

We would like to chop  $I$  (satisfying  $\|\cdot\|_{L_t^\infty L_x^3(I)} < \infty$ ) into subintervals  $I_j$  such that  $\|\cdot\|_{L_t^\infty L_x^3(I_j)}$  is as small as wanted. Unfortunately this is impossible because the  $L_t^\infty$ -norm is pathological. Instead we will apply this process to  $\|\cdot\|_{L_t^{\infty-} L_x^{3+}}$ . This creates slight variations almost everywhere in the process of the construction of  $g_i$ . Details with respect to these slight perturbations have been omitted for the sake of readability: they are left to the reader, who should ignore the  $+$  and  $-$  signs at the first reading.

We define

$$X(I) := D^{\frac{1}{2}-s_p} L_t^{\infty-} L_x^{3+}(I) \cap D^{\frac{1}{2}-s_p} W(I) \cap S(I) \cap L_t^\infty \dot{H}^{s_p}(I) \times L_t^\infty \dot{H}^{s_p-1}(I). \quad (5-15)$$

Let  $g_1$  be a smooth function, defined on the set of nonnegative real numbers, nondecreasing, and such that  $h_1 := g_1 - 2$  satisfies the following properties:  $h_1(0) = -1$ ,  $h$  is nondecreasing, and  $h_1(x) = 0$  if  $|x| \geq 1$ . It is not difficult to see that (1-2) and (1-10) are satisfied.

Observe that

$$|h_1(x)| \lesssim \frac{1}{|x|^{\frac{p-1}{2}-}} \quad (5-16)$$

and

$$|h'_1(x)| \lesssim \frac{1}{|x|^{\frac{p+1}{2}-}}. \quad (5-17)$$

Let  $u_{[\cdot]}$  and  $v_{[\cdot]}$  be solutions to the equations

$$\begin{cases} \partial_{tt} u_{[\cdot]} - \Delta u_{[\cdot]} = -|u_{[\cdot]}|^{p-1} u_{[\cdot]} g_1(|u_{[\cdot]}|), \\ u_{[\cdot]}(0) = u_0 \in \tilde{H}^2, \\ \partial_t u_{[\cdot]}(0) = u_1 \in \tilde{H}^1 \end{cases} \quad (5-18)$$

and

$$\begin{cases} \partial_{tt} v_{[\cdot]} - \Delta v_{[\cdot]} = -2|v_{[\cdot]}|^{p-1} v_{[\cdot]}, \\ v_{[\cdot]}(0) = u_0, \\ \partial_t v_{[\cdot]}(0) = u_1. \end{cases} \quad (5-19)$$

*Step 1a.* We claim that  $\|v_{[\cdot]}\|_{X(\mathbb{R})} < \infty$ . Indeed, since we assumed that Conjecture 1.1 is true, we can divide  $\mathbb{R}$  into subintervals  $(I_j = [t_j, t_{j+1}])_{1 \leq j \leq l}$  such that

$$\|v_{[\cdot]}\|_{S(I_j)} = \eta \quad \text{and} \quad \|v_{[\cdot]}\|_{S(I_l)} \leq \eta,$$

with  $\eta \ll 1$ . Then

$$\begin{aligned}
\|v_{[1]}\|_{X(I_{j+1})} &\lesssim \|(v_{[1]}(t_j), \partial_t v_{[1]}(t_j))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} + \|D^{sp-\frac{1}{2}}(|v_{[1]}|^{p-1}v_{[1]})\|_{\tilde{W}(I_{j+1})} \\
&\lesssim \|(v_{[1]}(t_j), \partial_t v_{[1]}(t_j))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} + \|D^{sp-\frac{1}{2}}v_{[1]}\|_{W(I_{j+1})}\|v_{[1]}\|_{S(I_{j+1})}^{p-1} \\
&\lesssim \|v_{[1]}\|_{X(I_j)} + \eta^{p-1}\|v_{[1]}\|_{X(I_{j+1})}.
\end{aligned} \tag{5-20}$$

Notice that  $l \lesssim 1$ : this follows from [Conjecture 1.1](#), [Condition 1.3](#) and the inequality

$$\begin{aligned}
\|(u_0, u_1)\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} &\leq \sup_{t \in I_{\max, g_1}} \|(u(t), \partial_t u(t))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} \leq C_2(\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1}) \\
&\lesssim 1,
\end{aligned} \tag{5-21}$$

following from [Condition 1.3](#) and the assumption  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq 1$ . (At this stage, we only need to know that  $\|(u_0, u_1)\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} \leq \|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq 1$  and apply [Conjecture 1.1](#). Therefore the introduction of  $\sup_{t \in I_{\max, g_1}} \|(u(t), \partial_t u(t))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)}$  in (5-21) is redundant. This is done on purpose. Indeed, we will use [Condition 1.3](#) in other parts of the argument: see (5-33).)

Now by a standard continuity argument and iteration on  $j$  we have

$$\|v_{[1]}\|_{X(\mathbb{R})} \lesssim 1 \tag{5-22}$$

*Step 1b.* We control  $\|u_{[1]} - v_{[1]}\|_{X([- \tilde{t}, \tilde{t}])}$ , for  $\tilde{t} \ll 1$  to be chosen later. By time reversal symmetry it is enough to control  $\|u_{[1]} - v_{[1]}\|_{X([0, \tilde{t}])}$ . To this end we consider  $w_{[1]} := u_{[1]} - v_{[1]}$ . We get

$$\partial_{tt} w_{[1]} - \Delta w_{[1]} = -|w_{[1]} + v_{[1]}|^{p-1}(v_{[1]} + w_{[1]})g_1(v_{[1]} + w_{[1]}) + 2|v_{[1]}|^{p-1}v_{[1]}.$$

Let  $\eta' \ll 1$ . By (5-22), we can divide  $[0, \tilde{t}]$  into subintervals  $(J_k = [t'_k, t'_{k+1}])_{1 \leq k \leq m}$  that satisfy

$$\|D^{sp-\frac{1}{2}}v_{[1]}\|_{L_t^\infty L_x^{3+}(J_k)} = \eta' \quad \text{or} \quad \|D^{sp-\frac{1}{2}}v_{[1]}\|_{W(J_k)} = \eta' \quad \text{for } 1 \leq k < m, \tag{5-23}$$

$$\|D^{sp-\frac{1}{2}}v_{[1]}\|_{W(J_k)} \leq \eta' \quad \text{and} \quad \|D^{sp-\frac{1}{2}}v_{[1]}\|_{L_t^\infty L_x^{3+}(J_k)} \leq \eta' \quad \text{for } 1 \leq k \leq m. \tag{5-24}$$

We have

$$\|w_{[1]}\|_{X(J_{k+1})} \lesssim \|(w_{[1]}(t'_k), \partial_t w_{[1]}(t'_k))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} + A_1 + A_2,$$

where

$$\begin{aligned}
A_1 &:= \|D^{sp-\frac{1}{2}}(2|v_{[1]}|^{p-1}v_{[1]} - 2|v_{[1]} + w_{[1]}|^{p-1}(v_{[1]} + w_{[1]}))\|_{\tilde{W}(J_{k+1})}, \\
A_2 &:= \|D^{sp-\frac{1}{2}}(h_1(|v_{[1]} + w_{[1]}|)|v_{[1]} + w_{[1]}|^{p-1}(v_{[1]} + w_{[1]}))\|_{L_t^1 L_x^{\frac{3}{2}}(J_{k+1})}.
\end{aligned} \tag{5-25}$$

By the fractional Leibnitz rule applied to  $q(x) := |x|^{p-1}xh(x)$ , (5-16), (5-17), Sobolev embedding and Hölder in time we have

$$\begin{aligned}
A_2 &\lesssim \| |v_{[1]} + w_{[1]}| ^{\frac{p-1}{2}+} \|_{L_t^{1+} L_x^{3-}(J_{k+1})} \| D^{s_{p-\frac{1}{2}}}(v_{[1]} + w_{[1]}) \|_{L_t^{\infty-} L_x^{3+}(J_{k+1})} \\
&\lesssim \| v_{[1]} + w_{[1]} \|_{L_t^{\frac{p-1}{2}+} L_x^{\frac{3(p-1)+}{2}(J_{k+1})}} \| D^{s_{p-\frac{1}{2}}}(v_{[1]} + w_{[1]}) \|_{L_t^{\infty-} L_x^{3+}(J_{k+1})} \\
&\lesssim \tilde{t} \| D^{s_{p-\frac{1}{2}}}(v_{[1]} + w_{[1]}) \|_{L_t^{\infty-} L_x^{3+}(J_{k+1})}^{\frac{p+1}{2}+} \lesssim \tilde{t}(\eta')^{\frac{p+1}{2}+} + \tilde{t} \| D^{s_{p-\frac{1}{2}}} w_{[1]} \|_{L_t^{\infty-} L_x^{3+}(J_{k+1})}^{\frac{p+1}{2}+} \\
&\lesssim \tilde{t}(\eta')^{\frac{p+1}{2}+} + \tilde{t} \| w_{[1]} \|_{X(J_{k+1})}^{\frac{p+1}{2}+}.
\end{aligned} \tag{5-26}$$

For  $A_1$  we follow [Kenig and Merle 2011, p. 9]:

$$\begin{aligned}
A_1 &\lesssim (\| v_{[1]} \|_{S(J_{k+1})}^{p-1} + \| w_{[1]} \|_{S(J_{k+1})}^{p-1}) \| D^{s_{p-\frac{1}{2}}} w_{[1]} \|_{W(J_{k+1})} \\
&\quad + (\| v_{[1]} \|_{S(J_{k+1})}^{p-2} + \| w_{[1]} \|_{S(J_{k+1})}^{p-2}) (\| D^{s_{p-\frac{1}{2}}} v_{[1]} \|_{W(J_{k+1})} + \| D^{s_{p-\frac{1}{2}}} w_{[1]} \|_{W(J_{k+1})}) \| w_{[1]} \|_{S(J_{k+1})} \\
&\lesssim (\eta')^{p-1} \| w_{[1]} \|_{X(J_{k+1})} + \| w_{[1]} \|_{X(J_{k+1})}^p + (\eta')^{p-2} \| w_{[1]} \|_{X(J_{k+1})}^2 + \eta' \| w_{[1]} \|_{X(J_{k+1})}^{p-1}.
\end{aligned} \tag{5-27}$$

This follows from (1-31) and (1-27). Therefore we have

$$\begin{aligned}
\| w_{[1]} \|_{X(J_{k+1})} &\lesssim \| w_{[1]} \|_{X(J_k)} + (\eta')^{\frac{p+1}{2}+} \tilde{t} + \tilde{t} \| w_{[1]} \|_{X(J_{k+1})}^{\frac{p+1}{2}+} \\
&\quad + (\eta')^{p-1} \| w_{[1]} \|_{X(J_{k+1})} + \| w_{[1]} \|_{X(J_{k+1})}^p + (\eta')^{p-2} \| w_{[1]} \|_{Y(J_{k+1})}^2 + \eta' \| w_{[1]} \|_{X(J_{k+1})}^{p-1}.
\end{aligned} \tag{5-28}$$

Let  $C$  be the constant determined by (5-28). By induction, we have

$$\| w_{[1]} \|_{X(J_k)} \leq (2C)^k \tilde{t}, \tag{5-29}$$

provided that for  $1 \leq k \leq m-1$  we have

$$\begin{aligned}
C(\eta')^{\frac{p+1}{2}+} \tilde{t} &\ll C(2C)^k \tilde{t}, \\
C \tilde{t} ((2C)^k \tilde{t})^{\frac{p+1}{2}+} &\ll (2C)^k \tilde{t}, \\
C(\eta')^{p-1} (2C)^{k+1} \tilde{t} &\ll C(2C)^k \tilde{t}, \\
C ((2C)^k \tilde{t})^p &\ll C(2C)^k \tilde{t}, \\
C(\eta')^{p-2} ((2C)^{k+1} \tilde{t})^2 &\ll C(2C)^k \tilde{t}, \\
\eta' ((2C)^{k+1} \tilde{t})^{p-1} &\ll C(2C)^k.
\end{aligned} \tag{5-30}$$

These inequalities are satisfied if  $\eta' \ll 1$  and

$$\tilde{t} \ll 1 \tag{5-31}$$

since  $k \leq m-1$  and, by (5-22),  $m \lesssim 1$ . We conclude that

$$\| w_{[1]} \|_{X([0, \tilde{t}])} \lesssim 1. \tag{5-32}$$

*Step 1c.* We control  $\|u_{[1]}\|_{X([-T, T])}$ . By time reversal symmetry, it is enough to control  $\|u_{[1]}\|_{X([0, T])}$ . Recall that  $T \leq 1$ . We chop  $T \leq 1$  into subintervals  $(J_{k'} = [a_{k'}, b_{k'}])_{1 \leq k' \leq l'}$  such that  $|J_{k'}| = \tilde{t}$  for  $1 \leq k' < l'$  and  $|J_{l'}| \leq \tilde{t}$ . Notice that, by [Condition 1.3](#), we have

$$\begin{aligned} \|(u(a_{k'}), \partial_t u(a_{k'}))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} &\leq \sup_{t \in I_{\max, g_1}} \|(u(t), \partial_t u(t))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} \\ &\leq C_2 (\|u_0, u_1\|_{\tilde{H}^2 \times \tilde{H}^1}) \lesssim 1, \end{aligned} \quad (5-33)$$

taking advantage of the assumption  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq 1$ . For each  $k'$  let  $v_{[1, k']}$  be the solution of

$$\begin{cases} \partial_{tt} v_{[1, k']} - \Delta v_{[1, k']} = -|v_{[1, k']}|^{p-1} v_{[1, k']}, \\ v_{[1, k']}(a_{k'}) = u_{[1]}(a_{k'}), \\ \partial_t v_{[1, k']}(a_{k'}) = \partial_t u_{[1]}(a_{k'}); \end{cases} \quad (5-34)$$

in particular,  $v_{[1, k']} = v_{[1]}$ . By slightly modifying the proof of [Step 1b](#) and letting  $v_{[1, k']}$  play the role of  $v_{[1]}$ , this leads, by [\(5-33\)](#), to

$$\|v_{[1, k']}\|_{X(\mathbb{R})} \lesssim 1 \quad (5-35)$$

and

$$\|w_{[1, k']}\|_{X(J_{k'})} \lesssim 1, \quad (5-36)$$

with  $w_{[1, k']} := u_{[1]} - v_{[1, k']}$ . Therefore  $\|u_{[1]}\|_{X(J_{k'})} \lesssim 1$ , and summing over  $J_{k'}$  we have

$$\|u_{[1]}\|_{X([0, T])} \lesssim 1. \quad (5-37)$$

*Step 1d.* We control  $\|(u_{[1]}, \partial_t u_{[1]})\|_{L_t^\infty \tilde{H}^2([-1, 1]) \times L_t^\infty \tilde{H}^1([-1, 1])}$  and  $\|u_{[1]}\|_{S([-1, 1])}$ . We get from [\(5-37\)](#)

$$\|u_{[1]}\|_{S([-1, 1])} \lesssim 1. \quad (5-38)$$

To conclude Step 1: By [Proposition 1.8](#) and [\(5-38\)](#) we have

$$\|(u_{[1]}, \partial_t u_{[1]})\|_{L_t^\infty \tilde{H}^2([-1, 1]) \times L_t^\infty \tilde{H}^1([-1, 1])} \lesssim 1. \quad (5-39)$$

Therefore

$$\max(\|u_{[1]}\|_{S([-1, 1])}, \|(u_{[1]}, \partial_t u_{[1]})\|_{L_t^\infty \tilde{H}^2([-1, 1]) \times L_t^\infty \tilde{H}^1([-1, 1])}) \lesssim 1. \quad (5-40)$$

We let  $C'_1$  in the statement of [Lemma 5.1](#) be equal to 1. We can assume without the loss of generality that the constant implicit in  $\lesssim$  in [\(5-40\)](#) is larger than 1; let  $C_1$  in the statement of [Lemma 5.1](#) be this constant. Then  $C'_1$  and  $C_1$  satisfy [\(5-2\)](#) and [\(5-3\)](#).

Step 2: Construction of  $g_i$  from  $g_{i-1}$ .

Recall that  $[-T, T] \subset [-i, i]$  and  $\|(u_0, u_1)\|_{\tilde{H}^2 \times \tilde{H}^1} \leq i$ . In view of [\(5-5\)](#) it is enough to construct  $g_i$  for  $|x| > AC_{i-1}$ . It is clear that, by choosing  $C'_i$  large enough, we can construct find a function  $\tilde{g}_i$  defined on  $[AC_{i-1}, C'_i]$  such that  $g_i$ , defined by

$$g_i(x) := \begin{cases} g_{i-1}(x) & \text{if } |x| \leq AC_{i-1}, \\ \tilde{g}_i(x) & \text{if } C'_i \geq |x| \geq AC_{i-1}, \\ i+1 & \text{if } |x| \geq C'_i \end{cases} \quad (5-41)$$



is smooth and slowly increasing; also it satisfies (1-2), (1-10), and

$$\int_{AC_{i-1}}^{C'_i} \frac{1}{yg_i^2(y)} dy \geq i. \quad (5-42)$$

It remains to determine  $C_i$  in the statement of Lemma 5.1. To do that we slightly modify the reasoning in Step 1.

We sketch the argument. Let  $h_i(x) := g_i(x) - (i+1)$ . Then  $h_i(x) = 0$  if  $|x| > C'_i$ . It is not difficult to see that

$$|h_i(x)| \lesssim_i \frac{1}{|x|^{\frac{p-1}{2}+}}, \quad (5-43)$$

$$|h'_i(x)| \lesssim_i \frac{1}{|x|^{\frac{p+1}{2}+}}. \quad (5-44)$$

Let  $u_{[i]}$  and  $v_{[i]}$  be the solutions of the equations

$$\begin{cases} \partial_{tt} u_{[i]} - \Delta u_{[i]} = -|u_{[i]}|^{p-1} u_{[i]} g_i(|u_{[i]}|), \\ u_{[i]}(0) := u_0, \\ \partial_t u_{[i]}(0) := u_1 \end{cases} \quad (5-45)$$

and

$$\begin{cases} \partial_{tt} v_{[i]} - \Delta v_{[i]} = -(i+1)|v_{[i]}|^{p-1} v_{[i]}, \\ v_{[i]}(0) := u_0, \\ \partial_t v_{[i]}(0) := u_1 \end{cases} \quad (5-46)$$

*Step 2a.* We have

$$\|v_{[i]}\|_{X(\mathbb{R})} \lesssim_i 1, \quad (5-47)$$

by adapting the proof of Step 1a. Notice, in particular, that we can use Conjecture 1.1 and control  $\|v_{[i]}\|_{S(\mathbb{R})}$  since  $w_{[i]} := (i+1)^{\frac{1}{p-1}} v_{[i]}$  satisfies  $\partial_{tt} w_{[i]} - \Delta w_{[i]} = -|w_{[i]}|^{p-1} w_{[i]}$ .

*Step 2b.* We have  $\|u_{[i]} - v_{[i]}\|_{X([0, \tilde{t}])} \lesssim_i 1$  for  $\tilde{t} \ll 1$ , by adapting the proof of Step 1b. The dependance on  $i$  basically comes from (5-43), (5-44) and (5-46).

*Step 2c.* We prove that  $\|u_{[i]}\|_{X([-T, T])} \lesssim_{i,p} 1$ . By time reversal symmetry, it is enough to control  $\|u_{[i]}\|_{X([0, T])}$ . Recall that  $T \leq i$ . We chop  $[0, T]$  into subintervals  $(J_{k'} = [a_{k'}, b_{k'}])_{1 \leq k' \leq l'}$  such that  $|J_{k'}| = \tilde{t}$  for  $1 \leq k' < l'$  and  $|J_{l'}| \leq \tilde{t}$  (with  $\tilde{t}$  defined in Step 2b). By Condition 1.3 and the assumption  $\|(u_0, u_1)\|_{\dot{H}^2 \times \dot{H}^1} \leq i$ , we have

$$\|(u_{[i]}(a_{k'}), \partial_t u_{[i]}(a_{k'}))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} \leq \sup_{t \in I_{\max, g_i}} \|(u_{[i]}(t), \partial_t u_{[i]}(t))\|_{\dot{H}^{sp}(\mathbb{R}^3) \times \dot{H}^{sp-1}(\mathbb{R}^3)} \lesssim_i 1. \quad (5-48)$$

We introduce

$$\begin{cases} \partial_{tt} v_{[i, k']} - \Delta v_{[i, k']} = -(i+1)|v_{[i, k']}|^{p-1} v_{[i, k']}, \\ v_{[i, k']}(a_{k'}) = u_{[i]}(a_{k'}), \\ \partial_t v_{[i, k']}(a_{k'}) = \partial_t u_{[i]}(a_{k'}) \end{cases} \quad (5-49)$$

and, by using (5-48), we can prove that

$$\|u_{[i]}\|_{S([-i,i])} \lesssim_i 1. \quad (5-50)$$

*Step 2d.* By using Proposition 1.8 and (5-50) we get

$$\max \left( \|u_{[i]}\|_{S([-i,i])}, \|(u_{[i]}, \partial_t u_{[i]})\|_{L_t^\infty \tilde{H}^2([-i,i]) \times L_t^\infty \tilde{H}^1([-i,i])} \right) \lesssim_i 1. \quad (5-51)$$

We can assume without loss of generality that the constant implicit in  $\lesssim$  is larger than  $i$  and  $C'_i$ . Let  $C_i$  be this constant; (5-2) and (5-3) are satisfied.

This concludes Step 2, and the proof of Lemma 5.1.  $\square$

### Acknowledgements

The author would like to thank Terence Tao for suggesting this problem and for valuable discussions related to this work.

### References

- [Bahouri and Chemin 2006] H. Bahouri and J.-Y. Chemin, “On global well-posedness for defocusing cubic wave equation”, *Int. Math. Res. Not.* **2006** (2006), Art. ID 54873. [MR 2008b:35178](#)
- [Bahouri and Gérard 1999] H. Bahouri and P. Gérard, “High frequency approximation of solutions to critical nonlinear wave equations”, *Amer. J. Math.* **121**:1 (1999), 131–175. [MR 2000i:35123](#) [Zbl 0919.35089](#)
- [Bahouri and Shatah 1998] H. Bahouri and J. Shatah, “Decay estimates for the critical semilinear wave equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **15**:6 (1998), 783–789. [MR 99h:35136](#) [Zbl 0924.35084](#)
- [Christ and Weinstein 1991] F. M. Christ and M. I. Weinstein, “Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation”, *J. Funct. Anal.* **100**:1 (1991), 87–109. [MR 92h:35203](#) [Zbl 0743.35067](#)
- [Gallagher and Planchon 2003] I. Gallagher and F. Planchon, “On global solutions to a defocusing semi-linear wave equation”, *Rev. Mat. Iberoamericana* **19**:1 (2003), 161–177. [MR 2004k:35265](#) [Zbl 1036.35142](#)
- [Ginibre and Velo 1989] J. Ginibre and G. Velo, “Scattering theory in the energy space for a class of nonlinear wave equations”, *Comm. Math. Phys.* **123**:4 (1989), 535–573. [MR 90i:35172](#) [Zbl 0698.35112](#)
- [Ginibre and Velo 1995] J. Ginibre and G. Velo, “Generalized Strichartz inequalities for the wave equation”, *J. Funct. Anal.* **133**:1 (1995), 50–68. [MR 97a:46047](#) [Zbl 0849.35064](#)
- [Grillakis 1990] M. G. Grillakis, “Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity”, *Ann. of Math. (2)* **132**:3 (1990), 485–509. [MR 92c:35080](#) [Zbl 0736.35067](#)
- [Grillakis 1992] M. G. Grillakis, “Regularity for the wave equation with a critical nonlinearity”, *Comm. Pure Appl. Math.* **45**:6 (1992), 749–774. [MR 93e:35073](#) [Zbl 0785.35065](#)
- [Kapitanski 1994] L. Kapitanski, “Global and unique weak solutions of nonlinear wave equations”, *Math. Res. Lett.* **1**:2 (1994), 211–223. [MR 95f:35158](#) [Zbl 0841.35067](#)
- [Kenig and Merle 2006] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case”, *Invent. Math.* **166**:3 (2006), 645–675. [MR 2007g:35232](#)
- [Kenig and Merle 2011] C. E. Kenig and F. Merle, “Nondispersive radial solutions to energy supercritical non-linear wave equations, with applications”, *Amer. J. Math.* **133**:4 (2011), 1029–1065. [MR 2823870](#) [Zbl 05947418](#)
- [Kenig et al. 1993] C. E. Kenig, G. Ponce, and L. Vega, “Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle”, *Comm. Pure Appl. Math.* **46**:4 (1993), 527–620. [MR 94h:35229](#) [Zbl 0808.35128](#)
- [Kenig et al. 2000] C. E. Kenig, G. Ponce, and L. Vega, “Global well-posedness for semi-linear wave equations”, *Comm. Partial Differential Equations* **25**:9-10 (2000), 1741–1752. [MR 2001h:35128](#)

- [Lindblad and Sogge 1995] H. Lindblad and C. D. Sogge, “On existence and scattering with minimal regularity for semilinear wave equations”, *J. Funct. Anal.* **130**:2 (1995), 357–426. [MR 96i:35087](#) [Zbl 0846.35085](#)
- [Rauch 1981] J. Rauch, “I: The  $u^5$  Klein–Gordon equation; II: Anomalous singularities for semilinear wave equations”, pp. 335–364 in *Nonlinear partial differential equations and their applications* (Paris, 1978/1979), edited by H. Brezis and J. L. Lions, Res. Notes in Math. **53**, Pitman, Boston, MA, 1981. [MR 83a:35066](#)
- [Roy 2007] T. Roy, “Global well-posedness for the radial defocusing cubic wave equation on  $\mathbb{R}^3$  and for rough data”, *Electron. J. Differential Equations* (2007), No. 166, 22 pp. [MR 2008m:35247](#)
- [Roy 2009a] T. Roy, “Adapted linear-nonlinear decomposition and global well-posedness for solutions to the defocusing cubic wave equation on  $\mathbb{R}^3$ ”, *Discrete Contin. Dyn. Syst.* **24**:4 (2009), 1307–1323. [MR 2010j:35327](#) [Zbl 1211.47023](#)
- [Roy 2009b] T. Roy, “Global existence of smooth solutions of a 3D log-log energy-supercritical wave equation”, *Anal. PDE* **2**:3 (2009), 261–280. [MR 2011k:35145](#) [Zbl 1195.35222](#)
- [Shatah and Struwe 1994] J. Shatah and M. Struwe, “Well-posedness in the energy space for semilinear wave equations with critical growth”, *Internat. Math. Res. Notices* **7** (1994), 303–309. [MR 95e:35132](#) [Zbl 0830.35086](#)
- [Struwe 1988] M. Struwe, “Globally regular solutions to the  $u^5$  Klein–Gordon equation”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **15**:3 (1988), 495–513. [MR 90j:35142](#) [Zbl 0728.35072](#)
- [Tao 2006a] T. Tao, *Nonlinear dispersive equations: Local and global analysis*, CBMS Regional Conference Series in Mathematics **106**, Amer. Math. Soc., Providence, 2006. [MR 2008i:35211](#) [Zbl 1106.35001](#)
- [Tao 2006b] T. Tao, “Spacetime bounds for the energy-critical nonlinear wave equation in three spatial dimensions”, *Dyn. Partial Differ. Equ.* **3**:2 (2006), 93–110. [MR 2007c:35116](#)
- [Tao 2007] T. Tao, “Global regularity for a logarithmically supercritical defocusing nonlinear wave equation for spherically symmetric data”, *J. Hyperbolic Differ. Equ.* **4**:2 (2007), 259–265. [MR 2009b:35294](#) [Zbl 1124.35043](#)
- [Zhang 2006] X. Zhang, “On the Cauchy problem of 3-D energy-critical Schrödinger equations with subcritical perturbations”, *J. Differential Equations* **230**:2 (2006), 422–445. [MR 2007h:35325](#)

Received 26 Apr 2010. Revised 17 Jul 2010. Accepted 16 Aug 2010.

TRISTAN ROY: [triroy@ias.edu](mailto:triroy@ias.edu)

School of Mathematics, Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, Institute For Advanced Study

# Analysis & PDE

[msp.berkeley.edu/apde](http://msp.berkeley.edu/apde)

## EDITORS

### EDITOR-IN-CHIEF

Maciej Zworski  
University of California  
Berkeley, USA

### BOARD OF EDITORS

Michael Aizenman	Princeton University, USA <a href="mailto:aizenman@math.princeton.edu">aizenman@math.princeton.edu</a>	Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>
Luis A. Caffarelli	University of Texas, USA <a href="mailto:caffarel@math.utexas.edu">caffarel@math.utexas.edu</a>	Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Nigel Higson	Pennsylvania State University, USA <a href="mailto:higson@math.psu.edu">higson@math.psu.edu</a>
Vaughan Jones	University of California, Berkeley, USA <a href="mailto:vfr@math.berkeley.edu">vfr@math.berkeley.edu</a>	Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>

## PRODUCTION

[contact@msp.org](mailto:contact@msp.org)

Silvio Levy, Scientific Editor

Sheila Newbery, Senior Production Editor

---

See inside back cover or [msp.berkeley.edu/apde](http://msp.berkeley.edu/apde) for submission instructions.

---

The subscription price for 2012 is US \$140/year for the electronic version, and \$240/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

---

Analysis & PDE, at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2012 by Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 5    No. 1    2012

---

A characterization of two weight norm inequalities for maximal singular integrals with one doubling measure	1
MICHAEL LACEY, ERIC T. SAWYER and IGNACIO URIARTE-TUERO	
Energy identity for intrinsically biharmonic maps in four dimensions	61
PETER HORNUNG and ROGER MOSER	
The wave equation on asymptotically anti de Sitter spaces	81
ANDRÁS VASY	
Small data scattering and soliton stability in $\dot{H}^{-1/6}$ for the quartic KdV equation	145
HERBERT KOCH and JEREMY L. MARZUOLA	
A remark on barely $\dot{H}^{s_p}$ -supercritical wave equations	199
TRISTAN ROY	