

ANALYSIS & PDE

Volume 6

No. 1

2013

HSI-WEI SHIH

**SOME RESULTS ON SCATTERING FOR LOG-SUBCRITICAL AND
LOG-SUPERCritical NONLINEAR WAVE EQUATIONS**

SOME RESULTS ON SCATTERING FOR LOG-SUBCRITICAL AND LOG-SUPERCRITICAL NONLINEAR WAVE EQUATIONS

HSI-WEI SHIH

We consider two problems in the asymptotic behavior of semilinear second order wave equations. First, we consider the $\dot{H}_x^1 \times L_x^2$ scattering theory for the energy log-subcritical wave equation

$$\square u = |u|^4 u g(|u|)$$

in \mathbb{R}^{1+3} , where g has logarithmic growth at 0. We discuss the solution with general (respectively spherically symmetric) initial data in the logarithmically weighted (respectively lower regularity) Sobolev space. We also include some observation about scattering in the energy subcritical case. The second problem studied involves the energy log-supercritical wave equation

$$\square u = |u|^4 u \log^\alpha(2 + |u|^2) \quad \text{for } 0 < \alpha \leq \frac{4}{3}$$

in \mathbb{R}^{1+3} . We prove the same results of global existence and $(\dot{H}_x^1 \cap \dot{H}_x^2) \times H_x^1$ scattering for this equation with a slightly higher power of the logarithm factor in the nonlinearity than that allowed in previous work by Tao (*J. Hyperbolic Differ. Equ.*, **4**:2 (2007), 259–265).

1. Introduction

Consider the semilinear wave equation

$$\begin{aligned} \square u &:= -\partial_t^2 u + \Delta u = f(u) \quad \text{on } \mathbb{R} \times \mathbb{R}^3, \\ u(0, x) &= u_0(x), \\ \partial_t u(0, x) &= u_1(x), \end{aligned} \tag{1}$$

where f is a complex-valued function. Let the potential function $F : \mathbb{C} \rightarrow \mathbb{R}$ be a real-valued function such that

$$2F_{\bar{z}}(z) = f(z), \tag{2}$$

with $F(0) = 0$ and u being the solution to (1) with initial data $u_0 \in \dot{H}_x^1 \cap \{\phi : \int_{\mathbb{R}^3} F(\phi) dx < \infty\}$ and $u_1 \in L_x^2$. We can easily verify that the equation has conserved energy

$$E(u)(t) := \int_{\mathbb{R}^3} \frac{1}{2} |u_t(t, x)|^2 + \frac{1}{2} |\nabla u(t, x)|^2 + F(u(t, x)) dx. \tag{3}$$

MSC2010: 35L15.

Keywords: scattering, log-subcritical, radial Sobolev inequality.

The main goal of the paper is to study the $\dot{H}_x^1 \times L_x^2$ scattering theory for log-subcritical wave equations with finite energy initial data, where the energy is defined by (3). In this paper, the term *log-subcritical wave equation* refers to (1) with f defined by

$$f(z) := \begin{cases} |z|^4 z g(|z|), & |z| \neq 0, \\ 0, & |z| = 0, \end{cases} \quad (4)$$

where $g : (0, \infty) \rightarrow \mathbb{R}$ is smooth, nonincreasing, and satisfies

$$g(x) := \begin{cases} -\log x, & 0 < x < \frac{1}{3}, \\ \sim 1, & \frac{1}{3} \leq x < 1, \\ 1, & x \geq 1. \end{cases} \quad (5)$$

We also prove global existence in the case of spherical symmetry for *log-supercritical wave equations*, by which we mean equations of the form

$$\square u = |u|^4 u \log^\alpha(2 + |u|^2) \quad (6)$$

In this paper, we will allow $0 < \alpha \leq \frac{4}{3}$, extending the range $0 < \alpha \leq 1$ allowed in [Tao 2007]. We also assume that the initial data is in the energy space, the set of data for which the energy (3) is finite.

Remark 1.1. We can easily compute that the potential function of log-subcritical wave equations (1), (4), and (5) is

$$F_{\text{sub}}(z) = \begin{cases} -\frac{1}{6}|z|^6(\log(|z|) - \frac{1}{6}), & 0 < |z| < \frac{1}{3}, \\ \sim \frac{1}{6}|z|^6, & \frac{1}{3} \leq |z| < 1, \\ \frac{1}{6}|z|^6, & |z| \geq 1, \end{cases} \quad (7)$$

and the potential function of the log-supercritical wave equations (6) is

$$F_{\text{sup}}(z) \sim |z|^6 \log^\alpha(2 + |z|^2). \quad (8)$$

We quickly recall some common terminology associated to the scaling properties of (1). Consider $f(z) = |z|^{p-1}z$ and let u be the solution of (1). By scaling, $\lambda^{2/(1-p)}u(t/\lambda, x/\lambda)$ is also a solution with initial data $\lambda^{2/(1-p)}u_0(t_0/\lambda, x/\lambda)$ and $\lambda^{(1+p)/(1-p)}u_1(t_0/\lambda, x/\lambda)$. Hence the scaling of u preserves the homogeneous Sobolev norm $\|u_0\|_{\dot{H}^{s_c}(\mathbb{R}^3)} + \|u_1\|_{\dot{H}^{s_c-1}(\mathbb{R}^3)}$ if

$$s_c := \frac{3}{2} - \frac{2}{p-1}, \quad \text{or equivalently} \quad p = 1 + \frac{4}{3-2s_c}.$$

Definition 1.2. For $f(z) = |z|^{p-1}z$ and a given value s , we call (1) an \dot{H}_x^s -critical (subcritical, supercritical) nonlinear wave equation if p equals (is less than, is greater than) $1 + 4/(3-2s)$. In particular, when $s = 1$, we call (1) an energy critical (subcritical, supercritical) nonlinear wave equation if $p = 5$ ($p < 5$, $p > 5$).

The results of global existence and uniqueness for the energy-critical ($\square u = |u|^4 u$) and energy-subcritical ($\square u = |u|^{p-1}u$, where $p < 5$) wave equations are already established by [Brenner and von Wahl 1981; Struwe 1988; Grillakis 1990; 1992; Shatah and Struwe 1993; 1994; Kapitanski 1994; Ginibre and Velo 1985]. It is natural to consider the decay of the solution, which we expect to behave linearly

as $t \rightarrow \pm\infty$. The decay estimate and scattering theory (see section 2 for definition) of critical wave equations are shown in [Bahouri and Shatah 1998]; see also [Bahouri and Gérard 1999; Ginibre and Velo 1989; Nakanishi 1999]. Hidano [2001] (see also [Ginibre and Velo 1987]), by the property of conformal invariance, proved that the solutions for certain subcritical wave equations ($\frac{5}{2} < p \leq 3$) scatter in the weighted Sobolev space $\Sigma := X \times Y$, where

$$X := H_x^1(\mathbb{R}^3) \cap \{\phi : |x|\nabla\phi \in L_x^2(\mathbb{R}^3)\}, \quad Y := L_x^2(\mathbb{R}^3) \cap \{\phi : |x|\phi \in L_x^2(\mathbb{R}^3)\}.$$

However, for energy subcritical equations, the $\dot{H}_x^1 \times L_x^2$ scattering theory¹ still remains open. In this paper, we consider the solutions to the log-subcritical wave equations (1), (4), and (5) with finite energy initial data. The global existence result is established in [Grillakis 1990; 1992; Kapitanski 1994; Nakanishi 1999]. We will prove that the solutions with a class of initial data scatter in $\dot{H}_x^1 \times L_x^2$. This class of data is contained in logarithmically weighted Sobolev spaces $X_1 \times Y_1$, where

$$\begin{aligned} X_1 &:= \dot{H}_x^1(\mathbb{R}^3) \cap \{\phi : \log^\gamma(1+|x|)\nabla\phi \in L_x^2(\mathbb{R}^3)\}, \\ Y_1 &:= L_x^2(\mathbb{R}^3) \cap \{\phi : \log^\gamma(1+|x|)\phi \in L_x^2(\mathbb{R}^3)\} \end{aligned} \quad (9)$$

for some $\gamma > \frac{1}{2}$. For initial data in these spaces, we show that the potential energy of the solution decays logarithmically for all large times. After dividing the time interval suitably, this decay helps us to control the key spacetime norm $\|f(u)\|_{L_t^1 L_x^2}$. This spacetime bound implies scattering (we will sketch the proof in Section 2; see also [Bahouri and Shatah 1998]). Our proof of the spacetime bound involves establishing a decay rate for certain constant-time norms of the solution and a bootstrap scheme motivated by that in [Tao 2007]. We rely heavily on ideas from [Bahouri and Shatah 1998].

The second part of this paper considers the solution of log-subcritical wave equations with spherically symmetric data. We prove that the solution u with initial data in $X_2 \times Y_2$ scatters in $\dot{H}_x^1 \times L_x^2$, where

$$X_2 := \dot{H}_x^1(\mathbb{R}^3) \cap \left(\bigcup_{\delta>0} \dot{H}_x^{1-\delta}(\mathbb{R}^3) \right), \quad Y_2 := L_x^2(\mathbb{R}^3) \cap \left(\bigcup_{\delta>0} \dot{H}_x^{-\delta}(\mathbb{R}^3) \right). \quad (10)$$

Our proof again uses the ideas from [Tao 2007] and the classical Morawetz inequality; see [Morawetz 1968]. However, we need a slightly sharpened version of the bootstrap argument. We also give remarks for some specific energy subcritical wave equations (see page 15 and following).

The third part of this paper studies global existence for log-supercritical wave equations. The global regularity of energy supercritical wave equations ($\square u = |u|^{p-1}u$, where $p > 5$) is still open. In [Tao 2007], the author considered the log-supercritical wave equation

$$\square u = u^5 \log^\alpha(2+u^2) \quad (11)$$

with spherically symmetric initial data and established a global regularity result for $0 < \alpha \leq 1$. For general initial data, the same result for loglog-supercritical wave equations

$$\square u = u^5 \log^c(\log(10+u^2))$$

¹ $\dot{H}_x^1 \times L_x^2$ scattering is defined in Definition 2.1.

with $0 < c < \frac{8}{225}$ is obtained in [Roy 2009]. In the present paper, we extend the result in [Tao 2007] to the range $0 < \alpha \leq \frac{4}{3}$, again for spherically symmetric data. This improvement is attained by employing the potential energy bound in place of the kinetic energy bound used in [Tao 2007] for pointwise control.

2. Definitions, notation, and preliminaries

Throughout this paper, we use $M \lesssim N$ to denote the estimate $M \leq CN$ for some absolute constant C (which can vary from line to line).

We use $L_t^q L_x^r$ to denote the spacetime norm

$$\|u\|_{L_t^q L_x^r(I \times \mathbb{R}^3)} := \left(\int_I \left(\int_{\mathbb{R}^3} |u(t, x)|^r dx \right)^{q/r} dt \right)^{1/q}$$

with the usual modifications when q or r is equal to infinity.

Definition 2.1. We say that a global solution $u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$ to (1) scatters in $\dot{H}_x^1 \times L_x^2$ (or $\dot{H}_x^1 \times L_x^2$ scattering) as $t \rightarrow +\infty$ ($-\infty$) if there exists a linear solution v^+ (v^-) with initial data in $\dot{H}_x^1 \times L_x^2$ such that

$$\begin{aligned} \|u(t, x) - v^+(t, x)\|_{\dot{H}_x^1 \times L_x^2} &\rightarrow 0 \quad \text{as } t \rightarrow +\infty \\ (\|u(t, x) - v^-(t, x)\|_{\dot{H}_x^1 \times L_x^2} &\rightarrow 0 \quad \text{as } t \rightarrow -\infty). \end{aligned}$$

Remark 2.2. We will sketch here that the *spacetime bound*,

$$\|f(u)\|_{L_t^1 L_x^2((t_0, \infty) \times \mathbb{R}^3)} < \infty \quad (12)$$

for some $t_0 > 0$, of the solution u to (1) implies the $\dot{H}_x^1 \times L_x^2$ scattering (as $t \rightarrow \infty$). Let

$$u \in C_t^1(\mathbb{R}, \dot{H}_x^1(\mathbb{R}^3)) \cap C_t^0(\mathbb{R}, L_x^2(\mathbb{R}^3))$$

be the solution to (1) and let v satisfy $\square v = 0$ with initial data $v_0 \in \dot{H}_x^1(\mathbb{R}^3)$, $v_1 \in L_x^2(\mathbb{R}^3)$ (to be chosen shortly). By Duhamel's formula,

$$u(t, x) = \cos(t\sqrt{-\Delta})u_0(x) + \frac{\sin(t\sqrt{-\Delta})}{\sqrt{-\Delta}}u_1(x) - \int_0^t \frac{\sin((t-\tau)\sqrt{-\Delta})}{\sqrt{-\Delta}}f(u(\tau))d\tau \quad (13)$$

and

$$v(t, x) = \cos(t\sqrt{-\Delta})v_0(x) + \frac{\sin(t\sqrt{-\Delta})}{\sqrt{-\Delta}}v_1(x), \quad (14)$$

where the operators $\cos(t\sqrt{-\Delta})$ and $\sin(t\sqrt{-\Delta})/\sqrt{-\Delta}$ are defined by

$$(\cos(t\sqrt{-\Delta})\phi)^\wedge(\xi) = \cos(t|\xi|)\hat{\phi}(\xi)$$

and

$$\left(\frac{\sin(t\sqrt{-\Delta})}{\sqrt{-\Delta}}\phi \right)^\wedge(\xi) = \frac{\sin(t|\xi|)}{|\xi|}\hat{\phi}(\xi).$$

Hence, that the solution u scatters and asymptotically approaches v in $\dot{H}_x^1 \times L_x^2$ means that

$$\left\| \cos(t\sqrt{-\Delta})(u_0 - v_0) + \frac{\sin(t\sqrt{-\Delta})}{\sqrt{-\Delta}}(u_1 - v_1) - \int_0^t \frac{\sin((t-\tau)\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau \right\|_{\dot{H}_x^1 \times L_x^2} \rightarrow 0 \quad (15)$$

as $t \rightarrow \infty$. From basic trigonometric identities, we can verify that (15) is implied by

$$\left\| (u_0 - v_0) + \int_0^t \frac{\sin(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau \right\|_{\dot{H}_x^1} \rightarrow 0$$

and

$$\left\| (u_1 - v_1) + \int_0^t \cos(-\tau\sqrt{-\Delta}) f(u(\tau)) d\tau \right\|_{L_x^2} \rightarrow 0$$

as $t \rightarrow \infty$. Therefore, if

$$\left(\int_0^t \frac{\sin(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau, \int_0^t \cos(-\tau\sqrt{-\Delta}) f(u(\tau)) d\tau \right) \quad (16)$$

converges in $\dot{H}_x^1 \times L_x^2$ as $t \rightarrow \infty$, and we take

$$\begin{aligned} v_0(x) &:= u_0(x) - \int_0^\infty \frac{\sin(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau, \\ v_1(x) &:= u_1(x) - \int_0^\infty \cos(-\tau\sqrt{-\Delta}) f(u(\tau)) d\tau, \end{aligned}$$

in (14), we then have, by (13), (14), and elementary trigonometric formulas,

$$\begin{aligned} \|u - v\|_{\dot{H}_x^1 \times L_x^2} &= \left\| -\int_0^t \frac{\sin((t-\tau)\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau + \int_0^\infty \frac{\cos(t\sqrt{-\Delta}) \sin(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau \right. \\ &\quad \left. + \int_0^\infty \frac{\sin(t\sqrt{-\Delta}) \cos(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau \right\|_{\dot{H}_x^1 \times L_x^2} \\ &= \left\| \int_t^\infty \frac{\sin(t-\tau)\sqrt{-\Delta}}{\sqrt{-\Delta}} f(u(\tau)) d\tau \right\|_{\dot{H}_x^1 \times L_x^2}. \end{aligned} \quad (17)$$

It remains to show two things:

- (i) Our initial data v_0, v_1 are well-defined, that is, that (16) does indeed converge in $\dot{H}_x^1 \times L_x^2$.
- (ii) The right side of (17) converges to 0 as $t \rightarrow \infty$.

The claim (i) can be shown in several ways, for example, by showing that

$$\lim_{N \rightarrow \infty} \left\| \int_N^\infty \frac{\sin(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) d\tau \right\|_{\dot{H}_x^1} = 0 \quad \text{and} \quad \lim_{N \rightarrow \infty} \left\| \int_N^\infty \cos(-\tau\sqrt{-\Delta}) f(u(\tau)) d\tau \right\|_{L_x^2} = 0,$$

where $N \in \mathbb{N}$. These two equalities follow from the dominated convergence theorem once we show that

$$\int_0^\infty \left\| \frac{\sin(-\tau\sqrt{-\Delta})}{\sqrt{-\Delta}} f(u(\tau)) \right\|_{\dot{H}_x^1}(\tau) d\tau < \infty \quad \text{and} \quad \int_0^\infty \left\| \cos(-\tau\sqrt{-\Delta}) f(u(\tau)) \right\|_{L_x^2}(\tau) d\tau < \infty.$$

But this follows quickly from (12) and the Plancherel theorem.

Claim (ii) has already been established in the discussion of claim (i). This concludes the argument that the finiteness of (12) implies scattering.

Definition 2.3. We say that the pair (q, r) is admissible if $2 \leq q, r \leq \infty$, $(q, r) \neq (2, \infty)$ and

$$\frac{1}{q} + \frac{1}{r} \leq \frac{1}{2}. \quad (18)$$

Theorem 2.4 (Strichartz estimates for wave equation [Strichartz 1977; Kapitanski 1989; Ginibre and Velo 1995; Lindblad and Sogge 1995; Keel and Tao 1998]). *Let I be a time interval and let $u : I \times \mathbb{R}^3 \rightarrow \mathbb{C}$ be a Schwartz solution to the wave equation $\square u = G$ with initial data $u(t_0) = u_0$, $\partial_t u(t_0) = u_1$ for some $t_0 \in I$. Then we have the estimates*

$$\|u\|_{L_t^q L_x^r(I \times \mathbb{R}^3)} + \|u\|_{C_t^0 \dot{H}_x^\sigma(I \times \mathbb{R}^3)} + \|\partial_t u\|_{C_t^0 \dot{H}_x^{\sigma-1}(I \times \mathbb{R}^3)} \lesssim \|u_0\|_{\dot{H}_x^\sigma(\mathbb{R}^3)} + \|u_1\|_{\dot{H}_x^{\sigma-1}(\mathbb{R}^3)} + \|G\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}'}(I \times \mathbb{R}^3)}, \quad (19)$$

where (q, r) and (\tilde{q}, \tilde{r}) are admissible pairs and obey the scaling condition

$$\frac{1}{q} + \frac{3}{r} = \frac{3}{2} - \sigma = \frac{1}{\tilde{q}'} + \frac{3}{\tilde{r}'} - 2, \quad (20)$$

and where \tilde{q}' and \tilde{r}' are conjugate to \tilde{q} and \tilde{r} , respectively. In addition, if u is a spherically symmetric solution, we allow $(q, r) = (2, \infty)$.

We define the Strichartz space $S_\sigma(I)$ for any time interval I , as the closure of the Schwartz function on $I \times \mathbb{R}^3$ under the norm

$$\|u\|_{S_\sigma(I)} := \sup_{(q,r) \text{ admissible}} \|u\|_{L_t^q L_x^r(I \times \mathbb{R}^3)}, \quad (21)$$

where (q, r) satisfies (20).

Morawetz inequality [Morawetz 1968]. *Let I be any time interval and $u : I \times \mathbb{R}^3 \rightarrow \mathbb{C}$ be the solution to (1) with finite energy E . Let F be the potential function as in (2). Then*

$$\int_I \int_{\mathbb{R}^3} \frac{F(u)}{|x|} dx dt \lesssim E. \quad (22)$$

Spherically symmetric solutions. In the last part of this section, we assume that u is the spherically symmetric solution to the log-subcritical wave equations (4), (5) (or log-supercritical wave equation (6)) and F is the corresponding potential function. We obtain the following a priori estimate for the solution.

Lemma 2.5 (pointwise estimate for spherically symmetric solution [Ginibre et al. 1992; Tao 2007]). *Let I be any time interval and let $u : I \times \mathbb{R}^3 \rightarrow \mathbb{C}$ be the spherically symmetric solution to the log-subcritical wave equations (4), (5) (or log-supercritical wave equation (6)) with finite energy E and vanishing at ∞ . Let F be the potential function. Then, for any $t \in I$,*

$$|x|^2 (F(u)^{1/2} |u|)(t, x) \lesssim E \quad (23)$$

Proof. We tackle the log-subcritical case; the proof for the log-supercritical case is similar and easier. Define $\phi(z) := (F(z))^{1/2}z$ and $r := |x|$. From (7), we can compute that, for fixed t ,

$$|\partial_r(\phi(u(t, x)))| \lesssim |u|^3 |\partial_r u|(t, x) \chi_{\{|u| \geq 1/3\}}(x) + |u|^3 (-\log |u|)^{1/2} |\partial_r u|(t, x) \chi_{\{|u| < 1/3\}}(x),$$

where χ is the characteristic function on \mathbb{R}^3 . Then, by the fundamental theorem of calculus, Hölder's inequality, and energy conservation,

$$\begin{aligned} |\phi(u(t, x))| &\lesssim \left| \int_r^\infty \left[|u|^3 |\partial_r u| \chi_{\{|u| \geq 1/3\}} + |u|^3 (-\log |u|)^{1/2} |\partial_r u| \chi_{\{|u| < 1/3\}} \right](t, s) ds \right| \\ &\lesssim \left(\int_r^\infty \frac{|u|^6}{s^2} s^2 \chi_{\{|u| \geq 1/3\}}^2 ds \right)^{1/2} \left(\int_r^\infty \frac{|\partial_r u|^2}{s^2} \chi_{\{|u| \geq 1/3\}} s^2 ds \right)^{1/2} \\ &\quad + \left(\int_r^\infty \frac{|u|^6 (-\log |u|)}{s^2} s^2 \chi_{\{|u| < 1/3\}}^2 ds \right)^{1/2} \left(\int_r^\infty \frac{|\partial_r u|^2}{s^2} \chi_{\{|u| < 1/3\}} s^2 ds \right)^{1/2} \\ &\lesssim \frac{1}{r^2} \left(\int_{\mathbb{R}^3} F(u) dx \right)^{1/2} E^{1/2} \lesssim \frac{1}{r^2} E. \quad \square \end{aligned}$$

Inserting (23) into (22), we obtain that, for any time interval I ,

$$\int_I \int_{\mathbb{R}^3} F^{5/4}(u) |u|^{1/2} dx dt \leq \int_I \int_{\mathbb{R}^3} \frac{F(u)}{|x|} \cdot \sup_{x \in \mathbb{R}^3} (|x| F^{1/4}(u) |u|^{1/2}) dx dt \lesssim E^{3/2}. \quad (24)$$

This implies

$$\int_I \int_{\{|u| \leq 1/3\}} |u|^8 (-\log |u|)^{5/4} dx dt + \int_I \int_{\{|u| > 1/3\}} |u|^8 dx dt \lesssim E^{3/2} \quad (\text{log-subcritical case}) \quad (25)$$

and

$$\int_I \int_{\mathbb{R}^3} |u|^8 \log^{5\alpha/4}(2 + |u|^2) dx dt \lesssim E^{3/2} \quad (\text{log-supercritical case}). \quad (26)$$

3. Log-subcritical wave equations

In this section, we consider the scattering theory for log-subcritical wave equations. We can take advantage of time reversal symmetry, and it suffices to prove that the solution u scatters in $\dot{H}_x^1 \times L_x^2$ as $t \rightarrow \infty$.

Throughout this section, we use the notation

$$A = \{(t, x) \in (0, \infty) \times \mathbb{R}^3 : |u| < \frac{1}{3}\}, \quad B = \{(t, x) \in (0, \infty) \times \mathbb{R}^3 : |u| \geq \frac{1}{3}\},$$

and for any interval I ,

$$A_I = A \cap (I \times \mathbb{R}^3), \quad B_I = B \cap (I \times \mathbb{R}^3). \quad (27)$$

General initial data in log-weighted Sobolev spaces.

Theorem 3.1. *Let $\gamma > \frac{1}{2}$ and let u be the solution to the log-subcritical wave equations (1), (4), and (5) with initial data*

$$u_0(x) \in X_1, \quad u_1(x) \in Y_1, \quad (28)$$

where X_1 and Y_1 are defined by (9). Then u scatters in $\dot{H}_x^1 \times L_x^2$.

Proof. We need some decay estimates for the equation with initial data satisfying (28).

Lemma 3.2. *Let γ and u be as in Theorem 3.1. There exists $T = T(\|u_0\|_{X_1}, \|u_1\|_{Y_1}, \gamma) \gg 1$ such that, for $\tau > T$,*

$$\int_{\mathbb{R}^3} F(u(\tau, x)) dx \lesssim \frac{1}{\log^{2\gamma} \tau}, \quad (29)$$

where $F(z) = F_{\text{sub}}(z)$ is defined by (7).

Proof. We essentially follow the proof of Lemma 2.1 in [Bahouri and Shatah 1998], with some changes. Define

$$e[u](t, x) := \frac{1}{2} |\partial_t u(t, x)|^2 + \frac{1}{2} |\nabla u(t, x)|^2 + F(u(t, x)).$$

We claim that there exists $C_\gamma = C_\gamma(\|u_0\|_{X_1}, \|u_1\|_{Y_1}, \gamma) \gg 1$ such that for $s > C_\gamma$,

$$\int_{|x|>s} e[u](0, x) dx \lesssim \frac{1}{\log^{2\gamma} s}. \quad (30)$$

We prove this claim in the Appendix and continue the proof of this lemma here. Choose T such that $T > \max(C_\gamma^2, \log^{4\gamma} T)$. We aim to show that (29) holds for all $\tau > T$.

Define the truncated forward light cone by

$$K_a^b(c) := \{(t, x) : a \leq t \leq b, |x| \leq t + c, 0 \leq a < b \leq \infty\}$$

and the boundary of the truncated cone by

$$M_a^b(c) := \partial K_a^b(c) = \{(t, x) : a \leq t \leq b, |x| = t + c, 0 \leq a < b \leq \infty\}.$$

Fix $\tau > T$ and let $s = \sqrt{\tau} > C_\gamma$. For any $t_1 > 0$, the energy conservation law on the exterior of the truncated forward light cone $K_0^{t_1}(s)$ implies that

$$\int_{|x|>s+t_1} e[u](t_1) dx + \frac{1}{\sqrt{2}} \text{flux}(0, t_1, s) = \int_{|x|>s} e[u](0) dx \lesssim \frac{1}{\log^{2\gamma} s}, \quad (31)$$

where

$$\text{flux}(a, b, c) := \int_{M_a^b(c)} \left\{ \frac{1}{2} \left| u_t + \frac{x \cdot \nabla u}{|x|} \right|^2 + F(u) \right\} d\sigma.$$

Hence

$$\int_{|x|>s+\tau} F(u(\tau)) dx \leq \int_{|x|>s+\tau} e[u](\tau) dx \lesssim \frac{1}{\log^{2\gamma} s} \lesssim \frac{1}{\log^{2\gamma} \tau}, \quad (32)$$

and it suffices to show that

$$\int_{|x| \leq s+\tau} F(u(\tau)) dx \lesssim \frac{1}{\log^{2\gamma} \tau}. \quad (33)$$

Define $w(t, x) = u(t - s, x)$. The bound (33) is equivalent to

$$\int_{|x| \leq s+\tau} F(w(s + \tau)) dx \lesssim \frac{1}{\log^{2\gamma} \tau}.$$

Set $w_t := \partial_t w$. Multiplying the equation $f(w) - \square w = 0$ by $tw_t + x \cdot \nabla w + w$, we get

$$\partial_t(tQ_0 + w_t w) - \operatorname{div}(tP_0) + R_0 = 0, \quad (34)$$

where

$$\begin{aligned} Q_0 &= e[w] + w_t \left(\frac{x}{t} \cdot \nabla w \right), \\ P_0 &= \frac{x}{t} \left(\frac{w_t^2 - |\nabla w|^2}{2} - F(w) \right) + \nabla w \left(w_t + \frac{x}{t} \cdot \nabla w + \frac{w}{t} \right), \\ R_0 &= |w|^6 g(|w|) - 4F(w), \end{aligned}$$

with g defined by (5). Define the horizontal sections of the forward solid cone by

$$D(t) := \{|x| \in \mathbb{R}^3 : |x| \leq t\}.$$

Fix $0 < T_1 < T_2$ and integrate (34) on $K_{T_1}^{T_2}(0)$. By the divergence theorem, we have

$$\begin{aligned} \int_{D(T_2)} (T_2 Q_0 + w_t w) dx - \int_{D(T_1)} (T_1 Q_0 + w_t w) dx - \frac{1}{\sqrt{2}} \int_{M_{T_1}^{T_2}(0)} \left(t Q_0 + w_t w + t P_0 \frac{x}{|x|} \right) d\sigma + \int_{K_{T_1}^{T_2}(0)} R_0 dx dt \\ =: L_1 + L_2 + L_3 + L_4 = 0. \end{aligned} \quad (35)$$

Now, following the same steps as in [Bahouri and Shatah 1998], we define $v(y) := w(|y|, y)$. Since L_3 is the integral on $M_{T_1}^{T_2}(0)$, using spherical coordinates, we obtain that

$$L_3 = - \int_{T_1}^{T_2} \int_{S^2} r \left(v_r + \frac{v}{r} \right)^2 r^2 dr d\omega + \frac{1}{2} \int_{S^2} T_2^2 v^2(T_2 \omega) d\omega - \frac{1}{2} \int_{S^2} T_1^2 v^2(T_1 \omega) d\omega, \quad (36)$$

$$\begin{aligned} L_1 = \int_{D(T_2)} \left\{ T_2 \left(\frac{|w_t|^2}{2} + \frac{1}{2} \left(w_r + \frac{1}{r} w \right)^2 + \frac{1}{2r^2} |\nabla_\omega w|^2 + F(w) \right) + r \left(w_r + \frac{1}{r} w \right) w_t \right\} dx \\ - \frac{1}{2} \int_{S^2} T_2^2 v^2(T_2 \omega) d\omega, \end{aligned} \quad (37)$$

and

$$\begin{aligned} L_2 = - \int_{D(T_1)} \left\{ T_1 \left(\frac{|w_t|^2}{2} + \frac{1}{2} \left(w_r + \frac{1}{r} w \right)^2 + \frac{1}{2r^2} |\nabla_\omega w|^2 + F(w) \right) + r \left(w_r + \frac{1}{r} w \right) w_t \right\} dx \\ + \frac{1}{2} \int_{S^2} T_1^2 v^2(T_1 \omega) d\omega. \end{aligned} \quad (38)$$

Since $L_4 \geq 0$, plugging (36), (37) and (38) into (35), we deduce that

$$T_2 \int_{D(T_2)} F(w) dx \leq C T_1 E + \int_{T_1}^{T_2} \int_{S^2} T_2 \left(v_r + \frac{v}{r} \right)^2 r^2 dr d\omega,$$

where C is a constant and E is the energy. Therefore,

$$\int_{D(T_2)} F(w(T_2)) dx \leq C \frac{T_1}{T_2} E + \int_{T_1}^{T_2} \int_{S^2} \left(v_r + \frac{v}{r} \right)^2 r^2 dr d\omega. \quad (39)$$

For any $T_1 \geq s$, by (31), the second term in the right-hand side of (39) is controlled by

$$\int_{T_1}^{T_2} \int_{S^2} \left(v_r + \frac{v}{r}\right)^2 r^2 dr d\omega \lesssim \int_{M_{T_1}^{T_2}(0)} \left\{ \frac{1}{2} \left| w_t + \frac{x \cdot \nabla w}{|x|} \right|^2 \right\} d\sigma \lesssim \frac{1}{\log^{2\gamma} s} \lesssim \frac{1}{\log^{2\gamma} \tau}.$$

Now, choosing $T_2 = \tau + s$ and $T_1 = (\tau + s)/\log^{2\gamma} \tau > \sqrt{\tau} = s$, (39) implies

$$\int_{D(\tau+s)} F(w(\tau + s, x)) dx \lesssim \frac{1}{\log^{2\gamma} \tau}. \quad (40)$$

Combining (32) and (40), the lemma is proved. \square

Before we prove Theorem 3.1, let's observe the following fact. Let I be any time interval with length $3 < |I| < \infty$. By Hölder's inequality, we have that, for $0 < \delta < 2$,

$$\begin{aligned} & \| |u|^4 u (-\log |u|) \|_{L_t^1 L_x^2(I \times \mathbb{R}^3)} \\ & \leq \| u^{3-\delta} (-\log |u|)^{(3-\delta)/6} \|_{L_t^\infty L_x^{6/(3-\delta)}(I \times \mathbb{R}^3)} \| u^2 \|_{L_t^{2/(2-\delta)} L_x^{6/\delta}(I \times \mathbb{R}^3)} \| u^\delta (-\log |u|)^{(3+\delta)/6} \|_{L_t^{2/\delta} L_x^\infty(I \times \mathbb{R}^3)} \\ & = \| u (-\log |u|)^{1/6} \|_{L_t^\infty L_x^6(I \times \mathbb{R}^3)}^{3-\delta} \| u \|_{L_t^{4/(2-\delta)} L_x^{12/\delta}(I \times \mathbb{R}^3)}^2 \| u^\delta (-\log |u|)^{(3+\delta)/6} \|_{L_t^{2/\delta} L_x^\infty(I \times \mathbb{R}^3)} \\ & \leq \| u (-\log |u|)^{1/6} \|_{L_t^\infty L_x^6(I \times \mathbb{R}^3)}^{3-\delta} \| u \|_{L_t^{4/(2-\delta)} L_x^{12/\delta}(I \times \mathbb{R}^3)}^2 \| u^\delta (-\log |u|)^{(3+\delta)/6} \|_{L_t^\infty L_x^\infty(I \times \mathbb{R}^3)} |I|^{\delta/2}. \end{aligned}$$

If $|u| \leq \frac{1}{3}$, we can estimate that

$$\| u^\delta (-\log |u|)^{(3+\delta)/6} \|_{L_t^\infty L_x^\infty(I \times \mathbb{R}^3)} \lesssim \left(\frac{1}{\delta} \right)^{1/2+\delta/6}.$$

Letting $\delta = 2/\log |I|$, we obtain

$$\| |u|^4 u (-\log |u|) \|_{L_t^1 L_x^2(I \times \mathbb{R}^3)} \lesssim \| u (-\log |u|)^{1/6} \|_{L_t^\infty L_x^6(I \times \mathbb{R}^3)}^{3-\delta} \| u \|_{L_t^{4/(2-\delta)} L_x^{12/\delta}(I \times \mathbb{R}^3)}^2 \log^{1/2} |I|. \quad (41)$$

To complete the proof of Theorem 3.1, by Remark 2.2, it suffices to show that

$$\| f(u) \|_{L_t^1 L_x^2((T, \infty) \times \mathbb{R}^3)} < \infty \quad \text{for some } T < \infty.$$

Let $J = (3^i, \infty)$, where i is sufficiently large and to be determined later. Then

$$\| f(u) \|_{L_t^1 L_x^2(J \times \mathbb{R}^3)} \lesssim \| |u|^4 u (-\log |u|) \|_{L_t^1 L_x^2(A_J)} + \| |u|^4 u \|_{L_t^1 L_x^2(B_J)} =: M_1 + M_2.$$

Since $(2 + \delta, 6(2 + \delta)/\delta)$ is an admissible pair satisfying (20) for $\sigma = 1$, from Hölder's inequality and Lemma 3.2,

$$M_2 \leq \| u \|_{L_t^\infty L_x^6(B_J)}^{3-\delta} \| u \|_{L_t^{2+\delta} L_x^{6(2+\delta)/\delta}(B_J)}^{2+\delta} \lesssim \frac{1}{(\log(3^i))^{(3-\delta)/3\gamma}} \| u \|_{S_1(J)}^{2+\delta}. \quad (42)$$

On the other hand, define interval J_k by subdividing J according to $J = \bigcup_{k=1}^\infty (3^{2k-1}i, 3^{2k}i) =: \bigcup_{k=1}^\infty J_k$. Define $\delta_k := 2/\log |J_k|$. By (41), Lemma 3.2, and the fact that the admissible pairs $(4/(2 - \delta_k), 12/\delta_k)$

satisfy (20) for $\sigma = 1$, we have

$$\begin{aligned} M_1 &\leq \sum_{k=1}^{\infty} \|u^5(-\log|u|)\|_{L_t^1 L_x^2(J_k \times \mathbb{R}^3)} \lesssim \sum_{k=1}^{\infty} \left(\frac{1}{(\log 3^{2^{k-1}i})^{(3-\delta_k)/3\gamma}} (\log 3^{2^k i})^{1/2} \right) \|u\|_{L_t^{4/(2-\delta_k)} L_x^{12/\delta_k}(J_k \times \mathbb{R}^3)}^2 \\ &\lesssim \left(\sum_{k=1}^{\infty} i^{1/2-(3-\delta_k)/3\gamma} \cdot 2^{(k-1)(1/2-(3-\delta_k)/3\gamma)} \right) \|u\|_{S_1(J)}^2. \end{aligned}$$

Since $\gamma > \frac{1}{2}$, we can choose i sufficiently large such that $((3-\delta_k)/3\gamma - \frac{1}{2}) > c > 0$ for all k . Hence

$$M_1 \lesssim i^{-c} \sum_{k=1}^{\infty} 2^{-(k-1)c} \|u\|_{S_1(J)}^2. \quad (43)$$

Combining (42) and (43), for $\epsilon_0 > 0$ sufficiently small, we can choose i sufficiently large such that

$$\|f(u)\|_{L_t^1 L_x^2(J \times \mathbb{R}^3)} \leq \epsilon_0 (\|u\|_{S_1(J)}^2 + \|u\|_{S_1(J)}^{2+\delta}).$$

By the Strichartz estimate (19), we have

$$\|u\|_{S_1(J)} \leq CE^{1/2} + \epsilon_0 (\|u\|_{S_1(J)}^2 + \|u\|_{S_1(J)}^{2+\delta}).$$

From a continuity argument, we conclude that

$$\|u\|_{S_1(J)} \leq 2CE^{1/2}.$$

This implies that

$$\|f(u)\|_{L_t^1 L_x^2(J \times \mathbb{R}^3)} < \infty. \quad \square$$

Spherically symmetric initial data in lower regularity Sobolev spaces. In this subsection, we consider the solutions to the log-subcritical wave equations with spherically symmetric initial data. If the finite energy initial data are in any lower regularity Sobolev spaces, we obtain the $\dot{H}_x^1 \times L_x^2$ scattering. The spirit of the proof follows from [Tao 2007] and a slightly sharpened bootstrap argument in Lemmas 3.5 and 3.6.

Throughout this subsection, for given $\delta > 0$, we denote

$$Z(t) := \|u(t, x)\|_{\dot{H}_x^{1-\delta}(\mathbb{R}^3)} + \|\partial_t u(t, x)\|_{\dot{H}_x^{-\delta}(\mathbb{R}^3)}. \quad (44)$$

It is easy to show that $Z(t) > 0$ for any time t .²

Theorem 3.3. *Let u be the solution to the log-subcritical wave equations (1), (4), (5) with spherically symmetric initial data*

$$u_0(x) \in X_2, \quad u_1(x) \in Y_2, \quad (45)$$

where X_2 and Y_2 are defined by (10). Then u scatters in $\dot{H}_x^1 \times L_x^2$.

To prove Theorem 3.3, we need some intermediate lemmas.

²If $Z(t_0) = 0$ for some t_0 , it is easy to prove that the solution u has energy $E(t_0) = 0$ and, hence, $E(t) = 0$ for any time t , by energy conservation. This implies the solution $u(t, x) \equiv 0$ for all t .

Lemma 3.4. *Let $I = [a, b]$ be any interval where $0 \leq a < b \leq \infty$ and let u be the solution to the log-subcritical wave equations (1), (4), (5) with spherically symmetric initial data*

$$u(a, x) = u_0(x) \in \dot{H}_x^1 \cap \dot{H}_x^{1-\delta}, \quad \partial_t u(a, x) = u_1(x) \in L_x^2 \cap \dot{H}_x^{-\delta}$$

for some fixed $0 < \delta < \frac{1}{2}$. Then there exists $0 < \epsilon(\delta) \ll 1$ such that for $0 < \epsilon < \epsilon(\delta)$,

$$\|u\|_{S_{1-\delta}(I)} \lesssim Z(a) + (\|u\|_{S_{1-\delta}(I)}^{1+\epsilon/(2\delta)} + \|u\|_{S_{1-\delta}(I)}) \left(\|u\|_{L_{t,x}^8(A_I)}^{5/32} \left\| |u|(-\log|u|)^{5/32} \right\|_{L_{t,x}^8(A_I)}^{4-\epsilon/(2\delta)-\epsilon} + \|u\|_{L_{t,x}^8(B_I)}^4 \right) \left(\frac{1}{\epsilon} \right)^{7/16}, \quad (46)$$

where the constant hidden in (46) is independent of the interval I and ϵ .

Proof. By the Strichartz estimate (19),

$$\|u\|_{S_{1-\delta}(I)} \lesssim Z(a) + \|f(u)\|_{L_t^{2/(2-\delta)} L_x^{2/(1+\delta)}(I \times \mathbb{R}^3)}. \quad (47)$$

Consider that

$$\|f(u)\|_{L_t^{2/(2-\delta)} L_x^{2/(1+\delta)}(I \times \mathbb{R}^3)} \lesssim \| -|u|^4 u(\log(|u|)) \|_{L_t^{2/(2-\delta)} L_x^{2/(1+\delta)}(A_I)} + \| |u|^4 u \|_{L_t^{2/(2-\delta)} L_x^{2/(1+\delta)}(B_I)} =: N_1 + N_2$$

with A_I and B_I as in (27). By Hölder's inequality,

$$N_2 \leq \|u\|_{L_t^{2/(1-\delta)} L_x^{2/\delta}(B_I)} \|u\|_{L_{t,x}^8(B_I)}^4 \leq \|u\|_{S_{1-\delta}(I)} \|u\|_{L_{t,x}^8(B_I)}^4. \quad (48)$$

On the other hand, choosing $\epsilon(\delta)$ sufficiently small such that for $0 < \epsilon < \epsilon(\delta)$,

$$0 < \frac{1}{p} := \frac{8\delta + \epsilon - 8\delta^2 + 2\epsilon\delta}{8(2\delta + \epsilon)} \leq \frac{1}{2}, \quad 0 < \frac{1}{q} := \frac{\delta}{2} + \frac{\epsilon(1-2\delta)}{8(2\delta + \epsilon)} \leq \frac{1}{2}, \quad \frac{3}{8} \approx \frac{12 + 5\epsilon/(2\delta) + 5\epsilon}{32} < \frac{7}{16}.$$

It is clear that (p, q) is an admissible pair satisfying (20) for $\sigma = 1 - \delta$. By Hölder's inequality and interpolation theory, we can estimate that

$$\begin{aligned} N_1 &\leq \| |u|^{5-\epsilon} (-\log|u|)^{5(4-\epsilon/(2\delta)-\epsilon)/32} \|_{L_t^{2/(2-\delta)} L_x^{2/(1+\delta)}(A_I)} \| |u|^\epsilon (-\log|u|)^{(12+5\epsilon/(2\delta)+5\epsilon)/32} \|_{L_{t,x}^\infty(A_I)} \\ &\leq \|u\|_{L_t^p L_x^q(A_I)}^{1+\epsilon/(2\delta)} \|u(-\log|u|)^{5/32}\|_{L_{t,x}^8(A_I)}^{4-\epsilon/(2\delta)-\epsilon} \| |u|^\epsilon (-\log|u|)^{(12+5\epsilon/(2\delta)+5\epsilon)/32} \|_{L_{t,x}^\infty(A_I)} \end{aligned} \quad (49)$$

$$\lesssim \|u\|_{L_t^p L_x^q(A_I)}^{1+\epsilon/(2\delta)} \| |u|(-\log|u|)^{5/32}\|_{L_{t,x}^8(A_I)}^{4-\epsilon/(2\delta)-\epsilon} \left(\frac{1}{\epsilon} \right)^{upnfrac{12+5\epsilon/(2\delta)+5\epsilon}{32}}. \quad (50)$$

The last factor of (50) comes from maximizing the last factor on the right of (49) using calculus. We note that the constant hidden in the last inequality is independent of ϵ . By (48) and (50), we have

$$\|f(u)\|_{L_t^{2/(2-\delta)} L_x^{2/(1+\delta)}(I \times \mathbb{R}^3)} \lesssim \|u\|_{S_{1-\delta}(I)}^{1+\epsilon/(2\delta)} \| |u|(-\log|u|)^{5/32}\|_{L_{t,x}^8(A_I)}^{4-\epsilon/(2\delta)-\epsilon} \left(\frac{1}{\epsilon} \right)^{7/16} + \|u\|_{S_{1-\delta}(I)} \|u\|_{L_{t,x}^8(B_I)}^4.$$

From (47),

$$\|u\|_{S_{1-\delta}(I)} \lesssim Z(a) + \|u\|_{S_{1-\delta}(I)}^{1+\epsilon/(2\delta)} \| |u|(-\log|u|)^{5/32}\|_{L_{t,x}^8(A_I)}^{4-\epsilon/(2\delta)-\epsilon} \left(\frac{1}{\epsilon} \right)^{7/16} + \|u\|_{S_{1-\delta}(I)} \|u\|_{L_{t,x}^8(B_I)}^4 \lesssim \text{RHS of (46)}.$$

One can check that all constants hidden in the inequalities above are independent of the interval I and ϵ . Hence, Lemma 3.4 is proved. \square

Lemma 3.5 (continuity argument). *Let $I (= [a, b])$ and u satisfy the assumptions of Lemma 3.4, C be the constant hidden in (46) and $0 < \epsilon(\delta)$ be chosen in Lemma 3.4. Let $\epsilon_0 = 1/(100C)$ and $0 < \epsilon < \epsilon(\delta)$ such that $Z(a)^{\epsilon/(2\delta)} \geq \frac{1}{2}$ and $(2C)^{\epsilon/(2\delta)} \leq 2$. We define*

$$Q(I) := \left(\| |u| (-\log |u|)^{5/32} \|_{L_{t,x}^8(A_I)}^{4-\epsilon/(2\delta)-\epsilon} + \| u \|_{L_{t,x}^8(B_I)}^{4-\epsilon/(2\delta)-\epsilon} \right).$$

If $\| u \|_{L_{t,x}^8(B_I)} \leq 1$ and $Q(I) \leq \epsilon_0(\epsilon^{7/16}/(Z(a)^{\epsilon/(2\delta)}))$, we have

$$\| u \|_{S_{1-\delta}(I)} \leq 2CZ(a).$$

Proof. We prove this lemma by contradiction. For $0 \leq t \leq b - a$, from the dominated convergence theorem, we have that the function $\Phi(t) := \| u \|_{S_{1-\delta}([a, a+t])}$ is nondecreasing and continuous in $[0, b - a]$ and $\Phi(0) = 0$. By the hypothesis and (46), we have

$$\Phi(t) \leq CZ(a) + \frac{1}{100}(\Phi(t))^{1+\epsilon/(2\delta)} + \Phi(t) \left(\frac{1}{Z(a)^{\epsilon/(2\delta)}} \right) \quad (51)$$

for all $t \in [0, b - a]$. Assume for contradiction that there exists $t_0 \in [0, b - a]$ such that $\Phi(t_0) = 2CZ(a)$. If $2CZ(a) < 1$, (51) implies that

$$2CZ(a) = \Phi(t_0) \leq CZ(a) + \frac{1}{50}(2CZ(a)) \left(\frac{1}{Z(a)^{\epsilon/(2\delta)}} \right) \leq \frac{11}{10}CZ(a).$$

On the other hand, if $2CZ(a) \geq 1$, (51) implies that

$$2CZ(a) = \Phi(t_0) \leq CZ(a) + \frac{1}{50}(2CZ(a))^{1+\epsilon/(2\delta)} \left(\frac{1}{Z(a)^{\epsilon/(2\delta)}} \right) \leq \frac{11}{10}CZ(a).$$

We get contradictions in both situations, and the lemma is proved. \square

Lemma 3.6 (finite division). *Let $I (= [a, b])$ and u satisfy the assumptions of Lemma 3.4 and C be the constant hidden in (46). We denote $Z_i = (2C)^i Z(a)$, where $i = 0, 1, 2, \dots$. For any $\epsilon_0 > 0$, we can choose $\epsilon \ll 1$ and finitely many numbers $a = T_0 < T_1 < T_2 < \dots < T_N < T_{N+1} = b$, where $N = N(\epsilon_0, \epsilon, \delta, E, Z_0, C)$, such that for $I_j := [T_j, T_{j+1}]$,*

$$Q(I_j) = \epsilon_0 \left(\frac{\epsilon^{7/16}}{Z_j^{\epsilon/(2\delta)}} \right) \quad (52)$$

for $0 \leq j \leq N - 1$ and $Q(I_N) \leq \epsilon_0(\epsilon^{7/16}/Z_N^{\epsilon/(2\delta)})$.

Proof. We observe that

$$\sum_{i=0}^{\infty} \left[\epsilon_0 \left(\frac{\epsilon^{7/16}}{Z_i^{\epsilon/(2\delta)}} \right) \right]^{8/(4-\epsilon/(2\delta)-\epsilon)} \gtrsim_{\epsilon_0, Z_0} \left\{ \epsilon^{7/(8-\epsilon/(\delta)-2\epsilon)} \sum_{i=0}^{\infty} \frac{1}{(2C)^{8i\epsilon/(8\delta-\epsilon-2\delta\epsilon)}} \right\} \rightarrow \infty \quad \text{as } \epsilon \rightarrow 0.$$

Therefore, by (25), we can choose ϵ sufficiently small such that

$$3 \left(\int_A \int |u|^8 (-\log |u|)^{5/4} dx dt + \int_B \int |u|^8 dx dt \right) < \sum_{i=0}^K \left[\epsilon_0 \left(\frac{\epsilon^{7/16}}{Z_i^{\epsilon/(2\delta)}} \right) \right]^{8/(4-\epsilon/(2\delta)-\epsilon)} \quad (53)$$

for some $K = K(\epsilon_0, \epsilon, \delta, E, Z_0, C)$.

Fix this ϵ . If $Q(I) < \epsilon_0(\epsilon^{7/16}/(Z_0^{\epsilon/(2\delta)}))$, we say $T_1 = b$ and the lemma is proved. Otherwise, we can choose $0 < T_1 < b$ such that (52) holds for $j = 0$. Again, if $Q([T_1, b]) < \epsilon_0(\epsilon^{7/16}/(Z_1^{\epsilon/(2\delta)}))$, we say $T_2 = b$. Otherwise, we can choose $T_1 < T_2 < b$ such that (52) holds for $j = 1$. By continuing this process, we can choose $a < T_1 < T_2 < \dots$ such that (52) holds for $j = 0, 1, \dots$. It suffices to show that this process will stop in at most $K + 1$ steps. Indeed, assume that there are more than $K + 1$ subintervals satisfying (52). Since

$$Q(I_j)^{8/(4-\epsilon/(2\delta)-\epsilon)} \leq 3 \left(\iint_{A_{I_j}} |u|^8 (-\log |u|)^{5/4} dx dt + \iint_{B_{I_j}} |u|^8 dx dt \right),$$

for $j = 0, 1, \dots$, by our construction of I_j , we have

$$\begin{aligned} \sum_{j=0}^{K+1} \epsilon_0 \left(\frac{\epsilon^{7/16}}{Z_j^{\epsilon/(2\delta)}} \right) &= \sum_{j=0}^{K+1} Q(I_j)^{8/(4-\epsilon/(2\delta)-\epsilon)} \\ &\leq \sum_{i=0}^{K+1} 3 \left(\iint_{A_{I_j}} |u|^8 (-\log |u|)^{5/4} dx dt + \iint_{B_{I_j}} |u|^8 dx dt \right) \\ &\leq 3 \left(\iint_A |u|^8 (-\log |u|)^{5/4} dx dt + \iint_B |u|^8 dx dt \right). \end{aligned}$$

This contradicts (53), and the lemma is proved. \square

Corollary 3.7. *Let I and u satisfy the assumptions of Lemma 3.4 and C be the constant hidden in (46). If $\|u\|_{L_{t,x}^8(B_I)} \leq 1$, $u \in L_{t,x}^{8/(1+2\delta)}(I \times \mathbb{R}^3)$.*

Proof. Let $\epsilon(\delta)$ be chosen in Lemma 3.4 and $0 < \epsilon < \epsilon(\delta)$ satisfy Lemma 3.6, $Z(a)^{\epsilon/(2\delta)} \geq \frac{1}{2}$ and $(2C)^{\epsilon/(2\delta)} \leq 2$. Let $\{I_j\}_{j=0}^N$ be the subintervals constructed by Lemma 3.6 such that (52) holds for $0 \leq j \leq N$.

We claim that

$$\|u\|_{S_{1-\delta}(I_j)} \leq 2C Z_j \quad \text{for } 0 \leq j \leq N, \quad (54)$$

where $Z_j = (2C)^j Z(a)$. Indeed, by Lemma 3.5, (54) holds for $j = 0$. Again, if (54) holds for $j = k - 1$, we have $Z(T_k) \leq \|u\|_{S_{1-\delta}(I_{k-1})} \leq Z_k$. Since $Z_k^{\epsilon/(2\delta)} \geq Z(a)^{\epsilon/(2\delta)} \geq \frac{1}{2}$, applying Lemma 3.5 on the interval I_k , we obtain (54) for $j = k$. By induction on j , the claim is proved and this implies

$$\|u\|_{L_{t,x}^{8/(1+2\delta)}(I \times \mathbb{R}^3)} \leq \sum_{j=0}^{N+1} \|u\|_{S_{1-\delta}(I_j)} \leq \sum_{j=0}^{N+1} (2C)^j Z_0 < \infty. \quad \square$$

Corollary 3.8. *Let u be the solution to the log-subcritical wave equations (1), (4), (5) with spherically symmetric initial data*

$$u(0, x) = u_0(x) \in \dot{H}_x^1 \cap \dot{H}_x^{1-\delta}, \quad \partial_t u(0, x) = u_1(x) \in L_x^2 \cap \dot{H}_x^{-\delta}$$

for some fixed $0 < \delta < \frac{1}{2}$. Then $u \in L_{t,x}^{8/(1+2\delta)}(\mathbb{R}_+ \times \mathbb{R}^3)$.

Proof. By (25), we can choose finitely many numbers $0 = S_0 < S_1 < \dots < S_{M-1} < S_M = \infty$ such that $\|u\|_{L_{t,x}^8(B_{[S_k, S_{k+1}]})} \leq 1$ for $0 \leq k \leq M$. By Corollary 3.7 and energy conservation, we have

$$(u(S_k, x), \partial_t u(S_k, x)) \in (\dot{H}_x^1 \cap \dot{H}_x^{1-\delta}) \times (L_x^2 \cap \dot{H}_x^{-\delta})$$

and $\|u\|_{L_{t,x}^{8/(1+2\delta)}([S_k, S_{k+1}] \times \mathbb{R}^3)} < \infty$ for $0 \leq k \leq M$. Hence

$$\|u\|_{L_{t,x}^{8/(1+2\delta)}(\mathbb{R}_+ \times \mathbb{R}^3)} \leq \sum_{k=0}^M \|u\|_{L_{t,x}^{8/(1+2\delta)}([S_k, S_{k+1}] \times \mathbb{R}^3)} < \infty. \quad \square$$

To finish the proof of Theorem 3.3, by Remark 2.2, it suffices to show that $\|f(u)\|_{L_t^1 L_x^2((T, \infty) \times \mathbb{R}^3)} < \infty$ for some $0 < T < \infty$. Since the initial data satisfy (45), we can choose some $0 < \delta < \frac{1}{2}$ such that $u_0 \in \dot{H}_x^1(\mathbb{R}^3) \cap \dot{H}_x^{1-\delta}(\mathbb{R}^3)$ and $u_1 \in L_x^2(\mathbb{R}^3) \cap \dot{H}_x^{-\delta}(\mathbb{R}^3)$. Observe that

$$\begin{aligned} \|f(u)\|_{L_t^1 L_x^2((T, \infty) \times \mathbb{R}^3)} &\lesssim \| |u|^5 (\log(|u|)) \|_{L_t^1 L_x^2(A_T)} + \| |u|^5 \|_{L_t^1 L_x^2(B_T)} \\ &\lesssim \|u\|_{L_{t,x}^{8/(1+2\delta)}(A_T)}^{4/(1+2\delta)} \|u\|_{L_t^2 L_x^\infty(A_T)} \|u\|_{L_{t,x}^{8\delta/(1+2\delta)}(A_T)}^{8\delta/(1+2\delta)} (\log(|u|)) \|_{L_{t,x}^\infty(A_T)} + \|u\|_{L_{t,x}^8(B_T)}^4 \|u\|_{L_t^2 L_x^\infty(B_T)} \\ &\lesssim \|u\|_{L_t^2 L_x^\infty((T, \infty) \times \mathbb{R}^3)} \left[\left(\frac{1+2\delta}{8\delta} \right) \|u\|_{L_{t,x}^{8/(1+2\delta)}(A_T)}^{4/(1+2\delta)} + \|u\|_{L_{t,x}^8(B_T)}^4 \right], \end{aligned}$$

where $A_T := A \cap ((T, \infty) \times \mathbb{R}^3)$ and $B_T := B \cap ((T, \infty) \times \mathbb{R}^3)$. The last inequality above is from the fact that $|u|^{8\delta/(1+2\delta)} (\log(|u|)) \lesssim (1+2\delta)/(8\delta)$ for $|u| \leq \frac{1}{3}$. By Corollary 3.8 and (25), for sufficiently small $\epsilon > 0$, we can choose $T = T(\epsilon)$ sufficiently large such that

$$\left(\frac{1+2\delta}{8\delta} \right) \|u\|_{L_{t,x}^{8/(1+2\delta)}(A_T)}^{4/(1+2\delta)} + \|u\|_{L_{t,x}^8(B_T)}^4 < \epsilon.$$

Hence, by the Strichartz inequality [Klainerman and Machedon 1993],

$$\|u\|_{L_t^2 L_x^\infty((T, \infty) \times \mathbb{R}^3)} \leq CE^{1/2} + \epsilon C \|u\|_{L_t^2 L_x^\infty((T, \infty) \times \mathbb{R}^3)}.$$

Again for $\epsilon < 1/(2C)$, we have $\|u\|_{L_t^2 L_x^\infty((T, \infty) \times \mathbb{R}^3)} < 2CE^{1/2}$ and this implies $\|f(u)\|_{L_t^1 L_x^2((T, \infty) \times \mathbb{R}^3)} < \infty$.

Energy subcritical nonlinear wave equations with specific spherically symmetric initial data. In the last part of this section, we will discuss an observation, for energy subcritical nonlinear wave equations, inspired by the proof of Theorem 3.3. For given $0 < \delta < \frac{1}{2}$, let $(u_0, u_1) \in (\dot{H}_x^1(\mathbb{R}^3) \cap \dot{H}_x^{1-\delta}(\mathbb{R}^3)) \times (L_x^2(\mathbb{R}^3) \cap \dot{H}_x^{-\delta}(\mathbb{R}^3))$ be spherically symmetric functions. In this subsection, we consider the energy-subcritical nonlinear wave equation

$$\square u = |u|^{4-\epsilon} u, \quad u(0, x) = u_0(x), \quad \partial_t u(0, x) = u_1(x), \quad (55)$$

where we allow ϵ to depend on the given data (u_0, u_1) . That is, we find a relation (R) (see Definition 3.10) among ϵ , the energy E , and $Z(0)$ as in (44), the lower regularity norm of the initial data, for which the solution scatters. We remark that relation (R) holds for data large in both the energy and $\dot{H}_x^{1-\delta}$ norms provided that ϵ is taken sufficiently small (depending on the size of these norms). In [Lindblad and Sogge

1995], scattering was established in $\dot{H}^{1-\delta}$ for the $\dot{H}^{1-\delta}$ critical nonlinear wave equation from small data. Our remarks here are related to that work, for example, relation (R) quantifies the extent to which large data can be allowed. Also, we will prove scattering in \dot{H}^1 , rather than $\dot{H}^{1-\delta}$.

In order to prove that u scatters in $\dot{H}_x^1 \times L_x^2$, It suffices to show that $\|u^{5-\epsilon}\|_{L_t^1 L_x^2((T,\infty)\times\mathbb{R}^3)} < \infty$ for some $T < \infty$. By the Strichartz estimate and Hölder's inequality,

$$\begin{aligned} \|u\|_{L_t^2 L_x^\infty((T,\infty)\times\mathbb{R}^3)} &\leq CE^{1/2} + C\|u^{5-\epsilon}\|_{L_t^1 L_x^2((T,\infty)\times\mathbb{R}^3)} \\ &\leq CE^{1/2} + C\|u\|_{L_t^2 L_x^\infty((T,\infty)\times\mathbb{R}^3)} \|u\|_{L_{t,x}^{8-2\epsilon}((T,\infty)\times\mathbb{R}^3)}^{4-\epsilon}. \end{aligned}$$

Following similar arguments as in the proof of Theorem 3.3, we only need to show that

$$\|u\|_{L_{t,x}^{8-2\epsilon}((T,\infty)\times\mathbb{R}^3)} < \infty \quad \text{for some } T < \infty.$$

Let $\epsilon_0(\delta) := 8\delta/(1+2\delta)$ (so that $\dot{H}^{1-\delta}$ is the scale invariant norm for (55) with $\epsilon = \epsilon_0(\delta)$). We restrict to the case $0 < \epsilon < \epsilon_0(\delta)$.

In this case, (55) is $\dot{H}^{1-\delta}$ -supercritical nonlinear wave equation. We denote

$$\gamma_\epsilon = \frac{3\epsilon}{16\delta - \frac{5}{2}\delta\epsilon - \frac{5}{4}\epsilon}, \quad \kappa_\epsilon = \frac{8 - \frac{5}{4}\epsilon}{4 - \gamma_\epsilon - \epsilon}, \quad \frac{1}{\alpha_\epsilon} = \frac{1+2\delta}{8} + \frac{3(1-2\delta)}{8(1+\gamma_\epsilon)}, \quad \frac{1}{\beta_\epsilon} = \frac{1+2\delta}{8} - \frac{1-2\delta}{8(1+\gamma_\epsilon)}.$$

Note that

- (i) as $\epsilon \rightarrow \epsilon_0(\delta)$, $\gamma_\epsilon \rightarrow 4 - \epsilon$ and $\kappa_\epsilon \rightarrow \infty$;
- (ii) $(\alpha_\epsilon, \beta_\epsilon)$ is an admissible pair satisfying (20) for $\sigma = 1 - \delta$.

Remark 3.9. Let u be the spherically symmetric solution to the energy-subcritical nonlinear wave equation (55) with energy E . We observe that Lemma 2.5 holds for u . Hence, for any interval $I = [a, b]$ where $0 \leq a < b \leq \infty$, (24) implies

$$\int_I \int_{\mathbb{R}^3} |u(t, x)|^{8-5\epsilon/4} dx dt \leq C_1 E^{3/2}, \quad (56)$$

where we can choose the constant C_1 to be independent of ϵ . Moreover, by the Strichartz estimate,

$$\|u\|_{S_{1-\delta}(I)} \leq CZ(a) + C\|u^{5-\epsilon}\|_{L_t^2/(2-\delta)L_x^{2/(1+\delta)}(I\times\mathbb{R}^3)}. \quad (57)$$

Definition 3.10. Given $0 < \delta < \frac{1}{2}$, let $0 < \epsilon < \epsilon_0(\delta)$, u be the solution to (55) with energy E and lower regularity norm $Z(0) > 0$. We say that the triple $(E, Z(0), \epsilon)$ satisfies the relation (R) if

$$C_1 E^{3/2} \leq \left(\frac{1}{2(2C)^{1+\gamma_\epsilon} Z(0)^{\gamma_\epsilon}} \right)^{\kappa_\epsilon} \frac{1}{1 - (2C)^{-\gamma_\epsilon \kappa_\epsilon}}$$

Lemma 3.11. Given $0 < \delta < \frac{1}{2}$ and $0 < \epsilon < \epsilon_0(\delta)$, let

$$(u_0, u_1) \in (\dot{H}_x^1(\mathbb{R}^3) \cap \dot{H}_x^{1-\delta}(\mathbb{R}^3)) \times (L_x^2(\mathbb{R}^3) \cap \dot{H}_x^{-\delta}(\mathbb{R}^3))$$

be spherically symmetric functions such that the triple $(E, Z(0), \epsilon)$ satisfies (R) and u is the solution to (55). Then $u \in L_{t,x}^{8/(1+2\delta)}(\mathbb{R}_+ \times \mathbb{R}^3)$.

Proof. Since $(E, Z(0), \epsilon)$ satisfies (R), by (56) and an argument similar to that in proof of Lemma 3.6, we can choose finitely many numbers $0 = T_0 < T_1 < \dots < T_N < T_{N+1} = \infty$ such that

$$\|u\|_{L_{t,x}^{8-5\epsilon/4}([T_i, T_{i+1}] \times \mathbb{R}^3)}^{4-\gamma_\epsilon-\epsilon} = \frac{1}{2(2C)^{1+\gamma_\epsilon} ((2C)^i Z(0))^{\gamma_\epsilon}} \quad (58)$$

for $0 \leq i \leq N-1$ and

$$\|u\|_{L_{t,x}^{8-5\epsilon/4}([T_N, T_{N+1}] \times \mathbb{R}^3)}^{4-\gamma_\epsilon-\epsilon} \leq \frac{1}{2(2C)^{1+\gamma_\epsilon} ((2C)^N Z(0))^{\gamma_\epsilon}}.$$

We claim that

$$Z(T_i) < (2C)^i Z(0) \quad (59)$$

and

$$\|u\|_{S_{1-\delta}([T_i, T_{i+1}])} < (2C)^{i+1} Z(0) \quad (60)$$

for $0 \leq i \leq N$.

Observe that (59) is clearly true for $i = 0$ and $Z(T_i) \leq \|u\|_{S_{1-\delta}([T_{i-1}, T_i])}$ for $1 \leq i \leq N$. Hence it suffices to show that (60) holds and then (59) is automatically true.

A similar proof to that of Lemma 3.5 applies here. Assume (60) is true for $i \leq j-1$. We aim to prove (60) for $i = j$. (Note that (59) follows from our assumption when $i = j$.) Let $\phi(t) = \|u\|_{S_{1-\delta}([T_j, T_j+t])}$. Then ϕ is a continuous and nondecreasing function on $[0, T_{j+1} - T_j]$ and $\phi(0) = 0$. Assume for contradiction that there exists $t_0 \in [0, T_{j+1} - T_j]$ such that $\phi(t_0) = (2C)^{j+1} Z(0)$. By Hölder's inequality, (57), (58), and (59), we have

$$\begin{aligned} (2C)^{j+1} Z(0) &= \phi(t_0) \leq CZ(T_j) + C\|u^{5-\epsilon}\|_{L_t^{2/(2-\delta)}L_x^{2/(1+\delta)}([T_j, T_j+t_0] \times \mathbb{R}^3)} \\ &\leq CZ(T_j) + C\|u\|_{L_t^\alpha L_x^{\beta\epsilon}([T_j, T_j+t_0] \times \mathbb{R}^3)}^{1+\gamma_\epsilon} \|u\|_{L_{t,x}^{8-5\epsilon/4}([T_j, T_j+t_0] \times \mathbb{R}^3)}^{4-\gamma_\epsilon-\epsilon} \\ &\leq CZ(T_j) + C\|u\|_{S_{1-\delta}([T_j, T_j+t_0])}^{1+\gamma_\epsilon} \|u\|_{L_{t,x}^{8-5\epsilon/4}([T_j, T_j+t_0] \times \mathbb{R}^3)}^{4-\gamma_\epsilon-\epsilon} \\ &\leq C(2C)^j Z(0) + \frac{1}{4[(2C)^{j+1} Z(0)]^{\gamma_\epsilon}} \|u\|_{S_{1-\delta}([T_j, T_j+t_0])}^{1+\gamma_\epsilon} \\ &< \frac{1}{2}(2C)^{j+1} Z(0) + \frac{1}{4[(2C)^{j+1} Z(0)]^{\gamma_\epsilon}} \times [(2C)^{j+1} Z(0)]^{1+\gamma_\epsilon} \\ &= \frac{3}{4}(2C)^{j+1} Z(0). \end{aligned}$$

The contradiction implies that (60) holds for $i = j$. By an inductive argument on i , the claim is proved. To finish proving this lemma, we have

$$\|u\|_{L_{t,x}^{8/(1+2\delta)}(\mathbb{R}_+ \times \mathbb{R}^3)} \leq \sum_{i=0}^{N+1} \|u\|_{S_{1-\delta}([T_i, T_{i+1}])} \leq \sum_{i=0}^{N+1} (2C)^i Z(0) < \infty \quad \square$$

Corollary 3.12. *Let $\delta, \epsilon, u_0, u_1$ and u satisfy the assumptions of Lemma 3.11. Then u scatters in $\dot{H}_x^1 \times L_x^2$.*

Proof. By the above discussion, it suffices to show

$$\|u\|_{L_{t,x}^{8-2\epsilon}([T, \infty) \times \mathbb{R}^3)} < \infty$$

for some $T < \infty$, since $0 < \epsilon < \epsilon_0(\delta)$ is equivalent to $8/(1+2\delta) < 8-2\epsilon$. The proof of $L_{t,x}^{8-2\epsilon}$ spacetime bound is straightforward by (56), Lemma 3.11, and interpolation theory. \square

4. Log-supercritical wave equation

For spherically symmetric log-supercritical nonlinear wave equation (1), (6) with finite energy E , we observe that the potential energy bound provides slightly better pointwise control, (26), of the solution than the one from the kinetic energy bound³; see [Ginibre et al. 1992; Tao 2007]. In this section, we consider a slightly more supercritical wave equation than the equation in [Tao 2007] and prove the same global regularity result by using (26).

Theorem 4.1. *Define*

$$\tilde{H}_x^2(\mathbb{R}^3) := \dot{H}_x^1(\mathbb{R}^3) \cap \dot{H}_x^2(\mathbb{R}^3).$$

Let $0 < \alpha \leq \frac{4}{3}$ and (u_0, u_1) be smooth, compactly supported, and spherically symmetric initial data with energy E . Then there exists a global smooth solution to

$$\square u = |u|^4 u \log^\alpha(2 + |u|^2), \quad u(0, x) = u_0(x), \quad \partial_t u(0, x) = u_1(x). \quad (61)$$

Furthermore, we have the universal bound of $\tilde{H}_x^2 \times H_x^1$ norm, which depends on both the energy E and $\tilde{H}_x^2 \times H_x^1$ norm of the initial data, of the solution u ; this implies that the solution u scatters in $\tilde{H}_x^2(\mathbb{R}^3) \times H_x^1(\mathbb{R}^3)$.⁴

Remark 4.2. This theorem was proved in [Tao 2007] for $\alpha = 1$, and it is easy to get the same result for $\alpha < 1$ from that argument. We take advantage of (26) to extend the range of α up to $\frac{4}{3}$. In the remainder of this section, we will essentially follow Tao's argument to prove Theorem 4.1 using (26) and sketch the proof of $\tilde{H}_x^2 \times H_x^1$ scattering. We will skip the argument providing an explicit $\tilde{H}_x^2 \times H_x^1$ universal bound here; see [Tao 2007] for details.

We will use a well-known global continuation result (for a proof see [Sogge 1995], for example).

Theorem 4.3 (classical existence theory). *Let $u : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{C}$ be a classical solution⁵ to (61) satisfying*

$$\|u\|_{L_t^\infty L_x^\infty([0, T] \times \mathbb{R}^3)} < \infty.$$

Then there is $\delta > 0$ such that one can extend the solution u to $[0, T + \delta] \times \mathbb{R}^3$.

Proof of Theorem 4.1. By time reversal symmetry, it suffices to consider the global existence and scattering theory of u on $\mathbb{R}_+ \times \mathbb{R}^3$.

³The kinetic energy bound can only provide $\int_I \int_{\mathbb{R}^3} |u|^8 \log^\alpha(2 + |u|^2) dx dt \lesssim E^{3/2}$.

⁴The definition of $\tilde{H}_x^2 \times H_x^1$ scattering for the solution u is similar to Definition 2.1, but the $\dot{H}_x^1 \times L_x^2$ -norm is replaced by the $\tilde{H}_x^2 \times H_x^1$ -norm.

⁵We call u a classical solution to (1) if u solves (1) and is smooth and compactly supported for each time.

By the Sobolev embedding theorem, for a classical solution u to (61) on $[0, T] \times \mathbb{R}^3$, we have

$$\|u\|_{L_t^\infty L_x^\infty([0, T] \times \mathbb{R}^3)} \lesssim \sum_{j=1}^2 \|\nabla_x^j u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)}. \quad (62)$$

Hence, applying classical existence theory (Theorem 4.3), in order to show global existence, it suffices to prove that for any fixed $0 < T \leq \infty$, we have

$$\sum_{j=1}^2 \|\nabla_x^j u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} < \infty,$$

provided that u is the classical solution to (61) on $[0, T] \times \mathbb{R}^3$.

Let $I = [a, b] \subseteq [0, T]$ be any interval. We define

$$\begin{aligned} M_I &:= \int_I \int_{\mathbb{R}^3} |u(t, x)|^8 \log^{5\alpha/4}(2 + |u(t, x)|^2) dx dt, \\ N_I &:= \sum_{j=0}^1 \|\nabla_x^j u\|_{L_t^2 L_x^\infty(I \times \mathbb{R}^3)} + \|\nabla_{t,x} \nabla_x^j u\|_{L_t^\infty L_x^2(I \times \mathbb{R}^3)}, \\ D_I &:= \|\nabla_{t,x} u(a)\|_{H_x^1(\mathbb{R}^3)}. \end{aligned}$$

In addition, we set $D = \|\nabla_{t,x} u(0)\|_{H_x^1(\mathbb{R}^3)}$.

From the Strichartz inequality, Hölder's inequality, and (62), we have

$$\begin{aligned} N_I &\leq C \|\nabla_{t,x} u(a)\|_{H_x^1(\mathbb{R}^3)} + C \sum_{j=0}^1 \|\nabla_x^j (|u|^4 u \log^\alpha(2 + |u|^2))\|_{L_t^1 L_x^2(I \times \mathbb{R}^3)} \\ &\leq C D_I + C \sum_{j=0}^1 \| |u|^4 |\nabla_x^j u| \log(2 + |u|^2) \|_{L_t^1 L_x^2(I \times \mathbb{R}^3)} \\ &\leq C D_I + C \| |u|^4 \log^{5\alpha/8}(2 + |u|^2) \|_{L_t^2 L_x^2(I \times \mathbb{R}^3)} \left(\sum_{j=0}^1 \|\nabla_x^j u\|_{L_t^2 L_x^\infty(I \times \mathbb{R}^3)} \|\log^{3\alpha/8}(2 + |u|^2)\|_{L_t^\infty L_x^\infty(I \times \mathbb{R}^3)} \right) \\ &\leq C D_I + C \|u \log^{5\alpha/32}(2 + |u|^2)\|_{L_t^8 L_x^8(I \times \mathbb{R}^3)}^4 \left(\sum_{j=0}^1 \|\nabla_x^j u\|_{L_t^2 L_x^\infty(I \times \mathbb{R}^3)} \|\log(2 + |u|^2)\|_{L_t^\infty L_x^\infty(I \times \mathbb{R}^3)}^{3\alpha/8} \right) \\ &\leq C D_I + C M_I^{1/2} N_I \log^{3\alpha/8}(2 + \|u\|_{L_t^\infty L_x^2(I \times \mathbb{R}^3)}^2) \\ &\leq C D_I + C M_I^{1/2} N_I \log^{1/2}(2 + N_I^2). \end{aligned}$$

From the result in [Tao 2007, Corollary 3.2], for any $\epsilon_0 > 0$,

$$\sum_{i=0}^k \frac{\epsilon_0}{\log(2 + (2C)^i D)} \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

Hence, for any fixed ϵ_0 , the finiteness of $M_{[0, T]}$ from (26) implies that we can choose finitely many

numbers $0 = T_0 < T_1 < \dots < T_K < T_{K+1} = T$, with K depending on D , E , and ϵ_0 , such that

$$M_i := \int_{T_i}^{T_{i+1}} \int_{\mathbb{R}^3} |u(t, x)|^8 \log^{5\alpha/4}(2 + |u(t, x)|^2) dx dt = \frac{\epsilon_0}{\log(2 + (2C)^i D)}$$

for $0 \leq i \leq K - 1$ and $M_K \leq \epsilon_0 / \log(2 + (2C)^K D)$.

Choosing $\epsilon_0 = 1/(100C)^2$, by iteration and continuity arguments, we claim that $N_{[T_i, T_{i+1}]} < (2C)^{i+1} D$ for $0 \leq i \leq K$.⁶ Indeed, assume that this claim is false for some $i = j$. Then there exists $t_0 \in (T_j, T_{j+1})$ such that $N_{[T_j, t_0]} = (2C)^{j+1} D$. We have

$$\begin{aligned} (2C)^{j+1} D &\leq C(2C)^j D + CM_j^{1/2} N_{[T_j, t_0]} \log^{1/2}(2 + N_{[T_j, t_0]}^2) \\ &\leq \frac{1}{2}(2C)^{j+1} D + \frac{\log^{1/2}(2 + (2C)^{j+1} D)}{100 \log^{1/2}(2 + (2C)^j D)} \times (2C)^{j+1} D \\ &\leq \frac{3}{4}(2C)^{j+1} D. \end{aligned}$$

Thus the claim is proved by contradiction. This implies

$$\sum_{j=1}^2 \|\nabla_x^j u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} \leq N_{[0, T]} \leq \sum_{i=0}^K N_{[T_i, T_{i+1}]} < \sum_{i=0}^K (2C)^{i+1} D < \infty.$$

The universal bound only depends on D and E^7 , indicating the global existence.

Now we sketch the proof of $\tilde{H}_x^2 \times H_x^1$ scattering. From a similar argument as the one discussed in Remark 2.2, in order to prove $\tilde{H}_x^2(\mathbb{R}^3) \times H_x^1(\mathbb{R}^3)$ scattering, it suffices to show that

$$\| |u|^4 u \log^\alpha(2 + |u|^2) \|_{L_t^1 H_x^1(\mathbb{R}_+ \times \mathbb{R}^3)} < \infty. \quad (63)$$

By the above discussion, the universal bound is independent of T . Hence we have $N_{\mathbb{R}_+} < \infty$. By Hölder's inequality,

$$\| |u|^4 u \log^\alpha(2 + |u|^2) \|_{L_t^1 H_x^1(\mathbb{R}_+ \times \mathbb{R}^3)} \lesssim M_{\mathbb{R}_+}^{1/2} N_{\mathbb{R}_+} \log^{1/2}(2 + N_{\mathbb{R}_+}^2) < \infty. \quad \square$$

Appendix: Proof of (30)

Since (u_0, u_1) lies in $X_1 \times Y_1$, defined in (9), we have

$$\|u_0\|_{X_1}^2 \geq \int_{\mathbb{R}^3} |\nabla u_0|^2 \log^{2\gamma}(1 + |x|) dx \gtrsim (\log^{2\gamma} s) \int_{|x|>s} |\nabla u_0|^2 dx.$$

Hence

$$\int_{|x|>s} |\nabla u_0|^2 dx \lesssim \frac{\|u_0\|_{X_1}^2}{\log^{2\gamma} s}. \quad (64)$$

Similarly,

$$\int_{|x|>s} |u_1|^2 dx \lesssim \frac{\|u_1\|_{Y_1}^2}{\log^{2\gamma} s}. \quad (65)$$

⁶See the similar arguments in Lemma 3.5 and Corollary 3.8 or Proposition 3.1 in [Tao 2007].

⁷In fact, from corollary 3.2 in [Tao 2007], we have $N_{\mathbb{R}_+} \lesssim (2 + D)^{(2+D)^{O(E)}}$.

Now, consider

$$\begin{aligned} \int_{|x|>s} F(u_0(x)) dx &= \int_{\{|x|>s\} \cap \{|u_0|<1/3\}} F(u_0(x)) dx + \int_{\{|x|>s\} \cap \{|u_0|\geq 1/3\}} F(u_0(x)) dx \\ &\lesssim \int_{\{|x|>s\} \cap \{|u_0|<1/3\}} |u_0|^6 (-\log |u_0|) dx + \int_{\{|x|>s\} \cap \{|u_0|\geq 1/3\}} |u_0|^6 dx =: I + II. \end{aligned}$$

Let

$$\begin{aligned} I &= \int_{\{|x|>s\} \cap \{|u_0(x)|<1/|x|^{2/3}\}} |u_0|^6 (-\log |u_0|) dx + \int_{\{|x|>s\} \cap \{1/|x|^{2/3} \leq |u_0(x)| \leq 1/3\}} |u_0|^6 (-\log |u_0|) dx \\ &=: I_1 + I_2. \end{aligned}$$

When s is sufficiently large,

$$\begin{aligned} I_1 &\lesssim \int_{\{|x|>s\} \cap \{|u_0|<1/|x|^{2/3}\}} |u_0|^{11/2} \left(\sup_{|u_0|<s^{-2/3}} |u_0|^{1/2} (-\log |u_0|) \right) dx \\ &\lesssim \int_{|x|>s} |x|^{-11/3} dx \lesssim s^{-2/3} \lesssim \frac{1}{\log^{2\gamma} s}. \end{aligned} \quad (66)$$

Now we aim to prove that $I_2 + II \lesssim \frac{1}{\log^{2\gamma} s}$ for s sufficiently large. For $\alpha \in \mathbb{R}$, define

$$Q(\alpha) := \int_{\mathbb{R}^3} \left| \frac{u_0 \log^\alpha(2+|x|)}{2+|x|} \right|^2 dx.$$

We claim that

$$Q(\alpha) \leq C(\|u_0\|_{X_1}, E, \alpha) \quad \text{for } \alpha \leq \gamma, \quad (67)$$

where E is the energy. Indeed, if $\alpha \leq 0$, by Hölder's inequality and Hardy's inequality,

$$\begin{aligned} Q(\alpha) &= \int_{|x|<3} \left| \frac{u_0 \log^\alpha(2+|x|)}{2+|x|} \right|^2 dx + \int_{|x|\geq 3} \left| \frac{u_0 \log^\alpha(2+|x|)}{2+|x|} \right|^2 dx \\ &\lesssim_\alpha \int_{|x|<3} |u_0|^2 dx + \int_{\mathbb{R}^3} \left| \frac{u_0}{|x|} \right|^2 dx \lesssim \|u_0\|_{\dot{H}_x^1(\mathbb{R}^3)}^2 + \left(\int_{\mathbb{R}^3} F(u_0) dx \right)^{1/3} \leq C(E, \alpha). \end{aligned} \quad (68)$$

Again, if $0 < \alpha \leq \gamma$,

$$\begin{aligned} Q(\alpha) &= \int_{\mathbb{R}^3} \left| \frac{u_0 \log^\alpha(2+|x|)}{2+|x|} \right|^2 dx \lesssim_\alpha \int_{|x|<3} |u_0|^2 dx + \int_{\mathbb{R}^3} \left| \frac{u_0 \log^\alpha(2+|x|)}{|x|} \right|^2 dx \\ &\lesssim \left(\int_{|x|<3} |u_0|^6 dx \right)^{1/3} + \int_{\mathbb{R}^3} |\nabla(u_0 \log^\alpha(2+|x|))|^2 dx \\ &\lesssim_\alpha \left(\int_{\mathbb{R}^3} F(u) dx \right)^{1/3} + \int_{\mathbb{R}^3} |\nabla u_0 \log^\alpha(2+|x|)|^2 dx + \int_{\mathbb{R}^3} \left| \frac{u_0 \log^{\alpha-1}(2+|x|)}{2+|x|} \right|^2 dx \\ &\lesssim E^{1/3} + \int_{|x|<3} |\nabla u_0|^2 dx + \int_{|x|\geq 3} |\nabla u_0 \log^\gamma(1+|x|)|^2 dx + Q(\alpha-1) \\ &\lesssim E^{1/3} + E + \|u_0\|_{X_1} + Q(\alpha-1). \end{aligned}$$

By an inductive argument and (68), the claim is proved.

Fix $s \gg 1$. Let χ be the smooth radial function which equals 1 on $\{|x| > s\}$, 0 on $\{|x| < s/2\}$, $0 \leq \chi \leq 1$ and $|\nabla \chi| \lesssim 1/s$. Then we have $|\nabla \chi| \lesssim 1/|x|$. By the Sobolev embedding theorem and Hardy's inequality,

$$\begin{aligned} & \log^{6\gamma} s \int_{|x|>s} |u_0|^6 dx \\ & \leq \int_{|x|>s} |u_0|^6 \log^{6\gamma}(|x|) dx \leq \int_{\mathbb{R}^3} (\chi |u_0| \log^\gamma(2+|x|))^6 dx \lesssim \left(\int_{\mathbb{R}^3} |\nabla(\chi u_0 \log^\gamma(2+|x|))|^2 dx \right)^3 \\ & \lesssim_\gamma \left(\int_{\mathbb{R}^3} |\nabla \chi u_0 \log^\gamma(2+|x|)|^2 dx + \int_{\mathbb{R}^3} |\chi \nabla u_0 \log^\gamma(2+|x|)|^2 dx + \int_{\mathbb{R}^3} \left| \frac{\chi u_0 \log^{\gamma-1}(2+|x|)}{2+|x|} \right|^2 dx \right)^3 \\ & =: (J_1 + J_2 + J_3)^3. \end{aligned}$$

We can compute that

$$\begin{aligned} J_2 & \lesssim_\gamma \int_{\mathbb{R}^3} |\nabla u_0 \log^\gamma(1+|x|)|^2 dx + \int_{|x|<3} |\nabla u_0|^2 dx \leq \|u_0\|_{X_1}^2 + E, \\ J_3 & \lesssim C(\|u_0\|_{X_1}, E, \gamma), \quad \text{by (67)}. \end{aligned}$$

Since $\nabla \chi \lesssim 1/|x|$,

$$\begin{aligned} J_1 & \lesssim \int_{|x|>s/2} \left| \frac{u_0 \log^\gamma(2+|x|)}{|x|} \right|^2 dx \lesssim \int_{|x|>s/2} \left| \frac{u_0 \log^\gamma(2+|x|)}{2+|x|} \right|^2 dx \\ & \lesssim C(\|u_0\|_{X_1}, E, \gamma). \end{aligned}$$

Hence $\log^{6\gamma} s \int_{|x|>s} |u_0|^6 dx \leq C(\|u_0\|_{X_1}, E, \gamma)$ for sufficiently large s . Then we deduce

$$II \leq \int_{|x|>s} |u_0|^6 dx \lesssim \frac{1}{\log^{6\gamma} s} \leq \frac{1}{\log^{2\gamma} s}. \quad (69)$$

Similarly,

$$\begin{aligned} \log^{6\gamma-1} s \int_{\{|x|>s\} \cap \{1/|x|^{2/3} \leq |u_0| \leq 1/3\}} |u_0|^6 (-\log |u_0|) dx & \lesssim \log^{6\gamma-1} s \int_{|x|>s} |u_0|^6 \log(|x|) dx \\ & \lesssim \int_{|x|>s} |u_0|^6 \log^{6\gamma}(|x|) dx \lesssim C(\|u_0\|_{X_1}, E, \gamma). \end{aligned}$$

Therefore,

$$I_2 \lesssim \frac{1}{\log^{6\gamma-1} s} \leq \frac{1}{\log^{2\gamma} s}. \quad (70)$$

Combining (64), (65), (66), (69), and (70), we obtain (30).

Acknowledgements

The author thanks Markus Keel and Terence Tao for illuminating conversations about the problems considered here.

References

- [Bahouri and Gérard 1999] H. Bahouri and P. Gérard, “High frequency approximation of solutions to critical nonlinear wave equations”, *Amer. J. Math.* **121**:1 (1999), 131–175. MR 2000i:35123 Zbl 0919.35089
- [Bahouri and Shatah 1998] H. Bahouri and J. Shatah, “Decay estimates for the critical semilinear wave equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **15**:6 (1998), 783–789. MR 99h:35136 Zbl 0924.35084
- [Brenner and von Wahl 1981] P. Brenner and W. von Wahl, “Global classical solutions of nonlinear wave equations”, *Math. Z.* **176**:1 (1981), 87–121. MR 82m:35103 Zbl 0457.35059
- [Ginibre and Velo 1985] J. Ginibre and G. Velo, “The global Cauchy problem for the nonlinear Klein–Gordon equation”, *Math. Z.* **189**:4 (1985), 487–505. MR 86f:35149 Zbl 0549.35108
- [Ginibre and Velo 1987] J. Ginibre and G. Velo, “Conformal invariance and time decay for nonlinear wave equations. I, II”, *Ann. Inst. H. Poincaré Phys. Théor.* **47**:3 (1987), 221–261, 263–276. MR 89e:35021 Zbl 0644.35067
- [Ginibre and Velo 1989] J. Ginibre and G. Velo, “Scattering theory in the energy space for a class of nonlinear wave equations”, *Comm. Math. Phys.* **123**:4 (1989), 535–573. MR 90i:35172 Zbl 0698.35112
- [Ginibre and Velo 1995] J. Ginibre and G. Velo, “Generalized Strichartz inequalities for the wave equation”, *J. Funct. Anal.* **133**:1 (1995), 50–68. MR 97a:46047 Zbl 0849.35064
- [Ginibre et al. 1992] J. Ginibre, A. Soffer, and G. Velo, “The global Cauchy problem for the critical nonlinear wave equation”, *J. Funct. Anal.* **110**:1 (1992), 96–130. MR 94d:35105 Zbl 0813.35054
- [Grillakis 1990] M. G. Grillakis, “Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity”, *Ann. of Math. (2)* **132**:3 (1990), 485–509. MR 92c:35080 Zbl 0736.35067
- [Grillakis 1992] M. G. Grillakis, “Regularity for the wave equation with a critical nonlinearity”, *Comm. Pure Appl. Math.* **45**:6 (1992), 749–774. MR 93e:35073 Zbl 0785.35065
- [Hidano 2001] K. Hidano, “Scattering problem for the nonlinear wave equation in the finite energy and conformal charge space”, *J. Funct. Anal.* **187**:2 (2001), 274–307. MR 2002k:35242 Zbl 1005.35066
- [Kapitanski 1989] L. V. Kapitanski, “Some generalizations of the Strichartz–Brenner inequality”, *Algebra i Analiz* **1**:3 (1989), 127–159. In Russian. translated in *Leningr. Math. J.* **1**:3 (1990), 693–726. MR 90h:46063 Zbl 0732.35118
- [Kapitanski 1994] L. Kapitanski, “Global and unique weak solutions of nonlinear wave equations”, *Math. Res. Lett.* **1**:2 (1994), 211–223. MR 95f:35158 Zbl 0841.35067
- [Keel and Tao 1998] M. Keel and T. Tao, “Endpoint Strichartz estimates”, *Amer. J. Math.* **120**:5 (1998), 955–980. MR 2000d:35018 Zbl 0922.35028
- [Klainerman and Machedon 1993] S. Klainerman and M. Machedon, “Space-time estimates for null forms and the local existence theorem”, *Comm. Pure Appl. Math.* **46**:9 (1993), 1221–1268. MR 94h:35137 Zbl 0803.35095
- [Lindblad and Sogge 1995] H. Lindblad and C. D. Sogge, “On existence and scattering with minimal regularity for semilinear wave equations”, *J. Funct. Anal.* **130**:2 (1995), 357–426. MR 96i:35087 Zbl 0846.35085
- [Morawetz 1968] C. S. Morawetz, “Time decay for the nonlinear Klein–Gordon equations”, *Proc. Roy. Soc. Ser. A* **306** (1968), 291–296. MR 38 #2455 Zbl 0157.41502
- [Nakanishi 1999] K. Nakanishi, “Unique global existence and asymptotic behaviour of solutions for wave equations with non-coercive critical nonlinearity”, *Comm. Partial Differential Equations* **24**:1-2 (1999), 185–221. MR 2000f:35104 Zbl 0926.35022
- [Roy 2009] T. Roy, “Global existence of smooth solutions of a 3D log-log energy-supercritical wave equation”, *Anal. PDE* **2**:3 (2009), 261–280. MR 2603799 Zbl 1195.35222
- [Shatah and Struwe 1993] J. Shatah and M. Struwe, “Regularity results for nonlinear wave equations”, *Ann. of Math. (2)* **138**:3 (1993), 503–518. MR 95f:35164 Zbl 0836.35096
- [Shatah and Struwe 1994] J. Shatah and M. Struwe, “Well-posedness in the energy space for semilinear wave equations with critical growth”, *Internat. Math. Res. Notices* **7** (1994), 303–309. MR 95e:35132 Zbl 0830.35086
- [Sogge 1995] C. D. Sogge, *Lectures on nonlinear wave equations*, Monographs in Analysis **2**, International Press, Boston, 1995. MR 2000g:35153 Zbl 1089.35500

- [Strichartz 1977] R. S. Strichartz, “Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations”, *Duke Math. J.* **44**:3 (1977), 705–714. MR 58 #23577 Zbl 0372.35001
- [Struwe 1988] M. Struwe, “Globally regular solutions to the u^5 Klein–Gordon equation”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **15**:3 (1988), 495–513 (1989). MR 90j:35142 Zbl 0728.35072
- [Tao 2007] T. Tao, “Global regularity for a logarithmically supercritical defocusing nonlinear wave equation for spherically symmetric data”, *J. Hyperbolic Differ. Equ.* **4**:2 (2007), 259–265. MR 2009b:35294 Zbl 1124.35043

Received 27 May 2011. Revised 22 Nov 2011. Accepted 20 Mar 2012.

HSI-WEI SHIH: shihx029@umn.edu

School of Mathematics, University of Minnesota, 127 Vincent Hall, 206 Church St. SE, Minneapolis, MN 55455, United States
<http://math.umn.edu/~shihx029/>

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Michael Aizenman	Princeton University, USA aizenman@math.princeton.edu	Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr
Luis A. Caffarelli	University of Texas, USA caffarel@math.utexas.edu	Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Charles Fefferman	Princeton University, USA cf@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Nigel Higson	Pennsylvania State University, USA higson@math.psu.edu
Vaughan Jones	University of California, Berkeley, USA vfr@math.berkeley.edu	Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr
László Lempert	Purdue University, USA lempert@math.purdue.edu	Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Igor Rodnianski	Princeton University, USA irod@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2013 is US \$160/year for the electronic version, and \$310/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 6 No. 1 2013

Some results on scattering for log-subcritical and log-supercritical nonlinear wave equations	1
HSI-WEI SHIH	
Localisation and compactness properties of the Navier–Stokes global regularity problem	25
TERENCE TAO	
A variational principle for correlation functions for unitary ensembles, with applications	109
DORON S. LUBINSKY	
Relative Kähler–Ricci flows and their quantization	131
ROBERT J. BERMAN	
Resolvent estimates for elliptic quadratic differential operators	181
MICHAEL HITRIK, JOHANNES SJÖSTRAND and JOE VIOLA	
Bilinear Hilbert transforms along curves I: the monomial case	197
XIAOCHUN LI	
A gluing formula for the analytic torsion on singular spaces	221
MATTHIAS LESCH	



2157-5045(2013)6:1;1-H