

ANALYSIS & PDE

Volume 7

No. 1

2014

MICHAEL DABKOWSKI, ALEXANDER KISELEV,
LUIS SILVESTRE AND VLAD VICOL

**GLOBAL WELL-POSEDNESS OF SLIGHTLY SUPERCRITICAL
ACTIVE SCALAR EQUATIONS**

GLOBAL WELL-POSEDNESS OF SLIGHTLY SUPERCRITICAL ACTIVE SCALAR EQUATIONS

MICHAEL DABKOWSKI, ALEXANDER KISELEV, LUIS SILVESTRE AND VLAD VICOL

The paper is devoted to the study of slightly supercritical active scalars with nonlocal diffusion. We prove global regularity for the surface quasigeostrophic (SQG) and Burgers equations, when the diffusion term is supercritical by a symbol with roughly logarithmic behavior at infinity. We show that the result is sharp for the Burgers equation. We also prove global regularity for a slightly supercritical two-dimensional Euler equation. Our main tool is a nonlocal maximum principle which controls a certain modulus of continuity of the solutions.

1. Introduction

Active scalars play an important role in fluid mechanics. An active scalar equation is given by

$$\partial_t \theta + (u \cdot \nabla) \theta + \mathcal{L} \theta = 0, \quad \theta(x, 0) = \theta_0(x), \quad (1-1)$$

where \mathcal{L} is typically some dissipative operator, such as fractional dissipation, and u is the flow velocity that is determined by θ . A common setting is either on \mathbb{R}^d or \mathbb{T}^d . Active scalar equations are nonlinear, and most active scalars of interest are nonlocal. This makes the analysis of these equations challenging. The best known active scalar equations are the two-dimensional Euler equation in vorticity form (for which $u = \nabla^\perp (-\Delta)^{-1} \theta$), the surface quasigeostrophic (SQG) equation ($d = 2$, $u = \nabla^\perp (-\Delta)^{-1/2} \theta$), and the one-dimensional Burgers equation ($u = \theta$). The two-dimensional Euler and Burgers equations are classical in fluid mechanics, while the SQG equation was first considered in the mathematical literature by Constantin, Majda, and Tabak [Constantin et al. 1994], and since then has attracted significant attention, in part due to certain similarities with three-dimensional Euler and Navier–Stokes equations.

Observe that for the SQG and Burgers equations the drift velocity u and the advected scalar θ are of the same order of regularity, while for the two-dimensional Euler equation u is more regular by a derivative. The two-dimensional Euler equation has global regular solutions, and can be thought of as a critical case. For the Burgers and SQG equations, fractional dissipation $\mathcal{L} = \Lambda^\alpha$, where $\Lambda = (-\Delta)^{1/2}$ is the Zygmund operator, have often been considered. Both of these equations possess the L^∞ maximum

Dabkowski was supported in part by a research grant of the University of Wisconsin–Madison Graduate School. Kiselev was partially supported by the NSF grants DMS-1104415, DMS-1159133, and the Guggenheim fellowship. Silvestre was partially supported by NSF grants DMS-1001629, DMS-1065979, and the Sloan fellowship. Vicol was partially supported by an AMS–Simons travel award and by the NSF grant DMS-1211828.

MSC2010: 35Q35, 76U05.

Keywords: surface quasigeostrophic equation, active scalars, global regularity, finite time blow-up, supercritical dissipation, nonlocal maximum principle, nonlocal dissipation, SQG equation, Burgers equation.

principle [Resnick 1995; Córdoba and Córdoba 2004], and this makes $\alpha = 1$ critical with respect to the natural scaling of the equations. It has been known for a while that in the subcritical case $\alpha > 1$, global regular solutions exist for sufficiently smooth initial data (see [Resnick 1995] for the SQG equation; the analysis for the Burgers equation is very similar; see, for example, [Kiselev et al. 2008]). The critical case $\alpha = 1$ has been especially interesting for the SQG equation since it is well motivated physically, with the $\Lambda\theta$ term modeling so called the Ekman boundary layer pumping effect; see, for example, [Pedlosky 1987]. The global regularity in the critical case has been settled independently by Kiselev, Nazarov, and Volberg [Kiselev et al. 2007] and Caffarelli and Vasseur [2010]. A third proof of the same result was provided by Kiselev and Nazarov [2009], and a fourth by Constantin and Vicol [2012]. All these proofs are quite different. The method of [Caffarelli and Vasseur 2010] is inspired by DeGiorgi iterative estimates, while the duality approach of [Kiselev and Nazarov 2009] uses an appropriate set of test functions and estimates on their evolution. The proof in [Constantin and Vicol 2012] takes advantage of a new nonlinear maximum principle, which gives a nonlinear bound on a linear operator. The method of [Kiselev et al. 2007], on the other hand, is based on a technique which may be called a nonlocal maximum principle. The idea is to prove that the evolution (1-1) preserves a certain modulus of continuity ω of the solution. In the critical SQG case, the control is strong enough to give a uniform bound on $\|\nabla\theta\|_{L^\infty}$, which is sufficient for global regularity.

In the supercritical case, until recently the only results available (for large initial data) have been on conditional regularity and finite time regularization of solutions. It was shown by Constantin and Wu [2008] that if the solution is C^δ with $\delta > 1 - \alpha$, it is smooth (see also [Silvestre 2011] for drift velocity that is not divergence free). Dong and Pavlovic [2009] improved this result to $\delta = 1 - \alpha$. Finite time regularization has been proved by Silvestre [2010] for α sufficiently close to 1, and for the whole dissipation range $0 < \alpha < 1$ by Dabkowski [2011] (with an alternative proof of the latter result given in [Kiselev 2011]). The issue of global regularity in the case $\alpha \in (0, 1)$ remains an outstanding open problem. A small advance into the supercritical regime was made in [Dabkowski et al. 2012], where the SQG equation with velocity given by

$$u = \nabla^\perp \Lambda^{-1} m(\Lambda)\theta$$

was considered. Here m is a Fourier multiplier which may grow at infinity at any rate slower than double logarithm. The method of [Dabkowski et al. 2012] was based on the technique of [Kiselev et al. 2007]. The main issue is that even with very slow growth of m , the equation loses scaling, which plays an important role in every proof of regularity for the critical case. [Dabkowski et al. 2012] was partly inspired by the slightly supercritical Navier–Stokes regularity result of Tao [2009], and partly by recent work on generalized Euler and SQG models [Chae et al. 2011; Chae et al. 2010].

In this paper, we analyze a slightly supercritical SQG equation and the Burgers equation equation. As opposed to [Dabkowski et al. 2012], we keep the velocity definition the same as for classical SQG and Burgers equations, and instead treat supercritical diffusion. We also consider nonlocal diffusion terms more general than the fractional Laplacian, including cases where the L^∞ maximum principle does not hold. We show, roughly, that symbols supercritical by a logarithm or less lead to global regular solutions

for both equations. Our main technique is the control of an appropriate family of moduli of continuity of the solutions. For the Burgers equation, when the conditions we impose on the diffusion in order to obtain global regularity are not satisfied, we prove that some smooth initial data leads to finite time blow-up; see also [Alibaud et al. 2007; Dong et al. 2009; Kiselev et al. 2008]. In this respect, our well-posedness result is sharp. For the SQG equation, the global regularity proof is more sophisticated than for the Burgers equation. The upgrade from the double logarithmic supercriticality of [Dabkowski et al. 2012] to the logarithmic one is made possible by exploiting the structure of nonlinearity, in particular the ∇^\perp in $u = \nabla^\perp \Lambda^{-1} \theta$. This idea is based on [Kiselev 2011], where this structure was exploited to prove finite time regularization for power supercritical SQG equations. We note that Xue and Zheng [2012] observed a similar improvement from log log to log in the context of supercritical velocity.

We also consider the slightly supercritical two-dimensional Euler equation, and generalize the results of [Chae et al. 2011] on global regularity of solutions.

General diffusion of integral type arises from probabilistic models which involve discontinuous Lévy processes. Indeed, the classical Lévy–Khintchine representation formula shows that very general integral diffusion arises as the generator of Lévy processes. This type of diffusion has many applications in the physical sciences; see, for example, [Klafter and Sokolov 2005] and the references therein.

Below, we state main results proved in the paper. In Section 2, we provide some basic background results on the nonlocal maximum principles. Section 3 is devoted to proving global regularity for the slightly supercritical SQG equation with nonlocal diffusions. The Burgers case is handled in Section 4. In Section 5, we consider the case of dissipation given by Fourier multipliers. Some natural multipliers can lead to nonpositive convolution kernels for the corresponding nonlocal diffusion, and we generalize our technique to this case. Section 6 is devoted to the slightly supercritical two-dimensional Euler equation.

To state our main results, we need to introduce some notation.

1A. Assumptions on m . Let $m : (0, \infty) \rightarrow [0, \infty)$ be a nonincreasing smooth function which is singular at the origin, that is, $\lim_{r \rightarrow 0} m(r) = \infty$, and satisfies the following conditions:

- (i) There exists a sufficiently large positive constant $C_0 > 0$ such that

$$rm(r) \leq C_0 \quad \text{for all } r \in (0, r_0) \tag{1-2}$$

for some $r_0 > 0$. This condition is natural in the present context, since otherwise the dissipative operator defined below is subcritical, which is not the purpose of this paper.

- (ii) There exists some $\alpha > 0$ such that

$$r^\alpha m(r) \text{ is nonincreasing.} \tag{1-3}$$

This assumption is slightly stronger than just having $m(r)$ be nonincreasing.

Throughout this paper we also denote by m the radially symmetric function $m : \mathbb{R}^d \setminus \{0\} \rightarrow [0, \infty)$ such that $m(y) = m(|y|)$ for each $y \in \mathbb{R}^d \setminus \{0\}$. Note that the above conditions allow for m to be identically zero on the complement of a ball.

The examples of functions m which are relevant to this paper are those that are less singular than r^{-1} near $r = 0$. These functions yield dissipative nonlocal operators (cf. (1-5) below) that are less smoothing than Λ , which makes the corresponding SQG and Burgers equations *supercritical*. The main examples we have in mind are

$$m(r) = \frac{1}{r^a} \quad \text{and} \quad m(r) = \frac{1}{r(\log(2/r))^a} \quad (1-4)$$

with $0 < a \leq 1$ and $0 < r \leq 1$, coupled with enough regularity and decay for $r > 1$. The first class corresponds to power supercriticality. The second class, supercritical by a logarithm, is relevant for the global well-posedness results we prove. It is not hard to verify that the functions in (1-4) verify (1-2)–(1-3) on $(0, 1]$, and that they can be suitably extended on $[1, \infty)$.

1B. Dissipative nonlocal operators. Associated to any such function m , we consider the nonlocal operator

$$\mathcal{L}\theta(x) = \int_{\mathbb{R}^d} (\theta(x) - \theta(x+y)) \frac{m(y)}{|y|^d} dy. \quad (1-5)$$

Above and throughout the rest of the paper the integral is meant in principal value sense, but we omit the P.V. in front of the integral. For example, when $m(r) = r^{-\alpha} C_{d,\alpha}$ for a suitable normalizing constant, $C_{d,\alpha}$, $\mathcal{L} = \Lambda^\alpha$. The nonlocal operators \mathcal{L} we consider here are dissipative because m is singular at the origin: due to (1-3), we have that $m(r) \geq m(1)r^{-\alpha}$ for some $\alpha > 0$ when $r \leq 1$, so that \mathcal{L} is at least as dissipative as Λ^α . It is now evident that when $\lim_{r \rightarrow 0} rm(r) = 0$, the corresponding nonlocal operator \mathcal{L} is less smoothing than Λ . We emphasize that, for $\theta \in C^\infty(\mathbb{T}^d)$, the P.V. integral in (1-5) converges only if m is subquadratic near $r = 0$, that is,

$$\int_0^1 rm(r) dr < \infty$$

holds. In our case, the above condition is satisfied in view of assumption (1-2). Convergence near infinity is not an issue due to assumption (1-3).

All results in this paper can be generalized to a more general class of dissipative operators. Namely, for each function m that satisfies (1-2)–(1-3), consider the class of smooth radially symmetric kernels $K : \mathbb{R}^d \setminus \{0\} \rightarrow (0, \infty)$ which satisfy

$$\frac{m(y)}{C|y|^d} \leq K(y) \leq \frac{Cm(y)}{|y|^d} \quad (1-6)$$

for some constant $C > 0$ and all $y \neq 0$. Associated to each such kernel K we may consider the dissipative nonlocal operator

$$\mathcal{L}\theta(x) = \int_{\mathbb{R}^d} (\theta(x) - \theta(x+y)) K(y) dy, \quad (1-7)$$

which generalizes the definition in (1-5). As we will see, such generalization will be useful when working with dissipative operators generated by Fourier multipliers. Moreover, as we will see later, conditions on K can be relaxed further.

1C. Main results. The generalized dissipative SQG equation reads

$$\partial_t \theta + u \cdot \nabla \theta + \mathcal{L} \theta = 0, \quad (1-8)$$

$$u = \nabla^\perp \Lambda^{-1} \theta, \quad (1-9)$$

where \mathcal{L} is as defined in (1-5) and m is as described above. The main result of this paper with respect to the dissipative SQG equation is the following.

Theorem 1.1 (global regularity for slightly supercritical SQG). *Assume that θ_0 is smooth and periodic, and m satisfies an additional assumption*

$$\lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^1 m(r) dr = \infty. \quad (1-10)$$

Then there exists a unique, global in time, C^∞ smooth solution θ of the initial value problem associated to (1-8)–(1-9).

In analogy, one may consider the generalized dissipative Burgers equation

$$\partial_t \theta - \theta \theta_x + \mathcal{L} \theta = 0, \quad (1-11)$$

where \mathcal{L} and m are as before, and $d = 1$. Then we prove

Theorem 1.2 (global regularity for fractal Burgers). *Assume that θ_0 is smooth and periodic, and m is such that (1-2)–(1-3) hold and*

$$\lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^1 m(r) dr = \infty. \quad (1-12)$$

Then there exists a unique, global in time, C^∞ smooth solution θ of the initial value problem associated to (1-11).

In addition, in the case of the Burgers equation we prove that condition (1-12) is sharp.

Theorem 1.3 (finite time blow-up for fractal Burgers). *Assume that m is such that (1-2)–(1-3) hold, and in addition we have*

$$r |m'(r)| \leq C m(r) \quad (1-13)$$

for $r > 0$ and some constant $C \geq 1$. Furthermore, suppose that

$$\lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^1 m(r) dr < \infty \quad (1-14)$$

holds. Then there exists an initial datum $\theta_0 \in C^\infty(\mathbb{T})$, and $T > 0$ such that $\lim_{t \rightarrow T} \|\theta_x(t)\|_{L^\infty} = \infty$, that is, we have finite time blow-up arising from smooth initial data.

A natural class of dissipation terms is associated with Fourier multiplier operators. This representation is closely related to the form (1-7). As noted above, when $m(r) = r^{-\alpha} C_{d,\alpha}$ for a suitable constant $C_{d,\alpha}$, $\mathcal{L} = \Lambda^\alpha$, corresponding to the Fourier multiplier with symbol $P(\zeta) = |\zeta|^\alpha$. One may generalize this statement as follows. Let $P(\zeta)$ be a sufficiently nice Fourier multiplier (see Lemmas 5.1 and 5.2 for

precise assumptions on P), and let $K(y)$ be the convolution kernel associated to the multiplier P , that is, $\hat{\mathcal{L}}\theta(\zeta) = P(\zeta)\hat{\theta}(\zeta)$, where \mathcal{L} is the operator defined in (1-7). Then there exists a positive constant C that depends only on P , such that (1-6) holds for all sufficiently small y , with $m(y) = P(1/\zeta)$. This turns out to be sufficient to prove an analog of Theorem 1.1 (and Theorem 1.2).

Theorem 1.4 (global regularity for slightly supercritical SQG). *Let P be a radially symmetric Fourier multiplier that is smooth away from zero, nondecreasing, satisfies $P(0) = 0$, $P(\zeta) \rightarrow \infty$ as $|\zeta| \rightarrow \infty$, as well as conditions (5-3)–(5-4), and (5-9). Suppose also*

$$P(|\zeta|) \leq C|\zeta| \quad (1-15)$$

for all $|\zeta|$ sufficiently large,

$$|\zeta|^{-\alpha} P(|\zeta|) \text{ is nondecreasing} \quad (1-16)$$

for some $\alpha > 0$, and

$$\lim_{\epsilon \rightarrow 0} \int_{\epsilon}^1 P(|\zeta|^{-1}) d|\zeta| = \infty. \quad (1-17)$$

Then, for any θ_0 that is smooth and periodic, the Cauchy problem for the dissipative SQG equation (5-1)–(5-2) has a unique global in time smooth solution.

In particular, Theorem 1.4 proves global regularity of solutions for dissipative terms given by multipliers with behavior $P(\zeta) \sim |\zeta|(\log |\zeta|)^{-a}$ for large ζ , where $0 \leq a \leq 1$. The details of the assumptions on P and more discussion can be found in Section 5 below.

2. Pointwise moduli of continuity

Definition 2.1 (modulus of continuity). *We call a function $\omega: [0, \infty) \rightarrow [0, \infty)$ a modulus of continuity if $\omega(0) = 0$, ω is nondecreasing, continuous, concave, piecewise C^2 with one sided derivatives, and additionally satisfies $\omega'(0+) < \infty$ and $\omega''(0+) = -\infty$. We say that a smooth function f obeys the modulus of continuity ω if $|f(x) - f(y)| < \omega(|x - y|)$ for all $x \neq y$.*

We recall that if $f \in C^\infty(\mathbb{T}^2)$ obeys the modulus ω , then $\|\nabla f\|_{L^\infty} < \omega'(0+)$ [Kiselev et al. 2007]. In addition, observe that a function $f \in C^\infty(\mathbb{T}^2)$ automatically obeys any modulus of continuity $\omega(\xi)$ that lies above the function $\min\{\xi \|\nabla f\|_{L^\infty}, 2\|f\|_{L^\infty}\}$.

The following lemma gives the modulus of continuity of the Riesz transform of a given function.

Lemma 2.2 (modulus of continuity under a Riesz transform). *Assume that θ obeys the modulus of continuity ω , and that the drift velocity is given by the constitutive law $u = \nabla^\perp \Lambda^{-1}\theta$. Then u obeys the modulus of continuity Ω defined as*

$$\Omega(\xi) = A \left(\int_0^\xi \frac{\omega(\eta)}{\eta} d\eta + \xi \int_\xi^\infty \frac{\omega(\eta)}{\eta^2} d\eta \right) \quad (2-1)$$

for some positive universal constant $A > 0$.

Moreover, for any two points x, y with $|x - y| = \xi > 0$, we have

$$\left| (u(x) - u(y)) \cdot \frac{x - y}{|x - y|} \right| \leq \tilde{\Omega}(\xi) + \Omega^\perp(\xi), \quad (2-2)$$

where

$$\tilde{\Omega}(\xi) = A \left(\omega(\xi) + \xi \int_\xi^\infty \frac{\omega(\eta)}{\eta^2} d\eta \right) \quad (2-3)$$

and

$$\Omega^\perp(\xi) = A \int_0^{\xi/4} \int_{\xi/4}^{3\xi/4} (\theta(\eta, v) - \theta(-\eta, v) - \theta(\eta, -v) + \theta(-\eta, -v)) \frac{v}{((\xi/2 - \eta)^2 + v^2)^{3/2}} d\eta dv, \quad (2-4)$$

where A is a universal constant.

The proof of (2-1) may be found in [Kiselev et al. 2007, Appendix], while (2-2) was obtained in [Kiselev 2011, Lemma 5.2], to which we refer for further details.

Lemma 2.3 (dissipation control). *Let \mathcal{L} be defined as in (1-7), with K satisfying (1-6). Assume that $\theta \in C^\infty(\mathbb{T}^2)$ obeys a concave modulus of continuity ω . Suppose that there exist two points x, y with $|x - y| = \xi > 0$ such that $\theta(x) - \theta(y) = \omega(\xi)$. Then we have*

$$\mathcal{L}\theta(x) - \mathcal{L}\theta(y) \geq \mathfrak{D}(\xi) + \mathfrak{D}^\perp(\xi) \quad (2-5)$$

where

$$\begin{aligned} \mathfrak{D}(\xi) = & \frac{1}{A} \int_0^{\xi/2} (2\omega(\xi) - \omega(\xi + 2\eta) - \omega(\xi - 2\eta)) \frac{m(2\eta)}{\eta} d\eta \\ & + \frac{1}{A} \int_{\xi/2}^\infty (2\omega(\xi) - \omega(\xi + 2\eta) + \omega(2\eta - \xi)) \frac{m(2\eta)}{\eta} d\eta \end{aligned} \quad (2-6)$$

and

$$\begin{aligned} \mathfrak{D}^\perp(\xi) = & \frac{1}{A} \int_0^{\xi/4} \int_{\xi/4}^{3\xi/4} (2\omega(2\eta) - \theta(\eta, v) + \theta(-\eta, v) - \theta(\eta, -v) + \theta(-\eta, -v)) \\ & \times \frac{m(\sqrt{(\xi/2 - \eta)^2 + v^2})}{(\xi/2 - \eta)^2 + v^2} d\eta dv \end{aligned} \quad (2-7)$$

with some sufficiently large universal constant $A > 0$ (that we can, for simplicity of presentation, choose to be the same as in Lemma 2.2). The corresponding lower bound in one dimension includes only the \mathfrak{D} term.

Note that $\mathfrak{D} \geq 0$ due to the concavity of ω , while $\mathfrak{D}^\perp \geq 0$ since θ obeys the modulus of continuity ω . The above lemma may be obtained along the lines of [Kiselev 2011, Section 5], where it was obtained for $\mathcal{L} = \Lambda$. However, some modifications are necessary for more general diffusions we consider, and we provide a proof in the Appendix below. We conclude this section by establishing a bound for $\Omega^\perp(\xi)$ in terms of $\mathfrak{D}^\perp(\xi)$.

Lemma 2.4 (connection between Ω^\perp and \mathcal{D}^\perp). *Let m be as in Section 1A, and assume θ obeys the modulus of continuity ω . For $\Omega^\perp(\xi)$ and $\mathcal{D}^\perp(\xi)$ defined via (2-4) and (2-7), respectively, we have*

$$m(\xi)\Omega^\perp(\xi) \leq A^2\mathcal{D}^\perp(\xi) \quad (2-8)$$

for all $\xi > 0$.

Proof of Lemma 2.4. To prove (2-8), first observe that since θ obeys ω , we have $\theta(\eta, \nu) - \theta(-\eta, \nu) \leq \omega(2\eta)$ and also $\theta(\eta, -\nu) - \theta(-\eta, -\nu) \leq \omega(2\eta)$. Therefore we have that

$$\begin{aligned} & |\theta(\eta, \nu) - \theta(-\eta, \nu) - \theta(\eta, -\nu) + \theta(-\eta, -\nu)| \\ & \leq 2\omega(2\eta) - \theta(\eta, \nu) + \theta(-\eta, \nu) - \theta(\eta, -\nu) + \theta(-\eta, -\nu) \end{aligned} \quad (2-9)$$

holds, for any $(\eta, \nu) \in \mathbb{R}^2$.

Next, we claim that, for any $0 < \nu \leq \xi/4$ and any $|\eta - \xi/2| \leq \xi/4$, we have

$$\frac{\nu m(\xi)}{((\xi/2 - \eta)^2 + \nu^2)^{3/2}} \leq \frac{m(\sqrt{(\xi/2 - \eta)^2 + \nu^2})}{(\xi/2 - \eta)^2 + \nu^2}. \quad (2-10)$$

To prove (2-10), we observe that in this range for (η, ν) we have $\sqrt{(\xi/2 - \eta)^2 + \nu^2} \leq \xi$, and due to the monotonicity of m , it follows that $m(\xi) \leq m(\sqrt{(\xi/2 - \eta)^2 + \nu^2})$. Since $\nu \leq \sqrt{(\xi/2 - \eta)^2 + \nu^2}$, (2-10) holds. Recalling the definitions of Ω^\perp and \mathcal{D}^\perp , it is clear that (2-8) follows directly from (2-9) and (2-10), concluding the proof of the lemma. \square

3. Global regularity for slightly supercritical SQG

Proof of Theorem 1.1. The local well-posedness for smooth solutions to SQG-type equations is by now standard. In particular, we have.

Proposition 3.1 (local existence of a smooth solution). *Given a periodic $\theta_0 \in C^\infty$, there exists $T > 0$ and a periodic solution $\theta(\cdot, t) \in C^\infty$ of (1-8)–(1-9). Moreover, the smooth solution may be continued beyond T as long as $\|\nabla\theta\|_{L^1(0,T;L^\infty)} < \infty$.*

The local in time propagation of C^∞ regularity may in fact even be obtained in the absence of dissipation. Since in (1-8)–(1-9) we have a dissipative term, one may actually show local C^∞ regularization of sufficiently regular initial data. The proof may be obtained in analogy to the usual supercritical SQG equation [Dong 2010], since, in view of (1-3), \mathcal{L} is smoothing more than Λ^α for some $\alpha > 0$. The presence of the general dissipation \mathcal{L} instead of the usual Λ^α does not introduce substantial difficulties.

The main difficulty in proving Theorem 1.1 is the supercriticality of the dissipation in (1-8)–(1-9). Thus, as opposed to the critical case [Kiselev et al. 2007], here we *cannot* construct a single modulus of continuity $\omega(\xi)$ preserved by the equation, and then use the scaling $\omega_B(\xi) = \omega(B\xi)$ to obtain a family of moduli of continuity such that each initial data obeys a modulus in this family. Instead, we will separately construct a modulus of continuity $\omega_B(\xi)$ for each initial data, and each such modulus will be preserved by the equation for all times; see also [Dabkowski et al. 2012].

Before constructing the aforementioned family of moduli, let us recall the breakthrough scenario of [Kiselev et al. 2007].

Lemma 3.2 (breakthrough scenario). *Assume ω is a modulus of continuity such that $\omega(0+) = 0$ and $\omega''(0+) = -\infty$. Suppose that the initial data θ_0 obeys ω . If the solution $\theta(x, t)$ violates ω at some positive time, there must exist $t_1 > 0$ and $x \neq y \in \mathbb{T}^2$ such that*

$$\theta(x, t_1) - \theta(y, t_1) = \omega(|x - y|),$$

and $\theta(x, t)$ obeys ω for every $0 \leq t < t_1$.

Let us consider the breakthrough scenario for a modulus of continuity ω . A simple computation [Kiselev 2011] combined with Lemma 2.2 and Lemma 2.3 yields

$$\begin{aligned} \partial_t(\theta(x, t) - \theta(y, t))|_{t=t_1} &= u \cdot \nabla \theta(y, t_1) - u \cdot \nabla \theta(x, t_1) + \mathcal{L}\theta(y, t_1) - \mathcal{L}\theta(x, t_1) \\ &\leq \left| (u(x, t_1) - u(y, t_1)) \cdot \frac{x - y}{|x - y|} \right| \omega'(\xi) + \mathcal{L}\theta(y, t_1) - \mathcal{L}\theta(x, t_1) \\ &\leq \min\{\Omega(\xi), \tilde{\Omega}(\xi) + \Omega^\perp(\xi)\} \omega'(\xi) - (\mathcal{D}(\xi) + \mathcal{D}^\perp(\xi)), \end{aligned} \quad (3-1)$$

where Ω , $\tilde{\Omega}$, Ω^\perp , \mathcal{D} , and \mathcal{D}^\perp are given in (2-1), (2-3), (2-4), (2-6), and (2-7), respectively. If we can show that the expression on the right side of (3-1) must be strictly negative, we obtain a contradiction: ω cannot be broken, and hence it is preserved by the evolution (1-8).

3A. Construction of the family of moduli of continuity. We now construct a family of moduli of continuity ω_B , such that, given any periodic C^∞ function θ_0 , there exists $B \geq 1$ such that θ_0 obeys ω_B .

Fix a sufficiently small positive constant $\kappa > 0$, to be chosen precisely later in terms of the constant A of (2-3) and the function m . For any $B \geq 1$, we define $\delta(B)$ to be the unique solution of

$$m(\delta(B)) = \frac{B}{\kappa}. \quad (3-2)$$

Since m is continuous, nonincreasing, $m(r) \rightarrow +\infty$ as $r \rightarrow 0+$, and (1-3) holds, such a solution $\delta(B)$ exists for any $B \geq 1$ (if κ is small enough). For convenience we can ensure that $\delta(B) \leq r_0/4$ for any $B \geq 1$ by using (1-2) and choosing $\kappa < r_0/(4C_0)$. Note that $\delta(B)$ is a nonincreasing function of B .

We let $\omega_B(\xi)$ be the continuous function with $\omega_B(0) = 0$ and

$$\omega'_B(\xi) = B - \frac{B^2}{2C_\alpha \kappa} \int_0^\xi \frac{3 + \ln(\delta(B)/\eta)}{\eta m(\eta)} d\eta \quad \text{for } 0 < \xi < \delta(B), \quad (3-3)$$

$$\omega'_B(\xi) = \gamma m(2\xi) \quad \text{for } \xi > \delta(B), \quad (3-4)$$

where $C_\alpha = (1 + 3\alpha)/\alpha^2$ and $\gamma > 0$ is a constant to be chosen later in terms of κ , A , and the function m (through C_0 , α , r_0 of assumptions (1-2)–(1-3)). We emphasize that neither κ nor γ will depend on B .

Let us now verify that the above defined function ω_B is indeed a modulus of continuity in the sense of Definition 2.1. First notice that by construction $\omega'_B(0+) = B$ and $\omega_B(\xi) \leq B\xi$ for all $0 < \xi \leq \delta(B)$. To verify that ω_B is nondecreasing, since m is nonnegative, we only need to check that $\omega'_B > 0$ for $\xi < \delta(B)$.

This is equivalent to verifying that $\omega'_B(\delta(B)-) > 0$. Using (1-3) and the change of variables $\delta(B)/\eta \mapsto \xi$, we may estimate

$$\int_0^{\delta(B)} \frac{3 + \ln(\delta(B)/\eta)}{\eta m(\eta)} d\eta \leq \int_0^{\delta(B)} \frac{3 + \ln(\delta(B)/\eta)}{\eta^{1-\alpha} \delta(B)^\alpha m(\delta(B))} d\eta = \frac{1}{m(\delta(B))} \int_1^\infty \frac{3 + \ln \xi}{\xi^{1+\alpha}} d\xi = \frac{C_\alpha}{m(\delta(B))},$$

where $C_\alpha = (1 + 3\alpha)/\alpha^2$ may be computed explicitly. The above estimate and (3-2)–(3-3) imply that

$$\omega'_B(\delta(B)-) \geq B - \frac{C_\alpha B^2}{2C_{\alpha\kappa} m(\delta(B))} = \frac{B}{2}, \quad (3-5)$$

which concludes the proof that $\omega'_B > 0$.

In order to verify that $\omega''_B(0+) = -\infty$, we use (1-2) and (3-3) to obtain

$$\omega''_B(\xi) = -\frac{B^2}{2C_{\alpha\kappa}\xi m(\xi)} \left(3 + \ln \frac{\delta(B)}{\xi} \right) \leq -\frac{B^2}{2C_{\alpha\kappa}C_0} \left(3 + \ln \frac{\delta(B)}{\xi} \right),$$

which is strictly negative for $0 < \xi < \delta(B)$, and also converges to $-\infty$ as $\xi \rightarrow 0+$.

Since m is nonincreasing, the concavity may only fail at $\xi = \delta(B)$. By (3-2) and (3-5) we have

$$\omega'_B(\delta(B)+) = \gamma m(2\delta(B)) \leq \gamma m(\delta(B)) = \frac{\gamma B}{\kappa} \leq \frac{B}{2} \leq \omega'_B(\delta(B)-)$$

provided that $2\gamma \leq \kappa$, and therefore ω_B is concave on $(0, \infty)$. It will also be useful to observe that, due to the concavity of ω_B and the mean value theorem, we have

$$\omega_B(\delta(B)) \geq \delta(B)\omega'_B(\delta(B)-) \geq \frac{\delta(B)B}{2}. \quad (3-6)$$

3B. Each initial data obeys a modulus. In order to show that given any $\theta_0 \in C^\infty(\mathbb{T}^2)$ there exists $B \geq 1$ such that θ_0 obeys $\omega_B(\xi)$, it is enough to find a B such that $\omega_B(\xi) > \min\{\xi\|\nabla\theta_0\|_{L^\infty}, 2\|\theta_0\|_{L^\infty}\}$ for all $\xi > 0$. Letting $a = 2\|\theta_0\|_{L^\infty}/\|\nabla\theta_0\|_{L^\infty}$, due to the concavity of ω_B , it is sufficient to find $B \geq 1$ such that $\omega_B(a) > 2\|\theta_0\|_{L^\infty}$. First, by choosing B large enough, we can ensure that $a > \delta(B)$. Then we have that

$$\omega_B(a) = \omega_B(\delta(B)) + \int_{\delta(B)}^a \omega'_B(\eta) d\eta \geq \gamma \int_{\delta(B)}^a m(2\eta) d\eta \rightarrow \infty \quad \text{as } \delta(B) \rightarrow 0 \quad (3-7)$$

due to the assumption (1-10). Therefore each initial data obeys a modulus from the family $\{\omega_B\}_{B \geq 1}$.

3C. The moduli are preserved by the evolution. It is left to verify that the above constructed family of moduli of continuity satisfy

$$\min\{\Omega_B(\xi), \tilde{\Omega}_B(\xi) + \Omega_B^\perp(\xi)\} \omega'_B(\xi) - (\mathcal{D}_B(\xi) + \mathcal{D}_B^\perp(\xi)) < 0 \quad (3-8)$$

for any $\xi > 0$ and $B \geq 1$. Here Ω_B and others are defined just as Ω and others, but with ω replaced by ω_B .

The case $\xi \geq \delta(B)$. First we observe that, by Lemma 2.4 and the fact that m is nonincreasing, we have

$$\omega'_B(\xi)\Omega_B^\perp(\xi) = \gamma m(2\xi)\Omega_B^\perp(\xi) \leq \gamma m(\xi)\Omega_B^\perp(\xi) \leq \gamma A^2 \mathcal{D}_B^\perp(\xi) \leq \mathcal{D}_B^\perp(\xi)$$

for all $\xi \geq \delta(B)$ if we choose $\gamma \leq 1/A^2$. In view of (3-8), it is left to prove that

$$\tilde{\Omega}_B(\xi)\omega'_B(\xi) - \mathcal{D}_B(\xi) < 0$$

for all $\xi \geq \delta(B)$. In order to do this we claim that, for all $\xi > \delta(B)$, we have

$$\omega_B(2\xi) \leq c_\alpha \omega_B(\xi), \quad (3-9)$$

where $c_\alpha = 1 + (3/2)^{-\alpha}$ and $\alpha > 0$ is as in assumption (1-3). Note that, by definition, $1 < c_\alpha < 2$. We postpone the proof of (3-9) to the end of this subsection. Using Lemma 2.3, (3-9), and the concavity and the monotonicity of ω_B , we may bound $-\mathcal{D}_B$ as

$$\begin{aligned} -\mathcal{D}_B(\xi) &\leq \frac{1}{A} \int_{\xi/2}^{\infty} (\omega_B(\xi + 2\eta) - \omega_B(2\eta - \xi) - \omega_B(2\xi) - (2 - c_\alpha)\omega_B(\xi)) \frac{m(2\eta)}{\eta} d\eta \\ &\leq -\frac{2 - c_\alpha}{A} \omega_B(\xi) \int_{\xi/2}^{\xi} \frac{m(2\eta)}{\eta} d\eta \leq -\frac{2 - c_\alpha}{A} \omega_B(\xi) m(2\xi). \end{aligned} \quad (3-10)$$

We emphasize that, for the upper bound (3-10), only the contribution from $\eta \in (\xi/2, \xi)$ was used.

On the other hand, integrating by parts, the contribution from $\tilde{\Omega}_B$ may be rewritten as

$$\frac{\tilde{\Omega}_B(\xi)}{A} = \omega_B(\xi) + \xi \int_{\xi}^{\infty} \frac{\omega_B(\eta)}{\eta^2} d\eta = 2\omega_B(\xi) + \gamma \xi \int_{\xi}^{\infty} \frac{m(2\eta)}{\eta} d\eta.$$

Using (1-3), we may bound

$$\int_{\xi}^{\infty} \frac{m(2\eta)}{\eta} d\eta \leq \xi^\alpha m(2\xi) \int_{\xi}^{\infty} \frac{1}{\eta^{1+\alpha}} d\eta \leq \frac{m(2\xi)}{\alpha},$$

where $\alpha > 0$ is given. Hence we obtain

$$\frac{\tilde{\Omega}_B(\xi)}{A} \leq 2\omega_B(\xi) + \frac{\gamma \xi m(2\xi)}{\alpha}. \quad (3-11)$$

Now, for $\delta(B) \leq \xi \leq 2\delta(B)$, by (1-3) we have

$$\frac{\gamma \xi m(2\xi)}{\alpha} \leq \frac{\gamma}{\alpha} (2\delta(B))^{1-\alpha} \delta(B)^\alpha m(\delta(B)) \leq \frac{2\gamma}{\alpha} \delta(B) \frac{B}{\kappa} \leq \frac{B\delta(B)}{2} \leq \omega_B(\delta(B)) \leq \omega_B(\xi) \quad (3-12)$$

by (3-6), if γ is small. On the other hand, for $\xi > 2\delta(B)$ we have $\xi - \delta(B) \geq \xi/2$ and therefore

$$\omega_B(\xi) \geq \gamma \int_{\delta(B)}^{\xi} m(2\eta) d\eta \geq \gamma m(2\xi)(\xi - \delta(B)) \geq \frac{\gamma \xi m(2\xi)}{2}.$$

Combining the above estimates with (3-11) leads to a bound

$$\tilde{\Omega}_B(\xi) \leq A \left(2 + \frac{2}{\alpha} \right) \omega_B(\xi). \quad (3-13)$$

From (3-4) and the bounds (3-10) and (3-13), we hence obtain

$$\widetilde{\Omega}_B(\xi)\omega'_B(\xi) - \mathfrak{D}_B(\xi) \leq \left(A\gamma \frac{2\alpha + 2}{\alpha} - \frac{2 - c_\alpha}{A} \right) \omega_B(\xi)m(2\xi) < 0$$

for all $\xi \geq \delta(B)$, if we set γ small enough, depending only on A , C , α , and c_α .

Proof of estimate (3-9). To verify (3-9) for $\delta(B) \leq \xi \leq 2\delta(B)$ is straightforward, since, by the mean value theorem and (1-3), similarly to (3-12), we obtain

$$\begin{aligned} \omega_B(2\xi) &\leq \omega_B(\xi) + \xi\omega'_B(\xi) = \omega_B(\xi) + \gamma\xi m(2\xi) \\ &\leq \omega_B(\xi) + \frac{2\gamma}{\kappa} B\delta(B) \leq \omega_B(\xi) + (c_\alpha - 1)\omega_B(\delta(B)), \end{aligned}$$

by choosing γ small enough.

Now, for $\xi > 2\delta(B)$, by (3-4) and (3-6) we have

$$\begin{aligned} c_\alpha\omega_B(\xi) - \omega_B(2\xi) &= (c_\alpha - 1)\omega_B(\delta(B)) + (c_\alpha - 1)\gamma \int_{\delta(B)}^{\xi} m(2\eta) d\eta - \gamma \int_{\xi}^{2\xi} m(2\eta) d\eta \\ &\geq (c_\alpha - 1) \frac{B\delta(B)}{2} - \gamma \int_{2\xi - \delta(B)}^{2\xi} m(2\eta) d\eta + \gamma \int_{\delta(B)}^{\xi} ((c_\alpha - 1)m(2\eta) - m(2\eta + 2\xi - 2\delta(B))) d\eta. \end{aligned}$$

We next note that for $\xi \geq 2\delta(B)$, due to the monotonicity of m , we have

$$\gamma \int_{2\xi - \delta(B)}^{2\xi} m(2\eta) d\eta \leq \gamma\delta(B)m(\delta(B)) = \frac{\gamma}{\kappa} B\delta(B) \leq (c_\alpha - 1) \frac{B\delta(B)}{2}$$

by letting γ be small enough. We next verify that

$$(c_\alpha - 1)m(2\eta) \geq m(2\eta + 2\xi - 2\delta(B))$$

holds for all $\eta \in (\delta(B), \xi)$. Using (1-3) and recalling that $c_\alpha = 1 + (3/2)^{-\alpha}$, the above inequality follows once we check that

$$(3/2)^{-\alpha}(2\eta + 2\xi - 2\delta(B))^\alpha \geq (2\eta)^\alpha$$

holds for all $\eta \in (\delta(B), \xi)$. But since $\xi > 2\delta(B)$, we have

$$\frac{\eta + \xi - \delta(B)}{\eta} \geq 1 + \frac{\xi - \delta(B)}{\xi} \geq \frac{3}{2}. \quad \square$$

The case $0 < \xi \leq \delta(B)$. For small values of ξ , we prefer to bound the contribution from the advective term using Ω_B instead of $\widetilde{\Omega}_B + \Omega_B^\perp$. It is sufficient to prove that

$$\Omega_B(\xi)\omega'_B(\xi) - \mathfrak{D}_B(\xi) < 0.$$

Using the concavity of ω_B and the mean value theorem, we may estimate

$$-\mathfrak{D}_B(\xi) \leq \frac{1}{A} \int_0^{\xi/2} (\omega_B(\xi + 2\eta) + \omega_B(\xi - 2\eta) - 2\omega_B(\xi)) \frac{m(2\eta)}{\eta} d\eta \leq \frac{C}{A} \omega_B''(\xi) \int_0^{\xi/2} \eta m(2\eta) d\eta.$$

From (1-3) we obtain $\eta^\alpha m(2\eta) \geq (\xi/2)^\alpha m(\xi)$ for $\eta \in (0, \xi/2)$. Since $\omega_B''(\xi) < 0$, we may further bound

$$-\mathcal{D}_B(\xi) \leq \frac{C}{A} \omega_B''(\xi) \xi^\alpha m(\xi) \int_0^{\xi/2} \eta^{1-\alpha} d\eta \leq \frac{C}{A} \omega_B''(\xi) \xi^2 m(\xi). \quad (3-14)$$

The contribution from the advecting velocity is bounded as

$$\begin{aligned} \frac{\Omega_B(\xi)}{A} &= \int_0^\xi \frac{\omega_B(\eta)}{\eta} d\eta + \xi \int_\xi^{\delta(B)} \frac{\omega_B(\eta)}{\eta^2} d\eta + \xi \int_{\delta(B)}^\infty \frac{\omega_B(\eta)}{\eta^2} d\eta \\ &\leq B\xi + B\xi \ln \frac{\delta(B)}{\xi} + \xi \left(\frac{\omega_B(\delta(B))}{\delta(B)} + \gamma \int_{\delta(B)}^\infty \frac{m(2\eta)}{\eta} d\eta \right). \end{aligned} \quad (3-15)$$

Here we used that $\omega_B(\eta) \leq B\eta$ for $\eta \in (0, \delta(B))$. Using (1-2)–(1-3) and (3-2), we bound

$$\int_{\delta(B)}^\infty \frac{m(2\eta)}{\eta} d\eta \leq \frac{m(2\delta(B))}{\alpha} \leq \frac{B}{\alpha\kappa} \leq \frac{B}{\gamma}$$

for $\gamma \leq \alpha\kappa$. Therefore, (3-15) gives

$$\Omega_B(\xi) \leq AB\xi \left(3 + \log \frac{\delta(B)}{\xi} \right). \quad (3-16)$$

From (3-3) and the bounds (3-14) and (3-16), we obtain

$$\begin{aligned} \Omega_B(\xi) \omega_B'(\xi) - \mathcal{D}_B(\xi) &\leq AB^2\xi \left(3 + \log \frac{\delta(B)}{\xi} \right) + \frac{C}{A} \xi^2 m(\xi) \omega_B''(\xi) \\ &\leq AB^2\xi \left(3 + \log \frac{\delta(B)}{\xi} \right) \left(1 - \frac{C}{2A^2 C_{\alpha\kappa}} \right) < 0 \end{aligned} \quad (3-17)$$

for any $\xi \in (0, \delta(B))$, if we choose κ small enough. Here we used the explicit expression of ω_B'' for small ξ . Note that the choice of κ is independent of γ and B , which is essential in order to avoid a circular argument. This concludes the proof of Theorem 1.1. \square

4. Global regularity vs. finite time blow-up for slightly supercritical Burgers

In this section we prove Theorems 1.2 (global regularity) and 1.3 (finite time blow-up).

Proof of Theorem 1.2. Due to evident similarities to the SQG proof given in Section 3 above, we only sketch the necessary modifications. See also [Kiselev et al. 2008] for more details.

First, we note that a modulus of continuity ω_B is preserved by (1-11) if

$$\omega_B(\xi) \omega_B'(\xi) - \mathcal{D}_B(\xi) < 0, \quad (4-1)$$

where \mathcal{D}_B is defined as by (2-6), with ω replaced by ω_B . We will consider exactly the same family of moduli of continuity ω_B as in the SQG case, defined via (3-3)–(3-4).

We need to verify that (4-1) holds for any $\xi > 0$. In the case $\xi \geq \delta(B)$, by using (3-10), we have

$$\omega_B(\xi) \omega_B'(\xi) - \mathcal{D}_B(\xi) \leq \omega_B(\xi) \gamma m(2\xi) - \frac{C}{4} \omega_B(\xi) m(2\xi) < 0 \quad (4-2)$$

if $\gamma \leq C/8$. On the other hand, for $\xi \in (0, \delta(B))$, we use (3-14) and obtain

$$\begin{aligned} \omega_B(\xi)\omega'_B(\xi) - \mathcal{D}_B(\xi) &\leq B\xi\omega'_B(\xi) + C\xi^2m(\xi)\omega''_B(\xi) \\ &\leq B^2\xi - \frac{CB^2\xi}{2C_\alpha\kappa} \left(3 + \ln \frac{\delta(B)}{\xi} \right) \leq B^2\xi \left(1 - \frac{3C}{2C_\alpha\kappa} \right) < 0 \end{aligned} \quad (4-3)$$

if $\kappa \leq 3C/(2C_\alpha)$. This concludes the proof of Theorem 1.2. \square

Proof of Theorem 1.3. The proof will use ideas from [Dong et al. 2009], which builds an appropriate Lyapunov functional to show blow-up. Throughout this section we assume that (1-14) holds, that is,

$$\int_0^1 m(r) dr = A < \infty. \quad (4-4)$$

Let $\theta_0 \in C^\infty$ be periodic and odd, with $\theta_0(0) = 0$. For simplicity we may take θ_0 to be $\mathbb{T} = [-\pi, \pi]$ periodic, and consider that $r_0 = 1$ in (1-2). It is clear that the proof carries over for any period length and for any value of $r_0 > 0$, with obvious modifications. Assume the solution $\theta(x, t)$ of (1-11) corresponding to this initial data lies in $C(0, T; W^{1,\infty})$ for some $T > 0$, and is hence C^∞ smooth on $[0, T]$. The Burgers equation preserves oddness of a smooth solution so that we have $\theta(0, t) = 0$ for $t \in [0, T]$, and also $\theta(x, t) = -\theta(-x, t)$ for all $x \in \mathbb{T}$ and $t \in [0, T]$.

Let $w(x)$ be defined as the *odd* function with

$$w(x) = 1 - x \quad \text{for } x \in (0, 1), \quad (4-5)$$

$$w(x) = 0 \quad \text{for } x \geq 1. \quad (4-6)$$

Associated to this function w we define the Lyapunov functional

$$L(t) = \int_0^\infty \theta(x, t)w(x) dx = \int_0^1 \theta(x, t)w(x) dx. \quad (4-7)$$

Then, due to the maximum principle $\|\theta(\cdot, t)\|_{L^\infty} \leq \|\theta_0\|_{L^\infty}$, which holds on $[0, T]$, and the definition of $w(x)$, we have

$$L(t) \leq \|\theta_0\|_{L^\infty} \quad (4-8)$$

for all $t \in [0, T]$. We will next show, using our assumption that $\theta \in C(0, T; W^{1,\infty})$, that if T is sufficiently large, the bound (4-8) is violated. This shows that our assumption has been wrong, and θ has finite time blow-up in the $W^{1,\infty}$ norm, concluding the proof of Theorem 1.3.

To proceed, we first need the following lemma.

Lemma 4.1. *If \mathcal{L} is a diffusive operator defined by (1-5) with m satisfying (4-4) in addition to our usual assumptions and w is given by (4-5),*

$$\int_{\mathbb{R}} |\mathcal{L}w(x)| dx < \infty.$$

Proof. It is sufficient to estimate the integral over positive x , since $\mathcal{L}w(x)$ is odd.

The case $x \geq 1$. Here we have $w(x) = 0$ and w is odd, hence

$$\begin{aligned} \mathcal{L}w(x) &= \int_{\mathbb{R}} (w(x) - w(y)) \frac{m(x-y)}{|x-y|} dy = - \int_0^1 w(y) \frac{m(x-y)}{|x-y|} dy - \int_{-1}^0 w(y) \frac{m(x-y)}{|x-y|} dy \\ &= - \int_0^1 w(y) \frac{m(x-y)}{|x-y|} dy + \int_0^1 w(y) \frac{m(x+y)}{|x+y|} dy \\ &= - \int_0^1 (1-y) \left(\frac{m(x-y)}{|x-y|} - \frac{m(x+y)}{|x+y|} \right) dy. \end{aligned} \quad (4-9)$$

Using the mean value theorem and the monotonicity of m , we estimate

$$\left| \frac{m(x-y)}{|x-y|} - \frac{m(x+y)}{|x+y|} \right| \leq 2y \sup_{r \in [x-y, x+y]} \frac{r|m'(r)| + m(r)}{r^2} \leq 4Cy \frac{m(x-y)}{|x-y|^2}.$$

But the above bound is only convenient when $x \geq 2$, and in this range we obtain

$$\begin{aligned} \int_2^\infty |\mathcal{L}w(x)| dx &\leq 4C \int_2^\infty \int_0^1 (1-y) y \frac{m(x-y)}{|x-y|^2} dy dx \\ &\leq 4C \int_2^\infty \int_0^1 (1-y) y \frac{m(1)}{|x-1|^2} dy dx \leq Cm(1). \end{aligned} \quad (4-10)$$

On the other hand, when $x \in [1, 2]$, it is convenient to work with (4-9) directly. By the monotonicity of m , we have that

$$|\mathcal{L}w(x)| \leq \int_0^1 (1-y) \left(\frac{m(x-y)}{|x-y|} + \frac{m(x+y)}{|x+y|} \right) dy \leq 2 \int_0^1 m(1-y) dy = 2A \quad (4-11)$$

for any $x \in [1, 2]$, by using (4-4). Therefore, $\int_1^2 |\mathcal{L}w(x)| dx \leq 2A$, and by using (4-10), we obtain that

$$\int_1^\infty |\mathcal{L}w(x)| dx \leq 2A + CC_0 =: C_1. \quad (4-12)$$

The case $0 < x < 1$. Here we have $w(x) = 1 - x$, and therefore

$\mathcal{L}w(x)$

$$\begin{aligned} &= \int_{1-x}^\infty (1-x) \frac{m(y)}{y} dy + \int_{-x}^{1-x} y \frac{m(y)}{|y|} dy + \int_{-x-1}^{-x} (2+y) \frac{m(y)}{|y|} dy + \int_{-\infty}^{-x-1} (1-x) \frac{m(y)}{|y|} dy \\ &=: T_1(x) + T_2(x) + T_3(x) + T_4(x). \end{aligned} \quad (4-13)$$

Using condition (1-3), we may easily bound T_1 and T_4 . More precisely, using a change of variables $y \rightarrow -y$ in T_4 , we may write

$$\begin{aligned} |T_1(x) + T_4(x)| &= (1-x) \int_{1-x}^{1+x} \frac{m(y)}{y} dy + 2(1-x) \int_{1+x}^\infty \frac{y^\alpha m(y)}{y^{1+\alpha}} dy \\ &\leq (1-x) \int_{1-x}^{1+x} \frac{m(1-x)}{1-x} dy + 2(1-x)(1+x)^\alpha m(1+x) \int_{1+x}^\infty \frac{1}{y^{1+\alpha}} dy \\ &\leq 2x m(1-x) + \frac{2(1-x)m(1)}{\alpha}, \end{aligned}$$

and therefore

$$\int_0^1 |T_1(x) + T_4(x)| dx \leq 2 \int_0^1 xm(1-x) dx + \frac{2C_0}{\alpha} \int_0^1 (1-x) dx \leq 2A + \frac{C_0}{\alpha}. \quad (4-14)$$

To bound T_2 , we recall that m is even and hence $T_2(x) = \int_x^{1-x} m(y) dy$, which in turn implies

$$\int_0^1 |T_2(x)| dx \leq \int_0^{1/2} \int_x^{1-x} m(y) dy dx + \int_{1/2}^1 \int_{1-x}^x m(y) dy dx \leq \int_0^1 m(y) dy = A. \quad (4-15)$$

Lastly, due to the monotonicity of m , we have that

$$\begin{aligned} |T_3(x)| &\leq 2 \int_x^1 \frac{m(y)}{y} dy + 2 \int_1^{x+1} \frac{m(y)}{y} dy + \int_x^{x+1} m(y) dy \\ &\leq 2 \int_x^1 \frac{m(y)}{y} dy + 2 \int_1^2 \frac{m(1)}{y} dy + \int_x^1 m(y) dy + \int_1^2 m(y) dy \\ &\leq 2 \int_x^1 \frac{m(y)}{y} dy + 2m(1) \log 2 + A + m(1), \end{aligned}$$

and therefore

$$\begin{aligned} \int_0^1 |T_3(x)| dx &\leq 2 \int_0^1 \int_x^1 \frac{m(y)}{y} dy dx + 3m(1) + A \\ &\leq 2 \int_0^1 \int_0^y \frac{m(y)}{y} dx dy + 3C_0 + A \leq 3(C_0 + A). \end{aligned} \quad (4-16)$$

Summarizing (4-14), (4-15), and (4-16), we obtain that

$$\int_0^1 |\mathcal{L}w(x)| dx \leq 6A + 3C_0 + \frac{C_0}{\alpha} =: C_2. \quad (4-17)$$

This concludes the proof. \square

Coming back to our Lyapunov functional $L(t)$, using the evolution (1-11) and integrating by parts, we obtain

$$\begin{aligned} \frac{d}{dt} L(t) &= \int_0^\infty \theta_t(x, t) w(x) dx = \int_0^\infty \left(\partial_x \frac{\theta(x, t)^2}{2} - \mathcal{L}\theta(x, t) \right) w(x) dx \\ &= -\frac{1}{2} \int_0^1 \theta(x, t)^2 w_x(x) dx - \int_0^\infty \theta(x, t) \mathcal{L}w(x) dx. \end{aligned} \quad (4-18)$$

Here we employed the identity $\int_0^\infty \mathcal{L}\theta(\cdot, t) w = \int_0^\infty \theta(\cdot, t) \mathcal{L}w$. This equality can be derived by using the oddness of *both* θ and w , the evenness of m , and Lemma 4.1, ensuring $\mathcal{L}w(x) \in L^1$ (see [Dong et al. 2009, (2.8)] for more details). Also, the integration by parts in the first term of (4-18) is justified, since, by our assumption, θ vanishes as $C|x|$ when $x \rightarrow 0$, with $C = \sup_{[0, T]} \|\nabla\theta(\cdot, t)\|_{L^\infty}$.

Now, since $w_x = -1$ for $0 < x < 1$, and using the Cauchy–Schwartz inequality, we obtain

$$\begin{aligned} L(t)^2 &= \left(\int_0^1 \theta(x, t) w(x) dx \right)^2 \leq \int_0^1 \theta(x, t)^2 dx \int_0^1 w(x)^2 dx \\ &= \frac{1}{3} \int_0^1 \theta(x, t)^2 dx = -\frac{1}{3} \int_0^1 \theta(x, t)^2 w_x(x) dx. \end{aligned}$$

Therefore, by (4-18) on $[0, T]$ we have

$$\frac{d}{dt} L(t) \geq \frac{3}{2} L(t)^2 - \int_0^\infty |\theta(x, t)| |\mathcal{L}w(x)| dx \geq L(t)^2 - \|\theta_0\|_{L^\infty} \int_0^\infty |\mathcal{L}w(x)| dx. \quad (4-19)$$

By Lemma 4.1, we then have

$$\frac{d}{dt} L(t) \geq L(t)^2 - (C_1 + C_2) \|\theta_0\|_{L^\infty}. \quad (4-20)$$

But (4-20) implies that $L(t)$ blows up in finite time provided that

$$0 < L(0)^2 - (C_1 + C_2) \|\theta_0\|_{L^\infty} = \left(\int_0^1 (1-x)\theta_0(x) dx \right)^2 - (C_1 + C_2) \|\theta_0\|_{L^\infty}.$$

It is easy to design initial data satisfying this condition, and thus leading to finite time blow-up. This completes the proof of Theorem 1.3. \square

5. Global regularity with dissipative Fourier multiplier

In this section we establish a connection between the global regularity results obtained for (1-8)–(1-9) when the dissipative nonlocal operators \mathcal{L} are replaced by dissipative Fourier multiplier operators, an approach that has been more standard in fluid dynamics. More precisely, we will replace $\mathcal{L}\theta(x)$ by

$$(P(\zeta)\hat{\theta}(\zeta))^\vee(x)$$

for a nice enough radially symmetric Fourier multiplier symbol P , and consider the global regularity for the slightly supercritical SQG equation

$$\partial_t \theta + u \cdot \nabla \theta + (P\hat{\theta})^\vee = 0, \quad (5-1)$$

$$u = \nabla^\perp \Lambda^{-1} \theta. \quad (5-2)$$

The setting can be either \mathbb{T}^2 or \mathbb{R}^2 with decaying initial data. In the latter case, an additional argument is needed for Lemma 3.2 to remain valid due to lack of compactness; see [Dong and Du 2008]. We will focus on the periodic case. Note that working on \mathbb{T}^d is equivalent to working on \mathbb{R}^d with $\theta(x, t)$ extended periodically. We will henceforth pursue this strategy, thinking of the Fourier multiplier P and its corresponding convolution kernel K in \mathbb{R}^d .

Intuitively, the Fourier multiplier corresponds to a nonlocal operator \mathcal{L} as defined in (1-5), with $m(y)$ that is comparable to $P(1/|y|)$. We make this connection more precise in the following two lemmas.

Lemma 5.1 (dissipative operator associated to Fourier multiplier — upper bound). *Let $P(\zeta) = P(|\zeta|)$ be a radially symmetric function which is smooth away from zero, nonnegative, nondecreasing, with $P(0) = 0$ and $P(\zeta) \rightarrow \infty$ as $|\zeta| \rightarrow \infty$. In addition assume the following:*

(i) *P satisfies the doubling condition:*

$$P(2|\zeta|) \leq c_D P(|\zeta|) \quad (5-3)$$

for some doubling constant $c_D \geq 1$.

(ii) *P satisfies the Hörmander–Mikhlin condition:*

$$|\partial_\zeta^k P(\zeta)| |\zeta|^{-|k|} \leq c_H P(\zeta) \quad (5-4)$$

for some constant $c_H \geq 1$, and for all multi-indices $k \in \mathbb{Z}^d$ with $|k| \leq N$, with N depending only on the dimension d and on the doubling constant c_D .

(iii) *P has subquadratic growth at ∞ , that is,*

$$\int_0^1 P(|\zeta|^{-1}) |\zeta|^d |\zeta|^{-1} d|\zeta| < \infty. \quad (5-5)$$

Then the Fourier multiplier operator with symbol $P(\zeta)$ is given as a nonlocal operator defined as the principal value of

$$(P(\cdot) \hat{\theta}(\cdot))^\vee(x) = \int_{\mathbb{R}^d} (\theta(x) - \theta(x+y)) K(y) dy \quad (5-6)$$

and the radially symmetric kernel K satisfying

$$|K(y)| \leq C |y|^{-d} P(|y|^{-1}) \quad (5-7)$$

for all $y \neq 0$, for some positive constant $C > 0$. Similarly $|\nabla K(y)| \leq C |y|^{-d-1} P(|y|^{-1})$ for $y \neq 0$.

Proof. As in Littlewood–Paley theory, consider smooth, radially symmetric functions φ , supported on $1/2 \leq |\zeta| \leq 2$, such that

$$1 = \sum_{j \in \mathbb{Z}} \varphi(2^{-j} \zeta) \quad (5-8)$$

holds for $\zeta \in \mathbb{R}^d \setminus \{0\}$. We write $\varphi_j(\zeta) = \varphi(2^{-j} \zeta)$ and note that $P(\zeta) \varphi_j(\zeta)$ is smooth and compactly supported with $P(0) \varphi_j(0) = 0$. Hence

$$K_j(y) = - \int_{\mathbb{R}^d} P(\zeta) \varphi_j(\zeta) e^{iy \cdot \zeta} d\zeta$$

is a family of L^1 kernels, which are smooth at the origin, radially symmetric, and have zero mean on \mathbb{R}^d . Thus we may write (in order to avoid principal value integrals we use double differences)

$$(P(\cdot) \varphi_j(\cdot) \hat{\theta}(\cdot))^\vee(x) = \int_{\mathbb{R}^d} K_j(y) (2\theta(x) - \theta(x-y) - \theta(x+y)) dy.$$

For $y \neq 0$, let $j_0 = [\log_2 |y|^{-1}]$ and fix $N > d + \log_2 c_D$ to be an even integer. By (5-3)–(5-4) we have

$$\begin{aligned}
\sum_j |K_j(y)| &= \sum_{j < j_0} \|K_j\|_{L^\infty} + \sum_{j \geq j_0} \left| \int_{\mathbb{R}^d} P(\zeta) \varphi_j(\zeta) e^{i\zeta \cdot y} d\zeta \right| \\
&\leq \sum_{j < j_0} \|\hat{K}_j\|_{L^1} + \sum_{j \geq j_0} |y|^{-N} \left| \int_{\mathbb{R}^d} (-\Delta)^{N/2} (P(\zeta) \varphi_j(\zeta)) e^{i\zeta \cdot y} d\zeta \right| \\
&\leq \sum_{j < j_0} \int_{\mathbb{R}^d} P(\zeta) \varphi(2^{-j}\zeta) d\zeta + C|y|^{-N} \sum_{j \geq j_0} 2^{-jN} \int_{2^{j-1} \leq |\zeta| \leq 2^{j+1}} P(\zeta) d\zeta \\
&\leq P(2^{j_0}) \sum_{j < j_0} \int_{\mathbb{R}^d} \varphi(2^{-j}\zeta) d\zeta + C|y|^{-N} \sum_{j \geq j_0} 2^{-j(N-d)} P(2^{j+1}) \\
&\leq CP(2^{j_0}) \sum_{j < j_0} 2^{jd} + C|y|^{-N} 2^{-j_0(N-d)} \sum_{j \geq j_0} 2^{-(j-j_0)(N-d)} P(2^{j_0}) c_D^{j-j_0} \\
&\leq CP(|y|^{-1})|y|^{-d} + CP(|y|^{-1})|y|^{-d} \sum_{j \geq j_0} 2^{-(j-j_0)(N-d-\log_2 c_D)},
\end{aligned}$$

which shows that the sum $K(y) = \sum_j K_j(y)$ converges absolutely for all $y \neq 0$, and proves (5-7). The purpose of condition (5-5) is now evident. For a smooth function θ (say at least of class C^2), in order to make sense of the integral

$$\int_{|y| \leq 1} K(y) (\theta(x-y) + \theta(x+y) - 2\theta(x)) dy,$$

in view of (5-7), we need to assume that $\int_{|y| \leq 1} P(|y|^{-1})|y|^{-d+2} dy < +\infty$, which is equivalent to (5-5). The bound for $|\nabla K|(y)$ is analogous, and we omit further details. \square

Lemma 5.2 (dissipative operator associated to Fourier multiplier — lower bound). *Let the Fourier multiplier P and its associated kernel K be as in Lemma 5.1. Assume additionally that*

(v) P satisfies

$$(-\Delta)^{(d+2)/2} P(\zeta) \geq c_H^{-1} P(\zeta) |\zeta|^{-d-2} \quad (5-9)$$

for all $|\zeta|$ sufficiently large (say larger than $c_0 > 0$), for some constant $c_H \geq 1$.

Then the kernel K corresponding to P (see (5-6)) may be bounded from below as

$$K(y) \geq C^{-1} |y|^{-d} P(|y|^{-1}) \quad (5-10)$$

for all sufficiently small $|y|$, for some sufficiently large constant $C > 0$.

Proof. From our assumptions, the symbol P is a C^{d+2} smooth function except perhaps at the origin. Without loss of generality we can assume P to be smooth at the origin as well. Otherwise, we can write $P = \tilde{P} + R$ where \tilde{P} is a C^{d+2} function everywhere with $\tilde{P}(\zeta) = P(\zeta)$ for all $|\zeta| > c_0$, $\tilde{P}(0) = 0$, and $(-\Delta)^{N/2} \tilde{P}$ is bounded in \mathbb{R}^d . The remainder R is a bounded compactly supported function with $R(0) = 0$. Therefore, the Fourier multiplier operator with symbol P is the sum of the operators with multipliers \tilde{P}

and R . For \tilde{P} we apply the proof below and obtain a kernel satisfying (5-10), and for the remainder R we have $(R\hat{\theta})^\vee = R^\vee * \theta$ and R^\vee is a bounded, mean zero L^1 kernel. Thus, adding R^\vee will not destroy the estimate (5-10) for small enough y .

If P is smooth near $\zeta = 0$, we have that $Q(\zeta) = (-\Delta)^{(d+2)/2} P(\zeta) \in L^1(\mathbb{R}^d)$. Indeed, $\int_{|\zeta| \leq 1} |Q(\zeta)| d\zeta$ is finite since P is smooth, while by (5-4) and (5-5) we have

$$\int_{|\zeta| \geq 1} |Q(\zeta)| d\zeta \leq c_H \int_{|\zeta| \geq 1} |\zeta|^{-(d+2)} P(\zeta) d\zeta = c_H \int_1^\infty |\zeta|^{-3} P(|\zeta|) d|\zeta| = c_H \int_0^1 r P(r^{-1}) dr < \infty.$$

We may hence define the function M , the inverse Fourier transform of $-Q$, as

$$M(y) = - \int_{\mathbb{R}^d} Q(\zeta) e^{i\zeta \cdot y} d\zeta = - \int_{\mathbb{R}^d} Q(\zeta) \cos(y \cdot \zeta) d\zeta,$$

where we have used the fact that Q is radially symmetric and real. Moreover, note that Q has zero mean, since in view of Lemma 5.1 we have the bound $|Q^\vee(x)| \leq |x|^{d+2} |P^\vee(x)| \leq C|x|^2 P(|x|^{-1}) \rightarrow 0$ as $|x| \rightarrow 0$, since P is subquadratic at infinity; cf. (5-5). Thus we may rewrite $M(y)$ as

$$M(y) = \int_{\mathbb{R}^d} Q(\zeta) (1 - \cos(y \cdot \zeta)) d\zeta = \int_{\mathbb{R}^d} Q(\zeta) (1 - \cos(\zeta_1 |y|)) d\zeta \quad (5-11)$$

by using that Q is radially symmetric. In order to appeal to (5-9), we further split

$$M(y) = \int_{|\zeta| \leq c_0} Q(\zeta) (1 - \cos(\zeta_1 |y|)) d\zeta + \int_{|\zeta| > c_0} Q(\zeta) (1 - \cos(\zeta_1 |y|)) d\zeta. \quad (5-12)$$

For all $|y| \leq c_0^{-1}$, the first integral in (5-12) can be estimated from below by $-C_Q |y|^2$, where $C_Q = \int_{|\zeta| \leq c_0} |Q(\zeta)| d\zeta$. Then, using (5-9), for $|y| \leq c_0^{-1}$ we obtain

$$\begin{aligned} M(y) &\geq -C_Q |y|^2 + c_H^{-1} \int_{|\zeta| \geq c_0} |\zeta|^{-(d+2)} P(\zeta) (1 - \cos(\zeta_1 |y|)) d\zeta \\ &\geq -C_Q |y|^2 + c_H^{-1} |y|^2 \int_{|z| \geq c_0 |y|} |z|^{-(d+2)} P(z|y|^{-1}) (1 - \cos(z_1)) dz \\ &\geq -C_Q |y|^2 + c_H^{-1} |y|^2 \int_{2 \geq |z| \geq 1} |z|^{-(d+2)} P(z|y|^{-1}) (1 - \cos(z_1)) dz \\ &\geq -C_Q |y|^2 + c_H^{-1} 2^{-(d+2)} |y|^2 P(|y|^{-1}) \int_{2 \geq |z| \geq 1} (1 - \cos z_1) dz \\ &\geq -C_Q |y|^2 + 2C^{-1} |y|^2 P(|y|^{-1}) \end{aligned} \quad (5-13)$$

for some sufficiently large constant $C > 0$ that depends only on c_H and d . The assumption that $P(\zeta) \rightarrow \infty$ as $|\zeta| \rightarrow \infty$ combined with (5-13) shows that

$$M(y) \geq C^{-1} |y|^2 P(|y|^{-1})$$

holds for all sufficiently small $|y|$.

To conclude, we note that since $\widehat{M} = -Q = -(-\Delta)^{(d+2)/2}P$, we have that $K(y) = -P^\vee(y) = |y|^{-(d+2)}M(y)$ in the sense of tempered distributions, and hence we obtain that, for sufficiently small $|y|$, the bound $K(y) \geq C^{-1}|y|^{-d}P(|y|^{-1})$ holds, concluding the proof of the lemma. \square

Remark 5.3 (examples of symbols P). The conditions (5-3)–(5-5) that were assumed on the symbol P in order to obtain the upper bound for the associated kernel are fairly common assumptions in Fourier analysis. For all symbols of interest to us in this paper, condition (5-9) also naturally holds. The dimension relevant to the SQG equation is $d = 2$. When $P(\zeta) = |\zeta|(\log(|\zeta|))^{-a}$ for sufficiently large $|\zeta|$ and $0 < a \leq 1$, corresponding to (1-4), one may verify that $(-\Delta)^2 P(\zeta)|\zeta|^4/P(\zeta) \rightarrow 1$ as $|\zeta| \rightarrow \infty$, so that we may take $c_H = 2$ in (5-9) if c_0 is sufficiently large. Thus condition (5-9) is not restrictive for the class of symbols we have in mind.

The proof of Theorem 1.4 combines the estimates in Lemmas 5.1 and 5.2 above with the argument given in Section 3. One complication arises due to the fact that (5-10) only holds for small enough $|y|$. In fact, for the class of multipliers P that we consider, positivity of the kernel K is not assured. Because of that, the L^∞ maximum principle is no longer available. However, there is an easy substitute which is sufficiently strong for our purpose.

Lemma 5.4. *Assume that a smooth function $\theta(x, t)$ solves (5-2). Suppose that the kernel $K(y)$ corresponding to the multiplier P via (5-6) satisfies $|K(y)| \leq C|y|^{-d}P(|y|^{-1})$ for all y and $K(y) \geq C^{-1}|y|^{-d}P(|y|^{-1})$ for all $|y| \leq 2\sigma$, where σ, C are positive constants. Then there exists $M_* = M_*(P, \theta_0)$ such that $\|\theta(x, t)\|_{L^\infty} \leq M$ for all $t \geq 0$.*

Proof. Letting $M(t) = \|\theta(\cdot, t)\|_{L^\infty}$, we prove that there exists $M_* \geq M(0)$, sufficiently large, such that $M(t) \leq M_*$ for all $t \geq 0$. If not, then, for any fixed M_* , there exists a $t_* > 0$ such that $M(t_*) = M_*$ and $\partial_t M(t_*) \geq 0$. For this fixed t_* let \bar{x} be a point of maximum for $\theta(\cdot, t_*)$. We have

$$\begin{aligned} \mathcal{L}\theta(\bar{x}) &\geq \int_{\sigma \leq |y| \leq \infty} (\theta(\bar{x}) - \theta(\bar{x} + y))K(y) dy \\ &\geq cM_* \int_{\sigma \leq |y| \leq 2\sigma} \frac{P(|y|^{-1})}{|y|^d} dy - C\|\theta\|_{L^2(\mathbb{T}^d)} \left(\int_{\sigma \leq |y|} \frac{P(|y|^{-1})^2}{|y|^{2d}} dy \right)^{1/2} \\ &\geq cM_*P((2\sigma)^{-1}) - C\|\theta_0\|_{L^2(\mathbb{T}^d)}P(\sigma^{-1})\sigma^{-d/2}. \end{aligned} \quad (5-14)$$

We used that $P \geq 0$ implies $\|\theta(\cdot, t)\|_{L^2(\mathbb{T}^d)} \leq \|\theta_0\|_{L^2(\mathbb{T}^d)}$ in the above calculation. The estimate (5-14) proves that $\partial_t M(t_*)$ must be negative if M_* is large enough, depending only on P (through σ and other constants) and θ_0 . It follows that $M(t)$ will never exceed the larger of these bound or $\|\theta_0\|_{L^\infty}$. \square

Proof of Theorem 1.4. The first two lemmas of this section show that, for the multiplier P satisfying (5-3)–(5-5) and (5-9), we have that $(P\hat{\theta})^\vee(x) = \int (\theta(x) - \theta(x + y))K(y) dy$, with K being radial and smooth away from zero. Moreover, K satisfies $|K(y)| \leq C|y|^{-d}P(|y|^{-1})$ for all y and $K(y) \geq c|y|^{-d}P(|y|^{-1})$ for all $|y| \leq 2\sigma$, where C, c, σ are positive constants depending only on P .

Consider a smooth radially decreasing function $\varphi_0(y)$ that is identically 1 on $|y| \leq \sigma$ and vanishes identically on $|y| \geq 2\sigma$. We decompose

$$K(y) = K(y)\varphi_0(y) + K(y)(1 - \varphi_0(y)) =: K_1(y) + K_2(y),$$

so that

$$(P\hat{\theta})^\vee(x) = \int_{\mathbb{R}^d} (\theta(x) - \theta(x+y))K_1(y) + \int_{\mathbb{R}^d} (\theta(x) - \theta(x+y))K_2(y) =: \mathcal{L}_1\theta(x) + \mathcal{L}_2\theta(x).$$

The nonlocal operator \mathcal{L}_1 is of type (1-7), since by letting

$$m(r) = C^{-1}P(r^{-1})\varphi_0(r), \quad (5-15)$$

we have that

$$K_1(y) \geq m(|y|)|y|^{-d}$$

for all y and some $C > 0$. It is clear that the above defined m satisfies properties (1-2)–(1-3) and (1-10) in view of assumptions (1-15)–(1-17) imposed on P . Therefore, for \mathcal{L}_1 , we will be able to directly use the estimate in Lemma 2.3, which relies only on lower bounds for the kernel associated to \mathcal{L}_1 .

On the other hand, we observe that $K_2 \in L^1(\mathbb{R}^d)$ since $K_2(y) = 0$ for $y \leq \sigma$, and we have

$$|K_2(y)| \leq |K(y)| \leq CP(|y|^{-1})|y|^{-d} \leq \sigma^\alpha P(\sigma^{-1})|y|^{-d-\alpha}$$

for any $|y| \geq \sigma$, by using (1-16). Let us fix the constant $C_2 = \|K_2\|_{L^1(\mathbb{R}^d)}$. Then if $\theta(\cdot, t)$ obeys the modulus of continuity $\omega(\xi)$, it is clear that

$$|\mathcal{L}_2\theta(x, t) - \mathcal{L}_2\theta(y, t)| \leq 2C_2 \min\{\omega(\xi), M_*\}, \quad (5-16)$$

where M_* is the L^∞ norm bound from Lemma 5.4, holds for all $x, y \in \mathbb{R}^d$, where $|x - y| = \xi$.

Now the argument of Section 3 goes through with minor changes. We provide an outline of the argument to verify this. First, similarly to (3-7) we may prove that for B large enough (now depending on σ as well) we have $\omega_B(\sigma) \geq 3M_* \geq 3\|\theta(\cdot, t)\|_{L^\infty}$, so that the modulus of continuity can only be broken at values of $\xi \in (0, \sigma)$. Let \mathcal{D}_B and \mathcal{D}_B^\perp be the bounds obtained from the dissipative operator \mathcal{L}_1 via Lemma 2.3. Note that the only contribution from the integral defining \mathcal{D}_B that is used in the estimates is for $\eta \in (0, \xi)$ (see (3-10) and (3-14)), and for us $\xi < \sigma$ so all the bounds on the dissipation given in the proof of Theorem 1.1 require no modification. Therefore, provided κ and γ are chosen sufficiently small, we have

$$\min\{\Omega_B(\xi), \tilde{\Omega}_B(\xi) + \Omega_B^\perp(\xi)\}\omega'_B(\xi) - (\frac{1}{2}\mathcal{D}_B(\xi) + \mathcal{D}_B^\perp(\xi)) < 0$$

for any $B \geq 1$ and $\xi \in (0, \sigma)$, exactly as in the proof of Theorem 1.1. The proof is hence completed once we show that the contribution of \mathcal{L}_2 is controlled:

$$2C_2 \min\{\omega_B(\xi), 2\|\theta\|_{L^\infty}\} \leq \frac{1}{2}\mathcal{D}_B(\xi) \quad (5-17)$$

for any $\xi \in (0, \sigma)$ and any $B \geq 1$. The range $\xi \in (0, \delta(B))$ is clear, since here $\omega_B(\xi) \leq B\xi$ and by (3-14) we have

$$\mathcal{D}_B(\xi) \geq -\frac{C}{A}\xi^2 m(\xi)\omega_B''(\xi) = \frac{CB^2}{2\kappa C_\alpha A}\xi \left(3 + \ln \frac{\delta(B)}{\xi}\right) \geq \frac{3C}{2\kappa C_\alpha A}B\xi \geq 4C_2 B\xi \geq 4C_2 \omega_B(\xi)$$

by letting κ be small enough (independent of $B \geq 1$).

We next consider the range $\xi \in (\delta(B), \sigma)$. In view of (3-10), we have $\mathcal{D}_B(\xi) \geq C\omega_B(\xi)m(2\xi)$, where $C = (2 - c_\alpha)/A$. Since $P(\zeta) \rightarrow \infty$ as $|\zeta| \rightarrow \infty$, we have that $Cm(2\xi) \geq 4C_2$, for all $\xi \in (\delta(B), \kappa)$, for some $\kappa > 0$. If $\kappa \geq \sigma$, the proof is completed, but this cannot be guaranteed, so we have to also consider the case $\kappa < \sigma$. For $\xi \in (\kappa, \sigma)$, we have

$$\mathcal{D}_B(\xi) \geq C\omega_B(\xi)m(2\xi) \geq C\omega_B(\kappa)m(\sigma) \geq Cm(\sigma)\gamma \int_{\delta(B)}^{\kappa} m(2\eta) d\eta. \quad (5-18)$$

By making B large enough, we can ensure that the right hand side of (5-18) is larger than $2M_*$. \square

6. Global well-posedness for a two-dimensional Euler-type equation with more singular velocity

In this section we address the issue of global regularity for the *inviscid* active scalar equation

$$\partial_t \theta - u \cdot \nabla \theta = 0, \quad (6-1)$$

$$u = \nabla^\perp \Lambda^{-2} P(\Lambda) \theta, \quad (6-2)$$

where the multiplier $P(\zeta) = P(|\zeta|)$ is a radially symmetric function which is smooth, nondecreasing, with $P(0) = 0$ and $P(\zeta) \rightarrow \infty$ as $|\zeta| \rightarrow \infty$. In addition, we assume that P satisfies a doubling property

$$P(2|\zeta|) \leq c_D P(|\zeta|) \quad (6-3)$$

for some doubling constant $c_D \geq 1$,

$$|\zeta|^{-\alpha} P(|\zeta|) \text{ is nonincreasing} \quad (6-4)$$

for some $\alpha \in (0, 1)$, and a Hörmander–Mikhlin type condition

$$|\partial_\zeta^k P(\zeta)| |\zeta|^{|k|} \leq c_H P(\zeta) \quad (6-5)$$

holds for some constant $c_H \geq 1$ and for all multi-indices $k \in \mathbb{Z}^d$ with $|k| \leq N$, where N depends only on the dimension d and on the doubling constant c_D . Condition (6-4) is quite natural in view of (6-7) below, while conditions (6-3) and (6-5) are standard in Fourier analysis. We remark that, while finalizing this paper, we learned of [Elgindi 2014], which proves a result very similar to the one proved in this section under slightly less restrictive assumptions on P .

Using the technique of Lemma 5.1, one may show using (6-3) and (6-5) that the convolution kernel K corresponding to the operator $\nabla^\perp \Lambda^{-2} P(\Lambda)$, that is, to the Fourier multiplier $i\zeta^\perp |\zeta|^{-2} P(|\zeta|)$, satisfies the estimates

$$|K(x)| \leq C|x|^{-d+1} P(|x|^{-1}), \quad |\nabla K(x)| \leq C|x|^{-d} P(|x|^{-1}), \quad |\nabla \Delta K(x)| \leq C|x|^{-d-2} P(|x|^{-1}) \quad (6-6)$$

for all $x \neq 0$. Moreover, we note that K integrates to 0 around the unit sphere, and hence convolution with K annihilates constants.

The study of Euler equations with more singular velocities, (6-1)–(6-2), was recently initiated by Chae, Constantin, and Wu [Chae et al. 2011]. They prove the global regularity for the loglog-Euler equation; namely, they prove global regularity in the case that arises when $P(\zeta) = [\ln(1 + \ln(1 + |\zeta|^2))]^\gamma$, for $\gamma \in [0, 1]$. Their approach relies on estimates for the Fourier localized gradient of the velocity for a particular class of symbols. Our aim here is to provide a proof of global regularity for a slightly more general class of symbols P , via the modulus of continuity method. The main result of this section is the following.

Theorem 6.1 (global regularity for the P -Euler equation). *Let P be a smooth radially symmetric function which is smooth and nondecreasing with $P(0) = 0$ and $P(\zeta) \rightarrow \infty$ as $|\zeta| \rightarrow \infty$ and satisfies assumptions (6-3)–(6-5). If θ_0 is periodic and smooth and we assume that*

$$\int_1^M \frac{dr}{r \ln(2r) P(r)} \rightarrow \infty, \quad \text{as } M \rightarrow \infty, \quad (6-7)$$

the P -Euler equation (6-1)–(6-2) has a global in time smooth solution.

Remark 6.2 (integral formulation). In fact, our proof provides a stronger result if we state the constitutive law relating u and θ in terms of an integrodifferential operator instead of a Fourier multiplier

$$u(x) = \int_{\mathbb{R}^d} \theta(x + y) K(y) dy,$$

where K is any kernel which satisfies the hypothesis (6-6) for any function P for which (6-3), (6-4), and (6-7) hold, but not necessarily (6-5).

In the previous sections, we constructed autonomous families of moduli of continuity preserved by the dynamics of the respective equations. In the inviscid case, we will construct a single modulus of continuity and then scale it autonomously. The following lemma makes the above observation precise.

Lemma 6.3 (modulus of continuity under pure transport). *Let u be a Lipschitz vector field and let θ solve the transport equation*

$$\partial_t \theta + u \cdot \nabla \theta = 0. \quad (6-8)$$

If $\theta_0 = \theta(\cdot, 0)$ has some modulus of continuity $\omega(\xi)$, then $\theta(\cdot, t)$ has the modulus of continuity $\omega(B(t)\xi)$, where $B(t)$ is given by

$$B(t) = \exp\left(\int_0^t \|\nabla u(\cdot, s)\|_{L^\infty} ds\right).$$

Equivalently, $B(t)$ solves $B(0) = 1$ and $\dot{B}(t) = \|\nabla u(\cdot, t)\|_{L^\infty} B(t)$.

Proof. The solution to the transport equation can be obtained by following the flow of the vector field backwards. Indeed, $\theta(x, t) = \theta_0(X(t))$ where X solves the ordinary differential equation

$$\dot{X}(s) = u(X(s), t - s), \quad X(0) = x.$$

If $X(t)$ and $Y(t)$ are two such trajectories starting at x and y , respectively, from Grönwall's inequality

$$|X(t) - Y(t)| \leq \exp\left(\int_0^t \|\nabla u(\cdot, s)\|_{L^\infty}\right) |x - y| = B(t)|x - y|.$$

Therefore,

$$|\theta(x, t) - \theta(y, t)| \leq |\theta_0(X(t)) - \theta_0(Y(t))| \leq \omega(B(t)|x - y|), \quad \square$$

which concludes the proof of the lemma.

Proof of Theorem 6.1. Let us consider an initial data θ_0 whose Lipschitz L^∞ and L^2 norms are bounded by an arbitrary constant A . Applying Lemma 6.3 with $\omega(\xi) = A\xi$, we obtain that $\theta(\cdot, t)$ obeys the modulus of continuity $AB(t)\xi$, that is, it is Lipschitz continuous with Lipschitz constant

$$\|\nabla\theta(\cdot, t)\|_{L^\infty} \leq AB(t), \quad (6-9)$$

where $B(0) = 1$ and $\dot{B} = \|\nabla u(\cdot, t)\|_{L^\infty} B(t)$.

By the maximum principle, $\|\theta(\cdot, t)\|_{L^\infty} \leq \|\theta_0\| \leq A$ for any time t . Moreover, since u is divergence-free, $\|\theta(\cdot, t)\|_{L^2} \leq \|\theta_0\|_{L^2} \leq A$ for any time t . In order to combine the last two estimates, we have to estimate the Lipschitz norm of u at time t . Let $\varphi(y)$ be a radially nonincreasing nonnegative function that is constant 1 on $|y| \leq 1/2$ and vanishes for $|y| \geq 1$. For some $r \in (0, 1)$ to be chosen later, we split the integral defining ∇u into three pieces to estimate

$$\begin{aligned} |\nabla u(x)| &= \left| \int_{\mathbb{R}^d} \nabla K(y) \theta(x+y) dy \right| \\ &\leq \int_{\mathbb{R}^d} |\nabla(\varphi(y/r)K(y))| |\theta(x+y) - \theta(x)| dy + \int_{\mathbb{R}^d} |\nabla((1-\varphi(y/r))\varphi(y)K(y))| |\theta(x+y)| dy \\ &\quad + \left| \int_{\mathbb{R}^d} \nabla((1-\varphi(y))K(y)) \theta(x+y) dy \right|. \end{aligned}$$

Using the bounds on K and its derivatives obtained in (6-6) and the fact that θ is Lipschitz with constant given by (6-9), we may further bound

$$\begin{aligned} |\nabla u(x)| &\leq C \int_{|y| \leq r} \frac{P(|y|^{-1})}{|y|^d} |\theta(x+y) - \theta(x)| dy + C \int_{r/2 \leq |y| \leq 1} \frac{P(|y|^{-1})}{|y|^d} |\theta(x+y)| dy \\ &\quad + \int_{\mathbb{R}^d} |(-\Delta)\nabla((1-\varphi(y))K(y))| |(-\Delta)^{-1}\theta(x+y)| dy \quad (6-10) \end{aligned}$$

$$\begin{aligned} &\leq CAB(t) \int_0^r P(\rho^{-1}) d\rho + C\|\theta_0\|_{L^\infty} \int_{r/2}^1 \frac{P(\rho^{-1})}{\rho} d\rho \\ &\quad + C\|(-\Delta)^{-1}\theta\|_{L^\infty(\mathbb{R}^d)} \int_{|y| \geq 1/2} \frac{P(|y|^{-1})}{|y|^{d+2}} dy \\ &\leq CAB(t)rP(r^{-1}) + CAP(r^{-1}) \ln \frac{2}{r} + CAP(2). \quad (6-11) \end{aligned}$$

In the last inequality above, we have additionally used two facts: first, that by (6-4) we have

$$\int_0^r P(\rho^{-1}) d\rho \leq CrP(r^{-1});$$

and second, that since θ is periodic and has zero mean on the torus, we can use the Sobolev inequality and estimate

$$\|(-\Delta)^{-1}\theta\|_{L^\infty(\mathbb{R}^d)} = \|(-\Delta)^{-1}\theta\|_{L^\infty(\mathbb{T}^d)} \leq C\|\theta\|_{L^2(\mathbb{T}^d)} \leq CA.$$

By choosing $r = B(t)^{-1}$ in (6-11), which is allowed since $B(0) = 1$ and $\dot{B} \geq 0$, we arrive at

$$\|\nabla u(\cdot, t)\|_{L^\infty} \leq CA(1 + P(B(t))(1 + \ln 2B(t))).$$

Finally we rewrite the differential equation for $B(t)$ as

$$\dot{B}(t) = \|\nabla u(\cdot, t)\|_{L^\infty} B(t) \leq CA(1 + P(B(t))(1 + \ln 2B(t)))B(t).$$

Clearly this ODE has a global in time solution *if and only if*

$$\int_1^\infty \frac{1}{r \ln(2r) P(r)} dr = \infty$$

holds, which finishes the proof. \square

Appendix: Estimate on the dissipative operator at points of modulus breakdown

Here we prove Lemma 2.3. Our argument parallels that of [Kiselev 2011], but is slightly simpler and more general, as we use the integral representation of the diffusion generator \mathcal{L} instead of generalized Poisson kernels employed in [Kiselev 2011]. We remark that a more general argument that allows one to also handle the Cordoba–Cordoba–Fontelos model has recently been given by Tam Do [2013].

Proof of Lemma 2.3. Due to translation invariance and radial symmetry, we may assume without loss of generality that

$$x = (\xi/2, 0) \quad \text{and} \quad y = (-\xi/2, 0).$$

For a point $(\eta, \nu) \in \mathbb{R}^2$, we write $K(\eta, \nu)$ for the dissipation kernel corresponding to \mathcal{L} . Then we have

$$\begin{aligned} \mathcal{L}\theta\left(\frac{\xi}{2}, 0\right) - \mathcal{L}\theta\left(-\frac{\xi}{2}, 0\right) \\ = \int_{\mathbb{R}} \int_{\mathbb{R}} \left(\theta\left(\frac{\xi}{2}, 0\right) - \theta\left(-\frac{\xi}{2}, 0\right) - \theta\left(\frac{\xi}{2} + \eta, \nu\right) + \theta\left(-\frac{\xi}{2} + \eta, \nu\right) \right) K(\eta, \nu) d\eta d\nu. \end{aligned} \quad (\text{A-1})$$

Note that since θ obeys the modulus of continuity ω , one may bound $\mathcal{L}\theta(\xi/2, 0) - \mathcal{L}\theta(-\xi/2, 0)$ from below by the expression on the right side of (A-1), with $K(\eta, \nu)$ replaced by $m(\sqrt{\eta^2 + \nu^2})(\eta^2 + \nu^2)^{-1}$. We will henceforth write K as a shortcut for the latter expression, and assume without loss of generality

that K is radially nonincreasing and nonnegative, since so is m in (1-6).

$$\begin{aligned}
& \mathcal{L}\theta\left(\frac{\xi}{2}, 0\right) - \mathcal{L}\theta\left(-\frac{\xi}{2}, 0\right) \\
&= \int_{\mathbb{R}} \int_{\mathbb{R}} \left(\omega(\xi) - \theta\left(\frac{\xi}{2} + \eta, v\right) + \theta\left(-\frac{\xi}{2} - \eta, v\right) \right) K(\eta, v) \, d\eta \, dv \\
&= \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} \left(\omega(\xi) - \theta\left(\frac{\xi}{2} + \eta, v\right) + \theta\left(-\frac{\xi}{2} - \eta, v\right) \right) K(\eta, v) \, d\eta \, dv \\
&\quad + \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} \left(\omega(\xi) - \theta\left(-\frac{\xi}{2} - \eta, v\right) + \theta\left(\frac{\xi}{2} + \eta, v\right) \right) K(-\xi - \eta, v) \, d\eta \, dv \\
&= \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} \omega(\xi) (K(\eta, v) + K(-\xi - \eta, v)) \\
&\quad - \left(\theta\left(\frac{\xi}{2} + \eta, v\right) - \theta\left(-\frac{\xi}{2} - \eta, v\right) \right) (K(\eta, v) - K(\xi + \eta, v)) \, d\eta \, dv \\
&= \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} \omega(\xi) (K(\eta, v) + K(-\xi - \eta, v)) - \omega(\xi + 2\eta) (K(\eta, v) - K(\xi + \eta, v)) \, d\eta \, dv \\
&\quad + \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} \omega(\xi) (K(\eta, v) + K(-\xi - \eta, v)) \\
&\quad + \left(\omega(\xi + 2\eta) - \theta\left(\frac{\xi}{2} + \eta, v\right) + \theta\left(-\frac{\xi}{2} - \eta, v\right) \right) (K(\eta, v) - K(\xi + \eta, v)) \, d\eta \, dv \\
&=: T^{\parallel} + T^{\perp}.
\end{aligned}$$

Note that

$$K(\eta, v) - K(\xi + \eta, v) \geq 0$$

for $\eta \geq -\xi/2$ due to the monotonicity of K (or that of its lower bound). Hence, using that θ obeys the modulus of continuity ω , we see that $T^{\perp} \geq 0$. To obtain a useful lower bound for T^{\perp} , we only retain the singular piece centered about $\eta = 0$. Changing variables $\eta + \xi/2 \mapsto \eta$, we have

$$T^{\perp} = \int_{\mathbb{R}} \int_0^{\infty} (\omega(2\eta) - \theta(\eta, v) + \theta(-\eta, v)) \left(K\left(\eta - \frac{\xi}{2}, v\right) - K\left(\eta + \frac{\xi}{2}, v\right) \right) \, d\eta \, dv. \quad (\text{A-2})$$

When $|v| \leq \xi/4$ and $|\eta - \xi/2| \leq \xi/4$, using that m is radially nonincreasing, we have that

$$\begin{aligned}
K(\eta - \xi/2, v) - K(\eta + \xi/2, v) &= \frac{m(\sqrt{(\eta - \xi/2)^2 + v^2})}{(\eta - \xi/2)^2 + v^2} - \frac{m(\sqrt{(\eta + \xi/2)^2 + v^2})}{(\eta + \xi/2)^2 + v^2} \\
&\geq m\left(\sqrt{(\eta - \xi/2)^2 + v^2}\right) \left(\frac{1}{(\eta - \xi/2)^2 + v^2} - \frac{1}{(\eta + \xi/2)^2 + v^2} \right) \\
&\geq m\left(\sqrt{(\eta - \xi/2)^2 + v^2}\right) \frac{1}{2((\eta - \xi/2)^2 + v^2)} = \frac{K(\eta - \xi/2, v)}{2}. \quad (\text{A-3})
\end{aligned}$$

Inserting estimate (A-3) into expression (A-2) and recalling that θ obeys ω , we obtain

$$\begin{aligned} T^\perp &\geq \frac{1}{2} \int_{-\xi/4}^{\xi/4} \int_{\xi/4}^{3\xi/4} (\omega(2\eta) - \theta(\eta, v) + \theta(-\eta, v)) K\left(\eta - \frac{\xi}{2}, v\right) d\eta dv \\ &= \frac{1}{2} \int_0^{\xi/4} \int_{\xi/4}^{3\xi/4} (2\omega(2\eta) - \theta(\eta, v) + \theta(-\eta, v) - \theta(\eta, -v) + \theta(-\eta, -v)) K\left(\eta - \frac{\xi}{2}, v\right) d\eta dv = \frac{\mathfrak{D}^\perp}{2}. \end{aligned}$$

On the other hand, the dissipation contribution from the direction parallel to $x - y$ may be rewritten as

$$\begin{aligned} T^\parallel &= \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} \omega(\xi)(K(\eta, v) + K(-\xi - \eta, v)) - \omega(\xi + 2\eta)(K(\eta, v) - K(\xi + \eta, v)) d\eta dv \\ &= \int_{\mathbb{R}} \int_{-\infty}^{-\xi/2} (\omega(\xi) + \omega(-\xi - 2\eta)) K(\eta, v) d\eta dv + \int_{\mathbb{R}} \int_{-\xi/2}^{\infty} (\omega(\xi) - \omega(\xi + 2\eta)) K(\eta, v) d\eta dv \\ &= \int_{-\infty}^{-\xi/2} (\omega(\xi) + \omega(-\xi - 2\eta)) \tilde{K}(\eta) d\eta + \int_{-\xi/2}^{\infty} (\omega(\xi) - \omega(\xi + 2\eta)) \tilde{K}(\eta) d\eta \\ &= \int_0^{\xi/2} (2\omega(\xi) - \omega(\xi + 2\eta) - \omega(\xi - 2\eta)) \tilde{K}(\eta) d\eta + \int_{\xi/2}^{\infty} (2\omega(\xi) - \omega(\xi + 2\eta) + \omega(2\eta - \xi)) \tilde{K}(\eta) d\eta, \end{aligned}$$

where we have denoted

$$\tilde{K}(\eta) = \int_{\mathbb{R}} K(\eta, v) dv.$$

Since ω is concave, the proof of the lemma is concluded once we establish the existence of a positive constant C such that

$$\tilde{K}(\eta) \geq \frac{Cm(2\eta)}{\eta}$$

for all $\eta > 0$. But this is immediate since m is nonincreasing, and hence

$$\int_{\mathbb{R}} K(\eta, v) dv \geq \int_{-\eta}^{\eta} K(\eta, v) dv \geq Cm(2\eta) \int_{-\eta}^{\eta} \frac{dv}{\eta^2 + v^2} \geq \frac{Cm(2\eta)}{\eta}. \quad \square$$

Remark A.1 (one-dimensional version). It is clear that this proof also holds in the one-dimensional case relevant for the Burgers equation. In fact this case is simpler since there is no need to introduce \tilde{K} .

References

- [Alibaud et al. 2007] N. Alibaud, J. Droniou, and J. Vovelle, “Occurrence and non-appearance of shocks in fractal Burgers equations”, *J. Hyperbolic Differ. Equ.* **4**:3 (2007), 479–499. MR 2008i:35198 Zbl 1144.35038
- [Caffarelli and Vasseur 2010] L. A. Caffarelli and A. Vasseur, “Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation”, *Ann. of Math. (2)* **171**:3 (2010), 1903–1930. MR 2011m:35281 Zbl 1204.35063
- [Chae et al. 2010] D. Chae, P. Constantin, and J. Wu, “Dissipative models generalizing the 2D Navier–Stokes and the SQG equations”, preprint, 2010. To appear in *Indiana Univ. Math. J.* arXiv 1011.0171v1
- [Chae et al. 2011] D. Chae, P. Constantin, and J. Wu, “Inviscid models generalizing the two-dimensional Euler and the surface quasi-geostrophic equations”, *Arch. Ration. Mech. Anal.* **202**:1 (2011), 35–62. MR 2012j:76017 Zbl 06101950
- [Constantin and Vicol 2012] P. Constantin and V. Vicol, “Nonlinear maximum principles for dissipative linear nonlocal operators and applications”, *Geom. Funct. Anal.* **22**:5 (2012), 1289–1321. MR 2989434 Zbl 1256.35078

- [Constantin and Wu 2008] P. Constantin and J. Wu, “Regularity of Hölder continuous solutions of the supercritical quasi-geostrophic equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **25**:6 (2008), 1103–1110. MR 2009h:35328 Zbl 1149.76052
- [Constantin et al. 1994] P. Constantin, A. J. Majda, and E. Tabak, “Formation of strong fronts in the 2–D quasigeostrophic thermal active scalar”, *Nonlinearity* **7**:6 (1994), 1495–1533. MR 95i:76107 Zbl 0809.35057
- [Córdoba and Córdoba 2004] A. Córdoba and D. Córdoba, “A maximum principle applied to quasi-geostrophic equations”, *Comm. Math. Phys.* **249**:3 (2004), 511–528. MR 2005f:76011 Zbl 02158321
- [Dabkowski 2011] M. Dabkowski, “Eventual regularity of the solutions to the supercritical dissipative quasi-geostrophic equation”, *Geom. Funct. Anal.* **21**:1 (2011), 1–13. MR 2012e:35190 Zbl 1210.35185
- [Dabkowski et al. 2012] M. Dabkowski, A. Kiselev, and V. Vicol, “Global well-posedness for a slightly supercritical surface quasi-geostrophic equation”, *Nonlinearity* **25**:5 (2012), 1525–1535. MR 2921019 Zbl 1239.35121
- [Do 2013] T. Do, “On a 1D transport equation with nonlocal velocity and supercritical dissipation”, preprint, 2013. arXiv 1304.4296
- [Dong 2010] H. Dong, “Dissipative quasi-geostrophic equations in critical Sobolev spaces: Smoothing effect and global well-posedness”, *Discrete Contin. Dyn. Syst.* **26**:4 (2010), 1197–1211. MR 2011c:35443 Zbl 1186.35158
- [Dong and Du 2008] H. Dong and D. Du, “Global well-posedness and a decay estimate for the critical dissipative quasi-geostrophic equation in the whole space”, *Discrete Contin. Dyn. Syst.* **21**:4 (2008), 1095–1101. MR 2009b:76175 Zbl 1141.35436
- [Dong and Pavlović 2009] H. Dong and N. Pavlović, “Regularity criteria for the dissipative quasi-geostrophic equations in Hölder spaces”, *Comm. Math. Phys.* **290**:3 (2009), 801–812. MR 2010k:35053 Zbl 1185.35187
- [Dong et al. 2009] H. Dong, D. Du, and D. Li, “Finite time singularities and global well-posedness for fractal Burgers equations”, *Indiana Univ. Math. J.* **58**:2 (2009), 807–821. MR 2010i:35320 Zbl 1166.35030
- [Elgindi 2014] T. M. Elgindi, “Osgood’s lemma and some results for the slightly supercritical 2D Euler equations for incompressible flow”, *Archive for Rational Mechanics and Analysis* **211**:3 (2014), 965–990. MR 3158812
- [Kiselev 2011] A. Kiselev, “Nonlocal maximum principles for active scalars”, *Adv. Math.* **227**:5 (2011), 1806–1826. MR 2012k:35414 Zbl 1244.35022
- [Kiselev and Nazarov 2009] A. Kiselev and F. Nazarov, “A variation on a theme of Caffarelli and Vasseur”, *Zap. Nauchn. Sem. (POMI)* **370**:40 (2009), 58–72. MR 2012c:35350
- [Kiselev et al. 2007] A. Kiselev, F. Nazarov, and A. Volberg, “Global well-posedness for the critical 2D dissipative quasi-geostrophic equation”, *Invent. Math.* **167**:3 (2007), 445–453. MR 2008f:35308 Zbl 1121.35115
- [Kiselev et al. 2008] A. Kiselev, F. Nazarov, and R. Shterenberg, “Blow up and regularity for fractal Burgers equation”, *Dyn. Partial Differ. Equ.* **5**:3 (2008), 211–240. MR 2009k:35264 Zbl 1186.35020
- [Klafter and Sokolov 2005] J. Klafter and I. Sokolov, “Anomalous diffusion spreads its wings”, *Physics World* **18**:8 (2005), 29–32.
- [Pedlosky 1987] J. Pedlosky, *Geophysical Fluid Dynamics*, 2nd ed., Springer, New York, 1987. Zbl 0429.76001
- [Resnick 1995] S. Resnick, *Dynamical problems in nonlinear advective partial differential equations*, Ph.D. thesis, University of Chicago, 1995, Available at <http://search.proquest.com/docview/304242616>.
- [Silvestre 2010] L. Silvestre, “Eventual regularization for the slightly supercritical quasi-geostrophic equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27**:2 (2010), 693–704. MR 2011c:35603 Zbl 1187.35186
- [Silvestre 2011] L. Silvestre, “On the differentiability of the solution to an equation with drift and fractional diffusion”, preprint, 2011. To appear in *Indiana Univ. Math. J.* arXiv 1012.2401v2
- [Tao 2009] T. Tao, “Global regularity for a logarithmically supercritical hyperdissipative Navier–Stokes equation”, *Anal. PDE* **2**:3 (2009), 361–366. MR 2011b:35382 Zbl 1190.35177
- [Xue and Zheng 2012] L. Xue and Z. Zheng, “Note on the well-posedness of a slightly supercritical surface quasi-geostrophic equation”, *Journal of Differential Equations* **253**:2 (2012), 795–813. MR 2921214

MICHAEL DABKOWSKI: mgdabkow@umich.edu

Department of Mathematics, University of Michigan, 2074 East Hall, 530 Church Street, Ann Arbor, MI 48109, United States

ALEXANDER KISELEV: kiselev@math.wisc.edu

Department of Mathematics, University of Wisconsin, 480 Lincoln Dr., Madison, WI 53705, United States

LUIS SILVESTRE: luis@math.uchicago.edu

Department of Mathematics, The University of Chicago, Chicago, IL 60637, United States

VLAD VICOL: vvicol@math.princeton.edu

Department of Mathematics, Princeton University, Fine Hall, Washington Road, Princeton, NJ 08544, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 1 2014

PAUL LAURAIN and TRISTAN RIVIÈRE	1
Global well-posedness of slightly supercritical active scalar equations MICHAEL DABKOWSKI, ALEXANDER KISELEV, LUIS SILVESTRE and VLAD VICOL	43
The nonlinear Schrödinger equation ground states on product spaces SUSANNA TERRACINI, NIKOLAY TZVETKOV and NICOLA VISCIGLIA	73
Orthonormal systems in linear spans ALLISON LEWKO and MARK LEWKO	97
A partial data result for the magnetic Schrödinger inverse problem FRANCIS J. CHUNG	117
Sharp polynomial decay rates for the damped wave equation on the torus NALINI ANANTHARAMAN and MATTHIEU LÉAUTAUD	159
The J -flow on Kähler surfaces: a boundary case HAO FANG, MIJIA LAI, JIAN SONG and BEN WEINKOVE	215
A priori estimates for complex Hessian equations SŁAWOMIR DINEW and SŁAWOMIR KOŁODZIEJ	227
The Aharonov–Bohm effect in spectral asymptotics of the magnetic Schrödinger operator GREGORY ESKIN and JAMES RALSTON	245



2157-5045(2014)7:1;1-F