

ANALYSIS & PDE

Volume 7

No. 2

2014

JEAN DOLBEAULT, MARIA J. ESTEBAN AND ARI LAPTEV

SPECTRAL ESTIMATES ON THE SPHERE

SPECTRAL ESTIMATES ON THE SPHERE

JEAN DOLBEAULT, MARIA J. ESTEBAN AND ARI LAPTEV

In this article we establish optimal estimates for the first eigenvalue of Schrödinger operators on the d -dimensional unit sphere. These estimates depend on L^p norms of the potential, or of its inverse, and are equivalent to interpolation inequalities on the sphere. We also characterize a semiclassical asymptotic regime and discuss how our estimates on the sphere differ from those on the Euclidean space.

1. Introduction

Let Δ be the Laplace–Beltrami operator on the unit d -dimensional sphere \mathbb{S}^d . Our first result is concerned with the sharp estimate of the first negative eigenvalue $\lambda_1 = \lambda_1(-\Delta - V)$ of the Schrödinger operator $-\Delta - V$ on \mathbb{S}^d (with potential $-V$) in terms of L^p -norms of V .

The literature on spectral estimates for the negative eigenvalues of Schrödinger operators on manifolds is limited. P. Federbusch [1969] and O. S. Rothaus [1981] established a link between logarithmic Sobolev inequalities and the ground state energy of Schrödinger operators. The Rozenbljum–Lieb–Cwikel inequality (case $\gamma = 0$ with standard notations: see below) on manifolds has been studied in [Levin and Solomyak 1997, Section 5]; we may also refer to [Lieb 1976] for the semiclassical regime, and to [Levin 2006; Ouhabaz and Poupaud 2010] for more recent results in this direction. A. Ilyin, in two articles [1993; 2012] on Lieb–Thirring type inequalities (see also [Levin 2006; Ouhabaz and Poupaud 2010] for other results on manifolds), considers Schrödinger operators on unit spheres restricted to the space of functions orthogonal to constants and uses the original method of E. Lieb and W. Thirring [1976]. The exclusion of the zero mode of the Laplace–Beltrami operator results in *semiclassical* estimates similar to those for negative eigenvalues of Schrödinger operators in Euclidean spaces.

The results in this paper are somewhat complementary. We show that if the L^p -norm of V is smaller than an explicit value, the first eigenvalue $\lambda_1(-\Delta - V)$ cannot satisfy the semiclassical inequality and thus it is impossible to obtain standard Lieb–Thirring type inequalities for the whole negative spectrum. However, we show that if the L^p -norm of the potential is large, the first eigenvalue behaves semiclassically and the best constant in the inequality asymptotically coincides with the best constants $L_{\gamma,d}^1$ of the corresponding inequality in the Euclidean space of the same dimension (see below). In this regime the first eigenfunction is concentrated around some point on \mathbb{S}^d and can be identified with an eigenfunction of the Schrödinger operator on the tangent space, up to a small error. In [Appendix A](#), we illustrate the

MSC2010: primary 35P15, 58J50, 81Q10, 81Q35; secondary 47A75, 26D10, 46E35, 58E35, 81Q20.

Keywords: spectral problems, partial differential operators on manifolds, quantum theory, estimation of eigenvalues, Sobolev inequality, interpolation, Gagliardo–Nirenberg–Sobolev inequalities, logarithmic Sobolev inequality, Schrödinger operator, ground state, one bound state Keller–Lieb–Thirring inequality.

transition between the small L^p -norm regime and the asymptotic, semiclassical regime by numerically computing the optimal estimates for the eigenvalue $\lambda_1(-\Delta - V)$ in terms of the norms $\|V\|_{L^p(\mathbb{S}^d)}$.

In order to formulate our first theorem, let us introduce the measure $d\omega$ induced by the Lebesgue measure on $\mathbb{S}^d \subset \mathbb{R}^{d+1}$ and the uniform probability measure $d\sigma = d\omega/|\mathbb{S}^d|$ with $|\mathbb{S}^d| = \omega(\mathbb{S}^d)$. We shall denote by $\|\cdot\|_{L^q(\mathbb{S}^d)}$ the quantity $\|u\|_{L^q(\mathbb{S}^d)} = (\int_{\mathbb{S}^d} |u|^q d\sigma)^{1/q}$ for any $q > 0$ (including the case $q \in (0, 1)$, for which $\|\cdot\|_{L^q(\mathbb{S}^d)}$ is no longer a norm, but is only a quasinorm). Because of the normalization of $d\sigma$, when making comparisons with corresponding results in the Euclidean space, we will need the constant

$$\kappa_{q,d} := |\mathbb{S}^d|^{1-2/q}.$$

The well-known optimal constant $L_{\gamma,d}^1$ in the one bound state Keller–Lieb–Thirring inequality is defined as follows: for any function ϕ on \mathbb{R}^d , if $\lambda_1(-\Delta - \phi)$ denotes the lowest negative eigenvalue of the Schrödinger operator $-\Delta - \phi$ (with potential $-\phi$) when it exists, and 0 otherwise, we have

$$|\lambda_1(-\Delta - \phi)|^\gamma \leq L_{\gamma,d}^1 \int_{\mathbb{R}^d} \phi_+^{\gamma+d/2} dx, \tag{1}$$

provided $\gamma \geq 0$ if $d \geq 3$, $\gamma > 0$ if $d = 2$, and $\gamma \geq 1/2$ if $d = 1$. Notice that only the positive part ϕ_+ of ϕ is involved in the right-hand side of the above inequality. Assuming that $\gamma > 1 - d/2$ if $d = 1$ or 2 , we shall consider the exponents

$$q = 2 \frac{2\gamma + d}{2\gamma + d - 2} \quad \text{and} \quad p = \frac{q}{q - 2} = \gamma + \frac{d}{2},$$

which are therefore such that

$$2 < q = \frac{2p}{p - 1} \leq 2^*$$

with $2^* := 2d/(d - 2)$ if $d \geq 3$, and $q = 2p/(p - 1) \in (2, +\infty)$ if $d = 1$ or 2 . To simplify notation, we adopt the convention $2^* := \infty$ if $d = 1$ or 2 . It is also convenient to introduce the notation

$$\alpha_* := \frac{1}{4}d(d - 2).$$

In [Section 2](#) we shall prove the following result.

Theorem 1. *Let $d \geq 1$ and $p \in (\max\{1, d/2\}, +\infty)$. Then there exists a convex increasing function $\alpha : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $\alpha(\mu) = \mu$ for any $\mu \in [0, (d/2)(p - 1)]$ and $\alpha(\mu) > \mu$ for any $\mu \in ((d/2)(p - 1), +\infty)$, such that*

$$|\lambda_1(-\Delta - V)| \leq \alpha(\|V\|_{L^p(\mathbb{S}^d)}) \tag{2}$$

for any nonnegative $V \in L^p(\mathbb{S}^d)$. Moreover, for large values of μ , we have

$$\alpha(\mu)^{p-d/2} = L_{p-d/2,d}^1 (\kappa_{q,d}\mu)^p (1 + o(1)).$$

The estimate (2) is optimal in the sense that there exists a nonnegative function V such that $\mu = \|V\|_{L^p(\mathbb{S}^d)}$ and $|\lambda_1(-\Delta - V)| = \alpha(\mu)$ for any $\mu \in ((d/2)(p - 1), +\infty)$. If $\mu \leq (d/2)(p - 1)$, equality in (2) is achieved by constant potentials.

If $p = d/2$ and $d \geq 3$, then (2) is satisfied with $\alpha(\mu) = \mu$ only for $\mu \in [0, \alpha_*]$. If $d = p = 1$, then (2) is also satisfied for some nonnegative, convex function α on \mathbb{R}^+ such that $\mu \leq \alpha(\mu) \leq \mu + \pi^2 \mu^2$ for any $\mu \in (0, +\infty)$, equality in (2) is achieved and $\alpha(\mu) = \pi^2 \mu^2 (1 + o(1))$ as $\mu \rightarrow +\infty$.

Since $\lambda_1(-\Delta - V)$ is nonpositive for any nonnegative, nontrivial V , inequality (2) is a lower estimate. We have indeed found that

$$0 \geq \lambda_1(-\Delta - V) \geq -\alpha(\|V\|_{L^p(\mathbb{S}^d)}).$$

If V changes sign, the above inequality still holds if V is replaced by the positive part V_+ of V , provided the lowest eigenvalue is negative. We can then write

$$|\lambda_1(-\Delta - V)| \leq \alpha(\|V_+\|_{L^p(\mathbb{S}^d)}) \quad \text{for all } V \in L^p(\mathbb{S}^d). \quad (3)$$

The expression of $L_{\gamma,d}^1$ is not explicit (except in the case $d = 1$: see [Lieb and Thirring 1976, page 290]), but can be given in terms of an optimal constant in some Gagliardo–Nirenberg–Sobolev inequality (see [Lieb and Thirring 1976] and (9)–(10) in Section 2.1). In case $d = p = 1$, notice that $L_{1/2,1}^1 = \frac{1}{2}$ (see Section B.2 in Appendix B) and $\kappa_{\infty,1} = 2\pi$ so that our formula in the asymptotic regime $\mu \rightarrow +\infty$ is consistent with the other cases.

The reader is invited to check that Theorem 1 can be reformulated in a more standard language of spectral theory as follows. We recall that $\gamma = p - d/2$ and that $d\omega$ is the standard measure induced on the unit sphere \mathbb{S}^d by the Lebesgue measure on \mathbb{R}^{d+1} .

Corollary 2. *Let $d \geq 1$ and consider a nonnegative function V . For $\mu = \|V\|_{L^{\gamma+d/2}(\mathbb{S}^d)}$ large, we have*

$$|\lambda_1(-\Delta - V)|^\gamma \lesssim L_{\gamma,d}^1 \int_{\mathbb{S}^d} V^{\gamma+d/2} d\omega \quad (4)$$

if either $\gamma > \max\{0, 1 - d/2\}$ or $\gamma = 1/2$ and $d = 1$. However, if $\mu = \|V\|_{L^{\gamma+d/2}(\mathbb{S}^d)} \leq \frac{1}{4}d(2\gamma + d - 2)$, we have

$$|\lambda_1(-\Delta - V)|^{\gamma+d/2} \leq \int_{\mathbb{S}^d} V^{\gamma+d/2} d\omega \quad (5)$$

for any $\gamma \geq \max\{0, 1 - d/2\}$ and this estimate is optimal.

Here the notation $f \lesssim g$ as $\mu \rightarrow +\infty$ means that $f \leq c(\mu)g$ with $\lim_{\mu \rightarrow \infty} c(\mu) = 1$. The limit case $\gamma = \max\{0, 1 - d/2\}$ in (5) is covered by approximations. We may also notice that optimality in (5) is achieved by constant potentials. Let us give some details.

If we consider a sequence of constant functions $(V_n)_{n \in \mathbb{N}}$ uniformly converging towards 0, for instance $V_n = 1/n$, we get that

$$\lim_{n \rightarrow \infty} \frac{|\lambda_1(-\Delta - V_n)|^\gamma}{\int_{\mathbb{S}^d} V_n^{\gamma+d/2} d\omega} = +\infty,$$

which clearly forbids the possibility of an inequality of the same type as (4) for small values of $\int_{\mathbb{S}^d} V^{\gamma+d/2} d\omega$. This is however compatible with the results of Ilyin in dimension $d = 2$. In [Ilyin 2012, Theorem 2.1], the author states that if P is the orthogonal projection defined by $Pu := u - \int_{\mathbb{S}^2} u d\omega$, the negative eigenvalues $\lambda_k(P(-\Delta - V)P)$ satisfy the semiclassical inequality

$$\sum_k |\lambda_k(P(-\Delta - V)P)| \leq \frac{3}{8} \int_{\mathbb{S}^2} V^2 d\omega.$$

Another way of seeing that inequalities like (4) are incompatible with small potentials is based on the following observation. Inequality (5) shows that

$$|\lambda_1(-\Delta - V)| \leq \left(\int_{\mathbb{S}^2} V^2 d\omega \right)^{1/2}$$

if the L^2 -norm of V is smaller than 1. Since such an inequality is sharp, the semiclassical Lieb–Thirring inequalities for the Schrödinger operator on the sphere \mathbb{S}^2 are therefore impossible for small potentials and can be achieved only in a semiclassical asymptotic regime, that is, when the norm $\|V\|_{L^2(\mathbb{S}^2)}$ is large.

Our second main result is concerned with the estimates from below for the first eigenvalue of Schrödinger operators with positive potentials. In this case, by analogy with (1), it is convenient to introduce the constant $L^1_{-\gamma,d}$ with $\gamma > d/2$, which is the optimal constant in the inequality

$$\lambda_1(-\Delta + \phi)^{-\gamma} \leq L^1_{-\gamma,d} \int_{\mathbb{R}^d} \phi^{d/2-\gamma} dx, \tag{6}$$

where ϕ is any positive potential on \mathbb{R}^d and $\lambda_1(-\Delta + \phi)$ denotes the lowest positive eigenvalue if it exists, or $+\infty$ otherwise. Inequality (6) is less standard than (1); we refer to [Dolbeault et al. 2006, Theorem 12] for a statement and a proof. As in Theorem 1, we shall also introduce exponents p and q such that

$$q = 2 \frac{2\gamma - d}{2\gamma - d + 2} \quad \text{and} \quad p = \frac{q}{2 - q} = \gamma - \frac{d}{2},$$

so that p (respectively $q = 2p/p + 1$) takes arbitrary values in $(0, +\infty)$ (respectively $(0, 2)$). With these notations, we have the counterpart of Theorem 1 in the case of positive potentials.

Theorem 3. *Let $d \geq 1$, $p \in (0, +\infty)$. There exists a concave increasing function $v : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $v(\beta) = \beta$ for any $\beta \in [0, (d/2)(p + 1)]$ if $p > 1$, $v(\beta) \leq \beta$ for any $\beta > 0$ and $v(\beta) < \beta$ for any $\beta \in ((d/2)(p + 1), +\infty)$, such that*

$$\lambda_1(-\Delta + W) \geq v(\beta) \quad \text{with} \quad \beta = \|W^{-1}\|_{L^p(\mathbb{S}^d)}^{-1}, \tag{7}$$

for any positive potential W such that $W^{-1} \in L^p(\mathbb{S}^d)$. Moreover, for large values of β , we have

$$v(\beta)^{-(p+d/2)} \lesssim L^1_{-(p+d/2),d} (\kappa_{q,d} \beta)^{-p}.$$

The estimate (7) is optimal in the sense that there exists a nonnegative potential W such that $\beta^{-1} = \|W^{-1}\|_{L^p(\mathbb{S}^d)}$ and $\lambda_1(-\Delta + W) = v(\beta)$ for any positive β and p . If $\beta \leq (d/2)(p + 1)$ and $p > 1$, equality in (7) is achieved by constant potentials.

Again the expression of $L^1_{-\gamma,d}$ is not explicit when $d \geq 2$ but can be given in terms of an optimal constant in some Gagliardo–Nirenberg–Sobolev inequality; see [Dolbeault et al. 2006] and (17)–(18) in Section 4.

We can rewrite Theorem 3 in terms of $\gamma = p + d/2$ and explicit integrals involving W .

Corollary 4. *Let $d \geq 1$ and $\gamma > d/2$. For $\beta = \|W^{-1}\|_{L^{\gamma-d/2}(\mathbb{S}^d)}^{-1}$ large, we have*

$$(\lambda_1(-\Delta + W))^{-\gamma} \lesssim L^1_{-\gamma,d} \int_{\mathbb{S}^d} W^{d/2-\gamma} d\omega.$$

However, if $\gamma \geq d/2 + 1$ and if $\beta = \|W^{-1}\|_{L^{\gamma-d/2}(\mathbb{S}^d)}^{-1} \leq \frac{1}{4}d(2\gamma - d + 2)$, we have

$$(\lambda_1(-\Delta + W))^{d/2-\gamma} \leq \int_{\mathbb{S}^d} W^{d/2-\gamma} d\omega,$$

and this estimate is optimal.

This paper is organized as follows. Section 2 contains various results on interpolation inequalities; the most important one for our purpose is stated in Lemma 5. Theorem 1, Corollary 2 and, various spectral estimates for Schrödinger operators with negative potentials are established in Section 3. Section 4 deals with the case of positive potentials and contains the proofs of Theorem 3 and Corollary 4. Section 5 is devoted to the threshold case ($q = 2$, that is, $p, \gamma \rightarrow +\infty$) of exponential estimates for eigenvalues, or, in terms of interpolation inequalities, to logarithmic Sobolev inequalities. Finally, numerical and technical results have been collected in two appendices.

2. Interpolation inequalities and consequences for negative potentials

2.1. Inequalities in the Euclidean space. Let us start with some considerations on inequalities in the Euclidean space, which play a crucial role in the semiclassical regime.

We recall that we denote by 2^* the Sobolev critical exponent $2d/(d - 2)$ if $d \geq 3$ and consider Sobolev’s inequality on \mathbb{R}^d , $d \geq 3$,

$$\|v\|_{L^{2^*}(\mathbb{R}^d)}^2 \leq S_d \|\nabla v\|_{L^2(\mathbb{R}^d)}^2 \quad \text{for all } v \in \mathcal{D}^{1,2}(\mathbb{R}^d) \tag{8}$$

where S_d is the optimal constant and $\mathcal{D}^{1,2}(\mathbb{R}^d)$ is the Beppo Levi space obtained by completion of smooth compactly supported functions with respect to the norm $v \mapsto \|\nabla v\|_{L^2(\mathbb{R}^d)}$. See Section B.4 for details and comments on the expression of S_d .

Assume now that $d \geq 1$ and recall that $2^* = +\infty$ if $d = 1$ or 2 . In the subcritical case, that is, $q \in (2, 2^*)$, let

$$K_{q,d} := \inf_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^q(\mathbb{R}^d)}^2}$$

be the optimal constant in the Gagliardo–Nirenberg–Sobolev inequality

$$K_{q,d} \|v\|_{L^q(\mathbb{R}^d)}^2 \leq \|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^2(\mathbb{R}^d)}^2 \quad \text{for all } v \in H^1(\mathbb{R}^d). \tag{9}$$

The optimal constant $L^1_{\gamma,d}$ in the one bound state Keller–Lieb–Thirring inequality is such that

$$L^1_{\gamma,d} := (K_{q,d})^{-p} \quad \text{with } p = \gamma + \frac{d}{2}, \quad q = 2 \frac{2\gamma + d}{2\gamma + d - 2}. \tag{10}$$

See [Section B.5](#) for a proof and references and [\[Lieb and Thirring 1976\]](#) for a detailed discussion. Also see [\[Barnes 1976\]](#) for numerical values of $K_{q,d}$.

We shall also define the exponent

$$\vartheta := d \frac{q-2}{2q}$$

which plays an important role in the scale invariant form of the Gagliardo–Nirenberg–Sobolev interpolation inequalities associated to $K_{q,d}$: see [Section B.1](#) for details.

2.2. Interpolation inequalities on the sphere. Using the inverse stereographic projection (see [Section B.3](#)), it is possible to relate interpolation inequalities on \mathbb{R}^d with interpolation inequalities on \mathbb{S}^d . In this section we consider the case of the sphere. Notice that $\alpha_* = d/(q-2)$ when $q = 2^* = 2d/(d-2)$, $d \geq 3$.

Lemma 5. *Let $q \in (2, 2^*)$. There exists a concave increasing function $\mu : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the properties*

$$\begin{aligned} \mu(\alpha) &= \alpha && \text{for all } \alpha \in \left[0, \frac{d}{q-2}\right], \\ \mu(\alpha) &< \alpha && \text{for all } \alpha \in \left(\frac{d}{q-2}, +\infty\right), \\ \mu(\alpha) &= \frac{K_{q,d}}{\kappa_{q,d}} \alpha^{1-\vartheta} (1 + o(1)) && \text{as } \alpha \rightarrow +\infty, \end{aligned}$$

and such that

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \mu(\alpha) \|u\|_{L^q(\mathbb{S}^d)}^2 \quad \text{for all } u \in H^1(\mathbb{S}^d). \tag{11}$$

If $d \geq 3$ and $q = 2^*$, the inequality also holds for any $\alpha > 0$ with $\mu(\alpha) = \min\{\alpha, \alpha_*\}$.

The remainder of this section is mostly devoted to the proof of [Lemma 5](#). A fundamental tool is a rigidity result proved by M.-F. Bidaut-Véron and L. Véron [\[1991, Theorem 6.1\]](#) for $q > 2$, which goes as follows. Any positive solution of

$$-\Delta f + \alpha f = f^{q-1} \tag{12}$$

has a unique solution $f \equiv \alpha^{1/(q-2)}$ for any $0 < \alpha \leq d/(q-2)$. A straightforward consequence of this rigidity result is the following interpolation inequality [\[Bidaut-Véron and Véron 1991, Corollary 6.2\]](#):

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\sigma \geq \frac{d}{q-2} \left[\left(\int_{\mathbb{S}^d} |u|^q d\sigma \right)^{2/q} - \int_{\mathbb{S}^d} |u|^2 d\sigma \right] \quad \text{for all } u \in H^1(\mathbb{S}^d, d\sigma). \tag{13}$$

Inequality [\(13\)](#) holds for any $q \in [1, 2) \cup (2, 2^*]$ if $d \geq 3$ and for any $q \in [1, 2) \cup (2, \infty)$ if $d = 1$ or 2 . An alternative proof of [\(13\)](#) has been established in [\[Beckner 1993\]](#) for $q > 2$ using previous results by Lieb [\[1983\]](#) and the Funk–Hecke formula [\[Funk 1915; Hecke 1917\]](#). The whole range $p \in [1, 2) \cup (2, 2^*)$ was covered in the case of the ultraspherical operator [\[Bentaleb and Fahlaoui 2009; 2010\]](#). Also see [\[Bakry and Ledoux 1996; Ledoux 2000\]](#) for the *carré du champ* method, and [\[Dolbeault et al. 2013\]](#) for an elementary proof. Inequality [\(13\)](#) is *tight* as defined by D. Bakry [\[2006, Section 2\]](#), in the sense that equality is achieved only by constants.

Remark 6. Inequality (13) is equivalent to

$$\inf_{u \in H^1(\mathbb{S}^d) \setminus \{0\}} \frac{(q-2) \|\nabla u\|_{L^2(\mathbb{S}^d)}^2}{\|u\|_{L^q(\mathbb{S}^d)}^2 - \|u\|_{L^2(\mathbb{S}^d)}^2} = d.$$

Although we will not make use of them in this paper, we may notice that the following properties hold true:

(i) If $q < 2^*$, the above infimum is not achieved in $H^1(\mathbb{S}^d) \setminus \{0\}$, but

$$\lim_{\varepsilon \rightarrow 0^+} \frac{(q-2) \|\nabla u_\varepsilon\|_{L^2(\mathbb{S}^d)}^2}{\|u_\varepsilon\|_{L^q(\mathbb{S}^d)}^2 - \|u_\varepsilon\|_{L^2(\mathbb{S}^d)}^2} = d$$

if $u_\varepsilon := 1 + \varepsilon\varphi$, where φ is a nontrivial eigenfunction of the Laplace–Beltrami operator corresponding to the first nonzero eigenvalue (see Section 2.3).

(ii) If $q = 2^*$, $d \geq 3$, there are nontrivial optimal functions for (13), due to the conformal invariance. Alternatively, these solutions can be constructed from the family of Aubin–Talenti optimal functions for Sobolev’s inequality, using the inverse stereographic projection.

(iii) If $\alpha > \alpha_*$ and $q = 2^*$, $d \geq 3$, there are no optimal functions for (11), since otherwise $\alpha \mapsto \mu(\alpha)$ would not be constant on (α_*, α) : see Proposition 7 below.

2.3. Properties of the function $\alpha \mapsto \mu(\alpha)$ in the subcritical case. Assume that $q \in (2, 2^*)$. For any $\alpha > 0$, consider

$$\mathcal{Q}_\alpha[u] := \frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^d)}^2}{\|u\|_{L^q(\mathbb{S}^d)}^2} \quad \text{for all } u \in H^1(\mathbb{S}^d, d\sigma).$$

It is a standard result of the calculus of variations that

$$\inf_{\substack{u \in H^1(\mathbb{S}^d, d\sigma) \\ \int_{\mathbb{S}^d} |u|^q d\sigma = 1}} \mathcal{Q}_\alpha[u] := \mu(\alpha)$$

is achieved by a minimizer $u \in H^1(\mathbb{S}^d, d\sigma)$ which solves the Euler–Lagrange equations

$$-\Delta u + \alpha u - \mu(\alpha) u^{q-1} = 0. \tag{14}$$

Indeed, we know that there is a Lagrange multiplier associated to the constraint $\int_{\mathbb{S}^d} |u|^q d\sigma = 1$, and multiplying (14) by u and integrating on \mathbb{S}^d , we can identify it with $\mu(\alpha)$. As a corollary, we have shown that (11) holds. The fact that the Lagrange multiplier can be identified so easily is a consequence of the fact that all terms in (11) are two-homogeneous.

We can now list some basic properties of the function $\alpha \mapsto \mu(\alpha)$.

(1) For any $\alpha > 0$, $\mu(\alpha)$ is positive, since the infimum is achieved by a nonnegative function u and $u = 0$ is incompatible with the constraint $\int_{\mathbb{S}^d} |u|^q d\sigma = 1$. By taking a constant test function, we see that $\mu(\alpha) \leq \alpha$ for all $\alpha > 0$. The function $\alpha \mapsto \mu(\alpha)$ is monotone nondecreasing since for a given $u \in H^1(\mathbb{S}^d, d\sigma) \setminus \{0\}$, the function $\alpha \mapsto \mathcal{Q}_\alpha[u]$ is monotone increasing. It is actually strictly monotone.

Indeed, if $\mu(\alpha_1) = \mu(\alpha_2)$ with $\alpha_1 < \alpha_2$, one can notice that $\mathcal{Q}_{\alpha_1}[u_2] < \mu(\alpha_1)$ if u_2 is a minimizer of \mathcal{Q}_{α_2} satisfying the constraint $\int_{\mathbb{S}^d} |u_2|^q d\sigma = 1$, which provides an obvious contradiction.

(2) We have

$$\mu(\alpha) = \alpha \quad \text{for all } \alpha \in \left(0, \frac{d}{q-2}\right].$$

Indeed, if u is a solution of (14), $f = \mu(\alpha)^{1/(q-2)}u$ solves (12) and is therefore a constant function if $\alpha \leq d/(q-2)$ according to [Bidaut-Véron and Véron 1991, Theorem 6.1], and so is u as well. Because of the normalization constraint $\|u\|_{L^q(\mathbb{S}^d)} = 1$, we get that $u = 1$, which proves the statement.

On the contrary, we have

$$\mu(\alpha) < \alpha \quad \text{for all } \alpha > \frac{d}{q-2}.$$

Let us prove this. Let φ be a nontrivial eigenfunction of the Laplace–Beltrami operator corresponding to the first nonzero eigenvalue:

$$-\Delta\varphi = d\varphi.$$

If $x = (x_1, x_2, \dots, x_d, z)$ are cartesian coordinates of $x \in \mathbb{R}^{d+1}$ so that $\mathbb{S}^d \subset \mathbb{R}^{d+1}$ is characterized by the condition $\sum_{i=1}^d x_i^2 + z^2 = 1$, a simple choice of such a function φ is $\varphi(x) = z$. By orthogonality with respect to the constants, we know that $\int_{\mathbb{S}^d} \varphi d\sigma = 0$. We may now Taylor expand \mathcal{Q}_α around $u = 1$ by considering $u = 1 + \varepsilon\varphi$ as $\varepsilon \rightarrow 0$ and obtain that

$$\mu(\alpha) \leq \mathcal{Q}_\alpha[1 + \varepsilon\varphi] = \frac{(d + \alpha)\varepsilon^2 \int_{\mathbb{S}^d} |\varphi|^2 d\sigma + \alpha}{\left(\int_{\mathbb{S}^d} |1 + \varepsilon\varphi|^q d\sigma\right)^{2/q}} = \alpha + [d + \alpha(2 - q)]\varepsilon^2 \int_{\mathbb{S}^d} |\varphi|^2 d\sigma + o(\varepsilon^2).$$

By taking ε small enough, we get $\mu(\alpha) < \alpha$ for all $\alpha > d/(q-2)$. Optimizing on the value of $\varepsilon > 0$ (not necessarily small) provides an interesting test function: see Section A.1.

(3) The function $\alpha \mapsto \mu(\alpha)$ is concave, because it is the minimum of a family of affine functions.

2.4. More estimates on the function $\alpha \mapsto \mu(\alpha)$. We first consider the critical case $q = 2^*$, $d \geq 3$. As in the subcritical case $q < 2^*$, we have $\mu(\alpha) = \alpha$ for $\alpha \leq \alpha^*$. For $\alpha > \alpha^*$, the function $\alpha \mapsto \mu(\alpha)$ is constant.

Proposition 7. *With the notations of Lemma 5, if $d \geq 3$ and $q = 2^*$, then*

$$\mu(\alpha) = \alpha_* \quad \text{for all } \alpha > \alpha_* = \frac{d}{q-2} = \frac{1}{4}d(d-2).$$

Proof. Consider the Aubin–Talenti optimal functions for Sobolev’s inequality and, more specifically, let us choose the functions

$$v_\varepsilon(x) := \left(\frac{\varepsilon}{\varepsilon^2 + |x|^2}\right)^{\frac{d-2}{2}} \quad \text{for all } x \in \mathbb{R}^d \text{ and all } \varepsilon > 0,$$

which are such that $\|v_\varepsilon\|_{L^{2^*}(\mathbb{R}^d)} = \|v_1\|_{L^{2^*}(\mathbb{R}^d)}$ is independent of ε . With standard notations (see Section B.3), let $N \in \mathbb{S}^d$ be the north pole. Using the stereographic projection Σ , that is, for the functions defined for

any $y \in \mathbb{S}^d \setminus \{\mathbf{N}\}$ by

$$u_\varepsilon(y) = \left(\frac{|x|^2 + 1}{2} \right)^{\frac{d-2}{2}} v_\varepsilon(x) \quad \text{with } x = \Sigma(y),$$

we find that $\|u_\varepsilon\|_{L^{2^*}(\mathbb{S}^d)} = \|v_1\|_{L^{2^*}(\mathbb{R}^d)}$ for any $\varepsilon > 0$, so that

$$\mu(\alpha) \leq \mathcal{Q}_\alpha[u_\varepsilon] = \frac{\|\nabla v_\varepsilon\|_{L^2(\mathbb{R}^d)}^2 + (\alpha - \alpha_*) \int_{\mathbb{R}^d} |v_\varepsilon|^2 (2/(1+|x|^2))^2 dx}{\kappa_{2^*,d} \|v_\varepsilon\|_{L^{2^*}(\mathbb{R}^d)}^2} = \alpha_* + 4|\mathbb{S}^d|^{1-2/d} (\alpha - \alpha_*) \frac{\delta(d, \varepsilon)}{\|v_1\|_{L^{2^*}(\mathbb{R}^d)}^2},$$

where we have used the fact that $\kappa_{2^*,d} S_d = 1/\alpha_*$ (see [Section B.4](#)) and

$$\delta(d, \varepsilon) := \int_0^\infty \left(\frac{\varepsilon}{\varepsilon^2 + r^2} \right)^{d-2} \frac{r^{d-1}}{(1+r^2)^2} dr = \varepsilon^2 \int_0^\infty \left(\frac{1}{1+s^2} \right)^{d-2} \frac{s^{d-1}}{(1+\varepsilon^2 s^2)^2} ds.$$

One can check that $\lim_{\varepsilon \rightarrow 0_+} \delta(d, \varepsilon) = 0$ since

$$\delta(d, \varepsilon) \leq \varepsilon^2 \int_0^\infty \frac{s^{d-1}}{(1+s^2)^{d-2}} ds \quad \text{if } d \geq 5 \quad \text{and} \quad \delta(d, \varepsilon) \leq \varepsilon c_d \int_0^{+\infty} \frac{ds}{(1+s^2)^2} \quad \text{if } d = 3 \text{ or } 4,$$

with $c_3 = 1$ and $c_4 = 3\sqrt{3}/16$. □

The next step is devoted to a lower estimate for the function $\alpha \mapsto \mu(\alpha)$ in the subcritical case, which shows that $\lim_{\alpha \rightarrow +\infty} \mu(\alpha) = +\infty$ in contrast with the critical case.

Proposition 8. *With the notations of [Lemma 5](#), if $d \geq 3$ and $q \in (2, 2^*)$, then, for any $\alpha > \alpha_*$, we have*

$$\alpha > \mu(\alpha) \geq \alpha_*^\vartheta \alpha^{1-\vartheta}$$

with $\vartheta = d(q-2)/2q$. For every $s \in (2, 2^*)$, if $d \geq 3$, or every $s \in (2, +\infty)$ if $d = 1$ or 2 , such that $s > q$, we also have that

$$\alpha > \mu(\alpha) \geq \left(\frac{d}{s-2} \right)^\theta \alpha^{1-\theta}$$

for any $\alpha > d/(s-2)$ and $\theta = \theta(s, q, d) := s(q-2)/(q(s-2))$.

Proof. The first case can be seen as a limit case of the second one as $s \rightarrow 2^*$ and $\vartheta = \theta(2^*, q, d)$. Using Hölder's inequality, we can estimate $\|u\|_{L^q(\mathbb{S}^d)}$ by

$$\|u\|_{L^q(\mathbb{S}^d)} \leq \|u\|_{L^s(\mathbb{S}^d)}^\theta \|u\|_{L^2(\mathbb{S}^d)}^{1-\theta}$$

and get the result using

$$\mathcal{Q}_\alpha[u] \geq \left(\frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^d)}^2}{\|u\|_{L^s(\mathbb{S}^d)}^2} \right)^\theta \left(\frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^d)}^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \right)^{1-\theta} \geq \left(\frac{d}{s-2} \right)^\theta \alpha^{1-\theta}. \quad \square$$

Proposition 9. *With the notations of [Lemma 5](#), for every $q \in (2, 2^*)$, we have*

$$\limsup_{\alpha \rightarrow +\infty} \alpha^{\vartheta-1} \mu(\alpha) \leq \frac{K_{q,d}}{\kappa_{q,d}}.$$

Proof. Let v be an optimal function for $K_{q,d}$ and define for any $x \in \mathbb{R}^d$ the function

$$v_\alpha(x) := v(2\sqrt{\alpha - \alpha_*}x)$$

with $\alpha_* = \frac{1}{4}d(d-2)$ and $\alpha > \alpha_*$, so that

$$\int_{\mathbb{R}^d} |\nabla v_\alpha|^2 dx = 2^{2-d}(\alpha - \alpha_*)^{1-d/2} \int_{\mathbb{R}^d} |\nabla v|^2 dx,$$

$$\int_{\mathbb{R}^d} |v_\alpha|^q \left(\frac{2}{1+|x|^2}\right)^{d-(d-2)q/1} dx = 2^{-(d-2)q/2}(\alpha - \alpha_*)^{-d/2} \int_{\mathbb{R}^d} |v|^q \left(1 + \frac{|x|^2}{4(\alpha - \alpha_*)}\right)^{-d+(d-2)q/2} dx.$$

Now we observe that the function $u_\alpha(y) := ((|x|^2 + 1)/2)^{(d-2)/2}v_\alpha(x)$, where $y = \Sigma^{-1}(x)$ and Σ is the stereographic projection (see Section B.3), is such that

$$\mathcal{Q}_\alpha[u_\alpha] = \frac{1}{\kappa_{q,d}} \frac{\int_{\mathbb{R}^d} |\nabla v_\alpha|^2 dx + (\alpha - \alpha_*) \int_{\mathbb{R}^d} |v_\alpha|^2 (2/(1+|x|^2))^2 dx}{\left[\int_{\mathbb{R}^d} |v_\alpha|^q (2/(1+|x|^2))^{d-(d-2)q/2} dx\right]^{2/q}}.$$

Passing to the limit as $\alpha \rightarrow +\infty$, we get

$$\lim_{\alpha \rightarrow +\infty} \int_{\mathbb{R}^d} |v|^q \left(1 + \frac{|x|^2}{4(\alpha - \alpha_*)}\right)^{-d+(d-2)q/2} dx = \int_{\mathbb{R}^d} |v|^q dx$$

by Lebesgue’s theorem of dominated convergence. The limit also holds with q replaced by 2. This proves that

$$\mathcal{Q}_\alpha[u_\alpha] = (\alpha - \alpha_*)^{1-d/2+d/q} \left(\frac{K_{q,d}}{\kappa_{q,d}} + o(1)\right) \quad \text{as } \alpha \rightarrow +\infty,$$

which concludes the proof because $\vartheta = d(q-2)/(2q)$. □

2.5. The semiclassical regime: behavior of the function $\alpha \mapsto \mu(\alpha)$ as $\alpha \rightarrow +\infty$. Assume $q \in (2, 2^*)$. If we combine the results of Propositions 8 and 9, we know that $\mu(\alpha) \sim \alpha^{1-\vartheta}$ as $\alpha \rightarrow +\infty$ if $d \geq 3$. If $d = 1$ or 2, we know that $\lim_{\alpha \rightarrow +\infty} \mu(\alpha) = +\infty$ with a growth at least equivalent to $\alpha^{2/q-\varepsilon}$ with $\varepsilon > 0$, arbitrarily small, according to Proposition 8, and at most equivalent to $\alpha^{1-\vartheta}$ by Proposition 9. To complete the proof of Lemma 5, it remains to determine the precise behavior of $\mu(\alpha)$ as $\alpha \rightarrow +\infty$.

Proposition 10. *With the notations of Lemma 5, for every $q \in (2, 2^*)$, with $\vartheta = d(q-2)/(2q)$ we have*

$$\mu(\alpha) = \frac{K_{q,d}}{\kappa_{q,d}} \alpha^{1-\vartheta} (1 + o(1)) \quad \text{as } \alpha \rightarrow +\infty.$$

Proof. Suppose by contradiction that there is a positive constant η and a sequence $(\alpha_n)_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow +\infty} \alpha_n = +\infty$ and

$$\lim_{n \rightarrow +\infty} \alpha_n^{\vartheta-1} \mu(\alpha_n) \leq \frac{K_{q,d}}{\kappa_{q,d}} - \eta. \tag{15}$$

Consider a sequence $(u_n)_{n \in \mathbb{N}}$ of functions in $H^1(\mathbb{S}^d)$ such that $\mathcal{Q}_{\alpha_n}[u_n] = \mu(\alpha_n)$ and $\|u_n\|_{L^q(\mathbb{S}^d)} = 1$ for any $n \in \mathbb{N}$. From (15), we know that

$$\alpha_n \|u_n\|_{L^2(\mathbb{S}^d)}^2 \leq \mathcal{Q}_{\alpha_n}[u_n] = \mu(\alpha_n) \leq \alpha_n^{1-\vartheta} \left(\frac{K_{q,d}}{\kappa_{q,d}} - \eta\right) (1 + o(1)) \quad \text{as } n \rightarrow +\infty,$$

that is,

$$\limsup_{n \rightarrow +\infty} \alpha_n^\vartheta \|u_n\|_{L^2(\mathbb{S}^d)}^2 \leq \frac{K_{q,d}}{\kappa_{q,d}} - \eta.$$

The normalization $\|u_n\|_{L^q(\mathbb{S}^d)} = 1$ for any $n \in \mathbb{N}$ and the limit $\lim_{n \rightarrow +\infty} \|u_n\|_{L^2(\mathbb{S}^d)} = 0$ mean that the sequence $(u_n)_{n \in \mathbb{N}}$ concentrates: there exists a sequence $(y_i)_{i \in \mathbb{N}}$ of points in \mathbb{S}^d (eventually finite) and two sequences of positive numbers $(\zeta_i)_{i \in \mathbb{N}}$ and $(r_{i,n})_{i,n \in \mathbb{N}}$ such that $\lim_{n \rightarrow +\infty} r_{i,n} = 0$, $\sum_{i \in \mathbb{N}} \zeta_i = 1$, and $\int_{\mathbb{S}^d \cap B(y_i, r_{i,n})} |u_{i,n}|^q d\sigma = \zeta_i + o(1)$, where $u_{i,n} \in H^1(\mathbb{S}^d)$, $u_{i,n} = u_n$ on $\mathbb{S}^d \cap B(y_i, r_{i,n})$, and

$$\text{supp } u_{i,n} \subset \mathbb{S}^d \cap B(y_i, 2r_{i,n}).$$

Here $o(1)$ means that uniformly with respect to i , the remainder term converges towards 0 as $n \rightarrow +\infty$. Using a computation similar to those of the proof of Proposition 9, we can *blow up* each function $u_{i,n}$ and prove

$$(\alpha_n - \alpha_*)^{\vartheta-1} \int_{\mathbb{S}^d} (|\nabla u_{i,n}|^2 + \alpha_n |u_{i,n}|^2) d\sigma \geq \frac{K_{q,d}}{\kappa_{q,d}} \zeta_i^{2/q} + o(1) \quad \text{for all } i.$$

Let us choose an integer N such that $(\sum_{i=1}^N \zeta_i)^{2/q} > 1 - \kappa_{q,d} \eta / (2K_{q,d})$. Then we find that

$$\begin{aligned} (\alpha_n - \alpha_*)^{\vartheta-1} \int_{\mathbb{S}^d} (|\nabla u_n|^2 + \alpha_n |u_n|^2) d\sigma &\geq \frac{K_{q,d}}{\kappa_{q,d}} \sum_1^N \zeta_i^{2/q} + o(1) \geq \frac{K_{q,d}}{\kappa_{q,d}} \left(\sum_1^N \zeta_i \right)^{2/q} + o(1) \\ &\geq \frac{K_{q,d}}{\kappa_{q,d}} - \frac{\eta}{2} + o(1), \end{aligned}$$

a contradiction with (15). □

For details on the behavior of $K_{q,d}$ as q varies, see Proposition 15. Collecting all results of this section completes the proof of Lemma 5.

3. Spectral estimates for the Schrödinger operator on the sphere

This section is devoted to the proof of Theorem 1. As a consequence of the results of Lemma 5, the function $\alpha \mapsto \mu(\alpha)$ is invertible, of inverse $\mu \mapsto \alpha(\mu)$, if $d = 1, 2$ or $d \geq 3$ and $q < 2^*$, and we have the inequality

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\sigma - \mu \left(\int_{\mathbb{S}^d} |u|^q d\sigma \right)^{\frac{2}{q}} \geq -\alpha(\mu) \int_{\mathbb{S}^d} |u|^2 d\sigma \quad \text{for all } u \in H^1(\mathbb{S}^d, d\sigma) \text{ and all } \mu > 0. \quad (16)$$

Moreover, the function $\mu \mapsto \alpha(\mu)$ is monotone increasing, convex, and satisfies $\alpha(\mu) = \mu$ for any $\mu \in (0, d/(q-2)]$ and $\alpha(\mu) > \mu$ for any $\mu > d/(q-2)$.

Consider the Schrödinger operator $-\Delta - V$ for some function $V \in L^p(\mathbb{S}^d)$ and the corresponding energy functional

$$\mathcal{E}[u] := \int_{\mathbb{S}^d} |\nabla u|^2 d\sigma - \int_{\mathbb{S}^d} V |u|^2 d\sigma.$$

Let

$$\lambda_1(-\Delta - V) := \inf_{\substack{u \in H^1(\mathbb{S}^d, d\sigma) \\ \int_{\mathbb{S}^d} |u|^2 d\sigma = 1}} \mathcal{E}[u].$$

By Hölder’s inequality, we have

$$\mathcal{E}[u] \geq \int_{\mathbb{S}^d} |\nabla u|^2 d\sigma - \|V_+\|_{L^p(\mathbb{S}^d)} \|u\|_{L^q(\mathbb{S}^d)}^2$$

with $1/p + 2/q = 1$. From Section 2, with $\mu = \|V_+\|_{L^p(\mathbb{S}^d)}$, we deduce

$$\mathcal{E}[u] \geq -\alpha(\mu) \|u\|_{L^2(\mathbb{S}^d)}^2 \quad \text{for all } u \in H^1(\mathbb{S}^d, d\sigma) \text{ and all } V \in L^p(\mathbb{S}^d),$$

which amounts to a Keller–Lieb–Thirring inequality on the sphere (3), or equivalently,

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\sigma - \int_{\mathbb{S}^d} V|u|^2 d\sigma + \alpha(\|V_+\|_{L^p(\mathbb{S}^d)}) \int_{\mathbb{S}^d} |u|^2 d\sigma \geq 0 \quad \text{for all } u \in H^1(\mathbb{S}^d, d\sigma) \text{ and all } V \in L^p(\mathbb{S}^d).$$

Notice that this inequality simultaneously contains (3) and (16), by optimizing either on u or on V .

Optimality in (3) still needs to be proved. This can be done by taking an arbitrary $\mu \in (0, \infty)$ and considering an optimal function for (16), for which we have

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\sigma - \mu \left(\int_{\mathbb{S}^d} |u|^q d\sigma \right)^{\frac{2}{q}} = \alpha(\mu) \int_{\mathbb{S}^d} |u|^2 d\sigma.$$

Because the above expression is homogeneous of degree two, there is no restriction to assume that $\int_{\mathbb{S}^d} |u|^q d\sigma = 1$, and since the solution is optimal, it solves the Euler–Lagrange equation

$$-\Delta u - Vu = \alpha(\mu)u$$

with $V = \mu u^{q-2}$, such that

$$\|V_+\|_{L^p(\mathbb{S}^d)} = \mu \|u\|_{L^q(\mathbb{S}^d)}^{q/p} = \mu.$$

Hence such a function V realizes the equality in (3).

Taking into account Lemma 5 and (10), this completes the proof of Theorem 1 in the general case. The case $d = 1$ and $\gamma = 1/2$ has to be treated specifically. Using $u \equiv 1$ as a test function, we know that $|\lambda_1(-\Delta - V)| \leq \mu = \int_{\mathbb{S}^1} V dx$. On the other hand, consider $u \in H^1(\mathbb{S}^1)$ such that $\|u\|_{L^2(\mathbb{S}^1)} = 1$. Since $H^1(\mathbb{S}^1)$ is embedded into $C^{0,1/2}(\mathbb{S}^1)$, there exists $x_0 \in \mathbb{S}^1 \approx [0, 2\pi)$ such that $u(x_0) = 1$ and

$$|u(x)|^2 - 1 = 2 \int_{x_0}^x u(y)u'(y) dy = 2 \int_{x_0+2\pi}^x u(y)u'(y) dy$$

can be estimated by

$$\begin{aligned} ||u(x)|^2 - 1| &\leq 2 \int_{x_0}^x |u(y)||u'(y)| dy = 2 \int_{x_0+2\pi}^x |u(y)||u'(y)| dy \\ &\leq \int_0^{2\pi} |u(y)||u'(y)| dy \leq \left(\int_0^{2\pi} |u(y)|^2 dy \int_0^{2\pi} |u'(y)|^2 dy \right)^{1/2} \end{aligned}$$

using the Cauchy–Schwarz inequality, that is,

$$||u(x)|^2 - 1| \leq 2\pi \|u'\|_{L^2(\mathbb{S}^1)},$$

since $\|u'\|_{L^2(\mathbb{S}^1)}^2 = (1/(2\pi)) \int_0^{2\pi} |u'(y)|^2 dy$ and $\|u\|_{L^2(\mathbb{S}^1)}^2 = (1/(2\pi)) \int_0^{2\pi} |u(y)|^2 dy = 1$ (recall that $d\sigma$ is a probability measure). Thus we get

$$|u(x)|^2 \leq 1 + 2\pi \|u'\|_{L^2(\mathbb{S}^1)},$$

from which it follows that

$$\lambda_1(-\Delta - V) \geq \|u'\|_{L^2(\mathbb{S}^1)}^2 - \mu(1 + 2\pi \|u'\|_{L^2(\mathbb{S}^1)}) \geq -\mu - \pi^2 \mu^2.$$

This shows that $\mu \leq \alpha(\mu) \leq \mu + \pi^2 \mu^2$. By the Arzelà–Ascoli theorem, the embedding of $H^1(\mathbb{S}^1)$ into $C^{0,1/2}(\mathbb{S}^1)$ is compact. When $d = 1$ and $\gamma = 1/2$, the proof of the asymptotic behavior of $\alpha(\mu)$ as $\mu \rightarrow +\infty$ can then be completed as in the other cases.

4. Spectral inequalities in the case of positive potentials

In this section we address the case of Schrödinger operators $-\Delta + W$ where W is a positive potential on \mathbb{S}^d and we derive estimates from below for the first eigenvalue of such operators. In order to do so, we first study interpolation inequalities in the Euclidean space \mathbb{R}^d , like those studied in [Section 2](#) (for $q > 2$).

For this purpose, let us define for $q \in (0, 2)$ the constant

$$K_{q,d}^* := \inf_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^q(\mathbb{R}^d)}^2}{\|v\|_{L^2(\mathbb{R}^d)}^2},$$

that is, the optimal constant in the Gagliardo–Nirenberg–Sobolev inequality

$$K_{q,d}^* \|v\|_{L^2(\mathbb{R}^d)}^2 \leq \|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^q(\mathbb{R}^d)}^2 \quad \text{for all } v \in H^1(\mathbb{R}^d) \quad (17)$$

(with the convention that the right-hand side is infinite if $|v|^q$ is not integrable).

The optimal constant $L_{-\gamma,d}^1$ in [\(6\)](#) is such that

$$L_{-\gamma,d}^1 := (K_{q,d}^*)^{-\gamma} \quad \text{with } q = 2 \frac{2\gamma - d}{2\gamma - d + 2}. \quad (18)$$

See [Section B.6](#) for a proof. Let us define the exponent

$$\delta := \frac{2q}{2d - q(d - 2)}.$$

Lemma 11. *Let $q \in (0, 2)$ and $d \geq 1$. Then there exists a concave increasing function $v : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the properties*

$$\begin{aligned} v(\beta) &\leq \beta \quad \text{for all } \beta > 0 \quad \text{and} \quad v(\beta) < \beta \quad \text{for all } \beta \in \left(\frac{d}{2-q}, +\infty\right), \\ v(\beta) &= \beta \quad \text{for all } \beta \in \left[0, \frac{d}{2-q}\right] \quad \text{if } q \in [1, 2) \quad \text{and} \quad \lim_{\beta \rightarrow 0^+} \frac{v(\beta)}{\beta} = 1 \quad \text{if } q \in (0, 1), \\ v(\beta) &= K_{q,d}^* (\kappa_{q,d} \beta)^\delta (1 + o(1)) \quad \text{as } \beta \rightarrow +\infty, \end{aligned}$$

such that

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \beta \|u\|_{L^q(\mathbb{S}^d)}^2 \geq v(\beta) \|u\|_{L^2(\mathbb{S}^d)}^2 \quad \text{for all } u \in H^1(\mathbb{S}^d). \tag{19}$$

Proof. Inequality (19) is obtained by minimizing the left-hand side the constraint $\|u\|_{L^2(\mathbb{S}^d)} = 1$: there is a minimizer which satisfies

$$-\Delta u + \beta u^{q-1} - v(\beta)u = 0.$$

Case $q \in (1, 2)$. The proof is very similar to that of Lemma 5, so we leave it to the reader. Written for the optimal value of $v(\beta)$, inequality (19) is optimal in the following sense:

- (i) If $0 < \beta \leq d/(2 - q)$, equality is achieved by constants. See [Dolbeault et al. 2013] for rigidity results on \mathbb{S}^d .
- (ii) If $\beta = d/(2 - q)$, the sequence $(u_n)_{n \in \mathbb{N}}$ with $u_n := 1 + (1/n)\varphi$, where φ is an eigenfunction of the Laplace–Beltrami operator, is a minimizing sequence of the quotient to the left-hand side of (19) divided by the right-hand side which converges to the optimal value of $v(\beta) = \beta = d/(2 - q)$, that is,

$$\lim_{n \rightarrow \infty} \frac{\|\nabla u_n\|_{L^2(\mathbb{S}^d)}^2}{\|u_n\|_{L^2(\mathbb{S}^d)}^2 - \|u_n\|_{L^q(\mathbb{S}^d)}^2} = \frac{d}{2 - q}.$$

- (iii) If $\beta > d/(2 - q)$, there exists a nonconstant positive function $u \in H^1(\mathbb{S}^d) \setminus \{0\}$ such that equality holds in (19).

Case $q \in (0, 1]$. In this case, since \mathbb{S}^d is compact, the case $q \leq 1$ does not differ from the case $q \in (1, 2)$ as far as the existence of $v(\beta)$ is concerned. The only difference is that there is no known rigidity result for $q < 1$. However, we can prove that

$$\lim_{\beta \rightarrow 0_+} \frac{v(\beta)}{\beta} = 1.$$

Indeed, let us notice that $v(\beta) \leq \beta$ (use constants as test functions). On the other hand, let $u_\beta = c_\beta + v_\beta$ be a minimizer for $v(\beta)$ such that $c_\beta = \int_{\mathbb{S}^d} u_\beta \, d\sigma$ and, as a consequence, $\int_{\mathbb{S}^d} v_\beta \, d\sigma = 0$. Without loss of generality we can set $\int_{\mathbb{S}^d} |c_\beta + v_\beta|^2 \, d\sigma = c_\beta^2 + \int_{\mathbb{S}^d} |v_\beta|^2 \, d\sigma = 1$. Using the Poincaré inequality, we know that $\|\nabla v_\beta\|_{L^2(\mathbb{S}^d)}^2 \geq d \|v_\beta\|_{L^2(\mathbb{S}^d)}^2$, and hence

$$d \|v_\beta\|_{L^2(\mathbb{S}^d)}^2 + \beta \|c_\beta + v_\beta\|_{L^q(\mathbb{S}^d)}^2 \leq \|\nabla v_\beta\|_{L^2(\mathbb{S}^d)}^2 + \beta \|c_\beta + v_\beta\|_{L^q(\mathbb{S}^d)}^2 = v(\beta) \leq \beta,$$

which shows that $\lim_{\beta \rightarrow 0_+} \|v_\beta\|_{L^2(\mathbb{S}^d)} = 0$ and $\lim_{\beta \rightarrow 0_+} c_\beta = 1$. As a consequence, $\|c_\beta + v_\beta\|_{L^q(\mathbb{S}^d)}^2 = c_\beta^2(1 + o(1))$ as $\beta \rightarrow 0_+$ and we obtain that

$$\beta(1 + o(1)) = \beta c_\beta^2(1 + o(1)) \leq v(\beta),$$

which concludes the proof.

Asymptotic behavior of $v(\beta)$. Finally, the asymptotic behavior of $v(\beta)$ when β is large can be investigated using concentration-compactness methods similar to those used in the proofs of Propositions 8, 9, and 10. Details are left to the reader. □

Proof of Theorem 3. By Hölder’s inequality we have

$$\|u\|_{L^q(\mathbb{S}^d)}^2 = \left(\int_{\mathbb{S}^d} W^{-q/2} (W|u|^2)^{q/2} d\sigma \right)^{2/q} \leq \|W^{-1}\|_{L^{q/(2-q)}(\mathbb{S}^d)} \int_{\mathbb{S}^d} W|u|^2 d\sigma.$$

Using (19), we get

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\sigma + \int_{\mathbb{S}^d} W|u|^2 d\sigma \geq \int_{\mathbb{S}^d} |\nabla u|^2 d\sigma + \|W^{-1}\|_{L^p(\mathbb{S}^d)}^{-1} \|u\|_{L^q(\mathbb{S}^d)}^2 \geq \nu(\|W^{-1}\|_{L^p(\mathbb{S}^d)}^{-1}) \int_{\mathbb{S}^d} |u|^2 d\sigma$$

with $p = q/(2 - q)$, which proves (7). Then Theorem 3 is an easy consequence of Lemma 11. □

5. The threshold case: $q = 2$

The limiting case $q = 2$ in the interpolation inequality (13) corresponds to the logarithmic Sobolev inequality

$$\int_{\mathbb{S}^d} |u|^2 \log \frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} d\sigma \leq \frac{2}{d} \int_{\mathbb{S}^d} |\nabla u|^2 d\sigma \quad \text{for all } u \in H^1(\mathbb{S}^d, d\sigma),$$

which has been studied, for example, in [Beckner 1993; Brouttelande 2003b; 2003a]. For earlier results on the sphere, see [Federbush 1969; Rothaus 1981; Mueller and Weissler 1982] and the references therein (in particular for the circle). Now, if we consider inequality (11), in the limiting case $q = 2$ we obtain the following interpolation inequality.

Lemma 12. *For any $p > \max\{1, d/2\}$, there exists a concave nondecreasing function $\xi : (0, +\infty) \rightarrow \mathbb{R}$ with the properties*

$$\xi(\alpha) = \alpha \quad \text{for all } \alpha \in (0, \alpha_0) \quad \text{and} \quad \xi(\alpha) < \alpha \quad \text{for all } \alpha > \alpha_0$$

for some $\alpha_0 \in [(d/2)(p - 1), (d/2)p]$, and

$$\xi(\alpha) \sim \alpha^{1-d/(2p)} \quad \text{as } \alpha \rightarrow +\infty$$

such that

$$\int_{\mathbb{S}^d} |u|^2 \log \frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} d\sigma + p \log \frac{\xi(\alpha)}{\alpha} \|u\|_{L^2(\mathbb{S}^d)}^2 \leq p \|u\|_{L^2(\mathbb{S}^d)}^2 \log \left(1 + \frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2}{\alpha \|u\|_{L^2(\mathbb{S}^d)}^2} \right)$$

for all $u \in H^1(\mathbb{S}^d)$. (20)

Proof. Consider Hölder’s inequality: $\|u\|_{L^r(\mathbb{S}^d)} \leq \|u\|_{L^2(\mathbb{S}^d)}^\theta \|u\|_{L^q(\mathbb{S}^d)}^{1-\theta}$, with $2 \leq r < q$ and $\theta = \frac{2}{r} \frac{q-r}{q-2}$. To emphasize the dependence of θ in r , we shall write $\theta = \theta(r)$. By taking the logarithm of both sides of the inequality, we find that

$$\frac{1}{r} \log \int_{\mathbb{S}^d} |u|^r d\sigma \leq \frac{\theta(r)}{2} \log \int_{\mathbb{S}^d} |u|^2 d\sigma + \frac{1-\theta(r)}{q} \log \int_{\mathbb{S}^d} |u|^q d\sigma.$$

The inequality becomes an equality when $r = 2$, so that we may differentiate at $r = 2$ and get, with $q = 2p/(p - 1) < 2^*$, that is, $p = q/(q - 2)$, the *logarithmic Hölder inequality*

$$\int_{\mathbb{S}^d} |u|^2 \log \frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} d\sigma \leq p \|u\|_{L^2(\mathbb{S}^d)}^2 \log \frac{\|u\|_{L^q(\mathbb{S}^d)}^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \quad \text{for all } u \in H^1(\mathbb{S}^d).$$

We may now use inequality (11) to estimate

$$\frac{\|u\|_{L^q(\mathbb{S}^d)}^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \leq \frac{\alpha}{\mu(\alpha)} \left(1 + \frac{1}{\alpha} \frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \right),$$

where $\mu = \mu(\alpha)$ is the constant which appears in Lemma 5. Thus we get

$$\int_{\mathbb{S}^d} |u|^2 \log \frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} d\sigma + p \log \frac{\mu(\alpha)}{\alpha} \|u\|_{L^2(\mathbb{S}^d)}^2 \leq p \|u\|_{L^2(\mathbb{S}^d)}^2 \log \left(1 + \frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2}{\alpha \|u\|_{L^2(\mathbb{S}^d)}^2} \right),$$

which proves that the inequality

$$\int_{\mathbb{S}^d} |u|^2 \log \frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} d\sigma + p \log \xi(\alpha) \|u\|_{L^2(\mathbb{S}^d)}^2 \leq p \|u\|_{L^2(\mathbb{S}^d)}^2 \log \left(\alpha + \frac{\|\nabla u\|_{L^2(\mathbb{S}^d)}^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \right)$$

holds for some optimal constant $\xi(\alpha) \geq \mu(\alpha)$, which is therefore concave, and such that $\lim_{\alpha \rightarrow +\infty} \xi(\alpha) = +\infty$. This establishes (20). The fact that equality is achieved for every $\alpha > 0$ follows from the method of [Dolbeault and Esteban 2012, Proposition 3.3].

Testing (20) with constant functions, we find that $\xi(\alpha) \leq \alpha$ for any $\alpha > 0$. On the other hand, $\xi(\alpha) \geq \mu(\alpha) = \alpha$ for any $\alpha \leq d/(q - 2) = (d/2)(p - 1)$. Testing (20) with $u = 1 + \varepsilon\varphi$, we find that $\xi(\alpha) < \alpha$ if $\alpha > (d/2)p$.

By Proposition 10, we know that $\xi(\alpha) \geq \mu(\alpha) \sim \alpha^{1-\vartheta}$ with $\vartheta = d(q - 2)/(2q) = d/(2p)$ as $\alpha \rightarrow +\infty$. As in the proof of Propositions 9 and 10, let us consider an optimal function u_α for (20). Then we have

$$\begin{aligned} & p \log \frac{\xi(\alpha)}{\alpha} \\ &= p \log \left(1 + \frac{1}{\alpha} \|\nabla u_\alpha\|_{L^2(\mathbb{S}^d)}^2 \right) - \int_{\mathbb{S}^d} |u_\alpha|^2 \log |u_\alpha|^2 d\sigma \sim \frac{p}{\alpha} \|\nabla u_\alpha\|_{L^2(\mathbb{S}^d)}^2 - \int_{\mathbb{S}^d} |u_\alpha|^2 \log |u_\alpha|^2 d\sigma \end{aligned}$$

as $\alpha \rightarrow +\infty$ and u_α concentrates at a single point like in the case $q > 2$ so that, after a stereographic projection which transforms u_α into v_α , the function v_α is, up to higher order terms, optimal for the Euclidean logarithmic Sobolev inequality

$$\int_{\mathbb{R}^d} |v|^2 \log \frac{|v|^2}{\|v\|_{L^2(\mathbb{R}^d)}^2} dx + \frac{d}{2} \log(\pi \varepsilon e^2) \|v\|_{L^2(\mathbb{R}^d)}^2 \leq \varepsilon \|\nabla v\|_{L^2(\mathbb{R}^d)}^2,$$

which holds for any $\varepsilon > 0$ and any $v \in H^1(\mathbb{R}^d)$. Here we have of course $\varepsilon = p/\alpha$ and we find that

$$p \log \frac{\xi(\alpha)}{\alpha} = \frac{d}{2} \log \left(\pi \frac{p}{\alpha} e^2 \right) (1 + o(1)) \quad \text{as } \alpha \rightarrow +\infty. \quad \square$$

Corollary 13. *With the notations of Lemma 12, for any $\alpha > 0$, we have*

$$\frac{\alpha}{p} \int_{\mathbb{S}^d} |u|^2 \log \frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} d\sigma + \alpha \log \frac{\xi(\alpha)}{\alpha} \|u\|_{L^2(\mathbb{S}^d)}^2 \leq \|\nabla u\|_{L^2(\mathbb{S}^d)}^2 \quad \text{for all } u \in H^1(\mathbb{S}^d).$$

Proof. This is a straightforward consequence of Lemma 12 using the fact that $\log(1+x) \leq x$ for any $x > 0$. \square

As in the case $q \neq 2$, Corollary 13 provides some spectral estimates. Let $u \in H^1(\mathbb{S}^d)$ be such that $\|u\|_{L^2(\mathbb{S}^d)} = 1$. A straightforward optimization with respect to an arbitrary function W shows that

$$\inf_W \left[\int_{\mathbb{S}^d} W |u|^2 d\sigma + \mu \log \int_{\mathbb{S}^d} e^{-W/\mu} d\sigma \right] = -\mu \int_{\mathbb{S}^d} |u|^2 \log |u|^2 d\sigma,$$

with the optimality case achieved by W such that

$$|u|^2 = \frac{e^{-W/\mu}}{\int_{\mathbb{S}^d} e^{-W/\mu} d\sigma}.$$

Notice that, up to the addition of a constant, we can always assume that $\int_{\mathbb{S}^d} e^{-W/\mu} d\sigma = 1$, which uniquely determines the optimal W . Now, by Corollary 13 applied with $\mu = \alpha/p$, we find that

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\sigma + \int_{\mathbb{S}^d} W |u|^2 d\sigma \geq \alpha \log \frac{\xi(\alpha)}{\alpha} - \frac{\alpha}{p} \log \int_{\mathbb{S}^d} e^{-pW/\alpha} d\sigma.$$

This leads us to the following statement.

Corollary 14. *Let $d \geq 1$. With the notations of Lemma 12, we have the estimate*

$$e^{-\lambda_1(-\Delta-W)/\alpha} \leq \frac{\alpha}{\xi(\alpha)} \left(\int_{\mathbb{S}^d} e^{-pW/\alpha} d\sigma \right)^{1/p}$$

for any function W such that $e^{-pW/\alpha}$ is integrable. This estimate is optimal in the sense that there exists a nonnegative function W for which the inequality becomes an equality.

Appendix A. Further estimates and numerical results

A.1. A refined upper estimate. Let $q \in (2, 2^*)$. For $\alpha > d/(q-2)$, we can give an upper estimate of the optimal constant $\mu(\alpha)$ in inequality (11) of Lemma 5. Consider functions which depend only on z , with the notations of Section 2.3. Then (11) is equivalent to an inequality that can be written as

$$F_\alpha[f] := \frac{\int_{-1}^1 |f'|^2 v dv_d + \alpha \int_{-1}^1 |f|^2 dv_d}{\left(\int_{-1}^1 |f|^q dv_d \right)^{2/q}} \geq \mu(\alpha),$$

where dv_d is the probability measure defined by

$$v_d(z) dz = dv_d(z) := Z_d^{-1} v^{d/2-1} dz \quad \text{with } v(z) := 1-z^2, \quad Z_d := \sqrt{\pi} \frac{\Gamma(d/2)}{\Gamma((d+1)/2)}.$$




Figure 1. In the case $q > 2$, the optimal constant is given by $\mu = \alpha$ for $\alpha \leq d/(q - 2)$ and the curve $\mu = \mu(\alpha)$ for $\alpha > d/(q - 2)$. An upper estimate is given by the curve $\mu = \mu_+(\alpha)$ obtained by optimizing the function $h_\alpha(\varepsilon)$ in terms of $\varepsilon \in (0, 1)$, while a lower estimate, namely $\mu = \mu_-(\alpha) = \alpha_*^\vartheta \alpha^{1-\vartheta}$, has been established in Proposition 8. The asymptotic regime is governed by $\mu(\alpha) \sim \mu_{\text{asyp}}(\alpha) = K_{q,d} \kappa_{q,d}^{-1} \alpha^{1-\vartheta}$ as $\alpha \rightarrow +\infty$ according to Lemma 5. The above plot shows the various curves in the special case $d = 3$ and $q = 3$.

See [Dolbeault et al. 2013] for details. To get an estimate, it is enough to take a well chosen test function. Consider $f_\varepsilon(z) := 1 + \varepsilon\varphi(z)$ and as in Section 2.3 we can choose $\varphi(z) = z$. Then one can optimize $h_\alpha(\varepsilon) = F_\alpha[f_\varepsilon]$ with respect to $\varepsilon \in (0, 1)$, and observe that $\int_{-1}^1 |f'_\varepsilon|^2 v dv_d = d\varepsilon^2 \int_{-1}^1 z^2 dv_d$, so that $h_\alpha(\varepsilon)$ can be written as

$$h_\alpha(\varepsilon) = \frac{\alpha + (d + \alpha)\varepsilon^2 \int_{-1}^1 z^2 dv_d}{\left(\int_{-1}^1 |1 + \varepsilon z|^q dv_d\right)^{2/q}} \geq \mu(\alpha).$$

When $\varepsilon \rightarrow 0_+$, we recover that $h_\alpha(\varepsilon) - \alpha \sim [d - \alpha(q - 2)]\varepsilon^2 \int_{-1}^1 z^2 dv_d < 0$ if $\alpha > d/(q - 2)$, but a better estimate can be achieved simply by considering $\mu_+(\alpha) := \inf_{\varepsilon \in (0,1)} h_\alpha(\varepsilon)$ so that $\mu(\alpha) \leq \mu_+(\alpha) < \alpha$. The function $\alpha \mapsto \mu_+(\alpha)$ can be computed explicitly (using hypergeometric functions) and is shown in Figure 1.

A.2. Numerical results. In this section, we illustrate the various estimates obtained in this paper by numerical computations done in the special case $d = 3$ and $q = 3$. See Figure 1 for the computation of the curve $\alpha \mapsto \mu(\alpha)$ and how it behaves compared to the theoretical estimates obtained in this paper. We emphasize that our upper and lower estimates $\alpha \mapsto \mu_\pm(\alpha)$ bifurcate from the line $\mu = \alpha$ precisely at $\alpha = d/(q - 2)$ if $q \in (2, 2^*)$ (and at $\alpha = d/(2 - q)$ if $q \in (1, 2)$). The curve corresponding to the asymptotic regime is also plotted, but gives relevant information only as $\alpha \rightarrow \infty$.

The convergence towards the asymptotic regime is illustrated in Figure 2 which shows the convergence of $\mu(\alpha)/\mu_{\text{asyp}}(\alpha)$ towards 1 as $\alpha \rightarrow +\infty$ in the special case $d = 3$ and $q = 3$. In terms of spectral properties, for large potentials, eigenvalues of the Schrödinger operator can be estimated according to




Figure 2. The asymptotic regime corresponding to $\alpha \rightarrow +\infty$ has the interesting feature that, up to a dependence in $\alpha^{1-\vartheta}$ and a normalization factor proportional to $\kappa_{q,d}$, the optimal constant $\mu(\alpha)$ behaves like the optimal constant in the Euclidean space, as has been established in Proposition 10.

Theorem 1 by the Euclidean Keller–Lieb–Thirring constant that has been numerically computed for instance in [Barnes 1976].

Appendix B. Constants on the Euclidean space

B.1. Scaling of the Gagliardo–Nirenberg–Sobolev inequality. Let $q > 2$ and denote by $K_{GN}(q)$ the optimal constant in the Gagliardo–Nirenberg–Sobolev inequality, given by

$$K_{GN}(q) := \inf_{u \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\|\nabla u\|_{L^2(\mathbb{R}^d)}^{2\vartheta} \|u\|_{L^2(\mathbb{R}^d)}^{2(1-\vartheta)}}{\|u\|_{L^q(\mathbb{R}^d)}^2} \quad \text{with } \vartheta = \vartheta(q, d) = d \frac{q-2}{2q}.$$

An optimization of the quotient in the definition of $K_{q,d}$, which has been defined in Section 2, allows us to relate this constant with $K_{GN}(q)$. Indeed, if we optimize $\mathcal{N}[u] := \int_{\mathbb{R}^d} |\nabla u|^2 dx + \int_{\mathbb{R}^d} |u|^2 dx$ under the scaling $\lambda \mapsto u_\lambda(x) := \lambda^{d/q} u(\lambda x)$, we find that

$$\mathcal{N}[u_\lambda] = \lambda^{2(1-\vartheta)} \int_{\mathbb{R}^d} |\nabla u|^2 dx + \lambda^{-2\vartheta} \int_{\mathbb{R}^d} |u|^2 dx$$

achieves its minimum at

$$\lambda_\star = \sqrt{\frac{\vartheta}{1-\vartheta}} \frac{\|u\|_{L^2(\mathbb{R}^d)}}{\|\nabla u\|_{L^2(\mathbb{R}^d)}},$$

so that

$$\mathcal{N}[u_{\lambda_\star}] = \vartheta^{-\vartheta} (1-\vartheta)^{-(1-\vartheta)} \|\nabla u\|_{L^2(\mathbb{R}^d)}^{2\vartheta} \|u\|_{L^2(\mathbb{R}^d)}^{2(1-\vartheta)},$$

thus proving that $K_{q,d}$ can be computed in terms of $K_{GN}(q)$ as

$$K_{q,d} = \vartheta^{-\vartheta} (1-\vartheta)^{-(1-\vartheta)} K_{GN}(q).$$

B.2. Asymptotic regimes in Gagliardo–Nirenberg–Sobolev inequalities. Let $q > 2$ and consider the constant $K_{q,d}$ as above. To handle the case of dimension $d = 1$, we may observe that, for any smooth compactly supported function u on \mathbb{R} , we can write either

$$|u(x)|^2 = 2 \left| \int_{-\infty}^x u(y)u'(y) dy \right| \leq \|u\|_{L^2(-\infty,x)}^2 + \|u'\|_{L^2(-\infty,x)}^2 \quad \text{for all } x \in \mathbb{R}$$

or

$$|u(x)|^2 = 2 \left| \int_x^{+\infty} u(y)u'(y) dy \right| \leq \|u\|_{L^2(x,+\infty)}^2 + \|u'\|_{L^2(x,+\infty)}^2 \quad \text{for all } x \in \mathbb{R},$$

thus proving that

$$|u(x)|^2 \leq \frac{1}{2} (\|u\|_{L^2(\mathbb{R})}^2 + \|u'\|_{L^2(\mathbb{R})}^2) \quad \text{for all } x \in \mathbb{R},$$

that is, the Agmon inequality

$$\frac{\|u\|_{L^2(\mathbb{R})}^2 + \|u'\|_{L^2(\mathbb{R})}^2}{\|u\|_{L^\infty(\mathbb{R})}^2} \geq 2,$$

and hence $K_{\infty,1} \geq 2$. Equality is achieved by the function $u(x) = e^{-|x|}$, $x \in \mathbb{R}$, and we have shown that

$$K_{\infty,1} = 2.$$

Proposition 15. Assume that $q > 2$. For all $d \geq 1$,

$$\lim_{q \rightarrow 2_+} K_{q,d} = 1$$

and, for all $d \geq 3$,

$$\lim_{q \rightarrow 2^*} K_{q,d} = S_d,$$

where S_d is the best constant in inequality (8). If $d = 1$, then $\lim_{q \rightarrow +\infty} K_{q,1} = K_{\infty,1}$.

Proof. For any $v \in H^1(\mathbb{R}^d)$ and $d \geq 3$, we have

$$\lim_{q \rightarrow 2^*} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^q(\mathbb{R}^d)}^2} \geq \lim_{q \rightarrow 2^*} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^q(\mathbb{R}^d)}^2} = \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^{2^*}(\mathbb{R}^d)}^2} \geq S_d,$$

thus proving that $\lim_{q \rightarrow 2^*} K_{q,d} \geq S_d$. On the other hand, we may use the Aubin–Talenti function

$$\bar{u}(x) = (1 + |x|^2)^{-(d-2)/2} \quad \text{for all } x \in \mathbb{R}^d \tag{21}$$

as a test function for $K_{q,d}$ if $d \geq 5$, that is,

$$K_{q,d} \leq \vartheta^{-\vartheta} (1 - \vartheta)^{-(1-\vartheta)} \frac{\|\nabla \bar{u}\|_{L^2(\mathbb{R}^d)}^{2\vartheta} \|\bar{u}\|_{L^2(\mathbb{R}^d)}^{2(1-\vartheta)}}{\|\bar{u}\|_{L^q(\mathbb{R}^d)}^2}$$

and observe that the right-hand side converges to S_d , since $\lim_{q \rightarrow 2^*} \vartheta(q, d) = 1$. If $d = 3$ or 4 , standard additional truncations are needed. The case corresponding to $q \rightarrow \infty$, $d = 1$ is dealt with as above.

Now we investigate the limit as $q \rightarrow 2_+$. For any $v \in H^1(\mathbb{R}^d)$, we have

$$\lim_{q \rightarrow 2_+} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^q(\mathbb{R}^d)}^2} \geq \lim_{q \rightarrow 2_+} \frac{\|v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^q(\mathbb{R}^d)}^2} = 1,$$

thus proving that $\lim_{q \rightarrow 2_+} K_{q,d} \geq 1$, and for any $v \in H^1(\mathbb{R}^d)$, the right-hand side in

$$K_{q,d} \leq \vartheta^{-\vartheta} (1 - \vartheta)^{-(1-\vartheta)} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^{2\vartheta} \|v\|_{L^2(\mathbb{R}^d)}^{2(1-\vartheta)}}{\|v\|_{L^q(\mathbb{R}^d)}^2}$$

converges to 1 as $q \rightarrow 2_+$. This completes the proof. □

B.3. Stereographic projection. On $\mathbb{S}^d \subset \mathbb{R}^{d+1}$, we can introduce the coordinates $y = (\rho\phi, z) \in \mathbb{R}^d \times \mathbb{R}$ such that $\rho^2 + z^2 = 1$, $z \in [-1, 1]$, $\rho \geq 0$, and $\phi \in \mathbb{S}^{d-1}$, and consider the stereographic projection

$$\Sigma : \mathbb{S}^d \setminus \{N\} \rightarrow \mathbb{R}^d$$

defined by $\Sigma(y) = x$, where, using the above notations, $x = r\phi$ with $r = \sqrt{(1+z)/(1-z)}$ for any $z \in [-1, 1)$. In this setting, the *north pole* N corresponds to $z = 1$ (and is formally sent at infinity) while the *equator* (corresponding to $z = 0$) is sent onto the unit sphere $\mathbb{S}^{d-1} \subset \mathbb{R}^d$. Hence $x \in \mathbb{R}^d$ is such that $r = |x|$, $\phi = x/|x|$, and we have the useful formulae

$$z = \frac{r^2 - 1}{r^2 + 1} = 1 - \frac{2}{r^2 + 1}, \quad \rho = \frac{2r}{r^2 + 1}.$$

With these notations in hand, we can transform any function u on \mathbb{S}^d into a function v on \mathbb{R}^d using

$$u(y) = \left(\frac{r}{\rho}\right)^{\frac{d-2}{2}} v(x) = \left(\frac{r^2 + 1}{2}\right)^{\frac{d-2}{2}} v(x) = (1 - z)^{-(d-2)/2} v(x),$$

and a painful but straightforward computation shows that, with $\alpha_* = \frac{1}{4}d(d - 2)$,

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\omega + \alpha_* \int_{\mathbb{S}^d} |u|^2 d\omega = \int_{\mathbb{R}^d} |\nabla v|^2 dx \quad \text{and} \quad \int_{\mathbb{S}^d} |u|^q d\omega = \int_{\mathbb{R}^d} |v|^q \left(\frac{2}{1 + |x|^2}\right)^{d-(d-2)q/2} dx.$$

As a consequence, Inequalities (11) and (19) are transformed, respectively, into

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla v|^2 dx + 4(\alpha - \alpha_*) \int_{\mathbb{R}^d} |v|^2 \frac{dx}{(1 + |x|^2)^2} \\ \geq \mu(\alpha) \kappa_{q,d} \left[\int_{\mathbb{R}^d} |v|^q \left(\frac{2}{1 + |x|^2}\right)^{d-(d-2)q/2} dx \right]^{2/q} \quad \text{for all } v \in \mathcal{D}^{1,2}(\mathbb{R}^d) \end{aligned}$$

if $q \in (2, 2^*)$ and $\alpha \geq \alpha_*$, and

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla v|^2 dx + \beta \kappa_{q,d} \left[\int_{\mathbb{R}^d} |v|^q \left(\frac{2}{1 + |x|^2}\right)^{d-(d-2)q/2} dx \right]^{2/q} \\ \geq 4(\nu(\beta) + \alpha_*) \int_{\mathbb{R}^d} |v|^2 \frac{dx}{(1 + |x|^2)^2} \quad \text{for all } v \in \mathcal{D}^{1,2}(\mathbb{R}^d) \end{aligned}$$

if $q \in (1, 2)$ and $\beta > 0$.

B.4. Sobolev’s inequality: expression of the constant and references. The proof that Sobolev’s inequality (8) becomes an equality if and only if $u = \bar{u}$ given by (21) up to a multiplication by a constant, a translation, and a scaling is due to T. Aubin [1976] and G. Talenti [1976]. However, G. Rosen [1971] showed (by linearization) that the function given by (21) is a local minimum when $d = 3$ and computed the critical value.

Much earlier, G. Bliss [1930] (also see [Hardy and Littlewood 1930]) established that, among radial functions, the inequality

$$\left(\int_{\mathbb{R}^d} |f|^p |x|^{r+1-d-p} dx \right)^{\frac{2}{p}} \leq C_{\text{Bliss}} \int_{\mathbb{R}^d} |\nabla f|^2 |x|^{1-d} dx$$

holds when $r = p/2 - 1$. With the change of variables $f(x) = v(|x|^{-1/(d-2)}x/|x|)$, the inequality is changed into

$$\left(\int_{\mathbb{R}^d} |v|^{2d/(d-2)} dx \right)^{\frac{d-2}{d}} \leq \frac{C_{\text{Bliss}}}{(d-2)^{2(d-1)/d}} \int_{\mathbb{R}^d} |\nabla v|^2 dx$$

if $p = 2^*$, and it is a straightforward consequence of [Bliss 1930] that the equality is achieved with $v = \bar{u}$.

According to the duplication formula (see, for instance, [Abramowitz and Stegun 1964]) for the Γ function, we know that

$$\Gamma(x)\Gamma(x + \frac{1}{2}) = 2^{1-2x} \sqrt{\pi} \Gamma(2x).$$

As a consequence, the best constant in Sobolev’s inequality (8) can be written either as

$$S_d = \frac{4}{d(d-2)|\mathbb{S}^d|^{2/d}},$$

where the surface of the d -dimensional unit sphere is given by $|\mathbb{S}^d| = 2\pi^{(d+1)/2} / \Gamma(\frac{d+1}{2})$ (see, for instance, [Beckner 1993]), or as

$$S_d = \frac{1}{\pi d(d-2)} \left(\frac{\Gamma(d)}{\Gamma(d/2)} \right)^{\frac{2}{d}}$$

according to [Aubin 1976; Bliss 1930; Rosen 1971; Talenti 1976]. This last expression can easily be recovered using the fact that optimality in (8) is achieved by \bar{u} defined in (21), while the first one, namely $1/S_d = \frac{1}{4}d(d-2)\kappa_{2^*,d}$, is an easy consequence of the stereographic projection and the computations of Section B.3 with $\alpha = \alpha_*$ and $q = 2^*$.

B.5. A proof of (10). Assume that $q > 2$ and let us relate the optimal constant $L_{\gamma,d}^1$ in the one bound state Keller–Lieb–Thirring inequality (1) with the optimal constant $K_{q,d}$ in the Gagliardo–Nirenberg–Sobolev inequality (9). In this case, recall that $p = q/(q-2) = \gamma + d/2$. For any nonnegative function ϕ defined on \mathbb{R}^d such that $\|\phi\|_{L^p(\mathbb{R}^d)} = K_{q,d}$, using Hölder’s inequality, we can write that

$$\int_{\mathbb{R}^d} (|\nabla v|^2 - \phi|v|^2) dx \geq \|\nabla v\|_{L^2(\mathbb{R}^d)}^2 - \|\phi\|_{L^p(\mathbb{R}^d)} \|v\|_{L^q(\mathbb{R}^d)}^2$$

for any $v \in H^1(\mathbb{R}^d)$. Using (9), namely

$$\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 - K_{q,d} \|v\|_{L^q(\mathbb{R}^d)}^2 \geq -\|v\|_{L^2(\mathbb{R}^d)}^2,$$

this proves that

$$|\lambda_1(-\Delta - \phi)| \leq 1 \quad \text{for all } \phi \in L^p(\mathbb{R}^d) \quad \text{such that} \quad \|\phi\|_{L^p(\mathbb{R}^d)} = K_{q,d}. \quad (22)$$

Next one can observe that inequality (1) can be rephrased as

$$L_{\gamma,d}^1 = \sup_{\phi \in L^p(\mathbb{S}^d)} \sup_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} (\mathcal{R}[v, \phi])^\gamma \quad \text{with } \mathcal{R}[v, \phi] := \frac{\int_{\mathbb{R}^d} (\phi|v|^2 - |\nabla v|^2) dx}{\|v\|_{L^2(\mathbb{R}^d)}^2 \|\phi\|_{L^p(\mathbb{R}^d)}^{2p/(2p-d)}},$$

where $p = \gamma + d/2$ so that the exponent $2p/(2p-d)$ is precisely the one for which we get the scaling invariance of \mathcal{R} . Indeed, with $v_\lambda(x) := v(\lambda x)$ and $\phi_\lambda(x) := \phi(\lambda x)$, we get that $\mathcal{R}[v_\lambda, \lambda^2 \phi_\lambda] = \mathcal{R}[v, \phi]$ for any $\lambda > 0$. Hence we find that

$$\sup_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} \mathcal{R}[v, \phi] = \frac{|\lambda_1(-\Delta - \phi)|}{\|\phi\|_{L^p(\mathbb{R}^d)}^{2p/(2p-d)}} = \sup_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} \mathcal{R}[v_\lambda, \lambda^2 \phi_\lambda] = \frac{|\lambda_1(-\Delta - \lambda^2 \phi_\lambda)|}{\|\lambda^2 \phi_\lambda\|_{L^p(\mathbb{R}^d)}^{2p/(2p-d)}},$$

and if we choose λ such that

$$\lambda^{(2p-d)/p} \|\phi\|_{L^p(\mathbb{R}^d)} = \|\lambda^2 \phi_\lambda\|_{L^p(\mathbb{R}^d)} = K_{q,d},$$

we obtain

$$\frac{|\lambda_1(-\Delta - \phi)|}{\|\phi\|_{L^p(\mathbb{R}^d)}^{2p/(2p-d)}} \leq \frac{1}{K_{q,d}^{2p/(2p-d)}}$$

using (22), which proves that $L_{\gamma,d}^1 \leq (K_{q,d})^{-p}$ with $p = \gamma + d/2$. Since optimality can be preserved at each step, this actually proves (10). See [Keller 1961; Lieb and Thirring 1976; Veling 2002; 2003; Benguria and Loss 2004; Dolbeault et al. 2006] for further details.

In the Euclidean case, notice that the equivalence can be extended to the case of systems on the one hand and to Lieb–Thirring inequalities on the other hand: see [Lieb and Thirring 1976; Lieb 1984; Dolbeault et al. 2006].

B.6. A proof of (18). As in [Dolbeault et al. 2006], we can also relate $L_{-\gamma,d}^1$ and $K_{q,d}^*$ when $q = 2(2\gamma - d)/(2\gamma - d + 2)$ takes values in $(0, 2)$. The method is similar to that of Section B.5. For any function $v \in H^1(\mathbb{R}^d)$ such that v^q is integrable and any positive potential ϕ such that ϕ^{-1} is in $L^p(\mathbb{R}^d)$ with $p = q/(2 - q)$, we can use Hölder’s inequality as in the proof of Theorem 3 and get

$$\int_{\mathbb{R}^d} (|\nabla v|^2 + \phi|v|^2) dx \geq \|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \frac{\|v\|_{L^q(\mathbb{R}^d)}^2}{\|\phi^{-1}\|_{L^p(\mathbb{R}^d)}}.$$

Using (17), namely $\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^q(\mathbb{R}^d)}^2 \geq K_{q,d}^* \|v\|_{L^2(\mathbb{R}^d)}^2$, this proves that

$$\lambda_1(-\Delta + \phi) \geq K_{q,d}^* \quad \text{for all } \phi \in L^p(\mathbb{R}^d) \quad \text{such that} \quad \|\phi^{-1}\|_{L^p(\mathbb{R}^d)} = 1.$$

Inequality (6) can be rephrased as

$$L_{-\gamma,d}^1 = \sup_{\phi \in L^p(\mathbb{S}^d)} \sup_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} (\mathcal{R}[v, \phi])^{-\gamma} \quad \text{with } \mathcal{R}[v, \phi] := \frac{\int_{\mathbb{R}^d} (|\nabla v|^2 + \phi|v|^2) dx}{\|v\|_{L^2(\mathbb{R}^d)}^2} \|\phi^{-1}\|_{L^p(\mathbb{R}^d)}^{p/\gamma}$$

with $\gamma = p + d/2$. The same scaling as in Section B.5 applies: with $v_\lambda(x) := v(\lambda x)$ and $\phi_\lambda(x) := \phi(\lambda x)$, we get that $\mathcal{R}[v_\lambda, \lambda^2 \phi_\lambda] = \mathcal{R}[v, \phi]$ for any $\lambda > 0$, and hence

$$L_{-\gamma,d}^1 = (K_{q,d}^*)^{-\gamma},$$

which completes the proof of (18).

Acknowledgements

Dolbeault and Esteban were partially supported by ANR grants *CBDif* and *NoNAP*. They thank the Mittag–Leffler Institute, where part of this research was carried out, for hospitality.

References

- [Abramowitz and Stegun 1964] M. Abramowitz and I. A. Stegun (editors), *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series **55**, US Government Printing Office, Washington, DC, 1964. Reprinted by Dover, New York, 1974. [MR 29 #4914](#) [Zbl 0171.38503](#)
- [Aubin 1976] T. Aubin, “Problèmes isopérimétriques et espaces de Sobolev”, *J. Differential Geometry* **11**:4 (1976), 573–598. [MR 56 #6711](#) [Zbl 0371.46011](#)
- [Bakry 2006] D. Bakry, “Functional inequalities for Markov semigroups”, pp. 91–147 in *Probability measures on groups: recent directions and trends* (Mumbai, 2002), edited by S. G. Dani and P. Graczyk, Tata Inst. Fund. Res. **18**, Narosa, New Delhi, 2006. [MR 2007g:60086](#) [Zbl 1148.60057](#)
- [Bakry and Ledoux 1996] D. Bakry and M. Ledoux, “Sobolev inequalities and Myers’s diameter theorem for an abstract Markov generator”, *Duke Math. J.* **85**:1 (1996), 253–270. [MR 97h:53034](#) [Zbl 0870.60071](#)
- [Barnes 1976] J. F. Barnes, “Appendix A: Numerical studies”, pp. 295–301 in [Lieb and Thirring 1976].
- [Beckner 1993] W. Beckner, “Sharp Sobolev inequalities on the sphere and the Moser–Trudinger inequality”, *Ann. of Math. (2)* **138**:1 (1993), 213–242. [MR 94m:58232](#) [Zbl 0826.58042](#)
- [Benguria and Loss 2004] R. D. Benguria and M. Loss, “Connection between the Lieb–Thirring conjecture for Schrödinger operators and an isoperimetric problem for ovals on the plane”, pp. 53–61 in *Partial differential equations and inverse problems* (Santiago, 2003), edited by C. Conca et al., Contemp. Math. **362**, Amer. Math. Soc., Providence, RI, 2004. [MR 2005f:81057](#) [Zbl 1087.81042](#)
- [Bentaleb and Fahlaoui 2009] A. Bentaleb and S. Fahlaoui, “Integral inequalities related to the Tchebychev semigroup”, *Semigroup Forum* **79**:3 (2009), 473–479. [MR 2010k:47081](#) [Zbl 1192.47039](#)
- [Bentaleb and Fahlaoui 2010] A. Bentaleb and S. Fahlaoui, “A family of integral inequalities on the circle \mathbb{S}^1 ”, *Proc. Japan Acad. Ser. A Math. Sci.* **86**:3 (2010), 55–59. [MR 2011c:42043](#) [Zbl 1204.47046](#)
- [Bidaut-Véron and Véron 1991] M.-F. Bidaut-Véron and L. Véron, “Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations”, *Invent. Math.* **106**:3 (1991), 489–539. [MR 93a:35045](#) [Zbl 0755.35036](#)
- [Bliss 1930] G. A. Bliss, “An integral inequality”, *J. London Math. Soc.* **5**:1 (1930), 40–46. [MR 1574997](#) [JFM 56.0434.02](#)
- [Brouttelande 2003a] C. Brouttelande, “The best-constant problem for a family of Gagliardo–Nirenberg inequalities on a compact Riemannian manifold”, *Proc. Edinb. Math. Soc. (2)* **46**:1 (2003), 117–146. [MR 2004b:58025](#) [Zbl 1031.58009](#)
- [Brouttelande 2003b] C. Brouttelande, “On the second best constant in logarithmic Sobolev inequalities on complete Riemannian manifolds”, *Bull. Sci. Math.* **127**:4 (2003), 292–312. [MR 2004d:58027](#) [Zbl 1036.58015](#)

- [Dolbeault and Esteban 2012] J. Dolbeault and M. J. Esteban, “Extremal functions for Caffarelli–Kohn–Nirenberg and logarithmic Hardy inequalities”, *Proc. Roy. Soc. Edinburgh Sect. A* **142**:4 (2012), 745–767. MR 2966111 Zbl 1267.26018
- [Dolbeault et al. 2006] J. Dolbeault, P. Felmer, M. Loss, and E. Paturel, “Lieb–Thirring type inequalities and Gagliardo–Nirenberg inequalities for systems”, *J. Funct. Anal.* **238**:1 (2006), 193–220. MR 2008g:35026 Zbl 1104.35021
- [Dolbeault et al. 2013] J. Dolbeault, M. J. Esteban, M. Kowalczyk, and M. Loss, “Sharp interpolation inequalities on the sphere: new methods and consequences”, *Chin. Ann. Math. Ser. B* **34**:1 (2013), 99–112. MR 3011461 Zbl 1263.26029
- [Federbush 1969] P. Federbush, “Partially alternate derivation of a result of Nelson”, *J. Math. Phys.* **10** (1969), 50–52. Zbl 0165.58301
- [Funk 1915] P. Funk, “Beiträge zur Theorie der Kugelfunktionen”, *Mathematische Annalen* **77**:1 (1915), 136–152. MR 1511852 JFM 45.0702.01
- [Hardy and Littlewood 1930] G. H. Hardy and J. E. Littlewood, “Notes on the theory of series, XII: On certain inequalities connected with the calculus of variations”, *J. London Math. Soc.* **5**:1 (1930), 34–39. MR 1574995 JFM 56.0434.01
- [Hecke 1917] E. Hecke, “Über orthogonal-invariante Integralgleichungen”, *Math. Ann.* **78**:1 (1917), 398–404. MR 1511908 JFM 46.0632.02
- [Ilyin 1993] A. A. Ilyin, “Lieb–Thirring inequalities on the N -sphere and in the plane, and some applications”, *Proc. London Math. Soc.* (3) **67**:1 (1993), 159–182. MR 94d:35129 Zbl 0789.58079
- [Ilyin 2012] A. A. Ilyin, “Lieb–Thirring inequalities on some manifolds”, *J. Spectr. Theory* **2**:1 (2012), 57–78. MR 2879309 Zbl 06033370
- [Keller 1961] J. B. Keller, “Lower bounds and isoperimetric inequalities for eigenvalues of the Schrödinger equation”, *J. Math. Phys.* **2** (1961), 262–266. MR 22 #11847 Zbl 0099.06901
- [Ledoux 2000] M. Ledoux, “The geometry of Markov diffusion generators”, *Ann. Fac. Sci. Toulouse Math.* (6) **9**:2 (2000), 305–366. MR 2002a:58045 Zbl 0980.60097
- [Levin 2006] D. Levin, “On some new spectral estimates for Schrödinger-like operators”, *Cent. Eur. J. Math.* **4**:1 (2006), 123–137. MR 2007b:35074 Zbl 1128.35076
- [Levin and Solomyak 1997] D. Levin and M. Solomyak, “The Rozenblum–Lieb–Cwikel inequality for Markov generators”, *J. Anal. Math.* **71** (1997), 173–193. MR 98j:47090 Zbl 0910.47017
- [Lieb 1976] E. H. Lieb, “Bounds on the eigenvalues of the Laplace and Schroedinger operators”, *Bull. Amer. Math. Soc.* **82**:5 (1976), 751–753. MR 53 #11679 Zbl 0329.35018
- [Lieb 1983] E. H. Lieb, “Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities”, *Ann. of Math.* (2) **118**:2 (1983), 349–374. MR 86i:42010 Zbl 0527.42011
- [Lieb 1984] E. H. Lieb, “On characteristic exponents in turbulence”, *Comm. Math. Phys.* **92**:4 (1984), 473–480. MR 86c:35114 Zbl 0598.76054
- [Lieb and Thirring 1976] E. H. Lieb and W. E. Thirring, “Inequalities for the moments of the eigenvalues of the Schrödinger Hamiltonian and their relation to Sobolev inequalities”, pp. 269–303 in *Studies in mathematical physics: essays in honor of Valentine Bargmann*, edited by E. H. Lieb et al., Princeton University Press, 1976. Reprinted as pp. 205–239 in *The stability of matter: from atoms to stars* (Selecta of Elliott H. Lieb), edited by W. Thirring, Springer, Berlin, 2005. Zbl 0342.35044
- [Mueller and Weissler 1982] C. E. Mueller and F. B. Weissler, “Hypercontractivity for the heat semigroup for ultraspherical polynomials and on the n -sphere”, *J. Funct. Anal.* **48**:2 (1982), 252–283. MR 83m:47036 Zbl 0506.46022
- [Ouhabaz and Poupaud 2010] E. M. Ouhabaz and C. Poupaud, “Remarks on the Cwikel–Lieb–Rozenblum and Lieb–Thirring estimates for Schrödinger operators on Riemannian manifolds”, *Acta Appl. Math.* **110**:3 (2010), 1449–1459. MR 2011c:58043 Zbl 1192.58018
- [Rosen 1971] G. Rosen, “Minimum value for c in the Sobolev inequality $\phi^3 \leq c \nabla \phi^3$ ”, *SIAM J. Appl. Math.* **21** (1971), 30–32. MR 44 #6927 Zbl 0201.38704
- [Rothaus 1981] O. S. Rothaus, “Logarithmic Sobolev inequalities and the spectrum of Schrödinger operators”, *J. Funct. Anal.* **42**:1 (1981), 110–120. MR 83f:58080b Zbl 0471.58025
- [Talenti 1976] G. Talenti, “Best constant in Sobolev inequality”, *Ann. Mat. Pura Appl.* (4) **110** (1976), 353–372. MR 57 #3846 Zbl 0353.46018

[Veling 2002] E. J. M. Veling, “Lower bounds for the infimum of the spectrum of the Schrödinger operator in \mathbb{R}^N and the Sobolev inequalities”, *J. Inequal. Pure Appl. Math.* **3**:4 (2002), Article ID #63. MR 2003g:35039 Zbl 1127.35323

[Veling 2003] E. J. M. Veling, “Corrigendum on the paper: “Lower bounds for the infimum of the spectrum of the Schrödinger operator in \mathbb{R}^N and the Sobolev inequalities” [J. Inequal. Pure Appl. Math. **3**:4 (2002), Article ID #63]”, *J. Inequal. Pure Appl. Math.* **4**:5 (2003), Article ID #109. MR 2048612 Zbl 02107677

Received 7 Jan 2013. Accepted 13 Jun 2013.

JEAN DOLBEAULT: dolbeaul@ceremade.dauphine.fr

Ceremade CNRS UMR 7534, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris 16, France

MARIA J. ESTEBAN: esteban@ceremade.dauphine.fr

Ceremade CNRS UMR 7534, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris 16, France

ARI LAPTEV: a.laptev@imperial.ac.uk

Department of Mathematics, Imperial College London, Huxley Building, 180 Queen’s Gate, London SW7 2AZ, United Kingdom

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 2 2014

Two-phase problems with distributed sources: regularity of the free boundary DANIELA DE SILVA, FAUSTO FERRARI and SANDRO SALSA	267
Miura maps and inverse scattering for the Novikov–Veselov equation PETER A. PERRY	311
Convexity of average operators for subsolutions to subelliptic equations ANDREA BONFIGLIOLI, ERMANNO LANCONELLI and ANDREA TOMMASOLI	345
Global uniqueness for an IBVP for the time-harmonic Maxwell equations PEDRO CARO and TING ZHOU	375
Convexity estimates for hypersurfaces moving by convex curvature functions BEN ANDREWS, MAT LANGFORD and JAMES MCCOY	407
Spectral estimates on the sphere JEAN DOLBEAULT, MARIA J. ESTEBAN and ARI LAPTEV	435
Nondispersive decay for the cubic wave equation ROLAND DONNINGER and ANIL ZENGINOĞLU	461
A non-self-adjoint Lebesgue decomposition MATTHEW KENNEDY and DILIAN YANG	497
Bohr’s absolute convergence problem for \mathcal{H}_p -Dirichlet series in Banach spaces DANIEL CARANDO, ANDREAS DEFANT and PABLO SEVILLA-PERIS	513