

# ANALYSIS & PDE

Volume 7

No. 4

2014

FRANCESC PERERA, ANDREW TOMS,  
STUART WHITE AND WILHELM WINTER

**THE CUNTZ SEMIGROUP AND STABILITY OF CLOSE  
 $C^*$ -ALGEBRAS**



# THE CUNTZ SEMIGROUP AND STABILITY OF CLOSE $C^*$ -ALGEBRAS

FRANCESC PERERA, ANDREW TOMS, STUART WHITE AND WILHELM WINTER

We prove that separable  $C^*$ -algebras which are completely close in a natural uniform sense have isomorphic Cuntz semigroups, continuing a line of research developed by Kadison–Kastler, Christensen, and Khoshkam. This result has several applications: we are able to prove that the property of stability is preserved by close  $C^*$ -algebras provided that one algebra has stable rank one; close  $C^*$ -algebras must have affinely homeomorphic spaces of lower-semicontinuous quasitraces; strict comparison is preserved by sufficient closeness of  $C^*$ -algebras. We also examine  $C^*$ -algebras which have a positive answer to Kadison’s Similarity Problem, as these algebras are completely close whenever they are close. A sample consequence is that sufficiently close  $C^*$ -algebras have isomorphic Cuntz semigroups when one algebra absorbs the Jiang–Su algebra tensorially.

## 1. Introduction

Kadison and Kastler [1972] introduced a metric  $d$  on the  $C^*$ -subalgebras of a given  $C^*$ -algebra by equipping the unit balls of the subalgebras with the Hausdorff metric (in norm). They conjectured that sufficiently close  $C^*$ -subalgebras of  $\mathcal{B}(\mathcal{H})$  should be isomorphic, and this conjecture was recently established by Christensen et al. [2012] when one  $C^*$ -algebra is separable and nuclear. The one-sided version of this result—that a sufficiently close near inclusion of a nuclear separable  $C^*$ -algebra into another  $C^*$ -algebra gives rise to a true inclusion—was later proved by Hirshberg, Kirchberg, and White [Hirshberg et al. 2012]. These results and others (see [Christensen et al. 2010; Cameron et al. 2012]) have given new momentum to the perturbation theory of operator algebras.

The foundational paper [Kadison and Kastler 1972] was concerned with structural properties of close algebras, showing that the type decomposition of a von Neumann algebra transfers to nearby algebras. We continue this theme here asking “Which properties or invariants of  $C^*$ -algebras are preserved by small perturbations?” With the proof of the Kadison–Kastler conjecture, the answer for nuclear separable  $C^*$ -algebras is “all of them”. Here we consider general separable  $C^*$ -algebras where already, there are some results. Sufficiently close  $C^*$ -algebras have isomorphic lattices of ideals [Phillips 1973/74] and algebras whose stabilisations are sufficiently close have isomorphic K-theories [Kirchberg 1996]. This was extended to the Elliott invariant consisting of K-theory, traces, and their natural pairing, in [Christensen et al. 2010]. A natural next step is to consider the Cuntz semigroup of (equivalence classes of) positive

---

Research partially supported by EPSRC (grants No. EP/G014019/1 and No. EP/I019227/1), by the DFG (SFB 878), by NSF (DMS-0969246), by the DGI MICIIN (grant No. MTM2011-28992-C02-01), and by the Comissionat per Universitats i Recerca de la Generalitat de Catalunya. Andrew Toms is partially supported by the 2011 AMS Centennial Fellowship.

*MSC2010:* 46L05, 46L35, 46L85.

*Keywords:*  $C^*$ -algebras, perturbation, Cuntz semigroup, stability, quasitraces, traces.

elements (in the stabilisation) of a  $C^*$ -algebra, due both to its exceptional sensitivity in determining non-isomorphism [Toms 2008], classification results using the semigroup [Robert 2012] and the host of  $C^*$ -algebraic properties that can be formulated as order-theoretic properties of the semigroup; for example there is strong evidence to suggest that the behaviour of the Cuntz semigroup characterises important algebraic regularity properties of simple separable nuclear  $C^*$ -algebras [Matui and Sato 2012; Winter 2010; 2012]. We prove that algebras whose stabilisations are sufficiently close do indeed have isomorphic Cuntz semigroups, a surprising fact given the sensitivity of a Cuntz class to perturbations of its representing positive element. This is in stark contrast with the case of Murray–von Neumann equivalence classes of projections, where classes are stable under perturbations of the representing projection of size strictly less than one. The bridge between these two situations is that we can arrange for the representing positive element of a Cuntz class to be almost a projection in trace. We exploit this fact through the introduction of what we call *very rapidly increasing sequences* of positive contractions, increasing sequences where each element almost acts as a unit on its predecessor.

The Kadison–Kastler metric  $d$  is equivalent to a complete version  $d_{cb}$  (given by applying  $d$  to the stabilisations) if and only if Kadison’s Similarity Problem has a positive solution [Christensen et al. 2010; Cameron et al. 2013]; the latter is known to hold in considerable generality, for instance in the case of  $\mathcal{L}$ -stable algebras [Johanesová and Winter 2012]. We show how this result, and a number of other similarity results for  $C^*$ -algebras, can be put in a common framework using Christensen’s [1977] property  $D_k$  and, building on [Christensen et al. 2010], make a more careful study of automatic complete closeness and its relation to property  $D_k$ . We prove that if an algebra  $A$  has  $D_k$  for some  $k$ , then  $d(A \otimes \mathcal{K}, B \otimes \mathcal{K}) \leq C(k)d(A, B)$ , where  $C(k)$  is a constant independent of  $A$  and  $B$ ; as a consequence, sufficiently close  $C^*$ -algebras have isomorphic Cuntz semigroups provided one algebra is  $\mathcal{L}$ -stable.

Stability is perhaps the most basic property one could study in perturbation theory, yet proving its permanence under small perturbations has seen very little progress. We take a significant step here by proving that stability is indeed preserved if one of the algebras considered has stable rank one. The proof is an application of our permanence result for the Cuntz semigroup. Another application is our proof that stably close  $C^*$ -algebras have affinely homeomorphic spaces of lower semicontinuous 2-quasitraces. This extends and improves results from [Christensen et al. 2010], showing that the affine isomorphism between the trace spaces of stably close  $C^*$ -algebras obtained there is weak\*-weak\*-continuous.

The paper is organized as follows: Section 2 contains the preliminaries on the Cuntz semigroup and the Kadison–Kastler metric; Section 3 establishes the permanence of the Cuntz semigroup under complete closeness; Section 4 discusses property  $D_k$  and proves our permanence result for stability; Section 5 proves permanence for quasitraces.

## 2. Preliminaries

Throughout the paper we write  $A^+$  for the positive elements of a  $C^*$ -algebra  $A$ ,  $A_1$  for the unit ball of  $A$  and  $A_1^+$  for the positive contractions in  $A$ .

In the next two subsections we review the definition and basic properties of the Cuntz semigroup. A complete account can be found in the survey [Ara et al. 2011].

**The Cuntz semigroup.** Let  $A$  be a  $C^*$ -algebra. Let us consider on  $(A \otimes \mathcal{K})^+$  the relation  $a \preceq b$  if  $v_n b v_n^* \rightarrow a$  for some sequence  $(v_n)$  in  $A \otimes \mathcal{K}$ . Let us write  $a \sim b$  if  $a \preceq b$  and  $b \preceq a$ . In this case we say that  $a$  is *Cuntz equivalent* to  $b$ . Let  $\text{Cu}(A)$  denote the set  $(A \otimes \mathcal{K})^+ / \sim$  of Cuntz equivalence classes. We use  $\langle a \rangle$  to denote the class of  $a$  in  $\text{Cu}(A)$ . It is clear that  $\langle a \rangle \leq \langle b \rangle \Leftrightarrow a \preceq b$  defines an order on  $\text{Cu}(A)$ . We also endow  $\text{Cu}(A)$  with an addition operation by setting  $\langle a \rangle + \langle b \rangle := \langle a' + b' \rangle$ , where  $a'$  and  $b'$  are orthogonal and Cuntz equivalent to  $a$  and  $b$  respectively (the choice of  $a'$  and  $b'$  does not affect the Cuntz class of their sum). The semigroup  $W(A)$  is then the subsemigroup of  $\text{Cu}(A)$  of Cuntz classes with a representative in  $\bigcup_n M_n(A)_+$ .

Alternatively,  $\text{Cu}(A)$  can be defined to consist of equivalence classes of countably generated Hilbert modules over  $A$  [Coward et al. 2008]. The equivalence relation boils down to isomorphism in the case that  $A$  has stable rank one, but is rather more complicated in general and as we do not require the precise definition of this relation in the sequel, we omit it. We note, however, that the identification of these two approaches to  $\text{Cu}(A)$  is achieved by associating the element  $\langle a \rangle$  to the class of the Hilbert module  $\overline{a\ell_2(A)}$ .

**The category  $\mathbf{Cu}$ .** The semigroup  $\text{Cu}(A)$  is an object in a category of ordered Abelian monoids denoted by  $\mathbf{Cu}$  introduced in [Coward et al. 2008] with additional properties. Before stating them, we require the notion of order-theoretic compact containment. Let  $T$  be a preordered set with  $x, y \in T$ . We say that  $x$  is *compactly contained* in  $y$ , denoted by  $x \ll y$ , if for any increasing sequence  $(y_n)$  in  $T$  with supremum  $y$ , we have  $x \leq y_{n_0}$  for some  $n_0 \in \mathbb{N}$ . An object  $S$  of  $\mathbf{Cu}$  enjoys the following properties (see [Coward et al. 2008; Ara et al. 2011]), which we use repeatedly in the sequel. In particular the existence of suprema in property P3 is a crucial in our construction of a map between the Cuntz semigroups of stably close  $C^*$ -algebras.

- P1.  $S$  contains a zero element.
- P2. The order on  $S$  is compatible with addition:  $x_1 + x_2 \leq y_1 + y_2$  whenever  $x_i \leq y_i, i \in \{1, 2\}$ .
- P3. Every countable upward directed set in  $S$  has a supremum.
- P4. For each  $x \in S$ , the set  $x_{\ll} = \{y \in S \mid y \ll x\}$  is upward directed with respect to both  $\leq$  and  $\ll$ , and contains a sequence  $(x_n)$  such that  $x_n \ll x_{n+1}$  for every  $n \in \mathbb{N}$  and  $\sup_n x_n = x$ .
- P5. The operation of passing to the supremum of a countable upward directed set and the relation  $\ll$  are compatible with addition: if  $S_1$  and  $S_2$  are countable upward directed sets in  $S$ , then  $S_1 + S_2$  is upward directed and  $\sup(S_1 + S_2) = \sup S_1 + \sup S_2$ , and if  $x_i \ll y_i$  for  $i \in \{1, 2\}$ , then  $x_1 + x_2 \ll y_1 + y_2$ .

We say that a sequence  $(x_n)$  in  $S \in \mathbf{Cu}$  is *rapidly increasing* if  $x_n \ll x_{n+1}$  for all  $n$ . We take the scale  $\Sigma(\text{Cu}(A))$  to be the subset of  $\text{Cu}(A)$  obtained as suprema of increasing sequences from  $A^+$ .

For objects  $S$  and  $T$  from  $\mathbf{Cu}$ , the map  $\phi: S \rightarrow T$  is a morphism in the category  $\mathbf{Cu}$  if

- M1.  $\phi$  is order-preserving;
- M2.  $\phi$  is additive and maps 0 to 0;
- M3.  $\phi$  preserves the suprema of increasing sequences;
- M4.  $\phi$  preserves the relation  $\ll$ .

**The Kadison–Kastler metric.** Let us recall the definition of the metric on the collection of all  $C^*$ -subalgebras of a  $C^*$ -algebra introduced in [Kadison and Kastler 1972].

**Definition 2.1.** Let  $A, B$  be  $C^*$ -subalgebras of a  $C^*$ -algebra  $C$ . Define a metric  $d$  on all such pairs as follows:  $d(A, B) < \gamma$  if and only if for each  $x$  in the unit ball of  $A$  or  $B$ , there is  $y$  in the unit ball of the other algebra such that  $\|x - y\| < \gamma$ .

In this definition, we typically take  $C = \mathcal{B}(\mathcal{H})$  for a Hilbert space  $\mathcal{H}$ . The complete, or stabilised, version of the Kadison–Kastler metric is defined by  $d_{cb}(A, B) = d(A \otimes \mathcal{K}, B \otimes \mathcal{K})$  inside  $C \otimes \mathcal{K}$  (here  $\mathcal{K}$  is the  $C^*$ -algebra of compact operators on  $\ell^2(\mathbb{N})$ ); the notion  $d_{cb}$  is used for this metric as  $d_{cb}(A, B) \leq \gamma$  is equivalent to the condition that  $d(M_n(A), M_n(B)) \leq \gamma$  for every  $n$ .

We repeatedly use the standard fact that if  $d(A, B) < \gamma$ , then given a positive contraction  $a \in A_1^+$ , there exists a positive contraction  $b \in B_1^+$  with  $\|a - b\| < 2\alpha$ . One way of seeing this is to use the hypothesis  $d(A, B) < \gamma$  to approximate  $a^{1/2}$  by some  $c \in B_1$  with  $\|a^{1/2} - c\| < \gamma$ . Then take  $b = cc^*$  so that

$$\|a - b\| \leq \|a^{1/2}(a^{1/2} - c)\| + \|(a^{1/2} - c^*)c\| < 2\gamma.$$

There is also a one-sided version of closeness introduced by Christensen [1980], which is referred to as a  $\gamma$ -near inclusion:

**Definition 2.2.** Let  $A, B$  be  $C^*$ -subalgebras of a  $C^*$ -algebra  $C$  and let  $\gamma > 0$ . Write  $A \subseteq_\gamma B$  if for every  $x$  in the unit ball of  $B$ , there is  $y \in B$  such that  $\|x - y\| \leq \gamma$  (note that  $y$  need not be in the unit ball of  $B$ ). Write  $A \subset_\gamma B$  if there exists  $\gamma' < \gamma$  with  $A \subseteq_{\gamma'} B$ . As with the Kadison–Kastler metric, we also use complete, or stabilised, near inclusions: write  $A \subseteq_{cb, \gamma} B$  when  $A \otimes M_n \subseteq_\gamma B \otimes M_n$  for all  $n$ , and  $A \subset_{cb, \gamma} B$  when there exists  $\gamma' < \gamma$  with  $A \subseteq_{cb, \gamma'} B$ .

### 3. Very rapidly increasing sequences and the Cuntz semigroup

We start by noting that, for close  $C^*$ -algebras of real rank zero, an isomorphism between their Cuntz semigroups can be deduced from existing results in the literature. For a  $C^*$ -algebra  $A$ , let  $V(A)$  be the Murray and von Neumann semigroup of equivalence classes of projections in  $\bigcup_{n=1}^\infty A \otimes M_n$  and write  $\Sigma(V(A)) = \{[p] \in V(A) \mid p = p^2 = p^* \in A\}$ . This is a local semigroup in the sense that if  $p, q, p'$  and  $q'$  are projections in  $A$  with  $p'q' = 0$  and  $p \sim p', q \sim q'$ , then  $[p] + [q] = [p' + q'] \in \Sigma(V(A))$ . Recall that, if  $A$  has real rank zero, then the work of Zhang [1990] shows that  $V(A)$  has the Riesz refinement property. By definition, this means that whenever  $x_1, \dots, x_n, y_1, \dots, y_m \in V(A)$  satisfy  $\sum_i x_i = \sum_j y_j$ , then there exist  $z_{i,j} \in V(A)$  with  $\sum_j z_{i,j} = x_i$  and  $\sum_i z_{i,j} = y_j$  for each  $i, j$ . The case  $m = n = 2$  of this can be found as [Ara and Pardo 1996, Lemma 2.3], and the same proof works in general.

The Cuntz semigroup of a  $C^*$ -algebra of real rank zero is completely determined by its semigroup of projections (see [Perera 1997] when  $A$  additionally has stable rank one and [Antoine et al. 2011] for the general case). We briefly recall how this is done. An *interval* in  $V(A)$  is a nonempty, order hereditary and upward directed subset  $I$  of  $V(A)$ , which is said to be countably generated provided there is an increasing sequence  $(x_n)$  in  $V(A)$  such that  $I = \{x \in V(A) \mid x \leq x_n \text{ for some } n\}$ . The set of countably generated intervals is denoted by  $\Lambda_\sigma(V(A))$ , and it has a natural semigroup structure. Namely, if  $I$  and  $J$  have

generating sequences  $(x_n)$  and  $(y_n)$  respectively, then  $I + J$  is the interval generated by  $(x_n + y_n)$ . Given a positive element  $a$  in  $A \otimes \mathcal{K}$  in a  $\sigma$ -unital  $C^*$ -algebra of real rank zero  $A$ , put  $I(a) = \{[p] \in V(A) \mid p \preceq a\}$ . The correspondence  $[a] \mapsto I(a)$  defines an ordered semigroup isomorphism  $\text{Cu}(A) \cong \Lambda_\sigma(V(A))$ .

**Theorem 3.1.** *Let  $A$  and  $B$  be  $\sigma$ -unital  $C^*$ -subalgebras of a  $C^*$ -algebra  $C$ , with  $d(A, B) < 1/8$ . If  $A$  has real rank zero, then  $B$  also has real rank zero and  $\text{Cu}(A) \cong \text{Cu}(B)$ .*

*Proof.* That  $B$  has real rank zero follows from [Christensen et al. 2010, Theorem 6.3]. We know from [Phillips and Raeburn 1979, Theorem 2.6] that there is an isomorphism of local semigroups  $\Phi_1: \Sigma(V(A)) \rightarrow \Sigma(V(B))$  (with inverse, say,  $\Psi_1$ ). This is defined as  $\Phi_1[p] = [q]$ , where  $q$  is a projection in  $B$  such that  $\|p - q\| < 1/8$ . Given  $p \in M_n(A)$ , by [Zhang 1990, Theorem 3.2] we can find projections  $\{p_i\}_{i=1, \dots, n}$  in  $A$  such that  $[p] = \sum_i [p_i]$ . Now extend  $\Phi_1$  to  $\Phi: V(A) \rightarrow V(B)$  by  $\Phi([p]) = \sum_i \Phi_1([p_i])$ . Let us check that  $\Phi$  is well defined. If  $[p] = \sum_i [p_i] = \sum_j [q_j]$  for projections  $p_i$  and  $q_j$  in  $A$ , then use refinement to find elements  $a_{ij} \in V(A)$  such that  $[p_i] = \sum_j a_{ij}$  and  $[q_j] = \sum_i a_{ij}$ . We may also clearly choose projections  $z_{ij}, z'_{ij} \in A$  such that  $a_{ij} = [z_{ij}] = [z'_{ij}]$ , and such that  $z_{ij} \perp z_{ik}$  if  $j \neq k$ , and  $z'_{ij} \perp z'_{il}$  if  $i \neq l$ . Then

$$\sum_i \Phi_1([p_i]) = \sum_i \sum_j \Phi_1([z_{ij}]) = \sum_i \sum_j \Phi_1([z'_{ij}]) = \sum_j \sum_i \Phi_1([z'_{ij}]) = \sum_j \Phi_1([q_j]).$$

Clearly  $\Phi$  is additive and  $\Phi|_{\Sigma(V(A))} = \Phi_1$ . Using  $\Psi_1$ , we construct an additive map  $\Psi: V(B) \rightarrow V(A)$ , with  $\Psi|_{\Sigma(V(B))} = \Psi_1$ . Since  $\Psi_1 \circ \Phi_1 = \text{id}_{\Sigma(V(A))}$ , it follows that  $\Psi \circ \Phi = \text{id}_{V(A)}$ . Similarly  $\Phi \circ \Psi = \text{id}_{V(B)}$ .

Since  $\text{Cu}(A) \cong \Lambda_\sigma(V(A))$  and  $\text{Cu}(B) \cong \Lambda_\sigma(V(B))$ , it follows that  $\text{Cu}(A)$  is isomorphic to  $\text{Cu}(B)$ .  $\square$

We turn now to very rapidly increasing sequences. These provide the key tool we use to transfer information between close algebras at the level of the Cuntz semigroup.

**Definition 3.2.** Let  $A$  be a  $C^*$ -algebra. We say that a rapidly increasing sequence  $(a_n)_{n=1}^\infty$  in  $A_1^+$  is *very rapidly increasing* if given  $\varepsilon > 0$  and  $n \in \mathbb{N}$ , there exists  $m_0 \in \mathbb{N}$  such that for  $m \geq m_0$ , there exists  $v \in A_1$  with  $\|(va_m v^*)a_n - a_n\| < \varepsilon$ . Say that a very rapidly increasing sequence  $(a_n)_{n=1}^\infty$  in  $(A \otimes \mathcal{K})_+^1$  represents  $x \in \text{Cu}(A)$  if  $\sup_n \langle a_n \rangle = x$ .

The following two functions are used in the sequel to manipulate very rapidly increasing sequences. Given  $a \in A_+$  and  $\varepsilon > 0$ , write  $(a - \varepsilon)_+$  for  $h_\varepsilon(a)$ , where  $h_\varepsilon$  is the continuous function  $h_\varepsilon(t) = \max(0, t - \varepsilon)$ . For  $0 \leq \beta < \gamma$ , let  $g_{\beta, \gamma}$  be the piecewise linear function on  $\mathbb{R}$  given by

$$g_{\beta, \gamma}(t) = \begin{cases} 0 & \text{if } t \leq \beta, \\ \frac{t - \beta}{\gamma - \beta} & \text{if } \beta < t < \gamma, \\ 1 & \text{if } t \geq \gamma. \end{cases} \tag{3-1}$$

With this notation, the standard example of a very rapidly increasing sequence is given by

$$(g_{2^{-(n+1)}, 2^{-n}}(a))_{n=1}^\infty \quad \text{for } a \in A_1^+.$$

This sequence represents  $\langle a \rangle$ . In this way every element of the Cuntz semigroup of  $A$  is represented by a very rapidly increasing sequence from  $(A \otimes \mathcal{K})_1^+$ . In the next few lemmas we develop properties of very rapidly increasing sequences, starting with a technical observation.

**Lemma 3.3.** *Let  $A$  be a  $C^*$ -algebra and let  $a, b \in A_1^+$  and  $v \in A_1$  satisfy  $\|v^*bva - a\| \leq \delta$  for some  $\delta > 0$ . Suppose that  $0 < \beta < 1$  and  $\gamma \geq 0$  satisfy  $\gamma + \delta\beta^{-1} < 1$ , then  $\langle (a - \beta)_+ \rangle \leq \langle (b - \gamma)_+ \rangle$  in  $\text{Cu}(A)$ .*

*Proof.* Let  $p \in A^{**}$  denote the spectral projection of  $a$  for the interval  $[\beta, 1]$ . When  $p = 0$ , then  $(a - \beta)_+ = 0$  and the result is trivial, so we may assume that  $p \neq 0$ . Then  $ap$  is invertible in  $pA^{**}p$  with inverse  $x$  satisfying  $\|x\| \leq \beta^{-1}$ . Compressing  $(v^*bva - a)$  by  $p$  and multiplying by  $x$ , we have  $\|pv^*bvp - p\| \leq \delta\beta^{-1}$ . Thus

$$\|pv^*(b - \gamma)_+vp - p\| \leq \|(b - \gamma)_+ - b\| + \|pv^*bvp - p\| \leq \gamma + \delta\beta^{-1},$$

and so

$$pv^*(b - \gamma)_+vp \geq (1 - (\gamma + \delta\beta^{-1}))p.$$

As  $p$  acts as a unit on  $(a - \beta)_+$ , we have

$$\begin{aligned} (a - \beta)_+ &= (a - \beta)_+^{1/2} p (a - \beta)_+^{1/2} \\ &\leq (1 - (\gamma + \delta\beta^{-1}))^{-1} (a - \beta)_+^{1/2} pv^*(b - \gamma)_+vp (a - \beta)_+^{1/2} \\ &= (1 - (\gamma + \delta\beta^{-1}))^{-1} (a - \beta)_+^{1/2} v^*(b - \gamma)_+v (a - \beta)_+^{1/2}. \end{aligned}$$

Thus  $(a - \beta)_+ \preceq (b - \gamma)_+$ . □

The next lemma encapsulates the fact that the element of the Cuntz semigroup represented by a very rapidly increasing sequence  $(a_n)_{n=1}^\infty$  of contractions depends only on the behaviour of parts of the  $a_n$  with spectrum near 1.

**Lemma 3.4.** *Let  $(a_n)_{n=1}^\infty$  be a very rapidly increasing sequence in  $A_1^+$ . Then for each  $\lambda < 1$ , the sequence  $\langle (a_n - \lambda)_+ \rangle_{n=1}^\infty$  has the property that for each  $n \in \mathbb{N}$ , there is  $m_0 \in \mathbb{N}$  such that, for  $m \geq m_0$ , we have  $\langle (a_n - \lambda)_+ \rangle \ll \langle (a_m - \lambda)_+ \rangle$ . Furthermore,*

$$\sup_n \langle (a_n - \lambda)_+ \rangle = \sup_n \langle a_n \rangle. \tag{3-2}$$

*Proof.* Fix  $n \in \mathbb{N}$  and  $0 < \varepsilon < \lambda$  and take  $0 < \delta$  small enough that  $\lambda + \varepsilon^{-1}\delta < 1$ . As  $(a_n)_{n=1}^\infty$  is very rapidly increasing, there exists  $m_0$  such that for  $m \geq m_0$ , there exists  $v \in A_1$  with  $\|(v^*a_m v)a_n - a_n\| < \delta$ .

Lemma 3.3 gives

$$\langle (a_n - \varepsilon)_+ \rangle \leq \langle (a_m - \lambda)_+ \rangle,$$

so that  $\langle (a_n - \lambda)_+ \rangle \ll \langle (a_m - \lambda)_+ \rangle$ , as  $\varepsilon < \lambda$ . This shows that  $\langle (a_r - \lambda)_+ \rangle_{r=1}^\infty$  is upward directed and that

$$\langle (a_n - \delta)_+ \rangle \leq \sup_r \langle (a_r - \lambda)_+ \rangle,$$

for all  $n$  and all  $\varepsilon > 0$ , from which (3-2) follows. □

We can modify elements sufficiently far down a very rapidly increasing sequences with contractions so that they almost act as units for positive contractions dominated in the Cuntz semigroup by the sequence.

**Lemma 3.5.** *Let  $A$  be a  $C^*$ -algebra.*

- (1) *Suppose that  $a, b \in A_1^+$  satisfy  $a \lesssim b$ . Then for all  $\varepsilon > 0$ , there exists  $v \in A$  with  $\|v^*bva - a\| \leq \varepsilon$  and  $\|v^*bv\| \leq 1$ .*
- (2) *Let  $(a_n)_{n=1}^\infty$  be a very rapidly increasing sequence in  $A_1^+$  and suppose  $a \in A_1^+$  satisfies  $\langle a \rangle \ll \sup \langle a_n \rangle$ . Then, for every  $\varepsilon > 0$ , there exists  $m_0 \in \mathbb{N}$  such that, for  $m \geq m_0$ , there exists  $v \in A_1$  with  $\|(v^*a_m v)a - a\| < \varepsilon$ .*

*Proof.* (1). Fix  $\varepsilon > 0$  and find  $r > 0$  so that  $\|a^{1+r} - a\| \leq \varepsilon/2$ . Now  $a^r \lesssim b$ , so there exists  $w \in A$  with  $\|a^r - w^*bw\| \leq \varepsilon/4$ . Thus  $\|w^*bw\| \leq 1 + \varepsilon/4$ , and so, writing  $v = (1 + \varepsilon/4)^{-1/2}w$ , we have  $\|v^*bv\| \leq 1$  and  $\|w^*bw - v^*bv\| \leq \varepsilon/4$ . As such  $\|a^r - v^*bv\| \leq \varepsilon/2$  and so

$$\|v^*bva - a\| \leq \|v^*bv - a^r\| \|a\| + \|a^{1+r} - a\| \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

as claimed.

(2) As  $\langle a \rangle \ll \sup \langle a_n \rangle$ , there exists some  $m_1 \in \mathbb{N}$  with  $a \lesssim a_{m_1} \sim a_{m_1}^2$ . Fix  $\varepsilon > 0$  and by part (1), find  $w \in A$  with  $\|(w^*a_{m_1}^2 w)a - a\| < \varepsilon/2$  and  $\|w^*a_{m_1}^2 w\| \leq 1$ . Now set  $\varepsilon' = \varepsilon/(2\|w\|)$  and, as  $(a_n)_{n=1}^\infty$  is very rapidly increasing, find some  $m_0 > m_1$  such that for  $m \geq m_0$  there exists  $t \in A_1$  with  $\|(t^*a_m t)a_{m_1} - a_{m_1}\| \leq \varepsilon'$ . Given such  $m$  and  $t$ , we have

$$\begin{aligned} \|(w^*a_{m_1} t^* a_m t a_{m_1} w)a - a\| &\leq \|w^*a_{m_1}\| \|(t^*a_m t)a_{m_1} - a_{m_1}\| \|w\| \|a\| + \|(w^*a_{m_1}^2 w)a - a\| \\ &\leq \|w\| \varepsilon' + \frac{1}{2} \varepsilon = \varepsilon, \end{aligned}$$

as  $\|w^*a_{m_1}\| \leq 1$ . Thus we can take  $v = t a_{m_1} w \in A_1$ . □

It follows immediately from part (2) above that two very rapidly increasing sequences representing the same element of the Cuntz semigroup can be intertwined to a single very rapidly increasing sequence.

**Proposition 3.6.** *Let  $(a_n)_{n=1}^\infty, (a'_n)_{n=1}^\infty$  be very rapidly increasing sequences in a  $C^*$ -algebra  $A$  representing the same element  $x \in \text{Cu}(A)$ . Then these sequences can be intertwined after telescoping to form a very rapidly increasing sequence which also represents  $x$ , i.e., there exists  $m_1 < m_2 < \dots$  and  $n_1 < n_2 < \dots$  such that  $(a_{m_1}, a'_{n_1}, a_{m_2}, a'_{n_2}, \dots)$  is a very rapidly increasing sequence.*

Given a rapidly increasing sequence in  $A_1^+$ , we can use the functions  $g_{\beta,\gamma}$  from (3-1) to push the spectrum of the elements of the sequence out to 1 and extract a very rapidly increasing sequence representing the same element of the Cuntz semigroup.

**Lemma 3.7.** *Let  $A$  be a  $C^*$ -algebra and  $(a_n)_{n=1}^\infty$  be a rapidly increasing sequence in  $A_1^+$ . There exists a sequence  $(m_n)_{n=1}^\infty$  in  $\mathbb{N}$  such that the sequence  $(g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n))_{n=1}^\infty$  is very rapidly increasing and*

$$\sup_n \langle g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n) \rangle = \sup_n \langle a_n \rangle.$$

*In particular, every element of the scale  $\Sigma(\text{Cu}(A))$  can be expressed as a very rapidly increasing sequence of elements from  $A_1^+$ .*

*Proof.* We will construct the  $m_n$  so that  $a_{n-1} \lesssim g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n)$  and for each  $1 \leq r < n$ , there exists  $v \in A_1$  with

$$\|(v^* g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n)v)g_{2^{-(m_r+1)}, 2^{-m_r}}(a_r) - g_{2^{-(m_r+1)}, 2^{-m_r}}(a_r)\| < 2^{-n}.$$

Fix  $n \in \mathbb{N}$  and suppose  $m_1, \dots, m_{n-1}$  have been constructed with these properties. As  $(g_{2^{-(m+1)}, 2^{-m}}(a_n))_{m=1}^\infty$  is a very rapidly increasing sequence representing  $\langle a_n \rangle$ , and  $\langle a_{n-1} \rangle \ll \langle a_n \rangle$ , there exists  $\tilde{m}_n$  such that  $\langle a_{n-1} \rangle \ll \langle (g_{2^{-(m+1)}, 2^{-m}}(a_n)) \rangle$  for  $m \geq \tilde{m}_n$ . Further, for  $1 \leq r < n$ ,

$$\langle g_{2^{-(m_r+1)}, 2^{-m_r}}(a_r) \rangle \ll \langle a_r \rangle \ll \sup_m \langle (g_{2^{-(m+1)}, 2^{-m}}(a_n)) \rangle$$

and so the required  $m_n$  can be found using part (2) of Lemma 3.5.

The resulting sequence  $(g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n))_{n=1}^\infty$  is very rapidly increasing by construction. Since  $a_{n-1} \lesssim g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n) \lesssim a_n$  for all  $n$ , we have  $\sup_n \langle g_{2^{-(m_n+1)}, 2^{-m_n}}(a_n) \rangle = \sup_n \langle a_n \rangle$ .  $\square$

We now consider the situation where we have two close  $C^*$ -algebras acting on the same Hilbert space. The following lemma ensures that we can produce a well defined map between the Cuntz semigroups.

**Lemma 3.8.** *Let  $A, B$  be  $C^*$ -algebras acting on the same Hilbert space and suppose that  $a \in A_1^+$  and  $b \in B_1^+$  satisfy  $\|a - b\| < 2\alpha$  for some  $\alpha < 1/27$ . Suppose that  $(a_n)_{n=1}^\infty$  is a very rapidly increasing sequence in  $A_1^+$  with  $\langle a \rangle \ll \sup \langle a_n \rangle$ . Then, there exists  $n_0 \in \mathbb{N}$  with the property that for  $n \geq n_0$  and  $b_n \in B_1^+$  with  $\|b_n - a_n\| < 2\alpha$ , we have*

$$\langle (b - 18\alpha)_+ \rangle \ll \langle (b_n - \gamma)_+ \rangle \ll \langle (b_n - 18\alpha)_+ \rangle$$

in  $\text{Cu}(B)$ , for all  $\gamma$  with  $18\alpha < \gamma < 2/3$ .

*Proof.* Fix  $\gamma$  with  $2/3 > \gamma > 18\alpha$  (which is possible as  $\alpha < 1/27$ ). By taking  $\varepsilon = 2\alpha - \|a - b\|$  in Lemma 3.5(2), there exists  $n_0 \in \mathbb{N}$  such that for  $n \geq n_0$ , there exists  $v \in A_1$  with

$$\|(v^* a_n v)a - a\| < 2\alpha - \|a - b\|.$$

Fix such an  $n \geq n_0$  and  $v \in A_1$ , and take  $b_n \in B_1^+$  with  $\|a_n - b_n\| < 2\alpha$  and choose some  $w \in B_1$  with  $\|w - v\| < \alpha$ . We have

$$\|w^* b_n w - v^* a_n v\| \leq 2\|w - v\| + \|b_n - a_n\| < 4\alpha$$

so that

$$\begin{aligned} \|(w^* b_n w)b - b\| &\leq \|((w^* b_n w) - 1)(b - a)\| + \|(w^* b_n w - v^* a_n v)a\| + \|(v^* a_n v)a - a\| \\ &\leq \|b - a\| + 4\alpha + \|(v^* a_n v)a - a\| \\ &\leq 6\alpha. \end{aligned}$$

Taking  $\delta = 6\alpha$ ,  $\beta = 18\alpha$  and  $2/3 > \gamma' > \gamma > 18\alpha$  so that  $\gamma' + \delta\beta^{-1} < 1$ , Lemma 3.3 gives

$$\langle (b - 18\alpha)_+ \rangle \leq \langle (b_n - \gamma')_+ \rangle \ll \langle (b_n - \gamma)_+ \rangle \ll \langle (b_n - 18\alpha)_+ \rangle. \quad \square$$

**Proposition 3.9.** Let  $A, B$  be  $C^*$ -algebras acting on the same Hilbert space with the property that there exists  $\alpha < 1/27$  such that for each  $a \in A_1$  there exists  $b \in B_1$  with  $\|a - b\| < \alpha$ . Then there is a well defined, order-preserving map  $\Phi : \Sigma(\text{Cu}(A)) \rightarrow \Sigma(\text{Cu}(B))$  given by

$$\Phi(\sup\langle a_n \rangle) = \sup\langle (b_n - 18\alpha)_+ \rangle,$$

whenever  $(a_n)_{n=1}^\infty$  is a very rapidly increasing sequence in  $A_1^+$  and  $b_n \in B_1^+$  have  $\|a_n - b_n\| < 2\alpha$  for all  $n \in \mathbb{N}$ . Moreover, if  $d(A, B) < \alpha$  for  $\alpha < 1/42$ , then  $\Phi$  is a bijection with inverse  $\Psi : \Sigma(\text{Cu}(B)) \rightarrow \Sigma(\text{Cu}(A))$  obtained from interchanging the roles of  $A$  and  $B$  in the definition of  $\Phi$ .

*Proof.* Suppose first that  $\alpha < 1/27$ . To see that  $\Phi$  is well defined, we apply [Lemma 3.8](#) repeatedly. Firstly, given a very rapidly increasing sequence  $(a_n)_{n=1}^\infty$  in  $A_1^+$  representing an element  $x \in \Sigma(\text{Cu}(A))$  and a sequence  $(b_n)_{n=1}^\infty$  in  $B_1^+$  with  $\|a_n - b_n\| < 2\alpha$  for all  $n$ , [Lemma 3.8](#) shows that the sequence  $((b_n - 18\alpha)_+)_{n=1}^\infty$  is upward directed. Indeed, for each  $m$ , take  $a = a_m$  and  $b = b_m$  in [Lemma 3.8](#) so that  $\langle (b_m - 18\alpha)_+ \rangle \ll \langle (b_n - 18\alpha)_+ \rangle$  for all sufficiently large  $n$ . As such,  $\sup_n \langle (b_n - 18\alpha)_+ \rangle$  exists in  $\Sigma(\text{Cu}(B))$ .

Secondly, this supremum does not depend on the choice of  $(b_n)_{n=1}^\infty$ . Consider two sequences  $(b_n)_{n=1}^\infty$  and  $(b'_n)_{n=1}^\infty$  satisfying  $\|b_n - a_n\| < 2\alpha$  and  $\|b'_n - a_n\| < 2\alpha$  for all  $n$ . For each  $n$ , [Lemma 3.8](#) shows that there exists  $m_0$  such that, for  $m \geq m_0$ , we have

$$\langle (b_n - 18\alpha)_+ \rangle \ll \langle (b'_m - 18\alpha)_+ \rangle, \quad \text{and} \quad \langle (b'_n - 18\alpha)_+ \rangle \ll \langle (b_m - 18\alpha)_+ \rangle.$$

Thus  $\sup_n \langle (b_n - 18\alpha)_+ \rangle = \sup_n \langle (b'_n - 18\alpha)_+ \rangle$ .

Thirdly, for two very rapidly increasing sequences  $(a'_n)_{n=1}^\infty$  and  $(a_n)_{n=1}^\infty$  in  $A_1^+$  with  $\sup_n \langle a'_n \rangle \leq \sup_n \langle a_n \rangle$ , and sequences  $(b'_n)_{n=1}^\infty$  and  $(b_n)_{n=1}^\infty$  in  $B_1^+$  with  $\|b'_n - a'_n\|, \|b_n - a_n\| < 2\alpha$  for all  $n$ , [Lemma 3.8](#) gives  $\sup_n \langle (b'_n - 18\alpha)_+ \rangle \leq \sup_n \langle (b_n - 18\alpha)_+ \rangle$ . In particular, when  $(a'_n)_{n=1}^\infty$  and  $(a_n)_{n=1}^\infty$  represent the same element of  $\Sigma(\text{Cu}(A))$ , this shows that the map  $\Phi$  given in the proposition is well defined. In general, this third observation also shows that  $\Phi$  is order-preserving.

Now suppose that  $d(A, B) < \alpha < 1/42$  and let  $\Psi : \Sigma(\text{Cu}(B)) \rightarrow \Sigma(\text{Cu}(A))$  be the order-preserving map obtained by interchanging the roles of  $A$  and  $B$  above. Take  $x \in \Sigma(\text{Cu}(A))$  and fix a very rapidly increasing sequence  $(a_n)_{n=1}^\infty$  in  $A_1^+$  representing  $x$ . Fix a sequence  $(b_n)_{n=1}^\infty$  in  $B_1^+$  with  $\|a_n - b_n\| < 2\alpha$  for all  $n$ . For each  $n$ , [Lemma 3.8](#) gives  $m > n$  with

$$\langle (b_n - 18\alpha)_+ \rangle \ll \langle (b_m - \gamma)_+ \rangle \ll \langle (b_m - 18\alpha)_+ \rangle,$$

for any  $\gamma$  with  $18\alpha < \gamma < 2/3$ . Passing to a subsequence if necessary, we can assume this holds for  $m = n + 1$  and hence  $((b_n - 18\alpha)_+)_{n=1}^\infty$  is a rapidly increasing sequence. By [Lemma 3.7](#), there exists a sequence  $(m_n)_{n=1}^\infty$  in  $\mathbb{N}$  so that, defining  $b'_n = g_{2^{-(m_n+1)}, 2^{-m_n}}((b_n - 18\alpha)_+)$ , we have a very rapidly increasing sequence  $(b'_n)_{n=1}^\infty$  in  $B_1^+$  with

$$\sup_n \langle b'_n \rangle = \sup_n \langle (b_n - 18\alpha)_+ \rangle = \Phi(x).$$

Choose a sequence  $(c_n)_{n=1}^\infty$  in  $A_1^+$  with  $\|c_n - b'_n\| < 2\alpha$  for each  $n$  so that the definition of  $\Psi$  gives  $\Psi(\Phi(x)) = \sup\langle (c_n - 18\alpha)_+ \rangle$ . We now show that  $x \leq \Psi(\Phi(x)) \leq x$ .

Fix  $0 < \beta < 1$  with  $\alpha(18 + 24\beta^{-1}) < 1$ . This choice can be made as  $\alpha < 1/42$ . Fix  $n \in \mathbb{N}$ . As  $\langle (b_n - 18\alpha)_+ \rangle \ll \sup_r \langle b'_r \rangle$ , [Lemma 3.5 \(2\)](#) provides  $m_0 \in \mathbb{N}$  such that for  $m \geq m_0$ , there exists  $w \in B_1$  with

$$\|(w^* b'_m w)(b_n - 18\alpha)_+ - (b_n - 18\alpha)_+\| < 2\alpha - \|a_n - b_n\|. \tag{3-3}$$

Take  $v \in A_1$  with  $\|v - w\| < \alpha$ . Then

$$\|(v^* c_m v - w^* b'_m w)\| \leq \|c_m - b'_m\| + 2\|v - w\| < 4\alpha. \tag{3-4}$$

Combining the estimates [\(3-3\)](#), [\(3-4\)](#) and noting that  $\|w^* b'_m w - 1\| \leq 1$  as  $w$  is a contraction, gives

$$\begin{aligned} \|(v^* c_m v) a_n - a_n\| &\leq \|((v^* c_m v) - 1)(a_n - b_n)\| + \|(v^* c_m v - w^* b'_m w) b_n\| \\ &\quad + \|(w^* b'_m w - 1)(b_n - (b_n - 18\alpha)_+)\| + \|(w^* b'_m w)(b_n - 18\alpha)_+ - (b_n - 18\alpha)_+\| \\ &< \|a_n - b_n\| + 4\alpha + 18\alpha + (2\alpha - \|a_n - b_n\|) = 24\alpha. \end{aligned}$$

Taking  $\gamma = 18\alpha$ ,  $\delta = 24\alpha$ , [Lemma 3.3](#) gives  $\langle (a_n - \beta)_+ \rangle \leq \langle (c_m - 18\alpha)_+ \rangle \leq \Psi(\Phi(x))$ . As  $n$  was arbitrary,  $\sup_n \langle (a_n - \beta)_+ \rangle \leq \Psi(\Phi(x))$ . As  $\beta < 1$ , [Lemma 3.4](#) gives  $\sup_n \langle (a_n - \beta)_+ \rangle = \sup_n \langle a_n \rangle = x$ , so that  $x \leq \Psi(\Phi(x))$ .

For the reverse inequality, fix  $k \in \mathbb{N}$  and apply [Lemma 3.8](#) (with the roles of  $A$  and  $B$  reversed,  $b'_k$  playing the role of  $a$ ,  $(b'_n)_{n=1}^\infty$  the role of  $(a_n)$ ) and  $\gamma = 1/2$  (so  $18\alpha < \gamma < 2/3$ ) to find some  $n \in \mathbb{N}$  such that  $\langle (c_k - 18\alpha)_+ \rangle \leq \langle (c_n - 1/2)_+ \rangle$ . Now, just as in the proof of [Lemma 3.8](#), there is  $z \in B_1$  with  $\|(z^* b_{n+1} z) b_n - b_n\| \leq 6\alpha$ . Let  $p \in B^{**}$  be the spectral projection of  $b_n$  for  $[18\alpha, 1]$ , so that, just as in the proof of [Lemma 3.3](#),  $\|z^* b_{n+1} z p - p\| \leq 1/3$ . Fix  $y \in A_1$  with  $\|y - z\| \leq \alpha$ . Since  $p$  is a unit for  $(b_n - 18\alpha)_+$ , it is a unit for  $b'_n = g_{2^{-(m+1)}, 2^{-m}}((b_n - 18\alpha)_+)$ , giving the estimate

$$\begin{aligned} \|y^* a_{n+1} y c_n - c_n\| &\leq \|y^* a_{n+1} y c_n - z^* b_{n+1} z c_n\| + \|(z^* b_{n+1} z - 1)(c_n - b'_n)\| + \|(z^* b_{n+1} z) b'_n - b'_n\| \\ &\leq 4\alpha + 2\alpha + \frac{1}{3} = 6\alpha + \frac{1}{3}. \end{aligned}$$

Take  $\delta = 6\alpha + 1/3$ ,  $\beta = 1/2$  and  $\gamma = 0$ , so that  $\gamma + \beta^{-1}\delta = 2/3 + 12\alpha < 1$ . Thus [Lemma 3.3](#) gives

$$\langle (c_n - 1/2)_+ \rangle \leq \langle a_{n+1} \rangle,$$

and hence

$$\langle (c_k - 18\alpha)_+ \rangle \leq \langle a_{n+1} \rangle \leq x.$$

Taking the supremum over  $k$  gives  $\Psi(\Phi(x)) \leq x$ . □

**Theorem 3.10.** *Let  $A$  and  $B$  be  $C^*$ -algebras acting on the same Hilbert space with  $d_{cb}(A, B) < \alpha < 1/42$ . Then  $(\text{Cu}(A), \Sigma(\text{Cu}(A)))$  is isomorphic to  $(\text{Cu}(B), \Sigma(\text{Cu}(B)))$ . Moreover, an order-preserving isomorphism  $\Phi : \text{Cu}(A) \rightarrow \text{Cu}(B)$  can be defined by  $\Phi(\sup \langle a_n \rangle) = \sup \langle (b_n - 18\alpha)_+ \rangle$ , whenever  $(a_n)_{n=1}^\infty$  is a very rapidly increasing sequence in  $(A \otimes \mathcal{K})_1^+$  and  $(b_n)_{n=1}^\infty$  is a sequence in  $(B \otimes \mathcal{K})_1^+$  with  $\|a_n - b_n\| < 2\alpha$  for all  $n \in \mathbb{N}$ .*

*Proof.* We have  $d(A \otimes \mathcal{K}, B \otimes \mathcal{K}) < \alpha < 1/42$ . By definition,

$$\Sigma(\text{Cu}(A \otimes \mathcal{K})) = \text{Cu}(A) \quad \text{and} \quad \Sigma(\text{Cu}(A \otimes \mathcal{K})) = \text{Cu}(B).$$

By applying Proposition 3.9 to  $A \otimes \mathcal{K}$  and  $B \otimes \mathcal{K}$ , we obtain mutually inverse order-preserving bijections  $\Phi : \text{Cu}(A \otimes \mathcal{K}) \rightarrow \text{Cu}(B \otimes \mathcal{K})$  and  $\Psi : \text{Cu}(B \otimes \mathcal{K}) \rightarrow \text{Cu}(A \otimes \mathcal{K})$ , given by  $\Phi(\text{sup}\langle a_n \rangle) = \text{sup}\langle (b_n - 18\alpha)_+ \rangle$ , whenever  $(a_n)_{n=1}^\infty$  is a very rapidly increasing sequence in  $(A \otimes \mathcal{K})_1^+$  and  $(b_n)_{n=1}^\infty$  is a sequence in  $(B \otimes \mathcal{K})_1^+$  with  $\|a_n - b_n\| < 2\alpha$  for all  $n \in \mathbb{N}$ . Given a very rapidly increasing sequence  $(a_n)_{n=1}^\infty$  in  $A_1^+$  representing an element  $x \in \Sigma(\text{Cu}(A))$ , we can find a sequence  $(b_n)_{n=1}^\infty$  in  $B_1^+$  with  $\|a_n - b_n\| < 2\alpha$ , so that  $\Phi(x) = \text{sup}\langle (b_n - 18\alpha)_+ \rangle \in \Sigma(\text{Cu}(B))$ . Since  $\Phi$  and  $\Phi^{-1}$  are order-preserving bijections, they also preserve the relation  $\ll$  of compact containment and suprema of countable upward directed sets, as these notions are determined by the order relation  $\leq$ . Further, taking  $a_n = b_n = 0$  for all  $n$  shows that  $\Phi(0_{\text{Cu}(A)}) = 0_{\text{Cu}(B)}$ . Finally, note that  $\Phi$  preserves addition: given very rapidly increasing sequences  $(a_n)_{n=1}^\infty$  and  $(a'_n)_{n=1}^\infty$  in  $(A \otimes \mathcal{K})_1^+$  representing  $x$  and  $y$  in  $\text{Cu}(A)$ , the sequence  $(a_n \oplus a'_n)$  is very rapidly increasing in  $M_2(A \otimes \mathcal{K}) \cong A \otimes \mathcal{K}$ . If  $(b_n)_{n=1}^\infty, (b'_n)_{n=1}^\infty$  have  $\|a_n - b_n\|, \|a'_n - b'_n\| < 2\alpha$  for all  $n$ , then

$$\|(a_n \oplus a'_n) - (b_n \oplus b'_n)\| < 2\alpha,$$

and has

$$((b_n \oplus b'_n) - 18\alpha)_+ = (b_n - 18\alpha)_+ \oplus (b'_n - 18\alpha)_+.$$

In this way we see that  $\Phi(x + y) = \Phi(x) + \Phi(y)$ . □

In particular properties of a  $C^*$ -algebra which are determined by its Cuntz semigroup transfer to completely close  $C^*$ -algebras. One of the most notable of these properties is that of strict comparison.

**Corollary 3.11.** *Let  $A$  and  $B$  be  $C^*$ -algebras acting on the same Hilbert space with  $d_{\text{cb}}(A, B) < 1/42$  and suppose that  $A$  has strict comparison. Then so too does  $B$ .*

#### 4. $\mathcal{L}$ -stability and automatic complete closeness

Given a  $C^*$ -algebra  $A \subset \mathcal{B}(\mathcal{H})$ , [Cameron et al. 2013] shows that the metrics  $d(A, \cdot)$  and  $d_{\text{cb}}(A, \cdot)$  are equivalent if and only if  $A$  has a positive answer to Kadison’s [1955] similarity problem. The most useful reformulation of the similarity property for working with close  $C^*$ -algebras is due to Christensen [1982, Theorem 3.1] and Kirchberg [1996]. Combining these, it follows that a  $C^*$ -algebra  $A$  has a positive answer to the similarity problem if and only if  $A$  has Christensen’s [1980] property  $D_k$  for some  $k$ .

**Definition 4.1.** Given an operator  $T \in \mathcal{B}(\mathcal{H})$ , we write  $\text{ad}(T)$  for the derivation  $\text{ad}(T)(x) = xT - Tx$ . A  $C^*$ -algebra  $A$  has *property  $D_k$*  for some  $k > 0$  if, for every nondegenerate representation  $\pi : A \rightarrow \mathcal{B}(\mathcal{H})$ , the inequality

$$d(T, \pi(A)') \leq k \|\text{ad}(T)|_{\pi(A)}\| \tag{4-1}$$

holds for all  $T \in \mathcal{B}(\mathcal{H})$ . A von Neumann algebra  $A$  is said to have the property  $D_k^*$  if the inequality (4-1) holds for all unital normal representations  $\pi$  on  $\mathcal{H}$  and all  $T \in \mathcal{B}(\mathcal{H})$ .

By taking weak\*-limit points, it follows that if  $A$  is a weak\*-dense  $C^*$ -subalgebra of a von Neumann algebra  $\mathcal{M}$  and  $A$  has property  $D_k$ , then  $\mathcal{M}$  has property  $D_k^*$ .

That property  $D_k$  converts near containments to completely bounded near containments originates in [Christensen 1980, Theorem 3.1]. The version we give below improves on the bounds  $\gamma' = 6k\gamma$  from there and  $\gamma' = (1 + \gamma)^{2k} - 1$  from [Christensen et al. 2010, Corollary 2.12].

**Proposition 4.2.** Suppose that  $A$  has property  $D_k$  for some  $k > 0$ . Then for  $\gamma > 0$ , every near inclusion  $A \subseteq_\gamma B$  (or  $A \subset_\gamma B$ ) with  $A$  and  $B$  acting nondegenerately on the same Hilbert space, gives rise to a completely bounded near inclusion  $A \subseteq_{\text{cb}, \gamma'} B$  (or  $A \subset_{\text{cb}, \gamma'}$ ), where  $\gamma' = 2k\gamma$ .

*Proof.* Suppose  $A \subseteq_\gamma B$  is a near inclusion of  $C^*$ -algebras acting nondegenerately on  $\mathcal{H}$  and fix  $n \in \mathbb{N}$ . Let  $C = C^*(A, B)$  and let  $\pi : C \rightarrow \mathcal{B}(\mathcal{H})$  be the universal representation of  $C$ . Then  $\pi(A)''$  has property  $D_k^*$  so that  $\pi(A)'' \subseteq_{\text{cb}, 2k\gamma} \pi(B)''$ , by [Cameron et al.  $\geq$  2014, Proposition 2.2.4]. By definition, for  $n \in \mathbb{N}$  we have  $\pi(A)'' \otimes M_n \subseteq_{2k\gamma} \pi(B)'' \otimes M_n$ . As  $\pi$  is the universal representation of  $C$ , the Hahn–Banach argument used to deduce [Christensen 1980, equation (3)] from [ibid., equation (2)] gives  $A \otimes M_n \subseteq_{2k\gamma} B \otimes M_n$ , as required. The result when we work with strict near inclusions  $A \subset_\gamma B$  follows immediately.  $\square$

$C^*$ -algebras with no bounded traces (such as stable algebras) were shown to have the similarity property in [Haagerup 1983]. Using the property  $D_k$  version of this fact, the previous proposition gives automatic complete closeness when one algebra has no bounded traces. The argument below which transfers the absence of bounded traces to a nearby  $C^*$ -algebra essentially goes back to [Kadison and Kastler 1972, Lemma 9]. We use more recent results in order to get better estimates.

**Corollary 4.3.** Suppose that  $A$  and  $B$  are  $C^*$ -algebras which act nondegenerately on the same Hilbert space and satisfy  $d(A, B) < \gamma$  for  $\gamma < (2 + 6\sqrt{2})^{-1}$ . Suppose that  $A$  has no bounded traces (for example if  $A$  is stable). Then  $B$  has no bounded traces, and therefore  $A \subset_{\text{cb}, 3\gamma} B$ ,  $B \subset_{\text{cb}, 3\gamma} A$  and  $d_{\text{cb}}(A, B) < 6\gamma$ .

*Proof.* Suppose  $d(A, B) < (2 + 6\sqrt{2})^{-1}$  and  $\tau : B \rightarrow \mathbb{C}$  is a bounded trace. Let  $\pi : B \rightarrow \mathcal{B}(\mathcal{H})$  be the GNS-representation of  $B$  corresponding to  $\tau$ . Then there is a larger Hilbert space  $\tilde{\mathcal{H}}$  and a representation  $\tilde{\pi} : C^*(A, B) \rightarrow \mathcal{B}(\tilde{\mathcal{H}})$  such that  $\pi$  is a direct summand of  $\tilde{\pi}|_B$ . That is, the projection  $p$  from  $\tilde{\mathcal{H}}$  onto  $\mathcal{H}$  is central in  $\tilde{\pi}(B)$  and  $\pi(b) = p\tilde{\pi}(b)p$  for all  $b \in B$ . Then, by [Kadison and Kastler 1972, Lemma 5], we have  $d(\tilde{\pi}(A)'', \tilde{\pi}(B)'') \leq d(A, B)$ , and hence there is a projection  $q \in \tilde{\pi}(A)''$  with  $\|p - q\| \leq \gamma/\sqrt{2}$  by [Khoshkam 1984, Lemma 1.10(ii)]. If  $q$  is an infinite projection in  $\tilde{\pi}(A)''$ , then as  $d(A, B) < (2 + 6\sqrt{2})^{-1}$ , one can follow the argument of [Christensen et al. 2010, Lemma 6.1] (using the estimate  $\|p - q\| < \gamma/\sqrt{2}$  in place of  $\|p - q\| < 2\gamma$ ) to see that  $p$  is infinite in  $\tilde{\pi}(B)''$ , giving a contradiction. If  $q$  is finite, then  $q\tilde{\pi}(A)''q$  has a finite trace  $\rho$  and  $\rho \circ \tilde{\pi}|_A$  defines a bounded trace on  $A$ , and again we have a contradiction. Thus  $B$  has no bounded traces.

Theorem 2.4 of [Christensen 1977] shows that a properly infinite von Neumann algebra has property  $D_{3/2}^*$ . As such, every  $C^*$ -algebra with no bounded traces has property  $D_{3/2}$ . Since  $A$  and  $B$  both have property  $D_{3/2}$ , Proposition 4.2 gives  $A \subset_{\text{cb}, 3\gamma} B$  and  $B \subset_{\text{cb}, 3\gamma} A$ , whence  $d_{\text{cb}}(A, B) < 6\gamma$ .  $\square$

**Corollary 4.4.** Suppose that  $A$  and  $B$  are  $C^*$ -algebras which act nondegenerately on the same Hilbert space and satisfy  $d(A, B) < 1/252$  and suppose that  $A$  has no bounded traces (for example, when  $A$  is stable). Then

$$(\text{Cu}(A), \Sigma(\text{Cu}(A))) \cong (\text{Cu}(B), \Sigma(\text{Cu}(B))).$$

*Proof.* Combine [Corollary 4.3](#) with [Theorem 3.10](#), noting that  $6d(A, B) < 1/42$ . □

We can use the Cuntz semigroup to show that stability transfers to close  $C^*$ -algebras provided one algebra has stable rank one. To detect stability for a  $\sigma$ -unital  $C^*$ -algebra we use the following criterion from [\[Ortega et al. 2012, Lemma 5.4\]](#) which reformulates the earlier characterisation from [\[Hjelmborg and Rørdam 1998\]](#).

**Lemma 4.5.** *Let  $A$  be a  $\sigma$ -unital  $C^*$ -algebra and let  $c \in A$  be a strictly positive element. Then,  $A$  is stable if and only if for every  $\epsilon > 0$ , there is  $b \in A^+$  such that  $(c - \epsilon)_+ \perp b$  and  $(c - \epsilon)_+ \precsim b$ .*

Following [\[Robert and Santiago 2010\]](#), we say that a  $C^*$ -algebra  $A$  has *weak cancellation* provided  $\text{Cu}(A)$  satisfies the property that  $x + z \ll y + z$  implies  $x \leq y$ . It was proved in [\[Rørdam and Winter 2010, Theorem 4.3\]](#) that if  $A$  has stable rank one, then  $W(A)$  has the property defining weak cancellation. When  $A$  has stable rank one, so too does  $A \otimes \mathcal{K}$  [\[Rieffel 1983, Theorem 3.6\]](#), and so  $A$  has weak cancellation.

**Lemma 4.6.** *Let  $A$  be a  $\sigma$ -unital  $C^*$ -algebra with weak cancellation. Then  $A$  is stable if and only if  $\text{Cu}(A) = \Sigma(\text{Cu}(A))$ .*

*Proof.* If  $A$  is stable, we have  $\Sigma(\text{Cu}(A)) = \text{Cu}(A)$ . Indeed, given  $n \in \mathbb{N}$ , choose an automorphism  $\theta_n : \mathcal{K} \otimes \mathcal{K} \rightarrow \mathcal{K} \otimes \mathcal{K}$  with  $\theta_n(\mathcal{K} \otimes e_{11}) = K \otimes M_n$  and let  $\psi : A \rightarrow A \otimes \mathcal{K}$  be an isomorphism. Then  $(\psi^{-1} \otimes \text{id}_{\mathcal{K}})(\text{id}_A \otimes \theta_n)(\psi \otimes \text{id}_{\mathcal{K}})$  is an automorphism of  $A \otimes \mathcal{K}$  which maps  $A \otimes e_{11}$  onto  $A \otimes M_n$ . In this way the class of a positive element in  $A \otimes M_n$  lies in the scale  $\Sigma(\text{Cu}(A))$ . For  $x \in (A \otimes \mathcal{K})_+$  and  $\epsilon > 0$ , we have  $(x - \epsilon)_+ \in \bigcup_{n=1}^{\infty} (A \otimes M_n)$ , and hence  $\langle (x - \epsilon)_+ \rangle \in \Sigma(\text{Cu}(A))$ . Since the scale is defined to be closed under suprema, it follows that  $\text{Cu}(A) = \Sigma(\text{Cu}(A))$ .

Conversely, let  $c \in A$  be a strictly positive element so that  $\Sigma(\text{Cu}(A)) = \{x \in \text{Cu}(A) : x \leq \langle c \rangle\}$  and let  $\epsilon > 0$  be given. The hypothesis ensures that  $2\langle c \rangle \leq \langle c \rangle$ , and so we can find  $\delta > 0$  such that  $2\langle (c - \epsilon/4)_+ \rangle \ll \langle (c - \delta)_+ \rangle$ . Now write

$$(c - \delta)_+ = (c - \delta)_{+g_{\epsilon/2, \epsilon}(c)} + (c - \delta)_+(1_{M(A)} - g_{\epsilon/2, \epsilon}(c)),$$

and observe that

$$\langle (c - \delta)_{+g_{\epsilon/2, \epsilon}(c)} \rangle \leq \langle g_{\epsilon/2, \epsilon}(c) \rangle = \langle (c - \epsilon/2)_+ \rangle \ll \langle (c - \epsilon/4)_+ \rangle.$$

We now have that

$$2\langle (c - \epsilon/4)_+ \rangle \ll \langle (c - \delta)_+ \rangle \leq \langle (c - \epsilon/2)_+ \rangle + \langle (c - \delta)_+(1_{M(A)} - g_{\epsilon/2, \epsilon}(c)) \rangle$$

and so weak cancellation enables us to conclude that

$$\langle (c - \epsilon/4)_+ \rangle \leq \langle (c - \delta)_+(1 - g_{\epsilon/2, \epsilon}(c)) \rangle.$$

Let  $b = (c - \delta)_+(1 - g_{\epsilon/2, \epsilon}(c))$ . It is clear that  $b \perp (c - \epsilon)_+$  and that  $(c - \epsilon)_+ \leq (c - \epsilon/4)_+ \precsim b$ . Thus we may invoke [Lemma 4.5](#) to conclude that  $A$  is stable. □

**Theorem 4.7.** *Let  $A$  and  $B$  be  $\sigma$ -unital  $C^*$ -algebras with  $A$  stable and  $d(A, B) < 1/252$  and suppose either  $A$  or  $B$  has stable rank one. Then  $B$  is stable.*

*Proof.* By [Corollary 4.4](#), we have an isomorphism

$$(\text{Cu}(A), \Sigma(\text{Cu}(A))) \cong (\text{Cu}(B), \Sigma(\text{Cu}(B))).$$

Since  $A$  is stable  $\text{Cu}(A) = \Sigma(\text{Cu}(A))$ . Our isomorphism condition now tells us that  $\text{Cu}(B) = \Sigma(\text{Cu}(B))$ . If  $B$  has stable rank one, then it has weak cancellation, whereas if  $A$  has stable rank one,  $A$  has weak cancellation and, as weak cancellation is a property of the Cuntz semigroup, so too does  $B$ . The result now follows from [Lemma 4.6](#).  $\square$

We now turn to the situation in which one  $C^*$ -algebra is  $\mathcal{L}$ -stable. Christensen [\[1977\]](#) shows that McDuff  $\text{II}_1$  factors have property  $D_{5/2}$ , and hence via the estimates of [\[Pisier 1998\]](#), have similarity length at most 5. (In fact, McDuff factors, and more generally  $\text{II}_1$  factors with Murray and von Neumann’s property  $\Gamma$  have length 3 [\[Christensen 2001\]](#), but at present we do not know how to use this fact to obtain better estimates for automatic complete closeness of close factors with property  $\Gamma$ .)

Analogous results have been established in a  $C^*$ -setting, in particular,  $\mathcal{L}$ -stable  $C^*$ -algebras [\[Johanesová and Winter 2012\]](#) and  $C^*$ -algebras of the form  $A \otimes B$ , where  $B$  is nuclear and has arbitrarily large unital matrix subalgebras [\[Pop 2004\]](#) have similarity degree (and hence length) at most 5. See also [\[Li and Shen 2008\]](#). Here we show how to use the original von Neumann techniques from [\[Christensen 1977\]](#) to show that a class of algebras generalising both these examples have property  $D_{5/2}$  (recapturing the upper bound 5 on the length). A similar result has been obtained independently by Hadwin and Li [\[2014, Corollary 1\]](#) working in terms of the similarity degree as opposed to property  $D_k$ . Once we have this  $D_k$  estimate, [Proposition 4.2](#) applies. In particular we obtain uniform estimates on the cb-distance  $d_{\text{cb}}(A, B)$  in terms of  $d(A, B)$  when  $A$  is  $\mathcal{L}$ -stable.

Given a von Neumann algebra  $\mathcal{M} \subset \mathcal{B}(\mathcal{H})$  and  $x \in \mathcal{B}(\mathcal{H})$ , write  $\overline{\text{co}}_{\mathcal{M}}^w(x)$  for the weak\*-closed convex hull of  $\{uxu^* : u \in \mathcal{U}(\mathcal{M})\}$ . If  $\mathcal{M}$  is injective, then by Schwartz’s property P,  $\overline{\text{co}}_{\mathcal{M}}^w(x) \cap \mathcal{M}$  is nonempty for all  $x \in \mathcal{B}(\mathcal{H})$ . Note that for a nondegenerately represented  $C^*$ -algebra  $A \subset \mathcal{B}(\mathcal{H})$ , we have  $\|\text{ad}(T)|_A\| = \|\text{ad}(T)|_{A'}\|$ . We say that an inclusion  $A \subset C$  of  $C^*$ -algebras is nondegenerate if the inclusion map is nondegenerate.

**Proposition 4.8.** Let  $C$  be a  $C^*$ -algebra and  $A, B \subset C$  be commuting nondegenerate  $C^*$ -subalgebras which generate  $C$ . Suppose  $B$  is nuclear and has no nonzero finite-dimensional representations. Then  $C$  has property  $D_{5/2}$ , and hence similarity length at most 5.

*Proof.* Suppose  $C$  is nondegenerately represented on  $\mathcal{H}$  and fix  $x \in \mathcal{B}(\mathcal{H})$ . The nondegeneracy assumption ensures that  $A$  and  $B$  are nondegenerately represented on  $\mathcal{H}$ . Note that  $C''$  has no finite type I part as  $B$  has no nonzero finite-dimensional representations. Let  $p$  be the central projection in  $C''$  so that  $C''p$  is type  $\text{II}_1$  and  $C''(1-p)$  is properly infinite. Fix a unital type  $\text{I}_\infty$  subalgebra  $\mathcal{M}_0 \subset (1-p)C''(1-p)$  and let  $\mathcal{M} = (\mathcal{M}_0 \cup pB)''$  which is injective. By Schwartz’s property P, there exists  $y \in \overline{\text{co}}_{\mathcal{M}}^w(x) \cap (\mathcal{M} \cup \{p\})'$ . As in Theorems 2.3 and 2.4 of [\[Christensen 1977\]](#),  $\|y - x\| \leq \|\text{ad}(x)|_{C''}\|$  and  $\|\text{ad}(y)|_{C''}\| \leq \|\text{ad}(x)|_{C''}\|$ . Write  $y_1 = yp$  and  $y_2 = y(1-p)$ . If  $p \neq 1$ , then the properly infinite algebra  $\mathcal{M}_0$  lies in  $C''(1-p) \cap \{y_2, y_2^*\}'$  and so by [Corollary 2.2](#) of the same reference,  $\|\text{ad}(y_2)|_{C''(1-p)}\| = 2d(y_2, C'(1-p))$ . Take  $x_2 \in C''(1-p)$  with  $\|x_2 - y_2\| = \|\text{ad}(y_2)|_{C''(1-p)}\|/2 \leq \|\text{ad}(x)|_{C''}\|/2$ .

If  $p \neq 0$ , then we argue exactly as in the proof of [Christensen 1977, Proposition 2.8] to produce first  $z_1 \in A'p$  with  $\|y_1 - z_1\| \leq \|\text{ad}(y_1)|_{C''p}\|/2 \leq \|\text{ad}(x)|_{C''}\|/2$ . Continuing with Christensen's proof, as  $B''p$  and  $A''p$  commute,  $\overline{\text{co}}_{B''p}^w(z_1)$  is contained in  $A'p$  and hence there exists  $x_1 \in \overline{\text{co}}_{B''p}^w(z_1) \cap B'p$  with

$$\|x_1 - z_1\| \leq \|\text{ad}(z_1)|_{B'p}\| \leq \|\text{ad}(z_1 - y_1)|_{B'p}\| \leq 2\|z_1 - y_1\| \leq \|\text{ad}(x)|_{C''}\|.$$

Then

$$\|y_1 - x_1\| \leq \|y_1 - z_1\| + \|z_1 - x_1\| \leq \frac{3}{2}\|\text{ad}(x)|_{C''}\|.$$

If  $p = 0$ , take  $x_1 = 0$  and the same inequality holds. The element  $x_1 + x_2 \in C'$  has

$$\begin{aligned} \|x - (x_1 + x_2)\| &\leq \|x - y\| + \|(y_1 - x_1) + (y_2 - x_2)\| \\ &\leq \|\text{ad}(x)|_{C''}\| + \max(\|y_1 - x_2\|, \|y_2 - x_2\|) \leq \frac{5}{2}\|\text{ad}(x)|_{C''}\|. \end{aligned}$$

Therefore  $C$  has property  $D_{5/2}$ , and so by [Pisier 1998, Remark 4.7] has length at most 5. □

**Corollary 4.9.** *Let  $A$  be a  $\mathcal{L}$ -stable  $C^*$ -algebra. Then  $A$  has property  $D_{5/2}$  and length at most 5.*

The main result of [Christensen et al. 2010]—a reference we will abbreviate to [CSSW] for the remainder of this section—is that the similarity property transfers to close  $C^*$ -algebras. This work is carried out with estimates depending on the length and length constant of  $A$ , but it is equally possible to carry out this work entirely in terms of property  $D_k$  so it can be applied to  $\mathcal{L}$ -stable algebras. Our objective is to obtain a version of [CSSW, Corollary 4.6] replacing the hypothesis that  $A$  has length at most  $\ell$  and length constant at most  $K$  with the formally weaker hypothesis that  $A$  has property  $D_k$  (if  $A$  has the specified length and length constants, then it has property  $D_k$  for  $k = K\ell/2$ , conversely if  $A$  has property  $D_k$ , then it has length at most  $\lfloor 2k \rfloor$ , but a length constant estimate is not known in this case; see [Pisier 1998, Remark 4.7]). This enables us to use Corollary 4.9 obtain an isomorphism between the Cuntz semigroups of sufficiently close  $C^*$ -algebras when one algebra is  $\mathcal{L}$ -stable. To achieve a  $D_k$  version of [CSSW, Section 4], we adjust the hypotheses in Lemma 4.1, Theorem 4.2 and Theorem 4.4 of that reference in turn, starting with Lemma 4.1. We begin by isolating a technical observation.

**Lemma 4.10.** *Let  $\mathcal{M}$  be a finite von Neumann algebra with a faithful tracial state acting in standard form on  $\mathcal{H}$  and let  $J$  be the conjugate linear modular conjugation operator inducing an isometric antisomorphism  $x \mapsto JxJ$  of  $\mathcal{M}$  onto  $\mathcal{M}' \cong \mathcal{M}^{\text{op}}$ . Suppose that  $\mathcal{S}$  is another von Neumann algebra acting nondegenerately on  $\mathcal{H}$  with  $\mathcal{M}' \subset_\gamma \mathcal{S}$ . If  $\mathcal{M}$  has property  $D_k^*$ , then  $\mathcal{M}' \subset_{\text{cb}, 2k\gamma} \mathcal{S}$ .*

*Proof.* As  $J$  is isometric,  $\mathcal{M} \subset_\gamma J\mathcal{S}J$ , so that  $\mathcal{M} \subset_{\text{cb}, 2k\gamma} J\mathcal{S}J$  by Proposition 4.2. Now, for each  $n \in \mathbb{N}$ , let  $J_n$  denote the isometric conjugate linear operator of component-wise complex conjugation on  $\mathbb{C}^n$  so that  $J \otimes J_n$  is a conjugate linear isometry on  $\mathcal{H} \otimes \mathbb{C}^n$ . We can conjugate the near inclusion  $\mathcal{M} \otimes M_n \subset_{2k\gamma} J\mathcal{S}J \otimes M_n$  by  $J \otimes J_n$  to obtain  $\mathcal{M}' \otimes M_n \subset_{2k\gamma} \mathcal{S} \otimes M_n$ , as required. □

The next lemma is the modification of [CSSW, Lemma 4.1]. The expression for  $\beta$  below is a slight improvement over that of the original.

**Lemma 4.11.** *Let  $\mathcal{M}$  and  $\mathcal{N}$  be von Neumann algebras of type  $II_1$  faithfully and nondegenerately represented on  $\mathcal{H}$  with common centre  $Z$  which admits a faithful state. Suppose  $d(\mathcal{M}, \mathcal{N}) = \alpha$  and  $\mathcal{M}$  has property  $D_k^*$ . If  $\alpha$  satisfies*

$$24(12\sqrt{2}k + 4k + 1)\alpha < \frac{1}{200},$$

*then  $d(\mathcal{M}', \mathcal{N}') < 2\beta + 1200k\alpha(1 + \beta)$ , where  $\beta = 96k\alpha(600k + 1)$ .*

*Proof.* This amounts to showing that the hypothesis in [CSSW, Lemma 4.1] that  $\mathcal{M}$  contains an weak\*-dense  $C^*$ -algebra  $A$  of length at most  $\ell$  and length constant at most  $K$  can be replaced by the statement that  $\mathcal{M}$  has property  $D_k^*$  (and that the specified expressions on  $\beta$  are valid). The hypothesis that  $\mathcal{M}$  has such a weak\*-dense  $C^*$ -algebra is initially used to see that  $\mathcal{M}$  has property  $D_k$  at the beginning of the lemma and then applied to a unital normal representation to obtain [CSSW, equation (4.5)]. As such property  $D_k^*$  suffices for this estimate.

The other use of this hypothesis comes on p. 385 in the last paragraph of the lemma, to obtain [CSSW, equation (4.28)]. Using the notation of this paragraph, the von Neumann algebra  $T_{\mathcal{M}}$  is a cutdown of  $\mathcal{M}$  acting as  $\mathcal{M} \otimes I_{\mathcal{G}}$  on  $\mathcal{H} \otimes \mathcal{G}$  by the projection  $e_{i_0, i_0}$  from the commutant of  $\mathcal{M}$  on this space. Since  $e_{i_0, i_0}$  is unitarily equivalent in this commutant to a projection of the form  $e \otimes g_0$ , where  $e$  is a projection from the commutant of  $\mathcal{M}$  on  $\mathcal{H}$  of full central support and  $g_0$  is a minimal projection in  $\mathcal{B}(\mathcal{G})$ , it follows that  $e_{i_0, i_0}$  has full central support in the commutant of  $\mathcal{M}$  on  $\mathcal{H} \otimes \mathcal{G}$ . As such  $T_{\mathcal{M}}$  is isomorphic to  $\mathcal{M}$ , so has property  $D_k^*$ . Thus Lemma 4.10 can be applied to the near inclusion  $T'_{\mathcal{M}} \subset_{48(600k\alpha + \alpha)} T'_{\mathcal{N}_2}$  from [CSSW, equation (4.25)] giving

$$T'_{\mathcal{M}} \subset_{cb, 96k(600k\alpha + \alpha)} T'_{\mathcal{N}_2}.$$

It then follows that

$$T_{\mathcal{M}} \overline{\otimes} \mathcal{B}(\ell^2(\Lambda)) \subset_{96k(600k\alpha + \alpha)} T_{\mathcal{N}} \overline{\otimes} \mathcal{B}(\ell^2(\Lambda)),$$

which is precisely [CSSW, equation (4.28)] with our new estimate for  $\beta$  replacing that of the original. We then deduce that  $d(\mathcal{M}', \mathcal{N}') \leq 2\beta + 1200k\alpha(1 + \beta)$  in just the same way that [CSSW, equation (4.30)] is obtained from [CSSW, equation (4.28)]. □

Now we adjust Theorem 4.2 of [CSSW]. The resulting constant  $\beta$  is obtained by taking  $\alpha = 11\gamma$  in the previous lemma. Note that there is an unfortunate omission in the value of  $\beta$  in Theorem 4.2 of [CSSW], which should be given by taking  $\alpha = 11\gamma$  in Lemma 4.1 of [CSSW], so should be  $K((1 + 316800k\gamma + 528\gamma)^\ell - 1)$ ; this has no knock-on consequences to Theorem 4.4 of [CSSW], where the correct value of  $\beta$  is used.

**Lemma 4.12.** *Let  $A$  and  $B$  be  $C^*$ -algebras acting on a Hilbert space and suppose that  $d(A, B) = \gamma$ . Suppose  $A$  has property  $D_k$  and  $24(12\sqrt{2}k + 4k + 1)\gamma < 1/2200$ . Then*

$$d(A', B') \leq 10\gamma + 2\beta + 13200k\gamma(1 + \beta),$$

*where  $\beta = 1056k(600k\gamma + \gamma)$ .*

*Proof.* This amounts to replacing the hypothesis that  $A$  has length at most  $\ell$  and length constant at most  $K$  with the condition that  $A$  has property  $D_k$  in Theorem 4.2 of [CSSW]. The length hypothesis on  $A$

is used to show that certain  $\text{II}_1$  von Neumann closures of  $A$  satisfy [CSSW, Lemma 4.1], but since the weak\*-closure of a  $C^*$ -algebra with property  $D_k$  has property  $D_k^*$ , Lemma 4.11 can be used in place of that lemma. Note that in the proof of [CSSW, Theorem 4.2] the reference to injective von Neumann algebras having property  $D_1$  is incorrect (it is an open question whether  $\prod_{n=1}^\infty M_n$  has the similarity property). The correct statement is that these algebras have property  $D_1^*$ , which is all that is used.  $\square$

Finally we can convert Theorem 4.4 of [CSSW]. Note the typo in the statement of this theorem; the definition of  $\tilde{k}$  should be  $k/(1 - 2\eta - 2k\gamma)$  rather than  $k/(1 - 2\eta - k\gamma)$ . The same change should be made in Corollary 4.6 of [CSSW].

**Proposition 4.13.** Let  $A$  and  $B$  be  $C^*$ -subalgebras of some  $C^*$ -algebra  $C$  with  $d(A, B) < \gamma$  and suppose  $A$  has property  $D_k$ . Write  $\beta = 1056(600k\gamma + \gamma)$  and  $\eta = 10\gamma + 2\beta + 13200k\gamma(1 + \beta)$  and suppose that

$$24(12\sqrt{2}k + 4k + 1)\gamma < \frac{1}{2200}, \quad 2\eta + 2k\gamma < 1. \tag{4-2}$$

Then  $d_{\text{cb}}(A, B) \leq 4\tilde{k}\gamma$ , where

$$\tilde{k} = \frac{k}{1 - 2\eta - 2k\gamma}.$$

*Proof.* We check that  $B$  has property  $D_{\tilde{k}}$ . This amounts to weakening the hypothesis of [CSSW, Theorem 4.4] in just the same way as the preceding lemmas. Applying Lemma 4.12 in place of Theorem 4.2 of [CSSW] in the proof of their Theorem 4.4 shows that under the hypotheses of this proposition  $B$  has property  $D_{\tilde{k}}$ , where

$$\tilde{k} = \frac{k}{1 - 2\eta - 2k\gamma}.$$

This is valid as property  $D_k$  descends to quotients so, following the proof of [CSSW, Theorem 4.4], the algebra  $\rho(A)$  inherits property  $D_k$  allowing the use of Lemma 4.12 above in place of [CSSW, Theorem 4.2]. Note that one should take care with issues of degeneracy here. In particular, the representation  $\pi$  of  $B$  in the proof of Theorem 4.4 of [CSSW] should be assumed nondegenerate.

Proposition 4.2 now shows that  $B \subset_{\text{cb}, 2\tilde{k}} A$  and  $A \subset_{\text{cb}, 2k} B$ . Therefore

$$d_{\text{cb}}(A, B) \leq 2 \max(2\tilde{k}\gamma, 2k\gamma) = 4\tilde{k}\gamma. \quad \square$$

**Corollary 4.14.** Let  $A$  be a  $C^*$ -algebra generated by two commuting nondegenerate  $C^*$ -subalgebras one of which is nuclear and has no finite-dimensional irreducible representations. Suppose that  $A \subset \mathcal{B}(\mathcal{H})$  and  $B$  is another  $C^*$ -subalgebra of  $\mathcal{B}(\mathcal{H})$  with  $d(A, B) < \gamma$  for  $\gamma < 1/6422957$ . Then  $d_{\text{cb}}(A, B) < 1/42$  and  $(\text{Cu}(A), \Sigma(\text{Cu}(A))) \cong (\text{Cu}(B), \Sigma(\text{Cu}(B)))$ .

*Proof.* By Proposition 4.8,  $A$  has property  $D_k$  for  $k = 5/2$  so in Proposition 4.13,  $\beta = 1585056\gamma$  and  $\eta = 3203122\gamma + 52306848000\gamma^2$ , so that  $2\eta + 2k\gamma < 10^{11}\gamma < 1$  for  $\gamma < 10^{-11}$ . The bound on  $\gamma$  ensures that (4-2) holds so that Proposition 4.13 applies. Further this bound gives

$$\frac{4k\gamma}{1 - 2\eta - 2k\gamma} < \frac{1}{42},$$

and so the result follows from Proposition 4.13 and Theorem 3.10.  $\square$

In particular,  $C^*$ -algebras sufficiently close to  $\mathcal{L}$ -stable algebras are automatically completely close and have the Cuntz semigroup of a  $\mathcal{L}$ -stable algebra. The question of whether the property of  $\mathcal{L}$ -stability transfers to sufficiently close subalgebras raised in [Christensen et al. 2012] remains open.

**Corollary 4.15.** *Let  $A$  be a  $\mathcal{L}$ -stable  $C^*$ -algebra and suppose that  $B$  is another  $C^*$ -algebra acting on the same Hilbert space as  $A$  with  $d(A, B) < 1/6422957$ . Then  $d_{cb}(A, B) < 1/42$ , and  $(Cu(A), \Sigma(Cu(A)))$  is isomorphic to  $(Cu(B), \Sigma(Cu(B)))$ . In particular,  $B$  has the Cuntz semigroup of a  $\mathcal{L}$ -stable algebra.*

### 5. Quasitraces

In this section we use our isomorphism between the Cuntz semigroups of completely close  $C^*$ -algebras to give an affine homeomorphism between the lower semicontinuous quasitraces on such algebras. This isomorphism is compatible with the affine isomorphism of the trace spaces of close  $C^*$ -algebras constructed in [Christensen et al. 2010, Section 5].

Given a  $C^*$ -algebra  $A$ , write  $T(A)$  for the cone of lower semicontinuous traces on  $A$  and  $QT_2(A)$  for the cone of lower semicontinuous 2-quasitraces on  $A$ . Precisely, a trace  $\tau$  on  $A$  is a linear function  $\tau : A_+ \rightarrow [0, \infty]$  vanishing at 0 and satisfying the trace identity  $\tau(xx^*) = \tau(x^*x)$  for all  $x \in A$ . A 2-quasitrace is a function  $\tau : A_+ \rightarrow [0, \infty]$  vanishing at 0 which satisfies the trace identity and which is linear on commuting elements of  $A_+$ . Write  $T_s(A)$  for the simplex of tracial states on  $A$  and  $QT_{2,s}(A)$  for the bounded 2-quasitraces on  $A$  of norm one. Lower semicontinuous traces and 2-quasitraces on  $A$  extend uniquely to lower semicontinuous traces and 2-quasitraces respectively on  $A \otimes \mathcal{K}$ ; see [Blanchard and Kirchberg 2004, Remark 2.27(viii)].

In [Elliott et al. 2011, Section 4], Elliott, Robert and Santiago extend earlier work of Blackadar and Handelman, setting out how functionals on  $Cu(A)$  arise from elements of  $QT_2(A)$ . Precisely, a functional on  $Cu(A)$  is a map  $f : Cu(A) \rightarrow [0, \infty]$  which is additive, order-preserving, has  $f(0) = 0$  and preserves the suprema of increasing sequences. Given  $\tau \in QT_2(A)$ , the expression  $d_\tau(\langle a \rangle) = \lim_{n \rightarrow \infty} \tau(a^{1/n})$  gives a well defined functional on  $Cu(A)$ , where we abuse notation by using  $\tau$  to denote the extension of the original lower semicontinuous 2-quasitrace to  $A \otimes \mathcal{K}$ . Alternatively, one can define  $d_\tau$  by  $d_\tau(\langle a \rangle) = \lim_{n \rightarrow \infty} \tau(a_n)$ , where  $(a_n)_{n=1}^\infty$  is any very rapidly increasing sequence from  $(A \otimes \mathcal{K})_1^+$  representing  $\langle a \rangle$ . Conversely, given a functional  $f$  on  $Cu(A)$ , a lower semicontinuous 2-quasitrace on  $A \otimes \mathcal{K}$  (and hence on  $A$ ) is given by  $\tau_f(a) = \int_0^\infty f(\langle (a-t)_+ \rangle) dt$ . With this notation, the assignments  $\tau \mapsto d_\tau$  and  $f \mapsto \tau_f$  are mutually inverse (see [ibid., Proposition 4.2]).

The topology on  $QT_2(A)$  is specified by saying that a net  $(\tau_i)$  in  $QT_2(A)$  converges to  $\tau \in QT_2(A)$  if and only if

$$\limsup_i \tau_i(\langle (a - \varepsilon)_+ \rangle) \leq \tau(a) \leq \liminf_i \tau_i(a)$$

for all  $a \in A_+$  and  $\varepsilon > 0$ . With this topology  $QT_2(A)$  is a compact Hausdorff space [ibid., Theorem 4.4] and  $T(A)$  is compact in the induced topology [ibid., Theorem 3.7]. In a similar fashion, the cone of functionals on  $Cu(A)$  is topologised by defining  $\lambda_i \rightarrow \lambda$  if and only if

$$\limsup_i \lambda_i(\langle (a - \varepsilon)_+ \rangle) \leq \lambda(\langle a \rangle) \leq \liminf_i \lambda_i(\langle a \rangle)$$

for all  $a \in (A \otimes \mathcal{K})_+$  and  $\varepsilon > 0$ . Theorem 4.4 of [ibid.] shows that the affine map  $\tau \mapsto d_\tau$  is a homeomorphism between the cone  $QT_2(A)$  and the cone of functionals on the Cuntz semigroup.

**Theorem 5.1.** (1) *Let  $A, B$  be  $C^*$ -algebras acting nondegenerately on a Hilbert space, with  $d_{cb}(A, B) < 1/42$ . The isomorphism  $\Phi : (\text{Cu}(A), \Sigma(\text{Cu}(A))) \rightarrow (\text{Cu}(B), \Sigma(\text{Cu}(B)))$  given by Theorem 3.10 induces an affine homeomorphism*

$$\widehat{\Phi} : QT_2(B) \rightarrow QT_2(A)$$

satisfying

$$d_{\widehat{\Phi}(\tau)}(x) = d_\tau(\Phi(x)) \tag{5-1}$$

for all  $x \in \text{Cu}(A)$  and  $\tau \in QT_2(B)$ .

(2) *Suppose additionally that  $A$  and  $B$  are unital and  $d_{cb}(A, B) < \gamma < 1/2200$ . Then  $\widehat{\Phi}$  is compatible with the map  $\Psi : T_s(B) \rightarrow T_s(A)$  given in Lemma 5.4 of [Christensen et al. 2010]. Precisely, for  $\tau \in T_s(B)$ , we have  $\widehat{\Phi}(\tau) \in T_s(A) \subset QT_2(A)$  and  $\widehat{\Phi}(\tau) = \Psi(\tau)$ .*

*Proof.* Part (1) of the theorem is a consequence of Theorem 3.10 and [Elliott et al. 2011, Proposition 4.2]: given  $\tau \in QT_2(B)$ , define  $\widehat{\Phi}(\tau)$  to be the lower semicontinuous 2-quasitrace induced by the functional  $d_\tau \circ \Phi$  on  $\text{Cu}(A)$ . It is immediate from the construction that the map  $\widehat{\Phi}$  is affine, bijective and the identity (5-1) holds.

To show that  $\widehat{\Phi}$  is continuous, we use the homeomorphism between the cone of lower semicontinuous quasitraces and functionals on the Cuntz semigroup in [Elliott et al. 2011, Theorem 4.4]. Consider a net  $(\tau_i)$  in  $QT_2(B)$  with  $\tau_i \rightarrow \tau$ . Fix  $a \in A_+$ , then

$$d_\tau(\Phi(\langle a \rangle)) \leq \liminf_i d_{\tau_i}(\Phi(\langle a \rangle)),$$

as  $d_{\tau_i} \rightarrow d_\tau$ . Now take  $\varepsilon > 0$  and fix a contraction  $b \in (B \otimes K)_+$  with  $\Phi(\langle a \rangle) = \langle b \rangle$ . As  $(\langle (b - 1/n)_+ \rangle)_{n=1}^\infty$  is very rapidly increasing with supremum  $\langle b \rangle$ , there exists  $n \in \mathbb{N}$  with  $\Phi(\langle (a - \varepsilon)_+ \rangle) \leq \langle (b - 1/n)_+ \rangle$ . As

$$\limsup_i d_{\tau_i}(\langle (b - \frac{1}{n})_+ \rangle) \leq d_\tau(\langle b \rangle),$$

it follows that

$$\limsup_i d_{\widehat{\Phi}(\tau_i)}(\langle (a - \varepsilon)_+ \rangle) \leq d_{\widehat{\Phi}(\tau)}(\langle a \rangle) \leq \liminf_i d_{\widehat{\Phi}(\tau_i)}(\langle a \rangle).$$

Thus  $d_{\widehat{\Phi}(\tau_i)} \rightarrow d_{\widehat{\Phi}(\tau)}$  and so, using the homeomorphism between  $QT_2(A)$  and functionals on  $\text{Cu}(A)$ , we have  $\widehat{\Phi}(\tau_i) \rightarrow \widehat{\Phi}(\tau)$ . Therefore  $\widehat{\Phi}$  is continuous, and hence a homeomorphism between  $QT_2(B)$  and  $QT_2(A)$ .

For the second part we first need to review the construction of the map  $\Psi$  from [Christensen et al. 2010], which we again abbreviate [CSSW]. Suppose  $d_{cb}(A, B) < \gamma < 1/2200$ . Write  $C = C^*(A, B)$  and let  $C \subset \mathcal{B}(\mathcal{H})$  be the universal representation of  $C$  so that  $\mathcal{M} = A''$  and  $\mathcal{N} = B''$  are isometrically isomorphic to  $A^{**}$  and  $B^{**}$  respectively. Note that the Kaplansky density argument of [Kadison and Kastler 1972, Lemma 5] gives  $d_{cb}(\mathcal{M}, \mathcal{N}) \leq d_{cb}(A, B)$ . Following the proof of [CSSW, Lemma 5.4] we can find a unitary  $u \in (Z(\mathcal{M}) \cup Z(\mathcal{N}))''$  such that  $Z(u\mathcal{M}u^*) = Z(\mathcal{N})$  and  $\|u - 1_C\| \leq 5\gamma$ . We write

$A_1 = uAu^*$  and  $\mathcal{M}_1 = u\mathcal{M}u^*$ . There is now a projection  $z_{\text{fin}} \in Z(\mathcal{M}_1) = Z(\mathcal{N})$  which simultaneously decomposes  $\mathcal{M}_1 = \mathcal{M}_1 z_{\text{fin}} \oplus \mathcal{M}_1(1 - z_{\text{fin}})$  and  $\mathcal{N} = \mathcal{N} z_{\text{fin}} \oplus \mathcal{N}(1 - z_{\text{fin}})$  into the finite and properly infinite parts respectively (see [CSSW, Lemma 3.5] or [Kadison and Kastler 1972]). Given a tracial state  $\tau$  on  $B$ , there is a unique extension  $\tau''$  to  $\mathcal{N}$ , which then factors uniquely through the centre valued trace  $\text{Tr}_{\mathcal{N} z_{\text{fin}}}$  on  $\mathcal{N} z_{\text{fin}}$ . That is,  $\tau''(x) = (\phi_\tau \circ \text{Tr}_{\mathcal{N} z_{\text{fin}}})(x z_{\text{fin}})$  for some state  $\phi_\tau$  on  $\mathcal{N} z_{\text{fin}}$ . The map  $\Psi$  in [CSSW] is then given by defining  $\Psi(\tau)(y) = (\phi_\tau \circ \text{Tr}_{\mathcal{M}_1 z_{\text{fin}}})(uyu^* z_{\text{fin}})$  for  $y \in A$ .

Now fix  $\tau \in T_s(B)$ . For  $m \in \mathbb{N}$  and  $a \in (A \otimes M_m)_1^+$ , consider the standard very rapidly increasing sequence  $(g_{2^{-(n+1)}, 2^{-n}}(a))_{n=1}^\infty$  which represents  $\langle a \rangle$ . Let  $p_n \in \mathcal{M} \otimes M_m$  be the spectral projection for  $a$  for  $[2^{-(n+1)}, 1]$ , so that the alternating sequence

$$g_{2^{-2}, 2^{-1}}(a), p_1, g_{2^{-3}, 2^{-2}}(a), p_2, g_{2^{-4}, 2^{-3}}(a), p_3, \dots$$

is very rapidly increasing. Then

$$d_{\Psi(\tau)}(\langle a \rangle) = \sup_n (\Psi(\tau))(g_{2^{-n}, 2^{-(n+1)}}(a)) = \sup_n \Psi(\tau)''(p_n). \tag{5-2}$$

Choose  $b_n \in (B \otimes M_m)_1^+$  with  $\|g_{2^{-(n+1)}, 2^{-n}}(a) - b_n\| \leq 2\gamma$  and projections  $q_n \in \mathcal{N} \otimes M_m$  with  $\|p_n - q_n\| \leq 2\gamma$  by a standard functional calculus argument [Christensen 1974/75, Lemma 2.1]. Note that  $d_{\text{cb}}(\mathcal{M}_1, \mathcal{N}) \leq 11\gamma$  and the algebras  $(\mathcal{M}_1 \otimes M_m)(z_{\text{fin}} \otimes 1_m)$  and  $(\mathcal{N}_1 \otimes M_m)(z_{\text{fin}} \otimes 1_m)$  have the same centre. Since

$$\|(u \otimes 1_m)p(u \otimes 1_m)^*(z_{\text{fin}} \otimes 1_m) - q(z_{\text{fin}} \otimes 1_m)\| < \frac{1}{2},$$

Lemma 3.6 of [CSSW] applies to show that

$$(\text{Tr}_{\mathcal{M}_1 z_{\text{fin}}} \otimes \text{tr}_m)((u \otimes 1_m)p_n(u \otimes 1_m)^*(z_{\text{fin}} \otimes 1_m)) = (\text{Tr}_{\mathcal{N} z_{\text{fin}}} \otimes \text{tr}_m)(q(z_{\text{fin}} \otimes 1_m)).$$

This ensures that  $\Psi(\tau)''(p_n) = \tau''(q_n)$  for all  $n$ .

As each  $(q_n - 18\gamma)_+ = q_n$ , the sequence

$$(b_1 - 18\gamma)_+, q_1, (b_2 - 18\gamma)_+, q_2, (b_3 - 18\gamma)_+, q_3, \dots$$

is upwards directed by Lemma 3.8 and the supremum of this sequence defines  $\Phi(\langle a \rangle)$ . We then have

$$d_\tau(\Phi(\langle a \rangle)) = \sup_n \tau''(q_n). \tag{5-3}$$

Indeed,  $d_\tau(\Phi(\langle a \rangle))$  is given by  $\sup \tau(c_n)$ , where  $(c_n)_{n=1}^\infty$  is any very rapidly increasing sequence in  $(B \otimes \mathcal{K})_+$  representing  $\Phi(\langle a \rangle)$ . But, working in  $\text{Cu}(\mathcal{N})$ , Proposition 3.6 shows that any such very rapidly increasing sequence  $(c_n)_{n=1}^\infty$  can be intertwined with the very rapidly increasing sequence  $(q_n)_{n=1}^\infty$  after telescoping, and this establishes (5-3). Combining (5-2) and (5-3), we have

$$d_{\Psi(\tau)}(\langle a \rangle) = d_{\widehat{\Phi}(\tau)}(\langle a \rangle) \tag{5-4}$$

for all  $m \in \mathbb{N}$  and  $a \in (A \otimes M_m)_+$ . As functionals on the Cuntz semigroup preserve suprema, (5-4) holds for all  $a \in (A \otimes \mathcal{K})_+$ , whence  $\Psi(\tau) = \widehat{\Phi}(\tau)$ . □

The homeomorphism between the lower semicontinuous quasitraces can be used to establish the weak\*-continuity of the map between the tracial state spaces of close unital  $C^*$ -algebras from [CSSW, Section 5] resolving a point left open there. In particular this shows that the map defined in [CSSW] provides an isomorphism between the Elliott invariants of completely close algebras.

For any closed two-sided ideal  $I \trianglelefteq A$ , the subcone  $T_I(A)$  of  $T(A)$  consists of those  $\tau \in T(A)$  such that the closed two-sided ideal generated by  $\{x \in A_+ : \tau(x) < \infty\}$  is  $I$ . Proposition 3.11 of [Elliott et al. 2011] shows that the relative topology on  $T_I(A)$  is the topology of pointwise convergence on the positive elements of the Pedersen ideal of  $I$ . In particular,  $T_s(A) \subset T_A(A)$ . In particular, the induced topology on  $T_s(A)$  is just the weak\*-topology.

**Corollary 5.2.** *Suppose that  $A$  and  $B$  are unital  $C^*$ -algebras acting nondegenerately on a Hilbert space with  $d_{cb}(A, B) < 1/42$  and  $d(A, B) < 1/2200$ . Then the affine isomorphism  $\Psi : T_s(B) \rightarrow T_s(A)$  between tracial state spaces in [Christensen et al. 2010, Section 5] is a homeomorphism with respect to the weak\*-topologies.*

We end with two further corollaries of Theorem 5.1.

**Corollary 5.3.** *Let  $A$  and  $B$  be unital  $C^*$ -algebras acting nondegenerately on the same Hilbert space with  $d_{cb}(A, B) < 1/2200$ . Suppose every bounded 2-quasitrace on  $A$  is a trace, then the same property holds for  $B$ .*

*Proof.* Given  $\tau \in QT_{2,s}(B)$ , its image  $\widehat{\Phi}(\tau)$  lies in  $QT_{2,s}(A) = T_s(A)$ . By Theorem 5.1 (2) (applied with  $A$  and  $B$  interchanged)

$$\tau = \widehat{\Phi}^{-1}(\widehat{\Phi}(\tau)) = \Psi^{-1}(\widehat{\Phi}(\tau)) \in T_s(B),$$

as claimed. □

The question of whether exactness transfers to (completely) close  $C^*$ -algebras raised in [Christensen et al. 2010] remains open, but we do at least obtain the following corollary.

**Corollary 5.4.** *Let  $A$  and  $B$  be unital  $C^*$ -algebras acting nondegenerately on the same Hilbert space with  $d_{cb}(A, B) < 1/2200$  and suppose  $A$  is exact. Then every bounded 2-quasitrace on  $B$  is a trace.*

*Proof.* This is immediate from Haagerup’s result [1991] that bounded 2-quasitraces on exact  $C^*$ -algebras are traces and the previous corollary. □

We end by noting that the isomorphism between the Cuntz semigroups of completely close algebras in Theorem 3.10 can also be used to directly recapture an isomorphism between the Elliott invariants in significant cases. Let  $\mathbf{Cu}_{\mathbb{T}}$  be the functor  $A \mapsto \mathbf{Cu}(A \otimes C(\mathbb{T}))$  mapping the category of  $C^*$ -algebras into the category  $\mathbf{Cu}$  introduced in [Coward et al. 2008] and let  $\mathbf{Ell}$  be the Elliott invariant functor taking values in the category  $\mathbf{Inv}$  whose objects are the 4-tuples arising from the Elliott invariant. Let  $\mathcal{C}$  be the subcategory of separable, unital, simple finite and  $\mathcal{L}$ -stable algebras  $A$  with  $QT_2(A) = T(A)$  (for example if  $A$  is exact). Then, building on work from [Brown et al. 2008; Brown and Toms 2007], Theorem 4.2 of [Antoine et al. 2014] provides functors  $F : \mathbf{Inv} \rightarrow \mathbf{Cu}$  and  $G : \mathbf{Cu} \rightarrow \mathbf{Inv}$  such that there are natural equivalences of functors  $F \circ \mathbf{Ell}|_{\mathcal{C}} \cong \mathbf{Cu}_{\mathbb{T}}|_{\mathcal{C}}$  and  $G \circ \mathbf{Cu}_{\mathbb{T}}|_{\mathcal{C}} \cong \mathbf{Ell}|_{\mathcal{C}}$  (a similar result for simple unital ASH algebras which are not type I and have slow dimension growth can be found in [Tikuisis 2011]). Note that

in Theorem 4.2 of [Antoine et al. 2014] there is an implicit nuclearity hypothesis, which is only actually used in order to see  $QT_2(A) = T(A)$ ; therefore the result holds in the generality stated. Thus if  $A$  and  $B$  are  $\mathcal{L}$ -stable  $C^*$ -algebras with  $d_{cb}(A, B)$  sufficiently small, and  $A$  is simple, separable, unital finite and has  $QT_2(A) = T(A)$ , then  $B$  enjoys all these properties. Further, since tensoring by an abelian algebra does not increase the complete distance between  $A$  and  $B$  (see [Christensen 1980, Theorem 3.2] for this result in the context of near inclusions—the same proof works for the metric  $d_{cb}$ ),  $\text{Cu}_{\mathbb{T}}(A) \cong \text{Cu}_{\mathbb{T}}(B)$  by Theorem 3.10. Thus  $\text{Ell}(A) \cong \text{Ell}(B)$ .

**Acknowledgements.** This work was initiated at the Centre de Recerca Matemàtica (Bellaterra) during the Programme “The Cuntz Semigroup and the Classification of  $C^*$ -algebras” in 2011. The authors would like to thank the CRM for the financial support for this programme and the conducive research environment.

## References

- [Antoine et al. 2011] R. Antoine, J. Bosa, and F. Perera, “Completions of monoids with applications to the Cuntz semigroup”, *Internat. J. Math.* **22**:6 (2011), 837–861. [MR 2012d:46146](#) [Zbl 1239.46042](#)
- [Antoine et al. 2014] R. Antoine, M. Dadarlat, F. Perera, and L. Santiago, “Recovering the Elliott invariant from the Cuntz semigroup”, *Trans. Amer. Math. Soc.* **366**:6 (2014), 2907–2922. [MR 3180735](#) [Zbl 06303118](#)
- [Ara and Pardo 1996] P. Ara and E. Pardo, “Refinement monoids with weak comparability and applications to regular rings and  $C^*$ -algebras”, *Proc. Amer. Math. Soc.* **124**:3 (1996), 715–720. [MR 96f:46124](#) [Zbl 0849.16009](#)
- [Ara et al. 2011] P. Ara, F. Perera, and A. S. Toms, “ $K$ -theory for operator algebras: Classification of  $C^*$ -algebras”, pp. 1–71 in *Aspects of operator algebras and applications*, edited by P. Ara et al., *Contemp. Math.* **534**, Amer. Math. Soc., Providence, RI, 2011. [MR 2012g:46092](#) [Zbl 1219.46053](#)
- [Blanchard and Kirchberg 2004] E. Blanchard and E. Kirchberg, “Non-simple purely infinite  $C^*$ -algebras: The Hausdorff case”, *J. Funct. Anal.* **207**:2 (2004), 461–513. [MR 2005b:46136](#) [Zbl 1048.46049](#)
- [Brown and Toms 2007] N. P. Brown and A. S. Toms, “Three applications of the Cuntz semigroup”, *Int. Math. Res. Not.* **2007**:19 (2007), Art. ID rnm068. [MR 2009a:46104](#) [Zbl 1134.46040](#)
- [Brown et al. 2008] N. P. Brown, F. Perera, and A. S. Toms, “The Cuntz semigroup, the Elliott conjecture, and dimension functions on  $C^*$ -algebras”, *J. Reine Angew. Math.* **621** (2008), 191–211. [MR 2010a:46125](#) [Zbl 1158.46040](#)
- [Cameron et al. 2012] J. Cameron, E. Christensen, A. M. Sinclair, R. R. Smith, S. A. White, and A. D. Wiggins, “Type  $\text{II}_1$  factors satisfying the spatial isomorphism conjecture”, *Proc. Natl. Acad. Sci. USA* **109**:50 (2012), 20338–20343. [MR 3023668](#)
- [Cameron et al. 2013] J. Cameron, E. Christensen, A. M. Sinclair, R. R. Smith, S. A. White, and A. D. Wiggins, “A remark on the similarity and perturbation problems”, *C. R. Math. Acad. Sci. Soc. R. Canada* **35**:2 (2013), 70–76. [MR 3114459](#) [Zbl 06308103](#)
- [Cameron et al.  $\geq$  2014] J. Cameron, E. Christensen, A. M. Sinclair, R. R. Smith, S. A. White, and A. D. Wiggins, “Kadison–Kastler stable factors”, To appear in *Duke Math. J.* [arXiv 1209.4116](#)
- [Christensen 1974/75] E. Christensen, “Perturbations of type I von Neumann algebras”, *J. London Math. Soc.* (2) **9** (1974/75), 395–405. [MR 50 #10839](#) [Zbl 0319.46052](#)
- [Christensen 1977] E. Christensen, “Perturbations of operator algebras, II”, *Indiana Univ. Math. J.* **26**:5 (1977), 891–904. [MR 58 #23628b](#) [Zbl 0395.46045](#)
- [Christensen 1980] E. Christensen, “Near inclusions of  $C^*$ -algebras”, *Acta Math.* **144**:3-4 (1980), 249–265. [MR 81h:46070](#) [Zbl 0469.46044](#)
- [Christensen 1982] E. Christensen, “Extensions of derivations, II”, *Math. Scand.* **50**:1 (1982), 111–122. [MR 83m:46092](#) [Zbl 0503.47032](#)
- [Christensen 2001] E. Christensen, “Finite von Neumann algebra factors with property  $\Gamma$ ”, *J. Funct. Anal.* **186**:2 (2001), 366–380. [MR 2003h:46090](#) [Zbl 1020.46017](#)

- [Christensen et al. 2010] E. Christensen, A. Sinclair, R. R. Smith, and S. White, “Perturbations of  $C^*$ -algebraic invariants”, *Geom. Funct. Anal.* **20**:2 (2010), 368–397. [MR 2012b:46099](#) [Zbl 1205.46036](#)
- [Christensen et al. 2012] E. Christensen, A. M. Sinclair, R. R. Smith, S. A. White, and W. Winter, “Perturbations of nuclear  $C^*$ -algebras”, *Acta Math.* **208**:1 (2012), 93–150. [MR 2910797](#) [Zbl 1252.46047](#)
- [Coward et al. 2008] K. T. Coward, G. A. Elliott, and C. Ivanescu, “The Cuntz semigroup as an invariant for  $C^*$ -algebras”, *J. Reine Angew. Math.* **623** (2008), 161–193. [MR 2010m:46101](#) [Zbl 1161.46029](#)
- [Elliott et al. 2011] G. A. Elliott, L. Robert, and L. Santiago, “The cone of lower semicontinuous traces on a  $C^*$ -algebra”, *Amer. J. Math.* **133**:4 (2011), 969–1005. [MR 2012f:46120](#) [Zbl 1236.46052](#)
- [Haagerup 1983] U. Haagerup, “Solution of the similarity problem for cyclic representations of  $C^*$ -algebras”, *Ann. of Math.* (2) **118**:2 (1983), 215–240. [MR 85d:46080](#) [Zbl 0543.46033](#)
- [Haagerup 1991] U. Haagerup, “Quasi-traces on exact  $C^*$ -algebras are traces”, preprint, 1991. updated version at [ArXiv 1403.7653](#).
- [Hadwin and Li 2014] D. Hadwin and W. Li, “The similarity degree of some  $C^*$ -algebras”, *Bull. Aust. Math. Soc.* **89**:1 (2014), 60–69. [MR 3163005](#) [Zbl 06276420](#)
- [Hirshberg et al. 2012] I. Hirshberg, E. Kirchberg, and S. White, “Decomposable approximations of nuclear  $C^*$ -algebras”, *Adv. Math.* **230**:3 (2012), 1029–1039. [MR 2921170](#) [Zbl 1256.46019](#)
- [Hjelmborg and Rørdam 1998] J. v. B. Hjelmborg and M. Rørdam, “On stability of  $C^*$ -algebras”, *J. Funct. Anal.* **155**:1 (1998), 153–170. [MR 99g:46079](#) [Zbl 0912.46055](#)
- [Johanesová and Winter 2012] M. Johanesová and W. Winter, “The similarity problem for  $\mathcal{L}$ -stable  $C^*$ -algebras”, *Bull. Lond. Math. Soc.* **44**:6 (2012), 1215–1220. [MR 3007654](#) [Zbl 1264.46042](#)
- [Kadison 1955] R. V. Kadison, “On the orthogonalization of operator representations”, *Amer. J. Math.* **77** (1955), 600–620. [MR 17,285c](#) [Zbl 0064.36605](#)
- [Kadison and Kastler 1972] R. V. Kadison and D. Kastler, “Perturbations of von Neumann algebras, I: Stability of type”, *Amer. J. Math.* **94** (1972), 38–54. [MR 45 #5772](#) [Zbl 0239.46070](#)
- [Khoshkam 1984] M. Khoshkam, “Perturbations of  $C^*$ -algebras and  $K$ -theory”, *J. Operator Theory* **12**:1 (1984), 89–99. [MR 86c:46084](#) [Zbl 0571.46039](#)
- [Kirchberg 1996] E. Kirchberg, “The derivation problem and the similarity problem are equivalent”, *J. Operator Theory* **36**:1 (1996), 59–62. [MR 97f:46108](#) [Zbl 0865.46055](#)
- [Li and Shen 2008] W. Li and J. Shen, “A note on approximately divisible  $C^*$ -algebras”, preprint, 2008. [arXiv 0804.0465](#)
- [Matui and Sato 2012] H. Matui and Y. Sato, “Strict comparison and  $\mathcal{L}$ -absorption of nuclear  $C^*$ -algebras”, *Acta Math.* **209**:1 (2012), 179–196. [MR 2979512](#) [Zbl 1277.46028](#)
- [Ortega et al. 2012] E. Ortega, F. Perera, and M. Rørdam, “The corona factorization property, stability, and the Cuntz semigroup of a  $C^*$ -algebra”, *Int. Math. Res. Not.* **2012**:1 (2012), 34–66. [MR 2012m:46068](#) [Zbl 1245.46049](#)
- [Perera 1997] F. Perera, “The structure of positive elements for  $C^*$ -algebras with real rank zero”, *Internat. J. Math.* **8**:3 (1997), 383–405. [MR 98i:46058](#) [Zbl 0881.46044](#)
- [Phillips 1973/74] J. Phillips, “Perturbations of  $C^*$ -algebras”, *Indiana Univ. Math. J.* **23** (1973/74), 1167–1176. [MR 49 #5861](#) [Zbl 0264.46059](#)
- [Phillips and Raeburn 1979] J. Phillips and I. Raeburn, “Perturbations of AF-algebras”, *Canad. J. Math.* **31**:5 (1979), 1012–1016. [MR 81e:46042a](#) [Zbl 0373.46074](#)
- [Pisier 1998] G. Pisier, “The similarity degree of an operator algebra”, *Algebra i Analiz* **10**:1 (1998), 132–186. Translated in *St. Petersburg Math. J.* **10**:1 (1999), 103–146. [MR 99c:46066](#) [Zbl 0911.47038](#)
- [Pop 2004] F. Pop, “The similarity problem for tensor products of certain  $C^*$ -algebras”, *Bull. Austral. Math. Soc.* **70**:3 (2004), 385–389. [MR 2005g:46111](#) [Zbl 1066.46047](#)
- [Rieffel 1983] M. A. Rieffel, “Dimension and stable rank in the  $K$ -theory of  $C^*$ -algebras”, *Proc. London Math. Soc.* (3) **46**:2 (1983), 301–333. [MR 84g:46085](#) [Zbl 0533.46046](#)
- [Robert 2012] L. Robert, “Classification of inductive limits of 1-dimensional NCCW complexes”, *Adv. Math.* **231**:5 (2012), 2802–2836. [MR 2970466](#) [Zbl 1268.46041](#)

- [Robert and Santiago 2010] L. Robert and L. Santiago, “Classification of  $C^*$ -homomorphisms from  $C_0(0, 1]$  to a  $C^*$ -algebra”, *J. Funct. Anal.* **258**:3 (2010), 869–892. [MR 2010j:46108](#) [Zbl 1192.46057](#)
- [Rørdam and Winter 2010] M. Rørdam and W. Winter, “The Jiang–Su algebra revisited”, *J. Reine Angew. Math.* **642** (2010), 129–155. [MR 2011i:46074](#) [Zbl 1209.46031](#)
- [Tikuisis 2011] A. Tikuisis, “The Cuntz semigroup of continuous functions into certain simple  $C^*$ -algebras”, *Internat. J. Math.* **22**:8 (2011), 1051–1087. [MR 2012h:46102](#) [Zbl 1232.46058](#)
- [Toms 2008] A. S. Toms, “On the classification problem for nuclear  $C^*$ -algebras”, *Ann. of Math. (2)* **167**:3 (2008), 1029–1044. [MR 2009g:46119](#) [Zbl 1181.46047](#)
- [Winter 2010] W. Winter, “Decomposition rank and  $\mathcal{K}$ -stability”, *Invent. Math.* **179**:2 (2010), 229–301. [MR 2011a:46092](#) [Zbl 1194.46104](#)
- [Winter 2012] W. Winter, “Nuclear dimension and  $\mathcal{K}$ -stability of pure  $C^*$ -algebras”, *Invent. Math.* **187**:2 (2012), 259–342. [MR 2885621](#) [Zbl 1280.46041](#)
- [Zhang 1990] S. Zhang, “Diagonalizing projections in multiplier algebras and in matrices over a  $C^*$ -algebra”, *Pacific J. Math.* **145**:1 (1990), 181–200. [MR 92h:46088](#) [Zbl 0673.46049](#)

Received 4 Mar 2013. Revised 13 Jun 2013. Accepted 23 Jul 2013.

FRANCESC PERERA: [perera@mat.uab.cat](mailto:perera@mat.uab.cat)

*Department of Mathematics, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain*

ANDREW TOMS: [atoms@purdue.edu](mailto:atoms@purdue.edu)

*Department of Mathematics, Purdue University, Room 720, Mathematical Sciences Building, 150 N. University St., West Lafayette, IN 47907-2067, United States*

STUART WHITE: [stuart.white@glasgow.ac.uk](mailto:stuart.white@glasgow.ac.uk)

*School of Mathematics and Statistics, University of Glasgow, University Gardens, Glasgow, Q12 8QW, Scotland*

WILHELM WINTER: [wwinter@uni-muenster.de](mailto:wwinter@uni-muenster.de)

*Mathematisches Institut der WWU Münster, Einsteinstraße 62, 48149, Münster, Germany*

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>	Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Vaughan Jones	U.C. Berkeley & Vanderbilt University <a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a>	Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>
Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 7 No. 4 2014

---

- The global stability of the Minkowski spacetime solution to the Einstein-nonlinear system in wave coordinates 771  
JARED SPECK
- Dispersion for the Schrödinger equation on the line with multiple Dirac delta potentials and on delta trees 903  
VALERIA BANICA and LIVIU I. IGNAT
- The Cuntz semigroup and stability of close  $C^*$ -algebras 929  
FRANCESC PERERA, ANDREW TOMS, STUART WHITE and WILHELM WINTER
- Wave and Klein–Gordon equations on hyperbolic spaces 953  
JEAN-PHILIPPE ANKER and VITTORIA PIERFELICE
- Probabilistic global well-posedness for the supercritical nonlinear harmonic oscillator 997  
AURÉLIEN POIRET, DIDIER ROBERT and LAURENT THOMANN