

# ANALYSIS & PDE

Volume 7

No. 8

2014

AAPPO KAURANEN AND PEKKA KOSKELA

**BOUNDARY BLOW-UP UNDER SOBOLEV MAPPINGS**





## BOUNDARY BLOW-UP UNDER SOBOLEV MAPPINGS

AAPO KAURANEN AND PEKKA KOSKELA

We prove that for mappings in  $W^{1,n}(\mathbb{B}^n, \mathbb{R}^m)$ , continuous up to the boundary and with modulus of continuity satisfying a certain divergence condition, the image of the boundary of the unit ball has zero  $n$ -Hausdorff measure. For Hölder continuous mappings we also prove an essentially sharp generalised Hausdorff dimension estimate.

### 1. Introduction

Throughout this paper  $\mathbb{B}^n$  denotes the unit ball in  $\mathbb{R}^n$  and  $W^{1,n}(\mathbb{B}^n, \mathbb{R}^m)$  is the Sobolev space of  $L^n(\mathbb{B}^n, \mathbb{R}^m)$ -functions  $f : \mathbb{B}^n \rightarrow \mathbb{R}^m$  with weak first-order derivatives in  $L^n(\mathbb{B}^n)$ .

If  $f : \mathbb{B}^2 \rightarrow \Omega \subset \mathbb{R}^2$  is a conformal mapping, then the boundary of  $\Omega$  can have positive Lebesgue measure even if  $f$  extends continuously up to the boundary of the disk. If one requires more, for example uniform Hölder continuity, then  $\partial\Omega$  is necessarily of Lebesgue measure zero. In fact, Jones and Makarov proved [1995, Theorem C.1] that  $\partial\Omega$  has measure zero if  $f$  satisfies  $|f(z) - f(w)| \leq \psi(|z - w|)$  in  $\mathbb{B}^2$  for  $\psi : [0, \infty) \rightarrow [0, \infty)$  with

$$\int_0 \left| \frac{\log \psi(t)}{\log t} \right|^2 \frac{dt}{t} = \infty. \tag{1}$$

This condition is very sharp: if the integral in (1) converges then [Jones and Makarov 1995, Section 6] provides us with a simply connected domain  $\Omega$  and a conformal mapping  $f : \mathbb{B}^2 \rightarrow \Omega$  such that the boundary of  $\Omega$  has positive Lebesgue measure and  $f$  has the modulus of continuity  $\psi$ .

Our first result gives a surprisingly general extension of the conformal setting; notice that each uniformly continuous conformal mapping  $f : \mathbb{B}^2 \rightarrow \Omega$  belongs to  $W^{1,2}(\mathbb{B}^2, \mathbb{R}^2)$ .

**Theorem 1.1.** *Let  $f \in W^{1,n}(\mathbb{B}^n, \mathbb{R}^m)$  be a continuous mapping that satisfies*

$$|f(z) - f(w)| \leq \psi(|z - w|) \tag{2}$$

*for all  $z, w \in \overline{\mathbb{B}^n}$ , where  $\psi : (0, \infty) \rightarrow (0, \infty)$  is an allowable modulus of continuity with*

$$\int_0 \left| \frac{\log \psi(t)}{\log t} \right|^n \frac{dt}{t} = \infty. \tag{3}$$

*Then  $\mathcal{H}^n(f(\partial\mathbb{B}^n)) = 0$ .*

---

The authors were partially supported by the Academy of Finland grants 131477 and 263850.  
 MSC2010: 26B10, 26B35, 46E35.

*Keywords:* Sobolev mapping, Hausdorff measure, modulus of continuity.

Recall that every uniformly continuous map defined on  $\mathbb{B}^n$  has a continuous extension to all of  $\overline{\mathbb{B}^n}$ . In the above,  $f$  on  $\partial\mathbb{B}^n$  refers to this extension,  $\mathcal{H}^n(A)$  denotes the  $n$ -dimensional Hausdorff measure of a set  $A$ , and the definition of an *allowable modulus of continuity* is given in Definition 2.2 of Section 2. For example, both  $\psi(t) = Ct^\gamma$ ,  $0 < \gamma \leq 1$ , and

$$\psi_{l,s}(t) = \exp\left(-C \frac{(\log(C_l/t))^{(n-1)/n}}{(\log^{(l)}(C_l/t))^{s/n} \left(\prod_{k=2}^{l-1} \log^{(k)}(C_l/t)\right)^{1/n}}\right)$$

are allowable, where  $l \geq 2$  is an integer and  $s > 0$ . Notice that  $\psi_{l,s}$  satisfies (3) if and only if  $s \leq 1$ . Here  $C > 0$ ,  $\log^{(k)} t$  is the  $k$ -times iterated logarithm and  $C_l$  can be any constant with  $\log^{(l)}(C_l/2) \geq 1$ .

Let us look at the special case  $n = m = 2$  of Theorem 1.1 in the Hölder continuous setting:  $\psi(t) = Ct^\gamma$ , where  $0 < \gamma \leq 1$ . Consider a space-filling (Peano) curve, i.e., a continuous mapping  $g$  from the unit circle onto a square. In one of the standard constructions,  $g$  is Hölder continuous with exponent  $\gamma = \frac{1}{2}$ ; see, for example, [Buckley 1996, Theorem 3]. If one takes, say, the Poisson extension  $f$  of  $g$  to the unit disk, then  $f$  is also Hölder continuous. It is easy to check by hand that the partial derivatives of  $f$  do not belong to  $L^2(\mathbb{B}^2)$ . By Theorem 1.1, no Hölder continuous (or even continuous with control function satisfying (3)) extension  $f$  of a space filling curve can satisfy  $|Df| \in L^2(\mathbb{B}^2)$ .

In the Hölder continuous case, Jones and Makarov actually proved that the Hausdorff dimension of  $f(\partial\mathbb{B}^2)$  is strictly less than two for conformal  $f$ . Contrary to the area zero results, this dimension estimate is truly conformal in the following sense:

**Example 1.2.** Let  $p > 1$ . There exists a locally Hölder continuous homeomorphism  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  with  $f \in W_{\text{loc}}^{1,2}(\mathbb{B}^2, \mathbb{R}^2)$ , which maps  $\partial\mathbb{B}^2$  onto a set of positive  $\mathcal{H}^g$ -measure for the gauge function  $g(t) = t^2(\log(1/t))^p$ .

This construction can be found in Section 4. Here  $\mathcal{H}^g$  denotes the generalised Hausdorff measure with the function  $g$  as the dimension gauge. The precise definitions are given in Section 2.

Our second result gives a rather optimal positive result.

**Theorem 1.3.** Fix  $\gamma \in (0, 1]$ ,  $C > 0$ , and let  $g(t) = t^n \log(1/t)$ . Suppose that  $f \in W^{1,n}(\mathbb{B}^n, \mathbb{R}^m)$  satisfies

$$|f(z) - f(w)| \leq C|z - w|^\gamma$$

for all  $z, w \in \mathbb{B}^n$ . Then  $\mathcal{H}^g(f(\partial\mathbb{B}^n)) = 0$ .

Jones and Makarov proved their result via harmonic measure and hence this technique does not work in the setting of Theorem 1.1. An alternate approach, relying on the conformal (quasi)invariance of the (quasi)hyperbolic metric, was given in [Koskela and Rohde 1997]; see also [Nieminen 2006]. Furthermore, Malý and Martio [1995] established Theorem 1.1 in the Hölder continuous case via a technique that we have not been able to push further.

Let us briefly describe the idea of the proof of Theorem 1.1. We consider a Whitney decomposition  $\mathcal{W}$  of  $\mathbb{B}^n$  and assign to each  $Q \in \mathcal{W}$  a vector  $f_Q \in \mathbb{R}^m$  and a radius  $r_Q$ . The vector  $f_Q$  will simply be the “average” of  $f$  over  $Q$  and  $r_Q$  the maximum of  $|f_Q - f_{\tilde{Q}}|$  over all neighbours  $\tilde{Q}$  of  $Q$ . Then the  $n$ -integrability of the weak derivatives of  $f$  guarantees, via the Poincaré inequality, that the sequence

$\{r_Q\}_{Q \in \mathcal{W}}$  belongs to  $l^n$ . We realise  $f(\partial \mathcal{B}^n)$  as (a part of) the closure of  $\{f_Q\}_{Q \in \mathcal{W}}$  in  $\mathbb{R}^m$ . Those  $f(\omega)$ ,  $\omega \in \partial \mathcal{B}^n$ , for which one can find a sequence of  $Q \in \mathcal{W}$  with  $|f_Q - f(\omega)| \lesssim r_Q$  are easily handled. For the remaining  $\omega \in \partial \mathcal{B}^n$  we modify our centres  $f_Q$  and radii  $r_Q$ , while still retaining the  $l^n$ -condition, so that suitably blown-up balls cover these points sufficiently many times. This is where the nonintegrability condition (3) kicks in. One cannot fully follow the above idea, and so our proof, given below in Section 3, is more complicated.

Our approach is flexible and applies to many related problems. In order to avoid extra technicalities we do not record such applications here. Let us simply mention that the dimension gap phenomenon from [Hencl et al. 2012] can be shown to extend from conformal mappings to general Sobolev mappings [Koskela and Zapadinskaya 2014].

### 2. Preliminaries

Let us first agree on some basic notation. Given a number  $a > 0$ , we write  $\lfloor a \rfloor$  for the largest integer less than or equal to  $a$ . Similarly,  $\lceil a \rceil$  is the smallest integer greater than or equal to  $a$ . If  $A$  is a finite set,  $\#A$  is the number of elements in  $A$ . If  $A \subset \mathbb{R}^n$  has finite and strictly positive Lebesgue measure and  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is a Lebesgue integrable function, we denote the average  $(1/|A|) \int_A f$  of  $f$  over the set  $A$  by  $f_A$  or  $f_A$ , where  $|A|$  is the  $n$ -dimensional Lebesgue measure of the set  $A$ . For  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $f_A$  is then defined via the component functions of  $f$ . Given a point  $x \in \mathbb{R}^n$  and a nonnegative number  $r$ ,  $B(x, r)$  denotes the open ball with centre  $x$  and radius  $r$  and  $Q(x, r)$  denotes the cube  $\{y \in \mathbb{R}^n : \max\{|x_i - y_i|\}_{i=1,2,\dots,n} \leq r\}$ . If  $B = B(x, r)$  is a ball and  $a$  is a positive number, the notation  $aB$  stands for the ball  $B(x, ar)$ . We denote the radius of a ball  $B$  by  $r(B)$ . When we write  $L = L(\cdot)$ , we mean that the positive constant  $L$  depends only on the parameters listed inside the parentheses. Finally,  $C$  denotes a positive constant, which may depend only on  $n$  and  $m$ , the dimensions of the domain space and the image space, and may differ from occurrence to occurrence.

We write  $\mathcal{H}^h(A)$  for the generalised Hausdorff measure of a set  $A \subset \mathbb{R}^n$ , given by

$$\mathcal{H}^h(A) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^h(A), \quad \text{where } \mathcal{H}_\delta^h(A) = \inf \left\{ \sum_{i=1}^\infty h(\text{diam } U_i) : A \subset \bigcup_{i=1}^\infty U_i, \text{diam } U_i \leq \delta \right\}$$

and  $h$  is a dimension gauge (a nondecreasing function with  $\lim_{t \rightarrow 0^+} h(t) = h(0) = 0$  and with  $h(t) > 0$  for all  $t > 0$ ). If  $h(t) = t^a$  for some  $a \geq 0$  we simply write  $\mathcal{H}^a$  for  $\mathcal{H}^h$  and call it the  $a$ -dimensional Hausdorff measure.

A sequence of pairs  $(c_i, U_i)_{i=1}^\infty$ , where  $c_i \geq 0$  and  $U_i \subset \mathbb{R}^n$ , that satisfies  $\chi_A(x) \leq \sum_{i=1}^\infty c_i \chi_{U_i}(x)$  for all  $x \in \mathbb{R}^n$  is called a weighted cover of the set  $A$ . We also need a generalised weighted Hausdorff content of a set  $A \subset \mathbb{R}^n$ , given by

$$\lambda_\infty^h(A) = \inf \left\{ \sum_{i=1}^\infty c_i h(\text{diam } U_i) : (c_i, U_i)_{i=1}^\infty \text{ is a weighted cover of } A \right\}.$$

Here also  $h$  is a gauge function. Again we write  $\lambda_\infty^h = \lambda_\infty^a$  if  $h(t) = t^a$ .

**Lemma 2.1.** *Let  $E \subset \mathbb{R}^n$  be bounded. Let  $h$  be a continuous gauge function with  $h(2t) \leq ch(t)$  for some  $c > 0$ . Then  $\mathcal{H}_\infty^h(E) \leq c\lambda_\infty^h(E)$ .*

*Proof.* The lemma follows from Corollary 8.2 and the proof of Theorem 9.7 of [Howroyd 1994]; see also [Federer 1969, 2.10.24]. □

Recall that for each open subset  $U$  of  $\mathbb{R}^n$  there exists a Whitney decomposition  ${}^{\circ}\mathcal{W}$  given by  $U = \bigcup_{i=1}^\infty Q_i$ , where  $Q_i \in {}^{\circ}\mathcal{W}$  are cubes with mutually parallel sides, pairwise disjoint interiors and each of edge-length  $2^k$  for some integer  $k$ , such that the relation

$$\frac{1}{4} \leq \frac{\text{diam } Q_i}{\text{dist}(Q_i, \partial\Omega)} \leq 1 \tag{4}$$

holds for all  $i = 1, 2, \dots$ . We write  $Q_1 \smile Q_2$  if the Whitney cubes  $Q_1 \neq Q_2$  share at least one point (the so-called neighbour cubes). We have

$$\frac{1}{4} \leq \frac{\text{diam } Q}{\text{diam } \tilde{Q}} \leq 4$$

whenever  $Q \smile \tilde{Q}$ . Therefore, the total number  $\#\{\tilde{Q} : \tilde{Q} \smile Q\}$  of all neighbours of a fixed cube  $Q$  does not exceed  $C$ . See [Stein 1970] for details.

Let  $\omega \in \partial\mathcal{B}^n$ . By  $(Q_j(\omega))_{j=1}^\infty$  we mean the sequence of all Whitney cubes in a fixed Whitney decomposition of  $\mathcal{B}^n$  intersecting the radius  $[0, \omega]$ . This sequence starts with a central cube and tends to  $\omega$ . For a point  $x \in [0, \omega]$ , we denote the number of Whitney cubes intersecting the segment  $[0, x]$  by  $\#q(0, x)$ . It is easy to see that

$$c_1 \leq \frac{\#q(0, x)}{\log(1/(1 - |x|))} \leq c_2 \tag{5}$$

whenever  $\#q(0, x) > c_3$ , where  $c_i > 0, i = 1, 2, 3$  are constants that may depend on  $n$ .

Finally, we define the allowable moduli of continuity:

**Definition 2.2.** A continuously differentiable increasing bijection  $\psi : (0, \infty) \rightarrow (0, \infty)$  is an *allowable modulus of continuity* if there exists  $t_0 < 1$  and  $\beta > 0$  such that for every  $t \leq t_0$  the following conditions hold:

$$\log \frac{1}{\psi^{-1}(t)} \text{ is differentiable and } \frac{(\psi^{-1})'(t)}{\psi^{-1}(t)} t \text{ is a decreasing function;} \tag{6}$$

$$\log \frac{1}{\psi^{-1}(t)} \leq \beta \log \frac{1}{\psi^{-1}(\sqrt{t})}; \tag{7}$$

$$\frac{(\log \psi(t))' t \log t}{\log \psi(t)} \text{ is a monotone function.} \tag{8}$$

**Remark 2.3.** (i) One could replace the monotonicity conditions in (6) and (8) with a *pseudomonotonicity* condition (e.g., there exists a constant  $C > 0$  such that  $u(t) \leq Cu(s)$  if  $t \leq s$ ). This would only affect the constants in the proofs.

(ii) The conditions (6) and (7) mean that the function  $\log(1/\psi^{-1}(t))$  is a function of logarithmic type in the sense of [Nieminen 2006, Definition 4.2].

### 3. Proofs

*Proof of Theorem 1.1.* We may assume that  $m, n \geq 2$ . Let  $f \in W^{1,n}(\mathbb{B}^n, \mathbb{R}^m)$  and  $\psi$  be as in the statement of Theorem 1.1. Denote  $\psi^{-1}(t)$  by  $u(t)$ . It follows from our assumptions (3), (6), (7), (8) and [Nieminen 2006, Remark 5.3.] that

$$\int_0^1 \left( \frac{u(t)}{u'(t)} \right)^{n-1} \frac{dt}{t^n} = \infty. \tag{9}$$

We define  $\alpha(t) = u(t)/u'(t)$  and  $\lambda(k) = 2^{-k}/\alpha(2^{-k})$  for  $k \in \mathbb{N}$ . By (6),  $\lambda$  is increasing for large  $k$ . For simplicity we assume  $\lambda$  to be increasing.

Let  $\mathcal{W}$  be a fixed Whitney decomposition of  $\mathbb{B}^n$ . For each cube  $Q \in \mathcal{W}$  we define a corresponding centre  $f_Q$  and a corresponding radius  $r_Q = \max\{|f_Q - f_{\tilde{Q}}| : Q \sim \tilde{Q}\}$ , which determine a family of balls on the image side indexed by  $\mathcal{W}$ :

$$\mathcal{B} = \{(Q, B(f_Q, r_Q)) : Q \in \mathcal{W}, r_Q > 0\}.$$

To simplify our notation we abbreviate  $(Q, B(f_Q, r_Q))$  to  $B(f_Q, r_Q)$  in what follows.

We assign two new weighted collections of balls to each element in  $\mathcal{B}$ . Given  $B = B(x, r) \in \mathcal{B}$ , we define concentric subballs  $S_i(B) = B(x, r/2^i)$  for all  $i \in \mathbb{N}$  and assign the weight  $w_{S_i(B)} = 2^i$  to each  $S_i(B)$ . We set  $\mathcal{S}_B = \{S_i(B) : i \in \mathbb{N}\}$ . Then

$$\sum_{B' \in \mathcal{S}_B} w_{B'} r(B')^n = \sum_{i=1}^{\infty} w_{S_i(B)} r(S_i(B))^n = \sum_{i=1}^{\infty} 2^i \frac{r(B)^n}{2^{ni}} \leq r(B)^n.$$

The second collection is defined in a similar way. If  $B = B(x, r)$  is a ball in  $\mathcal{B}$ , we choose the smallest number  $k_0(r) \in \mathbb{N}$  such that  $2^{-k_0(r)} \leq r$ . Next, for each  $k = k_0(r), k_0(r) + 1, \dots$ , we choose  $R_k(B) = B(x, \alpha(2^{-k}))$  and set  $\mathcal{R}_B = \{R_k(B) : k = k_0(r), k_0(r) + 1, \dots\}$ . The weights we assign this time are  $w_{R_k(B)} = \lambda(k)$  for all  $k = k_0(r), k_0(r) + 1, \dots$ . Similarly to the above,

$$\begin{aligned} \sum_{B' \in \mathcal{R}_B} w_{B'} r(B')^n &= \sum_{k=k_0(r)}^{\infty} w_{R_k(B)} r(R_k(B))^n = \sum_{k=k_0(r)}^{\infty} (\alpha(2^{-k}))^n \lambda(k) \\ &\leq \sum_{k=k_0(r)}^{\infty} (\alpha(2^{-k}))^n \frac{\lambda(k)^n}{\lambda(0)^{n-1}} = \frac{1}{\lambda(0)^{n-1}} \sum_{k=k_0(r)}^{\infty} 2^{-nk} \leq \frac{2 \cdot 2^{-nk_0(r)}}{\lambda(0)^{n-1}} \leq \frac{2}{\lambda(0)^{n-1}} r(B)^n. \end{aligned}$$

Finally, we define our weighted collection of balls by setting  $\mathcal{F} = \bigcup_{B \in \mathcal{B}} (\mathcal{S}_B \cup \mathcal{R}_B)$ .

Let us now estimate the weighted sums of the  $n$ -th powers of the radii of the balls in  $\mathcal{F}$ . Let  $N(Q) = Q \cup \bigcup_{\tilde{Q} \sim Q} \tilde{Q}$  be the union of  $Q \in \mathcal{W}$  and all neighbours  $\tilde{Q}$  of  $Q$ . For neighbouring cubes  $Q$  and  $\tilde{Q}$ , we obtain, via the Hölder and Poincaré inequalities, that

$$\begin{aligned} |f_Q - f_{\tilde{Q}}| &\leq \int_Q |f - f_{N(Q)}| + \int_{\tilde{Q}} |f - f_{N(Q)}| \leq C \int_{N(Q)} |f - f_{N(Q)}| \leq C \left( \int_{N(Q)} |f - f_{N(Q)}|^n \right)^{1/n} \\ &\leq C \left( \int_{N(Q)} |Df|^n \right)^{1/n}. \end{aligned}$$

Hence, we have the estimate

$$r_Q^n = \max\{|f_Q - f_{\tilde{Q}}|^n : Q \smile \tilde{Q}\} \leq C \int_{N(Q)} |Df|^n$$

for each  $Q \in \mathcal{W}$  and some constant  $C > 0$ . Next, using the fact that the inequality  $\sum_{Q \in \mathcal{W}} \chi_{N(Q)}(y) \leq C$  holds for every  $y \in \mathbb{R}^n$ , we estimate

$$\begin{aligned} \sum_{B \in \mathcal{F}} w_{Br}(B)^n &\leq C(\lambda(0)) \sum_{B \in \mathcal{B}} r(B)^n = C(\lambda(0)) \sum_{Q \in \mathcal{W}} r_Q^n \leq C(\lambda(0)) \sum_{Q \in \mathcal{W}} \int_{N(Q)} |Df|^n \\ &\leq C_1 \int_{\bigcup_{Q \in \mathcal{W}} N(Q)} |Df|^n \leq C_1 \int_{\mathbb{B}^n} |Df|^n < \infty, \end{aligned} \tag{10}$$

where  $C_1 > 0$  is some constant depending on  $n, m$  and  $\lambda(0)$  only.

We may assume that there is at least one  $Q \in \mathcal{W}$  with  $r_Q > 0$ ; otherwise  $f(\partial\mathbb{B}^n)$  is a singleton. Let  $\omega \in \partial\mathbb{B}^n$ . We consider the radius  $[0, \omega]$  and the sequence  $(Q_j(\omega))_{j=1}^\infty$ . We fix a large integer  $l_0 = l_0(\omega, f) \in \mathbb{N}$  so that there are elements of the sequence  $(f_{Q_j(\omega)})_{j=1}^\infty$  outside  $B(f(\omega), 2^{-l_0+1})$  if  $(f_{Q_j(\omega)})_{j=1}^\infty$  contains at least one element different from  $f(\omega)$ . If such an integer does not exist there necessarily is some  $Q = Q_\omega \in \mathcal{W}$  with  $f_Q = f(\omega)$  and  $r_Q > 0$ . In this case, we choose  $l_0 = l_0(\omega, f) \in \mathbb{N}$  so that  $2^{-l_0} < r_{Q_\omega}$ . In both cases we also require that  $2^{-l_0+1} < t_0$ . This allows us to use the properties (6) and (7).

For the purposes of our ‘‘porosity argument’’, we would like to make the number  $l_0$  independent of the point  $\omega$ . This is done by considering the decomposition

$$\partial\mathbb{B}^n = \bigcup_{l \in \mathbb{N}} E_l, \quad \text{where } E_l = \{\omega \in \partial\mathbb{B}^n : l_0(\omega, f) \leq l\}.$$

Setting  $F_l = f(E_l)$ , we then have  $f(\partial\mathbb{B}^n) = \bigcup_{l \in \mathbb{N}} F_l$ .

Let us fix  $l_0 \in \mathbb{N}$ . Our aim is to prove that  $\mathcal{H}_\infty^n(F_{l_0}) = 0$ .

Fix  $x \in F_{l_0}$ . Take any  $\omega \in E_{l_0}$  such that  $x = f(\omega)$  and define the sequence of concentric annuli  $A_l(x) = B(x, 2^{-l+1}) \setminus B(x, 2^{-l})$  with  $l = l_0, l_0 + 1, \dots$ . Next, we assign a suitable set  $P_l(x)$  of cubes from  $\mathcal{W}$  to each annulus  $A_l(x)$ ,  $l = l_0, l_0 + 1, \dots$ . If  $f_{Q_j(\omega)} = x$  for all  $j \in \mathbb{N}$ , we put  $P_l(x) = \{Q_\omega\}$  for each  $l \geq l_0$ , where  $Q_\omega$  is the cube defined earlier. Otherwise, all the sets  $P_l(x)$  with  $l \geq l_0$  consist of elements from  $(Q_j(\omega))_{j=1}^\infty$ . If an annulus  $A_l(x)$  with some  $l \geq l_0$  contains no centres from  $(f_{Q_j(\omega)})_{j=1}^\infty$  we define  $P_l(x) = \{Q_m(\omega)\}$ , where an integer  $m \in \mathbb{N}$  is chosen so that  $f_{Q_{m-1}(\omega)} \notin B(x, 2^{-l+1})$  but  $f_{Q_m(\omega)} \in B(x, 2^{-l})$ ; if, in contrast, there is at least one centre  $f_{Q_j(\omega)}$  in  $A_l(x)$  we take  $P_l(x) = \{Q_k(\omega) : k = m_1, \dots, m_2\}$ , where  $m_1, m_2 \in \mathbb{N}$  are such that  $f_{Q_{m_1-1}(\omega)} \notin B(x, 2^{-l+1})$ ,  $f_{Q_{m_2+1}(\omega)} \in B(x, 2^{-l})$  and  $f_{Q_k(\omega)} \in A_l(x)$  for all  $k = m_1, \dots, m_2$ . Moreover, it is possible to choose the sets  $P_l(x)$  above so that the inequality  $k_1 \leq k_2$  is valid whenever  $Q_{k_1}(\omega) \in P_{l_1}(x)$ ,  $Q_{k_2}(\omega) \in P_{l_2}(x)$  and  $l_1 < l_2$ .

Denoting

$$\theta_l(x) = \begin{cases} 1 & \text{if } \sharp P_l(x) \leq \tilde{c}_0 \lambda(l), \\ 0 & \text{otherwise} \end{cases}$$

for  $l \geq l_0$  and a constant  $\tilde{c}_0 > \lambda^{-1}(0)$ , which we will specify later, we would like to prove that there exists an integer  $l_1 \geq 2l_0$  such that

$$\sum_{k=l_0}^l \theta_k(x) \geq \frac{l}{2} \tag{11}$$

for each  $l \geq l_1$ . In other words, at least half of the annuli do not contain too many centres from  $(f_{Q_j(\omega)})_{j=1}^\infty$ . There is nothing to prove if  $f_{Q_j(\omega)} = x$  for all  $j \in \mathbb{N}$ ; otherwise, the proof is by contradiction:

Let us assume that (11) does not hold for some  $l \geq 2l_0$ . Take the smallest number  $J \in \mathbb{N}$  such that  $f_{Q_j(\omega)} \in B(x, 2^{-l})$  for all  $j > J$  and let  $\omega' \in [0, \omega]$  be the point of  $Q_J(\omega) \cap [0, \omega]$  which is closest to  $\omega$ . Now, the assumption on the continuity of  $f$  and the properties of our Whitney decomposition imply

$$2^{-l} \leq |f_{Q_J(\omega)} - x| = |f_{Q_J(\omega)} - f(\omega)| \leq \int_{Q_J} |f(y) - f(\omega)| dy \leq \psi(2(1 - |\omega'|)).$$

That is,

$$\frac{u(2^{-l})}{2} \leq 1 - |\omega'|.$$

Next, we connect this estimate to the number of Whitney cubes that precede  $Q_J$  in  $(Q_j(\omega))_{j=1}^\infty$ .

Using (5), we observe that

$$\log \frac{2}{u(2^{-l})} \geq \log \frac{1}{1 - |\omega'|} \geq \frac{1}{c_2} \#q(0, \omega').$$

In the calculation above we may have to adjust the choice of  $l_0$  to ensure  $\#q(0, \omega') > c_3$  (see (5)). Finally, we obtain a lower bound for  $\#q(0, \omega')$  using the assumption that we have at least  $\lfloor l/2 \rfloor - l_0 + 2$  annuli  $A_k(x)$  with  $\theta_k(x) = 0$ . We notice that the sets  $P_k(x)$  with  $\theta_k(x) = 0$  contain different cubes for different  $k$ , and if  $k \leq l$  then the cubes in  $P_k(x)$  precede  $Q_J(\omega)$  in  $(Q_j(\omega))_{j=1}^\infty$ . We have

$$\begin{aligned} c_2 \log \frac{2}{u(2^{-l})} &\geq \#q(0, \omega') \geq \sum_{\substack{k=l_0, \dots, l \\ \theta_k(x)=0}} \#P_k(x) \geq \sum_{k=l_0}^{\lfloor l/2 \rfloor + 1} \tilde{c}_0 \lambda(k) \geq \tilde{c}_0 \sum_{k=l_0}^{\lfloor l/2 \rfloor + 1} \frac{2^{-k} u'(2^{-k})}{u(2^{-k})} \\ &\geq \tilde{c}_0 \left( \log \frac{1}{u(2^{-l/2})} - \log \frac{1}{u(2^{-l_0})} \right) \geq \tilde{c}_0 \beta^{-1} \log \frac{1}{u(2^{-l})} - \tilde{c}_0 \log \frac{1}{u(2^{-l_0})}. \end{aligned}$$

Choosing  $\tilde{c}_0 > c_2 \beta$ , this cannot hold when  $l$  is large enough. Thus there is a number  $l_1 = l_1(\tilde{c}_0, l_0, u)$  such that (11) holds for all  $l \geq l_1$ .

Our next step is to prove that, if  $\theta_k(x) = 1$  for some  $k$  and  $P_k(x) = \{Q_1, \dots, Q_m\}$ , then it is possible to find a collection of balls  $\{B_1, \dots, B_{m'}\}$  from the families  $\mathcal{S}_{B(f_{Q_i}, r_{Q_i})}$  or  $\mathcal{R}_{B(f_{Q_i}, r_{Q_i})}$  having radii at least a constant times  $\alpha(2^{-k})$  and such that  $\sum_{i=1}^{m'} w_{B_i}$  is at least a constant times  $\lambda(k)$ . Moreover, we choose different balls for different  $k$ .

Let us fix  $k \geq l_0$  such that  $\theta_k(x) = 1$ . Suppose first that the annulus  $A_k(x)$  contains no centres from  $(f_{Q_j(\omega)})_{j=1}^\infty$ . Then the set  $P_k(x)$  consists of a single cube  $Q \in \mathcal{W}$  with  $f_Q \in B(x, 2^{-k})$ . The definitions of  $r_Q$  and  $l_0$  imply that  $r_Q > 2^{-k}$  and hence  $k \geq k_0(r_Q)$ . Thus, we may choose the ball  $R_k(B(f_Q, r_Q))$ , which, by definition, has radius  $\alpha(2^{-k})$  and weight  $\lambda(k)$ . In addition, the centre of this ball lies in  $B(x, 2^{-k})$ .

Assume now that the annulus  $A_k(x)$  contains at least one of the centres from  $(f_{Q_j(\omega)})_{j=1}^\infty$ . Then, by the definitions of  $P_k(x)$  and  $r_Q$ ,

$$\sum_{Q \in P_k(x)} 2r_Q \geq 2^{-k}.$$

Since  $\#P_k(x) \leq \tilde{c}_0\lambda(k)$ , we observe that

$$\sum_{\substack{Q \in P_k(x) \\ 2r_Q \geq \alpha(2^{-k})/2\tilde{c}_0}} 2r_Q \geq \frac{2^{-k}}{2}.$$

For each  $Q \in P_k(x)$  with  $2r_Q \geq \alpha(2^{-k})/2\tilde{c}_0$  we choose a number  $n_Q \in \mathbb{N}$  so that

$$2^{n_Q-1} \frac{\alpha(2^{-k})}{2\tilde{c}_0} \leq 2r_Q < 2^{n_Q} \frac{\alpha(2^{-k})}{2\tilde{c}_0}$$

and pick a ball  $\tilde{B} = S_{n_Q}(B(f_Q, r_Q)) = B(f_Q, r_Q/2^{n_Q}) \in \mathcal{G}_{B(f_Q, r_Q)}$ . By the definition of  $S_i(B)$ , we have  $w_{\tilde{B}} = 2^{n_Q}$  and

$$r(\tilde{B}) = \frac{r_Q}{2^{n_Q}} \geq \frac{\alpha(2^{-k})}{8\tilde{c}_0}.$$

For the sum of the weights  $\sum_Q 2^{n_Q}$  of all the balls obtained in such a manner, we observe that

$$\frac{\alpha(2^{-k})}{2\tilde{c}_0} \sum_{\substack{Q \in P_k(x) \\ 2r_Q \geq \alpha(2^{-k})/2\tilde{c}_0}} 2^{n_Q} > \sum_{\substack{Q \in P_k(x) \\ 2r_Q \geq \alpha(2^{-k})/2\tilde{c}_0}} 2r_Q \geq \frac{2^{-k}}{2}.$$

Hence, we have a collection of balls  $\{B_1, \dots, B_m\} \subset \mathcal{F}$  with weights sum  $\sum_{i=1}^m w_{B_i} > \tilde{c}_0\lambda(k)$  and of radii at least  $\alpha(2^{-k})/8\tilde{c}_0$ . Moreover, all these balls have their centres in the annulus  $A_k(x)$  and hence in the ball  $B(x, 2^{-k+1})$ .

We have proved that there exists a number  $l_1 = l_1(l_0, \tilde{c}_0)$  such that, for each  $\omega \in E_{l_0}$  and  $l \geq l_1$ , among the numbers  $l_0, \dots, l$  there are at least  $\lceil l/2 \rceil$  integers  $k \in \{l_0, \dots, l\}$  such that  $\theta_k(x) = 1$ . For these  $k$  we are able to find a finite collection of balls  $\{B_i\}_{i \in I} \subset \mathcal{F}$  with weight-sum  $\sum_{i \in I} w_{B_i}$  at least  $\lambda(k)$  and of radii at least  $\alpha(2^{-k})/8\tilde{c}_0$ , so that the centres of the balls  $B_i, i \in I$ , lie in the ball  $B(x, 2^{-k+1})$ . Here  $\tilde{c}_0$  is a positive constant depending only on  $\beta, n$  and  $\lambda(0)$ , and the balls are different for a fixed  $\omega$  and different  $k$ .

Fix  $l \geq l_1$ . We modify our family  $\mathcal{F}$  according to  $l$ . If  $B \in \mathcal{F}$  and there is  $k \in \{l_0 + 1, \dots, l\}$  such that  $\alpha(2^{-k})/8\tilde{c}_0 \leq r(B) < \alpha(2^{-k+1})/8\tilde{c}_0$ , we replace  $B$  with the ball  $\tilde{B} = (\lambda(k)/\lambda(l))B$  and set  $w_{\tilde{B}} = (\lambda(l)/\lambda(k))^n w_B$ . The radius of  $\tilde{B}$  satisfies  $r(\tilde{B}) \geq (\lambda(k)/\lambda(l))\alpha(2^{-k})/8\tilde{c}_0 = 2^{-k}/8\tilde{c}_0\lambda(l)$  and the equality  $w_{\tilde{B}}r(\tilde{B})^n = w_B r(B)^n$  holds. Similarly, we replace a ball  $B$  with  $r(B) \geq \alpha(2^{-l_0})/8\tilde{c}_0$  with the ball  $\tilde{B} = (\lambda(l_0)/\lambda(l))B$  and set  $w_{\tilde{B}} = (\lambda(l)/\lambda(l_0))^n w_B$ . Again, we have  $r(\tilde{B}) \geq 2^{-l_0}/8\tilde{c}_0\lambda(l)$  and  $w_{\tilde{B}}r(\tilde{B})^n = w_B r(B)^n$ . Finally,  $\mathcal{F}_l$  is the collection of balls obtained in this manner from the balls in  $\mathcal{F}$ . For this family of balls, we notice (see (10)) that

$$\sum_{B \in \mathcal{F}_l} w_B r(B)^n \leq \sum_{B \in \mathcal{F}} w_B r(B)^n < \infty. \tag{12}$$

If  $\omega \in E_{l_0}$ ,  $x = f(\omega)$  and  $k \in \{l_0, \dots, l\}$  is such that  $\theta_k(x) = 1$ , then there is a collection  $\{B_i\}_{i \in I} \subset \mathcal{F}$  with the properties mentioned above. If for some  $i \in I$  the ball  $B_i$  is replaced by the ball  $\tilde{B}_i = (\lambda(k_i)/\lambda(l))B_i$  while creating  $\mathcal{F}_l$ , we necessarily have  $k_i \leq k$ . Therefore, the inequalities

$$\sum_{i \in I} w_{\tilde{B}_i} = \sum_{i \in I} \left(\frac{\lambda(l)}{\lambda(k_i)}\right)^n w_{B_i} \geq \left(\frac{\lambda(l)}{\lambda(k)}\right)^n \sum_{i \in I} w_{B_i} \geq \left(\frac{\lambda(l)}{\lambda(k)}\right)^n \lambda(k) = \lambda(l)^n \frac{1}{\lambda(k)^{n-1}}$$

and  $r(\tilde{B}_i) \geq 2^{-k_i}/(8\tilde{c}_0\lambda(l)) \geq 2^{-k}/(8\tilde{c}_0\lambda(l))$  hold (by (6),  $\lambda$  is increasing). Since, for each  $i \in I$ , the centre of a ball  $\tilde{B}_i$  is contained in  $B(x, 2^{-k+1})$ , we have  $x \in 16\tilde{c}_0\lambda(l)\tilde{B}_i$ . Hence, we observe that

$$\sum_{B \in \mathcal{F}_l} w_B \chi_{16\tilde{c}_0\lambda(l)B}(y) \geq \sum_{\substack{k=l_0, \dots, l \\ \theta_k(y)=1}} \lambda(l)^n \frac{1}{\lambda(k)^{n-1}} \geq \frac{\lambda(l)^n}{4} \sum_{k=l_1}^l \frac{1}{\lambda(k)^{n-1}} \geq \frac{\lambda(l)^n}{4} G_l$$

for each  $y \in F_{l_0}$ , where  $G_l = \sum_{k=l_1}^l 1/\lambda(k)^{n-1}$ . That is,  $(4w_B/(\lambda(l)^n G_l), 16\tilde{c}_0\lambda(l)B)_{B \in \mathcal{F}_l}$  is a weighted cover of the set  $F_{l_0}$ . We observe also that the diameters of all balls in this cover are at least  $2^{-l}$ . This information will be used in the proof of Theorem 1.3 below.

Finally, using the weighted cover obtained above and (12), we estimate the weighted Hausdorff  $n$ -content  $\lambda_\infty^n(F_{l_0})$ :

$$\begin{aligned} \lambda_\infty^n(F_{l_0}) &\leq \frac{4}{\lambda(l)^n G_l} \sum_{B \in F_l} w_B (\text{diam}(16\tilde{c}_0\lambda(l)B))^n \leq \frac{4^{2n+1}\tilde{c}_0^n}{G_l} \sum_{B \in F_l} w_B (\text{diam } B)^n \\ &\leq \frac{2^{5n+2}\tilde{c}_0^n}{G_l} \sum_{B \in F_l} w_B r(B)^n \leq \frac{A}{G_l}, \end{aligned}$$

where the constant  $A$  depends on  $\beta, n, m, \|f\|_{W^{1,n}(\mathbb{B}^n, \mathbb{R}^m)}$  and  $\lambda(0)$  but not on  $l_0$  or  $l$ .

Now Lemma 2.1 implies  $\mathcal{H}_\infty^n(F_{l_0}) \leq CA/G_l$ . Here  $C$  depends only on the dimension  $n$ . Hence, we are done as soon as we can show that  $G_l \rightarrow \infty$  as  $l \rightarrow \infty$ . Towards this end, we have

$$G_l = \sum_{k=l_1}^l \frac{1}{\lambda(k)^{n-1}} = \sum_{k=l_1}^l \frac{u(2^{-k})^{n-1}}{2^{-k(n-1)}u'(2^{-k})^{n-1}} \geq \int_{2^{-l}}^{2^{-l_1}} \left(\frac{u(t)}{u'(t)}\right)^{n-1} \frac{dt}{t^n},$$

and the right-hand side diverges as  $l \rightarrow \infty$  by the assumptions on the modulus of continuity. □

The proof of Theorem 1.3 is similar to the proof of Theorem 1.1. We only point out the required changes.

*Proof of Theorem 1.3.* Let  $f$  be as in statement of the theorem. Our notation will be the same as in previous proof. That is,  $\alpha(t) = \gamma t$  and  $\lambda(k) = 1/\gamma$ .

Fix a small  $\varepsilon > 0$ . Then there exists a  $\delta > 0$  such that

$$\int_{\mathbb{B}^n \setminus B(0, 1-\delta)} |Df|^n \leq \varepsilon. \tag{13}$$

Let  $\mathcal{W}^\delta$  be the set of the cubes in  $\mathcal{W}$  which are contained in  $\mathbb{B}^n \setminus B(0, 1-\delta)$  and whose neighbour cubes are also contained in  $\mathbb{B}^n \setminus B(0, 1-\delta)$ . We define our collection of balls to be  $\mathcal{B}^\delta = \{B(f_Q, r_Q) : Q \in \mathcal{W}^\delta\}$ .

Then, proceeding as in the previous proof, we define  $\mathcal{F}^\delta$  analogously to  $\mathcal{F}$  and obtain the estimate (see (10))

$$\sum_{B \in \mathcal{F}^\delta} w_B r(B)^n \leq C_1 \varepsilon. \tag{14}$$

Let  $\omega \in \partial \mathcal{B}^n$ . We define the number  $l_0 = l_0(\omega, f, \delta)$  as in the previous proof, but instead of all cubes in  $(Q_j(\omega))_{j=1}^\infty$  we consider only those which are contained in  ${}^{\circ}W^\delta$ . Again, we split  $\partial \mathcal{B}^n$  into sets  $E_l = \{\omega \in \partial \mathcal{B}^n : l_0(\omega) \leq l\}$  and consider a fixed  $f(E_l)$ . With the same method as earlier we find for large  $l$  a collection of balls  $\mathcal{F}_l^\delta$  with weights such that  $(8w_B \gamma / (l - l_1), (16\tilde{c}_0/\gamma)B)_{B \in \mathcal{F}_l^\delta}$  is a weighted cover of the set  $f(E_l)$ , the radii of the balls  $(16\tilde{c}_0/\gamma)B$  are at least  $2^{-l}$  and

$$\sum_{B \in \mathcal{F}_l^\delta} w_B r(B)^n \leq C_1 \varepsilon.$$

We may assume that our  $\varepsilon > 0$  is so small that all balls in our weighted cover have radii smaller than  $\frac{1}{2}$ . With this weighted cover, we obtain

$$\begin{aligned} \lambda_\infty^g(f(E_l)) &\leq \frac{4\gamma}{l-l_1} \sum_{B \in \mathcal{F}_l^\delta} w_B \left( \text{diam} \left( \frac{16\tilde{c}_0}{\gamma} B \right) \right)^n \log \frac{1}{\text{diam}((16\tilde{c}_0/\gamma)B)} \\ &\leq \frac{4\gamma}{l-l_1} \sum_{B \in \mathcal{F}_l^\delta} w_B \left( \text{diam} \left( \frac{16\tilde{c}_0}{\gamma} B \right) \right)^n \log 2^l \leq \frac{2^{2+5n} \tilde{c}_0^n}{\gamma^{n-1}} \frac{l}{l-l_1} \sum_{B \in \mathcal{F}_l^\delta} w_B r(B)^n \leq \frac{2^{3+5n} \tilde{c}_0^n C_1}{\gamma^{n-1}} \varepsilon. \end{aligned}$$

Here we assumed  $l$  to be so large that  $l/(l - l_1) \leq 2$ . Lemma 2.1 implies  $\mathcal{H}_\infty^g(f(E_l)) \leq A\varepsilon$ . Here  $A$  depends on  $\gamma, n$  and  $m$  but not on  $l'$  or  $l$ ; therefore, we have  $\mathcal{H}_\infty^g(f(\partial \mathcal{B}^n)) \leq A\varepsilon$ ; see [Howroyd 1994, Corollary 8.2] or [Federer 1969, 2.10.22]. Letting  $\varepsilon$  tend to zero gives  $\mathcal{H}_\infty^g(f(\partial \mathcal{B}^n)) = 0$ , which implies  $\mathcal{H}^g(f(\partial \mathcal{B}^n)) = 0$ . □

### 4. Example

In this section, we work in  $\mathbb{R}^2$  and use the notation  $\|x\| = \max\{|x_1|, |x_2|\}$ . Let  $p > \frac{1}{2}$ . We will construct a locally Hölder continuous mapping  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  that belongs to  $W_{\text{loc}}^{1,2}(\mathbb{R}^2, \mathbb{R}^2)$  and maps  $\partial \mathcal{B}^2$  onto a set of positive  $\mathcal{H}^g$ -measure, where  $g(t) = t^2(\log(1/t))^{2p}$ .

The mapping is a composition of two locally Hölder continuous mappings. The second mapping is defined in [Herron and Koskela 2003, Proposition 5.1]. It is a homeomorphism  $h : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  that is the identity mapping outside  $[0, 1]^2$  and maps a small Cantor set  $\mathcal{C} \subset [0, 1]^2$  onto a large Cantor set  $\mathcal{C}' \subset [0, 1]^2$  with positive  $\mathcal{H}^g$ -measure. It was checked in [Koskela et al. 2009] that this mapping belongs to  $W_{\text{loc}}^{1,2}(\mathbb{R}^2, \mathbb{R}^2)$  if  $p > \frac{1}{2}$ .

Next, we elaborate on the construction of  $h$  and prove that it is Hölder continuous in  $[0, 1]^2$ . Let  $\sigma < \frac{1}{2}$ . We use the notation  $2r_k = \sigma^k$  and  $2R_k = \frac{1}{2}\sigma^{k-1}$  for  $k \in \mathbb{N}$ . The set  $\mathcal{C}$  is defined as follows: In the first generation we have one square  $Q_0 = [0, 1]^2$  with side length  $2r_0$ . We split this square into four subsquares  $P_{1i}, i = 1, 2, 3, 4$ , of side length  $2R_1$ . We define  $Q_{1i}$  to be the square of side length  $2r_1$  centred at the centre of  $P_{1i}$ . Then  $P_{1i}$  and  $Q_{1i}$  generate the frame  $A_{1i} = P_{1i} \setminus Q_{1i}$ . Next, we divide all squares  $Q_{1i}$  into

squares  $P_{2j}$ ,  $j = 1, \dots, 4^2$ . Then we define  $Q_{2j}$  and  $A_{2j}$  as in the first step. We proceed inductively. Thus, we obtain for all  $k \in \mathbb{N}$  sets  $Q_{ki}$ ,  $P_{ki}$  and  $A_{ki}$ , where  $i = 1, \dots, 2^{2k}$ , and we set  $\mathcal{C} = \bigcap_k \bigcup_i Q_{ki}$ .

The set  $\mathcal{C}'$  and sets  $Q'_{ki}$ ,  $P'_{ki}$  and  $A'_{ki}$  with  $k \in \mathbb{N}$  and  $i = 1, \dots, 2^{2k}$  are defined in the same way, using  $2r'_1 = \frac{1}{2}(\log 4)^{-p}$ ,  $2R'_2 = r'_1$ , and  $2r'_k = (\log 4)^{-p}2^{-k}k^{-p}$  and  $2R'_k = (\log 4)^{-p}2^{-k}(k - 1)^{-p}$  for other  $k \in \mathbb{N}$ .

The mapping  $h$  is defined so that it maps the frame  $A_{ki}$  to the frame  $A'_{ki}$  via a “radial” stretching and is continuous in  $[0, 1]^2$ . The radial stretching which maps  $A = \{x : r_k \leq \|x\| \leq R_k\}$  to  $A' = \{x : r'_k \leq \|x\| \leq R'_k\}$  is

$$\rho(x) = (a\|x\| + b)\frac{x}{\|x\|}, \quad \text{where } a = \frac{R'_k - r'_k}{R_k - r_k} \text{ and } b = \frac{R_k r'_k - R'_k r_k}{R_k - r_k}.$$

If  $x, y \in A$  then  $\|x - y\| \leq 2R_k = \frac{1}{2}\sigma^{k-1}$  and

$$a \leq \frac{4\sigma}{1 - 2\sigma}(2\sigma)^{-k} \leq C(\sigma)\sigma^{-(1-\beta)k} \leq C(\sigma)\|x - y\|^{\beta-1},$$

where  $\beta = \log 2 / \log(1/\sigma)$ . Similarly,

$$\frac{|b|}{|r_k|} \leq \frac{4}{1 - 2\sigma}(2\sigma)^{-k} \leq C(\sigma)\|x - y\|^{\beta-1}.$$

The mapping  $\rho$  is Hölder continuous with exponent  $\beta$ , as

$$\|\rho(x) - \rho(y)\| \leq Ca\|x - y\| + 2\frac{|b|}{|r_k|}\|x - y\| \leq C(\sigma)\|x - y\|^\beta.$$

If  $x \in A_{ki}$  and  $y \in Q_{k+1,j} \subset P_{ki}$ , then  $\|x - y\| \geq R_{k+1} - r_{k+1} = C(\sigma)\sigma^k$  and  $\|h(x) - h(y)\| \leq 2R'_k \leq 2^{-k}$ . These imply

$$\frac{\|h(x) - h(y)\|}{\|x - y\|^\beta} \leq C(\sigma).$$

The  $\beta$ -Hölder continuity of  $h$  easily follows from the continuity estimates obtained above.

The first mapping  $G : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a (locally Hölder continuous) quasiconformal mapping for which  $\mathcal{C} \subset G(\partial\mathbb{B}^2)$ . Such a mapping was constructed in [Gehring and Väisälä 1973].

Finally, the composition  $h \circ G : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a homeomorphism with  $h \circ G(\partial\mathbb{B}^2) \supset \mathcal{C}'$ . Moreover, it is locally Hölder continuous and  $h \circ G \in W_{loc}^{1,2}(\mathbb{R}^2, \mathbb{R}^2)$  by quasiconformality of  $G$  and the change of variable formula; see, for example, [Astala et al. 2009, Section 3.8].

### References

[Astala et al. 2009] K. Astala, T. Iwaniec, and G. Martin, *Elliptic partial differential equations and quasiconformal mappings in the plane*, Princeton Mathematical Series **48**, Princeton University Press, NJ, 2009. MR 2010j:30040 Zbl 1182.30001

[Buckley 1996] S. M. Buckley, “Space-filling curves and related functions”, *Irish Math. Soc. Bull.* 36 (1996), 9–18. MR 97b:26004 Zbl 0847.26004

[Federer 1969] H. Federer, *Geometric measure theory*, Die Grundlehren der mathematischen Wissenschaften **153**, Springer, New York, 1969. MR 41 #1976 Zbl 0176.00801

[Gehring and Väisälä 1973] F. W. Gehring and J. Väisälä, “Hausdorff dimension and quasiconformal mappings”, *J. London Math. Soc.* (2) **6** (1973), 504–512. MR 48 #2380 Zbl 0258.30020

- [Hencl et al. 2012] S. Hencl, P. Koskela, and T. Nieminen, “Dimension gap under conformal mappings”, *Adv. Math.* **230**:3 (2012), 1423–1441. MR 2921185 Zbl 1253.30034
- [Herron and Koskela 2003] D. A. Herron and P. Koskela, “Mappings of finite distortion: gauge dimension of generalized quasicircles”, *Illinois J. Math.* **47**:4 (2003), 1243–1259. MR 2005a:30031 Zbl 1050.30012
- [Howroyd 1994] J. D. Howroyd, *On the theory of Hausdorff measures in metric spaces*, Ph.D. thesis, University College London, 1994, <http://smithsold.gold.ac.uk/jhowroyd/jdhthesis.pdf>.
- [Jones and Makarov 1995] P. W. Jones and N. G. Makarov, “Density properties of harmonic measure”, *Ann. of Math. (2)* **142**:3 (1995), 427–455. MR 96k:30027 Zbl 0842.31001
- [Koskela and Rohde 1997] P. Koskela and S. Rohde, “Hausdorff dimension and mean porosity”, *Math. Ann.* **309**:4 (1997), 593–609. MR 98k:28004 Zbl 0890.30013
- [Koskela and Zapadinskaya 2014] P. Koskela and A. Zapadinskaya, “Dimension gap under Sobolev mappings”, preprint, 2014.
- [Koskela et al. 2009] P. Koskela, A. Zapadinskaya, and T. Zürcher, “Generalized dimension distortion under planar Sobolev homeomorphisms”, *Proc. Amer. Math. Soc.* **137**:11 (2009), 3815–3821. MR 2011b:30051 Zbl 1177.30022
- [Malý and Martio 1995] J. Malý and O. Martio, “Lusin’s condition (N) and mappings of the class  $W^{1,n}$ ”, *J. Reine Angew. Math.* **458** (1995), 19–36. MR 95m:26024 Zbl 0812.30007
- [Nieminen 2006] T. Nieminen, “Generalized mean porosity and dimension”, *Ann. Acad. Sci. Fenn. Math.* **31**:1 (2006), 143–172. MR 2007j:28006 Zbl 1099.30009
- [Stein 1970] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series **30**, Princeton University Press, Princeton, N.J., 1970. MR 44 #7280 Zbl 0207.13501

Received 30 Jul 2013. Revised 27 Aug 2014. Accepted 22 Oct 2014.

AAPO KAURANEN: [aapo.p.kauranen@jyu.fi](mailto:aapo.p.kauranen@jyu.fi)

*Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), FI-40014 Jyväskylä, Finland*

PEKKA KOSKELA: [pekka.j.koskela@jyu.fi](mailto:pekka.j.koskela@jyu.fi)

*Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), FI-40014 Jyväskylä, Finland*

# Analysis & PDE

msp.org/apde

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
zworski@math.berkeley.edu  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 7 No. 8 2014

---

Quantized slow blow-up dynamics for the corotational energy-critical harmonic heat flow	1713
PIERRE RAPHAËL and REMI SCHWEYER	
Existence and orbital stability of the ground states with prescribed mass for the $L^2$ -critical and supercritical NLS on bounded domains	1807
BENEDETTA NORIS, HUGO TAVARES and GIANMARIA VERZINI	
Boundary blow-up under Sobolev mappings	1839
AAPO KAURANEN and PEKKA KOSKELA	
Global gauges and global extensions in optimal spaces	1851
MIRCEA PETRACHE and TRISTAN RIVIÈRE	
Concentration of small Willmore spheres in Riemannian 3-manifolds	1901
PAUL LAURAIN and ANDREA MONDINO	
Hole probabilities of $SU(m + 1)$ Gaussian random polynomials	1923
JUNYAN ZHU	
Stochastic homogenization of viscous Hamilton–Jacobi equations and applications	1969
SCOTT N. ARMSTRONG and HUNG V. TRAN	
Global regularity for a slightly supercritical hyperdissipative Navier–Stokes system	2009
DAVID BARBATO, FRANCESCO MORANDIN and MARCO ROMITO	



2157-5045(2014)7:8;1-8