

ANALYSIS & PDE

Volume 8

No. 1

2015

TUOMO KUUSI, GIUSEPPE MINGIONE AND YANNICK SIRE

NONLOCAL SELF-IMPROVING PROPERTIES

NONLOCAL SELF-IMPROVING PROPERTIES

TUOMO KUUSI, GIUSEPPE MINGIONE AND YANNICK SIRE

Solutions to nonlocal equations with measurable coefficients are higher differentiable.

Specifically, we consider nonlocal integrodifferential equations with measurable coefficients whose model is given by

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} [u(x) - u(y)][\eta(x) - \eta(y)]K(x, y) dx dy = \int_{\mathbb{R}^n} f \eta dx \quad \text{for all } \eta \in C_c^\infty(\mathbb{R}^n),$$

where the kernel $K(\cdot)$ is a measurable function and satisfies the bounds

$$\frac{1}{\Lambda |x - y|^{n+2\alpha}} \leq K(x, y) \leq \frac{\Lambda}{|x - y|^{n+2\alpha}}$$

with $0 < \alpha < 1$, $\Lambda > 1$, while $f \in L_{loc}^q(\mathbb{R}^n)$ for some $q > 2n/(n + 2\alpha)$. The main result states that there exists a positive, universal exponent $\delta \equiv \delta(n, \alpha, \Lambda, q)$ such that for every weak solution u the self-improving property

$$u \in W^{\alpha,2}(\mathbb{R}^n) \implies u \in W_{loc}^{\alpha+\delta,2+\delta}(\mathbb{R}^n)$$

holds. This differentiability improvement is a genuinely nonlocal phenomenon and does not appear in the local case, where solutions to linear equations in divergence form with measurable coefficients are known to be higher integrable but are not, in general, higher differentiable.

The result is achieved by proving a new version of the Gehring lemma involving certain families of lifted reverse Hölder-type inequalities in \mathbb{R}^{2n} and which is implied by delicate covering and exit-time arguments. In turn, such reverse Hölder inequalities are based on the concept of dual pairs, that is, pairs (μ, U) of measures and functions in \mathbb{R}^{2n} which are canonically associated to solutions. We also allow for more general equations involving as a source term an integrodifferential operator whose kernel does not necessarily have to be of order α .

1. Introduction	58
2. Preliminaries and notation	64
3. The Caccioppoli inequality	66
4. The dual pair (μ, U) and reverse inequalities	75
5. Level set estimates for dual pairs	82
6. Self-improving inequalities	107
Acknowledgments	113
References	113

MSC2010: primary 35D10; secondary 35R11.

Keywords: elliptic equations, fractional differentiability, nonlocal operators.

1. Introduction

A basic and fundamental result in the theory of linear and nonlinear elliptic equations is given by the higher integrability of solutions. This falls in the realm of so-called self-improving properties. The result was first pioneered by Meyers [1963] and Elcrat and Meyers [1975], and then extended in various directions and in several different contexts; see for instance [Bojarski and Iwaniec 1983; Fusco and Sbordone 1990; Giusti 2003; Kinnunen and Lewis 2000]. Modern proofs of this property in the nonlinear case rely on the so-called Gehring lemma [Gehring 1973; Iwaniec 1998]. In the simplest possible instance the result in question asserts that distributional $W^{1,2}(\Omega)$ -solutions u to linear elliptic equations

$$-\operatorname{div}(A(x)Du) = f \in L_{\text{loc}}^{\frac{2n}{n+2}+\delta_0}(\Omega), \quad \delta_0 > 0,$$

actually belong to a better Sobolev space:

$$u \in W_{\text{loc}}^{1,2+\delta}(\Omega), \tag{1-1}$$

for some positive $\delta \leq \delta_0$. Here $\Omega \subset \mathbb{R}^n$ is an open subset and $n \geq 2$. The matrix $A(x)$ is supposed to be elliptic and with bounded and measurable entries, that is,

$$\Lambda^{-1}|\xi|^2 \leq \langle A(x)\xi, \xi \rangle \quad \text{and} \quad |A(x)| \leq \Lambda \tag{1-2}$$

hold whenever $\xi \in \mathbb{R}^n$, $x \in \Omega$, where $\Lambda > 1$. The number $\delta > 0$ appearing in (1-1) is universal in the sense that, essentially, it depends neither on the solution u nor the specific equation considered. It rather depends only on n , Λ , that is, on the ellipticity rate of the equation considered. The key point here is the measurability of the coefficients; when $A(\cdot)$ has more regular entries, higher regularity of solutions follows from the corresponding result for equations with constant coefficients, via perturbation. This is the reason why the result in (1-1) lies deep in the core of regularity theory, and allows for a proof of several other regularity results; see for instance [Giusti 2003].

We are interested in studying self-improving properties of solutions to nonlocal problems. To outline the results in a special yet meaningful model case, let us consider weak solutions $u \in W^{\alpha,2}(\mathbb{R}^n)$ of the nonlocal equation

$$\mathcal{E}_K(u, \eta) = \langle f, \eta \rangle \quad \text{for every test function } \eta \in C_c^\infty(\mathbb{R}^n), \tag{1-3}$$

where $f \in L_{\text{loc}}^{2+\delta_0}(\mathbb{R}^n)$ and

$$\mathcal{E}_K(u, \eta) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} [u(x) - u(y)][\eta(x) - \eta(y)]K(x, y) dx dy.$$

The measurable kernel is assumed to satisfy the uniform ellipticity assumptions

$$\frac{1}{\Lambda|x-y|^{n+2\alpha}} \leq K(x, y) \leq \frac{\Lambda}{|x-y|^{n+2\alpha}} \tag{1-4}$$

for every $x, y \in \mathbb{R}^n$, where $\alpha \in (0, 1)$ and $\Lambda \geq 1$. We recall that the fractional Sobolev space $W^{s,\gamma}$, for $\gamma \geq 1$ and $s \in (0, 1)$, is given by the subspace of $L^\gamma(\mathbb{R}^n)$ -functions u for which the Gagliardo seminorm

$$[u]_{s,\gamma}^\gamma := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^\gamma}{|x - y|^{n+\gamma s}} dx dy \quad (1-5)$$

is finite (see for instance [Di Nezza et al. 2012; Maz'ya 2011]).

In view of (1-1), a natural question to begin with is whether or not the inclusion

$$u \in W_{\text{loc}}^{\alpha,2+\delta}(\mathbb{R}^n) \quad (1-6)$$

holds for some $\delta > 0$, possibly depending only on the ellipticity parameters of the equation and not on the solution itself. For the definition of local fractional Sobolev spaces, see Section 2. This has been answered in a very interesting paper of Bass and Ren [2013], who consider the function

$$\Gamma(x) := \left(\int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dy \right)^{1/2}, \quad (1-7)$$

and prove that $\Gamma \in L^{2(1+\delta)}(\mathbb{R}^n)$ for some positive δ depending only on n, α, Λ and δ_0 . Then (1-6) follows by characterizations of Bessel potential spaces [Dorronsoro 1985; Stein 1961]. In this paper we provide a stronger and surprising result. Indeed, we see that *for nonlocal problems the self-improvement property extends to the differentiability scale*. This means that there exists some positive $\delta \in (0, 1 - \alpha)$, depending only on n, α, Λ , such that

$$u \in W_{\text{loc}}^{\alpha+\delta,2+\delta}(\mathbb{R}^n). \quad (1-8)$$

This phenomenon is purely nonlocal, and has no parallel in the regularity theory of local equations, where, in order to get fractional Sobolev differentiability of Du , a similar fractional regularity must be assumed on the coefficient matrix $A(x)$, as for instance established in [Kuusi and Mingione 2012; Mingione 2003].

In the classical local case, measurability is, in general, not sufficient to get any gradient differentiability. To see this already in the one-dimensional case, $n = 1$, it is sufficient to consider the equation

$$\frac{d}{dx} \left(a(x) \frac{du}{dx} \right) = 0, \quad \frac{1}{\Lambda} \leq a(x) \leq \Lambda, \quad (1-9)$$

and to note that

$$x \mapsto \int_0^x \frac{dt}{a(t)}$$

is a solution with $a(\cdot)$ being any measurable function satisfying nothing but the inequalities in (1-9). It is then easy to build similar multidimensional examples.

We remark that the differentiability gain is in fact the main information in (1-8), since a standard application of the fractional Sobolev embedding theorem gives that, if $u \in W^{\alpha+\delta,2}$ for some $\delta > 0$, then (1-8) holds for some other number δ . Our results actually cover a more general class of equations than the one in (1-3) and provide a full nonlocal analog of the classical higher integrability results valid in the local case. The precise statements are in the next section. Our results are a consequence of a new,

fractional version of the Gehring lemma for fractional Sobolev functions that replaces the classical one valid in the local case.

We finally remark that, in recent times, there has been much attention to the regularity of solutions to nonlocal problems, especially in the basic case of kernels with measurable coefficients; see for instance [Bass and Kassmann 2005; Bjorland et al. 2012; Cabré and Cinti 2014; Cabré and Roquejoffre 2013; Caffarelli et al. 2011; Caffarelli and Silvestre 2011; Felsinger and Kassmann 2013].

1A. Higher differentiability results. A rather general statement concerning higher integrability for weak solutions to local problems involves nonhomogeneous equations such as

$$-\operatorname{div}(A(x)Du) = -\operatorname{div} g + f \quad \text{in } \Omega, \quad (1-10)$$

where the matrix $A(\cdot)$ has measurable coefficients and satisfies (1-2). Indeed, assuming that $g \in L_{\text{loc}}^{2+\delta_0}(\Omega, \mathbb{R}^n)$ and $f \in L_{\text{loc}}^{2n/(n+2)+\delta_0}(\Omega)$ hold for some $\delta_0 > 0$, it follows that there exists another positive number $\delta < \delta_0$ such that (1-1) holds. The exponent $2n/(n+2)$ is nothing but the conjugate of the Sobolev embedding exponent of $W^{1,2}$, that is, $2n/(n-2)$.

A first nonlocal analog of (1-10) is given by

$$\mathcal{E}_K(u, \eta) = \mathcal{E}_K(g, \eta) + \int_{\mathbb{R}^n} f \eta \, dx \quad \text{for all } \eta \in C_c^\infty(\mathbb{R}^n), \quad (1-11)$$

considering weak solutions $u \in W^{\alpha,2}(\mathbb{R}^n)$. The assumptions are the natural counterpart of the local ones; indeed, we take $g \in W^{\alpha+\delta_0,2}(\mathbb{R}^n)$ and

$$f \in L_{\text{loc}}^{2_*+\delta_0}(\mathbb{R}^n) \quad (1-12)$$

for some $\delta_0 > 0$. The exponent 2_* is the conjugate of the relevant fractional Sobolev embedding exponent, that is,

$$2_* := \frac{2n}{n+2\alpha}, \quad 2^* := \frac{2n}{n-2\alpha}, \quad \frac{1}{2_*} + \frac{1}{2^*} = 1. \quad (1-13)$$

The terminology is motivated by the fractional version of the classical Sobolev embedding theorem, that is, $W^{\alpha,2} \subset L^{2^*}$. On the other hand, we recall that the essence of the structure of (1-10) lies in the fact that the right-hand side contains terms of all possible integer orders. A full extension to the fractional case then leads us to consider right-hand sides of *arbitrary fractional order*, not necessarily equal to the order of the considered nonlocal elliptic operator on the left-hand side. Moreover, since higher integrability of solutions still holds when considering monotone quasilinear equations, we will also examine nonlinear integrodifferential equations. Specifically, we will consider general equations of the type

$$\mathcal{E}_K^\varphi(u, \eta) = \mathcal{E}_H(g, \eta) + \int_{\mathbb{R}^n} f \eta \, dx \quad \text{for all } \eta \in C_c^\infty(\mathbb{R}^n). \quad (1-14)$$

The form $\mathcal{E}_K^\varphi(\cdot)$ is then defined by

$$\mathcal{E}_K^\varphi(u, \eta) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \varphi(u(x) - u(y))[\eta(x) - \eta(y)]K(x, y) \, dx \, dy,$$

where the Borel function $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ satisfies

$$|\varphi(t)| \leq \Lambda|t| \quad \text{and} \quad \varphi(t)t \geq t^2 \quad \text{for all } t \in \mathbb{R}, \quad (1-15)$$

making in fact \mathcal{E}_K^φ a coercive form in $W^{\alpha,2}$, and thereby (1-14) an elliptic equation. While we assume (1-4) for $K(\cdot)$, the measurable kernel $H(\cdot)$ is now assumed to satisfy

$$|H(x, y)| \leq \frac{\Lambda}{|x - y|^{n+2\beta}} \quad (1-16)$$

for $\beta \in (0, 1)$. In particular, β is also allowed to be larger than α . Here the function f is still assumed to satisfy (1-12), while the assumptions on g sharply match the structure in (1-14). We actually consider two different cases; the first one is when $2\beta \geq \alpha$. In this situation we assume the existence of a positive number $\delta_0 > 0$ such that

$$g \in W^{2\beta-\alpha+\delta_0,2}(\mathbb{R}^n). \quad (1-17)$$

Needless to say, we also assume that $2\beta - \alpha + \delta_0 \in (0, 1)$ to give (1-17) a sense in terms of the seminorm (1-5); this in particular implies that $\beta < \frac{1}{2}(1 + \alpha)$. In the case $0 < 2\beta < \alpha$ we instead do not consider any differentiability on g , but only integrability:

$$g \in L^{p_0+\delta_0}(\mathbb{R}^n), \quad p_0 := \frac{2n}{n + 2(\alpha - 2\beta)}. \quad (1-18)$$

We then have the following *main result* of the paper:

Theorem 1.1. *Let $u \in W^{\alpha,2}(\mathbb{R}^n)$ be a solution to (1-14) under the assumptions (1-4) and (1-12)–(1-18). Then there exists a positive number $\delta \in (0, 1 - \alpha)$, depending only on $n, \alpha, \Lambda, \beta, \delta_0$, but otherwise independent of the solution u and of the kernels $K(\cdot), H(\cdot)$, such that $u \in W_{\text{loc}}^{\alpha+\delta,2+\delta}(\mathbb{R}^n)$.*

Equation (1-11) is covered by taking $\alpha = \beta$. The optimality of the assumptions on f and g can be checked by considering the model equation $(-\Delta)^\alpha u = (-\Delta)^\beta g + f$, and using Fourier analysis. They sharply relate to the fractional Sobolev embedding theorem. As in the case of the classical, local Gehring lemma, explicit estimates on the exponent δ for Theorem 1.1 can be given by tracing back the dependence of the constants in the proof.

1B. Dual pairs (μ, U) and sketch of the proof. In order to get (1-8) we introduce here a new approach and develop a method aimed at exploiting the *hidden cancellation properties* which are intrinsic in the definition of the nonlocal seminorm (1-5). To this aim, we introduce dual pairs of measures and functions (μ, U) in \mathbb{R}^{2n} , proving that a version of the Gehring lemma applies to them; see Section 1C below. A natural choice would be to consider the measure generated by the density $|x - y|^{-n}$, but this would not yield a finite measure. We therefore consider a perturbation of it, i.e., the measure defined by

$$\mu(A) := \int_A \frac{dx dy}{|x - y|^{n-2\varepsilon}} \quad (1-19)$$

for suitably small $\varepsilon > 0$, whenever $A \subset \mathbb{R}^{2n}$ is a measurable subset. This is a locally finite, doubling Borel measure in \mathbb{R}^{2n} . Accordingly, for $x \neq y$, we introduce the function

$$U(x, y) := \frac{|u(x) - u(y)|}{|x - y|^{\alpha + \varepsilon}}. \quad (1-20)$$

The main point here is that the measure μ and the function U are in duality when $u \in W^{\alpha, 2}$ in the sense that for a function $u \in L^2(\mathbb{R}^n)$ we have that $U \in L^2(\mathbb{R}^{2n}; \mu)$ if and only if $u \in W^{\alpha, 2}(\mathbb{R}^n)$. This motivates in fact the following:

Definition 1.2. Let $u \in W^{\alpha, 2}(\mathbb{R}^n)$ and let $\varepsilon \in (0, \frac{1}{2}\alpha)$. The pair (μ, U) defined in (1-19)–(1-20) is called a dual pair generated by the function u .

We then look at the higher μ -integrability for U , proving that

$$U \in L_{\text{loc}}^{2+\delta}(\mathbb{R}^{2n}; \mu) \quad (1-21)$$

for some $\delta > 0$. Now, by the very definition of U , we have that (1-21) implies the higher differentiability of u , that is, (1-8); see Section 6. This is the effect of the cancellations hidden in the definition of fractional norm in (1-5) we mentioned above. In order to prove (1-21), we shall prove decay estimates for the μ -measure of the level sets of U . The first step consists of deriving suitable energy estimates (i.e., Caccioppoli-type inequalities) for U ; see Theorem 3.2. We obtain a kind of reverse Hölder-type inequality, that is,

$$\left(\int_B U^2 d\mu \right)^{1/2} \lesssim \sum_{k=1}^{\infty} 2^{-k(\alpha - \varepsilon)} \left(\int_{2^k B} U^q d\mu \right)^{1/q} + \text{terms involving } g, f, \quad (1-22)$$

with $q < 2$; see Proposition 4.4. The estimate in (1-22) holds whenever $\mathcal{B} \equiv B \times B$ and $B \subset \mathbb{R}^n$ is a ball. Notice that if we discard from the sum above all the terms but the first one we formally obtain a reverse Hölder-type inequality similar to those that hold for solutions to local problems.

The inequality (1-22) does not seem to be sufficient to proceed, since in order to prove estimates on level sets in \mathbb{R}^{2n} we need information on every ball $\mathcal{B} \subset \mathbb{R}^{2n}$, not only those of diagonal type $B \times B$. To overcome such an apparently decisive lack of information, we have to introduce an extremely delicate localization technique. Consider the level set $\{U > \lambda\}$; we use a Calderón–Zygmund-type exit-time argument in order to cover the level set with (almost disjoint) diagonal balls $B \times B$ and disjoint “off-diagonal” dyadic cubes \mathcal{K} :

$$\{U > \lambda\} \subset \bigcup B \times B \cup \bigcup \mathcal{K},$$

on which, for a suitably large number L , we have

$$\left(\int_{B \times B} U^2 d\mu \right)^{1/2} \approx \lambda \quad \text{and} \quad \left(\int_{\mathcal{K}} U^2 d\mu \right)^{1/2} \approx L\lambda;$$

see Sections 5A and 5F. We call the cubes \mathcal{K} off-diagonal because they are “far” from the diagonal, in the sense that their distance from the diagonal is larger than their side length. The number L is introduced to make the decomposition along the diagonal predominant with respect to the decomposition outside the

diagonal. Indeed, the exit-time balls $B \times B$ will tend to be “larger” than the cubes \mathcal{K} , since they have been obtained via an exit time at a lower level λ , as shown by the first formula in the previous display.

Surprisingly enough, the fact that a cube \mathcal{K} is off-diagonal allows us to prove that a reverse inequality of the type (1-22) also holds on \mathcal{K} (see Lemma 5.8). This inequality, however, incorporates certain correction terms involving diagonal cubes once again. This introduces serious difficulties, since this time such cubes do not come from any exit-time argument, and there is no a priori control on them. Matching the resulting reverse inequalities with those in (1-22) is not an easy task and indeed requires an involved covering/combinatorial argument. See Sections 5I and 5J, and in particular Lemma 5.12.

The final outcome of this lengthy procedure is an inequality on level sets of U — see Proposition 5.1 — that implies the *higher integrability* of U , together with the new reverse Hölder-type inequality reported in (1-24) below. This holds for some $\delta > 0$ that does not depend on the solution u . See Theorem 6.1. We have therefore proved (1-21). We also remark that treating the complete problem of Theorem 1.1 up to the sharp interpolation range described by (1-17) requires additional ideas. As a matter of fact, the exit-time arguments have to be adapted in order to realize a direct analog of the so-called good- λ inequality principle: i.e., no maximal operator is used here. In particular, we employ a simultaneous level set analysis by using the composite quantity $\Psi(\cdot)$ in (5-1), where the number M (appearing in the definition of $\Psi(\cdot)$) is used to adapt the size of the levels at the exit time. This must eventually match with the specific form of the energy estimates available for solutions.

Finally, we would like to remark that, although we are here dealing with the case of scalar, linear growth nonlocal equations, our approach is only based on energy inequalities, and therefore can be extended to more general nonlinear operators of nonlocal type; see for example [Di Castro et al. 2014a; 2014b]. This will be the object of future works.

1C. The fractional Gehring lemma for dual pairs. The classical Gehring lemma does not simply deal with solutions to equations, but, more generally, with self-improving properties of reverse Hölder-type inequalities. At the core of our approach lies in fact a new, fractional version of the Gehring lemma valid for general fractional Sobolev functions, and not only for solutions to nonlocal equations. Here is a version of it.

Theorem 1.3 (fractional Gehring lemma). *Let $u \in W^{\alpha,2}(\mathbb{R}^n)$ for $\alpha \in (0, 1)$. Let $\varepsilon \in (0, \alpha/2)$ and let (μ, U) be the dual pair generated by u in the sense of (1-19)–(1-20) and Definition 1.2. Assume that the following reverse Hölder-type inequality with the tail holds for every $\sigma \in (0, 1)$ and for every ball $B \subset \mathbb{R}^n$:*

$$\left(\int_B U^2 d\mu \right)^{1/2} \leq \frac{c(\sigma)}{\varepsilon^{1/q-1/2}} \left(\int_{2B} U^q d\mu \right)^{1/q} + \frac{\sigma}{\varepsilon^{1/q-1/2}} \sum_{k=2}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k B} U^q d\mu \right)^{1/q}, \quad (1-23)$$

where $q \in (1, 2)$ is a fixed exponent, $\mathcal{B} = B \times B$ and $c(\sigma)$ is a nonincreasing function depending on σ . Then there exists a positive number $\delta \in (0, 1 - \alpha)$, depending only on $n, \alpha, \varepsilon, q$ and the function $c(\cdot)$, such that $U \in L_{\text{loc}}^{2+\delta}(\mathbb{R}^{2n}; \mu)$ and $u \in W_{\text{loc}}^{\alpha+\delta, 2+\delta}(\mathbb{R}^n)$. Moreover, the following inequality holds whenever

$B \subset \mathbb{R}^n$, again for a constant c depending only on $n, \alpha, \varepsilon, q$ and the function $c(\cdot)$:

$$\left(\int_B U^{2+\delta} d\mu \right)^{1/(2+\delta)} \leq c \sum_{k=1}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k B} U^2 d\mu \right)^{1/2}. \quad (1-24)$$

In the literature there are several extensions of Gehring's lemma in general settings, for instance in metric spaces equipped with a doubling Borel measure, but [Theorem 1.3](#) is completely different. Indeed, its central feature is actually that global higher integrability information is reconstructed from reverse inequalities that do not hold on every ball in \mathbb{R}^{2n} , but only on diagonal ones. This is a crucial loss of information that makes [Theorem 1.3](#) hold not for any function $U \in L^2(\mathbb{R}^{2n}; \mu)$, but rather only for dual pairs (μ, U) . Moreover, the presence of the infinite series on the right-hand side of (1-23) gives to this inequality a delicate nonlocal character that adds relevant technical complications. [Theorem 1.3](#) is a particular case of a more general result; we prefer to report this form again to make the basic ideas more transparent. A more comprehensive version including additional functions F and G on the right-hand side of (1-23) can be proved as well; see [Theorem 6.1](#) below.

The results of this paper have been announced in the preliminary research report [[Kuusi et al. 2014](#)].

2. Preliminaries and notation

In what follows we denote by c a general positive constant, possibly varying from line to line; special occurrences will be denoted by $c_1, c_2, \bar{c}_1, \bar{c}_2$ or the like. All such constants will always be *greater than or equal to one*; moreover, relevant dependencies on parameters will be emphasized using parentheses, i.e., $c_1 \equiv c_1(n, \Lambda, p, \alpha)$ means that c_1 depends only on n, Λ, p, α . We denote by

$$B(x_0, r) \equiv B_r(x_0) := \{x \in \mathbb{R}^n : |x - x_0| < r\}$$

the open ball with center x_0 and radius $r > 0$; when not important, or clear from the context, we shall omit denoting the center by writing $B_r \equiv B(x_0, r)$; moreover, with B being a generic ball with radius r , we will denote by σB the ball concentric to B having radius σr , $\sigma > 0$. Unless otherwise stated, different balls in the same context will have the same center. With $\mathcal{O} \subset \mathbb{R}^k$ being a measurable set with positive μ -measure and with h being a measurable map, we shall denote by

$$(h)_{\mathcal{O}} \equiv \int_{\mathcal{O}} h d\mu := \frac{1}{\mu(\mathcal{O})} \int_{\mathcal{O}} h d\mu$$

its integral average. We shall need to consider integrals and functions in $\mathbb{R}^n \times \mathbb{R}^n$. In this respect, instead of dealing with the usual balls in \mathbb{R}^{2n} , we prefer to deal with balls generated by a different metric, that is, that relative to the norm (in \mathbb{R}^{2n}) defined by

$$\|(x_0, y_0)\| := \max\{|x_0|, |y_0|\}, \quad (2-1)$$

where $|\cdot|$ denotes the standard Euclidean norm in \mathbb{R}^n and $x_0, y_0 \in \mathbb{R}^n$. These balls are denoted by $\mathcal{B}(x_0, y_0, \varrho)$, and are of course of the form

$$\mathcal{B}(x_0, y_0, \varrho) := B(x_0, \varrho) \times B(y_0, \varrho).$$

In the case $x_0 = y_0$ we shall also use the shorter notation $\mathcal{B}(x_0, x_0, \varrho) \equiv \mathcal{B}(x_0, \varrho)$. With obvious meaning, these will be called diagonal balls. Moreover, with $\mathcal{B}(x_0, \varrho)$ being a fixed ball, we shall also denote $\mathcal{B} \equiv \mathcal{B}(x_0, x_0, \varrho)$ when no ambiguity shall arise, and $s\mathcal{B} := \mathcal{B}(x_0, s\varrho)$ for $s > 0$. Needless to say, since they are metric balls, and actually equivalent to the standard ones in \mathbb{R}^{2n} , we can apply to them several tools that are available for the usual balls. For instance, we shall later on apply the classical Vitali covering lemma. It follows that

$$B_{\mathbb{R}^{2n}}((x_0, y_0), \varrho) = \{(x_0, y_0) \in \mathbb{R}^{2n} : |(x_0, y_0)| < \varrho\} \subset \mathcal{B}(x_0, y_0, \varrho).$$

Accordingly, we shall denote

$$\text{Diag} := \{(x, x) \in \mathbb{R}^{2n} : x \in \mathbb{R}^n\}.$$

If A is a finite set, the symbol $\#A$ denotes the number of its elements. We shall very often use the elementary inequality

$$2^{2\beta k} \sum_{j=k-1}^{\infty} 2^{-2\beta j} \leq \frac{8}{\beta} \quad \text{for } \beta \in (0, 1] \text{ and } k \geq 1. \quad (2-2)$$

Finally, the local fractional Sobolev spaces are defined via the Gagliardo seminorm

$$[u]_{s,\gamma}(\Omega) := \left(\int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{\gamma}}{|x - y|^{n+\gamma s}} dx dy \right)^{1/\gamma} \quad (2-3)$$

for $\gamma \geq 1$ and $s \in (0, 1)$. A function $u \in L^{\gamma}_{\text{loc}}(\mathbb{R}^n)$ belongs to $W^{s,\gamma}_{\text{loc}}(\mathbb{R}^n)$ if $[u]_{s,\gamma}(\Omega)$ is finite whenever Ω is an open bounded subset of \mathbb{R}^n .

The following two lemmas report some classical Poincaré–Sobolev-type inequalities valid in the fractional setting; the proof of the first is exactly the one in [Mingione 2003], for the second we refer to [Kassmann 2009]. See also [Di Nezza et al. 2012; Maz'ya 2011].

Lemma 2.1 (fractional Poincaré inequality). *Let $v \in L^p(B)$, with $B \subset \mathbb{R}^n$ being a ball of radius r , and let α be a real number such that $n + p\alpha \geq 0$; then the following inequality holds:*

$$\int_B |v - (v)_B|^p dx \leq cr^{p\alpha} \int_B \int_B \frac{|v(x) - v(y)|^p}{|x - y|^{n+p\alpha}} dx dy.$$

This inequality in particular applies when $v \in W^{\alpha,p}(B)$, and in this case the quantity on the right-hand side is finite.

Lemma 2.2 (fractional Poincaré–Sobolev inequality). *Let $v \in W^{\alpha,p}(B)$, for $\alpha \in (0, 1)$, where $B \subset \mathbb{R}^n$ is a ball of radius r , or a cube of diameter r . If $p\alpha < n$, then the following inequality holds for a constant c depending only on n, α :*

$$\left(\int_B |v - (v)_B|^{p^*} dx \right)^{1/p^*} \leq cr^{\alpha} \left(\int_B \int_B \frac{|v(x) - v(y)|^p}{|x - y|^{n+p\alpha}} dx dy \right)^{1/p},$$

where $p^* := np/(n - p\alpha)$.

With 2_* being the exponent defined in (1-13), an immediate consequence of the previous lemma is the following inequality, that we report since it will be used several times:

$$\left(\int_B |v - (v)_B|^2 dx \right)^{1/2} \leq cr^\alpha \left(\int_B \int_B \frac{|v(x) - v(y)|^{2_*}}{|x - y|^{n+2_*\alpha}} dx dy \right)^{1/2_*}. \quad (2-4)$$

Moreover, if v is compactly supported in B , then $v - (v)_B$ above can be replaced by v .

3. The Caccioppoli inequality

3A. Preliminary reformulation of the assumptions. We start with the assumptions made on g , that is, (1-17)–(1-18). In order to give a unified proof for the two cases $2\beta \geq \alpha$ and $2\beta < \alpha$, and to simplify certain computations, we shall make a few preliminary reductions and will restate the assumptions in a more convenient way. First of all let us consider the case $2\beta \geq \alpha$, when (1-17) is in force. Let us notice that, eventually reducing the value of δ_0 , and in particular taking $\delta_0 \leq \alpha/40$, (1-17) implies the existence of exponents p , γ and $\delta_1 > 0$, such that $g \in W^{\gamma(1+\delta_1), p(1+\delta_1)}(\mathbb{R}^n)$ and

$$2\beta > \gamma > 2\beta - \alpha, \quad 2 > p > \frac{2n}{n + 2(\gamma - 2\beta + \alpha)}, \quad \delta_1 \leq \frac{\alpha}{4n}. \quad (3-1)$$

Indeed, let us set $\gamma = 2\beta - \alpha + \delta_0/2$ and recall that $W^{2\beta - \alpha + \delta_0, 2}$ embeds in $W^{\gamma(1+\delta_1), p(1+\delta_1)}$ whenever $2\beta - \alpha + \delta_0 - n/2 = \gamma(1 + \delta_1) - n/[p(1 + \delta_1)]$. A lengthy computation then shows that any choice of p as above and $\delta_1 \leq 1$ satisfying the inequalities

$$\frac{(1 + \delta_1)\delta_0}{n + 2\gamma(1 + \delta_1)} < \delta_1 < \frac{(2 + \delta_1)\delta_0}{n + 2\gamma(1 + \delta_1)}$$

matches the conditions in (3-1). We now consider the case $2\beta < \alpha$, when (1-18) is in force. In this case we can instead assume the existence of numbers $p > 1$ and $\delta_1 > 0$ such that

$$g \in L_{\text{loc}}^{p(1+\delta_1)}(\mathbb{R}^n), \quad p > \frac{2n}{n + 2(\alpha - 2\beta)}. \quad (3-2)$$

Let us now unify the previous conditions. In the case $2\beta \geq \alpha$ we clearly have that

$$\int_B \int_B \frac{|g(x) - g(y)|^{p(1+\delta_1)}}{|x - y|^{n+p(1+\delta_1)^2\gamma}} dx dy + \int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy < \infty \quad (3-3)$$

for every ball $B \subset \mathbb{R}^n$. This comes by the definition of the space $W^{\gamma(1+\delta_1), p(1+\delta_1)}$. On the other hand, when $2\beta < \alpha$, then assumptions (1-18) do not involve any number γ . Thanks to the lower bound on p in (1-18), we can find a negative number γ , such that $|\gamma| \in (0, \frac{1}{10})$ is small enough to still verify (3-1). In this case we note that

$$\begin{aligned} \int_B \int_B \frac{|g(x) - g(y)|^{p(1+\delta_1)}}{|x - y|^{n+p(1+\delta_1)^2\gamma}} dx dy &\leq \int_B \int_B \frac{(|g(x)| + |g(y)|)^{p(1+\delta_1)}}{|x - y|^{n+p(1+\delta_1)^2\gamma}} dx dy \\ &\leq \frac{cr^{-p(1+\delta_1)^2\gamma}}{-\gamma} \int_B |g|^{p(1+\delta_1)} dx < \infty, \end{aligned} \quad (3-4)$$

where r denotes the radius of B ; a similar estimate follows for the second quantity in (3-3). Summarizing, in the rest of the paper we shall always assume that (3-1) and (3-3) hold. In the case $2\beta < \alpha$ the number γ is negative.

Remark 3.1. We shall denote by c_b a constant that depends on $n, \alpha, \Lambda, p, \beta, \gamma$ and exhibits the blow-up behavior

$$\lim_{p \rightarrow 2n/(n+2(\gamma-2\beta+\alpha))} c_b = \infty. \quad (3-5)$$

3B. The Caccioppoli estimate. The Caccioppoli-type inequality stated in the next theorem is an essential tool in the proof of Theorem 1.1.

Theorem 3.2. Let $u \in W^{\alpha,2}(\mathbb{R}^n)$ be a solution to (1-14) under the assumptions of Theorem 1.1; in particular, (3-1) and (3-3) are in force. Let $B \equiv B(x_0, r) \subset \mathbb{R}^n$ be a ball, and let $\psi \in C_c^\infty(B(x_0, \frac{3}{4}r))$ be a cutoff function such that $0 \leq \psi \leq 1$ and $|D\psi| \leq c(n)/r$. Then the Caccioppoli-type inequality

$$\begin{aligned} & \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy \\ & \leq \frac{c}{r^{2\alpha}} \int_B |u(x)|^2 dx + c \int_{\mathbb{R}^n \setminus B} \frac{|u(y)|}{|x_0-y|^{n+2\alpha}} dy \int_B |u(x)| dx + cr^{n+2\alpha} \left(\int_B |f(x)|^{2^*} dx \right)^{2/2^*} \\ & \quad + c_b r^{n+2(\gamma-2\beta+\alpha)} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x)-g(y)|^p}{|x-y|^{n+p\gamma}} dx dy \right)^{1/p} \right]^2 \end{aligned} \quad (3-6)$$

holds for a constant $c \equiv c(n, \Lambda, \alpha)$ which is independent of p , and a constant $c_b \equiv c_b(n, \Lambda, \alpha, \beta, \gamma, p)$. The constant c_b exhibits the behavior described in (3-5); moreover, all the terms appearing on the right-hand side of (3-6) are finite.

Proof. In the weak formulation

$$\begin{aligned} & \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \varphi(u(x) - u(y))[\eta(x) - \eta(y)]K(x, y) dx dy \\ & = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} [g(x) - g(y)][\eta(x) - \eta(y)]H(x, y) dx dy + \int_{\mathbb{R}^n} f \eta dx, \end{aligned} \quad (3-7)$$

we choose $\eta = u\psi^2$, where $\psi \in C_c^\infty(B)$ is the cutoff function coming from the statement. By a density argument, η is an admissible test function. Then we have

$$\begin{aligned} I_1 + I_2 + I_3 & := \int_B \int_B \varphi(u(x) - u(y))[u(x)\psi^2(x) - u(y)\psi^2(y)]K(x, y) dx dy \\ & \quad + \int_{\mathbb{R}^n \setminus B} \int_B \varphi(u(x) - u(y))u(x)\psi^2(x)K(x, y) dx dy \\ & \quad - \int_B \int_{\mathbb{R}^n \setminus B} \varphi(u(x) - u(y))u(y)\psi^2(y)K(x, y) dx dy \end{aligned}$$

$$\begin{aligned}
&= \int_B \int_B [g(x) - g(y)][u(x)\psi^2(x) - u(y)\psi^2(y)]H(x, y) dx dy \\
&\quad + \int_{\mathbb{R}^n \setminus B} \int_B [g(x) - g(y)]u(x)\psi^2(x)H(x, y) dx dy \\
&\quad + \int_B \int_{\mathbb{R}^n \setminus B} [g(y) - g(x)]u(y)\psi^2(y)H(x, y) dx dy \\
&\quad + \int_B f(x)u(x)\psi^2(x) dx =: J_1 + J_2 + J_3 + J_4. \tag{3-8}
\end{aligned}$$

We proceed in estimating the various pieces stemming from this identity.

Estimation of I_1 . Let us first consider the case in which $\psi(x) \geq \psi(y)$. Then we write

$$\begin{aligned}
&\varphi(u(x) - u(y))[u(x)\psi^2(x) - u(y)\psi^2(y)] \\
&\quad = \varphi(u(x) - u(y))[u(x) - u(y)]\psi^2(x) + \varphi(u(x) - u(y))u(y)[\psi^2(x) - \psi^2(y)]. \tag{3-9}
\end{aligned}$$

Applying Young's inequality and recalling the first inequality in (1-15), we have

$$\begin{aligned}
\varphi(u(x) - u(y))u(y)[\psi^2(x) - \psi^2(y)] &= \varphi(u(x) - u(y))u(y)[\psi(x) - \psi(y)][\psi(x) + \psi(y)] \\
&\geq -2|\varphi(u(x) - u(y))||u(y)||\psi(x) - \psi(y)|\psi(x) \\
&\geq -\frac{1}{2}|u(x) - u(y)|^2\psi^2(x) - 2\Lambda^2u^2(y)[\psi(x) - \psi(y)]^2.
\end{aligned}$$

Combining the content of the last two displays, and using this time the second inequality in (1-15), yields

$$\varphi(u(x) - u(y))[u(x)\psi^2(x) - u(y)\psi^2(y)] \geq \frac{1}{2}[u(x) - u(y)]^2\psi^2(x) - 2\Lambda^2u^2(y)[\psi(x) - \psi(y)]^2.$$

Now we consider the case in which $\psi(y) \geq \psi(x)$, and we similarly write

$$\begin{aligned}
&\varphi(u(x) - u(y))[u(x)\psi^2(x) - u(y)\psi^2(y)] \\
&\quad = \varphi(u(x) - u(y))[u(x) - u(y)]\psi^2(y) + \varphi(u(x) - u(y))u(x)[\psi^2(x) - \psi^2(y)].
\end{aligned}$$

Proceeding similarly to the case $\psi(x) \geq \psi(y)$, we arrive at

$$\varphi(u(x) - u(y))[u(x)\psi^2(x) - u(y)\psi^2(y)] \geq \frac{1}{2}[u(x) - u(y)]^2\psi^2(y) - 2\Lambda^2u^2(x)[\psi(x) - \psi(y)]^2.$$

In any case, using also (1-4), we conclude that

$$I_1 \geq \frac{1}{c} \int_B \int_B \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} \max\{\psi^2(x), \psi^2(y)\} dx dy - c \int_B \int_B |u(x)|^2 \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy,$$

where c depends on Λ . Moreover, by noticing that

$$[u(x)\psi(x) - u(y)\psi(y)]^2 \leq 2[u(x)(\psi(x) - \psi(y))]^2 + 2[\psi(y)(u(x) - u(y))]^2$$

and integrating, we conclude that

$$I_1 \geq \frac{1}{c} \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy - c \int_B \int_B |u(x)|^2 \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy. \tag{3-10}$$

Estimation of I_2 and I_3 . The estimation of the terms I_2 and I_3 is similar. Indeed, as for I_2 , we start by observing that a direct computation yields

$$[u(x) - u(y)]u(x)\psi^2(x)K(x, y) \geq -\Lambda \frac{|u(x)||u(y)|\psi^2(x)}{|x - y|^{n+2\alpha}},$$

and therefore, by (1-15) we obtain (we can assume without loss of generality that $u(x) \neq u(y)$) that

$$\begin{aligned} \varphi(u(x) - u(y))u(x)\psi^2(x)K(x, y) &\geq -\Lambda \left| \frac{\varphi(u(x) - u(y))}{u(x) - u(y)} \right| \cdot \frac{|u(x)||u(y)|\psi^2(x)}{|x - y|^{n+2\alpha}} \\ &\geq -\Lambda^2 \frac{|u(x)||u(y)|\psi^2(x)}{|x - y|^{n+2\alpha}}. \end{aligned}$$

Similarly, we obtain

$$-\varphi(u(x) - u(y))u(y)\psi^2(y)K(x, y) \geq -\Lambda^2 \frac{|u(x)||u(y)|\psi^2(y)}{|x - y|^{n+2\alpha}}.$$

We then estimate

$$\begin{aligned} I_2 + I_3 &\geq -c \int_{\mathbb{R}^n \setminus B} \int_B \frac{|u(x)||u(y)|\psi^2(x)}{|x - y|^{n+2\alpha}} dx dy \\ &\geq -c \sup_{z \in \text{supp } \psi} \int_{\mathbb{R}^n \setminus B} \frac{|u(y)|}{|z - y|^{n+2\alpha}} dy \int_B |u(x)|\psi^2(x) dx \\ &\geq -c \int_{\mathbb{R}^n \setminus B} \frac{|u(y)|}{|x_0 - y|^{n+2\alpha}} dy \int_B |u(x)|\psi^2(x) dx. \end{aligned} \quad (3-11)$$

Here we have used the fact that, since ψ is supported in $B(x_0, \frac{3}{4}r)$, we have

$$\frac{|x_0 - y|}{|z - y|} \leq 1 + \frac{|x_0 - z|}{|z - y|} \leq 4 \quad (3-12)$$

whenever $z \in \text{supp } \psi$ and $y \in \mathbb{R}^n \setminus B$.

Estimation of J_4 . The fractional Sobolev inequality yields

$$\begin{aligned} J_4 &\leq cr^n \left(\int_B |u(x)\psi(x)|^{2^*} dx \right)^{1/2^*} \left(\int_B |f(x)|^{2^*} dx \right)^{1/2^*} \\ &\leq cr^{n/2+\alpha} \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy \right)^{1/2} \left(\int_B |f(x)|^{2^*} dx \right)^{1/2^*}, \end{aligned}$$

so that, applying Young's inequality with $\sigma \in (0, 1)$, we have

$$J_4 \leq \frac{c}{\sigma} r^{n+2\alpha} \left(\int_B |f(x)|^{2^*} dx \right)^{2/2^*} + \sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy. \quad (3-13)$$

The constant c depends only on n, α .

Estimation of J_1 . We write

$$u(x)\psi^2(x) - u(y)\psi^2(y) = [u(x)\psi(x) - u(y)\psi(y)]\psi(y) + u(x)\psi(x)[\psi(x) - \psi(y)].$$

Therefore, using that $\psi \leq 1$ together with (1-16), we have

$$\begin{aligned} J_1 &\leq \Lambda \int_B \int_B \frac{|g(x) - g(y)|}{|x - y|^{2\beta}} |u(x)\psi(x) - u(y)\psi(y)| \frac{dx dy}{|x - y|^n} \\ &\quad + \Lambda \int_B \int_B \frac{|g(x) - g(y)|}{|x - y|^{2\beta}} |u(x)\psi(x)| |\psi(x) - \psi(y)| \frac{dx dy}{|x - y|^n} \\ &=: J_{1,1} + J_{1,2}. \end{aligned}$$

In turn, we estimate $J_{1,1}$ and $J_{1,2}$ separately. Recalling (3-1), we now set

$$t := 1 - \frac{2\beta - \gamma}{\alpha} \quad \text{and} \quad s := \frac{n}{\alpha} \left[\frac{1}{p} - \frac{1}{2} \right]. \quad (3-14)$$

Observe that $0 < t \leq 1$ if and only if $2\beta - \alpha < \gamma \leq 2\beta$. Then we notice that

$$\begin{aligned} 2\beta \geq \gamma \quad \text{and} \quad 2 > p > \frac{2n}{n + 2(\gamma - 2\beta + \alpha)} &\implies 2 > p > \frac{2n}{n + 2\alpha} = 2_* \\ &\implies 0 < s < 1, \end{aligned} \quad (3-15)$$

and moreover

$$p > \frac{2n}{n + 2(\gamma - 2\beta + \alpha)} \implies 0 < s < t. \quad (3-16)$$

We also record the identity $\alpha t = \gamma - (2\beta - \alpha)$. Let us now write

$$J_{1,1} = cr^n \int_B \int_B \left[r^{\alpha t} \frac{|g(x) - g(y)|}{|x - y|^{2\beta - \alpha + t\alpha}} \right] \left[\frac{|u(x)\psi(x) - u(y)\psi(y)|}{|x - y|^\alpha} \right]^{1-s} \left[r^{-\alpha t/s} \frac{|u(x)\psi(x) - u(y)\psi(y)|}{|x - y|^{\alpha(1-t/s)}} \right]^s$$

The definitions in (3-14) imply that

$$\frac{1-s}{2} + \frac{s}{2_*} + \frac{1}{p} = 1;$$

therefore, applying Hölder's inequality with the corresponding choice of the exponents, we have

$$\begin{aligned} J_{1,1} &\leq cr^{n+\alpha t} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n+p(2\beta - \alpha + t\alpha)}} dx dy \right)^{1/p} \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy \right)^{(1-s)/2} \\ &\quad \times \left(r^{-2_*\alpha t/s} \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^{2_*}}{|x - y|^{n+2_*\alpha(1-t/s)}} dx dy \right)^{s/2_*}. \end{aligned} \quad (3-17)$$

Before going on, let us estimate the last integral:

$$\begin{aligned}
\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^{2^*}}{|x-y|^{n+2^*\alpha(1-t/s)}} dx dy &\leq 2^{2^*-1} \int_B \int_B \frac{|u(x)\psi(x)|^{2^*}}{|x-y|^{n+2^*\alpha(1-t/s)}} dx dy \\
&\leq \frac{cr^{-2^*\alpha(1-t/s)}}{t-s} \int_B |u(x)\psi(x)|^{2^*} dx \\
&\leq \frac{cr^{2^*\alpha t/s}}{t-s} \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy \right)^{2^*/2}. \quad (3-18)
\end{aligned}$$

Plugging the inequality into (3-17) yields

$$J_{1.1} \leq cr^{n/2+\alpha t} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x-y|^{n+p(2\beta-\alpha+t\alpha)}} dx dy \right)^{1/p} \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy \right)^{1/2}.$$

Using Young's inequality, and keeping in mind that $\alpha t = \gamma - (2\beta - \alpha)$, leads to

$$J_{1.1} \leq \frac{c}{\sigma} r^{n+2(\gamma-2\beta+\alpha)} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x-y|^{n+p\gamma}} dx dy \right)^{2/p} + \sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy$$

whenever $\sigma \in (0, 1)$. The constant c depends only on $n, \alpha, \Lambda, \beta, \gamma, p$. We then continue with the estimation of $J_{1.2}$. Upon setting $\eta := \frac{1}{2}(1 - \alpha)$, using Hölder's inequality with conjugate exponents $(2^*, 2_*)$ we have

$$\begin{aligned}
J_{1.2} &\leq c \|D\psi\|_{L^\infty} r^n \int_B \int_B \frac{|g(x) - g(y)|}{|x-y|^{2\beta-1+\eta}} \frac{|u(x)\psi(x)|}{|x-y|^{-\eta}} \frac{dx dy}{|x-y|^n} \\
&\leq c \|D\psi\|_{L^\infty} r^n \left(\int_B \int_B \frac{|g(x) - g(y)|^{2^*}}{|x-y|^{2_*(2\beta-1+\eta)}} \frac{dx dy}{|x-y|^n} \right)^{1/2_*} \left(\int_B \int_B \frac{|u(x)\psi(x)|^{2^*}}{|x-y|^{-2^*\eta}} \frac{dx dy}{|x-y|^n} \right)^{1/2^*}.
\end{aligned}$$

In turn, by Lemma 2.2 (see also the remark following it) we have

$$\begin{aligned}
\int_B \int_B \frac{|u(x)\psi(x)|^{2^*}}{|x-y|^{-2^*\eta}} \frac{dx dy}{|x-y|^n} &\leq \frac{cr^{2^*\eta}}{1-\alpha} \int_B |u(x)\psi(x)|^{2^*} dx \\
&\leq cr^{2^*(\eta+\alpha)} \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy \right)^{2^*/2}
\end{aligned}$$

and, recalling that $p > 2_*$ by (3-15), we proceed with

$$\begin{aligned}
\int_B \int_B \frac{|g(x) - g(y)|^{2^*}}{|x-y|^{2_*(2\beta-1+\eta)}} \frac{dx dy}{|x-y|^n} &= \int_B \int_B \left(\frac{|g(x) - g(y)|}{|x-y|^\gamma} \right)^{2^*} \frac{1}{|x-y|^{2_*(2\beta-1+\eta-\gamma)}} \frac{dx dy}{|x-y|^n} \\
&\leq \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x-y|^{n+p\gamma}} dx dy \right)^{2_*/p} \\
&\quad \times \left(\int_B \int_B \frac{1}{|x-y|^{\frac{2_*(2\beta-1+\eta-\gamma)p}{p-2_*}}} \frac{dx dy}{|x-y|^n} \right)^{1-2_*/p} \\
&\leq \frac{cr^{-2_*(2\beta-1+\eta-\gamma)}}{\gamma - 2\beta + \alpha} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x-y|^{n+p\gamma}} dx dy \right)^{2_*/p},
\end{aligned}$$

where of course we used that $2\beta - 1 + \eta - \gamma = 2\beta - \frac{1}{2} - \frac{1}{2}\alpha - \gamma < 2\beta - \alpha - \gamma < 0$ due to $\eta := \frac{1}{2}(1 - \alpha)$ and (3-1). Connecting the estimates in the last three displays yields

$$J_{1.2} \leq c \|D\psi\|_{L^\infty} r^{n/2 + \gamma - 2\beta + \alpha + 1} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n + p\gamma}} dx dy \right)^{1/p} \\ \times \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n + 2\alpha}} dx dy \right)^{1/2}.$$

Again using Young's inequality, we conclude that

$$J_{1.2} \leq \frac{c}{\sigma} r^2 \|D\psi\|_{L^\infty}^2 r^{n + 2(\gamma - 2\beta + \alpha)} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n + p\gamma}} dx dy \right)^{2/p} \\ + \sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n + 2\alpha}} dx dy,$$

which holds whenever $\sigma \in (0, 1)$. Gathering together the estimates found for $J_{1.1}$ and $J_{1.2}$, and using that $r^2 \|D\psi\|_{L^\infty}^2 \leq c(n)$, gives

$$J_1 \leq \frac{c}{\sigma} r^{n + 2(\gamma - 2\beta + \alpha)} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n + p\gamma}} dx dy \right)^{2/p} \\ + 2\sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n + 2\alpha}} dx dy. \quad (3-19)$$

The constant c depends on $n, \alpha, \Lambda, \beta, \gamma, p$.

Estimation of J_2 and J_3 . The estimation of the two terms is completely similar, and we therefore confine ourselves to estimating J_2 . Using (1-16) we have

$$J_2 \leq \Lambda \int_{\mathbb{R}^n \setminus B} \int_B \frac{|g(x) - (g)_B|}{|x - y|^{n + 2\beta}} |u(x)| \psi^2(x) dx dy + \Lambda \int_{\mathbb{R}^n \setminus B} \int_B \frac{|g(y) - (g)_B|}{|x - y|^{n + 2\beta}} |u(x)| \psi^2(x) dx dy \\ =: J_{2.1} + J_{2.2}.$$

In turn we estimate the two resulting terms. Using that $p \geq 2_*$ by (3-15), we have

$$J_{2.1} \leq c \sup_{z \in \text{supp } \psi} \int_{\mathbb{R}^n \setminus B} \frac{dy}{|z - y|^{n + 2\beta}} \int_B |g(x) - (g)_B| |u(x)| \psi(x) dx \\ \leq cr^n \sup_{z \in \text{supp } \psi} \int_{\mathbb{R}^n \setminus B} \frac{dy}{|z - y|^{n + 2\beta}} \left(\int_B |g(x) - (g)_B|^{2_*} dx \right)^{1/2_*} \left(\int_B |u(x)\psi(x)|^{2_*} dx \right)^{1/2_*} \\ \leq cr^n \sup_{z \in \text{supp } \psi} \int_{\mathbb{R}^n \setminus B} \frac{dy}{|z - y|^{n + 2\beta}} \left(\int_B |g(x) - (g)_B|^p dx \right)^{1/p} \left(\int_B |u(x)\psi(x)|^{2_*} dx \right)^{1/2_*} \\ \leq cr^{n/2 + \gamma - 2\beta + \alpha} \sup_{z \in \text{supp } \psi} \int_{\mathbb{R}^n \setminus B} \frac{r^{2\beta} dy}{|z - y|^{n + 2\beta}} \cdot \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n + p\gamma}} dx dy \right)^{1/p} \\ \times \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n + 2\alpha}} dx dy \right)^{1/2}.$$

Therefore, using Young's inequality, we have

$$J_{2.1} \leq \frac{c}{\sigma} r^{n+2(\gamma-2\beta+\alpha)} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{2/p} + \sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy,$$

where we have also used that $\psi \equiv 0$ outside $B(x_0, \frac{3}{4}r)$, and therefore (3-12), to estimate

$$\sup_{z \in \text{supp } \psi} \int_{\mathbb{R}^n \setminus B} \frac{r^{2\beta} dy}{|z - y|^{n+2\beta}} \leq c(n, \beta).$$

In order to estimate $J_{2.2}$ we need another splitting over annuli. Recalling again that $\psi \leq 1$ and that $\psi \equiv 0$ outside $B(x_0, \frac{3}{4}r)$, we have

$$\begin{aligned} J_{2.2} &\leq c \sum_{j=0}^{\infty} \int_{2^{j+1}B \setminus 2^jB} \int_B \frac{|g(y) - (g)_B|}{|x - y|^{n+2\beta}} |u(x)| \psi^2(x) dx dy \\ &\leq cr^n \sum_{j=0}^{\infty} (2^j r)^{-2\beta} \int_{2^{j+1}B} |g(y) - (g)_B| dy \int_B |u(x)\psi(x)| dx \\ &\leq cr^n \sum_{j=0}^{\infty} (2^j r)^{-2\beta} \left(\int_{2^{j+1}B} |g(y) - (g)_B|^p dy \right)^{1/p} \int_B |u(x)\psi(x)| dx. \end{aligned} \quad (3-20)$$

The estimation of $J_{2.2}$ needs again a splitting; we start with the telescoping summation

$$\begin{aligned} &\left(\int_{2^{j+1}B} |g(y) - (g)_B|^p dy \right)^{1/p} \\ &\leq \left(\int_{2^{j+1}B} |g(y) - (g)_{2^{j+1}B}|^p dy \right)^{1/p} + \sum_{k=0}^j |(g)_{2^{k+1}B} - (g)_{2^k B}| \\ &\leq \left(\int_{2^{j+1}B} |g(y) - (g)_{2^{j+1}B}|^p dy \right)^{1/p} + \sum_{k=0}^j \left(\int_{2^{k+1}B} |g(y) - (g)_{2^k B}|^p dy \right)^{1/p} \\ &\leq 2 \sum_{k=0}^{j+1} \left(\int_{2^k B} |g(y) - (g)_{2^k B}|^p dy \right)^{1/p}. \end{aligned} \quad (3-21)$$

Then an application of the fractional Poincaré inequality in Lemma 2.1 yields

$$\left(\int_{2^{j+1}B} |g(y) - (g)_B|^p dy \right)^{1/p} \leq c \sum_{k=0}^{j+1} (2^k r)^\gamma \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p}.$$

Merging the content of the last display with the one of (3-20) gives

$$J_{2.2} \leq cr^n \left[\sum_{j=0}^{\infty} \sum_{k=0}^{j+1} (2^j r)^{-2\beta} (2^k r)^\gamma \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \right] \int_B |u(x)\psi(x)| dx.$$

We now manipulate the content of the square brackets above using discrete Fubini's theorem:

$$\begin{aligned}
& \sum_{j=0}^{\infty} \sum_{k=0}^{j+1} (2^j r)^{-2\beta} (2^k r)^\gamma \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \\
&= r^{\gamma-2\beta} \left(\int_B \int_B \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \sum_{j=0}^{\infty} 2^{-2\beta j} \\
&\quad + r^{\gamma-2\beta} \sum_{k=1}^{\infty} 2^{\gamma k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \sum_{j=k-1}^{\infty} 2^{-2\beta j} \\
&\leq \frac{c r^{\gamma-2\beta}}{\beta} \sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p}.
\end{aligned}$$

We remark that in the previous display we have used the elementary inequality in (2-2). All in all we have, by using also Hölder's inequality and Lemma 2.1, that

$$\begin{aligned}
J_{2.2} &\leq c r^{n+\gamma-2\beta} \sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \left(\int_B |u(x)\psi(x)|^{2^*} dx \right)^{1/2^*} \\
&\leq c r^{n/2+\gamma-2\beta+\alpha} \sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \\
&\quad \times \left(\int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy \right)^{1/2}.
\end{aligned}$$

Finally, using Young's inequality we conclude that

$$\begin{aligned}
J_{2.2} &\leq \frac{c}{\sigma} r^{n+2(\gamma-2\beta+\alpha)} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \right]^2 \\
&\quad + \sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy
\end{aligned}$$

whenever $\sigma \in (0, 1)$. Connecting the inequalities found for $J_{1.2}$ and $J_{2.2}$, and again recalling that J_3 can be estimated in a completely similar way, we have

$$\begin{aligned}
J_2 + J_3 &\leq \frac{c}{\sigma} r^{n+2(\gamma-2\beta+\alpha)} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \right]^2 \\
&\quad + 4\sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy. \quad (3-22)
\end{aligned}$$

The constant c depends on $n, \Lambda, \alpha, \beta, \gamma, p$.

Reabsorbing terms. Inserting the estimates for the terms I_i and J_i into (3-8), we conclude that

$$\begin{aligned} & \frac{1}{c} \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy \\ & \leq 7\sigma \int_B \int_B \frac{|u(x)\psi(x) - u(y)\psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy + c \int_B \int_B |u(x)|^2 \frac{|\psi(x) - \psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy \\ & \quad + c \int_{\mathbb{R}^n \setminus B} \frac{|u(y)|}{|x_0 - y|^{n+2\alpha}} dy \int_B |u(x)|\psi^2(x) dx + \frac{c}{\sigma} r^{n+2\alpha} \left(\int_B |f(x)|^{2_*} dx \right)^{2/2_*} \\ & \quad + \frac{c_b}{\sigma} r^{n+2(\gamma-2\beta+\alpha)} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x-y|^{n+p\gamma}} dx dy \right)^{1/p} \right]^2. \end{aligned}$$

The constant c depends only on n, α, Λ , and the constant c_b depends only on $n, \Lambda, \alpha, \beta, \gamma, p$. Now, taking $\sigma = 1/(14c)$ and reabsorbing terms finishes the proof, together with the estimate

$$\begin{aligned} \int_B \int_B |u(x)|^2 \frac{|\psi(x) - \psi(y)|^2}{|x-y|^{n+2\alpha}} dx dy & \leq \|D\psi\|_{\infty} \int_B |u(x)|^2 \int_{B_{2r}(x)} |x-y|^{-n+2-2\alpha} dy dx \\ & = \frac{c(n)}{1-\alpha} \|D\psi\|_{\infty} r^{2-2\alpha} \int_B |u(x)|^2 dx \\ & \leq \frac{c(n)}{1-\alpha} \frac{1}{r^{2\alpha}} \int_B |u(x)|^2 dx. \end{aligned}$$

The finiteness of the terms appearing on the right in (3-6) follows directly from the fact that $u \in W^{\alpha,2}(\mathbb{R}^n)$ and from Section 4C below. This completes the proof of Theorem 3.2. \square

Remark 3.3. In the statement of Theorem 3.2, one can replace u with $u - (u)_B$ by testing with $(u - (u)_B)\psi^2$ instead of $u\psi^2$.

Remark 3.4. All the constants denoted by c that appear in Theorem 3.2 blow up as $\alpha \rightarrow 0$ or as $\alpha \rightarrow 1$. The blow-up of the constant c_b is more peculiar, and it is as in (3-5). This appears for instance in estimate (3-18), as in this case $s \rightarrow t$; see (3-16). In terms of (1-12) the blow-up of c_b occurs for instance when $\delta_0 \rightarrow 0$. Moreover, the constant c_b blows up also when $\beta \rightarrow 0$ and $\gamma \rightarrow 2\beta - \alpha$.

4. The dual pair (μ, U) and reverse inequalities

4A. A doubling measure. With ε initially satisfying the condition $0 < \varepsilon < \frac{1}{2}\alpha$, we consider the locally finite measure μ on $\mathbb{R}^n \times \mathbb{R}^n$ introduced in (1-19). We summarize its basic properties:

Proposition 4.1. *With μ being defined as in (1-19):*

- Whenever $\mathcal{B} = B \times B$ and $B \subset \mathbb{R}^n$ is a ball with radius r , we have

$$\mu(\mathcal{B}) = \frac{c_{\varepsilon}(n)r^{n+2\varepsilon}}{\varepsilon}, \quad (4-1)$$

where $c_{\varepsilon}(n)$ denotes a constant depending only on n, ε , and it satisfies $1/c(n) \leq c_{\varepsilon}(n) \leq c(n)$ for another constant $c(n)$ depending only on n .

- (doubling diagonal property) Whenever $A \geq 1$, we have

$$\sup_{\tilde{x} \in \mathbb{R}^n, \varrho > 0} \frac{\mu(\mathcal{B}(\tilde{x}, A\varrho))}{\mu(\mathcal{B}(\tilde{x}, \varrho))} = A^{n+2\varepsilon}. \quad (4-2)$$

- For every $A \geq 1$, there exists a constant $c_d \equiv c_d(n, A)$ such that

$$\frac{\mu(\mathcal{B}(\tilde{x}, \varrho))}{\mu(K_1 \times K_2)} \leq \frac{c_d}{\varepsilon} \quad (4-3)$$

whenever $K_1, K_2 \subset \mathcal{B}(\tilde{x}, \varrho) \subset \mathbb{R}^n$ are cubes with sides parallel to the coordinate axes and such that $|K_1| = |K_2| = \varrho^n / A^n$.

- (standard doubling property) There exists a constant c , depending only on n , such that

$$\sup_{\tilde{x}, \tilde{y} \in \mathbb{R}^n, \varrho > 0} \frac{\mu(\mathcal{B}(\tilde{x}, \tilde{y}, 2\varrho))}{\mu(\mathcal{B}(\tilde{x}, \tilde{y}, \varrho))} \leq \frac{c}{\varepsilon}. \quad (4-4)$$

Proof. The proof of (4-1) follows directly from the definition in (1-19) and a scaling argument, while (4-2) follows from (4-1). The proof of (4-3) is slightly less direct. First, observe that $K_1 \times K_2 \subset \mathcal{B}(\tilde{x}, \varrho)$ and moreover that $|x - y| < 2\varrho$ whenever $x \in K_1$ and $y \in K_2$. Therefore we can estimate

$$\begin{aligned} \mu(\mathcal{B}(\tilde{x}, \varrho)) &= \frac{c(n)\varrho^{n+2\varepsilon}}{\varepsilon} \leq \frac{c(n)A^{2n}}{\varepsilon} \frac{1}{\varrho^{n-2\varepsilon}} \int_{K_1} \int_{K_2} dx dy \\ &\leq \frac{c(n)A^{2n}}{\varepsilon} \int_{K_1} \int_{K_2} \frac{dx dy}{|x - y|^{n-2\varepsilon}} = \frac{c(n)A^{2n}}{\varepsilon} \mu(K_1 \times K_2), \end{aligned}$$

and the proof of (4-3) is complete. The proof of (4-4) is similar to the one of (4-3); this estimate will not be used in the rest of the paper. \square

4B. Diagonal reverse Hölder-type inequalities. For $(x, y) \in \mathbb{R}^{2n}$, we define the functions

$$U(x, y) := \frac{|u(x) - u(y)|}{|x - y|^{\alpha+\varepsilon}}, \quad G(x, y) := \frac{|g(x) - g(y)|}{|x - y|^{\gamma+2\varepsilon/p}}, \quad F(x, y) := |f(x)|, \quad (4-5)$$

the first two being defined when $x \neq y$. According to Definition 1.2 the function u generates the dual pair (μ, U) . From now on, we shall always assume the following restriction on the number ε :

$$0 < \varepsilon < \min\{\frac{1}{2}\alpha, \frac{1}{4}|\gamma|(1 + \delta_1), \frac{1}{4}(2\beta - \gamma)p\}. \quad (4-6)$$

Lemma 4.2. *With the definitions in (4-5), it follows that*

$$U \in L^2(\mathbb{R}^{2n}; \mu) \quad \text{and} \quad F \in L_{\text{loc}}^{2_*+\delta_f}(\mathbb{R}^{2n}; \mu), \quad \text{with } \delta_f \in [0, \delta_0]. \quad (4-7)$$

Moreover, assuming (4-6) it follows that

$$G \in L_{\text{loc}}^{p+\delta_g}(\mathbb{R}^{2n}; \mu), \quad \text{where } \delta_g \in [0, p\delta_1]. \quad (4-8)$$

Proof. The first inclusion in (4-7) is a direct consequence of the definition in (4-5). As for F , for a ball $\mathcal{B} = B \times B$, where $B \subset \mathbb{R}^n$ has radius $r > 0$, we have

$$\int_{\mathcal{B}} F^{2_*+\delta_0} d\mu = \int_B \int_B \frac{|f(x)|^{2_*+\delta_0}}{|x-y|^{n-2\varepsilon}} dx dy \leq \frac{cr^{2\varepsilon}}{\varepsilon} \int_B |f|^{2_*+\delta_0} dx.$$

This clearly implies that $F \in L_{\text{loc}}^{2_*+\delta_f}(\mathbb{R}^{2n}; \mu)$ as long as $\delta_f \leq \delta_0$. To prove that $G \in L_{\text{loc}}^{p+\delta_g}(\mathbb{R}^{2n}; \mu)$, let us start with the case $2\beta \geq \alpha$, when $\gamma > 0$. By using (4-6) we have

$$\begin{aligned} \int_{\mathcal{B}} G^{p+p\delta_1} d\mu &= \int_B \int_B \frac{|g(x) - g(y)|^{p(1+\delta_1)}}{|x-y|^{n+\gamma p(1+\delta_1)+2\varepsilon\delta_1}} dx dy \\ &\leq cr^{\delta_1[\gamma p(1+\delta_1)-2\varepsilon]} \int_B \int_B \frac{|g(x) - g(y)|^{p(1+\delta_1)}}{|x-y|^{n+\gamma p(1+\delta_1)^2}} dx dy. \end{aligned} \quad (4-9)$$

The last quantity is finite since we are assuming $g \in W^{\gamma(1+\delta_1), p(1+\delta_1)}$, so it follows that $G \in L_{\text{loc}}^{p+\delta_g}(\mathbb{R}^{2n}; \mu)$. We finally treat the case $2\beta < \alpha$. In this case, we have $2\varepsilon\delta_1 < 2\varepsilon < |\gamma|p(1+\delta_1) = -\gamma p(1+\delta_1)$, so that $\gamma p(1+\delta_1) + 2\varepsilon\delta_1 < 0$. We can therefore estimate

$$\begin{aligned} \int_{\mathcal{B}} G^{p+p\delta_1} d\mu &\leq \int_B \int_B \frac{(|g(x)| + |g(y)|)^{p(1+\delta_1)}}{|x-y|^{n+\gamma p(1+\delta_1)+2\varepsilon\delta_1}} dx dy \\ &\leq \frac{cr^{-[\gamma p(1+\delta_1)+2\varepsilon\delta_1]}}{-[\gamma p(1+\delta_1) + 2\varepsilon\delta_1]} \int_B |g|^{p(1+\delta_1)} dx < \infty, \end{aligned} \quad (4-10)$$

and (4-8) follows again since when $2\beta < \alpha$ we are precisely assuming that $g \in L_{\text{loc}}^{p(1+\delta_1)}(\mathbb{R}^n)$; see (3-2). \square

We are now going to state a few inequalities for later use. Let $v \in W^{\tilde{\sigma}, q}(B)$ for $\tilde{\sigma} \in (0, 1)$ and $q \geq 1$; then the fractional Sobolev inequality

$$\int_B |v - (v)_B|^2 dx \leq cr^{2\tilde{\sigma}} \left(\int_B \int_B \frac{|v(x) - v(y)|^q}{|x-y|^{n+\tilde{\sigma}q}} dx dy \right)^{2/q} \quad (4-11)$$

holds as a consequence of (2-4), provided $q \geq 2n/(n+2\tilde{\sigma})$ and $\tilde{\sigma} > 0$. With $\varepsilon \in (0, \frac{1}{2}\alpha)$ we study the compatibility of the conditions

$$\tilde{\sigma} := \alpha + \varepsilon - \frac{2\varepsilon}{q} \quad \text{and} \quad q \geq \frac{2n}{n+2\tilde{\sigma}} \quad (4-12)$$

in inequality (4-11); this gives $q \geq (2n+4\varepsilon)/(n+2\alpha+2\varepsilon)$. Recalling the definition of the function U in (4-5), and using (4-1), we gain

$$r^{2\tilde{\sigma}} \left(\int_B \int_B \frac{|u(x) - u(y)|^q}{|x-y|^{n+\tilde{\sigma}q}} dx dy \right)^{2/q} = \frac{c_\varepsilon(n)^{2/q} r^{2\alpha+2\varepsilon}}{\varepsilon^{2/q}} \left(\int_B U^q d\mu \right)^{2/q},$$

with $c_\varepsilon(n)$ defined in (4-1). We therefore have the following:

Lemma 4.3. *Let $\varepsilon \in (0, \frac{1}{2}\alpha)$, and let q be defined by*

$$q := \frac{2n+4\varepsilon}{n+2\alpha+2\varepsilon} < 2. \quad (4-13)$$

Then the inequality

$$\int_B |u - (u)_B|^2 dx \leq \frac{cr^{2(\alpha+\varepsilon)}}{\varepsilon^{2/q}} \left(\int_B U^q d\mu \right)^{2/q} \quad (4-14)$$

holds for a constant c depending only on n and α , whenever B is a ball with radius r and $\mathcal{B} = B \times B$. The same inequality continues to hold when the ball B is replaced by a cube Q with sides of length r , and consequently \mathcal{B} is replaced by $Q \times Q$.

We are now ready for the main result of this section:

Proposition 4.4 (diagonal reverse Hölder-type inequality). *Let $u \in W^{\alpha,2}(\mathbb{R}^n)$ be a solution to (1-14) under the assumptions of Theorem 1.1; in particular, (3-1) and (3-3) are in force. Assume that ε satisfies (4-6). Then the following reverse Hölder-type inequality with tail holds whenever $\mathcal{B} \subset \mathbb{R}^{2n}$ is a diagonal ball and $\sigma \in (0, 1)$:*

$$\begin{aligned} \left(\int_{\mathcal{B}} U^2 d\mu \right)^{1/2} &\leq \frac{c}{\sigma \varepsilon^{1/q-1/2}} \left(\int_{2\mathcal{B}} U^q d\mu \right)^{1/q} + \frac{\sigma}{\varepsilon^{1/q-1/2}} \sum_{k=1}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} \\ &+ \frac{c[\mu(\mathcal{B})]^\eta}{\varepsilon^{1/2^*-1/2}} \left(\int_{2\mathcal{B}} F^{2^*} d\mu \right)^{1/2^*} + \frac{c_b[\mu(\mathcal{B})]^\theta}{\varepsilon^{1/p-1/2}} \sum_{k=1}^{\infty} 2^{-k(2\beta-\gamma-2\varepsilon/p)} \left(\int_{2^k \mathcal{B}} G^p d\mu \right)^{1/p}, \end{aligned} \quad (4-15)$$

where θ and η denote the positive exponents

$$\theta := \frac{\gamma - 2\beta + \alpha + \varepsilon(2/p - 1)}{n + 2\varepsilon} \quad \text{and} \quad \eta := \frac{\alpha - \varepsilon}{n + 2\varepsilon}. \quad (4-16)$$

The constant c depends only on n, α, Λ , while the number $q \in (1, 2)$ has been defined in (4-13). The constant c_b depends on $n, \alpha, \Lambda, \beta, \gamma, p$ and exhibits the behavior described in (3-5). The infinite sums on the right side of (4-15) are finite.

Proof. In the rest of the proof all the constants depend at least on n, α, Λ . We write $\mathcal{B} \equiv B(x_0, r) \times B(x_0, r)$ and apply Theorem 3.2; we choose a cutoff function $\psi \in C_0^\infty(\frac{3}{4}\mathcal{B})$ such that $0 \leq \psi \leq 1$, $|D\psi| \leq c(n)/r$ and $\psi \equiv 1$ on $\frac{1}{2}\mathcal{B}$. Inequality (3-6) remains valid upon replacing u by $u - (u)_B$; see Remark 3.3. Indeed, notice that for such a function all the integrals on the right-hand side of (3-6) are finite. For this see Section 4C and (4-19) below. All in all, we have

$$\begin{aligned} I_4 &:= \int_B \int_B \frac{|[u(x) - (u)_B]\psi(x) - [u(y) - (u)_B]\psi(y)|^2}{|x - y|^{n+2\alpha}} dx dy \\ &\leq cr^{-2\alpha} \int_B |u(x) - (u)_B|^2 dx + c \int_{\mathbb{R}^n \setminus B} \frac{|u(y) - (u)_B|}{|x_0 - y|^{n+2\alpha}} dy \int_B |u(x) - (u)_B| dx \\ &\quad + cr^{2\alpha} \left(\int_B |f(x)|^{2^*} dx \right)^{2/2^*} \\ &\quad + c_b r^{2(\gamma-2\beta+\alpha)} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta)k} \left(\int_{2^k B} \int_{2^k B} \frac{|g(x) - g(y)|^p}{|x - y|^{n+p\gamma}} dx dy \right)^{1/p} \right]^2 \\ &=: J_5 + J_6 + J_7 + J_8. \end{aligned} \quad (4-17)$$

We start by rewriting I_4 as

$$I_4 = \frac{1}{|B|} \int_B \frac{|[u(x) - (u)_B]\psi(x) - [u(y) - (u)_B]\psi(y)|^2}{|x - y|^{2(\alpha+\varepsilon)}} d\mu(x, y)$$

so that, with the current choice of ψ , we have

$$\frac{r^{2\varepsilon}}{\varepsilon} \int_{B/2} U^2 d\mu \leq \frac{c(n)}{|B|} \int_{B/2} U^2 d\mu \leq cI_4.$$

We estimate J_5 with the aid of (4-14):

$$J_5 \leq \frac{cr^{2\varepsilon}}{\varepsilon^{2/q}} \left(\int_B U^q d\mu \right)^{2/q}.$$

To estimate J_6 we split the term in annuli, and proceed somewhat as in (3-21). As a matter of fact, we will prove that this term is finite; indeed, we have

$$\begin{aligned} \int_{\mathbb{R}^n \setminus B} \frac{|u(y) - (u)_B|}{|x_0 - y|^{n+2\alpha}} dy &= \sum_{j=0}^{\infty} \int_{2^{j+1}B \setminus 2^jB} \frac{|u(y) - (u)_B|}{|x_0 - y|^{n+2\alpha}} dy \\ &\leq c \sum_{j=0}^{\infty} (2^j r)^{-2\alpha} \int_{2^{j+1}B} |u(y) - (u)_B| dy. \end{aligned} \quad (4-18)$$

In turn, we again split every integral in the previous sum, similarly to (3-21), and using Hölder's inequality we estimate

$$\int_{2^{j+1}B} |u(y) - (u)_B| dy \leq 2 \sum_{k=0}^{j+1} \left(\int_{2^k B} |u(y) - (u)_{2^k B}|^q dy \right)^{1/q}.$$

Each of the previous integrals can be then estimated with the aid of the fractional Poincaré inequality of Lemma 2.1:

$$\begin{aligned} \int_{2^k B} |u(y) - (u)_{2^k B}|^q dy &\leq c(2^k r)^{q(\alpha+\varepsilon)-2\varepsilon} \int_{2^k B} \int_{2^k B} \frac{|u(x) - u(y)|^q}{|x - y|^{n+q\tilde{\sigma}}} dx dy \\ &= \frac{c(2^k r)^{q(\alpha+\varepsilon)}}{\varepsilon} \int_{2^k B} U^q d\mu, \end{aligned}$$

where $\tilde{\sigma}$ is as in (4-12) and c remains independent of ε . As a consequence, we obtain

$$\int_{2^{j+1}B} |u(y) - (u)_B| dy \leq \frac{c}{\varepsilon^{1/q}} \sum_{k=0}^{j+1} (2^k r)^{\alpha+\varepsilon} \left(\int_{2^k B} U^q d\mu \right)^{1/q}.$$

Connecting the content of the last display to that of (4-18) yields

$$\int_{\mathbb{R}^n \setminus B} \frac{|u(y) - (u)_B|}{|x_0 - y|^{n+2\alpha}} dy \leq \frac{cr^{-\alpha+\varepsilon}}{\varepsilon^{1/q}} \sum_{j=0}^{\infty} \sum_{k=0}^{j+1} 2^{-2\alpha j} 2^{k(\alpha+\varepsilon)} \left(\int_{2^k B} U^q d\mu \right)^{1/q}.$$

Reversing the order of summation gives

$$\begin{aligned}
& \sum_{j=0}^{\infty} \sum_{k=0}^{j+1} 2^{-2\alpha j} 2^{k(\alpha+\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} \\
&= \left(\int_{\mathcal{B}} U^q d\mu \right)^{1/q} \sum_{j=0}^{\infty} 2^{-2\alpha j} + \sum_{k=1}^{\infty} 2^{k(\alpha+\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} \sum_{j=k-1}^{\infty} 2^{-2\alpha j} \\
&\leq \frac{c}{\alpha} \sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q}.
\end{aligned}$$

Observe that we have once again used the elementary inequality in (2-2) (with $\beta = \alpha$). All in all, combining the content of the last two displays yields

$$\int_{\mathbb{R}^n \setminus \mathcal{B}} \frac{|u(y) - (u)_{\mathcal{B}}|}{|x_0 - y|^{n+2\alpha}} dy \leq \frac{cr^{-\alpha+\varepsilon}}{\varepsilon^{1/q}} \sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q}, \quad (4-19)$$

so that, via another application of (4-14), we have

$$J_6 \leq \frac{cr^{2\varepsilon}}{\varepsilon^{2/q}} \sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} \left(\int_{\mathcal{B}} U^q d\mu \right)^{1/q}.$$

With $\sigma \in (0, 1)$, using Young's inequality we finally conclude that

$$J_6 \leq \frac{cr^{2\varepsilon}}{\sigma^2 \varepsilon^{2/q}} \left(\int_{\mathcal{B}} U^q d\mu \right)^{2/q} + \frac{\sigma^2 r^{2\varepsilon}}{\varepsilon^{2/q}} \left[\sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} \right]^2.$$

For the estimation of J_7 we observe that

$$\begin{aligned}
\int_{\mathcal{B}} |f(x)|^{2^*} dx &= \int_{\mathcal{B}} \int_{\mathcal{B}} |f(x)|^{2^*} dx dy \\
&\leq \frac{c}{r^{n-2\varepsilon+2\varepsilon}} \int_{\mathcal{B}} \int_{\mathcal{B}} |f(x)|^{2^*} dx dy \\
&\leq \frac{c}{r^{n+2\varepsilon}} \int_{\mathcal{B}} \int_{\mathcal{B}} \frac{|f(x)|^{2^*}}{|x-y|^{n-2\varepsilon}} dx dy \\
&\leq \frac{c}{\varepsilon} \int_{\mathcal{B}} F^{2^*} d\mu.
\end{aligned}$$

Here we have used (4-1) to perform the last estimation and the very definition of the measure μ . By the definition of J_7 it then follows that

$$J_7 \leq \frac{cr^{2\alpha}}{\varepsilon^{2/2^*}} \left(\int_{\mathcal{B}} F^{2^*} d\mu \right)^{2/2^*}.$$

Next, the definitions of $G(\cdot)$ and μ imply

$$J_8 \leq \frac{cr^{2(\gamma-2\beta+\alpha+2\varepsilon/p)}}{\varepsilon^{2/p}} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta+2\varepsilon/p)k} \left(\int_{2^k \mathcal{B}} G^p d\mu \right)^{1/p} \right]^2.$$

Finally, connecting the estimates found for I_4 and J_5, \dots, J_8 to (4-17) yields

$$\begin{aligned} \frac{r^{2\varepsilon}}{\varepsilon} \int_{\mathcal{B}/2} U^2 d\mu &\leq \frac{cr^{2\varepsilon}}{\sigma^2 \varepsilon^{2/q}} \left(\int_{\mathcal{B}} U^q d\mu \right)^{2/q} + \frac{\sigma^2 r^{2\varepsilon}}{\varepsilon^{2/q}} \left[\sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} \right]^2 \\ &\quad + \frac{cr^{2\alpha}}{\varepsilon^{2/2_*}} \left(\int_{\mathcal{B}} F^{2_*} d\mu \right)^{2/2_*} + \frac{cr^{2(\gamma-2\beta+\alpha+2\varepsilon/p)}}{\varepsilon^{2/p}} \left[\sum_{k=0}^{\infty} 2^{(\gamma-2\beta+2\varepsilon/p)k} \left(\int_{2^k \mathcal{B}} G^p d\mu \right)^{1/p} \right]^2, \end{aligned}$$

from which (4-15) follows immediately (since the ball \mathcal{B} is arbitrary, and we can switch from \mathcal{B} to $2\mathcal{B}$). The right-hand side terms in (4-15) involving infinite sums are finite; we check this in the next remark. \square

Remark 4.5. A computation based on the definitions in (4-16) gives

$$\frac{2_* \eta}{1 - 2_* \eta} = \frac{2n(\alpha - \varepsilon)}{n^2 + 4\varepsilon n + 4\alpha\varepsilon} \leq \frac{2}{n}$$

and

$$\frac{p\theta}{1 - p\theta} = \frac{p(\gamma - 2\beta + \alpha) + \varepsilon(2 - p)}{n - p(\gamma - 2\beta + \alpha) + \varepsilon p} \leq \frac{3}{n - p(\gamma - 2\beta + \alpha) + \varepsilon p} =: \Lambda_\theta.$$

4C. The tails are finite. We here observe that all the terms on the right-hand sides of (3-6) and (4-15) are finite, obviously confining ourselves to those involving infinite sums. We start with the terms involving u . The second term appearing on the right-hand side of (3-6) is seen to be finite by estimating

$$\int_{\mathbb{R}^n \setminus \mathcal{B}} \frac{|u(y)|}{|x_0 - y|^{n+2\alpha}} dy \leq \int_{\mathbb{R}^n \setminus \mathcal{B}} \frac{|u(y) - (u)_B|}{|x_0 - y|^{n+2\alpha}} dy + \int_{\mathbb{R}^n \setminus \mathcal{B}} \frac{|(u)_B|}{|x_0 - y|^{n+2\alpha}} dy.$$

The last integral in this display is obviously finite, while the finiteness of the second one can be obtained as in (4-19). In fact, by (2-2) and since $\varepsilon \in (0, \frac{1}{2}\alpha)$, the right-hand side of (4-19) can be further estimated as

$$\begin{aligned} \sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^q d\mu \right)^{1/q} &\leq \sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^2 d\mu \right)^{1/2} \\ &\leq c(\varepsilon, \alpha) \left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy \right)^{1/2}. \end{aligned}$$

This also proves the finiteness of the first infinite sum appearing on the right-hand side of (4-15). We now come to the terms involving g , proving that the last series appearing in (4-15) is finite. The finiteness of the last series appearing in (3-6) is therefore implied by looking at the estimate for the term J_8 in the proof of Proposition 4.4. We start with the case $2\beta \geq \alpha$, where, using (4-9), we have

$$2^{-k(2\beta-\gamma-2\varepsilon/p)} \left(\int_{2^k \mathcal{B}} G^p d\mu \right)^{1/p} \leq c 2^{-k(2\beta-\gamma-\delta_1\gamma+n/[p(1+\delta_1)])} [g]_{\gamma(1+\delta_1), p(1+\delta_1)}$$

with $c \equiv c(n, \beta, \gamma, p, \delta_1, r)$, and since by (3-1) we have $\gamma < 2\beta$ and $\delta_1 \gamma p(1 + \delta_1) \leq n$, the convergence of the series follows. In the case $2\beta < \alpha$ we instead use (4-10) to have the inequality

$$2^{-k(2\beta - \gamma - 2\varepsilon/p)} \left(\int_{2^k \mathcal{B}} G^p d\mu \right)^{1/p} \leq c 2^{-k(2\beta + n/[p(1+\delta_1)])} \|g\|_{L^{p(1+\delta_1)}(\mathbb{R}^n)},$$

which again implies the convergence of the series in question.

5. Level set estimates for dual pairs

In this section we prove a level set estimate which is at the core of the proof of our higher differentiability and integrability results. Let us first define a few functionals. With θ and η as in (4-16), for every $\mathcal{B} \equiv \mathcal{B}(x, \varrho) \subset \mathbb{R}^{2n}$ we define

$$\begin{aligned} \Psi_{H,M}(\mathcal{B}(x, \varrho)) := & \left(\int_{\mathcal{B}(x, \varrho)} U^2 d\mu \right)^{1/2} + \frac{H[\mu(\mathcal{B}(x, \varrho))]^\eta}{\varepsilon^{1/2_* - 1/2}} \left(\int_{\mathcal{B}(x, \varrho)} F^{2_*} d\mu \right)^{1/2_*} \\ & + \frac{M[\mu(\mathcal{B}(x, \varrho))]^\theta}{\varepsilon^{1/p - 1/2}} \left(\int_{\mathcal{B}(x, \varrho)} G^p d\mu \right)^{1/p}, \end{aligned} \quad (5-1)$$

where $H, M \geq 1$ and $\mathcal{B}(x, \varrho) \subset \mathbb{R}^{2n}$. We also define the functionals

$$\Upsilon_0(\mathcal{B}(x, \varrho)) := \left(\int_{\mathcal{B}(x, \varrho)} F^{2_* + \delta_f} d\mu \right)^{1/(2_* + \delta_f)} + \left(\int_{\mathcal{B}(x, \varrho)} G^{p + \delta_g} d\mu \right)^{1/(p + \delta_g)}, \quad (5-2)$$

$$\Upsilon_1(\mathcal{B}(x, \varrho)) := \sum_{k=0}^{\infty} 2^{-k(\alpha - \varepsilon)} \left(\int_{\mathcal{B}(x, 2^k \varrho)} U^q d\mu \right)^{1/q} \quad (5-3)$$

and

$$\Upsilon_{2,M}(\mathcal{B}(x, \varrho)) := \frac{M[\mu(\mathcal{B}(x, \varrho))]^\theta}{\varepsilon^{1/p - 1/2}} \sum_{k=0}^{\infty} 2^{-k(2\beta - \gamma - 2\varepsilon/p)} \left(\int_{\mathcal{B}(x, 2^k \varrho)} G^p d\mu \right)^{1/p}. \quad (5-4)$$

We shall denote

$$\Psi(\mathcal{B}(x, \varrho)) := \Psi_{1,1}(\mathcal{B}(x, \varrho)),$$

and shall often use the abbreviations

$$\Psi_{H,M}(\mathcal{B}(x, \varrho)) \equiv \Psi_{H,M}(x, \varrho), \quad \Upsilon_0(\mathcal{B}(x, \varrho)) \equiv \Upsilon_0(x, \varrho),$$

and so forth. Finally, we can define

$$\text{ADD}(\mathcal{B}(x, \varrho)) \equiv \text{ADD}(x, \varrho) := \Psi(x, \varrho) + \Upsilon_0(x, \varrho) + \Upsilon_1(x, \varrho) + \Upsilon_{2,1}(x, \varrho). \quad (5-5)$$

The aim of this section is to prove the following:

Proposition 5.1. *Let $u \in W^{\alpha,2}(\mathbb{R}^n)$ be a solution to (1-14) under the assumptions of Theorem 1.1; in particular, (3-1) and (3-3) are in force. Let μ be the measure defined in (1-19), with ε satisfying (4-6).*

Consider a ball $\mathcal{B}(x_0, 2\varrho_0) \subset \mathbb{R}^{2n}$ such that $\varrho_0 \leq 1$, and related concentric balls

$$\mathcal{B}(x_0, \varrho_0) \subset \mathcal{B}(x_0, t) \subset \mathcal{B}(x_0, s) \subset \mathcal{B}(x_0, \frac{3}{2}\varrho_0) \quad (5-6)$$

for $\varrho_0 < t < s < \frac{3}{2}\varrho_0$. There exists a constant $c_s \equiv c_s(n, \alpha, \Lambda)$ independent of ε and p , and constants $c_f \equiv c_f(n, \alpha, \Lambda, \varepsilon) > 1$, $c_g \equiv c_g(n, \alpha, \Lambda, \beta, \gamma, p, \varepsilon) \geq 1$, $\kappa_f \equiv \kappa_f(n, \alpha, \Lambda, \varepsilon) \in (0, 1)$, $\kappa_g \equiv \kappa_g(n, \alpha, \Lambda, \beta, p, \varepsilon) \in (0, 1)$, such that the inequality

$$\begin{aligned} \frac{1}{\lambda^2} \int_{\mathcal{B}(x_0, t) \cap \{U > \lambda\}} U^2 d\mu &\leq \frac{c_s}{\varepsilon^{3(2-q)/q} \lambda^q} \int_{\mathcal{B}(x_0, s) \cap \{U > \lambda\}} U^q d\mu \\ &+ \frac{c_f \lambda_0^{(2_* + \delta_f)2_* \eta / (1 - 2_* \eta)}}{\lambda^{(1 + \eta \delta_f)2_* / (1 - 2_* \eta)}} \int_{\mathcal{B}(x_0, s) \cap \{F > \kappa_f \lambda\}} F^{2_*} d\mu \frac{c_g \lambda_0^{(p + \delta_g)p\theta / (1 - p\theta)}}{\lambda^{(1 + \theta \delta_g)p / (1 - p\theta)}} \int_{\mathcal{B}(x_0, s) \cap \{G > \kappa_g \lambda\}} G^p d\mu \end{aligned} \quad (5-7)$$

holds whenever $\lambda \geq \lambda_0$, where

$$\lambda_0 := \frac{c_a}{\varepsilon} \left(\frac{\varrho_0}{s - t} \right)^{2n} \text{ADD}(x_0, 2\varrho_0). \quad (5-8)$$

The constant c_a introduced in the last display depends on $n, \alpha, \Lambda, \beta, \gamma$, but is still independent of ε .

Remark 5.2. Unlike κ_f, c_f , the constants κ_g, c_g exhibit the following behavior:

$$\lim_{p \rightarrow 2n/[n+2(\gamma-2\beta+\alpha)]} \frac{1}{\kappa_g} = \lim_{\gamma \rightarrow 2\beta} \frac{1}{\kappa_g} = \infty = \lim_{p \rightarrow 2n/[n+2(\gamma-2\beta+\alpha)]} c_g = \lim_{\gamma \rightarrow 2\beta} c_g. \quad (5-9)$$

The proof of [Proposition 5.1](#) is rather delicate and falls into twelve steps. It will take the rest of this section.

5A. Diagonal balls and Vitali's covering. The proof starts with an exit-time argument for the functional $\Psi_{H,M}(\cdot)$, aimed at covering the ‘‘diagonal’’ level set of U . The constants $H, M \geq 1$ shall be fixed in due course of the proof, and the whole argument is independent of their particular values until the moment these are fixed. They will be used to give a different weight to the integrals of F^{2_*} and G^p : at the exit time, the averages of F^{2_*}, G^p will be smaller than the one of U^2 provided H, M are chosen to be large enough, respectively. Let us consider concentric diagonal balls as in (5-6). Let $\kappa \in (0, 1]$ be a free parameter to be chosen in the course of the proof, and define

$$\tilde{\lambda}_0 := \kappa^{-1} \sup_{\frac{s-t}{40^n} \leq \varrho \leq \frac{\varrho_0}{2}} \sup_{x \in \mathcal{B}(x_0, t)} \{ \Psi_{H,M}(x, \varrho) + \Upsilon_0(x, \varrho) + \Upsilon_1(x, \varrho) + \Upsilon_{2,M}(x, \varrho) \}. \quad (5-10)$$

All the foregoing steps of proofs are independent of the specific choice of κ until we fix κ in (5-55) below. For the same κ (to be defined later) and for $\lambda \geq \tilde{\lambda}_0$, define further the ‘‘diagonal level set’’

$$D_{\kappa\lambda} := \left\{ (x, x) \in \mathcal{B}(x_0, t) : \sup_{0 < \varrho < \frac{s-t}{40^n}} \Psi_{H,M}(x, \varrho) > \kappa\lambda \right\}. \quad (5-11)$$

Since, by the definition in (5-10), we have

$$\Psi_{H,M}(x, \varrho) \leq \kappa \tilde{\lambda}_0 \leq \kappa\lambda \quad \text{if } (x, x) \in \mathcal{B}(x_0, t) \text{ and } \varrho \in [(s-t)/40^n, \varrho_0/2], \quad (5-12)$$

we can find for all $(x, x) \in D_{\kappa\lambda}$ an exit radius $\varrho(x) \in (0, (s-t)/40^n)$ such that

$$\Psi_{H,M}(x, \varrho(x)) \geq \kappa\lambda, \quad \text{while} \quad \sup_{\varrho(x) < \varrho < \frac{s-t}{40^n}} \Psi_{H,M}(x, \varrho) \leq \kappa\lambda. \quad (5-13)$$

Collect the enlarged balls into a covering $\{\mathcal{B}(x, 2\varrho(x)) : (x, x) \in D_{\kappa\lambda}\}$. Balls of the type $\mathcal{B}(x, t, \varrho)$ are, as explained in [Section 2](#), metric balls with respect to the metric (2-1). We therefore apply Vitali's covering theorem to find a countable set J_D and related diagonal points $\{(x_j, x_j)\}_{j \in J_D}$ such that

$$\bigcup_{(x,x) \in D_{\kappa\lambda}} \mathcal{B}(x, 2\varrho(x)) \subset \bigcup_{j \in J_D} \mathcal{B}(x_j, 10\varrho(x_j)) \subset \mathcal{B}(x_0, s) \quad (5-14)$$

and

$$\{\mathcal{B}(x_j, 2\varrho(x_j))\}_{j \in J_D} \quad \text{is a family of mutually disjoint balls.} \quad (5-15)$$

Implicit in (5-14) is the fact that, since $\varrho(x_j) \leq (s-t)/40^n$ and $x_j \in \mathcal{B}(x_0, t)$ for every $x_j \in J_D$, then $\mathcal{B}(x_j, 10\varrho(x_j)) \subset \mathcal{B}(x_0, s)$. By (5-12)–(5-13) and the doubling property in (4-2), it follows that

$$\begin{aligned} \sum_{j \in J_D} \int_{\mathcal{B}(x_j, 10\varrho(x_j))} U^2 d\mu &\leq \sum_{j \in J_D} \mu(\mathcal{B}(x_j, 10\varrho(x_j))) [\Psi_{H,M}(\mathcal{B}(x_j, 10\varrho(x_j)))]^2 \\ &\leq 10^{n+2\varepsilon} \kappa^2 \lambda^2 \sum_{j \in J_D} \mu(\mathcal{B}(x_j, \varrho(x_j))). \end{aligned} \quad (5-16)$$

We shall denote in short

$$\mathcal{B}_j := \mathcal{B}(x_j, \varrho(x_j)), \quad \sigma \mathcal{B}_j := \mathcal{B}(x_j, \sigma \varrho(x_j)), \quad \sigma > 0. \quad (5-17)$$

Finally, since we are assuming that $\varrho_0 \leq 1$, by (4-1) we observe that

$$\mu(\mathcal{B}(x_0, 2\varrho_0)) \leq \frac{c2^{n+2\varepsilon}}{\varepsilon} =: L \equiv L(n, \varepsilon). \quad (5-18)$$

5B. Dyadic cubes, and two constants. This section has a very technical nature, and reports a few facts that are true independently of the specify context we are working in. In order to cover the off-diagonal level sets of U , we need a more elaborate argument based on classical Calderón–Zygmund coverings. To this aim, we start by recalling basic properties of dyadic cubes in \mathbb{R}^{2n} . They differ from the usual ones since they are “centered” at x_0 and the size is adapted to the size of the starting ball $\mathcal{B}(x_0, s)$. Define

$$k_0 := \left\lceil -\log_2 \left(\frac{s-t}{n10^{10n}} \right) \right\rceil + 1, \quad (5-19)$$

where $\lceil \cdot \rceil$ denotes the integer part of a given number, with the (unnecessarily large) constant 10^{10n} having also a symbolic meaning. Let Δ_k , $k \geq k_0$, be the disjoint collection—centered at x_0 —of half-open cubes of side length 2^{-k} whose closures are touching $\bar{\mathcal{B}}(x_0, \frac{1}{2}(s+t))$, i.e.,

$$\Delta_k := \{x_0 + 2^{-k}v + [0, 2^{-k})^n : v \in \mathbb{Z}^n, (x_0 + 2^{-k}v + [0, 2^{-k})^n \cap \bar{\mathcal{B}}(x_0, \frac{1}{2}(s+t)) \neq \emptyset\}.$$

Notice that, with such a definition, by using (5-19) it follows that $k \geq k_0$ implies

$$B(x_0, t) \subset \bigcup_{K \in \Delta_k} K \subset B(x_0, s). \quad (5-20)$$

The cubes defined above are, up to a translation aimed at centering everything at x_0 , the standard dyadic cubes in \mathbb{R}^n . Let us recall a few basic properties. Let Δ be the family of all cubes from the families Δ_k , that is, $\Delta := \{K \in \Delta_k : k \geq k_0\}$. Defined this way, every cube K in Δ_{k+1} , $k \geq k_0$, has only one predecessor $\tilde{K} \in \Delta_k$ such that $K \subset \tilde{K}$. Moreover, if $K_1 \in \Delta_{k_1}$ and $K_2 \in \Delta_{k_2}$ with $k_0 \leq k_1 < k_2$ and also $K_1 \cap K_2 \neq \emptyset$, then $K_2 \subset K_1$. Starting from the previous cubes, we fix the notation for the corresponding ones in \mathbb{R}^{2n} . We set, again for $k \geq k_0$,

$$\Xi_k := \{\mathcal{K} \equiv K_1 \times K_2 : K_1, K_2 \in \Delta_k\}, \quad \Xi := \bigcup_{k \geq k_0} \Xi_k,$$

while the diagonal cubes build up the family

$$\tilde{\Xi}_k := \{\mathcal{K} \equiv K \times K : K \in \Delta_k\}. \quad (5-21)$$

With the above definition, it follows from (5-20) that

$$B(x_0, t) \subset \bigcup_{\mathcal{K} \in \Xi_k} \mathcal{K} \subset B(x_0, s) \quad (5-22)$$

whenever $k \geq k_0$. Notice that, by defining the product cubes as above, we are actually once again considering dyadic cubes in \mathbb{R}^{2n} , with the same properties of the cubes from Δ_k . We also notice that if $\Xi \ni \mathcal{K} = K_1 \times K_2$ then $\tilde{\mathcal{K}} = \tilde{K}_1 \times \tilde{K}_2$ is its unique predecessor. Finally, let $\mathcal{K} \in \Xi$; then there exist $K_1, K_2 \in \Delta_k$ such that $\mathcal{K} = K_1 \times K_2$; in this case we let

$$k(\mathcal{K}) = k. \quad (5-23)$$

Next, again with $\mathcal{K} = K_1 \times K_2$, we define the *cube projections* as

$$P_1(\mathcal{K}) \equiv P_1\mathcal{K} := K_1 \times K_1 \quad \text{and} \quad P_2(\mathcal{K}) \equiv P_2\mathcal{K} := K_2 \times K_2$$

whenever $K_1, K_2 \in \Delta_k$. In order to shorten the notation, we shall also write $P_h(\mathcal{K}) = P_h\mathcal{K}$ for $h = 1, 2$. It hence follows that

$$P_1(K_1 \times K_2) = P_2(K_2 \times K_1). \quad (5-24)$$

For a given cube $\mathcal{K} \equiv K_1 \times K_2$ we define

$$\tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) := \text{dist}(K_1, K_2), \quad (5-25)$$

and its symmetric (or mirror-reflected) cube with respect to the diagonal Diag , is defined by

$$\text{Symm}(\mathcal{K}) = \text{Symm}(K_1 \times K_2) := K_2 \times K_1. \quad (5-26)$$

For future convenience we collect a few basic facts that are a direct consequence of the definitions above, and in particular of (5-23)–(5-26).

Proposition 5.3. *Let $\mathcal{K} = K_1 \times K_2 \in \Xi$. The following facts are true:*

- $P_1\mathcal{K}, P_2\mathcal{K} \in \Xi$.
- $\mu(P_1\mathcal{K}) = \mu(P_2\mathcal{K})$ and $k(\mathcal{K}) = k(P_1\mathcal{K}) = k(P_2\mathcal{K})$.
- If $\Xi \ni \mathcal{H} \subset \mathcal{K}$, then $k(\mathcal{K}) \leq k(\mathcal{H})$.
- If $\tilde{\mathcal{K}}$ is the predecessor of \mathcal{K} , then

$$\tilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}}) \leq \tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}). \quad (5-27)$$

- The following relations hold:

$$\text{dist}(P_1\mathcal{K}, P_2\mathcal{K}) = \sqrt{2} \tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}), \quad (5-28)$$

$$\text{dist}(\mathcal{K}, \text{Diag}) = \frac{1}{2} \text{dist}(P_1\mathcal{K}, P_2\mathcal{K}) = \frac{1}{\sqrt{2}} \tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}), \quad (5-29)$$

$$\text{dist}(\mathcal{K}, P_1\mathcal{K}) = \text{dist}(\mathcal{K}, P_2\mathcal{K}) = \text{dist}(K_1, K_2) = \tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}), \quad (5-30)$$

$$\tilde{\text{dist}}(P_1 \text{Symm}(\mathcal{K}), P_2 \text{Symm}(\mathcal{K})) = \tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}).$$

- Let $F : (x, y) \in \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a locally μ -integrable function which is symmetric, i.e., $F(x, y) = F(y, x)$ holds for every $x, y \in \mathbb{R}^n$. Then

$$\int_{\mathcal{K}} F d\mu = \int_{\text{Symm}(\mathcal{K})} F d\mu$$

whenever $\mathcal{K} \in \Xi$. In particular, $\mu(\mathcal{K}) = \mu(\text{Symm}(\mathcal{K}))$ and, moreover, $k(\mathcal{K}) = k(\text{Symm}(\mathcal{K}))$.

In the next two lemmas we introduce the ε -independent constants c_{dd} and \tilde{c}_d , and these will be used very often throughout.

Lemma 5.4. *There exists a constant c_{dd} , depending only on n , and in particular independent of ε , such that for $h \in \{1, 2\}$ we have the inequality*

$$c_{dd} \geq \sup_{\mathcal{K} \in \Xi} \left\{ \frac{1}{\varepsilon} \left(\frac{\tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})}{2^{-k(\mathcal{K})}} \right)^{n-2\varepsilon} \frac{\mu(\mathcal{K})}{\mu(P_h\mathcal{K})} \right\} + \sup_{\mathcal{K} \in \Xi, \tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k(\mathcal{K})}} \varepsilon \left\{ \left(\frac{\tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})}{2^{-k(\mathcal{K})}} \right)^{2\varepsilon-n} \frac{\mu(P_h\mathcal{K})}{\mu(\mathcal{K})} \right\} + 1. \quad (5-31)$$

Proof. Indeed, observe that using the definition of the measure μ together with (5-25) (and assuming without loss of generality that $\tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) > 0$) we have

$$\mu(P_h\mathcal{K}) = \frac{c(n)}{\varepsilon} 2^{-k(\mathcal{K})(n+2\varepsilon)} \quad \text{and} \quad \mu(\mathcal{K}) \leq \frac{2^{-2k(\mathcal{K})n}}{\tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})^{n-2\varepsilon}}.$$

This allows us to bound the first quantity in (5-31) in a universal way:

$$\frac{1}{\varepsilon} \left(\frac{\tilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})}{2^{-k(\mathcal{K})}} \right)^{n-2\varepsilon} \frac{\mu(\mathcal{K})}{\mu(P_h\mathcal{K})} \leq c(n).$$

On the other hand, again by (5-25), if $x \in K_1$ and $y \in K_2$ then

$$\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \leq |x - y| \leq 2\sqrt{n}[2^{-k(\mathcal{K})} + \widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})],$$

so that the very definition of the measure μ yields

$$\mu(\mathcal{K}) \geq \frac{2^{-2k(\mathcal{K})n}}{(2\sqrt{n})^{n-2\varepsilon}[2^{-k(\mathcal{K})} + \widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})]^{n-2\varepsilon}}.$$

Then we have

$$\varepsilon \left(\frac{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})}{2^{-k(\mathcal{K})}} \right)^{2\varepsilon-n} \frac{\mu(P_h\mathcal{K})}{\mu(\mathcal{K})} \leq c(n) \left(\frac{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})}{2^{-k(\mathcal{K})}} \right)^{2\varepsilon-n} \frac{[2^{-k(\mathcal{K})} + \widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})]^{n-2\varepsilon}}{2^{-2k(\mathcal{K})n+k(\mathcal{K})(n+2\varepsilon)}} \leq c(n),$$

where we have used that $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k(\mathcal{K})}$. We have therefore proved that (5-31) holds for a constant c_{dd} depending only on n . \square

The second constant is presented in the next lemma.

Lemma 5.5. *There exists a constant \tilde{c}_d , depending only on n , in particular independent of ε , such that the following inequality holds:*

$$\sup \left\{ \frac{\mu(\tilde{\mathcal{K}})}{\mu(\mathcal{K})} : \tilde{\mathcal{K}} \text{ is the predecessor of } \mathcal{K}, \widetilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}}) \geq 2^{-k(\mathcal{K})} \right\} \leq \tilde{c}_d. \quad (5-32)$$

Proof. Let us consider a dyadic cube $\mathcal{K} = K_1 \times K_2 \subset \mathbb{R}^{2n}$, with $\tilde{\mathcal{K}}$ being its predecessor, and such that $\widetilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}}) \geq 2^{-k(\mathcal{K})}$. The triangle inequality gives

$$|x - y| \leq 2\sqrt{n}2^{-k(\mathcal{K})+1} + \text{dist}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}}) \leq 8\sqrt{n} \text{dist}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}})$$

whenever $(x, y) \in K_1 \times K_2$. By the very definition of μ and (5-25), and finally using the inequality in the previous line when performing the final estimation, we get

$$\begin{aligned} \mu(\tilde{\mathcal{K}}) &\leq \widetilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}})^{-(n-2\varepsilon)} |\tilde{K}_1 \times \tilde{K}_2| \\ &= 4^n \widetilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}})^{-(n-2\varepsilon)} |K_1 \times K_2| \\ &\leq c(n)\mu(\mathcal{K}), \end{aligned}$$

and the proof of the lemma is complete. \square

5C. Off-diagonal cubes and Calderón–Zygmund coverings. We start by reporting an adaptation of the classical Calderón–Zygmund decomposition lemma. The argument is completely similar to the classical one and for a proof we refer for instance to [Stein 1993], taking into account that the measure μ is doubling and absolutely continuous with respect to the Lebesgue measure.

Theorem 5.6. *Let Q_0 be a cube in \mathbb{R}^{2n} and let \tilde{U} be a nonnegative function in $L^1(Q_0)$. Let $\tilde{\lambda}$ be a real number such that*

$$\int_{Q_0} \tilde{U} d\mu \leq \tilde{\lambda}.$$

There exists a countable, but possibly finite, family of pairwise disjoint dyadic cubes $\{Q_i\}$, with sides parallel to those of Q_0 , such that

$$\tilde{\lambda} < \int_{Q_i} \tilde{U} d\mu \quad \text{and} \quad \int_{\tilde{Q}_i} \tilde{U} d\mu \leq \tilde{\lambda} \quad \text{for every } Q_i,$$

where \tilde{Q}_i denotes the predecessor of Q_i , and

$$\tilde{U} \leq \tilde{\lambda} \quad \text{a.e. in } Q_0 \setminus \bigcup_i Q_i.$$

We now start to cover the off-diagonal part of the level set of U . To this end, let us consider the cubes from the family Ξ_{k_0} and, accordingly, the quantity

$$\lambda_1 := \max \left\{ \tilde{\lambda}_0, \sup_{\mathcal{K} \in \Xi_{k_0}} \left(\int_{\mathcal{K}} U^2 d\mu \right)^{1/2} \right\}. \quad (5-33)$$

We recall that the numbers $\tilde{\lambda}_0$ and k_0 have been determined in (5-10) and (5-19), respectively. Let us observe that (5-22) implies that the family $\{\mathcal{K}\}_{\mathcal{K} \in \Xi_{k_0}}$ forms a disjoint covering of $\mathcal{B}(x_0, t)$. With $\lambda \geq \lambda_1$ we now apply Theorem 5.6 with the choice $Q_0 \equiv \mathcal{K}_0$, for every single cube $\mathcal{K}_0 \in \Xi_{k_0}$; we therefore obtain a family of disjoint dyadic cubes $Q_i(\mathcal{K}_0)$ such that

$$\lambda^2 < \int_{Q_i(\mathcal{K}_0)} U^2 d\mu \quad \text{and} \quad \int_{\tilde{Q}_i(\mathcal{K}_0)} U^2 d\mu \leq \lambda^2 \quad \text{for every } Q_i,$$

where, as usual, $\tilde{Q}_i(\mathcal{K}_0)$ denotes the predecessor of $Q_i(\mathcal{K}_0)$, and

$$U \leq \lambda \quad \text{a.e. in } \mathcal{K}_0 \setminus \bigcup_i Q_i(\mathcal{K}_0).$$

Putting all such families of cubes together, we get a countable family

$$\mathcal{U}_\lambda := \bigcup_{\mathcal{K}_0 \in \Xi_{k_0}} \{Q_i(\mathcal{K}_0)\} \equiv \{\mathcal{K}\}$$

of disjoint dyadic cubes \mathcal{K} which are such that

$$\lambda^2 < \int_{\mathcal{K}} U^2 d\mu \quad \text{and} \quad \int_{\tilde{\mathcal{K}}} U^2 d\mu \leq \lambda^2 \quad \text{for every } \mathcal{K} \in \mathcal{U}_\lambda, \quad (5-34)$$

where $\tilde{\mathcal{K}}$ denotes the predecessor of \mathcal{K} , and such that

$$U \leq \lambda \quad \text{a.e. in } \mathcal{B}(x_0, t) \setminus \bigcup_{\mathcal{K} \in \mathcal{U}_\lambda} \mathcal{K}. \quad (5-35)$$

Remark 5.7. The symmetry of the function U and Proposition 5.3 imply that

$$\int_{\mathcal{K}} U^2 d\mu = \int_{\text{Symm}(\mathcal{K})} U^2 d\mu$$

whenever $\mathcal{K} \in \Xi$. It then follows that $\mathcal{K} \in \mathcal{U}_\lambda$ if and only if $\text{Symm}(\mathcal{K}) \in \mathcal{U}_\lambda$.

5D. First removal of nearly diagonal cubes. In this step we are going to show that, in order to cover the level sets of U^2 , it is sufficient to restrict our attention to those dyadic cubes that are “far” from the diagonal in a suitably quantified sense. Specifically, the word far refers to the fact that for such cubes it happens that their distance to the diagonal is larger than their size. These are really the relevant cubes to analyze, since we shall see that the remaining ones can be covered by the balls considered in (5-14)–(5-15). We therefore start by considering the family of *nearly diagonal* cubes

$$\mathcal{U}_\lambda^d := \{\mathcal{K} \in \mathcal{U}_\lambda : \widetilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}}) < 2^{-k(\mathcal{K})}, \tilde{\mathcal{K}} \text{ is the predecessor of } \mathcal{K}\}.$$

With $\mathcal{K} \in \mathcal{U}_\lambda^d$, consider now a point $(\tilde{x}, \tilde{x}) \in \text{Diag}$ such that $\text{dist}((\tilde{x}, \tilde{x}), \tilde{\mathcal{K}}) = \text{dist}(\text{Diag}, \tilde{\mathcal{K}})$ and a diagonal ball $\mathcal{B}(\tilde{x}, \varrho) \subset \mathbb{R}^{2n}$ with radius ϱ greater than or equal to

$$\frac{5\sqrt{n} \widetilde{\text{dist}}(P_1\tilde{\mathcal{K}}, P_2\tilde{\mathcal{K}})}{2} + 5\sqrt{n}2^{-k(\mathcal{K})+1}.$$

Keeping (5-29) in mind and applying it to $\tilde{\mathcal{K}}$, it follows that $\tilde{\mathcal{K}} \subset \mathcal{B}(\tilde{x}, \varrho)$. Ultimately, we can find a diagonal ball $\mathcal{B} \equiv \mathcal{B}(\tilde{x}, 24\sqrt{n}2^{-k(\mathcal{K})})$ such that $\mathcal{K} \subset \mathcal{B}$. Notice that, in this case, by using (4-3) from Proposition 4.1 and recalling that $(\tilde{x}, \tilde{x}) \in \text{Diag}$, we conclude there exists a constant c_d , which depends only on n , such that

$$1 \leq \frac{\mu(\mathcal{B})}{\mu(\mathcal{K})} \leq \frac{c_d}{\varepsilon} \equiv \frac{c_d(n)}{\varepsilon}. \quad (5-36)$$

Therefore, if $\mathcal{K} \in \mathcal{U}_\lambda^d$, then the lower bound in (5-34) yields

$$\lambda^2 < \int_{\mathcal{K}} U^2 d\mu \leq \frac{\mu(\mathcal{B})}{\mu(\mathcal{K})} \int_{\mathcal{B}} U^2 d\mu \leq \frac{c_d}{\varepsilon} \int_{\mathcal{B}} U^2 d\mu.$$

Assuming that the number $\kappa \in (0, 1]$ introduced in (5-10) satisfies

$$\kappa \in (0, \kappa_0], \quad \kappa_0 := \frac{\varepsilon^{1/2}}{\sqrt{2c_d}}, \quad (5-37)$$

all in all we have proved that

$$\text{for all } \mathcal{K} \in \mathcal{U}_\lambda^d, \text{ there exists } \mathcal{B}^{\mathcal{K}} \equiv B^{\mathcal{K}} \times B^{\mathcal{K}} \quad \text{such that} \quad \int_{\mathcal{B}^{\mathcal{K}}} U^2 d\mu > \kappa^2 \lambda^2 \text{ and } \mathcal{K} \subset \mathcal{B}^{\mathcal{K}}.$$

This means that, if \tilde{x} is the center of $\mathcal{B}^{\mathcal{K}}$, by the exit-time condition (5-13) it follows that $(\tilde{x}, \tilde{x}) \in D_{\kappa\lambda}$ and then $\mathcal{B}^{\mathcal{K}} \subset \mathcal{B}(\tilde{x}, \varrho(\tilde{x}))$. By (5-14) it hence follows that

$$\bigcup_{\mathcal{K} \in \mathcal{U}_\lambda^d} \mathcal{K} \subset \bigcup_{j \in J_D} 10\mathcal{B}_j. \quad (5-38)$$

Notice that here, in order to find the ball $\mathcal{B}^{\mathcal{K}}$ and apply the exit-time condition in (5-13), we have used that the radius of the diagonal ball $\mathcal{B} \equiv \mathcal{B}(\tilde{x}, 24\sqrt{n}2^{-k(\mathcal{K})})$ is smaller than $(s-t)/40^n$. In turn, this is a consequence of the fact that $k(\mathcal{K}) \geq k_0$ and of the fact that k_0 is large enough, as prescribed in (5-19).

5E. Off-diagonal reverse Hölder inequalities. As we saw in the previous section, \mathcal{U}_λ^d has already been covered by the diagonal cover. Thus, we shall now only consider so-called off-diagonal cubes:

$$\mathcal{U}_\lambda^{nd} := \{\mathcal{K} \in \mathcal{U}_\lambda : \widetilde{\text{dist}}(P_1\widetilde{\mathcal{K}}, P_2\widetilde{\mathcal{K}}) \geq 2^{-k(\mathcal{K})}, \widetilde{\mathcal{K}} \text{ is the predecessor of } \mathcal{K}\}. \quad (5-39)$$

We notice that (5-27) implies

$$\mathcal{K} \in \mathcal{U}_\lambda^{nd} \implies \widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k(\mathcal{K})}.$$

The goal is thus to sort and estimate suitable off-diagonal sums of the measures of cubes belonging to \mathcal{U}_λ^{nd} . The following lemma is our basic tool. It roughly says that, for nondiagonal cubes, reverse Hölder inequalities hold automatically, and independently of the fact that the function solves an equation. The price to pay is the appearance of certain diagonal correction terms, and this is eventually treated by some combinatorial lemmas.

Lemma 5.8 (off-diagonal reverse inequality). *Let $k \geq k_0$, and suppose that $\mathcal{K} \in \Xi_k$. There exists a constant $c_{nd} \equiv c_{nd}(n, \alpha)$, independent of ε , such that whenever $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k}$, the inequality*

$$\begin{aligned} \left(\int_{\mathcal{K}} U^2 d\mu \right)^{1/2} &\leq c_{nd} \left(\int_{\mathcal{K}} U^q d\mu \right)^{1/q} + \frac{c_{nd}}{\varepsilon^{1/q}} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{\alpha+\varepsilon} \left(\int_{P_1\mathcal{K}} U^q d\mu \right)^{1/q} \\ &\quad + \frac{c_{nd}}{\varepsilon^{1/q}} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{\alpha+\varepsilon} \left(\int_{P_2\mathcal{K}} U^q d\mu \right)^{1/q} \end{aligned}$$

holds, with the number q being defined in (4-13). In particular, this inequality holds whenever $\mathcal{K} \in \mathcal{U}_\lambda^{nd}$.

Proof. Let $\mathcal{K} \equiv K_1 \times K_2 \in \Xi_k$, and find points $x_1 \in \bar{K}_1$ and $y_1 \in \bar{K}_2$ such that $\text{dist}(K_1, K_2) = |x_1 - y_1|$. By the triangle inequality we obtain, whenever $x, y \in \mathcal{K}$,

$$\begin{aligned} |x - y| &\leq \text{dist}(K_1, K_2) + |x_1 - x| + |y_1 - y| \\ &\leq \text{dist}(K_1, K_2) + 2\sqrt{n}2^{-k} \\ &\leq 3\sqrt{n}\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) = 3\sqrt{n}\text{dist}(K_1, K_2). \end{aligned}$$

Therefore we have

$$1 \leq \frac{|x - y|}{\text{dist}(K_1, K_2)} \leq 3\sqrt{n} \quad \text{for all } (x, y) \in \mathcal{K}, \quad (5-40)$$

where the first inequality is a trivial consequence of the definition of $\text{dist}(K_1, K_2)$. Next, thanks to (5-40), the very definition of μ yields

$$\mu(\mathcal{K}) \approx \frac{4^{-nk}}{\text{dist}(K_1, K_2)^{n-2\varepsilon}}, \quad (5-41)$$

with the constant involved being independent of ε , but just depending on n . By using (5-40) and (5-41) we then have

$$\begin{aligned} \left(\int_{\mathcal{K}} U^2 d\mu \right)^{1/2} &= \left(\frac{1}{\mu(\mathcal{K})} \int_{K_1} \int_{K_2} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2\alpha}} dx dy \right)^{1/2} \\ &\leq c \left(\frac{\text{dist}(K_1, K_2)^{n-2\varepsilon-(n+2\alpha)}}{4^{-nk(\mathcal{K})}} \int_{K_1} \int_{K_2} |u(x) - u(y)|^2 dx dy \right)^{1/2} \\ &\leq c \text{dist}(K_1, K_2)^{-(\alpha+\varepsilon)} \left(\int_{K_1} \int_{K_2} |u(x) - u(y)|^2 dx dy \right)^{1/2}, \end{aligned} \quad (5-42)$$

where c depends only on n . We further estimate the integral on the right using Minkowski's inequality:

$$\begin{aligned} \left(\int_{K_1} \int_{K_2} |u(x) - u(y)|^2 dx dy \right)^{1/2} &\leq \left(\int_{K_1} |u(x) - (u)_{K_1}|^2 dx \right)^{1/2} \\ &\quad + \left(\int_{K_2} |u(x) - (u)_{K_2}|^2 dx \right)^{1/2} + |(u)_{K_1} - (u)_{K_2}|. \end{aligned}$$

By using the fractional Poincaré inequality of Lemma 4.3 applied to cubes, and recalling that $P_h \mathcal{K} = K_h \times K_h$ for $h \in \{1, 2\}$, we deduce that

$$\left(\int_{K_h} |u(x) - (u)_{K_h}|^2 dx \right)^{1/2} \leq \frac{c 2^{-k(\alpha+\varepsilon)}}{\varepsilon^{1/q}} \left(\int_{P_h \mathcal{K}} U^q d\mu \right)^{1/q}, \quad h \in \{1, 2\},$$

with the implied constant c depending only on n and α . Finally, by Hölder's inequality, and using (5-40) and (5-41) repeatedly, we get

$$\begin{aligned} |(u)_{K_1} - (u)_{K_2}| &\leq \int_{K_1} \int_{K_2} |u(x) - u(y)| dx dy \\ &\leq \left(\int_{K_1} \int_{K_2} |u(x) - u(y)|^q dx dy \right)^{1/q} \\ &\leq c \left(\frac{1}{\text{dist}(K_1, K_2)^{n-2\varepsilon} \mu(\mathcal{K})} \int_{K_1} \int_{K_2} |u(x) - u(y)|^q dx dy \right)^{1/q} \\ &\leq c \left(\int_{\mathcal{K}} |u(x) - u(y)|^q d\mu \right)^{1/q} \\ &\leq c \text{dist}(K_1, K_2)^{\alpha+\varepsilon} \left(\int_{\mathcal{K}} U^q d\mu \right)^{1/q}, \end{aligned}$$

with $c \equiv c(n)$. Combining the content of the last four displays and recalling the definition in (5-25) finishes the proof. \square

We remark that the previous lemma works for any function $u \in W^{\alpha,2}$ and does not require that u solves any equation; moreover, the lemma works for every positive integer k . Applying it in the present situation gives the next result:

Corollary 5.9. *Let $k \geq k_0$ be an integer, and suppose that $\mathcal{K} \in \Xi_k$ is such that $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k}$. Assume that*

$$\left(\int_{\mathcal{K}} U^2 d\mu \right)^{1/2} \geq \lambda$$

and that the number κ introduced in (5-10) satisfies

$$\kappa \in (0, \kappa_1], \quad \kappa_1 := \frac{\varepsilon^{1/q}}{2^{1/q} 3c_{nd}}, \quad (5-43)$$

where $c_{nd} \equiv c_{nd}(n, \alpha)$ has been defined in Lemma 5.8. Then we have

$$\begin{aligned} \mu(\mathcal{K}) \leq & \frac{3^q c_{nd}^q}{\lambda^q} \int_{\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu + \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} \frac{\mu(\mathcal{K})}{\mu(P_1\mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_1\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu \\ & + \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} \frac{\mu(\mathcal{K})}{\mu(P_2\mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_2\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu. \end{aligned} \quad (5-44)$$

In particular, the inequality (5-44) holds whenever $\mathcal{K} \in \mathcal{U}_\lambda^{nd}$.

Proof. Appealing to Lemma 5.8, and using the elementary inequality $(a + b + c)^q \leq 3^{q-1}(a^q + b^q + c^q)$ valid for all nonnegative numbers $a, b, c \in \mathbb{R}$, we get

$$\frac{\lambda^q}{3^{q-1} c_{nd}^q} \leq \int_{\mathcal{K}} U^q d\mu + \frac{1}{\varepsilon} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{q(\alpha+\varepsilon)} \left(\int_{P_1\mathcal{K}} U^q d\mu + \int_{P_2\mathcal{K}} U^q d\mu \right).$$

To estimate the integrals appearing on the right-hand side, we note that by (5-43) we have

$$\int_E U^q d\mu \leq \kappa_1^q \lambda^q + \frac{1}{\mu(E)} \int_{E \cap \{U > \kappa\lambda\}} U^q d\mu$$

with $E \in \{\mathcal{K}, P_1\mathcal{K}, P_2\mathcal{K}\}$ so that, recalling that $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k}$, we gain

$$\begin{aligned} \frac{\lambda^q}{3^{q-1} c_{nd}^q} \leq & \frac{3\kappa_1^q \lambda^q}{\varepsilon} + \frac{1}{\mu(\mathcal{K})} \int_{\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu + \frac{1}{\varepsilon \mu(P_1\mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_1\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu \\ & + \frac{1}{\varepsilon \mu(P_2\mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_2\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu. \end{aligned}$$

Now (5-44) follows by inserting (5-43) in the last estimate and reabsorbing terms. \square

5F. Families of off-diagonal cubes. With \mathcal{U}_λ^{nd} as defined in (5-39), consider now the families

$$\mathcal{M}_\lambda^h := \left\{ \mathcal{K} \in \mathcal{U}_\lambda^{nd} : \int_{P_h\mathcal{K}} U^q d\mu \leq (10n)^{n+2} \kappa^q \lambda^q \right\} \quad (5-45)$$

and

$$\mathcal{N}_\lambda^h := \left\{ \mathcal{K} \in \mathcal{U}_\lambda^{nd} : \int_{P_h\mathcal{K}} U^q d\mu > (10n)^{n+2} \kappa^q \lambda^q \right\} \quad (5-46)$$

for $h \in \{1, 2\}$, with the number κ defined as in (5-10) and q defined as in (4-13). Furthermore, define

$$\mathcal{M}_\lambda := \mathcal{M}_\lambda^1 \cap \mathcal{M}_\lambda^2 \quad \text{and} \quad \mathcal{N}_\lambda := \mathcal{N}_\lambda^1 \cup \mathcal{N}_\lambda^2, \quad (5-47)$$

so that we have the decomposition into disjoint families

$$\mathcal{U}_\lambda^{nd} = \mathcal{M}_\lambda \cup \mathcal{N}_\lambda. \quad (5-48)$$

Lemma 5.10 (soft off-diagonal summation). *The inequality*

$$\sum_{\mathcal{K} \in \mathcal{M}_\lambda} \mu(\mathcal{K}) \leq \frac{6^q c_{nd}^q}{\lambda^q} \int_{\mathcal{B}(x_0, s) \cap \{U > \kappa \lambda\}} U^q d\mu \quad (5-49)$$

holds whenever the number κ in (5-10) satisfies

$$\kappa \in (0, \kappa_2], \quad \kappa_2 := \frac{\varepsilon^{1/q}}{8^{1/q} 3c_{nd} (10n)^{(n+2)/q}}. \quad (5-50)$$

The constant $c_{nd} \equiv c_{nd}(n, \alpha)$ was defined in Lemma 5.8 and appears in Corollary 5.9; it is independent of ε .

Proof. It is sufficient to prove that if $\mathcal{K} \in \mathcal{M}_\lambda$, then

$$\mu(\mathcal{K}) \leq \frac{6^q c_{nd}^q}{\lambda^q} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu. \quad (5-51)$$

After this, (5-49) follows, since the initial family \mathcal{U}_λ is disjoint and (5-22) holds. For the proof of (5-51), notice that if $\mathcal{K} \in \mathcal{M}_\lambda^h$, then we have, for $h \in \{1, 2\}$, that

$$\begin{aligned} & \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} \frac{\mu(\mathcal{K})}{\mu(P_h \mathcal{K})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\ & \leq \mu(\mathcal{K}) \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} \int_{P_h \mathcal{K}} U^q d\mu \leq \mu(\mathcal{K}) \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} (10n)^{n+2} \kappa^q \lambda^q \leq \frac{\mu(\mathcal{K})}{8}. \end{aligned} \quad (5-52)$$

Using this last estimate for $h \in \{1, 2\}$ in combination with (5-44), and reabsorbing terms, gives (5-51); the proof is therefore complete. \square

It remains to study the family \mathcal{N}_λ defined in (5-47). To this aim, we introduce the family of diagonal cubes defined by

$$P_h \mathcal{N}_\lambda := \{P_h \mathcal{K} : \mathcal{K} \in \mathcal{N}_\lambda^h\}, \quad h \in \{1, 2\}.$$

Keeping (5-24) and Remark 5.7 in mind, we have that

$$\mathcal{K} \in \mathcal{N}_\lambda^1 \iff \text{Symm}(\mathcal{K}) \in \mathcal{N}_\lambda^2 \quad (5-53)$$

whenever $\mathcal{K} \in \mathfrak{E}$. Now, let us make a remark: consider $T \in P_1 \mathcal{N}_\lambda$, so that $T = P_1(\mathcal{K})$ for some $\mathcal{K} \in \mathcal{N}_\lambda^1$. Therefore $T = P_2(\text{Symm}(\mathcal{K}))$ by (5-24), and by (5-53) we have $\text{Symm}(\mathcal{K}) \in \mathcal{N}_\lambda^2$. We conclude that

$T \in P_2\mathcal{N}_\lambda$ and eventually that $P_1\mathcal{N}_\lambda \subset P_2\mathcal{N}_\lambda$. In a similar way it follows that $P_2\mathcal{N}_\lambda \subset P_1\mathcal{N}_\lambda$. We therefore conclude that $P_1\mathcal{N}_\lambda = P_2\mathcal{N}_\lambda = P_1\mathcal{N}_\lambda \cup P_2\mathcal{N}_\lambda$. Let $P\mathcal{N}_\lambda$ be a disjoint subfamily of $P_1\mathcal{N}_\lambda \cup P_2\mathcal{N}_\lambda$ such that

$$\bigcup_{\mathcal{H} \in P\mathcal{N}_\lambda} \mathcal{H} = \bigcup_{\mathcal{K} \in P_1\mathcal{N}_\lambda \cup P_2\mathcal{N}_\lambda} \mathcal{K}. \quad (5-54)$$

Note that, since all the cubes of the family $P\mathcal{N}_\lambda$ are themselves dyadic cubes, such an extracted disjoint covering always exists. We remark that a straightforward consequence of the definitions is that all cubes from $P\mathcal{N}_\lambda$ obviously belong to $P_1\mathcal{N}_\lambda \cup P_2\mathcal{N}_\lambda$ and are therefore diagonal cubes.

5G. Determining κ . We here determine the parameter κ in (5-10). By choosing

$$\kappa := \min\{\kappa_0, \kappa_1, \kappa_2\} \equiv \min \left\{ \frac{\varepsilon^{1/2}}{\sqrt{2c_d}}, \frac{\varepsilon^{1/q}}{2^{1/q}3c_{nd}}, \frac{\varepsilon^{1/q}}{8^{1/q}3c_{nd}(10n)^{(n+2)/q}} \right\}, \quad (5-55)$$

conditions (5-37), (5-43) and (5-50) are all satisfied, so the content and the results of Sections 5D–5F are at our disposal. Recalling that c_d in (5-36) (coming from Proposition 4.1) depends only on n , and that c_{nd} from Lemma 5.8 depends only on n, α , we conclude there exists a new constant c_κ such that

$$\kappa \geq \varepsilon^{1/q}/c_\kappa, \quad c_\kappa \equiv c_\kappa(n, \alpha). \quad (5-56)$$

5H. Further removal of nearly diagonal cubes. We recall that our final goal is to estimate the measure of the level sets of U . Since the nearly diagonal part has already been covered, we proceed in excluding from the subsequent analysis those cubes covered by the balls in (5-14)–(5-15). Therefore, we introduce

$$\mathcal{N}_{\lambda,d} := \left\{ \mathcal{K} \in \mathcal{N}_\lambda : \mathcal{K} \subset \bigcup_{j \in J_D} 10\mathcal{B}_j \right\} \quad (5-57)$$

and, accordingly,

$$\mathcal{N}_{\lambda,nd} := \mathcal{N}_\lambda \setminus \mathcal{N}_{\lambda,d} \quad \text{and} \quad \mathcal{N}_{\lambda,nd}^h := \mathcal{N}_{\lambda,nd} \cap \mathcal{N}_\lambda^h, \quad \text{for } h \in \{1, 2\}. \quad (5-58)$$

We observe that the main difficulty in handling the cubes from the family $P\mathcal{N}_\lambda$ stems from the fact that they do not belong to the family \mathcal{U}_λ , i.e., they do not come from an exit-time argument and therefore no control is available on the values taken by U^2 on such cubes. This will be bypassed via a very delicate combinatorial argument. The next lemma is instrumental to that.

Lemma 5.11. *Let $\mathcal{K} \in \mathcal{N}_{\lambda,nd}$ be such that $P_h\mathcal{K} \subset \mathcal{H}$ for some $\mathcal{H} \in P\mathcal{N}_\lambda$ and some $h \in \{1, 2\}$. Then $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k(\mathcal{H})}$.*

Proof. First, let us consider a cube $\mathcal{H} \in P\mathcal{N}_\lambda$; take the diagonal ball $\mathcal{B}(\mathcal{H}) \equiv \mathcal{B}(x_\mathcal{H}, 2^{-(k(\mathcal{H})+1)})$, $(x_\mathcal{H}, x_\mathcal{H})$ being the center of \mathcal{H} . It follows that

$$\mathcal{B}(\mathcal{H}) \subset \mathcal{H} \subset \sqrt{n}\mathcal{B}(\mathcal{H}). \quad (5-59)$$

Therefore we have by Hölder's inequality and the definition of $P\mathcal{N}_\lambda$ that

$$\begin{aligned}
(10n)^{(n+2)/q} \kappa \lambda &< \left(\int_{\mathcal{H}} U^q d\mu \right)^{1/q} \\
&\leq \left(\frac{\mu(10n\mathcal{B}(\mathcal{H}))}{\mu(\mathcal{B}(\mathcal{H}))} \int_{10n\mathcal{B}(\mathcal{H})} U^q d\mu \right)^{1/q} \\
&\leq (10n)^{(n+2)/q} \left(\int_{10n\mathcal{B}(\mathcal{H})} U^2 d\mu \right)^{1/2}.
\end{aligned} \tag{5-60}$$

By the definition of $D_{\kappa\lambda}$ in (5-11) it follows that $(x_{\mathcal{H}}, x_{\mathcal{H}}) \in D_{\kappa\lambda}$, and then the exit-time condition (5-13) gives $\mathcal{B}(\mathcal{H}) \subset \mathcal{B}(x_{\mathcal{H}}, \varrho(x_{\mathcal{H}}))$. We are using that the radius of the ball $10n\mathcal{B}(\mathcal{H})$ is smaller than $(s-t)/40^n$. In turn, this is a consequence of the fact that $k(\mathcal{H}) + 1 \geq k_0$ and of (5-19). Then (5-14) implies

$$10n\mathcal{B}(\mathcal{H}) \subset \bigcup_{j \in J_D} 10\mathcal{B}_j. \tag{5-61}$$

Now, in order to prove the lemma, assume by contradiction that $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) < 2^{-k(\mathcal{H})}$ and let $\mathcal{B}(\mathcal{H})$ be the ball determined in (5-59), and for which (5-61) holds. We are going to show that

$$\mathcal{K} \subset 10n\mathcal{B}(\mathcal{H}), \tag{5-62}$$

and this then contradicts the assumption $\mathcal{K} \in \mathcal{N}_{\lambda, nd}$ by (5-61). In order to show (5-62), we observe that Proposition 5.3 and the fact that $P_h\mathcal{K} \subset \mathcal{H}$ give

$$\text{dist}(\mathcal{K}, \mathcal{H}) \leq \text{dist}(\mathcal{K}, P_h\mathcal{K}) = \widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \leq 2^{-k(\mathcal{H})}.$$

Again by Proposition 5.3 we have $k(P_h\mathcal{K}) = k(\mathcal{K})$ and $k(\mathcal{K}) \geq k(\mathcal{H})$. Therefore, since $\mathcal{H} \subset \sqrt{n}\mathcal{B}(\mathcal{H})$ and the radius of $\mathcal{B}(\mathcal{H})$ is $2^{-(k(\mathcal{H})+1)}$, then (5-62) must hold. The proof of the lemma is complete. \square

5I. Summation in $\mathcal{N}_{\lambda, nd}$. The aim of this section is to prove the following:

Lemma 5.12 (hard off-diagonal summation). *There exists a constant c , depending only on n, α , such that the estimate*

$$\sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \mu(\mathcal{K}) \leq \frac{c}{\lambda^q} \int_{\mathcal{B}(x_0, s) \cap \{U > \kappa\lambda\}} U^q d\mu \tag{5-63}$$

holds, where κ has been determined in (5-55).

Proof. Step 1: Classifying cubes. Here we classify the cubes from $\mathcal{N}_{\lambda, nd}$ according to their projections, thereby partitioning $\mathcal{N}_{\lambda, nd}$ into suitable disjoint subfamilies. For every $\mathcal{H} \in P\mathcal{N}_\lambda$, set

$$\mathcal{N}_{\lambda, nd}^h(\mathcal{H}) := \{\mathcal{K} \in \mathcal{N}_{\lambda, nd} : P_h\mathcal{K} \subset \mathcal{H}\}, \quad h \in \{1, 2\}.$$

Since $P\mathcal{N}_\lambda$ is a disjoint covering of $P_1\mathcal{N}_\lambda \cup P_2\mathcal{N}_\lambda = P_1\mathcal{N}_\lambda = P_2\mathcal{N}_\lambda$, we have the decomposition in mutually disjoint families

$$\mathcal{N}_{\lambda, nd}^h = \bigcup_{\mathcal{H} \in P\mathcal{N}_\lambda} \mathcal{N}_{\lambda, nd}^h(\mathcal{H}). \tag{5-64}$$

This means that for $\mathcal{H}_1, \mathcal{H}_2 \in P\mathcal{N}_\lambda$ it follows that $\mathcal{N}_{\lambda,nd}^h(\mathcal{H}_1) \cap \mathcal{N}_{\lambda,nd}^h(\mathcal{H}_2) \neq \emptyset$ implies $\mathcal{H}_1 = \mathcal{H}_2$. Indeed, assume that a cube \mathcal{K} lies in $\mathcal{N}_{\lambda,nd}^h(\mathcal{H}_1) \cap \mathcal{N}_{\lambda,nd}^h(\mathcal{H}_2)$ and that $\mathcal{H}_1 \neq \mathcal{H}_2$; then we would have that $P_h\mathcal{K} \subset \mathcal{H}_1 \cap \mathcal{H}_2$ against the fact that \mathcal{H}_1 and \mathcal{H}_2 have a nonempty intersection, being elements of the disjoint covering $P\mathcal{N}_\lambda$. Next, let us recall that for every $\mathcal{K} \in \mathcal{N}_{\lambda,nd}^h(\mathcal{H})$ we have $k(\mathcal{K}) = k(P_h\mathcal{K}) \geq k(\mathcal{H})$, and this leads us to define the classes

$$[\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_i := \{\mathcal{K} \in \mathcal{N}_{\lambda,nd}^h(\mathcal{H}) : k(\mathcal{K}) = i + k(\mathcal{H})\}$$

for $h \in \{1, 2\}$ and for every integer $i \geq 0$. Therefore, the decomposition in mutually disjoint families

$$\mathcal{N}_{\lambda,nd}^h(\mathcal{H}) = \bigcup_{i \geq 0} [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_i$$

holds, in the sense that $[\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_i \cap [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_j \neq \emptyset$ implies that $i = j$. Next, take $\mathcal{H} \in P\mathcal{N}_\lambda$; by [Lemma 5.11](#) we have that if $\mathcal{K} \in \mathcal{N}_{\lambda,nd}^h(\mathcal{H})$, that is, if $P_h\mathcal{K} \subset \mathcal{H}$, then it follows that $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k(\mathcal{H})}$, and this finally leads us to classify elements of $[\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_i$ in the following way:

$$[\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_{i,j} := \{\mathcal{K} \in [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_i : 2^{j-k(\mathcal{H})} \leq \widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) < 2^{j+1-k(\mathcal{H})}\}$$

for $h \in \{1, 2\}$ and for integers $i, j \geq 0$. Again, we have the decomposition

$$\mathcal{N}_{\lambda,nd}^h(\mathcal{H}) = \bigcup_{i,j \geq 0} [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_{i,j}, \quad (5-65)$$

and these are disjoint classes in the sense that, if $[\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_{i_1,j_1} \cap [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_{i_2,j_2} \neq \emptyset$, then $(i_1, j_1) = (i_2, j_2)$. All in all, in view of [\(5-64\)](#) and [\(5-65\)](#), we have the decomposition into mutually disjoint classes

$$\mathcal{N}_{\lambda,nd}^h = \bigcup_{\mathcal{H} \in P\mathcal{N}_\lambda} \bigcup_{i,j \geq 0} [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_{i,j}. \quad (5-66)$$

Step 2: Sums and further partitions. Let us fix $\mathcal{H} \in P\mathcal{N}_\lambda$; our aim here is to prove that the following inequality holds for $h \in \{1, 2\}$:

$$\frac{1}{\varepsilon} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda,nd}^h(\mathcal{H})} \frac{\mu(\mathcal{K})}{\mu(P_h\mathcal{K})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_h\mathcal{K} \cap \{U > \kappa\lambda\}} U^q d\mu \leq \frac{c(n)}{\alpha^2} \int_{\mathcal{H} \cap \{U > \kappa\lambda\}} U^q d\mu. \quad (5-67)$$

We start by recalling that, by the very definitions in [\(5-46\)](#) and [\(5-47\)](#), and again [\(5-27\)](#), we have that $\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K}) \geq 2^{-k(\mathcal{K})}$ as soon as $\mathcal{K} \in \mathcal{N}_{\lambda,nd}^h$; [\(5-31\)](#) yields

$$\frac{1}{\varepsilon} \frac{\mu(\mathcal{K})}{\mu(P_h\mathcal{K})} \leq c_{dd} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \right)^{n-2\varepsilon}$$

for $h \in \{1, 2\}$, and, moreover, if $\mathcal{K} \in [\mathcal{N}_{\lambda,nd}^h(\mathcal{H})]_{i,j}$, we also have that

$$\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} = \frac{1}{2^i} \frac{2^{-k(\mathcal{H})}}{\widetilde{\text{dist}}(P_1\mathcal{K}, P_2\mathcal{K})} \leq \frac{1}{2^{i+j}}.$$

Using the inequalities in the last two displays we can estimate

$$\begin{aligned}
& \frac{1}{\varepsilon} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}^h(\mathcal{H})} \frac{\mu(\mathcal{K})}{\mu(P_h \mathcal{K})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\
& \leq c_{dd} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}^h(\mathcal{H})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{n+q(\alpha+\varepsilon)-2\varepsilon} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\
& = c_{dd} \sum_{i, j=0}^{\infty} \sum_{\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j}} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{n+q(\alpha+\varepsilon)-2\varepsilon} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\
& \leq c(n) \sum_{i, j=0}^{\infty} \left(\frac{1}{2^{i+j}} \right)^{n+q(\alpha+\varepsilon)-2\varepsilon} \sum_{\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j}} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu. \tag{5-68}
\end{aligned}$$

In order to evaluate the last sum we have to further decompose $[\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j}$. For each integer $i \geq 0$, \mathcal{H} contains precisely $4^{ni} = 2^{2ni}$ disjoint cubes from $\Xi_{i+k(\mathcal{H})}$ and exactly 2^{ni} disjoint cubes from $\widetilde{\Xi}_{i+k(\mathcal{H})}$; see the definition in (5-21) and in the preceding display. As a consequence, it contains at most 2^{ni} disjoint (diagonal) cubes from the class $\widetilde{\Xi}_{i+k(\mathcal{H})} \cap (P_1 \mathcal{N}_{\lambda} \cup P_2 \mathcal{N}_{\lambda})$. We anyway consider all the diagonal cubes $\widetilde{\Xi}_{i+k(\mathcal{H})}$ from \mathcal{H} , and relabel them as

$$\{\widetilde{\mathcal{H}} \in \widetilde{\Xi}_{i+k(\mathcal{H})} : \widetilde{\mathcal{H}} \subset \mathcal{H}\} = \{\mathcal{H}_i^m : 1 \leq m \leq 2^{ni}\}, \tag{5-69}$$

so that

$$\sum_{m=1}^{2^{ni}} \int_{\mathcal{H}_i^m \cap \{U > \kappa \lambda\}} U^q d\mu \leq \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu. \tag{5-70}$$

Now, let us concentrate one moment on the elements of $[\mathcal{N}_{\lambda, nd}^1(\mathcal{H})]_{i, j}$; a similar argument then applies to $[\mathcal{N}_{\lambda, nd}^2(\mathcal{H})]_{i, j}$. For any $\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^1(\mathcal{H})]_{i, j}$, there is the unique cube from the diagonal class (5-21), which we denote by $\mathcal{H}_i^m(\mathcal{K})$, such that $P_1 \mathcal{K} = \mathcal{H}_i^m(\mathcal{K})$. Now, note that for $h \in \{1, 2\}$ one can split $[\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j}$ into subsets

$$[\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j, m} := \{\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j} : P_h \mathcal{K} = \mathcal{H}_i^m\}, \quad m \in \{1, \dots, 2^{ni}\}.$$

Since $\mathcal{N}_{\lambda, nd}^1$ is a family of dyadic cubes, if $\mathcal{K}_1, \mathcal{K}_2 \in [\mathcal{N}_{\lambda, nd}^1(\mathcal{H})]_{i, j, m}$ and $\mathcal{K}_1 \neq \mathcal{K}_2$, then $P_2 \mathcal{K}_1 \cap P_2 \mathcal{K}_2 = \emptyset$, i.e., the second components are disjoint (otherwise the two cubes would coincide). A similar argument holds when looking at $\mathcal{N}_{\lambda, nd}^2$. It then follows that

$$\#[\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i, j, m} \leq c(n) 2^{n(i+j)}, \quad h \in \{1, 2\}, \tag{5-71}$$

for every choice of $i, j \geq 0$ and $m \in \{1, \dots, 2^{ni}\}$. We use now use (5-70)–(5-71) to estimate

$$\begin{aligned} \sum_{\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i,j}} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu &= \sum_{m=1}^{2^{ni}} \sum_{\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i,j,m}} \int_{\mathcal{H}_i^m \cap \{U > \kappa \lambda\}} U^q d\mu \\ &\leq c(n) 2^{n(i+j)} \sum_{m=1}^{2^{ni}} \int_{\mathcal{H}_i^m \cap \{U > \kappa \lambda\}} U^q d\mu \\ &\leq c(n) 2^{n(i+j)} \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu. \end{aligned}$$

Using also (2-2) it follows that

$$\begin{aligned} \sum_{i,j=0}^{\infty} \left(\frac{1}{2^{i+j}} \right)^{n+q(\alpha+\varepsilon)-2\varepsilon} \sum_{\mathcal{K} \in [\mathcal{N}_{\lambda, nd}^h(\mathcal{H})]_{i,j}} \int_{P_h \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\ \leq c(n) \sum_{i,j=0}^{\infty} \left(\frac{1}{2^{i+j}} \right)^{q(\alpha+\varepsilon)-2\varepsilon} \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu \\ \leq \frac{c(n)}{[q(\alpha+\varepsilon)-2\varepsilon]^2} \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu \\ \leq \frac{c(n)}{\alpha^2} \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu. \end{aligned}$$

Notice that we have used that, since $q > 1$ and $\varepsilon < \frac{1}{2}\alpha$, we have $q(\alpha+\varepsilon)-2\varepsilon > \frac{1}{2}\alpha$. Combining the inequality in the last display with (5-68) yields (5-67).

Step 3: Summation. Let now $\mathcal{K} \in \mathcal{N}_{\lambda, nd}^1$. There are then two cases: either $\mathcal{K} \in \mathcal{M}_{\lambda}^2$ or $\mathcal{K} \in \mathcal{N}_{\lambda}^2$ (the relevant definitions are in (5-45), (5-46) and (5-58)). Now, if $\mathcal{K} \in \mathcal{M}_{\lambda}^2$, then using (5-44) and (5-52), and reabsorbing terms, we obtain that

$$\mu(\mathcal{K}) \leq \frac{6^q c_{nd}^q}{\lambda^q} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu + \frac{6^q c_{nd}^q}{\varepsilon \lambda^q} \frac{\mu(\mathcal{K})}{\mu(P_1 \mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_1 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu.$$

If, on the other hand, $\mathcal{K} \in \mathcal{N}_{\lambda}^2$, then using (5-44) we get

$$\begin{aligned} \mu(\mathcal{K}) &\leq \frac{3^q c_{nd}^q}{\lambda^q} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu + \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} \frac{\mu(\mathcal{K})}{\mu(P_1 \mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_1 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\ &\quad + \frac{3^q c_{nd}^q}{\varepsilon \lambda^q} \frac{\mu(\mathcal{K})}{\mu(P_2 \mathcal{K})} \left(\frac{2^{-k}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha+\varepsilon)} \int_{P_2 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu. \end{aligned}$$

A similar reasoning holds if $\mathcal{K} \in \mathcal{N}_{\lambda, nd}^2$. Summing up over the cubes $\mathcal{K} \in \mathcal{N}_{\lambda, nd} = \mathcal{N}_{\lambda, nd}^1 \cup \mathcal{N}_{\lambda, nd}^2$ then yields

$$\begin{aligned} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \mu(\mathcal{K}) &\leq \frac{6^q c_{nd}^q}{\lambda^q} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\ &+ \frac{6^q c_{nd}^q}{\varepsilon \lambda^q} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}^1} \frac{\mu(\mathcal{K})}{\mu(P_1 \mathcal{K})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha + \varepsilon)} \int_{P_1 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\ &+ \frac{6^q c_{nd}^q}{\varepsilon \lambda^q} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}^2} \frac{\mu(\mathcal{K})}{\mu(P_2 \mathcal{K})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha + \varepsilon)} \int_{P_2 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu. \end{aligned} \quad (5-72)$$

Observe that a key point in the previous inequality, due to the argument at the beginning of Step 3, is that terms involving integrals over $P_h \mathcal{K}$ appear on the right-hand side if and only if $\mathcal{K} \in \mathcal{N}_{\lambda, nd}^h$, for $h \in \{1, 2\}$. By the symmetry of U and μ , by (5-53) and subsequent remarks, and using Proposition 5.3, we have that if $\mathcal{K} \in \mathcal{N}_{\lambda, nd}^2$, then $\text{Symm}(\mathcal{K}) \in \mathcal{N}_{\lambda, nd}^1$, and vice versa; moreover, again by Proposition 5.3, we have

$$\int_{P_2 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu = \int_{P_1 \text{Symm}(\mathcal{K}) \cap \{U > \kappa \lambda\}} U^q d\mu.$$

Hence the last two terms in (5-72) coincide. Therefore, also recalling (5-64), (5-72) can be rewritten as

$$\begin{aligned} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \mu(\mathcal{K}) &\leq \frac{c}{\lambda^q} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \\ &+ \frac{c}{\varepsilon \lambda^q} \sum_{\mathcal{H} \in P \mathcal{N}_{\lambda}} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}^1(\mathcal{H})} \frac{\mu(\mathcal{K})}{\mu(P_1 \mathcal{K})} \left(\frac{2^{-k(\mathcal{K})}}{\widetilde{\text{dist}}(P_1 \mathcal{K}, P_2 \mathcal{K})} \right)^{q(\alpha + \varepsilon)} \int_{P_1 \mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu \end{aligned}$$

for a constant c depending on n, α . To estimate the last term we make use of (5-67), and this yields

$$\sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \mu(\mathcal{K}) \leq \frac{c}{\lambda^q} \sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu + \frac{c}{\lambda^q} \sum_{\mathcal{H} \in P \mathcal{N}_{\lambda}} \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu.$$

At this stage (5-63) follows, observing that

$$\sum_{\mathcal{K} \in \mathcal{N}_{\lambda, nd}} \int_{\mathcal{K} \cap \{U > \kappa \lambda\}} U^q d\mu + \sum_{\mathcal{H} \in P \mathcal{N}_{\lambda}} \int_{\mathcal{H} \cap \{U > \kappa \lambda\}} U^q d\mu \leq 2 \int_{\mathcal{B}(x_0, s) \cap \{U > \kappa \lambda\}} U^q d\mu.$$

This is in turn true since the families $P \mathcal{N}_{\lambda}$ and $\mathcal{N}_{\lambda, nd}$ are made of mutually disjoint cubes and all their members are contained in $\mathcal{B}(x_0, s)$ (since these families are contained in Ξ and (5-22) holds). The proof of Lemma 5.12 is complete. \square

5J. Conclusion of the off-diagonal analysis. The next lemma summarizes the decomposition results in the off-diagonal case:

Lemma 5.13 (off-diagonal level set inequality). *The inequality*

$$\int_{\mathcal{B}(x_0,t) \cap \{U > \lambda\}} U^2 d\mu \leq 10^{n+2} \kappa^2 \lambda^2 \sum_{j \in J_D} \mu(\mathcal{B}_j) + c \lambda^{2-q} \int_{\mathcal{B}(x_0,s) \cap \{U > \kappa \lambda\}} U^q d\mu \quad (5-73)$$

holds for a constant c depending only on n, α , while the number κ has been defined in (5-55) and exhibits the dependence displayed in (5-56).

Proof. We have the decompositions in disjoint classes $\mathcal{U}_\lambda = \mathcal{U}_\lambda^d \cup \mathcal{U}_\lambda^{nd}$ and $\mathcal{U}_\lambda^{nd} = \mathcal{M}_\lambda \cup \mathcal{N}_{\lambda,d} \cup \mathcal{N}_{\lambda,nd}$, and we recall that all the cubes from \mathcal{U}_λ^{nd} are mutually disjoint. Moreover, by (5-38) and (5-57) it follows that

$$\bigcup_{\mathcal{K} \in \mathcal{U}_\lambda^d} \mathcal{K} \cup \bigcup_{\mathcal{K} \in \mathcal{N}_{\lambda,d}} \mathcal{K} \subset \bigcup_{j \in J_D} 10\mathcal{B}_j.$$

Therefore

$$\bigcup_{\mathcal{K} \in \mathcal{U}_\lambda} \mathcal{K} \subset \bigcup_{j \in J_D} 10\mathcal{B}_j \cup \bigcup_{\mathcal{K} \in \mathcal{M}_\lambda} \mathcal{K} \cup \bigcup_{\mathcal{K} \in \mathcal{N}_{\lambda,nd}} \mathcal{K}.$$

Keeping this in mind, and recalling (5-35), we start by estimating

$$\int_{\mathcal{B}(x_0,t) \cap \{U > \lambda\}} U^2 d\mu \leq \sum_j \int_{10\mathcal{B}_j \cap \{U > \lambda\}} U^2 d\mu + \sum_{\mathcal{K} \in \mathcal{M}_\lambda \cup \mathcal{N}_{\lambda,nd}} \int_{\mathcal{K} \cap \{U > \lambda\}} U^2 d\mu.$$

By (5-34) it follows that, if $\mathcal{K} \in \mathcal{M}_\lambda \cup \mathcal{N}_{\lambda,nd} \subset \mathcal{U}_\lambda^{nd}$, then

$$\int_{\mathcal{K}} U^2 d\mu \leq \frac{\mu(\tilde{\mathcal{K}})}{\mu(\mathcal{K})} \int_{\tilde{\mathcal{K}}} U^2 d\mu \leq \tilde{c}_d \lambda^2.$$

Note that we have used (5-32), since $\mathcal{K} \in \mathcal{U}_\lambda^{nd}$ implies by the definition in (5-39) that $\text{dist}(P_1 \tilde{\mathcal{K}}, P_2 \tilde{\mathcal{K}}) \geq 2^{-k(\mathcal{K})}$. Therefore we conclude that

$$\mathcal{K} \in \mathcal{M}_\lambda \cup \mathcal{N}_{\lambda,nd} \implies \int_{\mathcal{K} \cap \{U > \lambda\}} U^2 d\mu \leq \tilde{c}_d \lambda^2 \mu(\mathcal{K}).$$

Using this last inequality together with (5-16) yields

$$\int_{\mathcal{B}(x_0,t) \cap \{U > \lambda\}} U^2 d\mu \leq 10^{n+2\epsilon} \kappa^2 \lambda^2 \sum_{j \in J_D} \mu(\mathcal{B}_j) + \tilde{c}_d \lambda^2 \sum_{\mathcal{K} \in \mathcal{M}_\lambda \cup \mathcal{N}_{\lambda,nd}} \mu(\mathcal{K}),$$

and (5-73) follows by just using Lemmas 5.10 and 5.12. \square

Remark 5.14. An interesting point of Lemma 5.13 is that it does not make use of the fact that u is a solution. All the estimates just rely on the fact that u belongs to the Sobolev space $W^{\alpha,2}$. This is ultimately linked to the fact that the analysis in Sections 5B–5J is made in a zone where the kernel of the operator, that is, $|x - y|^{-(n+2\alpha)}$, is not very singular. The ultimate outcome is that the whole issue reduces to

estimating $\sum_j \mu(\mathcal{B}_j)$. Therefore, it remains to perform the analysis close to the diagonal, and this will be done in the next section.

5K. Diagonal estimates. Whenever \mathcal{B}_j is a ball from the covering determined in (5-14)–(5-15), from (5-13) it follows that $\Psi_{H,M}(\mathcal{B}_j) \geq \kappa\lambda$. By the very definition of $\Psi_{H,M}(\cdot)$ in (5-1) it then follows that at least one of the following three inequalities must hold:

$$\left(\int_{\mathcal{B}_j} U^2 d\mu \right)^{1/2} \geq \frac{\kappa\lambda}{3}, \quad (5-74)$$

$$\frac{H[\mu(\mathcal{B}_j)]^\eta}{\varepsilon^{1/2_*-1/2}} \left(\int_{\mathcal{B}_j} F^{2_*} d\mu \right)^{1/2_*} \geq \frac{\kappa\lambda}{3}, \quad (5-75)$$

$$\frac{M[\mu(\mathcal{B}_j)]^\theta}{\varepsilon^{1/p-1/2}} \left(\int_{\mathcal{B}_j} G^p d\mu \right)^{1/p} \geq \frac{\kappa\lambda}{3}, \quad (5-76)$$

where κ has been defined in (5-55). We now examine the occurrence of each of the three cases separately.

Occurrence of (5-74) (and estimate of the tail at the exit time). When (5-74) holds, using (4-15) we have

$$\begin{aligned} \kappa\lambda &\leq \frac{c}{\sigma \varepsilon^{1/q-1/2}} \left(\int_{2\mathcal{B}_j} U^q d\mu \right)^{1/q} \\ &+ \frac{\sigma}{\varepsilon^{1/q-1/2}} \sum_{k=1}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}_j} U^q d\mu \right)^{1/q} + \frac{c_1 [\mu(\mathcal{B}_j)]^\eta}{\varepsilon^{1/2_*-1/2}} \left(\int_{2\mathcal{B}_j} F^{2_*} d\mu \right)^{1/2_*} \\ &+ \frac{c_2 [\mu(\mathcal{B}_j)]^\theta}{\varepsilon^{1/p-1/2}} \sum_{k=1}^{\infty} 2^{-k(2\beta-\gamma-2\varepsilon/p)} \left(\int_{2^k \mathcal{B}_j} G^p d\mu \right)^{1/p} \end{aligned} \quad (5-77)$$

for all $\sigma \in (0, 1]$. The constants c_1, c depend only on n, α, Λ , while $c_2 := 3c_b$ and therefore it depends on $n, \alpha, \Lambda, \beta, \gamma, p$ and exhibits the behavior described in (3-5). With $\mathcal{B}_j \equiv \mathcal{B}(x_j, \varrho(x_j))$ we determine the integer $m \geq 0$ such that

$$2^{-m} \varrho_0/2 \leq \varrho(x_j) < 2^{-m+1} \varrho_0/2. \quad (5-78)$$

Notice that since $\varrho(x_j) < (s-t)/40^n$, we have $m \geq 3$ and moreover $(s-t)/40^n \leq \varrho_0/40^n \leq 2^{m-1} \varrho(x_j)$, so that (5-10) implies

$$\Upsilon_0(2^{m-1} \mathcal{B}_j) + \Upsilon_1(2^{m-1} \mathcal{B}_j) + \Upsilon_{2,M}(2^{m-1} \mathcal{B}_j) \leq \kappa \tilde{\lambda}_0. \quad (5-79)$$

On the other hand, the terms indexed before m can be estimated using Hölder's inequality and the exit-time condition in (5-13) as

$$\left(\int_{2^k \mathcal{B}_j} U^q d\mu \right)^{1/q} \leq \Psi_{H,M}(2^k \mathcal{B}_j) \leq \kappa\lambda \quad \text{if } 1 \leq k \leq m-1.$$

By using the inequalities in the last two displays we then have

$$\begin{aligned}
\sum_{k=1}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}_j} U^q d\mu \right)^{1/q} &= \sum_{k=1}^{m-2} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}_j} U^q d\mu \right)^{1/q} \\
&\quad + 2^{-(m-1)(\alpha-\varepsilon)} \sum_{k=0}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^{k+m-1} \mathcal{B}_j} U^q d\mu \right)^{1/q} \\
&\leq \kappa \lambda \sum_{k=1}^{m-2} 2^{-k(\alpha-\varepsilon)} + 2^{-(m-1)(\alpha-\varepsilon)} \Upsilon_1(2^{m-1} \mathcal{B}_j) \\
&\leq \kappa \lambda \sum_{k=1}^{m-2} 2^{-k(\alpha-\varepsilon)} + 2^{-(m-1)(\alpha-\varepsilon)} \kappa \tilde{\lambda}_0 \\
&\leq \kappa \lambda \sum_{k=1}^{\infty} 2^{-k(\alpha-\varepsilon)} \leq \frac{4\kappa\lambda}{\alpha-\varepsilon} \leq \frac{8\kappa\lambda}{\alpha},
\end{aligned}$$

where we have used (2-2) and that $\varepsilon < \frac{1}{2}\alpha$. In a completely similar way, again using (5-79), we have

$$\frac{c_2[\mu(\mathcal{B}_j)]^\theta}{\varepsilon^{1/p-1/2}} \sum_{k=1}^{\infty} 2^{-k(2\beta-\gamma-2\varepsilon/p)} \left(\int_{2^k \mathcal{B}_j} G^p d\mu \right)^{1/p} \leq \frac{4c_2\kappa\lambda}{(2\beta-\gamma-2\varepsilon/p)M} \leq \frac{8c_2\kappa\lambda}{(2\beta-\gamma)M},$$

where we also used the upper bound on ε in (4-6). By (5-78) and the fact that $m \geq 3$ we gain that $2\varrho(x_j) \leq \frac{1}{2}\varrho_0$ so that (5-13) and Hölder's inequality yield

$$\frac{c_1[\mu(\mathcal{B}_j)]^\eta}{\varepsilon^{1/2_*-1/2}} \left(\int_{2\mathcal{B}_j} F^{2_*} d\mu \right)^{1/2_*} \leq \frac{c_1\Psi_{H,M}(2\mathcal{B}_j)}{H} \leq \frac{c_1\kappa\lambda}{H}.$$

By merging the inequalities in the last three displays with (5-77) we obtain

$$\kappa\lambda \leq \frac{c}{\sigma\varepsilon^{1/q-1/2}} \left(\int_{2\mathcal{B}_j} U^q d\mu \right)^{1/q} + \frac{\sigma}{\varepsilon^{1/q-1/2}} \frac{8\kappa\lambda}{\alpha} + \frac{c_1\kappa\lambda}{H} + \frac{8c_2\kappa\lambda}{(2\beta-\gamma)M}. \quad (5-80)$$

We recall that up to now the parameters $H, M \geq 1$ in the definition in (5-1) have not yet been chosen, and neither has $\sigma \in (0, 1)$. Hence, taking

$$\sigma := \frac{\varepsilon^{1/q-1/2}\alpha}{56}, \quad H := 6c_1, \quad M := \frac{56c_2}{2\beta-\gamma}, \quad (5-81)$$

and reabsorbing terms in (5-80), we conclude that

$$\kappa\lambda \leq \frac{c}{\varepsilon^{2/q-1}} \left(\int_{2\mathcal{B}_j} U^q d\mu \right)^{1/q} \implies \mu(\mathcal{B}_j) \leq \frac{c}{\varepsilon^{2-q}(\kappa\lambda)^q} \int_{2\mathcal{B}_j} U^q d\mu,$$

where c depends on n, α, Λ . Now, select a number $\kappa_3 > 0$; also using (4-2), we estimate

$$\begin{aligned} \frac{c}{\varepsilon^{2-q}(\kappa\lambda)^q} \int_{2\mathcal{B}_j} U^q d\mu &\leq \frac{c}{\varepsilon^{2-q}(\kappa\lambda)^q} \int_{2\mathcal{B}_j \cap \{U \leq \kappa_3 \kappa \lambda\}} U^q d\mu + \frac{c}{\varepsilon^{2-q}(\kappa\lambda)^q} \int_{2\mathcal{B}_j \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu \\ &\leq \frac{\tilde{c} \mu(\mathcal{B}_j) \kappa_3^q}{\varepsilon^{2-q}} + \frac{\tilde{c}}{\varepsilon^{2-q}(\kappa\lambda)^q} \int_{2\mathcal{B}_j \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu, \end{aligned} \quad (5-82)$$

again for \tilde{c} depending only on n, α, Λ . By choosing

$$\kappa_3 \leq \left(\frac{\varepsilon^{2-q}}{2\tilde{c}} \right)^{1/q}, \quad (5-83)$$

we arrive at

$$\mu(\mathcal{B}_j) \leq \frac{c_3}{(\kappa\lambda)^q} \int_{2\mathcal{B}_j \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu, \quad \text{where } c_3 := \frac{2\tilde{c}}{\varepsilon^{2-q}}, \quad (5-84)$$

and \tilde{c} is independent of ε and only depends on n, α, Λ .

Occurrence of (5-75)–(5-76). In case of (5-75), we have

$$\left(\frac{\kappa\lambda}{3} \right)^{2_*} \leq \frac{H^{2_*} [\mu(\mathcal{B}_j)]^{2_*\eta-1}}{\varepsilon^{1-2_*/2}} \int_{\mathcal{B}_j} F^{2_*} d\mu,$$

which readily implies

$$\mu(\mathcal{B}_j) \leq \left(\frac{3H}{\varepsilon^{1/2_*-1/2} \kappa\lambda} \right)^{2_*/(1-2_*\eta)} \left(\int_{\mathcal{B}_j} F^{2_*} d\mu \right)^{1/(1-2_*\eta)}.$$

Observe that by the definitions given in (4-16) we have that $2_*\eta < \frac{1}{2}$. With $\kappa_4 \in (0, 1)$ being a positive number to be chosen in a few lines, we further split the support of the right-hand side integral as already done in (5-82):

$$\begin{aligned} \left(\int_{\mathcal{B}_j} F^{2_*} d\mu \right)^{1/(1-2_*\eta)} &\leq \left[\int_{\mathcal{B}_j \cap \{F > \kappa_4 \kappa \lambda\}} F^{2_*} d\mu + (\kappa_4 \kappa \lambda)^{2_*} \mu(\mathcal{B}_j) \right]^{1/(1-2_*\eta)} \\ &\leq 2^{2_*\eta/(1-2_*\eta)} \left(\int_{\mathcal{B}_j \cap \{F > \kappa_4 \kappa \lambda\}} F^{2_*} d\mu \right)^{1/(1-2_*\eta)} \\ &\quad + [2(L+1)]^{2_*/(1-2_*\eta)} (\kappa_4 \kappa \lambda)^{2_*/(1-2_*\eta)} \mu(\mathcal{B}_j). \end{aligned}$$

Observe that, in view of $\mathcal{B}_j \subset \mathcal{B}(x_0, 2\varrho_0)$ and (5-18), we have estimated

$$[\mu(\mathcal{B}_j)]^{1/(1-2_*\eta)} \leq [\mu(\mathcal{B}(x_0, 2\varrho_0))]^{2_*\eta/(1-2_*\eta)} \mu(\mathcal{B}_j) \leq L^{2_*\eta/(1-2_*\eta)} \mu(\mathcal{B}_j). \quad (5-85)$$

We now take $\kappa_4 \in (0, 1)$ in order to satisfy

$$\left[\frac{6H(L+1)\kappa_4}{\varepsilon^{1/2_*-1/2}} \right]^{2_*/(1-2_*\eta)} \leq \frac{1}{2} \implies \kappa_4 \leq \left(\frac{1}{2} \right)^{(1-2_*\eta)/2_*} \frac{\varepsilon^{1/2_*-1/2}}{6H(L+1)}. \quad (5-86)$$

Using this choice and combining the content of the last four displays (and recalling that $2_*\eta/(1-2_*\eta) \leq 1$) then yields that

$$\mu(\mathcal{B}_j) \leq 4 \left(\frac{3H}{\varepsilon^{1/2_*-1/2}\kappa\lambda} \right)^{2_*/(1-2_*\eta)} \left(\int_{\mathcal{B}_j \cap \{F > \kappa_4\kappa\lambda\}} F^{2_*} d\mu \right)^{1/(1-2_*\eta)}.$$

Now, by means of (5-78)–(5-79), we have

$$\begin{aligned} \int_{\mathcal{B}_j \cap \{F > \kappa_4\kappa\lambda\}} F^{2_*} d\mu &\leq (\kappa_4\kappa\lambda)^{2_*} \int_{\mathcal{B}_j \cap \{F > \kappa_4\kappa\lambda\}} \left(\frac{F}{\kappa_4\kappa\lambda} \right)^{2_*+\delta_f} d\mu \\ &\leq \frac{\mu(2^{m-1}\mathcal{B}_j)}{(\kappa_4\kappa\lambda)^{\delta_f}} \int_{2^{m-1}\mathcal{B}_j} F^{2_*+\delta_f} d\mu \\ &\leq \frac{\mu(\mathcal{B}(x_0, 2Q_0))}{(\kappa_4\kappa\lambda)^{\delta_f}} [\Upsilon_0(2^{m-1}\mathcal{B}_j)]^{2_*+\delta_f} \leq \frac{L\tilde{\lambda}_0^{2_*+\delta_f}}{(\kappa_4\kappa\lambda)^{\delta_f}}, \end{aligned} \quad (5-87)$$

and hence

$$\mu(\mathcal{B}_j) \leq \frac{c_4\tilde{\lambda}_0^{(2_*+\delta_f)2_*/(1-2_*\eta)}}{(\kappa_4\kappa\lambda)^{(1+\eta\delta_f)2_*/(1-2_*\eta)}} \int_{\mathcal{B}_j \cap \{F > \kappa_4\kappa\lambda\}} F^{2_*} d\mu, \quad (5-88)$$

where

$$c_4 := 4 \left[\frac{3H(L+1)}{\varepsilon^{1/2_*-1/2}} \right]^{2_*/(1-2_*\eta)}, \quad (5-89)$$

and H has been defined in (5-81). A similar argument can be used in case (5-76) holds. Specifically, we have

$$\mu(\mathcal{B}_j) \leq \left(\frac{3M}{\varepsilon^{1/p-1/2}\kappa\lambda} \right)^{p/(1-p\theta)} \left(\int_{\mathcal{B}_j} G^p d\mu \right)^{1/(1-p\theta)},$$

and then

$$\left(\int_{\mathcal{B}_j} G^p d\mu \right)^{\frac{1}{1-p\theta}} \leq 2^{p\theta/(1-p\theta)} \left(\int_{\mathcal{B}_j \cap \{G > \kappa_5\kappa\lambda\}} G^p d\mu \right)^{\frac{1}{1-p\theta}} + [2(L+1)]^{p/(1-p\theta)} (\kappa_5\kappa\lambda)^{p/(1-p\theta)} \mu(\mathcal{B}_j).$$

This time we select a number $\kappa_5 \in (0, 1)$ such that

$$\kappa_5 \leq \left(\frac{1}{2} \right)^{(1-p\theta)/p} \frac{\varepsilon^{1/p-1/2}}{6M(L+1)} \quad (5-90)$$

and recall Remark 4.5 in order to get

$$\mu(\mathcal{B}_j) \leq 2^{\Lambda_\theta+1} \left(\frac{3M}{\varepsilon^{1/p-1/2}\kappa\lambda} \right)^{p/(1-p\theta)} \left(\int_{\mathcal{B}_j \cap \{G > \kappa_5\kappa\lambda\}} G^p d\mu \right)^{1/(1-p\theta)}.$$

We then estimate as in (5-87), thereby obtaining

$$\int_{\mathcal{B}_j \cap \{G > \kappa_5\kappa\lambda\}} G^p d\mu \leq \frac{\mu(\mathcal{B}(x_0, 2Q_0))}{(\kappa_5\kappa\lambda)^{\delta_g}} [\Upsilon_0(2^{m-1}\mathcal{B}_j)]^{p+\delta_g} \leq \frac{L\tilde{\lambda}_0^{p+\delta_g}}{(\kappa_5\kappa\lambda)^{\delta_g}},$$

and we conclude that

$$\mu(\mathcal{B}_j) \leq \frac{c_5 \tilde{\lambda}_0^{(p+\delta_g)p\theta/(1-p\theta)}}{(\kappa_5 \kappa \lambda)^{(1+\theta\delta_g)p/(1-p\theta)}} \int_{\mathcal{B}_j \cap \{G > \kappa_5 \kappa \lambda\}} G^p d\mu, \quad (5-91)$$

where

$$c_5 := 2^{\Lambda_\theta + 1} \left[\frac{3M(L+1)}{\varepsilon^{1/p-1/2}} \right]^{p/(1-p\theta)}. \quad (5-92)$$

All in all, taking (5-84), (5-88) and (5-91) into account, we obtain

$$\begin{aligned} \mu(\mathcal{B}_j) \leq & \frac{c_3}{(\kappa \lambda)^q} \int_{2\mathcal{B}_j \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu + \frac{c_4 \tilde{\lambda}_0^{(2_* + \delta_f)2_*\eta/(1-2_*\eta)}}{(\kappa_4 \kappa \lambda)^{(1+\eta\delta_f)2_*/(1-2_*\eta)}} \int_{\mathcal{B}_j \cap \{F > \kappa_4 \kappa \lambda\}} F^{2_*} d\mu \\ & + \frac{c_5 \tilde{\lambda}_0^{(p+\delta_g)p\theta/(1-p\theta)}}{(\kappa_5 \kappa \lambda)^{(1+\theta\delta_g)p/(1-p\theta)}} \int_{\mathcal{B}_j \cap \{G > \kappa_5 \kappa \lambda\}} G^p d\mu. \end{aligned}$$

Since $\{2\mathcal{B}_j\}_j$ is a disjoint family and all members belong to $\mathcal{B}(x_0, s)$, we have that

$$\begin{aligned} \sum_{j \in J_D} \mu(\mathcal{B}_j) \leq & \frac{c_3}{(\kappa \lambda)^q} \int_{\mathcal{B}(x_0, s) \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu + \frac{c_4 \tilde{\lambda}_0^{(2_* + \delta_f)2_*\eta/(1-2_*\eta)}}{(\kappa_4 \kappa \lambda)^{(1+\eta\delta_f)2_*/(1-2_*\eta)}} \int_{\mathcal{B}(x_0, s) \cap \{F > \kappa_4 \kappa \lambda\}} F^{2_*} d\mu \\ & + \frac{c_5 \tilde{\lambda}_0^{(p+\delta_g)p\theta/(1-p\theta)}}{(\kappa_5 \kappa \lambda)^{(1+\theta\delta_g)p/(1-p\theta)}} \int_{\mathcal{B}(x_0, s) \cap \{G > \kappa_5 \kappa \lambda\}} G^p d\mu. \end{aligned} \quad (5-93)$$

The constants c_3, c_4, c_5 have been defined in (5-84), (5-89) and (5-92), respectively, while the numbers $\kappa, \kappa_3, \kappa_4, \kappa_5 \in (0, 1)$ must be taken in order to satisfy (5-55), (5-83), (5-86) and (5-90), respectively.

5L. Conclusion of the proof. We start by combining (5-73) and (5-93). Using the elementary estimate

$$\int_{\mathcal{B}(x_0, t) \cap \{U > \kappa_3 \kappa \lambda\}} U^2 d\mu \leq \lambda^{2-q} \int_{\mathcal{B}(x_0, t) \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu + \int_{\mathcal{B}(x_0, t) \cap \{U > \lambda\}} U^2 d\mu,$$

(5-73) and (5-93) yield, after a few elementary manipulations, the estimate

$$\begin{aligned} \int_{\mathcal{B}(x_0, t) \cap \{U > \kappa_3 \kappa \lambda\}} U^2 d\mu \leq & \frac{c}{(\kappa_3 \kappa)^{2-q}} (\kappa_3 \kappa \lambda)^{2-q} \int_{\mathcal{B}(x_0, s) \cap \{U > \kappa_3 \kappa \lambda\}} U^q d\mu \\ & + \frac{c_4 \tilde{\lambda}_0^{(2_* + \delta_f)2_*\eta/(1-2_*\eta)}}{\kappa_4^2 (\kappa_4 \kappa \lambda)^{(1+\eta\delta_f)2_*/(1-2_*\eta)-2}} \int_{\mathcal{B}(x_0, s) \cap \{F > \kappa_4 \kappa \lambda\}} F^{2_*} d\mu \\ & + \frac{c_5 \tilde{\lambda}_0^{(p+\delta_g)p\theta/(1-p\theta)}}{\kappa_5^2 (\kappa_5 \kappa \lambda)^{(1+\theta\delta_g)p/(1-p\theta)-2}} \int_{\mathcal{B}(x_0, s) \cap \{G > \kappa_5 \kappa \lambda\}} G^p d\mu. \end{aligned} \quad (5-94)$$

The constant c appearing above depends on n, α, Λ , but is still independent of ε , and we have also used the fact that $\kappa, \kappa_3 \in (0, 1)$. We can therefore reformulate estimate (5-94) as

$$\begin{aligned} \int_{\mathcal{B}(x_0, t) \cap \{U > \lambda\}} U^2 d\mu &\leq \frac{c\lambda^{2-q}}{(\kappa_3 \kappa \varepsilon)^{2-q}} \int_{\mathcal{B}(x_0, s) \cap \{U > \lambda\}} U^q d\mu + \frac{c_6 \tilde{\lambda}_0^{(2_* + \delta_f) 2_* \eta / (1 - 2_* \eta)}}{\lambda^{(1 + \eta \delta_f) 2_* / (1 - 2_* \eta) - 2}} \int_{\mathcal{B}(x_0, s) \cap \{F > \kappa_4 \lambda / \kappa_3\}} F^{2_*} d\mu \\ &\quad + \frac{c_7 \tilde{\lambda}_0^{(p + \delta_g) p \theta / (1 - p \theta)}}{\lambda^{(1 + \theta \delta_g) p / (1 - p \theta) - 2}} \int_{\mathcal{B}(x_0, s) \cap \{G > \kappa_5 \lambda / \kappa_3\}} G^p d\mu. \end{aligned} \quad (5-95)$$

The constant $c \equiv c(n, \alpha, \Lambda)$ is independent of ε , while

$$c_6 \equiv c_6(n, \alpha, \Lambda, L, \varepsilon) \quad \text{and} \quad c_7 \equiv c_7(n, \alpha, \Lambda, \beta, \gamma, p, L, \varepsilon);$$

the constant c_7 exhibits a blow-up behavior with respect to p as described in (3-5). Since estimate (5-94) holds for $\lambda \geq \lambda_1$ — and λ_1 has been defined in (5-33) — we have that (5-95) holds whenever $\lambda \geq \kappa \kappa_3 \lambda_1$. We remark that the previous inequality holds for a choice of $\kappa, \kappa_3, \kappa_4, \kappa_5 \in (0, 1)$ that satisfy (5-55), (5-83), (5-86) and (5-90), respectively. In order to conclude with (5-7) we now need to estimate a few constants. We are primarily interested in an explicit dependence on ε in the second integral appearing in (5-95). We therefore look at (5-55) and (5-83), and we infer that we can in fact choose κ, κ_3 in order to have

$$\kappa_3 \kappa \approx \frac{\varepsilon^{3/q-1}}{c_*}, \quad (5-96)$$

for a constant c_* which is now independent of ε , but just depends on n, α, Λ . We next find an upper bound for the numbers $\tilde{\lambda}_0$ and λ_1 introduced in (5-10) and (5-33), respectively; this will allow us to verify estimate (5-7) in the range dictated by (5-8). Let us notice that if $x \in \mathcal{B}(x_0, t)$ and $(s - t)/40^n \leq \varrho \leq \frac{1}{2}\varrho_0$, then $\mathcal{B}(x, \varrho) \subset \mathcal{B}(x_0, 2\varrho_0)$. Therefore, recalling (4-2), whenever \tilde{U} is a μ -integrable function we can estimate

$$\int_{\mathcal{B}(x, \varrho)} \tilde{U} d\mu \leq \frac{\mu(\mathcal{B}(x_0, 2\varrho_0))}{\mu(\mathcal{B}(x, \varrho))} \int_{\mathcal{B}(x_0, 2\varrho_0)} \tilde{U} d\mu \leq c \left(\frac{\varrho_0}{s - t} \right)^{n+2\varepsilon} \int_{\mathcal{B}(x_0, 2\varrho_0)} \tilde{U} d\mu \quad (5-97)$$

for a constant c depending on n but independent of ε . Applying the inequality in the last display to $U^2, G^p, F^{2_*}, G^{p+\delta_g}$ and $F^{2_*+\delta_f}$ — and eventually on different balls $2^k \mathcal{B}(x, \varrho) \subset 2^k \mathcal{B}(x_0, 2\varrho_0)$ — yields

$$\begin{aligned} &\kappa^{-1} \{ \Psi_{H,M}(x, \varrho) + \Upsilon_0(x, \varrho) + \Upsilon_1(x, \varrho) + \Upsilon_{2,M}(x, \varrho) \} \\ &\leq \frac{c}{\varepsilon^{1/q}} \left(\frac{\varrho_0}{s - t} \right)^{n+2\varepsilon} \{ \Psi_{H,M}(x_0, 2\varrho_0) + \Upsilon_0(x_0, 2\varrho_0) + \Upsilon_1(x_0, 2\varrho_0) + \Upsilon_{2,M}(x_0, 2\varrho_0) \} \\ &\leq \frac{c}{\varepsilon^{1/q}} \left(\frac{\varrho_0}{s - t} \right)^{n+2\varepsilon} \text{ADD}(x_0, 2\varrho_0). \end{aligned} \quad (5-98)$$

In order, we have also used (5-56), (5-81) to get rid of the presence of M and H and that $\varrho_0/(s - t)$ is bounded away from zero. We recall that the functional $\text{ADD}(\cdot)$ has been introduced in (5-5). We now obtain an upper bound for λ_1 defined in (5-33). The quantity appearing on the right-hand side of (5-98) provides an upper bound on $\tilde{\lambda}_0$. In a similar way, if $\mathcal{K} = K_1 \times K_2 \in \mathfrak{E}_{k_0}$, with k_0 as in (5-19), then

$\mathcal{K} \subset \mathcal{B}(x_0, s) \subset \mathcal{B}(x_0, 2\varrho_0)$ and therefore we have

$$\mu(\mathcal{K}) \geq \frac{c}{\varrho_0^{n-2\varepsilon}} \int_{K_1} \int_{K_2} dx dy = \frac{c(s-t)^{2n}}{\varrho_0^{n-2\varepsilon}}.$$

Hence, as for (5-97), we have

$$\int_{\mathcal{K}} \tilde{U} d\mu \leq \frac{\mu(\mathcal{B}(x_0, 2\varrho_0))}{\mu(\mathcal{K})} \int_{\mathcal{B}(x_0, 2\varrho_0)} \tilde{U} d\mu \leq \frac{c}{\varepsilon} \left(\frac{\varrho_0}{s-t} \right)^{2n} \int_{\mathcal{B}(x_0, 2\varrho_0)} \tilde{U} d\mu. \quad (5-99)$$

By using (5-98)-(5-99), and recalling that $\varepsilon < 1$, we get

$$\lambda_1 \leq \frac{c}{\varepsilon} \left(\frac{\varrho_0}{s-t} \right)^{2n} \text{ADD}(x_0, 2\varrho_0)$$

where c depends only on $n, \alpha, \Lambda, \beta, p, \gamma, \varepsilon$. Summarizing the content of these manipulations, we can finally arrive at (5-7), with the restriction on λ described in (5-8). Specifically, we use (5-96) to estimate the constant in front of the second integral appearing in (5-95), and the bounds found for $\tilde{\lambda}_0$ and λ_1 to conclude with the admissible range of values $\lambda \geq \lambda_0$ described via (5-8). Needless to say, we are taking $\kappa_f := \kappa_4/\kappa_3$ and $\kappa_g := \kappa_5/\kappa_3$.

6. Self-improving inequalities

This section is dedicated to the proof of a fractional reverse Hölder-type inequality on diagonal balls with increasing supports, that is, the estimate (6-1) below. This will eventually imply Theorem 1.1 at the end of the section.

Theorem 6.1 (reverse Hölder-type inequality). *Let $u \in W^{\alpha,2}(\mathbb{R}^n)$ be a solution to (1-14) under the assumptions of Theorem 1.1; in particular, (3-1) and (3-3) are in force. Define the functions U, F and G as in (4-5). Then there exist positive constants $\varepsilon \in (0, 1 - \alpha)$, $\delta \in (0, 1)$ and $c_8 \geq 1$, depending on $n, \alpha, \Lambda, \beta, p, \gamma, \delta_1$, such that whenever $\mathcal{B} \equiv \mathcal{B}(x_0, \varrho_0) \subset \mathbb{R}^{2n}$ we have the inequality*

$$\begin{aligned} \left(\int_{\mathcal{B}} U^{2+\delta} d\mu \right)^{1/(2+\delta)} &\leq c_8 \sum_{k=1}^{\infty} 2^{-k(\alpha-\varepsilon)} \left(\int_{2^k \mathcal{B}} U^2 d\mu \right)^{1/2} \\ &+ c_8 \varrho_0^{\alpha-\varepsilon} \left(\int_{2\mathcal{B}} F^{2_*+\delta_0} d\mu \right)^{1/(2_*+\delta_0)} + c_8 \varrho_0^{\gamma-2\beta+\alpha+\varepsilon(2/p-1)} \left(\int_{2\mathcal{B}} G^{p(1+\delta_1)} d\mu \right)^{1/[p(1+\delta_1)]} \\ &+ c_8 \varrho_0^{\gamma-2\beta+\alpha+\varepsilon(2/p-1)} \sum_{k=1}^{\infty} 2^{-k(2\beta-\gamma-2\varepsilon/p)} \left(\int_{2^k \mathcal{B}} G^p d\mu \right)^{1/p}. \end{aligned} \quad (6-1)$$

All the terms on the right-hand side of this inequality are finite.

Proof. Step 1: Determining the exponents. Let us observe that, whenever $\varepsilon \in (0, \frac{1}{2}\alpha)$, we have

$$\frac{8\varepsilon}{n+2\varepsilon} < \frac{2\varepsilon(n+2\alpha)}{n(\alpha-\varepsilon)}.$$

Therefore, we can always find two positive numbers $\varepsilon \in (0, \frac{1}{2}\alpha)$ and $\delta_f > 0$, satisfying (4-6) and $\delta_f \leq \delta_0$, respectively, such that

$$\frac{8\varepsilon}{n+2\varepsilon} < \delta_f \leq \frac{2\varepsilon(n+2\alpha)}{n(\alpha-\varepsilon)} \quad \text{and} \quad \varepsilon < 1-\alpha. \tag{6-2}$$

We recall that $F \in L_{\text{loc}}^{2_*+\delta_f}(\mathbb{R}^n; \mu)$ by (4-7). Next, we determine the positive number $\delta > 0$ by imposing different restrictions on it; we start by assuming that

$$\delta \leq \frac{4\varepsilon(n+2\alpha)}{n^2+4\varepsilon(n+\alpha)} \quad \text{and} \quad \delta \leq \frac{(\gamma-2\beta+\alpha)\delta_g}{4n}. \tag{6-3}$$

Let us briefly discuss a few consequences of the two conditions above, starting with the first one. Specifically, we start by showing that

$$\delta \leq \delta_f \left[\frac{(n+2\alpha)(n+2\varepsilon)}{n^2+4\varepsilon(n+\alpha)} \right] - \frac{4\varepsilon(n+2\alpha)}{n^2+4\varepsilon(n+\alpha)}. \tag{6-4}$$

Indeed, using the first inequality in (6-3), we have

$$\begin{aligned} \delta &\leq \frac{4\varepsilon(n+2\alpha)}{n^2+4\varepsilon(n+\alpha)} = \frac{8\varepsilon}{n+2\varepsilon} \frac{(n+2\alpha)(n+2\varepsilon)}{n^2+4\varepsilon(n+\alpha)} - \frac{4\varepsilon(n+2\alpha)}{n^2+4\varepsilon(n+\alpha)} \\ &\leq \delta_f \left[\frac{(n+2\alpha)(n+2\varepsilon)}{n^2+4\varepsilon(n+\alpha)} \right] - \frac{4\varepsilon(n+2\alpha)}{n^2+4\varepsilon(n+\alpha)}. \end{aligned}$$

Next, the definition in (4-16) and the fact that $\varepsilon < \frac{1}{2}\alpha$ gives that $1 > \theta > (\gamma-2\beta+\alpha)/(n+\alpha)$. Then the fact that the function $t \rightarrow t/(1-t)$ is increasing in the interval $(0, 1)$ allows to estimate

$$\frac{\gamma-2\beta+\alpha}{2n} \leq \frac{\gamma-2\beta+\alpha}{n-\gamma+2\beta} \leq \frac{\theta}{1-\theta} < \frac{p\theta}{1-p\theta},$$

so that, from the second inequality in (6-3), it follows that

$$\delta < \frac{(\gamma-2\beta+\alpha)\delta_g}{4n} \leq \frac{\delta_g}{2} \frac{p\theta}{1-p\theta}. \tag{6-5}$$

Finally, for $t \in (0, 1)$, we define the function

$$S(t) := \frac{2c_s(n+4)}{4\alpha t^6} \geq \frac{2c_s}{(2-q)t^{3(2-q)/q}}, \tag{6-6}$$

where c_s is the constant introduced in Proposition 5.1 and q has been introduced in (4-13); in the last estimation we have used that $\varepsilon \in (0, \frac{1}{2}\alpha)$. We then impose the last restriction on δ , that is,

$$\delta S(\varepsilon) \leq \frac{1}{4}. \tag{6-7}$$

All in all, the choices made in (6-3) and (6-7) allow us to determine δ as a positive number depending only on $n, \alpha, \Lambda, \beta, p, \gamma, \delta_1$, as required in the statement of Theorem 6.1.

Step 2: Reverse Hölder-type inequalities. In this step, by applying Proposition 5.1 with the numbers $\varepsilon, \delta, \delta_f$ as chosen in Step 1, we are going to prove that $U \in L_{\text{loc}}^{2_*+\delta}(\mathbb{R}^{2n}; \mu)$. The finiteness of the terms on the right-hand side of (6-1) has already been discussed in Section 4C. First of all, we show that we can

reduce to the case $\varrho_0 = 1$ and $\mathcal{B} = B(0, 1) \times B(0, 1)$; this eventually allows us to apply [Proposition 5.1](#). Indeed, notice that the rescaled functions

$$\tilde{u}(x) := u(x_0 + \varrho_0 x), \quad \tilde{g}(x) := \varrho_0^{2\alpha-2\beta} g(x_0 + \varrho_0 x), \quad \tilde{f}(x) := \varrho_0^{2\alpha} f(x_0 + \varrho_0 x),$$

still solve (1-14). Therefore, applying (6-1) in this case and in $B(0, 1) \times B(0, 1)$, and scaling back to the original functions and to the original diagonal ball \mathcal{B} , leads to (6-1) in the general case. We now pass to the proof of (6-1) when $\mathcal{B} = B(0, 1) \times B(0, 1)$. We define the truncated function $U_m := \min\{U, m\}$ for m being a positive integer, and the measure $d\nu = U^2 d\mu$. Moreover, we use the abbreviation $\mathcal{B}_s := \mathcal{B}(0, s)$. With the aim of applying [Proposition 5.1](#), we then consider balls

$$\mathcal{B} \equiv \mathcal{B}_1 \subset \mathcal{B}_t \subset \mathcal{B}_s \subset \mathcal{B}_2$$

as in (5-6), while λ_0 is accordingly defined as in (5-8). We shall derive uniform higher integrability for the functions U_m and will recover the final result by letting $m \rightarrow \infty$. With $\delta \in (0, 1)$ being the number determined in Step 1, by Cavalieri's principle we have that

$$\begin{aligned} \int_{\mathcal{B}_t} U_m^\delta U^2 d\mu &= \int_{\mathcal{B}_t} U_m^\delta d\nu \\ &= \delta \int_0^\infty \lambda^{\delta-1} \nu(\mathcal{B}_t \cap \{U_m > \lambda\}) d\lambda \\ &= \delta \int_0^m \lambda^{\delta-1} \int_{\mathcal{B}_t \cap \{U > \lambda\}} U^2 d\mu d\lambda \\ &\leq \lambda_0^\delta \int_{\mathcal{B}_t} U^2 d\mu + \delta \int_{\lambda_0}^m \lambda^{\delta-1} \int_{\mathcal{B}_t \cap \{U > \lambda\}} U^2 d\mu d\lambda. \end{aligned} \quad (6-8)$$

The second-last integral appearing in this display can be easily estimated by recalling the definition of λ_0 in (5-8) and that $\varrho_0/(s-t) \geq 1$, and using (4-2):

$$\lambda_0^\delta \int_{\mathcal{B}_t} U^2 d\mu \leq \mu(\mathcal{B}_2) \lambda_0^\delta \int_{2\mathcal{B}} U^2 d\mu \leq c\mu(\mathcal{B}_1) \lambda_0^{2+\delta}. \quad (6-9)$$

We proceed with the remaining term in (6-8); using (5-7) we gain

$$\begin{aligned} \delta \int_{\lambda_0}^m \lambda^{\delta-1} \int_{\mathcal{B}_t \cap \{U > \lambda\}} U^2 d\mu d\lambda &\leq \frac{c_s \delta}{\varepsilon^{3(2-q)/q}} \int_{\lambda_0}^m \lambda^{\delta+1-q} \int_{\mathcal{B}_s \cap \{U > \lambda\}} U^q d\mu d\lambda \\ &\quad + c_f \delta \int_{\lambda_0}^m \frac{\lambda_0^{(2_*+\delta_f)2_*\eta/(1-2_*\eta)}}{\lambda^{(1+\eta\delta_f)2_*/(1-2_*\eta)-1-\delta}} \int_{\mathcal{B}_s \cap \{F > \kappa_f \lambda\}} F^{2_*} d\mu d\lambda \\ &\quad + c_g \delta \int_{\lambda_0}^m \frac{\lambda_0^{(p+\delta_g)p\theta/(1-p\theta)}}{\lambda^{(1+\theta\delta_g)p/(1-p\theta)-1-\delta}} \int_{\mathcal{B}_s \cap \{G > \kappa_g \lambda\}} G^p d\mu d\lambda \\ &=: \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3. \end{aligned} \quad (6-10)$$

Using (6-6)–(6-7) and Fubini’s theorem, we get

$$\begin{aligned}
\mathcal{J}_1 &\leq \frac{c_s \delta}{\varepsilon^{3(2-q)/q}} \int_0^\infty \lambda^{\delta+1-q} \int_{\mathcal{B}_s \cap \{U_m > \lambda\}} U^q \, d\mu \, d\lambda \\
&= \frac{c_s \delta}{(\delta+2-q)\varepsilon^{3(2-q)/q}} \int_{\mathcal{B}_s} U_m^{2+\delta-q} U^q \, d\mu \\
&\leq \delta S(\varepsilon) \int_{\mathcal{B}_s} U_m^\delta U^2 \, d\mu \leq \frac{1}{4} \int_{\mathcal{B}_s} U_m^\delta U^2 \, d\mu.
\end{aligned} \tag{6-11}$$

We next estimate \mathcal{J}_2 . Changing variables, using Fubini’s theorem, and recalling the dependence $\kappa_f \equiv \kappa_f(n, \alpha, \Lambda, \varepsilon)$, we have

$$\begin{aligned}
&\int_{\lambda_0}^m \lambda^{\delta+1-(1+\eta\delta_f)2_*/(1-2_*\eta)} \int_{\mathcal{B}_s \cap \{F > \kappa_f \lambda\}} F^{2_*} \, d\mu \, d\lambda \\
&\leq c \int_0^\infty \lambda^{\delta+1-(1+\eta\delta_f)2_*/(1-2_*\eta)} \int_{\mathcal{B}_s \cap \{F > \lambda\}} F^{2_*} \, d\mu \, d\lambda \\
&= \frac{c\mu(\mathcal{B}_2)}{\delta+2-(1+\eta\delta_f)2_*/(1-2_*\eta)} \int_{\mathcal{B}_2} F^{\delta+2-(1+\eta\delta_f)2_*/(1-2_*\eta)+2_*} \, d\mu \\
&\leq \frac{c\mu(\mathcal{B}_2)}{\delta} \int_{\mathcal{B}_2} F^{\delta+2-(1+\eta\delta_f)2_*/(1-2_*\eta)+2_*} \, d\mu,
\end{aligned} \tag{6-12}$$

again for a constant depending on n, α, Λ and ε . In writing the last inequality we have used that (6-2) is in force and the fact that

$$\delta_f \leq \frac{2\varepsilon(n+2\alpha)}{n(\alpha-\varepsilon)} \iff 2 - \frac{(1+\eta\delta_f)2_*}{1-2_*\eta} \geq 0.$$

The last integral appearing in (6-12) is finite if $\delta+2-(1+\eta\delta_f)2_*/(1-2_*\eta)+2_* \leq 2_* + \delta_f$, and a lengthy computation shows that this is equivalent to (6-4). Therefore, using Hölder’s inequality, we can estimate

$$\begin{aligned}
\mathcal{J}_2 &\leq c\mu(\mathcal{B}_2)\lambda_0^{(2_*+\delta_f)2_*/(1-2_*\eta)} \left(\int_{\mathcal{B}_2} F^{2_*+\delta_f} \, d\mu \right)^{\frac{\delta+2-(1+\eta\delta_f)2_*/(1-2_*\eta)+2_*}{2_*+\delta_f}} \\
&\leq c\mu(\mathcal{B}_1)\lambda_0^{(2_*+\delta_f)2_*/(1-2_*\eta)+\delta+2-(1+\eta\delta_f)2_*/(1-2_*\eta)+2_*} \\
&= c\mu(\mathcal{B}_1)\lambda_0^{2+\delta},
\end{aligned} \tag{6-13}$$

where c depends only on n, α, Λ and ε . We finally come to the estimation of \mathcal{J}_3 . For this we notice that the definitions of p and θ give, independently of ε , that

$$p \geq \frac{2n}{n+2(\gamma-2\beta+\alpha)} \iff \frac{p}{1-p\theta} \geq 2, \tag{6-14}$$

and then, recalling that $\kappa_g \equiv \kappa_g(n, \alpha, \Lambda, \varepsilon, \gamma, \beta, p)$, we have

$$\begin{aligned} \int_{\lambda_0}^m \lambda^{\delta+1-(1+\theta\delta_g)p/(1-p\theta)} \int_{\mathcal{B}_s \cap \{G > \kappa_g \lambda\}} G^p d\mu d\lambda &\leq \int_{\lambda_0}^{\infty} \lambda^{\delta+1-(1+\theta\delta_g)p/(1-p\theta)} d\lambda \int_{\mathcal{B}_s} G^p d\mu \\ &\leq \frac{c\lambda_0^{\delta+2-(1+\theta\delta_g)p/(1-p\theta)} \mu(\mathcal{B}_2)}{(1+\theta\delta_g)p/(1-p\theta) - \delta - 2} \int_{\mathcal{B}_2} G^p d\mu \\ &\leq \frac{c\lambda_0^{\delta+2-(1+\theta\delta_g)p/(1-p\theta)} \mu(\mathcal{B}_2)}{\theta\delta_g p/(1-p\theta) - \delta} \left(\int_{\mathcal{B}_2} G^{p+\delta_g} d\mu \right)^{p/(p+\delta_g)} \\ &\leq \frac{c}{\delta} \lambda_0^{\delta+2-(1+\theta\delta_g)p/(1-p\theta)+p} \mu(\mathcal{B}_2). \end{aligned}$$

Observe that in order to perform the last two estimations we have also used (6-14) and (6-5), respectively. Therefore we can estimate as in (6-13), that is,

$$\mathcal{J}_3 \leq c\mu(\mathcal{B}_2)\lambda_0^{(p+\delta_g)p\theta/(1-p\theta)+\delta+2-(1+\theta\delta_g)p/(1-p\theta)+p} = c\mu(\mathcal{B}_2)\lambda_0^{2+\delta}, \quad (6-15)$$

with $c \equiv c(n, \alpha, \Lambda, \varepsilon, \gamma, \beta, p)$. Connecting (6-11), (6-13) and (6-15) to (6-10), and combining the resulting inequality with (6-8) and (6-9), we get

$$\int_{\mathcal{B}_t} U_m^\delta U^2 d\mu \leq \frac{1}{4} \int_{\mathcal{B}_s} U_m^\delta U^2 d\mu + c\mu(\mathcal{B}_1)\lambda_0^{2+\delta}.$$

By recalling the definition of λ_0 in (5-8), and using several times the doubling property of μ , after a few elementary manipulations we come to

$$\left(\int_{\mathcal{B}_t} U_m^\delta U^2 d\mu \right)^{1/(2+\delta)} \leq \frac{1}{2} \left(\int_{\mathcal{B}_s} U_m^\delta U^2 d\mu \right)^{1/(2+\delta)} + \frac{c}{\varepsilon} \left(\frac{\varrho_0}{s-t} \right)^{2n} \text{ADD}(2\mathcal{B}).$$

We can therefore rewrite the above inequality as

$$\phi(t) \leq \frac{1}{2}\phi(s) + \frac{c}{\varepsilon} \left(\frac{\varrho_0}{s-t} \right)^{2n} \text{ADD}(2\mathcal{B})$$

for a constant $c \equiv c(n, \alpha, \Lambda, \varepsilon, \gamma, \beta, p)$ which is still independent of $m \in \mathbb{N}$, and where, obviously, we have set

$$\phi(\varrho) := \left(\int_{\mathcal{B}_\varrho} U_m^\delta U^2 d\mu \right)^{1/(2+\delta)}$$

for $\varrho \in [\varrho_0, \frac{3}{2}\varrho_0]$. We are therefore in position to apply the standard iteration Lemma 6.2 below, which gives, after returning to the full notation,

$$\left(\int_{\mathcal{B}} U_m^\delta U^2 d\mu \right)^{1/(2+\delta)} \leq c \text{ADD}(2\mathcal{B}).$$

The previous inequality holds for a constant $c \equiv c(n, \alpha, \Lambda, \varepsilon, \gamma, \beta, p)$ which is independent of $m \in \mathbb{N}$. Therefore, letting $m \rightarrow \infty$ yields

$$\left(\int_{\mathcal{B}} U^{2+\delta} d\mu \right)^{1/(2+\delta)} \leq c \text{ADD}(2\mathcal{B}).$$

At this point (6-1) follows by recalling the definition of $\text{ADD}(2\mathcal{B})$ in (5-5) and using a few elementary manipulations involving Hölder’s inequality. In particular, we use the fact that $2_* + \delta_f \leq 2_* + \delta_0$ and $p + \delta_g \leq p(1 + \delta_1)$; see Lemma 4.2. □

Lemma 6.2. *Let $\phi: [\varrho_0, \frac{3}{2}\varrho_0] \rightarrow [0, \infty)$ be a function such that*

$$\phi(t) \leq \frac{1}{2}\phi(s) + \frac{A}{(s-t)^\gamma}$$

whenever $\varrho_0 < t < s < \frac{3}{2}\varrho_0$, where A and γ are positive constants. Then the inequality

$$\phi(\varrho_0) \leq \frac{cA}{\varrho_0^\gamma}$$

holds for a constant $c \equiv c(\gamma)$.

For a proof of this lemma, see for instance [Giusti 2003, Chapter 6].

Proof of Theorem 1.1. The proof is now a simple consequence of Theorem 6.1, that gives that $U \in L^{2+\delta}(\mathcal{B}; \mu)$ whenever $\mathcal{B} = B \times B$ and $B \subset \mathbb{R}^n$ is a ball (that for simplicity we take to be centered at the origin). We now translate this information in terms of fractional norms of the original function u . In fact, this means that, whenever $B \subset \mathbb{R}^n$ is a ball centered at the origin, we have

$$\int_{B \times B} U^{2+\delta} d\mu = \int_B \int_B \frac{|u(x) - u(y)|^{2+\delta}}{|x - y|^{n+(2+\delta)\alpha+\varepsilon\delta}} dx dy < \infty.$$

Rewriting the last integral, we find

$$\int_B \int_B \frac{|u(x) - u(y)|^{2+\delta}}{|x - y|^{n+(2+\delta)[\alpha+\varepsilon\delta/(2+\delta)]}} dx dy < \infty$$

whenever $B \subset \mathbb{R}^n$ is a ball, and this means that $u \in W_{\text{loc}}^{\alpha+\varepsilon\delta/(2+\delta), 2+\delta}(\mathbb{R}^n)$; observe that since $\varepsilon < 1 - \alpha$ then $\alpha + \varepsilon\delta/(2 + \delta) < 1$. We have therefore improved the regularity of u both in the fractional and in the differentiability scale, and Theorem 1.1 follows by suitably renaming (via embedding theorems) the number δ considered in its statement. □

Proof of Theorem 1.3. The proof is just a consequence of the arguments developed to prove Theorem 6.1. In fact the only thing needed there is Proposition 4.4, whose content is now considered as an assumption in (1-23), provided that we take $F = G = 0$; the rest of the argument then remains unchanged. □

Acknowledgments

We wish to thank Vladimir Maz'ya for a useful discussion and Paolo Baroni for a careful reading of a preliminary version of the manuscript. We also thank the referees for their extremely valuable work and their careful reading of the first draft of the paper; their comments led to an improved version.

References

- [Bass and Kassmann 2005] R. F. Bass and M. Kassmann, “Hölder continuity of harmonic functions with respect to operators of variable order”, *Comm. Partial Differential Equations* **30**:7-9 (2005), 1249–1259. MR 2006i:31005 Zbl 1087.45004
- [Bass and Ren 2013] R. F. Bass and H. Ren, “Meyers inequality and strong stability for stable-like operators”, *J. Funct. Anal.* **265**:1 (2013), 28–48. MR 3049880 Zbl 1295.47047
- [Bjorland et al. 2012] C. Bjorland, L. Caffarelli, and A. Figalli, “Non-local gradient dependent operators”, *Adv. Math.* **230**:4-6 (2012), 1859–1894. MR 2927356 Zbl 1252.35099
- [Bojarski and Iwaniec 1983] B. Bojarski and T. Iwaniec, “Analytical foundations of the theory of quasiconformal mappings in \mathbf{R}^n ”, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **8**:2 (1983), 257–324. MR 85h:30023 Zbl 0548.30016
- [Cabré and Cinti 2014] X. Cabré and E. Cinti, “Sharp energy estimates for nonlinear fractional diffusion equations”, *Calc. Var. Partial Differential Equations* **49**:1-2 (2014), 233–269. MR 3148114 Zbl 1282.35399
- [Cabré and Roquejoffre 2013] X. Cabré and J.-M. Roquejoffre, “The influence of fractional diffusion in Fisher-KPP equations”, *Comm. Math. Phys.* **320**:3 (2013), 679–722. MR 3057187 Zbl 06179645
- [Caffarelli and Silvestre 2011] L. Caffarelli and L. Silvestre, “The Evans-Krylov theorem for nonlocal fully nonlinear equations”, *Ann. of Math. (2)* **174**:2 (2011), 1163–1187. MR 2831115 Zbl 1232.49043
- [Caffarelli et al. 2011] L. Caffarelli, C. H. Chan, and A. Vasseur, “Regularity theory for parabolic nonlinear integral operators”, *J. Amer. Math. Soc.* **24**:3 (2011), 849–869. MR 2012c:45024 Zbl 1223.35098
- [Di Castro et al. 2014a] A. Di Castro, T. Kuusi, and G. Palatucci, “Local behavior of fractional p -minimizers”, preprint, 2014, Available at <http://cvgmt.sns.it/paper/2379/>.
- [Di Castro et al. 2014b] A. Di Castro, T. Kuusi, and G. Palatucci, “Nonlocal Harnack inequalities”, *J. Funct. Anal.* **267**:6 (2014), 1807–1836. MR 3237774 Zbl 06330975
- [Di Nezza et al. 2012] E. Di Nezza, G. Palatucci, and E. Valdinoci, “Hitchhiker’s guide to the fractional Sobolev spaces”, *Bull. Sci. Math.* **136**:5 (2012), 521–573. MR 2944369 Zbl 1252.46023
- [Dorronsoro 1985] J. R. Dorronsoro, “A characterization of potential spaces”, *Proc. Amer. Math. Soc.* **95**:1 (1985), 21–31. MR 86k:46046 Zbl 0577.46035
- [Felsinger and Kassmann 2013] M. Felsinger and M. Kassmann, “Local regularity for parabolic nonlocal operators”, *Comm. Partial Differential Equations* **38**:9 (2013), 1539–1573. MR 3169755 Zbl 1277.35090
- [Fusco and Sbordone 1990] N. Fusco and C. Sbordone, “Higher integrability of the gradient of minimizers of functionals with nonstandard growth conditions”, *Comm. Pure Appl. Math.* **43**:5 (1990), 673–683. MR 91b:49015 Zbl 0727.49021
- [Gehring 1973] F. W. Gehring, “The L^p -integrability of the partial derivatives of a quasiconformal mapping”, *Acta Math.* **130** (1973), 265–277. MR 53 #5861 Zbl 0258.30021
- [Giusti 2003] E. Giusti, *Direct methods in the calculus of variations*, World Scientific Publishing Co., River Edge, NJ, 2003. MR 2004g:49003 Zbl 1028.49001
- [Iwaniec 1998] T. Iwaniec, “The Gehring lemma”, pp. 181–204 in *Quasiconformal mappings and analysis* (Ann Arbor, MI, 1995), edited by P. Duren et al., Springer, New York, 1998. MR 99e:30012 Zbl 0888.30017
- [Kassmann 2009] M. Kassmann, “A priori estimates for integro-differential operators with measurable kernels”, *Calc. Var. Partial Differential Equations* **34**:1 (2009), 1–21. MR 2010b:35474 Zbl 1158.35019
- [Kinnunen and Lewis 2000] J. Kinnunen and J. L. Lewis, “Higher integrability for parabolic systems of p -Laplacian type”, *Duke Math. J.* **102**:2 (2000), 253–271. MR 2001b:35152 Zbl 0994.35036

- [Kuusi and Mingione 2012] T. Kuusi and G. Mingione, “Universal potential estimates”, *J. Funct. Anal.* **262**:10 (2012), 4205–4269. [MR 2900466](#) [Zbl 1252.35097](#)
- [Kuusi et al. 2014] T. Kuusi, G. Mingione, and Y. Sire, “A fractional Gehring lemma, with applications to nonlocal equations”, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **25**:4 (2014), 345–358. [MR 3283394](#)
- [Maz’ya 2011] V. Maz’ya, *Sobolev spaces with applications to elliptic partial differential equations*, 2nd ed., Grundlehren der Mathematischen Wissenschaften **342**, Springer, Heidelberg, 2011. [MR 2012a:46056](#) [Zbl 1217.46002](#)
- [Meyers 1963] N. G. Meyers, “An L^p -estimate for the gradient of solutions of second order elliptic divergence equations”, *Ann. Scuola Norm. Sup. Pisa* (3) **17** (1963), 189–206. [MR 28 #2328](#) [Zbl 0127.31904](#)
- [Meyers and Elcrat 1975] N. G. Meyers and A. Elcrat, “Some results on regularity for solutions of non-linear elliptic systems and quasi-regular functions”, *Duke Math. J.* **42** (1975), 121–136. [MR 54 #5618](#) [Zbl 0347.35039](#)
- [Mingione 2003] G. Mingione, “The singular set of solutions to non-differentiable elliptic systems”, *Arch. Ration. Mech. Anal.* **166**:4 (2003), 287–301. [MR 2004b:35070](#) [Zbl 1142.35391](#)
- [Stein 1961] E. M. Stein, “The characterization of functions arising as potentials”, *Bull. Amer. Math. Soc.* **67** (1961), 102–104. [MR 23 #A1051](#)
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, 1993. [MR 95c:42002](#) [Zbl 0821.42001](#)

Received 19 Feb 2014. Accepted 22 Oct 2014.

TUOMO KUUSI: tuomo.kuusi@aalto.fi

Institute of Mathematics, Aalto University, P.O. Box 11100, FI-00076 Aalto, Finland

GIUSEPPE MINGIONE: giuseppe.mingione@math.unipr.it

Dipartimento di Matematica e Informatica, Università di Parma, Parco Area delle Scienze 53/a, Campus, I-43100 Parma, Italy

YANNICK SIRE: yannick.sire@univ-amu.fr

Institut de Mathématique de Marseille, CMI-UMR CNRS 7353, Université Aix-Marseille, 9 rue F. Joliot Curie, 13453 Marseille Cedex 13, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 8 No. 1 2015

Hölder continuity and bounds for fundamental solutions to nondivergence form parabolic equations	1
SEIICHIRO KUSUOKA	
Eigenvalue distribution of optimal transportation	33
BO'AZ B. KLARTAG and ALEXANDER V. KOLESNIKOV	
Nonlocal self-improving properties	57
TUOMO KUUSI, GIUSEPPE MINGIONE and YANNICK SIRE	
Symbol calculus for operators of layer potential type on Lipschitz surfaces with VMO normals, and related pseudodifferential operator calculus	115
STEVE HOFMANN, MARIUS MITREA and MICHAEL E. TAYLOR	
Criteria for Hankel operators to be sign-definite	183
DIMITRI R. YAFAEV	
Nodal sets and growth exponents of Laplace eigenfunctions on surfaces	223
GUILLAUME ROY-FORTIN	