

# ANALYSIS & PDE

Volume 8

No. 7

2015

MOSTAFA FAZLY, JUN-CHENG WEI AND XINGWANG XU

**A POINTWISE INEQUALITY FOR  
THE FOURTH-ORDER LANE-EMDEN EQUATION**

## A POINTWISE INEQUALITY FOR THE FOURTH-ORDER LANE–EMDEN EQUATION

MOSTAFA FAZLY, JUN-CHENG WEI AND XINGWANG XU

We prove the pointwise inequality

$$-\Delta u \geq \left( \frac{2}{(p+1) - c_n} \right)^{\frac{1}{2}} |x|^{a/2} u^{(p+1)/2} + \frac{2}{n-4} \frac{|\nabla u|^2}{u} \quad \text{in } \mathbb{R}^n,$$

where  $c_n := 8/(n(n-4))$ , for positive bounded solutions of the fourth-order Hénon equation, that is,

$$\Delta^2 u = |x|^a u^p \quad \text{in } \mathbb{R}^n$$

for some  $a \geq 0$  and  $p > 1$ . Motivated by Moser’s proof of Harnack’s inequality as well as Moser iteration-type arguments in the regularity theory, we develop an iteration argument to prove the above pointwise inequality. As far as we know this is the first time that such an argument is applied towards constructing pointwise inequalities for partial differential equations. An interesting point is that the coefficient  $2/(n-4)$  also appears in the fourth-order  $Q$ -curvature and the Paneitz operator. This, in particular, implies that the scalar curvature of the conformal metric with conformal factor  $u^{4/(n-4)}$  is positive.

### 1. Introduction

We are interested in proving an a priori pointwise estimate for positive solutions of the fourth-order Hénon equation

$$\Delta^2 u = |x|^a u^p \quad \text{in } \mathbb{R}^n, \tag{1-1}$$

where  $p > 1$  and  $a \geq 0$ . Let us first mention that, for the case  $a = 0$ , it is known that (1-1) only admits  $u = 0$  as a nonnegative solution when  $p$  is a subcritical exponent, that is,  $1 < p < (n+4)/(n-4)$  when  $n \geq 5$ , and  $1 < p$  when  $n \leq 4$ . Moreover, for the critical case  $p = (n+4)/(n-4)$ , all entire positive solutions are classified. See [Lin 1998; Wei and Xu 1999]. This is a counterpart of the standard Liouville theorem of Gidas and Spruck [1981a; 1981b] for the second-order Lane–Emden equation

$$-\Delta u = u^p \quad \text{in } \mathbb{R}^n, \tag{1-2}$$

stating that  $u = 0$  is the only nonnegative solution for (1-2) when  $p$  is a subcritical exponent, that is,  $1 < p < (n+2)/(n-2)$  when  $n \geq 3$ . Note also that, for the fourth-order Hénon equation, it is conjectured that  $u = 0$  is the only nonnegative solution of (1-1) when  $p$  is a subcritical exponent, that is, when

---

Fazly and Wei are supported by NSERC grants.

MSC2010: 35B45, 35B50, 35J30, 53C21, 35B08.

Keywords: semilinear elliptic equations, a priori pointwise estimate, Moser iteration-type arguments, elliptic regularity.

$1 < p < (n + 4 + 2a)/(n - 4)$  and  $n \geq 5$ ; see [Fazly and Ghoussoub 2014]. Therefore, throughout this note, when we are dealing with (1-1) we assume that  $p > (n + 4 + 2a)/(n - 4)$  and  $n \geq 5$ . For more information, see [Fazly and Ghoussoub 2014; Souplet 2009] and references therein.

Pointwise estimates have had tremendous impact on the theory of elliptic partial differential equations. In what follows, we list some of the celebrated pointwise inequalities for certain semilinear elliptic equations and systems. These inequalities have been used to tackle well-known conjectures and open problems. The following inequality has been one of the main techniques to solve De Giorgi’s conjecture (1978) for the Allen–Cahn equation and to analyze various semilinear equations and problems.

**Theorem 1.1** [Modica 1985]. *Let  $F \in C^2(\mathbb{R})$  be a nonnegative function and  $u$  be a bounded entire solution of*

$$\Delta u = F'(u) \quad \text{in } \mathbb{R}^n. \tag{1-3}$$

Then

$$|\nabla u|^2 \leq 2F(u) \quad \text{in } \mathbb{R}^n. \tag{1-4}$$

For the specific case  $F(u) = \frac{1}{4}(1 - u^2)^2$ , equation (1-3) is known as the Allen–Cahn equation. Note also that [Caffarelli et al. 1994] extended this inequality to quasilinear equations. We refer interested readers to [Farina and Valdinoci 2010; 2011; 2013; 2014; Castellaneta et al. 2012; Farina et al. 2008] regarding pointwise gradient estimates and certain improvements of (1-4). For the fourth-order counterpart of (1-3) with an arbitrary nonlinearity, a general inequality of the form (1-4) is not known. However, for a particular nonlinearity known as the fourth-order Lane–Emden equation, i.e.,

$$\Delta^2 u = u^p \quad \text{in } \mathbb{R}^n \tag{1-5}$$

it was shown by Wei and Xu [1999, Theorem 3.1] that the negative Laplacian of the positive solutions is nonnegative, that is,  $-\Delta u \geq 0$  in  $\mathbb{R}^n$ . Set  $v = -\Delta u$  and, from the fact that  $-\Delta u \geq 0$ , we can consider (1-5) as a special case (when  $q = 1$ ) of the Lane–Emden system

$$\begin{cases} -\Delta u = v^q & \text{in } \mathbb{R}^n, \\ -\Delta v = u^p & \text{in } \mathbb{R}^n, \end{cases} \tag{1-6}$$

where  $p \geq q \geq 1$ . Note that there is a significance difference between system (1-6) and equation (1-5), in the sense that this system has Hamiltonian structure while the equation has gradient structure. This system has been of great interest, at least in the past two decades. In particular, the Lane–Emden conjecture, stating that  $u = v = 0$  is the only nonnegative solution for this system when  $1/(p + 1) + 1/(q + 1) > (n - 2)/n$  has been studied extensively and various methods and techniques have been developed to tackle this conjecture. Among these methods, Souplet [2009] proved the following pointwise inequality for solutions of (1-6) and then used it to prove the Lane–Emden conjecture in four dimensions. Note that the particular case  $1 < p < 2$  was done by Phan [2012].

**Theorem 1.2** [Souplet 2009]. *Let  $u$  and  $v$  be nonnegative solutions of (1-6). Then*

$$\frac{u^{p+1}}{p + 1} \leq \frac{v^{q+1}}{q + 1} \quad \text{in } \mathbb{R}^n. \tag{1-7}$$

Applying this theorem, the following pointwise inequality holds for nonnegative solutions of (1-5):

$$-\Delta u \geq \sqrt{\frac{2}{p+1}} u^{(p+1)/2} \quad \text{in } \mathbb{R}^n. \tag{1-8}$$

Note also that Phan [2012], with similar methods to those [Souplet 2009], extended the pointwise inequality (1-7) to nonnegative solutions of the Hénon–Lane–Emden system

$$\begin{cases} -\Delta u = |x|^b v^q & \text{in } \mathbb{R}^n, \\ -\Delta v = |x|^a u^p & \text{in } \mathbb{R}^n, \end{cases} \tag{1-9}$$

where  $p \geq q \geq 1$ . Suppose that  $0 \leq a - b \leq (n - 2)(p - q)$ ; then

$$|x|^a \frac{u^{p+1}}{p+1} \leq |x|^b \frac{v^{q+1}}{q+1} \quad \text{in } \mathbb{R}^n. \tag{1-10}$$

The standard method to prove a pointwise inequality, as is used to prove (1-7) and (1-4), is to derive an appropriate equation — call it an auxiliary equation — for the function that is the difference between the right-hand and left-hand sides of the inequality. Then, whenever we have enough decay estimates on solutions of the auxiliary equation, maximum principles can be applied to prove that the difference function has a fixed sign. So, the key point here is to manipulate a suitable auxiliary equation.

In a more technical framework, to construct an auxiliary equation to prove (1-7) and (1-8), a few positive terms, including a gradient term of the form  $|\nabla u|^2 u^{t-2}$  for some number  $t$ , are not considered in [Souplet 2009]. To be more explicit, in order to prove (1-8), which is a particular case of (1-7), the difference function  $w(x) := \Delta u + \sqrt{2/(p+1)} u^{(p+1)/2}$  is considered. Straightforward calculations show that the following auxiliary equation holds:

$$\left( \sqrt{\frac{2}{p+1}} u^{(1-p)/2} \right) \Delta w = \Delta u + \sqrt{\frac{2}{p+1}} u^{(p+1)/2} + \frac{p-1}{2} \frac{|\nabla u|^2}{u}. \tag{1-11}$$

In order to show that  $\Delta w$  is nonnegative when  $w$  is nonnegative, via maximum principles for the above equation, the gradient term  $|\nabla u|^2/u$  is not considered in [Souplet 2009]. Note that (1-11) implies, in spirit, that the gradient term  $|\nabla u|^2/u$  should have an impact on the inequality, just like the Laplacian operator and the power term  $u^{(p+1)/2}$ . This is our motivation to attempt to include the gradient term in the inequality (1-8) that gives a lower bound on the Laplacian operator. Let us briefly mention that Modica, in his proof of (1-4), took advantage of similar gradient terms to construct an auxiliary equation. Following ideas provided by Modica [1985] and Souplet [2009], as we shall see in the proof of Proposition 3.1, we manage to keep most of the positive terms when looking for an auxiliary equation.

In this paper, we develop a Moser iteration-type argument to prove a lower bound for the negative Laplacian of positive bounded solutions of (1-1) that involves powers of  $u$  and the new term  $|\nabla u|^2/u$  with  $2/(n - 4)$  as the coefficient. The remarkable point is that the coefficient  $2/(n - 4)$  is exactly what we need in the estimate of the scalar curvature for the conformal metric  $g = u^{2/(n-4)} g_0$ .

Here is our main result:

**Theorem 1.3.** *Let  $u$  be a bounded positive solution of (1-1). Then the following pointwise inequality holds:*

$$-\Delta u \geq \sqrt{\frac{2}{(p+1)-c_n}} |x|^{a/2} u^{(p+1)/2} + \frac{2}{n-4} \frac{|\nabla u|^2}{u} \quad \text{in } \mathbb{R}^n, \quad (1-12)$$

where  $c_n := 8/(n(n-4))$  and  $0 \leq a \leq \inf_{k \geq 0} A_k$  ( $A_k$  is defined in (4-28)).

**Remark 1.4.** A natural question here is: what are the best constants in the inequality (1-12)?

Let us now put the inequality (1-12) in a more geometric text. By the conformal change  $g = u^{4/(n-4)} g_0$ , where  $g_0$  is the usual Euclidean metric, the new scalar curvature becomes

$$S_g = -\frac{4(n-1)}{n-2} u^{-(n+2)/(n-4)} \Delta(u^{(n-2)/(n-4)}).$$

An immediate consequence of (1-12) is that the conformal scalar curvature is positive. Note that this cannot be deduced from the inequality (1-8).

The idea of proving a lower bound for the negative of the Laplacian operator is also used in the context of nonlinear eigenvalue problems to prove certain regularity results; see, e.g., [Cowan et al. 2010]. Similar pointwise inequalities are used to prove Liouville theorems in the notion of stability in [Wei et al. 2013; Wei and Ye 2013] and references therein as well. We would like to mention that Gui [2008] proved a very interesting Hamiltonian identity for elliptic systems that may be regarded as a generalization of Modica's inequality. He used this identity to rigorously analyze the structure of level curves of saddle solutions of the Allen–Cahn equation as well as Young's law for the contact angles in triple junction formation. Note also that, as is shown by Farina [2004] for the Ginzburg–Landau system, the analog of Modica's estimate is false for systems in general. We refer interested readers to [Alikakos 2013] for a review of this topic and to [Fazly and Ghoussoub 2013] for De Giorgi-type results for systems.

Here is the organization of the paper. In Section 2, we provide certain standard elliptic estimates that are consequences of Sobolev embeddings and the regularity theory. Then, in Section 3 we develop a Moser iteration-type argument, following ideas provided by Modica [1985] and Souplet [2009]. Finally, in Section 4, we first give a certain maximum principle argument for a quasilinear equation that arises in the Moser iteration process. Then we apply the estimates and methods developed in the earlier sections. We suggest the reader ignores the weight function  $|x|^a$  in (1-1) when reading the paper for the first time.

## 2. Technical elliptic estimates

In this section, we provide some elliptic decay estimates that we use frequently later in the proofs. Deriving the right decay estimates for solutions of (1-1) plays a fundamental role in our proofs. Similar estimates have been also used in the literature to construct Liouville theorems and regularity results. We refer interested readers to [Fazly 2014; Fazly and Ghoussoub 2014; Phan 2012; Souplet 2009; Phan and Souplet 2012]. We start with the following standard estimate:

**Lemma 2.1** ( $L^p$ -estimate on  $B_R$ ). *Suppose that  $u$  is a nonnegative solution of (1-1); then, for any  $R > 1$  we have*

$$\int_{B_R} |x|^a u^p \leq C R^{n-(4p+a)/(p-1)},$$

where  $C = C(n, p, a) > 0$  is independent of  $R$ .

*Proof.* Consider the following test function  $\phi_R \in C_c^4(\mathbb{R}^n)$  with  $0 \leq \phi_R \leq 1$ :

$$\phi_R(x) = \begin{cases} 1 & \text{if } |x| < R, \\ 0 & \text{if } |x| > 2R, \end{cases}$$

where  $\|D^i \phi_R\|_\infty \leq C/R^i$  for  $1 \leq i \leq 4$ . For fixed  $m \geq 2$ , we have

$$|\Delta^2 \phi_R^m(x)| \leq \begin{cases} 0 & \text{if } |x| < R \text{ or } |x| > 2R, \\ C R^{-4} \phi_R^{m-4} & \text{if } R < |x| < 2R, \end{cases}$$

where  $C > 0$  is independent of  $R$ . For  $m \geq 2$ , multiply the equation by  $\phi_R^m$  and integrate to get

$$\int_{B_{2R}} |x|^a u^p \phi_R^m = \int_{B_{2R}} \Delta^2 u \phi_R^m = \int_{B_{2R}} u \Delta^2 \phi_R^m \leq C R^{-4} \int_{B_{2R} \setminus B_R} u \phi_R^{m-4}.$$

Applying Hölder's inequality, we get

$$\begin{aligned} \int_{B_{2R}} |x|^a u^p \phi_R^m &\leq C R^{-4} \left( \int_{B_{2R} \setminus B_R} |x|^{(-a/p)p'} \right)^{\frac{1}{p'}} \left( \int_{B_{2R} \setminus B_R} |x|^a u^p \phi_R^{(m-4)p} \right)^{\frac{1}{p}} \\ &\leq C R^{(n-(a/p)p')/p'-4} \left( \int_{B_{2R} \setminus B_R} |x|^a u^p \phi_R^{(m-4)p} \right)^{\frac{1}{p}}, \end{aligned}$$

where  $p' = p/(p-1)$ . Set  $m = (m-4)p$ , so that  $m = 4p/(p-1)$ , to get

$$\int_{B_{2R}} |x|^a u^p \phi_R^m \leq C R^{(n-(a/p)p')/p'-4} \left( \int_{B_{2R}} |x|^a u^p \phi_R^m \right)^{\frac{1}{p}}.$$

Therefore,

$$\int_{B_{2R}} |x|^a u^p \phi_R^m \leq C R^{(n-(a/p)p')-4p'}.$$

This finishes the proof. □

From Hölder's inequality we get the following:

**Corollary 2.2.** *Under the same assumptions as Lemma 2.1,*

$$\int_{B_R \setminus B_{R/2}} u \leq C R^{n-(a+4)/(p-1)},$$

where  $C = C(n, p, a) > 0$  is independent of  $R$ .

We now show that the operator  $-\Delta u$  has a sign. Then, we apply this to provide various elliptic estimates for derivatives of  $u$ . In addition, later on this helps us to start an iteration argument.

**Proposition 2.3.** *Let  $u$  be a positive solution of (1-1). Then,  $-\Delta u \geq 0$  in  $\mathbb{R}^n$ .*

*Proof.* Let  $v = -\Delta u$ . Ideas and methods applied in this proof are strongly motivated by the ones given in [Wei and Xu 1999]. Suppose that there is  $x_0 \in \mathbb{R}^n$  such that  $v(x_0) < 0$ . Without loss of generality we take  $x_0 = 0$ , i.e., if  $x_0 \neq 0$  set  $\omega(x) = v(x + x_0)$  and apply the same argument. We use the notation  $\bar{f}(r) = (1/|\partial B_r|) \int_{\partial B_r} f \, dS$  for the average of a function  $f(x)$  on the boundary of  $B_r$ . We refer interested readers to [Ni 1982] regarding the average function. Applying Hölder’s inequality,

$$\begin{cases} -\Delta_r \bar{u}(r) = \bar{v}(r) & \text{in } \mathbb{R}, \\ -\Delta_r \bar{v}(r) \geq r^a (\bar{u})^p & \text{in } \mathbb{R}, \end{cases} \tag{2-1}$$

where  $\Delta_r$  is the Laplacian operator in polar coordinates, i.e.,

$$\Delta_r \bar{f}(r) = r^{1-n} (r^{n-1} \bar{f}'(r))'.$$

It is straightforward to see that

$$\bar{v}'(r) = \frac{1}{|\partial B_r|} \int_{B_r} \Delta v = -\frac{1}{|\partial B_r|} \int_{B_r} |x|^a u^p \leq 0.$$

Therefore,  $\bar{v}(r) \leq \bar{v}(0) < 0$  for  $r > 0$ . Similarly, for  $\bar{u}'(r)$  we have

$$\bar{u}'(r) = -\frac{1}{|\partial B_r|} \int_{B_r} v = -r^{1-n} \int_0^r s^{n-1} \bar{v}(s) \, ds \geq -\bar{v}(0) r^{1-n} \int_0^r s^{n-1} \, ds = -\frac{\bar{v}(0)}{n} r.$$

From this, for any  $r \geq r_0$  we get

$$\bar{u}(r) \geq \alpha r^2, \tag{2-2}$$

where  $\alpha = -\bar{v}(0)/(2n) > 0$ . We now have a lower bound on  $\bar{u}(r)$ . Suppose instead that the following more general lower bound holds on  $\bar{u}(r)$ :

$$\bar{u}(r) \geq \frac{\alpha^{p^k}}{\beta^{s_k}} r^{t_k} \quad \text{for } r \geq r_k, \tag{2-3}$$

where  $s_0 := 0, t_0 := 2, \alpha := -\bar{v}(0)/(2n) > 0$  and  $\beta := 2p + a + n + 4 > 0$ . Note that (2-1) gives a relation between the two functions  $\bar{u}(r)$  and  $\bar{v}(r)$ . Therefore, the lower bound on  $\bar{u}(r)$  forces an upper bound on  $\bar{v}(r)$  and vice versa. In the light of this fact, we can construct an iteration argument to improve the bound (2-3). Integrating the second equation of (2-1) over  $[r_k, r]$  when  $r \geq r_k$ , we get

$$\begin{aligned} r^{n-1} \bar{v}'(r) &\leq r_k^{n-1} \bar{v}'(r_k) - \frac{\alpha^{p^{k+1}}}{\beta^{ps_k}} \int_{r_k}^r s^{n-1+a+pt_k} \, ds \\ &\leq -\frac{\alpha^{p^{k+1}}}{\beta^{ps_k} (pt_k + n + a)} (r^{pt_k+n+a} - r_k^{pt_k+n+a}) \quad \text{since } \bar{v}' < 0. \end{aligned}$$

Therefore,  $\bar{v}'(r) \leq -(\alpha^{p^{k+1}} / (\beta^{ps_k} (pt_k + n + a))) (r^{pt_k+a+1} - r_k^{pt_k+a+1})$  for all  $r \geq r_k$ , that is,

$$\bar{v}'(r) \leq -\frac{\alpha^{p^{k+1}}}{2\beta^{ps_k} (pt_k + n + a)} r^{pt_k+a+1} \quad \text{for all } r \geq 2^{1/(pt_k+a+1)} r_k.$$

Integrating the last inequality over  $[2^{1/(pt_k+a+1)}r_k, r]$  when  $r \geq 2^{1/(pt_k+a+1)}r_k = \tilde{r}_k$ , we obtain

$$\bar{v}(r) \leq \bar{v}(\tilde{r}_k) - \frac{\alpha^{p^{k+1}}}{2\beta^{ps_k}T_{k,n,a,p}}(r^{pt_k+a+2} - \tilde{r}_k^{pt_k+a+2}),$$

where  $T_{k,n,a,p} := (pt_k + n + a)(pt_k + 2 + a)$ . By similar discussions and by taking  $r$  large enough, that is,  $r \geq 2^{1/(pt_k+a+1)}2^{1/(pt_k+a+2)}r_k = \tilde{r}_k$ , we end up with

$$\bar{v}(r) \leq -\frac{\alpha^{p^{k+1}}}{4\beta^{ps_k}T_{k,n,a,p}}r^{pt_k+a+2}. \tag{2-4}$$

Applying (2-4) and integrating (2-1) again over  $[\tilde{r}_k, r]$  when  $r \geq \tilde{r}_k$ , we have

$$r^{n-1}\bar{u}'(r) = \tilde{r}_k^{n-1}\bar{u}'(\tilde{r}_k) - \int_{\tilde{r}_k}^r s^{n-1}\bar{v}(s) ds \geq \frac{\alpha^{p^{k+1}}}{4\beta^{ps_k}T_{k,n,a,p}} \int_{\tilde{r}_k}^r s^{pt_k+a+n+1} ds.$$

Therefore, the following new lower bound on  $\bar{u}(r)$  holds:

$$\bar{u}(r) \geq \frac{\alpha^{p^{k+1}}}{2^4\beta^{ps_k}\tilde{T}_{k,n,a,p}}r^{pt_k+a+n+4},$$

where

$$r \geq 2^{1/(pt_k+a+3)}2^{1/(pt_k+a+4)}\tilde{r}_k = 2^{\sum_{i=1}^4 1/(pt_k+a+i)}r_k$$

and

$$\begin{aligned} \tilde{T}_{k,n,a,p} &= (pt_k + n + a + 2)(pt_k + 4 + a)T_{k,n,a,p} \\ &= (pt_k + n + a)(pt_k + 2 + a)(pt_k + n + a + 2)(pt_k + 4 + a) \\ &\leq (pt_k + n + a + 4)^4. \end{aligned}$$

We now modify this estimate to make the coefficients similar to (2-3). After simplifying, we get

$$\bar{u}(r) \geq \frac{\alpha^{p^{k+1}}}{\beta^{ps_k}M_k}r^{pt_k+a+4} \quad \text{for } r \geq 2^{4/(pt_k+a+1)}r_k, \tag{2-5}$$

where  $M_k := 2^4(pt_k + n + a + 4)^4$ . In what follows, we put an upper bound on  $M_k$  that is expressed as a power of  $\beta$ . Note that

$$\frac{1}{2}\sqrt[4]{M_{k+1}} = pt_{k+1} + n + a + 4 = p(pt_k + n + a + 4) + n + a + 4 \leq (pt_k + n + a + 4)(p + 1) = \frac{1}{2}(p + 1)\sqrt[4]{M_k}.$$

From this we have  $M_{k+1} \leq (p + 1)^4 M_k$  and therefore  $M_k \leq (p + 1)^{4k} M_0$ , where  $M_0 = 2^4(2p + n + a + 4)^4$  because  $t_0 = 2$ . Since the constant  $\beta$  is defined as  $\beta = 2p + n + a + 4$ , we get the bound

$$M_k \leq \beta^{4k+4}. \tag{2-6}$$

From this, (2-3) and (2-5), and to complete the iteration process, we set

$$t_{k+1} := pt_k + a + 4 \quad \text{for } t_0 = 2, \tag{2-7}$$

$$s_{k+1} := ps_k + 4k + 4 \quad \text{for } s_0 = 0, \tag{2-8}$$



and, therefore,

$$\bar{u}(r) \geq \frac{\alpha^{p^{k+1}}}{\beta^{s_{k+1}}} r^{t_{k+1}} \quad \text{for } r \geq r_{k+1}, \tag{2-9}$$

where  $r_{k+1} := 2^{4/(pt_k+a+1)} r_k \geq 2^{\sum_{i=1}^4 1/(pt_k+a+i)} r_k$ . By direct calculations on these recursive sequences, we get the explicit sequences

$$\begin{aligned} t_k &= \frac{2p^{k+1} + (a+2)p^k - (a+4)}{p-1}, \\ s_k &= \frac{4p^{k+1} - 4p(k+1) + 4k}{(p-1)^2}, \\ r_k &= 2^{\sum_{i=0}^{k-1} 4/(pt_i+a+1)} r_0 \leq 2^{\sum_{i=0}^{\infty} 4/(pt_i+a+1)} r_0 =: r^* < \infty. \end{aligned}$$

Set  $R := \beta^{2/(p-1)} M$ , where  $M = \max\{\alpha^{-1}, m\}$  when  $m > 1$  is large enough to ensure  $m\beta^{2/(p-1)} \geq r^*$ . Therefore,  $R \geq r^* \geq r_k$  for any  $k$  and we have

$$\bar{u}(R) \geq M^{t_k-p^k} \beta^{2t_k/(p-1)-s_k}.$$

If we take  $k$  large enough, e.g.,  $k \geq (\ln(a+4) - \ln(a+2))/\ln p$ , then  $t_k > p^k$ . The fact that  $M > 1$  gives us

$$\bar{u}(R) \geq \beta^{2t_k/(p-1)-s_k} = \beta^{(2(a+2)p^k+4k(p-1)+4p-2(a+4))/(p-1)^2}.$$

Since we have assumed that  $a+2 > 0$  and  $\beta > 1$ , we get  $\bar{u}(R) \rightarrow \infty$  as  $k \rightarrow \infty$ . Note that  $0 < R < \infty$  is independent of  $k$ . This finishes the proof. □

We now apply [Proposition 2.3](#) to conclude that  $-\Delta u \geq 0$  and therefore we can consider (1-1) as a special case of the Hénon–Lane–Emden equation.

**Lemma 2.4** ( $L^1$ -estimates on  $B_R$ ). *Suppose that  $u$  is a nonnegative solution of (1-1); then, for any  $R > 1$  we have*

$$\int_{B_R} |\Delta u| \leq C R^{n-(2p+2+a)/(p-1)},$$

where  $C = C(n, p, a) > 0$  is independent of  $R$ .

*Proof.* Set  $v = -\Delta u$ . From [Proposition 2.3](#) we know that  $v \geq 0$ . Therefore, the pair  $(u, v)$  satisfies the system

$$\begin{cases} -\Delta u = v & \text{in } \mathbb{R}^n, \\ -\Delta v = |x|^a u^p & \text{in } \mathbb{R}^n, \end{cases} \tag{2-10}$$

which is a particular case of the Hénon–Lane–Emden system. From the estimates provided in [\[Fazly and Ghoussoub 2014, Lemma 2.1\]](#), we get the desired result. □

**Lemma 2.5** (an interpolation inequality on  $B_R$ ). *Let  $R > 1$  and  $z \in W^{2,1}(B_{2R})$ . Then*

$$\int_{B_R \setminus B_{R/2}} |Dz| \leq CR \int_{B_{2R} \setminus B_{R/4}} |\Delta z| + CR^{-1} \int_{B_{2R} \setminus B_{R/4}} |z|,$$

where  $C = C(n) > 0$  is independent of  $R$ .

**Corollary 2.6.** *Under the same assumptions as Lemma 2.1. The following estimate holds:*

$$\int_{B_R \setminus B_{R/2}} |Du| \leq CR^{n-(p+3+a)/(p-1)},$$

where  $C = C(n, p, a) > 0$  is independent of  $R$ .

**Lemma 2.7** ( $L^\tau$ -estimate on  $B_R$ ). *Let  $1 < \tau < \infty$  and  $z \in W^{2,\tau}(B_{2R})$ . Then*

$$\int_{B_R \setminus B_{R/2}} |D^2z|^\tau \leq C \int_{B_{2R} \setminus B_{R/4}} |\Delta z|^\tau + CR^{-2\tau} \int_{B_{2R} \setminus B_{R/4}} |z|^\tau,$$

where  $C = C(n, \tau) > 0$  does not depend on  $R$ .

**Lemma 2.8** ( $L^2$ -estimates on  $B_R$ ). *Suppose that  $u$  is a bounded nonnegative solution of (1-1); then, for any  $R > 1$  we have*

$$\int_{B_R} |\Delta u|^2 \leq C \int_{B_{2R}} |x|^a u^{p+1} + CR^{-2} \int_{B_{2R}} |\Delta u| + CR^{-4} \int_{B_{2R} \setminus B_R} u, \tag{2-11}$$

where  $C = C(n, p, a) > 0$  does not depend on  $R$ .

*Proof.* We proceed in two steps.

Step 1: Multiply both sides of (1-1) by  $u\phi^2$ , where  $\phi \in C_c^\infty(\mathbb{R}^n) \cap [0, 1]$  is a test function. Then, integrating by parts, we get

$$\begin{aligned} \int_{\mathbb{R}^n} |\Delta u|^2 \phi^2 &= \int_{\mathbb{R}^n} |x|^a u^{p+1} \phi^2 - 4 \int_{\mathbb{R}^n} \Delta u \nabla u \cdot \nabla \phi \phi - \int_{\mathbb{R}^n} u \Delta u (2|\nabla \phi|^2 + 2\phi \Delta \phi) \\ &\leq \int_{\mathbb{R}^n} |x|^a u^{p+1} \phi^2 + \delta \int_{\mathbb{R}^n} |\Delta u|^2 \phi^2 + C(\delta) \int_{\mathbb{R}^n} |\nabla u|^2 |\nabla \phi|^2 + C \int_{\mathbb{R}^n} |\Delta u| (|\nabla \phi|^2 + |\Delta \phi|) \end{aligned}$$

for some constant  $C > 0$ . Here, we have used Cauchy's inequality for  $0 < \delta < 1$ . Therefore, if we set  $\phi$  to be the standard test function, that is,  $\phi = 1$  in  $B_R$  and  $\phi = 0$  in  $\mathbb{R}^n \setminus B_{2R}$  with  $\|D_x^i \phi\|_{L^\infty(B_{2R} \setminus B_R)} \leq CR^{-i}$  for  $i = 1, 2$ , then we get

$$\int_{B_R} |\Delta u|^2 \leq \int_{B_{2R}} |x|^a u^{p+1} + CR^{-2} \int_{B_{2R} \setminus B_R} |\nabla u|^2 + CR^{-2} \int_{B_{2R} \setminus B_R} |\Delta u|, \tag{2-12}$$

where  $C = C(n, p, a) > 0$  does not depend on  $R$ .

Step 2: Multiply both sides of  $-\Delta u = v$  by  $u\phi^2$ , where  $\phi$  is the same test function as in Step 1. Integrating by parts again, we get

$$\int_{\mathbb{R}^n} |\nabla u|^2 \phi^2 = \int_{\mathbb{R}^n} uv\phi^2 - 2 \int_{\mathbb{R}^n} u \nabla u \cdot \nabla \phi \phi \leq \int_{\mathbb{R}^n} uv\phi^2 + \delta \int_{\mathbb{R}^n} |\nabla u|^2 \phi^2 + C(\delta) \int_{\mathbb{R}^n} |\nabla \phi|^2 u^2,$$

where we have also used Cauchy's inequality for  $0 < \delta < 1$ . So,

$$\int_{B_R} |\nabla u|^2 \leq C \int_{B_{2R}} |\Delta u| + CR^{-2} \int_{B_{2R} \setminus B_R} u, \tag{2-13}$$

where we have used the boundedness of  $u$ . From (2-12) and (2-13) we get

$$\int_{B_R} |\Delta u|^2 \leq \int_{B_{2R}} |x|^a u^{p+1} + CR^{-2} \int_{B_{2R}} |\Delta u| + CR^{-4} \int_{B_{2R} \setminus B_R} u. \tag{2-14}$$

This completes the proof. □

We now apply Lemma 2.1, Lemma 2.8 and Corollary 2.2 to get the following:

**Corollary 2.9.** *Suppose that the assumptions of Lemma 2.1 hold. Moreover, let  $u$  be bounded; then*

$$\int_{B_R} |\Delta u|^2 \leq CR^{n-(4p+a)/(p-1)}, \tag{2-15}$$

where  $C = C(n, p, a) > 0$  is independent of  $R$ .

**Lemma 2.10** (Sobolev inequalities on the sphere  $S^{n-1}$ ). *Fix  $n \geq 2$ , a positive integer  $i$  and  $1 < t < \tau \leq \infty$ . For  $z \in W^{i,t}(S^{n-1})$ ,*

$$\|z\|_{L^\tau(S^{n-1})} \leq C \|D_\theta^i z\|_{L^t(S^{n-1})} + C \|z\|_{L^1(S^{n-1})},$$

where

$$\begin{cases} \frac{1}{\tau} = \frac{1}{t} - \frac{i}{n-1} & \text{if } it + 1 < n, \\ \tau = \infty & \text{if } it + 1 > n, \end{cases}$$

and  $C = C(i, t, n, \tau) > 0$ .

### 3. Developing the iteration argument

In this section, we develop a counterpart of the Moser iteration argument [1961] for solutions of (1-1). We define a sequence of functions  $(w_k)_{k=-1}$  of the form

$$w_k := \Delta u + \alpha_k |\nabla u|^2 (u + \epsilon)^{-1} + \beta_k |x|^{a/2} u^{(p+1)/2},$$

where  $\alpha_k$  and  $\beta_k$  are certain nondecreasing sequences of nonnegative numbers with  $\alpha_{-1} = \beta_{-1} = 0$ .

Assuming that  $w_k \leq 0$ , that is, essentially, a lower bound on the negative Laplacian operator holds, we construct a differential inequality for  $w_{k+1}$  with  $\alpha_{k+1} \geq \alpha_k$  and  $\beta_{k+1} \geq \beta_k$ . Then, applying certain maximum principle arguments, we show that  $w_{k+1} \leq 0$ . Note that  $w_{k+1} \leq 0$  is stronger than  $w_k \leq 0$ , because it forces a stronger lower bound on the negative of the Laplacian operator.

We start by proving that  $w_{-1}$ , which is the Laplacian operator of  $u$ , is nonpositive; see Proposition 2.3. Then, using this fact and applying (1-9) and (1-10) when  $q = 1$  and  $b = 0$ , we get the following inequality for nonnegative solutions of the fourth-order Hénon equation (1-1):

$$-\Delta u \geq \sqrt{\frac{2}{p+1}} |x|^{a/2} u^{(p+1)/2} \quad \text{in } \mathbb{R}^n, \tag{3-1}$$

where  $0 \leq a \leq (n-2)(p-1)$ . Inequality (3-1) is the first step of the iteration argument, meaning that  $w_0 \leq 0$  for  $\alpha_0 = 0$  and  $\beta_0 = \sqrt{2/(p+1)}$ .

We now perform the iteration argument:

**Proposition 3.1.** *Let  $u$  be a positive classical solution of (1-1). Suppose that  $(\alpha_k)_{k=0}$  and  $(\beta_k)_{k=0}$  are sequences of numbers. Define the sequence of functions*

$$w_k := \Delta u + \alpha_k |\nabla u|^2 (u + \epsilon)^{-1} + \beta_k |x|^{a/2} u^{(p+1)/2}, \tag{3-2}$$

where  $\epsilon = \epsilon(k)$  is a positive constant. Suppose that  $w_k \leq 0$ ; then  $w_{k+1}$  satisfies the differential inequality

$$\begin{aligned} \Delta w_{k+1} - 2\alpha_{k+1} (u + \epsilon)^{-1} \nabla u \cdot \nabla w_{k+1} + \alpha_{k+1} w_{k+1} (u + \epsilon)^{-2} |\nabla u|^2 - \frac{1}{2} \beta_{k+1} (p+1) u^{(p-1)/2} |x|^{a/2} w_{k+1} \\ \geq I_{\epsilon, \beta_k}^{(1)} |x|^a u^p + \alpha_{k+1} I_{\alpha_k}^{(2)} |\nabla u|^4 (u + \epsilon)^{-3} + I_{a, \alpha_k, \beta_k}^{(4)} |x|^{a-2} u^{(p+1)/2} \\ + I_{\epsilon, \alpha_k, \beta_k}^{(3)} |x|^a u^{(p+1)/2} \left| \frac{\nabla u}{u} + \frac{a\beta_{k+1} (\frac{1}{2}(p+1) - \alpha_{k+1} u / (u + \epsilon))}{2I_{\epsilon, \alpha_k, \beta_k}^{(3)}} \frac{x}{|x|^2} \right|^2, \end{aligned} \tag{3-3}$$

where

$$\begin{aligned} I_{\epsilon, \alpha_k, \beta_k}^{(1)} &:= 1 - \frac{p+1}{2} \beta_{k+1}^2 + \frac{2}{n} \alpha_{k+1} \beta_k^2 \frac{u}{u + \epsilon}, \\ I_{\alpha_k}^{(2)} &:= \frac{2}{n} (\alpha_{k+1} + \alpha_k + 1)^2 - 2\alpha_{k+1} (\alpha_{k+1} + 1) + \alpha_{k+1}, \\ I_{\epsilon, \alpha_k, \beta_k}^{(3)} &:= \frac{4}{n} \alpha_{k+1} \beta_k (\alpha_{k+1} + \alpha_k + 1) \frac{u^2}{(u + \epsilon)^2} + \beta_{k+1} \alpha_{k+1} \frac{u^2}{(u + \epsilon)^2} \\ &\quad - (p+1) \beta_{k+1} \alpha_{k+1} \frac{u}{u + \epsilon} + \frac{p+1}{2} \left( \frac{p-1}{2} - \alpha_{k+1} \frac{u}{u + \epsilon} \right) \beta_{k+1}, \\ I_{a, \epsilon, \alpha_k, \beta_k}^{(4)} &:= \frac{a}{2} \beta_{k+1} \left( n + \frac{a}{2} - 2 \right) - \frac{a^2 \beta_{k+1}^2 (\frac{1}{2}(p+1) - \alpha_{k+1} u / (u + \epsilon))^2}{4I_{\epsilon, \alpha_k, \beta_k}^{(3)}}. \end{aligned}$$

*Proof.* For the sake of simplicity in calculations, set  $b := \frac{1}{2}a$  and  $q := \frac{1}{2}(p+1)$ . From (3-2), the function  $w_{k+1}$  is defined as

$$w_{k+1} := \Delta u + \alpha_{k+1} |\nabla u|^2 (u + \epsilon)^{-1} + \beta_{k+1} |x|^b u^q.$$

Taking Laplacian of  $w_{k+1}$  and using (1-1), we get

$$\Delta w_{k+1} = \Delta^2 u + \alpha_{k+1} \Delta (|\nabla u|^2 (u + \epsilon)^{-1}) + \beta_{k+1} \Delta (|x|^b u^q) = |x|^a u^p + I + J, \tag{3-4}$$

where  $I := \alpha_{k+1} \Delta (|\nabla u|^2 (u + \epsilon)^{-1})$  and  $J := \beta_{k+1} \Delta (|x|^b u^q)$ . In what follows, we simplify  $I$  and  $J$  as well as finding lower bounds for these terms. We start with  $J$ :

$$\begin{aligned} \frac{J}{\beta_{k+1}} &= \Delta (|x|^b u^q) = \Delta |x|^b u^q + \Delta u^q |x|^b + 2 \nabla |x|^b \cdot \nabla u^q \\ &= b(n + b - 2) |x|^{b-2} u^q + q(q-1) |x|^b u^{q-2} |\nabla u|^2 + q |x|^b u^{q-1} \Delta u + 2bq |x|^{b-2} u^{q-1} \nabla u \cdot x. \end{aligned}$$

From the definition of  $w_{k+1}$ , we have

$$\Delta u = w_{k+1} - \alpha_{k+1} |\nabla u|^2 (u + \epsilon)^{-1} - \beta_{k+1} |x|^b u^q. \tag{3-5}$$

Substitute this into the previous equation to simplify  $J$  as

$$\frac{J}{\beta_{k+1}} = qu^{q-1}|x|^b w_{k+1} - q\beta_{k+1}u^{2q-1}|x|^{2b} + \left( q(q-1) - q\alpha_{k+1}\frac{u}{u+\epsilon} \right) |x|^b u^{q-2} |\nabla u|^2 + b(n+b-2)|x|^{b-2}u^q + 2bq|x|^{b-2}u^{q-1}\nabla u \cdot x. \tag{3-6}$$

We now simplify  $I$ :

$$\begin{aligned} \frac{I}{\alpha_{k+1}} &= \Delta(|\nabla u|^2(u+\epsilon)^{-1}) = \sum_{i,j} \partial_{jj}(u_i^2(u+\epsilon)^{-1}) \\ &= 2(u+\epsilon)^{-1} \sum_{i,j} (\partial_{ij}u)^2 + 2(u+\epsilon)^{-1}\nabla u \cdot \nabla \Delta u - 4(u+\epsilon)^{-2} \sum_{i,j} \partial_i u \partial_j u \partial_{ij}u \\ &\quad - |\nabla u|^2(u+\epsilon)^{-2} \Delta u + 2|\nabla u|^4(u+\epsilon)^{-3}. \end{aligned}$$

Again substituting (3-5) into the term  $2(u+\epsilon)^{-1}\nabla u \cdot \nabla \Delta u$  that appears above, we get

$$\begin{aligned} \frac{I}{\alpha_{k+1}} &= 2(u+\epsilon)^{-1} \sum_{i,j} (\partial_{ij}u)^2 - 4(u+\epsilon)^{-2} \sum_{i,j} \partial_i u \partial_j u \partial_{ij}u + 2|\nabla u|^4(u+\epsilon)^{-3} - |\nabla u|^2(u+\epsilon)^{-3} \Delta u \\ &\quad + 2(u+\epsilon)^{-1}\nabla u \cdot \nabla w_{k+1} - 2\alpha_{k+1}(u+\epsilon)^{-1}\nabla u \cdot (|\nabla u|^2(u+\epsilon)^{-1}) \\ &\quad - 2\beta_{k+1}(u+\epsilon)^{-1}\nabla u \cdot \nabla(|x|^b u^q). \end{aligned}$$

Then, collecting similar terms, we obtain

$$\begin{aligned} \frac{I}{\alpha_{k+1}} - 2(u+\epsilon)^{-1}\nabla u \cdot \nabla w_{k+1} &= 2(u+\epsilon)^{-1} \sum_{i,j} (\partial_{ij}u)^2 - 4(\alpha_{k+1}+1)(u+\epsilon)^{-2} \sum_{i,j} \partial_i u \partial_j u \partial_{ij}u + 2(\alpha_{k+1}+1)|\nabla u|^4(u+\epsilon)^{-3} \\ &\quad - |\nabla u|^2(u+\epsilon)^{-2} \Delta u - 2\beta_{k+1}b|x|^{b-2}(u+\epsilon)^{-1}u^q \nabla u \cdot x - 2\beta_{k+1}q|x|^b u^{q-1}(u+\epsilon)^{-1}|\nabla u|^2. \end{aligned}$$

Completing the square, we get

$$\begin{aligned} \frac{I}{\alpha_{k+1}} - 2(u+\epsilon)^{-1}\nabla u \cdot \nabla w_{k+1} &= 2(u+\epsilon)^{-1} \sum_{i,j} (\partial_{ij}u - (\alpha_{k+1}+1)(u+\epsilon)^{-1}\partial_i u \partial_j u)^2 - 2\alpha_{k+1}(\alpha_{k+1}+1)|\nabla u|^4(u+\epsilon)^{-3} \\ &\quad - |\nabla u|^2(u+\epsilon)^{-2} \Delta u - 2\beta_{k+1}b|x|^{b-2}(u+\epsilon)^{-1}u^q \nabla u \cdot x - 2\beta_{k+1}q|x|^b u^{q-1}(u+\epsilon)^{-1}|\nabla u|^2. \tag{3-7} \end{aligned}$$

Note that, for any  $n \times n$  matrix  $A = (a_{i,j})$ , the Hilbert–Schmidt norm is defined by  $\|A\|_2 = \sqrt{\sum_{i,j} |a_{i,j}|^2} = \sqrt{\text{trace}(AA^*)}$ , where  $A^*$  denotes the conjugate transpose of  $A$ . From the Cauchy–Schwarz inequality, the following inequality holds:

$$|\text{trace } A|^2 = |(A, I)|^2 \leq \|A\|_2^2 \|I\|_2^2 = n \sum_{i,j} |a_{i,j}|^2. \tag{3-8}$$

Set  $a_{i,j} := \partial_{ij}u - (\alpha_{k+1} + 1)(u + \epsilon)^{-1}\partial_iu\partial_ju$  in (3-8) to get

$$\sum_{i,j}^n (\partial_{ij}u - (\alpha_{k+1} + 1)(u + \epsilon)^{-1}\partial_iu\partial_ju)^2 \geq \frac{1}{n}(\Delta u - (\alpha_{k+1} + 1)(u + \epsilon)^{-1}|\nabla u|^2)^2.$$

From this lower bound for the Hessian and (3-7), we get

$$\begin{aligned} \frac{I}{\alpha_{k+1}} - 2(u + \epsilon)^{-1}\nabla u \cdot \nabla w_{k+1} &\geq \frac{2}{n}(u + \epsilon)^{-1}(\Delta u - (\alpha_{k+1} + 1)(u + \epsilon)^{-1}|\nabla u|^2)^2 \\ &\quad - 2\alpha_{k+1}(\alpha_{k+1} + 1)|\nabla u|^4(u + \epsilon)^{-3} - |\nabla u|^2(u + \epsilon)^{-2}\Delta u + T_k, \end{aligned} \quad (3-9)$$

where

$$T_k := -2\beta_{k+1}b|x|^{b-2}(u + \epsilon)^{-1}u^q\nabla u \cdot x - 2\beta_{k+1}q|x|^bu^{q-1}(u + \epsilon)^{-1}|\nabla u|^2.$$

Note also that, from the assumption  $w_k \leq 0$ , we have the upper bound on the Laplacian operator  $\Delta u \leq -\alpha_k|\nabla u|^2(u + \epsilon)^{-1} - \beta_k|x|^bu^q$ . Elementary calculations show that, if  $t \leq t_* \leq 0$  and  $s \geq 0$ , then  $(t - s)^2 \geq t_*^2 - 2t_*s + s^2$ . Set the parameters as  $t = \Delta u$ ,  $t_* = -\alpha_k|\nabla u|^2(u + \epsilon)^{-1} - \beta_k|x|^bu^q$  and  $s = (\alpha_{k+1} + 1)(u + \epsilon)^{-1}|\nabla u|^2$  to get the following lower bound on the square term that appears in (3-9):

$$\begin{aligned} &(\Delta u - (\alpha_{k+1} + 1)(u + \epsilon)^{-1}|\nabla u|^2)^2 \\ &\geq (\alpha_k|\nabla u|^2(u + \epsilon)^{-1} + \beta_k|x|^bu^q)^2 + 2(\alpha_k|\nabla u|^2(u + \epsilon)^{-1} + \beta_k|x|^bu^q)(\alpha_{k+1} + 1)(u + \epsilon)^{-1}|\nabla u|^2 \\ &\quad + (\alpha_{k+1} + 1)^2(u + \epsilon)^{-2}|\nabla u|^4. \end{aligned} \quad (3-10)$$

Substitute (3-5) into the term  $-|\nabla u|^2(u + \epsilon)^{-2}\Delta u$  that appears in (3-9) to eliminate the Laplacian operator. Then, apply inequality (3-10) to simplify (3-9) as

$$\begin{aligned} &\frac{I}{\alpha_{k+1}} - 2(u + \epsilon)^{-1}\nabla u \cdot \nabla w_{k+1} \\ &\geq \frac{2}{n}(u + \epsilon)^{-1}((\alpha_{k+1} + \alpha_k + 1)^2|\nabla u|^4(u + \epsilon)^{-2} + \beta_k^2|x|^{2b}u^{2q} + 2\beta_k(\alpha_{k+1} + \alpha_k + 1)|x|^bu^q(u + \epsilon)^{-1}|\nabla u|^2) \\ &\quad - w_{k+1}(u + \epsilon)^{-2}|\nabla u|^2 - \alpha_{k+1}(2\alpha_{k+1} + 1)|\nabla u|^4(u + \epsilon)^{-3} + \beta_{k+1}|x|^bu^q(u + \epsilon)^{-2}|\nabla u|^2 + T_k. \end{aligned}$$

Collecting similar terms and using the value of  $T_k$ , we end up with

$$\begin{aligned} &\frac{I}{\alpha_{k+1}} - 2(u + \epsilon)^{-1}\nabla u \cdot \nabla w_{k+1} + w_{k+1}(u + \epsilon)^{-2}|\nabla u|^2 \\ &\geq \frac{2}{n}\beta_k^2|x|^{2b}u^{2q}(u + \epsilon)^{-1} + I_{\alpha_k}^{(2)}|\nabla u|^4(u + \epsilon)^{-3} + S_{\epsilon,\alpha_k,\beta_k}|\nabla u|^2u^{q-2}|x|^b - 2\beta_{k+1}b|x|^{b-2}(u + \epsilon)^{-1}u^q\nabla u \cdot x, \end{aligned}$$

where

$$\begin{aligned} I_{\alpha_k}^{(2)} &:= \frac{2}{n}(\alpha_{k+1} + \alpha_k + 1)^2 - 2\alpha_{k+1}(\alpha_{k+1} + 1) + \alpha_{k+1}, \\ S_{\epsilon,\alpha_k,\beta_k} &:= \frac{4}{n}\beta_k(\alpha_{k+1} + \alpha_k + 1)\frac{u^2}{(u + \epsilon)^2} + \beta_{k+1}\frac{u^2}{(u + \epsilon)^2} - 2\beta_{k+1}q\frac{u}{u + \epsilon}. \end{aligned}$$

Therefore, the following lower bound for  $I$  holds:

$$I \geq 2\alpha_{k+1}(u + \epsilon)^{-1} \nabla u \cdot \nabla w_{k+1} - \alpha_{k+1} w_{k+1} (u + \epsilon)^{-2} |\nabla u|^2 + \frac{2}{n} \alpha_{k+1} \beta_k^2 |x|^{2b} u^{2q} (u + \epsilon)^{-1} + I_{\alpha_k} |\nabla u|^4 (u + \epsilon)^{-3} + S_{\epsilon, \alpha_k, \beta_k} |\nabla u|^2 u^{q-2} |x|^b - 2\beta_{k+1} b |x|^{b-2} (u + \epsilon)^{-1} u^q \nabla u \cdot x. \tag{3-11}$$

Finally, applying this lower bound for  $I$  and the lower bound given for  $J$  in (3-6), from (3-3) we get

$$\begin{aligned} &\Delta w_{k+1} - 2\alpha_{k+1}(u + \epsilon)^{-1} \nabla u \cdot \nabla w_{k+1} + \alpha_{k+1}(u + \epsilon)^{-2} |\nabla u|^2 w_{k+1} - \beta_{k+1} q u^{q-1} |x|^b w_{k+1} \\ &\geq |x|^a u^p \left( 1 - q\beta_{k+1}^2 + \frac{2}{n} \alpha_{k+1} \beta_k^2 \frac{u}{u + \epsilon} \right) + \alpha_{k+1} I_{\alpha_k}^{(2)} |\nabla u|^4 (u + \epsilon)^{-3} \\ &\quad + \left( \alpha_{k+1} S_{\epsilon, \alpha_k, \beta_k} + \left( q(q - 1) - \alpha_{k+1} q \frac{u}{u + \epsilon} \right) \beta_{k+1} \right) |\nabla u|^2 u^{q-2} |x|^b \\ &\quad + 2b\beta_{k+1} \left( q - \alpha_{k+1} \frac{u}{u + \epsilon} \right) |x|^{b-2} u^{q-1} \nabla u \cdot x + b\beta_{k+1} (n + b - 2) |x|^{b-2} u^q. \end{aligned}$$

Completing the square finishes the proof. □

#### 4. Proof of Theorem 1.3 via iteration arguments

To apply the iteration argument, we need to develop a maximum principle argument for the equation

$$\Delta w - 2\alpha(u + \epsilon)^{-1} \nabla u \cdot \nabla w + \alpha w (u + \epsilon)^{-2} |\nabla u|^2 - \frac{1}{2} \beta (p + 1) |x|^{a/2} u^{(p-1)/2} w = f(x) \geq 0 \quad \text{in } \mathbb{R}^n \tag{4-1}$$

that appears in Proposition 3.1, where  $\alpha$  and  $\beta$  are positive constants,  $u$  is a solution of (1-1) and  $w, f \in C^\infty(\mathbb{R}^n)$ .

**Lemma 4.1.** *Suppose that  $w$  is a solution of the differential inequality (4-1), where  $u$  is a solution of (1-1) and*

$$w = \Delta u + \alpha(u + \epsilon)^{-1} |\nabla u|^2 + \beta |x|^{a/2} u^{(p+1)/2} \tag{4-2}$$

for positive constants  $\epsilon, \alpha$  and  $\beta$ . Then, assuming that  $p + 1 > 2\alpha$ ,

$$\Delta \tilde{w} \geq 0 \quad \text{on } \{w \geq 0\} \subset \mathbb{R}^n, \tag{4-3}$$

where  $\tilde{w} = (u + \epsilon)^t w$  for  $t = -\alpha$ .

*Proof.* Straightforward calculations show that

$$\Delta \tilde{w} = (u + \epsilon)^t \Delta w + 2t(u + \epsilon)^{t-1} \nabla u \cdot \nabla w + t(u + \epsilon)^{t-1} w \Delta u + t(t - 1)w(u + \epsilon)^{t-2} |\nabla u|^2.$$

We now add and subtract two terms,  $\frac{1}{2} \beta (p + 1) |x|^{a/2} u^{(p-1)/2} (u + \epsilon)^t w$  and  $t w (u + \epsilon)^{t-2} |\nabla u|^2$ , to the above identity and collect similar terms to get

$$\begin{aligned} \Delta \tilde{w} &= (u + \epsilon)^t \left( \Delta w + 2t(u + \epsilon)^{-1} \nabla u \cdot \nabla w - t w (u + \epsilon)^{-2} |\nabla u|^2 - \frac{1}{2} \beta (p + 1) |x|^{a/2} u^{(p-1)/2} w \right) \\ &\quad + \frac{1}{2} \beta (p + 1) |x|^{a/2} u^{(p-1)/2} (u + \epsilon)^t w + t w (u + \epsilon)^{t-2} |\nabla u|^2 + t(u + \epsilon)^{t-1} w \Delta u \\ &\quad + t(t - 1)w(u + \epsilon)^{t-2} |\nabla u|^2. \end{aligned}$$

From the fact that  $t = -\alpha$  and  $w$  satisfies (4-1), we get

$$\Delta \tilde{w} \geq \frac{1}{2}\beta(p+1)|x|^{a/2}u^{(p-1)/2}(u+\epsilon)^t w + t(u+\epsilon)^{t-1}w\Delta u + t^2w(u+\epsilon)^{t-1}\frac{|\nabla u|^2}{u+\epsilon}.$$

Note that we can eliminate the gradient term using (4-2), that is,

$$\alpha(u+\epsilon)^{-1}|\nabla u|^2 = w - \Delta u - \beta|x|^{a/2}u^{(p+1)/2}.$$

Therefore, after collecting similar terms we get

$$\begin{aligned} \Delta \tilde{w} &\geq \frac{t^2}{\alpha}w^2(u+\epsilon)^{t-1} + (u+\epsilon)^{t-1}wt\left(1 - \frac{t}{\alpha}\right)\Delta u \\ &\quad + \beta(u+\epsilon)^{t-1}|x|^{a/2}u^{(p-1)/2}w\left(\frac{(p+1)\epsilon}{2} + u\left(\frac{p+1}{2} - \frac{t^2}{\alpha}\right)\right) \\ &=: R_1 + R_2 + R_3. \end{aligned}$$

We claim that the above three terms,  $R_1$ ,  $R_2$  and  $R_3$ , are nonnegative when  $w \geq 0$ . From the fact that  $\alpha > 0$  one can see that  $R_1$  is nonnegative. From the definition of  $t = -\alpha < 0$ , we have  $t(1 - t/\alpha) = -2\alpha < 0$ . This together with Proposition 2.3, that is,  $\Delta u \leq 0$ , confirms that  $R_2$  is nonnegative. Positivity of  $R_3$  is an immediate consequence of the assumptions:  $\beta$  is positive and  $\frac{1}{2}(p+1) - t^2/\alpha = \frac{1}{2}(p+1) - \alpha$  is also positive. This finishes the proof.  $\square$

We now apply Lemma 4.1 to show that any solution  $w$  of (4-1) is negative.

**Lemma 4.2.** *Suppose that  $\tilde{w}$  and  $w$  as in Lemma 4.1. Let  $u$  be a bounded solution of (1-1); then  $w \leq 0$ .*

*Proof.* The methods and ideas that we apply in the proof are motivated by Souplet [2009]. Multiply (4-3) by  $\tilde{w}_+^s$ , where  $s > 0$  is a parameter that will be determined later. Then, integration by parts over  $B_R$  gives us

$$0 \leq \int_{B_R} \Delta \tilde{w} \tilde{w}_+^s = -s \int_{B_R} |\nabla \tilde{w}_+|^2 \tilde{w}_+^{s-1} + R^{n-1} \int_{S^{n-1}} \tilde{w}_r \tilde{w}_+^s. \tag{4-4}$$

Therefore,

$$\int_{B_R} |\nabla \tilde{w}_+|^2 \tilde{w}_+^{s-1} \leq \frac{1}{s(s+1)} R^{n-1} \int_{S^{n-1}} (\tilde{w}_+^{s+1})_r = C(s) R^{n-1} I'(R), \tag{4-5}$$

where

$$I(R) := \int_{S^{n-1}} \tilde{w}_+^{s+1} = \int_{S^{n-1}} (u+\epsilon)^{-(s+1)\alpha} w_+^{s+1}$$

and  $C(s)$  is a constant independent of  $R$ . Note that  $w$ , given as  $w = \Delta u + \alpha|\nabla u|^2(u+\epsilon)^{-1} + \beta|x|^{a/2}u^{(p+1)/2}$ , satisfies  $w \geq 0$  if and only if  $-\Delta u \leq \alpha|\nabla u|^2(u+\epsilon)^{-1} + \beta|x|^{a/2}u^{(p+1)/2}$ . Therefore,

$$w_+^{s+1} \leq C|\nabla u|^{2(s+1)}(u+\epsilon)^{-(s+1)} + C|x|^{(s+1)a/2}u^{(s+1)(p+1)/2}, \tag{4-6}$$



where  $C = C(\alpha, \beta, s)$ . Applying this upper bound for  $w_+$ , we can get an upper bound for  $I(R)$ :

$$\begin{aligned} I(R) &\leq C \int_{S^{n-1}} (u + \epsilon)^{-(s+1)(\alpha+1)} |\nabla u|^{2(s+1)} + CR^{(s+1)a/2} \int_{S^{n-1}} (u + \epsilon)^{-\alpha(s+1)} u^{(s+1)(p+1)/2} \\ &\leq C(\epsilon) \int_{S^{n-1}} |\nabla u|^{2(s+1)} + C(\epsilon)R^{(s+1)a/2} \int_{S^{n-1}} u^{(s+1)(p+1)/2} \\ &=: C(\epsilon)(I_1(R) + I_2(R)). \end{aligned} \tag{4-7}$$

In what follows, we show that there is a sequence  $R$  such that the two terms  $I_1(R)$  and  $I_2(R)$  decay to zero for a fixed  $\epsilon$ . We start with  $I_2(R)$ , which includes an integral of a positive power of  $u$  over the sphere. Due to the boundedness assumption on  $u$ , it is straightforward to relate this term to  $L^p$  estimates of  $u$  over the sphere. As a matter of fact, if  $(s + 1)(p + 1) > 2p$  then, from the boundedness of  $u$ , we have

$$\int_{S^{n-1}} u^{(s+1)(p+1)/2} \leq C(n) \|u\|_{L^p(S^{n-1})}^p \tag{4-8}$$

and for the case  $(s + 1)(p + 1) \leq 2p$  we can use Hölder’s inequality to get

$$\int_{S^{n-1}} u^{(s+1)(p+1)/2} \leq C(n, p) \|u\|_{L^p(S^{n-1})}^{(p+1)(s+1)/2}. \tag{4-9}$$

So, to prove a decay estimate for  $I_2(R)$  we need to construct a decay estimate for  $\|u\|_{L^p(S^{n-1})}$ . On the other hand, we apply Lemma 2.10 to get an upper bound for the first term in (4-7),  $I_1(R)$ . In fact, from Lemma 2.10 with  $i = 1$ ,  $\tau = 2(s + 1)$  and  $t = 2$ , we have

$$\begin{aligned} \|D_x u\|_{L^{2(s+1)}(S^{n-1})} &\leq C \|D_\theta D_x u\|_{L^2(S^{n-1})} + C \|D_x u\|_{L^1(S^{n-1})} \\ &\leq CR \|D_x^2 u\|_{L^2(S^{n-1})} + C \|D_x u\|_{L^1(S^{n-1})} \end{aligned} \tag{4-10}$$

for  $s = 2/(n - 3)$ . In order to get a decay estimate for  $I_1(R)$ , we need decay estimates for the two terms in the right-hand side of (4-10),  $\|D_x^2 u\|_{L^2(S^{n-1})}$  and  $\|D_x u\|_{L^1(S^{n-1})}$ .

We now apply the elliptic estimates given in Section 2 to provide decay estimates for  $\|u\|_{L^p(S^{n-1})}$ ,  $\|D_x u\|_{L^1(S^{n-1})}$  and  $\|D_x^2 u\|_{L^2(S^{n-1})}$ . To do so we first find appropriate upper bounds for these terms on the ball of radius  $R$ . Then we use certain measure-comparison arguments to construct decay estimates over the sphere. So, from Lemma 2.7 with  $\tau = 2$ , we get

$$\int_{R/2}^R \|D_x^2 u\|_{L^2(S^{n-1})}^2 r^{n-1} dr \leq C \int_{B_{2R} \setminus B_{R/4}} |\Delta u|^2 + CR^{-4} \int_{B_{2R} \setminus B_{R/4}} u^2. \tag{4-11}$$

We now apply Corollary 2.9 and Corollary 2.2 to get a decay estimate for the right-hand side of (4-11), namely,

$$\begin{aligned} R^{-4} \int_{B_{2R} \setminus B_{R/4}} u^2 &\leq CR^{-4} \int_{B_{2R} \setminus B_{R/4}} u \leq CR^{-4} R^{n-(a+4)/(p-1)} = CR^{n-(a+4p)/(p-1)}, \\ \int_{B_{2R} \setminus B_{R/4}} |\Delta u|^2 &\leq CR^{n-(a+4p)/(p-1)}, \end{aligned}$$

where  $C$  is independent from  $R$ . From this and (4-11), we obtain the desired decay estimate on the Hessian operator of  $u$ ,

$$\int_{R/2}^R \|D_x^2 u\|_{L^2(S^{n-1})}^2 r^{n-1} dr \leq C R^{n-(4p+a)/(p-1)}. \tag{4-12}$$

Similarly, from Corollary 2.6 and Lemma 2.1, we have

$$\int_{R/2}^R \|D_x u\|_{L^1(S^{n-1})} r^{n-1} dr \leq C R^{n-(p+3+a)/(p-1)}, \tag{4-13}$$

$$\int_{R/2}^R \|u\|_{L^p(S^{n-1})}^p r^{n-1} dr \leq C R^{n-(a+4)p/(p-1)}. \tag{4-14}$$

Now let's define the following sets. These sets are meant to facilitate our arguments towards construction of decay estimates for  $\|u\|_{L^p(S^{n-1})}$ ,  $\|D_x u\|_{L^1(S^{n-1})}$  and  $\|D_x^2 u\|_{L^2(S^{n-1})}$ . For a large number  $M$ , which will be determined later, define

$$\begin{aligned} \Gamma_1(R) &:= \{r \in (R/2, R) : \|u\|_{L^p(S^{n-1})}^p > M R^{-(a+4)p/(p-1)}\}, \\ \Gamma_2(R) &:= \{r \in (R/2, R) : \|D_x u\|_{L^1(S^{n-1})} > M R^{-(p+3+a)/(p-1)}\}, \\ \Gamma_3(R) &:= \{r \in (R/2, R) : \|D_x^2 u\|_{L^2(S^{n-1})}^2 > M R^{-(a+4)p/(p-1)}\}. \end{aligned}$$

We claim that  $|\Gamma_i(R)| \leq R/4$  for  $1 \leq i \leq 3$ : Using (4-12), we get

$$\begin{aligned} C &\geq R^{-n+(a+4p)/(p-1)} \int_{R/2}^R \|D_x^2 u\|_{L^2(S^{n-1})}^2 r^{n-1} dr \\ &\geq N R^{-n+(a+4p)/(p-1)} R^{n-1} \int_{R/2}^R \|D_x^2 u\|_{L^2(S^{n-1})}^2 dr \\ &\geq N M R^{-n+(a+4p)/(p-1)} R^{n-1} \int_{|\Gamma_3(R)|} R^{-(a+4p)/(p-1)} dr \\ &\geq N M R^{-n+(a+4p)/(p-1)} R^{n-1} |\Gamma_3(R)| R^{-(a+4p)/(p-1)} = N M |\Gamma_3(R)| R^{-1}, \end{aligned}$$

where  $N = (\frac{1}{2})^{n-1}$ . Therefore,  $|\Gamma_3(R)| \leq C R/NM$ . Now, choosing  $M$  to be large enough, that is,  $M > 4C/N$ , we get  $|\Gamma_3(R)| \leq R/4$ . Similarly, applying (4-13) and (4-14), one can show that  $|\Gamma_i(R)| \leq R/4$  for  $i = 1, 2$ . Hence,  $|\Gamma_i(R)| \leq R/4$  for  $1 \leq i \leq 3$  while  $\Gamma_i(R) \subset (R/2, R)$ . So, we can find a sequence of  $\tilde{R}$  such that

$$\tilde{R} \in (R/2, R) \setminus \bigcup_{i=1}^{i=3} \Gamma_i(R) \neq \emptyset. \tag{4-15}$$

Therefore, for the sequence  $\tilde{R}$ , we obtain

$$\|u\|_{L^p(S^{n-1})}^p \leq M R^{-(a+4)p/(p-1)}, \tag{4-16}$$

$$\|D_x u\|_{L^1(S^{n-1})} \leq M R^{-(p+3+a)/(p-1)}, \tag{4-17}$$

$$\|D_x^2 u\|_{L^2(S^{n-1})}^2 \leq M R^{-(a+4p)/(p-1)}. \tag{4-18}$$

Substituting (4-16) into (4-8) and (4-9), we get the decay estimate on  $I_2(R)$

$$\begin{aligned}
 I_2(R) &\leq C\chi\{(s+1)(p+1) > 2p\}R^{(s+1)/2a-(a+4)p/(p-1)} \\
 &\quad + C\chi\{(s+1)(p+1) \leq 2p\}R^{(s+1)a/2-(a+4)(p+1)(s+1)/(2(p-1))} \\
 &= C\chi\{(s+1)(p+1) > 2p\}R^{-\eta_1} + C\chi\{(s+1)(p+1) > 2p\}R^{-\eta_2}, \tag{4-19}
 \end{aligned}$$

where  $\chi$  is the characteristic function,  $\eta_1 := a(p/(p-1) - \frac{1}{2}(s+1)) + 4p/(p-1) > 0$  and  $\eta_2 := (s+1)(ap + 2(p+1))/(p+1) > 0$ . Note that we have used the fact that  $p/(p-1) - \frac{1}{2}(s+1) > 0$  because  $0 < s = 2/(n-3) \leq 1$  when  $n \geq 5$ . On the other hand, substituting (4-17) and (4-18) into the Sobolev embedding (4-10), we get

$$\|D_x u\|_{L^{2(s+1)}(S^{n-1})} \leq CR^{1-(a+4p)/(p-1)} + CR^{-(p+3+a)/(p-1)} = 2CR^{-(p+3+a)/(p-1)}. \tag{4-20}$$

From this and the definition of  $I_1(R)$ , we end up with the decay estimate on  $I_1(R)$

$$I_1(R) = \int_{S^{n-1}} |\nabla u|^{2(s+1)} \leq CR^{-2(p+3+a)(s+1)/(p-1)} = CR^{-\eta_3}, \tag{4-21}$$

where  $\eta_3 := 2(p+3+a)(s+1)/(p-1) > 0$ . Finally, from (4-21) and (4-19), we observe that

$$I(R) \leq CR^{-\eta} \quad \text{for all } R > 1,$$

where  $\eta := \min\{\eta_1, \eta_2, \eta_3\} > 0$ . So,  $I(R) \rightarrow 0$  as  $R \rightarrow \infty$ . Note that  $\tilde{R} \rightarrow \infty$  as  $R \rightarrow \infty$ . Since  $I(R)$  is a positive function and converges to zero, there is a sequence such that the functional  $I'(R)$  is nonpositive. Therefore, (4-5) yields

$$\int_{B_R} |\nabla \tilde{w}_+|^2 \tilde{w}_+^{s-1} \leq 0. \tag{4-22}$$

Hence,  $\tilde{w}_+$  has to be a constant. From the continuity of  $\tilde{w}$ , we have  $\tilde{w} \equiv C$ . Note that the constant  $C$  cannot be strictly positive. So,  $\tilde{w}_+ = 0$  and therefore  $w_+ = 0$ . This finishes the proof.  $\square$

Note that Lemma 4.1 and Lemma 4.2 imply an iteration argument for the sequence of functions, for  $k \geq -1$ ,

$$w_k = \Delta u + \alpha_k(u + \epsilon)^{-1} |\nabla u|^2 + \beta_k |x|^{a/2} u^{(p+1)/2} \tag{4-23}$$

as long as the right-hand side of (3-3) stays nonnegative. For the rest of this section, we construct sequences  $\{\alpha_k\}_{k=-1}$  and  $\{\beta_k\}_{k=-1}$  such that the right-hand side of (3-3) is nonnegative.

**Constructing sequences  $\alpha_k$  and  $\beta_k$ .** In this part, we define sequences  $\alpha_k$  and  $\beta_k$  needed for the iteration argument.

**Lemma 4.3.** *Suppose  $\alpha_0 = 0$  and define*

$$\alpha_{k+1} := \frac{4(\alpha_k + 1) - n + \sqrt{n(16\alpha_k^2 + 24\alpha_k + n + 8)}}{4(n-1)}. \tag{4-24}$$

*Then  $(\alpha_k)_k$  is a positive, bounded and increasing sequence that converges to  $\alpha := 2/(n-4)$  provided  $n > 4$  and  $p > 1$ . Moreover, for this choice of  $(\alpha_k)_k$ , the sequence  $I_{\alpha_k}^{(2)}$  of coefficients defined in Proposition 3.1 equals zero.*

*Proof.* It is straightforward to show that  $\alpha_k > 0$  for any  $k \geq 0$ . Also, direct calculations show that  $\alpha_k \rightarrow \alpha := 2/(n-4)$  provided  $\alpha_k$  is convergent. Note that  $\alpha_1 = (4-n+\sqrt{n^2+8n})/(4n-4) < 2/(n-4)$  and, by induction, one can see that  $\alpha_k \leq \alpha$  for all  $k \geq 0$ . Lastly, we show that  $\alpha_k$  is an increasing sequence: For any  $k$ ,

$$\begin{aligned} \alpha_{k+1} - \alpha_k &= \frac{\sqrt{n(16\alpha_k^2 + 24\alpha_k + n + 8)} - ((n-4) + 4a_k(n-2))}{4(n-1)} \\ &= \frac{8(n-1)(n-4)(2\alpha_k + 1)}{S_{n,k}} \left( \frac{2}{n-4} - \alpha_k \right), \end{aligned}$$

where  $S_{n,k} = \sqrt{n(16\alpha_k^2 + 24\alpha_k + n + 8)} + (n-4) + 4a_k(n-2) > 0$ . Therefore, from the fact that  $\alpha_k \leq \alpha = 2/(n-4)$ , we get the desired result. □

Similarly, we provide an explicit formula for the sequence  $\beta_k$ :

**Lemma 4.4.** *Suppose  $\beta_0 = \sqrt{2/(p+1)}$  and define*

$$\beta_{k+1} := \sqrt{\frac{2}{p+1} + \frac{4}{(p+1)n} \alpha_k \beta_k^2}, \tag{4-25}$$

where  $(\alpha_k)_k$  is as in [Lemma 4.3](#). Then  $(\beta_k)_k$  is a positive, bounded and increasing sequence that converges to  $\beta := \sqrt{2/((p+1) - c_n)}$ , where  $c_n = 8/(n(n-4))$  provided that  $n > 4$  and  $p > 1$ . Moreover, for this choice of  $(\alpha_k)_k$  and  $(\beta_k)_k$ , the sequence  $I_{0,\alpha_k,\beta_k}^{(1)}$  of coefficients defined in [Proposition 3.1](#) is strictly positive.

*Proof.* The sequence  $(\beta_k)_k$  for all  $k \geq 0$  is positive. Note that boundedness of the sequence  $(\alpha_k)_k$  forces the boundedness of the  $(\beta_k)_k$ , meaning that  $\beta_{k+1} \leq \sqrt{2/(p+1) + (4\alpha/((p+1)n))\beta_k^2}$  for any  $k$ . By straightforward calculations we get

$$\beta_{k+1}^2 \leq \frac{2}{p+1} \sum_{i=0}^{k+1} \left( \frac{4\alpha}{n(p+1)} \right)^i.$$

Note that  $4\alpha/(n(p+1)) = 8/(n(n-4)(p+1)) < 1$  provided that  $n > 4$  and  $p > 1$ . Therefore,  $\sum_{i=0}^{\infty} (4\alpha/(n(p+1)))^i < \infty$ . This proves the boundedness of  $(\beta_k)_k$ .

Since  $(\alpha_k)_{k=0}$  is an increasing sequence, the sequence  $(\beta_k)_{k=0}$  will be nondecreasing by induction. Note that

$$\beta_1 = \beta_0 \quad \text{and} \quad \beta_2 = \sqrt{\frac{2}{p+1} + \frac{8}{(p+1)^2 n} \frac{4-n+\sqrt{n^2+8n}}{4n-4}} > \beta_1 = \sqrt{\frac{2}{p+1}}.$$

Suppose that  $\beta_{k-1} \leq \beta_k$  for a certain index  $k \geq 2$ ; then we apply the fact that  $\alpha_k \geq \alpha_{k-1}$  to show  $\beta_k \leq \beta_{k+1}$ . This can be found as a consequence of

$$\beta_{k+1} - \beta_k = \frac{\beta_{k+1}^2 - \beta_k^2}{\beta_{k+1} + \beta_k} = \frac{4}{(p+1)n(\beta_{k+1} + \beta_k)} (\beta_k^2 \alpha_k - \beta_{k-1}^2 \alpha_{k-1}) \geq \frac{4\alpha_{k-1}(\beta_k + \beta_{k-1})}{(p+1)n(\beta_{k+1} + \beta_k)} (\beta_k - \beta_{k-1}).$$

So,  $(\beta_k)_k$  is convergent and converges to  $\beta := \sqrt{2n(n-4)/((p+1)(n-4)n-8)}$ . Since  $(p+1)n(n-4) > 8$  for  $p > 1$  and  $n > 4$ ,  $\beta$  is well-defined. □

Note that, based on the definition of the sequences  $\{\alpha_k\}_{k=-1}$  and  $\{\beta_k\}_{k=-1}$ , we concluded that  $I_{0,\alpha_k,\beta_k}^{(1)} > 0$  and  $I_{\alpha_k}^{(2)} = 0$ . In the next two lemmata we investigate the positivity of  $I_{\epsilon,\alpha_k,\beta_k}^{(3)}$  and  $I_{a,\epsilon,\alpha_k,\beta_k}^{(4)}$ , the sequences that appeared in (3-3) in Proposition 3.1.

**Lemma 4.5.** *Set  $\epsilon = 0$  in  $I_{\epsilon,\alpha_k,\beta_k}^{(3)}$ , which is defined in Proposition 3.1. Then*

$$I_{0,\alpha_k,\beta_k}^{(3)} \rightarrow I_{0,\alpha,\beta}^{(3)} := \frac{4}{n}\alpha\beta(2\alpha + 1) + \alpha\beta + \beta q(q - 3\alpha - 1) \tag{4-26}$$

as  $k \rightarrow \infty$ . The constant  $I_{0,\alpha,\beta}^{(3)}$  is positive provided  $p > (n + 4)/(n - 4)$  and  $n > 4$ .

*Proof.* Note that when  $p > (n + 4)/(n - 4)$  and  $n > 4$ , we have  $\frac{1}{2}(p + 1) > n/(n - 4)$ . As  $k \rightarrow \infty$ , from Lemma 4.3 and Lemma 4.4, the sequences  $\alpha_k \rightarrow \alpha := 2/(n - 4)$  and  $\beta_k \rightarrow \beta := \sqrt{2/((p + 1) - c_n)}$ . Therefore,

$$\begin{aligned} \frac{I_{0,\alpha,\beta}^{(3)}}{\beta} &= \frac{4}{n} \left( \frac{2}{n-4} \right) \left( \frac{4}{n-4} + 1 \right) + \frac{2}{n-4} + \frac{p+1}{2} \left( \frac{p-1}{2} - \frac{6}{n-4} \right) \\ &= \left( \frac{p+1}{2} \right)^2 - \left( \frac{p+1}{2} \right) \left( \frac{n+2}{n-4} \right) + \frac{2n}{(n-4)^2} \\ &= \left( \frac{p+1}{2} - \frac{n}{n-4} \right) \left( \frac{p+1}{2} - \frac{2}{n-4} \right) > 0. \quad \square \end{aligned}$$

Note that  $I_{a,\epsilon,\alpha_k,\beta_k}^{(4)}$  appears in (3-3) mainly because of the weight function  $|x|^a$ . In other words, we have  $I_{0,\epsilon,\alpha_k,\beta_k}^{(4)} = 0$  in the case of  $a = 0$ .

**Lemma 4.6.** *For any  $k \geq 0$ ,*

$$I_{0,\alpha_k,\beta_k}^{(3)} < \beta_{k+1} \left( \frac{1}{2}(p + 1) - \alpha_{k+1} \right)^2 \tag{4-27}$$

provided  $p > (n + 4)/(n - 4)$  and  $n > 4$ . Therefore, for any  $a \geq 0$  that satisfies the upper bound

$$a \leq A_k := \frac{2(n - 2)I_{0,\alpha_k,\beta_k}^{(3)}}{\beta_{k+1} \left( \frac{1}{2}(p + 1) - \alpha_{k+1} \right)^2 - I_{0,\alpha_k,\beta_k}^{(3)}}, \tag{4-28}$$

the sequence  $I_{a,0,\alpha_k,\beta_k}^{(4)}$  is positive for any  $k$ .

*Proof.* Basic calculations show that

$$\begin{aligned} &\beta_{k+1} \left( \frac{p+1}{2} - \alpha_{k+1} \right)^2 - I_{0,\alpha_k,\beta_k}^{(3)} \\ &= \beta_{k+1} \left( \frac{p+1}{2} - \alpha_{k+1} \right)^2 - \frac{4}{n}\alpha_{k+1}\beta_k(\alpha_{k+1} + \alpha_k + 1) - \alpha_{k+1}\beta_{k+1} - \beta_{k+1} \frac{p+1}{2} \left( \frac{p+1}{2} - 3\alpha_{k+1} - 1 \right) \\ &\geq \beta_{k+1} \left( \left( \frac{p+1}{2} - \alpha_{k+1} \right)^2 - \frac{4}{n}\alpha_{k+1}(\alpha_{k+1} + \alpha_k + 1) - \alpha_{k+1} - \frac{p+1}{2} \left( \frac{p+1}{2} - 3\alpha_{k+1} - 1 \right) \right) \\ &= \beta_{k+1} \left( \frac{n-4}{n}\alpha_{k+1}^2 - \frac{4}{n}\alpha_{k+1}^2 - \frac{4}{n}\alpha_{k+1} + \frac{p-1}{2}\alpha_{k+1} + \frac{p+1}{2} \right), \end{aligned}$$

where we have used the fact that  $\beta_k$  and  $\alpha_k$  are increasing sequences in the first and the second inequality, respectively. Therefore,

$$\begin{aligned} \beta_{k+1} \left( \frac{p+1}{2} - \alpha_{k+1} \right)^2 - I_{0, \alpha_k, \beta_k}^{(3)} &\geq \beta_{k+1} \left( \frac{n-4}{n} \alpha_{k+1}^2 + \alpha_{k+1} \left( \frac{p-1}{2} - \frac{4}{n} \alpha_{k+1} \right) + \frac{p+1}{2} - \frac{4}{n} \alpha_{k+1} \right) \\ &\geq \beta_{k+1} \left( \frac{n-4}{n} \alpha_{k+1}^2 + (\alpha_{k+1} + 1) \left( \frac{p-1}{2} - \frac{4}{n} \alpha \right) \right) \\ &> 0. \end{aligned}$$

Note that in the last inequality we have used the fact that

$$\frac{p-1}{2} - \frac{4}{n} \alpha = \frac{p-1}{2} - \frac{4}{n} \frac{2}{n-4} > \frac{4}{(n-4)n} (n-2) > 0,$$

since  $p > (n+4)/(n-4)$  and  $n > 4$ . □

**Remark 4.7.** It would be interesting if a counterpart of (1-12) could be proved for bounded solutions of the fourth-order semilinear equation  $\Delta^2 u = f(u)$  under certain assumptions on the arbitrary nonlinearity  $f \in C^1(\mathbb{R})$ . We expect that such an inequality could be established for some convex nonlinearity  $f$ .

### Appendix

We would like to mention that given the estimates in Lemma 2.1 and Lemma 2.4, one can provide a somewhat simpler proof of Proposition 2.3, as follows.

*Second proof of Proposition 2.3.* From Lemma 2.1, we have  $\int_{\mathbb{R}^n} |x|^{2-n+a} u^p dx < \infty$ . Hence, we define the function

$$w(x) = \frac{1}{n(n-2)\omega_n} \int_{\mathbb{R}^n} \frac{|y|^a u^p(y)}{|x-y|^{n-2}} dy.$$

It is clear that  $w(x) \geq 0$  and  $\Delta w = -|x|^a u^p$ . This implies that, for a solution  $u$  of (1-1), the function  $h(x) := w(x) + \Delta u(x)$  is a well-defined harmonic function on  $\mathbb{R}^n$ . Thus, for any  $x_0 \in \mathbb{R}^n$  and any  $R > 0$ , by the mean value theorem for harmonic functions we will have

$$h(x_0) := \int_{\partial B_R(x_0)} h d\sigma = \int_{\partial B_R(x_0)} (w + \Delta u) d\sigma \leq \int_{\partial B_R(x_0)} w d\sigma + \int_{\partial B_R(x_0)} |\Delta u| d\sigma. \tag{A-1}$$

Since  $w(x_0) < \infty$ , through Tonelli’s theorem, we can change the order of the integrations to see that the first integral on the right-hand side of (A-1) tends to zero as  $R \rightarrow \infty$  for all  $R$ . To be more precise, notice that, up to a constant multiple, the first integral can be written as

$$\int_{\mathbb{R}^n} \int_{\partial B_R(x_0)} \frac{d\sigma_x}{|x-y|^{n-2}} |y|^a u^p(y) dy.$$

Then we use the fact that  $\int_{\partial B_R(x_0)} 1/|x-y|^{n-2} d\sigma_x = |y-x_0|^{2-n}$  if  $|y-x_0| > R$  and equals  $R^{2-n}$  if  $|y-x_0| < R$ . Thus the integral will split into two parts. The outside part tends to zero as  $R \rightarrow \infty$  due to the fact that  $w(x_0) < \infty$ , while the inside part tends to zero due to the fact that, by Lemma 2.1,

$$R^{2-n} \int_{B_R(x_0)} |y|^a u^p(y) dy \leq R^{2-n} \int_{B_{R+|x_0|}(0)} |y|^a u^p dy \leq C R^{2-n} (R + |x_0|)^{n-(4p+a)/(p-1)}$$

tends to zero as  $R \rightarrow \infty$ . The second integral will tend to zero for some sequence of  $R$  by [Lemma 2.4](#) again. Apply the above inequality to this sequence to see that  $h(x_0) \leq 0$ . Since  $x_0$  is arbitrary, we have  $-\Delta u \geq 0$ .  $\square$

## References

- [Alikakos 2013] N. D. Alikakos, “On the structure of phase transition maps for three or more coexisting phases”, pp. 1–31 in *Geometric partial differential equations*, edited by A. Chambolle et al., CRM Series **15**, Ed. Norm., Pisa, 2013. [MR 3156886](#) [Zbl 1298.35054](#)
- [Caffarelli et al. 1994] L. Caffarelli, N. Garofalo, and F. Segàla, “A gradient bound for entire solutions of quasi-linear equations and its consequences”, *Comm. Pure Appl. Math.* **47**:11 (1994), 1457–1473. [MR 95k:35030](#) [Zbl 0819.35016](#)
- [Castellaneta et al. 2012] D. Castellaneta, A. Farina, and E. Valdinoci, “A pointwise gradient estimate for solutions of singular and degenerate PDE’s in possibly unbounded domains with nonnegative mean curvature”, *Commun. Pure Appl. Anal.* **11**:5 (2012), 1983–2003. [MR 2911121](#) [Zbl 1266.35097](#)
- [Cowan et al. 2010] C. Cowan, P. Esposito, and N. Ghoussoub, “Regularity of extremal solutions in fourth order nonlinear eigenvalue problems on general domains”, *Discrete Contin. Dyn. Syst.* **28**:3 (2010), 1033–1050. [MR 2011e:35085](#) [Zbl 1196.35152](#)
- [Farina 2004] A. Farina, “Two results on entire solutions of Ginzburg–Landau system in higher dimensions”, *J. Funct. Anal.* **214**:2 (2004), 386–395. [MR 2005g:35093](#) [Zbl 1183.35127](#)
- [Farina and Valdinoci 2010] A. Farina and E. Valdinoci, “A pointwise gradient estimate in possibly unbounded domains with nonnegative mean curvature”, *Adv. Math.* **225**:5 (2010), 2808–2827. [MR 2011j:35080](#) [Zbl 1200.35131](#)
- [Farina and Valdinoci 2011] A. Farina and E. Valdinoci, “A pointwise gradient bound for elliptic equations on compact manifolds with nonnegative Ricci curvature”, *Discrete Contin. Dyn. Syst.* **30**:4 (2011), 1139–1144. [MR 2012k:35138](#) [Zbl 1230.53037](#)
- [Farina and Valdinoci 2013] A. Farina and E. Valdinoci, “Pointwise estimates and rigidity results for entire solutions of nonlinear elliptic PDE’s”, *ESAIM Control Optim. Calc. Var.* **19**:2 (2013), 616–627. [MR 3049726](#) [Zbl 1273.35126](#)
- [Farina and Valdinoci 2014] A. Farina and E. Valdinoci, “Gradient bounds for anisotropic partial differential equations”, *Calc. Var. Partial Differential Equations* **49**:3–4 (2014), 923–936. [MR 3168616](#) [Zbl 1288.35131](#)
- [Farina et al. 2008] A. Farina, B. Sciunzi, and E. Valdinoci, “Bernstein and De Giorgi type problems: new results via a geometric approach”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **7**:4 (2008), 741–791. [MR 2009j:58020](#) [Zbl 1180.35251](#)
- [Fazly 2014] M. Fazly, “Liouville theorems for the polyharmonic Hénon–Lane–Emden system”, *Methods Appl. Anal.* **21**:2 (2014), 265–281. [MR 3251923](#) [Zbl 1302.35165](#)
- [Fazly and Ghoussoub 2013] M. Fazly and N. Ghoussoub, “De Giorgi type results for elliptic systems”, *Calc. Var. Partial Differential Equations* **47**:3–4 (2013), 809–823. [MR 3070565](#) [Zbl 1275.35094](#)
- [Fazly and Ghoussoub 2014] M. Fazly and N. Ghoussoub, “On the Hénon–Lane–Emden conjecture”, *Discrete Contin. Dyn. Syst.* **34**:6 (2014), 2513–2533. [MR 3177647](#) [Zbl 1285.35024](#)
- [Gidas and Spruck 1981a] B. Gidas and J. Spruck, “Global and local behavior of positive solutions of nonlinear elliptic equations”, *Comm. Pure Appl. Math.* **34**:4 (1981), 525–598. [MR 83f:35045](#) [Zbl 0465.35003](#)
- [Gidas and Spruck 1981b] B. Gidas and J. Spruck, “A priori bounds for positive solutions of nonlinear elliptic equations”, *Comm. Partial Differential Equations* **6**:8 (1981), 883–901. [MR 82h:35033](#) [Zbl 0462.35041](#)
- [Gui 2008] C. Gui, “Hamiltonian identities for elliptic partial differential equations”, *J. Funct. Anal.* **254**:4 (2008), 904–933. [MR 2009b:35112](#) [Zbl 1148.35023](#)
- [Lin 1998] C.-S. Lin, “A classification of solutions of a conformally invariant fourth order equation in  $\mathbb{R}^n$ ”, *Comment. Math. Helv.* **73**:2 (1998), 206–231. [MR 99c:35062](#) [Zbl 0933.35057](#)
- [Modica 1985] L. Modica, “A gradient bound and a Liouville theorem for nonlinear Poisson equations”, *Comm. Pure Appl. Math.* **38**:5 (1985), 679–684. [MR 87m:35088](#) [Zbl 0612.35051](#)
- [Moser 1961] J. Moser, “On Harnack’s theorem for elliptic differential equations”, *Comm. Pure Appl. Math.* **14** (1961), 577–591. [MR 28 #2356](#) [Zbl 0111.09302](#)

- [Ni 1982] W. M. Ni, “A nonlinear Dirichlet problem on the unit ball and its applications”, *Indiana Univ. Math. J.* **31**:6 (1982), 801–807. [MR 84b:35051](#) [Zbl 0515.35033](#)
- [Phan 2012] Q. H. Phan, “Liouville-type theorems and bounds of solutions for Hardy–Hénon elliptic systems”, *Adv. Differential Equations* **17**:7–8 (2012), 605–634. [MR 2963798](#) [Zbl 1262.35108](#)
- [Phan and Souplet 2012] Q. H. Phan and P. Souplet, “Liouville-type theorems and bounds of solutions of Hardy–Hénon equations”, *J. Differential Equations* **252**:3 (2012), 2544–2562. [MR 2860629](#) [Zbl 1233.35093](#)
- [Souplet 2009] P. Souplet, “The proof of the Lane–Emden conjecture in four space dimensions”, *Adv. Math.* **221**:5 (2009), 1409–1427. [MR 2010h:35088](#) [Zbl 1171.35035](#)
- [Wei and Xu 1999] J. Wei and X. Xu, “Classification of solutions of higher order conformally invariant equations”, *Math. Ann.* **313**:2 (1999), 207–228. [MR 2000a:58093](#) [Zbl 0940.35082](#)
- [Wei and Ye 2013] J. Wei and D. Ye, “Liouville theorems for stable solutions of biharmonic problem”, *Math. Ann.* **356**:4 (2013), 1599–1612. [MR 3072812](#) [Zbl 1277.35156](#)
- [Wei et al. 2013] J. Wei, X. Xu, and W. Yang, “On the classification of stable solutions to biharmonic problems in large dimensions”, *Pacific J. Math.* **263**:2 (2013), 495–512. [MR 3068555](#) [Zbl 1277.35155](#)

Received 9 Oct 2013. Revised 5 May 2015. Accepted 29 Jul 2015.

MOSTAFA FAZLY: [fazly@ualberta.ca](mailto:fazly@ualberta.ca)

*Department of Mathematical and Statistical Sciences, CAB 632 Central Academic Building, University of Alberta, Edmonton, AB T6G 2G1, Canada*

and

*Department of Mathematics and Computer Science, University of Lethbridge, Lethbridge, AB T1K 3M4, Canada*

JUN-CHENG WEI: [jcwei@math.ubc.ca](mailto:jcwei@math.ubc.ca)

*Department of Mathematics, The University of British Columbia, Vancouver, BC V6T 1Z2, Canada*

XINGWANG XU: [matxuxw@nus.edu.sg](mailto:matxuxw@nus.edu.sg)

*Department of Mathematics, National University of Singapore, Block S17 (SOC1), 10 Lower Kent Ridge Road, Singapore 119076, Singapore*



# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)  
University of California  
Berkeley, USA

## BOARD OF EDITORS

|                      |   |                       |  |
|----------------------|---|-----------------------|--|
| Nicolas Burq         | Université Paris-Sud 11, France<br><a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>             | Yuval Peres           | University of California, Berkeley, USA<br><a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>    |
| Sun-Yung Alice Chang | Princeton University, USA<br><a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>                         | Gilles Pisier         | Texas A&M University, and Paris 6<br><a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>                |
| Michael Christ       | University of California, Berkeley, USA<br><a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>         | Tristan Rivière       | ETH, Switzerland<br><a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>                                 |
| Charles Fefferman    | Princeton University, USA<br><a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>                               | Igor Rodnianski       | Princeton University, USA<br><a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>                  |
| Ursula Hamenstaedt   | Universität Bonn, Germany<br><a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>                           | Wilhelm Schlag        | University of Chicago, USA<br><a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>               |
| Vaughan Jones        | U.C. Berkeley & Vanderbilt University<br><a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a> | Sylvia Serfaty        | New York University, USA<br><a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>                         |
| Herbert Koch         | Universität Bonn, Germany<br><a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>                               | Yum-Tong Siu          | Harvard University, USA<br><a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>                          |
| Izabella Laba        | University of British Columbia, Canada<br><a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>                          | Terence Tao           | University of California, Los Angeles, USA<br><a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>             |
| Gilles Lebeau        | Université de Nice Sophia Antipolis, France<br><a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>                         | Michael E. Taylor     | Univ. of North Carolina, Chapel Hill, USA<br><a href="mailto:met@math.unc.edu">met@math.unc.edu</a>                |
| László Lempert       | Purdue University, USA<br><a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>                              | Gunther Uhlmann       | University of Washington, USA<br><a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>      |
| Richard B. Melrose   | Massachusetts Institute of Technology, USA<br><a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>                        | András Vasy           | Stanford University, USA<br><a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>                 |
| Frank Merle          | Université de Cergy-Pontoise, France<br><a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>                  | Dan Virgil Voiculescu | University of California, Berkeley, USA<br><a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>        |
| William Minicozzi II | Johns Hopkins University, USA<br><a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>                           | Steven Zelditch       | Northwestern University, USA<br><a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a> |
| Werner Müller        | Universität Bonn, Germany<br><a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>                         |                       |  |

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 8 No. 7 2015

---

|   |      |
|---|------|
| A pointwise inequality for the fourth-order Lane–Emden equation<br>MOSTAFA FAZLY, JUN-CHENG WEI and XINGWANG XU                             | 1541 |
| Convergence rates and Hölder estimates in almost-periodic homogenization of elliptic systems<br>ZHONGWEI SHEN                               | 1565 |
| Quantitative decay rates for dispersive solutions to the Einstein-scalar field system in spherical symmetry<br>JONATHAN LUK and SUNG-JIN OH | 1603 |
| A model for studying double exponential growth in the two-dimensional Euler equations<br>NETS KATZ and ANDREW TAPAY                         | 1675 |
| Asymptotics of Hadamard type for eigenvalues of the Neumann problem on $C^1$ -domains for elliptic operators<br>JOHAN THIM                  | 1695 |
| Scaling limit for the kernel of the spectral projector and remainder estimates in the pointwise Weyl law<br>YAIZA CANZANI and BORIS HANIN   | 1707 |
| On the continuous resonant equation for NLS, II: Statistical study<br>PIERRE GERMAIN, ZAHER HANI and LAURENT THOMANN                        | 1733 |
| Blow-up analysis of a nonlocal Liouville-type equation<br>FRANCESCA DA LIO, LUCA MARTINAZZI and TRISTAN RIVIÈRE                             | 1757 |