

# ANALYSIS & PDE

Volume 9

No. 2

2016

EMANUEL INDREI AND ANDREAS MINNE

**NONTRANSVERSAL INTERSECTION OF FREE AND FIXED  
BOUNDARIES FOR FULLY NONLINEAR ELLIPTIC OPERATORS  
IN TWO DIMENSIONS**



## NONTRANSVERSAL INTERSECTION OF FREE AND FIXED BOUNDARIES FOR FULLY NONLINEAR ELLIPTIC OPERATORS IN TWO DIMENSIONS

EMANUEL INDREI AND ANDREAS MINNE

In the study of classical obstacle problems, it is well known that in many configurations, the free boundary intersects the fixed boundary tangentially. The arguments involved in producing results of this type rely on the linear structure of the operator. In this paper, we employ a different approach and prove tangential touch of free and fixed boundaries in two dimensions for fully nonlinear elliptic operators. Along the way, several  $n$ -dimensional results of independent interest are obtained, such as BMO-estimates,  $C^{1,1}$ -regularity up to the fixed boundary, and a description of the behavior of blow-up solutions.

### 1. Introduction

Optimal interior regularity results have recently been obtained for solutions to fully nonlinear free boundary problems [Figalli and Shahgholian 2014; Indrei and Minne 2015] via methods inspired by [Andersson et al. 2013]. Under further thickness assumptions, these results imply  $C^1$ -regularity of the free boundary. However, a description of the dynamics on how the free boundaries intersect the fixed boundary has remained an open problem for at least a decade in the fully nonlinear setting (although partial results have been obtained in [Matevosyan and Markowich 2004] under strong density and growth assumptions involving the solutions and a homogeneity assumption on the operator). On the other hand, extensive work has been carried out to investigate this question for the classical problem

$$\begin{cases} \Delta u = \chi_{u>0} & \text{in } B_1^+, \\ u \geq 0 & \text{in } B_1^+, \\ u = 0 & \text{on } \{x_n = 0\} \end{cases}$$

and its variations [Apushkinskaya and Uraltseva 1995; Shahgholian and Uraltseva 2003; Matevosyan 2005; Andersson et al. 2006; Andersson 2007]. The conclusions have varied as a function of the boundary data, but in the homogeneous case, it has been shown that the free boundary touches the fixed boundary tangentially. Dynamics of this type have also been the object of study in the classical dam problem [Caffarelli and Gilardi 1980; Alt and Gilardi 1982], which is a mathematical model describing the filtration of water through a porous medium split into wet and dry parts via a free boundary.

The methods utilized in establishing the above-mentioned results strongly rely on the linear structure of the operator, e.g., in arguments involving Green's functions and monotonicity formulas. In particular, the Alt–Caffarelli–Friedman and Weiss monotonicity formulas are frequently applied — tools only available

---

*MSC2010:* primary 35JXX, 35QXX; secondary 49SXX.

*Keywords:* obstacle problem, tangential touch, fully nonlinear equations, nontransverse intersection, free boundary problem.

in the setting of linear operators in divergence form; see [Petrosyan et al. 2012, Chapter 8]. Thus the tangential touch problem for fully nonlinear operators requires a different approach.

In this article, we prove nontransversal intersection of free and fixed boundaries in two dimensions for the problem

$$\begin{cases} F(D^2u) = \chi_\Omega & \text{a.e. in } B_1^+, \\ u = 0 & \text{on } B_1', \end{cases}$$

where  $\Omega = (\{u \neq 0\} \cup \{\nabla u \neq 0\}) \cap \{x_2 > 0\} \subset \mathbb{R}_+^2$  and the free boundary is  $\mathbb{R}_+^2 \cap \partial\Omega$ . The starting point of our method is to first consider functions  $u \in W^{2,n}(B_1^+)$  satisfying

$$\begin{cases} F(D^2u) = 1 & \text{a.e. in } B_1^+ \cap \Omega, \\ |D^2u| \leq K & \text{a.e. in } B_1^+ \setminus \Omega, \\ u = 0 & \text{on } B_1', \end{cases} \tag{1}$$

where  $\Omega \subset B_1^+$  is an (unknown) open set,  $K > 0$ ,  $F$  is  $C^1$  and satisfies standard structural assumptions (see Section 3).

Since by assumption  $D^2u$  is bounded in the complement of  $\Omega$ , it follows that  $F(D^2u)$  is bounded in  $B_1^+$  and  $u$  is a strong  $L^n$ -solution to a fully nonlinear elliptic equation with bounded right-hand side [Caffarelli et al. 1996]. Under our structural assumptions on  $F$ , we have that  $L^n$ -solutions are also viscosity solutions, and it follows that  $u \in W_{loc}^{2,p}(B_1^+)$  for all  $p < \infty$  [Petrosyan et al. 2012]. If  $u \geq 0$  and  $\Omega = \{u \neq 0\}$ , then since  $D^2u = 0$  a.e. in the set  $\{u = 0\}$ , the Hessian condition in (1) is trivially satisfied; thus, (1) encodes the classical obstacle problem and likewise the equations  $F(D^2u) = \chi_{u \neq 0}$ ,  $F(D^2u) = \chi_{\nabla u \neq 0}$ , and  $F(D^2u) = \chi_{\{\nabla u \neq 0\} \cup \{u \neq 0\}}$  via the appropriate selection of  $\Omega$ .

A heuristic description of our strategy is as follows: We consider

$$M := \limsup_{|x| \rightarrow 0} \frac{1}{x_n} \sup_{e \in \mathbb{S}^{n-2} \cap e_n^\perp} \partial_e u(x).$$

By extending interior  $C^{1,1}$ -results (see Section 3), it follows that  $M$  is finite, and we extract information on the nature of blow-up solutions by considering possible values for  $M$ . In particular, if  $\{\nabla u \neq 0\} \cap \{x_n > 0\} \subset \Omega$  and the origin is a contact point, we show that either all blow-ups are of the form  $bx_n^2$  if  $M = 0$ , or there is a sequence producing a blow-up having the form  $ax_1x_n + bx_n^2$  if  $M \neq 0$  (Theorem 2.1).

We then show that in  $\mathbb{R}_+^2$ , if  $\Omega = (\{u \neq 0\} \cup \{\nabla u \neq 0\}) \cap \{x_2 > 0\}$  and  $ax_1x_n + bx_n^2$  is a blow-up solution, then  $\partial(\text{Int}\{u = 0\})$  stays away from the origin (Lemma 2.2) and this enables us to prove that blow-ups at the origin are unique (Theorem 2.4). Thereafter, a standard argument readily yields nontransversal intersection of the free and fixed boundaries at contact points (Theorem 2.5).

The rest of the paper is organized as follows: in the remainder of this section, we set up the problem and discuss relevant notation; Section 2 is the core of the paper where we rigorously develop the heuristics described above; Section 3 is devoted to the  $C^{1,1}$ -regularity up to the boundary of solutions, which follows as in [Indrei and Minne 2015] once a suitable BMO result is established. The results of Section 3 are used in Section 2. We have chosen to reverse the logical ordering of these sections in order to make the tangential touch section more accessible.

**Setup and notation.** We study fully nonlinear elliptic partial differential equations of the form

$$\begin{cases} F(D^2u, x) = f(x) & \text{a.e. in } B_1^+ \cap \Omega, \\ |D^2u| \leq K & \text{a.e. in } B_1^+ \setminus \Omega, \\ u = 0 & \text{on } B_1', \end{cases} \tag{2}$$

where  $u : B_1^+ \rightarrow \mathbb{R}$  is assumed to be in  $W^{2,n}(B_1^+)$ ,  $\Omega \subseteq B_1^+$  is an open set,  $B_1(x) = \{x \in \mathbb{R}^n : |x| < 1\}$ ,  $B_r^+(x) = B_r(x) \cap \{x_n > 0\}$ ,  $B_r'(x) = B_r(x) \cap \{x_n = 0\}$ , and  $B_r = B_r(0)$ .

Furthermore,  $F$  is assumed to satisfy the following structural conditions:

(H1)  $F(0, x) \equiv 0$ .

(H2)  $F$  is uniformly elliptic with ellipticity constants  $\lambda_0, \lambda_1 > 0$  such that

$$\mathcal{P}^-(M - N) \leq F(M, x) - F(N, x) \leq \mathcal{P}^+(M - N) \quad \forall x \in B_1^+,$$

where  $M$  and  $N$  are symmetric matrices and  $\mathcal{P}^\pm$  are the Pucci operators

$$\mathcal{P}^-(M) := \inf_{\lambda_0 \text{Id} \leq N \leq \lambda_1 \text{Id}} \text{Tr } NM \quad \text{and} \quad \mathcal{P}^+(M) := \sup_{\lambda_0 \text{Id} \leq N \leq \lambda_1 \text{Id}} \text{Tr } NM.$$

(H3)  $F(\cdot, x)$  is concave or convex for all  $x \in B_1^+$ .

(H4)  $|F(M, x) - F(M, y)| \leq \bar{C}(|M| + 1)|x - y|^{\bar{\alpha}}$

for some  $\bar{\alpha} \in (0, 1]$  and  $x, y \in B_1^+$ .

Moreover, let

$$\beta(x, x^0) := \sup_{M \in \mathcal{S}} \frac{|F(M, x) - F(M, x^0)|}{|M| + 1},$$

where  $\mathcal{S}$  is the space of  $n \times n$  symmetric real valued matrices.

Points in  $\mathbb{R}^n$  are generally denoted by  $x, x^0, y$  etc., while subscripts are used for components, i.e.,  $x = (x_1, \dots, x_n)$ , scalar sequences, and functions. The notation  $x'$  is used for  $(n-1)$ -dimensional vectors. Similarly,  $\nabla$  and  $\nabla'$  will be used, respectively, for the gradient and the gradient with respect to the first  $n - 1$  variables. We will also use the following:

- $\mathbb{R}_+^n$  is the upper half space  $\{x \in \mathbb{R}^n : x_n > 0\}$ ;
- $\Omega$  is an open set in  $\mathbb{R}_+^n$ ;
- $\Gamma$  is the set  $\mathbb{R}_+^n \cap \partial\Omega$ ;
- $\Gamma_i$  is the set  $\mathbb{R}_+^n \cap \partial \text{Int}\{u = 0\}$ ;
- $B_r(x^0)$  is the open ball  $\{x \in \mathbb{R}^n : |x - x^0| < r\}$ ;
- $B_r^+(x^0)$  is the truncated open ball  $\{x \in \mathbb{R}^n : |x - x^0| < r, x_n > 0\}$ ;
- $\partial B_r^+(x^0)$  is the topological boundary of  $B_r^+(x^0)$  in  $\mathbb{R}^n$ ;

- $B'_r$  is the ball  $\{x' \in \mathbb{R}^{n-1} : |x'| < r\}$ ;
- $\mathbb{S}^{n-1}$  is the  $(n-1)$ -sphere  $\{x \in \mathbb{R}^n : |x| = 1\}$ ;
- $e^\perp$  is the vector space orthogonal to  $e \in \mathbb{S}^{n-1}$ ;
- $C^{k,\alpha}(\Omega)$  denotes the usual Hölder space;
- $C^{k,\alpha}_{\text{loc}}(\Omega)$  denotes the local Hölder space;
- $W^{k,p}(\Omega)$  denotes the usual Sobolev space.

The term “blow-ups of  $u$  at  $x^0$ ” will be used for limits of the form

$$\lim_{j \rightarrow \infty} \frac{u(x^0 + r_j x)}{r_j^2},$$

where  $r_j$  is a sequence such that  $r_j \rightarrow 0^+$  as  $j \rightarrow \infty$ , and  $\text{Int}\{u = 0\} = \{u = 0\}^\circ$  means the interior of the set  $\{u = 0\} := \{x \in \mathbb{R}^n_+ : u(x) = 0\}$ . Finally,  $S(\psi)$  denotes the space of viscosity solutions corresponding to  $\psi$  and the ellipticity constants  $\lambda_0$  and  $\lambda_1$  in (H2); see [Caffarelli and Cabré 1995].

### 2. Main result

Our first result gives a natural dichotomy of blow-ups of solutions to (1) in any dimension.<sup>1</sup>

**Theorem 2.1** (blow-up alternative). *Let  $u$  be a solution to (1) and suppose  $\{\nabla u \neq 0\} \cap \{x_n > 0\} \subset \Omega$ ,  $0 \in \overline{\{u \neq 0\}}$ , and  $\nabla u(0) = 0$ . Then exactly one of the following holds:*

- (i) *All blow-ups of  $u$  at the origin are of the form  $u_0(x) = bx_n^2$  for some unique  $b > 0$ .*
- (ii) *There exists a blow-up of  $u$  at the origin of the form*

$$u_0(x) = ax_1x_n + bx_n^2$$

*for  $a \neq 0, b \in \mathbb{R}$ .*

*Proof.* Firstly, since  $u(x', 0) = 0$ , it follows that  $\partial_{x_i} u(x', 0) = 0$  for all  $i \in \{1, \dots, n - 1\}$ . By  $C^{1,1}$ -regularity (Theorem 3.1), there is a constant  $C > 0$  such that

$$\left| \frac{1}{x_n} \partial_{x_i} u(x', x_n) \right| = \left| \frac{1}{x_n} (\partial_{x_i} u(x', x_n) - \partial_{x_i} u(x', 0)) \right| \leq C, \quad x_n > 0.$$

Define

$$M := \limsup_{\substack{|x| \rightarrow 0 \\ x_n > 0}} \frac{1}{x_n} \sup_{e \in \mathbb{S}^{n-2} \cap e_n^\perp} \partial_e u(x).$$

In particular,  $0 \leq M \leq C < \infty$  and there exists a sequence  $x^j \rightarrow 0$  with  $x_n^j > 0$  and directions  $e_{x^j} \in \mathbb{S}^{n-2}$  such that

$$\lim_{j \rightarrow \infty} \frac{1}{x_n^j} \partial_{e_{x^j}} u(x^j) = M.$$

---

<sup>1</sup>Regularity results from Section 3 will be utilized in the proof of Theorem 2.1.

Moreover, there exists  $e \in \mathbb{S}^{n-2}$  such that (up to a subsequence)  $e_{x^j} \rightarrow e$ . Next note

$$\begin{aligned} \left| \frac{1}{x_n^j} \nabla' u(x^j) \cdot e - M \right| &\leq \left| \frac{1}{x_n^j} \nabla' u(x^j) \cdot (e - e_{x^j}) \right| + \left| \frac{1}{x_n^j} \nabla' u(x^j) \cdot e_{x^j} - M \right| \\ &\leq C |e - e_{x^j}| + \left| \frac{1}{x_n^j} \nabla' u(x^j) \cdot e_{x^j} - M \right| \rightarrow 0 \end{aligned}$$

as  $j \rightarrow \infty$ . Thus, up to a rotation,

$$\lim_{j \rightarrow \infty} \frac{1}{x_n^j} \partial_{x_1} u(x^j) = M.$$

Thanks to uniform boundedness, consider a sequence  $\{s_j\}$  such that  $s_j \rightarrow 0^+$  and the corresponding blow-up procedure so that

$$u_j(x) := \frac{u(s_j x)}{s_j^2} \rightarrow u_0(x)$$

in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}_+^n)$  for any  $\alpha \in [0, 1)$ , and  $u_0$  satisfies

$$\begin{cases} F(D^2 u_0) = 1 & \text{a.e. in } \mathbb{R}_+^n \cap \Omega_0, \\ |\nabla u_0| = 0 & \text{in } \mathbb{R}_+^n \setminus \Omega_0, \\ u = 0 & \text{on } \mathbb{R}_+^{n-1}, \end{cases} \tag{3}$$

where  $\Omega_0 = \{\nabla u_0 \neq 0\} \cap \{x_n > 0\}$  (via nondegeneracy). The definition of  $M$  implies

$$M \geq \lim_j \left| \frac{\partial_{x_i} u(s_j x)}{s_j x_n} \right| = \lim_j \left| \frac{\partial_{x_i} u_j(x)}{x_n} \right| = \left| \frac{\partial_{x_i} u_0(x)}{x_n} \right| \tag{4}$$

for all  $i \in \{1, \dots, n-1\}$ . In particular, let  $v = \partial_{x_1} u_0$  so that in  $\mathbb{R}_+^n$ ,

$$|v(x)| \leq M x_n. \tag{5}$$

If  $M = 0$ , then (4) implies  $\partial_{x_i} u_0 = 0$  for all  $i \in \{1, \dots, n-1\}$  so that  $u_0(x) = u_0(x_n)$ . However, since  $u_0(0) = |\nabla u_0(0)| = 0$ ,  $0 \in \{u_0 \neq 0\}$  and  $u_0$  satisfies (3), the uniform ellipticity of  $F$  readily implies

$$u_0(x) = b x_n^2$$

for some unique  $b > 0$ . This shows that if  $M = 0$ , then any blow-up at the origin is of the form in (i).

Now suppose  $M > 0$ . In order to prove (ii), we cook up a specific blow-up: let  $r_j := |x^j|$  (recall that  $\{x^j\}$  is the sequence achieving the lim sup in the definition of  $M$ ) so that as before

$$u_j(x) := \frac{u(r_j x)}{r_j^2} \rightarrow u_0(x)$$

in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}_+^n)$  for any  $\alpha \in [0, 1)$ , and  $u_0$  satisfies (3), (4), and (5). Set  $y^j = x^j / r_j \in \mathbb{S}^{n-1} \cap \{x_n > 0\}$ , and note that along a subsequence,  $y^j \rightarrow y \in \mathbb{S}^{n-1} \cap \{x_n \geq 0\}$ . Moreover, by the choice of the sequence  $\{x^j\}$

and the  $C^{1,\alpha}$ -convergence of  $u_j$  to  $u_0$ , if  $y_n > 0$ , then

$$\lim_j \frac{v(y^j)}{y_n^j} = \lim_j \frac{\partial_{x_1} u_j(y^j)}{y_n^j} = \lim_j \frac{\partial_{x_1} u(x^j)}{x_n^j} = M,$$

so that

$$v(y) = My_n, \tag{6}$$

and note that (6) also holds if  $y_n = 0$ . We consider several possibilities, keeping in mind that  $M > 0$ .

**Case 1:** If  $y \in \Omega_0$ , then by differentiating (3), we get the elliptic equation

$$a_{ij} \partial_{ij}(v(x) - Mx_n) = 0$$

for some measurable  $a_{ij}$ , and by (5), (6), and the maximum principle, it follows that  $v(x) = Mx_n$  in the connected component of  $\Omega_0$  containing  $y$ , say  $\Omega_0^y$ . If there exists  $x \in \partial\Omega_0^y \cap \{x_n > 0\}$ , then  $Mx_n = v(x) = 0$ , so we must have  $M = 0$ , a contradiction. Thus,  $v(x) = Mx_n$  in  $\mathbb{R}_+^n$  and by integrating,

$$u_0(x) = Mx_1x_n + h(x_2, \dots, x_n).$$

Now, the  $C^{2,\alpha}$ -estimate up to the boundary given by the Krylov–Safonov theorem (see, e.g., Theorem 3.3) applied to  $u_0(Rx)/R^2$  yields

$$\frac{|D^2u_0(x) - D^2u_0(y)|}{|x - y|^\alpha} \leq \frac{C}{R^\alpha}, \quad x \neq y \in B_R^+,$$

and taking  $R \rightarrow \infty$  implies that  $D^2u_0$  is a constant matrix, and thus  $h$  is a second-order polynomial. Since  $u_0$  vanishes on  $\{x_n = 0\}$ , it follows that

$$h(x_2, \dots, x_n) = x_n \sum_{i \neq n} \alpha_i x_i + bx_n^2,$$

and so up to a rotation,

$$u_0(x) = ax_1x_n + bx_n^2,$$

with  $a$  or  $b \neq 0$ .

**Case 2:** If  $y \in \partial\Omega_0 \cap \{x_n > 0\}$ , then  $My_n = v(y) = 0$ , a contradiction.

**Case 3:** If  $y \in \overline{\Omega}_0^c$ , then for all but finitely many  $j$ , we have  $y^j \in \Omega_0^c$  and since  $\{\nabla u_0 \neq 0\} \subset \Omega_0$ , it follows that  $v(y^j) = 0$  if  $j \geq N$  for some  $N \in \mathbb{N}$ . However,  $y_n^j > 0$  and so

$$0 = \lim_j \frac{v(y^j)}{y_n^j} = M,$$

a contradiction.

**Case 4:** If  $y \in \partial\Omega_0 \cap \{x_n = 0\}$ , by differentiating (3) in  $\Omega_0$ , it can be seen that for  $r > 0$  (to be picked later),  $v$  satisfies

$$Lv = 0 \text{ in } B_r(y)^+ \cap \Omega_0,$$

where  $L = F_{ij}(D^2u_0)\partial_{ij}$  is elliptic. Since  $u_0 \in C^{1,1}(B_r^+(y))$ , it follows that the  $F_{ij}(D^2u_0)$  are bounded and measurable on  $B_r^+(y)$ .

We know that  $Mx_n - v(x) \geq 0$  in  $\mathbb{R}_+^n$ , and if equality holds everywhere,  $u_0(x) = ax_1x_n + bx_n^2$  just as in Case 1. If there is a point  $z$  where strict inequality holds, that is,  $Mz_n - v(z) > 0$ , then we can choose a ball  $B_r^+(y)$  so that, by continuity of  $v$ , we have  $v(x) < Mx_n$  in a neighborhood  $B_s(z)$ , where  $z$  is a boundary point of  $B_r^+(y)$ . Note that this strict inequality necessarily occurs on  $\partial B_r^+(y) \cap \{x_n > 0\}$  since both  $v$  and  $Mx_n$  are zero on the hyperplane  $\{x_n = 0\}$ . Now choose a smooth nonnegative (but not identically zero) function  $\phi$  supported on  $B_s(z)$  small enough such that  $Mx_n - \phi(x) \geq v(x)$  and  $Mx_n - \phi(x) > 0$  in  $\mathbb{R}_+^n$  (this can be done since  $B_s(z)$  is some distance away from the hyperplane  $\{x_n = 0\}$ ). Then if

$$\begin{cases} Lw = 0 & \text{in } B_r^+(y), \\ w = Mx_n - \phi & \text{on } \partial B_r^+(y), \end{cases}$$

we have that  $w > 0$  in  $B_r^+(y)$  by the strong maximum principle since  $Mx_n - \phi(x) > 0$ . In particular,  $w > v = 0$  on  $\partial\Omega$ , and since  $v \leq w$  on  $\partial B_r^+(y)$ , the strong maximum principle again gives  $w > v$  in  $B_r^+(y) \cap \Omega$ . Note also by linearity that  $w = Mx_n - h$ , where  $h$  solves

$$\begin{cases} Lh = 0 & \text{in } B_r^+(y), \\ h = \phi & \text{on } \partial B_r^+(y). \end{cases}$$

Once more, the strong maximum principle shows that  $h > 0$  in  $B_r^+(y)$ , so the boundary Harnack comparison principle implies that  $cx_n \leq h(x)$  in  $B_{r/2}^+(y)$ , where  $c > 0$  depends on ellipticity constants and  $\phi$ . With this,

$$M = \lim_{j \rightarrow \infty} \frac{v(y^j)}{y_n^j} \leq \limsup_{\substack{x_n \rightarrow 0^+ \\ x \in B_{r/4}^+(y)}} \frac{w(x)}{x_n} \leq \lim_{\substack{x_n \rightarrow 0^+ \\ x \in B_{r/4}^+(y)}} \frac{Mx_n - cx_n}{x_n} = M - c,$$

a contradiction. □

The next lemma shows that in two dimensions, if (ii) in Theorem 2.1 occurs, then  $\Gamma_i = \mathbb{R}_+^2 \cap \partial \text{Int}\{u = 0\}$  stays away from the origin.

**Lemma 2.2.** *Let  $u$  be a solution to (1) with  $\Omega = (\{u \neq 0\} \cup \{\nabla u \neq 0\}) \cap \{x_2 > 0\} \subset \mathbb{R}_+^2$ . If there exists  $\{r_j\} \subset \mathbb{R}^+$  such that  $r_j \rightarrow 0$  as  $j \rightarrow \infty$  and*

$$u_j(x) := \frac{u(r_j x)}{r_j^2} \rightarrow u_0(x) = ax_1x_2 + bx_2^2$$

*in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}_+^2)$  as  $j \rightarrow \infty$  for  $a \neq 0, b \in \mathbb{R}$ , then there exists  $\delta \in (0, 1)$  such that  $B_\delta^+ \cap \Gamma_i = \emptyset$ .*

*Proof.* We may assume  $a > 0$ . Set  $v_j := \partial_1 u_j$  and let  $R > 2, \mu \in (0, \frac{1}{4})$ , and  $\delta \in (0, \frac{1}{4})$ . Then select  $j_0 = j_0(R, \mu, \delta) > 0$  such that for all  $j \geq j_0$ ,

$$|\nabla u_j(x)| > 0, \quad x \in B_R^+ \setminus B_\delta^+, \tag{7}$$

$$v_j(x) > 0, \quad x \in B_R^+ \cap \{x_2 \geq \mu\} \tag{8}$$

(the two-dimensional setting is crucial for (7)). Consider  $z \in \partial B_1 \cap \{x_2 = 0\}$  and note that

$$B_{3/4}^+(z) \subset B_R^+ \setminus B_\delta^+.$$

Thanks to (7),  $u_j$  satisfies  $F(D^2u_j) = 1$  in  $B_{3/4}^+(z)$  for all  $j \geq j_0$ .  $C^{2,\alpha}$ -estimates up to the boundary (see Theorem 3.3) imply

$$\sup_j \|u_j\|_{C^{2,\alpha}(B_{3/4}^+(z))} < \infty.$$

Thus, along a subsequence,  $v_j \rightarrow ax_2 =: v$  in  $C^{0,1}$  ( $C^{2,\alpha}$  is compactly contained in  $C^{1,1}$ ) and so

$$c_j := \sup_{\substack{x,y \in B_{3/4}^+(z) \\ x \neq y}} \frac{|(v_j(x) - v_j(y)) - (v(x) - v(y))|}{|x - y|} \rightarrow 0.$$

In particular, since  $v_j(x_1, 0) = v(x_1, 0) = 0$ , it follows that

$$\frac{|v_j(x) - ax_2|}{x_2} \leq c_j,$$

and so

$$v_j(x) \geq (a - c_j)x_2.$$

Now we select  $j$  large such that  $v_j(x) \geq 0$  on  $\partial B_1$ . Note that  $Lv_j = 0$  in  $B_1^+ \cap \Omega(u_j)$ , where  $L$  is an elliptic second-order operator obtained by differentiating (1). Indeed,  $u_j$  satisfies

$$\begin{cases} F(D^2u_j) = 1 & \text{a.e. in } B_{1/r_j}^+ \cap \Omega(u_j), \\ u_j = 0 & \text{on } B'_{1/r_j}, \end{cases}$$

where  $\Omega(u_j)$  is the dilated set  $\Omega/r_j$ , and without loss we may assume  $r_j < \frac{1}{2}$ .

Since  $v_j$  vanishes on  $\partial\Omega(u_j)$  and is nonnegative on  $\partial B_1^+$ , the maximum principle implies  $v_j > 0$  in  $B_1^+ \cap \Omega(u_j)$  (note that  $v_j$  is not identically zero by (8)). If  $\Gamma_i(u_j) \cap B_\delta^+ \neq \emptyset$ , consider a ball  $N$  in the interior of  $\{u_j = 0\} \cap B_\delta^+$ . For  $t \in \mathbb{R}$ , let  $N_t = N + te_1$ . Note that by taking  $t$  negative, we can move  $N_t$  to the left so that eventually  $N_t \subset B_1^+ \setminus B_\delta^+$ . Consider the strip  $S = \bigcup_{t \in \mathbb{R}} N_t$ . The next claim is that there exists a ball in the set  $(S \cap B_1^+) \setminus B_\delta^+$  such that  $u_j \neq 0$  in this ball: if not, then for each point  $z \in (S \cap B_1^+) \setminus B_\delta^+$ , there exists a sequence  $\{z_k\} \subset \{u_j = 0\}$  such that  $z_k \rightarrow z$ ; by continuity,  $u_j(z) = 0$ , so  $u_j = 0$  in  $(S \cap B_1^+) \setminus B_\delta^+$ , and therefore the gradient also vanishes there, a contradiction to (7). Denote the ball by  $E \subset \Omega(u_j)$  and note that  $u_j < 0$  on  $E$  since for each  $z \in E$ , there exists  $t_z > 0$  such that  $z + e_1 t_z \in \{u_j = 0\}$  and  $v_j > 0$  in  $B_1^+ \cap \Omega(u_j)$ . Thus,  $E \subset \Omega(u_j) \cap \{u_j < 0\}$ . Now move  $E$  to the right until the first time it touches  $\{u_j = 0\}$ , and let  $y$  be the contact point.

If  $\nabla u_j(y) = 0$ , we immediately obtain a contradiction via Hopf's lemma. Thus we may assume  $\nabla u_j(y) \neq 0$ , which implies  $y \in \Omega(u_j)$ , whence  $v_j(y) > 0$  (recall that  $v_j > 0$  in  $\Omega(u_j)$ ). By continuity,  $v_j > 0$  in  $B_r(y)$  for some  $r > 0$ , so in particular  $v_j(y + te_1) > 0$  for all  $t > 0$  small. Since  $\{y + te_1 : t \in (0, r)\} \subset \Omega(u_j)$ , we know  $t_* := \sup\{t > 0 : y + te_1 \in \Omega(u_j)\}$  is positive. Note that  $y + te_1$

will eventually enter  $N$  as  $t$  gets larger. However,

$$u_j(y + t_*e_1) - u_j(y) = \int_0^{t_*} v_j(y + se_1) ds > 0,$$

and this implies  $0 = u_j(y + t_*e_1) > u_j(y) = 0$ , a contradiction. Thus  $\Gamma_i(u_j) \cap B_\delta^+ = \emptyset$  and the result follows.  $\square$

Before proving uniqueness of blow-ups and tangential touch, we require one more lemma.

**Lemma 2.3.** *Let  $u$  be a solution to (1) with  $\Omega = (\{u \neq 0\} \cup \{\nabla u \neq 0\}) \cap \{x_n > 0\}$ . If  $s \in (0, 1]$  and  $(B_s^+ \setminus \Omega)^\circ = \emptyset$ , then  $|B_s^+ \setminus \Omega| = 0$ .*

*Proof.* Since  $u \in W^{2,n}(B_1^+)$ , it follows that  $D^2u = 0$  a.e. on  $B_s^+ \setminus \Omega$ . Let  $Z := \{D^2u = 0\} \cap (B_s^+ \setminus \Omega)$  and note that  $|Z| = |B_s^+ \setminus \Omega|$ . Thus if  $Z \subset (B_s^+ \setminus \Omega)^\circ$ , then the result follows. Let  $x^0 \in Z$  and suppose  $x^0 \notin (B_s^+ \setminus \Omega)^\circ$ . Then consider a sequence of points  $x^j \rightarrow x^0$  such that  $u(x^j) \neq 0$  and let  $r_j := |x^0 - x^j|$ . Nondegeneracy (see, e.g., [Indrei and Minne 2015, Lemma 3.1]) implies that for  $j$  large,

$$\sup_{\partial B_{r_j}(x^0)} \frac{u}{r_j^2} \geq c > 0,$$

or in other words,

$$\sup_{\partial B_1(0)} \frac{u(x^0 + r_j x)}{r_j^2} \geq c > 0.$$

Now for each  $j$  large enough, let  $y^j \in \partial B_1(0)$  be the element achieving the supremum in the previous expression; note that since

$$u(x^0) = |\nabla u(x^0)| = |D^2u(x^0)| = 0,$$

we have

$$u(x^0 + r_j y^j) = o(r_j^2),$$

a contradiction.  $\square$

Theorem 2.1, Lemma 2.2, and Lemma 2.3 imply uniqueness of blow-ups in two dimensions.

**Theorem 2.4** (uniqueness of blow-ups). *Let  $u$  be a solution to (1) with  $\Omega = (\{u \neq 0\} \cup \{\nabla u \neq 0\}) \cap \{x_2 > 0\} \subset \mathbb{R}_+^2$ . If  $0 \in \overline{\{u \neq 0\}}$  and  $\nabla u(0) = 0$ , then all blow-up limits  $u_0$  of  $u$  at the origin are of the form*

$$u_0(x) = ax_1x_2 + bx_2^2,$$

where  $a, b \in \mathbb{R}$  with at least one of them nonzero.

*Proof.* We divide the proof into two cases.

**Case 1:**  $0 \in \bar{\Gamma}_i$ . Lemma 2.2 implies the nonexistence of a blow-up  $u_0$  of  $u$  of the form

$$ax_1x_2 + bx_2^2,$$

$a \neq 0, b \in \mathbb{R}$ , from which it follows that (i) holds in Theorem 2.1. Note that  $b$  is uniquely determined by the equation.

**Case 2:**  $0 \notin \overline{\Gamma}_i$ . In this case, there exists  $\delta > 0$  such that  $\Gamma_i \cap B_\delta^+ = \emptyset$ . Since  $0 \in \overline{\{u \neq 0\}}$  (by assumption), it follows that  $B_\delta^+ \not\subset \{u = 0\}^\circ$  and as  $\Gamma_i \cap B_\delta^+ = \emptyset$ , we may conclude that  $\{u = 0\}^\circ \cap B_\delta^+ = \emptyset$ . Thus the hypotheses of Lemma 2.3 are satisfied, and by applying the lemma, we obtain that  $F(D^2u) = 1$  a.e. in  $B_\delta^+$ . Therefore  $u \in C^{2,\alpha}(B_{\delta/2}^+)$  and the blow-up limit  $u_0$  is uniquely given by

$$\begin{aligned} \lim_{r \rightarrow 0} \frac{u(rx)}{r^2} &= \lim_{r \rightarrow 0} \frac{u(0) + \nabla u(0) \cdot rx + \langle rx, D^2u(0)rx \rangle + o(r^2)}{r^2} \\ &= \langle x, D^2u(0)x \rangle = ax_1x_2 + bx_2^2. \end{aligned}$$

The last equality follows from the boundary condition. Furthermore,  $u_0$  solves the same equation as  $u$ , so

$$F(D^2u_0) = F(D^2u(0)) = 1$$

and thus  $a$  and  $b$  cannot both be zero due to (H1). □

If blow-ups are unique and of the form given above, it is rather standard to show that the free boundary touches the fixed boundary tangentially (see, e.g., [Petrosyan et al. 2012, Chapter 8]). The proof is included for completeness.

**Theorem 2.5** (tangential touch). *Let  $u$  be a solution to (1) with  $\Omega = (\{u \neq 0\} \cup \{\nabla u \neq 0\}) \cap \{x_2 > 0\} \subset \mathbb{R}_+^2$ . Then there exists a constant  $r_0 > 0$  and a modulus of continuity  $\omega_u(r)$  such that*

$$\Gamma(u) \cap B_{r_0}^+ \subset \{x : x_2 \leq \omega_u(|x|)|x|\}$$

if  $0 \in \overline{\Gamma(u)}$ , where  $\Gamma(u) := \partial\Omega \cap \mathbb{R}_+^2$ .

*Proof.* By Theorem 2.4, the blow-up of  $u$  at the origin is not identically zero and is given by  $u_0(x) = ax_1x_2 + bx_2^2$ . In particular,  $\Gamma(u_0) = \emptyset$ . It suffices to show that for any  $\epsilon > 0$ , there exists  $\rho_\epsilon = \rho_\epsilon(u) > 0$  such that

$$\Gamma(u) \cap B_{\rho_\epsilon}^+ \subset B_{\rho_\epsilon}^+ \setminus \mathcal{C}_\epsilon,$$

where  $\mathcal{C}_\epsilon := \{x_2 > \epsilon|x_1|\}$ . Suppose not. Then there exists a solution  $u$  to (1) satisfying the hypotheses of the theorem and  $\epsilon > 0$  such that for all  $k \in \mathbb{N}$ , there exists

$$x^k \in \Gamma(u) \cap B_{1/k}^+ \cap \mathcal{C}_\epsilon.$$

Let  $r_k := |x^k|$  and  $y^k := x^k/r_k \in \partial B_1 \cap \mathcal{C}_\epsilon$ . Note that along a subsequence

$$y^k \rightarrow y \in \partial B_1 \cap \mathcal{C}_\epsilon.$$

Define

$$u_k(x) := \frac{u(r_k x)}{r_k^2}$$

so that  $u_k \rightarrow u_0$  in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}_+^n)$  (along a subsequence). In particular,  $y \in \Gamma(u_0)$  by nondegeneracy, which contradicts that  $\Gamma(u_0) = \emptyset$ . □

### 3. $C^{1,1}$ -regularity up to the boundary

We now show BMO-estimates as well as  $C^{1,1}$ -regularity up to the fixed boundary of solutions to (2).

**Theorem 3.1** ( $C^{1,1}$ -regularity). *Let  $f \in C^\alpha(B_1^+)$  be a given function and  $\Omega \subseteq B_1^+$  a domain such that  $u : B_1^+ \rightarrow \mathbb{R}$  is a  $W^{2,n}$ -solution of (2). Assume  $F$  satisfies (H1)–(H4). Then there exists a constant  $C$  depending on  $\|u\|_{W^{2,n}(B_1^+)}$ ,  $\|f\|_{C^\alpha(B_1^+)}$ , and universal constants such that*

$$|D^2u| \leq C \quad \text{a.e. in } B_{1/2}^+.$$

There are three key tools needed to prove this theorem. The first two are  $C^{2,\alpha}$ - and  $W^{2,p}$ -estimates up to the boundary for the following classical fully nonlinear problem

$$\begin{cases} F(D^2u, x) = f(x) & \text{a.e. in } B_1^+, \\ u = 0 & \text{on } B_1', \end{cases} \quad (9)$$

and the last involves BMO-estimates. The  $C^{2,\alpha}$ - and  $W^{2,p}$ -estimates are well-known [Wang 1992; Safonov 1994; Winter 2009; Krylov 1982]. We have been unable to find a reference for the BMO-estimates and thus provide a proof, which is an adaptation of the interior case. For convenience, we record the following estimates; see, e.g., [Winter 2009, Theorem 4.3; Safonov 1994, Theorem 7.1]. Recall the definition of  $\beta$ ,

$$\beta(x, x^0) := \sup_{M \in \mathcal{S}} \frac{|F(M, x) - F(M, x^0)|}{|M| + 1}.$$

**Theorem 3.2** ( $W^{2,p}$ -regularity). *Let  $u$  be a  $W^{2,p}$ -viscosity solution to (9) and  $f \in L^p(B_1^+)$  for  $n \leq p \leq \infty$ . If  $\beta(x^0, y) \leq \beta_0$  in  $B_r^+(x^0) \cap B_1^+$  for all  $x^0 \in B_1^+$  and  $0 < r \leq r_0$ , where  $\beta_0$  and  $r_0$  are universal constants, then  $u \in W^{2,p}(B_{1/2}^+)$  and*

$$\|u\|_{W^{2,p}(B_{1/2}^+)} \leq C (\|u\|_{L^\infty(B_1^+)} + \|f\|_{L^p(B_1^+)}),$$

where  $C = C(n, \lambda_0, \lambda_1, \bar{\alpha}, \bar{C}, p) > 0$ .

**Theorem 3.3** ( $C^{2,\alpha}$ -regularity). *Let  $u$  be a  $W^{2,n}$ -viscosity solution to (9) and  $f \in C^{\bar{\alpha}}(B_1^+)$ . Then if  $\beta(x^0, y) \leq \beta_0$  in  $B_r^+(x^0) \cap B_1^+$  for all  $x^0 \in B_1^+$  and  $0 < r \leq r_0$ , where  $\beta_0$  and  $r_0$  are universal constants, then  $u \in C^{2,\alpha}(B_{1/2}^+)$  and*

$$\|u\|_{C^{2,\alpha}(B_{1/2}^+)} \leq C (\|u\|_{L^\infty(B_1^+)} + \|f\|_{C^{\bar{\alpha}}(B_1^+)}),$$

where  $C = C(n, \lambda_0, \lambda_1, \bar{\alpha}, \bar{C}) > 0$ .

The next results are technical tools utilized in the proof of the BMO-estimate (i.e., Proposition 3.6). The first is an approximation lemma; see, e.g., [Wang 1992, Lemma 1.4].

**Lemma 3.4** (approximation). *Let  $\epsilon > 0$ ,  $u \in W^{2,p}(B_1^+(x^0))$ , and let  $v$  solve*

$$\begin{cases} F(D^2v, x^0) = a & \text{in } B_{1/2}^+(x^0), \\ v = u & \text{on } \partial B_{1/2}^+(x^0). \end{cases}$$

Then there exists  $\delta > 0$  and  $\eta > 0$  such that if

$$\beta(x, x^0) := \sup_{M \in \mathcal{S}} \frac{|F(M, x) - F(M, x^0)|}{|M| + 1} \leq \eta$$

and  $|f(x) - a| \leq \delta$  a.e. for  $f(x) := F(D^2u(x), x)$  in  $B_1^+(x^0)$ , then

$$|u - v| \leq \epsilon \quad \text{in } B_{1/2}^+.$$

**Lemma 3.5.** *Let  $u \in W^{2,n}(B_1^+)$  satisfy  $|F(D^2u(x), x)| \leq \delta$  a.e. in  $B_1^+$  for  $\delta$  as in Lemma 3.4. Moreover, assume  $|u| \leq 1$  and that  $\beta(x, y)$  satisfies (H4). Then there exists a universal constant  $\rho > 0$  and second-order polynomials  $P_{k,x^0}$  for any  $k \in \mathbb{N}_0$  and  $x^0 \in B_{1/2}^+$  such that*

$$\begin{aligned} |D^2P_{k,x^0} - D^2P_{k-1,x^0}| &\leq C_0(n, \lambda_0, \lambda_1), \\ F(D^2P_{k,x^0}, x^0) &= 0, \\ |u(x) - P_{k,x^0}(x)| &\leq \rho^{2k} \quad \text{inside } B_{\min(\rho^k, 1)}^+(x^0). \end{aligned}$$

*Proof.* For  $k = 0$  and  $k = -1$ , the statement is true for  $P_{k,x^0}(x) \equiv 0$  by assumption (recall (H1)). If we assume it is true up to some  $k$ , define

$$\begin{aligned} u_k &:= \frac{u(\rho^k x + x^0) - P_{k,x^0}(\rho^k x + x^0)}{\rho^{2k}}, \\ F_k(M, x) &:= F(M + D^2P_{k,x^0}, \rho^k x + x^0), \quad x \in B_1 \cap \{x_n > -x_n^0/\rho^k\}. \end{aligned}$$

Then  $|F_k(D^2u_k, x)| = |F((D^2u)(\rho^k x + x^0), \rho^k x + x^0)| \leq \delta$  a.e. Also,

$$\begin{aligned} \beta_k(x, 0) &= \sup_{M \in \mathcal{S}} \frac{|F_k(M, x) - F_k(M, 0)|}{|M| + 1} \\ &= \sup_{M \in \mathcal{S}} \frac{|F(M + D^2P_{k,x^0}, \rho^k x + x^0) - F(M + D^2P_{k,x^0}, x^0)|}{|M| + 1} \\ &= \sup_{M \in \mathcal{S}} \frac{|F(M, \rho^k x + x^0) - F(M, x^0)|}{|M - D^2P_{k,x^0}| + 1} \\ &= \sup_{M \in \mathcal{S}} \frac{|F(M, \rho^k x + x^0) - F(M, x^0)|}{|M| + 1} \frac{|M| + 1}{|M - D^2P_{k,x^0}| + 1} \\ &\leq \beta(\rho^k x + x^0, x^0) \sup_{M \in \mathcal{S}} \frac{|M| + 1}{|M - D^2P_{k,x^0}| + 1} \\ &\leq \bar{C} \rho^{\bar{\alpha}k} \sup_{M \in \mathcal{S}} \frac{|M| + 1}{||M| - |D^2P_{k,x^0}|| + 1} \\ &\leq \bar{C} \rho^{\bar{\alpha}k} (|D^2P_{k,x^0}| + 1), \end{aligned}$$

where the last inequality follows from a calculation of the maximum of the function

$$\frac{x + 1}{|x - a| + 1}, \quad x, a \geq 0.$$

However, from the induction hypothesis,

$$|D^2 P_{k,x^0}| \leq \sum_{j=1}^k |D^2 P_{j-1,x^0} - D^2 P_{j,x^0}| \leq C_0 k,$$

so

$$\bar{C} \rho^{\bar{\alpha}k} (|D^2 P_{k,x^0}| + 1) \leq \bar{C} \rho^{\bar{\alpha}k} C_0 k \leq \eta$$

if  $\rho$  is chosen small enough (depending only on universal constants) and  $\eta$  as in Lemma 3.4. Thus  $|v_k - u_k| \leq \epsilon$  in  $B_{1/2} \cap \{x : x_n > -x_n^0/\rho^k\}$  by Lemma 3.4, where  $v_k$  solves

$$\begin{cases} F_k(D^2 v_k, 0) = 0 & \text{in } B_{1/2} \cap \{x : x_n > -x_n^0/\rho^k\}, \\ v_k = u_k & \text{on } \partial(B_{1/2} \cap \{x : x_n > -x_n^0/\rho^k\}). \end{cases}$$

Since

$$\|v_k\|_{L^\infty(B_{1/2} \cap \{x : x_n > -x_n^0/\rho^k\})} \leq \|u_k\|_{L^\infty(B_{1/2} \cap \{x : x_n > -x_n^0/\rho^k\})} \leq 1$$

by the maximum principle, Theorem 3.3 gives

$$\|v_k\|_{C^{2,\alpha}(B_{1/4} \cap \{x : x_n > -x_n^0/\rho^k\})} \leq C_0. \tag{10}$$

Now define  $\hat{P}_{k,x^0}$  as the second-order Taylor expansion of  $v_k$  at the origin, and note that  $F_k(D^2 \hat{P}_{k,x^0}, 0) = F_k(D^2 v_k(0), 0) = 0$ . Then

$$|v_k - \hat{P}_{k,x^0}| \leq C_0 \rho^{2+\alpha} \quad \text{in } B_\rho \cap \{x : x_n > -x_n^0/\rho^k\}$$

for  $\rho < \frac{1}{4}$ , which gives

$$|u_k - \hat{P}_{k,x^0}| \leq |u_k - v_k| + |v_k - \hat{P}_{k,x^0}| \leq \epsilon + C_0 \rho^{2+\alpha} \quad \text{in } B_\rho \cap \{x : x_n > -x_n^0/\rho^k\}.$$

For  $\rho^\alpha \leq \frac{1}{2C_0}$  and  $\epsilon \leq \frac{1}{2}\rho^2$ , we get

$$|u_k - \hat{P}_{k,x^0}| \leq \rho^2 \quad \text{in } B_\rho \cap \{x : x_n > -x_n^0/\rho^k\},$$

or, in other words,

$$|u - P_{k+1,x^0}| \leq \rho^{2(k+1)} \quad \text{in } B_{\rho^{k+1}}^+(x^0)$$

for

$$P_{k+1,x^0}(x) := P_{k,x^0}(x) + \rho^{2k} \hat{P}_{k,x^0}\left(\frac{x - x^0}{\rho^k}\right).$$

Also, since  $F_k(D^2 \hat{P}_{k,x^0}, 0) = 0$ , by (10) we have

$$\begin{aligned} F(D^2 P_{k+1,x^0}, x^0) &= F(D^2 P_{k,x^0} + D^2 \hat{P}_{k,x^0}, x^0) = F_k(D^2 \hat{P}_{k,x^0}, 0) = 0, \\ |D^2 P_{k+1,x^0} - D^2 P_{k,x^0}| &= |D^2 \hat{P}_{k,x^0}| = |D^2 v_k(0)| \leq C_0. \end{aligned} \quad \square$$

**Proposition 3.6** (BMO-estimate). *Let  $u$  be a solution to (9),  $f$  bounded, and  $P_{k,x^0}$  and  $\rho$  be as in Lemma 3.5. Then*

$$\int_{B_{\rho^k/2}^+(x^0)} |D^2 u(y) - D^2 P_{k,x^0}|^2 \leq C, \quad x^0 \in \bar{B}_{1/2}^+,$$

if  $\rho$  is smaller than a constant that depends only on  $\|u\|_{W^{2,p}(B_1)}$ ,  $f$ ,  $\bar{C}$  in (H4), and universal constants.

*Proof.* Let  $x^0 \in \bar{B}_{1/2}^+$  and define

$$v(x) := u\left(\frac{x}{R}\right) \quad \text{and} \quad G(M, x) := \frac{1}{R^2} F\left(R^2 M, \frac{x}{R}\right)$$

for  $R = R(\bar{C}, f, K, \delta)$  ( $\bar{C}$  as in (H4)) chosen so that  $|G(D^2 v, x)| \leq \delta$  in  $B_R^+$  for  $\delta$  as in Lemma 3.4. Note also that

$$\beta_G(x, y) := \sup_{M \in \mathcal{S}} \frac{|G(M, x) - G(M, y)|}{|M| + 1}$$

satisfies (H4). Then  $v$  solves

$$\begin{cases} G(D^2 v, x) = f(x/R)/R^2 & \text{a.e. in } B_R^+ \cap (R\Omega), \\ |D^2 v| \leq K/R^2 & \text{a.e. in } B_R^+ \setminus (R\Omega), \\ v = 0 & \text{on } B'_R, \end{cases}$$

and there is a polynomial  $\tilde{P}_{k,x^0}$  for which  $G(D^2 \tilde{P}_{k,x^0}, Rx^0) = 0$ , and a constant  $\tilde{\rho}$  such that

$$|v(x) - \tilde{P}_{k,x^0}(x)| \leq \tilde{\rho}^{2k}, \quad x \in B_{\tilde{\rho}^k}^+(Rx^0),$$

that is,

$$|u(x) - P_{k,x^0}(x)| \leq R^2 \tilde{\rho}^{2k}, \quad x \in B_{\rho^k}^+(x^0),$$

for  $P_{k,x^0}(x) := \tilde{P}_{k,x^0}(Rx)$  and  $\rho^k := \tilde{\rho}^k/R$ . Note also that

$$F(D^2 P_{k,x^0}, x^0) = F\left(R^2 D^2 \tilde{P}_{k,x^0}, \frac{Rx^0}{R}\right) = R^2 G(D^2 \tilde{P}_{k,x^0}, Rx^0) = 0.$$

In particular, for

$$u_k(x) := \frac{u(\rho^k x + x^0) - P_{k,x^0}(\rho^k x + x^0)}{\rho^{2k}},$$

$$F_k(M, x) := F(M + D^2 P_{k,x^0}, \rho^k x + x^0),$$

and  $\beta_k$  as in the proof of Lemma 3.5, we have  $|u_k| \leq R^2$ ,  $\beta_k(x, y) \leq \eta$  and  $|F_k(u_k, x)| \leq C$ . Therefore we can apply Theorem 3.2 to deduce

$$\|u_k\|_{W^{2,p}(B_{1/2} \cap \{x_n \geq -x^0/\rho^k\})} \leq C,$$

or

$$\int_{B_{\rho^k/2}^+(x^0)} |D^2 u(x) - D^2 P_{k,x^0}|^p dx \leq C. \quad \square$$

From this it is straightforward to show that if  $u$  is a function satisfying (2), there exists a second-order polynomial  $P_{r,x^0}(x)$  with  $F(D^2 P_{r,x^0}, x^0) = f(x^0)$  such that

$$\sup_{r \in (0, 1/4)} \int_{B_r^+(x^0)} |D^2 u(y) - D^2 P_{r,x^0}|^2 dy \leq C,$$

where  $x^0 \in \bar{B}_{1/2}^+(0)$ . The proof of  $C^{1,1}$ -regularity now follows as in [Indrei and Minne 2015] up to minor modifications (see also [Figalli and Shahgholian 2014]). The idea is that  $D^2 P_{r,x^0}(x)$  provides a suitable approximation to  $D^2 u(x^0)$  and one may consider two cases: first, if  $D^2 P_{r,x^0}(x)$  stays bounded in  $r$ , then one can show that  $D^2 u(x^0)$  is also bounded by a constant depending only on the initial ingredients; next, if  $D^2 P_{r,x^0}(x)$  blows up in  $r$ , one can show that the set

$$A_r(x^0) := \frac{(B_r^+(x^0) \setminus \Omega) - x^0}{r} = B_1 \setminus ((\Omega - x^0)/r) \cap \{y : y_n > -x_n^0/r\}$$

decays fast enough to ensure yet again a bound on  $D^2 u(x^0)$ .

### Acknowledgements

We thank Henrik Shahgholian for bringing this problem to our attention. Moreover, we thank John Andersson for his help: the technique developed in this paper evolved through our interaction with him. We also thank the G. S. Magnuson Foundation for supporting this work. E. Indrei acknowledges partial support from NSF Grants OISE-0967140 (PIRE), DMS-0405343, and DMS-0635983 administered by the Center for Nonlinear Analysis at Carnegie Mellon University. Moreover, the excellent research environment provided by KTH, CMU, the Hausdorff Research Institute for Mathematics, and Universität Bonn is kindly acknowledged. Lastly, we thank Herbert Koch and an anonymous referee for providing valuable feedback on a preliminary version of this paper.

### References

- [Alt and Gilardi 1982] H. W. Alt and G. Gilardi, “The behavior of the free boundary for the dam problem”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **9**:4 (1982), 571–626. MR 693780 Zbl 0521.76092
- [Andersson 2007] J. Andersson, “On the regularity of a free boundary near contact points with a fixed boundary”, *J. Differential Equations* **232**:1 (2007), 285–302. MR 2281197 Zbl 1106.35143
- [Andersson et al. 2006] J. Andersson, N. Matevosyan, and H. Mikayelyan, “On the tangential touch between the free and the fixed boundaries for the two-phase obstacle-like problem”, *Ark. Mat.* **44**:1 (2006), 1–15. MR 2237208 Zbl 1170.35560
- [Andersson et al. 2013] J. Andersson, E. Lindgren, and H. Shahgholian, “Optimal regularity for the no-sign obstacle problem”, *Comm. Pure Appl. Math.* **66**:2 (2013), 245–262. MR 2999297 Zbl 06132672
- [Apushkinskaya and Uraltseva 1995] D. E. Apushkinskaya and N. N. Uraltseva, “On the behavior of the free boundary near the boundary of the domain”, *Zap. Nauchn. Sem. POMI* **221** (1995), 5–19. In Russian; translated in *J. Math. Sci. New York* **87**:2 (1997), 3267–3276. MR 1359745 Zbl 0888.35141
- [Caffarelli and Cabré 1995] L. A. Caffarelli and X. Cabré, *Fully nonlinear elliptic equations*, American Mathematical Society Colloquium Publications **43**, American Mathematical Society, Providence, RI, 1995. MR 1351007 Zbl 0834.35002
- [Caffarelli and Gilardi 1980] L. A. Caffarelli and G. Gilardi, “Monotonicity of the free boundary in the two-dimensional dam problem”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **7**:3 (1980), 523–537. MR 597551 Zbl 0513.76090

- [Caffarelli et al. 1996] L. Caffarelli, M. G. Crandall, M. Kocan, and A. Święch, “On viscosity solutions of fully nonlinear equations with measurable ingredients”, *Comm. Pure Appl. Math.* **49**:4 (1996), 365–397. MR 1376656 Zbl 0854.35032
- [Figalli and Shahgholian 2014] A. Figalli and H. Shahgholian, “A general class of free boundary problems for fully nonlinear elliptic equations”, *Arch. Ration. Mech. Anal.* **213**:1 (2014), 269–286. MR 3198649 Zbl 1326.35137
- [Indrei and Minne 2015] E. Indrei and A. Minne, “Regularity of solutions to fully nonlinear elliptic and parabolic free boundary problems”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* (online publication May 11 2015).
- [Krylov 1982] N. V. Krylov, “Boundedly inhomogeneous elliptic and parabolic equations”, *Izv. Akad. Nauk SSSR Ser. Mat.* **46**:3 (1982), 487–523, 670. In Russian; translated in *Math. USSR Izv.* **20**:3 (1983), 459–492. MR 661144 Zbl 0529.35026
- [Matevosyan 2005] N. Matevosyan, “Tangential touch between free and fixed boundaries in a problem from superconductivity”, *Comm. Partial Differential Equations* **30**:7-9 (2005), 1205–1216. MR 2180300 Zbl 1081.35161
- [Matevosyan and Markowich 2004] N. Matevosyan and P. A. Markowich, “Behavior of the free boundary near contact points with the fixed boundary for nonlinear elliptic equations”, *Monatsh. Math.* **142**:1-2 (2004), 17–25. MR 2065018 Zbl 1064.35215
- [Petrosyan et al. 2012] A. Petrosyan, H. Shahgholian, and N. Uraltseva, *Regularity of free boundaries in obstacle-type problems*, Graduate Studies in Mathematics **136**, American Mathematical Society, Providence, RI, 2012. MR 2962060 Zbl 1254.35001
- [Safonov 1994] M. Safonov, “On the boundary value problems for fully nonlinear elliptic equations of second order”, Mathematics Research Report No. MRR 049-94, The Australian National University, Canberra, 1994, available at [http://www.math.umn.edu/~safon002/NOTES/BVP\\_94/BVP.pdf](http://www.math.umn.edu/~safon002/NOTES/BVP_94/BVP.pdf).
- [Shahgholian and Uraltseva 2003] H. Shahgholian and N. Uraltseva, “Regularity properties of a free boundary near contact points with the fixed boundary”, *Duke Math. J.* **116**:1 (2003), 1–34. MR 1950478 Zbl 1050.35157
- [Wang 1992] L. Wang, “On the regularity theory of fully nonlinear parabolic equations, II”, *Comm. Pure Appl. Math.* **45**:2 (1992), 141–178. MR 1139064 Zbl 0774.35042
- [Winter 2009] N. Winter, “ $W^{2,p}$  and  $W^{1,p}$ -estimates at the boundary for solutions of fully nonlinear, uniformly elliptic equations”, *Z. Anal. Anwend.* **28**:2 (2009), 129–164. MR 2486925 Zbl 1206.35116

Received 12 Jun 2015. Revised 6 Jan 2016. Accepted 9 Feb 2016.

EMANUEL INDREI: [egi@cmu.edu](mailto:egi@cmu.edu)

Center for Nonlinear Analysis, Carnegie Mellon University, Pittsburgh, PA 15213, United States

ANDREAS MINNE: [minne@kth.se](mailto:minne@kth.se)

Department of Mathematics, KTH Royal Institute of Technology, 100 44 Stockholm, Sweden

# Analysis & PDE

msp.org/apde

## EDITORS

### EDITOR-IN-CHIEF

Patrick Gérard  
patrick.gerard@math.u-psud.fr  
Université Paris Sud XI  
Orsay, France

### BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 9 No. 2 2016

---

Resonances for large one-dimensional “ergodic” systems FRÉDÉRIC KLOPP	259
On characterization of the sharp Strichartz inequality for the Schrödinger equation JIN-CHENG JIANG and SHUANGLIN SHAO	353
Future asymptotics and geodesic completeness of polarized $T^2$ -symmetric spacetimes PHILIPPE G. LEFLOCH and JACQUES SMULEVICI	363
Obstacle problem with a degenerate force term KAREN YERESSIAN	397
A counterexample to the Hopf–Oleinik lemma (elliptic case) DARYA E. APUSHKINSKAYA and ALEXANDER I. NAZAROV	439
Ground states of large bosonic systems: the Gross–Pitaevskii limit revisited PHAN THÀNH NAM, NICOLAS ROUGERIE and ROBERT SEIRINGER	459
Nontransversal intersection of free and fixed boundaries for fully nonlinear elliptic operators in two dimensions EMANUEL INDREI and ANDREAS MINNE	487
Correction to the article Scattering threshold for the focusing nonlinear Klein–Gordon equation SLIM IBRAHIM, NADER MASMOUDI and KENJI NAKANISHI	503



2157-5045(2016)9:2;1-A