

ANALYSIS & PDE

Volume 9

No. 5

2016

GELU POPESCU

**FREE PLURIHARMONIC FUNCTIONS ON NONCOMMUTATIVE
POLYBALLS**



FREE PLURIHARMONIC FUNCTIONS ON NONCOMMUTATIVE POLYBALLS

GELU POPESCU

We study free k -pluriharmonic functions on the noncommutative regular polyball \mathbf{B}_n , $\mathbf{n} = (n_1, \dots, n_k) \in \mathbb{N}^k$, which is an analogue of the scalar polyball $(\mathbb{C}^{n_1})_1 \times \dots \times (\mathbb{C}^{n_k})_1$. The regular polyball has a universal model $\mathcal{S} := \{\mathcal{S}_{i,j}\}$ consisting of left creation operators acting on the tensor product $F^2(H_{n_1}) \otimes \dots \otimes F^2(H_{n_k})$ of full Fock spaces. We introduce the class \mathcal{T}_n of k -multi-Toeplitz operators on this tensor product and prove that $\mathcal{T}_n = \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{\text{SOT}}$, where \mathcal{A}_n is the noncommutative polyball algebra generated by \mathcal{S} and the identity. We show that the bounded free k -pluriharmonic functions on \mathbf{B}_n are precisely the noncommutative Berezin transforms of k -multi-Toeplitz operators. The Dirichlet extension problem on regular polyballs is also solved. It is proved that a free k -pluriharmonic function has continuous extension to the closed polyball \mathbf{B}_n^- if and only if it is the noncommutative Berezin transform of a k -multi-Toeplitz operator in $\text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{\|\cdot\|}$.

We provide a Naimark-type dilation theorem for direct products $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$ of unital free semigroups, and use it to obtain a structure theorem which characterizes the positive free k -pluriharmonic functions on the regular polyball with operator-valued coefficients. We define the noncommutative Berezin (resp. Poisson) transform of a completely bounded linear map on $C^*(\mathcal{S})$, the C^* -algebra generated by $\mathcal{S}_{i,j}$, and give necessary and sufficient conditions for a function to be the Poisson transform of a completely bounded (resp. completely positive) map. In the last section of the paper, we obtain Herglotz–Riesz representation theorems for free holomorphic functions on regular polyballs with positive real parts, extending the classical result as well as the Korányi–Pukánszky version in scalar polydisks.

Introduction	1185
1. k -multi-Toeplitz operators on tensor products of full Fock spaces	1190
2. Bounded free k -pluriharmonic functions and the Dirichlet extension problem	1204
3. Naimark-type dilation theorem for direct products of free semigroups	1209
4. Berezin transforms of completely bounded maps in regular polyballs	1216
5. Herglotz–Riesz representations for free holomorphic functions with positive real parts	1227
References	1232

Introduction

A multivariable operator model theory and a theory of free holomorphic functions on polydomains which admit universal operator models have been recently developed in [Popescu 2013; 2016]. An important feature of these theories is that they are related, via noncommutative Berezin transforms, to the study of

Research supported in part by NSF grant DMS 1500922.

MSC2010: primary 47A13, 47A56; secondary 47B35, 46L52.

Keywords: noncommutative polyball, Berezin transform, Poisson transform, Fock space, multi-Toeplitz operator, Naimark dilation, completely bounded map, pluriharmonic function, free holomorphic function, Herglotz–Riesz representation.

the operator algebras generated by the universal models as well as to the theory of functions in several complex variables. These results played a crucial role in our work on the curvature invariant [Popescu 2015a], the Euler characteristic [Popescu 2014], and the group of free holomorphic automorphisms on noncommutative regular polyballs [Popescu 2015b].

The main goal of the present paper is to continue our investigation along these lines and to study the class of free k -pluriharmonic functions of the form

$$F(X) = \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+ \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} a(\alpha; \beta) X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*, \quad a(\alpha; \beta) \in \mathbb{C},$$

where the series converge in the operator norm topology for any $X = \{X_{i,j}\}$ in the regular polyball $\mathbf{B}_n(\mathcal{H})$ and any Hilbert space \mathcal{H} . The results of this paper will play an important role in the hyperbolic geometry of noncommutative polyballs [Popescu ≥ 2016]. To present our results we need some notation and preliminaries on regular polyballs and their universal models.

Throughout this paper, $B(\mathcal{H})$ stands for the algebra of all bounded linear operators on a Hilbert space \mathcal{H} . We let $B(\mathcal{H})^{n_1} \times_c \cdots \times_c B(\mathcal{H})^{n_k}$, where $n_i \in \mathbb{N} := \{1, 2, \dots\}$, be the set of all tuples $X := (X_1, \dots, X_k)$ in $B(\mathcal{H})^{n_1} \times \cdots \times B(\mathcal{H})^{n_k}$ with the property that the entries of $X_s := (X_{s,1}, \dots, X_{s,n_s})$ commute with the entries of $X_t := (X_{t,1}, \dots, X_{t,n_t})$ for any $s, t \in \{1, \dots, k\}, s \neq t$. Note that the operators $X_{s,1}, \dots, X_{s,n_s}$ do not necessarily commute. Let $\mathbf{n} := (n_1, \dots, n_k)$ and define the polyball

$$\mathbf{P}_n(\mathcal{H}) := [B(\mathcal{H})^{n_1}]_1 \times_c \cdots \times_c [B(\mathcal{H})^{n_k}]_1,$$

where

$$[B(\mathcal{H})^n]_1 := \{(X_1, \dots, X_n) \in B(\mathcal{H})^n : \|X_1 X_1^* + \cdots + X_n X_n^*\| < 1\}, \quad n \in \mathbb{N}.$$

If A is a positive invertible operator, we write $A > 0$. The *regular polyball* on the Hilbert space \mathcal{H} is defined by

$$\mathbf{B}_n(\mathcal{H}) := \{X \in \mathbf{P}_n(\mathcal{H}) : \Delta_X(I) > 0\},$$

where the *defect mapping* $\Delta_X : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ is given by

$$\Delta_X := (\text{id} - \Phi_{X_1}) \circ \cdots \circ (\text{id} - \Phi_{X_k})$$

and $\Phi_{X_i} : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ is the completely positive linear map defined by

$$\Phi_{X_i}(Y) := \sum_{j=1}^{n_i} X_{i,j} Y X_{i,j}^*, \quad Y \in B(\mathcal{H}).$$

Note that if $k = 1$ then $\mathbf{B}_n(\mathcal{H})$ coincides with the noncommutative unit ball $[B(\mathcal{H})^{n_1}]_1$. We remark that the scalar representation of the (abstract) regular polyball $\mathbf{B}_n := \{\mathbf{B}_n(\mathcal{H}) : \mathcal{H} \text{ is a Hilbert space}\}$ is $\mathbf{B}_n(\mathbb{C}) = \mathbf{P}_n(\mathbb{C}) = (\mathbb{C}^{n_1})_1 \times \cdots \times (\mathbb{C}^{n_k})_1$.

Let H_{n_i} be an n_i -dimensional complex Hilbert space with orthonormal basis $e_1^i, \dots, e_{n_i}^i$. We consider the *full Fock space* of H_{n_i} , defined by $F^2(H_{n_i}) := \mathbb{C}1 \oplus \bigoplus_{p \geq 1} H_{n_i}^{\otimes p}$, where $H_{n_i}^{\otimes p}$ is the (Hilbert) tensor product of p copies of H_{n_i} . Let $\mathbb{F}_{n_i}^+$ be the unital free semigroup on n_i generators $g_1^i, \dots, g_{n_i}^i$ and the

identity g_0^i . Set $e_\alpha^i := e_{j_1}^i \otimes \cdots \otimes e_{j_p}^i$ if $\alpha = g_{j_1}^i \cdots g_{j_p}^i \in \mathbb{F}_{n_i}^+$ and $e_\alpha^i := 1 \in \mathbb{C}$. The length of $\alpha \in \mathbb{F}_{n_i}^+$ is defined by $|\alpha| := 0$ if $\alpha = g_0^i$ and $|\alpha| := p$ if $\alpha = g_{j_1}^i \cdots g_{j_p}^i$ with $j_1, \dots, j_p \in \{1, \dots, n_i\}$. We define the *left creation operator* $S_{i,j}$ acting on the Fock space $F^2(H_{n_i})$ by setting $S_{i,j}e_\alpha^i := e_{g_j^i\alpha}^i$, $\alpha \in \mathbb{F}_{n_i}^+$, and the operator $\mathbf{S}_{i,j}$ acting on the Hilbert tensor product $F^2(H_{n_1}) \otimes \cdots \otimes F^2(H_{n_k})$ by setting

$$\mathbf{S}_{i,j} := \underbrace{I \otimes \cdots \otimes I}_{i-1 \text{ times}} \otimes S_{i,j} \otimes \underbrace{I \otimes \cdots \otimes I}_{k-i \text{ times}}$$

where $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$. We define $\mathbf{S} := (\mathbf{S}_1, \dots, \mathbf{S}_k)$, where $\mathbf{S}_i := (\mathbf{S}_{i,1}, \dots, \mathbf{S}_{i,n_i})$, or write $\mathbf{S} := \{\mathbf{S}_{i,j}\}$. The noncommutative Hardy algebra \mathbf{F}_n^∞ (resp. the polyball algebra \mathcal{A}_n) is the weakly closed (resp. norm closed) nonselfadjoint algebra generated by $\{\mathbf{S}_{i,j}\}$ and the identity. Similarly, we define the *right creation operator* $R_{i,j} : F^2(H_{n_i}) \rightarrow F^2(H_{n_i})$ by setting $R_{i,j}e_\alpha^i := e_{\alpha g_j^i}^i$ for $\alpha \in \mathbb{F}_{n_i}^+$, and the corresponding operator $\mathbf{R}_{i,j}$ acting on $F^2(H_{n_1}) \otimes \cdots \otimes F^2(H_{n_k})$. The polyball algebra \mathcal{R}_n is the norm closed nonselfadjoint algebra generated by $\{\mathbf{R}_{i,j}\}$ and the identity.

We proved in [Popescu 2016] (in a more general setting) that $X \in B(\mathcal{H})^{n_1} \times \cdots \times B(\mathcal{H})^{n_k}$ is a *pure* element in the regular polyball $\mathbf{B}_n(\mathcal{H})^-$, i.e., $\lim_{q_i \rightarrow \infty} \Phi_{X_i}^{q_i}(I) = 0$ in the weak operator topology, if and only if there is a Hilbert space \mathcal{D} and a subspace $\mathcal{M} \subset F^2(H_{n_1}) \otimes \cdots \otimes F^2(H_{n_k}) \otimes \mathcal{D}$ invariant under each operator $\mathbf{S}_{i,j} \otimes I$ such that $X_{i,j}^* = (\mathbf{S}_{i,j}^* \otimes I)|_{\mathcal{M}^\perp}$, under an appropriate identification of \mathcal{H} with \mathcal{M}^\perp . The k -tuple $\mathbf{S} := (\mathbf{S}_1, \dots, \mathbf{S}_k)$, where $\mathbf{S}_i := (\mathbf{S}_{i,1}, \dots, \mathbf{S}_{i,n_i})$, is an element in the regular polyball $\mathbf{B}_n(\bigotimes_{i=1}^k F^2(H_{n_i}))^-$ and plays the role of *left universal model* for the abstract polyball $\mathbf{B}_n^- := \{\mathbf{B}_n(\mathcal{H})^- : \mathcal{H} \text{ is a Hilbert space}\}$. The existence of the universal model will play an important role in this paper, since it will make the connection between noncommutative function theory, operator algebras, and complex function theory in several variables.

Brown and Halmos [1963] showed that a bounded linear operator T on the Hardy space $H^2(\mathbb{D})$ is a Toeplitz operator if and only if $S^*TS = T$, where S is the unilateral shift. Expanding on this idea, a study of noncommutative multi-Toeplitz operators on the full Fock space with n generators $F^2(H_n)$ was initiated in [Popescu 1989; 1995] and has had an important impact in multivariable operator theory and the structure of free semigroup algebras (see [Davidson and Pitts 1998; Davidson et al. 2001; 2005; Popescu 2006; 2009; Kennedy 2011; 2013]).

In Section 1, we introduce and study the class \mathcal{T}_n , $\mathbf{n} := (n_1, \dots, n_k) \in \mathbb{N}^k$, of k -multi-Toeplitz operators. A bounded linear operator T on the tensor product $F^2(H_{n_1}) \otimes \cdots \otimes F^2(H_{n_k})$ of full Fock spaces is called a k -multi-Toeplitz operator with respect to the *right universal model* $\mathbf{R} = \{\mathbf{R}_{i,j}\}$ if

$$\mathbf{R}_{i,s}^* T \mathbf{R}_{i,t} = \delta_{st} T, \quad s, t \in \{1, \dots, n_i\},$$

for every $i \in \{1, \dots, k\}$. We associate with each k -multi-Toeplitz operator T a formal power series in several variables and show that we can recapture T from its noncommutative ‘‘Fourier series’’. Moreover, we characterize the noncommutative formal power series which are Fourier series of k -multi-Toeplitz operators (see Theorems 1.5 and 1.6). Using these results, we prove that the set of all k -multi-Toeplitz operators on $\bigotimes_{i=1}^k F^2(H_{n_i})$ coincides with

$$\text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{-\text{SOT}} = \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{-\text{WOT}},$$

where \mathcal{A}_n is the noncommutative polyball algebra.

In Section 2, we characterize the bounded free k -pluriharmonic functions on regular polyballs. We prove that a function $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{H})$ is a bounded free k -pluriharmonic function if and only if there is a k -multi-Toeplitz operator $A \in \mathcal{T}_n$ such that $F(X) = \mathcal{B}_X[A]$ for $X \in \mathbf{B}_n(\mathcal{H})$, where \mathcal{B}_X is the noncommutative Berezin transform at X (see Section 1 for the definition). In this case, $A = \text{SOT-lim}_{r \rightarrow 1} F(rS)$ and there is a completely isometric isomorphism of operator spaces

$$\Phi : \mathbf{PH}^\infty(\mathbf{B}_n) \rightarrow \mathcal{T}_n, \quad \Phi(F) := A,$$

where $\mathbf{PH}^\infty(\mathbf{B}_n)$ is the operator space of all bounded free k -pluriharmonic functions on the polyball.

The Dirichlet extension problem [Hoffman 1962] on noncommutative regular polyballs is solved. We show that a mapping $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{H})$ is a free k -pluriharmonic function which has continuous extension (in the operator norm topology) to the closed polyball $\mathbf{B}_n(\mathcal{H})^-$, and write $F \in \mathbf{PH}^c(\mathbf{B}_n)$, if and only if there exists a k -multi-Toeplitz operator $A \in \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{-\|\cdot\|}$ such that $F(X) = \mathcal{B}_X[A]$ for $X \in \mathbf{B}_n(\mathcal{H})$. In this case, $A = \lim_{r \rightarrow 1} F(rS)$, where the convergence is in the operator norm, and the map

$$\Phi : \mathbf{PH}^c(\mathbf{B}_n) \rightarrow \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{-\|\cdot\|}, \quad \Phi(F) := A,$$

is a completely isometric isomorphism of operator spaces.

In Section 3, we provide a Naimark-type dilation theorem [1943] for direct products $\mathbf{F}_n^+ := \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$ of free semigroups. We show that a map $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$ is a positive semidefinite left k -multi-Toeplitz kernel on \mathbf{F}_n^+ if and only if there exists a k -tuple of commuting row isometries $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, on a Hilbert space $\mathcal{K} \supset \mathcal{E}$ — i.e., the nonselfadjoint algebra $\text{Alg}(V_i)$ commutes with $\text{Alg}(V_s)$ for any $i, s \in \{1, \dots, k\}$ with $i \neq s$ — such that

$$K(\sigma, \omega) = P_{\mathcal{E}} V_\sigma^* V_\omega|_{\mathcal{E}}, \quad \sigma, \omega \in \mathbf{F}_n^+,$$

and $\mathcal{K} = \bigvee_{\omega \in \mathbf{F}_n^+} V_\omega \mathcal{E}$. In this case, the minimal dilation is unique up to isomorphism. Here, we use the notation $V_\sigma := V_{1,\sigma_1} \cdots V_{k,\sigma_k}$ if $\sigma = (\sigma_1, \dots, \sigma_k) \in \mathbf{F}_n^+$, and $V_{i,\sigma_i} := V_{i,j_1} \cdots V_{i,j_p}$ if $\sigma_i = g_{j_1}^{i_1} \cdots g_{j_p}^{i_p} \in \mathbb{F}_{n_i}^+$ and $V_{i,g_0^i} := I$. For more information on kernels in various noncommutative settings we refer the reader to the work of Ball and Vinnikov [2003] (see also [Ball et al. 2016] and the references therein).

We prove a Schur-type result [1918], which states that a free k -pluriharmonic function F on the polyball \mathbf{B}_n is positive if and only if a certain right k -multi-Toeplitz kernel Γ_{F_r} associated with the mapping $S \mapsto F(rS)$ is positive semidefinite for any $r \in [0, 1)$. Our Naimark-type result for positive semidefinite right k -multi-Toeplitz kernels on \mathbf{F}_n^+ is used to provide a structure theorem for positive free k -pluriharmonic functions. We show that a free k -pluriharmonic function $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ with $F(0) = I$ is positive if and only if it has the form

$$F(X) = \sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} V_{\tilde{\alpha}}^* V_{\tilde{\beta}}|_{\mathcal{E}} \otimes X_\alpha X_\beta^*,$$

where $\mathbf{V} = (V_1, \dots, V_k)$ is a k -tuple of commuting row isometries on a space $\mathcal{K} \supset \mathcal{E}$ and $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_k)$ is the reverse of $\alpha = (\alpha_1, \dots, \alpha_k)$, i.e., $\tilde{\alpha}_i = g_{i_k}^{i_1} \cdots g_{i_1}^{i_k}$ if $\alpha_i = g_{i_1}^{i_1} \cdots g_{i_k}^{i_k} \in \mathbb{F}_{n_i}^+$. The general case, when $F(0) \geq 0$,

is also considered. As a consequence of these results, we obtain a structure theorem for positive k -harmonic functions on the regular polydisk included in $[B(\mathcal{H})]_1 \times_c \cdots \times_c [B(\mathcal{H})]_1$, which extends the corresponding classical result in scalar polydisks [Rudin 1969].

In Section 4, we define the *free pluriharmonic Poisson kernel* on the regular polyball \mathbf{B}_n by setting

$$\mathcal{P}(\mathbf{R}, \mathbf{X}) := \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mathbf{R}_{1, \tilde{\alpha}_1}^* \cdots \mathbf{R}_{k, \tilde{\alpha}_k}^* \mathbf{R}_{1, \tilde{\beta}_1} \cdots \mathbf{R}_{k, \tilde{\beta}_k} \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*$$

for any $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$, where the convergence is in the operator norm topology. Given a completely bounded linear map $\mu : \text{span}\{\mathcal{R}_n^* \mathcal{R}_n\} \rightarrow B(\mathcal{E})$, we introduce the *noncommutative Poisson transform* of μ to be the map $\mathcal{P}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ defined by

$$(\mathcal{P}\mu)(\mathbf{X}) := \hat{\mu}[\mathcal{P}(\mathbf{R}, \mathbf{X})], \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}),$$

where the completely bounded linear map

$$\hat{\mu} := \mu \otimes \text{id} : \text{span}\{\mathcal{R}_n^* \mathcal{R}_n\}^{-\|\cdot\|} \otimes_{\min} B(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$$

is uniquely defined by $\hat{\mu}(A \otimes Y) := \mu(A) \otimes Y$ for any $A \in \text{span}\{\mathcal{R}_n^* \mathcal{R}_n\}$ and $Y \in B(\mathcal{H})$. We remark that, in the particular case when $n_1 = \cdots = n_k = 1$, $\mathcal{H} = \mathcal{K} = \mathbb{C}$, $\mathbf{X} = (X_1, \dots, X_k) \in \mathbb{D}^k$, and μ is a complex Borel measure on \mathbb{T}^k (which can be seen as a bounded linear functional on $C(\mathbb{T}^k)$), we have that the noncommutative Poisson transform of μ coincides with the classical Poisson transform of μ [Rudin 1969].

In Section 4, we give necessary and sufficient conditions for a function $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ to be the noncommutative Poisson transform of a completely bounded linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$, where $C^*(\mathbf{R})$ is the C^* -algebra generated by the operators $\mathbf{R}_{i,j}$. In this case, we show that there exist a k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, of *doubly commuting row isometries* acting on a Hilbert space \mathcal{K} , i.e., $C^*(V_i)$ commutes with $C^*(V_j)$ if $i \neq j$, and bounded linear operators $W_1, W_2 : \mathcal{E} \rightarrow \mathcal{K}$ such that

$$F(\mathbf{X}) = (W_1^* \otimes I)[C_X(\mathbf{V})^* C_X(\mathbf{V})](W_2 \otimes I), \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}),$$

where

$$C_X(\mathbf{V}) := (I \otimes \Delta_X(I)^{1/2}) \prod_{i=1}^k (I - V_{i,1} \otimes X_{i,1}^* - \cdots - V_{i,n_i} \otimes X_{i,n_i}^*)^{-1}.$$

In particular, we obtain necessary and sufficient conditions for a function $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ to be the noncommutative Poisson transform of a completely positive linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$. In this case, we have the representation

$$F(\mathbf{X}) = (W^* \otimes I)[C_X(\mathbf{V})^* C_X(\mathbf{V})](W \otimes I), \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

In Section 5, we introduce the *noncommutative Herglotz–Riesz transform* of a completely positive linear map $\mu : \text{span}\{\mathcal{R}_n^* \mathcal{R}_n\} \rightarrow B(\mathcal{E})$ as the map $\mathbf{H}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ defined by

$$(\mathbf{H}\mu)(\mathbf{X}) := \hat{\mu} \left(2 \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \cdots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right)$$

for $X := (X_1, \dots, X_n) \in \mathbf{B}_n(\mathcal{H})$. The main result of this section provides necessary and sufficient conditions for a function f from the polyball $\mathbf{B}_n(\mathcal{H})$ to $B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ to admit a Herglotz–Riesz-type representation [Herglotz 1911; Riesz 1911], i.e.,

$$f(X) = (\mathbf{H}\mu)(X) + i\Im f(0), \quad X \in \mathbf{B}_n(\mathcal{H}),$$

where $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ is a completely positive linear map with the property that $\mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in \mathbf{F}_n^+$. In this case, we show that there exist a k -tuple $V = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, of doubly commuting row isometries on a Hilbert space \mathcal{K} and a bounded linear operator $W : \mathcal{E} \rightarrow \mathcal{K}$ such that

$$f(X) = (W^* \otimes I) \left(2 \prod_{i=1}^k (I - V_{i,1}^* \otimes X_{i,1} - \dots - V_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right) (W \otimes I) + i\Im f(0)$$

and $W^* V_\alpha^* V_\beta W = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in \mathbf{F}_n^+$.

We remark that, in the particular case when $n_1 = \dots = n_k = 1$, we obtain an operator-valued extension of the integral representation for holomorphic functions with positive real parts in polydisks [Korányi and Pukánszky 1963].

1. k -multi-Toeplitz operators on tensor products of full Fock spaces

In this section, we introduce the class \mathcal{T}_n of k -multi-Toeplitz operators on tensor products of full Fock spaces. We associate with each k -multi-Toeplitz operator T a formal power series in several variables and show that we can recapture T from its noncommutative Fourier series. Moreover, we characterize the noncommutative formal power series which are Fourier series of k -multi-Toeplitz operators and prove that $\mathcal{T}_n = \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{\text{SOT}}$, where \mathcal{A}_n is the noncommutative polyball algebra.

First, we recall (see [Popescu 1999; 2016]) some basic properties for a class of noncommutative Berezin-type transforms [1972] associated with regular polyballs. Let $X = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})^-$ with $X_i := (X_{i,1}, \dots, X_{i,n_i})$. We use the notation $X_{i,\alpha_i} := X_{i,j_1} \cdots X_{i,j_p}$ if $\alpha_i = g_{j_1}^i \cdots g_{j_p}^i \in \mathbb{F}_{n_i}^+$ and $X_{i,g_0^i} := I$. The noncommutative Berezin kernel associated with any element X in the noncommutative polyball $\mathbf{B}_n(\mathcal{H})^-$ is the operator

$$\mathbf{K}_X : \mathcal{H} \rightarrow F^2(H_{n_1}) \otimes \dots \otimes F^2(H_{n_k}) \otimes \overline{\Delta_X(I)(\mathcal{H})}$$

defined by

$$\mathbf{K}_X h := \sum_{\beta_i \in \mathbb{F}_{n_i}^+, i=1, \dots, k} e_{\beta_1}^1 \otimes \dots \otimes e_{\beta_k}^k \otimes \Delta_X(I)^{1/2} X_{1,\beta_1}^* \cdots X_{k,\beta_k}^* h, \quad h \in \mathcal{H},$$

where the defect operator $\Delta_X(I)$ was defined in the introduction. A very important property of the Berezin kernel is that $\mathbf{K}_X X_{i,j}^* = (\mathbf{S}_{i,j}^* \otimes I) \mathbf{K}_X$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$. The Berezin transform at $X \in \mathbf{B}_n(\mathcal{H})$ is the map $\mathcal{B}_X : B(\bigotimes_{i=1}^k F^2(H_{n_i})) \rightarrow B(\mathcal{H})$ defined by

$$\mathcal{B}_X[g] := \mathbf{K}_X^*(g \otimes I_{\mathcal{H}}) f \mathbf{K}_X, \quad g \in B\left(\bigotimes_{i=1}^k F^2(H_{n_i})\right).$$

If g is in the C^* -algebra $C^*(\mathcal{S})$ generated by $\mathcal{S}_{i,1}, \dots, \mathcal{S}_{i,n_i}$, where $i \in \{1, \dots, k\}$, we define the Berezin transform at $X \in \mathbf{B}_n(\mathcal{H})^-$ by

$$\mathcal{B}_X[g] := \lim_{r \rightarrow 1} \mathbf{K}_{rX}^*(g \otimes I_{\mathcal{H}})\mathbf{K}_{rX}, \quad g \in C^*(\mathcal{S}),$$

where the limit is in the operator norm topology. In this case, the Berezin transform at X is a unital completely positive linear map such that

$$\mathcal{B}_X(\mathcal{S}_{\alpha}\mathcal{S}_{\beta}^*) = X_{\alpha}X_{\beta}^*, \quad \alpha, \beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+,$$

where $\mathcal{S}_{\alpha} := \mathcal{S}_{1,\alpha_1} \cdots \mathcal{S}_{k,\alpha_k}$ if $\alpha := (\alpha_1, \dots, \alpha_k) \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$.

The Berezin transform will play an important role in this paper. More properties concerning noncommutative Berezin transforms and multivariable operator theory on noncommutative balls and polydomains can be found in [Popescu 1999; 2013; 2016]. For basic results on completely positive and completely bounded maps we refer the reader to [Paulsen 1986; Pisier 2001; Effros and Ruan 2000].

Definition 1.1. Let \mathcal{E} be a Hilbert space. A bounded linear operator $A \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ is called k -multi-Toeplitz with respect to the universal model $\mathbf{R} := (\mathbf{R}_1, \dots, \mathbf{R}_k)$, where $\mathbf{R}_i := (\mathbf{R}_{i,1}, \dots, \mathbf{R}_{i,n_i})$, if

$$(I_{\mathcal{E}} \otimes \mathbf{R}_{i,s}^*)A(I_{\mathcal{E}} \otimes \mathbf{R}_{i,t}) = \delta_{st}A, \quad s, t \in \{1, \dots, n_i\},$$

for every $i \in \{1, \dots, k\}$.

A few more notations are necessary. If $\omega, \gamma \in \mathbb{F}_n^+$, we say that $\gamma <_r \omega$ if there is $\sigma \in \mathbb{F}_n^+ \setminus \{g_0\}$ such that $\omega = \sigma\gamma$. In this case, we set $\omega \setminus_r \gamma := \sigma$. Similarly, we say that $\gamma <_l \omega$ if there is $\sigma \in \mathbb{F}_n^+ \setminus \{g_0\}$ such that $\omega = \gamma\sigma$ and set $\omega \setminus_l \gamma := \sigma$. We denote by $\tilde{\alpha}$ the reverse of $\alpha \in \mathbb{F}_n^+$, i.e., $\tilde{\alpha} = g_{i_k} \cdots g_{i_1}$ if $\alpha = g_{i_1} \cdots g_{i_k} \in \mathbb{F}_n^+$. Notice that $\gamma <_r \omega$ if and only if $\tilde{\gamma} <_l \tilde{\omega}$. In this case we have $(\omega \setminus_r \gamma)^\sim = \tilde{\omega} \setminus_l \tilde{\gamma}$. We say that ω is *right comparable* with γ , and write $\omega \sim_{rc} \gamma$, if any one of the conditions $\omega <_r \gamma$, $\gamma <_r \omega$ or $\omega = \gamma$ holds. In this case, we define

$$c_r^+(\omega, \gamma) := \begin{cases} \omega \setminus_r \gamma & \text{if } \gamma <_r \omega, \\ g_0 & \text{if } \omega <_r \gamma \text{ or } \omega = \gamma, \end{cases} \quad \text{and} \quad c_r^-(\omega, \gamma) := \begin{cases} \gamma \setminus_r \omega & \text{if } \omega <_r \gamma, \\ g_0 & \text{if } \gamma <_r \omega \text{ or } \omega = \gamma. \end{cases}$$

Let $\omega = (\omega_1, \dots, \omega_k)$ and $\gamma = (\gamma_1, \dots, \gamma_k)$ be in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$. We say that ω and γ are *right comparable*, and write $\omega \sim_{rc} \gamma$, if for each $i \in \{1, \dots, k\}$, any one of the conditions $\omega_i <_r \gamma_i$, $\gamma_i <_r \omega_i$ or $\omega_i = \gamma_i$ holds. In this case, we define

$$c_r^+(\omega, \gamma) := (c_r^+(\omega_1, \gamma_1), \dots, c_r^+(\omega_k, \gamma_k)) \quad \text{and} \quad c_r^-(\omega, \gamma) := (c_r^-(\omega_1, \gamma_1), \dots, c_r^-(\omega_k, \gamma_k)). \quad (1-1)$$

Similarly, we say that ω and γ are *left comparable*, and write $\omega \sim_{lc} \gamma$, if $\tilde{\omega} \sim_{rc} \tilde{\gamma}$. The definitions of $c_l^+(\omega, \gamma)$ and $c_l^-(\omega, \gamma)$ are now clear. Note that

$$c_r^+(\omega, \gamma)^\sim = c_l^+(\tilde{\omega}, \tilde{\gamma}) \quad \text{and} \quad c_r^-(\omega, \gamma)^\sim = c_l^-(\tilde{\omega}, \tilde{\gamma}).$$

For each $m \in \mathbb{Z}$, we set $m^+ := \max\{m, 0\}$ and $m^- := \max\{-m, 0\}$.

Lemma 1.2. Let $\alpha = (\alpha_1, \dots, \alpha_k)$ and $\beta = (\beta_1, \dots, \beta_k)$ be k -tuples in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$ such that $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+$ for $i \in \{1, \dots, k\}$ with $|\alpha_i| = m_i^-$, $|\beta_i| = m_i^+$ and $m_i \in \mathbb{Z}$. If $\gamma = (\gamma_1, \dots, \gamma_k)$ and $\omega = (\omega_1, \dots, \omega_k)$ are k -tuples in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, then the inner product

$$\langle S_{1,\alpha_1} \cdots S_{k,\alpha_k} S_{1,\beta_1}^* \cdots S_{k,\beta_k}^* (e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k), e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k \rangle$$

is different from zero if and only if $\omega \sim_{rc} \gamma$ and $(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k) = (c_r^+(\omega, \gamma); c_r^-(\omega, \gamma))$.

Proof. Under the conditions of the lemma, $S_{i,\alpha_i} S_{j,\beta_j}^* = S_{j,\beta_j}^* S_{i,\alpha_i}$ for any $i, j \in \{1, \dots, k\}$, $\alpha_i \in \mathbb{F}_{n_i}^+$ and $\beta_j \in \mathbb{F}_{n_j}^+$. Note that the inner product is different from zero if and only if $\beta_i \omega_i = \alpha_i \gamma_i$ for any $i \in \{1, \dots, k\}$. Let $m_i \in \mathbb{Z}$ and assume that $|\alpha_i| = m_i^- > 0$. Then $\beta_i = g_0^i$ and, consequently, $\omega_i = \alpha_i \gamma_i$. This shows that $\gamma_i <_r \omega_i$, $c_r^+(\omega_i, \gamma_i) = \alpha_i$ and $c_r^-(\omega_i, \gamma_i) = g_0^i$. In the case when $|\beta_i| = m_i^+ > 0$, we have $\alpha_i = g_0^i$ and $\beta_i \omega_i = \gamma_i$. Consequently, $\omega_i <_r \gamma_i$, $c_r^+(\omega_i, \gamma_i) = g_0^i$ and $c_r^-(\omega_i, \gamma_i) = \beta_i$. When $\alpha_i = \beta_i = g_0^i$, we have $\omega_i = \gamma_i$. Therefore, the scalar product above is different from zero if and only if $\omega \sim_{rc} \gamma$ and $(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k) = (c_r^+(\omega, \gamma); c_r^-(\omega, \gamma))$. \square

If $\beta_i, \gamma_i \in \mathbb{F}_{n_i}^+$ and, for each $i \in \{1, \dots, k\}$, $\beta_i <_\ell \gamma_i$ or $\beta_i = \gamma_i$, then we write $\beta \leq_\ell \gamma$.

Lemma 1.3. Given a k -tuple $\gamma = (\gamma_1, \dots, \gamma_k) \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, the sequence

$$\{S_{1,\alpha_1} \cdots S_{k,\alpha_k} S_{1,\beta_1}^* \cdots S_{k,\beta_k}^* (e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k)\}$$

consists of orthonormal vectors if $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+$, $i \in \{1, \dots, k\}$ with $m_i \in \mathbb{Z}$, $|\alpha_i| = m_i^-$, $|\beta_i| = m_i^+$ and $\beta \leq_\ell \gamma$.

Proof. First, note that $S_{1,\alpha_1} \cdots S_{k,\alpha_k} S_{1,\beta_1}^* \cdots S_{k,\beta_k}^* (e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k) \neq 0$ if and only if $S_{i,\beta_i}^* (e_{\gamma_i}^i) \neq 0$ for each $i \in \{1, \dots, k\}$, which is equivalent to $\beta_i <_\ell \gamma_i$ or $\beta_i = \gamma_i$. Therefore, $\beta \leq_\ell \gamma$.

Fix $i \in \{1, \dots, k\}$ and $\gamma_i \in \mathbb{F}_{n_i}^+$. We prove that the sequence $\{S_{i,\alpha_i} S_{i,\beta_i}^* e_{\gamma_i}^i\}$ consists of orthonormal vectors if $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+$ have the following properties:

(i) If $|\alpha_i| > 0$ then $\beta_i = g_0^i$, and if $|\beta_i| > 0$ then $\alpha_i = g_0^i$.

(ii) $\beta_i \leq_\ell \gamma_i$.

Indeed, let (α_i, β_i) and (α'_i, β'_i) be two distinct pairs with the above-mentioned properties. First, we consider the case when $g_0^i \neq \beta_i <_\ell \gamma_i$. Then $\alpha_i = g_0^i$ and, consequently, $S_{i,\alpha_i} S_{i,\beta_i}^* e_{\gamma_i}^i = e_{\gamma_i \setminus \beta_i}^i$. Similarly, if $g_0^i \neq \beta'_i <_\ell \gamma_i$ then $\alpha'_i = g_0^i$ and, consequently, $S_{i,\alpha'_i} S_{i,\beta'_i}^* e_{\gamma_i}^i = e_{\gamma_i \setminus \beta'_i}^i$. Since $(\alpha_i, \beta_i) \neq (\alpha'_i, \beta'_i)$, we must have $\beta_i \neq \beta'_i$, which implies $e_{\gamma_i \setminus \beta_i}^i \perp e_{\gamma_i \setminus \beta'_i}^i$. On the other hand, if $\beta'_i = g_0^i$ then $\alpha'_i \in \mathbb{F}_{n_i}^+$ and $S_{i,\alpha'_i} S_{i,\beta'_i}^* e_{\gamma_i}^i = e_{\alpha'_i} e_{\gamma_i} \perp e_{\gamma_i \setminus \beta_i}^i$. It follows that $S_{i,\alpha'_i} S_{i,\beta'_i}^* e_{\gamma_i}^i \perp S_{i,\alpha_i} S_{i,\beta_i}^* e_{\gamma_i}^i$.

The second case is when $\beta_i = g_0^i$. Then $\alpha_i \in \mathbb{F}_{n_i}^+$ and $S_{i,\alpha_i} S_{i,\beta_i}^* e_{\gamma_i}^i = e_{\alpha_i} e_{\gamma_i}$. As we saw above, $S_{i,\alpha'_i} S_{i,\beta'_i}^* e_{\gamma_i}^i$ is equal to either $e_{\alpha'_i} e_{\gamma_i}$ (when $\beta'_i = g_0^i$) or $e_{\gamma_i \setminus \beta'_i}^i$ (when $g_0^i \neq \beta'_i <_\ell \gamma_i$). In each case, we have $S_{i,\alpha'_i} S_{i,\beta'_i}^* e_{\gamma_i}^i \perp S_{i,\alpha_i} S_{i,\beta_i}^* e_{\gamma_i}^i$, which completes the proof of our assertion. Using this result one can easily complete the proof of the lemma. \square

We associate with each k -multi-Toeplitz operator $A \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ a formal power series

$$\varphi_A(\mathbf{S}) := \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \mathbf{S}_{1, \alpha_1} \cdots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \cdots \mathbf{S}_{k, \beta_k}^*,$$

where the coefficients are given by

$$\langle A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, \ell \rangle := \langle A(h \otimes x), \ell \otimes y \rangle, \quad h, \ell \in \mathcal{E}, \tag{1-2}$$

and $x := x_1 \otimes \cdots \otimes x_k, y = y_1 \otimes \cdots \otimes y_k$ with

$$\begin{cases} x_i = e_{\beta_i}^i & \text{and } y_i = 1 & \text{if } m_i \geq 0, \\ x_i = 1 & \text{and } y_i = e_{\alpha_i}^i & \text{if } m_i < 0, \end{cases} \tag{1-3}$$

for every $i \in \{1, \dots, k\}$.

The next result shows that a k -multi-Toeplitz operator is uniquely determined by its Fourier series.

Theorem 1.4. *If A, B are k -multi-Toeplitz operators on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$, then $A = B$ if and only if the corresponding formal Fourier series $\varphi_A(\mathbf{S})$ and $\varphi_B(\mathbf{S})$ are equal. Moreover, $Aq = \varphi_A(\mathbf{S})q$ for any vector-valued polynomial*

$$q = \sum_{\substack{\omega_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\omega_i| \leq p_i}} h_{(\omega_1, \dots, \omega_k)} \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k,$$

where $h_{(\omega_1, \dots, \omega_k)} \in \mathcal{E}$ and $(p_1, \dots, p_k) \in \mathbb{N}^k$.

Proof. Let $\boldsymbol{\omega} = (\omega_1, \dots, \omega_k)$ and $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k)$ be k -tuples in $\mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, and let $h, h' \in \mathcal{E}$. Since A is a k -multi-Toeplitz operator on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$, we have

$$\begin{aligned} \langle A(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k \rangle \\ = \langle A(I_{\mathcal{E}} \otimes \mathbf{R}_{1, \tilde{\gamma}_1} \cdots \mathbf{R}_{k, \tilde{\gamma}_k})(h \otimes 1), (I_{\mathcal{E}} \otimes \mathbf{R}_{1, \tilde{\omega}_1} \cdots \mathbf{R}_{k, \tilde{\omega}_k})(h' \otimes 1) \rangle \\ = \begin{cases} \langle A_{(c_r^+(\boldsymbol{\omega}, \boldsymbol{\gamma}); c_r^-(\boldsymbol{\omega}, \boldsymbol{\gamma}))} h, h' \rangle & \text{if } \boldsymbol{\omega} \sim_{rc} \boldsymbol{\gamma}, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

where $c_r^+(\boldsymbol{\omega}, \boldsymbol{\gamma})$ and $c_r^-(\boldsymbol{\omega}, \boldsymbol{\gamma})$ are defined by (1-1). Consequently,

$$A(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k) = \sum_{\substack{\boldsymbol{\omega} = (\omega_1, \dots, \omega_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+ \\ \boldsymbol{\omega} \sim_{rc} \boldsymbol{\gamma}}} A_{(c_r^+(\boldsymbol{\omega}, \boldsymbol{\gamma}); c_r^-(\boldsymbol{\omega}, \boldsymbol{\gamma}))} h \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k$$

is a vector in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. Hence, we deduce that, for each $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, the series

$$\sum_{\substack{\boldsymbol{\omega} \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+ \\ \boldsymbol{\omega} \sim_{rc} \boldsymbol{\gamma}}} A_{(c_r^+(\boldsymbol{\omega}, \boldsymbol{\gamma}); c_r^-(\boldsymbol{\omega}, \boldsymbol{\gamma}))}^* A_{(c_r^+(\boldsymbol{\omega}, \boldsymbol{\gamma}); c_r^-(\boldsymbol{\omega}, \boldsymbol{\gamma}))} \tag{1-4}$$

is WOT-convergent. Due to Lemma 1.3, given $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, the sequence $\{\mathbf{S}_{1, \alpha_1} \cdots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \cdots \mathbf{S}_{k, \beta_k}^* (e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k)\}$, where $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\}$ with $m_i \in \mathbb{Z}, |\alpha_i| = m_i^-,$

$|\beta_i| = m_i^+$ and $\beta \leq_\ell \gamma$, consists of orthonormal vectors. Note that, in this case, we also have $\alpha \sim_{rc} \beta$ and, consequently, $A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} = A_{(c_r^+(\alpha, \beta); c_r^-(\alpha, \beta))}$. Hence, and taking into account that the series (1-4) is WOT-convergent, we deduce that

$$\begin{aligned} \varphi_A(\mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k) &:= \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h \otimes \mathbf{S}_{1, \alpha_1} \dots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \dots \mathbf{S}_{k, \beta_k}^* (e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k) \end{aligned}$$

is a convergent series in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. Let $\gamma = (\gamma_1, \dots, \gamma_k)$ and $\omega = (\omega_1, \dots, \omega_k)$ be k -tuples in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$. According to Lemma 1.2, the inner product

$$\langle \mathbf{S}_{1, \alpha_1} \dots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \dots \mathbf{S}_{k, \beta_k}^* (e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k), e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_k}^k \rangle$$

is different from zero if and only if $\omega \sim_{rc} \gamma$ and $(\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k) = (c_r^+(\omega, \gamma); c_r^-(\omega, \gamma))$. Now, using (1-4), one can see that

$$\begin{aligned} &\langle \varphi_A(\mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_k}^k \rangle \\ &= \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \langle A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, h' \rangle \\ &\quad \times \langle \mathbf{S}_{1, \alpha_1} \dots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \dots \mathbf{S}_{k, \beta_k}^* (e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k), e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_k}^k \rangle \\ &= \begin{cases} \langle A_{(c_r^+(\omega, \gamma); c_r^-(\omega, \gamma))} h, h' \rangle & \text{if } \omega \sim_{rc} \gamma, \\ 0 & \text{otherwise,} \end{cases} \\ &= \langle A(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_k}^k \rangle \end{aligned}$$

for any $h, h' \in \mathcal{E}$, and $\gamma = (\gamma_1, \dots, \gamma_k)$ and $\omega = (\omega_1, \dots, \omega_k)$ in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, which shows that $Aq = \varphi_A(\mathbf{S})q$ for any vector-valued polynomial in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. Therefore, if the formal Fourier series $\varphi_A(\mathbf{S})$ and $\varphi_B(\mathbf{S})$ are equal, then $A = B$. □

When \mathcal{G} is a Hilbert space, $C_{(\alpha; \beta)} \in B(\mathcal{G})$, and the series

$$\Sigma_1 := \sum_{m \in \mathbb{Z}, m < 0} \sum_{\substack{\alpha, \beta \in \mathbb{F}_n^+ \\ |\alpha| = m^-, |\beta| = m^+}} C_{(\alpha; \beta)} \quad \text{and} \quad \Sigma_2 := \sum_{m \in \mathbb{Z}, m \geq 0} \sum_{\substack{\alpha, \beta \in \mathbb{F}_n^+ \\ |\alpha| = m^-, |\beta| = m^+}} C_{(\alpha; \beta)}$$

are convergent in the operator topology, we say that the series

$$\sum_{m \in \mathbb{Z}} \sum_{\substack{\alpha, \beta \in \mathbb{F}_n^+ \\ |\alpha| = m^-, |\beta| = m^+}} C_{(\alpha; \beta)} := \Sigma_1 + \Sigma_2$$

is convergent in the operator topology. In what follows, we show how we can recapture the k -multi-Toeplitz operators from their Fourier series. Moreover, we characterize the formal series which are Fourier series of k -multi-Toeplitz operators. Let \mathcal{P} denote the set of all vector-valued polynomials in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$,

i.e., each $p \in \mathcal{P}$ has the form

$$q = \sum_{\substack{\omega_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\omega_i| \leq p_i}} h_{(\omega_1, \dots, \omega_k)} \otimes e^1_{\omega_1} \otimes \dots \otimes e^k_{\omega_k},$$

where $h_{(\omega_1, \dots, \omega_k)} \in \mathcal{E}$ and $(p_1, \dots, p_k) \in \mathbb{N}^k$.

Theorem 1.5. *Let $\{A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)}\}$ be a family of operators in $B(\mathcal{E})$, where $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+$, $|\alpha_i| = m_i^-$, $|\beta_i| = m_i^+$, $m_i \in \mathbb{Z}$ and $i \in \{1, \dots, k\}$. Then*

$$\varphi(\mathbf{S}) := \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \mathbf{S}_{1, \alpha_1} \dots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \dots \mathbf{S}_{k, \beta_k}^*$$

is the formal Fourier series of a k -multi-Toeplitz operator on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ if and only if the following conditions are satisfied:

(i) For each $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k) \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, the series

$$\sum_{\substack{\omega \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+ \\ \omega \sim_{rc} \boldsymbol{\gamma}}} A_{(c_r^+(\omega, \boldsymbol{\gamma}); c_r^-(\omega, \boldsymbol{\gamma}))}^* A_{(c_r^+(\omega, \boldsymbol{\gamma}); c_r^-(\omega, \boldsymbol{\gamma}))}$$

is WOT-convergent.

(ii) If \mathcal{P} is the set of all vector-valued polynomials in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$, then

$$\sup_{r \in [0, 1)} \sup_{p \in \mathcal{P}, \|p\| \leq 1} \|\varphi(r\mathbf{S})p\| < \infty.$$

Moreover, if there is a k -multi-Toeplitz operator $A \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ such that $\varphi(\mathbf{S}) = \varphi_A(\mathbf{S})$, then the following statements hold:

(a)
$$\varphi(r\mathbf{S}) := \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes r^{\sum_{i=1}^k (|\alpha_i| + |\beta_i|)} \mathbf{S}_{1, \alpha_1} \dots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \dots \mathbf{S}_{k, \beta_k}^*$$

is convergent in the operator norm topology, and its sum, which does not depend on the order of the series, is an operator in

$$\text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{\|\cdot\|},$$

where \mathcal{A}_n is the polyball algebra.

(b) $A = \text{SOT-lim}_{r \rightarrow 1} \varphi(r\mathbf{S})$ and

$$\|A\| = \sup_{r \in [0, 1)} \|\varphi(r\mathbf{S})\| = \lim_{r \rightarrow 1} \|\varphi(r\mathbf{S})\| = \sup_{q \in \mathcal{P}, \|q\| \leq 1} \|\varphi(\mathbf{S})q\|.$$

Proof. First, we assume that $A \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ is a k -multi-Toeplitz operator and $\varphi(\mathbf{S}) = \varphi_A(\mathbf{S})$, where the coefficients $A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)}$ are given by (1-2) and (1-3). Note that (i) follows from the proof

of [Theorem 1.4](#). Moreover, from the same proof and [Lemma 1.3](#) we have $\varphi_A(\mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k)$ and, consequently, $\varphi_A(r\mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k)$, $r \in [0, 1)$, are vectors in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ and

$$\lim_{r \rightarrow 1} \varphi_A(r\mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k) = \varphi_A(\mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k) = A(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k) \tag{1-5}$$

for any $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k)$ in $\mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$ and $h \in \mathcal{E}$. Note also that, due to (i) and [Lemma 1.3](#), we have $\varphi_A(r_1\mathbf{S}_1, \dots, r_k\mathbf{S}_k)(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k) \in \mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ for any $r_i \in [0, 1)$, $i \in \{1, \dots, k\}$. Now, we show that the series

$$\sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \left(\prod_{i=1}^k r_i^{|\alpha_i| + |\beta_i|} \right) \mathbf{S}_{1, \alpha_1} \cdots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \cdots \mathbf{S}_{k, \beta_k}^*$$

is convergent in the operator norm topology and its sum is in $\text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{-\|\cdot\|}$, where \mathcal{A}_n is the polyball algebra. We denote the series above by $\varphi_A(r_1\mathbf{S}_1, \dots, r_k\mathbf{S}_k)$. Since A is a k -multi-Toeplitz operator, it is also a 1-multi-Toeplitz operator with respect to $R_k := (R_{k,1}, \dots, R_{k,n_k})$, the right creation operators on the Fock space $F^2(H_{n_k})$. Applying [Theorem 1.4](#) to 1-multi-Toeplitz operators, we deduce that A has a unique Fourier representation

$$\psi_A(S_k) := \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} C_{(\alpha_k; \beta_k)} \otimes S_{k, \alpha_k} S_{k, \beta_k}^*$$

where $C_{(\alpha_k; \beta_k)} \in B(\mathcal{E} \otimes \bigotimes_{i=1}^{k-1} F^2(H_{n_i}))$. Moreover, we can prove that, for any $r_k \in [0, 1)$,

$$\psi_A(r_k S_k) := \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} r_k^{|\alpha_k| + |\beta_k|} C_{(\alpha_k; \beta_k)} \otimes S_{k, \alpha_k} S_{k, \beta_k}^* \tag{1-6}$$

is convergent in the operator norm topology. Indeed, since $\psi_A(S_k)$ is the Fourier representation of the 1-multi-Toeplitz operator A with respect to $R_k := (R_{k,1}, \dots, R_{k,n_k})$, item (i) implies, in the particular case when $\gamma_k = g_0^k$, that $\sum_{\alpha_k \in \mathbb{F}_{n_k}^+} C_{(\alpha_k; g_0^k)} C_{(\alpha_k; g_0^k)}^*$ is WOT-convergent. Since A^* is also a 1-multi-Toeplitz operator, we can similarly deduce that the series $\sum_{\beta_k \in \mathbb{F}_{n_k}^+} C_{(g_0^k; \beta_k)} C_{(g_0^k; \beta_k)}^*$ is WOT-convergent. Since $S_{k,1}, \dots, S_{k,n_k}$ are isometries with orthogonal ranges, we have

$$\begin{aligned} \left\| \sum_{\alpha_k \in \mathbb{F}_{n_k}^+, |\alpha_k| = m} C_{(\alpha_k; g_0^k)} \otimes r_k^{|\alpha_k|} S_{k, \alpha_k} \right\| &= r_k^m \left\| \sum_{\alpha_k \in \mathbb{F}_{n_k}^+} C_{(\alpha_k; g_0^k)}^* C_{(\alpha_k; g_0^k)} \right\|^{1/2}, \\ \left\| \sum_{\beta_k \in \mathbb{F}_{n_k}^+, |\beta_k| = m} C_{(g_0^k; \beta_k)} \otimes r_k^{|\alpha_k|} S_{k, \beta_k}^* \right\| &= r_k^m \left\| \sum_{\alpha_k \in \mathbb{F}_{n_k}^+} C_{(g_0^k; \beta_k)} C_{(g_0^k; \beta_k)}^* \right\|^{1/2}, \end{aligned}$$

for any $m \in \mathbb{N}$. Now, it is clear that the series defining $\psi_A(r_k S_k)$ is convergent in the operator norm topology and, consequently, $\psi_A(r_k S_k)$ belongs to

$$\text{span} \left\{ f^*g : f, g \in B \left(\mathcal{E} \otimes \bigotimes_{i=1}^{k-1} F^2(H_{n_i}) \right) \otimes_{\min} \mathcal{A}_{n_k} \right\}^{-\|\cdot\|},$$

where \mathcal{A}_{n_k} is the noncommutative disc algebra generated by $S_{k,1}, \dots, S_{k,n_k}$ and the identity. For each $i \in \{1, \dots, k\}$, we set $\mathcal{E}_i := \mathcal{E} \otimes F^2(H_{n_1}) \otimes \dots \otimes F^2(H_{n_i})$.

The next step in our proof is to show that

$$\psi_A(r_k S_k) = \mathcal{B}_{r_k S_k}^{\text{ext}}[A] := (I_{\mathcal{E}_{k-1}} \otimes K_{r_k S_k}^*)(A \otimes I_{F^2(H_{n_k})})(I_{\mathcal{E}_{k-1}} \otimes K_{r_k S_k}), \tag{1-7}$$

where $K_{r_k S_k} : F^2(H_{n_k}) \rightarrow F^2(H_{n_k}) \otimes \mathcal{D}_{r_k S_k} \subset F^2(H_{n_k}) \otimes F^2(H_{n_k})$ is the noncommutative Berezin kernel defined by

$$K_{r_k S_k} \xi := \sum_{\beta_k \in \mathbb{F}_{n_k}^+} e_{\beta_k}^k \otimes \Delta_{r_k S_k}(I)^{1/2} S_{k, \beta_k}^* \xi, \quad \xi \in F^2(H_{n_k}),$$

and $\mathcal{D}_{r_k S_k} := \overline{\Delta_{r_k S_k}(I)(F^2(H_{n_k}))}$. Let $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k)$ and $\boldsymbol{\omega} = (\omega_1, \dots, \omega_k)$ be k -tuples in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, set $q := \max\{|\gamma_k|, |\omega_k|\}$, and define the operator

$$Q_q := \sum_{m_k \in \mathbb{Z}, |m_k| \leq q} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} C_{(\alpha_k; \beta_k)} \otimes S_{k, \alpha_k} S_{k, \beta_k}^*.$$

Since $\psi_A(S_k)p = Ap$ for any polynomial $p \in \mathcal{P}$, a careful computation reveals that

$$\begin{aligned} & \langle \mathcal{B}_{r_k S_k}^{\text{ext}}[A](h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_k}^k \rangle \\ &= \langle (A \otimes I_{F^2(H_{n_k})})(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_{k-1}}^{k-1} \otimes K_{r_k S_k}(e_{\gamma_k}^k)), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_{k-1}}^{k-1} \otimes K_{r_k S_k}(e_{\omega_k}^k) \rangle \\ &= \left\langle (A \otimes I_{F^2(H_{n_k})}) \left(\sum_{\alpha_k \in \mathbb{F}_{n_k}^+} h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_{k-1}}^{k-1} \otimes e_{\alpha_k}^k \otimes \Delta_{r_k S_k}(I)^{1/2} S_{k, \alpha_k}^*(e_{\gamma_k}^k) \right), \right. \\ & \quad \left. \sum_{\beta_k \in \mathbb{F}_{n_k}^+} h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_{k-1}}^{k-1} \otimes e_{\beta_k}^k \otimes \Delta_{r_k S_k}(I)^{1/2} S_{k, \beta_k}^*(e_{\omega_k}^k) \right\rangle \\ &= \left\langle \sum_{\alpha_k \in \mathbb{F}_{n_k}^+} A(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_{k-1}}^{k-1} \otimes e_{\alpha_k}^k) \otimes \Delta_{r_k S_k}(I)^{1/2} S_{k, \alpha_k}^*(e_{\gamma_k}^k), \right. \\ & \quad \left. \sum_{\beta_k \in \mathbb{F}_{n_k}^+} h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_{k-1}}^{k-1} \otimes e_{\beta_k}^k \otimes \Delta_{r_k S_k}(I)^{1/2} S_{k, \beta_k}^*(e_{\omega_k}^k) \right\rangle \\ &= \sum_{m=0}^{\infty} \sum_{p=0}^{\infty} \sum_{\substack{\alpha_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k|=m}} \sum_{\substack{\beta_k \in \mathbb{F}_{n_k}^+ \\ |\beta_k|=p}} \langle A(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_{k-1}}^{k-1} \otimes e_{\alpha_k}^k), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_{k-1}}^{k-1} \otimes e_{\beta_k}^k \rangle \\ & \quad \times \langle \Delta_{r_k S_k}(I)^{1/2} S_{k, \alpha_k}^*(e_{\gamma_k}^k), \Delta_{r_k S_k}(I)^{1/2} S_{k, \beta_k}^*(e_{\omega_k}^k) \rangle \\ &= \sum_{m=0}^q \sum_{p=0}^q \sum_{\substack{\alpha_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k|=m}} \sum_{\substack{\beta_k \in \mathbb{F}_{n_k}^+ \\ |\beta_k|=p}} \langle Q_q(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_{k-1}}^{k-1} \otimes e_{\alpha_k}^k), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_{k-1}}^{k-1} \otimes e_{\beta_k}^k \rangle \\ & \quad \times \langle \Delta_{r_k S_k}(I)^{1/2} S_{k, \alpha_k}^*(e_{\gamma_k}^k), \Delta_{r_k S_k}(I)^{1/2} S_{k, \beta_k}^*(e_{\omega_k}^k) \rangle \\ &= \sum_{m=0}^{\infty} \sum_{p=0}^{\infty} \sum_{\substack{\alpha_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k|=m}} \sum_{\substack{\beta_k \in \mathbb{F}_{n_k}^+ \\ |\beta_k|=p}} \langle Q_q(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_{k-1}}^{k-1} \otimes e_{\alpha_k}^k), h' \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_{k-1}}^{k-1} \otimes e_{\beta_k}^k \rangle \\ & \quad \times \langle \Delta_{r_k S_k}(I)^{1/2} S_{k, \alpha_k}^*(e_{\gamma_k}^k), \Delta_{r_k S_k}(I)^{1/2} S_{k, \beta_k}^*(e_{\omega_k}^k) \rangle \end{aligned}$$

$$\begin{aligned}
 &= \langle (Q_q \otimes I_{F^2(H_{n_k})})(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_{k-1}}^{k-1} \otimes K_{r_k S_k}(e_{\gamma_k}^k)), h' \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_{k-1}}^{k-1} \otimes K_{r_k S_k}(e_{\omega_k}^k) \rangle \\
 &= \langle \mathcal{B}_{r_k S_k}^{\text{ext}}[Q_q](h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k \rangle \\
 &= \sum_{m_k \in \mathbb{Z}, |m_k| \leq q} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} \langle (C_{(\alpha_k; \beta_k)} \otimes r_k^{|\alpha_k| + |\beta_k|} S_{k, \alpha_k} S_{k, \beta_k}^*)(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k \rangle \\
 &= \langle \psi_A(r_k S_k)(h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k), h' \otimes e_{\omega_1}^1 \otimes \cdots \otimes e_{\omega_k}^k \rangle
 \end{aligned}$$

for any $h, h' \in \mathcal{E}$. Consequently, (1-7) holds for any $r_k \in [0, 1)$. Hence, and using the fact the noncommutative Berezin kernel $K_{r_k S_k}$ is an isometry, we deduce that

$$\|\psi_A(r_k S_k)\| \leq \|A\|, \quad r_k \in [0, 1).$$

Moreover, one can show that

$$A = \text{SOT-}\lim_{r_k \rightarrow 1} \psi_A(r_k S_k).$$

Indeed, due to (i) (for 1-multi-Toeplitz operators), we have $\|\psi_A(r_k S_k)p - \psi_A(S_k)p\| \rightarrow 0$ as $r_k \rightarrow 1$ for any polynomial $p \in \mathcal{E}_{k-1} \otimes F^2(H_{n_k})$ with coefficients in \mathcal{E}_{k-1} . Since $\psi_A(S_k)p = Ap$ and $\|\psi_A(r_k S_k)\| \leq \|A\|$ for any $r_k \in [0, 1)$, an approximation argument proves our assertion.

Now, we prove that the coefficients $C_{(\alpha_k; \beta_k)} \in B(\mathcal{E} \otimes \bigotimes_{i=1}^{k-1} F^2(H_{n_i}))$ of the Fourier series $\psi_A(S_k)$ are 1-multi-Toeplitz operators with respect to $R_{k-1} := (R_{k-1,1}, \dots, R_{k-1, n_{k-1}})$. For each $i \in \{1, \dots, k-1\}$, $s, t \in \{1, \dots, n_i\}$, and any vector-valued polynomial $p \in \mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ with coefficients in \mathcal{E} , [Theorem 1.4](#) implies

$$\begin{aligned}
 \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} & [(I_{\mathcal{E}_{k-2}} \otimes R_{i,s}^*) C_{(\alpha_k; \beta_k)} (I_{\mathcal{E}_{k-2}} \otimes R_{i,t}) \otimes S_{k, \alpha_k} S_{k, \beta_k}^*](p) \\
 &= (I_{\mathcal{E}} \otimes R_{i,s}^*) \psi_A(S_k) (I_{\mathcal{E}} \otimes R_{i,t})(p) \\
 &= (I_{\mathcal{E}} \otimes R_{i,s}^*) A (I_{\mathcal{E}} \otimes R_{i,t})(p) \\
 &= \delta_{st} A(p) = \delta_{st} \psi_A(S_k)(p) \\
 &= \delta_{st} \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} (C_{(\alpha_k; \beta_k)} \otimes S_{k, \alpha_k} S_{k, \beta_k}^*)(p).
 \end{aligned}$$

Hence, we deduce that

$$(I_{\mathcal{E}_{k-2}} \otimes R_{i,s}^*) C_{(\alpha_k; \beta_k)} (I_{\mathcal{E}_{k-2}} \otimes R_{i,t}) = \delta_{st} C_{(\alpha_k; \beta_k)}$$

for any $i \in \{1, \dots, k-1\}$ and $s, t \in \{1, \dots, n_i\}$, which proves that $C_{(\alpha_k; \beta_k)}$ is a 1-multi-Toeplitz operator with respect to $R_{k-1} := (R_{k-1,1}, \dots, R_{k-1, n_{k-1}})$. Consequently, similarly to the first part of the proof, $C_{(\alpha_k; \beta_k)}$ has a Fourier representation

$$\psi_{(\alpha_k; \beta_k)}(S_{k-1}) := \sum_{m_{k-1} \in \mathbb{Z}} \sum_{\substack{\alpha_{k-1}, \beta_{k-1} \in \mathbb{F}_{n_{k-1}}^+ \\ |\alpha_{k-1}| = m_{k-1}^-, |\beta_{k-1}| = m_{k-1}^+}} C_{(\alpha_{k-1}, \alpha_k; \beta_{k-1}, \beta_k)} \otimes S_{k-1, \alpha_{k-1}} S_{k-1, \beta_{k-1}}^*, \tag{1-8}$$

where $C_{(\alpha_{k-1}, \alpha_k; \beta_{k-1}, \beta_k)} \in B(\mathcal{E}_{k-2})$. Moreover, as above, one can prove that, for any $r_{k-1} \in [0, 1)$, the series $\psi_{(\alpha_k; \beta_k)}(r_{k-1} S_{k-1})$ is convergent in the operator norm topology, and its limit is an element in

$$\text{span}\{f^*g : f, g \in B(\mathcal{E}_{k-2}) \otimes_{\min} \mathcal{A}_{n_{k-1}}\}^{-\|\cdot\|},$$

where $\mathcal{A}_{n_{k-1}}$ is the noncommutative disc algebra generated by $S_{k-1,1}, \dots, S_{k-1,n_{k-1}}$ and the identity. We also have

$$\lim_{r_{k-1} \rightarrow 1} \psi_{(\alpha_k; \beta_k)}(r_{k-1} S_{k-1})p = C_{(\alpha_k; \beta_k)}p$$

for any vector-valued polynomial $p \in \mathcal{E}_{k-2} \otimes F^2(H_{n_{k-1}})$. As in the first part of the proof, setting

$$\mathcal{B}_{r_{k-1} S_{k-1}}^{\text{ext}}[u] := (I_{\mathcal{E}_{k-2}} \otimes K_{r_{k-1} S_{k-1}}^*)(u \otimes I_{F^2(H_{n_{k-1}})})(I_{\mathcal{E}_{k-2}} \otimes K_{r_{k-1} S_{k-1}}), \quad u \in B(\mathcal{E}_{n-1}),$$

one can prove that

$$\psi_{(\alpha_k; \beta_k)}(r_{k-1} S_{k-1}) = \mathcal{B}_{r_{k-1} S_{k-1}}^{\text{ext}}[C_{(\alpha_k; \beta_k)}] \quad \text{and} \quad \|\psi_{(\alpha_k; \beta_k)}(r_{k-1} S_{k-1})\| \leq \|C_{(\alpha_k; \beta_k)}\| \quad (1-9)$$

for any $r_{k-1} \in [0, 1)$. Moreover, we can also show that

$$C_{(\alpha_k; \beta_k)} = \text{SOT-} \lim_{r_{k-1} \rightarrow 1} \psi_{(\alpha_k; \beta_k)}(r_{k-1} S_{k-1}).$$

Now, due to (1-6), (1-7), (1-8) and (1-9), we obtain

$$\begin{aligned} & ([\mathcal{B}_{r_{k-1} S_{k-1}}^{\text{ext}} \otimes \text{id}_{B(F^2(H_{n_k}))}] \circ \mathcal{B}_{r_k S_k}^{\text{ext}})[A] \\ &= \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} \mathcal{B}_{r_{k-1} S_{k-1}}^{\text{ext}}[C_{(\alpha_k; \beta_k)}] \otimes r_k^{|\alpha_k| + |\beta_k|} S_{k, \alpha_k} S_{k, \beta_k}^* \\ &= \sum_{m_k \in \mathbb{Z}} \sum_{m_{k-1} \in \mathbb{Z}} \sum_{\substack{\alpha_k, \beta_k \in \mathbb{F}_{n_k}^+ \\ |\alpha_k| = m_k^-, |\beta_k| = m_k^+}} \sum_{\substack{\alpha_{k-1}, \beta_{k-1} \in \mathbb{F}_{n_{k-1}}^+ \\ |\alpha_{k-1}| = m_{k-1}^-, |\beta_{k-1}| = m_{k-1}^+}} r_k^{|\alpha_k|} r_{k-1}^{|\alpha_{k-1}|} C_{(\alpha_{k-1}, \alpha_k; \beta_{k-1}, \beta_k)} \\ & \quad \otimes S_{k-1, \alpha_{k-1}} S_{k-1, \beta_{k-1}}^* \otimes S_{k, \alpha_k} S_{k, \beta_k}^*, \end{aligned}$$

where the series are convergent in the operator norm topology. Continuing this process, one can prove that there are some operators $C_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \in B(\mathcal{E})$ such that the series $\varphi(r_1 S_1, \dots, r_k S_k)$ given by

$$\sum_{m_k \in \mathbb{Z}} \cdots \sum_{m_1 \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} r_k^{|\alpha_k|} \cdots r_1^{|\alpha_1|} C_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes S_{1, \alpha_1} \cdots S_{k, \alpha_k} S_{1, \beta_1}^* \cdots S_{k, \beta_k}^*$$

is convergent in the operator norm topology and

$$\varphi(r_1 S_1, \dots, r_k S_k) = [\mathcal{B}_{r_1 S_1}^{\text{ext}} \otimes \text{id}_{B(\otimes_{i=2}^k F^2(H_{n_i}))}] \circ [\mathcal{B}_{r_2 S_2}^{\text{ext}} \otimes \text{id}_{B(\otimes_{i=3}^k F^2(H_{n_i}))}] \circ \cdots \circ \mathcal{B}_{r_k S_k}^{\text{ext}}[A]. \quad (1-10)$$

Since the noncommutative Berezin kernels $K_{r_i S_i}, i \in \{1, \dots, k\}$, are isometries, we deduce that

$$\|\varphi(r_1 S_1, \dots, r_k S_k)\| \leq \|A\|, \quad r_i \in [0, 1).$$

Note that the coefficients of the k -multi-Toeplitz operator $\varphi(r_1 \mathbf{S}_1, \dots, r_k \mathbf{S}_k)$ satisfy the relation

$$\langle r_k^{|m_k|} \dots r_1^{|m_1|} C_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, \ell \rangle = \langle \varphi(r_1 \mathbf{S}_1, \dots, r_k \mathbf{S}_k)(h \otimes x), (\ell \otimes y) \rangle, \tag{1-11}$$

where x, y are defined as in (1-3). Since A is a k -multi-Toeplitz operator, so is $Y_{r_k} := \mathcal{B}_{r_k \mathbf{S}_k}^{\text{ext}}[A] = \psi_A(r_k \mathbf{S}_k)$ and, iterating the argument, we deduce that

$$Y_{r_2, \dots, r_k} := [\mathcal{B}_{r_2 \mathbf{S}_2}^{\text{ext}} \otimes \text{id}_{B(\otimes_{i=3}^k F^2(H_{n_i}))}] \circ \dots \circ \mathcal{B}_{r_k \mathbf{S}_k}^{\text{ext}}[A]$$

is a k -multi-Toeplitz operator. In particular, Y_{r_2, \dots, r_k} is a 1-multi-Toeplitz operator with respect to $R_1 := (R_{1,1}, \dots, R_{1,n_1})$. Applying the first part of the proof to Y_{r_2, \dots, r_k} , we deduce that

$$\text{SOT-}\lim_{r_1 \rightarrow 1} [\mathcal{B}_{r_1 \mathbf{S}_1}^{\text{ext}} \otimes \text{id}_{B(\otimes_{i=2}^k F^2(H_{n_i}))}] [Y_{r_2, \dots, r_k}] = Y_{r_2, \dots, r_k}.$$

Continuing this process, we obtain

$$\text{SOT-}\lim_{r_k \rightarrow 1} \dots \text{SOT-}\lim_{r_1 \rightarrow 1} [\mathcal{B}_{r_1 \mathbf{S}_1}^{\text{ext}} \otimes \text{id}_{B(\otimes_{i=2}^k F^2(H_{n_i}))}] [Y_{r_2, \dots, r_k}] = A.$$

Consequently, using (1-10), (1-11) and (1-2), we deduce that

$$\langle C_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, \ell \rangle = \langle A(h \otimes x), \ell \otimes y \rangle = \langle A_{(\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k)} h, \ell \rangle,$$

which shows that $\varphi_A(r_1 \mathbf{S}_1, \dots, r_k \mathbf{S}_k) = \varphi(r_1 \mathbf{S}_1, \dots, r_k \mathbf{S}_k)$ for any $r_i \in [0, 1)$. Hence, we obtain

$$\varphi_A(r_1 \mathbf{S}_1, \dots, r_k \mathbf{S}_k) = \sum_{m_k \in \mathbb{Z}} \dots \sum_{m_1 \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} r_k^{|m_k|} \dots r_1^{|m_1|} \times A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \mathbf{S}_{1, \alpha_1} \dots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \dots \mathbf{S}_{k, \beta_k}^*,$$

where the series are convergent in the operator norm topology. Moreover, due to (1-10), we have

$$\|\varphi_A(r_1 \mathbf{S}_1, \dots, r_k \mathbf{S}_k)\| \leq \|A\|, \quad r_i \in [0, 1).$$

Due to (1-5), we have

$$\lim_{r \rightarrow 1} \varphi_A(r \mathbf{S})(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k) = A(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k).$$

Since $\|\varphi_A(r \mathbf{S}_1, \dots, r \mathbf{S}_k)\| \leq \|A\|$, an approximation argument shows that

$$\text{SOT-}\lim_{r \rightarrow 1} \varphi_A(r \mathbf{S}_1, \dots, r \mathbf{S}_k) = A. \tag{1-12}$$

Let $\epsilon > 0$ and choose a vector-valued polynomial $q \in \mathcal{P}$ with $\|q\| = 1$ and $\|Aq\| > \|A\| - \epsilon$. Due to (1-12), there is $r_0 \in (0, 1)$ such that $\|\varphi_A(r_0 \mathbf{S}_1, \dots, r_0 \mathbf{S}_k)q\| > \|A\| - \epsilon$. Hence, we deduce that $\sup_{r \in [0, 1)} \|\varphi_A(r \mathbf{S}_1, \dots, r \mathbf{S}_k)\| = \|A\|$.

Now, let $r_1, r_2 \in [0, 1)$ with $r_1 < r_2$. We already proved that $g(\mathbf{S}) := \varphi_A(r_2 \mathbf{S}_1, \dots, r_2 \mathbf{S}_k)$ is in $\text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{\|\cdot\|}$. Due to the von Neumann-type inequality [1951] from [Popescu 2016], we have $\|g(r \mathbf{S})\| \leq \|g(\mathbf{S})\|$ for any $r \in [0, 1)$. In particular, setting $r = r_1/r_2$, we deduce that

$$\|\varphi_A(r_1 \mathbf{S}_1, \dots, r_1 \mathbf{S}_k)\| \leq \|\varphi_A(r_2 \mathbf{S}_1, \dots, r_2 \mathbf{S}_k)\|.$$

It is clear that $\lim_{r \rightarrow 1} \|\varphi_A(rS_1, \dots, rS_k)\| = \|A\|$. On the other hand, since $Aq = \varphi_A(\mathbf{S})q$ for any vector-valued polynomial $q \in \mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$, we deduce that $\|A\| = \sup_{q \in \mathcal{P}, \|q\| \leq 1} \|\varphi(\mathbf{S})q\|$.

Now we prove the converse of the theorem. Let $\{A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)}\}$ be a family of operators in $B(\mathcal{E})$, where $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, |\alpha_i| = m_i^-, |\beta_i| = m_i^+, m_i \in \mathbb{Z}$ and $i \in \{1, \dots, k\}$, and assume that conditions (i) and (ii) hold. Note that, due to (i), $\varphi(\mathbf{S})p$ and $\varphi(r\mathbf{S})p, r \in [0, 1)$, are vectors in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ and

$$\lim_{r \rightarrow 1} \varphi(r\mathbf{S})p = \varphi(\mathbf{S})p$$

for any $p \in \mathcal{P}$. Since $\sup_{p \in \mathcal{P}, \|p\| \leq 1} \|\varphi(r\mathbf{S})p\| < \infty$, there exists a unique bounded linear operator $A_r \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ such that $A_r p = \varphi(r\mathbf{S})p$ for any $p \in \mathcal{P}$. If $f \in \mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ and $\{p_m\}$ is a sequence of polynomials $p_m \in \mathcal{P}$ such that $p_m \rightarrow f$ as $m \rightarrow \infty$, we set $A_r(f) := \lim_{m \rightarrow \infty} \varphi(r\mathbf{S})p_m$. Note that the definition is valid. On the other hand, note that

$$\sup_{p \in \mathcal{P}, \|p\| \leq 1} \|\varphi(\mathbf{S})p\| < \infty.$$

Indeed, this follows from the facts that $\lim_{r \rightarrow 1} \varphi(r\mathbf{S})p = \varphi(\mathbf{S})p$ and $\sup_{p \in \mathcal{P}, \|p\| \leq 1} \|\varphi(r\mathbf{S})p\| < \infty$. Consequently, there is a unique operator $A \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ such that $Ap = \varphi(\mathbf{S})p$ for any $p \in \mathcal{P}$. Since $\lim_{r \rightarrow 1} A_r p = \lim_{r \rightarrow 1} \varphi(r\mathbf{S})p = \varphi(\mathbf{S})p = Ap$ and $\sup_{r \in [0, 1)} \|A_r\| < \infty$, we deduce that $A = \text{SOT-}\lim_{r \rightarrow 1} A_r$.

Now we show that A is a k -multi-Toeplitz operator. First, note that $S_{1, \alpha_1} \cdots S_{k, \alpha_k} S_{1, \beta_1}^* \cdots S_{k, \beta_k}^*$ is a k -multi-Toeplitz operator for any $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\}$ with $m_i \in \mathbb{Z}, |\alpha_i| = m_i^-$ and $|\beta_i| = m_i^+$. It is enough to check this on monomials of the form $h \otimes e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k$. Consequently,

$$(I_{\mathcal{E}} \otimes R_{i,s}^*) \varphi(r\mathbf{S}) (I_{\mathcal{E}} \otimes R_{i,t}) p = \delta_{st} \varphi(r\mathbf{S}) p, \quad s, t \in \{1, \dots, n_i\},$$

for any $p \in \mathcal{P}$ and every $i \in \{1, \dots, k\}$. Hence, A_r has the same property. Taking $r \rightarrow 1$, we conclude that A is a k -multi-Toeplitz operator. On the other hand, if $x := x_1 \otimes \cdots \otimes x_k, y = y_1 \otimes \cdots \otimes y_k$ satisfy (1-3) and $h, \ell \in \mathcal{E}$, we have

$$\begin{aligned} \langle A(h \otimes x), \ell \otimes y \rangle &= \lim_{r \rightarrow 1} \langle A_r(h \otimes x), \ell \otimes y \rangle \\ &= \lim_{r \rightarrow 1} \langle \varphi(r\mathbf{S})(h \otimes x), \ell \otimes y \rangle \\ &= \lim_{r \rightarrow 1} \langle r^{\sum_{i=1}^k |\alpha_i| + |\beta_i|} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, \ell \rangle \\ &= \langle A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, \ell \rangle. \end{aligned}$$

Therefore,

$$\varphi(\mathbf{S}) := \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes S_{1, \alpha_1} \cdots S_{k, \alpha_k} S_{1, \beta_1}^* \cdots S_{k, \beta_k}^*$$

is the formal Fourier series of the k -multi-Toeplitz operator A on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. □

Theorem 1.6. *Let $\{A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)}\}$ be a family of operators in $B(\mathcal{E})$, where $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, |\alpha_i| = m_i^-, |\beta_i| = m_i^+, m_i \in \mathbb{Z}$ and $i \in \{1, \dots, k\}$, and let*

$$\varphi(\mathcal{S}) := \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes S_{1, \alpha_1} \cdots S_{k, \alpha_k} S_{1, \beta_1}^* \cdots S_{k, \beta_k}^*$$

be the associated formal Fourier series. Then $\varphi(\mathcal{S})$ is the formal Fourier series of a k -multi-Toeplitz operator A on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$ if and only if the series defining $\varphi(r\mathcal{S})$ is convergent in the operator norm topology for any $r \in [0, 1)$ and

$$\sup_{r \in [0, 1)} \|\varphi(r\mathcal{S})\| < \infty.$$

Moreover, if A is a k -multi-Toeplitz operator on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$, then $\varphi(r\mathcal{S}) = \mathcal{B}_{r\mathcal{S}}^{\text{ext}}[A]$ and

$$\text{SOT-}\lim_{r \rightarrow 1} \mathcal{B}_{r\mathcal{S}}^{\text{ext}}[A] = A, \quad \text{where } \mathcal{B}_{r\mathcal{S}}^{\text{ext}}[u] := (I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathcal{S}}^*)(u \otimes I_{\bigotimes_{i=1}^k F^2(H_{n_i})})(I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathcal{S}}), \quad u \in B(\mathcal{E}_k),$$

and $\mathbf{K}_{r\mathcal{S}}$ is the noncommutative Berezin kernel associated with $r\mathcal{S} \in \mathbf{B}_n(\bigotimes_{i=1}^k F^2(H_{n_i}))$.

Proof. Assume that $\varphi(\mathcal{S})$ is the formal Fourier series of a k -multi-Toeplitz operator A on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. Then [Theorem 1.5](#) implies that $\varphi(r\mathcal{S})$ is convergent in the operator norm topology and

$$\|A\| = \sup_{r \in [0, 1)} \|\varphi(r\mathcal{S})\|.$$

We recall that the noncommutative Berezin kernel associated with $r\mathcal{S} \in \mathbf{B}_n(\bigotimes_{i=1}^k F^2(H_{n_i}))$ is defined on $\bigotimes_{i=1}^k F^2(H_{n_i})$ with values in $\bigotimes_{i=1}^k F^2(H_{n_i}) \otimes \mathcal{D}_{r\mathcal{S}} \subset (\bigotimes_{i=1}^k F^2(H_{n_i})) \otimes (\bigotimes_{i=1}^k F^2(H_{n_i}))$, where

$$\mathcal{D}_{r\mathcal{S}} := \overline{\Delta_{r\mathcal{S}}(I) \left(\bigotimes_{i=1}^k F^2(H_{n_i}) \right)}.$$

Let $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k), \boldsymbol{\omega} = (\omega_1, \dots, \omega_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, set $q := \max\{|\gamma_1|, \dots, |\gamma_k|, |\omega_1|, \dots, |\omega_k|\}$, and define the operator

$$\Gamma_q := \sum_{m_1 \in \mathbb{Z}, |m_1| \leq q} \cdots \sum_{m_k \in \mathbb{Z}, |m_k| \leq q} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \mathbf{S}_{\boldsymbol{\alpha}} \mathbf{S}_{\boldsymbol{\beta}}^*,$$

where we use the notation $\mathbf{S}_{\boldsymbol{\alpha}} := S_{1, \alpha_1} \cdots S_{k, \alpha_k}$ if $\boldsymbol{\alpha} := (\alpha_1, \dots, \alpha_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$. We also set $e_{\boldsymbol{\alpha}} := e_{\alpha_1}^1 \otimes \cdots \otimes e_{\alpha_k}^k$. Note that

$$\begin{aligned} & \langle \mathcal{B}_{r\mathcal{S}}^{\text{ext}}[A](h \otimes e_{\boldsymbol{\gamma}}), h' \otimes e_{\boldsymbol{\omega}} \rangle \\ &= \langle (I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathcal{S}}^*)(A \otimes I_{\bigotimes_{i=1}^k F^2(H_{n_i})})(I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathcal{S}})(h \otimes e_{\boldsymbol{\gamma}}), h' \otimes e_{\boldsymbol{\omega}} \rangle \\ &= \left\langle (A \otimes I_{\bigotimes_{i=1}^k F^2(H_{n_i})}) \sum_{\boldsymbol{\alpha} \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+} h \otimes e_{\boldsymbol{\alpha}} \otimes \Delta_{r\mathcal{S}}(I)^{1/2} \mathbf{S}_{\boldsymbol{\alpha}}^*(e_{\boldsymbol{\gamma}}), \sum_{\boldsymbol{\beta} \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+} h' \otimes e_{\boldsymbol{\beta}} \otimes \Delta_{r\mathcal{S}}(I)^{1/2} \mathbf{S}_{\boldsymbol{\beta}}^*(e_{\boldsymbol{\omega}}) \right\rangle \end{aligned}$$

$$\begin{aligned}
 &= \sum_{\alpha \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+} \sum_{\beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+} \langle A(h \otimes e_\alpha) \otimes \Delta_{rS}(I)^{1/2} S_\alpha^*(e_\gamma), h' \otimes e_\beta \otimes \Delta_{rS}(I)^{1/2} S_\beta^*(e_\omega) \rangle \\
 &= \sum_{\alpha \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+} \sum_{\beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+} \langle A(h \otimes e_\alpha), h' \otimes e_\beta \rangle \langle \Delta_{rS}(I)^{1/2} S_\alpha^*(e_\gamma), \Delta_{rS}(I)^{1/2} S_\beta^*(e_\omega) \rangle \\
 &= \sum_{m_1 \in \mathbb{Z}, |m_1| \leq q} \dots \sum_{m_k \in \mathbb{Z}, |m_k| \leq q} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_k}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \langle \Gamma_q(h \otimes e_\alpha), h' \otimes e_\beta \rangle \langle \Delta_{rS}(I)^{1/2} S_\alpha^*(e_\gamma), \Delta_{rS}(I)^{1/2} S_\beta^*(e_\omega) \rangle \\
 &= \sum_{\alpha \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+} \sum_{\beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+} \langle \Gamma_q(h \otimes e_\alpha), h' \otimes e_\beta \rangle \langle \Delta_{rS}(I)^{1/2} S_\alpha^*(e_\gamma), \Delta_{rS}(I)^{1/2} S_\beta^*(e_\omega) \rangle \\
 &= ((I_{\mathcal{E}} \otimes K_{rS})(\Gamma_q \otimes I_{\bigotimes_{i=1}^k F^2(H_{n_i})})(I_{\mathcal{E}} \otimes K_{rS})(h \otimes e_\gamma), h' \otimes e_\omega) \\
 &= (\mathcal{B}_{rS}^{\text{ext}}[\Gamma_q](h \otimes e_\gamma), h' \otimes e_\omega) \\
 &= \sum_{m_1 \in \mathbb{Z}, |m_1| \leq q} \dots \sum_{m_k \in \mathbb{Z}, |m_k| \leq q} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_k}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \langle (A_{(\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k)} \otimes r^{\sum_{i=1}^k (|\alpha_i| + |\beta_i|)} S_\alpha S_\beta^*)(h \otimes e_\gamma), h' \otimes e_\omega \rangle \\
 &= \langle \varphi_A(rS_1, \dots, rS_k)(h \otimes e_\gamma), h' \otimes e_\omega \rangle.
 \end{aligned}$$

Consequently, we obtain

$$\mathcal{B}_{rS}^{\text{ext}}[A] = \varphi_A(rS_1, \dots, rS_k), \quad r \in [0, 1),$$

which proves the second part of the theorem.

To prove the converse, assume that $\{A_{(\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k)}\}$ is a family of operators in $B(\mathcal{E})$, where $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, |\alpha_i| = m_i^-, |\beta_i| = m_i^+, m_i \in \mathbb{Z}$ and $i \in \{1, \dots, k\}$, and let $\varphi(S)$ be the associated formal Fourier series. We also assume that $\varphi(rS)$ is convergent in the operator norm topology for each $r \in [0, 1)$ and that

$$M := \sup_{r \in [0, 1)} \|\varphi(rS)\| < \infty.$$

Note that $\varphi(rS)$ is a k -multi-Toeplitz operator and

$$\varphi(rS)(h \otimes e_{\gamma_1}^1 \otimes \dots \otimes e_{\gamma_k}^k) = \sum_{\substack{\omega = (\omega_1, \dots, \omega_k) \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+ \\ \omega \sim_{rc} \boldsymbol{\gamma}}} r^{\sum_{i=1}^k (|c_r^+(\omega, \boldsymbol{\gamma})| + |c_r^-(\omega, \boldsymbol{\gamma})|)} A_{(c_r^+(\omega, \boldsymbol{\gamma}); c_r^-(\omega, \boldsymbol{\gamma}))} h \otimes e_{\omega_1}^1 \otimes \dots \otimes e_{\omega_k}^k$$

is a vector in $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. Hence, we deduce that, for each $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_k) \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$,

$$\left\langle r^{\sum_{i=1}^k (c_r^+(\omega, \boldsymbol{\gamma}) + c_r^-(\omega, \boldsymbol{\gamma}))} \sum_{\substack{\omega \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+ \\ \omega \sim_{rc} \boldsymbol{\gamma}}} A_{(c_r^+(\omega, \boldsymbol{\gamma}); c_r^-(\omega, \boldsymbol{\gamma}))}^* A_{(c_r^+(\omega, \boldsymbol{\gamma}); c_r^-(\omega, \boldsymbol{\gamma}))} h, h \right\rangle \leq \|\varphi(rS)\|^2 \|h\|^2 \leq M \|h\|^2$$

for any $r \in [0, 1)$ and $h \in \mathcal{E}$. Taking $r \rightarrow 1$, we get condition (i) of [Theorem 1.5](#). Applying [Theorem 1.5](#), we deduce that $\varphi(S)$ is the Fourier series of a k -multi-Toeplitz operator. □

We remark that, due to [Theorem 1.6](#), the order of the series in the definition of $\varphi_A(r\mathbf{S}_1, \dots, r\mathbf{S}_k)$ (see [Theorem 1.5\(a\)](#)) is irrelevant.

Theorem 1.7. *Let $\mathbf{n} = (n_1, \dots, n_k) \in \mathbb{N}^k$ and let $\mathcal{T}_{\mathbf{n}}$ be the set of all k -multi-Toeplitz operators on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. Then*

$$\mathcal{T}_{\mathbf{n}} = \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_{\mathbf{n}}\}^{-\text{SOT}} = \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_{\mathbf{n}}\}^{-\text{WOT}},$$

where $\mathcal{A}_{\mathbf{n}}$ is the polyball algebra.

Proof. Let

$$\mathcal{G} := \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_{\mathbf{n}}\}^{\|\cdot\|}.$$

According to [Theorem 1.5](#), if $A \in \mathcal{T}_{\mathbf{n}}$ and $\varphi_A(\mathbf{S})$ is its Fourier series, then $\varphi_A(r\mathbf{S}) \in \mathcal{G}$ for any $r \in [0, 1)$ and $A = \text{SOT-lim } \varphi_A(r\mathbf{S})$. Consequently, $\mathcal{T}_{\mathbf{n}} \subseteq \bar{\mathcal{G}}^{\text{SOT}}$. Conversely, note that each monomial $\mathbf{S}_{\alpha}^* \mathbf{S}_{\beta}$, $\alpha, \beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, is a k -multi-Toeplitz operator. This shows that, for each $Y \in \mathcal{G}$,

$$(I_{\mathcal{E}} \otimes \mathbf{R}_{i,s}^*)Y(I_{\mathcal{E}} \otimes \mathbf{R}_{i,t}) = \delta_{st}Y, \quad s, t \in \{1, \dots, n_i\},$$

for every $i \in \{1, \dots, k\}$. Consequently, taking SOT-limits, we deduce that $\bar{\mathcal{G}}^{\text{SOT}} \subseteq \mathcal{T}_{\mathbf{n}}$, which proves that $\bar{\mathcal{G}}^{\text{SOT}} = \mathcal{T}_{\mathbf{n}}$.

Now, if $T \in \bar{\mathcal{G}}^{\text{WOT}}$, an argument as above shows that $T \in \mathcal{T}_{\mathbf{n}} = \bar{\mathcal{G}}^{\text{SOT}}$. Since $\bar{\mathcal{G}}^{\text{SOT}} \subseteq \bar{\mathcal{G}}^{\text{WOT}}$, we conclude that $\mathcal{T}_{\mathbf{n}} = \bar{\mathcal{G}}^{\text{SOT}} = \bar{\mathcal{G}}^{\text{WOT}}$. □

Corollary 1.8. *The set of all k -multi-Toeplitz operators on $\bigotimes_{i=1}^k F^2(H_{n_i})$ coincides with*

$$\text{span}\{\mathcal{A}_{\mathbf{n}}^* \mathcal{A}_{\mathbf{n}}\}^{-\text{SOT}} = \text{span}\{\mathcal{A}_{\mathbf{n}}^* \mathcal{A}_{\mathbf{n}}\}^{-\text{WOT}},$$

where $\mathcal{A}_{\mathbf{n}}$ is the polyball algebra.

2. Bounded free k -pluriharmonic functions and the Dirichlet extension problem

In this section, we show that the bounded free k -pluriharmonic functions on $\mathbf{B}_{\mathbf{n}}$ are precisely the noncommutative Berezin transforms of k -multi-Toeplitz operators and solve the Dirichlet extension problem for the regular polyball $\mathbf{B}_{\mathbf{n}}$.

Definition 2.1. A function F with operator-valued coefficients in $B(\mathcal{E})$ is called free k -pluriharmonic on the polyball $\mathbf{B}_{\mathbf{n}}$ if it has the form

$$F(X) = \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes X_{1, \alpha_1} \dots X_{k, \alpha_k} X_{1, \beta_1}^* \dots X_{k, \beta_k}^*,$$

where the series converge in the operator norm topology for any $X = (X_1, \dots, X_k) \in \mathbf{B}_{\mathbf{n}}(\mathcal{H})$, with $X_i := (X_{i,1}, \dots, X_{i,n_i})$, and any Hilbert space \mathcal{H} .

Due to the remark following [Theorem 1.6](#), one can prove that the order of the series in the definition above is irrelevant. Note that any free holomorphic function on \mathbf{B}_n is k -pluriharmonic. Indeed, according to [\[Popescu 2015b\]](#), any free holomorphic function on the polyball \mathbf{B}_n has the form

$$f(\mathbf{X}) = \sum_{m_1 \in \mathbb{N}} \cdots \sum_{m_k \in \mathbb{N}} \sum_{\substack{\alpha_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i}} A_{(\alpha_1, \dots, \alpha_k)} \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k}, \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}),$$

where the series converge in the operator norm topology. A function $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E} \otimes \mathcal{H})$ is called bounded if

$$\|F\| := \sup_{\mathbf{X} \in \mathbf{B}_n(\mathcal{H})} \|F(\mathbf{X})\| < \infty.$$

A free k -pluriharmonic function is bounded if its representation on any Hilbert space is bounded. Denote by $\mathbf{PH}_{\mathcal{E}}^{\infty}(\mathbf{B}_n)$ the set of all bounded free k -pluriharmonic functions on the polyball \mathbf{B}_n with coefficients in $B(\mathcal{E})$. For each $m = 1, 2, \dots$, we define the norms $\|\cdot\|_m : M_m(\mathbf{PH}_{\mathcal{E}}^{\infty}(\mathbf{B}_n)) \rightarrow [0, \infty)$ by setting

$$\|[F_{ij}]_m\|_m := \sup \|[F_{ij}(\mathbf{X})]_m\|,$$

where the supremum is taken over all n -tuples $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$ and any Hilbert space \mathcal{H} . It is easy to see that the norms $\|\cdot\|_m, m = 1, 2, \dots$, determine an operator space structure on $\mathbf{PH}_{\mathcal{E}}^{\infty}(\mathbf{B}_n)$, in the sense of Ruan (see, e.g., [\[Effros and Ruan 2000\]](#)).

Let \mathcal{T}_n be the set of all k -multi-Toeplitz operators on $\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})$. According to [Theorem 1.7](#), we have

$$\mathcal{T}_n = \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{\text{SOT}},$$

where \mathcal{A}_n is the polyball algebra. The main result of this section is the following characterization of bounded free k -pluriharmonic functions:

Theorem 2.2. *If $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$, then the following statements are equivalent:*

- (i) F is a bounded free k -pluriharmonic function;
- (ii) there exists $A \in \mathcal{T}_n$ such that

$$F(\mathbf{X}) = \mathcal{B}_X^{\text{ext}}[A] := (I_{\mathcal{E}} \otimes \mathbf{K}_X^*)(A \otimes I_{\mathcal{H}})(I_{\mathcal{E}} \otimes \mathbf{K}_X), \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

In this case, $A = \text{SOT-}\lim_{r \rightarrow 1} F(r\mathbf{S})$. Moreover, the map

$$\Phi : \mathbf{PH}_{\mathcal{E}}^{\infty}(\mathbf{B}_n) \rightarrow \mathcal{T}_n, \quad \Phi(F) := A,$$

is a completely isometric isomorphism of operator spaces.

Proof. Assume that F is a bounded free k -pluriharmonic function on \mathbf{B}_n and has the representation from [Definition 2.1](#). Then, for any $r \in [0, 1)$,

$$F(r\mathbf{S}) \in \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{\|\cdot\|}$$

and, due to the noncommutative von Neumann inequality [\[Popescu 1999\]](#), we have $\sup_{r \in [0, 1)} \|F(r\mathbf{S})\| = \|F\|_{\infty} < \infty$. According to [Theorem 1.6](#), $F(\mathbf{S})$ is the formal Fourier series of a k -multi-Toeplitz operator

$A \in B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$ and $A = \text{SOT-lim}_{r \rightarrow 1} F(r\mathbf{S}) \in \mathcal{T}_n$. Using the properties of the noncommutative Berezin kernel on polyballs, we have

$$F(r\mathbf{X}) = (I_{\mathcal{E}} \otimes \mathbf{K}_X^*)[F(r\mathbf{S}) \otimes I_{\mathcal{H}}](I_{\mathcal{E}} \otimes \mathbf{K}_X), \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

Since the map $Y \mapsto Y \otimes I_{\mathcal{H}}$ is SOT-continuous on bounded subsets of $B(\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i}))$, we deduce that

$$\text{SOT-lim}_{r \rightarrow 1} F(r\mathbf{X}) = (I_{\mathcal{E}} \otimes \mathbf{K}_X^*)[A \otimes I_{\mathcal{H}}](I_{\mathcal{E}} \otimes \mathbf{K}_X) = \mathcal{B}_X^{\text{ext}}[A].$$

Since F is continuous in the norm topology on $\mathbf{B}_n(\mathcal{H})$, we have $F(r\mathbf{X}) \rightarrow F(\mathbf{X})$ as $r \rightarrow 1$. Consequently, the relation above implies $F(\mathbf{X}) = \mathcal{B}_X^{\text{ext}}[A]$, which completes the proof that (i) implies (ii).

To prove that (ii) implies (i), let $A \in \mathcal{T}_n$ and $F(\mathbf{X}) := \mathcal{B}_X^{\text{ext}}[A]$ for $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$. Since A is a k -multi-Toeplitz operator, [Theorem 1.5](#) shows that it has a formal Fourier series

$$\varphi(\mathbf{S}) := \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \mathbf{S}_{1, \alpha_1} \cdots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \cdots \mathbf{S}_{k, \beta_k}^*$$

with the property that the series $\varphi(r\mathbf{S})$ is convergent in the operator norm topology to an operator in $\text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{-\|\cdot\|}$. Moreover, we have $A = \text{SOT-lim}_{r \rightarrow 1} \varphi(r\mathbf{S})$ and

$$\|A\| = \sup_{r \in [0, 1)} \|\varphi(r\mathbf{S})\|.$$

Hence, the map $\mathbf{X} \mapsto \varphi(\mathbf{X})$ is a k -pluriharmonic function on $\mathbf{B}_n(\mathcal{H})$. On the other hand, due to [Theorem 1.6](#), we have $\varphi(r\mathbf{S}) = \mathcal{B}_{r\mathbf{S}}^{\text{ext}}[A]$, where

$$\mathcal{B}_{r\mathbf{S}}^{\text{ext}}[u] := (I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathbf{S}}^*)(u \otimes I_{\bigotimes_{i=1}^k F^2(H_{n_i})})(I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathbf{S}}), \quad u \in B(\mathcal{E}),$$

and $\mathbf{K}_{r\mathbf{S}}$ is the noncommutative Berezin kernel associated with $r\mathbf{S} \in \mathbf{B}_n(\bigotimes_{i=1}^k F^2(H_{n_i}))$. Note that

$$\varphi(r\mathbf{X}) = \mathcal{B}_X^{\text{ext}}[\varphi(r\mathbf{S})] = (I_{\mathcal{E}} \otimes \mathbf{K}_X^*)[\varphi(r\mathbf{S}) \otimes I_{\mathcal{H}}](I_{\mathcal{E}} \otimes \mathbf{K}_X).$$

Now, using continuity of φ on $\mathbf{B}_n(\mathcal{H})$ and the fact that $A = \text{SOT-lim}_{r \rightarrow 1} \varphi(r\mathbf{S})$, we deduce that

$$\varphi(\mathbf{X}) = \text{SOT-lim}_{r \rightarrow 1} \varphi(r\mathbf{X}) = \mathcal{B}_X^{\text{ext}}[A] = F(\mathbf{X}), \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

To prove the last part of the theorem, let $[F_{ij}]_m \in M_m(\mathbf{PH}_{\infty}^{\infty}(\mathbf{B}_n))$ and use the noncommutative von Neumann inequality to obtain

$$\|[F_{ij}]_m\| = \sup_{\mathbf{X} \in \mathbf{B}_n(\mathcal{H})} \|[F_{ij}(\mathbf{X})]_m\| = \sup_{r \in [0, 1)} \|[F_{ij}(r\mathbf{S})]_m\|.$$

On the other hand, $A_{ij} := \text{SOT-lim}_{r \rightarrow 1} F_{ij}(r\mathbf{S})$ is a k -multi-Toeplitz operator and

$$F_{ij}(r\mathbf{S}) = (I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathbf{S}}^*)(A_{ij} \otimes I_{\bigotimes_{i=1}^k F^2(H_{n_i})})(I_{\mathcal{E}} \otimes \mathbf{K}_{r\mathbf{S}}).$$

Hence, we obtain

$$\sup_{r \in [0, 1)} \|[F_{ij}(r\mathbf{S})]_m\| \leq \|[A_{ij}]_m\|.$$

Since $[A_{ij}]_m := \text{SOT-}\lim_{r \rightarrow 1} [F_{ij}(rS)]_m$, we deduce that the inequality above is in fact an equality. This shows that Φ is a completely isometric isomorphisms of operator spaces. \square

As a consequence, we can obtain the following Fatou-type result concerning the boundary behaviour of bounded k -pluriharmonic functions.

Corollary 2.3. *If $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ is a bounded free k -pluriharmonic function and X is a pure element in $\mathbf{B}_n(\mathcal{H})^-$, then the limit*

$$\text{SOT-}\lim_{r \rightarrow 1} F(rX)$$

exists.

Proof. If X is a pure element in $\mathbf{B}_n(\mathcal{H})^-$, then the noncommutative Berezin kernel \mathbf{K}_X is an isometry (see [Popescu 2016]). Since F is free k -pluriharmonic function on \mathbf{B}_n , we have

$$F(rS) \in \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{-\|\cdot\|}$$

and $F(rS)$ converges in the operator norm topology. Consequently,

$$F(rX) = (I_{\mathcal{E}} \otimes \mathbf{K}_X^*)[F(rS) \otimes I_{\mathcal{H}}](I_{\mathcal{E}} \otimes \mathbf{K}_X).$$

Since F is bounded, Theorem 2.2 implies $\text{SOT-}\lim_{r \rightarrow 1} F(rS) = A \in \mathcal{T}_n$ and $\sup_{0 \leq r < 1} \|F(rS)\| < \infty$. Using these facts in the relation above, we conclude that $\text{SOT-}\lim_{r \rightarrow 1} F(rX)$ exists. \square

We denote by $\mathbf{PH}_{\mathcal{E}}^c(\mathbf{B}_n)$ the set of all free k -pluriharmonic functions on \mathbf{B}_n with operator-valued coefficients in $B(\mathcal{E})$, which have continuous extensions (in the operator norm topology) to the closed polyball $\mathbf{B}_n(\mathcal{H})^-$ for any Hilbert space \mathcal{H} . Throughout this section, we assume that \mathcal{H} is an infinite-dimensional Hilbert space. In what follows we solve the Dirichlet extension problem for the regular polyballs.

Theorem 2.4. *If $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$, then the following statements are equivalent:*

- (i) *F is a free k -pluriharmonic function on $\mathbf{B}_n(\mathcal{H})$ such that $F(rS)$ converges in the operator norm topology as $r \rightarrow 1$.*
- (ii) *There exists $A \in \mathcal{P} := \text{span}\{f^*g : f, g \in B(\mathcal{E}) \otimes_{\min} \mathcal{A}_n\}^{-\|\cdot\|}$ such that*

$$F(X) = \mathcal{B}_X^{\text{ext}}[A], \quad X \in \mathbf{B}_n(\mathcal{H}).$$

- (iii) *F is a free k -pluriharmonic function on $\mathbf{B}_n(\mathcal{H})$ which has a continuous extension (in the operator norm topology) to the closed ball $\mathbf{B}_n(\mathcal{H})^-$.*

In this case, $A = \lim_{r \rightarrow 1} F(rS)$, where the convergence is in the operator norm. Moreover, the map

$$\Phi : \mathbf{PH}_{\mathcal{E}}^c(\mathbf{B}_n) \rightarrow \mathcal{P}, \quad \Phi(F) := A,$$

is a completely isometric isomorphism of operator spaces.

Proof. Assume that (i) holds. Then F has a representation

$$F(X) = \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*,$$

where the series converge in the operator norm topology for any $X = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$. Since the series defining $F(rS)$ converges in the operator topology, we deduce that

$$A := \lim_{r \rightarrow 1} F(rS) \in \mathcal{P}. \tag{2-1}$$

On the other hand, we have

$$\mathcal{B}_X^{\text{ext}}[F(rS)] = (I_{\mathcal{E}} \otimes \mathbf{K}_X^*)[F(rS) \otimes I_{\mathcal{H}}](I_{\mathcal{E}} \otimes \mathbf{K}_X) = F(rX)$$

for any $r \in [0, 1)$ and $X \in \mathbf{B}_n(\mathcal{H})$. Hence, and using (2-1), we deduce that

$$\mathcal{B}_X^{\text{ext}}[A] = \lim_{r \rightarrow 1} F(rX) = F(X),$$

which proves (ii). Now we show that (ii) implies (i). Assuming (ii) and taking into account [Theorem 1.7](#), one can see that A is a k -multi-Toeplitz operator. As in the proof of [Theorem 2.2](#), the map defined by $F(X) := \mathcal{B}_X^{\text{ext}}[A]$, $X \in \mathbf{B}_n(\mathcal{H})$, is a bounded free k -pluriharmonic function. Moreover, we proved that

$$F(rS) = \mathcal{B}_{rS}^{\text{ext}}[A], \quad r \in [0, 1), \tag{2-2}$$

$F(rS) \in \mathcal{P}$ and also that $A = \text{SOT-}\lim_{r \rightarrow 1} F(rS)$ and $\|A\| = \sup_{r \in [0, 1)} \|F(rS)\|$. Since $A \in \mathcal{P}$, there is a sequence of polynomials q_m in $\mathbf{S}_{\alpha}^* \mathbf{S}_{\beta}$ such that $q_m \rightarrow A$ in norm as $m \rightarrow \infty$. For any $\epsilon > 0$, let $N \in \mathbb{N}$ be such that $\|A - q_m\| < \frac{1}{3}\epsilon$ for any $m \geq N$. Choose $\delta \in (0, 1)$ such that $\|\mathcal{B}_{rS}^{\text{ext}}[q_N] - q_N\| < \frac{1}{3}\epsilon$ for any $r \in (\delta, 1)$. Note that

$$\|\mathcal{B}_{rS}^{\text{ext}}[A] - A\| \leq \|\mathcal{B}_{rS}^{\text{ext}}[A - q_N]\| + \|\mathcal{B}_{rS}^{\text{ext}}[q_N] - q_N\| + \|q_N - A\| \leq \|A - q_N\| + \frac{2}{3}\epsilon < \epsilon$$

for any $r \in (\delta, 1)$. Therefore, $\lim_{r \rightarrow 1} \mathcal{B}_{rS}^{\text{ext}}[A] = A$ in the norm topology. Hence, and due to (2-2), we deduce that $\lim_{r \rightarrow 1} F(rS) = A$ in the norm topology, which shows that (i) holds. Since \mathcal{H} is infinite-dimensional, that (iii) implies (i) is clear.

It remains to prove that (ii) implies (iii). We assume that (ii) holds. Then there exists $A \in \mathcal{P}$ such that $F(X) = \mathcal{B}_X^{\text{ext}}[A]$ for all $X \in \mathbf{B}_n(\mathcal{H})$. Due to [Theorem 2.2](#), F is a bounded free k -pluriharmonic function on $\mathbf{B}_n(\mathcal{H})$. For any $Y \in \mathbf{B}_n(\mathcal{H})^-$, one can show, as in the proof that (ii) implies (i), that $\tilde{F}(Y) := \lim_{r \rightarrow 1} \mathcal{B}_{rY}^{\text{ext}}[A]$ exists in the operator norm topology. Since $\|\mathcal{B}_{rY}^{\text{ext}}[A]\| \leq \|A\|$ for any $r \in [0, 1)$, we deduce that $\|\tilde{F}(Y)\| \leq \|A\|$ for any $Y \in \mathbf{B}_n(\mathcal{H})^-$. Note also that \tilde{F} is an extension of F . Lastly, we show that \tilde{F} is continuous on $\mathbf{B}_n(\mathcal{H})^-$. To this end, let $\epsilon > 0$ and, due to the equivalence of (ii) and (i), we can choose $r_0 \in [0, 1)$ such that $\|A - F(r_0S)\| < \frac{1}{3}\epsilon$. Since $A - F(r_0S) \in \mathcal{P}$, we deduce that

$$\|\tilde{F}(Y) - F(r_0Y)\| = \|\lim_{r \rightarrow 1} \mathcal{B}_{rY}^{\text{ext}}[A] - F(r_0Y)\| \leq \limsup_{r \rightarrow 1} \|\mathcal{B}_{rY}^{\text{ext}}[A] - F(r_0Y)\| \leq \|A - F(r_0Y)\| < \frac{1}{3}\epsilon$$

for any $Y \in \mathbf{B}_n(\mathcal{H})^-$. Since F is continuous on $\mathbf{B}_n(\mathcal{H})$, there is $\delta > 0$ such that $\|F(r_0Y) - F(r_0W)\| < \frac{1}{3}\epsilon$ for any $W \in \mathbf{B}_n(\mathcal{H})^-$ with $\|W - Y\| < \delta$. Now note that

$$\|\tilde{F}(Y) - \tilde{F}(W)\| \leq \|\tilde{F}(Y) - F(r_0Y)\| + \|F(r_0Y) - F(r_0W)\| + \|F(r_0W) - \tilde{F}(W)\| < \epsilon$$

for any $W \in \mathbf{B}_n(\mathcal{H})^-$ with $\|W - Y\| < \delta$. □

3. Naimark-type dilation theorem for direct products of free semigroups

In this section, we provide a Naimark-type dilation theorem for direct products $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$ of unital free semigroups, and use it to obtain a structure theorem which characterizes the positive free k -pluriharmonic functions on the regular polyball with operator-valued coefficients.

Consider the unital semigroup $\mathbf{F}_n^+ := \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$ with neutral element $\mathbf{g} := (g_0^1, \dots, g_0^k)$. Let $\omega = (\omega_1, \dots, \omega_k)$ and $\gamma = (\gamma_1, \dots, \gamma_k)$ be in $\mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$. We say that ω and γ are left comparable, and write $\omega \sim_{lc} \gamma$, if for each $i \in \{1, \dots, k\}$, one of the conditions $\omega_i <_l \gamma_i$, $\gamma_i <_l \omega_i$ or $\omega_i = \gamma_i$ holds (see the definitions preceding Lemma 1.2). In this case, we define

$$c_l^+(\omega, \gamma) := (c_l^+(\omega_1, \gamma_1), \dots, c_l^+(\omega_k, \gamma_k)) \quad \text{and} \quad c_l^-(\omega, \gamma) := (c_l^-(\omega_1, \gamma_1), \dots, c_l^-(\omega_k, \gamma_k)),$$

where

$$c_l^+(\omega, \gamma) := \begin{cases} \omega \setminus_l \gamma & \text{if } \gamma <_l \omega, \\ g_0 & \text{if } \omega <_l \gamma \text{ or } \omega = \gamma, \end{cases} \quad \text{and} \quad c_l^-(\omega, \gamma) := \begin{cases} \gamma \setminus_r \omega & \text{if } \omega <_l \gamma, \\ g_0 & \text{if } \gamma <_l \omega \text{ or } \omega = \gamma. \end{cases}$$

We say that $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$ is a left k -multi-Toeplitz kernel if $K(\mathbf{g}, \mathbf{g}) = I_{\mathcal{E}}$ and

$$K(\sigma, \omega) = \begin{cases} K(c_l^+(\sigma, \omega); c_l^-(\sigma, \omega)) & \text{if } \sigma \sim_{lc} \omega, \\ 0 & \text{otherwise.} \end{cases}$$

The kernel K is positive semidefinite if, for each $m \in \mathbb{N}$, any choice of $h_1, \dots, h_m \in \mathcal{E}$, and any $\sigma^{(i)} := (\sigma_1^{(i)}, \dots, \sigma_k^{(i)}) \in \mathbf{F}_n^+$, it satisfies the inequality

$$\sum_{i,j=1}^m \langle K(\sigma^{(i)}, \sigma^{(j)})h_j, h_i \rangle \geq 0.$$

Definition 3.1. A map $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$ has a Naimark dilation if there exists a k -tuple of commuting row isometries $V = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, on a Hilbert space $\mathcal{K} \supset \mathcal{E}$, i.e., the nonselfadjoint algebra $\text{Alg}(V_i)$ commutes with $\text{Alg}(V_s)$ for any $i, s \in \{1, \dots, k\}$ with $i \neq s$, such that

$$K(\sigma, \omega) = P_{\mathcal{E}} V_{\sigma}^* V_{\omega}|_{\mathcal{E}}, \quad \sigma, \omega \in \mathbf{F}_n^+.$$

The dilation is called minimal if $\mathcal{K} = \bigvee_{\omega \in \mathbf{F}_n^+} V_{\omega} \mathcal{E}$.

Theorem 3.2. A map $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{H})$ is a positive semidefinite left k -multi-Toeplitz kernel on the direct product \mathbf{F}_n^+ of free semigroups if and only if it admits a Naimark dilation.

Proof. Let \mathcal{K}_0 be the vector space of all sums of tensor monomials $\sum_{\sigma \in \mathbf{F}_n^+} e_\sigma \otimes h_\sigma$, where $\{h_\sigma\}_{\sigma \in \mathbf{F}_n^+}$ is a finitely supported sequence of vectors in \mathcal{H} . Define the sesquilinear form $\langle \cdot, \cdot \rangle_{\mathcal{K}_0}$ on \mathcal{K}_0 by setting

$$\left\langle \sum_{\omega \in \mathbf{F}_n^+} e_\omega \otimes h_\omega, \sum_{\sigma \in \mathbf{F}_n^+} e_\sigma \otimes h'_\sigma \right\rangle_{\mathcal{K}_0} := \sum_{\omega, \sigma \in \mathbf{F}_n^+} \langle K(\sigma, \omega) h_\omega, h'_\sigma \rangle_{\mathcal{H}}, \quad h_\omega, h'_\sigma \in \mathcal{H}.$$

Since K is positive semidefinite, so is $\langle \cdot, \cdot \rangle_{\mathcal{K}_0}$. Set $\mathcal{N} := \{f \in \mathcal{K}_0 : \langle f, f \rangle = 0\}$ and define the Hilbert space obtained by completing $\mathcal{K}_0/\mathcal{N}$ with the induced inner product. For each $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$, define the operator $V_{i,j}$ on \mathcal{K}_0 by setting

$$V_{i,j} \left(\sum_{\sigma \in \mathbf{F}_n^+} e_\sigma \otimes h_\sigma \right) := \sum_{\sigma = (\sigma_1, \dots, \sigma_k) \in \mathbf{F}_n^+} e_{\sigma_1} \otimes \dots \otimes e_{\sigma_{i-1}} \otimes e_{g_j \sigma_i} \otimes e_{\sigma_{i+1}} \otimes \dots \otimes e_{\sigma_k} \otimes h_\sigma.$$

Note that if $p \in \{1, \dots, n_i\}$ then

$$\begin{aligned} & \left\langle V_{i,j} \left(\sum_{\omega \in \mathbf{F}_n^+} e_\omega \otimes h_\omega \right), V_{i,p} \left(\sum_{\sigma \in \mathbf{F}_n^+} e_\sigma \otimes h'_\sigma \right) \right\rangle_{\mathcal{K}_0} \\ &= \sum_{\omega, \sigma \in \mathbf{F}_n^+} \langle K(\sigma_1, \dots, \sigma_{i-1}, g_j \sigma_i, \sigma_{i+1}, \dots, \sigma_k; \omega_1, \dots, \omega_{i-1}, g_p \omega_i, \omega_{i+1}, \dots, \omega_k) h_\omega, h'_\sigma \rangle_{\mathcal{H}} \\ &= \begin{cases} \sum_{\omega, \sigma \in \mathbf{F}_n^+} \langle K(\sigma, \omega) h_\omega, h'_\sigma \rangle_{\mathcal{H}} & \text{if } j = p, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Hence and using the definition of $\langle \cdot, \cdot \rangle_{\mathcal{K}_0}$, we deduce that, for each $i \in \{1, \dots, k\}$, the operators $V_{i,1}, \dots, V_{i,n_i}$ can be extended by continuity to isometries on \mathcal{K} with orthogonal ranges. Note also that, if $i, s \in \{1, \dots, k\}$, $i \neq s$, $j \in \{1, \dots, n_i\}$ and $t \in \{1, \dots, n_s\}$, then

$$V_{i,j} V_{st} (e_{\sigma_1} \otimes \dots \otimes e_{\sigma_k} \otimes h) = e_{\sigma_1} \otimes \dots \otimes e_{\sigma_{i-1}} \otimes e_{g_j \sigma_i} \otimes e_{\sigma_{i+1}} \otimes \dots \otimes e_{\sigma_{s-1}} \otimes e_{\sigma_{gt} \sigma_s} \otimes e_{\sigma_{s+1}} \otimes \dots \otimes e_{\sigma_k} \otimes h$$

when $i < s$. This shows that $V_{ij} V_{st} = V_{st} V_{ij}$. Since

$$\langle e_g \otimes h, e_g \otimes h' \rangle_{\mathcal{K}} = \langle K(\mathbf{g}, \mathbf{g}) h, h' \rangle_{\mathcal{H}} = \langle h, h' \rangle_{\mathcal{H}}, \quad h, h' \in \mathcal{H},$$

we can embed \mathcal{H} into \mathcal{K} by setting $h = e_g \otimes h$. Note that, for any $\omega, \sigma \in \mathbf{F}_n^+$ and $h, h' \in \mathcal{H}$, we have

$$\langle V_\sigma^* V_\omega h, h' \rangle_{\mathcal{K}} = \langle V_\omega h, V_\sigma h' \rangle_{\mathcal{K}} = \langle e_\omega \otimes h, e_\sigma \otimes h' \rangle_{\mathcal{K}} = \langle K(\sigma, \omega) h, h' \rangle_{\mathcal{H}}.$$

Therefore, $K(\sigma, \omega) = P_{\mathcal{H}} V_\sigma^* V_\omega|_{\mathcal{H}}$ for any $\sigma, \omega \in \mathbf{F}_n^+$. Since any element in \mathcal{K}_0 is a linear combination of vectors $V_\sigma h$, where $\sigma \in \mathbf{F}_n^+$ and $h \in \mathcal{H}$, we deduce that $\mathcal{K} = \bigvee_{\omega \in \mathbf{F}_n^+} V_\omega \mathcal{H}$, which proves the minimality of the Naimark dilation.

Now we prove the converse. Let $V = (V_1, \dots, V_n)$ and $V_i = (V_{i,1}, \dots, V_{i,n_i})$ be k -tuples of commuting row isometries on a Hilbert space $\mathcal{K} \supset \mathcal{H}$. Define $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{H})$ by setting $K(\sigma, \omega) = P_{\mathcal{H}} V_\sigma^* V_\omega|_{\mathcal{H}}$ for any $\sigma, \omega \in \mathbf{F}_n^+$. Assume that $\sigma, \omega \in \mathbf{F}_n^+$ and $\sigma \sim_{lc} \omega$. Using the commutativity of the row isometries V_1, \dots, V_k , we can assume without loss of generality that there is $p \in \{1, \dots, k\}$ such that

$\omega_1 \leq_l \sigma_1, \dots, \omega_p \leq_l \sigma_p, \sigma_{p+1} \leq_l \omega_{p+1}, \dots, \sigma_k \leq_l \omega$. Since each $V_i = (V_{i,1}, \dots, V_{i,n_i})$ is an isometry, we have $V_{i,t}^* V_{i,s} = \delta_{ts} I$. Consequently, and using the commutativity of the row isometries, we deduce that

$$\begin{aligned} & \langle V_{1,\omega_1} \cdots V_{k,\omega_k} h, V_{1,\sigma_1} \cdots V_{k,\sigma_k} h' \rangle \\ &= \langle V_{2,\omega_2} \cdots V_{k,\omega_k} h, V_{1,\sigma_1 \setminus \omega_1} V_{2,\sigma_2} \cdots V_{k,\sigma_k} h' \rangle \\ &= \langle V_{2,\omega_2} \cdots V_{k,\omega_k} h, V_{2,\sigma_2} \cdots V_{k,\sigma_k} V_{1,\sigma_1 \setminus \omega_1} h' \rangle \\ &\quad \vdots \\ &= \langle V_{p+1,\omega_{p+1}} \cdots V_{k,\omega_k} h, V_{p+1,\sigma_{p+1}} \cdots V_{k,\sigma_k} V_{1,\sigma_1 \setminus \omega_1} \cdots V_{p,\sigma_p \setminus \omega_p} h' \rangle \\ &= \langle V_{p+1,\omega_{p+1} \setminus \sigma_{p+1}} V_{p+2,\omega_{p+2}} \cdots V_{k,\omega_k} h, V_{p+2,\sigma_{p+2}} \cdots V_{k,\sigma_k} V_{1,\sigma_1 \setminus \omega_1} \cdots V_{p,\sigma_p \setminus \omega_p} h' \rangle \\ &= \langle V_{p+2,\omega_{p+2}} \cdots V_{k,\omega_k} V_{p+1,\omega_{p+1} \setminus \sigma_{p+1}} h, V_{p+2,\sigma_{p+2}} \cdots V_{k,\sigma_k} V_{1,\sigma_1 \setminus \omega_1} \cdots V_{p,\sigma_p \setminus \omega_p} h' \rangle \\ &\quad \vdots \\ &= \langle V_{p+1,\omega_{p+1} \setminus \sigma_{p+1}} \cdots V_{k,\omega_k \setminus \sigma_k} h, V_{1,\sigma_1 \setminus \omega_1} \cdots V_{p,\sigma_p \setminus \omega_p} h' \rangle \\ &= \langle V_{1,\sigma_1 \setminus \omega_1}^* \cdots V_{p,\sigma_p \setminus \omega_p}^* V_{p+1,\omega_{p+1} \setminus \sigma_{p+1}} \cdots V_{k,\omega_k \setminus \sigma_k} h, h' \rangle \end{aligned}$$

for any $h, h' \in \mathcal{H}$. Therefore, for any $\sigma, \omega \in \mathbf{F}_n^+$, we have

$$\begin{aligned} K(\sigma, \omega) &= P_{\mathcal{H}} V_{\sigma}^* V_{\omega} |_{\mathcal{H}} = \begin{cases} P_{\mathcal{H}} V_{c_l^+(\sigma, \omega)}^* V_{c_l^-(\sigma, \omega)} |_{\mathcal{H}} & \text{if } \sigma \sim_{lc} \omega, \\ 0 & \text{otherwise,} \end{cases} \\ &= \begin{cases} K(c_l^+(\sigma, \omega); c_l^-(\sigma, \omega)) & \text{if } \sigma \sim_{lc} \omega, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

and $K(\mathbf{g}, \mathbf{g}) = I_{\mathcal{H}}$. This shows that K is a left k -multi-Toeplitz kernel on \mathbf{F}_n^+ . On the other hand, for any finitely supported sequence $\{h_{\omega}\}_{\omega \in \mathbf{F}_n^+}$ of elements in \mathcal{H} , we have

$$\sum_{\omega, \sigma \in \mathbf{F}_n^+} \langle K(\sigma, \omega) h_{\omega}, h_{\sigma} \rangle = \sum_{\omega, \sigma \in \mathbf{F}_n^+} \langle P_{\mathcal{H}} V_{\sigma}^* V_{\omega} |_{\mathcal{H}} h_{\omega}, h_{\sigma} \rangle = \left\| \sum_{\omega \in \mathbf{F}_n^+} V_{\omega} h_{\omega} \right\|^2 \geq 0.$$

Therefore, K is a positive semidefinite left k -multi-Toeplitz kernel on \mathbf{F}_n^+ . □

We remark that the Naimark dilation provided in [Theorem 3.2](#) is minimal. To prove the uniqueness of the minimal Naimark dilation, let $\mathbf{V}' = (V'_1, \dots, V'_n)$, $V'_i = (V'_{i,1}, \dots, V'_{i,n_i})$, be a k -tuple of commuting row isometries on a Hilbert space $\mathcal{K}' \supset \mathcal{H}$ such that $K(\sigma, \omega) = P_{\mathcal{H}}^{\mathcal{K}'} (\mathbf{V}'_{\sigma})^* \mathbf{V}'_{\omega} |_{\mathcal{H}}$ for any $\sigma, \omega \in \mathbf{F}_n^+$ and with the property that $\mathcal{K} = \bigvee_{\omega \in \mathbf{F}_n^+} V'_{\omega} \mathcal{H}$. For any $x, y \in \mathcal{H}$, we have

$$\langle V_{\omega} x, V_{\sigma} y \rangle_{\mathcal{K}} = \langle K(\sigma, \omega) x, y \rangle_{\mathcal{H}} = \langle P_{\mathcal{H}}^{\mathcal{K}'} (\mathbf{V}'_{\sigma})^* \mathbf{V}'_{\omega} x, y \rangle_{\mathcal{K}'} = \langle \mathbf{V}'_{\omega} x, \mathbf{V}'_{\sigma} y \rangle_{\mathcal{K}'}$$

Consequently, the map

$$W \left(\sum_{\sigma \in \mathbf{F}_n^+} V_{\sigma} h_{\sigma} \right) := \sum_{\sigma \in \mathbf{F}_n^+} V'_{\sigma} h_{\sigma},$$

where $\{h_{\sigma}\}_{\sigma \in \mathbf{F}_n^+}$ is any finitely supported sequence of vectors in \mathcal{H} , is well-defined. Due to the minimality of the spaces \mathcal{K} and \mathcal{K}' , the map extends to a unitary operator W from \mathcal{K} onto \mathcal{K}' . Note also that

$WV_{i,j} = V'_{i,j}W$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$, which completes the proof of the uniqueness of the minimal Naimark dilation.

We should mention that there is a dual Naimark-type dilation for positive semidefinite right k -multi-Toeplitz kernels. A kernel $\Gamma : F_n^+ \times F_n^+ \rightarrow B(\mathcal{E})$ is called *right k -multi-Toeplitz* if $\Gamma(\mathbf{g}, \mathbf{g}) = I_{\mathcal{E}}$ and

$$\Gamma(\sigma, \omega) = \begin{cases} \Gamma(c_r^+(\sigma, \omega); c_r^-(\sigma, \omega)) & \text{if } \sigma \sim_{rc} \omega, \\ 0 & \text{otherwise,} \end{cases}$$

where $c_r^+(\sigma, \omega), c_r^-(\sigma, \omega)$ are defined by (1-1). We say that Γ has a Naimark dilation if there exists a k -tuple $\mathbf{W} = (W_1, \dots, W_n)$, $W_i = (W_{i,1}, \dots, W_{i,n_i})$, of commuting row isometries on a Hilbert space $\mathcal{K} \supset \mathcal{E}$ such that $\Gamma(\tilde{\sigma}, \tilde{\omega}) = P_{\mathcal{E}} \mathbf{W}_{\tilde{\sigma}}^* \mathbf{W}_{\tilde{\omega}}|_{\mathcal{E}}$ for any $\sigma, \omega \in F_n^+$.

Theorem 3.3. *A map $\Gamma : F_n^+ \times F_n^+ \rightarrow B(\mathcal{H})$ is a positive semidefinite right k -multi-Toeplitz kernel on F_n^+ if and only if it admits a Naimark dilation. In this case, there is a minimal dilation which is uniquely determined up to isomorphism.*

Proof. We only sketch the proof, which is very similar to that of Theorem 3.2, pointing out the differences. First, \mathcal{K}_0 is the vector space of all sums of tensor monomials $\sum_{\sigma \in F_n^+} e_{\tilde{\sigma}} \otimes h_{\sigma}$, where $\{h_{\sigma}\}_{\sigma \in F_n^+}$ is a finitely supported sequence of vectors in \mathcal{H} , while the sesquilinear form $\langle \cdot, \cdot \rangle_{\mathcal{K}_0}$ on \mathcal{K}_0 is defined by setting

$$\left\langle \sum_{\omega \in F_n^+} e_{\tilde{\omega}} \otimes h_{\omega}, \sum_{\sigma \in F_n^+} e_{\tilde{\sigma}} \otimes h'_{\sigma} \right\rangle_{\mathcal{K}_0} := \sum_{\omega, \sigma \in F_n^+} \langle \Gamma(\sigma, \omega) h_{\omega}, h'_{\sigma} \rangle_{\mathcal{H}}.$$

For each $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$, we define the operator $W_{i,j}$ on \mathcal{K}_0 by setting

$$W_{i,j} \left(\sum_{\sigma \in F_n^+} e_{\tilde{\sigma}} \otimes h_{\sigma} \right) := \sum_{\sigma = (\sigma_1, \dots, \sigma_k) \in F_n^+} e_{\tilde{\sigma}_1} \otimes \dots \otimes e_{\tilde{\sigma}_{i-1}} \otimes e_{g_j \tilde{\sigma}_i} \otimes e_{\tilde{\sigma}_{i+1}} \otimes \dots \otimes e_{\tilde{\sigma}_k} \otimes h_{\sigma}.$$

Taking into account the relations

$$c_r^+(\sigma, \omega) \sim = c_l^+(\tilde{\sigma}, \tilde{\omega}) \quad \text{and} \quad c_r^-(\sigma, \omega) \sim = c_l^-(\tilde{\sigma}, \tilde{\omega}),$$

we deduce that

$$\begin{aligned} P_{\mathcal{H}} \mathbf{W}_{\sigma}^* \mathbf{W}_{\omega} |_{\mathcal{H}} &= \begin{cases} P_{\mathcal{H}} \mathbf{W}_{c_l^+(\sigma, \omega)}^* \mathbf{W}_{c_l^-(\sigma, \omega)} |_{\mathcal{H}} & \text{if } \sigma \sim_{lc} \omega, \\ 0 & \text{otherwise,} \end{cases} \\ &= \begin{cases} \Gamma(c_l^+(\sigma, \omega) \sim; c_l^-(\sigma, \omega) \sim) & \text{if } \sigma \sim_{lc} \omega, \\ 0 & \text{otherwise,} \end{cases} \\ &= \begin{cases} \Gamma(c_r^+(\tilde{\sigma}, \tilde{\omega}); c_r^-(\tilde{\sigma}, \tilde{\omega})) & \text{if } \tilde{\sigma} \sim_{rc} \tilde{\omega}, \\ 0 & \text{otherwise,} \end{cases} \\ &= \Gamma(\tilde{\sigma}, \tilde{\omega}) \end{aligned}$$

for any $\sigma, \omega \in F_n^+$. The rest of the proof is similar to that of Theorem 3.2. We leave it to the reader. \square

Let F be a free k -pluriharmonic on the polyball \mathbf{B}_n with operator-valued coefficients in $B(\mathcal{E})$ with representation

$$F(X) = \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*, \tag{3-1}$$

where the series converge in the operator norm topology for any $X = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$, with $X_i := (X_{i,1}, \dots, X_{i,n_i})$, and any Hilbert space \mathcal{H} . We associate to F the kernel $\Gamma_F : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$ given by

$$\Gamma_F(\sigma, \omega) := \begin{cases} A_{(c_r^+(\sigma, \omega); c_r^-(\sigma, \omega))} & \text{if } \sigma \sim_{rc} \omega, \\ 0 & \text{otherwise.} \end{cases} \tag{3-2}$$

One can easily see that Γ_F is a right k -multi-Toeplitz kernel on \mathbf{F}_n^+ . In what follows, we prove a Schur-type result for positive k -pluriharmonic functions in polyballs.

Theorem 3.4. *Let F be a k -pluriharmonic function on the regular polyball \mathbf{B}_n , with coefficients in $B(\mathcal{E})$. Then F is positive on \mathbf{B}_n if and only if the kernel Γ_{F_r} is positive semidefinite for any $r \in [0, 1)$, where F_r stands for the mapping $X \mapsto F(rX)$.*

Proof. For every $\gamma := (\gamma_1, \dots, \gamma_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, we set $e_\gamma := e_{\gamma_1}^1 \otimes \cdots \otimes e_{\gamma_k}^k$ and $S_\gamma := S_{1, \gamma_1} \cdots S_{k, \gamma_k}$. Let F be a k -pluriharmonic function with representation (3-1). Taking into account Lemma 1.2, for each $\gamma, \omega \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, $r \in [0, 1)$, and $h, h' \in \mathcal{E}$, we have

$$\begin{aligned} \langle F(rS)(h \otimes e_\gamma), h' \otimes e_\omega \rangle &= \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, \\ i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \langle A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, h' \rangle r^{\sum_{i=1}^k (|\alpha_i| + |\beta_i|)} \langle S_\alpha S_\beta^* e_\gamma, e_\omega \rangle \\ &= \begin{cases} r^{\sum_{i=1}^k (|\alpha_i| + |\beta_i|)} \langle A_{(c_r^+(\omega, \gamma); c_r^-(\omega, \gamma))} h, h' \rangle & \text{if } \omega \sim_{rc} \gamma, \\ 0 & \text{otherwise,} \end{cases} \\ &= \langle \Gamma_{F_r}(\omega, \gamma) h, h' \rangle. \end{aligned}$$

Hence, we deduce that the kernel Γ_{F_r} is positive semidefinite for any $r \in [0, 1)$ if and only if $F(rS) \geq 0$ for any $r \in [0, 1)$. Now, let $X \in \mathbf{B}_n(\mathcal{H})$ and let $r \in (0, 1)$ be such that $(1/r)X \in \mathbf{B}_n(\mathcal{H})$. Since the noncommutative Berezin transform $\mathcal{B}_{(1/r)X}$ is continuous in the operator norm and completely positive, so is $\text{id} \otimes \mathcal{B}_{(1/r)X}$. Consequently, we obtain

$$F(X) = (\text{id} \otimes \mathcal{B}_{(1/r)X})[F(rS)] \geq 0, \quad X \in \mathbf{B}_n(\mathcal{H}).$$

Note that if F is positive on \mathbf{B}_n then $F(rS) \geq 0$ for any $r \in [0, 1)$. □

Corollary 3.5. *Let $f : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ be a free holomorphic function. Then the following statements are equivalent:*

- (i) $\Re f \geq 0$ on the polyball \mathbf{B}_n .

- (ii) $\Re f(r\mathcal{S}) \geq 0$ for any $r \in [0, 1)$.
- (iii) The right k -multi Toeplitz kernel $\Gamma_{\Re f_r}$ is positive semidefinite for any $r \in [0, 1)$.

Let us define the free k -pluriharmonic Poisson kernel by setting

$$\mathcal{P}(Y, X) := \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_n^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} Y_{1, \tilde{\alpha}_1}^* \cdots Y_{k, \tilde{\alpha}_k}^* Y_{1, \tilde{\beta}_1} \cdots Y_{k, \tilde{\beta}_k} \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*$$

for any $X \in \mathbf{B}_n(\mathcal{H})$ and any $Y = (Y_1, \dots, Y_k)$ with $Y_i = (Y_{i,1}, \dots, Y_{i,n_i}) \in B(\mathcal{K})^{n_i}$ such that the series above is convergent in the operator norm topology. Let $\Omega \subset \mathbf{F}_n^+ \times \mathbf{F}_n^+$ be the set of all (α, β) where $\alpha = (\alpha_1, \dots, \alpha_k), \beta = (\beta_1, \dots, \beta_k) \in \mathbf{F}_n^+$ are such that $\alpha_i, \beta_i \in \mathbb{F}_n^+, |\alpha_i| = m_i^-, |\beta_i| = m_i^+$ for some $m_i \in \mathbb{Z}$.

Theorem 3.6. *A map $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ with $F(0) = I$ is a positive free k -pluriharmonic function on the regular polyball if and only if it has the form*

$$F(X) = \sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} V_{\alpha}^* V_{\beta} |_{\mathcal{E}} \otimes X_{\alpha} X_{\beta}^*,$$

where $V = (V_1, \dots, V_k)$ is a k -tuple of commuting row isometries on a space $\mathcal{K} \supset \mathcal{E}$ such that

$$\sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} V_{\alpha}^* V_{\beta} |_{\mathcal{E}} \otimes r^{|\alpha|+|\beta|} S_{\alpha} S_{\beta}^* \geq 0, \quad r \in [0, 1),$$

and the series is convergent in the operator topology.

Proof. Assume that F is a positive free k -pluriharmonic function which has the representation (3-1) and $F(0) = I$. Due to Theorem 3.4, $F(r\mathcal{S}) \geq 0$ and the right k -multi-Toeplitz kernel Γ_{F_r} is positive semidefinite for any $r \in [0, 1)$. Taking limits as $r \rightarrow \infty$, we deduce that Γ_F is positive semidefinite as well. According to Theorem 3.3, Γ_F has a Naimark-type dilation. Therefore, there is a k -tuple $V = (V_1, \dots, V_k)$ of commuting row isometries on a Hilbert space $\mathcal{K} \supset \mathcal{E}$ such that $\Gamma(\tilde{\sigma}, \tilde{\omega}) = P_{\mathcal{E}} V_{\sigma}^* V_{\omega} |_{\mathcal{E}}$ for any $\sigma, \omega \in \mathbf{F}_n^+$. Using (3-1) and (3-2), we deduce that

$$F(X) = \sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} V_{\alpha}^* V_{\beta} |_{\mathcal{E}} \otimes X_{\alpha} X_{\beta}^*,$$

where the convergence is in the norm topology. This shows, in particular, that $F(r\mathcal{S})$ is convergent.

To prove the converse, assume that $V = (V_1, \dots, V_k)$ is a k -tuple of commuting row isometries on a space $\mathcal{K} \supset \mathcal{E}$ such that

$$\sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} V_{\alpha}^* V_{\beta} |_{\mathcal{E}} \otimes r^{|\alpha|+|\beta|} S_{\alpha} S_{\beta}^* \geq 0, \quad r \in [0, 1), \tag{3-3}$$

and the convergence is in the operator norm topology. Let $X \in \mathbf{B}_n(\mathcal{H})$ and let $r \in (0, 1)$ be such that $(1/r)X \in \mathbf{B}_n(\mathcal{H})$. Since the noncommutative Berezin transform $\mathcal{B}_{(1/r)X}$ is continuous in the operator norm and completely positive, so is $\text{id} \otimes \mathcal{B}_{(1/r)X}$. Consequently, we obtain

$$F(X) := (\text{id} \otimes \mathcal{B}_{(1/r)X}) \left(\sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} V_{\alpha}^* V_{\beta} |_{\mathcal{E}} \otimes r^{|\alpha|+|\beta|} S_{\alpha} S_{\beta}^* \right) \geq 0, \quad X \in \mathbf{B}_n(\mathcal{H}). \quad \square$$

We remark that the condition (3-3) is equivalent to the condition that the kernel defined by the relation $\Gamma_{rV}(\tilde{\sigma}, \tilde{\omega}) := r^{|\sigma|+|\omega|} P_{\mathcal{E}} V_{\sigma}^* V_{\omega}|_{\mathcal{E}}$ for any $\sigma, \omega \in F_n^+$ is positive semidefinite. We should also mention that one can find a version of the theorem above when the condition $F(0) = I$ is dropped. In this case, $F(0) = A \otimes I$ with $A \geq 0$ and we set

$$G_{\epsilon} := [(A + \epsilon I_{\mathcal{E}})^{-1/2} \otimes I](F + \epsilon I_{\mathcal{E}} \otimes I)[(A + \epsilon I_{\mathcal{E}})^{-1/2} \otimes I], \quad \epsilon > 0.$$

Since G_{ϵ} is a positive k -pluriharmonic function with $G_{\epsilon}(0) = I$, we can apply Theorem 3.6 to get a family $V(\epsilon) = (V_1(\epsilon), \dots, V_k(\epsilon))$ of k -tuples of commuting row isometries on a space $\mathcal{K}_{\epsilon} \supset \mathcal{E}$ such that

$$F(X) = \lim_{\epsilon \rightarrow 0} \sum_{(\alpha, \beta) \in \Omega} (A + \epsilon I_{\mathcal{E}})^{1/2} [P_{\mathcal{E}} V_{\alpha}^*(\epsilon) V_{\beta}(\epsilon)|_{\mathcal{E}}] (A + \epsilon I_{\mathcal{E}})^{1/2} \otimes X_{\alpha} X_{\beta}^*,$$

where the convergence is in the operator norm topology.

Definition 3.7. A k -tuple $V = (V_1, \dots, V_k)$ of commuting row isometries $V_i = (V_{i,1}, \dots, V_{i,n_i})$ is called *pluriharmonic* if the free k -pluriharmonic Poisson kernel $\mathcal{P}(V, rS)$ is a positive operator for any $r \in [0, 1)$.

Proposition 3.8. Let $V = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, be a k -tuple of commuting row isometries. Then V is pluriharmonic in each of the following cases:

- (i) $k = 1$ and $n_1 \in \mathbb{N}$.
- (ii) V is doubly commuting, i.e., the C^* -algebra $C^*(V_i)$ commutes with $C^*(V_s)$ if $i, s \in \{1, \dots, k\}$ with $i \neq s$.
- (iii) $n_1 = \dots = n_k = 1$.

Proof. It is easy to see that V is pluriharmonic if the condition in (i) is satisfied. Under the condition (ii), the proof that V is pluriharmonic is similar to the proof of Theorem 4.2(i), when we replace the universal operator R with V . Now, we assume that $n_1 = \dots = n_k = 1$. Then $V = (V_1, \dots, V_k)$, where V_1, \dots, V_k are commuting isometries on a Hilbert space \mathcal{K} . It is well known [Sz.-Nagy et al. 2010] that there is a k -tuple $U = (U_1, \dots, U_k)$ of commuting unitaries on a Hilbert space $\mathcal{G} \supset \mathcal{K}$ such that $U_i|_{\mathcal{K}} = V_i$ for $i \in \{1, \dots, k\}$. Due to Fuglede’s theorem (see [Douglas 1998]), the unitaries are doubly commuting. Due to (ii), $\mathcal{P}(U, rS)$ is a well-defined positive operator for any $r \in [0, 1)$, where the convergence defining the free k -pluriharmonic Poisson kernel $\mathcal{P}(U, rS)$ is in the operator norm topology. On the other hand, we have

$$\mathcal{P}(V, rS) = (P_{\mathcal{K}} \otimes I) \mathcal{P}(U, rS)|_{\mathcal{K} \otimes \otimes_{i=1}^k F^2(H_{n_i})} \geq 0,$$

which completes the proof. □

Proposition 3.9. Let $V = (V_1, \dots, V_k)$ be a pluriharmonic tuple of commuting row isometries on a Hilbert space \mathcal{K} and let $\mathcal{E} \subset \mathcal{K}$ be a subspace. Then the map

$$F(X) := (P_{\mathcal{E}} \otimes I) \mathcal{P}(V, X)|_{\mathcal{E} \otimes \mathcal{H}}, \quad X \in B_n(\mathcal{H}),$$

is a positive free k -pluriharmonic function on the polyball B_n with operator-valued coefficients in $B(\mathcal{E})$ and $F(0) = I$.

Moreover, in the particular cases (i) and (iii) of [Proposition 3.8](#), each positive free k -pluriharmonic function F with $F(0) = I$ has the form above.

Proof. Since \mathbf{V} is a tuple of commuting row isometries, the free k -pluriharmonic Poisson kernel $\mathcal{P}(\mathbf{V}, r\mathbf{S})$ is a positive operator for any $r \in [0, 1)$ and so is the compression $(P_{\mathcal{E}} \otimes I)\mathcal{P}(\mathbf{V}, r\mathbf{S})|_{\mathcal{E} \otimes \bigotimes_{i=1}^k F^2(H_{n_i})}$. Let $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$ and let $r \in (0, 1)$ be such that $(1/r)\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$. Since the noncommutative Berezin transform $\text{id} \otimes \mathcal{B}_{(1/r)\mathbf{X}}$ is continuous in the operator norm and completely positive, we deduce that

$$F(\mathbf{X}) := (P_{\mathcal{E}} \otimes I)\mathcal{P}(\mathbf{V}, \mathbf{X})|_{\mathcal{E} \otimes \mathcal{H}} \geq 0, \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}),$$

where the convergence of $\mathcal{P}(\mathbf{V}, \mathbf{X})$ is in the operator norm topology. Therefore, F is a positive free k -pluriharmonic function on the polyball \mathbf{B}_n with operator-valued coefficients in $B(\mathcal{E})$ and $F(0) = I$. To prove the second part of this proposition, assume that F is a positive free k -pluriharmonic function with $F(0) = I$. According to [Theorem 3.6](#), F has the form

$$F(\mathbf{X}) = \sum_{(\alpha, \beta) \in \Omega} P_{\mathcal{E}} \mathbf{V}_{\alpha}^* \mathbf{V}_{\beta} |_{\mathcal{E}} \otimes \mathbf{X}_{\alpha} \mathbf{X}_{\beta}^*,$$

where $\mathbf{V} = (V_1, \dots, V_k)$ is a k -tuple of commuting row isometries on a space $\mathcal{K} \supset \mathcal{E}$ and the convergence of the series is in the operator norm topology. Since in the particular cases (i) and (ii) of [Proposition 3.8](#) \mathbf{V} is pluriharmonic, one can easily complete the proof. \square

We remark that the theorem above contains, in particular, a structure theorem for positive k -harmonic functions on the regular polydisk included in $[B(\mathcal{H})]_1 \times_c \dots \times_c [B(\mathcal{H})]_1$, which extends the corresponding classical result on scalar polydisks [[Rudin 1969](#)]. In the general case of the polyball it is unknown if all positive free k -pluriharmonic functions F with $F(0) = I$ have the form of [Proposition 3.9](#).

4. Berezin transforms of completely bounded maps in regular polyballs

We define a class of noncommutative Berezin transforms of completely bounded linear maps and give necessary and sufficient conditions for a function to be the Poisson transform of a completely bounded or completely positive map.

Let \mathcal{H} be a Hilbert space and identify the set $M_m(B(\mathcal{H}))$ of $m \times m$ matrices with entries from $B(\mathcal{H})$ with $B(\mathcal{H}^{(m)})$, where $\mathcal{H}^{(m)}$ is the direct sum of m copies of \mathcal{H} . Thus we have a natural C^* -norm on $M_m(B(\mathcal{H}))$. If \mathcal{X} is an operator space, i.e., a closed subspace of $B(\mathcal{H})$, we consider $M_m(\mathcal{X})$ as a subspace of $M_m(B(\mathcal{H}))$ with the induced norm. Let \mathcal{X}, \mathcal{Y} be operator spaces and let $u : \mathcal{X} \rightarrow \mathcal{Y}$ be a linear map. Define the map $u_m : M_m(\mathcal{X}) \rightarrow M_m(\mathcal{Y})$ by $u_m([x_{ij}]) := [u(x_{ij})]$. We say that u is completely bounded if $\|u\|_{\text{cb}} := \sup_{m \geq 1} \|u_m\| < \infty$. If $\|u\|_{\text{cb}} \leq 1$ then u is completely contractive; if u_m is an isometry for any $m \geq 1$ then u is completely isometric; and if u_m is positive for all m then u is called completely positive. For basic results concerning completely bounded maps and operator spaces we refer to [[Paulsen 1986](#); [Pisier 2001](#); [Effros and Ruan 2000](#)].

Let \mathcal{K} be a Hilbert space and let $\mu : B(\bigotimes_{i=1}^k F^2(H_{n_i})) \rightarrow B(\mathcal{K})$ be a completely bounded map. It is well known (see, e.g., [Paulsen 1986]) that there exists a completely bounded linear map

$$\hat{\mu} := \mu \otimes \text{id} : B\left(\bigotimes_{i=1}^k F^2(H_{n_i})\right) \otimes_{\min} B(\mathcal{H}) \rightarrow B(\mathcal{K}) \otimes_{\min} B(\mathcal{H})$$

such that $\hat{\mu}(f \otimes Y) := \mu(f) \otimes Y$ for $f \in B(\bigotimes_{i=1}^k F^2(H_{n_i}))$ and $Y \in B(\mathcal{H})$. Moreover, $\|\hat{\mu}\|_{\text{cb}} = \|\mu\|_{\text{cb}}$ and, if μ is completely positive, then so is $\hat{\mu}$. We introduce the *noncommutative Berezin transform* associated with μ as the map

$$\mathcal{B}_\mu : B\left(\bigotimes_{i=1}^k F^2(H_{n_i})\right) \times \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{K}) \otimes_{\min} B(\mathcal{H})$$

defined by

$$\mathcal{B}_\mu(A, X) := \hat{\mu}[C_X^*(A \otimes I_{\mathcal{H}})C_X], \quad A \in B\left(\bigotimes_{i=1}^k F^2(H_{n_i})\right), \quad X \in \mathbf{B}_n(\mathcal{H}),$$

where the operator $C_X \in B(\bigotimes_{i=1}^k F^2(H_{n_i}) \otimes \mathcal{H})$ is defined by

$$C_X := (I_{\bigotimes_{i=1}^k F^2(H_{n_i})} \otimes \Delta_X(I)^{1/2}) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1}$$

and the *defect mapping* $\Delta_X : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ is given by

$$\Delta_X := (\text{id} - \Phi_{X_1}) \circ \cdots \circ (\text{id} - \Phi_{X_k}),$$

where $\Phi_{X_i} : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ is the completely positive linear map defined by

$$\Phi_{X_i}(Y) := \sum_{j=1}^{n_i} X_{i,j} Y X_{i,j}^*, \quad Y \in B(\mathcal{H}).$$

We need to show that the operator $I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*$ is invertible. Let $\mathbf{Y} = (Y_1, \dots, Y_k)$ with $Y_i := (Y_{i,1}, \dots, Y_{i,n_i}) \in B(\mathcal{H})^{n_i}$. We introduce the spectral radius of \mathbf{Y} by setting

$$r(\mathbf{Y}) := \limsup_{(p_1, \dots, p_k) \in \mathbb{Z}_+^k} \left\| \sum_{\substack{\alpha_i \in \mathbb{F}_{n_i}^+, |\alpha_i|=p_i \\ i \in \{1, \dots, k\}}} Y_\alpha Y_\alpha^* \right\|^{\frac{1}{2(p_1 + \dots + p_k)}},$$

where $Y_\alpha := Y_{1,\alpha_1} \cdots Y_{k,\alpha_k}$ for $\alpha := (\alpha_1, \dots, \alpha_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$ and $Y_{i,\alpha_i} := Y_{i,j_1} \cdots Y_{i,j_p}$ for $\alpha_i = g_{j_1}^{i_1} \cdots g_{j_p}^{i_p} \in \mathbb{F}_{n_i}^+$. We remark that, when $k = 1$, we recover the spectral radius of an n_i -tuple of operators, i.e., $r(Y_i) = \lim_{p \rightarrow \infty} \left\| \sum_{\beta_i \in \mathbb{F}_{n_i}^+, |\beta_i|=p} Y_{i,\beta_i} Y_{i,\beta_i}^* \right\|^{1/2p}$. Note also that

$$r(Y_i) = r(\mathbf{R}_{i,1} \otimes Y_{i,1}^* + \cdots + \mathbf{R}_{i,n_i} \otimes Y_{i,n_i}^*)$$

and $r(Y_i) \leq r(\mathbf{Y})$ for any $i \in \{1, \dots, k\}$. Consequently, if $r(\mathbf{Y}) < 1$ then $r(Y_i) < 1$ and the spectrum of $\mathbf{R}_{i,1} \otimes Y_{i,1}^* + \cdots + \mathbf{R}_{i,n_i} \otimes Y_{i,n_i}^*$ is included in $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$. In particular, when $X \in \mathbf{B}_n(\mathcal{H})$, the

noncommutative von Neumann inequality [Popescu 1999] implies $r(X) \leq r(tS) = t$ for some $t \in (0, 1)$, which proves our assertion.

Proposition 4.1. *Let \mathcal{B}_μ be the noncommutative Berezin transform associated with a completely bounded linear map $\mu : B(\bigotimes_{i=1}^k F^2(H_{n_i})) \rightarrow B(\mathcal{K})$.*

(i) *If $X \in \mathbf{B}_n(\mathcal{H})$ is fixed, then*

$$\mathcal{B}_\mu(\cdot, X) : B\left(\bigotimes_{i=1}^k F^2(H_{n_i})\right) \rightarrow B(\mathcal{K}) \otimes_{\min} B(\mathcal{H})$$

is a completely bounded linear map with $\|\mathcal{B}_\mu(\cdot, X)\|_{\text{cb}} \leq \|\mu\|_{\text{cb}} \|C_X\|^2$.

(ii) *If μ is selfadjoint, then $\mathcal{B}_\mu(A^*, X) = \mathcal{B}_\mu(A, X)^*$. Moreover, if μ is completely positive then so is the map $\mathcal{B}_\mu(\cdot, X)$.*

(iii) *If $A \in B(\bigotimes_{i=1}^k F^2(H_{n_i}))$ is fixed, then the map*

$$\mathcal{B}_\mu(A, \cdot) : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{K}) \otimes_{\min} B(\mathcal{H})$$

is continuous and $\|\mathcal{B}_\mu(A, X)\| \leq \|\mu\|_{\text{cb}} \|A\| \|C_X\|^2$ for any $X \in \mathbf{B}_n(\mathcal{H})$.

Proof. Parts (i) and (ii) follow easily from the definition of the noncommutative Berezin transform associated with μ . To prove (iii), let $X, Y \in \mathbf{B}_n(\mathcal{H})$ and note that

$$\begin{aligned} \|\mathcal{B}_\mu(A, X) - \mathcal{B}_\mu(A, Y)\| &\leq \|\mu\| \|C_X^*(A \otimes I_{\mathcal{H}})(C_X - C_Y)\| + \|\mu\| \|(C_X^* - C_Y^*)(A \otimes I_{\mathcal{H}})C_Y\| \\ &\leq \|\mu\| \|A\| \|C_X - C_Y\| (\|C_X\| + \|C_Y\|). \end{aligned}$$

The continuity of the map $X \mapsto \mathcal{B}_\mu(A, X)$ will follow once we prove that $X \mapsto C_X$ is a continuous map on $\mathbf{B}_n(\mathcal{H})$. Note that

$$\|C_X - C_Y\| \leq \|\Delta_X(I)^{1/2}\| \left\| \prod_{i=1}^k (I - R_{X_i^*})^{-1} - \prod_{i=1}^k (I - R_{Y_i^*})^{-1} \right\| + \|\Delta_X(I)^{1/2} - \Delta_Y(I)^{1/2}\| \left\| \prod_{i=1}^k (I - R_{X_i^*})^{-1} \right\|,$$

where $R_{X_i^*} := I - R_{i,1} \otimes X_{i,1}^* - \dots - R_{i,n_i} \otimes X_{i,n_i}^*$. Since the maps $X \mapsto \prod_{i=1}^k (I - R_{X_i^*})^{-1}$ and $X \mapsto \Delta_X(I)^{1/2}$ are continuous on $\mathbf{B}_n(\mathcal{H})$ in the operator norm topology, our assertion follows. The inequality in (iii) is obvious. □

We remark that the noncommutative Poisson transform introduced in [Popescu 1999] is in fact a particular case of the noncommutative Berezin transform associated with a linear functional. Indeed, let τ be the linear functional on $B(\bigotimes_{i=1}^k F^2(H_{n_i}))$ defined by $\tau(A) := \langle A(1), 1 \rangle$. If $X \in \mathbf{B}_n(\mathcal{H})$ is fixed, then $\mathcal{B}_\tau(\cdot, X) : B(\bigotimes_{i=1}^k F^2(H_{n_i})) \rightarrow B(\mathcal{H})$ is a completely contractive linear map and

$$\langle \mathcal{B}_\tau(A, X)x, y \rangle = \langle C_X^*(A \otimes I_{\mathcal{H}})C_X(1 \otimes x), 1 \otimes y \rangle, \quad x, y \in \mathcal{H}.$$

Hence, we have

$$\mathcal{B}_\tau(A, X) = K_X^*(A \otimes I)K_X,$$

where K_X is the noncommutative Berezin kernel at X . Note also that if $A \in B(\bigotimes_{i=1}^k F^2(H_{n_i}))$ is fixed, then $\mathcal{B}_\tau(A, \cdot) : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{H})$ is a bounded continuous map and $\|\mathcal{B}_\tau(A, X)\| \leq \|A\|$ for any $X \in \mathbf{B}_n(\mathcal{H})$.

We mention that, if $n_1 = \dots = n_k = 1$, $\mathcal{H} = \mathbb{C}$ and $X = \lambda = (\lambda_1, \dots, \lambda_k) \in \mathbb{D}^k$, then we recover the Berezin transform of a bounded linear operator on the Hardy space $H^2(\mathbb{D}^k)$, i.e.,

$$\mathcal{B}_\tau(A, \lambda) = \prod_{i=1}^k (1 - |\lambda_i|^2) \langle Ak_\lambda, k_\lambda \rangle, \quad A \in B(H^2(\mathbb{D}^k)),$$

where $k_\lambda(z) := \prod_{i=1}^k (1 - \bar{\lambda}_i z_i)^{-1}$ and $z = (z_1, \dots, z_k) \in \mathbb{D}^k$.

Define the set

$$\Lambda := \{(\sigma, \omega) \in F_n^+ \times F_n^+ : \sigma \sim_{lc} \omega \text{ and } (\sigma, \omega) = (c_l^+(\sigma, \omega), c_l^-(\sigma, \omega))\}. \tag{4-1}$$

Set $\tilde{\Lambda} := \{(\tilde{\sigma}, \tilde{\omega}) : (\sigma, \omega) \in \Lambda\}$ and note that

$$\tilde{\Lambda} := \{(\tilde{\sigma}, \tilde{\omega}) \in F_n^+ \times F_n^+ : \tilde{\sigma} \sim_{rc} \tilde{\omega} \text{ and } (\tilde{\sigma}, \tilde{\omega}) = (c_r^+(\tilde{\sigma}, \tilde{\omega}), c_r^-(\tilde{\sigma}, \tilde{\omega}))\}.$$

Moreover, we have $\Lambda = \tilde{\Lambda}$. In the case $(\sigma, \omega) \in \Lambda$, one can easily see that $c_l^+(\sigma, \omega) = c_r^+(\sigma, \omega)$ and $c_l^-(\sigma, \omega) = c_r^-(\sigma, \omega)$.

In what follows, we introduce the *noncommutative Poisson transform* of a completely positive linear map on the operator system

$$\mathcal{R}_n^* \mathcal{R}_n := \text{span}\{R_\alpha^* R_\beta : \alpha, \beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+\},$$

where $R := (R_1, \dots, R_k)$ and $R_i := (R_{i,1}, \dots, R_{i,n_i})$ is the n_i -tuple of right creation operators (see Section 1). Regard $M_m(\mathcal{R}_n^* \mathcal{R}_n)$ as a subspace of $M_m(B(\bigotimes_{i=1}^k F^2(H_{n_i})))$. Let $M_m(\mathcal{R}_n^* \mathcal{R}_n)$ have the norm structure that it inherits from the (unique) norm structure on the C^* -algebra $M_m(B(\bigotimes_{i=1}^k F^2(H_{n_i})))$. We remark that

$$\mathcal{R}_n^* \mathcal{R}_n = \text{span}\{R_\alpha^* R_\beta : (\alpha, \beta) \in \Lambda\} = \text{span}\{R_{\tilde{\alpha}}^* R_{\tilde{\beta}} : (\alpha, \beta) \in \Lambda\},$$

where $\Lambda = \tilde{\Lambda}$ is given by (4-1). If $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ is a completely bounded linear map, then there exists a unique completely bounded linear map

$$\hat{\mu} := \mu \otimes \text{id} : \overline{\mathcal{R}_n^* \mathcal{R}_n}^{\|\cdot\|} \otimes_{\min} B(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$$

such that

$$\hat{\mu}(A \otimes Y) = \mu(A) \otimes Y, \quad A \in \mathcal{R}_n^* \mathcal{R}_n, Y \in B(\mathcal{H}).$$

Moreover, $\|\hat{\mu}\|_{cb} = \|\mu\|_{cb}$ and, if μ is completely positive, then so is $\hat{\mu}$.

We define the *free pluriharmonic Poisson kernel* by setting

$$\mathcal{P}(R, X) := \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} R_{1, \tilde{\alpha}_1}^* \dots R_{k, \tilde{\alpha}_k}^* R_{1, \tilde{\beta}_1} R_{k, \tilde{\beta}_k} \otimes X_{1, \alpha_1} \dots X_{k, \alpha_k} X_{1, \beta_1}^* \dots X_{k, \beta_k}^*$$

for any $X \in \mathbf{B}_n(\mathcal{H})$, where the convergence is in the operator norm topology. We need to show that the latter convergence holds. Indeed, note that, for each $i \in \{1, \dots, k\}$ and $r \in [0, 1)$, we have

$$\begin{aligned} W_i &:= \sum_{m_i \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+ \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mathbf{R}_{i, \tilde{\alpha}_i}^* \mathbf{R}_{i, \tilde{\beta}_i} \otimes r^{|\alpha_i| + |\beta_i|} \mathbf{S}_{i, \alpha_i} \mathbf{S}_{i, \beta_i}^* \\ &= \lim_{p_i \rightarrow \infty} \left(\sum_{\substack{\alpha_i \in \mathbb{F}_{n_i} \\ 0 < |\alpha_i| \leq p_i}} \mathbf{R}_{i, \tilde{\alpha}_i}^* \otimes r^{|\alpha_i|} \mathbf{S}_{i, \alpha_i} + \sum_{\substack{\beta_i \in \mathbb{F}_{n_i} \\ 0 \leq |\beta_i| \leq p_i}} \mathbf{R}_{i, \tilde{\beta}_i} \otimes r^{|\beta_i|} \mathbf{S}_{i, \beta_i}^* \right), \end{aligned}$$

where the limit is in the operator norm topology. One can easily see that

$$\begin{aligned} W_1 \cdots W_k &= \mathcal{P}(\mathbf{R}, r\mathbf{S}) \\ &:= \lim_{p_1 \rightarrow \infty} \cdots \lim_{p_k \rightarrow \infty} \sum_{\substack{m_1 \in \mathbb{Z} \\ |m_1| \leq p_1}} \cdots \sum_{\substack{m_k \in \mathbb{Z} \\ |m_k| \leq p_k}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mathbf{R}_{1, \tilde{\alpha}_1}^* \cdots \mathbf{R}_{k, \tilde{\alpha}_k}^* \mathbf{R}_{1, \tilde{\beta}_1} \cdots \mathbf{R}_{k, \tilde{\beta}_k} \\ &\quad \otimes r^{\sum_{i=1}^k (|\alpha_i| + |\beta_i|)} \mathbf{S}_{1, \alpha_1} \cdots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \cdots \mathbf{S}_{k, \beta_k}^*. \end{aligned}$$

Therefore, the series defining $\mathcal{P}(\mathbf{R}, r\mathbf{S})$, i.e.,

$$\sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mathbf{R}_{1, \tilde{\alpha}_1}^* \cdots \mathbf{R}_{k, \tilde{\alpha}_k}^* \mathbf{R}_{1, \tilde{\beta}_1} \cdots \mathbf{R}_{k, \tilde{\beta}_k} \otimes r^{\sum_{i=1}^k (|\alpha_i| + |\beta_i|)} \mathbf{S}_{1, \alpha_1} \cdots \mathbf{S}_{k, \alpha_k} \mathbf{S}_{1, \beta_1}^* \cdots \mathbf{S}_{k, \beta_k}^*$$

are convergent in the operator norm topology. We remark that, due to the fact that the operators W_1, \dots, W_k commute, the order of the limits above is irrelevant. Fix $X \in \mathbf{B}_n(\mathcal{H})$ and let $r \in (0, 1)$ be such that $(1/r)X$ is in $\mathbf{B}_n(\mathcal{H})$. Since the noncommutative Berezin transform $\mathcal{B}_{(1/r)X}$ is continuous in the operator norm, so is $\text{id} \otimes \mathcal{B}_{(1/r)X}$. Consequently, applying $\text{id} \otimes \mathcal{B}_{(1/r)X}$ to the relation above, we deduce that

$$\begin{aligned} &(\text{id} \otimes \mathcal{B}_{(1/r)X})[\mathcal{P}(\mathbf{R}, r\mathbf{S})] \\ &= \lim_{p_1 \rightarrow \infty} \cdots \lim_{p_k \rightarrow \infty} \sum_{\substack{m_1 \in \mathbb{Z} \\ |m_1| \leq p_1}} \cdots \sum_{\substack{m_k \in \mathbb{Z} \\ |m_k| \leq p_k}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mathbf{R}_{1, \tilde{\alpha}_1}^* \cdots \mathbf{R}_{k, \tilde{\alpha}_k}^* \mathbf{R}_{1, \tilde{\beta}_1} \cdots \mathbf{R}_{k, \tilde{\beta}_k} \\ &\quad \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*, \end{aligned}$$

where the limits are in the operator norm topology. This proves our assertion. Now, we introduce the *noncommutative Poisson transform* of a completely bounded linear map $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ to be the map $\mathcal{P}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ defined by

$$(\mathcal{P}\mu)(X) := \hat{\mu}[\mathcal{P}(\mathbf{R}, X)], \quad X \in \mathbf{B}_n(\mathcal{H}).$$

The next result contains some of the basic properties of the noncommutative Poisson kernel and the noncommutative Poisson transform.

Theorem 4.2. *Let $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ be a completely bounded linear map.*

(i) The map $X \mapsto \mathcal{P}(\mathbf{R}, X)$ is a positive k -pluriharmonic function on the polyball \mathbf{B}_n , with coefficients in $B(\otimes_{i=1}^k F^2(H_{n_i}))$, and has the factorization $\mathcal{P}(\mathbf{R}, X) = C_X^* C_X$, where

$$C_X := (I_{\otimes_{i=1}^k F^2(H_{n_i})} \otimes \Delta_X(I)^{1/2}) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1}.$$

(ii) The noncommutative Poisson transform $\mathcal{P}\mu$ is a free k -pluriharmonic function on the regular polyball \mathbf{B}_n that coincides with the Berezin transform $\mathcal{B}_\mu(I, \cdot)$.

(iii) If μ is a completely positive linear map, then $\mathcal{P}\mu$ is a positive free k -pluriharmonic function on \mathbf{B}_n .

Proof. The fact that $X \mapsto \mathcal{P}(\mathbf{R}, X)$ is a free k -pluriharmonic function on the polyball \mathbf{B}_n with coefficients in $\mathcal{R}_n^* \mathcal{R}_n$ was proved in the remarks preceding the theorem. Setting $\Lambda_i := \mathbf{R}_{i,1} \otimes r S_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes r S_{i,n_i}^*$ for each $i \in \{1, \dots, k\}$, we have

$$\begin{aligned} W_i &:= \sum_{m_i \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+ \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mathbf{R}_{i,\tilde{\alpha}_i}^* \mathbf{R}_{i,\tilde{\beta}_i} \otimes r^{|\alpha_i| + |\beta_i|} S_{i,\alpha_i} S_{i,\beta_i}^* \\ &= (I - \Lambda_i)^{-1} - I + (I - \Lambda_i^*)^{-1} \\ &= (I - \Lambda_i^*)^{-1} [(I - \Lambda_i) - (I - \Lambda_i^*)(I - \Lambda_i) + (I - \Lambda_i^*)] (I - \Lambda_i)^{-1} \\ &= (I - \Lambda_i^*)^{-1} \left[I_{\otimes_{i=1}^k F^2(H_{n_i})} \otimes \left(I_{\otimes_{i=1}^k F^2(H_{n_i})} - \sum_{j=1}^{n_i} r^2 S_{i,j} S_{i,j}^* \right) \right] (I - \Lambda_i)^{-1}. \end{aligned}$$

Recall that $\mathbf{R}_{i,s} \mathbf{R}_{j,t} = \mathbf{R}_{j,t} \mathbf{R}_{i,s}$ and $\mathbf{R}_{i,s} \mathbf{R}_{j,t}^* = \mathbf{R}_{j,t}^* \mathbf{R}_{i,s}$ for any $i, j \in \{1, \dots, k\}$ with $i \neq j$ and for any $s \in \{1, \dots, n_i\}$ and $t \in \{1, \dots, n_j\}$. Similar commutation relations hold for the universal model \mathcal{S} . Since $\mathcal{P}(\mathbf{R}, r\mathcal{S}) = W_1 \cdots W_k$ and W_1, \dots, W_k are commuting positive operators, we deduce that

$$\begin{aligned} \mathcal{P}(\mathbf{R}, r\mathcal{S}) &= \left(\prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes r S_{i,1} - \cdots - \mathbf{R}_{i,n_i}^* \otimes r S_{i,n_i})^{-1} \right) \\ &\quad \times (I \otimes \Delta_{r\mathcal{S}}(I)) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes r S_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes r S_{i,n_i}^*)^{-1} \end{aligned}$$

for any $r \in [0, 1)$, and $\mathcal{P}(\mathbf{R}, r\mathcal{S}) = C_{r\mathcal{S}}^* C_{r\mathcal{S}} \geq 0$. Now, let $X \in \mathbf{B}_n(\mathcal{H})$ and let $r \in (0, 1)$ be such that $(1/r)X \in \mathbf{B}_n(\mathcal{H})$. Since the noncommutative Berezin transform $\mathcal{B}_{(1/r)X}$ is continuous in the operator norm and completely positive, so is $\text{id} \otimes \mathcal{B}_{(1/r)X}$. Consequently, applying $\text{id} \otimes \mathcal{B}_{(1/r)X}$ to the relations above, we deduce that

$$\begin{aligned} \mathcal{P}(\mathbf{R}, X) &= (\text{id} \otimes \mathcal{B}_{(1/r)X})[\mathcal{P}(\mathbf{R}, r\mathcal{S})] \\ &= \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \cdots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} (I \otimes \Delta_X(I)) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1} \\ &= C_X^* C_X, \end{aligned}$$

which completes the proof of (i).

Using the results above and the continuity of $\hat{\mu}$ in the operator norm, we deduce that the noncommutative Berezin transform $\mathcal{B}_\mu(I, \cdot)$ associated with μ coincides with the Poisson transform $\mathcal{P}\mu$. Indeed, we have

$$\begin{aligned} \mathcal{B}_\mu(I, X) &= \hat{\mu}(C_X^* C_X) \\ &= \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mu(\mathbf{R}_{1, \tilde{\alpha}_1}^* \cdots \mathbf{R}_{k, \tilde{\alpha}_k}^* \mathbf{R}_{1, \tilde{\beta}_1} \cdots \mathbf{R}_{k, \tilde{\beta}_k}) \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^* \\ &= \hat{\mu}(\mathcal{P}(\mathbf{R}, X)) \\ &= (\mathcal{P}\mu)(X) \end{aligned}$$

for any $X \in \mathbf{B}_n(\mathcal{H})$, where the convergence is in the operator norm topology of $B(\mathcal{K} \otimes \mathcal{H})$. This proves (ii). Note also that the Poisson transform $\mathcal{P}\mu$ is a free k -pluriharmonic function on \mathbf{B}_n with coefficients in $B(\mathcal{E})$. If μ is completely positive, then so is $\hat{\mu}$. Using the fact that $\hat{\mu}(C_X^* C_X) = (\mathcal{P}\mu)(X)$, we deduce (iii). \square

Consider the particular case when $n_1 = \cdots = n_k = 1$, $\mathcal{H} = \mathcal{K} = \mathbb{C}$, $\mathbf{X} = (X_1, \dots, X_k)$ with $X_j = r_j e^{i\theta_j} \in \mathbb{D}$, and μ is a complex Borel measure on \mathbb{T}^k . Note that μ can be seen as a bounded linear functional on $C(\mathbb{T}^k)$. Consequently, there is a unique bounded linear functional $\hat{\mu}$ on the operator system generated by the monomials $S_1^{m_1^-} \cdots S_k^{m_k^-} S_1^{*m_1^+} \cdots S_k^{*m_k^+}$, where $m_1, \dots, m_k \in \mathbb{Z}$, and S_1, \dots, S_k are the unilateral shifts acting on the Hardy space $H^2(\mathbb{T}^k)$, such that

$$\hat{\mu}(S_1^{m_1^-} \cdots S_k^{m_k^-} S_1^{*m_1^+} \cdots S_k^{*m_k^+}) = \mu(e^{im_1^- \varphi_1} \cdots e^{im_k^- \varphi_k} e^{-im_1^+ \varphi_1} \cdots e^{-im_k^+ \varphi_k}), \quad m_1, \dots, m_k \in \mathbb{Z}.$$

Indeed, if p is any polynomial function of the form

$$p(z_1, \dots, z_k, \bar{z}_1, \dots, \bar{z}_k) = \sum a_{(m_1, \dots, m_k)} z_1^{m_1^-} \cdots z_k^{m_k^-} \bar{z}_1^{m_1^+} \cdots \bar{z}_k^{m_k^+}, \quad (z_1, \dots, z_k) \in \mathbb{D}^k,$$

where $a_{(m_1, \dots, m_k)} \in \mathbb{C}$, then, due to the noncommutative von Neumann inequality [Popescu 1999], we have

$$\begin{aligned} |\hat{\mu}(p(S_1, \dots, S_k, S_1^*, \dots, S_k^*))| &= |\mu(p(e^{i\varphi_1}, \dots, e^{i\varphi_k}, e^{-i\varphi_1}, \dots, e^{-i\varphi_k}))| \\ &\leq \|\mu\| \|p(S_1, \dots, S_k, S_1^*, \dots, S_k^*)\|. \end{aligned}$$

Therefore, $\hat{\mu}$ is a bounded linear functional on the operator system $\text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{\|\cdot\|}$. Note that the noncommutative Poisson transform of $\hat{\mu}$, i.e., $B_{\hat{\mu}}(I, \cdot)$, coincides with the classical Poisson transform of μ . Indeed, for any $z = (r_1 e^{i\theta_1}, \dots, r_k e^{i\theta_k}) \in \mathbb{D}^k$, we have

$$\begin{aligned} \mathcal{B}_{\hat{\mu}}(I, z) &= \sum_{(p_1, \dots, p_k) \in \mathbb{Z}^k} \hat{\mu}(S_1^{p_1^-} \cdots p_k^{p_k^-} S_1^{*p_1^+} \cdots S_k^{*p_k^+}) z_1^{p_1} \cdots z_k^{p_k} \\ &= \sum_{(p_1, \dots, p_k) \in \mathbb{Z}^k} \mu(\bar{\xi}_1^{p_1} \cdots \bar{\xi}^{p_k}) z_1^{p_1} \cdots z_k^{p_k} \\ &= \sum_{(p_1, \dots, p_k) \in \mathbb{Z}^k} \left(\int_{\mathbb{T}^k} \bar{\xi}_1^{p_1} \cdots \bar{\xi}^{p_k} d\mu(\xi) \right) z_1^{p_1} \cdots z_k^{p_k} \\ &= \int_{\mathbb{T}^k} \left(\sum_{(p_1, \dots, p_k) \in \mathbb{Z}^k} r_1^{|p_1|} \cdots r_k^{|p_k|} e^{ip_1(\theta_1 - \varphi_1)} \cdots e^{ip_k(\theta_k - \varphi_k)} \right) d\mu(\xi) = \int_{\mathbb{T}^k} P(z, \xi) d\mu(\xi), \end{aligned}$$

where

$$P(z, \zeta) = P_{r_1}(\theta_1 - \varphi_1) \cdots P_{r_k}(\theta_k - \varphi_k), \quad \zeta = (e^{i\varphi_1}, \dots, e^{i\varphi_k}) \in \mathbb{T}^k,$$

and $P_r(\theta - \varphi) = (1 - r^2)/(1 - 2r \cos(\theta - \varphi) + r^2)$ is the Poisson kernel of the unit disc (see [Rudin 1969]).

We recall that Λ denotes the set of all pairs $(\alpha, \beta) \in \mathbf{F}_n^+ \times \mathbf{F}_n^+$, where $\mathbf{F}_n^+ := \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, with the property that $\alpha \sim_{lc} \beta$ and $(\alpha, \beta) = (c_l^+(\alpha, \beta), c_l^-(\alpha, \beta))$. We remark that $(\alpha, \beta) \in \Lambda$ if and only if $(\tilde{\alpha}, \tilde{\beta}) \in \Lambda$. As before, we use the notation $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_k)$ if $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{F}_n^+$.

Throughout the rest of this section, we assume that \mathcal{E} is a separable Hilbert space.

Lemma 4.3. *Let $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ be a completely bounded linear map. For each $r \in [0, 1)$, define the linear map $\mu_r : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ by*

$$\mu_r(\mathbf{R}_\alpha^* \mathbf{R}_\beta) := r^{|\alpha|+|\beta|} \mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta), \quad (\alpha, \beta) \in \Lambda,$$

where $|\alpha| := |\alpha_1| + \cdots + |\alpha_k|$ if $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{F}_n^+$. Then

- (i) μ_r is a completely bounded linear map;
- (ii) $\|\mu\|_{cb} = \sup_{0 \leq r < 1} \|\mu_r\|_{cb} = \lim_{r \rightarrow 1} \|\mu_r\|_{cb}$;
- (iii) $\mu_r(A) \rightarrow \mu(A)$ in the operator norm topology as $r \rightarrow 1$ for any $A \in \mathcal{R}_n^* \mathcal{R}_n$;
- (iv) if μ is completely positive, then so is μ_r for any $r \in [0, 1)$.

Proof. Let

$$p(\mathbf{R}^*, \mathbf{R}) := \sum_{\substack{(\alpha, \beta) \in \Lambda' \subset \Lambda \\ \text{card}(\Lambda') < \aleph_0}} a_{(\alpha, \beta)} \mathbf{R}_\alpha^* \mathbf{R}_\beta, \quad a_{(\alpha, \beta)} \in \mathbb{C},$$

and $0 \leq r_1 < r_2 \leq 1$. Using the noncommutative von Neumann inequality [Popescu 1999], we deduce that

$$\|\mu_{r_1}(p(\mathbf{R}^*, \mathbf{R}))\| = \|\mu(p(r_1 \mathbf{R}^*, r_1 \mathbf{R}))\| = \left\| \mu_{r_2} \left(p \left(\frac{r_1}{r_2} \mathbf{R}^*, \frac{r_1}{r_2} \mathbf{R} \right) \right) \right\| \leq \|\mu_{r_2}\| \|p(\mathbf{R}^*, \mathbf{R})\|.$$

In particular, we have $\|\mu_r\| \leq \|\mu\|$ for any $r \in [0, 1)$. Similarly, passing to matrices over $\mathcal{R}_n^* \mathcal{R}_n$, one can show that $\|\mu_{r_1}\|_{cb} \leq \|\mu_{r_2}\|_{cb}$ if $0 \leq r_1 < r_2 \leq 1$, and $\|\mu_r\|_{cb} \leq \|\mu\|_{cb}$ for any $r \in [0, 1)$. Now, one can easily see that $\mu_r(A) \rightarrow \mu(A)$ in the operator norm topology as $r \rightarrow 1$ for any $A \in \mathcal{R}_n^* \mathcal{R}_n$, and $\|\mu\|_{cb} = \sup_{0 \leq r < 1} \|\mu_r\|_{cb}$. Hence, and using the fact that the function $r \mapsto \|\mu_r\|_{cb}$ is increasing for $r \in [0, 1)$, we deduce that $\lim_{r \rightarrow 1} \|\mu_r\|_{cb}$ exists and it is equal to $\|\mu\|_{cb}$.

To prove (iv), note that $\mu_r(p(\mathbf{R}^*, \mathbf{R})) = \mu(\mathcal{B}_{r, \mathbf{R}}[p(\mathbf{S}^*, \mathbf{S})])$. Since the noncommutative Berezin transform $\mathcal{B}_{r, \mathbf{R}}$ and μ are completely positive linear maps and $p(\mathbf{R}^*, \mathbf{R})$ is unitarily equivalent to $p(\mathbf{S}^*, \mathbf{S})$, we deduce that μ_r is a completely positive linear map for each $r \in [0, 1)$. □

Let F be a free k -pluriharmonic function on the polyball \mathbf{B}_n , with operator-valued coefficients in $B(\mathcal{E})$, with representation

$$F(\mathbf{X}) = \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} \otimes \mathbf{X}_{1, \alpha_1} \cdots \mathbf{X}_{k, \alpha_k} \mathbf{X}_{1, \beta_1}^* \cdots \mathbf{X}_{k, \beta_k}^*.$$

We associate to F and each $r \in [0, 1)$ the linear map $\nu_{F_r} : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ by setting

$$\nu_{F_r}(\mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}}) := r^{|\alpha|+|\beta|} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)}, \quad (\alpha, \beta) \in \Lambda. \tag{4-2}$$

We remark that ν_{F_r} is uniquely determined by the radial function $r \mapsto F(r\mathbf{S})$. Indeed, note that, if $x := x_1 \otimes \dots \otimes x_k, y = y_1 \otimes \dots \otimes y_k$ satisfy (1-3) and $h, \ell \in \mathcal{E}$, we have

$$\langle F(r\mathbf{S})(h \otimes x), \ell \otimes y \rangle = \langle r^{|\alpha|+|\beta|} A_{(\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_k)} h, \ell \rangle = \langle \nu_{F_r}(\mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}}) h, \ell \rangle, \quad (\alpha, \beta) \in \Lambda.$$

In what follows, we denote by $C^*(\mathbf{R})$ the C^* -algebra generated by the right creation operators $\mathbf{R}_{i,j}$, where $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$.

Theorem 4.4. *Let $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ be a free k -pluriharmonic function. Then the following statements are equivalent:*

- (i) *There exists a completely bounded linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ such that $F = \mathcal{P}\mu$.*
- (ii) *The linear maps $\{\nu_{F_r}\}_{r \in [0,1)}$ associated with F are completely bounded and $\sup_{0 \leq r < 1} \|\nu_{F_r}\|_{\text{cb}} < \infty$.*
- (iii) *There exist a k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, of doubly commuting row isometries acting on a Hilbert space \mathcal{K} and bounded linear operators $W_1, W_2 : \mathcal{E} \rightarrow \mathcal{K}$ such that*

$$F(\mathbf{X}) = (W_1^* \otimes I)[C_{\mathbf{X}}(\mathbf{V})^* C_{\mathbf{X}}(\mathbf{V})](W_2 \otimes I),$$

where

$$C_{\mathbf{X}}(\mathbf{V}) := (I \otimes \Delta_{\mathbf{X}}(I)^{1/2}) \prod_{i=1}^k (I - V_{i,1} \otimes X_{i,1}^* - \dots - V_{i,n_i} \otimes X_{i,n_i}^*)^{-1}.$$

Moreover, in this case we can choose μ such that $\|\mu\|_{\text{cb}} = \sup_{0 \leq r < 1} \|\nu_{F_r}\|_{\text{cb}}$.

Proof. Assume that (i) holds. Then

$$F(\mathbf{X}) = \sum_{m_1 \in \mathbb{Z}} \dots \sum_{m_k \in \mathbb{Z}} \sum_{\substack{\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+, i \in \{1, \dots, k\} \\ |\alpha_i| = m_i^-, |\beta_i| = m_i^+}} \mu(\mathbf{R}_{1, \tilde{\alpha}_1}^* \dots \mathbf{R}_{k, \tilde{\alpha}_k}^* \mathbf{R}_{1, \tilde{\beta}_1} \dots \mathbf{R}_{k, \tilde{\beta}_k}) \otimes X_{1, \alpha_1} \dots X_{k, \alpha_k} X_{1, \beta_1}^* \dots X_{k, \beta_k}^*$$

for any $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$, where the convergence is in the operator norm topology. Set $A_{(\alpha; \beta)} := \mu(\mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}})$ for any $(\alpha, \beta) \in \Lambda$. Consequently, for each $r \in [0, 1)$, we have

$$\nu_{F_r}(\mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}}) := r^{|\alpha|+|\beta|} \mu(\mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}}), \quad (\alpha, \beta) \in \Lambda.$$

We recall that $(\alpha, \beta) \in \Lambda$ if and only if $(\tilde{\alpha}, \tilde{\beta}) \in \Lambda$. Applying Lemma 4.3, we deduce that $\{\nu_{F_r}\}$ is a completely bounded map and

$$\|\mu|_{\mathcal{R}_n^* \mathcal{R}_n}\|_{\text{cb}} = \sup_{0 \leq r < 1} \|\nu_{F_r}\|_{\text{cb}} < \infty,$$

which proves that (i) implies (ii).

Now, we prove that (ii) implies (i). Assume that F is a free pluriharmonic function on \mathcal{B}_n with coefficients in $B(\mathcal{E})$ and such that condition (ii) holds. Let $\{q_j\}$ be a countable dense subset of $\mathcal{R}_n^* \mathcal{R}_n$. For instance, we can consider all the operators of the form

$$p(\mathbf{R}^*, \mathbf{R}) := \sum_{(\alpha, \beta) \in \Lambda: |\alpha| \leq m, |\beta| \leq m} a_{(\alpha, \beta)} \mathbf{R}_\alpha^* \mathbf{R}_\beta,$$

where $m \in \mathbb{N}$ and the coefficients $a_{(\alpha, \beta)}$ lie in some countable dense subset of the complex plane. For each j , we have $\|v_{F_r}(q_j)\| \leq M\|q_j\|$ for any $r \in [0, 1)$, where $M := \sup_{0 \leq r < 1} \|v_{F_r}\|_{cb}$.

Due to the Banach–Alaoglu theorem [Douglas 1998], the ball $[B(\mathcal{E})]_M^-$ is compact in the w^* -topology. Since \mathcal{E} is a separable Hilbert space, $[B(\mathcal{E})]_M^-$ is a metric space in the w^* -topology which coincides with the weak operator topology on $[B(\mathcal{E})]_M^-$. Consequently, the diagonal process guarantees the existence of a sequence $\{r_m\}_{m=1}^\infty$ such that $r_m \rightarrow 1$ and $\text{WOT-}\lim_{m \rightarrow 1} v_{F_{r_m}}(q_j)$ exists for each q_j . Fix $A \in \mathcal{R}_n^* \mathcal{R}_n$ and $x, y \in \mathcal{E}$ and let us prove that $\{\langle v_{F_{r_m}}(A)x, y \rangle\}_{m=1}^\infty$ is a Cauchy sequence. Let $\epsilon > 0$ and choose q_j so that $\|q_j - A\| < \epsilon/(3M\|x\|\|y\|)$. Now we choose N so that $|\langle (v_{F_{r_m}}(q_j) - v_{F_{r_k}}(q_j))x, y \rangle| < \frac{1}{3}\epsilon$ for any $m, k > N$. Due to the fact that

$$\begin{aligned} |\langle (v_{F_{r_m}}(A) - v_{F_{r_k}}(A))x, y \rangle| &\leq |\langle v_{F_{r_m}}(A - q_j)x, y \rangle| + |\langle (v_{F_{r_m}}(q_j) - v_{F_{r_k}}(q_j))x, y \rangle| + |\langle v_{F_{r_k}}(q_j - A)x, y \rangle| \\ &\leq 2M\|x\|\|y\|\|A - q_j\| + |\langle (v_{F_{r_m}}(q_j) - v_{F_{r_k}}(q_j))x, y \rangle|, \end{aligned}$$

we deduce that $|\langle (v_{F_{r_m}}(A) - v_{F_{r_k}}(A))x, y \rangle| < \epsilon$ for $m, k > N$. Therefore,

$$\Phi(x, y) := \lim_{m \rightarrow \infty} \langle v_{F_{r_m}}(A)x, y \rangle$$

exists for any $x, y \in \mathcal{E}$ and defines a functional $\Phi : \mathcal{E} \times \mathcal{E} \rightarrow \mathbb{C}$ which is linear in the first variable and conjugate linear in the second. Moreover, we have $|\Phi(x, y)| \leq M\|A\|\|x\|\|y\|$ for any $x, y \in \mathcal{E}$. Due to the Riesz representation theorem, there exists a unique bounded linear operator $B(\mathcal{E})$, which we denote by $\nu(A)$, such that $\Phi(x, y) = \langle \nu(A)x, y \rangle$ for $x, y \in \mathcal{E}$. Therefore,

$$\nu(A) = \text{WOT-}\lim_{r_m \rightarrow 1} v_{F_{r_m}}(A), \quad A \in \mathcal{R}_n^* \mathcal{R}_n,$$

and $\|\nu(A)\| \leq M\|A\|$. Note that $\nu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ is a completely bounded map. Indeed, if $[A_{ij}]_m$ is an $m \times m$ matrix over $\mathcal{R}_n^* \mathcal{R}_n$, then $[\nu(A_{ij})]_m = \text{WOT-}\lim_{r_k \rightarrow 1} [v_{F_{r_k}}(A_{ij})]_m$. Hence, $\|[\nu(A_{ij})]_m\| \leq M\|[A_{ij}]_m\|$ for all m , and so $\|\nu\|_{cb} \leq M$. Note also that $\nu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = A_{(\alpha; \beta)}$ for any $(\alpha, \beta) \in \Lambda$, where $A_{(\alpha; \beta)}$ are the coefficients of F . By Wittstock’s extension theorem [1981; 1984], there exists a completely bounded linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ which extends ν such that $\|\mu\|_{cb} = \|\nu\|_{cb}$. Since $F = \mathcal{P}\mu$, the proof of (i) is complete.

Now, we prove the equivalence of (i) with (iii). If (i) holds, then according to Theorem 8.4 from [Paulsen 1986] there exist a Hilbert space \mathcal{K} , a $*$ -representation $\pi : C^*(\mathbf{R}) \rightarrow B(\mathcal{K})$, and bounded operators $W_1, W_2 : \mathcal{E} \rightarrow \mathcal{K}$ with $\|\mu\| = \|W_1\|\|W_2\|$ such that

$$\mu(A) = W_1^* \pi(A) W_2, \quad A \in C^*(\mathbf{R}). \tag{4-3}$$

Set $V_{i,j} := \pi(\mathbf{R}_{i,j})$ for $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$ and note that $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, is a k -tuple of doubly commuting row isometries. Using [Theorem 4.2](#), one can easily see that the equality $F = \mathcal{P}\mu$ implies the one from (iii). Now, we prove that (iii) implies (i). Since the k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, consists of doubly commuting row isometries on a Hilbert space \mathcal{K} , the noncommutative von Neumann inequality [[Popescu 1999](#)] implies that the map $\pi : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ defined by

$$\pi(\mathbf{R}_\alpha \mathbf{R}_\beta^*) := V_\alpha V_\beta^*, \quad \alpha, \beta \in \mathbf{F}_n^+,$$

is a $*$ -representation of $C^*(\mathbf{R})$. Define $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ by setting $\mu(A) := W_1^* \pi(A) W_2$, $A \in C^*(\mathbf{R})$, and note that μ is a completely bounded linear map. Using the relation

$$F(\mathbf{X}) = (W_1^* \otimes I)[C_{\mathbf{X}}(\mathbf{V})^* C_{\mathbf{X}}(\mathbf{V})](W_2 \otimes I)$$

and the factorization $P(\mathbf{V}, \mathbf{X}) = C_{\mathbf{X}}(\mathbf{V})^* C_{\mathbf{X}}(\mathbf{V})$ (see also [Theorem 4.2](#)), we deduce that $F(\mathbf{X}) = \mathcal{P}\mu(\mathbf{X})$ for $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$. □

We introduce the space $\mathbf{PH}^1(\mathbf{B}_n)$ of all free k -pluriharmonic functions F on \mathbf{B}_n such that the linear maps $\{\nu_{F_r}\}_{r \in [0,1]}$ associated with F are completely bounded and set $\|F\|_1 := \sup_{0 \leq r < 1} \|\nu_{F_r}\|_{\text{cb}} < \infty$. As a consequence of [Theorem 4.4](#), one can see that $\|\cdot\|_1$ is a norm on $\mathbf{PH}^1(\mathbf{B}_n)$ and $(\mathbf{PH}^1(\mathbf{B}_n), \|\cdot\|_1)$ is a Banach space that can be identified with the Banach space $\text{CB}(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$ of all completely bounded linear maps from $\mathcal{R}_n^* \mathcal{R}_n$ to $B(\mathcal{E})$.

Corollary 4.5. *Let $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ be a free k -pluriharmonic function. Then the following statements are equivalent:*

- (i) *There exists a completely positive linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ such that $F = \mathcal{P}\mu$.*
- (ii) *The linear maps $\{\nu_{F_r}\}_{r \in [0,1]}$ associated with F are completely positive.*
- (iii) *There exist a k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, of doubly commuting row isometries acting on a Hilbert space $\mathcal{K} \supset \mathcal{E}$ and a bounded operator $W : \mathcal{E} \rightarrow \mathcal{K}$ such that*

$$F(\mathbf{X}) = (W^* \otimes I)[C_{\mathbf{X}}(\mathbf{V})^* C_{\mathbf{X}}(\mathbf{V})](W \otimes I).$$

Proof. The proof is similar to that of [Theorem 4.4](#). Note that for (i) implies (ii) we have to use [Lemma 4.3\(iv\)](#). For the converse, note that if ν_{F_r} , $r \in [0, 1)$, are completely positive linear maps then

$$\|\nu_{F_r}\|_{\text{cb}} = \|\nu_{F_r}(1)\| = \|\nu_{F_r}\| = \|A_{(\mathbf{g}; \mathbf{g})}\|,$$

where $\mathbf{g} = (g_0^1, \dots, g_0^k)$ is the identity element in \mathbf{F}_n^+ . As in the proof of [Theorem 4.4](#), we find a completely bounded map $\nu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ such that

$$\nu(A) = \text{WOT-}\lim_{r_m \rightarrow 1} \nu_{F_{r_m}}(A), \quad A \in \mathcal{R}_n^* \mathcal{R}_n.$$

Since ν_{F_r} , $r \in [0, 1)$, are completely positive linear maps, one can easily see that ν is completely positive. Using Arveson’s extension theorem [[1969](#)], we find a completely positive map $\mu : C^*(\mathbf{R}) \rightarrow \mathbb{C}$ which extends ν and such that $\|\mu\|_{\text{cb}} = \|\nu\|_{\text{cb}}$. We also have that $F = \mathcal{P}\mu$. Now, the proof that (iii) is equivalent

to (i) uses Stinespring’s representation theorem [1955] and is similar to the same equivalence from Theorem 4.4. We leave it to the reader. \square

An open question remains. Is any positive free k -pluriharmonic function on the regular polyball B_n the Poisson transform of a completely positive linear map? The answer is positive if $k = 1$ (see [Popescu 2009]) and also when $n_1 = \dots = n_k$ (see Section 3).

5. Herglotz–Riesz representations for free holomorphic functions with positive real parts

In this section, we introduce the noncommutative Herglotz–Riesz transform of a completely positive linear map $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ and obtain Herglotz–Riesz representation theorems for free holomorphic functions with positive real parts in regular polyballs.

Define the space

$$\mathbf{RH}(B_n) := \text{span}\{\Re f : f \in \text{Hol}_{\mathcal{E}}(B_n)\},$$

where $\text{Hol}_{\mathcal{E}}(B_n)$ is the set of all free holomorphic functions in the polyball B_n with coefficients in $B(\mathcal{E})$. Let $\tau : B(\bigotimes_{i=1}^k F^2(H_{n_i})) \rightarrow \mathbb{C}$ be the bounded linear functional defined by $\tau(A) = \langle A1, 1 \rangle$. We remark that the radial function associated with $\varphi \in \mathbf{RH}(B_n)$, i.e., $r \mapsto \varphi(r\mathbf{R})$ for $r \in [0, 1)$, uniquely determines the family $\{\nu_{\varphi_r}\}_{r \in [0,1)}$ of linear maps $\nu_{\varphi_r} : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ defined by (4-2). Indeed, note that

$$\begin{aligned} \nu_{\varphi_r}(\mathbf{R}_{\tilde{\alpha}}^*) &:= (\text{id} \otimes \tau)[(I \otimes \mathbf{R}_{\tilde{\alpha}}^*)\varphi(r\mathbf{R})], \\ \nu_{\varphi_r}(\mathbf{R}_{\tilde{\alpha}}) &:= (\text{id} \otimes \tau)[\varphi(r\mathbf{R})(I \otimes \mathbf{R}_{\tilde{\alpha}})], \end{aligned}$$

for any $\alpha = (\alpha_1, \dots, \alpha_k) \in F_n^+ := \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+$, where $\tilde{\alpha} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_k)$ and $\mathbf{R}_{\alpha} := \mathbf{R}_{1,\alpha_1} \dots \mathbf{R}_{k,\alpha_k}$, and $\nu_{\varphi_r}(\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}) = 0$ if $\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}$ is different from \mathbf{R}_{γ} or \mathbf{R}_{γ}^* for some $\gamma \in F_n^+$. Consider the space

$$\mathbf{RH}^1(B_n) := \{\varphi \in \mathbf{RH}(B_n) : \nu_{\varphi_r} \text{ is completely bounded and } \sup_{0 \leq r < 1} \|\nu_{\varphi_r}\|_{\text{cb}} < \infty\}.$$

If $\varphi \in \mathbf{RH}^1(B_n)$, we define $\|\varphi\|_1 := \sup_{0 \leq r < 1} \|\nu_{\varphi_r}\|_{\text{cb}}$. Denote by $\text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$ the space of all completely bounded linear maps $\lambda : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ such that $\lambda(\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}) = 0$ if $\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}$ is not equal to \mathbf{R}_{γ} or \mathbf{R}_{γ}^* for some $\gamma \in F_n^+$.

Theorem 5.1. $(\mathbf{RH}^1(B_n), \|\cdot\|_1)$ is a Banach space which can be identified with the Banach space $\text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$. Moreover, if $\varphi : B_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ is a function, then the following statements are equivalent:

- (i) φ is in $\mathbf{RH}^1(B_n)$.
- (ii) There is a unique completely bounded linear map $\mu_{\varphi} \in \text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$ such that $\varphi = \mathcal{P}\mu_{\varphi}$.
- (iii) There exist a k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, of doubly commuting row isometries on a Hilbert space \mathcal{K} and bounded linear operators $W_1, W_2 : \mathcal{E} \rightarrow \mathcal{K}$ such that

$$\varphi(\mathbf{X}) = (W_1^* \otimes I)[C_X(\mathbf{V})^* C_X(\mathbf{V})](W_2 \otimes I)$$

and $W_1^* \mathbf{V}_{\alpha}^* \mathbf{V}_{\beta} W_2 = 0$ if $\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}$ is not equal to \mathbf{R}_{γ} or \mathbf{R}_{γ}^* for some $\gamma \in F_n^+$.

Proof. Define the map $\Psi : \text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E})) \rightarrow \mathbf{RH}^1(\mathbf{B}_n)$ by $\Psi(\mu) := \mathcal{P}\mu$. To prove the injectivity of Ψ , let μ_1, μ_2 be in $\text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$ such that $\Psi(\mu_1) = \Psi(\mu_2)$. Due to the uniqueness of the representation of a free k -pluriharmonic function and the definition of the noncommutative Poisson transform of a completely bounded map on $\mathcal{R}_n^* \mathcal{R}_n$, we deduce that $\mu_1(\mathbf{R}_\alpha) = \mu_2(\mathbf{R}_\alpha)$ and $\mu_1(\mathbf{R}_\alpha^*) = \mu_2(\mathbf{R}_\alpha^*)$ for $\alpha \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$, and $\mu_1(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = \mu_2(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in \mathbf{F}_n^+$. Hence, we deduce that $\mu_1 = \mu_2$.

By [Theorem 4.4](#), for any $\varphi \in \mathbf{RH}^1(\mathbf{B}_n)$ there is a completely bounded linear map $\mu_\varphi \in \text{CB}(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$ such that $\varphi = \mathcal{P}\mu_\varphi$ and $\|\varphi\|_1 = \|\mu_\varphi\|_{\text{cb}}$. This proves that the map Ψ is surjective and $\|\mathcal{P}\mu_\varphi\|_1 = \|\mu_\varphi\|_{\text{cb}}$. Therefore, (i) is equivalent to (ii).

Now, the latter equivalence and [Theorem 4.2](#) imply

$$\varphi(X) = (\mathcal{P}\mu_\varphi)(X) = \hat{\mu}_\varphi(C_X^* C_X), \quad X \in \mathbf{B}_n(\mathcal{H}), \tag{5-1}$$

where

$$C_X := (I_{\otimes_{i=1}^k F^2(H_{n_i})} \otimes \Delta_X(I)^{1/2}) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1}.$$

Due to Wittstock’s extension theorem [\[1984\]](#), there exists a completely bounded map $\Phi : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ that extends μ_φ with $\|\mu_\varphi\|_{\text{cb}} = \|\Phi\|_{\text{cb}}$. According to [Theorem 8.4](#) from [\[Paulsen 1986\]](#), there exist a Hilbert space \mathcal{K} , a $*$ -representation $\pi : C^*(\mathbf{R}) \rightarrow B(\mathcal{K})$, and bounded operators $W_1, W_2 : \mathcal{E} \rightarrow \mathcal{K}$ with $\|\Phi\| = \|W_1\| \|W_2\|$ such that

$$\Phi(A) = W_1^* \pi(A) W_2, \quad A \in C^*(\mathbf{R}). \tag{5-2}$$

Set $V_{i,j} := \pi(\mathbf{R}_{i,j})$ for $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$ and note that $\mathbf{V} = (V_1, \dots, V_n)$ is a k -tuple of doubly commuting row isometries $V_i = (V_{i,1}, \dots, V_{i,n_i})$. Using [\(5-1\)](#) and [\(5-2\)](#), one can deduce (iii). The proof that (iii) implies (i) is similar to the proof of the same implication from [Theorem 4.4](#). \square

Consider now the subspace of free holomorphic functions $\mathbf{H}^1(\mathbf{B}_n) := \text{Hol}(\mathbf{B}_n) \cap \mathbf{PH}^1(\mathbf{B}_n)$ together with the norm $\|\cdot\|_1$. Using [Theorem 5.1](#), we can obtain the following weak analogue of the F. and M. Riesz theorem [\[Hoffman 1962\]](#) in our setting.

Corollary 5.2. *($\mathbf{H}^1(\mathbf{B}_n), \|\cdot\|_1$) is a Banach space which can be identified with the annihilator of \mathcal{R}_n in $\text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E}))$, i.e.,*

$$(\mathcal{R}_n)^\perp := \{\mu \in \text{CB}_0(\mathcal{R}_n^* \mathcal{R}_n, B(\mathcal{E})) : \mu(\mathbf{R}_\alpha) = 0 \text{ for all } \alpha \in \mathbf{F}_n^+, |\alpha| \geq 1\}.$$

Moreover, for each $f \in \mathbf{H}^1(\mathbf{B}_n)$, there is a unique completely bounded linear map $\mu_f \in (\mathcal{R}_n)^\perp$ such that $f = \mathcal{P}\mu_f$.

Given a completely bounded linear map $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$, we introduce the *noncommutative Fantappiè transform* of μ to be the map $\mathcal{F}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ defined by

$$(\mathcal{F}\mu)(X) := \hat{\mu} \left(\prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \cdots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} \right)$$

for $\mathbf{X} := (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$. We remark that the noncommutative Fantappiè transform is a linear map and $\mathcal{F}\mu$ is a free holomorphic function in the open polyball \mathbf{B}_n with coefficients in $B(\mathcal{E})$.

Let $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ be a completely positive linear map. We introduce the *noncommutative Herglotz–Riesz transform* of μ on the regular polyball to be the map $\mathbf{H}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ defined by

$$(\mathbf{H}\mu)(\mathbf{X}) := \hat{\mu} \left(2 \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \dots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right)$$

for $\mathbf{X} := (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$. Note that $(\mathbf{H}\mu)(\mathbf{X}) = 2(\mathcal{F}\mu)(\mathbf{X}) - \mu(I) \otimes I$.

Theorem 5.3. *Let f be a function from the polyball $\mathbf{B}_n(\mathcal{H})$ to $B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$. Then the following statements are equivalent:*

- (i) *f is a free holomorphic function with $\Re f \geq 0$ and the linear maps $\{v_{\Re f, r}\}_{r \in [0,1]}$ associated with $\Re f$ are completely positive.*
- (ii) *The function f admits a Herglotz–Riesz representation*

$$f(\mathbf{X}) = (\mathbf{H}\mu)(\mathbf{X}) + i\Im f(0),$$

where $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ is a completely positive linear map with the property that $\mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in \mathbf{F}_n^+$.

- (iii) *There exist a k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, of doubly commuting row isometries on a Hilbert space \mathcal{K} and a bounded linear operator $W : \mathcal{E} \rightarrow \mathcal{K}$ such that*

$$f(\mathbf{X}) = (W^* \otimes I) \left(2 \prod_{i=1}^k (I - V_{i,1}^* \otimes X_{i,1} - \dots - V_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right) (W \otimes I) + i\Im f(0)$$

and $W^* V_\alpha^* V_\beta W = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in \mathbf{F}_n^+$.

Proof. We prove that (i) implies (ii). Let f have the representation $f(\mathbf{X}) = \sum_{\alpha \in \mathbf{F}_n^+} A_\alpha(\mathbf{X}) \otimes X_\alpha$. Due to Corollary 4.5, there exists a completely positive linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ such that $\Re f = \mathcal{P}\mu$. Consequently, we have

$$\mu(I) = \frac{1}{2}(A_{(g)} + A_{(g)}^*), \quad \mu(\mathbf{R}_{\tilde{\alpha}}) = \frac{1}{2}A_{(\alpha)}^*, \quad \mu(\mathbf{R}_{\tilde{\alpha}}^*) = \frac{1}{2}A_{(\alpha)} \quad \text{for all } \alpha \in \mathbf{F}_n^+, |\alpha| \geq 1,$$

and $\mu(\mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}}) = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in \mathbf{F}_n^+$. Using the definition of the Herglotz–Riesz transform, we obtain

$$\begin{aligned} (\mathbf{H}\mu)(\mathbf{X}) &:= \hat{\mu} \left(2 \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \dots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right) \\ &= \sum_{\alpha \in \mathbf{F}_n^+} A_\alpha(\mathbf{X}) \otimes X_\alpha + A_{(g)} \otimes I - \frac{1}{2}(A_{(g)} + A_{(g)}^*) \otimes I \\ &= f(\mathbf{X}) - \frac{1}{2}(A_{(g)}^* - A_{(g)}) \otimes I \\ &= f(\mathbf{X}) - i\Im f(0), \end{aligned}$$

which proves (ii). Now we prove that (ii) implies (i). Assume that (ii) holds. Then

$$f(\mathbf{X}) = 2(\mathcal{F}\mu)(\mathbf{X}) - \mu(I) \otimes I - i\Im f(0)$$

is a free holomorphic function on the polyball \mathbf{B}_n . Taking into account that $\mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = 0$ if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in F_n^+$, and using [Theorem 4.2](#), we deduce that

$$\begin{aligned} & \frac{1}{2}(f(\mathbf{X}) + f(\mathbf{X})^*) \\ &= \frac{1}{2}((\mathbf{H}\mu)(\mathbf{X}) + (\mathbf{H}\mu)(\mathbf{X})^*) \\ &= \hat{\mu} \left(\prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \dots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I + \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \dots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1} \right) \\ &= \hat{\mu}(\mathcal{P}(\mathbf{R}, \mathbf{X})) \geq 0. \end{aligned}$$

Therefore, $\Re f$ is a free k -pluriharmonic function such that $\Re f = \mathcal{P}\mu$. Due to [Corollary 4.5](#), we deduce that the linear maps $\{v_{\Re f, r}\}_{r \in [0,1]}$ associated with $\Re f$ are completely positive.

Now, we prove that (ii) implies (iii). Assume that (ii) holds. According to Stinespring’s representation theorem [[1955](#)], there is a Hilbert space \mathcal{K} , a $*$ -representation $\pi : C^*(\mathbf{R}) \rightarrow B(\mathcal{K})$ and a bounded $W : \mathcal{E} \rightarrow \mathcal{K}$ with $\|\mu(I)\| = \|W\|^2$ such that $\mu(A) = W^* \pi(A) W$ for all $A \in C^*(\mathbf{R})$. Setting $V_{i,j} := \pi(\mathbf{R}_{ij})$ for all $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$, it is clear that the k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, consists of doubly commuting row isometries. Note that, if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in F_n^+$, then

$$W^* \mathbf{V}_\alpha^* \mathbf{V}_\beta W = W^* \pi(\mathbf{R}_\alpha^* \mathbf{R}_\beta) W = \mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = 0.$$

Now, one can easily see that the relation $f(\mathbf{X}) = (\mathbf{H}\mu)(\mathbf{X}) + i\Im f(0)$ leads to the representation in (iii), which completes the proof.

It remains to prove that (iii) implies (ii). To this end, assume that (iii) holds. Since the k -tuple $\mathbf{V} = (V_1, \dots, V_k)$, $V_i = (V_{i,1}, \dots, V_{i,n_i})$, consists of doubly commuting row isometries on a Hilbert space \mathcal{K} , the noncommutative von Neumann inequality [[Popescu 1999](#)] implies that the map $\pi : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ defined by

$$\pi(\mathbf{R}_\alpha \mathbf{R}_\beta^*) := \mathbf{V}_\alpha \mathbf{V}_\beta^*, \quad \alpha, \beta \in F_n^+,$$

is a $*$ -representation of $C^*(\mathbf{R})$. Define $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ by setting $\mu(A) := W^* \pi(A) W$. It is clear that μ is a completely positive linear map and (iii) implies

$$f(\mathbf{X}) = \hat{\mu} \left(2 \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \dots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right) + i\Im f(0).$$

Note also that

$$\mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = W^* \pi(\mathbf{R}_\alpha^* \mathbf{R}_\beta) W = W^* \mathbf{V}_\alpha^* \mathbf{V}_\beta W = 0$$

if $\mathbf{R}_\alpha^* \mathbf{R}_\beta$ is not equal to \mathbf{R}_γ or \mathbf{R}_γ^* for some $\gamma \in F_n^+$. This shows that (ii) holds. □

In the particular case when $n_1 = \dots = n_k = 1$, we obtain an operator-valued extension of Korányi–Pukánszky integral representation for holomorphic functions with positive real part on polydisks [Korányi and Pukánszky 1963].

In what follows, we say that f has a Herglotz–Riesz representation if Theorem 5.3(ii) is satisfied.

Theorem 5.4. *Let $f : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ be a function, where $\mathbf{n} = (n_1, \dots, n_k) \in \mathbb{N}^k$. If f admits a Herglotz–Riesz representation, then f is a free holomorphic function with $\Re f \geq 0$.*

Conversely, if f is a free holomorphic function such that $\Re f \geq 0$, then there is a unique completely positive linear map $\mu : \mathcal{R}_n^ + \mathcal{R}_n \rightarrow B(\mathcal{E})$ such that*

$$f(Y) = (\mathbf{H}\mu)(kY) + i\Im f(0), \quad Y \in \frac{1}{k}\mathbf{B}_n(\mathcal{H}).$$

Moreover,

$$f(X) = 2 \sum_{\alpha \in F_n^+} k^{|\alpha|} \mu(\mathbf{R}_\alpha^*) \otimes X_\alpha - \mu(I) \otimes I, \quad X \in \mathbf{B}_n(\mathcal{H}).$$

Proof. The forward implication was proved in Theorem 5.3. We prove the converse. Assume that f has the representation

$$f(X) = \sum_{\alpha \in F_n^+} A_{(\alpha)} \otimes X_\alpha, \quad X \in \mathbf{B}_n(\mathcal{H}). \tag{5-3}$$

First we consider the case when $\frac{1}{2}(A_{(\mathbf{g})} + A_{(\mathbf{g})}^*) = I_{\mathcal{E}}$. Since $\Re f \geq 0$ and $\Re f(0) = I$, Theorem 3.6 shows that there is a k -tuple $\mathbf{V} = (V_1, \dots, V_k)$ of commuting row isometries on a space $\mathcal{K} \supset \mathcal{E}$ such that

$$\Re f(X) = \sum_{(\sigma, \omega) \in F_n^+ \times F_n^+} P_{\mathcal{E}} \mathbf{V}_\alpha^* \mathbf{V}_\beta |_{\mathcal{E}} \otimes X_\alpha X_\beta^*.$$

The uniqueness of the representation of free k -pluriharmonic functions on \mathbf{B}_n implies

$$P_{\mathcal{E}} \mathbf{V}_\alpha^* \mathbf{V}_\beta |_{\mathcal{E}} = \begin{cases} \frac{1}{2} A_{(\alpha)} & \text{if } \alpha \in F_n^+, \beta = \mathbf{g}, \\ \frac{1}{2} A_{(\beta)}^* & \text{if } \beta \in F_n^+, \alpha = \mathbf{g}, \\ 0 & \text{otherwise.} \end{cases} \tag{5-4}$$

Set $T_{i,j} := (1/k)V_{i,j}$ for $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$. According to Proposition 1.9 from [Popescu 2016], the k -tuple $\mathbf{T} := (T_1, \dots, T_k)$, $T_i := (T_{i,1}, \dots, T_{i,n_i})$, is in the closed polyball $\mathbf{B}_n(\mathcal{K})$. Using Theorem 7.2 from [Popescu 2016], we find a k -tuple $\mathbf{W} := (W_1, \dots, W_k)$ of doubly commuting row isometries on a Hilbert space $\mathcal{G} \supset \mathcal{K}$ such that $W_{i,j}^*|_{\mathcal{K}} = T_{i,j}^*$ for all $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i\}$. Define the linear map $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ by setting

$$\mu(\mathbf{R}_{\tilde{\beta}} \mathbf{R}_\alpha^*) = P_{\mathcal{E}} [P_{\mathcal{K}} \mathbf{W}_{\tilde{\beta}} \mathbf{W}_\alpha^* |_{\mathcal{K}}] |_{\mathcal{E}}, \quad \alpha, \beta \in F_n^+.$$

Note that μ is a completely positive linear map with the property that $\mu(\mathbf{R}_{\tilde{\beta}}) = (1/2k^{|\beta|})A_{(\beta)}^*$ and $\mu(\mathbf{R}_\alpha^*) = (1/2k^{|\alpha|})A_{(\alpha)}$ if $\alpha, \beta \in F_n^+$ with $\alpha \neq \mathbf{g}$ and $\beta \neq \mathbf{g}$, and $\mu(I) = I_{\mathcal{E}}$. Consequently, (5-3) and (5-4) imply

$$f(X) = 2 \sum_{\alpha \in F_n^+} k^{|\alpha|} \mu(\mathbf{R}_\alpha^*) \otimes X_\alpha - \mu(I) \otimes I, \quad X \in \mathbf{B}_n(\mathcal{H}).$$

Setting $Y := (1/k)X$, we deduce that

$$\begin{aligned} f(Y) &= 2 \sum_{\alpha \in F_n^+} \mu(\mathbf{R}_\alpha^*) \otimes k^{|\alpha|} Y_\alpha - \mu(I) \otimes I \\ &= \hat{\mu} \left(2 \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes kY_{i,1} - \dots - \mathbf{R}_{i,n_i}^* \otimes kY_{i,n_i})^{-1} - I \right) \\ &= (\mathbf{H}\mu)(kY) \end{aligned}$$

for any $Y \in (1/k)\mathbf{B}_n(\mathcal{H})$, which completes the proof when $A_{(\mathbf{g})} = I_{\mathcal{E}}$. Now, we consider the case when $C_{(\mathbf{g})} := \frac{1}{2}(A_{(\mathbf{g})} + A_{(\mathbf{g})}^*) \geq 0$. For each $\epsilon > 0$, define the free holomorphic function

$$g_\epsilon := (C_{(\mathbf{g})} + \epsilon I)^{-1/2} [f + \epsilon I_{\mathcal{E}} \otimes I_{\mathcal{H}}] (C_{(\mathbf{g})} + \epsilon I)^{-1/2}$$

and note that $\Re g_\epsilon(0) = I$. Applying the first part of the proof to g_ϵ , we find a completely positive linear map $\mu_\epsilon : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ with the property that

$$\mu_\epsilon(\mathbf{R}_{\tilde{\beta}}) = \frac{1}{2k^{|\beta|}} (C_{(\mathbf{g})} + \epsilon I)^{-1/2} A_{(\beta)}^* (C_{(\mathbf{g})} + \epsilon I)^{-1/2}$$

and

$$\mu_\epsilon(\mathbf{R}_\alpha^*) = \frac{1}{2k^{|\alpha|}} (C_{(\mathbf{g})} + \epsilon I)^{-1/2} A_{(\alpha)} (C_{(\mathbf{g})} + \epsilon I)^{-1/2}$$

if $\alpha, \beta \in F_n^+$ with $\alpha \neq \mathbf{g}$ and $\beta \neq \mathbf{g}$, and $\mu_\epsilon(I) = I_{\mathcal{E}}$. Setting $v_\epsilon(\xi) := (C_{(\mathbf{g})} + \epsilon I)^{1/2} \mu_\epsilon(\xi) (C_{(\mathbf{g})} + \epsilon I)^{1/2}$ for all $\xi \in C^*(\mathbf{R})$, one can easily see that v_ϵ is a completely positive linear map with the property that $v_\epsilon(\mathbf{R}_{\tilde{\beta}}) = (1/2k^{|\beta|})A_{(\beta)}^*$ and $v_\epsilon(\mathbf{R}_\alpha^*) = (1/2k^{|\alpha|})A_{(\alpha)}$ if $\alpha, \beta \in F_n^+$ with $\alpha \neq \mathbf{g}$ and $\beta \neq \mathbf{g}$, and $v_\epsilon(I) = C_{(\mathbf{g})} + \epsilon I_{\mathcal{E}}$. Following the proofs of [Theorem 4.4](#) and [Corollary 4.5](#), we find a completely positive linear map $v : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ such that $v(\xi) = \text{WOT-}\lim_{\epsilon_k \rightarrow 0} v_{\epsilon_k}(\xi)$ for $\xi \in C^*(\mathbf{R})$, where $\{\epsilon_k\}$ is a sequence of positive numbers converging to zero. Consequently, we have $v(\mathbf{R}_{\tilde{\beta}}) = (1/2k^{|\beta|})A_{(\beta)}^*$ and $v(\mathbf{R}_\alpha^*) = (1/2k^{|\alpha|})A_{(\alpha)}$ if $\alpha, \beta \in F_n^+$ with $\alpha \neq \mathbf{g}$ and $\beta \neq \mathbf{g}$, and $v(I) = C_{(\mathbf{g})}$. As in the first part of this proof, one can easily see that

$$f(Y) = (\mathbf{H}v)(kY) + i\Im f(0), \quad Y \in \frac{1}{k}\mathbf{B}_n(\mathcal{H}),$$

and

$$f(X) = 2 \sum_{\alpha \in F_n^+} k^{|\alpha|} v(\mathbf{R}_\alpha^*) \otimes X_\alpha - v(I) \otimes I, \quad X \in \mathbf{B}_n(\mathcal{H}). \quad \square$$

References

[Arveson 1969] W. B. Arveson, “Subalgebras of C^* -algebras”, *Acta Math.* **123** (1969), 141–224. [MR 0253059](#) [Zbl 0194.15701](#)
 [Ball and Vinnikov 2003] J. A. Ball and V. Vinnikov, “Formal reproducing kernel Hilbert spaces: the commutative and noncommutative settings”, pp. 77–134 in *Reproducing kernel spaces and applications*, edited by D. Alpay, Operator Theory: Advances and Applications **143**, Birkhäuser, Basel, 2003. [MR 2019348](#) [Zbl 1084.46018](#)
 [Ball et al. 2016] J. A. Ball, G. Marx, and V. Vinnikov, “Noncommutative reproducing kernel Hilbert spaces”, preprint, 2016. [arXiv 1602.00760](#)

- [Berezin 1972] F. A. Berezin, “Ковариантные и контравариантные символы операторов”, *Izv. Akad. Nauk SSSR Ser. Mat.* **36**:5 (1972), 1134–1167. Translated as “Covariant and contravariant symbols of operators” in *Math. USSR Izv.* **6**:5 (1972), 1117–1151. [MR 350504](#) [Zbl 0247.47019](#)
- [Brown and Halmos 1963] A. Brown and P. R. Halmos, “Algebraic properties of Toeplitz operators”, *J. Reine Angew. Math.* **213** (1963), 89–102. [MR 0160136](#) [Zbl 0116.32501](#)
- [Davidson and Pitts 1998] K. R. Davidson and D. R. Pitts, “The algebraic structure of non-commutative analytic Toeplitz algebras”, *Math. Ann.* **311**:2 (1998), 275–303. [MR 1625750](#) [Zbl 0939.47060](#)
- [Davidson et al. 2001] K. R. Davidson, E. Katsoulis, and D. R. Pitts, “The structure of free semigroup algebras”, *J. Reine Angew. Math.* **533** (2001), 99–125. [MR 1823866](#) [Zbl 0967.47047](#)
- [Davidson et al. 2005] K. R. Davidson, J. Li, and D. R. Pitts, “Absolutely continuous representations and a Kaplansky density theorem for free semigroup algebras”, *J. Funct. Anal.* **224**:1 (2005), 160–191. [MR 2139108](#) [Zbl 1084.46042](#)
- [Douglas 1998] R. G. Douglas, *Banach algebra techniques in operator theory*, 2nd ed., Graduate Texts in Mathematics **179**, Springer, New York, 1998. [MR 1634900](#) [Zbl 0920.47001](#)
- [Effros and Ruan 2000] E. G. Effros and Z.-J. Ruan, *Operator spaces*, London Mathematical Society Monographs. New Series **23**, Oxford University Press, New York, 2000. [MR 1793753](#) [Zbl 0969.46002](#)
- [Herglotz 1911] G. Herglotz, “Über Potenzreihen mit positivem, reeltem Teil im Einheitskreis”, *Ber. Verh. Sächs. Gesellsch. Wiss. Leipzig Math. Phys. Kl.* **63** (1911), 501–511. [JFM 42.0438.02](#)
- [Hoffman 1962] K. Hoffman, *Banach spaces of analytic functions*, Prentice-Hall, Englewood Cliffs, NJ, 1962. [MR 0133008](#) [Zbl 0117.34001](#)
- [Kennedy 2011] M. Kennedy, “Wandering vectors and the reflexivity of free semigroup algebras”, *J. Reine Angew. Math.* **653** (2011), 47–73. [MR 2794625](#) [Zbl 1218.47136](#)
- [Kennedy 2013] M. Kennedy, “The structure of an isometric tuple”, *Proc. Lond. Math. Soc.* (3) **106**:5 (2013), 1157–1177. [MR 3066752](#) [Zbl 1278.47016](#)
- [Korányi and Pukánszky 1963] A. Korányi and L. Pukánszky, “Holomorphic functions with positive real part on polycylinders”, *Trans. Amer. Math. Soc.* **108** (1963), 449–456. [MR 0159029](#) [Zbl 0136.07103](#)
- [von Neumann 1951] J. von Neumann, “Eine Spektraltheorie für allgemeine Operatoren eines unitären Raumes”, *Math. Nachr.* **4** (1951), 258–281. [MR 0043386](#) [Zbl 0042.12301](#)
- [Neumark 1943] M. A. Neumark, “Положительно-определенные операторные функции на коммутативной группе”, *Izv. Akad. Nauk SSSR Ser. Mat.* **7**:5 (1943), 237–244. [MR 0010265](#) [Zbl 0061.25411](#)
- [Paulsen 1986] V. I. Paulsen, *Completely bounded maps and dilations*, Pitman Research Notes in Mathematics Series **146**, Longman Scientific & Technical, Harlow, 1986. [MR 868472](#) [Zbl 0614.47006](#)
- [Pisier 2001] G. Pisier, *Similarity problems and completely bounded maps*, 2nd ed., Lecture Notes in Mathematics **1618**, Springer, Berlin, 2001. [MR 1818047](#) [Zbl 0971.47016](#)
- [Popescu 1989] G. Popescu, “Multi-analytic operators and some factorization theorems”, *Indiana Univ. Math. J.* **38**:3 (1989), 693–710. [MR 1017331](#) [Zbl 0661.47020](#)
- [Popescu 1995] G. Popescu, “Multi-analytic operators on Fock spaces”, *Math. Ann.* **303**:1 (1995), 31–46. [MR 1348353](#) [Zbl 0835.47015](#)
- [Popescu 1999] G. Popescu, “Poisson transforms on some C^* -algebras generated by isometries”, *J. Funct. Anal.* **161**:1 (1999), 27–61. [MR 1670202](#) [Zbl 0933.46070](#)
- [Popescu 2006] G. Popescu, *Entropy and multivariable interpolation*, Memoirs of the American Mathematical Society **868**, American Mathematical Society, Providence, RI, 2006. [MR 2263661](#) [Zbl 1115.47016](#)
- [Popescu 2009] G. Popescu, “Noncommutative transforms and free pluriharmonic functions”, *Adv. Math.* **220**:3 (2009), 831–893. [MR 2483229](#) [Zbl 1169.47010](#)
- [Popescu 2013] G. Popescu, “Berezin transforms on noncommutative varieties in polydomains”, *J. Funct. Anal.* **265**:10 (2013), 2500–2552. [MR 3091823](#) [Zbl 1298.47019](#)
- [Popescu 2014] G. Popescu, “Euler characteristic on noncommutative polyballs”, *J. Reine Angew. Math.* (online publication December 2014).

- [Popescu 2015a] G. Popescu, “Curvature invariant on noncommutative polyballs”, *Adv. Math.* **279** (2015), 104–158. [MR 3345180](#) [Zbl 1325.47015](#)
- [Popescu 2015b] G. Popescu, “Holomorphic automorphisms of noncommutative polyballs”, preprint, 2015. To appear in *J. Operator Theory*. [arXiv 1502.07905](#)
- [Popescu 2016] G. Popescu, “Berezin transforms on noncommutative polydomains”, *Trans. Amer. Math. Soc.* **368**:6 (2016), 4357–4416. [MR 3453374](#) [Zbl 06551095](#)
- [Popescu \geq 2016] G. Popescu, “Hyperbolic geometry of noncommutative polyballs”, in preparation.
- [Riesz 1911] F. Riesz, “Sur certains systèmes singuliers d’équations intégrales”, *Ann. Sci. École Norm. Sup.* (3) **28** (1911), 33–62. [MR 1509135](#) [Zbl 42.0374.03](#)
- [Rudin 1969] W. Rudin, *Function theory in polydiscs*, W. A. Benjamin, New York, 1969. [MR 0255841](#) [Zbl 0177.34101](#)
- [Schur 1918] J. Schur, “Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind”, *J. Reine Angew. Math.* **148** (1918), 122–145. [MR 1580958](#) [JFM 46.0475.01](#)
- [Stinespring 1955] W. F. Stinespring, “Positive functions on C^* -algebras”, *Proc. Amer. Math. Soc.* **6** (1955), 211–216. [MR 0069403](#) [Zbl 0064.36703](#)
- [Sz.-Nagy et al. 2010] B. Sz.-Nagy, C. Foias, H. Bercovici, and L. Kérchy, *Harmonic analysis of operators on Hilbert space*, 2nd ed., Springer, New York, 2010. [MR 2760647](#) [Zbl 1234.47001](#)
- [Wittstock 1981] G. Wittstock, “Ein operatorwertiger Hahn–Banach Satz”, *J. Funct. Anal.* **40**:2 (1981), 127–150. [MR 609438](#) [Zbl 0495.46005](#)
- [Wittstock 1984] G. Wittstock, “On matrix order and convexity”, pp. 175–188 in *Functional analysis: surveys and recent results, III* (Paderborn, 1983), edited by K. D. Bierstedt and B. Fuchssteiner, North-Holland Mathematics Studies **90**, North-Holland, Amsterdam, 1984. [MR 761380](#) [Zbl 0589.46003](#)

Received 3 Dec 2015. Revised 29 Feb 2016. Accepted 12 Apr 2016.

GELU POPESCU: gelu.popescu@utsa.edu

Department of Mathematics, The University of Texas at San Antonio, One UTSA Circle, San Antonio, TX 78249, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 5 2016

Multidimensional entire solutions for an elliptic system modelling phase separation NICOLA SOAVE and ALESSANDRO ZILIO	1019
Time-periodic approximations of the Euler–Poisson system near Lane–Emden stars JUHI JANG	1043
Sharp weighted norm estimates beyond Calderón–Zygmund theory FRÉDÉRIC BERNICOT, DOROTHEE FREY and STEFANIE PETERMICHL	1079
Existence, uniqueness and optimal regularity results for very weak solutions to nonlinear elliptic systems MIROSLAV BULÍČEK, LARS DIENING and SEBASTIAN SCHWARZACHER	1115
On polynomial configurations in fractal sets KEVIN HENRIOT, IZABELLA ŁABA and MALABIKA PRAMANIK	1153
Free pluriharmonic functions on noncommutative polyballs GELU POPESCU	1185
Bohnenblust–Hille inequalities for Lorentz spaces via interpolation ANDREAS DEFANT and MIECZYŚLAW MASTYŁO	1235
On the negative spectrum of the Robin Laplacian in corner domains VINCENT BRUNEAU and NICOLAS POPOFF	1259