

ANALYSIS & PDE

Volume 9

No. 7

2016

DUONG H. PHONG, SEBASTIEN PICARD AND XIANGWEN ZHANG

**A SECOND ORDER ESTIMATE
FOR GENERAL COMPLEX HESSIAN EQUATIONS**



A SECOND ORDER ESTIMATE FOR GENERAL COMPLEX HESSIAN EQUATIONS

DUONG H. PHONG, SEBASTIEN PICARD AND XIANGWEN ZHANG

We consider the general complex Hessian equations with right-hand sides depending on gradients, which are motivated by the Fu–Yau equations arising from the study of Strominger systems. The second order estimate for the solution is crucial to solving the equation by the method of continuity. We obtain such an estimate for the χ -plurisubharmonic solutions.

1. Introduction

Let (X, ω) be a compact Kähler manifold of dimension $n \geq 2$. Let $u \in C^\infty(X)$ and consider a $(1, 1)$ -form $\chi(z, u)$ possibly depending on u and satisfying the positivity condition $\chi \geq \varepsilon\omega$ for some $\varepsilon > 0$. We define

$$g = \chi(z, u) + i\partial\bar{\partial}u, \quad (1-1)$$

and u is called χ -plurisubharmonic if $g > 0$ as a $(1, 1)$ form. In this paper, we are concerned with the complex Hessian equation

$$(\chi(z, u) + i\partial\bar{\partial}u)^k \wedge \omega^{n-k} = \psi(z, Du, u) \omega^n \quad (1-2)$$

for $1 \leq k \leq n$, where $\psi(z, v, u) \in C^\infty(T^{1,0}(X)^* \times \mathbb{R})$ is a given strictly positive function.

The complex Hessian equation can be viewed as an intermediate equation between the Laplace equation and the complex Monge–Ampère equation. It encompasses the most natural invariants of the complex Hessian matrix of a real valued function, namely the elementary symmetric polynomials of its eigenvalues. When $k = 1$, the equation (1-2) is quasilinear, and the estimates follow from the classical theory of quasilinear PDEs. The real counterparts of (1-2) for $1 < k \leq n$, with ψ not depending on the gradient of u , have been studied extensively in the literature (see the survey paper [Wang 2009] and more recent related work [Guan 2014]), as these equations appear naturally and play very important roles in both classical and conformal geometry. When the right-hand side ψ depends on the gradient of the solution, even the real case has been a long-standing problem due to substantial difficulties in obtaining a priori C^2 estimates. This problem has recently been solved by Guan, Ren and Wang [Guan et al. 2015] for convex solutions of real Hessian equations.

In the complex case, equation (1-2) with $\psi = \psi(z, u)$ has been extensively studied in recent years, due to its appearance in many geometric problems, including the J -flow [Song and Weinkove 2008] and

Work supported in part by the National Science Foundation under Grant DMS-1266033, DMS-1605968 and DMS-1308136.
MSC2010: 35J15, 35J60, 58J05, 35J96.

Keywords: complex Hessian equation, second order estimate.

quaternionic geometry [Alesker and Verbitsky 2010]. The related Dirichlet problem for (1-2) on domains in \mathbb{C}^n has been studied by Li [2004] and Błocki [2005]. The corresponding problem on compact Kähler or Hermitian manifolds has also been studied extensively; see, for example, [Dinew and Kołodziej 2014; Hou 2009; Kołodziej and Nguyen 2016; Lu and Nguyen 2015; Zhang 2010]. In particular, as a crucial step in the continuity method, C^2 estimates for complex Hessian type equations have been studied in various settings; see [Hou et al. 2010; Sun 2014; Székelyhidi 2015; Székelyhidi et al. 2015; Zhang 2015].

However, (1-2) with $\psi = \psi(z, Du, u)$ has been much less studied. An important case corresponding to $k = n = 2$, so that it is actually a Monge–Ampère equation in two dimensions, is central to the solution by Fu and Yau [2008; 2007] of a Strominger system on a toric fibration over a K3 surface. A natural generalization of this case to general dimension n was suggested by Fu and Yau [2008] and can be expressed as

$$((e^u + fe^{-u})\omega + ni\partial\bar{\partial}u)^2 \wedge \omega^{n-2} = \psi(z, Du, u)\omega^n, \quad (1-3)$$

where $\psi(z, v, u)$ is a function on $T^{1,0}(X)^* \times \mathbb{R}$ with a particular structure, and (X, ω) is a compact Kähler manifold. A priori estimates for this equation were obtained by the authors in [Phong et al. 2015].

In this paper, motivated by our previous work [Phong et al. 2015], we study a priori C^2 estimates for the equation (1-2) with general $\chi(z, u)$ and general right-hand side $\psi(z, Du, u)$. Building on the techniques developed in [Guan et al. 2015] (see also [Li et al. 2016] for real Hessian equations), we can prove the following theorem.

Theorem 1. *Let (X, ω) be a compact Kähler manifold of complex dimension n . Suppose $u \in C^4(X)$ is a solution of (1-2) with $g = \chi + i\partial\bar{\partial}u > 0$ and $\chi(z, u) \geq \varepsilon\omega$. Let $0 < \psi(z, v, u) \in C^\infty(T^{1,0}(X)^* \times \mathbb{R})$. Then we have the uniform second order derivative estimate*

$$|D\bar{D}u|_\omega \leq C, \quad (1-4)$$

where C is a positive constant depending only on $\varepsilon, n, k, \sup_X |u|, \sup_X |Du|$, and the C^2 norm of χ as a function of (u, z) , the infimum of ψ , and the C^2 norm of ψ as a function of (z, Du, u) , all restricted to the ranges in Du and u defined by the uniform upper bounds on $|u|$ and $|Du|$.

We remark that the above estimate is stated for χ -plurisubharmonic solutions, that is, $g = \chi + i\partial\bar{\partial}u > 0$. Actually, we only need to assume that g is in the Γ_{k+1} cone (see (3-11) below for the definition of the Garding cone Γ_k and also the discussion in Remark 2 at the end of the paper). However, a better condition would be $g \in \Gamma_k$, which is the natural cone for ellipticity. In fact, this is still an open problem even for real Hessian equations when $2 < k < n$. If $k = 2$, [Guan et al. 2015] removed the convexity assumption by investigating the structure of the operator. A simpler argument was given recently by Spruck and Xiao [2015]. However, the arguments are not applicable to the complex case due to the difference between the terms $|DDu|^2$ and $|D\bar{D}u|^2$ in the complex setting. When $k = 2$ in the complex setting, C^2 estimates for (1-3) were obtained in [Phong et al. 2015] without the plurisubharmonicity assumption, but the techniques rely on the specific right-hand side $\psi(z, Du, u)$ studied there.

We also note that if $k = n$, the condition $g = \chi + i\partial\bar{\partial}u > 0$ is the natural assumption for the ellipticity of equation (1-2). Thus, our result implies the a priori C^2 estimate for complex Monge–Ampère equations

with right-hand side depending on gradients:

$$(\chi(z, u) + i\partial\bar{\partial}u)^n = \psi(z, Du, u)\omega^n.$$

This generalizes the C^2 estimate for the equation studied by Fu and Yau [2008; 2007] mentioned above, which corresponds to $n = 2$ and a specific form $\chi(z, u)$ as well as a specific right-hand side $\psi(z, Du, u)$. For dimension $n \geq 2$ and $k = n$, the estimate was obtained by Guan and Ma, in unpublished notes, using a different method where the structure of the Monge–Ampère operator plays an important role.

Compared to the estimates when $\psi = \psi(z, u)$, the dependence on the gradient of u in (1-2) creates substantial new difficulties. The main obstacle is the appearance of terms such as $|DDu|^2$ and $|D\bar{D}u|^2$ when one differentiates the equation twice. We adapt the techniques used in [Guan et al. 2015] and [Li et al. 2016] for real Hessian equations to overcome these difficulties. Furthermore, we also need to handle properly some subtle issues when dealing with the third-order terms due to complex conjugacy.

2. Preliminaries

Let σ_k be the k -th elementary symmetric function; that is, for $1 \leq k \leq n$ and $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$,

$$\sigma_k(\lambda) = \sum_{1 < i_1 < \dots < i_k < n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_n}.$$

Let $\lambda(a_{\bar{j}i})$ denote the eigenvalues of a Hermitian symmetric matrix $(a_{\bar{j}i})$ with respect to the background Kähler metric ω . We define $\sigma_k(a_{\bar{j}i}) = \sigma_k(\lambda(a_{\bar{j}i}))$. This definition can be naturally extended to complex manifolds. Denoting by $A^{1,1}(X)$ the space of smooth real $(1, 1)$ -forms on a compact Kähler manifold (X, ω) , we define, for any $g \in A^{1,1}(X)$,

$$\sigma_k(g) = \binom{n}{k} \frac{g^k \wedge \omega^{n-k}}{\omega^n}.$$

Using the above notation, we can rewrite (1-2) as follows:

$$\sigma_k(g) = \sigma_k(\chi_{\bar{j}i} + u_{\bar{j}i}) = \psi(z, Du, u). \tag{2-1}$$

We use the notation

$$\sigma_k^{p\bar{q}} = \frac{\partial \sigma_k(g)}{\partial g_{\bar{q}p}}, \quad \sigma_k^{p\bar{q}, r\bar{s}} = \frac{\partial^2 \sigma_k(g)}{\partial g_{\bar{q}p} \partial \bar{g}_{\bar{s}r}}.$$

The symbol D indicates the covariant derivative with respect to the given metric ω . All norms and inner products are with respect to ω unless denoted otherwise. We denote by $\lambda_1, \lambda_2, \dots, \lambda_n$ the eigenvalues of $g_{\bar{j}i} = \chi_{\bar{j}i} + u_{\bar{j}i}$ with respect to ω , and use the ordering $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0$. Our calculations are carried out at a point z on the manifold X , and we use coordinates such that at this point $\omega = i \sum \delta_{\ell k} dz^k \wedge d\bar{z}^\ell$ and $g_{\bar{j}i}$ is diagonal. We also use the notation

$$\mathcal{F} = \sum_p \sigma_k^{p\bar{p}}.$$

Differentiating (2-1) yields

$$\sigma_k^{p\bar{q}} D_{\bar{j}} g_{\bar{q}p} = D_{\bar{j}} \psi. \tag{2-2}$$

Differentiating the equation a second time gives

$$\begin{aligned} \sigma_k^{p\bar{q}} D_i D_{\bar{j}} g_{\bar{q}p} + \sigma_k^{p\bar{q},r\bar{s}} D_i g_{\bar{q}p} D_{\bar{j}} g_{\bar{s}r} &= D_i D_{\bar{j}} \psi \\ &\geq -C(1 + |DDu|^2 + |D\bar{D}u|^2) + \sum_{\ell} \psi_{v_{\ell}} u_{\ell\bar{j}i} + \sum_{\ell} \psi_{\bar{v}_{\ell}} u_{\bar{\ell}\bar{j}i}. \end{aligned} \quad (2-3)$$

We denote by C a uniform constant which depends only on (X, ω) , $n, k, \|\chi\|_{C^2}, \inf \psi, \|u\|_{C^1}$ and $\|\psi\|_{C^2}$. We now compute the operator $\sigma_k^{p\bar{q}} D_p D_{\bar{q}}$ acting on $g_{\bar{j}i} = \chi_{\bar{j}i} + u_{\bar{j}i}$. Recalling that $\chi_{\bar{j}i}$ depends on u , we estimate

$$\begin{aligned} \sigma_k^{p\bar{q}} D_p D_{\bar{q}} g_{\bar{j}i} &= \sigma_k^{p\bar{q}} D_p D_{\bar{q}} D_i D_{\bar{j}} u + \sigma_k^{p\bar{q}} D_p D_{\bar{q}} \chi_{\bar{j}i} \\ &\geq \sigma_k^{p\bar{q}} D_p D_{\bar{q}} D_i D_{\bar{j}} u - C(1 + \lambda_1) \mathcal{F}. \end{aligned} \quad (2-4)$$

Commuting derivatives

$$\begin{aligned} D_p D_{\bar{q}} D_i D_{\bar{j}} u &= D_i D_{\bar{j}} D_p D_{\bar{q}} u - R_{\bar{q}i\bar{j}}^{\bar{a}} u_{\bar{a}p} + R_{\bar{q}p\bar{j}}^{\bar{a}} u_{\bar{a}i} \\ &= D_i D_{\bar{j}} g_{\bar{q}p} - D_i D_{\bar{j}} \chi_{\bar{q}p} - R_{\bar{q}i\bar{j}}^{\bar{a}} u_{\bar{a}p} + R_{\bar{q}p\bar{j}}^{\bar{a}} u_{\bar{a}i}. \end{aligned} \quad (2-5)$$

Therefore, by (2-3),

$$\begin{aligned} \sigma_k^{p\bar{q}} D_p D_{\bar{q}} g_{\bar{j}i} &\geq -\sigma_k^{p\bar{q},r\bar{s}} D_j g_{\bar{q}p} D_{\bar{j}} g_{\bar{s}r} + \sum \psi_{v_{\ell}} g_{\bar{j}i\ell} + \sum \psi_{\bar{v}_{\ell}} g_{\bar{j}i\bar{\ell}} \\ &\quad - C(1 + |DDu|^2 + |D\bar{D}u|^2 + (1 + \lambda_1) \mathcal{F}). \end{aligned} \quad (2-6)$$

We next compute the operator $\sigma_k^{p\bar{q}} D_p D_{\bar{q}}$ acting on $|Du|^2$, introducing the notation

$$|DDu|_{\sigma\omega}^2 = \sigma_k^{p\bar{q}} \omega^{m\bar{\ell}} D_p D_m u D_{\bar{q}} D_{\bar{\ell}} u, \quad |D\bar{D}u|_{\sigma\omega}^2 = \sigma_k^{p\bar{q}} \omega^{m\bar{\ell}} D_p D_{\bar{\ell}} u D_m D_{\bar{q}} u. \quad (2-7)$$

Then

$$\begin{aligned} \sigma_k^{p\bar{q}} |Du|_{\bar{q}p}^2 &= \sigma_k^{p\bar{q}} (D_p D_{\bar{q}} D_m u D^m u + D_m u D_p D_{\bar{q}} D^m u) + |DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2 \\ &= \sigma_k^{p\bar{q}} (D_m (g_{\bar{q}p} - \chi_{\bar{q}p}) D^m u + D_m u D^m (g_{\bar{q}p} - \chi_{\bar{q}p})) + \sigma_k^{p\bar{q}} R_{\bar{q}p}^{m\bar{\ell}} u_{\bar{\ell}} u_m \\ &\quad + |DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2. \end{aligned} \quad (2-8)$$

Using the differentiated equation, we obtain

$$\begin{aligned} \sigma_k^{p\bar{q}} |Du|_{\bar{q}p}^2 &\geq 2 \operatorname{Re} \langle Du, D\psi \rangle - C(1 + \mathcal{F}) + |DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2 \\ &\geq 2 \operatorname{Re} \left(\sum_{p,m} (D_p D_m u D_{\bar{p}} u + D_p u D_{\bar{p}} D_m u) \psi_{v_m} \right) - C(1 + \mathcal{F}) + |DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2. \end{aligned}$$

To simplify the expression, we introduce the notation

$$\langle D|Du|^2, D_{\bar{v}} \psi \rangle = \sum_m (D_m D_p u D^p u \psi_{v_m} + D_p u D_m D^p u \psi_{v_m}). \quad (2-9)$$

We obtain

$$\sigma_k^{p\bar{q}} |Du|_{\bar{q}p}^2 \geq 2 \operatorname{Re} \langle D|Du|^2, D_{\bar{v}} \psi \rangle - C(1 + \mathcal{F}) + |DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2. \quad (2-10)$$

We also compute

$$-\sigma_k^{p\bar{q}} u_{\bar{q}p} = \sigma_k^{p\bar{q}} (\chi_{\bar{q}p} - g_{\bar{q}p}) \geq \varepsilon \mathcal{F} - k \psi. \quad (2-11)$$

3. The C^2 estimate

In this section, we give the proof of the estimate stated in the theorem. When $k = 1$, (1-2) becomes

$$\Delta_\omega u + \text{Tr}_\omega \chi(z, u) = n\psi(z, Du, u), \tag{3-1}$$

where Δ_ω and Tr_ω are the Laplacian and trace with respect to the background metric ω . It follows that $\Delta_\omega u$ is bounded, and the desired estimate follows in turn from the positivity of the metric g . Henceforth, we assume that $k \geq 2$. Motivated by the idea from [Guan et al. 2015] for real Hessian equations, we apply the maximum principle to the test function

$$G = \log P_m + mN|Du|^2 - mMu, \tag{3-2}$$

where $P_m = \sum_j \lambda_j^m$. Here, m, M and N are large positive constants to be determined later. We may assume that the maximum of G is achieved at some point $z \in X$. After rotating the coordinates, we may assume that the matrix $g_{\bar{j}i} = \chi_{\bar{j}i} + u_{\bar{j}i}$ is diagonal.

Recall that if $F(A) = f(\lambda_1, \dots, \lambda_n)$ is a symmetric function of the eigenvalues of a Hermitian matrix $A = (a_{\bar{j}i})$, then at a diagonal matrix A with distinct eigenvalues, we have

$$F^{i\bar{j}} = \delta_{ij} f_i, \tag{3-3}$$

$$F^{i\bar{j}, r\bar{s}} w_{i\bar{j}k} w_{r\bar{s}\bar{k}} = \sum f_{ij} w_{i\bar{i}k} w_{j\bar{j}\bar{k}} + \sum_{p \neq q} \frac{f_p - f_q}{\lambda_p - \lambda_q} |w_{p\bar{q}k}|^2, \tag{3-4}$$

where

$$F^{i\bar{j}} = \frac{\partial F}{\partial a_{\bar{j}i}}, \quad F^{i\bar{j}, r\bar{s}} = \frac{\partial^2 F}{\partial a_{\bar{j}i} \partial a_{\bar{s}r}},$$

and $w_{i\bar{j}k}$ is an arbitrary tensor; see [Ball 1984]. Using these identities to differentiate G , we first obtain the critical equation

$$\frac{DP_m}{P_m} + mND|Du|^2 - mMDu = 0. \tag{3-5}$$

Differentiating G a second time and contracting with $\sigma_k^{p\bar{q}}$ yields

$$\begin{aligned} 0 \geq & \frac{m}{P_m} \left(\sum_j \lambda_j^{m-1} \sigma_k^{p\bar{p}} D_p D_{\bar{p}} g_{\bar{j}j} \right) - \frac{|DP_m|_\sigma^2}{P_m^2} + mN\sigma_k^{p\bar{p}} |Du|_{\bar{p}p}^2 - mM\sigma_k^{p\bar{p}} u_{\bar{p}p} \\ & + \frac{m}{P_m} \left((m-1) \sum_j \lambda_j^{m-2} \sigma_k^{p\bar{p}} |D_p g_{\bar{j}j}|^2 + \sigma_k^{p\bar{p}} \sum_{i \neq j} \frac{\lambda_i^{m-1} - \lambda_j^{m-1}}{\lambda_i - \lambda_j} |D_p g_{\bar{j}i}|^2 \right). \end{aligned} \tag{3-6}$$

Here, we use the notation $|\eta|_\sigma^2 = \sigma_k^{p\bar{q}} \eta_p \eta_{\bar{q}}$. Substituting (2-6), (2-10) and (2-11), we obtain

$$\begin{aligned}
 0 \geq & \frac{1}{P_m} \left(-C \sum_j \lambda_j^{m-1} (1 + |DDu|^2 + |D\bar{D}u|^2 + (1 + \lambda_1)\mathcal{F}) \right) \\
 & + \frac{1}{P_m} \left(\sum_j \lambda_j^{m-1} (-\sigma_k^{p\bar{q},r\bar{s}} D_j g_{\bar{q}p} D_{\bar{j}} g_{\bar{s}r} + \sum_\ell \psi_{v_\ell} g_{j\ell} + \sum_\ell \psi_{\bar{v}_\ell} g_{\bar{j}\bar{\ell}}) \right) \\
 & + \frac{1}{P_m} \left((m-1) \sum_j \lambda_j^{m-2} \sigma_k^{p\bar{p}} |D_p g_{\bar{j}j}|^2 + \sigma_k^{p\bar{p}} \sum_{i \neq j} \frac{\lambda_i^{m-1} - \lambda_j^{m-1}}{\lambda_i - \lambda_j} |D_p g_{ji}|^2 \right) \\
 & - \frac{|DP_m|_\sigma^2}{mP_m^2} + N(|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2) \\
 & + N\langle D|Du|^2, D_{\bar{v}}\psi \rangle + N\langle D_{\bar{v}}\psi, D|Du|^2 \rangle + (M\varepsilon - CN)\mathcal{F} - kM\psi. \tag{3-7}
 \end{aligned}$$

From the critical equation (3-5), we have

$$\frac{1}{P_m} \sum_{j,\ell} \lambda_j^{m-1} g_{j\ell} \psi_{v_\ell} = \frac{1}{m} \left\langle \frac{DP_m}{P_m}, D_{\bar{v}}\psi \right\rangle = -N\langle D|Du|^2, D_{\bar{v}}\psi \rangle + M\langle Du, D_{\bar{v}}\psi \rangle.$$

It follows that

$$\begin{aligned}
 \frac{1}{P_m} \sum_{j,\ell} (\psi_{v_\ell} g_{j\ell} + \psi_{\bar{v}_\ell} g_{\bar{j}\bar{\ell}}) + N\langle D|Du|^2, D_{\bar{v}}\psi \rangle + N\langle D_{\bar{v}}\psi, D|Du|^2 \rangle \\
 = M(\langle Du, D_{\bar{v}}\psi \rangle + \langle D_{\bar{v}}\psi, Du \rangle) \geq -CM.
 \end{aligned}$$

Using (3-4), one can obtain the well-known identity

$$-\sigma_k^{p\bar{q},r\bar{s}} D_j g_{\bar{q}p} D_{\bar{j}} g_{\bar{s}r} = -\sigma_k^{p\bar{p},q\bar{q}} D_j g_{\bar{p}p} D_{\bar{j}} g_{\bar{q}q} + \sigma_k^{p\bar{p},q\bar{q}} |D_j g_{\bar{p}q}|^2, \tag{3-8}$$

where

$$\sigma_k^{p\bar{p},q\bar{q}} = \frac{\partial}{\partial \lambda_p} \frac{\partial}{\partial \lambda_q} \sigma_k(\lambda).$$

We assume that $\lambda_1 \gg 1$, otherwise the C^2 estimate is complete. The main inequality (3-7) becomes

$$\begin{aligned}
 0 \geq & \frac{-C}{\lambda_1} (1 + |DDu|^2 + |D\bar{D}u|^2) + \frac{1}{P_m} \left(\sum_j \lambda_j^{m-1} (-\sigma_k^{p\bar{p},q\bar{q}} D_j g_{\bar{p}p} D_{\bar{j}} g_{\bar{q}q} + \sigma_k^{p\bar{p},q\bar{q}} |D_j g_{\bar{p}q}|^2) \right) \\
 & + \frac{1}{P_m} \left((m-1) \sum_j \lambda_j^{m-2} \sigma_k^{p\bar{p}} |D_p g_{\bar{j}j}|^2 + \sigma_k^{p\bar{p}} \sum_{i \neq j} \frac{\lambda_i^{m-1} - \lambda_j^{m-1}}{\lambda_i - \lambda_j} |D_p g_{ji}|^2 \right) \\
 & - \frac{|DP_m|_\sigma^2}{mP_m^2} + N(|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2) + (M\varepsilon - CN - C)\mathcal{F} - CM. \tag{3-9}
 \end{aligned}$$

The main objective is to show that the third-order terms on the right-hand side of (3-9) are nonnegative. To deal with this issue, we need a lemma from [Guan et al. 2015]; see also [Guan et al. 2012; Li et al. 2016].

Lemma 1 [Guan et al. 2015]. *Suppose $1 \leq \ell < k \leq n$, and let $\alpha = 1/(k - \ell)$. Let $W = (w_{\bar{q}p})$ be a Hermitian tensor in the Γ_k cone. Then for any $\theta > 0$,*

$$\begin{aligned}
 & -\sigma_k^{p\bar{p},q\bar{q}}(W)w_{\bar{p}pi}w_{\bar{q}q\bar{i}} + \left(1 - \alpha + \frac{\alpha}{\theta}\right) \frac{|D_i\sigma_k(W)|^2}{\sigma_k(W)} \\
 & \geq \sigma_k(W)(\alpha + 1 - \alpha\theta) \left| \frac{D_i\sigma_\ell(W)}{\sigma_\ell(W)} \right|^2 - \frac{\sigma_k}{\sigma_\ell}(W)\sigma_\ell^{p\bar{p},q\bar{q}}(W)w_{\bar{p}pi}w_{\bar{q}q\bar{i}}. \tag{3-10}
 \end{aligned}$$

Here the Γ_k cone is defined as

$$\Gamma_k = \{\lambda \in \mathbb{R}^n \mid \sigma_m(\lambda) > 0, m = 1, \dots, k\}. \tag{3-11}$$

We say a Hermitian matrix $W \in \Gamma_k$ if $\lambda(W) \in \Gamma_k$.

It follows from the above lemma that, by taking $\ell = 1$, we have

$$-\sigma_k^{p\bar{p},q\bar{q}}D_i g_{\bar{p}p}D_{\bar{i}}g_{\bar{q}q} + K|D_i\sigma_k|^2 \geq 0, \tag{3-12}$$

for $K > (1 - \alpha + \alpha/\theta)(\inf \psi)^{-1}$ if $2 \leq k \leq n$.

We denote

$$\begin{aligned}
 A_i &= \frac{\lambda_i^{m-1}}{P_m} (K|D_i\sigma_k|^2 - \sigma_k^{p\bar{p},q\bar{q}}D_i g_{\bar{p}p}D_{\bar{i}}g_{\bar{q}q}), \\
 B_i &= \frac{1}{P_m} \left(\sum_p \sigma_k^{p\bar{p},i\bar{i}} \lambda_p^{m-1} |D_i g_{\bar{p}p}|^2 \right), & C_i &= \frac{(m-1)\sigma_k^{i\bar{i}}}{P_m} \left(\sum_p \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 \right), \\
 D_i &= \frac{1}{P_m} \left(\sum_{p \neq i} \sigma_k^{p\bar{p}} \frac{\lambda_p^{m-1} - \lambda_i^{m-1}}{\lambda_p - \lambda_i} |D_i g_{\bar{p}p}|^2 \right), & E_i &= \frac{m\sigma_k^{i\bar{i}}}{P_m^2} \left| \sum_p \lambda_p^{m-1} D_i g_{\bar{p}p} \right|^2.
 \end{aligned}$$

Define $T_{j\bar{p}q} = D_j \chi_{\bar{p}q} - D_q \chi_{\bar{p}j}$. For any $0 < \tau < 1$, we can estimate

$$\begin{aligned}
 & \frac{1}{P_m} \left(\sum_p \lambda_p^{m-1} \sigma_k^{j\bar{j},i\bar{i}} |D_p g_{\bar{j}i}|^2 \right) \geq \frac{1}{P_m} \left(\sum_p \lambda_p^{m-1} \sigma_k^{p\bar{p},i\bar{i}} |D_i g_{\bar{p}p} + T_{p\bar{p}i}|^2 \right) \\
 & \geq \frac{1}{P_m} \left(\sum_p \lambda_p^{m-1} \sigma_k^{p\bar{p},i\bar{i}} ((1 - \tau)|D_i g_{\bar{p}p}|^2 - C_\tau |T_{p\bar{p}i}|^2) \right) \\
 & = (1 - \tau) \sum_i B_i - \frac{C_\tau}{P_m} \sum_p \lambda_p^{m-2} (\lambda_p \sigma_k^{p\bar{p},i\bar{i}}) |T_{p\bar{p}i}|^2.
 \end{aligned}$$

Now we use $\sigma_l(\lambda|i)$ and $\sigma_l(\lambda|ij)$ to denote the l -th elementary functions of

$$(\lambda|i) = (\lambda_1, \dots, \widehat{\lambda}_i, \dots, \lambda_n) \in \mathbb{R}^{n-1} \quad \text{and} \quad (\lambda|ij) = (\lambda_1, \dots, \widehat{\lambda}_i, \dots, \widehat{\lambda}_j, \dots, \lambda_n) \in \mathbb{R}^{n-2},$$

respectively. The following simple identities are used frequently:

$$\sigma_k^{i\bar{i}} = \sigma_{k-1}(\lambda|i) \quad \text{and} \quad \sigma_k^{p\bar{p},i\bar{i}} = \sigma_{k-2}(\lambda|pi).$$

Using the identity $\sigma_l(\lambda) = \sigma_l(\lambda|p) + \lambda_p\sigma_{l-1}(\lambda|p)$ for any $1 \leq p \leq n$, we obtain

$$\begin{aligned} \frac{1}{P_m} \left(\sum_p \lambda_p^{m-1} \sigma_k^{j\bar{j}, i\bar{i}} |D_p g_{ji}|^2 \right) &\geq (1 - \tau) \sum_i B_i - \frac{C_\tau}{P_m} \sum_p \lambda_p^{m-2} (\sigma_k^{i\bar{i}} - \sigma_{k-1}(\lambda|pi)) |T_{p\bar{p}i}|^2 \\ &\geq (1 - \tau) \sum_i B_i - \frac{C_\tau}{\lambda_1^2} \mathcal{F} \geq (1 - \tau) \sum_i B_i - \mathcal{F}. \end{aligned} \tag{3-13}$$

We use the notation C_τ for a constant depending on τ . To get the last inequality above, we assume that $\lambda_1^2 \geq C_\tau$; otherwise, we already have the desired estimate $\lambda_1 \leq C$. Similarly, we may estimate

$$\begin{aligned} \frac{1}{P_m} \sigma_k^{j\bar{j}} \sum_{i \neq p} \frac{\lambda_i^{m-1} - \lambda_p^{m-1}}{\lambda_i - \lambda_p} |D_j g_{\bar{p}i}|^2 &\geq \frac{1}{P_m} \sigma_k^{p\bar{p}} \sum_{p \neq i} \frac{\lambda_i^{m-1} - \lambda_p^{m-1}}{\lambda_i - \lambda_p} |D_i g_{\bar{p}p} + T_{p\bar{p}i}|^2 \\ &\geq \frac{1}{P_m} \sigma_k^{p\bar{p}} \sum_{p \neq i} \frac{\lambda_i^{m-1} - \lambda_p^{m-1}}{\lambda_i - \lambda_p} ((1 - \tau) |D_i g_{\bar{p}p}|^2 - C_\tau |T_{p\bar{p}i}|^2) \\ &\geq \sum_i (1 - \tau) D_i - \frac{C_\tau}{\lambda_1^2} \mathcal{F} \geq \sum_i (1 - \tau) D_i - \mathcal{F}. \end{aligned} \tag{3-14}$$

With the introduced notation in place, the main inequality becomes

$$\begin{aligned} 0 \geq \frac{-C(K)}{\lambda_1} (1 + |DDu|^2 + |D\bar{D}u|^2) - \tau \frac{|DP_m|_\sigma^2}{mP_m^2} \\ + \sum_i (A_i + (1 - \tau)B_i + C_i + (1 - \tau)D_i - (1 - \tau)E_i) \\ + N(|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2) + (M\varepsilon - CN - C)\mathcal{F} - CM. \end{aligned} \tag{3-15}$$

Using the critical equation (3-5), we have

$$\begin{aligned} \tau \frac{|DP_m|_\sigma^2}{mP_m^2} &= \tau m |ND|Du|^2 - MDu|_\sigma^2 \leq 2\tau m (N^2 |D|Du|^2|_\sigma^2 + M^2 |Du|_\sigma^2) \\ &\leq C\tau m N^2 (|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2) + C\tau m M^2 \mathcal{F}. \end{aligned} \tag{3-16}$$

We thus have

$$\begin{aligned} 0 \geq \frac{-C(K)}{\lambda_1} (1 + |DDu|^2 + |D\bar{D}u|^2) + (N - C\tau m N^2) (|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2) \\ + \sum_i (A_i + (1 - \tau)B_i + C_i + (1 - \tau)D_i - (1 - \tau)E_i) \\ + (M\varepsilon - C\tau m M^2 - CN - C)\mathcal{F} - CM. \end{aligned} \tag{3-17}$$

3.1. Estimating the third-order terms. In this subsection, we will adapt the argument in [Li et al. 2016] to estimate the third-order terms.

Lemma 2. *For sufficiently large m , the following estimates hold:*

$$P_m^2(B_1 + C_1 + D_1 - E_1) \geq P_m \lambda_1^{m-2} \sum_{p \neq 1} \sigma_k^{p\bar{p}} |D_1 g_{\bar{p}p}|^2 - \lambda_1^m \sigma_k^{1\bar{1}} \lambda_1^{m-2} |D_1 g_{\bar{1}1}|^2, \quad (3-18)$$

and for any fixed $i \neq 1$,

$$P_m^2(B_i + C_i + D_i - E_i) \geq 0. \quad (3-19)$$

Proof. Fix $i \in \{1, 2, \dots, n\}$. First, we compute

$$\begin{aligned} P_m(B_i + D_i) &= \sum_{p \neq i} \sigma_k^{p\bar{p}, i\bar{i}} \lambda_p^{m-1} |D_i g_{\bar{p}p}|^2 + \sum_{p \neq i} \sigma_k^{p\bar{p}} \frac{\lambda_p^{m-1} - \lambda_i^{m-1}}{\lambda_p - \lambda_i} |D_i g_{\bar{p}p}|^2 \\ &= \sum_{p \neq i} \lambda_p^{m-2} ((\lambda_p \sigma_k^{p\bar{p}, i\bar{i}} + \sigma_k^{p\bar{p}}) |D_i g_{\bar{p}p}|^2) + \left(\sum_{p \neq i} \sigma_k^{p\bar{p}} \sum_{q=0}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2 \right). \end{aligned}$$

Note that

$$\lambda_p \sigma_k^{p\bar{p}, i\bar{i}} + \sigma_k^{p\bar{p}} \geq \sigma_k^{i\bar{i}}.$$

To see this, we write

$$\begin{aligned} \lambda_p \sigma_k^{p\bar{p}, i\bar{i}} + \sigma_k^{p\bar{p}} &= \lambda_p \sigma_{k-2}(\lambda | pi) + \sigma_{k-1}(\lambda | p) \\ &= \sigma_{k-1}(\lambda | i) - \sigma_{k-1}(\lambda | ip) + \sigma_{k-1}(\lambda | p) \\ &= \sigma_{k-1}(\lambda | i) + \lambda_i \sigma_{k-2}(\lambda | ip) \geq \sigma_{k-1}(\lambda | i) = \sigma_k^{i\bar{i}}, \end{aligned}$$

where we used the standard identity $\sigma_l(\lambda) = \sigma_l(\lambda | p) + \lambda_p \sigma_{l-1}(\lambda | p)$ twice, to get the second and third equalities. Therefore

$$P_m(B_i + D_i) \geq \sigma_k^{i\bar{i}} \left(\sum_{p \neq i} \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 \right) + \left(\sum_{p \neq i} \sigma_k^{p\bar{p}} \sum_{q=0}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2 \right). \quad (3-20)$$

It follows that

$$\begin{aligned} P_m(B_i + C_i + D_i) &\geq m \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 + (m-1) \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{\bar{i}i}|^2 \\ &\quad + \sum_{p \neq i} \sigma_k^{p\bar{p}} \sum_{q=0}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2. \end{aligned} \quad (3-21)$$

Expanding out the definition of E_i ,

$$P_m^2 E_i = m \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_p^{2m-2} |D_i g_{\bar{p}p}|^2 + m \sigma_k^{i\bar{i}} \lambda_i^{2m-2} |D_i g_{\bar{i}i}|^2 + m \sigma_k^{i\bar{i}} \sum_p \sum_{q \neq p} \lambda_p^{m-1} \lambda_q^{m-1} D_i g_{\bar{p}p} D_{\bar{i}} g_{\bar{q}q}. \quad (3-22)$$

Therefore,

$$\begin{aligned}
 & P_m^2(B_i + C_i + D_i - E_i) \\
 & \geq \left(m\sigma_k^{i\bar{i}} \sum_{p \neq i} (P_m - \lambda_p^m) \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 - m\sigma_k^{i\bar{i}} \sum_{p \neq i} \sum_{q \neq p, i} \lambda_p^{m-1} \lambda_q^{m-1} D_i g_{\bar{p}p} D_{\bar{i}} g_{\bar{q}q} \right) \\
 & \quad + P_m \sum_{p \neq i} \sigma_k^{p\bar{p}} \sum_{q=0}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2 - 2m\sigma_k^{i\bar{i}} \operatorname{Re} \sum_{q \neq i} \lambda_i^{m-1} \lambda_q^{m-1} D_i g_{\bar{i}i} D_{\bar{i}} g_{\bar{q}q} \\
 & \quad \quad \quad + ((m-1)P_m - m\lambda_i^m) \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{\bar{i}i}|^2. \quad (3-23)
 \end{aligned}$$

We now estimate the expression on the second line above. First,

$$m\sigma_k^{i\bar{i}} \sum_{p \neq i} (P_m - \lambda_p^m) \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 = m\sigma_k^{i\bar{i}} \sum_{p \neq i} \sum_{q \neq p, i} \lambda_q^m \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 + m\sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^m \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2.$$

Next, we can estimate

$$\begin{aligned}
 -m\sigma_k^{i\bar{i}} \sum_{p \neq i} \sum_{q \neq p, i} \lambda_p^{m-1} \lambda_q^{m-1} D_i g_{\bar{p}p} D_{\bar{i}} g_{\bar{q}q} & \geq -m\sigma_k^{i\bar{i}} \sum_{p \neq i} \sum_{q \neq p, i} \frac{1}{2} (\lambda_p^{m-2} \lambda_q^m |D_i g_{\bar{p}p}|^2 + \lambda_p^m \lambda_q^{m-2} |D_i g_{\bar{q}q}|^2) \\
 & = -m\sigma_k^{i\bar{i}} \sum_{p \neq i} \sum_{q \neq p, i} \lambda_p^{m-2} \lambda_q^m |D_i g_{\bar{p}p}|^2. \quad (3-24)
 \end{aligned}$$

We arrive at

$$\begin{aligned}
 & P_m^2(B_i + C_i + D_i - E_i) \\
 & \geq m\sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^m \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 + P_m \sum_{p \neq i} \sigma_k^{p\bar{p}} \sum_{q=0}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2 \\
 & \quad - 2m\sigma_k^{i\bar{i}} \operatorname{Re} \left(\lambda_i^{m-1} D_i g_{\bar{i}i} \sum_{q \neq i} \lambda_q^{m-1} D_{\bar{i}} g_{\bar{q}q} \right) + ((m-1)P_m - m\lambda_i^m) \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{\bar{i}i}|^2. \quad (3-25)
 \end{aligned}$$

The next step is to extract good terms from the second summation on the first line. We fix $p \neq i$.

Case 1: $\lambda_i \geq \lambda_p$. Then $\sigma_k^{p\bar{p}} \geq \sigma_k^{i\bar{i}}$. Hence

$$P_m \sigma_k^{p\bar{p}} \sum_{q=1}^{m-3} \lambda_p^q \lambda_i^{m-2-q} \geq \lambda_i^m \sigma_k^{i\bar{i}} \sum_{q=1}^{m-3} \lambda_p^q \lambda_p^{m-2-q} = (m-3) \sigma_k^{i\bar{i}} \lambda_i^m \lambda_p^{m-2}. \quad (3-26)$$

Case 2: $\lambda_i \leq \lambda_p$. Then $\lambda_p \sigma_k^{p\bar{p}} = \lambda_i \sigma_k^{i\bar{i}} + (\sigma_k(\lambda|i) - \sigma_k(\lambda|p)) \geq \lambda_i \sigma_k^{i\bar{i}}$, and we obtain

$$P_m \sigma_k^{p\bar{p}} \sum_{q=1}^{m-3} \lambda_p^q \lambda_i^{m-2-q} \geq \lambda_p^m \sigma_k^{i\bar{i}} \sum_{q=1}^{m-3} \lambda_p^{q-1} \lambda_i^{m-1-q} \geq (m-3) \sigma_k^{i\bar{i}} \lambda_i^m \lambda_p^{m-2}. \quad (3-27)$$

Combining both cases, we have

$$\begin{aligned}
 P_m \sigma_k^{p\bar{p}} \sum_{q=0}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2 &= P_m \sigma_k^{p\bar{p}} \sum_{q=1}^{m-3} \lambda_p^q \lambda_i^{m-2-q} |D_i g_{\bar{p}p}|^2 + P_m \sigma_k^{p\bar{p}} \lambda_i^{m-2} |D_i g_{\bar{p}p}|^2 \\
 &\geq (m-3) \sigma_k^{i\bar{i}} \lambda_i^m \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 + P_m \sigma_k^{p\bar{p}} \lambda_i^{m-2} |D_i g_{\bar{p}p}|^2.
 \end{aligned}$$

Substituting this estimate into inequality (3-25), we obtain

$$\begin{aligned}
 &P_m^2 (B_i + C_i + D_i - E_i) \\
 &\geq (2m-3) \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^m \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 - 2m \sigma_k^{i\bar{i}} \operatorname{Re} \left(\lambda_i^{m-1} D_i g_{i\bar{i}} \sum_{p \neq i} \lambda_p^{m-1} D_i g_{\bar{p}p} \right) \\
 &\quad + P_m \lambda_i^{m-2} \sum_{p \neq i} \sigma_k^{p\bar{p}} |D_i g_{\bar{p}p}|^2 + ((m-1)P_m - m \lambda_i^m) \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{i\bar{i}}|^2. \tag{3-28}
 \end{aligned}$$

Choose $m \gg 1$ such that

$$m^2 \leq (2m-3)(m-2). \tag{3-29}$$

We can therefore estimate

$$\begin{aligned}
 &2m \sigma_k^{i\bar{i}} \operatorname{Re} \left(\lambda_i^{m-1} D_i g_{i\bar{i}} \sum_{p \neq i} \lambda_p^{m-1} D_i g_{\bar{p}p} \right) \\
 &\leq 2 \sigma_k^{i\bar{i}} \sum_{p \neq i} ((2m-3)^{1/2} \lambda_i^{m/2} \lambda_p^{(m-2)/2} |D_i g_{\bar{p}p}|) ((m-2)^{1/2} \lambda_i^{(m-2)/2} \lambda_p^{m/2} |D_i g_{i\bar{i}}|) \\
 &\leq (2m-3) \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^m \lambda_p^{m-2} |D_i g_{\bar{p}p}|^2 + (m-2) \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^{m-2} \lambda_p^m |D_i g_{i\bar{i}}|^2. \tag{3-30}
 \end{aligned}$$

We finally arrive at

$$\begin{aligned}
 P_m^2 (B_i + C_i + D_i - E_i) &\geq P_m \lambda_i^{m-2} \sum_{p \neq i} \sigma_k^{p\bar{p}} |D_i g_{\bar{p}p}|^2 + ((m-1)P_m - m \lambda_i^m) \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{i\bar{i}}|^2 \\
 &\quad - (m-2) \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^{m-2} \lambda_p^m |D_i g_{i\bar{i}}|^2. \tag{3-31}
 \end{aligned}$$

If we let $i = 1$, we obtain inequality (3-18). For any fixed $i \neq 1$, this inequality yields

$$\begin{aligned}
 P_m^2 (B_i + C_i + D_i - E_i) &\geq P_m \lambda_i^{m-2} \sum_{p \neq i} \sigma_k^{p\bar{p}} |D_i g_{\bar{p}p}|^2 + ((m-1)\lambda_1^m - \lambda_i^m) \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{i\bar{i}}|^2 \\
 &\quad + (m-1) \sum_{p \neq 1, i} \lambda_p^m \sigma_k^{i\bar{i}} \lambda_i^{m-2} |D_i g_{i\bar{i}}|^2 - (m-2) \sigma_k^{i\bar{i}} \sum_{p \neq i} \lambda_i^{m-2} \lambda_p^m |D_i g_{i\bar{i}}|^2 \\
 &\geq P_m \lambda_i^{m-2} \sum_{p \neq i} \sigma_k^{p\bar{p}} |D_i g_{\bar{p}p}|^2 \geq 0.
 \end{aligned}$$

This completes the proof of Lemma 2. □

We observed in (3-12) that $A_i \geq 0$. Lemma 2 implies that for any $i \neq 1$,

$$A_i + B_i + C_i + D_i - E_i \geq 0.$$

Thus we have shown that for $i \neq 1$, the third-order terms in the main inequality (3-17) are indeed nonnegative. The only remaining case is when $i = 1$. By adapting once again the techniques from [Guan et al. 2015], we obtain the following lemma.

Lemma 3. *Let $1 < k \leq n$. Suppose there exists $0 < \delta \leq 1$ such that $\lambda_\mu \geq \delta \lambda_1$ for some $\mu \in \{1, 2, \dots, k-1\}$. There exists a small $\delta' > 0$ such that if $\lambda_{\mu+1} \leq \delta' \lambda_1$, then*

$$A_1 + B_1 + C_1 + D_1 - E_1 \geq 0.$$

Proof. By Lemma 2, we have

$$P_m^2(A_1 + B_1 + C_1 + D_1 - E_1) \geq P_m^2 A_1 + P_m \lambda_1^{m-2} \sum_{p \neq 1} \sigma_k^{p\bar{p}} |D_1 g_{\bar{p}p}|^2 - \lambda_1^m \sigma_k^{1\bar{1}} \lambda_1^{m-2} |D_1 g_{\bar{1}1}|^2. \tag{3-32}$$

The key insight in [Guan et al. 2015], used also in [Li et al. 2016], is to extract a good term involving $|D_1 g_{\bar{1}1}|^2$ from A_1 . By the inequality in Lemma 1 (with $\theta = 1/2$), we have for $\mu < k$

$$\begin{aligned} P_m^2 A_1 &\geq \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} \left(\left(1 + \frac{\alpha}{2}\right) \left| \sum_p \sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p} \right|^2 - \sigma_\mu \sigma_\mu^{p\bar{p}, q\bar{q}} D_1 g_{\bar{p}p} D_{\bar{1}} g_{\bar{q}q} \right) \\ &= \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} \left(\sum_p \left(1 + \frac{\alpha}{2}\right) |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 + \sum_{p \neq q} \frac{\alpha}{2} \sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p} \sigma_\mu^{q\bar{q}} D_{\bar{1}} g_{\bar{q}q} \right. \\ &\quad \left. + \sum_{p \neq q} (\sigma_\mu^{p\bar{p}} \sigma_\mu^{q\bar{q}} - \sigma_\mu \sigma_\mu^{p\bar{p}, q\bar{q}}) D_1 g_{\bar{p}p} D_{\bar{1}} g_{\bar{q}q} \right) \\ &\geq \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} \left(\sum_p |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 - \sum_{p \neq q} |F^{pq} D_1 g_{\bar{p}p} D_{\bar{1}} g_{\bar{q}q}| \right), \end{aligned} \tag{3-33}$$

where we defined $F^{pq} = \sigma_\mu^{p\bar{p}} \sigma_\mu^{q\bar{q}} - \sigma_\mu \sigma_\mu^{p\bar{p}, q\bar{q}}$. Notice that if $\mu = 1$, then $F^{pq} = 1$. If $\mu \geq 2$, then the Newton–Maclaurin inequality implies

$$F^{pq} = \sigma_{\mu-1}^2(\lambda|pq) - \sigma_\mu(\lambda|pq) \sigma_{\mu-2}(\lambda|pq) \geq 0. \tag{3-34}$$

We split the sum involving F^{pq} in the following way:

$$\sum_{p \neq q} |F^{pq} D_1 g_{\bar{p}p} D_{\bar{1}} g_{\bar{q}q}| = \sum_{\substack{p \neq q \\ p, q \leq \mu}} F^{pq} |D_1 g_{\bar{p}p}| |D_{\bar{1}} g_{\bar{q}q}| + \sum_{(p, q) \in J} F^{pq} |D_1 g_{\bar{p}p}| |D_{\bar{1}} g_{\bar{q}q}|, \tag{3-35}$$

where J is the set of indices where at least one of $p \neq q$ is strictly greater than μ . The summation of terms in J can be estimated by

$$\begin{aligned} - \sum_{(p,q) \in J} F^{pq} |D_1 g_{\bar{p}p}| |D_{\bar{1}} g_{\bar{q}q}| &\geq - \sum_{(p,q) \in J} \sigma_\mu^{p\bar{p}} \sigma_\mu^{q\bar{q}} |D_1 g_{\bar{p}p}| |D_{\bar{1}} g_{\bar{q}q}| \\ &\geq -\epsilon \sum_{p \leq \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 - C \sum_{p > \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2. \end{aligned} \tag{3-36}$$

If $\mu = 1$, the first term on the right-hand side of (3-35) vanishes and this estimate applies to all terms on the right hand side of (3-35).

If $\mu \geq 2$, we have for $p, q \leq \mu$,

$$\sigma_{\mu-1}(\lambda|pq) \leq C \frac{\lambda_1 \cdots \lambda_{\mu+1}}{\lambda_p \lambda_q} \leq C \frac{\sigma_\mu^{p\bar{p}} \lambda_{\mu+1}}{\lambda_q}. \tag{3-37}$$

Using (3-34) and (3-37), for δ' small enough we can control

$$\begin{aligned} - \sum_{\substack{p \neq q \\ p, q \leq \mu}} F^{pq} |D_1 g_{\bar{p}p}| |D_{\bar{1}} g_{\bar{q}q}| &\geq - \sum_{\substack{p \neq q \\ p, q \leq \mu}} \sigma_{\mu-1}^2(\lambda|pq) |D_1 g_{\bar{p}p}| |D_{\bar{1}} g_{\bar{q}q}| \\ &\geq -C \lambda_{\mu+1}^2 \sum_{\substack{p \neq q \\ p, q \leq \mu}} \frac{\sigma_\mu^{p\bar{p}}}{\lambda_p} |D_1 g_{\bar{p}p}| \frac{\sigma_\mu^{q\bar{q}}}{\lambda_q} |D_{\bar{1}} g_{\bar{q}q}| \\ &\geq -C \sum_{p \leq \mu} \frac{\lambda_{\mu+1}^2}{\lambda_p^2} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 \\ &\geq -C \sum_{p \leq \mu} \frac{\delta'^2}{\delta^2} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 \geq -\epsilon \sum_{p \leq \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2. \end{aligned} \tag{3-38}$$

Combining all cases, we have

$$- \sum_{p \neq q} |F^{pq} D_1 g_{\bar{p}p} D_{\bar{1}} g_{\bar{q}q}| \geq -2\epsilon \sum_{p \leq \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 - C \sum_{p > \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2. \tag{3-39}$$

Using this inequality in (3-33) yields

$$\begin{aligned} P_m^2 A_1 &\geq \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} \left((1 - 2\epsilon) \sum_{p \leq \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 - C \sum_{p > \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2 \right) \\ &\geq (1 - 2\epsilon) \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} |\sigma_\mu^{1\bar{1}} D_1 g_{\bar{1}1}|^2 - C \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} \sum_{p > \mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2. \end{aligned} \tag{3-40}$$

We estimate

$$\begin{aligned}
 (1 - 2\epsilon) \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} |\sigma_\mu^{1\bar{1}} D_1 g_{\bar{1}1}|^2 &= (1 - 2\epsilon) \frac{P_m \lambda_1^{m-2} \sigma_k}{\lambda_1} \left(\frac{\lambda_1 \sigma_\mu^{1\bar{1}}}{\sigma_\mu} \right)^2 |D_1 g_{\bar{1}1}|^2 \\
 &\geq (1 - 2\epsilon) P_m \lambda_1^{m-2} \frac{\sigma_k}{\lambda_1} \left(1 - C \frac{\lambda_{\mu+1}}{\lambda_1} \right)^2 |D_1 g_{\bar{1}1}|^2 \\
 &\geq (1 - 2\epsilon) (1 - C\delta')^2 P_m \lambda_1^{m-2} \sigma_k^{1\bar{1}} |D_1 g_{\bar{1}1}|^2 \\
 &\geq (1 - 2\epsilon) (1 - C\delta')^2 (1 + \delta^m) \lambda_1^{2m-2} \sigma_k^{1\bar{1}} |D_1 g_{\bar{1}1}|^2.
 \end{aligned} \tag{3-41}$$

For δ' and ϵ small enough, we obtain

$$P_m^2 A_1 \geq \lambda_1^m \sigma_k^{1\bar{1}} \lambda_1^{m-2} |D_1 g_{\bar{1}1}|^2 - C \frac{P_m \lambda_1^{m-1} \sigma_k}{\sigma_\mu^2} \sum_{p>\mu} |\sigma_\mu^{p\bar{p}} D_1 g_{\bar{p}p}|^2. \tag{3-42}$$

We see that the $|D_1 g_{\bar{1}1}|^2$ term cancels from the inequality (3-32) and we are left with

$$P_m^2 (A_1 + B_1 + C_1 + D_1 - E_1) \geq P_m \lambda_1^{m-2} \sum_{p>\mu} \left(\sigma_k^{p\bar{p}} - C \frac{\lambda_1 \sigma_k (\sigma_\mu^{p\bar{p}})^2}{\sigma_\mu^2} \right) |D_1 g_{\bar{p}p}|^2. \tag{3-43}$$

For δ' small enough, the above expression is nonnegative. Indeed, for any $p > \mu$, we have

$$(\lambda_1 \sigma_\mu^{p\bar{p}})^2 \leq \frac{1}{\delta^2} (\lambda_\mu \sigma_\mu^{p\bar{p}})^2 \leq C \frac{(\sigma_\mu)^2}{\delta^2}. \tag{3-44}$$

Therefore,

$$C \frac{\lambda_1 \sigma_k (\sigma_\mu^{p\bar{p}})^2}{\sigma_\mu^2} \leq \frac{C}{\delta^2} \frac{\sigma_k}{\lambda_1}. \tag{3-45}$$

On the other hand, we notice that if $p > k$, then

$$\sigma_k^{p\bar{p}} \geq \lambda_1 \cdots \lambda_{k-1} \geq c_n \frac{\sigma_k}{\lambda_k} \geq \frac{c_n}{\delta'} \frac{\sigma_k}{\lambda_1}.$$

If $\mu < p \leq k$, then

$$\sigma_k^{p\bar{p}} \geq \frac{\lambda_1 \cdots \lambda_k}{\lambda_p} \geq c_n \frac{\sigma_k}{\lambda_p} \geq \frac{c_n}{\delta'} \frac{\sigma_k}{\lambda_1}.$$

It follows that for δ' small enough we have

$$\sigma_k^{p\bar{p}} \geq C \frac{\lambda_1 \sigma_k (\sigma_\mu^{p\bar{p}})^2}{\sigma_\mu^2}. \tag{3-46}$$

This completes the proof of [Lemma 3](#). □

3.2. Completing the proof. With Lemma 2 and Lemma 3 at our disposal, we claim that we may assume in inequality (3-17) that

$$A_i + B_i + C_i + D_i - E_i \geq 0, \quad \forall i = 1, \dots, n. \tag{3-47}$$

Indeed, first set $\delta_1 = 1$. If $\lambda_2 \leq \delta_2 \lambda_1$ for $\delta_2 > 0$ small enough, then by Lemma 3 we see that (3-47) holds. Otherwise, $\lambda_2 \geq \delta_2 \lambda_1$. If $\lambda_3 \leq \delta_3 \lambda_1$ for $\delta_3 > 0$ small enough, then by Lemma 3 we see that (3-47) holds. Otherwise, $\lambda_3 \geq \delta_3 \lambda_1$. Proceeding iteratively, we may arrive at $\lambda_k \geq \delta_k \lambda_1$. But in this case, the C^2 estimate follows directly from the equation as

$$C \geq \sigma_k \geq \lambda_1 \cdots \lambda_k \geq (\delta_k)^{k-1} \lambda_1. \tag{3-48}$$

Therefore we may assume (3-47), and inequality (3-17) becomes

$$0 \geq \frac{-C(K)}{\lambda_1} (1 + |DDu|^2 + |D\bar{D}u|^2) + (N - C\tau mN^2)(|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2) + (M\varepsilon - C\tau mM^2 - CN - C)\mathcal{F} - CM. \tag{3-49}$$

Since $\sigma_k^{i\bar{i}} \geq \sigma_k^{1\bar{1}} \geq \frac{k}{n} \frac{\sigma_k}{\lambda_1} \geq \frac{1}{C\lambda_1}$ for fixed i , we can estimate

$$|DDu|_{\sigma\omega}^2 + |D\bar{D}u|_{\sigma\omega}^2 \geq \frac{1}{C\lambda_1} (|DDu|^2 + |D\bar{D}u|^2) \geq \frac{1}{C\lambda_1} |DDu|^2 + \frac{\lambda_1}{C}. \tag{3-50}$$

This leads to

$$0 \geq \left(\frac{N}{C} - C\tau mN^2 - C(K)\right)\lambda_1 + \frac{1}{\lambda_1} \left(\frac{N}{C} - C\tau mN^2 - C(K)\right)(1 + |DDu|^2) + (M\varepsilon - C\tau mM^2 - CN - C)\mathcal{F} - CM.$$

By choosing τ small, for example $\tau = 1/(NM)$, we have

$$0 \geq \left(\frac{N}{C} - \frac{Cm}{M}N - C(K)\right)\lambda_1 + \frac{1}{\lambda_1} \left(\frac{N}{C} - \frac{Cm}{M}N - C(K)\right)(1 + |DDu|^2) + \left(M\varepsilon - \frac{Cm}{N}M - CN - C\right)\mathcal{F} - CM.$$

Taking N and M large enough, we can ensure that the coefficients of the first three terms are positive. For example, if we let $M = N^2$ for N large, then

$$\begin{aligned} \frac{N}{C} - \frac{Cm}{M}N - C(K) &= \frac{N}{C} - \frac{Cm}{N} - C(K) > 0, \\ M\varepsilon - \frac{Cm}{N}M - CN - C &= N^2\varepsilon - CmN - CN - C > 0. \end{aligned}$$

Thus, an upper bound of λ_1 follows. □

Remark 2. In the above estimate, we assume that $\lambda = (\lambda_1, \dots, \lambda_n) \in \Gamma_n$. Indeed, our estimate still works with $\lambda \in \Gamma_{k+1}$. It was observed in [Li et al. 2016, Lemma 7] that if $\lambda \in \Gamma_{k+1}$, then $\lambda_1 \geq \dots \geq \lambda_n > -K_0$ for some positive constant K_0 . Thus, we can replace λ by $\tilde{\lambda} = \lambda + K_0I$ in our test function G in (3-2).

Acknowledgements

The authors would like to thank Pengfei Guan for stimulating conversations and for sharing his unpublished notes. The authors are also very grateful to the referees for an exceptionally careful reading and for many helpful suggestions.

References

- [Alesker and Verbitsky 2010] S. Alesker and M. Verbitsky, “Quaternionic Monge–Ampère equation and Calabi problem for HKT-manifolds”, *Israel J. Math.* **176** (2010), 109–138. [MR 2653188](#) [Zbl 1193.53118](#)
- [Ball 1984] J. M. Ball, “Differentiability properties of symmetric and isotropic functions”, *Duke Math. J.* **51**:3 (1984), 699–728. [MR 757959](#) [Zbl 0566.73001](#)
- [Błocki 2005] Z. Błocki, “Weak solutions to the complex Hessian equation”, *Ann. Inst. Fourier (Grenoble)* **55**:5 (2005), 1735–1756. [MR 2172278](#) [Zbl 1081.32023](#)
- [Dinew and Kołodziej 2014] S. Dinew and S. Kołodziej, “A priori estimates for complex Hessian equations”, *Anal. PDE* **7**:1 (2014), 227–244. [MR 3219505](#) [Zbl 1297.32020](#)
- [Fu and Yau 2007] J.-X. Fu and S.-T. Yau, “A Monge–Ampère-type equation motivated by string theory”, *Comm. Anal. Geom.* **15**:1 (2007), 29–75. [MR 2301248](#) [Zbl 1122.35145](#)
- [Fu and Yau 2008] J.-X. Fu and S.-T. Yau, “The theory of superstring with flux on non-Kähler manifolds and the complex Monge–Ampère equation”, *J. Differential Geom.* **78**:3 (2008), 369–428. [MR 2396248](#) [Zbl 1141.53036](#)
- [Guan 2014] B. Guan, “Second-order estimates and regularity for fully nonlinear elliptic equations on Riemannian manifolds”, *Duke Math. J.* **163**:8 (2014), 1491–1524. [MR 3284698](#) [Zbl 1296.58012](#)
- [Guan et al. 2012] P. Guan, J. Li, and Y. Li, “Hypersurfaces of prescribed curvature measure”, *Duke Math. J.* **161**:10 (2012), 1927–1942. [MR 2954620](#) [Zbl 1254.53073](#)
- [Guan et al. 2015] P. Guan, C. Ren, and Z. Wang, “Global C^2 -estimates for convex solutions of curvature equations”, *Comm. Pure Appl. Math.* **68**:8 (2015), 1287–1325. [MR 3366747](#) [Zbl 1327.53009](#)
- [Hou 2009] Z. Hou, “Complex Hessian equation on Kähler manifold”, *Int. Math. Res. Not.* **2009**:16 (2009), 3098–3111. [MR 2533797](#) [Zbl 1177.32013](#)
- [Hou et al. 2010] Z. Hou, X.-N. Ma, and D. Wu, “A second order estimate for complex Hessian equations on a compact Kähler manifold”, *Math. Res. Lett.* **17**:3 (2010), 547–561. [MR 2653687](#) [Zbl 1225.32026](#)
- [Kołodziej and Nguyen 2016] S. Kołodziej and N. C. Nguyen, “Weak solutions of complex Hessian equations on compact Hermitian manifolds”, *Compos. Math.* (online publication September 2016).
- [Li 2004] S.-Y. Li, “On the Dirichlet problems for symmetric function equations of the eigenvalues of the complex Hessian”, *Asian J. Math.* **8**:1 (2004), 87–106. [MR 2128299](#) [Zbl 1068.32024](#)
- [Li et al. 2016] M. Li, C. Ren, and Z. Wang, “An interior estimate for convex solutions and a rigidity theorem”, *J. Funct. Anal.* **270**:7 (2016), 2691–2714. [MR 3464054](#) [Zbl 1341.35055](#)
- [Lu and Nguyen 2015] C. H. Lu and V.-D. Nguyen, “Degenerate complex Hessian equations on compact Kähler manifolds”, *Indiana Univ. Math. J.* **64**:6 (2015), 1721–1745. [MR 3436233](#) [Zbl 1341.32033](#)
- [Phong et al. 2015] D. H. Phong, S. Picard, and X. Zhang, “On estimates for the Fu–Yau generalization of a Strominger system”, preprint, 2015. [arXiv 1507.08193](#)
- [Song and Weinkove 2008] J. Song and B. Weinkove, “On the convergence and singularities of the J -flow with applications to the Mabuchi energy”, *Comm. Pure Appl. Math.* **61**:2 (2008), 210–229. [MR 2368374](#) [Zbl 1135.53047](#)
- [Spruck and Xiao 2015] J. Spruck and L. Xiao, “A note on starshaped compact hypersurfaces with a prescribed scalar curvature in space forms”, preprint, 2015. [arXiv 1505.01578](#)
- [Sun 2014] W. Sun, “On uniform estimate of complex elliptic equations on closed Hermitian manifolds”, preprint, 2014. [arXiv 1412.5001](#)

- [Székelyhidi 2015] G. Székelyhidi, “Fully non-linear elliptic equations on compact Hermitian manifolds”, preprint, 2015. [arXiv 1501.02762v3](#)
- [Székelyhidi et al. 2015] G. Székelyhidi, V. Tosatti, and B. Weinkove, “Gauduchon metrics with prescribed volume form”, preprint, 2015. [arXiv 1503.04491](#)
- [Wang 2009] X.-J. Wang, “The k -Hessian equation”, pp. 177–252 in *Geometric analysis and PDEs*, edited by A. Ambrosetti et al., Lecture Notes in Mathematics **1977**, Springer, 2009. [MR 2500526](#) [Zbl 1196.35093](#)
- [Zhang 2010] X. Zhang, “A priori estimates for complex Monge–Ampère equation on Hermitian manifolds”, *Int. Math. Res. Not.* **2010**:19 (2010), 3814–3836. [MR 2725515](#) [Zbl 1209.32025](#)
- [Zhang 2015] D. Zhang, “Hessian equations on closed Hermitian manifolds”, preprint, 2015. [arXiv 1501.03553](#)

Received 4 Jan 2016. Revised 29 Apr 2016. Accepted 28 May 2016.

DUONG H. PHONG: phong@math.columbia.edu

Department of Mathematics, Columbia University, New York, NY 10027, United States

SEBASTIEN PICARD: picard@math.columbia.edu

Department of Mathematics, Columbia University, New York, NY 10027, United States

XIANGWEN ZHANG: xiangwen@math.uci.edu

Department of Mathematics, University of California, Irvine, CA 92697, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

| | | | |
|----------------------|---|-----------------------|---|
| Nicolas Burq | Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr | Werner Müller | Universität Bonn, Germany mueller@math.uni-bonn.de |
| Massimiliano Berti | Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it | Gilles Pisier | Texas A&M University, and Paris 6 pisier@math.tamu.edu |
| Sun-Yung Alice Chang | Princeton University, USA chang@math.princeton.edu | Tristan Rivière | ETH, Switzerland riviere@math.ethz.ch |
| Michael Christ | University of California, Berkeley, USA mchrist@math.berkeley.edu | Igor Rodnianski | Princeton University, USA irod@math.princeton.edu |
| Charles Fefferman | Princeton University, USA cf@math.princeton.edu | Wilhelm Schlag | University of Chicago, USA schlag@math.uchicago.edu |
| Ursula Hamenstaedt | Universität Bonn, Germany ursula@math.uni-bonn.de | Sylvia Serfaty | New York University, USA serfaty@cims.nyu.edu |
| Vaughan Jones | U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu | Yum-Tong Siu | Harvard University, USA siu@math.harvard.edu |
| Vadim Kaloshin | University of Maryland, USA vadim.kaloshin@gmail.com | Terence Tao | University of California, Los Angeles, USA tao@math.ucla.edu |
| Herbert Koch | Universität Bonn, Germany koch@math.uni-bonn.de | Michael E. Taylor | Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu |
| Izabella Laba | University of British Columbia, Canada ilaba@math.ubc.ca | Gunther Uhlmann | University of Washington, USA gunther@math.washington.edu |
| Gilles Lebeau | Université de Nice Sophia Antipolis, France lebeau@unice.fr | András Vasy | Stanford University, USA andras@math.stanford.edu |
| Richard B. Melrose | Massachusetts Inst. of Tech., USA rbb@math.mit.edu | Dan Virgil Voiculescu | University of California, Berkeley, USA dvv@math.berkeley.edu |
| Frank Merle | Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr | Steven Zelditch | Northwestern University, USA zelditch@math.northwestern.edu |
| William Minicozzi II | Johns Hopkins University, USA minicozz@math.jhu.edu | Maciej Zworski | University of California, Berkeley, USA zvorski@math.berkeley.edu |
| Clément Mouhot | Cambridge University, UK c.mouhot@dpms.cam.ac.uk | | |

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 7 2016

| | |
|--|------|
| The final-state problem for the cubic-quintic NLS with nonvanishing boundary conditions | 1523 |
| ROWAN KILLIP, JASON MURPHY and MONICA VISAN | |
| Magnetic wells in dimension three | 1575 |
| BERNARD HELFFER, YURI KORDYUKOV, NICOLAS RAYMOND and SAN VŨ NGỌC | |
| An analytical and numerical study of steady patches in the disc | 1609 |
| FRANCISCO DE LA HOZ, ZINEB HASSAINIA, TAOUFIK HMIDI and JOAN MATEU | |
| Isolated singularities of positive solutions of elliptic equations with weighted gradient term | 1671 |
| PHUOC-TAI NGUYEN | |
| A second order estimate for general complex Hessian equations | 1693 |
| DUONG H. PHONG, SEBASTIEN PICARD and XIANGWEN ZHANG | |
| Parabolic weighted norm inequalities and partial differential equations | 1711 |
| JUHA KINNUNEN and OLLI SAARI | |
| A double well potential system | 1737 |
| JAEYOUNG BYEON, PIERO MONTECCHIARI and PAUL H. RABINOWITZ | |