

ANALYSIS & PDE

Volume 17

No. 2

2024

LIANG LI, RUIPENG SHEN AND LIJUAN WEI

**EXPLICIT FORMULA OF RADIATION FIELDS OF FREE WAVES
WITH APPLICATIONS ON CHANNEL OF ENERGY**

EXPLICIT FORMULA OF RADIATION FIELDS OF FREE WAVES WITH APPLICATIONS ON CHANNEL OF ENERGY

LIANG LI, RUIPENG SHEN AND LIJUAN WEI

We give a few explicit formulas regarding the radiation fields of linear free waves. We then apply these formulas on the channel-of-energy theory. We characterize all the radial weakly nonradiative solutions in all dimensions and give a few new exterior energy estimates.

1. Introduction

1A. Background and topics. The semilinear wave equation

$$\partial_t^2 u - \Delta u = \pm |u|^{p-1} u, \quad (x, t) \in \mathbb{R}^d \times \mathbb{R},$$

especially the energy critical case $p = 1 + 4/(d-2)$, has been extensively studied by many mathematicians in the past few decades. Please see, for example, [Kapitanski 1994; Lindblad and Sogge 1995] for local existence and well-posedness, and [Ginibre, Soffer and Velo 1992; Grillakis 1990; 1992; Kenig and Merle 2008; Nakanishi 1999a; 1999b; Shatah and Struwe 1993; 1994] for global existence, regularity, scattering and blow-up. Since the semilinear wave equation can be viewed as a small perturbation of the homogenous linear wave equation in many situations, especially when we consider the asymptotic behaviors of solutions as spatial variables or time tends to infinity, it is important to first understand the asymptotic behaviors of solutions to the homogenous linear wave equation, i.e., free waves. This work is concerned with two important tools to understand the asymptotic behaviors of free waves: radiation fields and the channel of energy. We first introduce some necessary notation. Throughout this work we consider the homogenous linear wave equation with initial data in the energy space

$$\begin{cases} \partial_t^2 u - \Delta u = 0, & (x, t) \in \mathbb{R}^d \times \mathbb{R}, \\ u|_{t=0} = u_0 \in \dot{H}^1(\mathbb{R}^d), \\ u_t|_{t=0} = u_1 \in L^2(\mathbb{R}^d). \end{cases} \quad (1)$$

In this work we also use the notation $\mathcal{S}_L(u_0, u_1)$ to represent the free wave u defined above. If it is necessary to mention the velocity u_t , we use the notation

$$\mathcal{S}_L(t) \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} = \begin{pmatrix} u(\cdot, t) \\ u_t(\cdot, t) \end{pmatrix} \in \dot{H}^1 \times L^2.$$

MSC2020: 35L05.

Keywords: wave equation, radiation field, channel of energy.

It is well known that the linear wave propagation preserves the $\dot{H}^1 \times L^2$ norm, i.e., the energy conservation law holds $(\nabla_{x,t} u = (\nabla u, u_t))$:

$$\int_{\mathbb{R}^d} |\nabla_{x,t} u(x, t)|^2 dx = \int_{\mathbb{R}^d} (|\nabla u_0|^2 + |u_1|^2) dx.$$

Now we make a brief review of radiation fields and the channel-of-energy method.

Radiation fields. The asymptotic behavior of free waves at the energy level can be characterized by the following theorem.

Theorem 1.1 (radiation field). *Assume that $d \geq 3$ and let u be a solution to the free wave equation $\partial_t^2 u - \Delta u = 0$, with initial data $(u_0, u_1) \in \dot{H}^1 \times L^2(\mathbb{R}^d)$. Then*

$$\lim_{t \rightarrow \pm\infty} \int_{\mathbb{R}^d} \left(|\nabla u(x, t)|^2 - |u_r(x, t)|^2 + \frac{|u(x, t)|^2}{|x|^2} \right) dx = 0$$

and there exist two functions $G_{\pm} \in L^2(\mathbb{R} \times \mathbb{S}^{d-1})$ so that

$$\begin{aligned} \lim_{t \rightarrow \pm\infty} \int_0^\infty \int_{\mathbb{S}^{d-1}} |r^{(d-1)/2} \partial_t u(r\theta, t) - G_{\pm}(r \mp t, \theta)|^2 d\theta dr &= 0, \\ \lim_{t \rightarrow \pm\infty} \int_0^\infty \int_{\mathbb{S}^{d-1}} |r^{(d-1)/2} \partial_r u(r\theta, t) \pm G_{\pm}(r \mp t, \theta)|^2 d\theta dr &= 0. \end{aligned}$$

In addition, the maps $(u_0, u_1) \rightarrow \sqrt{2}G_{\pm}$ are a bijective isometries from $\dot{H}^1 \times L^2(\mathbb{R}^d)$ to $L^2(\mathbb{R} \times \mathbb{S}^{d-1})$.

This has been known for more than 50 years, at least in the 3-dimensional case. Please see [Friedlander 1962; 1980], for example. The version of the radiation field theorem given above and a proof for all dimensions $d \geq 3$ can be found in [Duyckaerts, Kenig and Merle 2019]. In addition, there is also a 2-dimensional version of the radiation field theorem. The statement in dimension 2 can be given in almost the same way as in the higher-dimensional case, except that the limit

$$\lim_{t \rightarrow \pm\infty} \int_{\mathbb{R}^2} \frac{|u(x, t)|^2}{|x|^2} dx = 0$$

no longer holds. A proof by Radon transform for all dimensions $d \geq 2$ can be found in [Katayama 2013], where the statement of the theorem is slightly different. Throughout this work we call the function G_{\pm} radiation profiles and use the notation T_{\pm} for the linear map $(u_0, u_1) \rightarrow G_{\pm}$.

Channel of energy. The second tool is the channel-of-energy method, which plays an important role in the study of wave equations in the past decade. This method is first introduced in the 3-dimensional case in [Duyckaerts, Kenig and Merle 2011] and then in the 5-dimensional case in [Kenig, Lawrie and Schlag 2014]. This method was used in the proof of the soliton resolution conjecture of the energy critical wave equation with radial data in all odd dimensions $d \geq 3$ in [Duyckaerts, Kenig and Merle 2013; 2023]. It can also be used to show the nonexistence of minimal blow-up solutions in a compactness-rigidity argument in the energy super- or subcritical case. Please see, for example, [Duyckaerts, Kenig and Merle

2014; Shen 2013]. Roughly speaking, the channel-of-energy method discusses the amount of energy located in an exterior region as time tends to infinity:

$$\lim_{t \rightarrow \pm\infty} \int_{|x| > |t| + R} |\nabla_{x,t} u(x, t)|^2 dx.$$

Here the constant R is nonnegative. Since the energy travels at a finite speed, the energy in the exterior region $\{x : |x| > |t| + R\}$ decays as $|t|$ increases. Thus the limits above are always well-defined. We can also give the exact value of the limit in terms of the radiation field:

$$\lim_{t \rightarrow \pm\infty} \int_{|x| > |t| + R} |\nabla_{x,t} u(x, t)|^2 dx = 2 \int_R^\infty \int_{\mathbb{S}^{d-1}} |G_\pm(s, \theta)|^2 d\theta ds. \tag{2}$$

We first introduce a few already known results. We start with the odd dimensions.

Proposition 1.2 [Duyckaerts, Kenig and Merle 2012]. *Assume that $d \geq 3$ is an odd integer. All solutions to $\partial_t^2 u - \Delta u = 0$ satisfy*

$$\sum_{\pm} \lim_{t \rightarrow \pm\infty} \int_{|x| > |t|} |\nabla_{x,t} u(x, t)|^2 dx = \int_{\mathbb{R}^d} |\nabla_{x,t} u(x, 0)|^2 dx. \tag{3}$$

As a result, we have:

Corollary 1.3. *Assume that $d \geq 3$ is odd. Then $u \equiv 0$ is the only free wave u satisfying*

$$\lim_{t \rightarrow \pm\infty} \int_{|x| > |t|} |\nabla_{x,t} u(x, t)|^2 dx = 0.$$

In contrast, if $R > 0$, the subspace of $\dot{H}^1 \times L^2(\mathbb{R}^d)$ defined by

$$P(R) = \left\{ (u_0, u_1) \in \dot{H}^1 \times L^2(\mathbb{R}^d) : \lim_{t \rightarrow \pm\infty} \int_{|x| > R + |t|} |\nabla_{t,x} \mathbf{S}_L(u_0, u_1)(x, t)|^2 dx = 0 \right\} \tag{4}$$

does contain initial data (u_0, u_1) whose support is essentially bigger than $\{x : |x| \leq R\}$. The free waves u satisfying

$$\lim_{t \rightarrow \pm\infty} \int_{|x| > R + |t|} |\nabla_{t,x} u(x, t)|^2 dx = 0$$

are usually called (R -weakly) nonradiative solutions. If the dimension is odd, these solutions are well-understood in the radial case:

Theorem 1.4 [Kenig, Lawrie, Liu and Schlag 2015]. *In any odd dimension $d \geq 1$, every radial solution u to (1) satisfies*

$$\max_{\pm} \lim_{t \rightarrow \pm\infty} \int_{r > |t| + R} |\nabla_{x,t} u(r, t)|^2 r^{d-1} dr \geq \frac{1}{2} \|\Pi_{P_{\text{rad}}(R)}^\perp(u_0, u_1)\|_{\dot{H}^1 \times L^2(r \geq R; r^{d-1} dr)}. \tag{5}$$

Here

$$P_{\text{rad}}(R) \doteq \text{Span} \left\{ (r^{2k_1-d}, 0), (0, r^{2k_2-d}) : k_1, k_2 \in \mathbb{N}; 1 \leq k_1 \leq \frac{d+2}{4}, 1 \leq k_2 \leq \frac{d}{4} \right\}.$$

$\Pi_{P_{\text{rad}}(R)}^\perp$ is the orthogonal projection from $\dot{H}^1 \times L^2(r \geq R : r^{d-1} dr)$ onto the complement of the finite-dimensional subspace $P_{\text{rad}}(R)$.

Note the proof of Theorem 1.4 in [Kenig, Lawrie, Liu and Schlag 2015] uses the radial Fourier transform.

The case of even dimensions is much more complicated and subtle. Côte, Kenig and Schlag [2014] showed that in general the inequality

$$\sum_{\pm} \lim_{t \rightarrow \pm\infty} \int_{|x|>|t|} |\nabla_{x,t} u(x, t)|^2 dx \geq C \int_{\mathbb{R}^d} |\nabla_{x,t} u(x, 0)|^2 dx$$

does not hold for any positive constant C in even dimensions. But a similar inequality holds in the radial case for either initial data $(u_0, 0)$ if $d = 0 \pmod{4}$, or $(0, u_1)$ if $d = 2 \pmod{4}$. More precisely we have

$$\lim_{t \rightarrow \pm\infty} \int_{|x|>|t|} |\nabla_{x,t} \mathcal{S}_L(u_0, 0)(x, t)|^2 dx \geq \frac{1}{2} \int_{\mathbb{R}^d} |\nabla u_0(x)|^2 dx, \quad d = 4k, k \in \mathbb{N}, \quad (6)$$

$$\lim_{t \rightarrow \pm\infty} \int_{|x|>|t|} |\nabla_{x,t} \mathcal{S}_L(0, u_1)(x, t)|^2 dx \geq \frac{1}{2} \int_{\mathbb{R}^d} |u_1(x)|^2 dx, \quad d = 4k + 2, k \in \mathbb{N}. \quad (7)$$

In addition, Duyckaerts, Kenig and Merle [2021] showed that the only nonradiative solution is still the zero solution in even dimensions $d \geq 4$; i.e., Corollary 1.3 still holds for even dimensions $d \geq 4$, even in the nonradial case. Much less is known about the exterior energy estimate in the region $\{x : |x| > R + |t|\}$ with $R > 0$. Duyckaerts, Kenig, Martel and Merle [2022] proves the exterior energy estimate in dimensions 4 and 6 if the initial data are radial:

$$\lim_{t \rightarrow \pm\infty} \int_{|x|>|t|+R} |\nabla_{x,t} \mathcal{S}_L(u_0, 0)(x, t)|^2 dx \geq \frac{3}{10} \|\Pi_R^\perp u_0\|_{\dot{H}^1(\{x \in \mathbb{R}^4 : |x| > R\})}^2,$$

$$\lim_{t \rightarrow \pm\infty} \int_{|x|>|t|+R} |\nabla_{x,t} \mathcal{S}_L(0, u_1)(x, t)|^2 dx \geq \frac{3}{10} \|\pi_R^\perp u_1\|_{L^2(\{x \in \mathbb{R}^6 : |x| > R\})}^2.$$

Here Π_R^\perp is the orthogonal projection from $\dot{H}^1(\{x \in \mathbb{R}^4 : |x| > R\})$ onto the complement space of $\text{Span}\{|x|^{-2}\}$. While π_R^\perp is the orthogonal projection from $L^2(\{x \in \mathbb{R}^6 : |x| > R\})$ onto the complement space of $\text{Span}\{|x|^{-4}\}$.

1B. Main idea. According to (2) we may obtain exterior energy estimates conveniently from the radiation profiles G_\pm . Please note that G_- and G_+ are not independent of each other. In fact the map $T_+ \circ T_-^{-1} : G_- \rightarrow G_+$ is a bijective isometry. If we could find explicit expressions of the maps

$$T_+ \circ T_-^{-1} : G_- \rightarrow G_+, \quad T_-^{-1} : G_- \rightarrow (u_0, u_1), \quad \mathcal{S}_L \circ T_-^{-1} : G_- \rightarrow u,$$

then we would be able to:

(a) Understand how the asymptotic behavior in one time direction determines the behavior in the other time direction. This is known in the odd-dimensional case, as shown (although not stated explicitly) in the proof of Proposition 1.2 in [Duyckaerts, Kenig and Merle 2012]. In this work we will try to figure out the even-dimensional case.

(b) Characterize (weakly) nonradiative solutions, especially in the radial case. We first determine all the radiation profiles G_- so that

$$G_-(s, \theta) = G_+(s, \theta) = 0, \quad s > R \quad \iff \quad \lim_{t \rightarrow \pm\infty} \int_{|x|>|t|+R} |\nabla_{x,t} u(x, t)|^2 dx = 0;$$

then we may obtain all the nonradiative solutions (as well as their initial data) by applying the formula of T_-^{-1} . In particular we prove that radial nonradiative solutions in even dimensions can be characterized in the same way as in odd dimensions.

(c) Prove exterior energy estimates. We generalize the radial exterior energy estimates in 4 and 6 dimensions to all even dimensions; we also prove a nonradial exterior energy estimate in odd dimensions. In both applications (b) and (c) we follow the same roadmap:

$$\text{exterior energy} \quad \leftrightarrow \quad \text{radiation profile} \quad \leftrightarrow \quad \text{initial data.}$$

1C. Main results. Now we give the statement of our results. The details and proofs can be found in subsequent sections.

Theorem 1.5. *Let u be a finite-energy free wave with an even spatial dimension $d \geq 2$ and G_+ , G_- be the radiation profiles associated with u . Then we may give an explicit formula of the operator $T_+ \circ T_-^{-1} : G_- \rightarrow G_+$*

$$G_+(s, \theta) = (-1)^{d/2}(\mathcal{H}G_-)(-s, -\theta).$$

Here \mathcal{H} is the Hilbert transform in the first variable, i.e.,

$$(\mathcal{H}G_-)(-s, -\theta) = \text{p.v.} \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{G_-(\tau, -\theta)}{(-s) - \tau} d\tau.$$

Remark 1.6. A similar but simpler argument shows that if d is odd, then $T_+ \circ T_-^{-1} : G_- \rightarrow G_+$ can be explicitly given by

$$G_+(s, \theta) = (-1)^{(d-1)/2}G_-(-s, -\theta).$$

This can also be verified by assuming that the initial data is smooth and compactly supported, and considering the expression of G_- and G_+ in terms of (u_0, u_1) if d is odd. Please refer to [Duyckaerts, Kenig and Merle 2012]. Since we have $\mathcal{H}^2 = -1$. We may write the odd and even dimensions in a universal formula

$$G_+(s, \theta) = ((-\mathcal{H})^{d-1}G_-)(-s, -\theta).$$

Remark 1.7. It has been proved in Section 3.2 of [Duyckaerts, Kenig and Merle 2021] (in the language of Hankel and Laplace transforms) that the zero function is the only function $f \in L^2(\mathbb{R})$ satisfying

$$f(s) = 0, \quad s > 0, \quad (\mathcal{H}f)(s) = 0, \quad s < 0.$$

It immediately follows that:

Corollary 1.8. *Assume $d \geq 2$. Let Ω be a region in \mathbb{S}^{d-1} . If a finite-energy solution u to the homogenous linear wave equation satisfies*

$$\lim_{t \rightarrow \pm\infty} \int_{|t|}^{\infty} \int_{\pm\Omega} |\nabla_{t,x}u(r\theta, t)|^2 r^{d-1} d\theta dr = 0,$$

then we have

$$\lim_{t \rightarrow \pm\infty} \int_0^{\infty} \int_{\pm\Omega} |\nabla_{t,x}u(r\theta, t)|^2 r^{d-1} d\theta dr = 0.$$

This is an angle-localized version of Corollary 1.3.

Applications on channel of energy. By following the idea described above, we obtain the following results about the channel of energy.

Proposition 1.9 (radial weakly nonradiative solutions). *Let $d \geq 2$ be an integer and $R > 0$ be a constant. If the initial data $(u_0, u_1) \in \dot{H}^1 \times L^2$ are radial, then the corresponding solution to the homogeneous linear wave equation u is R -weakly nonradiative, i.e.,*

$$\lim_{t \rightarrow \pm\infty} \int_{|x| > |t| + R} |\nabla_{t,x} u(x, t)|^2 dx = 0,$$

if and only if the restriction of (u_0, u_1) in the region $\{x \in \mathbb{R}^d : |x| > R\}$ is contained in

$$P_{\text{rad}}(R) = \text{Span} \left\{ (r^{2k_1-d}, 0), (0, r^{2k_2-d}) : 1 \leq k_1 \leq \left\lfloor \frac{d+1}{4} \right\rfloor, 1 \leq k_2 \leq \left\lfloor \frac{d-1}{4} \right\rfloor \right\}.$$

Here the notation $\lfloor q \rfloor$ is the integer part of q . In particular, all radial R -weakly nonradiative solutions in dimension 2 are supported in $\{(x, t) : |x| \leq |t| + R\}$.

Remark 1.10. If d is odd, we have $\lfloor (d+1)/4 \rfloor = \lfloor (d+2)/4 \rfloor$ and $\lfloor (d-1)/4 \rfloor = \lfloor d/4 \rfloor$; thus our result here is the same as the already known result in odd dimension, as given in Theorem 1.4.

Proposition 1.11 (radial exterior estimates in even dimensions). *Let $d = 4k$ with $k \in \mathbb{N}$ and $R > 0$. If initial data $u_0 \in \dot{H}^1(\mathbb{R}^d)$ are radial, then the corresponding solution u to the homogenous linear wave equation with initial data $(u_0, 0)$ satisfies*

$$\lim_{t \rightarrow \infty} \int_{|x| > R+|t|} |\nabla u(x, t)|^2 dx = \lim_{t \rightarrow \infty} \int_{|x| > R+|t|} |u_t(x, t)|^2 dx \geq \frac{1}{4} \|\Pi_{Q_k(R)}^\perp u_0\|_{\dot{H}^1(\{x:|x|>R\})}^2.$$

Here $\Pi_{Q_k(R)}^\perp$ is the orthogonal projection from $\dot{H}^1(\{x \in \mathbb{R}^d : |x| > R\})$ onto the complement of the k -dimensional linear space

$$Q_k(R) = \text{Span} \left\{ \frac{1}{|x|^{4k-2k_1}} : 1 \leq k_1 \leq k \right\}.$$

Similarly if the dimension d satisfies $d = 4k + 2 \geq 2$, with $k \in \{0\} \cup \mathbb{N}$ and the initial data $u_1 \in L^2(\mathbb{R}^d)$ are radial, then the corresponding solution u to the homogenous linear wave equation with initial data $(0, u_1)$ satisfies

$$\lim_{t \rightarrow \infty} \int_{|x| > R+|t|} |\nabla u(x, t)|^2 dx = \lim_{t \rightarrow \infty} \int_{|x| > R+|t|} |u_t(x, t)|^2 dx \geq \frac{1}{4} \|\Pi_{Q'_k(R)}^\perp u_1\|_{L^2(\{x:|x|>R\})}^2.$$

Here $\Pi_{Q'_k(R)}^\perp$ is the orthogonal projection from $L^2(\{x \in \mathbb{R}^d : |x| > R\})$ onto the complement of the k -dimensional linear space

$$Q'_k(R) = \text{Span} \left\{ \frac{1}{|x|^{4k+2-2k_1}} : 1 \leq k_1 \leq k \right\}.$$

Remark 1.12. Given $u_0 \in \dot{H}^1(\mathbb{R}^{4k})$ or $u_1 \in L^2(\mathbb{R}^{4k+2})$, the orthogonal projection of u_0 or u_1 onto the finite-dimensional space $Q_k(R)$ or $Q'_k(R)$ gradually vanishes as $R \rightarrow 0^+$. Therefore if we make $R \rightarrow 0^+$ in Proposition 1.11, we immediately obtain (6) and (7).

Remark 1.13. At the same time as this work was done, Kenig et al. proved radial exterior estimates similar to Proposition 1.11 for even dimensions $d \geq 8$ by using the already-known result in dimension 4 and an induction argument.

Proposition 1.14 (nonradial exterior energy estimates). *Let $d \geq 3$ be an odd integer and $R > 0$ be a constant. Then the following identity holds for all $(u_0, u_1) \in \dot{H}^1 \times L^2(\mathbb{R}^d)$:*

$$\sum_{\pm} \lim_{t \rightarrow \pm\infty} \int_{|x| > R+|t|} |\nabla_{t,x} \mathbf{S}_L(t)(u_0, u_1)(x, t)|^2 dx = \|\Pi_{P(R)}^\perp(u_0, u_1)\|_{\dot{H}^1 \times L^2(\mathbb{R}^d)}^2.$$

Here $\Pi_{P(R)}^\perp$ is the orthogonal projection from $\dot{H}^1 \times L^2(\mathbb{R}^d)$ onto the complement of the closed linear space

$$P(R) = \left\{ (u_0, u_1) \in \dot{H}^1 \times L^2(\mathbb{R}^d) : \lim_{t \rightarrow \pm\infty} \int_{|x| > R+|t|} |\nabla_{t,x} \mathbf{S}_L(u_0, u_1)(x, t)|^2 dx = 0 \right\}.$$

Structure of this work. This work is organized as follows. In Section 2 we deduce an explicit formula of T_-^{-1} in all dimensions. Then in Section 3 we prove the explicit formula of $T_+ \circ T_-^{-1}$ given in Theorem 1.5. The rest of the paper is devoted to the applications in the channel of energy. We characterize radial weakly nonradiative solutions in Section 4, prove radial exterior energy estimate for all even dimensions in Section 5 and finally give a short proof of nonradial exterior energy estimate in odd-dimensional space in Section 6. The Appendix is concerned with Hilbert transform of a family of special functions, since the Hilbert transform is involved in the even dimensions.

Notation. In this work we use the notation $C(d)$ for a nonzero constant determined solely by the dimension d . It may represent different constants in different places. This avoids the trouble of keeping track of the constants when unnecessary.

2. From radiation profile to solution

Now we assume that $G_-(r, \theta)$ is smooth and compactly supported and give an explicit formula of the operator T_-^{-1} . We consider the odd dimensions first.

2A. Odd dimensions.

Lemma 2.1. *Assume that $d \geq 3$ is odd. Let G_- be a smooth function with $\text{supp } G_- \subset [-R, R] \times \mathbb{S}^{d-1}$. Then $(u_0, u_1) = T_-^{-1}G_-$ satisfies*

$$u_0(x) = \frac{1}{(2\pi)^{(d-1)/2}} \int_{\mathbb{S}^{d-1}} G_-^{(\mu-1)}(x \cdot \omega, \omega) d\omega, \tag{8}$$

$$u_1(x) = \frac{1}{(2\pi)^{(d-1)/2}} \int_{\mathbb{S}^{d-1}} G_-^{(\mu)}(x \cdot \omega, \omega) d\omega. \tag{9}$$

Here the notation $G_-^{(k)}$ represents the partial derivative

$$G_-^{(k)}(s, \theta) = \frac{\partial^k G_-(s, \theta)}{\partial s^k}.$$

Remark 2.2. This formula in 3-dimensional case was previously known. Please refer to [Friedlander 1973], for example.

Proof. Let $(u_0, u_1) = T_-^{-1}G_-$ and $u = S_L(u_0, u_1)$. Given a large time $t > 0$, we choose approximated data $(v_{0,t}, v_{1,t}) \approx (u(\cdot, -t), u_t(\cdot, -t))$ as

$$v_{1,t}(r\theta) = r^{-\mu}G_-(r-t, \theta), \quad r > 0, \theta \in \mathbb{S}^{d-1}, \tag{10}$$

$$v_{0,t}(r\theta) = -\chi\left(\frac{r}{t}\right) \int_r^{+\infty} r'^{-\mu}G_-(r'-t, \theta) dr', \quad r > 0, \theta \in \mathbb{S}^{d-1}. \tag{11}$$

Here $\mu = (d-1)/2$ and $\chi : \mathbb{R} \rightarrow [0, 1]$ is a smooth center cut-off function satisfying

$$\chi(s) = \begin{cases} 1, & s > \frac{1}{2}, \\ 0, & s < \frac{1}{4}. \end{cases}$$

It is clear that the data $(v_{0,t}, v_{1,t})$ are smooth and compactly supported in $\{x : |x| < R+t\}$. A straightforward calculation shows that

$$\begin{aligned} \int_0^\infty \int_{\mathbb{S}^{d-1}} |r^\mu v_{1,t}(r\theta) - G_-(r-t, \theta)|^2 d\theta dr &= 0, \\ \int_0^\infty \int_{\mathbb{S}^{d-1}} |r^\mu \partial_r v_{0,t}(r\theta) - G_-(r-t, \theta)|^2 d\theta dr &\lesssim \frac{1}{t}, \\ \int_{\mathbb{R}^d} (|\nabla v_{0,t}(x)|^2 - |\partial_r v_{0,t}(x)|^2) dx &\lesssim \frac{1}{t}. \end{aligned}$$

Thus by the radiation field we have

$$\lim_{t \rightarrow +\infty} \|(v_{0,t}, v_{1,t}) - (u(\cdot, -t), u_t(\cdot, -t))\|_{\dot{H}^1 \times L^2(\mathbb{R}^d)} = 0.$$

Since the linear propagation operator $S_L(t)$ preserves the $\dot{H}^1 \times L^2$ norm, we have

$$\lim_{t \rightarrow +\infty} \left\| \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} - S_L(t) \begin{pmatrix} v_{0,t} \\ v_{1,t} \end{pmatrix} \right\|_{\dot{H}^1 \times L^2(\mathbb{R}^d)} = 0. \tag{12}$$

Next we use the explicit expression of the linear propagation operator (see, for instance, [Evans 1998]) and write $v = S_L(v_0, v_1)$ in terms of (v_0, v_1) when the initial data are sufficiently smooth:

$$\begin{aligned} v(x, t) &= c_d \cdot \frac{\partial}{\partial t} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\mu-1} \left(t^{d-2} \int_{\mathbb{S}^{d-1}} v_0(x+t\omega) d\omega \right) + c_d \cdot \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\mu-1} \left(t^{d-2} \int_{\mathbb{S}^{d-1}} v_1(x+t\omega) d\omega \right) \\ &= c_d t^\mu \int_{\mathbb{S}^{d-1}} [((w \cdot \nabla)^\mu v_0)(x+t\omega) + ((w \cdot \nabla)^{\mu-1} v_1)(x+t\omega)] d\omega \\ &\quad + \sum_{0 \leq k < \mu} A_{d,k} t^k \int_{\mathbb{S}^{d-1}} ((w \cdot \nabla)^k v_0)(x+t\omega) d\omega + \sum_{0 \leq k < \mu-1} B_{d,k} t^{k+1} \int_{\mathbb{S}^{d-1}} ((w \cdot \nabla)^k v_1)(x+t\omega) d\omega. \end{aligned}$$

Here $c_d = 1/(2(2\pi)^{(d-1)/2})$, $A_{d,k}$, $B_{d,k}$ (and $A'_{d,k}$, $B'_{d,k}$ below) are all constants. We may differentiate and obtain

$$\begin{aligned} v_t(x, t) &= c_d t^\mu \int_{\mathbb{S}^{d-1}} [((w \cdot \nabla)^{\mu+1} v_0)(x+t\omega) + ((w \cdot \nabla)^\mu v_1)(x+t\omega)] d\omega \\ &\quad + \sum_{1 \leq k \leq \mu} A'_{d,k} t^{k-1} \int_{\mathbb{S}^{d-1}} ((w \cdot \nabla)^k v_0)(x+t\omega) d\omega + \sum_{0 \leq k \leq \mu-1} B'_{d,k} t^k \int_{\mathbb{S}^{d-1}} ((w \cdot \nabla)^k v_1)(x+t\omega) d\omega. \end{aligned}$$

Now we plug in $(v_0, v_1) = (v_{0,t}, v_{1,t})$ with large time t . We observe that

$$|(\omega \cdot \nabla)^k v_{j,t}(x + tw)| \lesssim t^{-\mu}, \quad j = 0, 1, k \geq 0, \tag{13}$$

and $(r = |x + t\omega|, \theta = (x + t\omega)/|x + t\omega|, k = \mu - 1, \mu)$

$$\begin{aligned} ((\omega \cdot \nabla)^{k+1} v_{0,t})(x + t\omega) &= (\omega \cdot \theta)^{k+1} r^{-\mu} G_-^{(k)}(r - t, \theta) + O(t^{-\mu-1}), \\ ((\omega \cdot \nabla)^k v_{1,t})(x + t\omega) &= (\omega \cdot \theta)^k r^{-\mu} G_-^{(k)}(r - t, \theta) + O(t^{-\mu-1}). \end{aligned}$$

Thus

$$\begin{pmatrix} w_{0,t} \\ w_{1,t} \end{pmatrix} = S_L(t) \begin{pmatrix} v_{0,t} \\ v_{1,t} \end{pmatrix}$$

satisfies

$$\begin{aligned} w_{0,t} &= c_d \int_{\mathbb{S}^{d-1}} (\omega \cdot \theta)^{\mu-1} (1 + \omega \cdot \theta) G_-^{(\mu-1)}(r - t, \theta) d\omega + O\left(\frac{1}{t}\right), \\ w_{1,t} &= c_d \int_{\mathbb{S}^{d-1}} (\omega \cdot \theta)^\mu (1 + \omega \cdot \theta) G_-^{(\mu)}(r - t, \theta) d\omega + O\left(\frac{1}{t}\right). \end{aligned}$$

Please note that the implicit constants in (13), $O(t^{-\mu-1})$ and $O(1/t)$ above, may depend on x but remain uniformly bounded if x is contained in a compact subset of \mathbb{R}^d . Next we observe the facts

$$\theta(\omega) = \omega + O\left(\frac{1}{t}\right), \quad r(\omega) - t = x \cdot \omega + O\left(\frac{1}{t}\right),$$

and further simplify the formulas

$$\begin{aligned} w_{0,t} &= 2c_d \int_{\mathbb{S}^{d-1}} G_-^{(\mu-1)}(x \cdot \omega, \omega) d\omega + O\left(\frac{1}{t}\right), \\ w_{1,t} &= 2c_d \int_{\mathbb{S}^{d-1}} G_-^{(\mu)}(x \cdot \omega, \omega) d\omega + O\left(\frac{1}{t}\right). \end{aligned}$$

Finally we make $t \rightarrow +\infty$, utilize (12) and obtain

$$\begin{aligned} u_0 &= 2c_d \int_{\mathbb{S}^{d-1}} G_-^{(\mu-1)}(x \cdot \omega, \omega) d\omega, \\ u_1 &= 2c_d \int_{\mathbb{S}^{d-1}} G_-^{(\mu)}(x \cdot \omega, \omega) d\omega. \end{aligned}$$

We plug in the value of c_d and finish the proof. □

Remark 2.3. An explicit formula of the free wave $u = S_L T_-^{-1} G_-$ can be given by

$$u(x, t) = \frac{1}{(2\pi)^{(d-1)/2}} \int_{\mathbb{S}^{d-1}} G_-^{(\mu-1)}(x \cdot \omega + t, \omega) d\omega.$$

This can be verified by a straightforward calculation. One may check

- The function u above is a smooth solution to the homogenous linear wave equation.
- The initial data of u are exactly those given in Lemma 2.1.

We may differentiate and obtain

$$\begin{aligned} u_t(x, t) &= \frac{1}{(2\pi)^{(d-1)/2}} \int_{\mathbb{S}^{d-1}} G_-^{(\mu)}(x \cdot \omega + t, \omega) d\omega, \\ \nabla u(x, t) &= \frac{1}{(2\pi)^{(d-1)/2}} \int_{\mathbb{S}^{d-1}} G_-^{(\mu)}(x \cdot \omega + t, \omega) \omega d\omega. \end{aligned}$$

2B. Even dimensions. The formula of T_-^{-1} in even dimensions is a little more complicated.

Lemma 2.4. *Assume that $d \geq 2$ is even and $G_- \in C_0^\infty(\mathbb{R} \times \mathbb{S}^{d-1})$. Then the operator T_-^{-1} is given explicitly by*

$$\begin{aligned} u_0(x) &= \frac{\sqrt{2}}{(2\pi)^{d/2}} \cdot \int_0^\infty \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2-1)}(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho, \\ u_1(x) &= \frac{\sqrt{2}}{(2\pi)^{d/2}} \cdot \int_0^\infty \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2)}(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho. \end{aligned}$$

Proof. Without loss of generality let us assume $\text{supp } G_- \subset [-R_1, R_1] \times \mathbb{S}^{d-1}$. It is sufficient to show that given any $R_2 > 0$, the formula above holds for almost every $x \in B(0, R_2)$. Let us use the notation $(u_0, u_1) = T_-^{-1}(G_-)$ and $u = \mathcal{S}_L(u_0, u_1)$. We consider the approximated data

$$\begin{aligned} v_{1,t}(r\theta) &= r^{-\mu} G_-(r-t, \theta), \\ v_{0,t}(r\theta) &= -\chi\left(\frac{r}{t}\right) \int_r^{+\infty} r'^{-\mu} G_-(r'-t, \theta) dr', \quad r > 0, \theta \in \mathbb{S}^{d-1}. \end{aligned}$$

and

$$\begin{pmatrix} w_{0,t} \\ w_{1,t} \end{pmatrix} = \mathcal{S}_L(t) \begin{pmatrix} v_{0,t} \\ v_{1,t} \end{pmatrix}.$$

Here χ is the center cut-off function as given in the previous subsection. A basic calculation shows

$$\lim_{t \rightarrow +\infty} \|(v_{0,t}, v_{1,t}) - (u(\cdot, -t), u_t(\cdot, -t))\|_{\dot{H}^1 \times L^2(\mathbb{R}^d)} = 0.$$

Thus

$$\lim_{t \rightarrow +\infty} \|(w_{0,t}, w_{1,t}) - (u_0, u_1)\|_{\dot{H}^1 \times L^2(\mathbb{R}^d)} = 0. \quad (14)$$

Let us first recall the explicit formula of $v = \mathcal{S}_L(v_0, v_1)$ in the even-dimensional case:

$$\begin{aligned} v(x, t) &= c_d \cdot \frac{\partial}{\partial t} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{(d-2)/2} \left(t^{d-1} \int_{\mathbb{B}^d} \frac{v_0(x+ty)}{\sqrt{1-|y|^2}} dy \right) + c_d \cdot \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{(d-2)/2} \left(t^{d-1} \int_{\mathbb{B}^d} \frac{v_1(x+ty)}{\sqrt{1-|y|^2}} dy \right) \\ &= c_d \cdot t^{d/2} \int_{\mathbb{B}^d} \frac{((y \cdot \nabla)^{d/2} v_0)(x+ty) + ((y \cdot \nabla)^{d/2-1} v_1)(x+ty)}{\sqrt{1-|y|^2}} dy \\ &\quad + \sum_{0 \leq k < d/2} A_{d,k} t^k \int_{\mathbb{B}^d} \frac{(y \cdot \nabla)^k v_0(x+ty)}{\sqrt{1-|y|^2}} dy + \sum_{0 \leq k < d/2-1} B_{d,k} t^{k+1} \int_{\mathbb{B}^d} \frac{(y \cdot \nabla)^k v_1(x+ty)}{\sqrt{1-|y|^2}} dy. \end{aligned}$$

Here \mathbb{B}_d is the unit ball in \mathbb{R}^d and $c_d = (2\pi)^{-d/2}$ is a constant. The notations $A_{d,k}$, $B_{d,k}$ (and $A'_{d,k}$, $B'_{d,k}$ below) represent constants. We differentiate and obtain

$$v_t(x, t) = c_d \cdot t^{d/2} \int_{\mathbb{B}^d} \frac{((y \cdot \nabla)^{d/2+1} v_0)(x + ty) + ((y \cdot \nabla)^{d/2} v_1)(x + ty)}{\sqrt{1 - |y|^2}} dy + \sum_{1 \leq k \leq d/2} A'_{d,k} t^{k-1} \int_{\mathbb{B}^d} \frac{(y \cdot \nabla)^k v_0(x + ty)}{\sqrt{1 - |y|^2}} dy + \sum_{0 \leq k \leq d/2-1} B'_{d,k} t^k \int_{\mathbb{B}^d} \frac{(y \cdot \nabla)^k v_1(x + ty)}{\sqrt{1 - |y|^2}} dy.$$

We plug in $(v_0, v_1) = (v_{0,t}, v_{1,t})$ and observe

$$|(y \cdot \nabla)^k v_{0,t}| \leq t^{-(d-1)/2}, \quad |(y \cdot \nabla)^k v_{1,t}| \leq t^{-(d-1)/2}.$$

This gives the approximation

$$w_{0,t}(x) = c_d \cdot t^{d/2} \int_{\mathbb{B}^d} \frac{((y \cdot \nabla)^{d/2} v_{0,t})(r\theta) + ((y \cdot \nabla)^{d/2-1} v_{1,t})(r\theta)}{\sqrt{1 - |y|^2}} dy + O(t^{-1/2}),$$

$$w_{1,t}(x) = c_d \cdot t^{d/2} \int_{\mathbb{B}^d} \frac{((y \cdot \nabla)^{d/2+1} v_{0,t})(r\theta) + ((y \cdot \nabla)^{d/2} v_{1,t})(r\theta)}{\sqrt{1 - |y|^2}} dy + O(t^{-1/2}).$$

Here $r = |x + ty|$ and $\theta = (x + ty)/|x + ty|$. Furthermore, we observe ($k = d/2, d/2 - 1$)

$$((y \cdot \nabla)^{k+1} v_{0,t})(r\theta) = (y \cdot \theta)^{k+1} r^{-(d-1)/2} G_-^{(k)}(r - t, \theta) + O(t^{-(d+1)/2}),$$

$$((y \cdot \nabla)^k v_{1,t})(r\theta) = (y \cdot \theta)^k r^{-(d-1)/2} G_-^{(k)}(r - t, \theta) + O(t^{-(d+1)/2}),$$

and write

$$w_{0,t}(x) = c_d \cdot t^{d/2} \int_{\mathbb{B}^d} \frac{(y \cdot \theta)^{d/2-1} (y \cdot \theta + 1) r^{-(d-1)/2} G_-^{(d/2-1)}(r - t, \theta)}{\sqrt{1 - |y|^2}} dy + O(t^{-1/2}),$$

$$w_{1,t}(x) = c_d \cdot t^{d/2} \int_{\mathbb{B}^d} \frac{(y \cdot \theta)^{d/2} (y \cdot \theta + 1) r^{-(d-1)/2} G_-^{(d/2)}(r - t, \theta)}{\sqrt{1 - |y|^2}} dy + O(t^{-1/2}).$$

Next we observe that if $|y| < 1 - (R_1 + R_2)/t$, then we have $r \leq t|y| + |x| < t - R_1$; thus $G_-^{(k)}(r - t, \theta) = 0$. As a result, we may restrict the domain of the integral to

$$\mathbb{B}_t = \left\{ y \in \mathbb{B}^d : |y| \geq 1 - \frac{R_1 + R_2}{t} \right\}.$$

Because in the region we have

$$\theta = \frac{y}{|y|} + O\left(\frac{1}{t}\right), \quad y \cdot \theta = 1 + O\left(\frac{1}{t}\right), \quad r = t + O(1),$$

we can simplify the formulas

$$w_{0,t}(x) = 2c_d \cdot t^{1/2} \int_{\mathbb{B}_t} \frac{G_-^{(d/2-1)}(r - t, y/|y|)}{\sqrt{1 - |y|^2}} dy + O(t^{-1/2}),$$

$$w_{1,t}(x) = 2c_d \cdot t^{1/2} \int_{\mathbb{B}_t} \frac{G_-^{(d/2)}(r - t, y/|y|)}{\sqrt{1 - |y|^2}} dy + O(t^{-1/2}).$$

Next we utilize the change of variables

$$y = \left(1 - \frac{\rho}{t}\right)\omega, \quad (\rho, \omega) \in (0, R_1 + R_2) \times \mathbb{S}^{d-1},$$

and the approximations

$$r - t = x \cdot \omega - \rho + O\left(\frac{1}{t}\right), \quad \sqrt{1 - |y|^2} = \left(1 + O\left(\frac{1}{t}\right)\right)\sqrt{\frac{2\rho}{t}}, \quad dy = \left(1 + O\left(\frac{1}{t}\right)\right)t^{-1} d\rho d\omega$$

to obtain

$$w_{0,t}(x) = \sqrt{2}c_d \cdot \int_0^{R_1+R_2} \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2-1)}(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho + O(t^{-1/2}),$$

$$w_{1,t}(x) = \sqrt{2}c_d \cdot \int_0^{R_1+R_2} \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2)}(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho + O(t^{-1/2}).$$

Finally we recall (14), let $t \rightarrow +\infty$ and conclude

$$u_0(x) = \sqrt{2}c_d \cdot \int_0^{R_1+R_2} \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2-1)}(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho,$$

$$u_1(x) = \sqrt{2}c_d \cdot \int_0^{R_1+R_2} \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2)}(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho. \quad \square$$

Remark 2.5. If $d \geq 4$, the convergence (14) implies that $(w_{0,t}, w_{1,t})$ converges to (u_0, u_1) in $L^{2d/(d-2)} \times L^2$ by Sobolev embedding. We may combine this convergence with the local uniform convergence given above to verify the identities above. This argument breaks down in dimension 2. We given another argument below in dimension 2. Given any test function $\varphi \in C_0^\infty(\mathbb{R}^2)$, integration by parts gives an identity

$$\int w_{0,t}(x) \nabla \varphi(x) dx = - \int \nabla w_{0,t}(x) \varphi(x) dx.$$

We recall the local uniform convergence of $w_{0,t}$ given above and the L^2 convergence of $\nabla w_{0,t} \rightarrow \nabla u_0$ and then obtain

$$\int \left(\sqrt{2}c_2 \cdot \int_0^\infty \int_{\mathbb{S}^1} \frac{G_-(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho \right) \nabla \varphi(x) dx = - \int \nabla u_0(x) \varphi(x) dx.$$

This finishes the proof. Finally we would like to mention that we have

$$\lim_{|x| \rightarrow +\infty} \sqrt{2}c_2 \cdot \int_0^\infty \int_{\mathbb{S}^1} \frac{G_-(x \cdot \omega - \rho, \omega)}{\sqrt{\rho}} d\omega d\rho = 0.$$

Corollary 2.6. If $G_- \in C_0^\infty(\mathbb{R} \times \mathbb{S}^{d-1})$, then $u = \mathbf{S}_L \mathbf{T}^{-1}(G_-)$ is given by

$$u(x, t) = \frac{\sqrt{2}}{(2\pi)^{d/2}} \int_0^\infty \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2-1)}(x \cdot \omega - \rho + t, \omega)}{\sqrt{\rho}} d\omega d\rho.$$

Thus

$$u_t(x, t) = \frac{\sqrt{2}}{(2\pi)^{d/2}} \int_0^\infty \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2)}(x \cdot \omega - \rho + t, \omega)}{\sqrt{\rho}} d\omega d\rho.$$

Proof. A basic calculation shows that $u(x, t)$ solves the free wave equation with initial data given in Lemma 2.4. □

2C. Universal formula. Now let us give a universal formula of T_-^{-1} for all dimensions. We first define two convolution operators ($1/\sqrt{\pi x}$ is understood as zero if $x < 0$)

$$Qf = \frac{1}{\sqrt{\pi x}} * f, \quad Q'f = \frac{1}{\sqrt{-\pi x}} * f.$$

Their Fourier symbols are

$$\frac{1 - i(\xi/|\xi|)}{2\sqrt{\pi}|\xi|} \quad \text{and} \quad \frac{1 + i(\xi/|\xi|)}{2\sqrt{\pi}|\xi|},$$

respectively. Let us also use the notation $\mathcal{D} = d/dx$ and recall that its Fourier symbol is $2\pi i\xi$. A simple calculation of symbols shows

$$Q^2\mathcal{D} = 1, \quad Q'^2\mathcal{D} = -1, \quad QQ'\mathcal{D} = \mathcal{H}. \tag{15}$$

As a result, we may understand Q as $\mathcal{D}^{-1/2}$ and rewrite $u = S_L T_-^{-1} G_-$ in the form

$$\begin{aligned} u(x, t) &= \frac{1}{(2\pi)^{(d-1)/2}} \int_{\mathbb{S}^{d-1}} (QG_-^{(d/2-1)})(x \cdot \omega + t, \omega) d\omega \\ &= \frac{1}{(2\pi)^\mu} \int_{\mathbb{S}^{d-1}} \mathcal{D}^{\mu-1} G_-(x \cdot \omega + t, \omega) d\omega. \end{aligned} \tag{16}$$

Here $\mu = (d - 1)/2$. This formula holds for both odd and even dimensions.

3. Between radiation profiles

In this section we give an explicit expression of the operator $T_+ \circ T_-^{-1}$ in the even-dimensional case, without the radial assumption.

Theorem 3.1. *Assume that $d \geq 2$ is an even integer. The operator $T_+ \circ T_-^{-1}$ can be explicitly given by the formula*

$$G_+(s, \theta) = (T_+ T_-^{-1} G_-)(s, \theta) = (-1)^{d/2} (\mathcal{H}G_-)(-s, -\theta).$$

Here \mathcal{H} is the Hilbert transform in the first variable, i.e.,

$$(\mathcal{H}G_-)(-s, -\theta) = \text{p.v.} \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{G_-(\tau, -\theta)}{-\tau - s} d\tau.$$

Proof. Since $T_+ \circ T_-^{-1}$ is a bijective isometry from $L^2(\mathbb{R} \times \mathbb{S}^{d-1})$ to itself. We only need to prove this formula for smooth and compactly supported data G_- . Without loss of generality let us assume $\text{supp } G_- \subset [-R_1, R_1] \times \mathbb{S}^{d-1}$. Let us also fix a positive constant $R_2 > 0$. If $(s, \theta) \in (-R_2, R_2) \times \mathbb{S}^{d-1}$, then we may apply Corollary 2.6 and obtain

$$(t + s)^{(d-1)/2} \partial_t u((t + s)\theta, t) = \sqrt{2} c_d (t + s)^{(d-1)/2} \int_0^\infty \int_{\mathbb{S}^{d-1}} \frac{G_-^{(d/2)}((t + s)\theta \cdot \omega - \rho + t, \omega)}{\sqrt{\rho}} d\omega d\rho.$$

Let $M \gg R_1 + R_2 + 1$ be a large constant. We may split the integral above into two parts:

$$J_1 = \sqrt{2}c_d(t+s)^{(d-1)/2} \int_0^\infty \int_{\theta \cdot \omega < -1+M/t} \frac{G_-^{(d/2)}((t+s)\theta \cdot \omega - \rho + t, \omega)}{\sqrt{\rho}} d\omega d\rho,$$

$$J_2 = \sqrt{2}c_d(t+s)^{(d-1)/2} \int_0^\infty \int_{\theta \cdot \omega \geq -1+M/t} \frac{G_-^{(d/2)}((t+s)\theta \cdot \omega - \rho + t, \omega)}{\sqrt{\rho}} d\omega d\rho.$$

We may find an upper bound of J_2 . In this region we have

$$(t+s)\theta \cdot \omega + t \geq M - R_2 \implies G_-((t+s)\theta \cdot \omega - \rho + t) = 0 \quad \text{if } \rho < \frac{M}{2}.$$

Thus we may integrate by parts and obtain

$$J_2 = C(d)(t+s)^{(d-1)/2} \int_0^\infty \int_{\theta \cdot \omega \geq -1+M/t} \frac{G_-((t+s)\theta \cdot \omega - \rho + t, \omega)}{\rho^{(d+1)/2}} d\omega d\rho.$$

Therefore when t is sufficiently large

$$\begin{aligned} |J_2| &\lesssim t^{(d-1)/2} \int_{\theta \cdot \omega \geq -1+M/t} \int_{(t+s)\theta \cdot \omega + t - R_1}^{(t+s)\theta \cdot \omega + t + R_1} \frac{|G_-((t+s)\theta \cdot \omega - \rho + t, \omega)|}{\rho^{(d+1)/2}} d\rho d\omega \\ &\lesssim t^{(d-1)/2} \int_{\theta \cdot \omega \geq -1+M/t} \int_{(t+s)\theta \cdot \omega + t - R_1}^{(t+s)\theta \cdot \omega + t + R_1} \frac{|G_-((t+s)\theta \cdot \omega - \rho + t, \omega)|}{|(t+s)\theta \cdot \omega + t|^{(d+1)/2}} d\rho d\omega \\ &\lesssim t^{(d-1)/2} \int_{\theta \cdot \omega \geq -1+M/t} \frac{1}{|t\theta \cdot \omega + t|^{(d+1)/2}} d\omega \lesssim \frac{1}{M}. \end{aligned}$$

In the integral region of J_1 , we have the approximation $\omega = -\theta + O(t^{-1/2})$. Thus we have

$$J_1 = \sqrt{2}c_d t^{(d-1)/2} \int_0^\infty \int_{\theta \cdot \omega < -1+M/t} \frac{G_-^{(d/2)}((t+s)\theta \cdot \omega - \rho + t, -\theta)}{\sqrt{\rho}} d\omega d\rho + O(t^{-1/2}).$$

Next we utilize the change of variables (please refer to Figure 1 for a geometrical meaning)

$$\omega = \left(-1 + \frac{\rho'}{t}\right)\theta + \sqrt{\left(\frac{\rho'}{t}\right)\left(2 - \frac{\rho'}{t}\right)}\varphi, \quad \rho' \in [0, M], \quad \varphi \in \mathbb{S}^{d-2} = \{\varphi \in \mathbb{S}^{d-1} : \varphi \perp \theta\},$$

$$d\omega = \left[1 + O\left(\frac{1}{t}\right)\right] \left(\frac{2\rho'}{t}\right)^{d/2-1} d\mathbb{S}^{d-2}(\varphi) \cdot \frac{d\rho'}{\sqrt{2\rho't}} = \left[1 + O\left(\frac{1}{t}\right)\right] (2\rho')^{(d-3)/2} t^{-(d-1)/2} d\mathbb{S}^{d-2}(\varphi) d\rho'$$

and obtain

$$J_1 = \frac{1}{2\pi^{d/2}} \int_0^\infty \int_0^M \int_{\mathbb{S}^{d-2}} G_-^{(d/2)}(\rho' - \rho - s, -\theta) \rho'^{(d-3)/2} \rho^{-1/2} d\varphi d\rho' d\rho + O(t^{-1/2}).$$

We observe that the integrand is independent of φ and integrate by parts

$$J_1 = \frac{(-1)^{d/2-1}}{\pi} \int_0^\infty \int_0^M \frac{G'_-(\rho' - \rho - s, -\theta)}{\sqrt{\rho\rho'}} d\rho' d\rho + O(t^{-1/2}).$$




Figure 1. Change of variables, where $r = \sqrt{(\rho'/t)(2 - \rho'/t)}$.

We next change the variables $\tau = \rho' - \rho$, $\eta = \rho' + \rho$, and write

$$\begin{aligned} J_1 &= \frac{(-1)^{d/2-1}}{\pi} \int_{-\infty}^M \int_{|\tau|}^{2M-\tau} \frac{G'_-(\tau - s, -\theta)}{\sqrt{\eta^2 - \tau^2}} d\eta d\tau + O(t^{-1/2}) \\ &= \frac{(-1)^{d/2-1}}{\pi} \int_{-\infty}^M G'_-(\tau - s, -\theta) [\ln(2M - \tau + \sqrt{4M^2 - 4M\tau}) - \ln |\tau|] d\tau + O(t^{-1/2}) \\ &= \frac{(-1)^{d/2-1}}{\pi} \int_{-R_1-R_2}^{R_1+R_2} G'_-(\tau - s, -\theta) [\ln(2M - \tau + \sqrt{4M^2 - 4M\tau}) - \ln |\tau|] d\tau + O(t^{-1/2}). \end{aligned}$$

The integrals above can be split into two parts:

$$\begin{aligned} I_1 &= \int_{-R_1-R_2}^{R_1+R_2} G'_-(\tau - s, -\theta) [\ln(2M - \tau + \sqrt{4M^2 - 4M\tau})] d\tau \\ &= \int_{-R_1-R_2}^{R_1+R_2} G'_-(\tau - s, -\theta) [\ln(2M - \tau + \sqrt{4M^2 - 4M\tau}) - \ln(4M)] d\tau \\ &= \int_{-R_1-R_2}^{R_1+R_2} G'_-(\tau - s, -\theta) O\left(\frac{1}{M}\right) d\tau = O\left(\frac{1}{M}\right) \end{aligned}$$

and

$$\begin{aligned} I_2 &= - \lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon < |\tau| < R_1+R_2} G'_-(\tau - s, -\theta) \ln |\tau| d\tau \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon < |\tau| < R_1+R_2} \frac{G_-(\tau - s, -\theta)}{\tau} d\tau = -\pi(\mathcal{H}G_-)(-s, -\theta). \end{aligned}$$

In summary we have

$$J_1 = (-1)^{d/2}(\mathcal{H}G_-)(-s, -\theta) + O\left(\frac{1}{M}\right) + O(t^{-1/2}).$$

Now we may combine J_1 and J_2

$$(t + s)^{(d-1)/2} \partial_t u((t + s)\theta, t) = (-1)^{d/2}(\mathcal{H}G_-)(-s, -\theta) + O\left(\frac{1}{M}\right) + O(t^{-1/2}).$$

Because the implicit constants in O 's do not depend on $s \in [-R_2, R_2]$ or $\theta \in \mathbb{S}^{d-1}$, we may let $t \rightarrow +\infty$ then $M \rightarrow +\infty$ to conclude

$$\lim_{t \rightarrow +\infty} \int_{-R_2}^{R_2} \int_{\mathbb{S}^{d-1}} |(t+s)^{(d-1)/2} \partial_t u((t+s)\theta, t) - (-1)^{d/2} (\mathcal{H}G_-)(-s, -\theta)|^2 d\theta ds = 0. \quad \square$$

4. Radial weakly nonradiative solutions

In this section we prove Proposition 1.9. First of all, we briefly show that any initial data in $P_{\text{rad}}(R)$ leads to an R -weakly nonradiative solution. By linearity we only need to consider the case $(u_0, u_1) = (r^{2k_1-d}, 0)$ or $(u_0, u_1) = (0, r^{2k_2-d})$. If $(u_0, u_1) = (r^{2k_1-d}, 0)$, then a basic calculation shows that if we choose $C_1, C_2, \dots, C_{k_1-1}$ inductively, the solution

$$u_{k_1}(x, t) = \frac{1}{|x|^{d-2k_1}} + \frac{C_1 t^2}{|x|^{d-2k_1+2}} + \dots + \frac{C_{k_1-1} t^{2k_1-2}}{|x|^{d-2}}$$

solves the linear wave equation with initial data $(|x|^{2k_1-d}, 0)$ in the region $\mathbb{R}^d \setminus \{0\}$. By finite speed of propagation, we have

$$S_L(u_0, u_1)(x, t) = u_{k_1}(x, t), \quad |x| > R + |t|.$$

A simple calculation shows that this is indeed a nonradiative solution. The case $(u_0, u_1) = (0, r^{2k_2-d})$ can be dealt with in the same manner by considering the solution

$$u_{k_2}(x, t) = \frac{t}{|x|^{d-2k_1}} + \frac{C_1 t^3}{|x|^{d-2k_1+2}} + \dots + \frac{C_{k_2-1} t^{2k_1-1}}{|x|^{d-2}}.$$

Thus it is sufficient to show initial data of any nonradiative solution are contained in the space $P_{\text{rad}}(R)$. We first consider the odd dimensions.

4A. Odd dimensions. Assume that $u = S_L(u_0, u_1)$ is a radial R -weakly nonradiative solution. Let $G_- = T_-(u_0, u_1)$. By radial assumption G_- is independent of the angle $\omega \in \mathbb{S}^{d-1}$. Let us first consider smooth functions G_- . We may calculate $(r > R, e_1 = (1, 0, \dots, 0) \in \mathbb{R}^d)$

$$u_0(re_1) = (2\pi)^{-\mu} \int_{\mathbb{S}^{d-1}} G_-^{(\mu-1)}(r\omega_1) d\omega = \frac{\sigma_{d-2}}{(2\pi)^\mu} \int_{-1}^1 G_-^{(\mu-1)}(r\omega_1) (1-\omega_1^2)^{\mu-1} d\omega_1.$$

Here ω_1 is the first variable of $\mathbb{R}^d \supset \mathbb{S}^{d-1}$ and σ_{d-2} is the area of the sphere \mathbb{S}^{d-2} . We may integrate by parts and rescale:

$$\begin{aligned} u_0(re_1) &= \frac{(-1)^{\mu-1} \sigma_{d-2}}{(2\pi)^\mu r^{\mu-1}} \int_{-1}^1 G_-(r\omega_1) [\partial_{\omega_1}^{\mu-1} (1-\omega_1^2)^{\mu-1}] d\omega_1 \\ &= \sum_{k=0}^{\lfloor (\mu-1)/2 \rfloor} \frac{A_{d,k}}{r^{\mu-1}} \int_{-1}^1 G_-(r\omega_1) \omega_1^{\mu-1-2k} d\omega_1 = \sum_{k=0}^{\lfloor (d-3)/4 \rfloor} \frac{A_{d,k}}{r^{d-2-2k}} \int_{-R}^R G_-(s) s^{(d-3)/2-2k} ds \\ &= \sum_{k=1}^{\lfloor (d+1)/4 \rfloor} \frac{A_{d,k}}{r^{d-2k}} \int_{-R}^R G_-(s) s^{(d+1)/2-2k} ds. \end{aligned}$$

Here the $A_{d,k}$ are nonzero constants. Similarly we have

$$\begin{aligned} u_1(re_1) &= (2\pi)^{-\mu} \int_{\mathbb{S}^{d-1}} G_-^{(\mu)}(r\omega_1) d\omega = \frac{\sigma_{d-2}}{(2\pi)^\mu} \int_{-1}^1 G_-^{(\mu)}(r\omega_1)(1-\omega_1^2)^{\mu-1} d\omega_1 \\ &= \frac{(-1)^\mu \sigma_{d-2}}{(2\pi)^\mu r^\mu} \int_{-1}^1 G_-(r\omega_1)[\partial_{\omega_1}^\mu (1-\omega_1^2)^{\mu-1}] d\omega_1 = \sum_{k=0}^{\lfloor(\mu-2)/2\rfloor} \frac{B_{d,k}}{r^\mu} \int_{-1}^1 G_-(r\omega_1)\omega_1^{\mu-2-2k} d\omega_1 \\ &= \sum_{k=1}^{\lfloor(d-1)/4\rfloor} \frac{B_{d,k}}{r^{d-2k}} \int_{-R}^R G_-(s)s^{(d-1)/2-2k} ds. \end{aligned}$$

Here the $B_{d,k}$ are nonzero constants. Since smooth functions are dense in $L^2([-R, R])$, we have:

Proposition 4.1. *There exist constants $\{A_{d,k}\}_{1 \leq k \leq \lfloor(d+1)/4\rfloor}$, $\{B_{d,k}\}_{1 \leq k \leq \lfloor(d-1)/4\rfloor}$, so that for any $G_- \in L^2(\mathbb{R})$ supported in $[-R, R]$, the initial data $(u_0, u_1) = T_-^{-1}G_-$ satisfy $(r > R)$*

$$\begin{aligned} u_0(r) &= \sum_{k=1}^{\lfloor(d+1)/4\rfloor} \left(A_{d,k} \int_{-R}^R G_-(s)s^{(d+1)/2-2k} ds \right) r^{-d+2k}, \\ u_1(r) &= \sum_{k=1}^{\lfloor(d-1)/4\rfloor} \left(B_{d,k} \int_{-R}^R G_-(s)s^{(d-1)/2-2k} ds \right) r^{-d+2k}. \end{aligned}$$

This clearly shows that if $u = S_L(u_0, u_1)$ is a radial R -weakly nonradiative solution, then $(u_0, u_1) \in P_{\text{rad}}(R)$.

4B. Even dimensions. The even dimensions involve the Hilbert transform, and thus are much more difficult to handle. The general idea is the same. If the initial data (u_0, u_1) are radial, then $G_\pm(s) = T_\pm(u_0, u_1)$ is independent to the angle. We also have $G_+(s) = (-1)^{d/2} \mathcal{H}G_-(-s)$. Thus $S_L(u_0, u_1)$ is R -weakly nonradiative if and only if G_- is contained in the space

$$P_{\text{rad}} = \{G_- \in L^2(\mathbb{R}) : G_-(s) = 0, s > R, (\mathcal{H}G_-)(s) = 0, s < -R\}.$$

Now recall the operators \mathcal{Q} , \mathcal{Q}' and \mathcal{D} defined in Section 2C. We claim:

Lemma 4.2. $\mathcal{Q}'P_{\text{rad}} = \dot{H}_0^{1/2}(-R, R)$. Here $\dot{H}_0^{1/2}(-R, R)$ is the completion of $C_0^\infty(-R, R)$ equipped with the $\dot{H}^{1/2}(\mathbb{R})$ norm.

Proof. In order to avoid technical difficulties, we use an approximation technique. Given any $G_- \in P_{\text{rad}}$, we may utilize a local smoothing kernel to generate a sequence G_k , so that:

- (a) $G_k \in P_{\text{rad}}(R + 1/k)$.
- (b) $G_k \in H^n(\mathbb{R})$ for all $n \geq 0$ and thus $G_k \in C^\infty(\mathbb{R})$.
- (c) G_k converges to G_- in $L^2(\mathbb{R})$.

Let us consider the properties of the function $g_k = \mathcal{Q}'G_k \in C^\infty(\mathbb{R})$. According to part (a), $G_k(s) = 0$ if $s > R + 1/k$. We may use the convolution expression of \mathcal{Q}' to obtain that g_k vanishes in the interval $(R + 1/k, +\infty)$. Similarly $g_k = \mathcal{Q}\mathcal{H}G_k$ vanishes in the interval $(-\infty, -R - 1/k)$. We recall that

$\mathcal{Q}' : L^2(\mathbb{R}) \rightarrow \dot{H}^{1/2}(\mathbb{R})$ is an isometry up to a constant. Thus $g_k \rightarrow g = \mathcal{Q}'G_-$ in $\dot{H}^{1/2}(\mathbb{R})$. This verifies $g \in \dot{H}_0^{1/2}(-R, R)$. We also need to show that given any $g \in \dot{H}_0^{1/2}(-R, R)$, then $\mathcal{Q}'^{-1}g \in \mathcal{P}_{\text{rad}}$. It is sufficient to consider $g \in C_0^\infty(-R, R)$ by smooth approximation. A simple calculation of Fourier symbols shows that $\mathcal{Q}'^{-1} = -\mathcal{Q}'\mathcal{D}$ and $\mathcal{H}\mathcal{Q}'^{-1} = \mathcal{Q}\mathcal{D}$. A combination of these identities with the convolution expressions of \mathcal{Q} and \mathcal{Q}' immediately verifies $\mathcal{Q}'^{-1}g \in \mathcal{P}_{\text{rad}}$. \square

We also need to use the following explicit formula of T_- for radial data.

Lemma 4.3. *Assume $G \in C^\infty(\mathbb{R})$ so that $|G(s)| \lesssim |s|^{-3/2}$ for $|s| \gg 1$. Then the corresponding radial free wave $u = S_L T_-^{-1}G$ satisfies*

$$u(r, t) = C(d) \cdot r^{1-d/2} \int_{-1}^1 \mathcal{Q}G(r\omega_1 + t) P_d(w_1) (1 - w_1^2)^{-1/2} dw_1. \quad (17)$$

Here P_d is an even or odd polynomial of degree $d/2 - 1$ defined by

$$\left(\frac{\partial}{\partial w_1} \right)^{d/2-1} (1 - w_1^2)^{(d-3)/2} = P_d(w_1) (1 - w_1^2)^{-1/2}.$$

Proof. If $G \in C_0^\infty(\mathbb{R})$, we use polar coordinates and integrate by parts:

$$\begin{aligned} u(r, t) &= C(d) \int_0^\infty \int_{\mathbb{S}^{d-1}} \frac{G^{(d/2-1)}(r\omega_1 - \rho + t)}{\sqrt{\rho}} d\omega d\rho \\ &= C(d) \int_0^\infty \int_{-1}^1 \frac{G^{(d/2-1)}(r\omega_1 - \rho + t)}{\sqrt{\rho}} (1 - w_1^2)^{(d-3)/2} dw_1 d\rho \\ &= C(d) \cdot r^{1-d/2} \int_0^\infty \int_{-1}^1 \frac{G(r\omega_1 - \rho + t)}{\sqrt{\rho}} P_d(w_1) (1 - w_1^2)^{-1/2} dw_1 d\rho \\ &= C(d) \cdot r^{1-d/2} \int_{-1}^1 \mathcal{Q}G(r\omega_1 + t) P_d(w_1) (1 - w_1^2)^{-1/2} dw_1. \end{aligned}$$

This verifies the formula if $G \in C_0^\infty(\mathbb{R})$. In order to deal with profile G without compact support, we use standard smooth cut-off techniques. More precisely, we may choose $G_k \in C_0^\infty(\mathbb{R})$ so that $G_k \rightarrow G$ in $L^2(\mathbb{R})$ and

$$|G_k(s) - G(s)| = 0, \quad s < k, \quad |G_k(s) - G(s)| \lesssim |s|^{-3/2}, \quad s \geq k.$$

Thus we have $\|\mathcal{Q}G - \mathcal{Q}G_k\|_{L^\infty} \lesssim 1/k$. This means we have the uniform convergence for all (r, t) in any compact subset of $\mathbb{R}^+ \times \mathbb{R}$:

$$\begin{aligned} u_k(r, t) &= \frac{C(d)}{r^{d/2-1}} \int_{-1}^1 \mathcal{Q}G_k(r\omega_1 + t) P_d(w_1) (1 - w_1^2)^{-1/2} dw_1 \\ &\Rightarrow \frac{C(d)}{r^{d/2-1}} \int_{-1}^1 \mathcal{Q}G(r\omega_1 + t) P_d(w_1) (1 - w_1^2)^{-1/2} dw_1. \end{aligned}$$

Combining this with the convergence $u_k \rightarrow u$ in \dot{H}^1 we finish the proof. \square

Remark 4.4. If $d \geq 4$ and $G \in L^2(\mathbb{R})$, then formula (17) still holds. This follows standard smooth approximation and/or cut-off techniques. Let $G_k \in C_0^\infty(\mathbb{R})$ so that $G_k \rightarrow G$ in $L^2(\mathbb{R})$. Thus $\mathcal{Q}G_k \rightarrow \mathcal{Q}G$ in $\dot{H}^{1/2}(\mathbb{R})$. Finally we observe the fact $P_d(w_1)(1 - w_1^2)^{-1/2} \in \dot{H}^{-1/2}(\mathbb{R})$, obtain a locally uniform convergence $u_k(r, t) \rightarrow u(r, t)$ and conclude the proof.

Now we are ready to give an expression of $u = S_L T_-^{-1} G_-$ when $G_- \in \mathcal{P}_{\text{rad}}(R)$.

Lemma 4.5. Assume $G_- \in \mathcal{P}_{\text{rad}}(R)$. Then the following identity holds:

$$u(r, t) = \frac{C(d)}{r^{d/2}} \int_{-R}^R \mathcal{Q}'G_-(s)W_d\left(\frac{s-t}{r}\right) ds.$$

Here $W_d(\sigma)$ is the Hilbert transform (the function below is understood as zero if $|w_1| > 1$)

$$W_d(\sigma) \doteq \mathcal{H}\left(\frac{P_d(w_1)}{\sqrt{1-w_1^2}}\right) = \mathcal{H}\left[\left(\frac{d}{dw_1}\right)^{d/2-1} (1-w_1^2)^{(d-3)/2}\right].$$

Proof. By Lemma 4.2, we have $\mathcal{Q}'G_- \in \dot{H}_0^{1/2}(-R, R)$. We claim that it is sufficient to consider the case $\mathcal{Q}'G_- \in C_0^\infty(-R, R)$. In fact, we may choose $G_k \in \mathcal{P}_{\text{rad}}(R)$ so that $\mathcal{Q}'G_k \in C_0^\infty(-R, R)$ so that

$$\mathcal{Q}'G_k \rightarrow \mathcal{Q}'G_- \quad \text{in } \dot{H}^{1/2}(-R, R) \quad \Rightarrow \quad G_k \rightarrow G_- \quad \text{in } L^2(\mathbb{R}).$$

Now we observe a few important facts: we have the embedding $\dot{H}_0^{1/2}(-R, R) \hookrightarrow L^p(-R, R)$ for all $1 \leq p < +\infty$ and

$$\frac{P_d(w_1)}{\sqrt{1-w_1^2}} \in L^p(\mathbb{R}) \quad \Rightarrow \quad W_d(\sigma) \in L^p(\mathbb{R}), \quad p \in (1, 2).$$

As a result, if the identity

$$u_k(r, t) = \frac{C(d)}{r^{d/2}} \int_{-R}^R \mathcal{Q}'G_k(s)W_d\left(\frac{s-t}{r}\right) ds, \quad k \geq 1,$$

holds, then we may make $k \rightarrow +\infty$ in the identity above and verify that a similar identity holds for u and G_- . In fact the left-hand side converges in the space $\dot{H}^1(\mathbb{R}^d)$ for any given time t , while the right-hand side converges uniformly for (r, t) in any compact subset of $\mathbb{R}^+ \times \mathbb{R}$. Now we assume $g = \mathcal{Q}'G_- \in C_0^\infty(-R, R)$. Then $G_- = \mathcal{Q}'^{-1}g = -\mathcal{Q}'\mathcal{D}g$ satisfies the assumption of Lemma 4.3. As a result we have

$$\begin{aligned} u(r, t) &= C(d) \cdot r^{1-d/2} \int_{-1}^1 \mathcal{Q}\mathcal{Q}'\mathcal{D}g(r\omega_1 + t)P_d(w_1)(1 - w_1^2)^{-1/2} d\omega_1 \\ &= C(d) \cdot r^{1-d/2} \int_{-1}^1 \mathcal{H}g(r\omega_1 + t)P_d(w_1)(1 - w_1^2)^{-1/2} d\omega_1 \\ &= \frac{C(d)}{r^{d/2-1}} \int_{-\infty}^\infty g(r\sigma + t)W_d(\sigma) d\sigma. \end{aligned}$$

Here we use the facts $\mathcal{Q}\mathcal{Q}'\mathcal{D} = \mathcal{H}$ and

$$\int \mathcal{H}f \cdot \overline{\mathcal{H}g} dx = \int f \cdot \bar{g} dx, \quad \mathcal{H}(\mathcal{H}g(r\omega_1 + t))(\sigma) = (\mathcal{H}^2g)(r\sigma + t) = -g(r\sigma + t).$$

Finally we apply change of variables $s = r\sigma + t$, recall the support of g and finish the proof. □

Now let us consider the Hilbert transform W_d . The key observation is the following technical lemma. This result has been known for many years; see [Solmon 1987], for instance. But we still give a brief proof in the Appendix for the purpose of completeness.

Lemma 4.6. *Assume that $P(x)$ is a polynomial of degree κ . Let W be the Hilbert transform*

$$W = \mathcal{H}\left(\frac{P(x)}{\sqrt{1-x^2}}\right).$$

Then $W(\sigma)$ is equal to a polynomial of degree $\kappa - 1$ if $\sigma \in (-1, 1)$. In particular, $W_2(\sigma) = 0$ for $\sigma \in (-1, 1)$; if $d \geq 4$, then the function $W_d(\sigma)$ is equal to an even or odd polynomial of degree $d/2 - 2$ in the interval $(-1, 1)$.

Proof of Proposition 1.11. According to Lemma 4.5, we have already obtained

$$u(r, t) = \frac{C(d)}{r^{d/2}} \int_{-R}^R \mathcal{Q}'G_-(s)W_d\left(\frac{s-t}{r}\right) ds.$$

Here $\mathcal{Q}'G_- \in \dot{H}_0^{1/2}(-R, R) \hookrightarrow L^p(-R, R)$ for all $1 < p < +\infty$. If we also have $r > |t| + R$, then

$$\left|\frac{s-t}{r}\right| < 1 \quad \text{for all } s \in (-R, R).$$

If $d = 2$, Lemma 4.6 immediately gives $u(r, t) \equiv 0$ if $r > |R| + t$ since we always have $W_2((s-t)/r) = 0$. In the higher-dimensional case $d \geq 4$, Lemma 4.6 guarantees that

$$W_d(s) = \sum_{l=1}^{\lfloor d/4 \rfloor} A_l s^{d/2-2l}, \quad -1 < s < 1,$$

is a polynomial. We plug this in the expression of u and obtain

$$u(r, t) = C(d) \sum_{l=1}^{\lfloor d/4 \rfloor} \frac{A_l}{r^{d-2l}} \int_{-R}^R \mathcal{Q}'G_-(s)(s-t)^{d/2-2l} ds, \quad r > R + |t|. \tag{18}$$

This immediately gives $(u_0, u_1) \in P_{\text{rad}}(R)$.

5. Exterior energy estimates of even dimensions

In this section we prove Proposition 1.11. It suffices to consider the case $d = 4k$. The proof of $d = 4k + 2$ is almost the same. Again we switch to the space of radiation profiles $G_- \in L^2(\mathbb{R} \times \mathbb{S}^{d-1})$. We start with:

Lemma 5.1. *The image of radial data in the form of $(u_0, 0)$ can be characterized by*

$$\begin{aligned} \{\mathcal{T}_-(u_0, 0) : u_0 \in \dot{H}_{\text{rad}}^1(\mathbb{R}^d)\} &= \{G_- \in L^2(\mathbb{R}) : \mathcal{H}G_-(-s) = -G_-(s)\} \\ &= \left\{ \frac{G(s) - \mathcal{H}G(-s)}{2} : G \in L^2(\mathbb{R}) \right\}. \end{aligned}$$

Proof. First of all, if $u_0 \in \dot{H}_{\text{rad}}^1(\mathbb{R}^d)$, then the free wave $u = \mathcal{S}_L(u_0, u_1)$ is radial and satisfies

$$u(x, t) = u(x, -t), \quad u_t(x, t) = -u_t(x, -t).$$

Therefore G_-, G_+ are radial, i.e., independent of ω and satisfy $G_+(s) = -G_-(s)$. We may apply Theorem 1.5 and obtain $G_+(s) = \mathcal{H}G_-(s)$. As a result, G_- satisfies the identity $\mathcal{H}G_-(s) = -G_-(s)$. Next, let us assume G_- satisfies this identity. Then we have

$$G_-(s) = \frac{G_-(s) - \mathcal{H}G_-(s)}{2} \in \left\{ \frac{G(s) - \mathcal{H}G(s)}{2} : G \in L^2(\mathbb{R}) \right\}.$$

Finally, if $G_-(s) = (G(s) - \mathcal{H}G(s))/2$, we show there exists $u_0 \in \dot{H}_{\text{rad}}^1(\mathbb{R}^d)$, so that $G_- = T_-(u_0, 0)$. In fact, we consider radial initial data $(u_0, u_1) = T_-^{-1}G$ and free wave $u = \mathcal{S}_L(u_0, u_1)$. We may reverse the time and obtain $u(x, -t) = \mathcal{S}_L(u_0, -u_1)(x, t)$. Thus

$$T_-(u_0, -u_1)(s) = -T_+(u_0, u_1)(s) = -\mathcal{H}G(s).$$

Therefore we have

$$T_-(2u_0, 0)(s) = T_-(u_0, u_1) + T_-(u_0, -u_1) = G(s) - \mathcal{H}G(s) = 2G_-(s),$$

which completes the proof. □

The key observation is the following:

Lemma 5.2. *Given $g \in L^2(\mathbb{R}^+)$, there exists a function G with $\|G\|_{L^2(\mathbb{R})} \leq 2\|g\|_{L^2(\mathbb{R}^+)}$ so that*

$$G(s) - \mathcal{H}G(-s) = 2g(s), \quad s > 0, \quad \left\| \frac{G(s) - \mathcal{H}G(-s)}{2} \right\|_{L^2(\mathbb{R})} \leq \sqrt{2}\|g\|_{L^2(\mathbb{R}^+)}.$$

Proof. Let us first find a function G with $\|G\|_{L^2(\mathbb{R})} \leq 2\|g\|_{L^2}$ so that

$$G(s) - \frac{G(s) + \mathcal{H}G(-s)}{2} = g(s), \quad s > 0.$$

We define a linear bounded operator T from $L^2(\mathbb{R}^+)$ to itself. In the formula below we extend the domain of G to \mathbb{R} by assuming $G(s) = 0$ if $s < 0$ before we apply the Hilbert transform:

$$(TG)(s) = \frac{G(s) + \mathcal{H}G(-s)}{2} = \frac{G(s)}{2} - \frac{1}{2\pi} \int_0^\infty \frac{G(\tau)}{s + \tau} d\tau, \quad s > 0.$$

We may further rewrite it as

$$TG = \frac{G}{2} - \frac{1}{2\pi} L^2 G.$$

Here L is the Laplace transform

$$LG(s) = \int_0^\infty G(\tau)e^{-s\tau} d\tau,$$

which is self-adjoint operator in $L^2(\mathbb{R}^+)$ with an operator norm $\sqrt{\pi}$. More details about the Laplace transform can be found in [Lax 2002]. As a result, we have

$$\begin{aligned} \|TG\|_{L^2(\mathbb{R}^+)}^2 &= \frac{1}{4} \langle G - \frac{1}{\pi} L^2 G, G - \frac{1}{\pi} L^2 G \rangle \\ &= \frac{1}{4} \|G\|_{L^2}^2 + \frac{1}{4\pi^2} \|L^2 G\|_{L^2}^2 - \frac{1}{4\pi} \langle G, L^2 G \rangle - \frac{1}{4\pi} \langle L^2 G, G \rangle \\ &\leq \frac{1}{4} \|G\|_{L^2}^2 + \frac{1}{4\pi} \|LG\|_{L^2}^2 - \frac{1}{2\pi} \langle LG, LG \rangle = \frac{1}{4} \|G\|_{L^2}^2 - \frac{1}{4\pi} \|LG\|_{L^2}^2. \end{aligned}$$

Thus the operator norm of T is less than or equal to $\frac{1}{2}$. This means that the function

$$G = \sum_{j=0}^{\infty} T^j g \in L^2(\mathbb{R}^+)$$

satisfies the equation $G - TG = g$ and $\|G\|_{L^2(\mathbb{R}^+)} \leq 2\|g\|_{L^2(\mathbb{R}^+)}$. Finally we naturally extend the domain of G to \mathbb{R} by defining $G(s) = 0$ if $s < 0$. We have

$$\frac{G(s) - \mathcal{H}G(-s)}{2} = \begin{cases} g(s), & s > 0, \\ -\frac{1}{2}\mathcal{H}G(-s), & s < 0. \end{cases}$$

Therefore we may find an upper bound of the L^2 norm

$$\left\| \frac{G(s) - \mathcal{H}G(-s)}{2} \right\|_{L^2(\mathbb{R})}^2 \leq \|g\|_{L^2(\mathbb{R}^+)}^2 + \frac{1}{4}\|\mathcal{H}G\|_{L^2(\mathbb{R})}^2 \leq 2\|g\|_{L^2(\mathbb{R}^+)}^2. \quad \square$$

Proof of Proposition 1.11. Let $G_- = T_-(u_0, 0)$ and $g(s)$ be its cut-off version:

$$g(s) = \begin{cases} G_-(s), & s > R, \\ 0, & s < R. \end{cases}$$

Then radiation field implies that the free wave $u = S_L(u_0, 0)$ satisfies

$$\lim_{t \rightarrow -\infty} \int_{|x| > R+|t|} |\nabla u(x, t)|^2 dx = \lim_{t \rightarrow -\infty} \int_{|x| > R+|t|} |u_t(x, t)|^2 dx = \sigma_{4k-1} \|g\|_{L^2(\mathbb{R}^+)}^2. \quad (19)$$

Here again σ_{4k-1} is the area of the sphere \mathbb{S}^{4k-1} . According to Lemmas 5.1 and 5.2, there exists a function $\tilde{u}_0 \in \dot{H}_{\text{rad}}^1(\mathbb{R}^{4k})$, so that

$$T_-(\tilde{u}_0, 0)(s) = g(s), \quad s > 0, \quad \|\tilde{u}_0\|_{\dot{H}^1(\mathbb{R}^{4k})}^2 \leq 4\sigma_{4k-1} \|g\|_{L^2(\mathbb{R}^+)}^2.$$

Therefore $T_-(u_0 - \tilde{u}_0, 0)$ vanishes if $s > R$. A combination of this fact with the time symmetry gives

$$\lim_{t \rightarrow \pm\infty} \int_{|x| > |t|+R} |\nabla_{t,x} S_L(u_0 - \tilde{u}_0, 0)(x, t)|^2 dx = 0.$$

As a result, we may apply Proposition 1.9 and conclude $u_0 - \tilde{u}_0 \in Q_k(R)$. This means

$$\|\Pi_{Q_k(R)}^\perp u_0\|_{\dot{H}^1(\{x:|x|>R\})}^2 \leq \|\tilde{u}_0\|_{\dot{H}^1(\{x:|x|>R\})}^2 \leq 4\sigma_{4k-1} \|g\|_{L^2(\mathbb{R}^+)}^2.$$

A combination of this inequality and identity (19) immediately verifies the conclusion of Proposition 1.11 in the negative time direction. The positive time direction follows the time symmetry.

6. Nonradial exterior energy estimates

In this section we give a short proof of Proposition 1.14. We start with:

Lemma 6.1. *Let $d \geq 3$ be an odd integer. Then*

$$\sum_{\pm} \lim_{t \rightarrow \pm\infty} \int_{|x| > R+|t|} |\nabla_{t,x} S_L(u_0, u_1)(x, t)|^2 dx = 2 \int_{|s| > R} \int_{\mathbb{S}^{d-1}} |T_-(u_0, u_1)(s, \theta)|^2 d\theta ds. \quad (20)$$

In particular, we have (see (4) for the definition of $P(R)$)

$$\mathbf{T}_-(P(R)) = \mathcal{P}(R) \doteq \{G_- \in L^2(\mathbb{R} \times \mathbb{S}^{d-1}) : \text{supp } G_- \subset [-R, R] \times \mathbb{S}^{d-1}\}.$$

Proof. Let u be the solution of linear wave equation with initial data (u_0, u_1) . Then by radiation field (Theorem 1.1) we have

$$\begin{aligned} \lim_{t \rightarrow -\infty} \int_{|x| > |t|+R} |\nabla_{t,x} u|^2 dx &= 2 \int_R^\infty \int_{\mathbb{S}^{d-1}} |G_-(s, \theta)|^2 d\theta ds, \\ \lim_{t \rightarrow -\infty} \int_{|x| < |t|-R} |\nabla_{t,x} u|^2 dx &= 2 \int_{-\infty}^{-R} \int_{\mathbb{S}^{d-1}} |G_-(s, \theta)|^2 d\theta ds. \end{aligned}$$

In addition, we may apply the energy conservation law, Proposition 1.2 and obtain

$$\begin{aligned} \lim_{t \rightarrow -\infty} \int_{|x| < |t|-R} |\nabla_{t,x} u|^2 dx &= \int_{\mathbb{R}^d} (|\nabla u_0|^2 + |u_1|^2) dx - \lim_{t \rightarrow -\infty} \int_{|x| > |t|-R} |\nabla_{t,x} u|^2 dx \\ &= \lim_{t \rightarrow +\infty} \int_{|x| > t+R} |\nabla_{t,x} u|^2 dx. \end{aligned}$$

Combining these identities we have

$$\sum_{\pm} \lim_{t \rightarrow \pm\infty} \int_{|x| > R+|t|} |\nabla_{t,x} u(x, t)|^2 dx = 2 \int_{|s| > R} \int_{\mathbb{S}^{d-1}} |G_-(s, \theta)|^2 d\theta ds.$$

Finally $(u_0, u_1) \in P(R)$ is equivalent to saying

$$\int_{|s| > R} \int_{\mathbb{S}^{d-1}} |G_-(s, \theta)|^2 d\theta ds = 0,$$

namely $\text{supp } G_- \subset [-R, R] \times \mathbb{S}^{d-1}$. □

Now we are ready to prove Proposition 1.14. Since $\sqrt{2}\mathbf{T}_-$ is a bijective isometry from $\dot{H}^1 \times L^2(\mathbb{R}^d)$ to $L^2(\mathbb{R} \times \mathbb{S}^{d-1})$, we have

$$\mathbf{\Pi}_{P(R)}^\perp(u_0, u_1) = \mathbf{T}_-^{-1} \mathbf{\Pi}_{\mathbf{T}_-(P(R))}^\perp \mathbf{T}_-(u_0, u_1).$$

We next use the expression of $\mathcal{P}(R) = \mathbf{T}_-(P(R))$:

$$\begin{aligned} \|\mathbf{\Pi}_{P(R)}^\perp(u_0, u_1)\|_{\dot{H}^1 \times L^2}^2 &= 2 \|\mathbf{\Pi}_{\mathcal{P}(R)}^\perp \mathbf{T}_-(u_0, u_1)\|_{L^2(\mathbb{R} \times \mathbb{S}^{d-1})}^2 \\ &= 2 \int_{|s| > R} \int_{\mathbb{S}^{d-1}} |\mathbf{T}_-(u_0, u_1)(s, \theta)|^2 d\theta ds. \end{aligned}$$

Combining this with (20) we finish the proof.

Appendix

In this section we give a brief proof of Lemma 4.6 for completeness. We first prove this lemma for two special cases, $P(x) = 1$ and $P(x) = 1 - x^2$. We start with $P(x) = 1$. A straightforward calculation gives

$$\begin{aligned}
\pi W(s) &= \text{p.v.} \int_{-1}^1 \frac{(1-x^2)^{-1/2}}{s-x} dx \\
&= \text{p.v.} \int_{-1}^1 \frac{(1-s^2)^{-1/2}}{s-x} dx + \int_{-1}^1 \frac{(1-x^2)^{-1/2} - (1-s^2)^{-1/2}}{s-x} dx \\
&= (1-s^2)^{-1/2} \ln \left| \frac{1+s}{1-s} \right| + \int_{-1}^1 \frac{(1-s^2) - (1-x^2)}{(s-x)\sqrt{1-x^2}\sqrt{1-s^2}(\sqrt{1-x^2} + \sqrt{1-s^2})} dx \\
&= (1-s^2)^{-1/2} \ln \left| \frac{1+s}{1-s} \right| + \frac{-s}{\sqrt{1-s^2}} \int_{-1}^1 \frac{1}{\sqrt{1-x^2}(\sqrt{1-x^2} + \sqrt{1-s^2})} dx.
\end{aligned}$$

Next we apply the change of variables $x = 2z/(1+z^2)$. We have

$$\sqrt{1-x^2} = \frac{1-z^2}{1+z^2} dx = \frac{2(1-z^2)}{(1+z^2)^2} dz.$$

Thus

$$\begin{aligned}
\int_{-1}^1 \frac{1}{\sqrt{1-x^2}(\sqrt{1-x^2} + \sqrt{1-s^2})} dx &= \int_{-1}^1 \frac{2 dz}{1-z^2 + \sqrt{1-s^2}(1+z^2)} \\
&= \frac{2}{s} \int_0^1 \left(\frac{1}{(1+\sqrt{1-s^2})/s-z} + \frac{1}{(1+\sqrt{1-s^2})/s+z} \right) dz \\
&= \frac{2}{s} \ln \left| \left(\frac{1+\sqrt{1-s^2}}{s} + 1/\frac{1+\sqrt{1-s^2}}{s} - 1 \right) \right| = \frac{1}{s} \ln \left| \frac{1+s}{1-s} \right|. \quad (21)
\end{aligned}$$

This immediately gives $W(s) = 0$ for $s \in (-1, 1)$. Next we consider the case $P(x) = 1-x^2$. In this case we calculate the Hilbert transform of $\sqrt{1-x^2}$:

$$\begin{aligned}
\pi W(s) &= \text{p.v.} \int_{-1}^1 \frac{\sqrt{1-x^2}}{s-x} dx \\
&= \text{p.v.} \int_{-1}^1 \frac{\sqrt{1-s^2}}{s-x} dx + \int_{-1}^1 \frac{\sqrt{1-x^2} - \sqrt{1-s^2}}{s-x} dx \\
&= \sqrt{1-s^2} \ln \left| \frac{1+s}{1-s} \right| + \int_{-1}^1 \frac{(1-x^2) - (1-s^2)}{(s-x)(\sqrt{1-x^2} + \sqrt{1-s^2})} dx \\
&= \sqrt{1-s^2} \ln \left| \frac{1+s}{1-s} \right| + s \int_{-1}^1 \frac{1}{\sqrt{1-x^2} + \sqrt{1-s^2}} dx \\
&= \sqrt{1-s^2} \ln \left| \frac{1+s}{1-s} \right| + \pi s + s \int_{-1}^1 \left(\frac{1}{\sqrt{1-x^2} + \sqrt{1-s^2}} - \frac{1}{\sqrt{1-x^2}} \right) dx \\
&= \sqrt{1-s^2} \ln \left| \frac{1+s}{1-s} \right| + \pi s - s\sqrt{1-s^2} \int_{-1}^1 \frac{1}{\sqrt{1-x^2}(\sqrt{1-x^2} + \sqrt{1-s^2})} dx = \pi s.
\end{aligned}$$

Here we use the integral (21) again.

Induction. Now we are ready to prove Lemma 4.6 by induction. It is clear that we only need to show the Hilbert transform of $f_\kappa(x) = x^\kappa(1-x^2)^{-1/2}$ is a polynomial of degree $\kappa - 1$ in the interval $(-1, 1)$. The

cases of $\kappa = 0, 2$ have been done. Now let us consider the case of $f_1(x) = x(1-x^2)^{-1/2}$. We observe that ($s \in (-1, 1)$)

$$\mathcal{H}f_1 = \mathcal{H} \frac{d}{dx} (-\sqrt{1-x^2}) = -\frac{d}{ds} \mathcal{H}(\sqrt{1-x^2}) = -1.$$

This proves the case $\kappa = 1$. Now let us assume that the cases $\kappa = 0, 1, 2, \dots, n$ are done and consider the case $\kappa = n + 1$. Here $n \geq 2$. We have

$$x^{n+1}(1-x^2)^{-1/2} = -x^{n-1}(1-x^2)^{1/2} + x^{n-1}(1-x^2)^{-1/2}.$$

The Hilbert transform of the second term in the right-hand side is known to be a polynomial of degree $n-2$. Thus we only need to consider the first term. We have

$$\begin{aligned} \frac{d}{ds} \mathcal{H}(x^{n-1}(1-x^2)^{1/2}) &= \mathcal{H} \frac{d}{dx} (x^{n-1}(1-x^2)^{1/2}) \\ &= \mathcal{H}\{-nx^n + (n-1)x^{n-2}\}(1-x^2)^{-1/2}. \end{aligned}$$

This is a polynomial of degree $n-1$ by induction hypothesis. A simple integration then finishes the proof of the case $\kappa = n + 1$. Generally speaking, the derivative with respect to s as given above is in the weak sense. But since the derivative is known to be a polynomial in $(-1, 1)$, we can integrate as usual.

Acknowledgement

Shen is financially supported by National Natural Science Foundation of China, projects 12071339 and 11771325.

References

- [Côte, Kenig and Schlag 2014] R. Côte, C. E. Kenig, and W. Schlag, “Energy partition for the linear radial wave equation”, *Math. Ann.* **358**:3-4 (2014), 573–607. MR Zbl
- [Duyckaerts, Kenig and Merle 2011] T. Duyckaerts, C. Kenig, and F. Merle, “Universality of blow-up profile for small radial type II blow-up solutions of the energy-critical wave equation”, *J. Eur. Math. Soc.* **13**:3 (2011), 533–599. MR Zbl
- [Duyckaerts, Kenig and Merle 2012] T. Duyckaerts, C. Kenig, and F. Merle, “Universality of the blow-up profile for small type II blow-up solutions of the energy-critical wave equation: the nonradial case”, *J. Eur. Math. Soc.* **14**:5 (2012), 1389–1454. MR Zbl
- [Duyckaerts, Kenig and Merle 2013] T. Duyckaerts, C. Kenig, and F. Merle, “Classification of radial solutions of the focusing, energy-critical wave equation”, *Cambridge J. Math.* **1**:1 (2013), 75–144. MR Zbl
- [Duyckaerts, Kenig and Merle 2014] T. Duyckaerts, C. Kenig, and F. Merle, “Scattering for radial, bounded solutions of focusing supercritical wave equations”, *Int. Math. Res. Not.* **2014**:1 (2014), 224–258. MR Zbl
- [Duyckaerts, Kenig and Merle 2019] T. Duyckaerts, C. Kenig, and F. Merle, “Scattering profile for global solutions of the energy-critical wave equation”, *J. Eur. Math. Soc.* **21**:7 (2019), 2117–2162. MR Zbl
- [Duyckaerts, Kenig and Merle 2021] T. Duyckaerts, C. Kenig, and F. Merle, “Decay estimates for nonradiative solutions of the energy-critical focusing wave equation”, *J. Geom. Anal.* **31**:7 (2021), 7036–7074. MR Zbl
- [Duyckaerts, Kenig and Merle 2023] T. Duyckaerts, C. Kenig, and F. Merle, “Soliton resolution for the radial critical wave equation in all odd space dimensions”, *Acta Math.* **230**:1 (2023), 1–92. MR
- [Duyckaerts, Kenig, Martel and Merle 2022] T. Duyckaerts, C. Kenig, Y. Martel, and F. Merle, “Soliton resolution for critical co-rotational wave maps and radial cubic wave equation”, *Comm. Math. Phys.* **391**:2 (2022), 779–871. MR Zbl

- [Evans 1998] L. C. Evans, *Partial differential equations*, Grad. Stud. Math. **19**, Amer. Math. Soc., Providence, RI, 1998. MR Zbl
- [Friedlander 1962] F. G. Friedlander, “On the radiation field of pulse solutions of the wave equation”, *Proc. Roy. Soc. London Ser. A* **269** (1962), 53–65. MR Zbl
- [Friedlander 1973] F. G. Friedlander, “An inverse problem for radiation fields”, *Proc. Lond. Math. Soc.* (3) **27**:3 (1973), 551–576. MR Zbl
- [Friedlander 1980] F. G. Friedlander, “Radiation fields and hyperbolic scattering theory”, *Math. Proc. Cambridge Philos. Soc.* **88**:3 (1980), 483–515. MR Zbl
- [Ginibre, Soffer and Velo 1992] J. Ginibre, A. Soffer, and G. Velo, “The global Cauchy problem for the critical nonlinear wave equation”, *J. Funct. Anal.* **110**:1 (1992), 96–130. MR Zbl
- [Grillakis 1990] M. G. Grillakis, “Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity”, *Ann. of Math.* (2) **132**:3 (1990), 485–509. MR Zbl
- [Grillakis 1992] M. G. Grillakis, “Regularity for the wave equation with a critical nonlinearity”, *Comm. Pure Appl. Math.* **45**:6 (1992), 749–774. MR Zbl
- [Kapitanski 1994] L. Kapitanski, “Weak and yet weaker solutions of semilinear wave equations”, *Comm. Partial Differential Equations* **19**:9-10 (1994), 1629–1676. MR Zbl
- [Katayama 2013] S. Katayama, “Asymptotic behavior for systems of nonlinear wave equations with multiple propagation speeds in three space dimensions”, *J. Differential Equations* **255**:1 (2013), 120–150. MR Zbl
- [Kenig and Merle 2008] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation”, *Acta Math.* **201**:2 (2008), 147–212. MR Zbl
- [Kenig, Lawrie and Schlag 2014] C. E. Kenig, A. Lawrie, and W. Schlag, “Relaxation of wave maps exterior to a ball to harmonic maps for all data”, *Geom. Funct. Anal.* **24**:2 (2014), 610–647. MR Zbl
- [Kenig, Lawrie, Liu and Schlag 2015] C. Kenig, A. Lawrie, B. Liu, and W. Schlag, “Channels of energy for the linear radial wave equation”, *Adv. Math.* **285** (2015), 877–936. MR Zbl
- [Lax 2002] P. D. Lax, *Functional analysis*, Wiley, New York, 2002. MR Zbl
- [Lindblad and Sogge 1995] H. Lindblad and C. D. Sogge, “On existence and scattering with minimal regularity for semilinear wave equations”, *J. Funct. Anal.* **130**:2 (1995), 357–426. MR Zbl
- [Nakanishi 1999a] K. Nakanishi, “Scattering theory for the nonlinear Klein–Gordon equation with Sobolev critical power”, *Int. Math. Res. Not.* **1999**:1 (1999), 31–60. MR Zbl
- [Nakanishi 1999b] K. Nakanishi, “Unique global existence and asymptotic behaviour of solutions for wave equations with non-coercive critical nonlinearity”, *Comm. Partial Differential Equations* **24**:1-2 (1999), 185–221. MR Zbl
- [Shatah and Struwe 1993] J. Shatah and M. Struwe, “Regularity results for nonlinear wave equations”, *Ann. of Math.* (2) **138**:3 (1993), 503–518. MR Zbl
- [Shatah and Struwe 1994] J. Shatah and M. Struwe, “Well-posedness in the energy space for semilinear wave equations with critical growth”, *Int. Math. Res. Not.* **1994**:7 (1994), 303–309. MR Zbl
- [Shen 2013] R. Shen, “On the energy subcritical, nonlinear wave equation in \mathbb{R}^3 with radial data”, *Anal. PDE* **6**:8 (2013), 1929–1987. MR Zbl
- [Solmon 1987] D. C. Solmon, “Asymptotic formulas for the dual Radon transform and applications”, *Math. Z.* **195**:3 (1987), 321–343. MR Zbl

Received 28 Nov 2021. Accepted 11 Aug 2022.

LIANG LI: 17864193561@163.com
Center for Applied Mathematics, Tianjin University, Tianjin, China

RUIPENG SHEN: srpgow@163.com
Center for Applied Mathematics, Tianjin University, Tianjin, China

LIJUAN WEI: lijuanwei8@163.com
Center for Applied Mathematics, Tianjin University, Tianjin, China

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 2 2024

On a spatially inhomogeneous nonlinear Fokker–Planck equation: Cauchy problem and diffusion asymptotics	379
FRANCESCA ANCESCHI and YUZHE ZHU	
Strichartz inequalities with white noise potential on compact surfaces	421
ANTOINE MOUZARD and IMMANUEL ZACHHUBER	
Curvewise characterizations of minimal upper gradients and the construction of a Sobolev differential	455
SYLVESTER ERIKSSON-BIQUE and ELEFTERIOS SOULTANIS	
Smooth extensions for inertial manifolds of semilinear parabolic equations	499
ANNA KOSTIANKO and SERGEY ZELIK	
Semiclassical eigenvalue estimates under magnetic steps	535
WAFAA ASSAAD, BERNARD HELFFER and AYMAN KACHMAR	
Necessary density conditions for sampling and interpolation in spectral subspaces of elliptic differential operators	587
KARLHEINZ GRÖCHENIG and ANDREAS KLOTZ	
On blowup for the supercritical quadratic wave equation	617
ELEK CSOBO, IRFAN GLOGIĆ and BIRGIT SCHÖRKHUBER	
Arnold’s variational principle and its application to the stability of planar vortices	681
THIERRY GALLAY and VLADIMÍR ŠVERÁK	
Explicit formula of radiation fields of free waves with applications on channel of energy	723
LIANG LI, RUIPENG SHEN and LIJUAN WEI	
On L^∞ estimates for Monge–Ampère and Hessian equations on nef classes	749
BIN GUO, DUONG H. PHONG, FREID TONG and CHUWEN WANG	