

A combination theorem for strong relative hyperbolicity

MAHAN MJ
LAWRENCE REEVES

We prove a combination theorem for trees of (strongly) relatively hyperbolic spaces and finite graphs of (strongly) relatively hyperbolic groups. This gives a geometric extension of Bestvina and Feighn’s Combination Theorem for hyperbolic groups and answers a question of Swarup. We also prove a converse to the main Combination Theorem.

[20F67](#); [57M50](#)

1 Introduction

In [4], Bestvina and Feighn proved a combination theorem for hyperbolic groups. Motivated by this, Swarup asked the analogous question [3] for relatively hyperbolic groups. Dahmani [9] and Alibegović [1] have proven combination theorems motivated by applications to convergence groups and limit groups (cf Sela [28]).

In this paper, we prove a *geometric combination theorem* (as opposed to a dynamical one) for trees of (strong) relatively hyperbolic metric spaces. We use Bestvina and Feighn’s Combination Theorem [4] directly in deducing the relevant combination theorem. The conditions we impose are quite different from those of [9] and [1]. Our main Theorems 4.5 and 4.7 are stated below:

Strong Combination Theorem and converse: Theorem 4.5 and Theorem 4.7 *Let X be a tree (T) of strongly relatively hyperbolic spaces satisfying:*

- (1) *the q (uasi)- i (sometrically)-embedded condition*
- (2) *the strictly type-preserving condition*
- (3) *the qi -preserving electrocution condition*
- (4) *the induced tree of coned-off spaces satisfies the hallways flare condition*
- (5) *the cone-bounded hallways strictly flare condition.*

Then X is strongly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces.

Conversely, if X be a tree (T) of strongly relatively hyperbolic spaces satisfying conditions (1)–(3) such that X is strongly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces, then the tree of spaces satisfies conditions (4)–(5).

Of the conditions given in the above Theorem, Condition (1) is taken directly from Bestvina and Feighn [4]. Condition (2) roughly says that the preimage of a horosphere-like subset (thought of as parabolic) in a vertex space (under the edge-space to vertex-space map) is either empty or a horosphere-like subset in the corresponding edge-space. This condition may be likened to the restriction to strictly type-preserving maps in the theory of Kleinian groups. Condition (2) ensures an induced tree of electrocuted spaces. Condition (3) says that the induced tree of spaces also satisfies the qi -embedded condition. Condition (4) is again taken directly from [4]. Condition (5) is the one essential new condition. It says roughly that a pair of geodesics whose vertices consist only of cone-points cannot lie close to each other for long. The notion of *fully quasiconvex subgroups* introduced by Dahmani [9] is related to Condition (3), the *qi-preserving electrocution condition*.

Note In this paper we adopt the convention that the are horosphere-like subsets are *coarsely proper*, ie no finite neighborhood of a horosphere-like subset is the whole space (cf Behrstock, Drutu and Mosher [2] which follows a similar convention). This excludes the trivial case that X is strongly hyperbolic relative to itself, or a net in X . This assumption translates into the context of groups. Hence we assume that if a group G is strongly hyperbolic relative to a collection of subgroups, then no subgroup H in the collection is of finite index in G .

As an immediate consequence of [Theorem 4.5](#), we have:

Strong Combination Theorem for graphs of groups: [Theorem 4.6](#) *Let G be a finite graph (Γ) of strongly relatively hyperbolic groups satisfying:*

- (1) *the qi -embedded condition*
- (2) *the strictly type-preserving condition*
- (3) *the qi -preserving electrocution condition*
- (4) *the induced tree of coned-off spaces satisfies the hallways flare condition*
- (5) *the cone-bounded hallways strictly flare condition.*

Then G is strongly hyperbolic relative to the family \mathcal{C} of maximal parabolic subgroups.

All these conditions are satisfied in the classical case of a 3–manifold fibering over the circle with fiber a punctured surface. The one condition that needs checking is the *hallways flare condition* for the induced tree (in fact line) of coned-off spaces. This fact is due to Bowditch [7, Section 6]. The verification involves using the associated singular structure coming from stable and unstable foliations. We shall give a slightly modified version, using an idea of Mosher [26] to show this. (See Section 4.3). In fact we prove the stronger Theorem:

Theorem 4.9 *Let $\Phi_1 \cdots \Phi_m$ be m pseudo-Anosov diffeomorphisms of Σ with different sets of stable and unstable foliations. Let $H = \pi_1(\Sigma)$. Then there is an $n \geq 1$ such that the diffeomorphisms $\Phi_1^n, \dots, \Phi_m^n$ generate a free group F and the group G given by the exact sequence*

$$1 \rightarrow H \rightarrow G \rightarrow F \rightarrow 1$$

is (strongly) hyperbolic relative to the maximal parabolic subgroups of the form $\mathcal{Z} \times F$.

We remark here that in Dahmani’s combination theorem [9] an essential condition is *acylindricity*. Again, in Alibegović’s combination theorem [1], an essential assumption is the *compact intersection property*. Both acylindricity and the compact intersection property prevent infinite (or even arbitrarily long chains of parabolics from occurring). To us, this seemed a bit unsatisfactory, as the original motivation for the Bestvina–Feighn result came from Thurston’s monster theorem (see Kapovich [16]), and we wanted a generalization of the Bestvina–Feighn theorem that would cover the case of hyperbolic 3–manifolds with parabolics, particularly hyperbolic 3–manifolds of finite volume fibering over the circle. The hypotheses in the present paper do allow for infinite chains of parabolics and covers the above case. Our emphasis here is geometric and so the main theorem is stated in terms of spaces rather than groups.

Acknowledgements The authors are grateful to G A Swarup, who was instrumental in bringing about this collaborative effort. The work was completed during a visit of the second author to RKM Vivekananda University in February 2006. We are also grateful to the referee for helpful comments and corrections. ¹

¹After the submission of this paper, we learnt of the paper [11] by Gautero, which gives a different proof of a result equivalent to Theorem 4.6. A couple of points of difference between our work and [11]: We use Bestvina–Feighn’s result [4] directly, whereas an alternate proof of the Combination Theorem of [4] is provided in [11]. However, in this paper, we provide in addition, a converse (Theorem 4.7) to the main Combination Theorem.

2 Relative hyperbolicity

In this section, we shall first recall certain notions of relative hyperbolicity due to Farb [10] and Gromov [14].

2.1 Electric geometry

Let X be a path metric space. A collection of closed subsets $\mathcal{H} = \{H_\alpha\}$ of X will be said to be *uniformly separated* if there exists $\epsilon > 0$ such that $d(H_1, H_2) \geq \epsilon$ for all distinct $H_1, H_2 \in \mathcal{H}$.

Definition 2.1 (Farb [10]) The *electric space* (or coned-off space) \widehat{X} corresponding to the pair (X, \mathcal{H}) is a metric space which consists of X and a collection of vertices v_α (one for each $H_\alpha \in \mathcal{H}$) such that each point of H_α is joined to (coned off at) v_α by an edge of length $1/2$. The sets H_α shall be referred to as *horosphere-like sets*.

A geodesic (resp. quasigeodesic) in \widehat{X} will be referred to as an electric geodesic (resp. quasigeodesic).

Definition 2.2 A path $\gamma: I \rightarrow X$ in a path metric space X is an *ambient K -quasigeodesic* if we have

$$L(\beta) \leq KL(A) + K$$

for any subsegment $\beta = \gamma|_{[a, b]}$ and any rectifiable path $A: [a, b] \rightarrow Y$ with the same endpoints. (Here L denotes length of path.)

$N_R(Z)$ will denote the R -neighborhood about the subset Z in X . $N_R^e(Z)$ will denote the R -neighborhood about the subset Z in the electric metric.

Much of what Farb proved in [10] goes through under considerably weaker assumptions than those of [10]. In [10] the theorems were proven in the particular context of a pair (X, \mathcal{H}) , where X is a Hadamard space of pinched negative curvature with the interiors of a family of horoballs \mathcal{H} removed. Then \mathcal{H} can be regarded as a collection of horospheres in X separated by a minimum distance from each other. In this situation, X is not a hyperbolic metric space itself, but is hyperbolic relative to a collection of separated *horospheres*. Alternately let H_h be the horoball corresponding to the horosphere $H \in \mathcal{H}$. Let $X_h = X \cup_{H \in \mathcal{H}} H_h$ be the entire Hadamard manifold of pinched negative curvature. Then the coned-off space \widehat{X} obtained by coning off the horospheres of X is essentially equivalent to coned-off space \widehat{X}_h obtained by coning off the horoballs of X_h .

We consider therefore a hyperbolic metric space X and a collection \mathcal{H} of (uniformly) C -quasiconvex uniformly separated subsets, ie there exists $D > 0$ such that for $H_1, H_2 \in \mathcal{H}$, $d_X(H_1, H_2) \geq D$. In this situation X is weakly hyperbolic relative to the collection \mathcal{H} in the sense that the coned-off space \widehat{X} is hyperbolic. The result in this form is due to Klarreich [17]. However, the property of *Bounded Horosphere Penetration (BHP)* or *Bounded Coset Penetration (BCP)* used by Farb [10] was not abstracted out in Klarreich's proof as it was not necessary. What is essential for BCP (or BHP) to go through has been abstracted out by Bowditch [7; 6] in the case that the collection \mathcal{H} is a collection of geodesics or horocycles in a Farey graph. (See also Bumagin [8].) But though these things are available at the level of folklore, an explicit statement seems to be lacking.

The crucial condition can be isolated as per the following definition [23]:

Definition 2.3 A collection \mathcal{H} of uniformly C -quasiconvex sets in a δ -hyperbolic metric space X is said to be *mutually D -cobounded* if for all $H_i, H_j \in \mathcal{H}$, $\pi_i(H_j)$ has diameter less than D , where π_i denotes a nearest point projection of X onto H_i . A collection is *mutually cobounded* if it is mutually D -cobounded for some D .

Mutual coboundedness was proven for horoballs by Farb in Lemma 4.7 of [10] and by Bowditch in stating that the projection of the link of a vertex onto another [6] has bounded diameter in the link. However, the comparability of intersection patterns in this context needs to be stated a bit more carefully. We give the general version of Farb's theorem below and refer to [10] and Klarreich [17] for proofs.

Lemma 2.4 (See Lemma 4.5 and Proposition 4.6 of Farb [10] and Mj [23].) *Given δ, C, D there exists Δ such that if X is a δ -hyperbolic metric space with a collection \mathcal{H} of C -quasiconvex D -separated sets. then we have the following:*

- (1) *Electric quasigeodesics electrically track hyperbolic geodesics: Given $P > 0$, there exists $K > 0$ with the property that if β is any electric P -quasigeodesic from x to y , and γ is the hyperbolic geodesic from x to y , then $\beta \subset N_K^e(\gamma)$.*
- (2) *γ lies in a hyperbolic K -neighborhood of $N_0(\beta)$, where $N_0(\beta)$ denotes the zero neighborhood of β in the electric metric.*
- (3) *Hyperbolicity: The electric space \widehat{X} is Δ -hyperbolic.*

Item (2) in the above Lemma is due to Klarreich [17], where the proof is given for β an electric geodesic, but the same proof goes through for electric quasigeodesics without backtracking.

The above Lemma motivates:

Definition 2.5 [10; 6] Let X be a geodesic metric space and \mathcal{H} be a collection of mutually disjoint uniformly separated subsets. Then X is said to be *weakly hyperbolic* relative to the collection \mathcal{H} , if the electric space \widehat{X} is hyperbolic.

We shall need to give a general definition of geodesics and quasigeodesics without backtracking.

Definition 2.6 Given a collection \mathcal{H} of C -quasiconvex, D -separated sets and a number ϵ we shall say that a geodesic (resp. quasigeodesic) γ is a geodesic (resp. quasigeodesic) *without backtracking* with respect to ϵ neighborhoods if γ does not return to $N_\epsilon(H)$ after leaving it, for any $H \in \mathcal{H}$. A geodesic (resp. quasigeodesic) γ is a geodesic (resp. quasigeodesic) *without backtracking* if it is a geodesic (resp. quasigeodesic) without backtracking with respect to ϵ neighborhoods for some $\epsilon \geq 0$.

Note For the above Lemma, the hypothesis is that \mathcal{H} consists of uniformly quasiconvex, mutually separated sets. Mutual coboundedness has not yet been used. We introduce it in the next Lemma.

Lemma 2.7 [23] Suppose X is a δ -hyperbolic metric space with a collection \mathcal{H} of C -quasiconvex K -separated D -mutually cobounded subsets. There exists $\epsilon_0 = \epsilon_0(C, K, D, \delta)$ such that the following holds:

Let β be an electric P -quasigeodesic without backtracking and γ a hyperbolic geodesic, both joining x, y . Then, given $\epsilon \geq \epsilon_0$ there is $D = D(P, \epsilon)$ such that we have:

- (1) *Similar intersection patterns 1: If precisely one of $\{\beta, \gamma\}$ meets an ϵ -neighborhood $N_\epsilon(H_1)$ of an electrocuted quasiconvex set $H_1 \in \mathcal{H}$, then the length (measured in the intrinsic path-metric on $N_\epsilon(H_1)$) from the entry point to the exit point is at most D .*
- (2) *Similar intersection patterns 2: If both $\{\beta, \gamma\}$ meet some $N_\epsilon(H_1)$ then the length (measured in the intrinsic path-metric on $N_\epsilon(H_1)$) from the entry point of β to that of γ is at most D ; similarly for exit points.*

Lemma 2.7 is essentially a paraphrasing of the BCP property [10] in terms of mutual coboundedness. The above Lemma motivates the following definition:

Definition 2.8 [10; 6] Let X be a geodesic metric space and \mathcal{H} be a collection of mutually disjoint uniformly separated subsets such that X is *weakly hyperbolic* relative to the collection \mathcal{H} . If any pair of electric quasigeodesics without backtracking starting and ending at the same point have *similar intersection patterns* with *horosphere-like sets* (elements of \mathcal{H}) then quasigeodesics are said to satisfy *Bounded Penetration* and X is said to be *strongly hyperbolic* relative to the collection \mathcal{H} .

We summarize the [Lemma 2.4](#) and [Lemma 2.7](#) as follows:

- If X is a hyperbolic metric space and \mathcal{H} a collection of uniformly quasiconvex separated subsets, then X is hyperbolic relative to the collection \mathcal{H} .
- If X is a hyperbolic metric space and \mathcal{H} a collection of uniformly quasiconvex mutually cobounded separated subsets, then X is hyperbolic relative to the collection \mathcal{H} and satisfies *Bounded Penetration*, ie hyperbolic geodesics and electric quasigeodesics have similar intersection patterns in the sense of [Lemma 2.7](#).

2.2 Partial electrocution

In this subsection, we indicate, following [\[22\]](#), a modification of Farb's [\[10\]](#) notion of *strong relative hyperbolicity* and his construction of an electric metric, described earlier. The modification we shall discuss is called *partial electrocution* and will be used in proving the converse to the Strong Combination Theorem ([Theorem 4.5](#)). Most of this discussion is taken from [\[22\]](#).

We start with a few motivating examples:

Partial electrocution of a horosphere $H = \mathbb{R}^{n-1} \times \mathbb{R}$ will be defined as putting the zero metric in the \mathbb{R}^{n-1} direction, and retaining the usual Euclidean metric in the other \mathbb{R} direction.

In the partially electrocuted case, instead of coning all of a horosphere down to a point we cone only horocyclic leaves of a foliation of the horosphere. Effectively, therefore, we have a cone-line rather a cone-point.

Let Y be a convex simply connected hyperbolic n -manifold. Let \mathcal{B} denote a collection of horoballs. Let X denote Y minus the interior of the horoballs in \mathcal{B} . Let \mathcal{H} denote the collection of boundary horospheres. Then each $H \in \mathcal{H}$ with the induced metric is isometric to a Euclidean product $E^{n-2} \times L$ for an interval $L \subset \mathbb{R}$. Partially electrocute each H by giving it the product of the zero metric with the Euclidean metric, ie on E^{n-2} give the zero metric and on L give the Euclidean metric. The resulting space is exactly what one would get by gluing to each H the mapping cylinder of the projection of H onto the L -factor.

This motivates the following scenario:

Definition 2.9 Let $(X, \mathcal{H}, \mathcal{G}, \mathcal{L})$ be an ordered quadruple such that the following holds:

- (1) X is (strongly) hyperbolic relative to a collection of subsets H_α , thought of as horospheres (and *not horoballs*).
- (2) For each H_α there is a uniform large-scale retraction $g_\alpha: H_\alpha \rightarrow L_\alpha$ to some (uniformly) δ -hyperbolic metric space L_α , ie there exist $\delta, K, \epsilon > 0$ such that for all H_α there exists a δ -hyperbolic L_α and a map $g_\alpha: H_\alpha \rightarrow L_\alpha$ with $d_{L_\alpha}(g_\alpha(x), g_\alpha(y)) \leq Kd_{H_\alpha}(x, y) + \epsilon$ for all $x, y \in H_\alpha$. Further, we denote the collection of such g_α 's as \mathcal{G} .

The *partially electrocuted space* or *partially coned-off space* corresponding to the quadruple $(X, \mathcal{H}, \mathcal{G}, \mathcal{L})$ is obtained from X by gluing in the (metric) mapping cylinders for the maps $g_\alpha: H_\alpha \rightarrow L_\alpha$.

In Farb's construction L_α is just a single point. However, the notions and arguments of Farb [10] or Klarreich [17] go through even in this setting. The metric, and geodesics and quasigeodesics in the partially electrocuted space will be referred to as the partially electrocuted metric d_{pel} , and partially electrocuted geodesics and quasigeodesics respectively. In this situation, we conclude as in Lemma 2.4:

Lemma 2.10 (X, d_{pel}) is a hyperbolic metric space and the sets L_α are uniformly quasiconvex.

Note 1 When K_α is a point, the last statement is a triviality.

Note 2 (X, d_{pel}) is strongly hyperbolic relative to the sets $\{L_\alpha\}$. In fact the space obtained by electrocuting the sets L_α in (X, d_{pel}) is just the space (X, d_e) obtained by (completely) electrocuting the sets $\{H_\alpha\}$ in X .

Note 3 The proof of Lemma 2.10 and other such results below follow Farb's [10] constructions. For instance, consider a hyperbolic geodesic η in a convex complete simply connected n -manifold X with pinched negative curvature. Let $H_i, i = 1 \cdots k$ be the partially electrocuted horoballs it meets. Let $N(\eta)$ denote the union of η and H_i 's. Let Y denote X minus the interiors of the H_i 's. The first step is to show that $N(\eta) \cap Y$ is quasiconvex in (Y, d_{pel}) . To do this one takes a hyperbolic R -neighborhood of $N(\eta)$ and projects (Y, d_{pel}) onto it, using the hyperbolic projection. It was shown by Farb in [10] that the projections of all horoballs are uniformly bounded

in hyperbolic diameter. (This is essentially mutual coboundedness). Hence, given K , choosing R large enough, any path that goes out of an R -neighborhood of $N(\eta)$ cannot be a K -partially electrocuted quasigeodesic. This is the one crucial step that allows the results of [10], in particular, Lemma 2.10 to go through in the context of partially electrocuted spaces.

As in Lemma 2.7, partially electrocuted quasigeodesics and geodesics without backtracking have the same intersection patterns with horospheres and boundaries of lifts of tubes as electric geodesics without backtracking. Further, since electric geodesics and hyperbolic quasigeodesics have similar intersection patterns with horoballs and lifts of tubes it follows that partially electrocuted quasigeodesics and hyperbolic quasigeodesics have similar intersection patterns with horospheres and boundaries of lifts of tubes. We state this formally below:

Lemma 2.11 *Given $K, \epsilon \geq 0$, there exists $C > 0$ such that the following holds:*

Let γ_{pel} and γ denote respectively a (K, ϵ) partially electrocuted quasigeodesic in (X, d_{pel}) and a hyperbolic (K, ϵ) -quasigeodesic in (Y, d) joining a, b . Then $\gamma \cap X$ lies in a (hyperbolic) C -neighborhood of (any representative of) γ_{pel} . Further, outside of a C -neighborhood of the horoballs that γ meets, γ and γ_{pel} track each other.

3 Trees of hyperbolic metric spaces

3.1 Trees of spaces: hyperbolic and relatively hyperbolic

We start with a notion closely related to one introduced in [4].

Definition 3.1 A tree (T) of hyperbolic (resp. strongly relatively hyperbolic) metric spaces satisfying the q(uasi) i(sometrically) embedded condition is a metric space (X, d) admitting a map $P: X \rightarrow T$ onto a simplicial tree T , such that there exist δ, ϵ and $K > 0$ satisfying the following:

- (1) For all vertices $v \in T$, $X_v = P^{-1}(v) \subset X$ with the induced path metric d_v is a δ -hyperbolic metric space (resp. a geodesic metric space X_v strongly hyperbolic relative to a collection $\mathcal{H}_{v\alpha}$). Further, the inclusions $i_v: X_v \rightarrow X$ are uniformly proper, ie for all $M > 0$, $v \in T$ and $x, y \in X_v$, there exists $N > 0$ such that $d(i_v(x), i_v(y)) \leq M$ implies $d_v(x, y) \leq N$.
- (2) Let e be an edge of T with initial and final vertices v_1 and v_2 respectively. Let X_e be the preimage under P of the midpoint of e . Then X_e with the induced path metric is δ -hyperbolic (resp. a geodesic metric space X_e strongly hyperbolic relative to a collection $\mathcal{H}_{e\alpha}$).

- (3) There exist maps $f_e: X_e \times [0, 1] \rightarrow X$, such that $f_e|_{X_e \times (0,1)}$ is an isometry onto the preimage of the interior of e equipped with the path metric.
- (4) $f_e|_{X_e \times \{0\}}$ and $f_e|_{X_e \times \{1\}}$ are (K, ϵ) -quasi-isometric embeddings into X_{v_1} and X_{v_2} respectively. $f_e|_{X_e \times \{0\}}$ and $f_e|_{X_e \times \{1\}}$ will occasionally be referred to as f_{v_1} and f_{v_2} respectively.
- (5) For a tree of strongly relatively hyperbolic spaces, we demand in addition, that the maps f_{v_i} above ($i = 1, 2$) are *strictly type-preserving*, ie $f_{v_i}^{-1}(H_{v_i\alpha})$, $i = 1, 2$ (for any $H_{v_i\alpha} \in \mathcal{H}_{v_i\alpha}$) is either empty or some $H_{e\alpha} \in \mathcal{H}_{e\alpha}$.
- (6) For a tree of strongly relatively hyperbolic spaces, we demand that the coned-off spaces are uniformly δ -hyperbolic.

Denote by d_v and d_e path metrics on X_v and X_e respectively. i_v, i_e will denote inclusion of X_v, X_e respectively into X .

For a tree of relatively hyperbolic spaces, the sets $H_{v\alpha}$ and $H_{e\alpha}$ will be referred to as *horosphere-like vertex sets and edge sets* respectively.

When (X, d) is a tree (T) of strongly relatively hyperbolic metric spaces, the *strictly type-preserving condition* (Condition 5 above) ensures that we obtain an induced tree (T) (the same tree T) of *coned-off, or electric spaces*. We demand further:

Qi-preserving electrocution condition The induced maps of the electric edge spaces into the electric vertex spaces $\widehat{f}_{v_i}: \widehat{X}_e \rightarrow \widehat{X}_{v_i}$ ($i = 1, 2$) are uniform quasi-isometries.

The resulting tree of coned-off spaces will be called the *induced tree of coned-off spaces*. The resulting space will be denoted as \widehat{X} .

Definition 3.2 A finite graph of (strongly) relatively hyperbolic groups is said to satisfy Condition C , if the associated tree of relatively hyperbolic Cayley graphs satisfies Condition C . Here C will be one of the following:

- (1) the qi-embedded condition
- (2) the strictly type-preserving condition
- (3) the qi-preserving electrocution condition
- (4) the induced tree of coned-off spaces satisfies the *hallways flare* condition (see [Definition 3.5](#))
- (5) the *cone-bounded hallways strictly flare* condition (see [Definition 3.6](#)).

Remark Strictly speaking, this induced tree exists for any collection of vertex and edge spaces satisfying the *strictly type-preserving condition*. Hyperbolicity is not essential for the existence of the induced tree of spaces.

The *cone locus* of \widehat{X} , the induced tree (T) of coned-off spaces, is the graph (in fact a forest) whose vertex set \mathcal{V} consists of the cone-points in the vertex set of \widehat{X} and whose edge-set \mathcal{E} consists of the cone-points in the edge set of \widehat{X} . The incidence relations are dictated by the incidence relations in T .

Note that connected components of the cone-locus can be naturally identified with subtrees of T . Each such connected component of the cone-locus will be called a *maximal cone-subtree*. The collection of *maximal cone-subtrees* will be denoted by \mathcal{T} and elements of \mathcal{T} will be denoted as T_α . Further, each maximal cone-subtree T_α naturally gives rise to a tree T_α of horosphere-like subsets depending on which cone-points arise as vertices and edges of T_α . The metric space that T_α gives rise to will be denoted as C_α and will be referred to as a *maximal cone-subtree of horosphere-like spaces*. The collection of C_α 's will be denoted as \mathcal{C} .

Note Each T_α thus appears in two guises:

- (1) as a subset of \widehat{X}
- (2) as the underlying tree of C_α

We shall have need for both these interpretations.

3.2 The Bestvina–Feighn flare condition

Next, we would like to recall the essential condition (due to Bestvina and Feighn [4]) ensuring hyperbolicity of a tree of spaces. We retain the terminology.

Definition 3.3 A disk $f: [-m, m] \times I \rightarrow X$ is a hallway of length $2m$ if it satisfies:

- (1) $f^{-1}(\cup X_v : v \in T) = \{-m, \dots, m\} \times I$.
- (2) f maps $i \times I$ to a geodesic in X_v for some vertex space.
- (3) f is transverse, relative to condition (1) to $\cup_e X_e$.

Definition 3.4 A hallway is ρ -thin if $d(f(i, t), f(i + 1, t)) \leq \rho$ for all i, t .

A hallway is λ -hyperbolic if

$$\lambda l(f(\{0\} \times I)) \leq \max\{l(f(\{-m\} \times I)), l(f(\{m\} \times I))\}.$$

A hallway is *essential* if the edge path in T resulting from projecting X onto T does not backtrack (and is therefore a geodesic segment in the tree T).

An essential hallway of length $2m$ is *cone-bounded* if $f(i \times \partial I)$ lies in the cone-locus for $i = \{-m, \dots, m\}$.

Definition 3.5 (Hallways flare condition) The tree of spaces, X , is said to satisfy the *hallways flare* condition if there are numbers $\lambda > 1$ and $m \geq 1$ such that for all ρ there is a constant $H(\rho)$ such that any ρ -thin essential hallway of length $2m$ and girth at least H is λ -hyperbolic.

Definition 3.6 (Cone-bounded hallways strictly flare condition) The tree of spaces, X , is said to satisfy the *hallways flare* condition if there are numbers $\lambda > 1$ and $m \geq 1$ such that any cone-bounded hallway of length $2m$ is λ -hyperbolic.

The main theorem of Bestvina and Feighn follows (though this is stated in [4] for groups, the proof nowhere requires uniform properness of the space). Bowditch [7] notes the equivalence of the hallways flare condition with the hyperbolicity of the tree of spaces.

Theorem 3.7 [4, pp 85-86; 7] *Let X be a tree of hyperbolic metric spaces satisfying the qi -embedded condition and the hallways flare condition. Then X is hyperbolic. Conversely, if X is hyperbolic, then hallways flare.*

Apart, from [Theorem 3.7](#) above, we shall need one more simple observation.

Lemma 3.8 *Suppose that \widehat{X} is hyperbolic. Then maximal cone-subtrees T_α are uniformly quasiconvex in \widehat{X} .*

Proof Let $P: X \rightarrow T$ be the natural projection map of the tree of spaces to the underlying subtree. Then P induces $P': \widehat{X} \rightarrow T$ as \widehat{X} may be regarded as (the same) tree (T) of coned-off spaces. P' is distance nonincreasing. Further, restricted to each T_α , P' is an isometry.

Also note that any path from $x \in H_{v_1}$ to $y \in H_{v_2}$ in \widehat{X} has length not less than $d_T(P'(x), P'(y))$, where d_T is the natural metric on T and $v_1, v_2 \in T_\alpha$.

Now suppose that $x, y \in T_\alpha \subset \widehat{X}$. Let $\gamma \subset T_\alpha \subset \widehat{X}$ be the geodesic in T_α joining x, y . It therefore follows that for any $x, y \in T_\alpha \subset \widehat{X}$ and any path A joining $x, y \in \widehat{X}$,

$$l(A) \leq d_T(P'(x), P'(y)) = l(\gamma).$$

Hence γ is quasi-isometrically (in fact isometrically) embedded in \widehat{X} and hence a geodesic in \widehat{X} . The Lemma follows. \square

4 The Combination Theorem

4.1 Weak Combination Theorem

We start with the following:

Theorem 4.1 (Weak Combination Theorem) *Let X be a tree (T) of strongly relatively hyperbolic spaces satisfying:*

- (1) *the qi-embedded condition*
- (2) *the strictly type-preserving condition*
- (3) *the qi-preserving electrocution condition*
- (4) *the induced tree of coned-off spaces satisfies the hallways flare condition.*

Then X is weakly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces.

Proof As usual let \widehat{X} denote the induced tree (T) of coned-off spaces, \mathcal{T} denote the family of maximal cone-subtrees $T_\alpha \subset \widehat{X}$. Let \widehat{X}^c denote \widehat{X} with the family of sets \mathcal{T} coned off (ie vertices v_α are introduced, one each for each T_α , and joined to points of the corresponding T_α by edges of length $1/2$).

Since vertex and edge-spaces are strongly relatively hyperbolic, then by item (6) in the definition of a tree of strongly relatively hyperbolic spaces, \widehat{X} is a tree of (uniformly) hyperbolic metric spaces.

By the *qi-preserving electrocution condition*, the induced tree of coned-off spaces satisfies the qi-embedded condition.

By the *hallways flare condition* and [Theorem 3.7](#), \widehat{X} is a hyperbolic metric space.

By [Lemma 3.8](#), the sets $T_\alpha \in \mathcal{T}$ are uniformly quasiconvex and uniformly separated.

Hence by [Lemma 2.4](#), \widehat{X} is weakly hyperbolic relative to the sets $T_\alpha \in \mathcal{T}$, ie \widehat{X}^c is a hyperbolic metric space.

Let \widehat{X}_1 denote the space obtained from X by coning off *maximal cone-subtrees of horosphere-like sets*. Then \widehat{X}_1 is quasi-isometric to \widehat{X}^c . To see this, one notes that \widehat{X}^c is obtained from X by first coning-off (or partially electrocuting) C_α 's, the maximal cone-subtrees of horosphere-like sets, to maximal cone-subtrees T_α . This gives rise to \widehat{X} . Further coning off the T_α 's gives \widehat{X}^c . On the other hand, \widehat{X}_1 is obtained from X by coning off (or completely electrocuting) the C_α 's to points in one step. The two constructions clearly give quasi-isometric spaces.

Hence \widehat{X}_1 is hyperbolic, ie X is weakly hyperbolic relative to the collection of sets $C_\alpha \in \mathcal{C}$. □

4.2 Strong Combination Theorem

Under the additional *cone-bounded hallways strictly flare condition*, we would now like to prove a stronger version of the combination [Theorem 4.1](#), ie X is *strongly hyperbolic* relative to the collection of $C_\alpha \in \mathcal{C}$:

By [Lemma 2.7](#) and [Theorem 4.1](#), it suffices to show that the sets $T_\alpha \subset \widehat{X}$ are mutually cobounded. Most of the rest of this subsection is devoted to proving *mutual coboundedness*.

The next Lemma follows easily from stability of quasigeodesics [[12](#)] [[14](#)]. (See, for instance Lemma 4.1.1 of [[18](#)].)

Lemma 4.2 *Give δ, C , there exist D, K, ϵ such that the following holds:*

Let (X, d) be a δ -hyperbolic metric space and Y a C -quasiconvex subset. Let π be a nearest-point retraction of X onto Y . Let $x, y \in X$ such that $d(\pi(x), \pi(y)) \geq D$. Then $[x, \pi(x)] \cup [\pi(x), \pi(y)] \cup [\pi(y), y]$ is a (K, ϵ) -quasigeodesic.

We use [Lemma 4.2](#) below:

Corollary 4.3 *Given δ, C , there exist D, M such that the following holds:*

Suppose that Y, Z are C -quasiconvex subsets of a δ -hyperbolic metric space (X, d) . Let π denote nearest point projection onto Y . If $\pi(Z)$ has diameter greater than D , then $\pi(Z)$ lies in an M -neighborhood of Z .

Proof Let $x, y \in Z$. By [Lemma 4.2](#), there exist D_0, K, ϵ (depending on δ, C) such that if $d(\pi(x), \pi(y)) \geq D_0$, then $[x, \pi(x)] \cup [\pi(x), \pi(y)] \cup [\pi(y), y] = \gamma$ is a (K, ϵ) -quasigeodesic. Since Z is C -quasiconvex, γ lies in an $M_0 = M_0(K, \epsilon, C, \delta)$ -neighborhood of Z .

Now, choose a, b in Y, Z respectively, such that $d(a, b) = d(Y, Z)$. From the previous paragraph, we deduce that if $z \in Z$ such that $d(\pi(z), a) \geq D_0$, then $\pi(z)$ lies in an M_0 -neighborhood of Z . Taking $D = 2D_0$ and $M = M_0 + D_0$, we are through. \square

The following Proposition deduces mutual coboundedness of maximal cone-subtrees T_α in \widehat{X} (obtained by partially electrocuting maximal cone-subtrees of horosphere-like sets) from hyperbolicity of \widehat{X} (established for instance in [Theorem 4.1](#)) and *cone-bounded hallways strictly flare condition*. This will be used in proving the Strong Combination Theorem ([Theorem 4.5](#)).

Proposition 4.4 *Suppose that the tree of coned-off spaces \widehat{X} is hyperbolic and that the cone-bounded hallways strictly flare condition is satisfied. Then there exists $D \geq 0$ such that the family of maximal cone-subtrees T_α in \widehat{X} is D -cobounded.*

Proof Suppose not. Then, by Corollary 4.3, there exists $M \geq 0$ such that for any $D \geq 0$, there exist maximal cone-subtrees T_1, T_2 and a connected subtree $T_3 \subset T_1$ such that T_3 has diameter greater than D and lies in an M -neighborhood of T_2 . Hence, there exist geodesic edge-paths $\gamma_1 \subset T_1$ and $\gamma_2 \subset T_2$ such that each γ_i has length greater than $(D - 2M)$ and such that they lie in an M -neighborhood of each other in \widehat{X} .

Next, $P(\gamma_i) \subset T$ (where T is the tree underlying \widehat{X} , the tree (T) of coned-off spaces and $P: \widehat{X} \rightarrow T$ is the natural projection). Also, $P(\gamma_i)$ ($i = 1, 2$) is abstractly isomorphic to γ_i as an edge-path, since $P: \widehat{X} \rightarrow T$ is an isometry restricted to each T_i . Then $P(\gamma_i)$ may be regarded as geodesic paths in T having lengths greater than $(D - 2M)$ and lying in an M -neighborhood of each other (since P does not increase distances). This means that $P(\gamma_i)$ (for $i = 1, 2$) must overlap over an interval of length at least $(D - 4M)$.

Let $\alpha_i \subset \gamma_i$ be paths having length at least $(D - 4M)$ with $P(\alpha_1) = P(\alpha_2)$. Then there exists $M_1 = M_1(M)$ and a cone-bounded hallway $\Delta: [-m, m] \times I \rightarrow \widehat{X}$ with $2m \geq (D - 4M)$ such that:

- (1) $\Delta([-m, m] \times \{0\}) = \alpha_1$
- (2) $\Delta([-m, m] \times \{1\}) = \alpha_2$
- (3) each $\Delta(j \times I)$ has length less than M_1 .

Since D , and hence m can be arbitrarily large, while M (and hence M_1) are fixed, it follows that for any given $\lambda > 1$, there exists a hallway Δ , which is not λ -hyperbolic. This violates the cone-bounded hallways strictly flare condition. Hence, by contradiction, there exists $D \geq 0$ such that the family of maximal cone-subtrees T_α in \widehat{X} is D -cobounded. □

We are now in a position to prove:

Theorem 4.5 (Strong Combination Theorem) *Let X be a tree (T) of strongly relatively hyperbolic spaces satisfying:*

- (1) *the qi-embedded condition*
- (2) *the strictly type-preserving condition*

- (3) the qi -preserving electrocution condition
- (4) the induced tree of coned-off spaces satisfies the hallways flare condition
- (5) the cone-bounded hallways strictly flare condition.

Then X is strongly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces.

Proof By [Theorem 4.1](#), we know that X is weakly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces.

This is equivalent to saying that \widehat{X} is weakly hyperbolic relative to the family \mathcal{T} of maximal cone-subtrees $T_\alpha \subset \widehat{X}$.

By the cone-bounded hallways strictly flare condition and [Proposition 4.4](#), we see that the family \mathcal{T} is mutually cobounded.

Hence by [Lemma 2.7](#), we conclude that \widehat{X} is strongly hyperbolic relative to the family \mathcal{T} of maximal cone-subtrees $T_\alpha \subset \widehat{X}$. Equivalently, X is strongly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces. \square

Recall that a finite graph of (strongly) relatively hyperbolic groups is said to satisfy a Condition C , if the associated tree of relatively hyperbolic Cayley graphs also satisfies Condition C . The resulting group will be denoted as G . A quotient of maximal cone-subtrees of horosphere-like spaces in this case, is called a maximal cone-subgraph of horosphere-like subgroups. Note that such a subgraph gives rise to a subgroup of G . We shall refer to such subgroups as maximal parabolic subgroups. Recall that we are using the convention that all parabolic subgroups are of infinite index. As an immediate consequence of [Theorem 4.5](#), we have the following:

Theorem 4.6 (Strong Combination Theorem for graphs of groups) *Let G be a finite graph (Γ) of strongly relatively hyperbolic groups satisfying:*

- (1) the qi -embedded condition
- (2) the strictly type-preserving condition
- (3) the qi -preserving electrocution condition
- (4) the induced tree of coned-off spaces satisfies the hallways flare condition
- (5) the cone-bounded hallways strictly flare condition.

Then G is strongly hyperbolic relative to the family \mathcal{C} of maximal parabolic subgroups.

4.3 Converse to the Strong Combination Theorem

Recall that the *partially electrocuted space* or *partially coned-off space* corresponding to a quadruple $(X, \mathcal{H}, \mathcal{G}, \mathcal{L})$ is obtained from X by gluing in the (metric) mapping cylinders for the maps $g_\alpha: H_\alpha \rightarrow L_\alpha$. Note that from [Theorem 4.5](#), it follows that \widehat{X} is obtained from X by partially electrocuting each C_α . Here:

- (1) $H_\alpha = C_\alpha$ and $\mathcal{H} = \mathcal{C}$
- (2) $L_\alpha = T_\alpha$ and $\mathcal{L} = \mathcal{T}$
- (3) $g_\alpha: C_\alpha \rightarrow T_\alpha$ collapses C_α , the tree of horosphere-like spaces to the underlying tree T_α .

Theorem 4.7 (Converse to Strong Combination Theorem) *Let X be a tree (T) of strongly relatively hyperbolic spaces satisfying:*

- (1) *the qi-embedded condition*
- (2) *the strictly type-preserving condition*
- (3) *the qi-preserving electrocution condition*
- (4) *X is strongly hyperbolic relative to the family \mathcal{C} of maximal cone-subtrees of horosphere-like spaces.*

Then the induced tree of coned-off spaces satisfies the hallways flare condition and the cone-bounded hallways strictly flare condition.

Proof As usual let C_α denote maximal cone subtrees (T_α) of horosphere-like sets. By [Lemma 2.10](#), the induced tree of coned-off spaces \widehat{X} , obtained by partially electrocuting each C_α to T_α . Then by the converse part of [Theorem 3.7](#), hallways, including cone-bounded hallways flare.

It remains to show that cone-bounded hallways *strictly* flare. Suppose not. Then there exists D_0 such that for all $N \in \mathbb{N}$, there exist cone-bounded hallways of length greater than N , bounded by "vertical" (parametrized) geodesics λ_1, λ_2 in distinct cone-subtrees T_1, T_2 respectively such that $d(\lambda_1(i), \lambda_2(i)) \leq D_0$ for all $i = 0, \dots, N$. Let μ_0, μ_N denote "horizontal" paths in the hallway joining $\lambda_1(i), \lambda_2(i)$ for $i = 0, N$. Hence, there exist points a_j, b_j ($j = 0, N$) lying on the corresponding cones $C_1 \cap \mu_j, C_2 \cap \mu_j$, respectively such that $d(a_j, b_j) \leq D_0$. Then we have two paths:

- σ_1 starts at a_0 , moves to $\lambda_1(0)$ (by a cone-edge of length $1/2$), proceeds to $\lambda_1(N)$ and exits to a_N (again by a cone-edge of length $1/2$).
- σ_2 starts at a_0 , moves to b_0 by a path of length $\leq D_0$, then to $\lambda_2(0)$ (by a cone-edge of length $1/2$), proceeds to $\lambda_2(N)$, exits to b_N (again by a cone-edge of length $1/2$) and then goes to a_N by a path of length $\leq D_0$.

The path σ_1 has length 1 in the electric metric on \widehat{X}^c and σ_2 has length $\leq 2D_0 + 1$. Since D_0 is fixed, both σ_1 and σ_2 are uniform quasigeodesics beginning and ending at the same point, but have manifestly different intersection patterns with C_1, C_2 . This contradicts [Lemma 2.7](#) and hence X cannot be strongly hyperbolic relative to the collection C_α . This final contradiction proves the Theorem. \square

4.4 Examples

The first class of examples are hyperbolic 3-manifolds fibering over the circle with fiber a punctured hyperbolic surface Σ . All the conditions of [Theorem 4.6](#) are satisfied in this case of a 3-manifold fibering over the circle with fiber a punctured surface. The one condition that needs checking is the *hallways flare condition* for the induced tree (in fact line) of coned-off spaces. This fact is due to Bowditch [[7](#), Section 6]. We give here a somewhat different argument based on work of Mosher [[26](#)].

In [[26](#)], Mosher constructs examples of exact sequences of hyperbolic groups of the form

$$1 \rightarrow H \rightarrow G \rightarrow F \rightarrow 1$$

where H is a closed surface group, G is hyperbolic and F is free. (This construction is modified by Bestvina, Feighn and Handel [[5](#)] to the case where H is a free group.)

We shall modify Mosher's argument slightly to make it work for punctured surfaces.

Let Φ be a pseudo-Anosov diffeomorphism of a punctured hyperbolic surface Σ . Taking a suitable power of Φ if necessary, we may assume that Φ fixes all the punctures. The stable and unstable foliations of Φ give rise to a piecewise Euclidean metric on Σ . This metric is incomplete at the punctures. Complete it to get a surface with boundary Σ_B , which may be thought of as the blow-up of Σ at the punctures. Equip Σ_B with a pseudometric which is zero on all the boundary components and equal to the piecewise Euclidean metric elsewhere. This metric is discontinuous at the boundary, but this is not important. This is essentially the *electric metric* on Σ_B .

Then any electric geodesic λ in Σ_B is the union of two types of segments:

- (1) geodesics in the piecewise Euclidean metric meeting the boundary at right angles. Let λ_{eu} denote this union.
- (2) segments lying along the boundary.

The total length of such an electric geodesic is the sum of the lengths of the Euclidean pieces. The projection of the union of the Euclidean pieces λ_{eu} onto the stable and unstable foliations will be denoted by λ_{eus} and λ_{euu} respectively. Abusing notation slightly, we assume that λ_{eu} , λ_{eus} , λ_{euu} denote the respective lengths also. Then $\max(\lambda_{\text{eus}}, \lambda_{\text{euu}}) \geq 1/2\lambda_{\text{eu}}$. Let $\Phi(\lambda_{\text{eu}})$ denote the image of λ_{eu} under Φ . If we assume that the stable and unstable foliations meet the boundary components of Σ_B at right angles, then it can be easily shown that for any given $k > 1$, there is an n (depending on the stretch factor of Φ) such that

$$\max(\Phi(\lambda_{\text{eus}}), \Phi^{-1}(\lambda_{\text{euu}})) \geq k\lambda_{\text{eu}}$$

and hence we have the inequality

$$\max(\Phi(\lambda_{\text{eu}}), \Phi^{-1}(\lambda_{\text{eu}})) \geq k\lambda_{\text{eu}}.$$

This proves the one condition that needed checking, viz. the *hallways flare condition* for the induced tree (in fact line) of coned-off spaces.

More generally, we may take any m (equal to two below for concreteness) pseudo-Anosov diffeomorphisms Φ, Ψ with different stable and unstable foliations. Then generalizing the above construction, we can prove the following generalization of an essential Lemma of Mosher: *3 out of 4 stretch*.

Lemma 4.8 *For any $k > 1$, there exists $n > 0$ such that for any electric geodesic λ in Σ_B , at least three of the four elements $\Phi^n, \Phi^{-n}, \Psi^n, \Psi^{-n}$ stretch λ by a factor of k .*

Thus we get an exact sequence of groups of the form

$$1 \rightarrow H \rightarrow G \rightarrow F \rightarrow 1$$

where H is a punctured surface group, and F is free.

The Cayley graph of G may thus be regarded as a tree T of hyperbolic spaces, where T arises as the Cayley graph of F . Also, the maximal parabolic subgroups here correspond exactly to the peripheral subgroups, ie the cusp groups. Maximal cone-subtrees are each isometric to T . Then a tree T of maximal parabolics would correspond to $\mathbb{Z} \times F$. [Lemma 4.8](#) shows that the induced tree of coned-off spaces is hyperbolic.

Thus, we obtain from [Theorem 4.6](#):

Theorem 4.9 Let $\Phi_1 \cdots \Phi_m$ be m (orientation-preserving) pseudo-Anosov diffeomorphisms of Σ with different sets of stable and unstable foliations. Let $H = \pi_1(\Sigma)$. Then there is an $n \geq 1$ such that the diffeomorphisms $\Phi_1^n, \dots, \Phi_m^n$ generate a free group F and the group G given by the exact sequence

$$1 \rightarrow H \rightarrow G \rightarrow F \rightarrow 1$$

is (strongly) hyperbolic relative to the maximal parabolic subgroups of the form $\mathbb{Z} \times F$.

For $n = 1$, we get back Bowditch's theorem [7].

4.5 Applications, consequences and problems

[Theorem 4.5](#) and [Theorem 4.6](#) open up the possibility of generalizing several theorems about hyperbolic groups to (strongly) relatively hyperbolic groups.

Cannon–Thurston maps In [19], the first author proved the existence of Cannon–Thurston maps for trees of hyperbolic metric spaces. In [21], he generalized this theorem to the relatively hyperbolic case under the additional assumption that the tree of spaces gives rise to a hyperbolic 3–manifold of bounded geometry whose core is incompressible away from cusps. In [24], Mj–Pal prove the existence of Cannon–Thurston maps for the situation discussed in this paper, viz. trees of (strongly) relatively hyperbolic trees of metric spaces that are (strongly) relatively hyperbolic.

Strongly relatively hyperbolic extensions of groups A Theorem of Mosher [25] says that if an exact sequence of groups of the form

$$1 \rightarrow H \rightarrow G \rightarrow Q \rightarrow 1$$

exists, where H is hyperbolic, then there exists a quasi-isometric section of Γ_Q into Γ_G exists. In particular, if G is hyperbolic, so is Q . The essential technique is to use the action of Q on the boundary ∂H of H . A fact (due to Gromov [14]) that is used is that the space of triples of points on the boundary of a hyperbolic group H is quasi-isometric to Γ_H . An analogous result is shown by Pal in [27].

Heights of groups In [13], Gitik, Mitra, Rips and Sageev show that quasiconvex subgroups of hyperbolic groups have finite height and finite width. A partial converse was obtained by the first author in [20] for groups splitting over subgroups. This converse was used by Swarup in [29] to prove a weak hyperbolization theorem. All three theorems should have analogues in the (strongly) relatively hyperbolic world. Hruska and Wise [15] have already shown the finiteness of height and width of quasiconvex subgroups of relatively hyperbolic groups.

References

- [1] **E Alibegović**, *A combination theorem for relatively hyperbolic groups*, Bull. London Math. Soc. 37 (2005) 459–466 [MR2131400](#)
- [2] **J Behrstock, C Drutu, L Mosher**, *Thick metric spaces, relative hyperbolicity, and quasi-isometric rigidity* [arXiv:math/0512592](#)
- [3] **M Bestvina**, *Geometric group theory problem list*, available at <http://math.utah.edu/~bestvina>
- [4] **M Bestvina, M Feighn**, *A combination theorem for negatively curved groups*, J. Differential Geom. 35 (1992) 85–101 [MR1152226](#)
- [5] **M Bestvina, M Feighn, M Handel**, *Laminations, trees, and irreducible automorphisms of free groups*, Geom. Funct. Anal. 7 (1997) 215–244 [MR1445386](#)
- [6] **B H Bowditch**, *Relatively hyperbolic groups*, preprint, Southampton (1997)
- [7] **B H Bowditch**, *The Cannon–Thurston map for punctured-surface groups*, Math. Z. 255 (2007) 35–76 [MR2262721](#)
- [8] **I Bumagin**, *On definitions of relatively hyperbolic groups*, from: “Geometric methods in group theory”, Contemp. Math. 372, Amer. Math. Soc. (2005) 189–196 [MR2139687](#)
- [9] **F Dahmani**, *Combination of convergence groups*, Geom. Topol. 7 (2003) 933–963 [MR2026551](#)
- [10] **B Farb**, *Relatively hyperbolic groups*, Geom. Funct. Anal. 8 (1998) 810–840 [MR1650094](#)
- [11] **F Gautero**, *Geodesics in trees of hyperbolic and relatively hyperbolic groups* [arXiv:0710.4079](#)
- [12] **E Ghys, P de la Harpe, editor**, *Sur les groupes hyperboliques d’après Mikhael Gromov*, Progress in Math. 83, Birkhäuser, Boston (1990) [MR1086648](#) Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988
- [13] **R Gitik, M Mitra, E Rips, M Sageev**, *Widths of subgroups*, Trans. Amer. Math. Soc. 350 (1998) 321–329 [MR1389776](#)
- [14] **M Gromov**, *Hyperbolic groups*, from: “Essays in group theory”, Math. Sci. Res. Inst. Publ. 8, Springer, New York (1987) 75–263 [MR919829](#)
- [15] **G C Hruska, D T Wise**, *Packing subgroups in relatively hyperbolic groups* [arXiv:math.GR/0609369](#)
- [16] **M Kapovich**, *Hyperbolic manifolds and discrete groups*, Progress in Math. 183, Birkhäuser, Boston (2001) [MR1792613](#)
- [17] **E Klarreich**, *Semiconjugacies between Kleinian group actions on the Riemann sphere*, Amer. J. Math. 121 (1999) 1031–1078 [MR1713300](#)

- [18] **M Mitra**, *Maps on boundaries of hyperbolic metric spaces*, PhD thesis, UC Berkeley (1997)
- [19] **M Mitra**, *Cannon–Thurston maps for trees of hyperbolic metric spaces*, J. Differential Geom. 48 (1998) 135–164 [MR1622603](#)
- [20] **M Mitra**, *Height in splittings of hyperbolic groups*, Proc. Indian Acad. Sci. Math. Sci. 114 (2004) 39–54 [MR2040599](#)
- [21] **M Mj**, *Cannon–Thurston maps for pared manifolds of bounded geometry* [arXiv:math.GT/0503581](#)
- [22] **M Mj**, *Cannon–Thurston maps for surface groups I: amalgamation geometry and split geometry* [arXiv:math.GT/0512539](#)
- [23] **M Mj**, *Cannon–Thurston maps, i -bounded geometry and a theorem of McMullen* [arXiv:math.GT/0511041](#)
- [24] **M Mj**, **A Pal**, *Relative hyperbolicity, trees of spaces and Cannon–Thurston maps* [arXiv:0708.3578](#)
- [25] **L Mosher**, *Hyperbolic extensions of groups*, J. Pure Appl. Algebra 110 (1996) 305–314 [MR1393118](#)
- [26] **L Mosher**, *A hyperbolic-by-hyperbolic hyperbolic group*, Proc. Amer. Math. Soc. 125 (1997) 3447–3455 [MR1443845](#)
- [27] **A Pal**, *Cannon–Thurston maps and relative hyperbolicity*, PhD thesis, Indian Statistical Institute, Calcutta (expected 2009)
- [28] **Z Sela**, *Diophantine geometry over groups. I. Makanin–Razborov diagrams*, Publ. Math. Inst. Hautes Études Sci. (2001) 31–105 [MR1863735](#)
- [29] **G A Swarup**, *Proof of a weak hyperbolization theorem*, Q. J. Math. 51 (2000) 529–533 [MR1806457](#)

RKM Vivekananda University
Belur Math, WB-711 202, India

University of Melbourne
Victoria 3010, Australia

brmahan@gmail.com, lawrencereeves@gmail.com

Proposed: Benson Farb
Seconded: Walter Neumann, Dave Gabai

Received: 30 March 2007
Revised: 10 April 2008