



Geometry & Topology

Volume 13 (2009)

Rigidity of polyhedral surfaces, II

REN GUO
FENG LUO

Rigidity of polyhedral surfaces, II

REN GUO
FENG LUO

We study the rigidity of polyhedral surfaces using variational principles. The action functionals are derived from the cosine laws. The main focus of this paper is on the cosine law for a nontriangular region bounded by three possibly disjoint geodesics. Several of these cosine laws were first discovered and used by Fenchel and Nielsen. By studying the derivative of the cosine laws, we discover a uniform approach to several variational principles on polyhedral surfaces with or without boundary. As a consequence, the work of Penner, Bobenko and Springborn and Thurston on rigidity of polyhedral surfaces and circle patterns are extended to a very general context.

A correction was submitted on 04 Dec 2024 and posted [online](#) on 25 Aug 2025.

52C26, 52B70, 58E30; 51M10, 57Q15

1 Introduction

1.1 Variational principles

We study geometry of polyhedral surfaces using variational principles in this paper. This can be considered as a continuation of the paper by the second author [9]. By a polyhedral surface we mean an isometric gluing of geometric polygons in \mathbb{E}^2 (Euclidean plane), \mathbb{H}^2 (hyperbolic plane) or \mathbb{S}^2 (the 2–sphere). We emphasize that the combinatorics, ie, the topological cellular decomposition associated to a polyhedral surface, is considered as an intrinsic part of the polyhedral surface. The investigation of the geometry of polyhedral surfaces has a long history. Recent resurgence of interest in this subject is mainly due to the work of William Thurston on geometrization of 3–manifolds and circle packing on surfaces since 1978. Thurston’s and Andreev’s works on circle packing are nonvariational. The variational approach to circle packing was introduced in a seminal paper by Colin de Verdière [4]. Since then, many works on variational principles on triangulated surfaces have appeared, for example Brägger [2], Rivin [14], Leibon [8], Bobenko and Springborn [1], Luo [10; 11] and others. A uniform approach to variational principles on triangulated surfaces, based on the derivative of the cosine law for triangles, was proposed in [9]. It is shown in [9] that almost all known variational principles on triangulated surfaces are covered by the cosine law for triangles and right-angled hexagons and their Legendre transformations.

In [9], compact triangles in spherical, Euclidean and hyperbolic geometry are the building blocks of polyhedral surfaces. Variational principles are established for compact triangles. The goal of this paper is to develop variational principles arising from the cosine laws for hyperbolic polygons bounded by three geodesics. Figure 1 is the list of all ten cases of triangles in the Klein model of the hyperbolic plane. In Figure 2, generalized hyperbolic triangles are drawn in the Poincaré model where a horocycle is represented by a circle passing through a vertex. Cosine laws for the cases $(1, 1, -1)$, $(-1, -1, 1)$, $(-1, -1, -1)$ were discovered in Fenchel and Nielsen's work [5]. R Penner [13] discovered the cosine law for the case $(0, 0, 0)$ (decorated ideal triangles).

We observe that there is a uniform way to write the cosine laws in all these cases (Lemma 3.1). Furthermore, there is a uniform formula for the derivative cosine laws (Lemma 3.5). From the derivative cosine laws, we are able to find the complete list of localized energy functionals as in [9]. These action functionals provide variational principles for cellular decomposed surfaces. All rigidity results obtained in this paper and the work of Thurston [17], Penner [13] and Bobenko and Springborn [1] can be deduced from those concave energy functionals.




Figure 1: Generalized hyperbolic triangles in Klein model of hyperbolic plane

1.2 Generalized hyperbolic triangles

A *decorated convex polygon* in the hyperbolic plane \mathbb{H}^2 is a finite area convex polygon P so that each ideal vertex of P is associated with a horodisk centered at the vertex. A *generalized hyperbolic triangle* (or simply a *generalized triangle*) Δ in \mathbb{H}^2 is a




Figure 2: Generalized hyperbolic triangles in Poincaré model of hyperbolic plane

decorated convex polygon in \mathbb{H}^2 bounded by three distinct geodesics L_1, L_2, L_3 and all other (if any) geodesics L_{ij} perpendicular to both L_i and L_j . The complete list of them is in Figure 2. We call $L_i \cap \Delta$ an *edge* of Δ . In the Klein model of \mathbb{H}^2 , there exists a Euclidean triangle $\tilde{\Delta}$ in \mathbb{R}^2 so that each edge of $\tilde{\Delta}$ corresponds to L_1, L_2 or L_3 .

The vertices of $\tilde{\Delta}$ are called (*generalized*) *vertices* of Δ . Note that if v is a vertex of $\tilde{\Delta}$ outside $\mathbb{H}^2 \cup \partial\mathbb{H}^2$, then v corresponds to the geodesic L_{ij} perpendicular to the two edges L_i and L_j adjacent to the vertex v . The *generalized angle* (or simply *angle*) $a(v)$ at a vertex v of Δ is defined as follows. Let L_i, L_j be the edges adjacent to v . If $v \in \mathbb{H}^2$, then $a(v)$ is the inner angle of Δ at v ; if $v \in \partial\mathbb{H}^2$, $a(v)$ is TWICE of the hyperbolic length of the intersection of the associated horocycle with the cusp bounded by L_i and L_j ; if $v \notin \mathbb{H}^2 \cup \partial\mathbb{H}^2$, then $a(v)$ is the distance between L_i and L_j . Note that a generalized angle is always positive.

As in Figure 3, for a generalized vertex u of Δ , let $B_u = \{u\}$ if $u \in \mathbb{H}^2$, B_u be the horodisk at u if $u \in \partial\mathbb{H}^2$ and B_u is the half plane missing Δ bounded by L_{ij} if $u \notin \mathbb{H}^2 \cup \partial\mathbb{H}^2$. The *generalized edge length* (or *edge length* for simplicity) of L_i is




Figure 3: Generalized vertices

defined as follows. The generalized length of the edge $L_i \cap \Delta$ with vertices u, v is the distance from B_u to B_v if $B_u \cap B_v = \emptyset$ and is the negative of the distance from $\partial B_u \cap L_i$ to $\partial B_v \cap L_i$ if $B_u \cap B_v \neq \emptyset$. Note that generalized edge length may be a negative number. See Figure 4.




Figure 4: Generalized edge lengths

The cosine law (Lemma 3.1) for generalized triangles relates the generalized angles with the generalized edge lengths. Given a generalized triangle Δ and a vertex v of Δ , the type ε of v is defined to be $\varepsilon = 1$ if $v \in \mathbb{H}^2$, $\varepsilon = 0$ if $v \in \partial \mathbb{H}^2$ and $\varepsilon = -1$ if $v \notin \mathbb{H}^2 \cup \partial \mathbb{H}^2$. In this way, generalized triangles are classified into ten types $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$ where $\varepsilon_i \in \{-1, 0, 1\}$ as in Figure 2.

1.3 The work of Penner and its generalization

Suppose (\tilde{S}, \tilde{T}) is a triangulated closed surface \tilde{S} with the set of vertices V , the set of edges E . We call $T = \{\sigma - V \mid \text{a simplex } \sigma \in \tilde{T}\}$ an *ideal triangulation* of the punctured surface $S = \tilde{S} - V$. We call V ideal vertices (or cusps) of the surface S . If the Euler characteristic of S is negative, a *decorated hyperbolic metric* (d, r) on S ,

introduced by Penner [13], is a complete hyperbolic metric d of finite area on S so that each ideal vertex v is assigned a positive number r_v . Let $T_c(S)$ be the Teichmüller space of complete hyperbolic metrics with cusps ends on S . Then the decorated Teichmüller space introduced in [13] is $T_c(S) \times \mathbb{R}_{>0}^V$.




Figure 5: Simplicial coordinates. a is a generalized angle and $\frac{a}{2}$ is the length of the horocyclic arc.

Given a decorated hyperbolic metric $(d, r) \in T_c(S) \times \mathbb{R}_{>0}^V$, using the ideal triangulation T , Penner defined a map $\Psi: T_c(S) \times \mathbb{R}_{>0}^V \rightarrow \mathbb{R}^E$ as follows. Given a metric (d, r) , each edge $e \in E$ is isotopic to a complete geodesic e^* and each triangle σ in T is isotopic to an ideal triangle σ^* in the metric d . Since assigning a positive number r_v to each cusp v is the same as associating a horodisk B centered at the cusp so that the length of ∂B is r_v , we see that each ideal triangle σ^* is naturally a decorated ideal triangle, ie, a type $(0, 0, 0)$ generalized hyperbolic triangle. The value of $\Psi(d, r)$ at an edge $e \in E$, is

$$\Psi(d, r)(e) = \frac{b + c - a}{2} + \frac{b' + c' - a'}{2},$$

where a, a' are the generalized angles facing e^* , and b, b', c, c' are the generalized angles adjacent to e^* as labeled in Figure 5.

An edge cycle $(e_1, t_1, e_2, t_2, \dots, e_k, t_k)$ in a triangulation T is an alternating sequence of edges e_i and faces t_i in T so that adjacent faces t_i and t_{i+1} share the same edge e_i for any $i = 1, \dots, k$ and $t_{k+1} = t_1$.

A beautiful theorem proved by Penner is the following.

Theorem 1.1 (Penner [13]) Suppose (S, T) is an ideally triangulated surface of negative Euler characteristic. Then for any vector $z \in \mathbb{R}_{\geq 0}^E$ so that $\sum_{i=1}^k z(e_i) > 0$ for any edge cycle $(e_1, t_1, \dots, e_k, t_k)$, there exists a unique decorated complete hyperbolic metric (d, r) on S so that $\Psi(d, r) = z$.

Using the derivative cosine law associated to the decorated ideal triangle and the associated energy function, we generalize Penner's theorem to the following.

Theorem 1.2 Suppose (S, T) is an ideally triangulated surface of negative Euler characteristic. Then Penner's map $\Psi: T_c(S) \times \mathbb{R}_{>0}^V \rightarrow \mathbb{R}^E$ is a smooth embedding whose image is the convex polytope

$$P(T) = \{z \in \mathbb{R}^E \mid \sum_{i=1}^k z(e_i) > 0 \text{ for any edge cycle } (e_1, t_1, \dots, e_k, t_k)\}.$$

We remark that, from the definition, the set $P(T)$ is convex. It is in fact a convex polytope defined by the finite set of linear inequalities $\sum_{i=1}^k z(e_i) > 0$ for those edge cycles $(e_1, t_1, \dots, e_k, t_k)$ where each edge appears at most twice (see Luo [11] and Guo [6]).

Results similar to Penner's work have been established recently for hyperbolic cone metric by Leibon [8] and hyperbolic metric with geodesic boundary in [11]. In fact Leibon defined the coordinate $\psi(e) = \frac{b+c-a}{2} + \frac{b'+c'-a'}{2}$, where edge e is shared by two hyperbolic triangles, a, a' are the inner angles facing e , and b, b', c, c' are the inner angles adjacent to e . In [11], the coordinate is $\psi(e) = \frac{b+c-a}{2} + \frac{b'+c'-a'}{2}$, where edge e is shared by two hyperbolic right-angled hexagons, a, a' are the lengths of boundary arcs e , and b, b', c, c' are the lengths of boundary arcs adjacent to e .

In [9], a one-parameter family of coordinates depending on a parameter $h \in \mathbb{R}$ is introduced for hyperbolic cone metrics and hyperbolic metrics with geodesic boundary. These coordinates generalized the ones in [8] and [11]. In fact, Leibon's coordinate is deformed as

$$\psi_h(e) = \int_0^{\frac{b+c-a}{2}} \cosh^h(t) dt + \int_0^{\frac{b'+c'-a'}{2}} \cosh^h(t) dt.$$

The coordinate in [11] is deformed as

$$\psi_h(e) = \int_0^{\frac{a+b-c}{2}} \cosh^h(t) dt + \int_0^{\frac{a'+b'-c'}{2}} \cosh^h(t) dt.$$

These two cases can be written in one formula as

$$\psi_h(e) = \int_0^{\frac{a+b-c}{2}} \cosh^h(\sqrt{\varepsilon}t) dt + \int_0^{\frac{a'+b'-c'}{2}} \cosh^h(\sqrt{\varepsilon}t) dt.$$

Compact hyperbolic triangles correspond to $\varepsilon = 1$. Hyperbolic right-angled hexagons correspond to $\varepsilon = -1$. Ideal hyperbolic triangles correspond to $\varepsilon = 0$. These 1-parameter family of deformations are derived from the associated 1-parameter family of 2-dimensional Schläefli formulas for the associated generalized hyperbolic triangles. However, there is only one 2-dimensional Schläefli formula for decorated ideal triangles. This is the reason that Penner's coordinate cannot be deformed.

The relationship between the edge invariant in [11] and Penner's map Ψ was recently established by Mondello [12].

1.4 Thurston–Andreev's circle packing and its generalizations

Thurston's work on circle packing can be summarized as follows. Suppose (Σ, T) is a triangulated closed surface so that V , E , F are sets of all vertices, edges and triangles in T . Fix a map $\Phi: E \rightarrow [\frac{\pi}{2}, \pi]$. According to Thurston [17], a hyperbolic circle packing metric with intersection angles Φ is a function $r: V \rightarrow \mathbb{R}_{>0}$ so that the associated edge length function $l: E \rightarrow \mathbb{R}_{>0}$ is defined as follows. In Figure 6, consider a topological triangle with vertices v_i, v_j, v_k . One can construct a hyperbolic triangle




Figure 6: Thurston–Andreev circle packing

$\Delta v_i v_j q_k$ such that the edges $v_i q_k$, $v_j q_k$ have lengths $r(v_i)$, $r(v_j)$ respectively and the angle at q_k is $\Phi(v_i v_j)$. Let $l(v_i v_j)$ be the length of edge $v_i v_j$ in the hyperbolic triangle $\Delta v_i v_j q_k$, which is a function of $r(v_i)$, $r(v_j)$ and given $\Phi(v_i v_j)$ via the cosine law. Similarly, one obtains the edge lengths $l(v_j v_k)$, $l(v_k v_i)$.

Under the assumption that $\Phi: E \rightarrow [\frac{\pi}{2}, \pi]$, Thurston observed that lengths $l(v_i v_j)$, $l(v_j v_k)$ and $l(v_k v_i)$ satisfy the triangle inequality for each triangle $\Delta v_i v_j v_k$ in F . Thus there exists a hyperbolic polyhedral metric on (Σ, T) whose edge length function

is l . Let $K: V \rightarrow \mathbb{R}$ be the discrete curvature of the polyhedral metric, which sends a vertex to 2π less the sum of all inner angles at the vertex.

Theorem 1.3 (Thurston [17]) *For any closed triangulated surface (Σ, T) and any $\Phi: E \rightarrow [\frac{\pi}{2}, \pi]$, a hyperbolic circle packing metric on (Σ, T) is determined by its discrete curvature, ie, the map from r to K is injective. Furthermore, the set of all K 's is an open convex polytope in \mathbb{R}^V .*




Figure 7: Generalized circle packing

Since there are many other cosine laws available, we may try to use these other cosine laws for generalized triangles of type $(\varepsilon, \varepsilon, \delta)$ instead of type $(1, 1, 1)$ used by Thurston. To state our result, let us fix the notation once and for all. Let

(1-1)
$$I_\delta = \begin{cases} \mathbb{R}_{>0} & \text{if } \delta = 0, -1, \\ (0, \pi] & \text{if } \delta = 1. \end{cases} \quad \overset{\circ}{I}_\delta = \begin{cases} \mathbb{R}_{>0} & \text{if } \delta = 0, -1, \\ (0, \pi) & \text{if } \delta = 1, \end{cases}$$

(1-2)
$$J_\sigma = \begin{cases} \mathbb{R}_{>0} & \text{if } \sigma = 1, -1, \\ \mathbb{R} & \text{if } \sigma = 0. \end{cases}$$

A *generalized circle packing metric of type $(\varepsilon, \varepsilon, \delta)$* on a triangulated surface (Σ, T) with weight $\Phi: E \rightarrow I_\delta$ is given by a radius function $r: V \rightarrow J_{\varepsilon\delta}$ so that the edge length function $l: E \rightarrow J_\varepsilon$ is obtained from the radius r and weight Φ by the cosine law applied to the generalized triangle of type $(\varepsilon, \varepsilon, \delta)$. In Figure 7, consider a triangle with vertices v_i, v_j, v_k in the triangulation. One can construct a generalized hyperbolic triangle $\triangle v_i v_j q_k$ of type $(\varepsilon, \varepsilon, \delta)$ such that the edges $v_i q_k, v_j q_k$ have lengths $r(v_i), r(v_j)$ respectively and the generalized angle at q_k is $\Phi(v_i v_j)$. Let $l(v_i v_j)$ be the length of edge $v_i v_j$ in the generalized hyperbolic triangle $\triangle v_i v_j q_k$, which is a function of

$r(v_i)$, $r(v_j)$ and given $\Phi(v_i v_j)$ via the cosine law. Similarly, one obtains the edge lengths $l(v_j v_k)$, $l(v_k v_i)$.

Depending on $\varepsilon \in \{0, -1, 1\}$, the numbers $l(v_i v_j)$, $l(v_j v_k)$, $l(v_k v_i)$ may not be the three edge lengths of a type $(\varepsilon, \varepsilon, \varepsilon)$ triangle. Let $\mathcal{M}_{\varepsilon, \delta}(\Phi(v_i v_j), \Phi(v_j v_k), \Phi(v_k v_i))$ be the set of all $(r(v_i), r(v_j), r(v_k)) \in J_{\varepsilon, \delta}^3$ such that there exists a type $(\varepsilon, \varepsilon, \varepsilon)$ triangle $\Delta v_i v_j v_k$ with edge lengths $l(v_i v_j)$, $l(v_j v_k)$, $l(v_k v_i)$. Therefore r can only take values in $\mathcal{N}_{\varepsilon, \delta}(\Phi)$ a subspace of $(J_{\varepsilon, \delta})^V$, where $\mathcal{N}_{\varepsilon, \delta}(\Phi)$ is the set of all $r: V \rightarrow J_{\varepsilon, \delta}$ such that $(r(v_i), r(v_j), r(v_k)) \in \mathcal{M}_{\varepsilon, \delta}(\Phi(v_i v_j), \Phi(v_j v_k), \Phi(v_k v_i))$, if v_i, v_j, v_k are vertices of a triangle.

The edge length function $l: E \rightarrow J_\varepsilon$ produces a polyhedral metric on (Σ, T) . We define the *generalized discrete curvature* of the polyhedral metric to be $\tilde{K}: V \rightarrow \mathbb{R}_{>0}$ sending a vertex to the sum of all generalized angles at the vertex. We remark that the generalized discrete curvature and the discrete curvature in [Theorem 1.3](#) differ by a sign and a constant.

We show that Thurston's circle packing theorem can be generalized to the following six cases corresponding to the generalized triangles of type $(-1, -1, 1)$, $(-1, -1, -1)$, $(-1, -1, 0)$, $(0, 0, 1)$, $(0, 0, -1)$, $(0, 0, 0)$ in [Figure 8](#). More precisely, we list the six cases below.

- (a) For the case of $(-1, -1, 1)$, where $\Phi: E \rightarrow (0, \pi]$, the edge length $l(v_i v_j)$ is obtained from the radii $r(v_i)$, $r(v_j)$ by the cosine law for the hyperbolic pentagon as in [Figure 8](#) (a).
- (b) For the case of $(-1, -1, -1)$, where $\Phi: E \rightarrow (0, \infty)$, the edge length $l(v_i v_j)$ is obtained from the radii $r(v_i)$, $r(v_j)$ by the cosine law for the right-angled hexagon as in [Figure 8](#) (b).
- (c) For the case of $(-1, -1, 0)$, where $\Phi: E \rightarrow (0, \infty)$, the edge length $l(v_i v_j)$ is obtained from the radii $r(v_i)$, $r(v_j)$ by the cosine law for the hexagon as in [Figure 8](#) (c).
- (d) For the case of $(0, 0, 1)$, where $\Phi: E \rightarrow (0, \pi]$, the edge length $l(v_i v_j)$ is obtained from the radii $r(v_i)$, $r(v_j)$ by the cosine law for the pentagon as in [Figure 8](#) (d).
- (e) For the case of $(0, 0, -1)$, where $\Phi: E \rightarrow (0, \infty)$, the edge length $l(v_i v_j)$ is obtained from the radii $r(v_i)$, $r(v_j)$ by the cosine law for the hexagon as in [Figure 8](#) (e).
- (f) For the case of $(0, 0, 0)$, where $\Phi: E \rightarrow (0, \infty)$, the edge length $l(v_i v_j)$ is obtained from the radii $r(v_i)$, $r(v_j)$ by the cosine law for the hexagon as in [Figure 8](#) (f).




Figure 8: Symmetric generalized triangles for generalized circle packing

Theorem 1.4 *Given a closed triangulated surface (Σ, T) and $\Phi: E \rightarrow I_\delta$ in the above six cases, the generalized $(\epsilon, \epsilon, \delta)$ type circle packing metric $r \in \mathcal{N}_{\epsilon, \delta}(\Phi)$ is determined by its generalized discrete curvature $\tilde{K}: V \rightarrow \mathbb{R}_{>0}$. In particular, the map from r to \tilde{K} is a smooth embedding. Furthermore, the set of all \tilde{K} 's is the space $\mathbb{R}_{>0}^V$.*

Our method of proof of [Theorem 1.4](#) also produces a new variational proof of the rigidity of circle packing in Thurston's theorem ([Theorem 1.3](#)) similar to the proof in [\[3\]](#). However, unlike the proof in [\[3\]](#) which uses Thurston's geometric argument and Maple, our proof is a straight forward calculation. We are not able to establish [Theorem 1.4](#) for the remaining two cases of $(1, 1, -1)$, $(1, 1, 0)$ in [Figure 9](#).




Figure 9: Generalized circle packing does not work in the two cases.

Generalized circle patterns have been considered by many authors including Bobenko and Springborn [1], Schlenker [15], Stephenson [16] and others. In particular, Stephenson's question [16, page 331] about disjoint circle patterns motivates us to consider the generalized circle packing of type $(1, 1, -1)$.

Furthermore, similar to the work of [3] on discrete curvature flow, for the case of $\varepsilon = -1$ in Theorem 1.4, there exists a corresponding generalized curvature flow. Indeed, let $r_i := r(v_i)$ be the radii at vertex $v_i \in V$ and \tilde{K}_i be the generalized discrete curvature at vertex $v_i \in V$. The generalized curvature flow is

$$\frac{dr_i(t)}{dt} = -\tilde{K}_i \frac{1}{2} e^{r_i} - \frac{1}{2} \varepsilon \delta e^{-r_i}.$$

From the proof of Theorem 1.4 in Section 4, we obtain that, when $(\varepsilon, \varepsilon, \delta)$ is one of the six cases in Theorem 1.4, the flow is a negative gradient flow of a strictly concave down function after a change of variables.

1.5 Bobenko–Springborn's circle pattern and its generalizations

In [1], Bobenko and Springborn generalized Thurston's circle packing pattern in the case of $\Phi \equiv \pi$ in a different setting. The energy functional in [1] was derived from a discrete integrable system. Let us recall briefly the framework in [1]. See Figure 10 (a). Let (Σ, G) be a cellular decomposition of a closed surface with the set of vertices V , edges E and 2-cells F . The dual cellular decomposition G^* has the set of vertices $V^* (\cong F)$ so that each 2-cell f in F contains exactly one vertex f^* in G^* . If v is a vertex of a 2-cell f , we denote it by $v < f$. For all pairs (v, f) where $v < f$, join v to f^* by an arc in f , denoted by (v, f^*) , so that (v, f^*) and (v', f^*) don't intersect in their interior. Then these arcs $\bigcup_{(v, f)} (v, f^*)$ decompose the surface Σ into a union of quadrilaterals of the form (v, v', f^*, f'^*) where $vv' \in E$, $v < f$, $v' < f'$. According to [1], this quadrilateral decomposition of a closed surface arises naturally from integrable systems and discrete Riemann surfaces. Now suppose $\theta : E \rightarrow (0, \pi)$ is given. For $r : V^* \rightarrow \mathbb{R}_{>0}$, called a *circle pattern metric*, and a quadrilateral (v, v', f^*, f'^*) , construct an \mathbb{E}^2 (or \mathbb{H}^2) triangle $\triangle f^* f'^* v$ so that the length of the edges vf^* , $v'f'^*$ are given by $r(f^*)$, $r(f'^*)$ and the angle at v is $\theta(vv')$.

In this way the quadrilateral (v, v', f^*, f'^*) is realized in \mathbb{E}^2 (or \mathbb{H}^2) as the isometric double of the triangle $\triangle f^* f'^* v$ across the edge $f^* f'^*$. Since the surface Σ is a gluing of the quadrilateral (v, v', f^*, f'^*) along edges, by isometrically gluing these quadrilaterals, one obtains a polyhedral metric on the surface Σ with cone points at the vertices V and V^* . The cone angles at $v \in V$ are prescribed by θ , the only variable curvatures are at $f^* \in V^*$. Bobenko–Springborn's rigidity result says:




Figure 10: Bobenko–Springborn’s circle pattern

Theorem 1.5 (Bobenko–Springborn [1, Theorems 3 and 4]) *For any cell decomposition (Σ, G) of a closed surface and any map $\theta: E \rightarrow (0, \pi)$, the circle pattern metric $r: V^* \rightarrow \mathbb{R}_{>0}$ is determined by its discrete curvature $K: V^* \rightarrow \mathbb{R}$ for hyperbolic polyhedral metrics and is determined up to scaling by $K: V^* \rightarrow \mathbb{R}$ for Euclidean polyhedral metrics.*

We remark that Bobenko and Springborn [1] proved that the set of all discrete curvatures K forms a convex polytope. In the hyperbolic geometric setting, the essential part of Bobenko and Springborn’s construction is to produce the hyperbolic triangle $\triangle f^* f'^* v$ with two prescribed edge lengths and the prescribed angle between the two edges. Since there are eight other generalized hyperbolic triangles of type $(\varepsilon, \varepsilon, \delta)$ as listed in Figure 2, we can use them to produce hyperbolic metrics. Our result below shows that the rigidity phenomena still hold for the other eight cases.

We assume the same setting as in [1] that (Σ, G) is a cellular decomposed surface so that V, E, F are the sets of all vertices, edges and 2–cells with dual (Σ, G^*) . Suppose $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$, and a map $\theta: E \rightarrow I_\delta$ is given (see (1-1) for the definition of I_δ).

Assume for an $r \in (J_{\varepsilon\delta})^{V^*}$ and any quadrilateral (v, v', f^*, f'^*) , we can construct a type $(\varepsilon, \varepsilon, \delta)$ generalized hyperbolic triangle $\triangle f^* f'^* v$ so that the lengths of vf^*, vf'^* are $r(f^*), r(f'^*)$ respectively and the generalized angle at v is $\theta(vv')$. Now realize the quadrilateral (v, v', f^*, f'^*) as the metric double of $\triangle f^* f'^* v$ across the edge $f^* f'^*$. The generalized curvature of the resulting circle pattern is concentrated at the vertices V^* . It is defined as follows. For $h \in \mathbb{R}$, define the generalized curvature

$K_h\colon V^* \rightarrow \mathbb{R}$ by

(1-3)
$$K_h(f^*) = \sum_{i=1}^m 2 \int_1^{a_i} \rho_\varepsilon^h(t) \, dt$$

where the a_i ’s are the generalized angles at the vertex f^* in the triangle $\triangle f^* f'^* v$ and $\rho_\varepsilon(t) = \int_0^t \cos(\sqrt{\varepsilon}x) \, dx$ (see the definition in [Section 2](#)).

The following theorem is only considering the rigidity of the generalized Bobenko–Springborn circle pattern. We do not have a result addressing the existence, ie, describing the imaging of the map in the following theorem. In the last two subsections about the generalized Penner’s map and the generalized Thurston–Andreev circle packing, we established both of rigidity and existence.

Theorem 1.6 *Under the setup of generalized circle pattern above, for any $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$ and $\theta\colon E \rightarrow \overset{\circ}{I}_\delta$, the map from $(J_{\varepsilon\delta})^{V^*}$ to \mathbb{R}^{V^*} sending r to K_h is a smooth embedding.*

Bobenko and Springborn’s circle pattern theorem ([Theorem 1.5](#)) in the hyperbolic geometry corresponds to $(\varepsilon, \varepsilon, \delta) = (1, 1, 1)$ and $h = 0$ in [Theorem 1.6](#). Bobenko and Springborn [\[1\]](#) also showed that the image of $\{K\}$ is an explicit open convex polytope in \mathbb{R}^{V^*} . It is an interesting question to investigate the images of $\{K_h\}$ in the generalized setting.

We remark that there is a discrete curvature flow for each case in [Theorem 1.6](#). Let $r_i := r(f_i^*)$. The flow is defined by

$$\frac{dr_i(t)}{dt} = -K_h(f_i^*) \left(\frac{1}{2} e^{r_i} - \frac{1}{2} \varepsilon \delta e^{-r_i} \right)^{1-h}.$$

From the proof of [Theorem 1.6](#) in [Section 5](#), we obtain that, for any $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$, the flow is a negative gradient flow of a strictly concave down function.

Similar situations have been considered before by Hazel. In [\[7\]](#), he considered the flow for cases when $h = 0$ and $(\varepsilon, \delta, \varepsilon) = (1, 1, 1)$, $(1, 1, -1)$ or $(1, 1, 0)$.

Acknowledgment We would like to thank referee for his/her careful reading and very helpful suggestions and comments.

2 A proof of Theorem 1.2

We give a proof of Theorem 1.2 in this section. The proof consists of two parts. In the first part we show that Penner's map Ψ is an embedding. Then we determine its image. As in [11], the rigidity follows from the following well-known fact together with the cosine law for decorated ideal triangles.

Lemma 2.1 *If X is an open convex set in \mathbb{R}^n and $f: X \rightarrow \mathbb{R}$ is smooth strictly convex, then the gradient $\nabla f: X \rightarrow \mathbb{R}^n$ is injective. Furthermore, if the Hessian of f is positive definite for all $x \in X$, then ∇f is a smooth embedding.*

To begin, recall that (S, T) is an ideally triangulated surface with sets of edges, triangles and cusps given by E, F, V . We assume that $\chi(S) < 0$.

Following Penner [13], we will produce a smooth parametrization of the decorated Teichmüller space $T_c(S) \times \mathbb{R}_{>0}^V$ by \mathbb{R}^E using the edge lengths. From the derivative cosine law for decorated ideal triangles, we will construct a smooth strictly concave down function H on \mathbb{R}^E so that its gradient is Penner's map Ψ . Then by Lemma 2.1, the map $\Psi: T_c(S) \times \mathbb{R}_{>0}^V \rightarrow \mathbb{R}^E$ is an embedding. To determine the image $\Psi(T_c(S) \times \mathbb{R}_{>0}^V)$, we study the degenerations of decorated ideal triangles. The strategy of the proof is the same as that in [11].

2.1 Penner's length parametrization of $T_c(S) \times \mathbb{R}_{>0}^V$

For each decorated hyperbolic metric $(d, r) \in T_c(S) \times \mathbb{R}_{>0}^V$, where $r = (r_1, \dots, r_{|V|})$, one replaces each edge $e \in E$ by the geodesic e^* in the metric d and constructs for each cusp $v_i \in V$ a horocyclic disk $B_{r_i}(v_i)$ centered at v_i whose circumference $\partial B_{r_i}(v_i)$ has length $r_i = r(v_i)$. Now, the *length coordinate* $l_{d,r} \in \mathbb{R}^E$ of (d, r) is defined as follows. Given $r: V \rightarrow \mathbb{R}_{>0}$, realize each triangle Δuvw in T by a decorated ideal hyperbolic triangle with generalized angles at u, v, w being $r(u), r(v), r(w)$. Then $l_{d,r}(e)$ is the generalized edge length of the edge $e = uv$ in the triangle Δuvw . In this way, Penner defined a length map

$$\begin{aligned} L: T_c(S) \times \mathbb{R}_{>0}^V &\rightarrow \mathbb{R}^E \\ (d, r) &\rightarrow l_{d,r}. \end{aligned}$$

Lemma 2.2 (Penner [13]) *The length map $L: T_c(S) \times \mathbb{R}_{>0}^V \rightarrow \mathbb{R}^E$ is a diffeomorphism.*

Proof By the cosine law for decorated ideal triangle, the map L satisfies for all $\lambda \in \mathbb{R}_{>0}$

$$(2-1) \quad L(d, \lambda r) = L(d, r) - (2 \ln \lambda)(1, 1, \dots, 1).$$

Thus, it suffices to deal with those decorated metrics (d, r) so that r are small, ie, $L(d, r) \in \mathbb{R}_{>0}^E$. In this case, Penner proved that $L|: L^{-1}(\mathbb{R}_{>0}^E) \rightarrow \mathbb{R}_{>0}^E$ is a diffeomorphism by a direct geometric construction using isometric gluing of decorated ideal triangles. By (2-1), it follows that L is a diffeomorphism. \square

2.2 Penner's map Ψ is a coordinate

Recall that for a decorated ideal triangle Δ with edges e_1, e_2, e_3 of lengths l_1, l_2, l_3 and opposite generalized angles $\theta_1, \theta_2, \theta_3$, the cosine law obtained by Penner [13] says

$$\frac{e^{l_i}}{2} = \frac{2}{\theta_j \theta_k}, \quad \frac{\theta_i^2}{4} = e^{l_i - l_j - l_k}.$$

where $\{i, j, k\} = \{1, 2, 3\}$. The derivative cosine law expressing l_i in terms of $(\theta_1, \theta_2, \theta_3)$ says

$$\frac{\partial l_i}{\partial \theta_i} = 0, \quad \frac{\partial l_i}{\partial \theta_j} = -\frac{1}{\theta_j}.$$

Let $x_i = \frac{1}{2}(\theta_j + \theta_k - \theta_i)$ (or $\theta_i = x_j + x_k$). We call x_i the *radius invariant* at the edge e_i in the triangle Δ .

Using the derivative cosine law, we have:

Lemma 2.3 *Under the same assumption as above, the differential 1-form $\omega = \sum_{i=1}^3 x_i dl_i$ is closed in \mathbb{R}^3 and its integration $W(l) = \int_0^l \omega$ is strictly concave down in \mathbb{R}^3 . Furthermore,*

$$(2-2) \quad \frac{\partial W}{\partial l_i} = x_i.$$

Proof Consider the matrix $H = [\partial l_a / \partial x_b]_{3 \times 3}$. The closeness of ω is equivalent to that H is symmetric. The strictly concavity of W will be a consequence of the negative definiteness of H . We establish these two properties for H as follows. Assume that indices $\{i, j, k\} = \{1, 2, 3\}$. By definition, $\partial / \partial x_i = \partial / \partial \theta_j + \partial / \partial \theta_k$. It follows from the derivative cosine law that

$$\frac{\partial l_i}{\partial x_i} = -\left(\frac{1}{\theta_j} + \frac{1}{\theta_k}\right),$$

$$\text{and} \quad \frac{\partial l_i}{\partial x_j} = -\frac{1}{\theta_k},$$

which is symmetric in i, j . This shows that the matrix H is symmetric. Furthermore, the negative matrix $-H$ is of the form $[m_{ab}]_{3 \times 3}$ where $m_{ab} = m_{ba} > 0$ and $m_{ii} = m_{ij} + m_{ik}$. The determinant of such a matrix can be calculated easily as $4m_{11}m_{22}m_{33} > 0$, and the determinant of a principal 2×2 submatrix is $m_{ii}m_{jj} - m_{ij}^2 = (m_{ij} + m_{ik})(m_{ij} + m_{jk}) - m_{ij}^2 > 0$. It follows that $-H$ is positive definite. \square

By the construction in Section 2.1, it suffices to show that the composition $\widetilde{\Psi} = \Psi \circ L^{-1}: \mathbb{R}^E \rightarrow \mathbb{R}^E$ is a smooth embedding. The map $\widetilde{\Psi}$ is constructed explicitly as follows. For each $l \in \mathbb{R}^E$ and each triangle $\sigma \in F$ realize σ by an ideal hyperbolic triangle together with horocycles centered at three vertices so that the generalized edge length of an edge e in σ is $l(e)$. Now isometrically glue these ideal hyperbolic triangles along edges so that the horocycles match. The result is a complete finite area hyperbolic metric on the surface S together with a horocycle at each cusp. For each edge $e \in E$, the value $\widetilde{\Psi}(l)(e)$ is equal to $\frac{b+c-a}{2} + \frac{b'+c'-a'}{2}$ where a, a', b, b', c, c' are generalized angles facing and adjacent to e in Figure 5. Thus

(2-3)
$$\widetilde{\Psi}(l)(e) = r_f(e) + r_{f'}(e)$$

where f, f' are the decorated ideal triangles sharing the edge e and $r_f(e), r_{f'}(e)$ are the radius invariants at the edge e in f, f' respectively.

Given a vector $l \in \mathbb{R}^E$, define the energy $H(l)$ of l to be

$$H(l) = \sum_{\{i,j,k\} \in F} W(l(e_i), l(e_j), l(e_k))$$

where the sum is over all triangles $\{i, j, k\}$ in F with edges e_i, e_j, e_k . By definition and Lemma 2.3, $H: \mathbb{R}^E \rightarrow \mathbb{R}$ is smooth and strictly concave down when Hessian is negative definite. Furthermore, by (2-2) and (2-3),

$$\frac{\partial H}{\partial l(e_i)} = \widetilde{\Psi}(l)(e_i)$$

ie, $\nabla H = \widetilde{\Psi}$. It follows from Lemma 2.1, that $\widetilde{\Psi}: \mathbb{R}^E \rightarrow \mathbb{R}^E$ is a smooth embedding. Therefore Ψ is a smooth embedding.

2.3 The image of Penner’s map

Let Ω be the convex subset of \mathbb{R}^E consisting of all $z \in \mathbb{R}^E$ such that $\sum_{i=1}^p z(e_{n_i}) > 0$ whenever $(e_{n_1}, t_{n_1}, e_{n_2}, t_{n_2}, \dots, e_{n_p}, t_{n_p}, e_{n_1})$ is an edge cycle. To show that $\widetilde{\Psi}(\mathbb{R}^E) = \Omega$, due to convexity of Ω , it suffices to prove that $\widetilde{\Psi}(\mathbb{R}^E)$ is both open and closed in Ω .




Figure 11: Edge cycles

First to see that $\widetilde{\Psi}(\mathbb{R}^E) \subset \Omega$, take an edge cycle $(e_{n_1}, t_{n_1}, e_{n_2}, t_{n_2}, \dots, e_{n_p}, t_{n_p}, e_{n_1})$ as shown in Figure 11 and take $l \in \mathbb{R}^E$.

Let the generalized angles in the decorated ideal triangle t_{n_i} in the metric $L^{-1}(l)$ be a_i, b_i, c_i , where b_i faces the edge e_{n_i} , c_i faces the edge $e_{n_{i+1}}$ and a_i is adjacent to $e_{n_i}, e_{n_{i+1}}$. Then the contribution to $\sum_{j=1}^p \widetilde{\Psi}(l)(e_{n_j})$ from $e_{n_i}, e_{n_{i+1}}$ in triangle t_{n_i} is given by $(a_i + b_i - c_i)/2 + (a_i + c_i - b_i)/2 = a_i$. Thus

(2-4)
$$\sum_{j=1}^p \widetilde{\Psi}(l)(e_{n_j}) = \sum_{j=1}^p a_i > 0$$

due to $a_i > 0$ for all i . It follows that $\widetilde{\Psi}(\mathbb{R}^E)$ is open in Ω since $\widetilde{\Psi}: \mathbb{R}^E \rightarrow \mathbb{R}^E$ was just proved to be an embedding.

It remains to prove that $\widetilde{\Psi}(\mathbb{R}^E)$ is closed in Ω . The closeness of $\widetilde{\Psi}(\mathbb{R}^E)$ in Ω requires to show that if a sequence $\{l^{(m)} \in \mathbb{R}^E\}_{m=0}^\infty$ satisfies $\lim_{m \rightarrow \infty} \widetilde{\Psi}(l^{(m)}) = z \in \Omega$, then $\{l^{(m)}\}_{m=0}^\infty$ contains a subsequence converging to a point in \mathbb{R}^E . Given a decorated hyperbolic metric $l^{(m)} \in \mathbb{R}^E$ on (S, T) and a generalized angle θ , let $\theta^{(m)} \in \mathbb{R}^{3F}$ be the generalized angles of the decorated ideal triangles in (S, T) in the metric $l^{(m)}$. By taking a subsequence if necessary, we may assume that $\lim_{m \rightarrow \infty} \widetilde{\Psi}(l^{(m)})$ converges in $[-\infty, \infty]^E$ and that for each angle θ_i , the limit $\lim_{m \rightarrow \infty} \theta_i^{(m)}$ exists in $[0, \infty]$.

Lemma 2.4 For all i , $\lim_{m \rightarrow \infty} \theta_i^{(m)} \in [0, \infty)$.

Proof If otherwise, suppose that $\lim_{m \rightarrow \infty} \theta_1^{(m)} = \infty$ for some angle θ_1 . Let e_1, e_2 be the edges adjacent to the angle θ_1 in the triangle t . Take an edge cycle $(e_{n_1}, t_{n_1}, e_{n_2}, t_{n_2}, \dots, e_{n_p}, t_{n_p}, e_{n_1})$ which contains (e_1, t, e_2) as a part. Then by the calculation as in (2-4),

$$\sum_{i=1}^p z(e_{n_i}) = \lim_{m \rightarrow \infty} \sum_{i=1}^p \widetilde{\Psi}(l^{(m)})(e_{n_i}) \geq \lim_{m \rightarrow \infty} \theta_1^{(m)} = \infty.$$

This contradicts the assumption that $z \in \Omega$. □

Now, we finish the proof by contradiction as follows. If $\lim_{m \rightarrow \infty} l^{(m)}$ were not in \mathbb{R}^E , there would exist an edge $e \in E$ so that $\lim_{m \rightarrow \infty} l^{(m)}(e) = \pm \infty$. Let t be a triangle adjacent to e . Let $\theta_1^{(m)}, \theta_2^{(m)}$ be the generalized angles in t adjacent to e in the metric $l^{(m)}$. By the cosine law,

(2-5)
$$\exp(l^{(m)}(e)) = \frac{4}{\theta_1^{(m)} \theta_2^{(m)}}$$

and $\theta_1^{(m)}, \theta_2^{(m)} \in (0, \infty)$.

Case 1 If $\lim_{m \rightarrow \infty} l^{(m)}(e) = -\infty$, then $\lim_{m \rightarrow \infty} \exp(l^{(m)}(e)) = 0$. By (2-5), one of $\lim_{m \rightarrow \infty} \theta_i^{(m)}$ must be ∞ . But this contradicts Lemma 2.4.

Case 2 If $\lim_{m \rightarrow \infty} l^{(m)}(e) = \infty$, then $\lim_{m \rightarrow \infty} \exp(l^{(m)}(e)) = \infty$. Since Lemma 2.4 shows that $\theta_1^{(m)}$ and $\theta_2^{(m)}$ are bounded, by (2-5), one of the limits $\lim_{m \rightarrow \infty} \theta_i^{(m)}$ must be zero for $i = 1$ or 2 . Say $\lim_{m \rightarrow \infty} \theta_1^{(m)} = 0$. Let e_1 be the other edge in the triangle t so that e_1, e are adjacent to the generalized angle θ_1 . Let θ_3 be the third angle in t , facing e . Then by the cosine law that $\exp(l^{(m)}(e_1)) = 4/(\theta_1^{(m)} \theta_3^{(m)})$ and Lemma 2.4 on the boundedness of $\theta_3^{(m)}$, we conclude that $\lim_{m \rightarrow \infty} l^{(m)}(e_1) = \infty$. To summarize, from $\lim_{m \rightarrow \infty} l^{(m)}(e) = \infty$ and any triangle t adjacent to e , we conclude that there is an edge e_1 and an angle θ_1 adjacent to e, e_1 in t so that $\lim_{m \rightarrow \infty} l^{(m)}(e_1) = \infty$ and $\lim_{m \rightarrow \infty} \theta_1^{(m)} = 0$.

Applying this procedure to e_1 and the triangle t_1 adjacent to e_1 from the side other than the side that t lies. We obtain the next angle, say θ_2 and edge e_2 in t_1 so that $\lim_{m \rightarrow \infty} l^{(m)}(e_2) = \infty$ and $\lim_{m \rightarrow \infty} \theta_2^{(m)} = 0$. Since there are only finite number of edges and triangles, this procedure will produce an edge cycle $(e_{n_1}, t_{n_1}, e_{n_2}, t_{n_2}, \dots, e_{n_k}, t_{n_k}, e_{n_1})$ in T so that

- (i) $\lim_{m \rightarrow \infty} l^{(m)}(e_{n_i}) = \infty$ for each i ,
- (ii) $\lim_{m \rightarrow \infty} \theta_i^{(m)} = 0$ for each i , where θ_i is the angle in triangle t_{n_i} adjacent to e_{n_i} and $e_{n_{i+1}}$.

By (2-4),

$$\sum_{i=1}^k z(e_{n_i}) = \lim_{m \rightarrow \infty} \sum_{i=1}^k \widetilde{\Psi}(l^{(m)})(e_{n_i}) = \lim_{m \rightarrow \infty} \sum_{i=1}^k \theta_i^{(m)} = 0.$$

This contradicts the assumption that $z \in \Omega$.

3 The derivative cosine law

We give a unified approach to all cosine laws and sine laws in this section. The derivatives of the cosine laws are also determined. Most of the proofs are straightforward checking and will be delayed to [Appendix B](#).

Assume that a generalized hyperbolic triangle of type $(\varepsilon_1, \varepsilon_2, \varepsilon_3) \in \{-1, 0, 1\}^3$ has generalized angles $\theta_1, \theta_2, \theta_3$ and opposite generalized edge lengths l_1, l_2, l_3 . There are ten different types of generalized hyperbolic triangles as shown in [Figure 2](#). The relationships between l_i 's and θ_i 's are expressed in the cosine law and the sine law. To state them, we introduce the two functions ρ_ε, τ_s depending on $\varepsilon, \theta, s, l \in \mathbb{R}$:

$$(3-1) \quad \rho_\varepsilon(\theta) = \int_0^\theta \cos(\sqrt{\varepsilon}x) dx = \frac{1}{\sqrt{\varepsilon}} \sin(\sqrt{\varepsilon}x),$$

$$(3-2) \quad \tau_s(l) = \frac{1}{2}e^l - \frac{1}{2}se^{-l}.$$

To be more explicit,

$$\begin{aligned} \rho_1(\theta) &= \sin(\theta), & \rho_0(\theta) &= \theta, & \rho_{-1}(\theta) &= \sinh(\theta), \\ \tau_1(l) &= \sinh(l), & \tau_0(l) &= \frac{1}{2}e^l, & \tau_{-1}(l) &= \cosh(l). \end{aligned}$$

A simple calculation shows

$$\begin{aligned} \rho'_\varepsilon(\theta) &:= \frac{\partial \rho_\varepsilon(\theta)}{\partial \theta} = \cos(\sqrt{\varepsilon}\theta), \\ \tau'_s(l) &:= \frac{\partial \tau_s(l)}{\partial l} = \frac{1}{2}e^l + \frac{1}{2}se^{-l}. \end{aligned}$$

Lemma 3.1 (The cosine laws and the sine laws) *For a generalized hyperbolic triangle of type $(\varepsilon_1, \varepsilon_2, \varepsilon_3) \in \{-1, 0, 1\}^3$ with generalized angles $\theta_1, \theta_2, \theta_3$ and opposite generalized edge lengths l_1, l_2, l_3 , for $\{i, j, k\} = \{1, 2, 3\}$, the following hold:*

$$(3-3) \quad \tau'_{\varepsilon_j \varepsilon_k}(l_i) = \frac{\rho'_{\varepsilon_i}(\theta_i) + \rho'_{\varepsilon_j}(\theta_j)\rho'_{\varepsilon_k}(\theta_k)}{\rho_{\varepsilon_j}(\theta_j)\rho_{\varepsilon_k}(\theta_k)},$$

$$(3-4) \quad 2\rho_{\varepsilon_i}^2\left(\frac{\theta_i}{2}\right) = \frac{\tau'_{\varepsilon_j \varepsilon_k}(l_i) - \frac{1}{2}\varepsilon_j e^{l_j-l_k} - \frac{1}{2}\varepsilon_k e^{l_k-l_j}}{\tau_{\varepsilon_k \varepsilon_i}(l_j)\tau_{\varepsilon_i \varepsilon_j}(l_k)},$$

$$(3-5) \quad \rho'_{\varepsilon_i}(\theta_i) = \frac{-\varepsilon_i \tau'_{\varepsilon_j \varepsilon_k}(l_i) + \tau'_{\varepsilon_k \varepsilon_i}(l_j)\tau'_{\varepsilon_i \varepsilon_j}(l_k)}{\tau_{\varepsilon_k \varepsilon_i}(l_j)\tau_{\varepsilon_i \varepsilon_j}(l_k)},$$

$$(3-6) \quad \frac{\rho_{\varepsilon_i}(\theta_i)}{\tau_{\varepsilon_j \varepsilon_k}(l_i)} = \frac{\rho_{\varepsilon_j}(\theta_j)}{\tau_{\varepsilon_k \varepsilon_i}(l_j)}.$$

Proof In [Appendix A](#), the cosine law and the sine law for each type of the ten generalized hyperbolic triangles are listed. We check directly that all the formulas there fit the uniform formulas (3-3)–(3-6). \square

Remark 3.2 The identity (3-6) is called the *sine law*. The formula (3-4) is stronger than the formula (3-5). They are equivalent if $\varepsilon_i = \pm 1$. If $\varepsilon_i = 0$, the formula (3-5) is trivial while the formula (3-4) expresses θ in term of lengths l .

For the six cases of generalized triangle without ideal vertices, there is a unified strategy to derive the cosine law (3-3) and (3-5) by using the hyperboloid model. This unified strategy is essentially given in [18, pages 74–82]. But this method does not work for the four cases of generalized triangle with ideal vertices.

The concepts of Gram matrix and angle Gram matrix of a generalized hyperbolic triangle are defined as follows. For a generalized hyperbolic triangle of type $(\varepsilon_1, \varepsilon_2, \varepsilon_3) \in \{-1, 0, 1\}^3$ with generalized angles $\theta_1, \theta_2, \theta_3$ and opposite generalized edge lengths l_1, l_2, l_3 , its Gram matrix is

$$G_l := - \begin{pmatrix} \varepsilon_1 & \tau'_{\varepsilon_1 \varepsilon_2}(l_3) & \tau'_{\varepsilon_3 \varepsilon_1}(l_2) \\ \tau'_{\varepsilon_1 \varepsilon_2}(l_3) & \varepsilon_2 & \tau'_{\varepsilon_2 \varepsilon_3}(l_1) \\ \tau'_{\varepsilon_3 \varepsilon_1}(l_2) & \tau'_{\varepsilon_2 \varepsilon_3}(l_1) & \varepsilon_3 \end{pmatrix}$$

and its angle Gram matrix is

$$G_\theta := - \begin{pmatrix} -1 & \rho'_{\varepsilon_3}(\theta_3) & \rho'_{\varepsilon_2}(\theta_2) \\ \rho'_{\varepsilon_3}(\theta_3) & -1 & \rho'_{\varepsilon_1}(\theta_1) \\ \rho'_{\varepsilon_2}(\theta_2) & \rho'_{\varepsilon_1}(\theta_1) & -1 \end{pmatrix}.$$

Lemma 3.3 *The determinants of G_l and G_θ satisfy the following:*

$$(3-7) \quad \det G_l = -(\tau_{\varepsilon_k \varepsilon_i}(l_j) \tau_{\varepsilon_i \varepsilon_j}(l_k) \rho_{\varepsilon_i}(\theta_i))^2$$

$$(3-8) \quad \det G_\theta = -(\rho_{\varepsilon_j}(\theta_j) \rho_{\varepsilon_k}(\theta_k) \tau_{\varepsilon_j \varepsilon_k}(l_i))^2$$

For a hyperbolic triangle, [Lemma 3.3](#) is well-known. We will prove it for a generalized triangle in [Appendix B](#).

By [Lemma 3.3](#) and the sine laws (3-6), we see

$$\begin{aligned} M &:= \frac{1}{\sqrt{-\det G_l}} \begin{pmatrix} \tau_{\varepsilon_2 \varepsilon_3}(l_1) & 0 & 0 \\ 0 & \tau_{\varepsilon_3 \varepsilon_1}(l_2) & 0 \\ 0 & 0 & \tau_{\varepsilon_1 \varepsilon_2}(l_3) \end{pmatrix} \\ &= \frac{1}{\sqrt{-\det G_\theta}} \begin{pmatrix} \rho_{\varepsilon_1}(\theta_1) & 0 & 0 \\ 0 & \rho_{\varepsilon_2}(\theta_2) & 0 \\ 0 & 0 & \rho_{\varepsilon_3}(\theta_3) \end{pmatrix}. \end{aligned}$$

Lemma 3.4 $M G_l M G_\theta = I$.

This lemma is a consequence of the cosine laws and the sine laws (3-3), (3-5) and (3-6). It is checked by direct calculation.

Let y_1, y_2, y_3 be three functions of variables x_1, x_2, x_3 . Let $A = (\frac{\partial y_i}{\partial x_j})_{3 \times 3}$ be the Jacobi matrix. Then the differentials dy_1, dy_2, dy_3 and dx_1, dx_2, dx_3 satisfy

$$\begin{pmatrix} dy_1 \\ dy_2 \\ dy_3 \end{pmatrix} = A \begin{pmatrix} dx_1 \\ dx_2 \\ dx_3 \end{pmatrix}.$$

Lemma 3.5 (The derivative cosine law) *For a generalized hyperbolic triangle of type $(\varepsilon_1, \varepsilon_2, \varepsilon_3) \in \{-1, 0, 1\}^3$ with generalized angles $\theta_1, \theta_2, \theta_3$ and opposite generalized edge lengths l_1, l_2, l_3 , the differentials of l 's and θ 's satisfy the following relations:*

$$(3-9) \quad \begin{pmatrix} dl_1 \\ dl_2 \\ dl_3 \end{pmatrix} = M G_l \begin{pmatrix} d\theta_1 \\ d\theta_2 \\ d\theta_3 \end{pmatrix},$$

$$(3-10) \quad \begin{pmatrix} d\theta_1 \\ d\theta_2 \\ d\theta_3 \end{pmatrix} = M G_\theta \begin{pmatrix} dl_1 \\ dl_2 \\ dl_3 \end{pmatrix}.$$

We will prove [Lemma 3.5](#) in [Appendix B](#).

In the rest of the section, we establish the existence of a generalized hyperbolic triangle of type $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$ with two given edge lengths l_1, l_2 and a generalized angle θ between them, where the generalized angles opposite to l_1, l_2 have type ε and θ has type δ . Recall $\overset{\circ}{I}_\delta$ and $J_{\varepsilon\delta}$ are in (1-1) and (1-2).

Fix type $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$. For a given $\theta \in \overset{\circ}{I}_\delta$, let's introduce the set $\mathcal{D}_{\varepsilon,\delta}(\theta) = \{(l_1, l_2) \in (J_{\varepsilon\delta})^2 \mid \text{there exists a generalized hyperbolic triangle of type } (\varepsilon, \varepsilon, \delta) \text{ with two edge lengths } l_1, l_2 \text{ so that the generalized angle between them is } \theta \text{ of type } \delta\}$.

Lemma 3.6 (1) If $\varepsilon = 1$ or 0 , then $\mathcal{D}_{\varepsilon,\delta}(\theta) = (J_{\varepsilon\delta})^2$.

(2) If $(\varepsilon, \delta) = (-1, 1)$, then

$$\mathcal{D}_{\varepsilon,\delta}(\theta) = \{(l_1, l_2) \in \mathbb{R}_{>0}^2 \mid \sinh l_1 \sinh l_2 - \cos \theta \cosh l_1 \cosh l_2 > 1\}.$$

(3) If $(\varepsilon, \delta) = (-1, 0)$, then

$$\mathcal{D}_{\varepsilon,\delta}(\theta) = \{(l_1, l_2) \in \mathbb{R}^2 \mid \theta > e^{-l_1} + e^{-l_2}\}.$$

(4) If $(\varepsilon, \delta) = (-1, -1)$, then

$$\mathcal{D}_{\varepsilon,\delta}(\theta) = \{(l_1, l_2) \in \mathbb{R}_{>0}^2 \mid \cosh \theta \sinh l_1 \sinh l_2 - \cosh l_1 \cosh l_2 > 1\}.$$

Proof We will construct a generalized hyperbolic triangle of type $(\varepsilon, \varepsilon, \delta)$ in each case as follows.

Denote by \mathbb{H}^2 the hyperbolic plane and by $\partial\mathbb{H}^2$ the ideal boundary of \mathbb{H}^2 .

If $(\varepsilon, \varepsilon, \delta) = (1, 1, 1)$, choose a point $O \in \mathbb{H}^2$. Draw two geodesics rays L_1, L_2 starting from the point O such that the angle between L_1, L_2 is $\theta \in (0, \pi)$. Let P_i be the point on L_i such that the length of the segment OP_i is $l_i > 0, i = 1, 2$. Then one obtains the hyperbolic triangle by joining P_1, P_2 by a geodesic segment.

If $(\varepsilon, \varepsilon, \delta) = (1, 1, 0)$, choose a point $O \in \partial\mathbb{H}^2$ and draw a horocycle centered at O . Let H_1, H_2 be two points on the horocycle such that the length of the horocyclic arc H_1H_2 is $\frac{\theta}{2} > 0$. For $i = 1, 2$, draw a geodesic L_i passing through O and H_i . Let P_i be the point on L_i such that the length of the segment of H_iP_i is $|l_i|$ and H_i is between O, P_i on L_i if and only if $l_i > 0$. Then one obtains the generalized hyperbolic triangle by joining P_1, P_2 by a geodesic segment.

If $(\varepsilon, \varepsilon, \delta) = (1, 1, -1)$, draw a geodesic segment G_1G_2 of length $\theta > 0$ with end points G_1, G_2 . For $i = 1, 2$, let L_i be a geodesic ray starting from G_i perpendicular to G_1G_2 and L_1, L_2 are in the same half plane bounded by the geodesic containing G_1G_2 . Let P_i be the point on L_i such that the length of the segment G_iP_i is $l_i > 0, i = 1, 2$. Then one obtains the generalized hyperbolic triangle by joining P_1, P_2 by a geodesic segment.

If $(\varepsilon, \varepsilon, \delta) = (0, 0, 1)$, choose a point $O \in \mathbb{H}^2$. Draw two geodesics rays L_1, L_2 starting from the point O such that the angle between L_1, L_2 is $\theta \in (0, \pi)$ and L_i ends at point $P_i \in \partial\mathbb{H}^2$. Join P_1, P_2 by a geodesic. Let H_i be the point on the geodesic containing L_i such the length of OH_i is $|l_i|$ and H_i is between O, P_i if and only if $l_i > 0$. Draw a horocycle centered at P_i passing through H_i . One obtains the generalized triangle OP_1P_2 decorated by two horocycles.

If $(\varepsilon, \varepsilon, \delta) = (0, 0, 0)$ or $(0, 0, -1)$, the construction is similar to the case of $(0, 0, 1)$ above.

If $(\varepsilon, \varepsilon, \delta) = (-1, -1, 1)$, choose a point $O \in \mathbb{H}^2$. Draw two geodesics rays L_1, L_2 starting from the point O such that the angle between L_1, L_2 is $\theta \in (0, \pi)$. For $i = 1, 2$, let P_i be the point on L_i such that the length of the segment OP_i is $l_i > 0$. Let M_i be the geodesic passing through P_i perpendicular to L_i . There is a generalized hyperbolic triangle with prescribed lengths l_1, l_2 and angle θ if and only if the distance between M_1, M_2 is positive. The distance l between M_1, M_2 can be calculated from the cosine law of generalized triangle of type $(-1, -1, 1)$:

$$\cosh l = \sinh l_1 \sinh l_2 - \cos \theta \cosh l_1 \cosh l_2.$$

This shows that the condition in part (2) is equivalent to $l > 0$.

If $(\varepsilon, \varepsilon, \delta) = (-1, -1, 0)$, choose a point $O \in \partial\mathbb{H}^2$ and draw a horocycle centered at O . Let H_1, H_2 be the two points on the horocycle such that the length of the horocyclic arc H_1H_2 is $\frac{\theta}{2} > 0$. For $i = 1, 2$, draw a geodesic L_i passing through O and H_i . Let P_i be the point on L_i such that the length of the segment H_iP_i is $|l_i|$ and H_i is between O, P_i if and only if $l_i > 0$. Let M_i be the geodesic passing through P_i perpendicular to L_i . There is a generalized hyperbolic triangle with prescribed lengths l_1, l_2 and generalized angle θ if and only if the distance l between M_1, M_2 is positive. The distance l between M_1, M_2 can be calculated from the cosine law of generalized triangle of type $(-1, -1, 0)$:

$$\cosh l = \frac{1}{2}\theta^2 e^{l_1+l_2} - \cosh(l_1 - l_2).$$

Now, one sees $\frac{1}{2}\theta^2 e^{l_1+l_2} - \cosh(l_1 - l_2) > 1$ if and only if $\theta > e^{-l_1} + e^{-l_2}$. This shows part (3) holds.

If $(\varepsilon, \varepsilon, \delta) = (-1, -1, -1)$, draw a geodesic segment G_1G_2 of length $\theta > 0$ with end points G_1, G_2 . For $i = 1, 2$, let L_i be a geodesic half line starting from G_i perpendicular to G_1G_2 and L_1, L_2 are in the same half plane bounded by the geodesic containing G_1G_2 . Let P_i be the point on L_i such that the length of the segment G_iP_i is $l_i > 0, i = 1, 2$. Let M_i be the geodesic passing through P_i perpendicular to L_i .

There is a right-angled hexagon with prescribed lengths l_1, l_2, θ if and only if the distance l between M_1, M_2 is positive. The distance l between M_1, M_2 can be calculated from the cosine law of a right-angled hexagon:

$$\cosh l = \cosh \theta \sinh l_1 \sinh l_2 - \cosh l_1 \cosh l_2.$$

This shows that the condition in part (4) is equivalent to $l > 0$. □

4 A proof of Theorem 1.4

We give a proof of the generalized circle packing theorem (Theorem 1.4) in this section.

Recall that (Σ, T) is a closed triangulated surface with V, E, F the sets all vertices, edges and triangles in T . Given a type $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$ and $\Phi: E \rightarrow I_\delta$, for each $r \in \mathcal{N}_{\varepsilon, \delta}(\Phi)$, a generalized $(\varepsilon, \varepsilon, \delta)$ circle packing on (Σ, T) is based on the following local construction.

As in Figure 12, consider a topological triangle with vertices v_i, v_j, v_k . One can construct a generalized hyperbolic triangle $\Delta v_i v_j q_k$ of type $(\varepsilon, \varepsilon, \delta)$ such that the edges $v_i q_k, v_j q_k$ have lengths $r(v_i) := r_i, r(v_j) := r_j$ respectively and the generalized angle at q_k is $\Phi(v_i v_j) =: \phi_k$. Let $l(v_i v_j) =: l_k$ be the length of edge $v_i v_j$ in the generalized hyperbolic triangle $\Delta v_i v_j q_k$ which is a function of r_i, r_j . Similarly, one obtains the edge lengths $l(v_j v_k) =: l_i, l(v_k v_i) =: l_j$. Let θ_i be the generalized angle of $\Delta v_i v_j v_k$ at the vertex v_i . With fixed (ϕ_i, ϕ_j, ϕ_k) , we consider θ_i as a function of (r_i, r_j, r_k) .




Figure 12: Data for generalized circle packing

Let’s recall the definition of $\mathcal{M}_{\varepsilon, \delta}(\Phi(v_i v_j), \Phi(v_j v_k), \Phi(v_k v_i))$ in Section 1.4 and definition of $\mathcal{D}_{\varepsilon, \delta}(\theta)$ in Section 3.

Lemma 4.1 When $\varepsilon = 0$ or -1 , we have

$$\mathcal{M}_{\varepsilon,\delta}(\phi_1, \phi_2, \phi_3) = \{(r_1, r_2, r_3) \in (J_{\varepsilon\delta})^3 \mid (r_i, r_j) \in \mathcal{D}_{\varepsilon,\delta}(\phi_k), \{i, j, k\} = \{1, 2, 3\}\}.$$

Proof Given $\phi_k \in \overset{\circ}{I}_\delta$, by Lemma 3.6, for $r_i, r_j \in \mathcal{D}_{\varepsilon,\delta}(\phi_k)$, there is generalized hyperbolic triangle $\triangle v_i v_j q_k$ of type $(\varepsilon, \varepsilon, \delta)$ with edge lengths r_i, r_j and angle ϕ_k of type δ between the two edges. We obtain the edge length $l_k = l(v_i v_j)$. There is a case which is not contained in Lemma 3.6: $\delta = 1, \phi_k = \pi$. It is easier since we have $l_k = r_i + r_j$.

For $\varepsilon = 0$, we get $l_i, l_j, l_k \in \mathbb{R}$. Thus there exists a decorated ideal triangle with three edge lengths l_i, l_j, l_k . For $\varepsilon = -1$, the inequality defining $\mathcal{D}_{-1,\delta}(\phi_k)$ guarantees that $l_i, l_j, l_k > 0$. Thus there is a right-angled hexagon with three edge lengths l_i, l_j, l_k . \square

4.1 A proof of Theorem 1.4 for $\varepsilon = 0$

We assume that indices i, j, k are distinct in this section. First by Lemma 3.6 and Lemma 4.1, we have $\mathcal{M}_{0,\delta}(\phi_i, \phi_j, \phi_k) = \mathbb{R}^3$. Therefore $\mathcal{N}_{0,\delta}(\Phi) = \mathbb{R}^V$.

The case $\varepsilon = 0$ is very simple. Indeed, for a fixed $\Phi: E \rightarrow I_\delta$, there exists a map $C: V \rightarrow \mathbb{R}_{>0}$ so that for all $r \in \mathcal{N}_{0,\delta}(\Phi) = \mathbb{R}^V$,

$$(4-1) \quad \tilde{K}(r)(v) = C(v)e^{-r(v)},$$

ie, the discrete curvature $\tilde{K}(r)$ is uniformly proportional to $e^{-r(v)}$. Thus, one sees easily that the generalized circle packing metric $r \in \mathbb{R}^V$ is determined by its generalized discrete curvature $\tilde{K}: V \rightarrow \mathbb{R}$ and $\{\tilde{K}(v) \mid v \in V\} = \mathbb{R}_{>0}^V$.

Indeed, in Figure 12, the generalized triangle $\triangle v_i v_j q_k$ has type $(0, 0, \delta)$, edge lengths r_i, r_j, l_k and inner angle ϕ_k opposite to l_k . By the cosine law (3-4), we have

$$2\rho_\delta^2\left(\frac{\phi_k}{2}\right) = \frac{\tau'_0(l_k)}{\tau_0(r_i)\tau_0(r_j)} = 2e^{l_k - r_i - r_j}.$$

Thus

$$(4-2) \quad e^{l_k} = \rho_\delta^2\left(\frac{\phi_k}{2}\right)e^{r_i + r_j}.$$

By the cosine law (3-4) for the decorated ideal triangle $\triangle v_i v_j v_k$ (or type $(0,0,0)$ generalized triangle) we have

$$(4-3) \quad \frac{\theta_k^2}{4} = e^{l_k - l_i - l_j}.$$

Combining (4-2) and (4-3) we obtain

$$\frac{\theta_k^2}{4} = e^{l_k - l_i - l_j} = \frac{\rho_\delta^2(\frac{\phi_k}{2})e^{r_i + r_j}}{\rho_\delta^2(\frac{\phi_i}{2})e^{r_j + r_k} \rho_\delta^2(\frac{\phi_j}{2})e^{r_k + r_i}} = \frac{\rho_\delta^2(\frac{\phi_k}{2})}{\rho_\delta^2(\frac{\phi_i}{2})\rho_\delta^2(\frac{\phi_j}{2})}e^{-2r_k},$$

hence

$$(4-4) \quad \theta_k = \frac{2\rho_\delta(\frac{\phi_k}{2})}{\rho_\delta(\frac{\phi_i}{2})\rho_\delta(\frac{\phi_j}{2})}e^{-r_k}.$$

Summing up (4-4) for all triangle having v_k as a vertex, we obtain (4-1) where $C(v)$ depends explicitly on δ .

4.2 Rigidity of circle packing for $\varepsilon = -1$ or $(\varepsilon, \delta) = (1, 1)$

In this subsection we prove Theorem 1.4 for the case $\varepsilon = -1$ and give a new proof of Thurston's rigidity Theorem 1.3.

In the variational framework, the natural parameter is $u = (u_1, u_2, u_3)$ where

$$(4-5) \quad u_i = - \int_{r_i}^{\infty} \frac{1}{\tau_{\varepsilon\delta}(t)} dt$$

for $\varepsilon = \pm 1$. Note that by the definition of τ_s , $\partial u_i / \partial r_i > 0$.

Lemma 4.2 Fix $\phi = (\phi_1, \phi_2, \phi_3) \in I_\delta^3$, the set $u(\mathcal{M}_{\varepsilon,\delta}(\phi_1, \phi_2, \phi_3))$ is an open convex polyhedron in \mathbb{R}^3 in the following cases:

- (i) $\varepsilon = -1$, or
- (ii) $(\varepsilon, \delta) = (1, 1)$ and $\phi_i \in [\frac{\pi}{2}, \pi]$, $i = 1, 2, 3$.

Proof (i) When $\varepsilon = -1$, we have figured out the set $\mathcal{M}_{\varepsilon,\delta}(\phi_1, \phi_2, \phi_3)$ in Lemma 4.1.

If the type of ϕ_k is $\delta = 1$, by the proof of Lemma 3.6 (2), we need $l_k > 0$ which is the same as

$$(4-6) \quad \cos \phi_k = \frac{-\cosh l_k + \sinh r_i \sinh r_j}{\cosh r_i \cosh r_j} < \frac{-1 + \sinh r_i \sinh r_j}{\cosh r_i \cosh r_j}.$$

In this case $u_i = - \int_{r_i}^{\infty} 1/\cosh t dt = 2 \arctan e^{r_i} - \pi \in (-\frac{\pi}{2}, 0)$. Then

$$(4-7) \quad \sinh r_i = -\frac{1}{\tan u_i}, \quad \cosh r_i = -\frac{1}{\sin u_i}.$$

By substituting (4-7) into (4-6), we obtain $\cos \phi_k < \cos(u_i + u_j)$. Since $\phi_k \in (0, \pi]$, $-u_i - u_j \in (0, \pi)$, we see that $l_k > 0$ if and only if $u_i + u_j > -\phi_k$. Then $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3)) = \{(u_1, u_2, u_3) \in \mathbb{R}_{<0}^3 \mid u_i + u_j > -\phi_k, \{i, j, k\} = \{1, 2, 3\}\}$. The last description shows that $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$ is a convex polyhedron.

If the type of ϕ_k is $\delta = 0$, we need $l_k > 0$ which is equivalent to

$$\frac{\phi_k^2}{2} = \frac{\cosh l_k + \cosh(r_i - r_j)}{e^{r_i + r_j}} > \frac{1 + \cosh(r_i - r_j)}{e^{r_i + r_j}} = \frac{1}{2}(e^{-r_i} + e^{-r_j})^2.$$

Since in this case $u_i = -\int_{r_i}^{\infty} 1/e^t dt = -e^{-r_i} < 0$, we see that $l_k > 0$ is the same as having $u_i + u_j > -\phi_k$. Thus $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3)) = \{(u_1, u_2, u_3) \in \mathbb{R}_{<0}^3 \mid l_i > 0 \text{ for all } i\} = \{(u_1, u_2, u_3) \in \mathbb{R}_{<0}^3 \mid u_i + u_j > -\phi_k, \{i, j, k\} = \{1, 2, 3\}\}$. The last description shows that $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$ is a convex polyhedron.

If the type of ϕ_k is $\delta = -1$, we need $l_k > 0$ which is the same as

$$\begin{aligned} \cosh \phi_k &= \frac{\cosh l_k + \cosh r_i \cosh r_j}{\sinh r_i \sinh r_j} \\ &> \frac{1 + \cosh r_i \cosh r_j}{\sinh r_i \sinh r_j} \\ &= \frac{1}{2} \left(\tanh \frac{r_i}{2} \tanh \frac{r_j}{2} + \frac{1}{\tanh \frac{r_i}{2} \tanh \frac{r_j}{2}} \right). \end{aligned}$$

Hence $l_k > 0$ if and only if

$$e^{-\phi_k} < \tanh \frac{r_i}{2} \tanh \frac{r_j}{2}.$$

It follows that $l_k > 0$ is the same as

$$\ln \tanh \frac{r_i}{2} + \ln \tanh \frac{r_j}{2} > -\phi_k.$$

Since in this case $u_i = -\int_{r_i}^{\infty} 1/\sinh t dt = \ln \tanh \frac{r_i}{2} < 0$, we see that $l_k > 0$ if and only if $u_i + u_j > -\phi_k$. Thus $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3)) = \{(u_1, u_2, u_3) \in \mathbb{R}_{<0}^3 \mid u_i + u_j > -\phi_k, \{i, j, k\} = \{1, 2, 3\}\}$. The last description shows that $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$ is a convex polyhedron.

(ii) When $\varepsilon = 1$, $\delta = 1$ and $\phi_k \in [\frac{\pi}{2}, \pi]$, Thurston observed $\mathcal{M}_{1,1}(\phi_1, \phi_2, \phi_3) = \mathbb{R}_{>0}^3$. Indeed, $\phi_i \in [\frac{\pi}{2}, \pi]$ implies $l_i > r_j$. Similarly, $l_j > r_i$. See Figure 12. Hence $l_i + l_j > r_j + r_i > l_k$. This shows that l_1, l_2, l_3 are the lengths of a hyperbolic triangle. In this case $u_i = -\int_{r_i}^{\infty} 1/\sinh t dt = \ln \tanh \frac{r_i}{2} < 0$. Thus $u(\mathcal{M}_{1,1}(\phi_1, \phi_2, \phi_3)) = \mathbb{R}_{<0}^3$. \square

Lemma 4.3 For $\varepsilon = \pm 1$ and fixed $(\phi_1, \phi_2, \phi_3) \in I_{\delta}^3$, the Jacobi matrix A of the function $(\theta_1, \theta_2, \theta_3) = (\theta_1(u), \theta_2(u), \theta_3(u))$: $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3)) \rightarrow \mathbb{R}^3$ is symmetric.

Proof Consider the $(\varepsilon, \varepsilon, \delta)$ -triangle $\Delta v_i v_j q_k$ in Figure 12 with edge lengths r_i, r_j, l_k and opposite generalized angles a_{ji}, a_{ij}, ϕ_k . By applying the derivative cosine law (3-10) to $\Delta v_i v_j q_k$, the third row of the matrix in (3-10) is

$$d\phi_k = \frac{-\rho_\delta(\phi_k)}{\sqrt{-\det G_\theta}} (\rho'_\varepsilon(a_{ji})dr_i + \rho'_\varepsilon(a_{ij})dr_j - dl_k).$$

Since ϕ_k is fixed, then $d\phi_k = 0$. We have $dl_k = \rho'_\varepsilon(a_{ji})dr_i + \rho'_\varepsilon(a_{ij})dr_j$. For $\varepsilon = \pm 1$, we have $u_i = -\int_{r_i}^\infty 1/\tau_{\varepsilon\delta}(t) dt$. Then $dr_i = \tau_{\varepsilon\delta}(r_i)du_i$. Thus

$$\begin{aligned} \begin{pmatrix} dl_1 \\ dl_2 \\ dl_3 \end{pmatrix} &= \begin{pmatrix} 0 & \rho'_\varepsilon(a_{23}) & \rho'_\varepsilon(a_{32}) \\ \rho'_\varepsilon(a_{13}) & 0 & \rho'_\varepsilon(a_{31}) \\ \rho'_\varepsilon(a_{12}) & \rho'_\varepsilon(a_{21}) & 0 \end{pmatrix} \begin{pmatrix} dr_1 \\ dr_2 \\ dr_3 \end{pmatrix} \\ (4-8) \quad &= \begin{pmatrix} 0 & \rho'_\varepsilon(a_{23}) & \rho'_\varepsilon(a_{32}) \\ \rho'_\varepsilon(a_{13}) & 0 & \rho'_\varepsilon(a_{31}) \\ \rho'_\varepsilon(a_{12}) & \rho'_\varepsilon(a_{21}) & 0 \end{pmatrix} \begin{pmatrix} \tau_{\varepsilon\delta}(r_1) & 0 & 0 \\ 0 & \tau_{\varepsilon\delta}(r_2) & 0 \\ 0 & 0 & \tau_{\varepsilon\delta}(r_3) \end{pmatrix} \begin{pmatrix} du_1 \\ du_2 \\ du_3 \end{pmatrix}. \end{aligned}$$

For the $(\varepsilon, \varepsilon, \varepsilon)$ -triangle $\Delta v_1 v_2 v_3$ with angles $\theta_1, \theta_2, \theta_3$ and edge lengths l_1, l_2, l_3 , by the derivative cosine law (3-9) and (4-8), we have

$$\begin{aligned} \begin{pmatrix} d\theta_1 \\ d\theta_2 \\ d\theta_3 \end{pmatrix} &= \frac{-1}{\sqrt{-\det G_l}} \begin{pmatrix} \tau_1(l_1) & 0 & 0 \\ 0 & \tau_1(l_2) & 0 \\ 0 & 0 & \tau_1(l_3) \end{pmatrix} \begin{pmatrix} -1 & \rho'_\varepsilon(\theta_3) & \rho'_\varepsilon(\theta_2) \\ \rho'_\varepsilon(\theta_3) & -1 & \rho'_\varepsilon(\theta_1) \\ \rho'_\varepsilon(\theta_2) & \rho'_\varepsilon(\theta_1) & -1 \end{pmatrix} \\ &\quad \begin{pmatrix} 0 & \rho'_\varepsilon(a_{23}) & \rho'_\varepsilon(a_{32}) \\ \rho'_\varepsilon(a_{13}) & 0 & \rho'_\varepsilon(a_{31}) \\ \rho'_\varepsilon(a_{12}) & \rho'_\varepsilon(a_{21}) & 0 \end{pmatrix} \begin{pmatrix} \tau_{\varepsilon\delta}(r_1) & 0 & 0 \\ 0 & \tau_{\varepsilon\delta}(r_2) & 0 \\ 0 & 0 & \tau_{\varepsilon\delta}(r_3) \end{pmatrix} \begin{pmatrix} du_1 \\ du_2 \\ du_3 \end{pmatrix} \\ &=: \frac{-1}{\sqrt{-\det G_l}} N \begin{pmatrix} du_1 \\ du_2 \\ du_3 \end{pmatrix}. \end{aligned}$$

To show the Jacobi matrix A of $(\theta_1(u), \theta_2(u), \theta_3(u))$ is symmetric, we only need to check that $N = (N_{ij})$ is symmetric. In fact

$$N_{ij} = \tau_1(l_i)\tau_{\varepsilon\delta}(r_j)(-\rho'_\varepsilon(a_{jk}) + \rho'_\varepsilon(\theta_j)\rho'_\varepsilon(a_{ji})).$$

By the cosine law (3-5),

$$\begin{aligned} N_{ij} = \tau_1(l_i)\tau_{\varepsilon\delta}(r_j) &\left(-\frac{\varepsilon\tau'_{\varepsilon\delta}(r_k) + \tau'_{\varepsilon\delta}(r_j)\tau'_1(l_i)}{\tau_{\varepsilon\delta}(r_j)\tau_1(l_i)} \right. \\ &\quad \left. + \frac{\varepsilon\tau'_1(l_j) + \tau'_1(l_i)\tau'_1(l_k)}{\tau_1(l_i)\tau_1(l_k)} \frac{\varepsilon\tau'_{\varepsilon\delta}(r_i) + \tau'_{\varepsilon\delta}(r_j)\tau'_1(l_k)}{\tau_{\varepsilon\delta}(r_j)\tau_1(l_k)} \right) \end{aligned}$$

$$= \frac{\varepsilon \tau'_1(l_k)(-\tau_1^2(l_k) + \tau'_{\varepsilon\delta}(r_j)\tau'_1(l_j) + \tau'_{\varepsilon\delta}(r_i)\tau'_1(l_i))}{\tau_1^2(l_k)} + \frac{(\tau_1'^2(l_k) - \tau_1^2(l_k))\tau'_{\varepsilon\delta}(r_j)\tau'_1(l_i) + \varepsilon^2\tau'_{\varepsilon\delta}(r_i)\tau'_1(l_j)}{\tau_1^2(l_k)}.$$

Since $\varepsilon = \pm 1$, $\tau_1(l_k) = \frac{1}{2}e^{l_k} - \frac{1}{2}e^{-l_k} = \sinh l_k$, we have $\tau_1'^2(l_k) - \tau_1^2(l_k) = 1 = \varepsilon^2$. Therefore N_{ij} is symmetric in the index i, j . \square

Lemma 4.4 Given $(\phi_1, \phi_2, \phi_3) \in I_\delta^3$, the Jacobi matrix A of

$$(\theta_1, \theta_2, \theta_3) = (\theta_1(u), \theta_2(u), \theta_3(u)): u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3)) \rightarrow \mathbb{R}^3$$

is negative definite in the following cases:

- (i) $\varepsilon = -1$ or
- (ii) (Colin de Verdière [4], Chow-Luo [3]) $(\varepsilon, \delta) = (1, 1)$ and $\phi_i \in [\frac{\pi}{2}, \pi]$, $i = 1, 2, 3$.

We remark that the case (ii) for $\phi_i = \pi$ was first proved by Colin de Verdière [4], and was proved for $\phi_i \in [\frac{\pi}{2}, \pi]$ in [3] using the Maple program. Our proof of (ii) is new.

Proof In the proof of Lemma 4.3, the Jacobi matrix $A = (-1/\sqrt{-\det G_I})N$. Hence it is sufficient to show N is positive definite. First $\det N > 0$. Indeed N is a product of four matrixes. The first one and forth one are diagonal matrixes with positive determinant. By Lemma 3.3, the determinant of the second matrix (negative of an angle Gram matrix) is positive. The determinant of the third matrix is $\rho'_\varepsilon(a_{12})\rho'_\varepsilon(a_{23})\rho'_\varepsilon(a_{31}) + \rho'_\varepsilon(a_{32})\rho'_\varepsilon(a_{21})\rho'_\varepsilon(a_{13}) > 0$. If $\varepsilon = -1$, the above inequality is obvious. If $\varepsilon = 1$, for $\{i, j, k\} = \{1, 2, 3\}$, ϕ_k, a_{ij}, a_{ji} are inner angles of a hyperbolic triangle as in Figure 12. By the assumption $\phi_k \in [\frac{\pi}{2}, \pi]$, we must have $a_{ij}(0, \frac{\pi}{2})$. Therefore $\rho'_1(a_{ij}) = \cos(a_{ij}) > 0$. Thus the above inequality holds.

Since the set $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$ is connected and the $\det N \neq 0$, to show N is positive definite, we only need to check N is positive definite at one point in $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$. We choose the point such that $r_1 = r_2 = r_3 = r$.

Now in Figure 12 and Figure 13, in the $(\varepsilon, \varepsilon, \delta)$ -triangle $\Delta v_i v_j q_k$ with edge lengths r, r, l_k and opposite generalized angles $a_{ij} = a_{ji}, \phi_k$, let h be the geodesic length realizing the distance between the edge $v_i v_j$ and the generalized vertex q_k . In the $(\varepsilon, 1, \delta)$ -triangle $\Delta v_i R q_k$, where $R q_k$ is perpendicular to $v_i v_j$, by the cosine law (3-5), we have

$$\cos \frac{\pi}{2} = \frac{-\tau'_{\varepsilon\delta}(r) + \tau'_\delta(h)\tau'_\varepsilon(\frac{l_k}{2})}{\tau_\delta(h)\tau_\varepsilon(\frac{l_k}{2})}.$$




Figure 13: Data for generalized circle packing

Thus $\tau'_\delta(h) = \tau'_{\varepsilon\delta}(r)/\tau'_\varepsilon(\frac{l_k}{2})$. Also in $\triangle v_i R q_k$, by the cosine law (3-5), we have

$$\rho'_\varepsilon(a_{ij}) = \frac{-\varepsilon\tau'_\delta(h) + \tau'_\varepsilon(\frac{l_k}{2})\tau'_{\varepsilon\delta}(r)}{\tau_\varepsilon(\frac{l_k}{2})\tau_{\varepsilon\delta}(r)} = \frac{\tau'_{\varepsilon\delta}(r)(-\varepsilon + \tau'^2_\varepsilon(\frac{l_k}{2}))}{\tau'_\varepsilon(\frac{l_k}{2})\tau(\frac{l_k}{2})\tau_{\varepsilon\delta}(r)}.$$

Since

$$-\varepsilon + \tau'^2_\varepsilon(l) = -\varepsilon + \left(\frac{1}{2}e^l + \frac{1}{2}\varepsilon e^{-l}\right)^2 = \left(\frac{1}{2}e^l - \frac{1}{2}\varepsilon e^{-l}\right)^2 = \tau^2_\varepsilon(l),$$

we have

(4-9)

$$\rho'_\varepsilon(a_{ij}) = \frac{\tau'_{\varepsilon\delta}(r)\tau_\varepsilon(\frac{l_k}{2})}{\tau_{\varepsilon\delta}(r)\tau'_\varepsilon(\frac{l_k}{2})}.$$

By substituting (4-9) into N , let

$$N_1 = \begin{pmatrix} -1 & \rho'_\varepsilon(\theta_3) & \rho'_\varepsilon(\theta_2) \\ \rho'_\varepsilon(\theta_3) & -1 & \rho'_\varepsilon(\theta_1) \\ \rho'_\varepsilon(\theta_2) & \rho'_\varepsilon(\theta_1) & -1 \end{pmatrix} \begin{pmatrix} 0 & \tau_\varepsilon(\frac{l_1}{2})/\tau'_\varepsilon(\frac{l_1}{2}) & \tau_\varepsilon(\frac{l_1}{2})/\tau'_\varepsilon(\frac{l_1}{2}) \\ \tau_\varepsilon(\frac{l_2}{2})/\tau'_\varepsilon(\frac{l_2}{2}) & 0 & \tau_\varepsilon(\frac{l_2}{2})/\tau'_\varepsilon(\frac{l_2}{2}) \\ \tau_\varepsilon(\frac{l_3}{2})/\tau'_\varepsilon(\frac{l_3}{2}) & \tau_\varepsilon(\frac{l_3}{2})/\tau'_\varepsilon(\frac{l_3}{2}) & 0 \end{pmatrix}.$$

Then N , N_1 differ by left and right multiplication by diagonal matrices of positive diagonal entries. We will show that the determinants of the 1×1 and 2×2 principal submatrices of N_1 are positive. This implies that the determinants of the 1×1 and 2×2 principal submatrices of N are positive. Thus N is positive definite.

Indeed, the determinant of the 1×1 principal submatrix of N_1 is

(4-10)

$$\rho'_\varepsilon(\theta_3)\frac{\tau_\varepsilon(\frac{l_2}{2})}{\tau'_\varepsilon(\frac{l_2}{2})} + \rho'_\varepsilon(\theta_2)\frac{\tau_\varepsilon(\frac{l_3}{2})}{\tau'_\varepsilon(\frac{l_3}{2})}.$$

The determinant of the 2×2 principal matrix of N_1 is

$$(4-11) \quad (\rho'_\varepsilon(\theta_3) - 1) \frac{\tau_\varepsilon(\frac{l_1}{2})}{\tau'_\varepsilon(\frac{l_1}{2})} \frac{\tau_\varepsilon(\frac{l_2}{2})}{\tau'_\varepsilon(\frac{l_2}{2})} + (\rho'_\varepsilon(\theta_2) + \rho'_\varepsilon(\theta_1)\rho'_\varepsilon(\theta_3)) \frac{\tau_\varepsilon(\frac{l_2}{2})}{\tau'_\varepsilon(\frac{l_2}{2})} \frac{\tau_\varepsilon(\frac{l_3}{2})}{\tau'_\varepsilon(\frac{l_3}{2})} \\ + (\rho'_\varepsilon(\theta_1) + \rho'_\varepsilon(\theta_2)\rho'_\varepsilon(\theta_3)) \frac{\tau_\varepsilon(\frac{l_1}{2})}{\tau'_\varepsilon(\frac{l_1}{2})} \frac{\tau_\varepsilon(\frac{l_3}{2})}{\tau'_\varepsilon(\frac{l_3}{2})}.$$

(i) If $\varepsilon = -1$, then $\rho'_\varepsilon(\theta_i) = \cosh \theta_i$. Each term in (4-10) and (4-11) is positive. Hence (4-10) and (4-11) are positive.

(ii) If $\varepsilon = 1$, then $\rho'_\varepsilon(\theta_i) = \cos \theta_i$. We see that the expression in (4-10) is positive is the same as

$$(4-12) \quad \cos \theta_3 \tanh \frac{l_2}{2} + \cos \theta_2 \tanh \frac{l_3}{2} > 0.$$

By the cosine law, (4-12) is equivalent to

$$(4-13) \quad \frac{-\cosh l_3 + \cosh l_1 \cosh l_2}{\sinh l_1 \sinh l_2} \frac{\sinh l_2}{1 + \cosh l_2} \\ + \frac{-\cosh l_2 + \cosh l_1 \cosh l_3}{\sinh l_1 \sinh l_3} \frac{\sinh l_3}{1 + \cosh l_3} > 0.$$

By calculation, (4-13) is equivalent to

$$(4-14) \quad \cosh l_1 > \frac{\cosh^2 l_2 + \cosh^2 l_3 + \cosh l_2 + \cosh l_3}{2 \cosh l_2 \cosh l_3 + \cosh l_2 + \cosh l_3}.$$

Since $l_1 > |l_2 - l_3|$, therefore $\cosh l_1 > \cosh(l_2 - l_3)$. To show (4-14) holds, it is enough to check

$$(4-15) \quad \cosh l_2 \cosh l_3 - \sinh l_2 \sinh l_3 \geq \frac{\cosh^2 l_2 + \cosh^2 l_3 + \cosh l_2 + \cosh l_3}{2 \cosh l_2 \cosh l_3 + \cosh l_2 + \cosh l_3}.$$

We simplify the notation by introducing $a := \cosh l_2 > 1$, $b := \cosh l_3 > 1$. Then (4-15) is rewritten as

$$(4-16) \quad ab - \sqrt{(a^2 - 1)(b^2 - 1)} \geq \frac{a^2 + b^2 + a + b}{2ab + a + b}.$$

(4-16) is equivalent to

$$(4-17) \quad ab - \frac{a^2 + b^2 + a + b}{2ab + a + b} \geq \sqrt{(a^2 - 1)(b^2 - 1)}.$$

Since $a > 1$, $b > 1$, the left hand side of (4-17) is positive. To show (4-17) holds, we square the two sides and simplify. We have

$$\begin{aligned} & (ab^4 + a^4b - a^3b^2 - a^2b^3) + (a^4 + b^4 - 2a^2b^2) + (a^3 + b^3 - ab^2 - a^2b) \geq 0 \\ \Leftrightarrow & (ab + 1)(a^3 + b^3 - a^2b - ab^2) + (a^2 - b^2)^2 \geq 0 \\ \Leftrightarrow & (ab + 1)(a + b)(a - b)^2 + (a^2 - b^2)^2 \geq 0. \end{aligned}$$

This shows that the expression in (4-10) is positive.

We see that the expression in (4-11) is positive is the same as

$$\begin{aligned} (4-18) \quad & -\sin^2 \theta_3 \tanh \frac{l_1}{2} \tanh \frac{l_2}{2} + (\cos \theta_2 + \cos \theta_1 \cos \theta_3) \tanh \frac{l_2}{2} \tanh \frac{l_3}{2} \\ & + (\cos \theta_1 + \cos \theta_2 \cos \theta_3) \tanh \frac{l_1}{2} \tanh \frac{l_3}{2} > 0. \end{aligned}$$

By the cosine law, (4-18) is equivalent to

$$\begin{aligned} (4-19) \quad & -\sin^2 \theta_3 \tanh \frac{l_1}{2} \tanh \frac{l_2}{2} + \cosh l_2 \sin \theta_1 \sin \theta_3 \tanh \frac{l_2}{2} \tanh \frac{l_3}{2} \\ & + \cosh l_1 \sin \theta_2 \sin \theta_3 \tanh \frac{l_1}{2} \tanh \frac{l_3}{2} > 0. \end{aligned}$$

By the sine law, (4-19) is equivalent to

$$\begin{aligned} & \left(\cosh l_1 \sinh l_2 \tanh \frac{l_1}{2} + \cosh l_2 \sinh l_1 \tanh \frac{l_2}{2} \right) \tanh \frac{l_3}{2} > \sinh l_3 \tanh \frac{l_1}{2} \tanh \frac{l_2}{2} \\ \Leftrightarrow & \frac{\cosh l_1 \sinh l_2}{\tanh \frac{l_2}{2}} + \frac{\cosh l_2 \sinh l_1}{\tanh \frac{l_1}{2}} > \frac{\sinh l_3}{\tanh \frac{l_3}{2}} \\ \Leftrightarrow & \cosh l_1 (1 + \cosh l_2) + \cosh l_2 (1 + \cosh l_1) > 1 + \cosh l_3 \\ \Leftrightarrow & (\cosh l_1 + \cosh l_2 - 1) + (2 \cosh l_1 \cosh l_2 - \cosh l_3) > 0. \end{aligned}$$

This is true since $\cosh l_1 + \cosh l_2 > 1$ and $2 \cosh l_1 \cosh l_2 > \cosh l_1 \cosh l_2 + \sinh l_1 \sinh l_2 = \cosh(l_1 + l_2) > \cosh l_3$. This shows that the expression in (4-11) is positive. \square

As a corollary of Lemma 4.3 and Lemma 4.4, we obtain the following result.

Corollary 4.5 Given $(\phi_1, \phi_2, \phi_3) \in I_\delta^3$, the differential 1-form $\sum_{i=1}^3 \theta_i du_i$ is closed in $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$. Furthermore, its integration $w(u) = \int^u \sum_{i=1}^3 \theta_i du_i$ is a strictly concave function defined in $u(\mathcal{M}_{\varepsilon, \delta}(\phi_1, \phi_2, \phi_3))$ if either

- (i) $\varepsilon = -1$ or
- (ii) $(\varepsilon, \delta) = (1, 1)$ and $\phi_i \in [\frac{\pi}{2}, \pi]$, $i = 1, 2, 3$.

Furthermore

$$(4-20) \quad \frac{\partial w}{\partial u_i} = \theta_i.$$

Let's prove the rigidity of circle packing for $\varepsilon = -1$, or $(\varepsilon, \delta) = (1, 1)$ and $\Phi: E \rightarrow [\frac{\pi}{2}, \pi]$. Fix $\Phi: E \rightarrow I_\delta$. For each $r \in \mathcal{N}_{\varepsilon, \delta}(\Phi)$, define the function u in terms of r as in (4-5) by

$$u(v) = - \int_{r(v)}^{\infty} \frac{1}{\tau_{\varepsilon \delta}(t)} dt.$$

The set of all values of u is $u(\mathcal{N}_{\varepsilon, \delta}(\Phi))$ which is an open convex set in \mathbb{R}^V due to Lemma 4.2. To be more precise, let us label vertices $V = \{v_1, \dots, v_n\}$ and denote by $\{i, j, k\} \in F$ a generalized triangle in F with vertices v_i, v_j, v_k . Let $u_i = u(v_i)$. Then $u(\mathcal{N}_{\varepsilon, \delta}(\Phi)) = \{u \in \mathbb{R}^V \mid (u_i, u_j, u_k) \in u(\mathcal{M}_{\varepsilon, \delta}(\phi_i, \phi_j, \phi_k)) \text{ if } \{i, j, k\} \in F\}$ is convex since it is the intersection of the convex set $\prod_{\{i, j, k\} \in F} u(\mathcal{M}_{\varepsilon, \delta}(\phi_i, \phi_j, \phi_k))$ with affine spaces.

We now use the function in Corollary 4.5 to introduce a function $W: u(\mathcal{N}_{\varepsilon, \delta}(\Phi)) \rightarrow \mathbb{R}$ by defining

$$W(u) = \sum_{\{i, j, k\} \in F} w(u_i, u_j, u_k)$$

where the sum is over all triangles in F with vertices $\{v_i, v_j, v_k\}$. By Corollary 4.5, W is smooth and strictly concave down in $u(\mathcal{N}_{\varepsilon, \delta}(\Phi))$ so that

$$\frac{\partial W}{\partial u_i} = \tilde{K}(v_i)$$

by (4-20) and the definition of \tilde{K} .

By Lemma 2.1, the map $\nabla W: u(\mathcal{N}_{\varepsilon, \delta}(\Phi)) \rightarrow \mathbb{R}^V$ is a smooth embedding. Thus the map from $\{r \in \mathcal{N}_{\varepsilon, \delta}(\Phi)\}$ to $\{\tilde{K} \in \mathbb{R}_{>0}^V\}$ is a smooth injective map.

4.3 The image of \tilde{K} for $\varepsilon = -1$

To prove that the image $X = \{\tilde{K}(v) \mid v \in V\} = \mathbb{R}_{>0}^V$ for $\varepsilon = -1$, we will show that X is both open and closed in $\mathbb{R}_{>0}^V$. By definition, $X \subset \mathbb{R}_{>0}^V$ and X is open due to the injectivity of the map from $\{r \in \mathcal{N}_{-1, \delta}(\Phi)\}$ to $\mathbb{R}_{>0}^V$. It remains to prove that X is closed in $\mathbb{R}_{>0}^V$. To this end, let us first establish the following lemma.

Lemma 4.6 *Let $\varepsilon = -1$ and (ϕ_i, ϕ_j, ϕ_k) be given so that $(\theta_i, \theta_j, \theta_k) = (\theta_i(r_i, r_j, r_k), \theta_j(r_i, r_j, r_k), \theta_k(r_i, r_j, r_k))$ are considered as functions of $r_i, r_j, r_k \in J_{-\delta}$. Then*

$$\lim_{r_k \rightarrow \infty} \theta_k(r_i, r_j, r_k) = 0$$

and the convergence is uniform.

Proof We only need to check that for constants $a, b, c \in J_{-\delta}$ the following holds.

- (1) If $\lim r_i = a, \lim r_j = b, \lim r_k = \infty$, then $\lim \theta_k = 0$.
- (2) If $\lim r_i = c, \lim r_j = \infty, \lim r_k = \infty$, then $\lim \theta_k = 0$.
- (3) If $\lim r_i = \infty, \lim r_j = \infty, \lim r_k = \infty$, then $\lim \theta_k = 0$.

The strategy of the proof is the same for all three cases. First, in $\triangle P_j P_k Q_i$, $\triangle P_k P_i Q_j$, $\triangle P_i P_j Q_k$ of type $(-1, -1, \delta)$, by the cosine law (3-4), the lengths l_i, l_j, l_k can be written as functions of r_i, r_j, r_k :

$$\begin{aligned}\cosh l_i &= 2\rho_\delta^2 \left(\frac{\phi_i}{2} \right) \tau_{-\delta}(r_j) \tau_{-\delta}(r_k) - \frac{1}{2} (e^{r_j - r_k} + e^{r_k - r_j}), \\ \cosh l_j &= 2\rho_\delta^2 \left(\frac{\phi_j}{2} \right) \tau_{-\delta}(r_k) \tau_{-\delta}(r_i) - \frac{1}{2} (e^{r_k - r_i} + e^{r_i - r_k}), \\ \cosh l_k &= 2\rho_\delta^2 \left(\frac{\phi_k}{2} \right) \tau_{-\delta}(r_i) \tau_{-\delta}(r_j) - \frac{1}{2} (e^{r_i - r_j} + e^{r_j - r_i}).\end{aligned}$$

When the limits of r_i, r_j, r_k are given, we can find the limits of l_i, l_j, l_k . Then in $\triangle P_i P_j P_k$ (of type $(-1, -1, -1)$), by the cosine law (3-5), write θ_k as a function of l_i, l_j, l_k :

$$\cosh \theta_k = \frac{\cosh l_k + \cosh l_i \cosh l_j}{\sinh l_i \sinh l_j}.$$

Then we find the limit of θ_k .

- (1) If $\lim r_i = a, \lim r_j = b, \lim r_k = \infty$, then

$$\begin{aligned}\lim \cosh l_i &= \lim \left(2\rho_\delta^2 \left(\frac{\phi_i}{2} \right) \tau_{-\delta}(b) \frac{1}{2} e^{r_k} - \frac{1}{2} e^{r_k - b} \right) \\ &= \lim \frac{1}{2} e^{r_k} \left(2\rho_\delta^2 \left(\frac{\phi_i}{2} \right) \tau_{-\delta}(b) - 1 \right).\end{aligned}$$

Since $\cosh l_i > 0$, we have $2\rho_\delta^2 \left(\frac{\phi_i}{2} \right) \tau_{-\delta}(b) - 1 > 0$. When $\lim r_k = \infty$, we have $\lim \cosh l_i = \infty$.

By symmetry $\lim \cosh l_j = \infty$. Furthermore $\lim \cosh l_k$ is finite. Therefore $\lim l_i = \lim l_j = \infty, \lim l_k$ is finite. Hence

$$\lim \cosh \theta_k = \lim \frac{\cosh l_k}{\sinh l_i \sinh l_j} + 1 = 1.$$

Therefore $\lim \theta_k = 0$.

(2) If $\lim r_i = c$, $\lim r_j = \infty$, $\lim r_k = \infty$, then

$$\begin{aligned}\lim \cosh l_i &= \lim \left(2\rho_\delta^2 \left(\frac{\phi_i}{2} \right) \frac{1}{4} e^{r_j+r_k} - \frac{1}{2} (e^{r_j-r_k} + e^{r_k-r_j}) \right) \\ &= \lim \frac{1}{2} e^{r_j+r_k} \left(\rho_\delta^2 \left(\frac{\phi_i}{2} \right) - e^{-2r_k} - e^{-2r_j} \right) \\ &= \lim \frac{1}{2} e^{r_j+r_k} \rho_\delta^2 \left(\frac{\phi_i}{2} \right) \\ &= \infty, \\ \lim \cosh l_j &= \lim \left(2\rho_\delta^2 \left(\frac{\phi_j}{2} \right) \tau_{-\delta}(c) \frac{1}{2} e^{r_k} - \frac{1}{2} e^{r_k-c} \right) \\ &= \lim e^{r_k} \left(\rho_\delta^2 \left(\frac{\phi_j}{2} \right) \tau_{-\delta}(c) - \frac{1}{2} e^{-c} \right) \\ &= \infty.\end{aligned}$$

Here we use the same argument as in (1).

By the same calculation of $\lim \cosh l_j$, we see that $\lim \cosh l_k = \lim e^{r_j} c_k = \infty$ for some constant c_k . Hence

$$\begin{aligned}\lim \cosh \theta_k &= \lim \frac{\cosh l_k}{\sinh l_i \sinh l_j} + 1 \\ &= \lim \frac{\cosh l_k}{\cosh l_i \cosh l_j} + 1 \\ &= \lim \frac{e^{r_j} c_k}{e^{r_k} c_j e^{r_j+r_k} c_i} + 1 \\ &= 1.\end{aligned}$$

Therefore $\lim \theta_k = 0$.

(3) If $\lim r_i = \infty$, $\lim r_j = \infty$, $\lim r_k = \infty$, by the same calculation of $\lim \cosh l_i$ in (2), we see that

$$\lim \cosh l_k = \lim e^{r_i+r_j} a_k, \quad \lim \cosh l_j = \lim e^{r_k+r_i} a_j, \quad \lim \cosh l_i = \lim e^{r_j+r_k} a_i$$

for some constants

$$\begin{aligned}\lim \cosh \theta_k &= \lim \frac{\cosh l_k}{\sinh l_i \sinh l_j} + 1 = \lim \frac{\cosh l_k}{\cosh l_i \cosh l_j} + 1 \\ &= \lim \frac{e^{r_i+r_j} a_k}{e^{r_k+r_i} a_j e^{r_j+r_k} a_i} + 1 = 1.\end{aligned}$$

Therefore $\lim \theta_k = 0$. □

To show that X is closed in $\mathbb{R}_{>0}^V$, take a sequence of radius $r^{(m)}$ in $\mathcal{N}_{\Phi,-1}$ such that $\lim_{m \rightarrow \infty} \tilde{K}^{(m)} \in \mathbb{R}_{>0}^V$. To prove closedness, it is sufficient to show that there is a subsequence, say $r^{(m)}$, so that $\lim_{m \rightarrow \infty} r^{(m)}$ is in $\mathcal{N}_{-1,\delta}(\Phi)$.

Suppose otherwise, there is a subsequence, say $r^{(m)}$ so that $\lim_{m \rightarrow \infty} r^{(m)}$ is in the boundary of $\mathcal{N}_{-1,\delta}(\Phi)$. For $\delta = \pm 1$, there are two possibilities that either for some $v \in V$, $\lim_{m \rightarrow \infty} r^{(m)}(v) = \infty$ or there is an edge e such that $l_e^{(m)} = 0$. For $\delta = 0$, although $r^{(m)}$'s are allowed to be negative, the limit of $r^{(m)}$ can not be $-\infty$ since when ϕ_k is given, the condition in [Lemma 3.6](#) (3)

$$\phi_k > \exp(r_i^{(m)}) + \exp(r_j^{(m)})$$

implies that $r_i^{(m)}$ is bounded away from $-\infty$. Therefore there are only those two possibilities as in case $\delta = \pm 1$.

In the first possibility, by [Lemma 4.6](#) we see each generalized angle incident to the vertex v converges to 0. Hence $\lim_{m \rightarrow \infty} \tilde{K}^{(m)}(v) = 0$. This contradicts the assumption that $\lim_{m \rightarrow \infty} \tilde{K}^{(m)} \in \mathbb{R}_{>0}^V$.

In the second possibility, in a hyperbolic right-angled hexagon with lengths l_i, l_j, l_k and opposite generalized angle $\theta_i, \theta_j, \theta_k$, by the cosine law we see that

$$\begin{aligned} \cosh \theta_j &= \frac{\cosh l_j + \cosh l_i \cosh l_k}{\sinh l_i \sinh l_k} \\ &> \frac{\cosh l_i \cosh l_k}{\sinh l_i \sinh l_k} \\ &\geq \frac{\cosh l_i}{\sinh l_i}. \end{aligned}$$

Hence we have $\lim_{l_i \rightarrow 0} \theta_j = \infty$. Hence the generalized discrete curvature containing θ_j converges to ∞ . This contradicts the assumption that $\lim_{m \rightarrow \infty} \tilde{K}^{(m)} \in \mathbb{R}_{>0}^V$.

Now we have finished the proof of [Theorem 1.4](#).

5 A proof of Theorem 1.6

The proof is based on constructing a strictly concave energy function on the space of all generalized hyperbolic triangles of type $(\varepsilon, \varepsilon, \delta)$ so that its gradient is the generalized angles. Then using [Lemma 2.1](#) on injectivity of gradient, we establish [Theorem 1.6](#).

5.1 An energy functional on the space of triangles

Fix a type $(\varepsilon, \varepsilon, \delta) \in \{-1, 0, 1\}^3$. Consider $(\varepsilon, \varepsilon, \delta)$ type generalized hyperbolic triangles whose edge lengths are l_1, l_2, l_3 and opposite angles $\theta_1, \theta_2, \theta_3$ so that the angle θ_i faces l_i and the type of θ_3 angle is δ . For a fixed angle θ_3 , all values of the two edge lengths (l_1, l_2) form the set $\mathcal{D}_{\varepsilon, \delta}(\theta_3)$. For the definition of $\mathcal{D}_{\varepsilon, \delta}(\theta_3)$, see [Section 3](#).

For $h \in \mathbb{R}$, make a change of variables (l_1, l_2) to (w_1, w_2) and (θ_1, θ_2) to (a_1, a_2) as follows. Let $i = 1, 2$,

$$(5-1) \quad a_i = \int_1^{\theta_i} \rho_\varepsilon^h(t) dt$$

where $\rho_\varepsilon(t) = \int_0^t \cos(\sqrt{\varepsilon}x) dx$ and

$$(5-2) \quad w_i = \int_1^{l_i} \tau_{\varepsilon\delta}^{h-1}(t) dt$$

where $\tau_{\varepsilon\delta}(t) = \frac{1}{2}e^t - \frac{1}{2}\varepsilon\delta e^{-t}$, as introduced in [Section 3](#). By the construction, the maps (l_1, l_2) to $w = (w_1, w_2)$ and (θ_1, θ_2) to $a = (a_1, a_2)$ are diffeomorphisms. Thus the cosine law relating l to θ can be considered, with θ_3 fixed, as a smooth map $a = a(w)$ defined on $w(\mathcal{D}_{\varepsilon, \delta}(\theta_3))$.

Lemma 5.1 *Under the above assumption, for a fixed angle θ_3 , and any $h \in \mathbb{R}$, the differential 1-form $a_1 dw_2 + a_2 dw_1$ is closed in $w(\mathcal{D}_{\varepsilon, \delta}(\theta_3))$. Furthermore, the integration*

$$F_{\theta_3, h}(w_1, w_2) = \int_{(1,1)}^{(w_1, w_2)} (a_1 dw_2 + a_2 dw_1)$$

is strictly concave down in $w(\mathcal{D}_{\varepsilon, \delta}(\theta_3))$. In particular

$$(5-3) \quad \frac{\partial F_{\theta_3, h}}{\partial w_i} = a_j = \int_1^{\theta_j} \rho_\varepsilon^h(t) dt$$

for $\{i, j\} = \{1, 2\}$.

Proof If θ_3 is fixed, then $d\theta_3 = 0$. By the derivative cosine law [\(3-9\)](#) we have

$$\begin{aligned} \begin{pmatrix} dl_1 \\ dl_2 \end{pmatrix} &= \frac{-1}{\sqrt{-\det G_l}} \begin{pmatrix} \tau_{\varepsilon\delta}(l_1) & 0 \\ 0 & \tau_{\varepsilon\delta}(l_2) \end{pmatrix} \begin{pmatrix} \varepsilon & \tau'_{\varepsilon\varepsilon}(l_3) \\ \tau'_{\varepsilon\varepsilon}(l_3) & \varepsilon \end{pmatrix} \begin{pmatrix} d\theta_1 \\ d\theta_2 \end{pmatrix} \\ &= \frac{-1}{\sqrt{-\det G_l}} \begin{pmatrix} \tau_{\varepsilon\delta}(l_1) & 0 \\ 0 & \tau_{\varepsilon\delta}(l_2) \end{pmatrix} \begin{pmatrix} \tau'_{\varepsilon\varepsilon}(l_3) & \varepsilon \\ \varepsilon & \tau'_{\varepsilon\varepsilon}(l_3) \end{pmatrix} \begin{pmatrix} d\theta_2 \\ d\theta_1 \end{pmatrix}. \end{aligned}$$

Since $dw_i = \tau_{\varepsilon\delta}^{h-1}(l_i)dl_i$, $da_i = \rho_{\varepsilon}^h(\theta_i)d\theta_i$, for $i = 1, 2$, then

$$\begin{aligned} \begin{pmatrix} dw_1 \\ dw_2 \end{pmatrix} &= \frac{-1}{\sqrt{-\det G_l}} \begin{pmatrix} \tau_{\varepsilon\delta}^h(l_1) & 0 \\ 0 & \tau_{\varepsilon\delta}^h(l_2) \end{pmatrix} \begin{pmatrix} \tau'_{\varepsilon\varepsilon}(l_3) & \varepsilon \\ \varepsilon & \tau'_{\varepsilon\varepsilon}(l_3) \end{pmatrix} \\ &\quad \begin{pmatrix} \rho_{\varepsilon}^{-h}(\theta_2) & 0 \\ 0 & \rho_{\varepsilon}^{-h}(\theta_1) \end{pmatrix} \begin{pmatrix} da_2 \\ da_1 \end{pmatrix} \\ &=: \frac{-1}{\sqrt{-\det G_l}} A \begin{pmatrix} da_2 \\ da_1 \end{pmatrix}. \end{aligned}$$

Since $A_{12} = \tau_{\varepsilon\delta}^h(l_1)\varepsilon\rho_{\varepsilon}^{-h}(\theta_1) = \tau_{\varepsilon\delta}^h(l_2)\varepsilon\rho_{\varepsilon}^{-h}(\theta_2) = A_{21}$ by the sine law (3-6), the matrix A is symmetric. Thus the differential 1-form $a_1dw_2 + a_2dw_1$ is closed. Therefore the function $F_{\theta_3,h}(w_1, w_2)$ is well defined.

The above calculation shows that the Hessian of the function $F_{\theta_3,h}(w_1, w_2)$ is the matrix $-\sqrt{-\det G_l}A^{-1}$. To show the function $F_{\theta_3,h}(w_1, w_2)$ is strictly concave down, we need to check that A^{-1} is positive definite. It is equivalent to show that A is positive definite. By forgetting the two diagonal matrices, it is enough to show

$$B = \begin{pmatrix} \tau'_{\varepsilon\varepsilon}(l_3) & \varepsilon \\ \varepsilon & \tau'_{\varepsilon\varepsilon}(l_3) \end{pmatrix}$$

is positive definite. Since $\tau'_{\varepsilon\varepsilon}(l_3) = 1/2(e^{l_3} + \varepsilon^2e^{-l_3}) > 0$ and

$$\det B = \tau'^2_{\varepsilon\varepsilon}(l_3) - \varepsilon^2 = \left(\frac{1}{2}e^{l_3} + \frac{1}{2}\varepsilon\varepsilon e^{-l_3}\right)^2 - \varepsilon^2 = \left(\frac{1}{2}e^{l_3} - \frac{1}{2}\varepsilon\varepsilon e^{-l_3}\right)^2 > 0,$$

the matrix B is positive definite. □

5.2 A proof of Theorem 1.6

Now the proof of Theorem 1.6 follows from the routine variational framework. Let us recall the set up in Section 1.5. Suppose (Σ, G) is a cell decomposed surface so that the sets of all vertices, edges and 2-cells are V , E , F respectively. The dual decomposition is G^* with vertices $V^*(\cong F)$. Elements in V^* are denoted by f^* where $f \in F$. A quadrilateral $(v, v', f^*, f'^*) \in V \times V \times V^* \times V^*$ in Σ satisfies $vv' \in E$, $f > vv'$ and $f' > vv'$.

Now fix a type $(\varepsilon, \varepsilon, \delta)$ and a function $\theta: E \rightarrow \overset{\circ}{I}_{\delta}$. The set of all circle pattern metrics is $\mathcal{E}_{\varepsilon,\delta}(\theta) = \{r \in (J_{\varepsilon\delta})^{V^*} \mid (r_i, r_j) \in \mathcal{D}_{\varepsilon,\delta}(\theta(v_i v_j)) \text{ whenever } f_i, f_j \text{ share an edge}\}$. For any circle pattern metric $r \in \mathcal{E}_{\varepsilon,\delta}(\theta)$ and any quadrilateral $(v, v', f^*, f'^*) \in V \times V \times V^* \times V^*$ where $vv' \in E$, $f > vv'$ and $f' > vv'$, construct a type $(\varepsilon, \varepsilon, \delta)$ generalized hyperbolic triangle $\triangle f^* f'^* v$ so that the lengths of $f^* v$ and $f'^* v$ are

$r(f^*)$, $r(f'^*)$, and the generalized angle at v is $\theta(vv')$ of type δ . Realize the quadrilateral (v, v', f^*, f'^*) geometrically as the metric double of $\triangle f^*vf'^*$ along the edge $f^*f'^*$. Now isometrically glue all these geometric quadrilaterals along edges. The result is a polyhedral surface. Recall that for $h \in \mathbb{R}$, the K_h -curvature of r

$$K_h\colon V^* \rightarrow \mathbb{R}$$

is defined by (1-3).

For $h \in \mathbb{R}$, make a change of parameter from $r \in \mathcal{E}_{\varepsilon,\delta}(\theta)$ to $w = w(r) \in w(\mathcal{E}_{\varepsilon,\delta}(\theta))$ so that $w(r)(x) = \int_1^{r(x)} \tau_{\varepsilon\delta}^{h-1}(s) \, ds$ is given by (5-2). We now use Lemma 5.1 to construct a smooth strictly concave function $W\colon w(\mathcal{E}_{\varepsilon,\delta}(\theta)) \rightarrow \mathbb{R}$ so that $\nabla W|_w$ is the generalized curvature K_h of r where $w = w(r)$. Then using Lemma 2.1, we see that Theorem 1.6 follows.

Here is the construction. For $w = w(r) \in w(\mathcal{E}_{\varepsilon,\delta}(\theta))$ and for each quadrilateral (v, v', f^*, f'^*) in Σ , we define the F-energy of it in r metric to be


$$F_{\theta(vv'),h}(w(r(f^*)), w(r(f'^*)))$$

where $F_{\theta,h}$ is given by Lemma 5.1. The function $W\colon w = w(r) \in w(\mathcal{E}_{\varepsilon,\delta}(\theta)) \rightarrow \mathbb{R}$ is the sum of F-energies of all quadrilateral (v, v', f^*, f'^*) in r metric. By the construction, W is smooth and strictly concave. By (5-3) we have

$$\nabla W|_w = K_h|_r$$

where $w = w(r)$. This ends the proof of Theorem 1.6.


Appendix A Formulas of cosine and sine laws



For $\{i, j, k\} = \{1, 2, 3\}$,


$$\begin{aligned} \cosh l_i &= \frac{\cos \theta_i + \cos \theta_j \cos \theta_k}{\sin \theta_j \sin \theta_k}, \\ \cos \theta_i &= \frac{-\cosh l_i + \cosh l_j \cosh l_k}{\sinh l_j \sinh l_k}, \\ \frac{\sin \theta_1}{\sinh l_1} &= \frac{\sin \theta_2}{\sinh l_2} = \frac{\sin \theta_3}{\sinh l_3}. \end{aligned}$$

For $\{i, j\} = \{1, 2\}$,




$$\begin{aligned}\sinh l_i &= \frac{\cos \theta_i + \cos \theta_j \cosh \theta_3}{\sin \theta_j \sinh \theta_3}, \\ \cosh l_3 &= \frac{\cosh \theta_3 + \cos \theta_1 \cos \theta_2}{\sin \theta_1 \sin \theta_2}, \\ \cos \theta_i &= \frac{-\sinh l_i + \sinh l_j \cosh l_3}{\cosh l_j \sinh l_3}, \\ \cosh \theta_3 &= \frac{\cosh l_3 + \sinh l_1 \sinh l_2}{\cosh l_1 \cosh l_2}, \\ \frac{\sin \theta_1}{\cosh l_1} &= \frac{\sin \theta_2}{\cosh l_2} = \frac{\sinh \theta_3}{\sinh l_3}.\end{aligned}$$

For $\{i, j\} = \{1, 2\}$,




$$\begin{aligned}\sinh l_i &= \frac{\cosh \theta_i + \cosh \theta_j \cos \theta_3}{\sinh \theta_j \sin \theta_3}, \\ \cosh l_3 &= \frac{\cos \theta_3 + \cosh \theta_1 \cosh \theta_2}{\sinh \theta_1 \sinh \theta_2}, \\ \cos \theta_i &= \frac{\sinh l_i + \sinh l_j \cosh l_3}{\cosh l_j \sinh l_3}, \\ \cos \theta_3 &= \frac{-\cosh l_3 + \sinh l_1 \sinh l_2}{\cosh l_1 \cosh l_2}, \\ \frac{\sinh \theta_1}{\cosh l_1} &= \frac{\sinh \theta_2}{\cosh l_2} = \frac{\sin \theta_3}{\sinh l_3}.\end{aligned}$$

For $\{i, j, k\} = \{1, 2, 3\}$,




$$\begin{aligned}\cosh l_i &= \frac{\cosh \theta_i + \cosh \theta_j \cosh \theta_k}{\sinh \theta_j \sinh \theta_k}, \\ \cosh \theta_i &= \frac{\cosh l_i + \cosh l_j \cosh l_k}{\sinh l_j \sinh l_k}, \\ \frac{\sinh \theta_1}{\sinh l_1} &= \frac{\sinh \theta_2}{\sinh l_2} = \frac{\sinh \theta_3}{\sinh l_3}.\end{aligned}$$




For $\{i, j\} = \{1, 2\}$,

$$\begin{aligned} \frac{e^{l_i}}{2} &= \frac{\cos \theta_i + \cos \theta_j}{\theta_3 \sin \theta_j}, & \cos \theta_i &= \frac{-e^{l_i} + e^{l_j} \cosh l_3}{e^{l_j} \sinh l_3}, \\ \cosh l_3 &= \frac{1 + \cos \theta_1 \cos \theta_2}{\sin \theta_1 \sin \theta_2}, & \frac{\theta_3^2}{2} &= \frac{\cosh l_3 - \cosh(l_1 - l_2)}{\frac{e^{l_1+l_2}}{4}}, \\ \frac{\sin \theta_1}{\frac{e^{l_1}}{2}} &= \frac{\sin \theta_2}{\frac{e^{l_2}}{2}} = \frac{\theta_3}{\sinh l_3}. \end{aligned}$$



$$\begin{aligned} \frac{e^{l_1}}{2} &= \frac{\cos \theta_1 + \cosh \theta_2}{\theta_3 \sinh \theta_2}, & \cos \theta_1 &= \frac{-e^{l_1} + e^{l_2} \sinh l_3}{e^{l_2} \cosh l_3}, \\ \frac{e^{l_2}}{2} &= \frac{\cosh \theta_2 + \cos \theta_1}{\theta_3 \sin \theta_1}, & \cosh \theta_2 &= \frac{e^{l_2} + e^{l_1} \sinh l_3}{e^{l_1} \cosh l_3}, \\ \sinh l_3 &= \frac{1 + \cos \theta_1 \cosh \theta_2}{\sin \theta_1 \sinh \theta_2}, & \frac{\theta_3^2}{2} &= \frac{\sinh l_3 + \sinh(l_2 - l_1)}{\frac{e^{l_1+l_2}}{4}}, \\ \frac{\sin \theta_1}{\frac{e^{l_1}}{2}} &= \frac{\sinh \theta_2}{\frac{e^{l_2}}{2}} = \frac{\theta_3}{\cosh l_3}. \end{aligned}$$



For $\{i, j\} = \{1, 2\}$,


$$\frac{e^{l_i}}{2} = \frac{\cosh \theta_i + \cosh \theta_j}{\theta_3 \sinh \theta_j},$$

$$\cosh \theta_i = \frac{e^{l_i} + e^{l_j} \cosh l_3}{e^{l_j} \sinh l_3},$$

$$\cosh l_3 = \frac{1 + \cosh \theta_1 \cosh \theta_2}{\sinh \theta_1 \sinh \theta_2},$$

$$\frac{\theta_3^2}{2} = \frac{\cosh l_3 + \cosh(l_1 - l_2)}{\frac{e^{l_1+l_2}}{4}},$$

$$\frac{\sinh \theta_1}{\frac{e^{l_1}}{2}} = \frac{\sinh \theta_2}{\frac{e^{l_2}}{2}} = \frac{\theta_3}{\sinh l_3}.$$



For $\{i, j\} = \{1, 2\}$,

$$\frac{e^{l_i}}{2} = \frac{1 + \cos \theta_3}{\theta_j \sin \theta_3},$$


$$\frac{\theta_i^2}{4} = \frac{e^{l_i} - e^{l_3-l_j}}{e^{l_j+l_3}},$$

$$\frac{e^{l_3}}{2} = \frac{1 + \cos \theta_3}{\theta_1 \theta_2},$$

$$\sin^2 \frac{\theta_3}{2} = e^{l_3-l_1-l_2},$$

$$\frac{\theta_1}{e^{l_1}} = \frac{\theta_2}{e^{l_2}} = \frac{\sin \theta_3}{e^{l_3}}.$$

For $\{i, j\} = \{1, 2\}$,



$$\frac{e^{l_i}}{2} = \frac{1 + \cosh \theta_3}{\theta_j \sinh \theta_3},$$


$$\frac{e^{l_3}}{2} = \frac{1 + \cosh \theta_3}{\theta_1 \theta_2},$$

$$\frac{\theta_i^2}{4} = \frac{e^{l_i} + e^{l_3 - l_j}}{e^{l_j + l_3}},$$

$$\sinh^2 \frac{\theta_3}{2} = e^{l_3 - l_1 - l_2},$$

$$\frac{\theta_1}{e^{l_1}} = \frac{\theta_2}{e^{l_2}} = \frac{\sinh \theta_3}{e^{l_3}}.$$

For $\{i, j, k\} = \{1, 2, 3\}$,



$$\frac{e^{l_i}}{2} = \frac{2}{\theta_j \theta_k},$$

$$\frac{\theta_i^2}{4} = e^{l_i - l_j - l_k},$$

$$\frac{\theta_1}{e^{l_1}} = \frac{\theta_2}{e^{l_2}} = \frac{\theta_3}{e^{l_3}}.$$

Appendix B Proofs of Lemma 3.3 and Lemma 3.5

For simplicity, we abuse the notation. Let

$$\begin{aligned} g(\theta_i) &:= \rho_{\varepsilon_i}(\theta_i), & g'(\theta_i) &:= \rho'_{\varepsilon_i}(\theta_i), \\ f(l_k) &:= \tau_{\varepsilon_i \varepsilon_j}(l_k), & f'(l_k) &:= \tau'_{\varepsilon_i \varepsilon_j}(l_k), \end{aligned}$$

for $\{i, j, k\} = \{1, 2, 3\}$, where g, f represent functions which depend on the type of generalized vertices.

In this simplified notation we have

$$\begin{aligned} -\det G_I &= \det \begin{pmatrix} \varepsilon_1 & f'(l_3) & f'(l_2) \\ f'(l_3) & \varepsilon_2 & f'(l_1) \\ f'(l_2) & f'(l_1) & \varepsilon_3 \end{pmatrix} \\ &= \varepsilon_1 \varepsilon_2 \varepsilon_3 + 2f'(l_1)f'(l_2)f'(l_3) - \varepsilon_1 f'^2(l_1) - \varepsilon_2 f'^2(l_2) - \varepsilon_3 f'^2(l_3). \end{aligned}$$

If $\varepsilon_3 = 0$, then $g(\theta_3) = \theta_3$, $f(l_1) = \frac{e^{l_1}}{2}$, $f(l_2) = \frac{e^{l_2}}{2}$. By the cosine law (3-4), we have

$$\frac{\theta_3^2}{2} = \frac{f'(l_3) - \frac{1}{2}\varepsilon_1 e^{l_1-l_2} - \frac{1}{2}\varepsilon_2 e^{l_2-l_1}}{f(l_1)f(l_2)}.$$

Thus the negative of the right hand side of (3-7) is

$$\begin{aligned} f^2(l_1)f^2(l_2)g^2(\theta_3) &= f^2(l_1)f^2(l_2)\theta_3^2 \\ &= f^2(l_1)f^2(l_2)\frac{2f'(l_3) - \varepsilon_1 e^{l_1-l_2} - \varepsilon_2 e^{l_2-l_1}}{f(l_1)f(l_2)} \\ &= f(l_1)f(l_2)(2f'(l_3) - \varepsilon_1 e^{l_1-l_2} - \varepsilon_2 e^{l_2-l_1}) \\ &= \frac{e^{l_1}}{2} \frac{e^{l_2}}{2} (2f'(l_3) - \varepsilon_1 e^{l_1-l_2} - \varepsilon_2 e^{l_2-l_1}) \\ &= 2 \frac{e^{l_1}}{2} \frac{e^{l_2}}{2} f'(l_3) - \varepsilon_1 \frac{e^{2l_1}}{4} - \varepsilon_2 \frac{e^{2l_2}}{4} \\ &= -\det G_l. \end{aligned}$$

If $\varepsilon_3 = \pm 1$, then $g^2(\theta_3) = \varepsilon_3(1 - g'^2(\theta_3))$. And we have

$$(B-1) \quad f^2(l_i) - f'^2(l_i) = \left(\frac{1}{2}e^l - \frac{1}{2}\varepsilon_j \varepsilon_k e^{-l} \right)^2 - \left(\frac{1}{2}e^l + \frac{1}{2}\varepsilon_j \varepsilon_k e^{-l} \right)^2 = -\varepsilon_j \varepsilon_k.$$

Thus the negative of the right hand side of (3-7) is

$$\begin{aligned} f^2(l_1)f^2(l_2)g^2(\theta_3) &= f^2(l_1)f^2(l_2)(1 - g'^2(\theta_3)) \\ &= \varepsilon_3(f^2(l_1)f^2(l_2) - (-\varepsilon_3 f'(l_3) + f'(l_1)f'(l_2))^2) \\ &= \varepsilon_3((f'^2(l_1) - \varepsilon_2 \varepsilon_3)(f'^2(l_2) - \varepsilon_3 \varepsilon_1) \\ &\quad - (-\varepsilon_3 f'(l_3) + f'(l_1)f'(l_2))^2) \\ &= \varepsilon_3(\varepsilon_1 \varepsilon_2 \varepsilon_3^2 + 2\varepsilon_3 f'(l_1)f'(l_2)f'(l_3) \\ &\quad - \varepsilon_1 \varepsilon_3 f'^2(l_1) - \varepsilon_2 \varepsilon_3 f'^2(l_2) - \varepsilon_3^2 f'^2(l_3)) \\ &= -\det G_l. \end{aligned}$$

The second equality is due to the cosine law, the third equality is due to (B-1) and the last equality is due to $\varepsilon_3 = \pm 1$.

And we have

$$\begin{aligned} -\det G_\theta &= \det \begin{pmatrix} -1 & g'(\theta_3) & g'(\theta_2) \\ g'(\theta_3) & -1 & g'(\theta_1) \\ g'(\theta_2) & g'(\theta_1) & -1 \end{pmatrix} \\ &= -1 + 2g'(\theta_1)g'(\theta_2)g'(\theta_3) + g'^2(\theta_1) + g'^2(\theta_2) + g'^2(\theta_3). \end{aligned}$$

Notice that we have

$$(B-2) \quad g'^2(\theta_i) + \varepsilon_i g^2(\theta_i) = 1.$$

Thus negative of the right hand side of (3-8) is

$$\begin{aligned} g^2(\theta_1)g^2(\theta_2)f^2(l_3) &= g^2(\theta_1)g^2(\theta_2)(f'^2(l_3) - \varepsilon_1\varepsilon_2) \\ &= (g'(\theta_3) + g'(\theta_1)g'(\theta_2))^2 - g^2(\theta_1)g^2(\theta_2)\varepsilon_1\varepsilon_2 \\ &= (g'(\theta_3) + g'(\theta_1)g'(\theta_2))^2 - (1 - g'^2(\theta_1))(1 - g'^2(\theta_2)) \\ &= -\det G_\theta. \end{aligned}$$

The second equality is due to the cosine law and the third equality is due to (B-2).

We can prove either one of the derivative cosine law (3-9) and (3-10). The other one will be a corollary due to Lemma 3.4. For example, if (3-9) is true, we have

$$\begin{pmatrix} dl_1 \\ dl_2 \\ dl_3 \end{pmatrix} = MG_l \begin{pmatrix} d\theta_1 \\ d\theta_2 \\ d\theta_3 \end{pmatrix}.$$

It is equivalent to

$$\begin{pmatrix} d\theta_1 \\ d\theta_2 \\ d\theta_3 \end{pmatrix} = (MG_l)^{-1} \begin{pmatrix} dl_1 \\ dl_2 \\ dl_3 \end{pmatrix}.$$

By Lemma 3.4, $(MG_l)^{-1} = MG_\theta$. Thus we obtained (3-10):

$$\begin{pmatrix} d\theta_1 \\ d\theta_2 \\ d\theta_3 \end{pmatrix} = MG_\theta \begin{pmatrix} dl_1 \\ dl_2 \\ dl_3 \end{pmatrix}.$$

In the following, we give a proof of (3-9). By the cosine law (3-3) we have

$$f'(l_i)g(\theta_j)g(\theta_k) = g'(\theta_i) + g'(\theta_j)g'(\theta_k).$$

After differentiating the two sides we have

$$f''(l_i)g(\theta_j)g(\theta_k)dl_i + f'(l_i)g'(\theta_j)g(\theta_k)d\theta_j + f'(l_i)g(\theta_j)g'(\theta_k)d\theta_k \\ = g''(\theta_i)d\theta_i + g''(\theta_j)g'(\theta_k)d\theta_j + g'(\theta_j)g''(\theta_k)d\theta_k$$

which is equivalent to

$$(B-3) \quad f''(l_i)g(\theta_j)g(\theta_k)dl_i \\ = g''(\theta_i)d\theta_i + (g''(\theta_j)g'(\theta_k) - f'(l_i)g'(\theta_j)g(\theta_k))d\theta_j \\ + (g'(\theta_j)g''(\theta_k) - f'(l_i)g(\theta_j)g'(\theta_k))d\theta_k.$$

By the cosine law (3-3), the coefficient of $d\theta_j$ in (B-3) is

$$g''(\theta_j)g'(\theta_k) - f'(l_i)g'(\theta_j)g(\theta_k) \\ = g''(\theta_j)g'(\theta_k) - \frac{g'(\theta_i) + g'(\theta_j)g'(\theta_k)}{g(\theta_j)g(\theta_k)}g'(\theta_j)g(\theta_k) \\ (B-4) \quad = \frac{1}{g(\theta_j)}((g(\theta_j)g''(\theta_j) - g'^2(\theta_j))g'(\theta_k) - g'(\theta_i)g'(\theta_j)).$$

For $g(\theta) = \sin \theta$, or $\sinh \theta$, or θ , we always have

$$g(\theta_j)g''(\theta_j) - g'^2(\theta_j) = -1.$$

Hence (B-4) is

$$\frac{1}{g(\theta_j)}(-g'(\theta_k) - g'(\theta_i)g'(\theta_j)) = -g(\theta_i)f(l_k)$$

due to the cosine law (3-3). By symmetry, the similar formula holds for the coefficient of $d\theta_k$. Hence (B-3) is

$$f''(l_i)g(\theta_j)g(\theta_k)dl_i = g''(\theta_i)d\theta_i - g(\theta_i)f(l_k)d\theta_j - g(\theta_i)f(l_j)d\theta_k.$$

By the definition of f , we have $f'' = f$. Thus

$$dl_i = \frac{-g(\theta_i)}{f(l_i)g(\theta_j)g(\theta_k)} \left(-\frac{g''(\theta_i)}{g(\theta_i)}d\theta_i + f(l_k)d\theta_j + f(l_j)d\theta_k \right) \\ = \frac{-g(\theta_i)}{f(l_i)g(\theta_j)g(\theta_k)}(\varepsilon_i d\theta_i + f(l_k)d\theta_j + f(l_j)d\theta_k).$$

This proves (3-9).

References

- [1] **A I Bobenko, B A Springborn**, *Variational principles for circle patterns and Koebe's theorem*, Trans. Amer. Math. Soc. 356 (2004) 659–689 [MR2022715](#)
- [2] **W Brägger**, *Kreispackungen und Triangulierungen*, Enseign. Math. (2) 38 (1992) 201–217 [MR1189006](#)
- [3] **B Chow, F Luo**, *Combinatorial Ricci flows on surfaces*, J. Differential Geom. 63 (2003) 97–129 [MR2015261](#)
- [4] **Y Colin de Verdière**, *Un principe variationnel pour les empilements de cercles*, Invent. Math. 104 (1991) 655–669 [MR1106755](#)
- [5] **W Fenchel, J Nielsen**, *Discontinuous groups of isometries in the hyperbolic plane*, de Gruyter Studies in Math. 29, Walter de Gruyter & Co., Berlin (2003) [MR1958350](#)
Edited and with a preface by A L Schmidt, Biography of the authors by B Fuglede
- [6] **R Guo**, *On parameterizations of Teichmüller spaces of surfaces with boundary*, to appear in J. Differential Geom. [arXiv:math.GT/0612221](#)
- [7] **G P Hazel**, *Triangulating Teichmüller space using the Ricci flow*, PhD thesis, University of California San Diego (2004) Available at www.math.ucsd.edu/~thesis/thesis/ghazel/ghazel.pdf
- [8] **G Leibon**, *Characterizing the Delaunay decompositions of compact hyperbolic surfaces*, Geom. Topol. 6 (2002) 361–391 [MR1914573](#)
- [9] **F Luo**, *Rigidity of polyhedral surfaces* [arXiv:math.GT/0612714](#)
- [10] **F Luo**, *A characterization of spherical polyhedral surfaces*, J. Differential Geom. 74 (2006) 407–424 [MR2269784](#)
- [11] **F Luo**, *On Teichmüller spaces of surfaces with boundary*, Duke Math. J. 139 (2007) 463–482 [MR2350850](#)
- [12] **G Mondello**, *Triangulated Riemann surfaces with boundary and the Weil–Petersson Poisson structure*, J. Differential Geom. 81 (2009) 391–436
- [13] **R C Penner**, *The decorated Teichmüller space of punctured surfaces*, Comm. Math. Phys. 113 (1987) 299–339 [MR919235](#)
- [14] **I Rivin**, *Euclidean structures on simplicial surfaces and hyperbolic volume*, Ann. of Math. (2) 139 (1994) 553–580 [MR1283870](#)
- [15] **J-M Schlenker**, *Circle patterns on singular surfaces*, Discrete Comput. Geom. 40 (2008) 47–102 [MR2429649](#)
- [16] **K Stephenson**, *Introduction to circle packing: The theory of discrete analytic functions*, Cambridge Univ. Press (2005) [MR2131318](#)
- [17] **W P Thurston**, *The geometry and topology of three-manifolds*, Princeton Univ. Math. Dept. Lecture Notes (1979) Available at <http://msri.org/publications/books/gt3m/>

- [18] **W P Thurston**, *Three-dimensional geometry and topology. Vol. 1*, Princeton Math. Series 35, Princeton University Press (1997) [MR1435975](#) Edited by S Levy

*School of Mathematics, University of Minnesota
Minneapolis, MN 55455, USA*

*The Center of Mathematical Science, Zhejiang University
Hangzhou, Zhejiang 310027, China*
and

*Department of Mathematics, Rutgers University
Piscataway, NJ 08854, USA*

guoxx170@math.umn.edu, fluo@math.rutgers.edu

Proposed: David Gabai

Seconded: Peter Teichner, Walter Neumann

Received: 5 November 2007

Accepted: 17 January 2009