

Some new examples with almost positive curvature

MARTIN KERIN

As a means to better understanding manifolds with positive curvature, there has been much recent interest in the study of nonnegatively curved manifolds which contain points at which all 2–planes have positive curvature. We show that there are generalisations of the well-known Eschenburg spaces and quotients of $\mathbf{S}^7 \times \mathbf{S}^7$ which admit metrics with this property.

53C20; 57S25, 57T15

It is an unfortunate fact that for a simply connected manifold which admits a metric of nonnegative curvature there are no known obstructions to admitting positive curvature. While there exist many examples of manifolds with nonnegative curvature, the known examples with positive curvature are very sparse (see Ziller [31] for a comprehensive survey of both situations). Other than the rank-one symmetric spaces, there are isolated examples in dimensions 6, 7, 12, 13 and 24 due to Wallach [27] and Berger [4], and two infinite families, one in dimension 7 (Eschenburg spaces; see Aloff and Wallach [2] and Eschenburg [9; 10]) and the other in dimension 13 (Bazaikin spaces [3]). In recent developments, two distinct metrics with positive curvature on a particular cohomogeneity-one manifold have been proposed by Grove, Verdiani and Ziller [15] and Darricott [8], while Petersen and Wilhelm [21] propose that the Gromoll–Meyer exotic 7–sphere admits positive curvature, which would be the first exotic sphere known to exhibit this property.

In this paper we are interested in the study of manifolds which lie “between” those with nonnegative and those with positive sectional curvature. It is hoped that the study of such manifolds will yield a better understanding of the differences between these two classes. Recall that a Riemannian manifold $(M, \langle \cdot, \cdot \rangle)$ is said to have *quasipositive curvature* (resp. *almost positive curvature*) if $(M, \langle \cdot, \cdot \rangle)$ has nonnegative sectional curvature and there is a point (resp. an open dense set of points) at which all 2–planes have positive sectional curvature.

Theorem A

- (i) *There exists a free circle action and a free \mathbf{S}^3 action on $\mathbf{S}^7 \times \mathbf{S}^7$ such that each of the respective quotients $M^{13} := \mathbf{S}^1 \backslash (\mathbf{S}^7 \times \mathbf{S}^7)$ and $N^{11} := \mathbf{S}^3 \backslash (\mathbf{S}^7 \times \mathbf{S}^7)$ admits a metric with almost positive curvature.*

- (ii) If M^{13} and N^{11} are equipped with the metrics from (i), then there exist Riemannian submersions $M^{13} \rightarrow \mathbf{CP}^3$ and $N^{11} \rightarrow \mathbf{S}^4$ such that in each case the fibre is \mathbf{S}^7 and the bundle is nontrivial but shares the same cohomology ring as the corresponding product.

We use the Pontrjagin class to distinguish M^{13} and N^{11} from the respective products in Theorem A(ii). Moreover, in each case the induced metric on the base is positively curved.

It has been conjectured by F Wilhelm that, if M is a positively curved Riemannian manifold, the dimension of the fibre of a Riemannian submersion $M \rightarrow B$ must be less than the dimension of the base. Theorem A shows that this is false when the hypothesis is weakened to almost positive curvature.

Other than the Gromoll–Meyer exotic 7–sphere (see Gromoll and Meyer [14], Wilhelm [29], Eschenburg and Kerin [11] and Petersen and Wilhelm [21]), the only other previously known examples of manifolds with almost positive or quasipositive curvature are given in Petersen and Wilhelm [22], Wilking [30], Tapp [25] and Kerin [18].

While Wilking [30] has shown that it is not possible in general to deform quasipositive curvature to positive curvature, it is still unknown whether this can be achieved in the simply connected case or whether one can always deform quasipositive curvature to almost positive curvature.

Theorem B Let $L_{p,q} \subset U(n + 1) \times U(n + 1)$, $n \geq 2$, be defined by

$$L_{p,q} = \{(\text{diag}(z^{p_1}, \dots, z^{p_{n+1}}), \text{diag}(z^{q_1}, z^{q_2}, A)) \mid z \in \mathbf{S}^1, A \in U(n - 1)\}$$

where $p_1, \dots, p_{n+1}, q_1, q_2 \in \mathbb{Z}$. When the two-sided action of $L_{p,q}$ on $U(n + 1)$ is free, denote the quotient $U(n + 1) // L_{p,q}$ by $E_{p,q}^{4n-1}$. All $E_{p,q}^{4n-1}$ admit a metric with quasipositive curvature.

Observe that to allow $p_i = p_j$ for all $1 \leq i < j \leq n + 1$ in Theorem B is equivalent to setting $p_i = 0$ for all i , since the centre of $U(n + 1)$ is given by multiples of the identity. Wilking [30] showed that these homogeneous spaces admit a metric with almost positive curvature whenever $q_1 q_2 < 0$, while Tapp [25] subsequently showed that $(q_1, q_2) \neq (0, 0)$ is sufficient to guarantee the existence of a metric with quasipositive curvature.

The biquotients $E_{p,q}^{4n-1}$ in Theorem B should be thought of as generalisations of the Eschenburg spaces, which arise when $n = 2$. In Eschenburg [9] it is shown that infinitely many Eschenburg spaces admit positive curvature, while in Kerin [18] it is shown that all Eschenburg spaces admit a metric with quasipositive curvature.

The paper is organised as follows. In [Section 1](#) we review some notation and geometric techniques for biquotients. In [Section 2](#) we review some facts about the Cayley numbers and the exceptional Lie group G_2 . In [Section 3](#) we describe the manifolds M^{13} and N^{11} of [Theorem A](#) as biquotients and as the total spaces of Riemannian submersions. We prove the curvature statements of [Theorem A\(i\)](#) in [Section 4](#), while proof of the topological statements in [\(ii\)](#) is postponed until [Section 6](#). [Section 5](#) is devoted to establishing [Theorem B](#).

Acknowledgments The majority of this work was completed as part of a PhD thesis at the University of Pennsylvania under the supervision of Wolfgang Ziller. His constant advice, support and friendship were inspiring. Thanks also to Jost Eschenburg for useful comments. Finally, I wish to thank Burkhard Wilking for suggesting some improvements.

1 Biquotient actions and metrics

In his Habilitation [\[9\]](#), Eschenburg studied biquotients in great detail. The following section provides a review of the material in [\[9\]](#) and establishes the basic language, notation and results which will be used throughout the remainder of the paper.

Let G be a compact Lie group, $U \subset G \times G$ a closed subgroup, and let U act on G via

$$(u_1, u_2) \star g = u_1 g u_2^{-1}, \quad g \in G, (u_1, u_2) \in U.$$

The action is free if and only if, for all nontrivial $(u_1, u_2) \in U$, u_1 is never conjugate to u_2 in G . The resulting manifold is called a *biquotient*.

Let $K \subset G$ be a closed subgroup, $\langle \cdot, \cdot \rangle$ be a left-invariant, right K -invariant metric on G , and $U \subset G \times K \subset G \times G$ act freely on G as above. Let $g \in G$. Define

$$U_L^g := \{(g u_1 g^{-1}, u_2) \mid (u_1, u_2) \in U\}.$$

Since U acts freely on G , so too does U_L^g , and $G//U$ is isometric to $G//U_L^g$. This follows from the fact that left-translation $L_g: G \rightarrow G$ is an isometry which satisfies $g u_1 g^{-1} (L_g g') u_2^{-1} = L_g (u_1 g' u_2^{-1})$. Therefore L_g induces an isometry of the orbit spaces $G//U$ and $G//U_L^g$.

Consider a Riemannian submersion $\pi: M^n \rightarrow N^{n-k}$. By O'Neill's formula for Riemannian submersions, π is curvature nondecreasing. Therefore $\sec_M \geq 0$ implies $\sec_N \geq 0$, and zero-curvature planes on N lift to horizontal zero-curvature planes on M . Because of the Lie bracket term in the O'Neill formula the converse is not true in

general, namely horizontal zero-curvature planes in M cannot be expected to project to zero-curvature planes on N .

Let $K \subset G$ be Lie groups, $\mathfrak{k} \subset \mathfrak{g}$ the corresponding Lie algebras, and $\langle \cdot, \cdot \rangle$ a nonnegatively curved left-invariant metric on G which is right-invariant under K . We can write $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ with respect to $\langle \cdot, \cdot \rangle$. Given $X \in \mathfrak{g}$ we will always use $X_{\mathfrak{k}}$ and $X_{\mathfrak{p}}$ to denote the \mathfrak{k} and \mathfrak{p} components of X respectively.

Recall that
$$G \cong (G \times K) / \Delta K$$

via $(g, k) \mapsto gk^{-1}$, where ΔK is the free diagonal action of K on the right of $G \times K$. Notice that the restriction of $\langle \cdot, \cdot \rangle$ to K is bi-invariant. Equip $G \times K$ with the metric $\langle\langle \cdot, \cdot \rangle\rangle := \langle \cdot, \cdot \rangle \oplus t \langle \cdot, \cdot \rangle|_{\mathfrak{k}}$, $t > 0$. Thus we may define a new left-invariant, right K -invariant metric $\langle \cdot, \cdot \rangle_1$ (with $\sec \geq 0$) on G via the Riemannian submersion

$$(1) \quad \begin{aligned} (G \times K, \langle\langle \cdot, \cdot \rangle\rangle) &\longrightarrow (G, \langle \cdot, \cdot \rangle_1) \\ (g, k) &\longmapsto gk^{-1} \end{aligned}$$

with
$$\langle X, Y \rangle_1 = \langle X, \Phi(Y) \rangle$$

where $\Phi(Y) = Y_{\mathfrak{p}} + \lambda Y_{\mathfrak{k}}$, $\lambda = t/(t + 1) \in (0, 1)$. Furthermore, it is clear that the metric tensor Φ is invertible with inverse described by $\Phi^{-1}(Y) = Y_{\mathfrak{p}} + (1/\lambda)Y_{\mathfrak{k}}$.

Suppose $\sigma = \text{Span}\{\Phi^{-1}(X), \Phi^{-1}(Y)\} \subset \mathfrak{g}$ is a zero-curvature plane with respect to the metric $\langle \cdot, \cdot \rangle_1$, ie $\sec_1(\sigma) = 0$. By the O’Neill formula σ must therefore lift to a horizontal zero-curvature plane $\tilde{\sigma} \subset \mathfrak{g} \oplus \mathfrak{k}$ with respect to $\langle\langle \cdot, \cdot \rangle\rangle$. It is easy to check that the horizontal lift of a vector $\Phi^{-1}(X) \in \mathfrak{g}$ to $\mathfrak{g} \oplus \mathfrak{k}$ is given by $(X, -(1/t)X_{\mathfrak{k}})$. Then clearly

$$\tilde{\sigma} = \text{Span} \left\{ \left(X, -\frac{1}{t}X_{\mathfrak{k}} \right), \left(Y, -\frac{1}{t}Y_{\mathfrak{k}} \right) \right\}.$$

But, since $\langle\langle \cdot, \cdot \rangle\rangle$ is a nonnegatively curved product metric, it follows immediately by considering the unnormalised curvature that $\tilde{\sigma}$ has zero-curvature if and only if $[X_{\mathfrak{k}}, Y_{\mathfrak{k}}] = 0$ and the plane $\text{Span}\{X, Y\} \subset \mathfrak{g}$ has zero-curvature with respect to the original metric $\langle \cdot, \cdot \rangle$, ie $\sec(X, Y) = 0$.

From Tapp [26], which generalizes similar results in Eschenburg [9] and Wilking [30], we know that if $\langle \cdot, \cdot \rangle$ is induced by a Riemannian submersion to G from a bi-invariant metric on some Lie group L , then in fact $\sec_1(\sigma) = 0$ if and only if $\sec(\tilde{\sigma}) = 0$ with respect to $\langle\langle \cdot, \cdot \rangle\rangle$, ie if and only if $\sec(X, Y) = 0$ and $[X_{\mathfrak{k}}, Y_{\mathfrak{k}}] = 0$. We will always be in this situation as throughout the paper we will use only the metrics described in [Example \(a\)](#) and [Example \(b\)](#) below.

Example (a) Suppose that (G, K) is a symmetric pair and that the initial metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_0$ is a bi-invariant metric on G . As in (1), equip G with a new metric

$$(2) \quad \langle X, Y \rangle_1 = \langle X, \Phi_1(Y) \rangle_0$$

where $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ with respect to $\langle \cdot, \cdot \rangle_0$ and $\Phi_1(Y) = Y_p + \lambda_1 Y_{\mathfrak{k}}$. Then

$$\sigma = \text{Span}\{\Phi_1^{-1}(X), \Phi_1^{-1}(Y)\} \subset \mathfrak{g}$$

has zero-curvature with respect to $\langle \cdot, \cdot \rangle_1$, ie $\text{sec}_1(\sigma) = 0$, if and only if

$$(3) \quad 0 = [X, Y] = [X_{\mathfrak{k}}, Y_{\mathfrak{k}}] = [X_p, Y_p].$$

The proof of this follows immediately from our previous discussion together with the fact that $[\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{k}$ whenever (G, K) is a symmetric pair.

Example (b) Let $G \supset K \supset H$ be a chain of subgroups and suppose that both (G, K) and (K, H) are symmetric pairs. Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ and $\mathfrak{k} = \mathfrak{h} \oplus \mathfrak{m}$ be the corresponding orthogonal decompositions with respect to the bi-invariant metric $\langle \cdot, \cdot \rangle_0$ on G . Start with the metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_1$ defined by Example (a). Now define the metric $\langle \cdot, \cdot \rangle_2$ on G as in (1), where K is replaced by H , $s > 0$ takes the role of t , and Ψ replaces Φ :

$$(4) \quad \langle X, Y \rangle_2 = \langle X, \Psi(Y) \rangle_1$$

$$(5) \quad = \langle X, \Phi_2(Y) \rangle_0$$

with $\Phi_2(Y) = Y_p + \lambda_1 Y_m + \lambda_1 \lambda_2 Y_{\mathfrak{h}}$, $\lambda_2 = s/(s + 1)$, and $\Psi(Y) = \Phi_1^{-1} \Phi_2(Y) = Y_p + Y_m + \lambda_2 Y_{\mathfrak{h}}$.

Let $\sigma = \text{Span}\{\Psi^{-1}(X), \Psi^{-1}(Y)\} \subset \mathfrak{g}$. Then, by our discussion prior to Example (a), $\text{sec}_2(\sigma) = 0$ if and only if $\text{sec}_1(X, Y) = 0$ and $[X_{\mathfrak{h}}, Y_{\mathfrak{h}}] = 0$. By again considering horizontal lifts it is not difficult to check that $\text{sec}_1(X, Y) = 0$ if and only if conditions (3) hold as for $\text{sec}_1(\Phi_1^{-1}(X), \Phi_1^{-1}(Y)) = 0$. Hence $\text{sec}_2(\sigma) = 0$ if and only if

$$(6) \quad 0 = [X, Y] = [X_{\mathfrak{k}}, Y_{\mathfrak{k}}] = [X_p, Y_p] = [X_m, Y_m] = [X_{\mathfrak{h}}, Y_{\mathfrak{h}}]$$

where we have used the fact that $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$ since (K, H) is a symmetric pair.

Now that we have described how to induce new metrics on G from old ones and derived zero-curvature conditions for these metrics, we proceed to consider biquotients $G//U$. Let $\Delta G = \{(g, g) \mid g \in G\}$. Then, if the two-sided action of U on G is free, $\Delta G \times U$ acts freely on $G \times G$ via

$$(7) \quad ((g, g), (u_1, u_2)) \star (g_1, g_2) = (gg_1u_1^{-1}, gg_2u_2^{-1})$$

with $(g, g) \in \Delta G$, $(u_1, u_2) \in U$, $(g_1, g_2) \in G \times G$, and there is a canonical diffeomorphism

$$(8) \quad \Delta G \backslash (G \times G) / U \cong G // U$$

induced by the map

$$\begin{aligned} G \times G &\longrightarrow G \\ (g_1, g_2) &\longmapsto g_1^{-1} g_2. \end{aligned}$$

Let K_1 and K_2 be arbitrary subgroups of G . We define left-invariant metrics, $\langle \cdot, \cdot \rangle_{K_1}$ and $\langle \cdot, \cdot \rangle_{K_2}$, on G as in (1). Equip $G \times G$ with a left-invariant, right $(K_1 \times K_2)$ -invariant product metric $((\cdot, \cdot)) = \langle \cdot, \cdot \rangle_{K_1} \oplus \langle \cdot, \cdot \rangle_{K_2}$. If $U \subset K_1 \times K_2$ then the $\Delta G \times U$ action is by isometries and $((\cdot, \cdot))$ induces a metric on $G // U$. Our goal is to determine when a plane tangent to $G // U$ has zero-curvature with respect to this induced metric.

By (8) and our choice of metric, the quotient map $(G \times G, ((\cdot, \cdot))) \longrightarrow G // U$ is a Riemannian submersion. O’Neill’s formula implies that a zero-curvature plane tangent to $G // U$ must lift to a horizontal zero-curvature plane with respect to $((\cdot, \cdot))$. As in the case of metrics on G , if $((\cdot, \cdot))$ is induced from a bi-invariant metric on some Lie group L , then Tapp [26] implies that horizontal zero-curvature planes with respect to $((\cdot, \cdot))$ must project to zero-curvature planes in $G // U$. For our purposes this will always be true since we will consider only metrics as in Example (a) and Example (b).

We must determine what it means for a plane to be horizontal with respect to $((\cdot, \cdot))$ and the $\Delta G \times U$ action. Since each $\Delta G \times U$ orbit passes through some point of the form $(g, e) \in G \times G$, where e is the identity element of G , we may restrict our attention to such points.

Recall that $((\cdot, \cdot))$ is left-invariant. Therefore, letting \mathfrak{u} denote the Lie algebra of U , the vertical subspace at $(g, e) \in G \times G$ is given by

$$\mathcal{V}_g = \{(\text{Ad}_{g^{-1}} X - Y_1, X - Y_2) \mid X \in \mathfrak{g}, (Y_1, Y_2) \in \mathfrak{u}\}$$

after left-translation to $(e, e) \in G \times G$. Note that this is independent of the choice of left-invariant metric on $G \times G$.

Thus, with respect to $((\cdot, \cdot))$, the horizontal subspace at (g, e) is

$$(9) \quad \mathcal{H}_g = \{(\Omega_1^{-1}(-\text{Ad}_{g^{-1}} X), \Omega_2^{-1}(X)) \mid \langle X, \text{Ad}_g Y_1 - Y_2 \rangle_0 = 0 \forall (Y_1, Y_2) \in \mathfrak{u}\}$$

where Ω_1 and Ω_2 are the metric tensors relating the left-invariant metrics $\langle \cdot, \cdot \rangle_{K_1}$ and $\langle \cdot, \cdot \rangle_{K_2}$ respectively to a fixed bi-invariant metric $\langle \cdot, \cdot \rangle_0$ on G , ie $\langle X, Y \rangle_{K_i} = \langle X, \Omega_i(Y) \rangle_0$, $i = 1, 2$. We recall that the metric tensors in Example (a) and Example (b) are given by Φ_1 and Φ_2 respectively, as shown in (2) and (5).

In particular, (9) shows that a horizontal 2-plane σ in $(G \times G, ((,)))$ must project to a 2-plane on each factor, denoted by $\check{\sigma}_1$ and $\check{\sigma}_2$ respectively. Moreover, since $((,))$ is a product metric, $\text{sec}(\sigma) = 0$ if and only if $\text{sec}(\check{\sigma}_i) = 0, i = 1, 2$. Thus, for product metrics involving the metrics described by Example (a) and Example (b), we may apply conditions (3) and (6) respectively in order to determine when a horizontal plane σ has zero-curvature.

2 The Cayley numbers, G_2 and its Lie algebra

We recall without proof some well known facts about Cayley numbers, the Lie group G_2 and its Lie algebra. More details may be found in Gluck, Warner and Ziller [13] and Murakami [20].

We may write the Cayley numbers as $\mathbf{Ca} = \mathbb{H} + \mathbb{H}\ell$, where \mathbb{H} is the algebra of quaternions. Thus we have a natural orthonormal basis

$$\{e_0 = 1, e_1 = i, e_2 = j, e_3 = k, e_4 = \ell, e_5 = i\ell, e_6 = j\ell, e_7 = k\ell\}$$

for \mathbf{Ca} . Note that this description of \mathbf{Ca} differs slightly from that given in Murakami [20], and accounts for the difference which occurs in the description of the Lie algebra \mathfrak{g}_2 in Theorem 2.2. Multiplication in \mathbf{Ca} is nonassociative and defined via

$$(10) \quad (a + b\ell)(c + d\ell) = (ac - \bar{d}b) + (da + b\bar{c})\ell, \quad a, b, c, d \in \mathbb{H}.$$

Hence we have the following multiplication table, where the order of multiplication is given by (row)*(column):

	$e_1 = i$	$e_2 = j$	$e_3 = k$	$e_4 = \ell$	$e_5 = i\ell$	$e_6 = j\ell$	$e_7 = k\ell$
$e_1 = i$	-1	k	$-j$	$i\ell$	$-\ell$	$-k\ell$	$j\ell$
$e_2 = j$	$-k$	-1	i	$j\ell$	$k\ell$	$-\ell$	$-i\ell$
$e_3 = k$	j	$-i$	-1	$k\ell$	$-j\ell$	$i\ell$	$-\ell$
$e_4 = \ell$	$-i\ell$	$-j\ell$	$-k\ell$	-1	i	j	k
$e_5 = i\ell$	ℓ	$-k\ell$	$j\ell$	$-i$	-1	$-k$	j
$e_6 = j\ell$	$k\ell$	ℓ	$-i\ell$	$-j$	k	-1	$-i$
$e_7 = k\ell$	$-j\ell$	$i\ell$	ℓ	$-k$	$-j$	i	-1

Table 1: Multiplication table for \mathbf{Ca}

Recall that the Lie group G_2 is the automorphism group of $\mathbf{Ca} \cong \mathbb{R}^8$. In fact G_2 is a connected subgroup of $SO(7) \subset SO(8)$, where $SO(8)$ acts on $\mathbf{Ca} \cong \mathbb{R}^8$ by orthogonal transformations and $SO(7)$ is that subgroup consisting of elements which leave $e_0 = 1$

fixed. $SO(8)$ also contains two copies of $Spin(7)$ which are not conjugate in $SO(8)$, and G_2 is the intersection of these two subgroups.

As our eventual goal is to prove [Theorem A\(i\)](#) and [\(ii\)](#), it is useful to recall the fact that G_2 appears in the descriptions of some interesting homogeneous spaces. The following statements are well-known and follow from applications of the triality principle for $SO(8)$. More details may be found in, for example, Adams [\[1, Theorem 5.5\]](#), Murakami [\[20\]](#) and Jacobson [\[17, page 79\]](#).

Theorem 2.1

- (i) $Spin(7)/G_2 = \mathbf{S}^7$, which inherits constant positive curvature from the bi-invariant metric on $Spin(7)$. Moreover, the isotropy representation is transitive on the collection of pairs of orthogonal unit tangent vectors.
- (ii) $Spin(8)/G_2 = \mathbf{S}^7 \times \mathbf{S}^7$ and $SO(8)/G_2 = (\mathbf{S}^7 \times \mathbf{S}^7)/\mathbb{Z}_2$, where $\mathbb{Z}_2 = \{\pm(\text{id}, \text{id})\}$.
- (iii) $G_2/SU(3) = \mathbf{S}^6$.

We now turn our attention to the Lie algebra of G_2 . The proof of the following theorem follows exactly as in Murakami [\[20\]](#) except that we use the basis and multiplication conventions for \mathbf{Ca} as in [Table 1](#). Recall that $\mathfrak{so}(n) = \{A \in M_n(\mathbb{R}) \mid A^t = -A\}$.

Theorem 2.2 *The Lie algebra of G_2 , denoted by \mathfrak{g}_2 , consists of matrices $A = (a_{ij}) \in \mathfrak{so}(7)$ which satisfy $a_{ij} + a_{ji} = 0$ and the following equations:*

$$\begin{aligned} a_{23} + a_{45} + a_{76} &= 0 \\ a_{12} + a_{47} + a_{65} &= 0 \\ a_{13} + a_{64} + a_{75} &= 0 \\ a_{14} + a_{72} + a_{36} &= 0 \\ a_{15} + a_{26} + a_{37} &= 0 \\ a_{16} + a_{52} + a_{43} &= 0 \\ a_{17} + a_{24} + a_{53} &= 0 \end{aligned}$$

Hence $\mathfrak{g}_2 \subset \mathfrak{so}(7)$ is 14-dimensional and consists of matrices of the form:

$$(11) \quad \begin{pmatrix} 0 & x_1 + x_2 & y_1 + y_2 & x_3 + x_4 & y_3 + y_4 & x_5 + x_6 & y_5 + y_6 \\ -(x_1 + x_2) & 0 & \alpha_1 & -y_5 & x_5 & -y_3 & x_3 \\ -(y_1 + y_2) & -\alpha_1 & 0 & x_6 & y_6 & -x_4 & -y_4 \\ -(x_3 + x_4) & y_5 & -x_6 & 0 & \alpha_2 & y_1 & -x_1 \\ -(y_3 + y_4) & -x_5 & -y_6 & -\alpha_2 & 0 & x_2 & y_2 \\ -(x_5 + x_6) & y_3 & x_4 & -y_1 & -x_2 & 0 & \alpha_1 + \alpha_2 \\ -(y_5 + y_6) & -x_3 & y_4 & x_1 & -y_2 & -(\alpha_1 + \alpha_2) & 0 \end{pmatrix}$$

Recall that G_2 is a rank 2 Lie group. Therefore an examination of the elements (11) of \mathfrak{g}_2 reveals that the maximal torus of G_2 is given by:

$$(12) \quad T^2 = \left\{ \left(\begin{array}{ccc} 1 & & \\ & R(\theta) & \\ & & R(\varphi) \\ & & & R(\theta + \varphi) \end{array} \right) \mid R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \right\}$$

3 Free isometric actions on SO(8)

Consider the rank one symmetric pair $(G, K) = (SO(8), SO(7))$ where

$$SO(7) \hookrightarrow SO(8) \\ A \mapsto \begin{pmatrix} 1 & \\ & A \end{pmatrix}$$

with Lie algebras $\mathfrak{g}, \mathfrak{k}$ respectively. Let $\langle X, Y \rangle_0 = -\text{tr}(XY)$ be a bi-invariant metric on G . With respect to $\langle \cdot, \cdot \rangle_0$ we thus have $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$. As in (2) we define a left-invariant, right K -invariant metric $\langle \cdot, \cdot \rangle_1$ on G by

$$(13) \quad \langle X, Y \rangle_1 = \langle X, \Phi_1(Y) \rangle_0$$

where $\Phi_1(Y) = Y_{\mathfrak{p}} + \lambda_1 Y_{\mathfrak{k}}$, $\lambda_1 \in (0, 1)$. Recall that from Example (a) we know that a plane

$$\sigma = \text{Span}\{\Phi_1^{-1}(X), \Phi_1^{-1}(Y)\} \subset \mathfrak{g}$$

has zero-curvature with respect to $\langle \cdot, \cdot \rangle_1$ if and only if

$$(14) \quad 0 = [X, Y] = [X_{\mathfrak{p}}, Y_{\mathfrak{p}}] = [X_{\mathfrak{k}}, Y_{\mathfrak{k}}].$$

We now equip $G \times G$ with the product metric $\langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_0$.

Consider an isometric action of $U := \mathbf{S}^1 \times G_2 \subset K \times G$ on $SO(8)$ defined by

$$(15) \quad A \mapsto \tilde{R}(\theta) \cdot A \cdot g^{-1}$$

where $A \in SO(8)$, $g \in G_2$, and

$$(16) \quad \tilde{R}(\theta) = \begin{pmatrix} I_{2 \times 2} & & & \\ & R(p_1\theta) & & \\ & & R(p_2\theta) & \\ & & & R(p_3\theta) \end{pmatrix}, \quad R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

From (8) we know that $\Delta G \backslash G \times G / U \cong G // U$ whenever the biquotient action of U on G is free.

Lemma 3.1 $\Delta G \times U$ acts freely and isometrically on $(G \times G, \langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_0)$ if and only if (p_1, p_2, p_3) is equal to $(0, 0, 1)$ (up to sign and permutations of the p_i).

Proof Recall that conjugation of either factor of U by elements of G is a diffeomorphism, and that a biquotient action is free if and only if nontrivial elements in each factor are never conjugate to one another in G . Thus we need only show that nontrivial elements of S^1 and T^2 are never conjugate in G if and only if $(p_1, p_2, p_3) = (0, 0, 1)$ up to sign and permutations of the p_i , where T^2 is the maximal torus of G_2 described in (12). This amounts to investigating when the sets of 2×2 blocks on each side are equal up to conjugation by an element of the Weyl group of $SO(8)$. We recall that the Weyl group of $SO(2n)$ acts via permutations of the 2×2 blocks and changing an even number of signs, where by a change of sign we mean $R(\theta) \mapsto R(-\theta)$. A simple calculation then yields the result. \square

Note that there are many other free $S^1 \times G_2$ actions on G . For example, there is a free S^1 action on the left of G/G_2 by matrices of the form

$$(17) \quad \begin{pmatrix} R(\theta) & & & \\ & R(\theta) & & \\ & & R(\theta) & \\ & & & R(k\theta) \end{pmatrix}$$

where $(k, 3) = 1$. However, it is clear that only the action in Lemma 3.1 is isometric with respect to the metric $\langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_0$ on $G \times G$.

It follows immediately from the long exact homotopy sequence for fibrations that a biquotient $Spin(8) // (S^1 \times G_2) = S^1 \backslash (S^7 \times S^7)$ must be simply connected. By the lifting criterion for covering spaces the action by U on $SO(8)$ described above lifts to some action by $S^1 \times G_2$ on $Spin(8)$. Therefore, together with Theorem 2.1, one might expect that the resulting simply connected biquotient $Spin(8) // (S^1 \times G_2) = S^1 \backslash (S^7 \times S^7)$ is a nontrivial finite cover of $SO(8) // (S^1 \times G_2)$. In fact the lemma below will demonstrate that this covering map is a diffeomorphism.

Lemma 3.2 $M^{13} := SO(8) // (S^1 \times G_2)$ is simply connected and hence a quotient of $S^7 \times S^7$ by an S^1 action.

Proof Consider a general embedding

$$S^1_q \hookrightarrow SO(8)$$

$$R(\theta) \mapsto \begin{pmatrix} R(q_1\theta) & & & \\ & R(q_2\theta) & & \\ & & R(q_3\theta) & \\ & & & R(q_4\theta) \end{pmatrix}$$

where $q = (q_1, q_2, q_3, q_4) \in \mathbb{Z}^4$ and where $R(u) \in \text{SO}(2)$. The long exact homotopy sequence for the fibration $\mathbf{S}_q^1 \times G_2 \rightarrow \text{SO}(8) \rightarrow \text{SO}(8) // \mathbf{S}_q^1 \times G_2$ yields

$$\dots \rightarrow \pi_1(\mathbf{S}_q^1 \times G_2) = \mathbb{Z} \rightarrow \pi_1(\text{SO}(8)) = \mathbb{Z}_2 \rightarrow \pi_1(\text{SO}(8) // \mathbf{S}_q^1 \times G_2) \rightarrow 0.$$

Thus to obtain the desired result we need only show that the map $\mathbb{Z} \rightarrow \mathbb{Z}_2$ is surjective.

Recall that the homomorphism $\iota_*: \pi_1(\mathbf{S}_q^1) \rightarrow \pi_1(\text{SO}(n))$ is determined by the weights $q = (q_1, \dots, q_m)$, $m = \lfloor n/2 \rfloor$, of the embedding, namely $\iota_*(1) = \sum q_i \pmod 2$. Therefore ι_* is onto exactly when $\sum q_i$ is odd. In our case we have $q = (0, 0, 0, 1)$, and so ι_* is a surjection. \square

Notice that the action of U on $\text{SO}(8)$ given in Lemma 3.1 may be enlarged to an isometric action by $\text{SO}(3) \times G_2$, and the resulting biquotient we call N^{11} . Now recall that for all n we have a 2-fold cover $\text{Spin}(n) \rightarrow \text{SO}(n)$ with $\pi_1(\text{Spin}(n)) = 0$ and $\pi_1(\text{SO}(n)) = \mathbb{Z}_2$. Thus, by the lifting criterion for covering spaces, the inclusion $\text{SO}(3) \hookrightarrow \text{SO}(8)$ must lift to $\text{Spin}(3) = \mathbf{S}^3 \hookrightarrow \text{Spin}(8)$. As in the case of $U = \mathbf{S}^1 \times G_2$ above we show that $N^{11} = \text{SO}(8) // (\text{SO}(3) \times G_2)$ is simply connected and hence diffeomorphic to $\text{Spin}(8) // \mathbf{S}^3 \times G_2 = \mathbf{S}^3 \setminus (\mathbf{S}^7 \times \mathbf{S}^7)$.

Lemma 3.3 $N^{11} = \text{SO}(8) // (\text{SO}(3) \times G_2)$ is simply connected and hence a quotient of $\mathbf{S}^7 \times \mathbf{S}^7$ by an \mathbf{S}^3 action.

Proof Consider the chain of embeddings $j \circ i: \mathbf{S}^1 = \text{SO}(2) \hookrightarrow \text{SO}(3) \hookrightarrow \text{SO}(8)$ given by enlarging \mathbf{S}^1 above to an $\text{SO}(3)$ in $\text{SO}(8)$. We thus have an induced homomorphism on fundamental groups $(j \circ i)_* = j_* \circ i_*: \mathbb{Z} \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$. But i_* and $(j \circ i)_*$ are simply the homomorphism ι_* from Lemma 3.2. Hence $i_*(1) = 1 \pmod 2$ and $(j \circ i)_*(1) = 1 \pmod 2$. This implies $j_*(1) = 1 \pmod 2$ and therefore j_* is a surjection. An examination of the long exact homotopy sequence of the fibration $\text{SO}(3) \times G_2 \rightarrow \text{SO}(8) \rightarrow N^{11}$ yields the result. \square

Recall that $G_2 \subset \text{Spin}(7)' \subset \text{SO}(8)$, where the second inclusion is via the spin embedding. By our choice of metric on $G \times G$ we may therefore enlarge the actions of $\mathbf{S}^1 \times G_2$ and $\text{SO}(3) \times G_2$ to isometric actions by $\mathbf{S}^1 \times \text{Spin}(7)'$ and $\text{SO}(3) \times \text{Spin}(7)'$ respectively. It was shown in Eschenburg [9, Theorem 75, Table 101] that these actions are free and the resulting biquotients are \mathbf{CP}^3 and \mathbf{S}^4 respectively. It follows immediately that there are fibre bundles

$$\begin{aligned} \mathbf{S}^7 &= \text{Spin}(7)' / G_2 \rightarrow M^{13} \rightarrow \mathbf{CP}^3 \\ \mathbf{S}^7 &= \text{Spin}(7)' / G_2 \rightarrow N^{11} \rightarrow \mathbf{S}^4 \end{aligned}$$

for which the projections are Riemannian submersions.

4 Almost positive curvature on M^{13} and N^{11}

We are now in a position to prove the curvature statements [Theorem A\(i\)](#). We will concentrate on the circle quotient of $S^7 \times S^7$, namely

$$M^{13} = SO(8) // (S^1 \times G_2) = G // U,$$

since the other case follows trivially.

Consider the inclusions $G = SO(8) \supset K = SO(7) \supset G_2$. With respect to the bi-invariant metric $\langle X, Y \rangle_0 = -\text{tr}(XY)$ on G we have

$$\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k} \quad \text{and} \quad \mathfrak{k} = \mathfrak{m} \oplus \mathfrak{g}_2$$

where

$$(18) \quad \mathfrak{p} = \left\{ \left(\begin{array}{c|c} 0 & -w^t \\ \hline w & 0 \end{array} \right) \mid w \in \mathbb{R}^7 \right\}$$

and, by [\(11\)](#):

$$(19) \quad \mathfrak{m} = \left\{ \left(\begin{array}{c|cccccccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & v_1 & v_2 & v_3 & v_4 & v_5 & v_6 \\ 0 & -v_1 & 0 & v_7 & v_6 & -v_5 & v_4 & -v_3 \\ 0 & -v_2 & -v_7 & 0 & -v_5 & -v_6 & v_3 & v_4 \\ 0 & -v_3 & -v_6 & v_5 & 0 & v_7 & -v_2 & v_1 \\ 0 & -v_4 & v_5 & v_6 & -v_7 & 0 & -v_1 & -v_2 \\ 0 & -v_5 & -v_4 & -v_3 & v_2 & v_1 & 0 & -v_7 \\ 0 & -v_6 & v_3 & -v_4 & -v_1 & v_2 & v_7 & 0 \end{array} \right) \mid v_i \in \mathbb{R} \right\}$$

Lemma 4.1 *Let $W_i \in \mathfrak{p}$, $i = 1, 2$, and $V_j \in \mathfrak{m}$, $j = 1, 2$, be orthonormal vectors with respect to $\langle \cdot, \cdot \rangle_0$. Then $\text{rank}(W_i) = 2$, $\text{rank}(V_j) = 6$, $\text{rank}([W_1, W_2]) = 2$ and $\text{rank}([V_1, V_2]) = 6$.*

Proof Suppose P, Q and Z are real $n \times n$ matrices. It is well-known (and not difficult to prove using the rank-nullity theorem) that:

- (i) If $\text{rank}(P) = n$, then $\text{rank}(PZ) = \text{rank}(Z)$.
- (ii) If $\text{rank}(Q) = n$, then $\text{rank}(ZQ) = \text{rank}(Z)$.

In particular, if P and Q are invertible matrices then, for $Z_0 = PZQ$, it follows that $\text{rank}(Z) = \text{rank}(Z_0)$.

Choose arbitrary pairs of orthonormal vectors in \mathfrak{p} and \mathfrak{m} and show that the conclusions of the lemma hold. The result now follows from the facts that the isotropy representation

of $G_2 \subset SO(8)$ splits irreducibly as $\mathfrak{p} \oplus \mathfrak{m}$ and, moreover, is transitive on the sets of pairs of orthonormal vectors in \mathfrak{p} and \mathfrak{m} respectively, ie G_2 acts transitively on the unit tangent bundle to S^6 . (Recall [Theorem 2.1\(i\)](#).) \square

Let $U = S^1 \times G_2 \subset K \times G$ be as in [Lemma 3.1](#). Thus, equipping $G \times G$ with the product metric $\langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_0$ as before, we may induce a metric on $G//U$ via the diffeomorphism

$$\Delta G \setminus G \times G / U \longrightarrow G // U.$$

As discussed in [Section 1](#), we may restrict our attention to points of the form $(A, I) \in G \times G$. Let $E_{78} \in \mathfrak{g}$ denote the vector spanning the Lie algebra of the S^1 factor of U . By [\(9\)](#) the horizontal subspace at (A, I) with respect to $\langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_0$ is given by

$$\mathcal{H}_A = \{(-\Phi_1^{-1}(\text{Ad}_{A^t} W), W) \mid W_{\mathfrak{g}_2} = 0, \langle W, \text{Ad}_A E_{78} \rangle_0 = 0\}.$$

Suppose that

$$\sigma = \text{Span}\{(-\Phi_1^{-1}(\text{Ad}_{A^t} X), X), (-\Phi_1^{-1}(\text{Ad}_{A^t} Y), Y)\} \subset \mathfrak{g} \oplus \mathfrak{g}$$

is a horizontal zero-curvature plane at $(A, I) \in G \times G$. Since we have equipped $G \times G$ with the product metric $\langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_0$, our discussion in [Section 1](#) shows that σ must project to zero-curvature planes $\check{\sigma}_i$, $i = 1, 2$, on each factor. Considering $\check{\sigma}_2$ we find:

Lemma 4.2 *Suppose $X, Y \in \mathfrak{g}$ are linearly independent vectors such that $X_{\mathfrak{g}_2} = Y_{\mathfrak{g}_2} = 0$ and $[X, Y] = 0$. Then it may be assumed without loss of generality that $X \in \mathfrak{p}$ and $Y \in \mathfrak{m}$.*

Proof Since $[X, Y] = 0$ and (G, K) is a symmetric pair, it follows that $[X_p, Y_p] = -[X_m, Y_m]$. Suppose that $[X_m, Y_m] \neq 0$. Then, by [Lemma 4.1](#), $[X_m, Y_m]$ is a matrix of rank 6. But $[X_p, Y_p]$ has rank no more than 2, so we have a contradiction. Therefore $[X_p, Y_p] = [X_m, Y_m] = 0$.

The equality $[X_p, Y_p] = 0$ implies that, since (G, K) is a rank one symmetric pair, we may assume $Y_p = 0$ without loss of generality. Hence $X \in \mathfrak{p} \oplus \mathfrak{m}$ and $Y \in \mathfrak{m}$. On the other hand, by [Theorem 2.1\(i\)](#) there are no independent commuting vectors in \mathfrak{m} . Then, without loss of generality, $X \in \mathfrak{p}$ and $Y \in \mathfrak{m}$. \square

The horizontal zero-curvature plane σ is thus determined by

$$(20) \quad X = \left(\begin{array}{c|c} 0 & -w^t \\ \hline w & 0 \end{array} \right) \in \mathfrak{p}, \quad Y = \left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & (v_{ij}) \end{array} \right) \in \mathfrak{m}$$

with $[X, Y] = 0$, where $w \in \mathbb{R}^7$ and $(v_{ij}) = (v_{ij} \mid 2 \leq i, j \leq 8)$.

By Lemma 4.1 we know that Y has rank 6, that is, the column space of Y is a six-dimensional subspace of \mathbb{R}^7 . The condition $[X, Y] = 0$ is equivalent to $w \in \mathbb{R}^7$ being perpendicular to each of the columns of Y . Therefore w , and hence X , is uniquely determined up to scaling by Y . Let $\psi: \mathfrak{m} \rightarrow \mathfrak{p}$ be the (unique) map assigning $X \in \mathfrak{p}$ to $Y \in \mathfrak{m}$ such that, by abusing the notation of (18) and (19), $X = \psi(Y)$ is given by

$$(21) \quad w^t = \psi(v_1, \dots, v_7) = (v_7, -v_2, v_1, -v_4, v_3, v_6, -v_5).$$

It is easy to check that $[\psi(Y), Y] = 0$, and ψ is clearly a linear isomorphism. Moreover, ψ is G_2 -equivariant since $\text{Ad}_g[X, Y] = [\text{Ad}_g X, \text{Ad}_g Y]$, for all $g \in G_2$, and by uniqueness. Therefore every horizontal zero-curvature plane σ is determined by a pair $(X, Y) = (\psi(Y), Y)$, with $X \in \mathfrak{p}$, $Y \in \mathfrak{m}$.

Given $[X, Y] = 0$, the conditions in (14) imply that $\check{\sigma}_1$ has zero curvature if and only if $[(\text{Ad}_{A^t} X)_\mathfrak{p}, (\text{Ad}_{A^t} Y)_\mathfrak{p}] = 0$. But (G, K) is a rank one symmetric pair and thus $\check{\sigma}_1$ has zero curvature if and only if $(\text{Ad}_{A^t} X)_\mathfrak{p}, (\text{Ad}_{A^t} Y)_\mathfrak{p}$ are linearly dependent, that is, if and only if either

$$(\text{Ad}_{A^t} X)_\mathfrak{p} = 0 \quad \text{or} \quad (\text{Ad}_{A^t} Y)_\mathfrak{p} = 0 \quad \text{or} \quad (\text{Ad}_{A^t} X)_\mathfrak{p} = s(\text{Ad}_{A^t} Y)_\mathfrak{p}$$

for some $s \in \mathbb{R} - \{0\}$.

Suppose that $(\text{Ad}_{A^t} X)_\mathfrak{p} = s(\text{Ad}_{A^t} Y)_\mathfrak{p}$, for some $s \in \mathbb{R} - \{0\}$. Then $\text{Ad}_{A^t}(X - sY) \perp \mathfrak{p}$. Since elements of $\mathfrak{p}^\perp = \mathfrak{k} = \mathfrak{so}(7)$ have vanishing determinant, it follows from the discussion on ranks in the proof of Lemma 4.1 that $\text{rank}(X - sY) = \text{rank Ad}_{A^t}(X - sY) \leq 6$. But X and Y describe a horizontal zero-curvature plane, so the vector $w \in \mathbb{R}^7$ determining X is orthogonal to the columns of sY , which itself has rank 6. Hence $\text{rank}(X - sY) = 7$, which is a contradiction. Therefore either $(\text{Ad}_{A^t} X)_\mathfrak{p} = 0$ or $(\text{Ad}_{A^t} Y)_\mathfrak{p} = 0$.

Before we continue, we recall some simple facts about $A \in \text{SO}(8)$. We may write A as

$$A = \left(\begin{array}{c|ccc} a_{11} & a_{12} & \cdots & a_{18} \\ a_{21} & & & \\ \vdots & & \tilde{A} & \\ a_{81} & & & \end{array} \right)$$

where \tilde{A} is a 7×7 matrix. Since $AA^t = I$, one can easily derive the identities

$$(22) \quad a_{11}^2 + \sum |a_{1k}|^2 = 1$$

$$(23) \quad a_{11}(a_{k1}) + \tilde{A}(a_{1k})^t = 0$$

$$(24) \quad (a_{k1})(a_{k1})^t + \tilde{A}\tilde{A}^t = I_{7 \times 7}$$

where (a_{1k}) and (a_{k1}) are row and column vectors respectively (with $2 \leq k \leq 8$).

Consider $X \in \mathfrak{p}$ as in (20) and suppose that $(\text{Ad}_{A^t} X)_{\mathfrak{p}} = 0$. A simple computation shows that this is equivalent to

$$a_{11} w^t \tilde{A} = (a_{k1})^t w (a_{1k}) = \langle (a_{k1}), w \rangle (a_{1k}).$$

If we multiply both sides by $(a_{1k})^t$, then the identity (23) above yields

$$\begin{aligned} -a_{11}^2 \langle (a_{k1}), w \rangle &= -a_{11}^2 w^t (a_{k1}) \\ &= \langle (a_{k1}), w \rangle (a_{1k})(a_{1k})^t \\ &= \langle (a_{k1}), w \rangle \sum |a_{1k}|^2. \end{aligned}$$

By (22), this reduces to $(a_{k1})^t w = \langle (a_{k1}), w \rangle = 0$. Hence $a_{11} w^t \tilde{A} = 0$, ie either $a_{11} = 0$ or $w^t \tilde{A} = 0$. However, if $w^t \tilde{A} = 0$, then $\text{Ad}_{A^t} X = 0$. Since $X \neq 0$, this gives a contradiction. Therefore the condition $(\text{Ad}_{A^t} X)_{\mathfrak{p}} = 0$ is satisfied if and only if $a_{11} = 0$ and $\langle (a_{21}, \dots, a_{81})^t, w \rangle = 0$. It is clear that the set of such w is six-dimensional. For reasons of dimension we will therefore always be able to find a pair (X, Y) describing a zero-curvature plane, since we need only ensure that $\text{Ad}_{A^t} X, \text{Ad}_{A^t} Y \perp \mathbf{S}^1$.

On the other hand, consider $Y \in \mathfrak{m}$ as in (20) such that $(\text{Ad}_{A^t} Y)_{\mathfrak{p}} = 0$. It is again simple to show that this is equivalent to

$$(a_{k1})^t (v_{ij}) \tilde{A} = 0.$$

If we multiply both sides by \tilde{A}^t , then (24) gives

$$\begin{aligned} 0 &= (a_{k1})^t (v_{ij}) (I_{7 \times 7} - (a_{k1})(a_{k1})^t) \\ &= (a_{k1})^t (v_{ij}) - ((a_{k1})^t (v_{ij})(a_{k1}))(a_{k1})^t \\ &= (a_{k1})^t (v_{ij}) \end{aligned}$$

where the last equality follows from $(a_{k1})^t (v_{ij})(a_{k1}) = 0$, since all entries on the diagonal of $\text{Ad}_{A^t} Y \in \mathfrak{so}(8)$ must be zero.

Let $X = \psi(Y) \in \mathfrak{p}$ be given by $w \in \mathbb{R}^7$. Now $(a_{k1})^t (v_{ij}) = 0$ implies that the vector $(a_{21}, \dots, a_{81})^t \in \mathbb{R}^7$ is perpendicular to each column of the rank 6 matrix (v_{ij}) . Hence either $a_{k1} = 0$, for all $2 \leq k \leq 8$, or $w = (a_{21}, \dots, a_{81})^t$.

If $a_{k1} = 0$, for all $2 \leq k \leq 8$, then by (22), (23) and (24) $A \in \text{O}(7) \subset \text{SO}(8)$. Conversely, given $A \in \text{O}(7)$, it is clear that Ad_{A^t} preserves the orthogonal decomposition $\mathfrak{p} \oplus \mathfrak{k}$. Thus $\text{Ad}_{A^t} X$, with $X \in \mathfrak{p}$, will always be orthogonal to \mathbf{S}^1 and, for dimension reasons, there will always be a $Y \in \mathfrak{m}$ such that $\text{Ad}_{A^t} Y \perp \mathbf{S}^1$. Therefore it is always possible to find a pair (X, Y) spanning a horizontal zero-curvature plane.

Finally, in the case $w = (a_{21}, \dots, a_{81})^t$, there is, up to scaling, a unique pair (X, Y) determined by the coordinates of A . Clearly X is determined by w while, by (21), Y is given by

$$(v_1, \dots, v_7) = (a_{41}, -a_{31}, a_{61}, -a_{51}, -a_{81}, a_{71}, a_{21}).$$

It is easy to see that the condition $\langle X, \text{Ad}_A E_{78} \rangle_0 = 0$ is given by the algebraic expression:

$$(25) \quad \sum_{k=2}^8 (a_{18}a_{k7} - a_{17}a_{k8})a_{k1} = 0$$

A straightforward computation shows that condition $\langle Y, \text{Ad}_A E_{78} \rangle_0 = 0$ is given by

$$(26) \quad \begin{aligned} 0 = & (m_{23} + m_{58} - m_{67})a_{41} - (m_{24} - m_{57} - m_{68})a_{31} \\ & + (m_{25} - m_{38} + m_{47})a_{61} - (m_{26} + m_{37} + m_{48})a_{51} \\ & - (m_{27} - m_{36} - m_{45})a_{81} + (m_{28} + m_{35} - m_{46})a_{71} \\ & + (m_{34} + m_{56} - m_{78})a_{21} \end{aligned}$$

where $m_{k\ell} = a_{k8}a_{\ell 7} - a_{k7}a_{\ell 8}$. It is clear that if both (25) and (26) hold, then there is a horizontal zero-curvature plane at (A, I) .

We have shown that a horizontal zero-curvature plane occurs at (A, I) if and only if one of the following conditions is satisfied:

- (a) $a_{11} = 0$
- (b) $A \in \text{O}(7) \subset \text{SO}(8)$
- (c) Equations (25) and (26) hold.

The locus of such points is clearly lower dimensional. Moreover, extending the U action to an action by $\text{SO}(3) \times G_2$ increases the number of conditions which must be satisfied in order for a zero-curvature plane to be horizontal. Theorem A(i) now follows immediately.

5 Generalised Eschenburg spaces

Consider the rank one symmetric pairs $(G, K) = (U(n + 1), U(1)U(n))$ and $(K, H) = (U(1)U(n), U(1)U(1)U(n - 1))$ where $n \geq 2$ and the inclusions $K \hookrightarrow G$ and $H \hookrightarrow K$

are given by

$$(z, B) \mapsto \begin{pmatrix} z & \\ & B \end{pmatrix}, \quad z \in \mathbf{U}(1), \quad B \in \mathbf{U}(n)$$

and
$$(z, w, C) \mapsto \left(z, \begin{pmatrix} w & \\ & C \end{pmatrix} \right), \quad z, w \in \mathbf{U}(1), \quad C \in \mathbf{U}(n-1)$$

respectively. Let $\mathfrak{g}, \mathfrak{k}$ and \mathfrak{h} be the Lie algebras of G, K and H respectively. Let $\langle X, Y \rangle_0 = -\operatorname{Re} \operatorname{tr}(XY)$ be a bi-invariant metric on G . With respect to $\langle \cdot, \cdot \rangle_0$ we thus have the orthogonal decompositions $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ and $\mathfrak{k} = \mathfrak{m} \oplus \mathfrak{h}$, where:

$$\mathfrak{p} = \left\{ \left(\begin{array}{c|c} 0 & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \mid x = \begin{pmatrix} x_2 \\ \vdots \\ x_{n+1} \end{pmatrix} \in \mathbb{C}^n \right\}$$

and
$$\mathfrak{m} = \left\{ \left(\begin{array}{c|c|c} 0 & & \\ \hline & 0 & -\bar{y}^t \\ \hline & y & 0 \end{array} \right) \mid y = \begin{pmatrix} y_3 \\ \vdots \\ y_{n+1} \end{pmatrix} \in \mathbb{C}^{n-1} \right\}$$

As in [Example \(a\)](#) and [Example \(b\)](#) we define a left-invariant, right K -invariant metric $\langle \cdot, \cdot \rangle_1$ on G by

$$(27) \quad \langle X, Y \rangle_1 = \langle X, \Phi_1(Y) \rangle_0$$

where $\Phi_1(Y) = Y_p + \lambda_1 Y_k$, $\lambda_1 \in (0, 1)$, and a left-invariant, right H -invariant metric $\langle \cdot, \cdot \rangle_2$ on G via

$$(28) \quad \langle X, Y \rangle_2 = \langle X, \Psi(Y) \rangle_1 = \langle X, \Phi_2(Y) \rangle_0$$

where $\Phi_2(Y) = Y_p + \lambda_1 Y_m + \lambda_1 \lambda_2 Y_h$, $\lambda_2 \in (0, 1)$, and $\Psi(Y) = \Phi_1^{-1} \Phi_2(Y) = Y_p + Y_m + \lambda_2 Y_h$.

Equip $G \times G$ with the left-invariant, right $(K \times H)$ -invariant product metric $\langle \cdot, \cdot \rangle_1 \oplus \langle \cdot, \cdot \rangle_2$.

Consider the subgroup $L_{p,q} \subset K \times H$ defined by

$$L_{p,q} = \{ (\operatorname{diag}(z^{p_1}, \dots, z^{p_{n+1}}), \operatorname{diag}(z^{q_1}, z^{q_2}, B)) \mid z \in \mathbf{S}^1, B \in \mathbf{U}(n-1) \}$$

where $p_1, \dots, p_{n+1}, q_1, q_2 \in \mathbb{Z}$. $L_{p,q}$ acts on G via

$$\begin{aligned} G &\longrightarrow G \\ A &\longmapsto \operatorname{diag}(z^{p_1}, \dots, z^{p_{n+1}}) A \operatorname{diag}(\bar{z}^{q_1}, \bar{z}^{q_2}, B^{-1}) \end{aligned}$$

where $z \in U(1)$ and $B \in U(n - 1)$. It is not difficult to show that this action is free if and only if

$$(29) \quad (p_{\sigma(1)} - q_1, p_{\sigma(2)} - q_2) = 1 \quad \text{for all } \sigma \in S_{n+1}.$$

We denote the resulting biquotients $G//L_{p,q}$ by $E_{p,q}^{4n-1}$ and remark that $n = 2$ gives the usual Eschenburg spaces (see Eschenburg [9, Section 41]).

Recall the canonical diffeomorphism

$$E_{p,q}^{4n-1} = G//L_{p,q} \cong \Delta G \backslash G \times G/L_{p,q}$$

given in (8). Now, since $L_{p,q} \subset K \times H$, there is a metric on $E_{p,q}^{4n-1}$ induced from the product metric on $G \times G$.

From (9) it is easy to show that the horizontal subspace at a point $(A, I) \in G \times G$ is given by

$$(30) \quad \mathcal{H}_A = \{(-\Phi_1^{-1}(\text{Ad}_{A^*} W), \Phi_2^{-1}(W)) \mid W_{\mathfrak{u}(n-1)} = 0, \langle W, \text{Ad}_A P - Q \rangle_0 = 0\}$$

where $A^* = \bar{A}^t$, $P = \text{diag}(ip_1, \dots, ip_{n+1})$, $Q = \text{diag}(iq_1, iq_2, 0, \dots, 0)$, and $\mathfrak{h} = \mathfrak{u}(1) \oplus \mathfrak{u}(1) \oplus \mathfrak{u}(n - 1)$ as before.

Proposition 5.1 *Suppose that*

$$\sigma = \text{Span}\{(-\Phi_1^{-1}(\text{Ad}_{A^*} \tilde{X}), \Phi_2^{-1}(\tilde{X})), (-\Phi_1^{-1}(\text{Ad}_{A^*} \tilde{Y}), \Phi_2^{-1}(\tilde{Y}))\}$$

is a horizontal zero-curvature plane at $(A, I) \in G \times G$. Then $X = \Phi_1^{-1}(\tilde{X})$ and $Y = \Phi_1^{-1}(\tilde{Y})$ can be written in one of the following forms:

- (i) $X \in \mathfrak{g}$ and $Y = \text{diag}(i, 0, \dots, 0)$
- (ii) $X \in \mathfrak{p} \oplus \mathfrak{h}$ and $Y = \text{diag}(0, i, 0, \dots, 0)$
- (iii) $X \in \mathfrak{p} \oplus \mathfrak{h}$ and $Y = \text{diag}(i, i, 0, \dots, 0)$
- (iv)

$$X = \left(\begin{array}{c|c} 0 & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \quad \text{and} \quad Y = \left(\begin{array}{c|c|c} i & & \\ \hline & i\beta & -\bar{y}^t \\ \hline & y & 0 \end{array} \right)$$

where $x_2 \neq 0$, $\beta = 1 - \sum_{j=3}^{n+1} |y_j|^2$, and $x_j = -ix_2 y_j$ for $j = 3, \dots, n + 1$

$$(v) \quad X = \left(\begin{array}{c|c} i\alpha & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \quad \text{and} \quad Y = \left(\begin{array}{c|c|c} 0 & & \\ \hline & i\beta & -\bar{y}^t \\ \hline & y & 0 \end{array} \right)$$

where $x = (0, x_3, \dots, x_{n+1})^t \neq 0 \in \mathbb{C}^n$ and $\sum_{j=3}^{n+1} x_j \bar{y}_j = 0$.

Proof From the discussion in Section 1 we know that the projections $\check{\sigma}_i, i = 1, 2$, onto the first and second factor must be two-dimensional zero-curvature planes with respect to $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ respectively.

Consider

$$\check{\sigma}_2 = \text{Span}\{\Phi_2^{-1}(\tilde{X}), \Phi_2^{-1}(\tilde{Y})\} = \text{Span}\{\Psi^{-1}(X), \Psi^{-1}(Y)\}$$

where $\Psi = \Phi_1^{-1}\Phi_2$. $\check{\sigma}_2$ has zero-curvature with respect to $\langle \cdot, \cdot \rangle_2$ if and only if the equalities in (6) hold, namely if and only if

$$0 = [X, Y] = [X_{\mathfrak{k}}, Y_{\mathfrak{k}}] = [X_{\mathfrak{p}}, Y_{\mathfrak{p}}] = [X_{\mathfrak{m}}, Y_{\mathfrak{m}}] = [X_{\mathfrak{h}}, Y_{\mathfrak{h}}].$$

Since (G, K) is a rank-one symmetric pair, $[X_{\mathfrak{p}}, Y_{\mathfrak{p}}] = 0$ if and only if $X_{\mathfrak{p}}$ and $Y_{\mathfrak{p}}$ are linearly dependent. Without loss of generality we may assume that $Y_{\mathfrak{p}} = 0$. Similarly, (K, H) being a rank-one symmetric pair implies that $[X_{\mathfrak{m}}, Y_{\mathfrak{m}}] = 0$ if and only if $X_{\mathfrak{m}}$ and $Y_{\mathfrak{m}}$ are linearly dependent. Without loss of generality we may assume that either $X_{\mathfrak{m}} = 0$ or $Y_{\mathfrak{m}} = 0$. Thus we have two possibilities:

$$(31) \quad X = X_{\mathfrak{p}} + X_{\mathfrak{m}} + X_{\mathfrak{h}} \quad \text{and} \quad Y = Y_{\mathfrak{h}}$$

$$(32) \quad X = X_{\mathfrak{p}} + X_{\mathfrak{h}} \quad \text{and} \quad Y = Y_{\mathfrak{m}} + Y_{\mathfrak{h}}.$$

Since σ is horizontal and Φ_1 simply scales $\mathfrak{k} = \mathfrak{m} \oplus \mathfrak{h}$ by $\lambda_1 \in (0, 1)$, then we must have $X_{\mathfrak{u}(n-1)} = Y_{\mathfrak{u}(n-1)} = 0$, where $\mathfrak{h} = \mathfrak{u}(1) \oplus \mathfrak{u}(1) \oplus \mathfrak{u}(n-1)$. Therefore in both cases above we have

$$X_{\mathfrak{h}} = \text{diag}(ia, ib, 0, \dots, 0), \quad Y_{\mathfrak{h}} = \text{diag}(ic, id, 0, \dots, 0), \quad \text{some } a, b, c, d \in \mathbb{R}.$$

Clearly $[X_{\mathfrak{h}}, Y_{\mathfrak{h}}] = 0$. Then $[X_{\mathfrak{k}}, Y_{\mathfrak{k}}] = [X_{\mathfrak{m}}, Y_{\mathfrak{h}}] + [X_{\mathfrak{h}}, Y_{\mathfrak{m}}]$. In the case of (31) our zero-curvature condition is thus $0 = [X_{\mathfrak{m}}, Y_{\mathfrak{h}}]$, while for case (32) we have $0 = [X_{\mathfrak{h}}, Y_{\mathfrak{m}}]$.

Consider general vectors $Z = \text{diag}(i\alpha, i\beta, 0, \dots, 0) \in \mathfrak{h}$ and $W \in \mathfrak{m}$. Then $0 = [Z, W]$ if and only if either $\beta = 0$ or $W = 0$. Applying this to case (31) yields (after rescaling) two cases:

$$(33) \quad X = X_{\mathfrak{p}} + X_{\mathfrak{m}} + X_{\mathfrak{h}} \quad \text{and} \quad Y = \text{diag}(i, 0, \dots, 0)$$

$$(34) \quad X = X_{\mathfrak{p}} + X_{\mathfrak{h}} \quad \text{and} \quad Y = Y_{\mathfrak{h}}.$$

On the other hand, case (32) yields the added possibility

$$(35) \quad X = \left(\begin{array}{c|c} i\alpha & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \in \mathfrak{p} \oplus \mathfrak{h} \quad \text{and} \quad Y = Y_m + Y_h.$$

As (33) is already of the form (i) in the proposition, we concentrate on cases (34) and (35).

The only zero-curvature condition remaining to us is $[X, Y] = 0$. Since $Y_p = 0$, this is equivalent to $[X_p, Y_{\mathfrak{k}}] = 0$. Consider the general vectors

$$U = \left(\begin{array}{c|c} 0 & -\bar{u}^t \\ \hline u & 0 \end{array} \right) \in \mathfrak{p} \quad \text{and} \quad V = \left(\begin{array}{c|c|c} i\gamma & & \\ \hline & i\delta & -\bar{v}^t \\ \hline & v & 0 \end{array} \right) \in \mathfrak{k}$$

where $u = (u_2, \dots, u_{n+1})^t \in \mathbb{C}^n$, $v = (v_3, \dots, v_{n+1})^t \in \mathbb{C}^{n-1}$, and $\gamma, \delta \in \mathbb{R}$. Then

$$(36) \quad [U, V] = 0 \iff i(\gamma - \delta)u_2 + \sum_{j=3}^{n+1} u_j \bar{v}_j = 0 \quad \text{and} \quad \gamma u_j = -i u_2 v_j, \quad j = 3, \dots, n+1.$$

Suppose $u_2 = 0$. Then (36) becomes $\gamma u_j = 0$, $j = 3, \dots, n+1$, and $\sum_{j=3}^{n+1} u_j \bar{v}_j = 0$. This is satisfied if and only if one of the two following possibilities holds:

$$(37) \quad u_j = 0 \quad \text{for all } j = 2, \dots, n+1,$$

or

$$(38) \quad u_2 = 0, \quad \gamma = 0, \quad \sum_{j=3}^{n+1} u_j \bar{v}_j = 0.$$

On the other hand, if we assume $u_2 \neq 0$ then (36) becomes

$$(39) \quad u_2 \neq 0, \quad \gamma u_j = -i u_2 v_j, \quad j = 3, \dots, n+1, \quad \delta = \frac{\gamma}{|u_2|^2} \left(|u_2|^2 - \sum_{j=3}^{n+1} |u_j|^2 \right).$$

Now, if we apply conditions (37), (38) and (39) to case (34) we arrive at one of the following (after rescaling where appropriate):

$$(40) \quad X = \text{diag}(ia, ib, 0, \dots, 0) \quad \text{and} \quad Y = \text{diag}(ic, id, 0, \dots, 0)$$

$$(41) \quad X \in \mathfrak{p} \oplus \mathfrak{h} \quad \text{and} \quad Y = \text{diag}(0, i, 0, \dots, 0)$$

$$(42) \quad X \in \mathfrak{p} \oplus \mathfrak{h} \quad \text{and} \quad Y = \text{diag}(i, i, 0, \dots, 0).$$

Since X and Y must span a two-plane, it is clear that $\text{diag}(i, 0, \dots, 0)$ must lie in the plane spanned by the X and Y given in (40). Therefore zero-curvature planes described by (34) fall into one of the classes given by (i), (ii) and (iii) of the proposition.

For case (35) conditions (37), (38) and (39) imply that X and Y must have one of the following forms (after rescaling):

$$(43) \quad X = \text{diag}(i, 0, \dots, 0) \quad \text{and} \quad Y \in \mathfrak{k}$$

or

$$X = \left(\begin{array}{c|c} i\alpha & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \quad \text{and} \quad Y = \left(\begin{array}{c|c|c} 0 & & \\ \hline & i\beta & -\bar{y}^t \\ \hline & y & 0 \end{array} \right)$$

where $x = (0, x_3, \dots, x_{n+1})^t \neq 0 \in \mathbb{C}^n$ and $\sum_{j=3}^{n+1} x_j \bar{y}_j = 0$, or finally

$$(44) \quad X = \left(\begin{array}{c|c} i\alpha & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \quad \text{and} \quad Y = \left(\begin{array}{c|c|c} i & & \\ \hline & i\beta & -\bar{y}^t \\ \hline & y & 0 \end{array} \right)$$

where $x_2 \neq 0$, $\beta = 1 - \sum_{j=3}^{n+1} |y_j|^2$, and $x_j = -ix_2 y_j$ for $j = 3, \dots, n + 1$.

Therefore, in order to complete the proof we may restrict our attention to horizontal zero-curvature planes for which X and Y are of the form (44).

Without loss of generality we may assume that the vectors $\Psi^{-1}(X)$ and $\Psi^{-1}(Y)$ spanning $\check{\sigma}_2$ are orthogonal. By (28) and since $Y \in \mathfrak{k}$ this is equivalent to $\langle X_{\mathfrak{h}}, Y_{\mathfrak{h}} \rangle_0 = 0$, where we recall that $\langle V, W \rangle_0 = -\text{Re tr}(VW)$. For (44) we get orthogonality if and

only if $\alpha = 0$. Hence, as desired, we may rewrite (44) as

$$X = \left(\begin{array}{c|c} 0 & -\bar{x}^t \\ \hline x & 0 \end{array} \right) \quad \text{and} \quad Y = \left(\begin{array}{c|c|c} i & & \\ \hline & i\beta & -\bar{y}^t \\ \hline & y & 0 \end{array} \right)$$

where $x_2 \neq 0$, $\beta = 1 - \sum_{j=3}^{n+1} |y_j|^2$, and $x_j = -ix_2 y_j$ for $j = 3, \dots, n + 1$. □

In order to simplify the statements and computations to follow we fix

$$A_t := \left(\begin{array}{cc|c} \cos t & -\sin t & \\ \sin t & \cos t & \\ \hline & & I_{n-1} \end{array} \right) \in \text{U}(n + 1), \quad \cos^2 t \neq \mathbb{Q}$$

for the remainder of the proof, where I_{n-1} denotes the $(n - 1) \times (n - 1)$ identity matrix.

Lemma 5.2 *If $p_1 \neq p_2$ and $p_1 + p_2 \neq q_1 + q_2$, Then a vector*

$$(-\Phi_1^{-1}(\text{Ad}_{A^*} W), \Phi_2^{-1}(W))$$

with $W = \text{diag}(i, 0, \dots, 0), \text{diag}(0, i, 0, \dots, 0)$ or $\text{diag}(i, i, 0, \dots, 0)$

cannot be horizontal at $(A_t, I) \in \text{G} \times \text{G}$.

Proof Consider $V = \text{diag}(i\theta, i\varphi, 0, \dots, 0)$. From (30) we see that a vector of the form $(-\Phi_1^{-1}(\text{Ad}_{A_t^*} V), \Phi_2^{-1}(V))$ is horizontal if and only if $\langle V, \text{Ad}_{A_t} P - Q \rangle_0 = 0$. Since $\langle X, Y \rangle_0 = -\text{Re tr}(XY)$, this is equivalent to the condition

$$\begin{aligned} \theta q_1 + \varphi q_2 &= \theta \sum_{k=1}^{n+1} |a_{1k}|^2 p_k + \varphi \sum_{k=1}^{n+1} |a_{2k}|^2 p_k \\ &= (\theta \cos^2 t + \varphi \sin^2 t) p_1 + (\theta \sin^2 t + \varphi \cos^2 t) p_2. \end{aligned}$$

If $(\theta, \varphi) = (1, 0)$ this becomes $q_1 = p_1 \cos^2 t + p_2 \sin^2 t$ which is equivalent to $(p_1 - p_2) \cos^2 t = q_1 - p_2$, that is $\cos^2 t \in \mathbb{Q}$. This is impossible by definition of A_t . Similarly, when $(\theta, \varphi) = (0, 1)$ we again have a contradiction. Finally, when $(\theta, \varphi) = (1, 1)$ we get $q_1 + q_2 = p_1 + p_2$, which contradicts our hypothesis. □

Recall that the projection $\check{\sigma}_1$ of a plane σ as in Proposition 5.1 onto the first factor is given by

$$\begin{aligned} \check{\sigma}_1 &= \text{Span}\{\Phi_1^{-1}(\text{Ad}_{A^*} \tilde{X}), \Phi_1^{-1}(\text{Ad}_{A^*} \tilde{Y})\} \\ &= \text{Span}\{\Phi_1^{-1}(\text{Ad}_{A^*}(\Phi_1 X)), \Phi_1^{-1}(\text{Ad}_{A^*}(\Phi_1 Y))\}. \end{aligned}$$

Equation (3) provides us with conditions for $\check{\sigma}_1$ to have zero-curvature with respect to $\langle \cdot, \cdot \rangle_1$ but, if we assume that $\check{\sigma}_2$ has zero-curvature, since (G, K) is a rank-one symmetric pair the conditions reduce to

$$(45) \quad [(\text{Ad}_{A^*}(\Phi_1 X))_{\mathfrak{p}}, (\text{Ad}_{A^*}(\Phi_1 Y))_{\mathfrak{p}}] = 0.$$

That is, the \mathfrak{p} components of $\text{Ad}_{A^*}(\Phi_1 X)_{\mathfrak{p}}$ and $\text{Ad}_{A^*}(\Phi_1 Y)_{\mathfrak{p}}$ must be linearly dependent. There are three possible cases:

$$\begin{aligned} (\text{Ad}_{A^*}(\Phi_1 X))_{\mathfrak{p}} &= 0 \\ (\text{Ad}_{A^*}(\Phi_1 Y))_{\mathfrak{p}} &= 0 \\ (\text{Ad}_{A^*}(\Phi_1 X))_{\mathfrak{p}} &= s (\text{Ad}_{A^*}(\Phi_1 Y))_{\mathfrak{p}} \end{aligned}$$

for some $s \in \mathbb{R} - \{0\}$.

Recall that $\tilde{W} = \Phi_1(W) = W_{\mathfrak{p}} + \lambda_1 W_{\mathfrak{k}}$, $\lambda_1 \in (0, 1)$, and that

$$\text{Ad}_{A^*} \tilde{W} = \sum_{k,\ell=1}^{n+1} \bar{a}_{k1} a_{\ell j} \tilde{w}_{k\ell}$$

where $\tilde{W} = (\tilde{w}_{ij})$ and $W \in \mathfrak{g}$. Then, since \mathfrak{p} is completely determined by the first row of vectors in \mathfrak{g} , we may abuse notation to write

$$(46) \quad (\text{Ad}_{A^*} \tilde{W})_{\mathfrak{p}} = \left(\sum_{k,\ell=1}^{n+1} \bar{a}_{k1} a_{\ell j} \tilde{w}_{k\ell} \mid j = 2, \dots, n+1 \right).$$

Lemma 5.3 *Let σ be a zero-curvature plane at $(A_t, I) \in G \times G$ as given by Proposition 5.1(iv). If $p_1 \neq p_2$ and $p_1 + p_2 \neq q_1 + q_2$ then σ cannot be horizontal.*

Proof By (46) we have

$$\begin{aligned} (\text{Ad}_{A_t^*}(\Phi_1 X))_{\mathfrak{p}} &= (-\cos^2 t \bar{x}_2 - \sin^2 t x_2, -\cos t \bar{x}_3, \dots, -\cos t \bar{x}_{n+1}) \\ (\text{Ad}_{A_t^*}(\Phi_1 Y))_{\mathfrak{p}} &= (i\lambda_1(\beta - 1) \cos t \sin t, -\lambda_1 \sin t \bar{y}_3, \dots, -\lambda_1 \sin t \bar{y}_{n+1}). \end{aligned}$$

If $(\text{Ad}_{A_t^*}(\Phi_1 X))_{\mathfrak{p}} = 0$ then $x_j = y_j = 0$ for all $j = 3, \dots, n+1$, since $x_j = -ix_2 y_j$, $j = 3, \dots, n+1$. Thus $\beta = 1$ and $Y = \text{diag}(i, i, 0, \dots, 0)$. By Lemma 5.2 Y cannot determine a horizontal vector.

If $(\text{Ad}_{A_t^*}(\Phi_1 Y))_p = 0$ then $Y = \text{diag}(i, i, 0, \dots, 0)$ and, again, Lemma 5.2 shows that σ cannot be horizontal.

Finally we examine the situation $(\text{Ad}_{A_t^*}(\Phi_1 X))_p = s(\text{Ad}_{A_t^*}(\Phi_1 Y))_p$ for some nonzero $s \in \mathbb{R}$. Then $\cos t x_j = s\lambda_1 \sin t y_j$ implies that $-i \cos t x_2 y_j = s\lambda_1 \sin t y_j$ for $j = 3, \dots, n+1$, since $x_j = -ix_2 y_j$ for $j = 3, \dots, n+1$. We have already shown that if σ is to be horizontal the y_j cannot all be zero. Therefore $x_2 = i(s\lambda_1 \sin t / \cos t) \in i\mathbb{R}$.

Now $(-\Phi_1^{-1}(\text{Ad}_{A_t^*}(\Phi_1 X)), \Phi_2^{-1}(\Phi_1 X))$ is a horizontal vector if and only if the equation $\langle X, \text{Ad}_{A_t} P - Q \rangle_0 = 0$ is satisfied; that is, if and only if

$$\begin{aligned} 0 &= \sum_{\ell=1}^{n+1} \text{Im} \left(a_{1\ell} \sum_{k=2}^{n+1} \bar{a}_{k\ell} x_k \right) p_\ell \\ &= \text{Im}(a_{11} \bar{a}_{21} x_2) p_1 + \text{Im}(a_{12} \bar{a}_{22} x_2) p_2 \\ &= \cos t \sin t (p_1 - p_2) \text{Im}(x_2) \end{aligned}$$

where again we recall that $\langle V, W \rangle_0 = -\text{Re tr}(VW)$. By hypothesis and definition of A_t we have $x_2 = i \text{Im}(x_2) = 0$, which contradicts the assumption that $x_2 \neq 0$. \square

Lemma 5.4 *Let σ be a plane at $(A_t, I) \in G \times G$ as determined by Proposition 5.1(v). Then σ does not have zero curvature at (A_t, I) .*

Proof Following (46) we write

$$\begin{aligned} (\text{Ad}_{A_t^*}(\Phi_1 X))_p &= (-i\lambda_1 \alpha \cos t \sin t, -\cos t \bar{x}_3, \dots, -\cos t \bar{x}_{n+1}) \\ (\text{Ad}_{A_t^*}(\Phi_1 Y))_p &= (i\lambda_1 \beta \cos t \sin t, -\lambda_1 \sin t \bar{y}_3, \dots, -\lambda_1 \sin t \bar{y}_{n+1}). \end{aligned}$$

If $(\text{Ad}_{A_t^*}(\Phi_1 X))_p = 0$ then $X = 0$, which is a contradiction since σ is two-dimensional. Similarly, $(\text{Ad}_{A_t^*}(\Phi_1 Y))_p = 0$ gives a contradiction. On the other hand, if there is some nonzero $s \in \mathbb{R}$ such that $(\text{Ad}_{A_t^*}(\Phi_1 X))_p = s(\text{Ad}_{A_t^*}(\Phi_1 Y))_p$ then we find $\cos t x_j = s\lambda_1 \sin t y_j$, $j = 3, \dots, n+1$. However, since $\sum_{j=3}^{n+1} x_j \bar{y}_j = 0$, this implies that $x_j = y_j = 0$ for all $j = 3, \dots, n+1$. But, by hypothesis, x_j cannot all be zero for planes of this type, and so we have a contradiction. Therefore, by the discussion following (45), σ cannot admit zero-curvature at (A_t, I) . \square

Proof of Theorem B If $p_1 = \dots = p_{n+1}$ then we may assume without loss of generality that $p_i = 0$, for all $i = 1, \dots, n+1$. Since the action is free we must have $(q_1, q_2) \neq (0, 0)$. Hence, by Tapp [25], we are done.

As discussed in Section 1, a permutation of the integers p_1, \dots, p_{n+1} induces a diffeomorphism $E_{p,q}^{4n-1} \rightarrow E_{p,q}^{4n-1}$. Therefore, if $p_i \neq p_j$ for some $i \neq j$, we may

assume that $p_1 \neq p_2$. If $p_1 + p_2 \neq q_1 + q_2$ then we are done by [Proposition 5.1](#) and [Lemmas 5.2, Lemma 5.3](#) and [5.4](#).

Finally, if $p_1 \neq p_2$ but $p_1 + p_2 = q_1 + q_2$, then the freeness condition (29) implies that either $(p_1, p_2, \dots, p_{n+1}) = (1, \dots, 1, -1, \dots, -1)$ and $(q_1, q_2) = (0, 0)$ (up to sign and permutations of the p_i), where $\#\{i \mid p_i = -1\} \in \{\lfloor (n+2)/2 \rfloor, n\}$, or there is some $p_k \notin \{p_1, p_2\}$, $k \in \{3, \dots, n+1\}$.

In the latter case we may permute and relabel the p_i such that $p_1 + p_2 \neq q_1 + q_2$, in which case we are done as above. In the former case we may assume that $p_1 = 1$, $p_2 = -1$ and $p_3 = -1$ (after reordering if necessary). Then, for

$$A_0 := \left(\begin{array}{ccc|c} 1/2 & 1/\sqrt{2} & 1/2 & \\ -1/2 & 1/\sqrt{2} & -1/2 & \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} & \\ \hline & & & I_{n-2} \end{array} \right) \in U(n+1),$$

we may repeat the approaches used in the proofs of [Lemmas 5.2, 5.3](#) and [5.4](#) to rule out the existence of any horizontal zero-curvature planes at $(A_0, I) \in G \times G$. Note that, unlike in the proof of [Lemma 5.4](#), it is necessary to show that a plane as determined by [Proposition 5.1\(v\)](#) cannot have zero curvature and be horizontal at the same time. \square

6 Topology of M^{13} and N^{11}

We turn now to the topological assertions of [Theorem A](#) regarding the biquotients $M^{13} = SO(8) // (S^1 \times G_2)$ and $N^{11} = SO(8) // (SO(3) \times G_2)$, namely that they have the same cohomology rings but are not homeomorphic to $CP^3 \times S^7$ and $S^4 \times S^7$ respectively.

Theorem 6.1 *The biquotients M^{13} and N^{11} have the same cohomology rings as $CP^3 \times S^7$ and $S^4 \times S^7$ respectively. In particular M^{13} and N^{11} are not manifolds known to admit positive curvature.*

Proof In [Section 3](#) we established that M^{13} and N^{11} are the total spaces of S^7 -bundles over CP^3 and S^4 respectively. Given an arbitrary fibration $S^7 \rightarrow E \rightarrow B$, where B is a simply connected, compact manifold with $\dim B \leq 7$, the Euler class e is trivial since $H^8(B; \mathbb{Z}) = 0$ and so the Gysin sequence splits into short exact sequences, from which it immediately follows that $H^j(E; \mathbb{Z}) = H^j(B \times S^7; \mathbb{Z})$, for all j . Now M^{13} and N^{11} are quotients of $S^7 \times S^7$ by S^1 and S^3 respectively. A quick glance

at the Serre spectral sequences of these fibrations shows that the ring structures of $H^*(M^{13}; \mathbb{Z})$ and $H^*(N^{11}; \mathbb{Z})$ agree with those of $H^*(\mathbf{CP}^3 \times \mathbf{S}^7; \mathbb{Z})$ and $H^*(\mathbf{S}^4 \times \mathbf{S}^7; \mathbb{Z})$ as desired. \square

In order to distinguish M^{13} and N^{11} from the products $\mathbf{CP}^3 \times \mathbf{S}^7$ and $\mathbf{S}^4 \times \mathbf{S}^7$, respectively, we want to analyse their Pontrjagin classes. There is a general procedure (developed in Borel and Hirzebruch [5], Eschenburg [10] and Singhof [23]) for computing the R–Pontrjagin class of a biquotient $G//U$, where R is a coefficient ring such that $H^*(G; R)$ and $H^*(U; R)$ have no torsion, and the action of U on G is (effectively) free. Let $\iota: U \rightarrow G \times G$ denote the embedding and assume that we have such an R.

We adopt the following notation: For a compact, connected Lie group L, let T_L denote the maximal torus and W_L the Weyl group. Let E_L be a contractible space on which L acts freely. The classifying space of L is the quotient $B_L := E_L/L$. A product of Lie groups L_1 and L_2 is written L_1L_2 .

Consider the following commutative diagram of fibrations

$$\begin{array}{ccc}
 G \times E_{GG} & \longrightarrow & G \times E_{GG} \\
 \downarrow & & \downarrow \\
 G \times_U E_{GG} & \xrightarrow{\varphi_G} & G \times_{GG} E_{GG} = B_{\Delta G} \\
 \varphi_U \downarrow & & \downarrow B_{\Delta} \\
 B_U & \xrightarrow{B_{\iota}} & B_{GG}
 \end{array}$$

where φ_G and φ_U are the respective classifying maps, and $\Delta: G \rightarrow GG$ denotes the diagonal embedding. Now, since projection onto the first factor in each case is a homotopy equivalence, we have $G \simeq G \times E_{GG}$ and $G//U \simeq G \times_U E_{GG}$. Thus, up to homotopy, we can consider the diagram above as:

(47)

$$\begin{array}{ccc}
 G & \longrightarrow & G \\
 \downarrow & & \downarrow \\
 G//U & \xrightarrow{\varphi_G} & B_G \\
 \varphi_U \downarrow & & \downarrow B_{\Delta} \\
 B_U & \xrightarrow{B_{\iota}} & B_{GG}
 \end{array}$$

We would like to use the Serre spectral sequences of the fibrations on the left and right, as well as the commutativity of the diagram, to compute the Pontrjagin class of (the tangent bundle of) $G//U$. We will need to know $H^*(G; R)$, $H^*(B_U; R)$, $H^*(B_{GG}; R)$ and

the homomorphisms $(B_U)^*: H^*(B_{GG}; \mathbb{R}) \rightarrow H^*(B_U; \mathbb{R})$ and $(\varphi_U)^*: H^*(B_U; \mathbb{R}) \rightarrow H^*(G//U; \mathbb{R})$.

Let L be a compact, connected Lie group and \mathbb{R} a ring such that $H^*(L; \mathbb{R})$ has no torsion. Then

$$(48) \quad H^*(L; \mathbb{R}) = \Lambda(y_1, \dots, y_r)$$

where $r = \text{rank}(L)$. Hence, from the Serre spectral sequence of the universal bundle $L \rightarrow E_L \rightarrow B_L$,

$$(49) \quad H^*(B_L; \mathbb{R}) = \mathbb{R}[\bar{y}_1, \dots, \bar{y}_r]$$

where \bar{y}_j denotes the transgression of y_j .

Let (t_1, \dots, t_r) be coordinates of the maximal torus T_L . By an abuse of notation we identify t_j with the element $t_j \in H^1(T_L; \mathbb{R})$. The corresponding transgression arising from the Serre spectral sequence for $T_L \rightarrow E_{T_L} \rightarrow B_{T_L}$ is $\bar{t}_j \in H^2(B_{T_L}; \mathbb{R})$. Since L does not have any torsion in its \mathbb{R} -cohomology we have

$$(50) \quad H^*(B_L; \mathbb{R}) = H^*(B_{T_L}; \mathbb{R})^{\mathcal{W}_L} = \mathbb{R}[\bar{t}_1, \dots, \bar{t}_r]^{\mathcal{W}_L}.$$

Thus we have explicit well-defined generators of $H^*(B_L; \mathbb{R})$ which we identify with \bar{y}_j , $j = 1, \dots, r$.

Suppose now that $h: L_1 \rightarrow L_2$ is a homomorphism of Lie groups. Then the commutative diagram

$$\begin{array}{ccc} L_1 & \xrightarrow{h} & L_2 \\ \uparrow & & \uparrow \\ T_{L_1} & \xrightarrow{h} & T_{L_2} \end{array}$$

induces a commutative diagram of classifying spaces

$$(51) \quad \begin{array}{ccc} B_{L_1} & \xrightarrow{Bh} & B_{L_2} \\ \uparrow & & \uparrow \\ B_{T_{L_1}} & \xrightarrow{Bh} & B_{T_{L_2}} \end{array}$$

which in turn induces the commutative diagram:

$$(52) \quad \begin{array}{ccc} H^*(B_{L_1}) & \xleftarrow{(B_h)^*} & H^*(B_{L_2}) \\ \downarrow & & \downarrow \\ H^*(B_{T_{L_1}}) & \xleftarrow{(B_h)^*} & H^*(B_{T_{L_2}}) \end{array}$$

In particular, by (50), we see that $(B_h)^*$ can be determined by simply understanding $h(T_{L_1}) \subset T_{L_2}$.

Consider again diagram (47). Recall that, since there is no torsion,

$$H^*(B_{GG}; \mathbb{R}) = H^*(B_G; \mathbb{R}) \otimes H^*(B_G; \mathbb{R}).$$

Then $H^*(B_{GG}; \mathbb{R})$ is generated by classes of the form $\bar{y}_j \otimes 1$ and $1 \otimes \bar{y}_j$, $j = 1, \dots, r = \text{rank}(G)$. Consider the diagonal embedding $\Delta: G \hookrightarrow GG$. In coordinates $\Delta|_{T_G}$ is given by $t_j \mapsto (t_j, t_j)$, $j = 1, \dots, r$. We have commutative diagrams as in (51) and (52). Now

$$\begin{aligned} \Delta^*: H^1(T_G; \mathbb{R}) \otimes H^1(T_G; \mathbb{R}) &\longrightarrow H^1(T_G; \mathbb{R}) \\ t_j \otimes 1 &\longmapsto t_j \\ 1 \otimes t_j &\longmapsto t_j \end{aligned}$$

which in turn implies

$$\begin{aligned} (B_\Delta)^*: H^2(B_{T_G}; \mathbb{R}) \otimes H^2(B_{T_G}; \mathbb{R}) &\longrightarrow H^2(B_{T_G}; \mathbb{R}) \\ \bar{t}_j \otimes 1 &\longmapsto \bar{t}_j \\ 1 \otimes \bar{t}_j &\longmapsto \bar{t}_j. \end{aligned}$$

Therefore, by (50),

$$\begin{aligned} (B_\Delta)^*: H^*(B_{GG}; \mathbb{R}) &\longrightarrow H^*(B_G; \mathbb{R}) \\ \bar{y}_j \otimes 1 &\longmapsto \bar{y}_j \\ 1 \otimes \bar{y}_j &\longmapsto \bar{y}_j. \end{aligned}$$

Since the diagram (47) is commutative we see that

$$(53) \quad \begin{aligned} \varphi_G^*(\bar{y}_j) &= \varphi_G^*((B_\Delta)^*(\bar{y}_j \otimes 1)) = \varphi_U^*((B_l)^*(\bar{y}_j \otimes 1)) \\ \varphi_G^*(\bar{y}_j) &= \varphi_G^*((B_\Delta)^*(1 \otimes \bar{y}_j)) = \varphi_U^*((B_l)^*(1 \otimes \bar{y}_j)). \end{aligned}$$

We remark that, by naturality of spectral sequences, one can deduce from (47) that

$$H^*(G//U; \mathbb{R}) = H^* \left(\frac{H^*(G; \mathbb{R}) \otimes H^*(B_U; \mathbb{R})}{\langle dy_j = \varphi_U^*((B_l)^*(\bar{y}_j \otimes 1)) - \varphi_U^*((B_l)^*(1 \otimes \bar{y}_j)) \rangle} \right).$$

Let us now focus on computing the Pontrjagin class of $G//U$. Let τ be the tangent bundle of $G//U$. In analogue with Singhof [23], we introduce the following vector bundles over $G//U$: First, let $\alpha_G := (GG/U) \times_G \mathfrak{g}$, where G acts on $(GG/U) \times \mathfrak{g}$ via

$$g \star ([g_1, g_2], X) = ([gg_1, gg_2], \text{Ad}_g X)$$

with $g, g_1, g_2 \in G, [g_1, g_2] = (g_1, g_2) \cdot U \in GG/U$, and $X \in \mathfrak{g}$. Second, let $\alpha_U := G \times_U \mathfrak{u}$, where $\mathfrak{u} \subset \mathfrak{g} \oplus \mathfrak{g}$ is the Lie algebra of U and U acts on $G \times \mathfrak{u}$ via

$$(u_1, u_2) \star (g, (Y_1, Y_2)) = (u_1 g u_2^{-1}, (\text{Ad}_{u_1} Y_1, \text{Ad}_{u_2} Y_2))$$

with $g \in G, (u_1, u_2) \in U$ and $(Y_1, Y_2) \in \mathfrak{u}$. Then, as in Singhof [23, Proposition 3.2]:

Lemma 6.2 $\tau \oplus \alpha_U = \alpha_G$.

Proof Recall that, since U acts on G via $(u_1, u_2) \star g = u_1 g u_2^{-1}, (u_1, u_2) \in U, g \in G$, the vertical subspace at $g \in G$ (after left-translating back to $e \in G$) is given by

$$\mathcal{V}_g = \{ \text{Ad}_{g^{-1}} Y_1 - Y_2 \mid (Y_1, Y_2) \in \mathfrak{u} \} \subset \mathfrak{g}.$$

Moreover, since the action of U on G is free, we have

$$(54) \quad \{ (Y_1, Y_2) \in \mathfrak{u} \mid \text{Ad}_{g^{-1}} Y_1 = Y_2 \text{ for some } g \in G \} = \{0\}.$$

Given $g \in G$, consider the maps

$$\begin{aligned} \psi_g: \mathfrak{g} &\longrightarrow \alpha_G \\ X &\longmapsto [[g^{-1}, e], X] \end{aligned}$$

and

$$\begin{aligned} f: \alpha_U &\longrightarrow \alpha_G \\ [g, (Y_1, Y_2)] &\longmapsto [[g^{-1}, e], \text{Ad}_{g^{-1}} Y_1 - Y_2]. \end{aligned}$$

It is easy to check that f is well-defined and that $f_{[g]}((\alpha_U)_{[g]}) = \psi_g(\mathcal{V}_g)$, where $(\alpha_U)_{[g]}$ is the fibre of the bundle α_U over $[g] = g \cdot U \in G//U$. Furthermore, (54) allows us to deduce that f is injective.

If we now equip G with a bi-invariant metric, then (after left-translation to the identity in G) the horizontal subspace at g is given by

$$\mathcal{H}_g = \{ X \mid X \in \mathfrak{g}, X \perp \mathcal{V}_g \} \subset \mathfrak{g}.$$

Note that U acts on the horizontal distribution by taking $X \in \mathcal{H}_g$ to $\text{Ad}_{u_2} X \in \mathcal{H}_{u_1 g u_2^{-1}}$, where $(u_1, u_2) \in U$. Let $\check{\mathcal{H}} = \mathcal{H}/U$ be the orbit space under the U action. In particular, $\tau \cong \check{\mathcal{H}}$ and we can represent the tangent space at $[g] \in G//U$ by

$$\check{\mathcal{H}}_{[g]} = \{[g, X] \mid X \in \mathfrak{g}, X \perp \mathcal{V}_g\}$$

where the g in $[g, X]$ serves only to keep track of the base point.

Recall that each point in α_G may be represented as $[[g_1, g_2], \text{Ad}_{g_2} X] = [[g_2^{-1} g_1, e], X]$, for some $g_1, g_2 \in G$ and $X \in \mathfrak{g}$. Consider the map

$$q: \alpha_G \longrightarrow \check{\mathcal{H}}_{[g]} = \tau$$

$$[[g_1, g_2], \text{Ad}_{g_2} X] \longmapsto [g_1^{-1} g_2, X^\perp]$$

where X^\perp is the component of $X \in \mathfrak{g}$ orthogonal to $\mathcal{V}_{g_1^{-1} g_2}$. Clearly q is surjective. The fact that q is well-defined follows from noticing that $(\text{Ad}_{u_2} X)^\perp = \text{Ad}_{u_2}(X^\perp) \in \mathcal{H}_{u_1 g u_2^{-1}}$, for all $X \in \mathfrak{g}$, $g \in G$ and $(u_1, u_2) \in U$.

Now $q([[g_1, g_2], \text{Ad}_{g_2} X]) = 0$ for some $[[g_1, g_2], \text{Ad}_{g_2} X] \in \alpha_G$ if and only if $X^\perp = 0 \in \mathcal{H}_{g_1^{-1} g_2}$, that is, $X \in \mathcal{V}_{g_1^{-1} g_2}$. Thus $[[g_1, g_2], \text{Ad}_{g_2} X] = [[g_2^{-1} g_1, e], X] \in f_{[g_1^{-1} g_2]}(\alpha_U)_{[g_1^{-1} g_2]}$ and we have $\ker q = f(\alpha_U)$. Therefore $\alpha_G/f(\alpha_U) = \tau$. In other words, $\alpha_G = \tau \oplus \alpha_U$ as desired. \square

Recall from Borel and Hirzebruch [5] that, if V is a representation of a Lie group L , then the homogeneous vector bundle $\alpha_L = P \times_L V$ associated to the L -principal bundle $P \longrightarrow B := P/L$ is the pullback under the classifying map $\varphi_L: B \longrightarrow B_L$ of the vector bundle $E_L \times_L V$ associated to the universal L -principal bundle $E_L \longrightarrow B_L$, ie $\alpha_L = \varphi_L^*(E_L \times_L V)$. In particular, from Borel and Hirzebruch [5] and Singhof [23] we know that the Pontrjagin class of the bundle $\alpha_L \longrightarrow B$ is given by

$$p(\alpha_L) = 1 + p_1(\alpha_L) + p_2(\alpha_L) + \dots = \varphi_L^*(a), \quad a := \prod_{\alpha_i \in \Delta_L^+} (1 + \bar{\alpha}_i^2)$$

where Δ_L^+ is the set of positive weights of the representation of L on V . We have identified $\alpha_i \in H^1(T_{L}; \mathbb{R})$ with $\bar{\alpha}_i \in H^2(B_{T_L}; \mathbb{R})$ via transgression, and so it follows that $a \in H^*(B_{T_L}; \mathbb{R})^{W_L} \cong H^*(B_L; \mathbb{R})$.

The vector bundles α_U and α_G are associated to principal U and G bundles respectively and, in this case, the weights of the defining representations are the roots of the corresponding Lie groups.

Since $p(V \oplus W) = p(V) \sim p(W)$, for vector bundles V and W over some manifold M , we have

$$p(\tau) p(\alpha_U) = p(\alpha_G).$$

By our discussion above and since inverses are well-defined in the polynomial algebra $H^*(B_U; \mathbb{R})$ it follows that

$$p(\tau) = \varphi_G^*(a)\varphi_U^*(b^{-1})$$

where $a := \prod_{\alpha_i \in \Delta_G^+} (1 + \bar{\alpha}_i^2)$ and $b := \prod_{\beta_j \in \Delta_U^+} (1 + \bar{\beta}_j^2)$. In particular,

$$\begin{aligned} p_1(\tau) &= p_1(\alpha_G) - p_1(\alpha_U) \\ (55) \quad &= \varphi_G^* \left(\sum_{\alpha_i \in \Delta_G^+} \bar{\alpha}_i^2 \right) - \varphi_U^* \left(\sum_{\beta_j \in \Delta_U^+} \bar{\beta}_j^2 \right). \end{aligned}$$

In our situation, the Lie groups $G = \text{SO}(8)$ and $U_k := \text{SO}(k) \times G_2$, with $k = 2, 3$, have no torsion in their cohomology for coefficients in $\mathbb{R} = \mathbb{Z}_p$, p an odd prime (see Mimura and Toda [19, Corollary 3.15, Theorem 5.12]). We use the process outlined above to compute the \mathbb{Z}_p -Pontrjagin classes of M^{13} and N^{11} . Before we continue we establish an easy lemma which will prove useful in the topological computations to follow.

Lemma 6.3 *Consider a triple (r_1, r_2, r_3) such that $\sum r_i = 0$. Let $\sigma_i(r)$ and $\sigma_i(r^2)$ denote the i -th elementary symmetric polynomials in r_1, r_2, r_3 and r_1^2, r_2^2, r_3^2 respectively. Then $\sigma_1(r^2) = -2\sigma_2(r)$ and $\sigma_2(r^2) = \sigma_2(r)^2$.*

Proof Since $\sigma_1(r) = \sum r_i = 0$ we have

$$\begin{aligned} 0 &= \sigma_1(r)^2 \\ &= (r_1^2 + r_2^2 + r_3^2) + 2(r_1r_2 + r_1r_3 + r_2r_3) \\ &= \sigma_1(r^2) + 2\sigma_2(r) \end{aligned}$$

as desired. On the other hand:

$$\begin{aligned} \sigma_2(r)^2 - \sigma_2(r^2) &= (r_1r_2 + r_1r_3 + r_2r_3)^2 - (r_1^2r_2^2 + r_1^2r_3^2 + r_2^2r_3^2) \\ &= 2(r_1^2r_2r_3 + r_1r_2^2r_3 + r_1r_2r_3^2) \\ &= 2r_1r_2r_3(r_1 + r_2 + r_3) \\ &= 0 \end{aligned} \quad \square$$

In [9, pages *vii* and 139], Eschenburg provides a beautiful diagram which explicitly describes the embedding of the root system G_2 into B_3 . Recall that B_3 is the root system corresponding to the Lie algebra $\mathfrak{so}(7)$ and is given by

$$B_3 = \{\pm t_i \mid 1 \leq i \leq 3\} \cup \{\pm(t_i \pm t_j) \mid 1 \leq i < j \leq 3\}.$$

The root system G_2 lies on a hypersurface in $\text{Span}\{B_3\}$ and is given by

$$G_2 = \{\pm s_i \mid 1 \leq i \leq 3\} \cup \{\pm(s_i - s_j) \mid 1 \leq i < j \leq 3\}$$

where $s_i = \frac{1}{3}(2t_i - t_j - t_k)$, $\{i, j, k\} = \{1, 2, 3\}$. Notice that $\sum s_i = 0$ and that $s_i - s_j = t_i - t_j \in B_3$. Furthermore, s_i is the projection of $t_i \in B_3$ and $-(t_j + t_k) \in B_3$ onto the hypersurface containing G_2 .

Since the Lie group G_2 is simply connected and has no centre, we see that the inclusions

$$\exp^{-1}(I) = \text{integral lattice of } G_2 \subset \text{root lattice of } G_2 \subset \text{weight lattice of } G_2$$

are in fact equalities. Therefore, by our above discussion of the roots of G_2 , the integral and weight lattices of G_2 are spanned by $\{s_i \mid 1 \leq i \leq 3\}$, $\sum s_i = 0$. Thus by an abuse of notation we may assume that $\{s_i \mid 1 \leq i \leq 3\}$, $\sum s_i = 0$, spans $H^1(T_{G_2}; \mathbb{Z}) = \text{Hom}(\Gamma, \mathbb{Z})$, where T_{G_2} is a maximal torus of G_2 and Γ is the integral lattice of G_2 .

Proposition 6.4 *Let p be an odd prime. The \mathbb{Z}_p first Pontrjagin classes of M^{13} and N^{11} are*

$$p_1(M^{13}) = 2\alpha^2 \quad \text{and} \quad p_1(N^{11}) = \beta$$

respectively, where α is a generator of $H^2(M^{13}; \mathbb{Z}_p) = \mathbb{Z}_p$ and β is a generator of $H^4(N^{11}; \mathbb{Z}_p) = \mathbb{Z}_p$.

Proof Let $G = \text{SO}(8)$ and let $U_k := \text{SO}(k) \times G_2$, $k = 2, 3$, act freely on G as described in Section 3 with quotients M^{13} and N^{11} respectively. Let $\iota_k: U_k \hookrightarrow GG$, $k = 2, 3$, denote the respective inclusions.

For p an odd prime, the \mathbb{Z}_p -cohomology of G and U_k is

$$\begin{aligned} H^*(G; \mathbb{Z}_p) &= \Lambda(y_1, y_2, y_3, y_4), & y_1 \in H^3, y_2, y_4 \in H^7, y_3 \in H^{11} \\ H^*(U_k; \mathbb{Z}_p) &= \Lambda(w_k, x_1, x_2), & w_k \in H^{2k-3}(\text{SO}(k)), x_i \in H^{8i-5}(G_2). \end{aligned}$$

Let T_G and T_U be the maximal tori of G and U_k respectively, with coordinates being given by (t_1, t_2, t_3, t_4) and (u, s_1, s_2, s_3) , $\sum s_i = 0$, respectively. By an abuse of notation (and our earlier discussion of the roots of G_2) we will identify t_i , u and s_j with the elements $t_i \in H^1(T_G)$ and $u, s_j \in H^1(T_U)$. The corresponding transgressions are $\bar{t}_i \in H^2(B_{T_G})$ and $\bar{u}, \bar{s}_j \in H^2(B_{T_U})$.

The Weyl group W_G acts on $H^*(B_{T_G})$ via permutations in \bar{t}_i and an even number of sign changes. Therefore a basis for $H^*(B_{T_G})^{W_G}$ is given by elementary symmetric polynomials $\sigma_i(\bar{t}^2) := \sigma_i(\bar{t}_1^2, \dots, \bar{t}_4^2)$, $i = 1, 2, 3$, and $\bar{t}_1\bar{t}_2\bar{t}_3\bar{t}_4$. Hence, by (50), the generators of $H^*(B_G)$ can be chosen as $\bar{y}_i := \sigma_i(\bar{t}^2) \in H^{4i}$, $i = 1, 2, 3$, and $\bar{y}_4 := \bar{t}_1\bar{t}_2\bar{t}_3\bar{t}_4 \in H^8$.

The Weyl group W_{U_2} is the dihedral group of order twelve. It acts trivially on u , while the action on the root system G_2 is by rotations of $\pi/3$ and by reflections through the horizontal axis. Therefore, given our description of the root system of G_2 above, W_{U_2} acts on $H^*(B_{T_U})$ via permutations in \bar{s}_i and a simultaneous sign change of all \bar{s}_i . On the other hand, $W_{U_3} = \mathbb{Z}_2 \times W_{U_2}$, where the \mathbb{Z}_2 factor acts trivially on the s_i and by a sign change on u . Thus elements of $H^*(B_{T_U})$ which are invariant under W_{U_k} are given by sums and products of \bar{u}^{k-1} and the elementary symmetric polynomials $\sigma_2(\bar{s}) := \sigma_2(\bar{s}_1, \bar{s}_2, \bar{s}_3)$ and $\sigma_i(\bar{s}^2) := \sigma_i(\bar{s}_1^2, \bar{s}_2^2, \bar{s}_3^2)$, $i = 1, 2, 3$. However, since $\sum s_i = 0$, Lemma 6.3 shows that a basis for $H^*(B_{T_U})^{W_{U_k}}$ is given by \bar{u}^{k-1} and the symmetric polynomials $\sigma_2(\bar{s})$ and $\sigma_3(\bar{s}^2)$. By (50), generators of $H^*(B_{U_k})$ are given by $\bar{w}_k := \bar{u}^{k-1} \in H^{2(k-1)}$, $\bar{x}_1 := \sigma_2(\bar{s}) \in H^4$ and $\bar{x}_2 := \sigma_3(\bar{s}^2) \in H^{12}$.

Consider $\iota_k: U_k \hookrightarrow GG$ as above. By (12), $\iota_k|_{T_U}: T_U \rightarrow T_{GG}$ has (in coordinates) the form $\iota_k|_{T_U}(u, s_1, s_2, s_3) = ((0, 0, 0, u), (0, s_1, s_2, -s_3))$. Hence

$$\begin{aligned}
 (\iota_k|_{T_U})^*: H^1(T_{GG}) &\longrightarrow H^1(T_U) \\
 t_i \otimes 1 &\longmapsto 0, & i = 1, 2, 3 \\
 t_4 \otimes 1 &\longmapsto u \\
 1 \otimes t_1 &\longmapsto 0 \\
 1 \otimes t_i &\longmapsto s_{i-1}, & i = 2, 3 \\
 1 \otimes t_4 &\longmapsto -s_3
 \end{aligned}$$

from which it follows:

$$\begin{aligned}
 (B_{\iota_k|_{T_U}})^*: H^2(B_{T_{GG}}) &\longrightarrow H^2(B_{T_U}) \\
 \bar{t}_i \otimes 1 &\longmapsto 0, & i = 1, 2, 3 \\
 \bar{t}_4 \otimes 1 &\longmapsto \bar{u} \\
 1 \otimes \bar{t}_1 &\longmapsto 0 \\
 1 \otimes \bar{t}_i &\longmapsto \bar{s}_{i-1}, & i = 2, 3 \\
 1 \otimes \bar{t}_4 &\longmapsto -\bar{s}_3
 \end{aligned}$$

Therefore, by (52) we have:

$$\begin{aligned}
 (B_{\iota_k})^*: H^*(B_{GG}) &\longrightarrow H^*(B_{U_k}) \\
 \bar{y}_1 \otimes 1 &\longmapsto \bar{u}^2 \\
 \bar{y}_i \otimes 1 &\longmapsto 0, & i = 2, 3, 4 \\
 1 \otimes \bar{y}_i &\longmapsto \sigma_i(\bar{s}^2), & i = 1, 2, 3 \\
 1 \otimes \bar{y}_4 &\longmapsto 0
 \end{aligned}
 \tag{56}$$

In particular, $(\mathbf{B}_{l_k})^*(1 \otimes \bar{y}_1) = -2\bar{x}_1$, $(\mathbf{B}_{l_k})^*(1 \otimes \bar{y}_2) = \bar{x}_1^2$ (by Lemma 6.3) and $(\mathbf{B}_{l_k})^*(1 \otimes \bar{y}_3) = \bar{x}_2$.

We are now in a position to compute the first Pontrjagin class of $G//U_k$. Recall that the positive roots of $G = \text{SO}(8)$ are $t_i \pm t_j$, $1 \leq i < j \leq 4$. Hence:

$$\begin{aligned} \sum_{\alpha_i \in \Delta_G^+} \bar{\alpha}_i^2 &= \sum_{1 \leq i < j \leq 4} ((\bar{t}_i - \bar{t}_j)^2 + (\bar{t}_i + \bar{t}_j)^2) \\ &= 2 \sum_{1 \leq i < j \leq 4} (\bar{t}_i^2 + \bar{t}_j^2) \\ &= 6 \sum_{i=1}^4 \bar{t}_i^2 \\ &= 6\bar{y}_1 \end{aligned}$$

Since $\text{SO}(2)$ has no (positive) roots and $\text{SO}(3)$ has only one positive root, namely u , we may denote the positive root of $\text{SO}(k)$ by $(k-2)u$, for $k = 2, 3$, respectively. From our earlier description of the roots of G_2 , the positive roots of $U_k = \text{SO}(k) \times G_2$ are

$$(k-2)u, s_1, s_2, -s_3, s_1 - s_3, s_2 - s_1, s_2 - s_3$$

where $\sum s_i = 0$. Then:

$$\begin{aligned} \sum_{\beta_j \in \Delta_{U_k}^+} \bar{\beta}_j^2 &= (k-2)\bar{u}^2 + \bar{s}_1^2 + \bar{s}_2^2 + \bar{s}_3^2 + (\bar{s}_1 - \bar{s}_3)^2 + (\bar{s}_2 - \bar{s}_1)^2 + (\bar{s}_2 - \bar{s}_3)^2 \\ &= (k-2)\bar{u}^2 + 3\sigma_1(\bar{s}^2) - 2\sigma_2(\bar{s}) \\ &= (k-2)\bar{u}^2 - 8\sigma_2(\bar{s}) \quad \text{by Lemma 6.3} \\ &= (k-2)\bar{u}^2 - 8\bar{x}_1 \end{aligned}$$

Note that, by (53) and (56),

$$\varphi_G^*(\bar{y}_1) = \varphi_{U_k}^*((\mathbf{B}_{l_k})^*(\bar{y}_1 \otimes 1)) = \varphi_{U_k}^*(\bar{u}^2)$$

and $-2\varphi_{U_k}^*(\bar{x}_1) = \varphi_{U_k}^*((\mathbf{B}_{l_k})^*(1 \otimes \bar{y}_1)) = \varphi_{U_k}^*((\mathbf{B}_{l_k})^*(\bar{y}_1 \otimes 1)) = \varphi_{U_k}^*(\bar{u}^2)$.

It follows now from (55) that:

$$\begin{aligned} p_1(G//U_k) &= p_1(\alpha_G) - p_1(\alpha_{U_k}) \\ &= 6\varphi_G^*(\bar{y}_1) - (k-2)\varphi_{U_k}^*(\bar{u}^2) + 8\varphi_{U_k}^*(\bar{x}_1) \\ &= 6\varphi_{U_k}^*(\bar{u}^2) - (k-2)\varphi_{U_k}^*(\bar{u}^2) - 4\varphi_{U_k}^*(\bar{u}^2) \\ &= (4-k)\varphi_{U_k}^*(\bar{u}^2) \in H^4(G//U_k) \end{aligned}$$

It remains only to show that $\varphi_{U_k}^*(\bar{u}^2)$ is a generator of $H^4(G//U_k)$. To achieve this, consider the Serre spectral sequence for the fibration $G \rightarrow G//U_k \rightarrow B_{U_k}$.

Suppose $k = 2$. Notice that $\langle \bar{w}_2 \rangle = H^2(B_{U_2}) = E_2^{2,0}$ will survive until E_∞ since $H^*(G)$ contains no elements of degree 1.

On the other hand, suppose now that $k = 3$. By (56) the generator $y_1 \in H^3(G) = E_2^{0,3} = E_4^{0,3}$ gets mapped under d_4 to

$$d_4(y_1) = (B_{I_3})^*(\bar{y}_1 \otimes 1) - (B_{I_3})^*(1 \otimes \bar{y}_1) = \bar{u}^2 + 2\bar{x}_1 \in E_2^{4,0} = E^4 = H^4(B_{U_3}).$$

The generators $\bar{w}_3 = \bar{u}^2$ and \bar{x}_1 of $H^4(B_{U_3})$ are both mapped to zero by d_4 . Thus, the $E_5^{4,0}$ term is a \mathbb{Z}_p generated by $\bar{w}_3 = \bar{u}^2$ and survives to E_∞ .

Recall that the classifying map $\varphi_{U_k}^*$ is the edge homomorphism

$$\varphi_{U_k}^*: H^i(B_{U_k}) = E_2^{i,0} \rightarrow E_\infty^{i,0} \hookrightarrow H^i(G//U_k).$$

By the discussion above $\varphi_{U_k}^*(\bar{w}_k) = \varphi_{U_k}^*(\bar{u}^{k-1}) \neq 0$, $k = 2, 3$. Hence $\varphi_{U_k}^*(\bar{w}_k)$ is a generator of $H^{2(k-1)}(G//U_k) = \mathbb{Z}_p$. If $k = 3$ we are done. When $k = 2$, we know from Theorem 6.1 that $H^4(M^{13})$ is generated by the square of a generator of $H^2(M^{13})$. That is, $\varphi_{U_2}^*(\bar{w}_2) = \varphi_{U_2}^*(\bar{u}^2)$ is a generator of $H^4(M^{13})$. \square

Remark 6.5 Since H^8 and H^{12} are trivial for each of the manifolds M^{13} and N^{11} , we have in fact computed their total Pontrjagin classes $p = 1 + p_1$ in \mathbb{Z}_p coefficients.

Remark 6.6 In terms of integral cohomology, the proposition tells us only that $p_1(M^{13})$ and $p_1(N^{11})$ are nontrivial and not divisible by any primes $p \geq 3$. Thus $p_1(M^{13})$ and $p_1(N^{11})$ have the form $\pm 2^\ell \in \mathbb{Z} = H^4$, for some $\ell \in \mathbb{N} \cup \{0\}$.

Corollary 6.7 N^{11} is not homeomorphic to $S^4 \times S^7$.

Proof By the previous remark, in integral coefficients $p_1(N^{11}) = \pm 2^\ell \in \mathbb{Z} = H^4(N^{11}; \mathbb{Z})$, for some $\ell \in \mathbb{N} \cup \{0\}$. However, since all (integral) Pontrjagin classes for spheres are trivial and integral Pontrjagin classes are homeomorphism invariants, N^{11} cannot be homeomorphic to a product of spheres. \square

Since $p_1(\mathbb{C}P^3 \times S^7) = 4\gamma^2$, where γ generates $H^2(\mathbb{C}P^3 \times S^7; \mathbb{Z})$, we are unable to distinguish M^{13} and $\mathbb{C}P^3 \times S^7$ using the proposition. We need to explicitly compute the integral Pontrjagin class of M^{13} . We can accomplish this by “hot-wiring” the technique for computing Pontrjagin classes in the absence of torsion in the cohomology groups.

Before we begin we establish two topological statements which will be used in the proof of [Theorem 6.10](#). From now on we will always assume that our cohomology groups have integral coefficients, and by spectral sequence we will always mean Serre spectral sequence.

Proposition 6.8 *The classifying space B_{G_2} of G_2 , has low dimensional integral cohomology groups $H^1 = H^2 = H^3 = H^5 = 0$ and $H^4 = \mathbb{Z}$ with generator $\bar{x} = \sigma_2(\bar{s})$, where $\sigma_2(\bar{s}) := \sigma_2(\bar{s}_1, \bar{s}_2, \bar{s}_3)$, $\sum \bar{s}_i = 0$, and $\bar{s}_i \in H^2(B_{T_{G_2}})$, $i = 1, 2, 3$, are the transgressions of the elements $s_i \in H^1(T_{G_2})$, $i = 1, 2, 3$, which span the integral lattice of G_2 .*

Proof Consider the universal bundle $G_2 \rightarrow E_{G_2} \rightarrow B_{G_2}$ where E_{G_2} is contractible. From Whitehead [[28](#), Theorem 5.17], we know that $H^j(G_2) = 0$, $j = 1, 2, 4, 5$, and $H^3(G_2) = \mathbb{Z}$. Let x be a generator of $H^3(G_2)$. Since E_{G_2} is contractible all entries in the spectral sequence for the fibration $G_2 \rightarrow E_{G_2} \rightarrow B_{G_2}$ must get killed off. Since $d_4: E_4^{0,3} \rightarrow E_4^{4,0}$ is the only possible nontrivial differential with domain $E_4^{0,3}$ it must map $x \in H^3(G_2)$ to a generator \bar{x} of $H^4(B_{G_2})$, and so $H^4(B_{G_2}) = \mathbb{Z}$. Similarly it is clear from the spectral sequence that $H^j(B_{G_2}) = 0$ for $j = 1, 2, 3, 5$.

Now consider the fibration $S^6 = G_2/SU(3) \rightarrow B_{SU(3)} \rightarrow B_{G_2}$. The spectral sequence associated to this fibration shows that $\bar{x} \in E_2^{4,0} = H^4(B_{G_2})$ survives to E_∞ . Thus, since there are no other nonzero entries on the corresponding diagonal in E_∞ , we see that $H^4(B_{G_2}) = H^4(B_{SU(3)})$. Recall that $H^*(B_{SU(3)})$ is a polynomial algebra generated by the elementary symmetric polynomials $\sigma_i(\bar{s}) = \sigma_i(\bar{s}_1, \bar{s}_2, \bar{s}_3)$, $i = 2, 3$, in the transgressions \bar{s}_j of $s_j \in H^1(T_{SU(3)})$, $j = 1, 2, 3$, where the s_j span the integral lattice of $SU(3)$. Note that $\sum s_j = 0$, $T_{G_2} = T_{SU(3)}$ and $\deg(\sigma_i(\bar{s})) = 2i$. Therefore $H^4(B_{G_2})$ is generated by $\sigma_2(\bar{s})$ as desired. □

Proposition 6.9 *The low dimensional integral cohomology groups of the manifold $SO(8)/G_2 = (S^7 \times S^7)/\mathbb{Z}_2$ are $H^j(SO(8)/G_2) = H^j(\mathbb{R}P^7)$, $0 \leq j \leq 6$.*

Proof Consider the spectral sequence for the fibration

$$\mathbb{R}P^7 = SO(7)/G_2 \rightarrow SO(8)/G_2 \rightarrow SO(8)/SO(7) = S^7.$$

Recall:

$$H^j(\mathbb{R}P^7) = \begin{cases} \mathbb{Z} & \text{if } j = 0, 7, \\ \mathbb{Z}_2 & \text{if } j = 2, 4, 6, \\ 0 & \text{if } j = 1, 3, 5. \end{cases}$$

It is clear that each $E_2^{0,j} = H^j(\mathbf{RP}^7)$, $j \leq 5$, survives to E_∞ . For $E_2^{0,6} = H^6(\mathbf{RP}^7) = \mathbb{Z}_2$ notice that there are no nontrivial homomorphisms $\mathbb{Z}_2 \rightarrow \mathbb{Z}$ and so the differential $d_7: E_7^{0,6} = \mathbb{Z}_2 \rightarrow E_7^{7,0} = \mathbb{Z}$ must be trivial. Therefore $E_2^{0,6} = H^6(\mathbf{RP}^7)$ also survives to E_∞ . Since there are no other nonzero entries on the corresponding diagonals we get the desired result. \square

We are now ready to complete the proof of [Theorem A\(i\)](#).

Theorem 6.10 *The first integral Pontrjagin class of $M^{13} = \text{SO}(8) // (\mathbf{S}^1 \times G_2)$ is given by*

$$|p_1(M^{13})| = 8z^2$$

where z is a generator of $H^2(M^{13}; \mathbb{Z}) = \mathbb{Z}$. In particular, M^{13} is not homeomorphic to $\mathbf{CP}^3 \times \mathbf{S}^7$.

Proof Consider once again diagram (47), with $G = \text{SO}(8)$ and $U = \mathbf{S}^1 \times G_2$ such that $G//U = M^{13}$. In the proof of [Proposition 6.4](#) we followed the usual techniques of Borel and Hirzebruch [5], Eschenburg [10] and Singhof [23] when there is no torsion in cohomology, namely we computed B_t and B_Δ and then used the fact that the diagram commutes in order to compute the \mathbb{Z}_p Pontrjagin class, for odd primes p . However, since $\text{SO}(8)$ and G_2 have torsion in integral cohomology, we need to adopt a different approach in order to compute the integral Pontrjagin class. Since $H^8(M^{13}) = H^{12}(M^{13}) = 0$ we can restrict our attention to the first integral Pontrjagin class $p_1(M^{13}) \in H^4(M^{13})$. The key idea to be taken from the proof of [Proposition 6.4](#) is that we computed the first Pontrjagin classes of some vector bundles over B_G and B_U , then pulled them back to M^{13} under the classifying maps φ_G and φ_U respectively. As it turns out, the first Pontrjagin classes of these vector bundles over B_G and B_U are the same in integral coefficients as in \mathbb{Z}_p coefficients $p \geq 3$. Our strategy, therefore, is to compute the maps $\varphi_U^*: H^4(B_U) \rightarrow H^4(M^{13})$ and $\varphi_G^*: H^4(B_G) \rightarrow H^4(M^{13})$ and pull back the respective first Pontrjagin classes.

As a first step in computing $\varphi_U^*: H^4(B_U) \rightarrow H^4(M^{13})$ we notice that $H^*(U) = H^*(\mathbf{S}^1) \otimes H^*(G_2)$ and $H^*(B_U) = H^*(B_{\mathbf{S}^1}) \otimes H^*(B_{G_2})$ since $H^*(\mathbf{S}^1)$ and $H^*(B_{\mathbf{S}^1})$ are torsion-free. Therefore

$$H^j(U) = \begin{cases} \mathbb{Z} & \text{if } j = 0, \\ \mathbb{Z} = \langle w \rangle & \text{if } j = 1, \\ \mathbb{Z} = \langle x \rangle & \text{if } j = 3, \\ 0 & \text{if } j = 2, 4, 5, \end{cases}$$

where w is a generator of $H^1(S^1)$ and x is a generator of $H^3(G_2)$, and applying Proposition 6.8,

$$H^j(B_U) = \begin{cases} \mathbb{Z} & \text{if } j = 0, \\ \mathbb{Z} = \langle \bar{w} \rangle & \text{if } j = 2, \\ \mathbb{Z} \oplus \mathbb{Z} = \langle \bar{w}^2 \rangle \oplus \langle \bar{x} \rangle & \text{if } j = 4, \\ 0 & \text{if } j = 1, 3, 5, \end{cases}$$

where \bar{w} is the transgression of w resulting from the spectral sequence for the universal bundle of S^1 and generates $H^2(B_{S^1})$ (hence generates $H^*(B_{S^1}) = \mathbb{Z}[\bar{w}]$), and \bar{x} is the transgression of x resulting from the spectral sequence for the universal bundle of G_2 and generates $H^4(B_{G_2})$.

Recall that $\varphi_U: G//U \rightarrow B_U$ is the classifying map since we have the following diagram of principal U -bundles

$$\begin{array}{ccc} U & \longrightarrow & U \\ \downarrow & & \downarrow \\ G \times E_U & \xrightarrow{\pi_2} & E_U \\ \downarrow & & \downarrow \\ G \times_U E_U & \xrightarrow{\pi_2} & B_U \end{array}$$

where π_2 denotes projection onto the second factor and $U \rightarrow E_U \rightarrow B_U$ is the universal bundle. Since E_U is contractible, projection onto the first factor gives homotopy equivalences $G \times E_U \simeq G$ and $G \times_U E_U \simeq G//U$. The resulting map $G//U \rightarrow B_U$ is φ_U . Hence φ_U is the classifying map. Therefore, up to homotopy, we may consider the following commutative diagram of fibrations:

$$\begin{array}{ccc} U & \longrightarrow & U \\ \downarrow & & \downarrow \\ G & \longrightarrow & E_U \\ \downarrow & & \downarrow \\ G//U & \xrightarrow{\varphi_U} & B_U \end{array}$$

Consider first the spectral sequence for the fibration on the left. Recall that $H^*(M^{13}) = H^*(\mathbf{CP}^3 \times \mathbf{S}^7)$. Hence:

$$H^j(G//U) = \begin{cases} \mathbb{Z} & \text{if } j = 0, \\ \mathbb{Z} = \langle z \rangle & \text{if } j = 2, \\ \mathbb{Z} = \langle z^2 \rangle & \text{if } j = 4, \\ 0 & \text{if } j = 1, 3, 5. \end{cases}$$

Since $G = \text{SO}(8)$ we have from Čadek, Mimura and Vanžura [7]:

$$H^j(G) = \begin{cases} \mathbb{Z} & \text{if } j = 0, \\ 0 & \text{if } j = 1, \\ \mathbb{Z}_2 = \langle r \rangle & \text{if } j = 2, \\ \mathbb{Z} = \langle y \rangle & \text{if } j = 3, \\ \mathbb{Z}_2 = \langle r^2 \rangle & \text{if } j = 4. \end{cases}$$

Since $H^1(G) = 0$ we see that $d_2: E_2^{0,1} = \langle w \rangle \rightarrow E_2^{2,0} = \langle z \rangle$ must have trivial kernel, ie $d_2(w) = kz$ for some $k \in \mathbb{Z}$, $k \neq 0$. Then $E_3^{0,2} = \langle z \rangle / \langle kz \rangle$ survives to E_∞ and since $H^2(G) = \mathbb{Z}_2$ we must therefore have $k = \pm 2$, ie $d_2(w) = \pm 2z$.

On the other hand, the spectral sequence shows that on the E_4 -page we have the differential $d_4: E_4^{0,3} = \langle x \rangle \rightarrow E_4^{0,4} = \langle z^2 \rangle / \langle 2z^2 \rangle$. However, since $H^3(G) = \mathbb{Z}$ and $H^4(G) = \mathbb{Z}_2$, we must have $d_4(x) = 0 \in \langle z^2 \rangle / \langle 2z^2 \rangle$.

Since E_U is contractible it is clear from the spectral sequence for the fibration on the right that $d_2: E_2^{0,1} = \langle w \rangle \rightarrow E_2^{2,0} = \langle \bar{w} \rangle$ is an isomorphism with $d_2(w) = \bar{w}$, and $d_4: E_4^{0,3} = \langle x \rangle \rightarrow E_4^{4,0} = \langle \bar{w}^2 \rangle \oplus \langle \bar{x} \rangle$ is given by $d_4(x) = \bar{x}$.

By naturality of the spectral sequence we thus have for the left-hand fibration that $d_2(w) = \varphi_U^*(\bar{w}) \in \langle z \rangle$ and $d_4(x) = \varphi_U^*(\bar{x}) \in \langle z^2 \rangle / \langle 2z^2 \rangle$. Therefore, since we have already shown that $d_2(w) = \pm 2z \in \langle z \rangle$ and $d_4(x) = 0 \in \langle z^2 \rangle / \langle 2z^2 \rangle$, we find

$$(57) \quad \begin{aligned} \varphi_U^*(\bar{w}) &= \pm 2z \in H^2(G//U) = \langle z \rangle \\ \varphi_U^*(\bar{x}) &= 2kz^2 \in H^4(G//U) = \langle z^2 \rangle, \text{ for some } k \in \mathbb{Z}. \end{aligned}$$

We now turn our attention to computing $\varphi_G^*: H^4(B_G) \rightarrow H^4(M^{13})$. In order to show that $\varphi_G: G//U \rightarrow B_G$ is the classifying map consider the commutative diagram of

principal G -bundles:

$$\begin{array}{ccc}
 G & \longrightarrow & G \\
 \downarrow & & \downarrow \\
 GG \times_U E_{GG} & \longrightarrow & GG \times_{GG} E_{GG} \\
 \downarrow & & \downarrow \\
 (\Delta G \backslash GG) \times_U E_{GG} & \xrightarrow{\varphi_G} & (\Delta G \backslash GG) \times_{GG} E_{GG}
 \end{array}$$

Since $GG \times_{GG} E_{GG} = E_{GG}$ and $(\Delta G \backslash GG) \times_{GG} E_{GG} = G \times_{GG} E_{GG} = B_{\Delta G}$ we see that the fibration on the right-hand side is the universal bundle for G . On the left-hand side we have $(\Delta G \backslash GG) \times_U E_{GG} = G \times_U E_{GG}$, and projection onto the first factor gives homotopy equivalences $GG \times_U E_{GG} \simeq GG/U$ and $G \times_U E_{GG} \simeq G//U$. Thus up to homotopy the diagram becomes

$$\begin{array}{ccc}
 G & \longrightarrow & G \\
 \downarrow & & \downarrow \\
 GG/U & \longrightarrow & E_{GG} \\
 \downarrow & & \downarrow \\
 G//U & \xrightarrow{\varphi_G} & B_G
 \end{array}$$

as desired. Recall that $H^3(G) = \langle y \rangle$. The cohomology of B_G is described in Brown [6] and Feshbach [12], but for our purposes we need only that:

$$H^j(B_G) = \begin{cases} \mathbb{Z} & \text{if } j = 0, \\ 0 & \text{if } j = 1, 2, \\ \mathbb{Z}_2 & \text{if } j = 3, \\ \mathbb{Z} & \text{if } j = 4, \\ \mathbb{Z}_2 & \text{if } j = 5. \end{cases}$$

Whilst proving Proposition 3.6 in [16] Grove and Ziller showed that, since $E = E_{GG}$ is contractible, in the spectral sequence for the bundle $G \rightarrow E \rightarrow B_G$ the differential $d_4: E_4^{0,3} = \langle 2y \rangle \rightarrow E_4^{4,0} = H^4(B_G)$ is an isomorphism, ie $2y$ gets mapped to a generator \bar{y} of $H^4(B_G) = \mathbb{Z}$. This follows from the facts that $E_2^{2,2} = \mathbb{Z}_2$ (by the Universal Coefficient Theorem) and that $d_2: E_2^{0,3} = \langle y \rangle \rightarrow E_2^{2,2} = \mathbb{Z}_2$ must be onto.

Therefore naturality of the spectral sequence implies that $d_4(2y) = \varphi_G^*(\bar{y})$ in the spectral sequence for the left-hand fibration $G \rightarrow GG/U \rightarrow G//U$, where $H^3(G) = \langle y \rangle$ and $H^4(B_G) = \langle \bar{y} \rangle$.

In order to determine the exact value of $\varphi_G^*(\bar{y}) \in H^4(G//U)$ we need to examine the spectral sequence for the left-hand fibration. First we must compute the cohomology of GG/U in low-dimensions. Recall that $GG/U = V_{8,6} \times SO(8)/G_2$, where $V_{8,6}$ is the Stiefel manifold $SO(8)/SO(2)$. From Čadek, Mimura and Vanžura [7] we find:

$$H^j(V_{8,6}) = \begin{cases} \mathbb{Z} & \text{if } j = 0, 2, \\ 0 & \text{if } j = 1, 3, 5, \\ \mathbb{Z}_2 & \text{if } j = 4. \end{cases}$$

In Proposition 6.9 we computed the low dimensional cohomology groups of $SO(8)/G_2$. From the general Künneth formula for cohomology (see Spanier [24, Theorem 11, Section 5]) it follows that:

$$H^j(GG/U) = \begin{cases} \mathbb{Z} & \text{if } j = 0, \\ 0 & \text{if } j = 1, 3, \\ \mathbb{Z} \oplus \mathbb{Z}_2 & \text{if } j = 2, \\ \mathbb{Z}_2^3 & \text{if } j = 4, \\ \mathbb{Z}_2 & \text{if } j = 5. \end{cases}$$

Since $H^4(GG/U) = \mathbb{Z}_2^3$, in the spectral sequence for $G \rightarrow GG/U \rightarrow G//U$ the differential $d_2: E_2^{0,3} = H^3(G) = \langle y \rangle \rightarrow E_2^{2,2} = \mathbb{Z}_2$ must be trivial, ie $E_2^{2,2} = \mathbb{Z}_2$ must survive to E_∞ . It thus follows that $E_2^{i,j} = E_3^{i,j} = E_4^{i,j}$ for $i \leq 5, j \leq 4$. Since $H^3(GG/U) = 0$ the differential $d_4: E_4^{0,3} = \langle y \rangle \rightarrow E_4^{4,0} = H^4(G//U) = \langle z^2 \rangle$ must be given by $d_4(y) = nz^2$ for some nonzero $n \in \mathbb{Z}$. On the other hand, since $H^4(GG/U) = \mathbb{Z}_2^3$, $E_4^{0,4} = E_4^{2,2} = \mathbb{Z}_2$ and $E_4^{1,3} = E_4^{3,1} = 0$, the filtration for the spectral sequence shows that $n = \pm 2$, ie $d_4(y) = \pm 2z^2$. But we have already shown that $d_4(2y) = \varphi_G^*(\bar{y})$. Therefore

$$\varphi_G^*(\bar{y}) = \pm 4z^2 \in H^4(G//U) = \langle z^2 \rangle.$$

Furthermore, while proving Lemma 5.4 in [16] Grove and Ziller showed that, by considering the spectral sequences of the fibrations $SO(8)/SO(3) \rightarrow B_{SO(3)} \rightarrow B_{SO(8)}$ and $SO(3)/SO(2) \rightarrow B_{SO(2)} \rightarrow B_{SO(3)}$, we can let $\bar{y} = \sigma_1(\bar{t}^2) = \sigma_1(\bar{t}_1^2, \bar{t}_2^2, \bar{t}_3^2, \bar{t}_4^2)$, where (t_1, \dots, t_4) are the coordinates of a maximal torus T_G of G and by abuse of notation we identify $t_i \in H^1(T_G)$ with $\bar{t}_i \in H^2(B_{T_G})$ via transgression.

We are now in a position to compute the first Pontrjagin class of $M^{13} = G//U$. We have already shown that, despite having torsion in cohomology, $H^4(B_G) \cong H^4(B_{T_G})^{\mathcal{W}_G}$

and $H^4(B_U) \cong H^4(B_{T_U})^{\mathcal{W}_U}$ since the generators are $\bar{y} = \sigma_1(\bar{t}^2)$ and $\bar{x} = \sigma_2(\bar{s})$ respectively. Therefore, in the present setting, Equation (55) for p_1 is still valid for integral coefficients.

As in the proof of Proposition 6.4, $p_1(\alpha_G) = 6\varphi_G^*(\bar{y})$. But $\varphi_G^*(\bar{y}) = \pm 4z^2$. Hence $p_1(\alpha_G) = \pm 24z^2 \in H^4(G//U)$.

Similarly, from the proof of Proposition 6.4 we have $p_1(\alpha_U) = -8\varphi_U^*(\bar{x})$. Thus, since $\varphi_U^*(\bar{x}) = 2kz^2$ by (57), $p_1(\alpha_U) = -16kz^2 \in H^4(G//U)$, for some $k \in \mathbb{Z}$.

Therefore, by (55),

$$\begin{aligned} p_1(\tau) &= p_1(\alpha_G) - p_1(\alpha_U) \\ &= 8(2k \pm 3)z^2 \in H^4(G//U). \end{aligned}$$

By Proposition 6.4 we know that $p_1(\tau) = p_1(G//U)$ is divisible only by 2. Therefore we must have $2k \pm 3 = \pm 1$, which implies $p_1(G//U) = \pm 8z^2$ as desired. \square

References

- [1] **J F Adams**, *Lectures on exceptional Lie groups*, (Z Mahmud, M Mimura, editors) Chicago Lectures in Math., Univ. of Chicago Press (1996) [MR1428422](#) With a foreword by J P May
- [2] **S Aloff, N R Wallach**, *An infinite family of distinct 7-manifolds admitting positively curved Riemannian structures*, Bull. Amer. Math. Soc. 81 (1975) 93–97 [MR0370624](#)
- [3] **Y V Bazaikin**, *On a family of 13-dimensional closed Riemannian manifolds of positive curvature*, Sibirsk. Mat. Zh. 37 (1996) 1219–1237, ii [MR1440379](#)
- [4] **M Berger**, *Les variétés riemanniennes homogènes normales simplement connexes à courbure strictement positive*, Ann. Scuola Norm. Sup. Pisa (3) 15 (1961) 179–246 [MR0133083](#)
- [5] **A Borel, F Hirzebruch**, *Characteristic classes and homogeneous spaces. I*, Amer. J. Math. 80 (1958) 458–538 [MR0102800](#)
- [6] **E H Brown, Jr**, *The cohomology of BSO_n and BO_n with integer coefficients*, Proc. Amer. Math. Soc. 85 (1982) 283–288 [MR652459](#)
- [7] **M Čadek, M Mimura, J Vanžura**, *The cohomology rings of real Stiefel manifolds with integer coefficients*, J. Math. Kyoto Univ. 43 (2003) 411–428 [MR2051031](#)
- [8] **O Dearricott**, *A 7-manifold with positive curvature*, Preprint (2008)
- [9] **J-H Eschenburg**, *Freie isometrische Aktionen auf kompakten Lie-Gruppen mit positiv gekrümmten Orbträumen*, Schrift. des Math. Inst. der Univ. Münster, 2. Serie 32, Univ. Münster Math. Inst. (1984) [MR758252](#)

- [10] **J-H Eschenburg**, *Cohomology of biquotients*, Manuscripta Math. 75 (1992) 151–166 [MR1160094](#)
- [11] **J-H Eschenburg, M Kerin**, *Almost positive curvature on the Gromoll–Meyer sphere*, Proc. Amer. Math. Soc. 136 (2008) 3263–3270 [MR2407092](#)
- [12] **M Feshbach**, *The integral cohomology rings of the classifying spaces of $O(n)$ and $SO(n)$* , Indiana Univ. Math. J. 32 (1983) 511–516 [MR703281](#)
- [13] **H Gluck, F Warner, W Ziller**, *The geometry of the Hopf fibrations*, Enseign. Math. (2) 32 (1986) 173–198 [MR874686](#)
- [14] **D Gromoll, W Meyer**, *An exotic sphere with nonnegative sectional curvature*, Ann. of Math. (2) 100 (1974) 401–406 [MR0375151](#)
- [15] **K Grove, L Verdiani, W Ziller**, *An exotic T_1S^4 with positive curvature*, Preprint (2008) Available at <http://www.math.upenn.edu/~wziller/papers/T1S4.pdf>
- [16] **K Grove, W Ziller**, *Lifting group actions and non-negative curvature*, to appear in Trans. Amer. Math. Soc. [arXiv:0801.0767](#)
- [17] **N Jacobson**, *Composition algebras and their automorphisms*, Rend. Circ. Mat. Palermo (2) 7 (1958) 55–80 [MR0101253](#)
- [18] **M Kerin**, *On the curvature of biquotients*, to appear in Math. Ann.
- [19] **M Mimura, H Toda**, *Topology of Lie groups. I, II*, Translations of Math. Monogr. 91, Amer. Math. Soc. (1991) [MR1122592](#) Translated from the 1978 Japanese edition by the authors
- [20] **S Murakami**, *Exceptional simple Lie groups and related topics in recent differential geometry*, from: “Differential geometry and topology (Tianjin, 1986–87)”, (B J Jiang, C-K Peng, Z X Hou, editors), Lecture Notes in Math. 1369, Springer, Berlin (1989) 183–221 [MR1001187](#)
- [21] **P Petersen, F Wilhelm**, *An exotic sphere with positive sectional curvature* [arXiv:0805.0812](#)
- [22] **P Petersen, F Wilhelm**, *Examples of Riemannian manifolds with positive curvature almost everywhere*, Geom. Topol. 3 (1999) 331–367 [MR1714915](#)
- [23] **W Singhof**, *On the topology of double coset manifolds*, Math. Ann. 297 (1993) 133–146 [MR1238411](#)
- [24] **E H Spanier**, *Algebraic topology*, Springer, New York (1981) [MR666554](#) Corrected reprint
- [25] **K Tapp**, *Quasi-positive curvature on homogeneous bundles*, J. Differential Geom. 65 (2003) 273–287 [MR2058263](#)
- [26] **K Tapp**, *Flats in Riemannian submersions from Lie groups*, Asian J. Math. 13 (2009) 459–464 [MR2653712](#)

- [27] **NR Wallach**, *Compact homogeneous Riemannian manifolds with strictly positive curvature*, Ann. of Math. (2) 96 (1972) 277–295 [MR0307122](#)
- [28] **G W Whitehead**, *Elements of homotopy theory*, Graduate Texts in Math. 61, Springer, New York (1978) [MR516508](#)
- [29] **F Wilhelm**, *An exotic sphere with positive curvature almost everywhere*, J. Geom. Anal. 11 (2001) 519–560 [MR1857856](#)
- [30] **B Wilking**, *Manifolds with positive sectional curvature almost everywhere*, Invent. Math. 148 (2002) 117–141 [MR1892845](#)
- [31] **W Ziller**, *Examples of Riemannian manifolds with non-negative sectional curvature*, from: “Surveys in differential geometry. Vol. XI”, (J Cheeger, K Grove, editors), Surv. Differ. Geom. 11, Int. Press, Somerville, MA (2007) 63–102 [MR2408264](#)

*Mathematisches Institut, Fachbereich Mathematik und Informatik der Universität Münster
Einsteinstrasse 62, D-48149 Münster, Germany*

m.kerin@math.uni-muenster.de

Proposed: Dmitri Burago

Seconded: Tobias H Colding, Steven Ferry

Received: 13 February 2009

Revised: 3 August 2010