

# SO(3)–Donaldson invariants of $\mathbb{C}P^2$ and mock theta functions

ANDREAS MALMENDIER

KEN ONO

We compute the Moore–Witten regularized  $u$ –plane integral on  $\mathbb{C}P^2$ , and we confirm the conjecture that it is the generating function for the SO(3)–Donaldson invariants of  $\mathbb{C}P^2$ . We also derive generating functions for the SO(3)–Donaldson invariants with  $2N_f$  massless monopoles using the geometry of certain rational elliptic surfaces ( $N_f \in \{0, 2, 3, 4\}$ ), and we show that the partition function for  $N_f = 4$  is nearly modular. Our results rely heavily on the theory of mock theta functions and harmonic Maass forms (for example, see Ono [40]).

57R57

## 1 Introduction and statement of results

In his plenary address at the Ramanujan Centenary Conference at the University of Illinois (Urbana–Champaign) in 1987, Freeman Dyson proclaimed his hope [15] that the “theory” of Ramanujan’s mock theta functions would someday play a role in mathematical physics.

The mock theta-functions give us tantalizing hints of a grand synthesis still to be discovered. Somehow it should be possible to build them into a coherent group-theoretical structure, analogous to the structure of modular forms which Hecke built around the old theta-functions of Jacobi. This remains a challenge for the future. My dream is that I will live to see the day when our young physicists, struggling to bring the predictions of superstring theory into correspondence with the facts of nature, will be led to enlarge their analytic machinery to include mock theta-functions. . . . But before this can happen, the purely mathematical exploration of the mock-modular forms and their mock-symmetries must be carried a great deal further.

—Freeman Dyson

The 2002 PhD thesis of Sander Zwegers [56] has provided Dyson’s desired “coherent group-theoretical structure”, and it turns out that Ramanujan’s mock theta functions are *holomorphic parts* of weight  $1/2$  *harmonic weak Maass forms* (see Section 5 for definitions). Zwegers’s thesis has sparked a flurry of recent activity involving such Maass forms. Indeed, harmonic Maass forms are now known to play a central role in the study of Ramanujan’s mock theta functions, as well as other important mathematical topics: Borcherds products, derivatives of modular  $L$ -functions, Gross–Zagier formulas and Faltings heights of CM cycles, partitions, and traces of singular moduli (see Bringmann and Ono [5; 6], Bringmann, Ono and Rhoades [7], Bruinier [8], Bruinier and Funke [9], Bruinier and Ono [10], Bruinier and Yang [12], Ono [40], Zagier [55] and Zwegers [56]).

Here we give an application, which combined with earlier important works of Göttsche and Zagier [19; 22], realizes Dyson’s original hope that mock theta functions would one day play a role in mathematical physics. We shall relate mock theta functions to Donaldson invariants of a smooth simply connected four-dimensional manifold; see Donaldson and Kronheimer [14].

There are two families of Donaldson invariants, corresponding to the  $SU(2)$ -gauge theory and the  $SO(3)$ -gauge theory with nontrivial Stiefel–Whitney class. In each case, the Donaldson invariants are graded homogeneous polynomials on the homology  $H_0(\mathbb{C}P^2) \oplus H_2(\mathbb{C}P^2)$ , where  $H_i(\mathbb{C}P^2)$  is considered to have degree  $(4-i)/2$ , defined using the fundamental homology classes of the corresponding moduli spaces of antiselfdual instantons arising in gauge theory. The Donaldson invariants depend by definition on the choice of a Riemannian metric. In the case  $b_2^+ > 1$ , the Donaldson invariants are independent of the metric as long as it is generic. In the case  $b_2^+ = 1$ , Kotschick and Morgan [27] showed that the Donaldson invariants depended on the metric but only through the chamber of its period point in the positive cone in  $H^2(X, \mathbb{R})$ . Crossing through a wall in  $H^2(X, \mathbb{R})$  defined by a class  $\xi \in H^2(X, \mathbb{Z})$  adds a certain wall-crossing term  $\delta_{\xi, k}^X$  to the Donaldson invariants. Kotschick and Morgan made the following conjecture:  $\delta_{\xi, k}^X$  is a polynomial in the multiplication by  $\xi$  and the quadratic form  $Q_X$  on  $H_2(X, \mathbb{Z})$  whose coefficients depend only on  $\xi^2$ , the instanton number  $k$  and the homotopy type on  $X$ .

We restrict ourselves to the simplest manifold to which Donaldson’s definition applies, the complex projective plane  $\mathbb{C}P^2$  with the Fubini–Study metric. In earlier work, Göttsche and Zagier [22] gave a formula for the Donaldson invariants of rational surfaces in terms of theta functions of indefinite lattices. These are Jacobi forms for special choices of the polarization on the boundary of the positive cone. As an application, Göttsche [19] derived closed expressions for the two families of the Donaldson invariants of  $\mathbb{C}P^2$  assuming the Kotschick–Morgan conjecture. In the recent

work of Göttsche et al [21], it was shown that the assumption of the Kotschick–Morgan conjecture was *not* necessary for the formula for the Donaldson invariants.

The present paper concerns deep conjectural relations between these works and constructions in theoretical physics which are presently the focus of much study. From the viewpoint of theoretical physics (see Witten [51]), these two families of Donaldson invariants and the related Seiberg–Witten invariants are the correlation functions of a supersymmetric topological gauge theory with gauge group SU(2) and SO(3) respectively. Using physical considerations, Witten [53] argued that one should be able to compute these correlation functions in a so called *low energy effective field theory* instead. This theory has the advantage of being an *abelian*  $\mathcal{N} = 2$  supersymmetric topological gauge theory. The data required to define the theory only involves line bundles of even (resp. odd) first Chern class on  $\mathbb{C}P^2$  if the gauge group is SU(2) (resp. SO(3)). The vacua of the low energy effective field theory are parametrized by the  $u$ –plane which is a certain analytically marked rational elliptic surface. Seiberg and Witten [42] argued further that the rational elliptic surface is the modular curve  $\mathbb{H}/\Gamma_0(4)$ , together with a meromorphic one-form. Moreover, Moore and Witten [37] obtained the correlation functions as a regularized integral over the  $u$ –plane. The integrand is a modular invariant function which is determined by the marked elliptic surface and the gauge group. The regularization procedure depends on interpreting the integrand as a total derivative, combined with constant term contributions from cusps. This integration by parts naturally introduces nonholomorphic modular forms of weight  $3/2$  (resp.  $1/2$ ) for the gauge group SU(2) (resp. SO(3)). Thus, the regularized  $u$ –plane integral defines a way of extracting certain contributions for each boundary component near the cusps at  $\tau = 0, 2, \infty$  of the rational elliptic surface. Moore and Witten observed [37] that the cuspidal contributions at  $\tau = 0, 2$  vanish trivially, which coincides with the mathematical statement that all Seiberg–Witten invariants on  $\mathbb{C}P^2$  vanish due to the presence of a Fubini–Study metric of positive scalar curvature; see Witten [52]. Moore and Witten went further and made the following comprehensive conjecture:

**Conjecture** (Moore and Witten [37]) *The contribution from the cusp at  $\tau = \infty$  to the regularized  $u$ –plane integral is the generating function for the Donaldson invariants of  $\mathbb{C}P^2$ .*

As evidence for this conjecture, in the case of the gauge group SU(2), Moore and Witten [37] computed the first 40 invariants and found them to be in agreement with the results of Ellingsrud and Göttsche [16]. However, the conjecture remains open.

**Remark** Moore and Witten [37] also showed that the general equality for SU(2) implies some interesting relations involving the classical Hurwitz class numbers which

arise in number theory. It is in this way that harmonic Maass forms make their first appearance. The holomorphic part of Zagier's weight  $3/2$  Maass–Eisenstein series, which first arose in Hirzebruch and Zagier [23] in connection with intersection theory for certain Hilbert modular surfaces, is the generating function for Hurwitz class numbers.

Our main result concerns the case of the  $SO(3)$ –gauge theory where little was known about the conjecture. We prove the following theorem:

**Theorem 1.1** *The conjecture of Moore and Witten in the case of the  $SO(3)$ –gauge theory on  $\mathbb{CP}^2$  is true.*

Donaldson theory can be generalized by introducing sections of a  $\text{spin}_{\mathbb{C}}$ –structure on  $\mathbb{CP}^2$  coupled to the  $SO(3)$ –gauge bundle. In the physics literature, these additional fields are called *massless monopoles*, and their number is denoted by  $2N_f$ , where  $0 \leq N_f \leq 4$ . The case where  $N_f = 0$  corresponds to the original  $SO(3)$ –Donaldson invariants<sup>1</sup>. The corresponding moduli spaces and invariants when  $N_f > 0$  have not been given much consideration in mathematics. However in physics, Seiberg and Witten [43] argued that including the monopoles changes the rational elliptic surface. For  $N_f > 0$  the modular elliptic surface for  $\Gamma_0(4)$  is replaced by rational elliptic surfaces determined by  $N_f$  vectors in the  $E_8$  root lattice. Rather nicely, the integrand of the  $u$ –plane integral remains a modular invariant when the gauge group is  $SO(3)$ . We classify all rational elliptic surfaces which appear as Seiberg–Witten curves for  $0 \leq N_f \leq 4$  (see Lemma 3.1). For  $N_f = 2, 3$ , it turns out that the elliptic surfaces for the Donaldson theory with massless monopoles are still modular elliptic surfaces related to the elliptic surface for  $N_f = 0$  by two–isogeny or a twist of the elliptic fibration. However, in the case where  $N_f = 1$  the surface is not modular elliptic (see Lemma 4.1 and Nahm [38] for a physical explanation). In the case  $N_f = 4$ , elliptic surfaces for the Donaldson theory with massive monopoles can be obtained from twisting the fibration of the modular elliptic surface in the case  $N_f = 2$ . We give explicit Weierstrass representations (see Lemmas 4.1 and 4.21) for the cases  $N_f = 2, 3, 4$ , and we compute the generating functions (see Equation (65) and Theorem 4.25) using the regularized  $u$ –plane integral. We conjecture:

**Conjecture** *The generating function in (65) is the generating function (24) for the  $SO(3)$ –Donaldson invariants of  $\mathbb{CP}^2$  with  $2N_f$  monopoles for  $N_f = 2, 3$ .*

Using the Atiyah–Singer index theorem and Theorem 1.1, we can also show that the conjecture passes a nontrivial test proving that the highest order contributions to

<sup>1</sup>The number of sections is even and  $N_f \leq 4$ ; otherwise the quantum theory is inconsistent [37].

each coefficient in the partition functions agree for  $N_f = 2, 3$ . In the  $N_f = 4$  case, instead of asymptotic freedom one has conformal invariance of the physical system. Thus, the partition function is a function of the complex coupling constant  $\tau$ . We explicitly compute the partition function. To this end, we shall define a power series in  $q = \exp(2\pi i \tau)$  (see Sections 4 and 7)

$$Q^+(q) = \frac{1}{q^{1/8}}(1 + 28 q^{1/2} + 39 q + 196 q^{3/2} + 161 q^2 + \dots).$$

Adding a nonholomorphic part  $Q^-(q)$  gives the Maass form  $Q(q) = Q^+(q) + Q^-(q)$  (see Section 4). In terms of the function  $Z(\tau) = \eta^3(\tau) Q(q)$ , we prove the following theorem:

**Theorem 1.2** *The regularized partition function of the massless  $N_f = 4$  low energy effective field theory on  $\mathbb{C}P^2$  is well defined, and is given by*

$$(1) \quad Z_{UP}^4 = \left[ \frac{1}{2} \frac{q}{\eta^4(\tau)} \frac{d}{dq} \left( \frac{q}{\eta^4(\tau)} \frac{d}{dq} \right) + g(\tau) \right] \frac{Z(\tau)}{\eta^4(\tau)},$$

where

$$g(\tau) = -\frac{1}{2^2 3^2} \left[ \left( \frac{\vartheta_2(\tau)}{\eta(\tau)} \right)^8 - \left( \frac{\vartheta_2(\tau)}{\eta(\tau)} \right)^4 \left( \frac{\vartheta_3(\tau)}{\eta(\tau)} \right)^4 + \left( \frac{\vartheta_3(\tau)}{\eta(\tau)} \right)^8 \right].$$

Here we discuss this result in the context of the theory of modular forms. Recall that the Donaldson polynomials arise as correlation functions of the  $\mathcal{N} = 2$  supersymmetric topological gauge theory with gauge group  $SU(2)$  or  $SO(3)$ . There are also quantum field theories on  $\mathbb{C}P^2$  with more supersymmetry. Roughly speaking, there are three choices for a  $\mathcal{N} = 4$  supersymmetric topological gauge theory, determined by what is referred to as *the topological twist*. One particular choice, called the *Vafa–Witten twist*, has been studied extensively in the physics literature [50], and its partition function has been computed in many examples. It turns out that the coefficient of  $q^k$  in the partition function equals the Euler characteristic of the instanton moduli space of instanton number  $k$ . The *S-duality* sends  $\tau \mapsto -1/\tau$  and exchanges electric and magnetic fields. In the  $\mathcal{N} = 4$  theory, the S-duality is a symmetry of the physical theory whence the partition function is in fact a weakly holomorphic weight 0 modular form. Indeed, Vafa and Witten even point out the strong relation with the nonholomorphic weight  $3/2$  Eisenstein series of Zagier which is perhaps the first half-integral weight harmonic Maass form which was seriously investigated. In the  $\mathcal{N} = 2$  theory, the S-duality is not a symmetry. Accordingly, we see that the partition function is not modular, but only a “piece” of a modular invariant, a fact which is central to our proof of Theorem 1.1. It has the property that its “trace”, in a very natural sense, gives rise to modular invariants

(see Equation (74)) which resemble some modular partition functions computed by Vafa and Witten [50, Section 42]. The partition function (1) is naturally related to the mock theta function

$$M(q^8) := q^{-1} \sum_{n=0}^{\infty} \frac{(-1)^{n+1} q^{8(n+1)^2} \prod_{k=1}^n (1 - q^{16k-8})}{\prod_{k=1}^{n+1} (1 + q^{16k-8})^2} = -q^7 + 2q^{15} - 3q^{23} + \dots$$

In Section 7 we prove that

$$Q(q) - 4M(q)$$

is a weight  $1/2$  weakly holomorphic modular form. Therefore, we then find that

$$\frac{1}{\eta(\tau)} \cdot (Q(q) - 4M(q))$$

is the desired modular invariant, a weight 0 modular form. This paper is organized as follows. In Section 2 we recall facts about  $\mathrm{SO}(3)$ –Donaldson invariants on  $\mathbb{C}\mathbb{P}^2$ , and we recall certain generating functions of Göttsche which play a central role in the proof of Theorem 1.1. In Section 3 we recall and classify the Seiberg–Witten curves, and in Section 4 we use their properties to define the  $u$ –plane integral, which in turn we employ to construct the vital generating series when  $N_f \in \{0, 2, 3, 4\}$ . The proof of Theorem 1.1 is then reduced to a criterion (see Theorem 4.26) involving these new generating functions, in the case where  $N_f = 0$ , and those of Göttsche. To verify this criterion, we employ the theory of harmonic Maass forms which is briefly recalled in Section 5. Specifically, we relate the relevant generating functions to weight  $1/2$  harmonic Maass forms which are described using a nonholomorphic Jacobi form constructed by Zwegers. In Section 6 we recall the construction of this Jacobi form, and in Section 7 we use it to represent the generating functions in terms of explicit harmonic Maass forms. In Section 8 we conclude with the proof of Theorem 1.1, and we give some numerical examples.

**Acknowledgements** The authors thank Kathrin Bringmann, Amanda Folsom, Lothar Göttsche, David Morrison, Marcos Mariño and Brian Rice for their comments on earlier versions of this paper. The authors thank Michael Griffin who simplified the deduction of Theorem 1.1 from Theorem 4.26. The authors also thank the referee for many helpful suggestions which improved the paper.

The second author thanks the generous support of the National Science Foundation, the Manasse family, and the Hilldale Foundation and the Candler Fund.

## 2 SO(3)–Donaldson invariants on $\mathbb{C}P^2$

In this section we recall basic facts about the SO(3)–Donaldson invariants on  $\mathbb{C}P^2$  (cf Freed and Uhlenbeck [17], Göttsche [20] and Kotschick [25]), and we recall a closed formula expression for these invariants which is due to Göttsche [19]. We also discuss briefly the SO(3)–Donaldson theory with massless monopoles.

### 2.1 The generating function

The Fubini–Study metric  $g$  on  $\mathbb{C}P^2$  is Kähler with Kähler form  $K = (i/2)g_{a\bar{b}} dz^a \wedge d\bar{z}^{\bar{b}}$ . We denote the first Chern class of the dual of the hyperplane bundle over  $\mathbb{C}P^2$  by  $H = K/\pi$ , so that  $\int_{\mathbb{C}P^2} H^2 = 1$ ,  $c_1(\mathbb{C}P^2) = 3H$ , and  $p_1(\mathbb{C}P^2) = 3H^2$ . The Poincaré dual  $\check{H}$  of  $H$  is a generator of the rank-one homology group  $H_2(\mathbb{C}P^2)$ . SO(3)–bundles on four-dimensional manifolds are classified by the second Stiefel–Whitney class  $w_2(P) \in H^2(\mathbb{C}P^2; \mathbb{Z}_2)$  and the first Pontrjagin class  $p_1(P) \in H^4(\mathbb{C}P^2)$ , such that

$$p_1(P)[\mathbb{C}P^2] \equiv w_2^2(P)[\mathbb{C}P^2] \pmod{4}.$$

Since  $\mathbb{C}P^2$  is simply connected there is an integer class  $\alpha \in H^2(\mathbb{C}P^2)$  whose mod-two reduction is  $w_2(P)$ . Then, there is a smooth complex two-dimensional vector bundle  $\xi \rightarrow \mathbb{C}P^2$  with the Chern classes  $c_1(\xi) = \alpha$  and  $c_2(\xi)$ , such that  $c_1^2(\xi) - 4c_2(\xi) = p_1(P)$ . The bundle  $\xi$  can be reduced to an SU(2)–bundle if and only if  $c_1(\xi) = 0$ . Thus, a SO(3)–bundle which does not arise as the associated bundle for the adjoint representation of a SU(2)–bundle has to satisfy  $w_2(P) \neq 0$ . From now on, we will always assume  $c_1(\xi) = -H$  and  $c_2(\xi) = kH^2$ . Once a Hermitian metric on  $\xi$  is fixed, let  $\mathfrak{A}$  be the set of compatible unitary connections  $d_A = d + A$  on  $\xi$ , and  $\mathfrak{G}$  the group of gauge transformations. The metric on  $\mathbb{C}P^2$  gives rise to a Hodge star operator  $*$ :  $\Lambda_{\mathbb{C}P^2}^p \rightarrow \Lambda_{\mathbb{C}P^2}^{4-p}$  with  $*^2 = (-1)^p$ . On  $p$ –forms, the adjoint operator of  $d_A$  is  $d_A^* = - * d_A *$ . Let  $\text{End}_0 \xi$  be the bundle of the traceless endomorphisms of  $\xi$ . In terms of the curvature  $F_A \in \Lambda_{\mathbb{C}P^2}^2(\text{End}_0 \xi)$  on  $\xi$ , the Pontrjagin number is given by

$$(2) \quad (c_1^2(\xi) - 4c_2(\xi))[\mathbb{C}P^2] = -\frac{1}{4\pi^2} \int_{\mathbb{C}P^2} \text{tr}(F_A \wedge F_A).$$

A connection is called *antiselfdual* if the curvature satisfies  $F_A^+ = 0$  (ie  $*F_A = -F_A$ ). A connection is called *reducible* if the bundle  $P \rightarrow \mathbb{C}P^2$  reduces to the direct sum  $\lambda \oplus \epsilon$  of a line bundle  $\lambda \rightarrow \mathbb{C}P^2$  and a trivial oriented real line bundle  $\epsilon \rightarrow \mathbb{C}P^2$ ; otherwise it is called *irreducible*. We denote by  $\mathfrak{A}^*$  the space of irreducible connections. Since the gauge group  $\mathfrak{G}$  acts freely on  $\mathfrak{A}^*$ , the space  $\mathfrak{B}^* = \mathfrak{A}^*/\mathfrak{G}$  is a Banach manifold.

The action functional of the Yang–Mills theory is

$$(3) \quad \int_{\mathbb{C}P^2} \text{tr} (F_A \wedge *F_A),$$

and has the Euler–Lagrange equation  $d_A^*F_A = 0$ . For  $k > 0$ , the Yang–Mills action (3) is minimized by the anti-selfdual connections. The moduli space of anti-selfdual irreducible connections with  $c_2(\xi)[\mathbb{C}P^2] = k$  modulo gauge transformations is denoted by  $\mathfrak{M}(-1, k)$ . The Atiyah–Singer complex of  $d_A$  in dimension four is the three term complex

$$(4) \quad \Lambda^0_{\mathbb{C}P^2}(\xi) \xrightarrow{d_A} \Lambda^1_{\mathbb{C}P^2}(\xi) \xrightarrow{\pi_+ d_A} \Lambda^{2+}_{\mathbb{C}P^2}(\xi),$$

where the map  $\pi_+$  is the orthogonal projection on the self-dual two-forms. It is a complex if  $F_A$  is anti-selfdual. The associated elliptic operator of Laplace type is a Fredholm operator, and the index of the elliptic complex equals  $\text{ind } d_A = 2(4k - 1) - 3(1 + b_2^+)$ . A local model of a regular neighborhood of  $[A] \in \mathfrak{M}(-1, k)$  is given by the intersection of the slice  $\ker d_A^* \in T_A \mathfrak{A}^*$ , which is locally orthogonal to pure gauge transformations, with  $\ker(\pi_+ d_A)$ , which is the linearization of  $F_A^+ = 0$ . Since  $c_1^2(\xi) - 4c_2(\xi) \not\equiv 0 \pmod 8$  the moduli spaces  $\mathfrak{M}(-1, k)$  of rank-two stable vector bundles on  $\mathbb{C}P^2$  are smooth, projective varieties of dimension  $2d_k = 8k - 8$ ; see Strømme [49].

**Remark** It is known that the moduli space  $\mathfrak{M}(c_1, c_2)$  of rank-two stable vector bundles  $\xi$  over  $\mathbb{C}P^2$  with Chern classes  $c_1, c_2$  only depends on the discriminant  $c_1^2 - 4c_2$ , with the discriminant being negative for stable bundles. The isomorphism between  $\mathfrak{M}(c_1, c_2)$  and  $\mathfrak{M}(c_1 - 2r, c_2 - r c_1 + r^2)$  is given by twisting  $\xi \mapsto \xi \otimes \mathcal{O}_{\mathbb{C}P^2}^*(r)$  by Klyachko [24].

On  $\mathbb{C}P^2 \times \mathfrak{B}^*$ , there exists a universal bundle with a  $\text{SO}(3)$ –connection (see Atiyah and Singer [4]); it is the vector bundle  $\mathfrak{P}$  with the structure group  $\text{SO}(3)$  in the commutative diagram:

$$(5) \quad \begin{array}{ccc} P \times \mathfrak{A}^* & \rightarrow & \mathfrak{P} = (P \times \mathfrak{A}^*)/\mathfrak{G} \\ \downarrow & & \downarrow \\ \mathbb{C}P^2 \times \mathfrak{A}^* & \rightarrow & \mathbb{C}P^2 \times \mathfrak{B}^* \end{array}$$

where the action of  $\mathfrak{G}$  and  $\text{SO}(3)$  on  $P \times \mathfrak{A}^*$  commutes. Using the connection  $A \in \mathfrak{A}^*$  on  $P$  and the canonical connection on  $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ , one defines a  $\mathfrak{G}$ –invariant connection on  $P \times \mathfrak{A}^* \rightarrow \mathbb{C}P^2 \times \mathfrak{A}^*$ . The connection and its curvature  $\mathfrak{F}$  descend to a connection and a curvature form on the quotient bundle. The Pontrjagin number of the universal



bundle can be decomposed into its components

$$(6) \quad p_1(\mathfrak{P}) = -\frac{1}{4\pi^2} \operatorname{tr}(\mathfrak{F} \wedge \mathfrak{F}) = -4 \sum_{r=0}^4 \mathfrak{W}^{r,4-r},$$

with  $\mathfrak{W}^{r,4-r} \in \Lambda_{\mathbb{C}P^2 \times \mathfrak{B}^*}^{r,4-r}$ . We evaluate the class  $-\frac{1}{4}p_1(\mathfrak{P})$  on the 2–cycle  $[\check{H}] \in H_2(\mathbb{C}P^2)$  and the 0–cycle  $[\text{pt}] \in H_0(\mathbb{C}P^2)$  to obtain

$$(7) \quad \mu(\check{H}) = \int_{\check{H}} \mathfrak{W}^{2,2} \in H^2(\mathfrak{B}^*; \mathbb{Q}), \quad \mu(\text{pt}) = \mathfrak{W}^{0,4}(\text{pt}) \in H^4(\mathfrak{B}^*; \mathbb{Q}).$$

By applying [14, Proposition 5.1.15] to  $\mathbb{C}P^2$  we obtain:

**Proposition 2.1** *The rational cohomology ring  $H^*(\mathfrak{B}^*; \mathbb{Q})$  for  $\mathbb{C}P^2$  is a polynomial algebra generated by the four-dimensional generator  $\mu(\text{pt})$  and the two-dimensional generator  $\mu(\check{H})$ . In particular, there is no odd-dimensional cohomology of  $\mathfrak{B}^*$ .*

Furthermore, it is known [49, Section 2] that for  $\mathfrak{M}(-1, k)$  there is a universal complex rank-two vector bundle  $\Xi \rightarrow \mathbb{C}P^2 \times \mathfrak{B}^*$  lifting the universal bundle  $\mathfrak{P}$  which is defined uniquely up to a linear bundle lifted from  $\mathfrak{B}^*$  [49].

**Lemma 2.2** *For  $\mathbb{C}P^2$ , it follows that as an element of  $H^*(\mathbb{C}P^2 \times \mathfrak{B}^*; \mathbb{Q})$  we have*

$$(8) \quad [p_1(\mathfrak{P})] = (1 - 4k) H^2 - 4H \wedge \mu(\check{H}) - 4\mu(\text{pt}).$$

*There is a unique choice for  $\Xi \rightarrow \mathbb{C}P^2 \times \mathfrak{B}^*$  such that*

$$(9) \quad [c_2(\Xi)] = k H^2 + H \wedge \mu(\check{H}) + \mu(\text{pt}), \quad [c_1(\Xi)] = -H.$$

**Proof** Since there is no odd-dimensional cohomology for  $\mathfrak{B}^*$  and  $\mathbb{C}P^2$ , it follows from the Künneth isomorphism that

$$H^4(\mathbb{C}P^2 \times \mathfrak{B}^*; \mathbb{Q}) \cong \bigoplus_{i+j=2} H^{2i}(\mathbb{C}P^2; \mathbb{Q}) \otimes H^{2j}(\mathfrak{B}^*; \mathbb{Q}).$$

Thus, as elements of  $H^*(\mathfrak{B}; \mathbb{Q})$  we have  $[\mathfrak{W}^{1,3}] = [\mathfrak{W}^{3,1}] = 0$  in Equation (6). If we denote the exterior derivatives on  $\mathbb{C}P^2$  and  $\mathfrak{B}^*$  by  $d$  and  $d$  respectively, it follows that  $(d + d)\mathfrak{W}^{\bullet,4-\bullet} = 0$ . For dimensional reason, we have  $d\mathfrak{W}^{4,0} = 0$ . However,  $H^0(\mathfrak{B}^*; \mathbb{Q}) \cong \mathbb{Q}$  is trivial. Hence,  $\mathfrak{W}^{4,0}$  is constant on each connected component of  $\mathfrak{B}^*$ . The restriction to  $\mathbb{C}P^2$  yields  $[-4\mathfrak{W}^{4,0}] = (1 - 4k)H^2$ . Similarly, since  $d\mathfrak{W}^{0,4} = 0$  it must be constant along  $\mathbb{C}P^2$ . From the definition (7), it follows that  $[-4\mathfrak{W}^{0,4}] = -4\mu(\text{pt})$ . Finally,  $[\mathfrak{W}^{2,2}] = H \wedge \mu(\check{H})$  since by Poincaré duality we have

$$\int_{\check{H}} \mathfrak{W}^{2,2} = \int_{\mathbb{C}P^2} H \wedge \mathfrak{W}^{2,2} = \mu(\check{H}) \int_{\mathbb{C}P^2} H \wedge H = \mu(\check{H}).$$

Since there exists a universal bundle  $\Xi \rightarrow \mathbb{C}P^2 \times \mathfrak{B}^*$  its Chern classes have to satisfy  $p_1(\mathfrak{B}) = c_1^2(\Xi) - 4c_2(\Xi)$ . Thus, we have  $c_1(\Xi) = -H + 2\delta\mu(\check{H})$  and  $c_2(\Xi) = kH^2 + (1 - \delta)H \wedge \mu(\check{H}) + \mu(\text{pt}) + \delta^2\mu(\check{H})^2$ . We normalize the universal bundle by setting  $\delta = 0$ . □

Classes of the de Rham cohomology of  $\mathfrak{M}(-1, k)$  are obtained from  $H^\bullet(\mathfrak{B}^*)$  by restriction which we will still denote by  $\mu(\check{H})$  and  $\mu(\text{pt})$ . To define the integration over the moduli space, a compactification of the moduli space of antiselfdual instantons is needed. Such a compactification was introduced by Donaldson [14], but since it is based on Uhlenbeck’s results it is usually called the Uhlenbeck compactification. It was shown that the  $\mu$ -map in (7) extends over the compactification  $\overline{\mathfrak{M}}(-1, k)$ . For  $2d_k = 4m + 2n$ , the *Donaldson invariants* are defined as

$$(10) \quad \Phi_{k,m,n} = \int_{\overline{\mathfrak{M}}(-1,k)} \mu(\text{pt})^m \wedge \mu(\check{H})^n.$$

This map  $\Phi$  can be extended to a linear function from the graded algebra  $\text{Sym}_*(H_0(\mathbb{C}P^2) \oplus H_2(\mathbb{C}P^2))$ , where the elements of  $H_i(\mathbb{C}P^2)$  have degree  $(4 - i)/2$  to  $\mathbb{Q}$ . It is known that the  $\text{SO}(3)$ -Donaldson invariants are topological invariants of  $\mathbb{C}P^2$ . For  $z \in \text{Sym}_*(H_0(\mathbb{C}P^2) \oplus H_2(\mathbb{C}P^2))$ , there is a definition of a geometric representative  $\mathcal{V}(z) \subset \mathfrak{M}(-1, k)$  for  $\mu(z)$  [28]. If  $\overline{\mathcal{V}}(z)$  is the closure of  $\mathcal{V}(z)$  in  $\overline{\mathfrak{M}}(-1, k)$ , then the Donaldson invariants are

$$(11) \quad \Phi_k(z) = \begin{cases} \#(\overline{\mathcal{V}}(z) \cap \overline{\mathfrak{M}}(-1, k)) & \text{if } 4m + 2n = 8(k - 1), \\ 0 & \text{otherwise.} \end{cases}$$

**Definition 2.3** The formal power series

$$(12) \quad \mathbf{Z}(p, S) = \sum_{k \geq 1} \sum_{m,n \geq 0} \Phi_{k,m,n} \frac{p^m}{m!} \frac{S^n}{n!}$$

is the generating function for the  $\text{SO}(3)$ -Donaldson invariants of  $\mathbb{C}P^2$ .

Using the blowup formula for the Donaldson invariants, Göttsche [19] and Göttsche, Nakajima and Yoshioka [21] derived a closed formula expression for  $\Phi_{k,m,n}$ . Göttsche’s work was based on earlier work with Ellingsrud [16] and Zagier [22], and it extended the results previously obtained by Kotschick and Lisca [26] up to an overall sign convention. We state [19, Theorem 3.5(1)] in terms of the Jacobi theta-functions  $\vartheta_2, \vartheta_3, \vartheta_4$  which are used in theoretical physics, and give a formulation that is in a convenient form with respect to  $\text{pt}$  and  $\check{H}$ . In this way, we obtain a closed formula expression for  $\Phi_{k,m,n}$ , which we shall later show equals the Moore–Witten prediction based on the  $u$ -plane integral.

**Theorem 2.4** (Göttsche [19]) *Assuming the notation and hypotheses above, then we have that the only nonvanishing coefficients in the generating function in (12) satisfy*

$$(13) \quad \Phi_{k,m,2n} = \sum_{l=0}^n \sum_{j=0}^l \frac{(-1)^{n+j}}{2^{l-3} 3^l} \frac{(2n)!}{(2n-2l)! j!(l-j)!} \\ \times \text{Coeff}_{q^0} \left( \frac{\vartheta_4^8(\tau) [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]^{m+j}}{[\vartheta_2(\tau) \vartheta_3(\tau)]^{2m+2n+3}} E_2^{l-j}(\tau) F_{2(n-l)}(\tau) \right),$$

where  $m, n \in \mathbb{N}_0$ ,  $2(k-1) = m+n$ ,  $\text{Coeff}_{q^0}$  is the constant term in a series expansion in  $q = \exp(2\pi i \tau)$ . The series  $F_l(\tau)$  are defined in (93), and the Jacobi  $\vartheta$ -functions are

$$\vartheta_2(\tau) = 2\Theta_2\left(\frac{\tau}{8}\right), \quad \vartheta_3(\tau) = \Theta_3\left(\frac{\tau}{8}\right), \quad \vartheta_4(\tau) = \Theta_4\left(\frac{\tau}{8}\right),$$

where  $\Theta_2, \Theta_3, \Theta_4$  are defined in (90)-(92). Also, we have that  $E_2(\tau)$  is the normalized Eisenstein series

$$E_2(\tau) := 1 - 24 \sum_{n=1}^{\infty} \sum_{d|n} dq^n.$$

**Remark** Equation (13) uses the original sign convention of [16; 26] for the Donaldson invariants in which the Donaldson invariants for  $d_k = 4$  are all negative.

**Proof** Table 1 summarizes the quantities used by Göttsche [19, Theorem 3.5(1)] and in this article.

Göttsche	present paper	Göttsche	present paper
$z$	$S$	$\theta(\tau)$	$\vartheta_4(\tau)$
$x$	$p$	$f(\tau)$	$\frac{1}{2\sqrt{i}} \vartheta_2(\tau) \vartheta_3(\tau)$
$n$	$2\beta + 1, \beta \geq 0$	$\Delta^2(2\tau)/(\Delta(\tau) \Delta(4\tau))$	$-16 \frac{\vartheta_4^8(\tau)}{[\vartheta_2(\tau) \vartheta_3(\tau)]^4}$
$a$	$2\alpha, \alpha \geq \beta + 1$	$G_2(2\tau)$	$-\frac{1}{24} E_2(\tau)$
$\tau$	$\frac{\tau - 1}{2}$	$e_3(2\tau)$	$\frac{1}{12} [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]$
$q$	$-q^{1/2}$	$\frac{-3i e_3(2\tau)}{f(\tau)^2}$	$\frac{\vartheta_2^4(\tau) + \vartheta_3^4(\tau)}{[\vartheta_2(\tau) \vartheta_3(\tau)]^2}$

Table 1

We use

$$\begin{aligned} &\left(\frac{n\sqrt{i}}{2f(\tau)}\right)^{2(n-l)}\left(-\frac{i}{2f(\tau)^2}(2G_2(2\tau)+e_3(2\tau))\right)^l \\ &= \frac{(-1)^{n+l}}{2^l 3^l} (2\beta + 1)^{2(n-l)} \frac{(-E_2(\tau) + [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)])^l}{[\vartheta_2(\tau)\vartheta_3(\tau)]^{2n}}. \end{aligned}$$

An expansion of the exponential in [19, Theorem 3.5(1)] then yields (13). □

### 2.2 Donaldson theory with monopoles

Since  $w_2(\mathbb{C}P^2) = H$ ,  $\mathbb{C}P^2$  is not a spin manifold. For the  $\text{spin}_{\mathbb{C}}$ -structure on  $\mathbb{C}P^2$  with the canonical class  $c = 3H$ , we denote by  $S_{\mathbb{C}}^{\pm} \rightarrow \mathbb{C}P^2$  the complex spinor bundles of positive and negative chirality. In other words, for the spinor decomposition  $T_{\mathbb{C}}\mathbb{C}P^2 = S_{\mathbb{C}}^{-} \otimes S_{\mathbb{C}}^{+}$  we have  $c = c_1(\det S_{\mathbb{C}}^{\pm}) = 3H$ . The vacua of the  $\text{SO}(3)$ -Donaldson theory with  $2N_f$  massless monopoles are given by a smooth connection  $A \in \mathfrak{A}$  and  $N_f$  complex sections  $\Phi_i \in C^{\infty}(\mathbb{C}P^2, S_{\mathbb{C}}^{+} \otimes \xi)$  such that

$$(14) \quad F_A^{+} = \sum_{i=0}^{N_f} [\Phi_i \otimes \bar{\Phi}_i]_{00}, \quad \not{D}_A \Phi_i = 0 \quad \text{for } 1 \leq i \leq N_f,$$

where  $\not{D}_A :: C^{\infty}(\mathbb{C}P^2, S_{\mathbb{C}}^{+} \otimes \xi) \rightarrow C^{\infty}(\mathbb{C}P^2, S_{\mathbb{C}}^{-} \otimes \xi)$  is the Dirac operator coupled to  $\xi$ . Here  $[\Phi \otimes \bar{\Phi}]_{00}$  is the double trace free component in  $\text{End } \xi \otimes \text{End}_{\mathbb{C}} S_{\mathbb{C}}^{+}$ , and we have used the identification  $\text{End}_{0,\mathbb{C}} S_{\mathbb{C}}^{+} \cong \Lambda_{\mathbb{C}P^2}^{2+}$ . Due to the positive scalar curvature of the Fubini–Study metric, the Weizenbock formula implies that  $\ker \not{D}_A$  always vanishes whence  $\Phi_i = 0$ . However, a reducible connection may induce a nonsurjective  $\not{D}_A$ , thus instigating the study of the vector bundle formed by the cokernel of  $\not{D}_A$  or equivalently  $\ker \not{D}_A^*$ , called the obstruction bundle. Since the kernel of  $\not{D}_A$  vanishes for the Fubini–Study metric on  $\mathbb{C}P^2$ , the obstruction bundle is the index bundle  $\text{Ind } \not{D}^* \rightarrow \mathfrak{B}^*$  in  $K(\mathfrak{B}^*)$ . The fibers of the obstruction bundle are obstruction spaces to the existence of a section for the Dirac operator.

**2.2.1 The index bundle of the twisted Dirac operator** In light of the isomorphism between  $\mathfrak{M}(c_1, c_2)$  and  $\mathfrak{M}(c_1 - 2r, c_2 - r, c_1 + r^2)$ , we will again restrict ourselves to the rank-two stable vector bundles  $\xi \rightarrow \mathbb{C}P^2$  with  $c_1(\xi) = -H, c_2(\xi) = kH^2$  and investigate the index bundle of  $\not{D}_A$  for the  $\text{spin}_{\mathbb{C}}$ -structure with the canonical class  $c = (2r + 1)H$ . The numerical index of  $\not{D}_A^*$  is computed by the Atiyah–Singer index theorem

$$\text{ind } \not{D}_A^* = - \int_{\mathbb{C}P^2} e^{c/2} \hat{A}(\mathbb{C}P^2) \text{ch}(\xi) = k - r,$$

where  $\widehat{A}(\mathbb{C}P^2) = 1 - p_1(\mathbb{C}P^2)/24$  and  $\text{ch}(\xi) = \text{rk}(\xi) + c_1(\xi) + c_1^2(\xi)/2 - c_2(\xi)$ . Furthermore, the Chern character of the index bundle can be computed from the family index theorem [4]:

**Theorem 2.5** *The Chern character of the index bundle is*

$$(15) \quad \text{ch}(\text{Ind } \mathcal{D}^*) = - \int_{\mathbb{C}P^2} e^{c/2} \widehat{A}(\mathbb{C}P^2) \text{ch}(\Xi),$$

where  $\Xi$  is the universal bundle introduced in (5).  $\Xi$  has the total Chern class  $c(\Xi) = 1 + c_1(\Xi) + c_2(\Xi)$ .

The next lemma was proved in Leness [30, Proposition 3.10]:

**Lemma 2.6** *We have*

$$(16) \quad \text{ch}(\Xi) = 2e^{-H/2} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} \left[ \left( k - \frac{1}{4} \right) H^2 + H \wedge \mu(\check{H}) + \mu(\text{pt}) \right]^n.$$

For  $c = (2r + 1)H$ , we have

$$\begin{aligned} \text{ch}(\text{Ind } \mathcal{D}^*)_{[2l]} &= - \frac{(-1)^l}{(2l)!} \left[ \left( r^2 - \frac{k+l/2}{2l+1} \right) \mu(\text{pt})^l - \frac{l}{2(2l+1)} \mu(\check{H})^2 \wedge \mu(\text{pt})^{l-1} \right], \\ \text{ch}(\text{Ind } \mathcal{D}^*)_{[2l+1]} &= \frac{(-1)^l r}{(2l+1)!} \mu(\check{H}) \wedge \mu(\text{pt})^l. \end{aligned}$$

The following was also proved in [30, Theorem 1.1]:

**Theorem 2.7** *It follows that*

$$(17) \quad c_k(\text{Ind } \mathcal{D}^*) = \sum_{i+2j+2l=k} f_{i,2j,2l} (2r)^i \mu(\check{H})^i \wedge \mu(\check{H})^{2j} \wedge \mu(\text{pt})^l,$$

where the coefficients  $f_{i,2j,2l}$  are given as

$$F(x, y, z) = \sum_{i,j,l} f_{i,2j,2l} x^i y^{2j} z^{2l} = \exp \left( \frac{1}{2} x J_1(z) + \frac{1}{4} y^2 J_2(z) + J_3(z) \right),$$

and the functions  $J_i(z)$  are given by

$$\begin{aligned} J_1(z) &= \frac{\tan^{-1}(z)}{z}, \\ J_2(z) &= \frac{z - \tan^{-1}(z)}{z^3}, \\ J_3(z) &= -\frac{r^2 - k}{2} \ln(1 + z^2) + \left( k - \frac{1}{4} \right) \left( \frac{\tan^{-1}(z)}{z} - 1 \right). \end{aligned}$$

**Lemma 2.8** For  $r = 0$ , we have for all  $l \in \mathbb{N}_0$  that  $c_{2l+1}(\text{Ind } \mathcal{D}^*) = 0$ .

**Proof** We include an alternative proof as convenience for the reader. We check that  $e^{c/2} \widehat{A}(\mathbb{C}P^2) = 1 + H/2$ . If  $x_1, x_2$  are the formal Chern roots of  $\mathfrak{E}$  it follows that  $x_1 + x_2 = c_1(\mathfrak{E}) = -H$ . We compute for  $n > 2$ :

$$\text{ch}(\mathfrak{E})_{[n]} + \frac{H}{2} \text{ch}(\mathfrak{E})_{[n-1]} = \frac{1}{n!} (x_1^n + x_2^n) - \frac{1}{2(n-1)!} (x_1 + x_2)(x_1^{n-1} + x_2^{n-1}).$$

We use that  $x_2 = -(x_1 + H)$ , then expand in terms of  $H$ , and truncate the series to the order  $O(H^3)$ . We obtain

$$\text{ch}(\mathfrak{E})_{[n]} + \frac{H}{2} \text{ch}(\mathfrak{E})_{[n-1]} = \frac{[1 + (-1)^n]}{n!} x_1^n + \frac{[1 + (-1)^{n-1} + 2(-1)^n]}{2(n-1)!} H \wedge x_1^{n-1}.$$

Thus,  $\text{ch}(\mathfrak{E})_{[n]} + (H/2) \text{ch}(\mathfrak{E})_{[n-1]} = 0$  if  $n$  is odd. So we have  $\text{ch}(\text{Ind } \mathcal{D}^*)_{[2l+1]} = 0$  for all  $l$ . It follows that  $c_{2l+1}(\text{Ind } \mathcal{D}^*) = 0$  for all  $l$ . □

**Remark** Since  $\ker \mathcal{D}_A$  vanishes and the rank of the index bundle  $\text{Ind } \mathcal{D}^*$  is  $k - r$ , the dimension of  $\ker \mathcal{D}_A^*$  is  $k - r$  and the determinant line bundle  $\text{DET Ind } \mathcal{D}^* = \Lambda^{k-r} \ker \mathcal{D}^* \rightarrow \mathfrak{B}^*$  is well defined. It follows from [4] that

$$c_1(\text{DET Ind } \mathcal{D}^*) = c_1(\text{Ind } \mathcal{D}^*).$$

Computing  $c_1(\text{Ind } \mathcal{D}^*)$  for  $r = 0$  and the universal bundle  $\mathfrak{E}$  with

$$\begin{aligned} c_1(\mathfrak{E}) &= -H + 2\delta\mu(\check{H}), \\ c_2(\mathfrak{E}) &= kH^2 + (1 - \delta)H \wedge \mu(\check{H}) + \mu(\text{pt}) + \delta^2\mu(\check{H})^2. \end{aligned}$$

It follows that

$$c_1(\text{DET Ind } \mathcal{D}^*) = c_1(\text{Ind } \mathcal{D}^*) = k\delta\mu(\check{H}).$$

Thus, the choice  $\delta = 0$  in Lemma 2.2 is equivalent to the vanishing of  $c_1(\text{DET Ind } \mathcal{D}^*)$ , also called the vanishing of the local anomaly in physics. On the other hand, the normalization used in [49] was different. For  $c = (2r + 1)H$  and  $\delta = -1$  we observe that  $\text{ind } \mathcal{D}^* = k - r^2$  and

$$c_1(\text{Ind } \mathcal{D}^*) = (r^2 + r - k)\mu(\check{H}).$$

If  $\alpha, \beta, \gamma$  equal  $c_1(\text{Ind } \mathcal{D}^*)$  for  $r = 1, 0, -1$  then  $\beta = \gamma = -k\mu(\check{H})$  and  $\gamma - \alpha = 2\mu(\check{H})$ .

**2.2.2 The vanishing locus of the obstruction** For  $r = 1$ , we set

$$c_{\text{top}}(\text{Ind } \mathcal{D}^*) = c_{k-1}(\text{Ind } \mathcal{D}^*).$$

For  $r = 0$ , we set

$$c_{\text{top}}(\text{Ind } \mathcal{D}^*) = \begin{cases} c_{k-1}(\text{Ind } \mathcal{D}^*) & \text{if } k \text{ is odd,} \\ c_k(\text{Ind } \mathcal{D}^*) & \text{if } k \text{ is even.} \end{cases}$$

We also set

$$(18) \quad c_{\text{top}}(\text{Ind } \mathcal{D}^* \oplus N_f) = c_{\text{top}}^{N_f}(\text{Ind } \mathcal{D}^*).$$

We denote the Poincaré dual of the top-dimensional Chern class  $c_{\text{top}}$  by  $\mathfrak{J}(N_f, k, r) \subset \mathfrak{M}(-1, k)$ . Hence,  $\mathfrak{J}(N_f, k, r)$  is a smooth submanifold of  $\mathfrak{M}(-1, k)$  of codimension  $N_f(k - r)$  if  $r = 1$  and of codimension  $2N_f \lfloor \frac{k}{2} \rfloor$  if  $r = 0$ . We interpret  $\mathfrak{J}(N_f, k, r)$  as the vanishing locus of the obstruction for the existence of  $N_f$  sections for the Dirac operator. On the other hand,  $\mathcal{V}(c_{\text{top}})$  is the geometric representative for the  $\mu$ -class used in the definition of the Donaldson invariants.

**2.2.3 The  $\mathcal{N} = 2$  invariants for  $N_f = 2, 3, 4$**  Using the Euler class, for  $N_f = 2, 3$  we define the invariants

$$(19) \quad \Phi_{k,m,n}^{N_f,c,0} = \int_{\overline{\mathfrak{M}}(-1,k)} \mu(\text{pt})^m \wedge \mu(\check{H})^n \wedge c_{\text{top}}(\text{Ind } \mathcal{D}^* \oplus N_f).$$

Using the closure  $\overline{\mathfrak{J}}(N_f, k)$  of the vanishing locus in the Uhlenbeck compactification  $\overline{\mathfrak{M}}(-1, k)$ , we define the invariants

$$(20) \quad \Phi_{k,m,n}^{N_f,c,1} = \int_{\overline{\mathfrak{J}}(N_f,k,r)} \mu(\text{pt})^m \wedge \mu(\check{H})^n.$$

For a  $z \in \text{Sym}_*(H_0(\mathbb{C}P^2) \oplus H_2(\mathbb{C}P^2))$ , we can write the invariants as the following intersection numbers

$$(21) \quad \begin{aligned} \Phi_k^{N_f,c,0}(z) &= \#(\overline{\mathcal{V}}(c_{\text{top}}) \cap \overline{\mathcal{V}}(z) \cap \overline{\mathfrak{M}}(-1, k)), \\ \Phi_k^{N_f,c,1}(z) &= \#(\overline{\mathfrak{J}}(N_f, k, r) \cap \overline{\mathcal{V}}(z) \cap \overline{\mathfrak{M}}(-1, k)). \end{aligned}$$

The intersections of  $\mathcal{V}(c_{\text{top}})$  and  $\mathfrak{J}$  with any compact cycle in  $Z \subset \mathfrak{M}(-1, k)$  will be cobordant. Hence, the intersection numbers will be the same. If  $Z \subset \mathcal{V}(z) \cap \mathfrak{M}(-1, k)$  is a noncompact cycle, both  $\mathcal{V}(c_{\text{top}}) \cap Z$  and  $\mathfrak{J} \cap Z$  can still be compact. However, the cobordism need not to be compactly supported. In this case, their closures in the enlarged Uhlenbeck compactification are no longer cobordant, and the intersection numbers  $\#(\mathcal{V}(c_{\text{top}}) \cap Z)$  and  $\#(\mathfrak{J} \cap Z)$  differ. An error term arises which we denote by

$$(22) \quad \mathcal{E}_k^{N_f,c}(z) = \Phi_k^{N_f,c,1}(z) - \Phi_k^{N_f,c,0}(z).$$

If  $\mathcal{V}(z)$  has dimension greater than or equal to four, then the closure of  $\mathcal{V}(z)$  in the Uhlenbeck compactification will intersect the lower strata, and there is a nonvanishing error term. The lower strata of  $\tilde{\mathfrak{M}}(-1, k)$  have the form  $\mathfrak{M}(-1, k - l) \times \Sigma$  where  $\Sigma \subset \text{Sym}^l(\mathbb{C}P^2)$  is a smooth stratum. Thus, the error term is a polynomial expression in the  $N_f = 0$  Donaldson invariants with smaller instanton numbers  $k' < k$  and

$$(23) \quad \mathcal{E}_k^{N_f, c}(z) = P(\{\Phi_{k'}(z')\}_{k', z'}),$$

where  $\text{deg } z' = k'$ .

**Definition 2.9** For  $N_f = 2, 3$ , the formal power series

$$(24) \quad \mathbf{Z}^{N_f, c}(p, S) = \sum_{k \geq 1} \sum_{m, n \geq 0} \Phi_{k, m, n}^{N_f, c, 1} \frac{p^m}{m!} \frac{S^n}{n!}$$

is the generating functions for the  $\text{SO}(3)$ -Donaldson invariants with  $2N_f$  massless monopoles of  $\mathbb{C}P^2$  for the  $\text{spin}_{\mathbb{C}}$ -structure with the canonical class  $c = (2r + 1)H$ .

**Lemma 2.10** For  $N_f = 2, 3$ ,  $r = 0$ ,  $a = 0, 1$ , and  $n = 2l + 1$ , we have  $\Phi_{k, m, 2l+1}^{N_f, c, a} = 0$  for all integers  $k, l$ .

**Proof** For  $a = 0$  it follows from Lemma 2.8 that  $c_{\text{top}}$  is a form of a degree divisible by four. The claim then follows from the dimension of the moduli space and the fact that  $\mu(\text{pt})$  and  $\mu(\check{H})$  have degree four and two. Since the dimension of the moduli space  $\mathfrak{M}(-1, k)$  equals  $8(k - 1)$  it follows  $4m + 2n \equiv 0 \pmod{4}$  whence  $n$  is even. The case  $a = 1$  follows similarly.  $\square$

We have the following Lemma:

**Lemma 2.11** Assume  $c = H$ ,  $r = 0$ . For  $k$  even and  $m + n + 2 = k$ , we have

$$(25) \quad \Phi_{k, m, 2n}^{2, c, 0} = \sum_{j+l=k} \left( \sum_{\substack{j_1+j_2=j \\ l_1+l_2=l}} f_{0, 2j_1, 2l_1} \cdot f_{0, 2j_2, 2l_2} \right) \Phi_{k, m+l, 2(n+j)}.$$

For  $k$  even and  $2m + 2n + 4 = k$ , we have

$$(26) \quad \Phi_{k, m, 2n}^{3, c, 0} = \sum_{j+l=3k/2} \left( \sum_{\substack{j_1+j_2+j_3=j \\ l_1+l_2+l_3=l}} f_{0, 2j_1, 2l_1} \cdot f_{0, 2j_2, 2l_2} \cdot f_{0, 2j_3, 2l_3} \right) \Phi_{k, m+l, 2(n+j)}.$$



**Proof** For  $N_f = 2$  and  $k = 2\kappa$ , we start by computing

$$c_{2\kappa}(\text{Ind } \mathcal{D}^*) = \sum_{j+l=\kappa} f_{0,2j,2l} \mu(\check{H})^{2j} \wedge \mu(\text{pt})^l.$$

It follows that

$$c_k^2(\text{Ind } \mathcal{D}^*) = \sum_{j+l=k} \left( \sum_{\substack{j_1+j_2=j \\ l_1+l_2=l}} f_{0,2j_1,2l_1} \cdot f_{0,2j_2,2l_2} \right) \mu(\check{H})^{2j} \wedge \mu(\text{pt})^l.$$

The case  $N_f = 3$  follows similarly. □

**Definition 2.12** For  $N_f = 4$ , the formal power series

$$(27) \quad \mathbf{Z}^{4,c}(q) = \sum_{k \geq 1} \Phi_{k,0,0}^{4,c,1} q^{k/2}$$

is the generating function for the SO(3)–Donaldson invariants with  $2N_f = 8$  massless monopoles of  $\mathbb{C}P^2$  for the  $\text{spin}_{\mathbb{C}}$ –structure with the canonical class  $c = (2r + 1) H$ .

**Remark** For  $c = 3 H$  and  $N_f = 4$ , the Euler class in Equation (18) is a top-dimensional form, ie, a form of degree  $8(k - 1)$  on the moduli space. The first four coefficients of (27) for  $r = 1$  were determined by Gorodenzev [18].

**Lemma 2.13** For  $c = H$ ,  $r = 0$ , we have  $\Phi_{2l,0,0}^{4,c,1} = 0$  for all integers  $l$ .

**Proof** For  $N_f = 4$  and  $c = H$ , the Euler class in Equation (18) is a form on  $\mathfrak{B}^*$  of degree  $8k$  when  $k$  is even and  $8(k - 1)$  when  $k$  is odd. Hence,  $c_{\text{top}} = 0$  when  $k$  is even and the top class is restricted to the moduli space of dimension  $8(k - 1)$ . The statement follows. □

### 2.3 The $\mathcal{N} = 4$ partition function

We will later show that the  $u$ –plane integral determines a partition function (27) for the  $\mathcal{N} = 2$  topological SO(3)–theory on  $\mathbb{C}P^2$ . The partition function satisfies a holomorphic anomaly equation analogous to the partition function for the  $\mathcal{N} = 4$  topological SO(3)–theory on  $\mathbb{C}P^2$  discussed in [50, Section 4.2]. Vafa and Witten showed that the holomorphic part of the partition function  $\mathcal{Z}$  is (cf [50, Equation (4.19)])

$$(28) \quad \mathcal{Z}^+(\tau) = \frac{1}{q^{1/4}} \sum_{k \geq 1} \chi(\overline{\mathfrak{M}}(-1, k)) q^k,$$

where  $\chi(\overline{\mathfrak{M}}(-1, k))$  is the Euler characteristic of the moduli space  $\overline{\mathfrak{M}}(-1, k)$ . The partition function  $\mathcal{Z}(\tau)$  satisfies the holomorphic anomaly equation

$$(29) \quad \frac{d}{d\bar{\tau}} \mathcal{Z}(\tau) = \frac{3}{16\pi i} \frac{\overline{\vartheta_2(2\tau)}}{\text{Im } \tau^{3/2}}.$$

They conclude that  $\mathcal{Z}(\tau) = \mathcal{Z}^+(\tau) + \mathcal{Z}^-(\tau)$  is a mock modular form of weight  $\frac{3}{2}$ :

$$\begin{aligned} \mathcal{Z}^+(\tau) &= \frac{1}{q^{1/4}} \sum_{\alpha \geq 1} 3\mathcal{H}_{4\alpha-1} q^\alpha, \\ \mathcal{Z}^-(\tau) &= \frac{3}{4\sqrt{\pi}} \sum_{l \geq 0} (2l+1) \Gamma\left(-\frac{1}{2}, \pi(2l+1)^2 \text{Im } \tau\right) q^{-(2n+1)^2/4}, \end{aligned}$$

where  $\Gamma(-1/2, t)$  is defined in (80) and  $\mathcal{H}_{4\alpha-1}$  are the Hurwitz class numbers. The comparison with Equation (28) implies  $\chi(\overline{\mathfrak{M}}(-1, k)) = 3 \mathcal{H}_{4k-1}$ . This formula was proved by Klyachko [24, Theorem 1.1]. It is well known that the generating function for the stable rank-two sheaves is equal to the product of the generating function (28) for the stable rank-two vector bundles and the contribution from the boundary. Yoshioka [54, Theorem 0.4] determined that the contribution from the boundary is equal to  $\eta^{-2\chi}(\tau)$ . He also determined directly a formula for  $\mathcal{Z}^+(\tau)/\eta^{2\chi}(\tau)$  [54, Theorem 0.1]. The first terms in the  $q$ -series expansion are

$$(30) \quad \frac{\mathcal{Z}^+(\tau)}{\eta^6(\tau)} = \frac{1}{q^{1/2}} \left( q + 9q^2 + 48q^3 + 203q^4 + 729q^5 + 2346q^6 + 6918q^7 + \dots \right).$$

The expansion

$$\begin{aligned} \frac{\mathcal{Z}^+(\tau)}{\eta^6(\tau)} &= \frac{1}{q^{1/2}} \left( 3\mathcal{H}_3 q + (18\mathcal{H}_3 + 3\mathcal{H}_7) q^2 + (81\mathcal{H}_3 + 18\mathcal{H}_7 + 3\mathcal{H}_{11}) q^3 \right. \\ &\quad \left. + (294\mathcal{H}_3 + 81\mathcal{H}_7 + 18\mathcal{H}_{11} + 3\mathcal{H}_{15}) q^4 + \dots \right) \end{aligned}$$

coincides with Equation (30).

### 3 The Seiberg–Witten curves

In this section we define and classify the Seiberg–Witten curves needed to define the regularized  $u$ -plane integral. A *Seiberg–Witten curve* is a rational elliptic surface with a section and an analytical marking whose fibers are of Kodaira-type  $I_n$  and  $I_n^*$  with  $0 \leq n \leq 4$ . Physics predicts that the Seiberg–Witten curve encodes the algebraic topology of the moduli spaces of  $\text{SO}(3)$ -Donaldson theory.

### 3.1 Weierstrass elliptic fibrations

An elliptic curve  $E$  in the Weierstrass form can be written as

$$(31) \quad y^2 = 4x^3 - g_2x - g_3,$$

where  $g_2$  and  $g_3$  are numbers such that the discriminant  $\Delta = g_2^3 - 27g_3^2$  does not vanish. In homogeneous coordinates  $[X : Y : W]$ , (31) becomes

$$WY^2 = 4X^3 - g_2XW^2 - g_3W^3.$$

One can check that the point  $P$  with the coordinates  $[0 : 1 : 0]$  is always a smooth point of the curve. We consider  $P$  the base point of the elliptic curve and the origin of the group law on  $E$ . The two types of singularities that can occur as Weierstrass cubics are a rational curve with a node, which appears when the discriminant vanishes and  $g_2, g_3 \neq 0$ , or a cusp when  $g_2 = g_3 = 0$ . Next, we look at a family of cubic curves over  $\mathbb{C}P^1$ . The family is parametrized by the base space  $\mathbb{C}P^1$  and a line bundle  $N \rightarrow \mathbb{C}P^1$ . The quantities  $g_2$  and  $g_3$  are promoted to global sections of  $N^{\otimes 4}$  and  $N^{\otimes 6}$  respectively; the discriminant becomes a section of  $N^{\otimes 12}$ . If the sections are generic enough so that they do not always lie in the discriminant locus, we obtain a Weierstrass fibration  $\pi: Z \rightarrow \mathbb{C}P^1$  with section. Each fiber comes equipped with the base point  $P$ , which defines a section  $\sigma$  of the elliptic fibration which does not pass through the nodes or cusps. The bundle  $N$  is the conormal bundle of the section  $\sigma$ . We will always assume  $N = \mathcal{O}_{\mathbb{C}P^1}(-1)$ . We will also assume that in the coordinate chart  $[u : 1] \in \mathbb{C}P^1$ , the discriminant  $\Delta$  is a polynomial of degree  $N_f + 2$  in  $u$ , and  $g_2$  and  $g_3$  are polynomials in  $u$  of degree at most 2 and 3 respectively. The space of all such Weierstrass elliptic surfaces has  $N_f + 1$  moduli. To see this, first consider the case where  $N_f = 4$ . From the seven parameters defining  $g_2$  and  $g_3$ , two can be eliminated by scaling and a shift in the  $u$ -plane. Furthermore, we can arrange the coefficient of  $g_2$  of degree two and the coefficient of  $g_3$  of degree three to be the modular invariants of an elliptic curve with periods 1 and  $\tau_0$ . The remaining four coefficients can be expressed in terms of four complex parameters [43]. In physics, they are usually denoted  $m_1, \dots, m_{N_f}$ , and called the *masses of the hypermultiplets*. We will explain their geometric meaning below (cf Fact 3.4). For  $0 \leq N_f \leq 3$ , we obtain  $4 - N_f$  additional constraints from the requirement that the discriminant has degree  $N_f + 2$ . A nontrivial elliptic fibration has to develop singular fibers; the classification of the singular fibers is part of Kodaira's classification theorem of all possible singular fibers of an elliptic fibration (cf Miranda [36]). For generic values of the masses, the polynomial  $\Delta$  has only  $N_f + 2$  simple zeros for  $|u| < \infty$ , where the elliptic fibration develops a node (ie a singular fiber of Kodaira-type  $I_1$ ). For special values for  $m_1, \dots, m_{N_f}$ , several singular fibers of Kodaira-type  $I_1$  can coalesce and form singular

fibers of Kodaira-type  $I_k$  with  $k \geq 2$ , where the discriminant has a zero of order  $k$ . The second chart over the base space is  $[1 : v] \in \mathbb{CP}^1$ . The intersection of the two charts is given by  $u = 1/v$  with  $v \neq 0$ . The Weierstrass coordinates transform according to  $x \mapsto v^2 x$  and  $y \mapsto v^3 y$ ; since  $g_2$  and  $g_3$  are sections of  $N^4$  and  $N^6$  respectively, they transform according to  $g_2 \mapsto v^4 g_2$  and  $g_3 \mapsto v^6 g_3$ . The discriminant  $\Delta \mapsto v^{12} \Delta$  becomes a polynomial in  $v$  of degree  $10 - N_f$ . From Kodaira's classification theorem it follows that the singular fiber  $E_\infty$  over  $u = \infty$  ( $v = 0$ ) is a cusp, a singular fiber of Kodaira-type  $I_{4-N_f}^*$ . In physics, a Weierstrass fibration of the type described is called a *Seiberg–Witten curve*.  $Z$  has surface singular points whenever all partial derivatives in  $u, x, y$  simultaneously vanish. Singular fibers of Kodaira-type  $I_1$  do not give rise to surface singularities, whereas all singular fibers of Kodaira-type  $I_n$ , with  $n \geq 2$ , and  $I_n^*$ , with  $n \geq 0$ , do. It is known [36, Section 4.6] that a Weierstrass fibration is rational (ie birational to  $\mathbb{CP}^2$ ), if  $g_2$  and  $g_3$  are polynomials in  $u$  of degree at most 4 and 6 respectively. The minimal resolution  $\hat{Z}$  of  $Z$  is the blow-up of  $\mathbb{CP}^2$  in nine points, and therefore has Picard number 10. The section  $\sigma$  uses up one dimension, and so the number of components of any singular fiber is at most nine since the components are always independent in the Neron–Severi group. However, not all configurations of singular fibers exist. Conversely, by contracting every component of the fiber which does not meet  $\sigma$ , we obtain the normal surface  $Z$ . A complete list of the possible configurations of singular fibers of the rational elliptic surfaces with sections that appear as Seiberg–Witten curves will be presented in Lemma 3.1. We also list the constraints on the moduli, which when substituted into the Weierstrass presentation in [43], realize the configuration of singular fibers and hence prove their existence.

### 3.2 The Mordell–Weil lattice

Let us fix a generic smooth fiber  $E$  of  $\pi: Z \rightarrow \mathbb{CP}^1$ . Let  $\text{MW}(\pi)$  be the group of sections of  $\pi$  which can be naturally identified with the rational points of  $E$  with the origin given by  $\sigma$ . The group  $\text{MW}(\pi)$  is equipped with a natural bilinear height pairing; see Oguiso and Shioda [39]. The Picard group of  $\hat{Z}$  is an unimodular lattice with signature  $(1, 9)$ . The section  $\sigma$  and the smooth fiber class  $E$  generate a two dimensional lattice that splits off the Picard lattice as  $\mathbb{I} \oplus -\mathbb{I}$ . The orthogonal complement  $(\sigma, E)^\perp$  is isomorphic to the even negative definite rank 8 unimodular lattice  $E_8$ ; see Miranda [35]. Each reducible fiber consists of a certain number of components that form a sublattice in the Picard lattice. In particular, a reducible fiber of type  $I_k$  and  $I_k^*$  generate a root lattice of type  $A_{k-1}$  and  $D_{k+4}$  respectively. We will denote the direct sum of the sublattices of the Picard group generated by the components of the fibers not meeting the section  $\sigma$  by  $\mathbf{T}$ . Thus,  $\mathbf{T}$  must have an embedding into the  $E_8$  lattice. It turns out that  $\mathbf{T}^\perp$  is isomorphic to the dual lattice of the free part

of the Mordell–Weil group  $[\text{MW}(\pi)/\text{MW}(\pi)_{\text{tor}}]^*$ . Oguiso and Shioda [39] proved a complete structure theorem of the Mordell–Weil group including the torsion subgroup. In the table of Lemma 3.1, we can therefore present  $\mathbf{T}$ ,  $\text{MW}(\pi)$  as well as their ranks, and the classification number assigned by Oguiso and Shioda.

**Lemma 3.1** *Table 2 gives a complete list of the singular fibers of the rational elliptic surfaces with sections that appear as the Seiberg–Witten curves (up to permutation of the masses).*

$N_f$	No	$r$	$\text{rk } \mathbf{T}$	$\# \text{ tor}$	$\mathbf{T}$	$\text{MW}(\pi)$	$E_\infty$	$E_{u,\text{sing}}$	mass constraints
4	9	4	4	1	$D_4$	$D_4^*$	$I_0^*$	$6I_1$	generic
4	18	3	5	1	$D_4 \oplus A_1$	$A_1^{*\oplus 3}$	$I_0^*$	$I_2, 4I_1$	$m_1 = m_2$
4	32	2	6	1	$D_4 \oplus A_2$	$A_1^* \oplus \langle \frac{1}{6} \rangle$	$I_0^*$	$I_3, 3I_1$	$m_1 = m_2 = m_3$
4	34	2	6	2	$D_4 \oplus A_1^{\oplus 2}$	$A_1^{*\oplus 2} \oplus \mathbb{Z}_2$	$I_0^*$	$2I_2, 2I_1$	$m_1 = m_2 = 0$
4	54	1	7	2	$D_4 \oplus A_3$	$\langle \frac{1}{4} \rangle \oplus \mathbb{Z}_2$	$I_0^*$	$I_4, 2I_1$	$m_1 = m_2 = m_3 = 0$
4	57	1	7	4	$D_4 \oplus A_1^{\oplus 3}$	$A_1^* \oplus (\mathbb{Z}_2)^2$	$I_0^*$	$3I_2$	$m_1 = m_2, m_3 = m_4 = 0$
3	16	3	5	1	$D_5$	$A_3^*$	$I_1^*$	$5I_1$	generic
3	30	2	6	1	$D_5 \oplus A_1$	$A_1^* \oplus \langle \frac{1}{4} \rangle$	$I_1^*$	$I_2, 3I_1$	$m_1 = m_2$
3	50	1	7	1	$D_5 \oplus A_2$	$\langle \frac{1}{12} \rangle$	$I_1^*$	$I_3, 2I_1$	$m_1 = m_2 = m_3$
3	52	1	7	2	$D_5 \oplus A_1^{\oplus 2}$	$\langle \frac{1}{4} \rangle \oplus \mathbb{Z}_2$	$I_1^*$	$2I_2, I_1$	$m_1 = m_2 = 0$
3	72	0	8	4	$D_5 \oplus A_3$	$\mathbb{Z}_4$	$I_1^*$	$I_4, I_1$	$m_1 = m_2 = m_3 = 0$
2	26	2	6	1	$D_6$	$A_1^{*\oplus 2}$	$I_2^*$	$4I_1$	generic
2	48	1	7	2	$D_6 \oplus A_1$	$A_1^* \oplus \mathbb{Z}_2$	$I_2^*$	$I_2, 2I_1$	$m_1 = m_2$
2	71	0	8	4	$D_6 \oplus A_1^{\oplus 2}$	$(\mathbb{Z}_2)^2$	$I_2^*$	$2I_2$	$m_1 = m_2 = 0$
1	46	1	7	1	$D_7$	$\langle \frac{1}{4} \rangle$	$I_3^*$	$3I_1$	-
0	64	0	8	2	$D_8$	$\mathbb{Z}_2$	$I_4^*$	$2I_1$	

Table 2

In Table 2, we have put  $r = \text{rk MW}(\pi)$ ,  $\# \text{ tor} = |\text{MW}(\pi)_{\text{tor}}|$  and  $\langle m \rangle$  for a rank 1 lattice  $\mathbb{Z}x$  with  $\langle x, x \rangle = m$ . The embedding of  $\mathbf{T}$  into  $-E_8$  is unique up to the action of the Weyl group. Table 3 shows configurations of singular fibers do not exist on a rational elliptic surface.

**Proof** Using the parametrization of Seiberg and Witten [43], one determines the zeros of  $g_2, g_3, \Delta$  and their degrees which will determine the singularities of Kodaira-type  $I_k$  and  $I_k^*$ . The obtained configuration of singular fibers appear in [39], and

$N_f$	$E_\infty$	$E_{u,\text{sing}}$	$N_f$	$E_\infty$	$E_{u,\text{sing}}$
4	$I_0^*$	$I_6$	2	$I_2^*$	$I_4$
4	$I_0^*$	$I_5, I_1$	2	$I_2^*$	$I_3, I_1$
4	$I_0^*$	$I_3, I_3$	1	$I_3^*$	$I_3$
4	$I_0^*$	$I_4, I_2$	1	$I_3^*$	$I_2, I_1$
4	$I_0^*$	$I_3, I_2, I_1$	0	$I_4^*$	$I_2$
3	$I_1^*$	$I_5$			
3	$I_1^*$	$I_3, I_2$			

Table 3

they give  $\mathbf{T}$  and  $\text{MW}(\pi)$ . The uniqueness of the embedding follows from the fact that the Seiberg–Witten curves by construction always contain a singular fiber of type  $I_k^*$  and [39, Theorem 3.3]. The impossible configurations are found in a list produced by Persson [41], and are further explained by Miranda [36].  $\square$

Let  $\mathbf{\Lambda}$  be the set of simple roots of the Mordell–Weil lattice (ie a basis with the property that every vector in the root lattice is a linear combination of elements of  $\mathbf{\Lambda}$  with all coefficients nonnegative). For each  $\alpha \in \mathbf{\Lambda}$ , we denote the corresponding section by  $s_\alpha \in \text{MW}(\pi)$ . The sections  $s_\alpha$  generate  $\text{MW}(\pi)$ , and we denote the union of all sections by  $\mathfrak{S} = \sum_{\alpha \in \mathbf{\Lambda}} s_\alpha$ .

### 3.3 Analytical marking and Seiberg–Witten one-form

We will denote by UP the base curve  $\mathbb{CP}^1$  with small open discs around the points with a singular fiber removed. In physics, UP is known as the *u-plane*. The restriction of  $\pi: Z \rightarrow \mathbb{CP}^1$  to UP, which we will still denote by  $Z$ , is a smooth four-dimensional manifold with a three-dimensional boundary with multiple components, and so  $\pi: Z \rightarrow \text{UP}$  is a proper, surjective, holomorphic map of smooth complex manifolds. We denote by  $\Omega_Z$  the canonical bundle of  $Z$ . The spaces  $H^0(\Omega_Z)$  is the space of global holomorphic two-forms on  $Z$ . We fix a smooth fiber  $E_u$  of the fibration, and thus have  $\dim H^1(E_u) = 2$ . Since a base point is given in each fiber by the section  $\sigma$ , we can choose a symplectic basis  $\{A_u, B_u\}$  of the homology  $H_1(E_u)$  with respect to the intersection form, called a *homological marking*. We cannot define  $A_u, B_u$  globally over UP. The cycles are transformed by monodromies around the points with singular fibers. However, we can define globally an analytical marking. An *analytical marking* is a choice of a nonzero holomorphic one-form on the smooth fiber  $E_u$ . We choose the canonical holomorphic differential  $dx/y \in H^0(\Omega_{E_u})$  in the coordinates of (31) where

$\Omega_{E_u}$  is the holomorphic cotangent bundle of  $E_u$ . Similarly, let  $\Omega_{UP}$  be the restriction of the canonical bundle on  $\mathbb{C}P^1$  to UP. The bundle  $\Omega_{Z/UP} = \Omega_Z \otimes (\pi^* \Omega_{UP})^{-1}$  restricts to the canonical bundle  $\Omega_{E_u}$  on each smooth fiber  $E_u$ . Thus, given the rational elliptic surface  $Z \rightarrow UP$  and the analytic marking, we can associate to it the holomorphic symplectic two-form  $\Omega_{SW} = du \wedge dx/y \in H^0(\Omega_Z)$ . The Picard–Fuchs equations on  $Z \rightarrow UP$  ask whether there is a meromorphic one-form  $\lambda_{SW} \in H^0(\Omega_{Z-\mathfrak{S}/UP})$ , called the *Seiberg–Witten differential*, such that  $\Omega_{SW}$  is the derivative of  $\lambda_{SW}$ :

**Definition 3.2** The Picard–Fuchs equations on  $Z \rightarrow UP$  for  $\lambda_{SW} \in H^0(\Omega_{Z-\mathfrak{S}/UP})$  are

$$(32) \quad d \int_{A_u} \lambda_{SW} = \int_{A_u} \Omega_{SW}, \quad d \int_{B_u} \lambda_{SW} = \int_{B_u} \Omega_{SW},$$

where  $d$  is the exterior derivative on UP.

In general,  $\lambda_{SW}$  has poles located at the intersection of sections and fibers, and it is only well-defined as a section of  $\Omega_{Z-\mathfrak{S}/UP}$ . We denote the period integrals of the elliptic fiber by  $2\omega du = \int_{A_u} \Omega_{SW}$  and  $2\omega' du = \int_{B_u} \Omega_{SW}$ , and the integrals of  $\lambda_{SW}$  by  $2\mathbf{a} = \int_{A_u} \lambda_{SW}$  and  $2\mathbf{a}_D = \int_{B_u} \lambda_{SW}$ . The modular parameter of the elliptic fiber  $E_u$  is  $\tau = \omega'/\omega \in \mathbb{H}$ . We have that  $\omega, \omega'$  are sections of a holomorphic rank-two vector bundle over UP, called the *period bundle*. The vector bundle is equipped with a flat connection  $\nabla: H^0(\Omega_{Z-\mathfrak{S}/UP}) \rightarrow H^0(\Omega_{UP}) \otimes H^0(\Omega_{Z-\mathfrak{S}/UP})$  which satisfies the equation  $\nabla \lambda_{SW} = \Omega_{SW}$ , and is known as the *Gauss–Manin connection*. The holonomy of the connection around the singular fibers determines a local system on UP and a representation of the fundamental group  $\pi_1(UP) \rightarrow SL_2(\mathbb{Z})$ ; see Atiyah [3] and Stiller [47]. The connection has regular singularities at the base points of the singular fibers; see Stiller [48] and Malmendier [33].

**Remark** The variable  $a$  used by Seiberg and Witten [42; 43] is related to  $\mathbf{a}$  by  $\mathbf{a} = 2\sqrt{2}\pi a$ .

For the Seiberg–Witten curves listed in Lemma 3.1, the following two facts were proved in [43] by an explicit computation. For their interpretation in terms of the Gauss–Manin connection on the period bundle, we refer to Shimizu [44, Lemma 3.1.5, Remark 3.2.6].

**Fact 3.3** The Picard–Fuchs equations in (32) have a solution  $\lambda_{SW} \in H^0(\Omega_{Z-\mathfrak{S}/UP})$  which is unique up to a holomorphic one-form on  $Z$ .

**Fact 3.4** For each  $s_\alpha \in MW(\pi)$  with  $\alpha \in \mathbf{\Lambda}$ , the residue  $m_\alpha = \text{Res}_{s_\alpha}(\lambda_{SW})$  is flat. Up to the permutation of the roots, the residues are the parameters  $m_1, \dots, m_{N_f}$  in the description of the rational elliptic surfaces in Lemma 3.1.

### 4 The $u$ -plane integral

In this section we define the regularized  $u$ -plane integral on  $\mathbb{CP}^2$ . The  $u$ -plane integral will depend on the choice of a Seiberg–Witten curve (ie a rational elliptic surface with a section and an analytical marking). To compute the  $u$ -plane integral, the integral has to be renormalized and then extended across the points in the  $u$ -plane with singular fibers. For  $\mathbb{CP}^2$ , we will show that the regularized  $u$ -plane integral receives contributions only from the point  $u = \infty$ , where the fiber is cuspidal. The regularized  $u$ -plane integral computes the generating function of a  $\mathcal{N} = 2$  supersymmetric, topological  $U(1)$ -gauge theories on  $\mathbb{CP}^2$ . In physics, the theory is called the *low energy effective field theory*. We compute the  $u$ -plane integrals on  $\mathbb{CP}^2$  for the Weierstrass presentations of the cases # 54, 57, 64, 71, 72 in [Lemma 3.1](#).

#### 4.1 Massless Seiberg–Witten curves for $N_f = 0, 2, 3$

**Lemma 4.1** *In [Lemma 3.1](#), the cases # 64, 71, 72 are the only elliptic modular surfaces. The Weierstrass presentations are given in [Table 4](#). The relation between the discriminant and the Dedekind eta-function is given by  $\Delta = \pi^{12} \eta^{24}(\tau)/\omega^{12}$ .  $B_\epsilon(u_{\text{sing}})$  is the  $\epsilon$ -disc around the point  $u_{\text{sing}}$  with singular fiber.*

$N_f = 0, \#64$				$u^{(0)} \in \text{UP}_\epsilon^{(0)} = \mathbb{CP}^1 - B_\epsilon(-1) - B_\epsilon(1) - B_\epsilon(\infty)$	
$E_{\text{sing}}$	$I_1$	$I_1$	$I_4^*$	$g_2^{(0)} = \frac{(u^{(0)})^2}{12} - \frac{1}{16}$	$\tau^{(0)} \in \mathbb{H}/\Gamma_0(4)$
$u_{\text{sing}}^{(0)}$	-1	1	$\infty$	$g_3^{(0)} = \frac{(u^{(0)})^3}{216} - \frac{u^{(0)}}{192}$	$u^{(0)} = \frac{1}{2} \frac{\vartheta_2^4 + \vartheta_3^4}{[\vartheta_2 \vartheta_3]^2}$
$\tau_{\text{sing}}^{(0)}$	0	2	$\infty$	$\Delta^{(0)} = \frac{1}{4096} [(u^{(0)})^2 - 1]$	$\omega^{(0)} = \sqrt{2} \pi \vartheta_2 \vartheta_3$
$N_f = 2, \#71$				$u^{(2)} \in \text{UP}_\epsilon^{(1)} = \mathbb{CP}^1 - B_\epsilon(-1) - B_\epsilon(1) - B_\epsilon(\infty)$	
$E_{\text{sing}}$	$I_2$	$I_2$	$I_2^*$	$g_2^{(2)} = \frac{(u^{(2)})^2}{12} + \frac{1}{4}$	$\tau^{(2)} = \frac{\tau^{(0)}}{2} \in \mathbb{H}/\Gamma(2)$
$u_{\text{sing}}^{(2)}$	-1	1	$\infty$	$g_3^{(2)} = \frac{(u^{(2)})^3}{216} - \frac{u^{(2)}}{24}$	$u^{(2)} = u^{(0)}$
$\tau_{\text{sing}}^{(2)}$	0	1	$\infty$	$\Delta^{(2)} = \frac{1}{64} [(u^{(2)})^2 - 1]^2$	$\omega^{(2)} = \omega^{(0)}$
$N_f = 3, \#72$				$u^{(3)} \in \text{UP}_\epsilon^{(3)} = \mathbb{CP}^1 - B_\epsilon(0) - B_\epsilon(1) - B_\epsilon(\infty)$	
$E_{\text{sing}}$	$I_4$	$I_1$	$I_1^*$	$g_2^{(3)} = \frac{(u^{(3)})^2}{12} - \frac{5u^{(3)}}{4} + \frac{11}{16}$	$\tau^{(3)} = -\frac{1}{\tau^{(0)}}$
$u_{\text{sing}}^{(3)}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\infty$	$g_3^{(3)} = \frac{(u^{(3)})^3}{216} + \frac{7(u^{(3)})^2}{48}$	$u^{(3)} = -\frac{2}{u^{(0)} - 1} - \frac{1}{2}$
$\tau_{\text{sing}}^{(3)}$	0	$-\frac{1}{2}$	$\infty$	$-\frac{29u^{(3)}}{96} + \frac{7}{64}$	$\omega^{(3)} = \frac{\pi}{2} (\vartheta_3^2 - \vartheta_4^2)$
				$\Delta^{(3)} = -\frac{1}{512} (2u^{(3)} - 1)(2u^{(3)} + 1)^4$	

Table 4



**Proof** Shioda [46] proved that elliptic modular surfaces have maximal Picard number. The Seiberg–Witten curves # 64, 71, 72 are the only extremal ones. The parametrization in the cases # 64, 71 was given by Seiberg and Witten [42; 43]. The cases # 64 and # 71 are related by a two-isogeny. We have obtained the parametrization for #72 by twisting the fibration in # 64. The Weierstrass presentations of any two surfaces that are related by a twist are related by “starring” one and “unstarring” another singular fiber of the first fibration to obtain the second. The Euler number of the singular fibers increase or decrease by six respectively. One multiplies  $g_2$  and  $g_3$  by  $(u + 1)^2$  and  $(u + 1)^3$  respectively to obtain the new  $g_2$  and  $g_3$  up to a change of the chart on  $\mathbb{C}P^1$ .  $\square$

**Remark** The rational elliptic surfaces in the cases # 54, 57 in Lemma 3.1 are said to be in *noncanonical form* since they can be obtained by twisting the elliptic fibrations # 64, 71 in Lemma 4.1. We will discuss the three cases in Lemma 4.3.

**Remark** We label the points in the  $u$ –plane with a singular fiber by an index  $s \in \{D, M, \infty\}$ . The two points in the  $u$ –plane with  $|u_{\text{sing}}| < \infty$  where the fiber is of Kodaira-type  $I_{K_s}$  are called the points where the *dyon* ( $u_{\text{sing}} = 1/2$  or  $1$ ) and *monopole* ( $u_{\text{sing}} = -1/2$  or  $-1$ ) becomes massless. There is a duality transformation  $\tau = A_s \cdot \tau_s$  with  $A_s \in \text{SL}_2(\mathbb{Z})$  such that approaching the singular point corresponds to  $\text{Im } \tau_s \rightarrow \infty$ . We set  $u_s(\tau_s) = u(\tau)$  and  $A_s \cdot \omega_s(\tau_s) = \omega(\tau)$ . Over  $u = \infty$ , the fiber is of Kodaira-type  $I_{4-N_f}^*$ , and approaching the singular point corresponds to  $\text{Im } \tau \rightarrow \infty$ . We set  $\tau_\infty = \tau$  and  $u_\infty = u$ . If  $N_f$  is fixed, we will often suppress the superindex ( $N_F$ ).

**Remark** The  $\epsilon$ –disc  $B_\epsilon(u_{\text{sing}})$  around the point  $u_{\text{sing}}$  with a singular fiber of the Kodaira-type  $I_{K_s}$  is constructed as follows: the boundary of the  $\epsilon$ –disc corresponds to the line in the  $\tau_s$ –coordinate with  $\text{Re } \tau_s \in [0; K_s]$  and  $\text{Im } \tau_s = -\ln \epsilon$ . The  $\epsilon$ –disc around the point  $u_{\text{sing}} = \infty$  with a singular fiber of the Kodaira-type  $I_{4-N_f}^*$  is constructed as follows: the boundary of the  $\epsilon$ –disc corresponds to the line in the  $\tau$ –coordinate with  $\text{Re } \tau \in [0; 4 - N_f]$  and  $\text{Im } \tau = -\ln \epsilon$ .

**Lemma 4.2** For  $|q_s| < 1$ , there are normally convergent power series expansions, where  $\alpha_{s,m}, \beta_{s,m} \in \mathbb{Q}$ , such that

$$\begin{aligned}
 (33) \quad u_s(\tau_s) &= \sum_{m \geq 0} \alpha_{s,m} q_s^{m/K_s}, \\
 \omega_s^2(\tau_s) &= \sum_{m \geq 0} \beta_{s,m} q_s^{m/K_s}.
 \end{aligned}$$

For  $|q| < 1$ , there are normally convergent power series expansions, where  $\alpha_m, \beta_m \in \mathbb{Q}$ , such that

$$(34) \quad \begin{aligned} [q^{1/(4-N_f)} u(\tau)] &= \sum_{m \geq 0} \alpha_m q^{m/(4-N_f)}, \\ [q^{-1/(4-N_f)} \omega^2(\tau)] &= \sum_{m \geq 0} \beta_m q^{m/(4-N_f)}. \end{aligned}$$

**Proof** The proof is a direct consequence of the explicit Weierstrass presentations in Lemma 4.1. □

**Lemma 4.3** *In the cases # 64, 71, 72, the Seiberg–Witten meromorphic one-form is*

$$(35) \quad \lambda_{\text{SW}} = \frac{(N_f + 2)u}{3} \frac{dx}{y} - \frac{\delta_{3,N_f}}{2} \frac{dx}{y} - 4(4 - N_f) \frac{x dx}{y},$$

and it has vanishing residues. The integrals over the A-cycle and B-cycle are

$$(36a) \quad \mathbf{a} = \frac{(N_f + 2)u}{3} \omega - \frac{\delta_{3,N_f}}{2} \omega + (4 - N_f) \frac{\pi^2 E_2(\tau)}{3\omega},$$

$$(36b) \quad \mathbf{a}_D = \frac{(N_f + 2)u}{3} \omega' - \frac{\delta_{3,N_f}}{2} \omega' + (4 - N_f) \frac{\pi^2 \tau E_2(\tau)}{3\omega} - 2\pi i \frac{4 - N_f}{\omega},$$

such that  $d\mathbf{a}/du = \omega$ ,  $d\mathbf{a}_D/du = \omega'$ , and

$$(37) \quad \omega' \mathbf{a} - \omega \mathbf{a}_D = 2\pi i (4 - N_f).$$

**Proof** Equation (35) was proved in [1]. The case  $N_f = 2$  follows from the case  $N_f = 0$  using the equation  $E_2(\tau/2) = 2E_2(\tau) - u\omega^2/\pi^2$ . Equations (36)(a) and (b) follow from the fact that the integrals of the meromorphic form  $x dx/y$  over the A-cycle and B-cycle equal  $\int_{A_u} x dx/y = -2\underline{\eta}$  and  $\int_{B_u} x dx/y = -2[\tau\underline{\eta} - \pi i/(2\omega)]$  with  $\underline{\eta} = \pi^2 E_2(\tau)/(12\omega)$ . □

**Remark** In the cases # 64, 71, 72, the Picard–Fuchs equations (32) are second order hypergeometric ordinary differential equations with three regular singularities [47].

### 4.2 The renormalization group

To describe the  $u$ -plane integral, we have to introduce the *two-observable*  $\widehat{T}$  of the low energy effective field theory for the families # 64, 71, 72 described in Lemma 4.1. To do so, we rescale (31) describing the elliptic fiber over  $u$  by  $\Lambda^6$  and introduce  $x_\Lambda = \Lambda^2 x$ ,  $y_\Lambda = \Lambda^3 y$ ,  $g_{\Lambda,2} = \Lambda^4 g_2$ ,  $g_{\Lambda,3} = \Lambda^6 g_3$ . Then, we observe that (31) is still satisfied in the rescaled quantities. We also rescale the coordinate of the base

space  $u_\Lambda = \Lambda^2 u$ . In physics, rescaling in this manner is called the action of *the renormalization group*. One of the important properties of the renormalization group is that the holomorphic symplectic two-form is not invariant but rescaled  $\Omega_{\Lambda,sw} = \Lambda \Omega_{sw}$ . We set  $\mathbf{a}_\Lambda = \Lambda \mathbf{a}$  and consider  $u_\Lambda$  a function of  $\mathbf{a}_\Lambda$ .

**Definition 4.4** The contact term of the renormalization group flow is

$$(38) \quad T = \frac{1}{4 - N_f} \left( \Lambda \frac{\partial u_\Lambda}{\partial \Lambda} \right)_{\mathbf{a}_\Lambda = \text{const}} \Big|_{\Lambda=1}.$$

**Lemma 4.5** We have that

$$T = -\frac{\pi^2 E_2(\tau)}{3\omega^2} + \frac{u}{3} + \frac{\delta_{3,N_f}}{2}.$$

**Proof** From the equation

$$d\mathbf{a}_\Lambda = \frac{\mathbf{a}_\Lambda}{\Lambda} d\Lambda + \Lambda \frac{d\mathbf{a}}{du} du = \frac{\mathbf{a}_\Lambda}{\Lambda} d\Lambda + \Lambda^2 \omega_\Lambda du$$

it follows that

$$\Lambda du_\Lambda = 2\Lambda^2 u d\Lambda + \Lambda^3 du = \left( 2u_\Lambda - \frac{\mathbf{a}_\Lambda}{\omega_\Lambda} \right) d\Lambda + \frac{\Lambda}{\omega_\Lambda} d\mathbf{a}_\Lambda,$$

whence

$$T = \frac{1}{4 - N_f} \left( 2u - \frac{\mathbf{a}}{\omega} \right) - \delta_{3,N_f} = -\frac{\pi^2 E_2(\tau)}{3\omega^2} + \frac{u}{3} + \frac{\delta_{3,N_f}}{2}. \quad \square$$

**Lemma 4.6** We have that  $T$  satisfies  $T = O(u^{-1})$  as  $u \rightarrow \infty$ .

**Proof** On the elliptic curve  $E_u$  in (31), we have the following relation between  $g_2, g_3$  and the Eisenstein series  $E_4(\tau), E_6(\tau)$  of weight four and six

$$\frac{g_3}{g_2} = \frac{(\pi^6 / (216\omega^6)) E_6(\tau)}{(\pi^4 / (12\omega^4)) E_4(\tau)} = \frac{\pi^2 E_6(\tau)}{18\omega^2 E_4(\tau)},$$

whence

$$(39) \quad T = -6 \frac{g_3 E_2(\tau) E_4(\tau)}{g_2 E_6(\tau)} + \frac{u}{3} + \frac{\delta_{3,N_f}}{2}.$$

It is easy to check from the presentation in Lemma 4.2 that near the singular fiber of type  $I_{4-N_f}^*$ , the behavior of  $q = \exp(2\pi i \tau)$  is as follows:

$$(40) \quad q = \frac{1}{u^{4-N_f}} (c_0 + O(u^{-1})) \quad (u \rightarrow \infty).$$

It follows from the presentation in [Lemma 4.1](#) that near the singular fiber of type  $I_{4-N_f}^*$ , we have the following behavior:

$$\frac{g_3}{g_2} = \frac{u}{18} + \frac{31}{12}\delta_{3,N_f} + O(u^{-1}) \quad (u \rightarrow \infty).$$

For  $N_f = 0, 1, 2$  the claim follows. Using the equation  $\Delta = \pi^{12} \eta^{24}(\tau)/\omega^{12}$ , we compute  $c_0 = -1/16$  in [\(40\)](#) for  $N_f = 3$  and case #72. It follows that

$$\begin{aligned} \frac{E_2(\tau) E_4(\tau)}{E_6(\tau)} &= \frac{(1 - 24c_0/u)(1 + 240c_0/u)}{(1 - 504c_0/u)} + O(u^{-2}) \\ &= 1 - \frac{45}{u} + O(u^{-2}) \quad (u \rightarrow \infty), \end{aligned}$$

and the claim follows from the expansion of [\(39\)](#) as  $u \rightarrow \infty$ . □

**Remark** The asymptotic behavior proved in [Lemma 4.6](#), together with the holomorphicity in  $u$ , was originally used by Moore and Witten [\[37\]](#) to define  $T$ . Losev, Nekrasov and Shatashvili [\[31\]](#) and Mariño and Moore [\[34\]](#) later showed that  $T$  can be described as the contact term of the renormalization group flow which allows a definition in terms of the geometry of the elliptic surface alone. However, for  $N_f = 3$  these two methods of determining  $T$  generally only agree up to a shift in  $u$ . The variable  $u^{(3)}$  defined in [Lemma 4.1](#) is chosen in a way that makes the two methods agree. The variable  $u^{(3)}$  relates to the variable  $u_{\text{SW}}^{(3)}$  used by Seiberg and Witten [\[43\]](#) by  $u^{(3)} = u_{\text{SW}}^{(3)} - 1/2$ .

Finally, to obtain an expression which is a modular invariant in  $\tau$ , we make the following definition.

**Definition 4.7** For the elliptic surfaces in the cases # 64, 71, 72 in [Lemma 4.1](#), the two-observable of the low energy effective field theory is

$$(41) \quad \hat{T} = -\frac{\pi^2 \hat{E}_2(\tau)}{3\omega^2} + \frac{u}{3} + \frac{\delta_{3,N_f}}{2},$$

where  $\hat{E}_2(\tau)$  is the nonholomorphic weight 2 Eisenstein series defined by

$$\hat{E}_2(\tau) = E_2(\tau) - \frac{3}{\pi \text{Im } \tau}.$$

### 4.3 The $u$ -plane integral

We briefly describe some features of the low energy effective field theory on  $\mathbb{CP}^2$ . In particular, we outline the physics argument on how the generating function is computed

using the language of determinant line bundles so that the reader can see how the idea arose in physics that elliptic integrals compute the Donaldson invariants. The bosonic fields of the classical field theory are a connection  $\mathfrak{a}$  on a line bundle  $L \rightarrow \mathbb{C}P^2$  with the curvature  $F_{\mathfrak{a}}$  and a complex valued function  $\varphi \in C^\infty(\mathbb{C}P^2; \mathbb{C})$ . A family of bosonic gauge-invariant actions is parametrized by a complex coupling  $\tau \in \mathbb{H}$  which is the modular parameter  $\tau$  of the elliptic fiber  $E_u$  of a Seiberg–Witten curve  $Z \rightarrow UP_\epsilon$  in [Lemma 4.1](#), and it is given by

$$(42) \quad \mathbf{S} = \frac{i\bar{\tau}}{16\pi} \int_{\mathbb{C}P^2} F_{\mathfrak{a}}^+ \wedge F_{\mathfrak{a}}^+ + \frac{i\tau}{16\pi} \int_{\mathbb{C}P^2} F_{\mathfrak{a}}^- \wedge F_{\mathfrak{a}}^- + \frac{i}{4} \int_{\mathbb{C}P^2} F_{\mathfrak{a}} \wedge \alpha + \frac{\text{Im } \tau}{8\pi} \int_{\mathbb{C}P^2} d\varphi \wedge *d\bar{\varphi},$$

where  $\alpha \in H^2(\mathbb{C}P^2)$  is an integer class whose mod-two reduction is  $w_2(\mathbb{C}P^2)$ . The critical points of the action are self-dual connections (ie  $F_{\mathfrak{a}}^- = 0$ ), and a constant  $\varphi$ . The field theory is connected to the SO(3)–Donaldson theory in [Section 2.1](#) through two global constraints: the topological condition  $c_1(L)[\check{H}] = (2c_2(\xi) + c_1^2(\xi))[\mathbb{C}P^2]$ , and the constraint  $\varphi = \mathbf{a}$ , where  $\mathbf{a}$  is the integral of  $\lambda_{\text{SW}}$  over the A-cycle in [Lemma 4.3](#). The first condition implies  $c_1(L) = [-\frac{1}{2\pi} F_{\mathfrak{a}}] = (2k + 1) H$ , the latter will be further explained below. If we view  $\tau$  as a function of  $\mathbf{a}$ , each critical point of the action in (42) is labeled by the discrete topological data  $k$  and the continuous modulus  $\mathbf{a}$ . The action  $\mathbf{S}$  at a critical point  $(k, \mathbf{a})$  is

$$\mathbf{S}^{(0)} = i\pi\bar{\tau} \left(k + \frac{1}{2}\right)^2 - i\pi \left(k + \frac{1}{2}\right).$$

The path integral for the supersymmetric extension of the action in (42) can be defined with mathematical rigor by the stationary phase approximation [51]. The quadratic approximation of the action around a critical point is the Hessian of the supersymmetric extension of  $\mathbf{S}$ . It determines a free field theory in the collected variations of the bosonic fields  $\tilde{\Phi}$  and fermionic fields  $\tilde{\Psi}$  of the form

$$\mathbf{S}^{(2)} = \int_{\mathbb{C}P^2} \text{vol} (\langle \tilde{\Phi}, \Delta_{(k,\mathbf{a})} \tilde{\Phi} \rangle + \langle \tilde{\Psi}, \mathcal{D}_{(k,\mathbf{a})} \tilde{\Psi} \rangle),$$

where  $\Delta_{(k,\mathbf{a})}$  is a family of second-order elliptic operators, and  $\mathcal{D}_{(k,\mathbf{a})}$  is a family of real skew-symmetric first-order operators depending on the moduli. We point out that the operators  $\Delta_{(k,\mathbf{a})}$  and  $\mathcal{D}_{(k,\mathbf{a})}$  describe the Hessian of the supersymmetric action that describes the gauge theory, but also the coupling to gravity. The functional integration over the fluctuations (ie the coordinates of the normal bundle at the critical points), is

an infinite-dimensional Gaussian integral. We define the semiclassical path integral<sup>2</sup>

$$\int [\mathcal{D}\tilde{\Phi}\mathcal{D}\tilde{\Psi}]e^{-S^{(2)}} \quad \text{to be} \quad \frac{\text{pfaff}' \mathcal{D}(k, \mathbf{a})}{\sqrt{\det' \Delta(k, \mathbf{a})}}.$$

The above expression is a section of the product of the Pfaffian line bundles over the moduli space. To proceed further, one has to be able to integrate this section over the moduli space. To do so, the line bundle needs to be flat. In physics, this is called the *vanishing of the local anomaly*. Since the topology of the moduli space is not trivial, the section can still have holonomy around nontrivial loops in the moduli space. To make sure that the bundle is globally trivial as well, the holonomy around all curves needs to be the identity. In physics, this is called the *vanishing of the global anomaly*. Finally, it is not enough to have a trivial line bundle. We also need a canonical trivialization (ie a trivializing section). All these points are in fact satisfied [32], and we regard the ratio of determinants as a function on the moduli space. We then carry out the integral over the continuous moduli, and we carry out the sum over the discrete moduli to obtain the semiclassical approximation of the *partition function*

$$(43) \quad \sum_{k \in \mathbb{Z}} \int d\mathbf{a} \wedge d\bar{\mathbf{a}} e^{-S^{(0)}} \int [\mathcal{D}\tilde{\Phi}\mathcal{D}\tilde{\Psi}]e^{-S^{(2)}} = \sum_{k \in \mathbb{Z}} \int d\mathbf{a} \wedge d\bar{\mathbf{a}} e^{-S^{(0)}} \frac{\text{pfaff}' \mathcal{D}_{k, \mathbf{a}}}{\sqrt{\det' \Delta_{k, \mathbf{a}}}}.$$

In the case which is relevant for the definition of the  $u$ -plane integral, physical considerations guarantee that the semiclassical approximation is in fact exact by Moore and Witten [37] and Witten [51], and (43) is the full quantum partition function. Moore and Witten [37] also showed that

$$(44) \quad \frac{\text{pfaff}' \mathcal{D}_{k, \mathbf{a}}}{\sqrt{\det' \Delta_{k, \mathbf{a}}}} = -\frac{4i}{(2\pi)^{3/2}} \frac{1}{\sqrt{\text{Im} \tau}} \frac{d\bar{\tau}}{d\bar{\mathbf{a}}} \left( \int_{\check{H}} F_{\alpha} \right) \frac{\hat{\Delta}^{1/8}}{\omega^{3/2}},$$

where the last term on the right side in (44) describes the coupling to gravity on  $\mathbb{C}P^2$ , which had already been determined by Witten [53]. We not only want to determine the partition function, but we also wish to construct the generating function with the insertion of the observables that correspond to the observables in (7) for the  $SO(3)$ -Donaldson theory on  $\mathbb{C}P^2$ . Moore and Witten [37] determined that  $\mu(\text{pt}) \mapsto 2u$  as a consequence of the choice  $\varphi = \mathbf{a}$ , and  $\mu(\check{H}) \mapsto \hat{T}$ , where  $\hat{T}$  was defined in (41).

<sup>2</sup>The Gaussian integral for the free field theory is actually well-defined and agrees with the ratio of the Pfaffians in the definition we give.

Thus, we obtain the *generating function*

$$(45) \quad \sum_{k \in \mathbb{Z}} \int d\mathbf{a} \wedge d\bar{\mathbf{a}} e^{-S^{(0)}} \int [D\tilde{\Phi} D\tilde{\Psi}] e^{-S^{(2)}} e^{2up+S^2\hat{T}}$$

$$= \sum_{k \in \mathbb{Z}} \int d\mathbf{a} \wedge d\bar{\mathbf{a}} e^{-S^{(0)}} \frac{\text{pfaff}' \not{D}_{k,\mathbf{a}}}{\sqrt{\det' \Delta_{k,\mathbf{a}}}} e^{2up+S^2\hat{T}}.$$

We use the formula

$$(46) \quad \sum_{k \in \mathbb{Z}} e^{i\pi(k+1/2)} \left(k + \frac{1}{2}\right) e^{-i\pi\bar{\tau}(k+1/2)^2} = i\overline{\eta^3(\tau)},$$

the substitution rule, (44) and (43) to make the following definition:

**Definition 4.8** The generating function of the massless  $N_f$  low energy effective field theory on  $\mathbb{C}P^2$  is

$$(47) \quad \tilde{\mathbf{Z}}_{\text{UP}_\epsilon}^{N_f}(p, S) = -\frac{8}{\sqrt{2\pi}} \int_{\text{UP}_\epsilon} \frac{du \wedge d\bar{u} d\bar{\tau} \Delta^{1/8}}{\sqrt{\text{Im } \tau} d\bar{u} \omega^{1/2}} e^{2up+S^2\hat{T}} \overline{\eta^3(\tau)}.$$

By  $\text{UP}_\epsilon$ , we denote the base curve  $\mathbb{C}P^1$  with small open  $\epsilon$ –discs around the points with the singular fiber of Kodaira-type  $I_{K_s}$  and the point  $u = \infty$  with the singular fiber of Kodaira-type  $I_{4-N_f}^*$ , removed. Also, we respectively let  $\tau, \Delta, \omega, \hat{T}$  denote the modular parameter, the discriminant, the period integral over the A-cycle, and the two-observable of the regular fiber defined for the families # 64, 71, 72 by (41).

**Lemma 4.9** *The generating function in (47) is well-defined.*

**Proof** For the integral to be well-defined we have to check two facts: the integral in (47) is locally well-defined since the poles of the integrand only appear at the singular locus which have been removed. Secondly, the integrand is written in terms which have monodromy in  $\tau$ . The  $\tau$ –dependent part of the integral is

$$(48) \quad \frac{\pi^{3/2}}{\sqrt{\text{Im } \tau}} \frac{d\bar{\tau}}{d\bar{u}} \frac{|\eta(\tau)|^6}{\omega^2} e^{S^2\hat{T}}.$$

It is easy to show that the term is invariant under any modular transformation  $\Gamma \subset \text{SL}_2(\mathbb{Z})$  which leaves  $u$  invariant. □

In summary, formula (42) is the bosonic part of the low energy effective field theory. A physics argument says that computing the nonabelian Donaldson invariants can be reduced to a saddle-point computation in an abelian theory given by the supersymmetric extension of (42) where the coupling constant is taken to be the  $\tau$ –parameter of a Seiberg–Witten family of curves.

### 4.4 The mock modular form

To compute the generating function, we integrate by parts using the “holomorphic part” of a weight  $1/2$  harmonic Maass form. The weight  $1/2$  harmonic Maass form is given in [Theorem 7.2](#), and the holomorphic part of a weight  $1/2$  harmonic Maass form is defined in [Equation \(82\)](#). We set  $Q(\tau) = Q^+(\tau) + Q^-(\tau)$ , where  $Q^+(\tau)$  with  $q = \exp(2\pi i \tau)$  is the holomorphic part of this Maass form. The holomorphic part has a series expansion of the form

$$\begin{aligned}
 (49) \quad Q^+(\tau) &= \frac{1}{q^{1/8}} \sum_{\alpha \geq 0} H_\alpha q^{\alpha/2} \\
 &= \frac{1}{q^{1/8}} (1 + 28q^{1/2} + 39q + 196q^{3/2} + 161q^2 + \dots).
 \end{aligned}$$

The “nonholomorphic part”  $Q^-$  is

$$\begin{aligned}
 (50) \quad Q^-(\tau) &= \frac{1}{q^{1/8}} \sum_{\alpha \geq 0} H_{-\alpha} q^{-\alpha} \\
 &= -\frac{2i}{\sqrt{\pi}} \sum_{l \geq 0} (-1)^l \Gamma\left(\frac{1}{2}, \pi \frac{(2l+1)^2}{2} \text{Im } \tau\right) q^{-(2l+1)^2/8},
 \end{aligned}$$

where  $\Gamma(1/2, t)$  is defined in [\(80\)](#). The nonholomorphic part  $Q^-$  has an exponential decay since

$$\Gamma\left(\frac{1}{2}, t\right) = \frac{e^{-t}}{\sqrt{t}} (1 + O(t^{-1})) \quad (t \rightarrow \infty).$$

Setting  $\zeta_{2k} = \exp(\pi i/k)$  it follows that  $\zeta_8^2 Q^+(\tau + 2) = Q^+(\tau)$  and  $\zeta_8 Q^-(\tau + 1) = Q^-(\tau)$ . Thanks to [Theorem 7.2](#), we know that

$$(51) \quad \sqrt{2}i \frac{d}{d\bar{\tau}} Q(\tau) = \frac{1}{\sqrt{\text{Im } \tau}} \overline{\eta^3(\tau)}.$$

We have the following analog of [\[37, \(9.18\)\]](#) for  $Q$ :

**Lemma 4.10** *The function*

$$\mathcal{E}_{1/2}^k[Q] = \sum_{j=0}^k (-1)^j \binom{k}{j} \frac{\Gamma(1/2)}{\Gamma(1/2 + j)} 2^{2j} 3^j E_2^{k-j}(\tau) \left(q \frac{d}{dq}\right)^j Q(\tau)$$

is modular of weight  $2k + 1/2$  for  $\Gamma(2) \cap \Gamma_0(4)$  and satisfies

$$(52) \quad \sqrt{2}i \frac{d}{d\bar{\tau}} \mathcal{E}_{1/2}^k[Q] = \frac{1}{\sqrt{\text{Im } \tau}} \widehat{E}_2^k(\tau) \overline{\eta^3(\tau)}.$$



**Proof** The proof of the modularity is very similar to the proof of the Lemma in [37, Section 9.1]. Let  $f$  be modular of weight  $(s, 0)$ . Cohen’s operator acting on  $f$  is given by

$$\mathcal{F}_r[f] = \sum_{j=0}^r \binom{r}{j} \frac{\Gamma(s+r)}{\Gamma(s+j)} \left(\frac{1}{2iy}\right)^{r-j} \left(\frac{d}{d\tau}\right)^j f.$$

If  $f$  is modular of weight  $(s, 0)$  for  $\Gamma(2) \cap \Gamma_0(4)$  then  $\mathcal{F}_r[f]$  is modular of weight  $(2r + s, 0)$  for  $\Gamma(2) \cap \Gamma_0(4)$ . One can easily check that

$$\mathcal{E}_{1/2}^k[Q] = \sum_{m=0}^k \binom{k}{m} \frac{\Gamma(1/2)}{\Gamma(1/2 + k - m)} \left(\frac{6i}{\pi}\right)^{k-m} \widehat{E}_2^m(\tau) \mathcal{F}_{k-m}[Q].$$

Moreover, we apply  $d/d\bar{\tau}$  to  $Q(\tau)$  in each summand. We use the fact that  $\overline{\eta^3(\tau)}$  does not depend on  $\tau$ . Using

$$\left(\frac{d}{d\tau}\right)^j \frac{1}{\sqrt{\text{Im } \tau}} = (-1)^j \frac{\Gamma(1/2 + j)}{\Gamma(1/2)} \frac{\sqrt{2i}}{(\tau - \bar{\tau})^{j+1/2}},$$

the claim follows. □

For the different coordinate patches around the cusps of Kodaira-type  $I_{4-N_f}^*$  and  $I_{K_s}$  in Lemma 4.2 we need to define the transforms of  $Q$  which we will use to integrate by parts in Equation (56). If  $F(\tau)$  is a solution to (51), then taking  $\tau \mapsto -1/\tau$  and  $\tau \mapsto \tau + n$ , implies that  $\tilde{F}(\tau) = (-1/\tau)/\sqrt{-i\tau}$  and  $\zeta_8^n F(\tau + n)$  are solutions as well. Their nonholomorphic parts equal  $Q^-(\tau)$  in Equation (50) since the transforms satisfy the same differential equation in  $\partial_{\bar{\tau}}$ .

**Lemma 4.11** *The function  $Q(\tau)$  satisfies*

$$\frac{1}{\sqrt{-i\tau}} Q\left(-\frac{1}{\tau}\right) = \frac{1}{\sqrt{-i\tau}} \zeta_8^2 Q\left(2 - \frac{1}{\tau}\right).$$

We have the following  $q$ -expansion for the holomorphic part:

$$\left[\frac{1}{\sqrt{-i\tau}} Q\left(-\frac{1}{\tau}\right)\right]^+ = \frac{1}{q^{1/8}} \left(\frac{5}{2} + \frac{111}{2}q + \frac{413}{2}q^2 + 819q^3 + \frac{4407}{2}q^4 + \dots\right).$$

**Proof** It is proved in Theorem 7.2 that

$$\begin{aligned} Q(\tau) &= -\frac{7}{2}A_{3,8}(\tau) + \frac{3}{2}A_{7,8}(\tau) - \frac{1}{2}B(\tau) + 4M(\tau) \\ &= \frac{1}{q^{1/8}}(1 + 28q^{1/2} + 39q + 196q^{3/2} + 161q^2 + \dots), \end{aligned}$$

where the functions  $A_{3,8}, A_{7,8}, B, M$  are defined in (98), (97), and (99). We have the transformation properties

$$\begin{aligned} \zeta_8 A_{3,8}(\tau + 1) &= -A_{3,8}(\tau), \\ \zeta_8 B(\tau + 1) &= B(\tau), \\ \zeta_8 A_{7,8}(\tau + 1) &= A_{7,8}(\tau), \\ \zeta_8 M(\tau + 1) &= M(\tau). \end{aligned}$$

The expansions of the modular forms in  $-1/\tau$  can be computed easily. For example, we have

$$\frac{1}{\sqrt{-i\tau}} A_{3,8} \left( -\frac{1}{\tau} \right) = -\frac{1}{2} \frac{\eta^8(\tau/2)}{\eta^7(\tau)} = \frac{1}{q^{1/8}} \left( -\frac{1}{2} + 4q^{1/2} - \frac{27}{2}q + 28q^{3/2} \dots \right),$$

and similar equations for  $B(\tau), A_{7,8}(\tau)$ . For the expansion of  $M$  we use Theorem 6.2 to show that

$$\begin{aligned} \frac{1}{\sqrt{-i\tau}} M \left( -\frac{1}{\tau} \right) &= -\frac{\sqrt{i}}{4q^{1/32}} \hat{\mu} \left( \frac{1}{2}, \frac{1}{4} - \frac{\tau}{8}; \frac{\tau}{4} \right) - \frac{1}{4\sqrt{i}q^{1/32}} \hat{\mu} \left( \frac{1}{2}, \frac{3}{4} - \frac{\tau}{8}; \frac{\tau}{4} \right) \\ &= \frac{1}{q^{1/8}} \left( \frac{1}{2}q^{1/4} - q^{1/2} + 2q^{3/4} - 3q + \dots \right). \end{aligned}$$

This proves the series expansion. The second series expansion then follows from  $\zeta_8^2 Q(\tau + 2) = Q(\tau)$ . □

**Definition 4.12** We set  $Q_\infty^{(0)}(\tau) = Q_\infty^{(2)}(\tau) = Q(\tau)$ , and

$$Q_D^{(0)}(\tau) = Q_M^{(0)}(\tau) = \frac{1}{\sqrt{-i\tau}} Q \left( -\frac{1}{\tau} \right).$$

Similarly, we define

$$Q_M^{(2)}(\tau) = \frac{1}{\sqrt{-i\tau}} Q \left( -\frac{1}{\tau} \right), \quad Q_D^{(2)}(\tau) = \frac{1}{\sqrt{-i\tau}} \zeta_8 Q \left( 1 - \frac{1}{\tau} \right).$$

For  $N_f = 3$ , we set

$$Q_\infty^{(3)}(\tau) = Q_M^{(0)}(\tau), \quad Q_M^{(3)}(\tau) = Q_\infty^{(0)}(\tau), \quad Q_D^{(3)}(\tau) = Q_D^{(0)}(\tau).$$

**Lemma 4.13** For  $\mathbf{Z}(\tau) = \eta^3(\tau)Q(\tau)$  and  $\rho(\tau) = \sqrt{2}\eta^2(2\tau)/\eta^2(\tau)$ , we have that

$$\mathbf{Z}(\tau) = \mathbf{Z}(\tau + 1) + 14\eta^4(\tau)\rho^4(\tau).$$

We have the following transformation properties under modular transformations

$$\mathbf{Z}(\tau + 2) = \mathbf{Z}(\tau), \quad \mathbf{Z}\left(-\frac{1}{\tau} + 1\right) = \tau^2 \mathbf{Z}(\tau + 1).$$

The expression

$$(53) \quad \frac{1}{\eta^4(\tau)} \sum_{k=0}^3 (-1)^k \mathbf{Z}(\tau + k) = 28\rho^4(\tau) = \frac{4q^{3/8}}{\eta(\tau)} \sum_{m=0}^{\infty} H_{2m+1} q^m$$

is a weight 0 modular form.

**Proof** The equation

$$\mathbf{Z}(\tau) = \mathbf{Z}(\tau + 1) + 14\eta^4(\tau)\rho^4(\tau)$$

follows from the proof of [Lemma 4.11](#). Using the remark following [Theorem 7.2](#) it follows that

$$\sum_{k=0}^3 (-1)^k \mathbf{Z}(\tau + k) = -14A_{3,8}(\tau)\eta^3(\tau) = 112 \frac{\eta(2\tau)^8}{\eta(\tau)^4}.$$

To prove that the indicated  $q$ -series is modular, we apply the following principle (see the [Section 5](#)): A harmonic weak Maass form whose nonholomorphic part is zero, is a weakly holomorphic modular form. To this end we now compute the contribution to the Fourier expansion of the “period integrals” to see that indeed they vanish thereby giving us a weight  $1/2$  weakly holomorphic modular form, which after division by  $\eta(\tau)$  then becomes a weakly holomorphic modular form of weight 0. To this end, notice that [\(50\)](#) and [\(49\)](#) imply that  $Q^\pm(\tau + 2) = \zeta_8^{-2} Q^\pm(\tau)$ . Since  $\eta^3(\tau + 2) = \zeta_8^2 \eta^3(\tau)$ , we have  $\mathbf{Z}(\tau + 2) = \mathbf{Z}(\tau)$ . To see that the nonholomorphic parts vanish, we check that for  $Q^-(\tau)$  in [Equation \(50\)](#) we have  $Q^-(\tau + 1) = \zeta_8^{-1} Q^-(\tau)$  whence  $\mathbf{Z}^-(\tau + 1) = \mathbf{Z}^-(\tau)$ . Thus, we obtain

$$\begin{aligned} 2[\mathbf{Z}(\tau) - \mathbf{Z}(\tau + 1)] &= 2[\mathbf{Z}^+(\tau) - \mathbf{Z}^+(\tau + 1)] \\ &= 2\eta^3(\tau) [Q^+(\tau) - \zeta_8 Q^+(\tau + 1)] \\ &= \frac{4\eta^3(\tau)}{q^{1/8}} \sum_{m=0}^{\infty} H_{2m+1} q^{m+1/2}. \end{aligned}$$

The S-duality equation for  $\mathbf{Z}(\tau)$  follows from the identity

$$(54) \quad \frac{1}{\sqrt{-i\tau}} \zeta_8 Q\left(1 - \frac{1}{\tau}\right) = -\zeta_8 Q(\tau + 1),$$

which can be proved using the results in [Section 6](#) and [Theorem 7.2](#). □

### 4.5 The regularization procedure

We now extend the integral by taking the limit  $\epsilon \rightarrow 0$  to include the contributions from the singularities. To do so we will have to *regularize* the  $u$ -plane integral. We will denote the result by

$$(55) \quad Z_{UP}^{N_f}(p, S) = -\frac{8}{\sqrt{2\pi}} \int_{\mathbb{C}P^1}^{\text{reg}} \frac{du \wedge d\bar{u}}{\sqrt{\text{Im } \tau}} \frac{d\bar{\tau}}{d\bar{u}} \frac{\Delta^{1/8}}{\omega^{1/2}} e^{2up+S^2\hat{T}} \overline{\eta^3(\tau)}.$$

**Remark** Equation (55) is the analog of [37, (9.1)] for the SO(3)-Donaldson theory and follows the same normalization.

We expand the exponential in the generating function in (47) as follows:

$$(56) \quad \begin{aligned} &\tilde{Z}_{UP_\epsilon}^{N_f}(p, S) \\ &= \sum_{m,n \geq 0} \frac{p^m}{m!} \frac{S^{2n}}{(2n)!} e^{S^2(\delta_3, N_f/2)} \sum_{k=0}^n (-1)^{k+1} \frac{2^{m+5/2} \pi^{2k+1}}{3^n} \frac{(2n)!}{k!(n-k)!} \\ &\quad \times \int_{UP_\epsilon} du \wedge d\bar{u} \frac{d\bar{\tau}}{d\bar{u}} \frac{u^{m+n-k} \Delta \eta^3(\tau)}{\omega^{2k+2}} \frac{\hat{E}_2^k(\tau) \overline{\eta^3(\tau)}}{\sqrt{\text{Im } \tau}}. \end{aligned}$$

The integrals in (56) are all of the form

$$(57) \quad \int_{UP_\epsilon} d\tau \wedge d\bar{\tau} \frac{u^{m+n-k} \Delta \eta^3(\tau)}{\omega^{2k}} \frac{\hat{E}_2^k(\tau) \overline{\eta^3(\tau)}}{\sqrt{\text{Im } \tau}}.$$

We use the substitution rule and Lemma 4.10 to write the integrand as  $d\tau \wedge d\bar{\tau} \partial_{\bar{\tau}}(\dots)$ . With  $\tau = x + iy$  and for an integrand  $f$ , we have

$$d\tau \wedge d\bar{\tau} \partial_{\bar{\tau}} f = -i dx \wedge dy (\partial_x + i\partial_y) f = -d(f dx + i f dy).$$

Thus, the integral (57) reduces to the following integral over the boundary of  $UP_\epsilon$  which is the union of the boundary of the  $\epsilon$ -discs around the points with singular fibers described in Section 4.1:

$$(58) \quad X_\epsilon := \sqrt{2i} \int_{\partial UP_\epsilon} \frac{u^{m+n-k} \Delta \eta^3(\tau)}{\omega^{2k}} \mathcal{E}_{1/2}^k[Q] dz.$$

The regularized  $u$ -plane integral is defined by

$$(59) \quad \int_{UP_\epsilon}^{\text{reg}} d\tau \wedge d\bar{\tau} \frac{u^{m+n-k} \Delta \eta^3(\tau)}{\omega^{2k}} \frac{\hat{E}_2^k(\tau) \overline{\eta^3(\tau)}}{\sqrt{\text{Im } \tau}} := \lim_{\epsilon \rightarrow 0} X_\epsilon.$$

Integrating each summand in (56) and taking the limit  $\epsilon \rightarrow 0$ , we obtain a well-defined procedure to compute (55).

### 4.6 The generating functions for $N_f = 0, 2, 3$

**Lemma 4.14** *In the notation of Lemma 4.2 and Definition 4.8, it follows that*

$$\begin{aligned} \tilde{Z}_{UP_\epsilon}^{N_f}(p, S) &= \sum_{m,n \geq 0} \frac{p^m}{m!} \frac{S^{2n}}{(2n)!} e^{S^2(\delta_{3,N_f/2})} \sum_{i=0}^n (-1)^i \frac{2^{m+4} \pi^{2i+2}}{3^n} \frac{(2n)!}{i!(n-i)!} \\ &\quad \times \sum_{\text{sing}} \text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}, \epsilon), \end{aligned}$$

with

$$(60) \quad \text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}, \epsilon) = \int_0^{K_s} dx_s q_s \frac{du_s}{dq_s} \frac{\eta^3(\tau_s) u_s^{m+n-i}}{\omega_s^{2i+2}} \mathcal{E}_{1/2}^i[Q_s] \Big|_{\tau_s = x_s - i \ln \epsilon}.$$

**Proof** We use the substitution rule and Lemma 4.10 to write each summand in the integrand as  $d\tau \wedge d\bar{\tau} \partial_{\bar{\tau}}(\dots)$ . Thus, the integral reduces to an integral over the boundary components of  $UP_\epsilon$ . To carry out the integration along the boundary, we switch to the variable  $\tau_s$  as described in Lemma 4.2. In the  $\tau_s$ –coordinate the boundary is a line parallel to the  $x_s$ –axis passed from the right to the left. We use  $2\pi i q_s \partial_{q_s} = \partial_{\tau_s}$  and the claim follows.  $\square$

**Lemma 4.15** *The limit  $\text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}) = \lim_{\epsilon \rightarrow 0} \text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}, \epsilon)$  is well-defined and equals*

$$(61) \quad \begin{aligned} &\text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}) \\ &= K_s \sum_{j=0}^i (-1)^j \binom{i}{j} \frac{\Gamma(1/2)}{\Gamma(1/2 + j)} 2^{2j} 3^j \\ &\quad \times \text{Coeff}_{q_s^0} \left[ q_s \frac{du_s}{dq_s} \frac{\eta^3(\tau_s) u_s^{m+n-i}}{\omega_s^{2i+2}} E_2^{i-j}(\tau_s) \left( q_s \frac{d}{dq_s} \right)^j Q_s^+(\tau_s) \right]. \end{aligned}$$

In particular, it follows that for all  $u_{\text{sing}} \neq \infty$  that

$$(62) \quad \text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}) = 0.$$

**Proof** We first consider the contribution to integral from a singular point  $u_{\text{sing}} = \infty$ . We write  $\mathcal{E}_{1/2}^k[Q] = q^{-1/8} (\sum_{\alpha \geq 0} H_{k,-\alpha}(y) q^{-\alpha} + \sum_{\alpha \geq 0} H_{k,\alpha} q^{\alpha/(4-N_f)})$  where  $H_{k,\alpha} \in \mathbb{Q}$ . For  $\alpha \in \mathbb{N}$ , the coefficients  $H_{k,-\alpha}(y)$  have the same exponential decay as the coefficient  $H_{-\alpha}(y)$  in Equation (50). Using Lemma 4.2 and (46), it follows,

for  $|q| < 1$ , that the integrand in (56) has a normally convergent power series expansion

$$(63) \quad q^{(m+n+2)/(4-N_f)+1} \frac{du}{dq} \frac{\eta^3(\tau)u^{m+n-i}}{\omega^{2i+2}} \mathcal{E}_{1/2}^i[\mathcal{Q}_\infty]$$

$$= \sum_{\beta \geq 0} C_\beta q^{\beta/(4-N_f)} \left( \sum_{\alpha \geq 0} H_{i,-\alpha}(y)q^{-\alpha} + \sum_{\alpha \geq 0} H_{i,\alpha}q^{\alpha/(4-N_f)} \right),$$

where the  $C_\beta \in \mathbb{Q}$ . For  $\alpha \in \mathbb{Z}$ , we have that  $\int_0^{4-N_f} dx q^{\alpha/(4-N_f)} = (4 - N_f)\delta_{\alpha,0}$ . Thus, the term-by-term integration in Equation (60) with respect to  $dx$  extracts the term proportional to  $q^0$ . The term contains the linear combination of products of the coefficients  $H_{k,-\alpha}(y)$  and  $C_\alpha$ , and it has exponential decay in  $y$ . It follows that  $\lim_{\epsilon \rightarrow 0} \text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}}, \epsilon)$  is well-defined, Equation (61) follows.

Secondly, we consider the contribution to the integral from a singular point  $u_{\text{sing}} \neq \infty$ . We write  $\mathcal{E}_{1/2}^k[\mathcal{Q}_s] = q_s^{-1/8} (\sum_{\alpha \geq 0} H_{s,k,-\alpha}(y_s)q_s^{-\alpha} + \sum_{\alpha \geq 0} H_{s,k,\alpha}q_s^{\alpha/K_s})$  where  $H_{s,k,\alpha} \in \mathbb{Q}$ . For  $\alpha \in \mathbb{N}$ , the coefficients  $H_{k,-\alpha}(y)$  have the same exponential decay as the coefficients  $H_{-\alpha}(y)$  in Equation (50). Using Lemma 4.2 and (46), it follows that for  $|q_s| < 1$  the integrand in (56) has a normally convergent power series expansion of the form

$$(64) \quad q_s \frac{du_s}{dq_s} \frac{\eta^3(\tau_s)u_s^{m+n-i}}{\omega_s^{2i+2}} \mathcal{E}_{1/2}^i[\mathcal{Q}_s]$$

$$= \sum_{\beta \geq 1} C_{s,\beta} q_s^{\beta/K_s} \left( \sum_{\alpha \geq 0} H_{s,i,-\alpha}(y_s)q_s^{-\alpha} + \sum_{\alpha \geq 0} H_{s,i,\alpha}q_s^{\alpha/K_s} \right),$$

with  $C_{s,\beta} \in \mathbb{Q}$ . For  $\alpha \in \mathbb{Z}$ , we have that  $\int_0^{K_s} dx_s q_s^{\alpha/K_s} = K_s \delta_{\alpha,0}$ . Thus, the term-by-term integration in Equation (60) with respect to  $dx_s$  receives only one contribution since the summation for  $\beta$  starts at 1. The term proportional to  $q_s^0$  contains the linear combinations of the coefficients  $H_{s,k,-\alpha}(y_s)$  and  $C_{s,\alpha}$ , but not  $H_{s,k,\alpha}$ . Thus, this contribution has an exponential decay in  $y_s$ . Hence  $\lim_{\epsilon \rightarrow 0} \text{Coeff}_{m,n,i}^0(u_{\text{sing}}, \epsilon) = 0$ .  $\square$

**Definition 4.16** The regularized generating function of the massless  $N_f$  low energy effective field theory on  $\mathbb{CP}^2$  is

$$\mathbf{Z}_{\text{UP}}^{N_f}(p, S) = \sum_{m,n \geq 0} \frac{p^m}{m!} \frac{S^{2n}}{(2n)!} e^{S^2(\delta_3, N_f/2)} \sum_{i=0}^n (-1)^i \frac{2^{m+4} \pi^{2i+2}}{3^n} \frac{(2n)!}{i!(n-i)!}$$

$$\times \text{Coeff}_{m,n,i}^{N_f}(u_{\text{sing}} = \infty).$$

**Theorem 4.17** We have that

$$(65) \quad \mathbf{Z}_{\text{UP}}^{N_f}(p, S) = \sum_{m,n \geq 0} \mathbf{D}_{m,2n}^{N_f} \frac{p^m}{m!} \frac{S^{2n}}{(2n)!} e^{(-p+S^2/3)\delta_3, N_f},$$

where the coefficients  $\mathbf{D}_{m,n}^{N_f}$  are as follows:

$$\begin{aligned}
 (66) \quad & \mathbf{D}_{m,2n}^0 \\
 &= \sum_{i=0}^n \sum_{j=0}^i \frac{(-1)^{i+j+1}}{2^{n-2j-1} 3^{n-j}} \frac{(2n)!}{(n-i)! j! (i-j)!} \Gamma(1/2) / \Gamma(j+1/2) \\
 & \quad \times \text{Coeff}_{q^0} \left[ \frac{\vartheta_4^9(\tau) [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]^{m+n-i}}{[\vartheta_2(\tau) \vartheta_3(\tau)]^{2m+2n+3}} E_2^{i-j}(\tau) \left( q \frac{d}{dq} \right)^j \mathcal{Q}_\infty^{(0)+}(\tau) \right], \\
 & \mathbf{D}_{m,2n}^2 \\
 &= \sum_{i=0}^n \sum_{j=0}^i \frac{(-1)^{i+j+1} 2^{-n+3j+2}}{3^{n-j}} \frac{(2n)!}{(n-i)! j! (i-j)!} \Gamma(1/2) / \Gamma(j+1/2) \\
 & \quad \times \text{Coeff}_{q^0} \left[ \frac{\vartheta_4^{10}(\tau) [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]^{m+n-i}}{\vartheta_2(\tau/2) [\vartheta_2(\tau) \vartheta_3(\tau)]^{2m+2n+3}} E_2^{i-j}\left(\frac{\tau}{2}\right) \left( q \frac{d}{dq} \right)^j \mathcal{Q}_\infty^{(2)+}\left(\frac{\tau}{2}\right) \right], \\
 & \mathbf{D}_{m,2n}^3 \\
 &= \sum_{i=0}^n \sum_{j=0}^i (-1)^{i+j} (-1)^{m+n-j} \frac{2^{3m+2n+2j+5}}{3^{n-j}} \frac{(2n)!}{j! (i-j)! (n-i)!} \frac{\Gamma(1/2t)}{\Gamma(j+1/2)} \\
 & \quad \times \text{Coeff}_{q^0} \left[ \frac{\vartheta_4^9(\tau) [\vartheta_3(\tau) \vartheta_4(\tau)]^{2m+2n-2i+3}}{[\vartheta_3^2(\tau) - \vartheta_4^2(\tau)]^{2m+2n+6}} E_2^{i-j}(\tau) \left( q \frac{d}{dq} \right)^j \mathcal{Q}_\infty^{(3)+}(\tau) \right].
 \end{aligned}$$

**Proof** In Lemma 4.14 and Lemma 4.15, we have already obtained an expansion of the coefficients. We use the explicit Weierstrass presentations from Lemma 4.1 to write each coefficient as a ratio of Jacobi  $\vartheta$ -functions. We use  $2\eta^3(\tau) = \vartheta_2(\tau) \vartheta_3(\tau) \vartheta_4(\tau)$  and  $q du/dq = -\vartheta_4^8(\tau) / [8 \vartheta_2^2(\tau) \vartheta_3^2(\tau)]$  for  $N_f = 0$ . For  $N_f = 2$ , we use  $\tau^{(2)} = \tau^{(0)}/2$ . For  $N_f = 3$ , we use  $q du/dq = \vartheta_2^8(\tau) \vartheta_3^2(\tau) \vartheta_4^2(\tau) / [\vartheta_3^2(\tau) - \vartheta_4^2(\tau)]^4$ .  $\square$

### 4.6.1 The invariants for $N_f = 0$

**Lemma 4.18** For all  $m, n \in \mathbb{N}$  with  $m + n \not\equiv 0 \pmod{2}$  it follows  $\mathbf{D}_{m,2n}^0 = 0$ .

**Proof** The claim follows from the fact that a factor

$$\vartheta_2^{2m+2n+3}(\tau) q^{1/8} = q^{(m+n+2)/4} (1 + O(\sqrt{q}))$$

is present in the denominator of the expression defining  $\mathbf{D}_{m,2n}^0$ .  $\square$

In Table 5 we list the first nonvanishing coefficients of the generating function in (65) for  $N_f = 0$ . They agree with the coefficients  $-\Phi_H^{\mathbb{P}^2, H}$  computed in [16, Theorem 4.4].

$p^m S^{2n}$	$\mathbf{D}_{m,2n}^0$	$\mathbf{D}_{m,2n}^0$
1	-1	$-\frac{1}{4}H_1 + 6H_0$
$S^4$	$-\frac{3}{16}$	$-\frac{49}{64}H_2 + \frac{9}{4}H_1 - \frac{2133}{64}H_0$
$p S^2$	$-\frac{5}{16}$	$-\frac{7}{64}H_2 + \frac{1}{4}H_1 - \frac{195}{64}H_0$
$p^2$	$-\frac{19}{16}$	$-\frac{1}{64}H_2 - \frac{1}{4}H_1 + \frac{411}{64}H_0$
$S^8$	$-\frac{232}{256}$	$-\frac{14641}{1024}H_3 + \frac{2401}{128}H_2 + \frac{44631}{1024}H_1 + \frac{108741}{128}H_0$
$p S^6$	$-\frac{152}{256}$	$-\frac{1331}{1024}H_3 - \frac{49}{128}H_2 + \frac{10341}{1024}H_1 - \frac{1749}{128}H_0$
$p^2 S^4$	$-\frac{136}{256}$	$-\frac{121}{1024}H_3 - \frac{91}{128}H_2 + \frac{2895}{1024}H_1 - \frac{3687}{128}H_0$
$p^3 S^2$	$-\frac{184}{256}$	$-\frac{11}{1024}H_3 - \frac{29}{128}H_2 + \frac{589}{1024}H_1 - \frac{753}{128}H_0$
$p^4$	$-\frac{680}{256}$	$-\frac{1}{1024}H_3 - \frac{7}{128}H_2 - \frac{505}{1024}H_1 + \frac{1725}{128}H_0$
$S^{12}$	$-\frac{69525}{4096}$	$-\frac{11390625}{16384}H_4 + \frac{44838675}{16384}H_2 + \frac{6075}{4}H_1 - \frac{76478175}{2048}H_0$
$p S^{10}$	$-\frac{26907}{4096}$	$-\frac{759375}{16384}H_4 - \frac{43923}{512}H_3 + \frac{4833213}{16384}H_2 + \frac{185733}{512}H_1 + \frac{5340591}{2048}H_0$
$p^2 S^8$	$-\frac{12853}{4096}$	$-\frac{50625}{16384}H_4 - \frac{9317}{512}H_3 + \frac{462707}{16384}H_2 + \frac{43587}{512}H_1 + \frac{1179489}{2048}H_0$
$p^3 S^6$	$-\frac{7803}{4096}$	$-\frac{3375}{16384}H_4 - \frac{363}{128}H_3 + \frac{861}{16384}H_2 + \frac{2829}{128}H_1 - \frac{69201}{2048}H_0$
$p^4 S^4$	$-\frac{6357}{4096}$	$-\frac{225}{16384}H_4 - \frac{99}{256}H_3 - \frac{21549}{16384}H_2 + \frac{1653}{256}H_1 - \frac{108639}{2048}H_0$
$p^5 S^2$	$-\frac{8155}{4096}$	$-\frac{15}{16384}H_4 - \frac{25}{512}H_3 - \frac{9475}{16384}H_2 + \frac{815}{512}H_1 - \frac{29265}{2048}H_0$
$p^6$	$-\frac{29557}{4096}$	$-\frac{1}{16384}H_4 - \frac{3}{512}H_3 - \frac{3021}{16384}H_2 - \frac{619}{512}H_1 + \frac{71649}{2048}H_0$

Table 5: First nonvanishing coefficients of the generating function in (65) for  $N_f = 0$

**4.6.2 The invariants for  $N_f = 2$**  In Table 6 we list the first coefficients of the generating function for  $N_f = 2$ .

**Lemma 4.19** For all  $m, n \in \mathbb{N}$  with  $m + n \not\equiv 0 \pmod{2}$  it follows  $\mathbf{D}_{m,2n}^2 = 0$ .

The lemma follows from the explicit formulas for these  $q$ -series. These series are given in terms of sums of products of derivatives of modular forms and harmonic Maass forms which are explicitly given in terms of theta functions and the  $\mu$ -function of Zwegers (see Section 6). We leave the proof of this lemma to the reader.

**4.6.3 The invariants for  $N_f = 3$**

**Lemma 4.20** For all  $m, n \in \mathbb{N}$  it follows that

$$(67) \quad \mathbf{D}_{m,2n}^3[Q_\infty^{(3)}(\tau)] = \mathbf{D}_{m,2n}^3[-Q(\tau)].$$



$p^m S^{2n}$	$\mathbf{D}_{m,2n}^2$	$\mathbf{D}_{m,2n}^2$
1	-3	$-\frac{1}{4}H_2 + \frac{27}{4}H_0$
$S^2$	0	$-\frac{11}{16}H_3 + \frac{77}{16}H_1$
$p$	0	$-\frac{1}{16}H_3 + \frac{7}{16}H_1$
$S^4$	$-\frac{21}{16}$	$-\frac{225}{64}H_4 + \frac{1043}{64}H_2 - \frac{567}{8}H_0$
$p S^2$	$-\frac{27}{16}$	$-\frac{15}{64}H_4 + \frac{61}{64}H_2 - \frac{9}{8}H_0$
$p^2$	$-\frac{53}{16}$	$-\frac{1}{64}H_4 - \frac{13}{64}H_2 + \frac{57}{8}H_0$
$S^6$	0	$-\frac{6859}{256}H_5 + \frac{22869}{256}H_3 + \frac{12555}{128}H_1$
$p S^4$	0	$-\frac{361}{256}H_5 + \frac{759}{256}H_3 + \frac{2217}{128}H_1$
$p^2 S^2$	0	$-\frac{19}{256}H_5 - \frac{115}{256}H_3 + \frac{659}{128}H_1$
$p^3$	0	$-\frac{1}{256}H_5 - \frac{33}{256}H_3 + \frac{129}{128}H_1$
$S^8$	$-\frac{3955}{256}$	$-\frac{279841}{1024}H_6 + \frac{664875}{1024}H_4 + \frac{366667}{256}H_2 + \frac{4203535}{1024}H_0$
$p S^6$	$-\frac{1925}{256}$	$-\frac{12167}{1024}H_6 + \frac{10125}{1024}H_4 + \frac{37709}{256}H_2 - \frac{195895}{1024}$
$p^2 S^4$	$-\frac{1219}{256}$	$-\frac{529}{1024}H_6 - \frac{2565}{1024}H_4 + \frac{5051}{256}H_2 - \frac{61409}{1024}H_0$
$p^3 S^2$	$-\frac{949}{256}$	$-\frac{23}{1024}H_6 - \frac{451}{256}H_4 + \frac{541}{256}H_2 - \frac{1735}{1024}H_0$
$p^4$	$-\frac{1811}{256}$	$-\frac{1}{1024}H_6 - \frac{53}{1024}H_4 - \frac{85}{256}H_2 + \frac{15151}{1024}H_0$

Table 6: First coefficients of the generating function for  $N_f = 2$

**Proof** We have the  $S$ -duality equation

$$(68) \quad \frac{1}{\sqrt{-i\tau}} \zeta_8 Q \left( 1 - \frac{1}{\tau} \right) = -\zeta_8 Q(\tau + 1).$$

In Lemma 4.13, we will show  $Q(\tau) = \zeta_8 Q(\tau + 1) + 14 \eta(\tau) \rho^4(\tau)$  where we have set  $\rho(\tau) = \sqrt{2} \eta^2(2\tau) / \eta^2(\tau)$ . It then follows that

$$(69) \quad \mathbf{D}_{m,2n}^3[-Q(\tau)] = \mathbf{D}_{m,2n}^3[-\zeta_8 Q(\tau + 1)] - 14 \mathbf{D}_{m,2n}^3[\eta(\tau) \rho^4(\tau)].$$

On the other hand, we have

$$\begin{aligned} \mathbf{D}_{m,2n}^3[-\zeta_8 Q(\tau + 1)] &= \mathbf{D}_{m,2n}^3 \left[ \frac{1}{\sqrt{-i\tau}} \left\{ Q \left( -\frac{1}{\tau} \right) - 14 \eta \left( -\frac{1}{\tau} \right) \rho^4 \left( -\frac{1}{\tau} \right) \right\} \right] \\ &= \mathbf{D}_{m,2n}^3 \left[ \frac{1}{\sqrt{-i\tau}} Q \left( -\frac{1}{\tau} \right) - 14 \eta(\tau) \frac{1}{\rho^4(\tau/2)} \right]. \end{aligned}$$

Hence, we obtain

$$\mathbf{D}_{m,2n}^3[-\mathcal{Q}(\tau)] = \mathbf{D}_{m,2n}^3[\mathcal{Q}_\infty^{(3)}(\tau)] - 14\mathbf{D}_{m,2n}^3\left[\eta(\tau)\left(\rho^4(\tau) + \frac{1}{\rho^4(\tau/2)}\right)\right].$$

One then checks that the last term on the right hand side vanishes. □

In Table 7 we list the first coefficients of the generating function for  $N_f = 3$ .

$p^m S^{2n}$	$\mathbf{D}_{m,2n}^3$	$\mathbf{D}_{m,2n}^3$
1	$-\frac{5}{4}$	$-\frac{1}{16}H_4 + \frac{3}{16}H_2 + \frac{3}{2}H_0$
$S^2$	$-\frac{95}{96}$	$-\frac{23}{128}H_6 + \frac{119}{384}H_4 + \frac{45}{32}H_2 + \frac{313}{128}H_0$
$p$	$\frac{45}{32}$	$-\frac{1}{128}H_6 + \frac{11}{128}H_4 - \frac{5}{32}H_2 - \frac{209}{128}H_0$
$S^4$	$-\frac{1787}{768}$	$-\frac{961}{1024}H_8 + \frac{851}{1024}H_6 + \frac{133}{24}H_4 + \frac{4587}{1024}H_2 - \frac{171}{128}H_0$
$p S^2$	$\frac{201}{256}$	$-\frac{31}{1024}H_8 + \frac{743}{3076}H_6 - \frac{5}{24}H_4 - \frac{4577}{3072}H_2 - \frac{991}{384}H_0$
$p^2$	$-\frac{489}{256}$	$-\frac{1}{1024}H_8 + \frac{19}{1024}H_6 - \frac{1}{8}H_4 + \frac{171}{1024}H_2 + \frac{277}{128}H_0$
$S^6$	$-\frac{189187}{18432}$	$-\frac{59319}{8192}H_{10} + \frac{12493}{8192}H_8 + \frac{70403}{2048}H_6 + \frac{3091945}{73728}H_4 + \frac{600451}{12288}H_2 - \frac{970759}{12288}H_0$
$p S^4$	$\frac{2211}{2048}$	$-\frac{1521}{8192}H_{10} + \frac{9579}{8192}H_8 - \frac{2065}{6144}H_6 - \frac{128731}{24576}H_4 - \frac{29563}{12288}H_2 + \frac{35039}{12288}H_0$
$p^2 S^2$	$-\frac{1627}{2048}$	$-\frac{39}{8192}H_{10} + \frac{1751}{24576}H_8 - \frac{2087}{6144}H_6 + \frac{4051}{24576}H_4 + \frac{7953}{4096}H_2 + \frac{40585}{12288}H_0$
$p^3$	$\frac{5843}{2048}$	$-\frac{1}{8192}H_{10} + \frac{27}{8192}H_8 - \frac{75}{2048}H_6 + \frac{1575}{8192}H_4 - \frac{825}{4096}H_2 - \frac{12987}{4096}H_0$

Table 7: First coefficients of the generating function for  $N_f = 3$

**Remark** In the case  $N_f = 3$ , using the variable  $u_{\text{SW}}^{(3)} = u^{(3)} + 1/2$  of [43] to define the  $u$ -plane integral leads to the generating function in Equation (65) with the term  $\exp(-p)$  removed.

### 4.7 The partition function for $N_f = 4$

Here we consider the case where  $N_f = 4$ . We set  $\alpha = \vartheta_2^4(\tau_0)$ ,  $\beta = \vartheta_3^4(\tau_0)$ .

**Lemma 4.21** *The Weierstrass presentation in the case # 57 is related to the Weierstrass presentation in the case # 71 in Lemma 4.1 by Table 8 and*

$$\begin{aligned} \Delta^{(4)} &= \frac{1}{256}\alpha^2\beta^2(\alpha - \beta)^2 \frac{1}{2^{12}}(u^{(4)})^2[u^{(4)} - 4\beta m^2]^2[u^{(4)} - 4\alpha m^2]^2 \\ &= \frac{\eta^{24}(\tau_0)}{2^{12}}u^2[u^{(4)} - 4\alpha m^2]^2[u^{(4)} - 4\beta m^2]^2. \end{aligned}$$

**Proof** The parametrization in the case #57 are obtained from #71 in Lemma 4.1 by transferring a star. This determines their Weierstrass presentations. □

$N_f = 4, \#57$					$UP_\epsilon^{(4)} = \mathbb{C}P^1 - B_\epsilon(0) - B_\epsilon(u_\pm) - \tilde{B}_\epsilon(\infty)$
$E_{\text{sing}}$	$I_2$	$I_2$	$I_2$	$I_0^*$	
$\tau_{\text{sing}}^{(4)}$	$\infty$	$0$	$1$	$\tau_0$	$u^{(4)} = \frac{8\alpha\beta m^2}{(\alpha - \beta)u^{(2)} + \alpha + \beta}$
$u_{\text{sing}}^{(2)}$	$\infty$	$-1$	$1$	$-\frac{\alpha + \beta}{\alpha - \beta}$	$u_\pm = 4\beta m^2, 4\alpha m^2$
$u_{\text{sing}}^{(4)}$	$0$	$u_-$	$u_+$	$\infty$	$\omega^{(4)}(\tau) = \frac{2\sqrt{2}\omega^{(2)}(\tau)}{\sqrt{\alpha - \beta}\sqrt{u^{(4)}}}$

Table 8

**Remark** The case #57 agrees with the case described in [43, Section 17.4] if we set  $m_1 = m_2 = m/2$  and  $m_3 = m_4 = 0$ .

**Remark** The  $\epsilon$ –disc  $\tilde{B}_\epsilon(\infty)$  around the point  $\infty$  with a singular fiber of the Kodaira-type  $I_0^*$  is constructed as follows: the boundary of the  $\epsilon$ –disc corresponds to the circle in the  $u$ –coordinate with  $u = Re^{i\theta}$ ,  $R = 1/\epsilon$ , and  $\theta \in [0; 2\pi]$ . It is easy to show that Lemmas 4.8, 4.14, 4.15 still hold for the elliptic families in Lemma 4.21 for all singular points with  $|u_{\text{sing}}| < \infty$ . Thus, the partition function receives contributions only from the singular point  $u = \infty$ .

**Definition 4.22** The partition function of the  $N_f = 4$  low energy effective field theory on  $\mathbb{C}P^2$  is

$$(70) \quad \tilde{Z}_{UP_\epsilon}^4(m, \tau_0) = -\frac{8}{\sqrt{2\pi}} f(m^2, \tau_0) \int_{UP_\epsilon} \frac{du \wedge d\bar{u} \, d\bar{\tau} \, \Delta^{1/8}}{\sqrt{\text{Im } \tau} \, d\bar{u} \, \omega^{1/2}} \overline{\eta^3(\tau)},$$

where  $f(m^2, \tau_0) = 1/(4^4 m^4 \eta^{12}(\tau_0))$  is a universal normalization factor. The regularized partition function of the massive  $N_f = 4$  low energy effective field theory on  $\mathbb{C}P^2$  is

$$Z_{UP}^4(m, \tau_0) = \lim_{\epsilon \rightarrow 0} \tilde{Z}_{UP_\epsilon}^4(m, \tau_0).$$

**Remark** The function  $f(m^2, \tau_0)$  is a universal normalization factor appearing in the  $u$ –plane integral which is familiar from the  $\mathcal{N} = 4$  case [29].

**Lemma 4.23** It follows that

$$(71) \quad \tilde{Z}_{UP_\epsilon}^4(m, \tau_0) = 16 f(m^2, \tau_0) \int_0^1 d\hat{\theta} \left( u \Delta^{1/6} \frac{Z(\tau)}{\eta^4(\tau)} \right) \Big|_{u=(1/\epsilon)e^{2\pi i \hat{\theta}}}.$$

**Proof** The integral reduces to an integral over the boundary component of  $UP_\epsilon$  at  $u = \infty$ . To carry out the integration along the boundary we set  $u = R e^{i\theta}$  (with  $\theta$  running clockwise from 0 to  $-2\pi$ ) and use

$$(72) \quad du \wedge d\bar{u} \partial_{\bar{u}} f = -d\left(f \frac{u}{R} dR + i f u d\theta\right). \quad \square$$

**Lemma 4.24** *The regularized partition function of the massive  $N_f = 4$  low energy effective field theory on  $\mathbb{C}P^2$  is*

$$(73) \quad \mathbf{Z}_{UP}^4(m, \tau_0) = \left[ \frac{1}{2} \frac{q}{\eta^4(\tau)} \frac{d}{dq} \left( \frac{q}{\eta^4(\tau)} \frac{d}{dq} \right) + g(\tau) \right] \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} \Big|_{\tau=\tau_0},$$

where

$$g(\tau) = -\frac{1}{2^2 3^2 \eta^8(\tau)} [\alpha^2 - \alpha\beta + \beta^2].$$

**Proof** In the case #57, we have

$$\frac{4u^{(4)}}{\eta^4(\tau_0)} \Delta^{1/6} = (u^{(4)})^2 - (\alpha + \beta) \frac{4m^2 u^{(4)}}{3} - (\alpha^2 - \alpha\beta + \beta^2) \frac{16m^4}{9} + m^4 O\left(\frac{m^2}{u^{(4)}}\right).$$

Using

$$u^{(2)} + \frac{\alpha + \beta}{\alpha - \beta} = \frac{8\alpha\beta m^2}{(\alpha - \beta)u^{(4)}},$$

we obtain the series expansion

$$\begin{aligned} \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} &= \frac{\mathbf{Z}(\tau_0)}{\eta^4(\tau_0)} + \left[ \eta^4(\tau) \frac{d\tau}{du^{(2)}} \right]_{\tau_0} \left[ \frac{1}{\eta^4(\tau)} \frac{d}{d\tau} \left( \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} \right) \right]_{\tau_0} \frac{8\alpha\beta m^2}{(\alpha - \beta)u^{(4)}} \\ &+ \frac{1}{2} \left[ \frac{d}{du^{(2)}} \left( \eta^4(\tau) \frac{d\tau}{du^{(2)}} \right) \right]_{\tau_0} \left[ \frac{1}{\eta^4(\tau)} \frac{d}{d\tau} \left( \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} \right) \right]_{\tau_0} \left( \frac{8\alpha\beta m^2}{(\alpha - \beta)u^{(4)}} \right)^2 \\ &+ \frac{1}{2} \left[ \eta^4(\tau) \frac{d\tau}{du^{(2)}} \right]_{\tau_0}^2 \left[ \frac{1}{\eta^4(\tau)} \frac{d}{d\tau} \left\{ \frac{1}{\eta^4(\tau)} \frac{d}{d\tau} \left( \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} \right) \right\} \right]_{\tau_0} \left( \frac{8\alpha\beta m^2}{(\alpha - \beta)u^{(4)}} \right)^2 \\ &+ O\left(\left[\frac{m^2}{u^{(4)}}\right]^3\right), \end{aligned}$$

where

$$2\pi i \eta^4(\tau) \frac{d\tau}{du^{(2)}} = -\frac{\pi^2}{4} \frac{\eta^4(\tau)}{\omega^{(2)}(\tau)^2 \Delta^{(2)}(\tau)^{1/2}} = -\frac{1}{4[\Delta^{(2)}]^{1/3}}.$$

A tedious calculation using  $2^4 \eta^{12}(\tau_0) = \alpha\beta(\alpha - \beta)$ , and

$$[\Delta^{(2)}]_{\tau_0} = \frac{1}{4} \frac{(\alpha\beta)^2}{(\alpha - \beta)^4}, \quad [\partial_{u^{(2)}} \Delta^{(2)}]_{\tau_0} = -\frac{1}{4} \frac{\alpha\beta(\alpha + \beta)}{(\alpha - \beta)^3},$$

then gives

$$\left[ \frac{u^{(4)}}{m^4 \eta^4(\tau_0)} \Delta^{1/6} \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} \right]_{[u^{(4)}]_0} = -\frac{4}{9}(\alpha^2 - \alpha\beta + \beta^2) \frac{\mathbf{Z}(\tau_0)}{\eta^4(\tau_0)} + 8\eta^8(\tau_0) \left[ \frac{q}{\eta^4(\tau)} \frac{d}{dq} \left\{ \frac{q}{\eta^4(\tau)} \frac{d}{dq} \left( \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)} \right) \right\} \right]_{\tau_0}.$$

The statement the follows from using the regularized integral

$$\lim_{\epsilon \rightarrow 0} \int_0^1 d\hat{\theta} u^k \Big|_{u=e^{2\pi i \hat{\theta}}/\epsilon} = \delta_{k,0}. \quad \square$$

The family of elliptic curves for the conformally invariant case corresponds to the massless  $N_f = 4$  low energy effective field theory on  $\mathbb{C}P^2$ .

**Theorem 4.25** *The regularized generating function of the massless  $N_f = 4$  low energy effective field theory on  $\mathbb{C}P^2$  is given by*

$$(74) \quad \mathbf{Z}_{\text{UP}}^4(0, \tau) = \left[ \frac{1}{2} \frac{q}{\eta^4(\tau)} \frac{d}{dq} \left( \frac{q}{\eta^4(\tau)} \frac{d}{dq} \right) + g(\tau) \right] \frac{\mathbf{Z}(\tau)}{\eta^4(\tau)},$$

where

$$g(\tau) = -\frac{1}{2^2 3^2} \left[ \left( \frac{\vartheta_2(\tau)}{\eta(\tau)} \right)^8 - \left( \frac{\vartheta_2(\tau)}{\eta(\tau)} \right)^4 \left( \frac{\vartheta_3(\tau)}{\eta(\tau)} \right)^4 + \left( \frac{\vartheta_3(\tau)}{\eta(\tau)} \right)^8 \right].$$

We have the following transformation properties under modular transformations:

$$\mathbf{Z}_{\text{UP}}^4(0, \tau + 2) = \mathbf{Z}_{\text{UP}}^4(0, \tau), \quad \mathbf{Z}_{\text{UP}}^4\left(0, -\frac{1}{\tau} + 1\right) = -\mathbf{Z}_{\text{UP}}^4(0, \tau + 1).$$

**Proof** The family of elliptic curves for the conformally invariant case is obtained from #57 in the limit  $m \rightarrow 0$ . We start from the generating function in the case #57 to obtain a duality group  $\Gamma(2)$ . In the limit  $m \rightarrow 0$ , the fibration becomes a rational elliptic surface with a constant  $j$ -invariant and two singularities of Kodaira-type  $I_0^*$ , one at  $u = 0$  and one at  $u = \infty$ . We have  $g_3 = \pi^6 E_6(\tau)/[216\omega^6]$ ,  $g_2 = \pi^4 E_4(\tau)/[12\omega^4]$ , and  $\omega = 2\pi/\sqrt{u}$  and  $\tau$  arbitrary. Higher terms proportional to  $p^k S^{2l}$  in the generating function  $\mathbf{Z}_{\text{UP}}^4(p, S)$  will be proportional to  $m^{2n}$  with  $n = k + l \geq 1$ . The reason is that the integrand contains higher powers of  $u$  or  $\hat{T}$ . The series expansion then introduces higher derivatives  $u^{(0)n} \partial_u^n$  which are proportional to  $m^{2n}$ . Thus, in the limit  $m \rightarrow 0$  only the constant term in the generating function  $\mathbf{Z}_{\text{UP}}^4(p, S)$  survives. The transformation under the shift follows from Lemma 4.13 and the fact that  $g(\tau)/\eta^4(\tau)$  is invariant under  $\tau \mapsto \tau + 2$ . The remaining transformation follows from Lemma 4.13 and the fact that  $g(1 + \tau) = g(1 - 1/\tau)$ .  $\square$

### 4.8 Criterion for Theorem 1.1

From a physics point of view, at a high energy scale the SO(3)–Donaldson theory is described by the low energy effective field theory. We show that the physics speculation by Moore and Witten on how to compute Donaldson invariants from the Seiberg–Witten curves can be turned into a mathematical statement which we then prove. The cuspidal contributions to the generating function of the low energy effective field theory should be equal to the generating function of the SO(3)–Donaldson theories with  $2N_f$  massless monopoles. We conjecture the following relationship between the Donaldson invariants defined in Equations (10) and (20) and the  $u$ –plane invariants given in Equation (65):

**Conjecture** Assume  $c = H$ . For  $N_f = 0, 2, 3$  and  $2m + 2n + 4 = (4 - N_f)k$ , we have

$$(75a) \quad \mathbf{D}_{m,2n}^0 = \Phi_{k,m,2n},$$

$$(75b) \quad \mathbf{D}_{m,2n}^{N_f} = 2^{(N_f+2)k/2} \Phi_{k,m,2n}^{N_f,c,1}.$$

For  $N_f = 0$ , this conjecture is the conjecture of Moore and Witten. Lemma 4.18 and Lemma 4.19 show that for  $N_f = 0, 2$  the invariants vanish on both sides for  $m + n \not\equiv 0 \pmod 2$ . For  $N_f = 0$ , the conjecture is equivalent to the assertion that the generating functions in (12) and in (65) are equal. We will prove Equation (75a) by proving:

**Theorem 4.26** Theorem 1.1 is equivalent to the vanishing of constant terms, for every pair of nonnegative integers  $m$  and  $n$ , of the series

$$(76) \quad \sum_{k=0}^n \sum_{j=0}^k (-1)^j \frac{(2n)!}{(n-k)!j!(k-j)!} \frac{\vartheta_4^8(\tau) [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]^m}{[\vartheta_2(\tau)\vartheta_3(\tau)]^{2m+2n+3}} E_2^{k-j}(\tau) \\ \times \left[ \frac{(-1)^{n+1}}{2^{k-3}3^k} \frac{(n-k)!}{(2n-2k)!} [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]^j F_{2(n-k)}(\tau) \right. \\ \left. + \frac{(-1)^{k+1}}{2^{n-2j-1}3^{n-j}} \frac{\Gamma(1/2)}{\Gamma(j+1/2)} \vartheta_4(\tau) [\vartheta_2^4(\tau) + \vartheta_3^4(\tau)]^{n-k} \left( q \frac{d}{dq} \right)^j Q^+(\tau) \right],$$

where series  $F_t(\tau)$  are defined in (93).

**Proof** In Theorem 4.17 we established explicit formula (66) for the contribution from the cusp at  $\tau = \infty$  in the regularized  $u$ –plane integral. Theorem 2.4 contained explicit formula (13) for the generating function of the Donaldson invariants of  $\mathbb{C}P^2$ . The difference of Equations (13) and (66) is the constant coefficient in Equation (76) where we have used that  $Q_\infty^{(0)+}(\tau) = Q^+(\tau)$ . Hence the vanishing of the constant coefficient term in Equation (76) is equivalent to Theorem 1.1. □

**Remark** For  $N_f = 2, 3$ , it is difficult to compute the geometrically defined invariants  $\Phi_{k,m,2n}^{N_f,c,1}$  directly. In contrast, the invariants  $\Phi_{k,m,2n}^{N_f,c,0}$  defined in Equation (19) are easy to compute by using Lemma 2.11. However, there is no known formula to compute the error term in Equation (22) to relate the latter to the former. However, we successfully performed the following check of the Equation (75b) for  $N_f = 2, 3$ .

The Donaldson invariants  $\Phi_{k,m,2n}^{N_f,c,1}$  and  $\Phi_{k,m,2n}^{N_f,c,0}$  are rational polynomials in the coefficients  $H_k$  of the holomorphic part of the Maass form  $Q(\tau)$ . By definition, the same statement is true for the invariants  $\mathbf{D}_{m,2n}^{N_f}$  as well. Geometrically, the invariants  $\Phi_{k,m,2n}^{N_f,c,1}$  and  $\Phi_{k,m,2n}^{N_f,c,0}$  differ by the contributions from the lower strata in the Uhlenbeck compactification. But the contribution from the highest stratum in the Uhlenbeck compactification agrees whence the coefficients of the highest  $H_k$  in  $\Phi_{k,m,2n}^{N_f,c,1}$  and  $\Phi_{k,m,2n}^{N_f,c,0}$  must agree. We proved the following weak version of a Moore–Witten type conjecture (75b) for the Donaldson invariants for  $N_f = 2, 3$ : In terms of the coefficients  $H_k$  of the holomorphic part of the weight  $1/2$  harmonic Maass form  $Q(\tau)$ , we have for  $c = H$ ,  $k$  even, and  $2m + 2n + 4 = (4 - N_f)k$  that

$$(77) \quad \text{Coeff}_{H_k}[\mathbf{D}_{m,2n}^{N_f} - 2^{(N_f+2)k/2}\Phi_{k,m,2n}^{N_f,c,0}] = 0.$$

The proof will appear somewhere else.

## 5 Harmonic Maass forms

We shall use Theorem 4.26 to prove Theorem 1.1. To this end, we make use of the theory of harmonic Maass forms, which we briefly recall here.

### 5.1 Definitions and facts

Following Bruinier and Funke [9], we define the notion of a harmonic weak Maass form of weight  $k \in \frac{1}{2}\mathbb{Z}$  as follows. We let  $\tau = u + iv \in \mathbb{H}$  with  $u, v \in \mathbb{R}$ , and throughout we let  $q := e^{2\pi i\tau}$ . We define the weight  $k$  hyperbolic Laplacian  $\Delta_k$  by

$$(78) \quad \Delta_k := -v^2 \left( \frac{\partial^2}{\partial u^2} + \frac{\partial^2}{\partial v^2} \right) + ikv \left( \frac{\partial}{\partial u} + i \frac{\partial}{\partial v} \right).$$

For odd integers  $d$ , define  $\epsilon_d$  by

$$(79) \quad \epsilon_d := \begin{cases} 1 & \text{if } d \equiv 1 \pmod{4}, \\ i & \text{if } d \equiv 3 \pmod{4}. \end{cases}$$

**Definition 5.1** If  $N$  is a positive integer (with  $4 \mid N$  if  $k \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$ ), then a *weight  $k$  harmonic weak Maass form on the congruence subgroup  $\Gamma_1(N)$*  is any smooth function  $M: \mathbb{H} \rightarrow \mathbb{C}$  satisfying the following:

- (1) For all  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1(N)$  and all  $\tau \in \mathbb{H}$ , we have

$$M\left(\frac{a\tau + b}{c\tau + d}\right) = \begin{cases} (c\tau + d)^k M(\tau) & \text{if } k \in \mathbb{Z}, \\ (c/d)^{2k} \epsilon_d^{-2k} (c\tau + d)^k M(\tau) & \text{if } k \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}. \end{cases}$$

Here  $\left(\frac{c}{d}\right)$  denotes the extended Legendre symbol, and  $\sqrt{\tau}$  is the principal branch of the holomorphic square root.

- (2) We have that  $\Delta_k M = 0$ .
- (3) There is a polynomial  $P_M = \sum_{n \leq 0} c^+(n)q^n \in \mathbb{C}[q^{-1}]$  such that

$$M(\tau) - P_M(\exp(2\pi i \tau)) = O(e^{-\epsilon v})$$

as  $v \rightarrow +\infty$  for some  $\epsilon > 0$ . Analogous conditions are required at all cusps.

**Remarks** (1) We refer to the polynomial  $P_M$  as the *principal part* of  $M(\tau)$  at  $\infty$ . Obviously, if  $P_M$  is nonconstant, then  $M(\tau)$  has exponential growth at  $\infty$ . Similar remarks apply at all cusps.

(2) Bruinier and Funke [9] give a slightly different definition for a harmonic weak Maass form. In place of Definition 5.1(3), they require that  $M(\tau)$  has at most linear exponential growth at cusps. Zagier’s weight  $3/2$  Maass–Eisenstein series is a harmonic weak Maass form in this relaxed sense.

(3) Since holomorphic functions on  $\mathbb{H}$  are harmonic, it follows that *weakly holomorphic modular forms*, those forms whose poles (if any) are supported at cusps, are harmonic weak Maass forms.

(4) For  $k \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$ , the transformation law in Definition 5.1(1) coincides with those in Shimura’s theory of half-integral weight modular forms [45].

In this paper we consider weight  $1/2$  harmonic weak Maass forms. Harmonic weak Maass forms have distinguished Fourier expansions which are described in terms of the incomplete Gamma-function  $\Gamma(\alpha; x)$

$$(80) \quad \Gamma(\alpha; x) := \int_x^\infty e^{-t} t^{\alpha-1} dt.$$



The following characterization is straightforward (for example, see [9, Section 3]). If  $f(\tau) \in H_{1/2}(N)$ , the space of weight  $1/2$  harmonic weak Maass forms on  $\Gamma_1(N)$ , for some  $N$ , then its Fourier expansion is of the form

$$(81) \quad f(\tau) = \sum_{n \gg -\infty} c_f^+(n)q^n + \sum_{n < 0} c_f^-(n)\Gamma(1/2, 4\pi|n|v)q^n.$$

One sees that  $f(\tau)$  naturally decomposes into two summands, its *holomorphic part*

$$(82) \quad f^+(\tau) := \sum_{n \gg -\infty} c_f^+(n)q^n,$$

and its *nonholomorphic part*

$$f^-(\tau) := \sum_{n < 0} c_f^-(n)\Gamma(1/2, 4\pi|n|v)q^n.$$

**Remark** A harmonic weak Maass form with trivial nonholomorphic part is a weakly holomorphic modular form.

Harmonic weak Maass forms are related to classical modular forms via properties of differential operators. One nontrivial relationship depends on the differential operator

$$(83) \quad \xi_w := 2iv^w \cdot \overline{\frac{\partial}{\partial \bar{\tau}}}.$$

This operator relates  $H_{1/2}(N)$  with  $S_{3/2}(N)$ , the space of weight  $3/2$  cusp forms on  $\Gamma_1(N)$ . The following lemma<sup>3</sup>, which is a straightforward refinement of a proposition of Bruinier and Funke [9, Proposition 3.2], plays a central role.

**Lemma 5.2** *If  $f \in H_{1/2}(N)$ , then*

$$\xi_{1/2}: H_{1/2}(N) \longrightarrow S_{3/2}(N).$$

Moreover, if

$$f(\tau) = \sum_{n \gg -\infty} c_f^+(n)q^n + \sum_{n < 0} c_f^-(n)\Gamma(1/2, 4\pi|n|v)q^n,$$

then we have that

$$\xi_{1/2}(f) = -(4\pi)^{1/2} \sum_{n=1}^{\infty} \overline{c_f^-(-n)}n^{1/2}q^n.$$

<sup>3</sup>The formula for  $\xi_{1/2}(f)$  corrects a typographical error in [9].

Thanks to [Lemma 5.2](#), we are in a position to relate the nonholomorphic parts of harmonic weak Maass forms, the expansions

$$f^-(\tau) := \sum_{n < 0} c_f^-(n) \Gamma(1/2, 4\pi|n|v) q^n,$$

with the “period integral” of the modular form  $\xi_{1/2}(f) \in S_{3/2}(N)$ . This observation was critical in Zwegers’s work on Ramanujan’s mock theta functions. To make this precise, we relate the Fourier expansion of the cusp form  $\xi_{1/2}(f)$  with  $f^-(\tau)$ . This connection is made by applying the simple integral identity

$$(84) \quad \int_{-\bar{\tau}}^{i\infty} \frac{e^{2\pi inz}}{(-i(\tau + z))^{1/2}} dz = i(2\pi n)^{-1/2} \cdot \Gamma(1/2, 4\pi nv) q^{-n}.$$

This identity follows by the direct calculation

$$\int_{-\bar{\tau}}^{i\infty} \frac{e^{2\pi inz}}{(-i(\tau + z))^{1/2}} dz = \int_{2iv}^{i\infty} \frac{e^{2\pi in(z-\tau)}}{(-iz)^{1/2}} dz = i(2\pi n)^{-1/2} \cdot \Gamma(1/2, 4\pi nv) q^{-n}.$$

By interchanging summation with integration, this identity then implies that the period integral

$$\int_{-\bar{\tau}}^{i\infty} \frac{\overline{\xi_{1/2}(f)(-\bar{z})}}{(-i(\tau + z))^{1/2}} dz$$

is proportional to  $\overline{f^-}$ , the nonholomorphic part of  $\overline{f}$ . In this way the nonholomorphic parts of weight 1/2 harmonic weak Maass forms are period integrals of weight 3/2 cusp forms. Zagier refers to  $\xi_{1/2}(f)$  as the *shadow* [55] of  $f^+$ . The holomorphic part of a weight 1/2 harmonic weak Maass form is called a *mock theta function* if  $\xi_{1/2}(f)$  is a linear combination of weight 3/2 theta series  $\vartheta(a, b; \delta\tau)$ , where

$$\vartheta(a, b; \tau) := \sum_{n \equiv a \pmod{b}} nq^{n^2}.$$

The mock theta functions which are relevant for [Theorem 1.1](#) are holomorphic parts of Maass forms whose shadows turn out to be proportional to

$$\eta(8\tau)^3 = \sum_{n=0}^{\infty} (-1)^n (2n + 1) q^{(2n+1)^2}.$$

## 6 Work of Zwegers

In his PhD thesis on mock theta functions [56], Zwegers constructs weight 1/2 harmonic weak Maass forms by making use of the transformation properties of functions which

were investigated earlier by Appell and Lerch. Here we briefly recall some of his results. For  $\tau \in \mathbb{H}$ , and  $u, v \in \mathbb{C} \setminus (\mathbb{Z}\tau + \mathbb{Z})$ , Zwegers defines the function

$$(85) \quad \mu(u, v; \tau) := \frac{a^{1/2}}{\theta(v; \tau)} \cdot \sum_{n \in \mathbb{Z}} \frac{(-b)^n q^{n(n+1)/2}}{1 - aq^n},$$

where  $a := e^{2\pi i u}$ ,  $b := e^{2\pi i v}$  and

$$(86) \quad \theta(v; \tau) := \sum_{v \in \mathbb{Z} + 1/2} (-1)^{v-1/2} b^v q^{v^2/2}.$$

Zwegers [56, Section 1.3] proves that  $\mu(u, v; \tau)$  satisfies the following important properties.

**Lemma 6.1** *Assuming the notation above, we have that*

$$\begin{aligned} \mu(u, v; \tau) &= \mu(v, u; \tau), \\ \mu(u + 1, v; \tau) &= -\mu(u, v; \tau), \\ a^{-1} b q^{-1/2} \mu(u + \tau, v; \tau) &= -\mu(u, v; \tau) + a^{-1/2} b^{1/2} q^{-1/8}, \\ \mu(u, v; \tau + 1) &= \zeta_8^{-1} \mu(u, v; \tau) \quad (\zeta_N := e^{2\pi i/N}), \\ (\tau/i)^{-1/2} e^{\pi i(u-v)^2/\tau} \mu\left(\frac{u}{\tau}, \frac{v}{\tau}; -\frac{1}{\tau}\right) &= -\mu(u, v; \tau) + \frac{1}{2} h(u - v; \tau), \end{aligned}$$

where

$$h(z; \tau) := \int_{-\infty}^{\infty} \frac{e^{\pi i x^2 \tau - 2\pi x z} dx}{\cosh \pi x}.$$

**Remark** The integral  $h(z; \tau)$  is known as a *Mordell integral*.

**Lemma 6.1** shows that  $\mu(u, v; \tau)$  is nearly a weight  $1/2$  Jacobi form, where  $\tau$  is the modular variable. Zwegers then uses  $\mu$  to construct weight  $1/2$  harmonic weak Maass forms. He achieves this by modifying  $\mu$  to obtain a function  $\hat{\mu}$  which he then uses as building blocks for such Maass forms. To make this precise, for  $\tau \in \mathbb{H}$  and  $z \in \mathbb{C}$ , let

$$(87) \quad R(z; \tau) := \sum_{v \in \mathbb{Z} + 1/2} \left( (-1)^{v-1/2} \{ \text{sgn}(v) - E((v + \text{Im}(z)/\text{Im}(\tau))\sqrt{2\text{Im}(\tau)}) \} e^{-2\pi i v z} q^{-v^2/2} \right),$$

where  $E(z)$  is the odd function

$$(88) \quad E(z) := 2 \int_0^z e^{-\pi u^2} du.$$

Using  $\mu$  and  $R$ , we let

$$(89) \quad \widehat{\mu}(u, v; \tau) := \mu(u, v; \tau) - \frac{1}{2}R(u - v; \tau).$$

Zwegers’s construction of weight  $1/2$  harmonic weak Maass forms depends on the following theorem (see Zwegers [56, Section 1.4] and Zagier [55]).

**Theorem 6.2** *Assuming the notation and hypotheses above, we have that*

$$\begin{aligned} \widehat{\mu}(u, v; \tau) &= \widehat{\mu}(v, u; \tau), \\ \widehat{\mu}(u + 1, v; \tau) &= a^{-1}bq^{-1/2}\widehat{\mu}(u + \tau, v; \tau) = -\widehat{\mu}(u, v; \tau), \\ \zeta_8\widehat{\mu}(u, v; \tau + 1) &= -(\tau/i)^{-1/2}e^{\pi i(u-v)^2/\tau}\widehat{\mu}\left(\frac{u}{\tau}, \frac{v}{\tau}; -\frac{1}{\tau}\right) = \widehat{\mu}(u, v; \tau). \end{aligned}$$

Theorem 6.2 gives the modular transformation properties for  $\widehat{\mu}$ . Since  $R$  is nonholomorphic, this theorem, combined with the following lemma, allows us to relate suitable specializations of  $\widehat{\mu}$  to weight  $1/2$  harmonic weak Maass forms whose shadows are linear combinations of weight  $3/2$  theta functions.

**Lemma 6.3** [56, Lemma 1.8] *The function  $R$  is real analytic and satisfies*

$$\frac{\partial R}{\partial \bar{u}}(u; \tau) = \sqrt{2}i v^{-1/2}e^{-2\pi a^2 v}\theta(\bar{u}; -\bar{\tau}),$$

where  $a := \text{Im}(u)/\text{Im}(\tau)$ . Moreover, we have that

$$\frac{\partial}{\partial \bar{\tau}}R(a\tau - b; \tau) = -\frac{i}{\sqrt{2}v}e^{-2\pi a^2 v} \sum_{v \in \mathbb{Z} + 1/2} (-1)^{v-1/2}(v + a)e^{-\pi i v^2 \bar{\tau} - 2\pi i v(a\bar{\tau} - b)}.$$

## 7 Some $q$ -series identities

Here we relate some important  $q$ -series to the  $\mu$ -function. For convenience, we first recall our notation for the following theta-functions which are simple Dedekind-eta

quotients:

$$(90) \quad \Theta_2(\tau) := \frac{\eta(16\tau)^2}{\eta(8\tau)} = \sum_{n=0}^{\infty} q^{(2n+1)^2} = q + q^9 + q^{25} + \dots,$$

$$(91) \quad \Theta_3(\tau) := \frac{\eta(8\tau)^5}{\eta(4\tau)^2\eta(16\tau)^2} = 1 + 2 \sum_{n=1}^{\infty} q^{4n^2} = 1 + 2q^4 + 2q^{16} + 2q^{36} + \dots,$$

$$(92) \quad \Theta_4(\tau) := \frac{\eta(4\tau)^2}{\eta(8\tau)} = 1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{4n^2} = 1 - 2q^4 + 2q^{16} - 2q^{36} + \dots.$$

These are related to the theta functions  $\vartheta_2(\tau)$ ,  $\vartheta_3(\tau)$  and  $\vartheta_4(\tau)$  by

$$\vartheta_2(\tau) = 2\Theta_2\left(\frac{\tau}{8}\right), \quad \vartheta_3(\tau) = \Theta_3\left(\frac{\tau}{8}\right), \quad \vartheta_4(\tau) = \Theta_4\left(\frac{\tau}{8}\right).$$

Thanks to [Theorem 4.26](#), the proof of [Theorem 1.1](#) is reduced to the properties of certain specific power series. For nonnegative even  $t$ , we define the series

$$(93) \quad F_t(q) = q^{-1/8} \sum_{\beta=0}^{\infty} \sum_{\alpha=\beta+1}^{\infty} (-1)^{\alpha+\beta} (2\beta+1)^t q^{(\alpha^2-\beta(\beta+1))/2}.$$

We shall work with the renormalizations

$$(94) \quad \mathcal{F}_t(q) := F_t(q^8) = \sum_{\beta=0}^{\infty} \sum_{\alpha=\beta+1}^{\infty} (-1)^{\alpha+\beta} (2\beta+1)^t q^{4\alpha^2-(2\beta+1)^2}.$$

We shall also require the renormalization of the series  $\mathcal{Q}^+(\tau)$  in [\(49\)](#):

$$(95) \quad \mathcal{Q}(q) := \mathcal{Q}^+(8\tau) = q^{-1} + 28q^3 + 39q^7 + 196q^{11} + 161q^{15} + 756q^{19} + \dots.$$

### 7.1 The $\mathcal{F}_t(q)$ series

Here we interpret the series  $\mathcal{F}_t(q)$  in terms of Zwegers’s  $\mu$ –function. To make this precise, we require the differential operator

$$D_\omega := \frac{1}{2\pi i} \cdot \frac{\partial}{\partial \omega}.$$

The main result in this section is the following theorem which expresses the  $q$ –series  $\mathcal{F}_t(q)$  in terms of the image of the iterated  $t$ –th derivative, with respect to  $\omega$  then evaluated at  $\omega = 0$ , of a certain specialization of the  $\mu$ –function.

**Theorem 7.1** *If  $t$  is a nonnegative even integer, then*

$$\frac{\mathcal{F}_t(q)}{\Theta_4(\tau)} = \frac{1}{2} D_\omega^t (\mu(4\tau + 2\omega, 4\tau; 8\tau))|_{\omega=0} = \frac{1}{2} D_\omega^t (\mu(2\omega, 4\tau; 8\tau))|_{\omega=2\tau}.$$

Moreover,  $\mathcal{F}_0(q)/\Theta_4(\tau)$  is the holomorphic part of the weight  $1/2$  harmonic weak Maass form  $\frac{1}{2}\hat{\mu}(4\tau, 4\tau; 8\tau)$  whose nonholomorphic part is the period integral of  $\eta(8\tau)^3$ .

**Remark** That  $t$  is even will be important in the proof of [Theorem 1.1](#) due to the close connection between  $D_\omega^t$  and the iterated heat operator.

**Proof** It is not difficult to show that

$$\begin{aligned} \mathcal{F}_t(q) &= - \sum_{x,y \geq 0} (2x+1)^t q^{16y^2+16y+16xy+4x+3} \\ &\quad + \sum_{x,y \geq 0} (2x+1)^t q^{16y^2+32y+16xy+12x+15} \\ &= - \sum_{y=0}^{\infty} (-1)^y q^{(2y+1)^2} \sum_{x=0}^{\infty} (2x+1)^t q^{2(2x+1)(2y+1)} \\ &= - \sum_{y=0}^{\infty} (-1)^y q^{(2y+1)^2} \sum_{x=0}^{\infty} \left( \frac{x^t - (-1)^x x^t}{2} \right) q^{2x(2y+1)} \\ &= -\frac{1}{2} \sum_{y=0}^{\infty} (-1)^y q^{(2y+1)^2} \sum_{x=0}^{\infty} x^t q^{2x(2y+1)} \\ &\quad + \frac{1}{2} \sum_{y=0}^{\infty} (-1)^y q^{(2y+1)^2} \sum_{x=0}^{\infty} (-1)^x x^t q^{2x(2y+1)}. \end{aligned}$$

Using this last expression, and by letting  $\rho := e^{2\pi i \omega}$ , it follows that

$$\begin{aligned} \mathcal{F}_t(q) &= -\frac{1}{2} \sum_{y=0}^{\infty} (-1)^y q^{(2y+1)^2} \cdot D_\omega^t \left( \sum_{x=0}^{\infty} \rho^x q^{x(4y+2)} \right) \Big|_{\omega=0} \\ &\quad + \frac{1}{2} \sum_{y=0}^{\infty} (-1)^y q^{(2y+1)^2} \cdot D_\omega^t \left( \sum_{x=0}^{\infty} (-\rho)^x q^{x(4y+2)} \right) \Big|_{\omega=0} \\ &= -\frac{1}{2} D_\omega^t \left( \sum_{y=0}^{\infty} \left( \frac{(-1)^y q^{(2y+1)^2}}{1 - \rho q^{4y+2}} + \frac{(-1)^{y+1} q^{(2y+1)^2}}{1 + \rho q^{4y+2}} \right) \right) \Big|_{\omega=0} \\ &= -D_\omega^t \left( \sum_{y=0}^{\infty} \frac{(-1)^y \rho q^{4y^2+8y+3}}{1 - \rho^2 q^{8y+4}} \right) \Big|_{\omega=0}. \end{aligned}$$

Since we have that

$$\begin{aligned} \sum_{y=0}^{\infty} \frac{(-1)^y \rho q^{4y^2+8y+3}}{1-\rho^2 q^{8y+4}} &= \frac{1}{2} \sum_{y=0}^{\infty} \frac{(-1)^y \rho q^{4y^2+8y+3}}{1-\rho^2 q^{8y+4}} + \frac{1}{2} \sum_{y=0}^{-\infty} \frac{(-1)^y \rho q^{4y^2-8y+3}}{1-\rho^2 q^{-8y+4}} \\ &= \frac{1}{2} \sum_{y=0}^{\infty} \frac{(-1)^y \rho q^{4y^2+8y+3}}{1-\rho^2 q^{8y+4}} - \frac{1}{2} \sum_{y=-1}^{-\infty} \frac{(-1)^y \rho q^{4y^2-1}}{1-\rho^2 q^{-8y-4}} \\ &= \frac{1}{2} \sum_{y=0}^{\infty} \frac{(-1)^y \rho q^{4y^2+8y+3}}{1-\rho^2 q^{8y+4}} - \frac{1}{2} \sum_{y=-1}^{-\infty} \frac{(-1)^y \rho^{-1} q^{4y^2+8y+3}}{\rho^{-2} q^{8y+4}-1} \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \frac{(-1)^n \rho q^{4n^2+8n+3}}{1-\rho^2 q^{8n+4}} + \frac{1}{2} \sum_{n=-1}^{-\infty} \frac{(-1)^n \rho^{-1} q^{4n^2+8n+3}}{1-\rho^{-2} q^{8n+4}}, \end{aligned}$$

the fact that  $t$  is even then implies that

$$(96) \quad \mathcal{F}_t(q) = -\frac{1}{2} D_{\omega}^t \left( \sum_{n \in \mathbb{Z}} \frac{(-1)^n \rho q^{4n^2+8n+3}}{1-\rho^2 q^{8n+4}} \right) \Big|_{\omega=0}.$$

Since we have that

$$\frac{1}{2} D_{\omega}^t (\mu(4\tau + 2\omega, 4\tau; 8\tau)) \Big|_{\omega=0} = \frac{1}{2} D_{\omega}^t (\mu(2\omega, 4\tau; 8\tau)) \Big|_{\omega=2\tau},$$

to complete the proof of the claimed identity, it suffices to compute the expansion of  $\mu(4\tau + 2\omega, 4\tau; 8\tau)$ . By (85), we have that

$$\mu(4\tau + 2\omega, 4\tau; 8\tau) = \frac{q^{-1}}{\theta(4\tau; 8\tau)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n \rho q^{4n^2+8n+3}}{1-\rho^2 q^{8n+4}}.$$

The claimed identity follows immediately from (86) and (96) since

$$q\theta(4\tau; 8\tau) = \sum_{m \in \mathbb{Z}} (-1)^m q^{4(m+1)^2} = - \sum_{m \in \mathbb{Z}} (-1)^m q^{4m^2} = -\frac{\eta(4\tau)^2}{\eta(8\tau)} = -\Theta_4(\tau).$$

That  $\mathcal{F}_0(q)/\Theta_4(\tau)$  is the holomorphic part of the weight  $1/2$  harmonic weak Maass form  $\frac{1}{2} \hat{\mu}(4\tau, 4\tau; 8\tau)$  follows, after a straightforward calculation, from Theorem 6.2. That the nonholomorphic part of this Maass form is a multiple of the period integral of  $\eta(8\tau)^3$  follows from the classical identity

$$\eta(8\tau)^3 = \sum_{n=0}^{\infty} (-1)^n (2n+1) q^{(2n+1)^2},$$

and the explicit Fourier expansion of the nonholomorphic part of  $\frac{1}{2} \hat{\mu}(4\tau, 4\tau; 8\tau)$ . This expansion is obtained using (87), Lemma 6.3, and the discussion after Lemma 5.2.  $\square$

### 7.2 The $\mathcal{Q}(q)$ series

Here we describe the  $q$ -series  $\mathcal{Q}(q)$  in terms of Zwegers’s  $\mu$ -function. To make this precise, we require some auxiliary weight  $1/2$  weakly holomorphic modular forms. Define modular forms  $A(\tau)$  and  $B(\tau)$  by

$$\begin{aligned}
 (97) \quad \mathcal{A}(\tau) &:= A(8\tau) = \sum_{n=-1}^{\infty} a(n)q^n := \frac{\eta(4\tau)^8}{\eta(8\tau)^7} = q^{-1} - 8q^3 + 27q^7 - \dots, \\
 \mathcal{B}(\tau) &:= B(8\tau) = \sum_{n=-1}^{\infty} b(n)q^n := \frac{\eta(8\tau)^5}{\eta(16\tau)^4} = q^{-1} - 5q^7 + 9q^{15} - \dots.
 \end{aligned}$$

We sieve on the Fourier expansion of  $\mathcal{A}(\tau)$  to define the modular forms

$$\begin{aligned}
 (98) \quad \mathcal{A}_{3,8}(\tau) &:= A_{3,8}(8\tau) = \sum_{n \equiv 3 \pmod{8}} a(n)q^n = -8q^3 - 56q^{11} + \dots, \\
 \mathcal{A}_{7,8}(\tau) &:= A_{7,8}(8\tau) = \sum_{n \equiv 7 \pmod{8}} a(n)q^n = q^{-1} + 27q^7 + 105q^{15} + \dots.
 \end{aligned}$$

In terms of these modular forms, we have the following theorem.

**Theorem 7.2** *The following  $q$ -series identity is true:*

$$\begin{aligned}
 \mathcal{Q}(q) &= -\frac{7}{2}\mathcal{A}_{3,8}(\tau) + \frac{3}{2}\mathcal{A}_{7,8}(\tau) - \frac{1}{2}\mathcal{B}(\tau) \\
 &\quad + 2iq^{-1}\mu\left(-16\tau, -8\tau - \frac{1}{2}; 32\tau\right) - 2iq^{-1}\mu\left(-16\tau, -24\tau - \frac{1}{2}; 32\tau\right).
 \end{aligned}$$

Moreover,  $\mathcal{Q}(q)$  is the holomorphic part of a weight  $1/2$  harmonic weak Maass form whose nonholomorphic part is the period integral of  $\eta(8\tau)^3$ .

**Remark** One can determine the image of  $\mathcal{Q}(q)$  under the inversion map  $\tau \rightarrow -1/\tau$  using the formula for  $\mathcal{Q}(q)$  above. One obtains this expansion by applying the third part of [Theorem 6.2](#) to two  $\mu$ -functions, and by applying the classical transformation

$$\eta(-1/\tau) = \sqrt{-i\tau} \cdot \eta(\tau)$$



to the weakly holomorphic modular forms

$$\begin{aligned} \mathcal{A}_{3,8}(\tau) &= -8 \cdot \frac{\eta(16\tau)^8}{\eta(8\tau)^7} = -8q^3 - 56q^{11} - 216q^{19} - \dots, \\ \mathcal{A}_{7,8}(\tau) &= \frac{\eta(8\tau)^5}{\eta(16\tau)^4} + 32 \cdot \frac{\eta(32\tau)^8}{\eta(8\tau)^3 \eta(16\tau)^4} = q^{-1} + 27q^7 + 105q^{15} + \dots, \\ \mathcal{B}(\tau) &= \frac{\eta(8\tau)^5}{\eta(16\tau)^4} = q^{-1} - 5q^7 + 9q^{15} - \dots. \end{aligned}$$

**Sketch of the proof of Theorem 7.2** This result follows from Theorem 6.2 in an argument which is analogous to the proof of Theorem 7.1. Namely, since  $\mathcal{A}_{3,8}(\tau)$ ,  $\mathcal{A}_{7,8}(\tau)$  and  $\mathcal{B}(\tau)$  are weight  $1/2$  modular forms, it follows that  $\mathcal{Q}(\tau)$  is the holomorphic part of a weight  $1/2$  harmonic weak Maass form whose nonholomorphic part is a multiple of the period integral of  $\eta(8\tau)^3$ . Therefore, it must be a linear combination of certain mock theta functions and weakly holomorphic modular forms of weight  $1/2$ . Instead of using Theorem 7.1 with  $t = 0$ , we work with the mock theta function

$$\begin{aligned} (99) \quad \mathcal{M}(q) &:= q^{-1} \sum_{n=0}^{\infty} \frac{(-1)^{n+1} q^{8(n+1)^2} \prod_{k=1}^n (1 - q^{16k-8})}{\prod_{k=1}^{n+1} (1 + q^{16k-8})^2} \\ &= -q^7 + 2q^{15} - 3q^{23} + \dots, \end{aligned}$$

and  $\mathcal{M}(q) = M(q^8)$ . Using Watson’s  $q$ -analog of Whipple’s theorem, we have that

$$\mathcal{M}(q) = -\frac{1}{2\Theta_2(\tau)} \cdot \sum_{n \in \mathbb{Z}} \frac{q^{16n^2-8n}}{1 + q^{16n-8}}.$$

Andrews [2] already interpreted<sup>4</sup> this particular  $q$ -series as the holomorphic part of a weight  $1/2$  harmonic weak Maass form whose nonholomorphic part is the period integral of  $\eta(8\tau)^3$ . Using the fact that

$$\sum_{n \in \mathbb{Z}} \frac{q^{16n^2-8n}}{1 + q^{16n-8}} = \sum_{n \in \mathbb{Z}} \frac{q^{16n^2-8n}(1 - q^{16n-8})}{1 - q^{32n-16}} = \sum_{n \in \mathbb{Z}} \frac{q^{16n^2-8n}}{1 - q^{32n-16}} - \sum_{n \in \mathbb{Z}} \frac{q^{16n^2+8n-8}}{1 - q^{32n-16}},$$

we find that

$$\mathcal{M}(q) = -\frac{iq^{-1}}{2} \cdot \left( \mu \left( -16\tau, -24\tau - \frac{1}{2}; 32\tau \right) - \mu \left( -16\tau, -8\tau - \frac{1}{2}; 32\tau \right) \right).$$

<sup>4</sup> We have reformulated Andrews’s work to be consistent with the terminology in this paper.

Here we have used the fact that

$$q\theta \left( -8\tau - \frac{1}{2}; 32\tau \right) = q^9\theta \left( -24\tau - \frac{1}{2}; 32\tau \right) = i\Theta_2(\tau).$$

Since two weight  $1/2$  harmonic weak Maass forms with the same nonholomorphic parts differ by a weakly holomorphic modular form, it then follows that  $\mathcal{Q}(q) - 4\mathcal{M}(q)$  is a weakly holomorphic modular form. Standard calculations in the theory of modular forms reveals that

$$\mathcal{Q}(q) - 4\mathcal{M}(q) = -\frac{7}{2}\mathcal{A}_{3,8}(\tau) + \frac{3}{2}\mathcal{A}_{7,8}(\tau) - \frac{1}{2}\mathcal{B}(\tau). \quad \square$$

**Remark** One also sees directly that the shadow of  $\mathcal{Q}(q)$  is proportional to  $\eta(8\tau)^3$  using the proven identity. One applies [Lemma 6.3](#) to this particular linear combination  $\hat{\mu}$  functions.

## 8 Proof of [Theorem 1.1](#) and some numerical examples

Here we prove [Theorem 1.1](#) by verifying the criterion in [Theorem 4.26](#). We also conclude this section with some numerical examples which illustrate the phenomenon which appears in the proof of the theorem.

### 8.1 Proof of [Theorem 1.1](#)

Thanks to [Theorem 4.26](#), it suffices to prove that the differences between certain  $q$ -series have vanishing constant term. We shall derive these conclusions by using differential operators. For brevity we describe the  $n = 0$  cases in detail, and then provide general remarks which are required to justify the remaining cases. By [Theorems 7.1](#) and [7.2](#), it follows that both

$$\begin{aligned} \mathcal{Q}(q) &= q^{-1} + 28q^3 + 39q^7 + 196q^{11} + 161q^{15} + 756q^{19} + \dots, \\ -\frac{4\mathcal{F}_0(q)}{\Theta_4(\tau)} &= 4q^3 + 12q^7 + 28q^{11} + \dots \end{aligned}$$

are the holomorphic parts of weight  $1/2$  harmonic weak Maass forms with equal nonholomorphic parts. Therefore, it follows that

$$\mathcal{Q}(q) + \frac{4\mathcal{F}_0(q)}{\Theta_4(\tau)} = q^{-1} + 24q^3 + 27q^7 + 168q^{11} + \dots$$

is a weakly holomorphic modular form. Standard calculations using the theory of modular forms reveals that

$$(100) \quad Z_0(q) := \mathcal{Q}(q) + \frac{4\mathcal{F}_0(q)}{\Theta_4(\tau)} = \frac{E^*(4\tau)}{\eta(8\tau)^3},$$

where  $E^*(4\tau)$  is the weight 2 Eisenstein series

$$(101) \quad E^*(\tau) := -E_2(\tau) + 2E_2(2\tau) = 1 + 24 \sum_{n=1}^{\infty} \sigma_{\text{odd}}(n)q^n,$$

where  $\sigma_{\text{odd}}(n)$  denotes the sum of the positive odd divisors of  $n$ . The  $n = 0$  cases of [Theorem 4.26](#) are equivalent to the claim that the constant terms of

$$(102) \quad \frac{\Theta_4(\tau)^9(16\Theta_2(\tau)^4 + \Theta_3(\tau)^4)^m Z_0(q)}{\Theta_2(\tau)^{2m+3}\Theta_3(\tau)^{2m+3}}$$

are 0 for every  $m \geq 0$ . In order to verify this claim, we will find it helpful to define

$$(103) \quad \widehat{Z}_0(q) := \frac{E^*(4\tau)}{\Theta_2(\tau)^2\Theta_3(\tau)^2} = \frac{Z_0(q)\eta(8\tau)^3}{\Theta_2(\tau)^2\Theta_3(\tau)^2}.$$

A calculation shows that

$$(104) \quad q \frac{d}{dq} \widehat{Z}_0(q) = \frac{\Theta_4(\tau)^9}{\Theta_2(\tau)\Theta_3(\tau)\eta(8\tau)^3}.$$

Using this notation, and noting that

$$16\Theta_2(\tau)^4 + \Theta_3(\tau)^4 = 1 + 24q^4 + 24q^2 + \dots = E^*(4\tau),$$

[Equation \(102\)](#) becomes

$$\widehat{Z}_0(q)^{m+1} \cdot q \frac{d}{dq} \widehat{Z}_0(q),$$

which clearly has a vanishing constant term. In fact, for each  $m, n \geq 0$  we find a similar phenomenon.

To this end, we make use of the combinatorial structure of series  $\mathcal{G}_k(q)$  and  $H_k(q)$  defined below. These series are obtained by iteratively applying differential operators to the harmonic Maass forms and modular forms in this paper, which in turn lead to a nice sequence of modular functions of level 8. The modular properties of these series follow easily from the results and methods of [[11](#), pages 53–55; [13](#)].

For every nonnegative  $k$ , define

$$(105) \quad \mathcal{G}_k(q) := \frac{\eta(8\tau)^3}{(\Theta_2(\tau)\Theta_3(\tau))^{2k+2}} \times \sum_{j=0}^k \binom{k}{j} \frac{\Gamma(1/2)(-12)^j E_2(8\tau)^{k-j}}{\Gamma(1/2+j)8^j} \left[ \frac{(-1)^j 4\mathcal{F}_2(j)(q)}{\Theta_4(\tau)} + \left(q \frac{d}{dq}\right)^j \mathcal{Q}(q) \right].$$

Theorem 4.26 is equivalent to the claim that the constant coefficient of

$$(106) \quad \left(q \frac{d}{dq} \widehat{Z}_0(q)\right) \widehat{Z}_0(q)^m \sum_{k=0}^n \binom{n}{k} \widehat{Z}_0(q)^{n-k} \mathcal{G}_k(\tau)$$

is zero for each nonnegative  $m$  and  $n$ . To prove the theorem, it suffices to show that  $\mathcal{G}_k(q)$  is a polynomial in  $\widehat{Z}_0(q)$ . To this end, we define  $M_0^*(\Gamma_0(8))$  to be the space of modular function on  $\Gamma_0(8)$  which are holomorphic away from infinity, and is a subspace of  $\mathbb{C}((q^2))$ . One can easily verify that  $M_0^*(\Gamma_0(8))$  is precisely the set of polynomials in  $\widehat{Z}_0(q)$ . In order to show that  $\mathcal{G}_k(\tau)$  is in  $M_0^*(\Gamma_0(8))$ , we first show that a similar function,  $\mathcal{H}_k(q)$  is in  $M_0^*(\Gamma_0(8))$ . We define the function

$$\mathcal{H}_k(q) := \frac{\eta(8\tau)^3}{(\Theta_2(\tau)\Theta_3(\tau))^{2k+2}} \sum_{j=0}^k \binom{k}{j} \frac{\Gamma(1/2)(-12)^j E_2(8\tau)^{k-j}}{\Gamma(1/2+j)8^j} \left(q \frac{d}{dq}\right)^j Z_0(q).$$

We can observe that  $\mathcal{H}_k(q)$  is modular on  $\Gamma_0(8)$  with weight 0 by comparing the summation to the expression  $\mathcal{Z}_k(q) := \mathcal{E}_{1/2}^k[Z_0(q^{1/8})]$  where  $\mathcal{E}_{1/2}^k[*]$  is defined as in Lemma 4.10. A calculation shows that  $(\Theta_2(\tau)\Theta_3(\tau))^{-2}$  is holomorphic away from infinity, which, combined with the fact that

$$\frac{\eta(8\tau)^3 Z_0(q)}{(\Theta_2(\tau)\Theta_3(\tau))^2} = \widehat{Z}_0(q),$$

shows that  $\mathcal{H}_k(\tau)$  is in  $M_0^*(\Gamma_0(8))$ . Hence we need only show that  $\mathcal{G}_k(q) - \mathcal{H}_k(q)$  is in  $M_0^*(\Gamma_0(8))$  as well. We observe that

$$\begin{aligned} &\mathcal{G}_k(q) - \mathcal{H}_k(q) \\ &= \frac{\eta(8\tau)^3}{(\Theta_2(\tau)\Theta_3(\tau))^{2k+2}} \\ &\quad \times \sum_{j=0}^k \binom{k}{j} \frac{\Gamma(1/2)(-12)^j E_2(8\tau)^{k-j}}{\Gamma(1/2+j)8^j} \left[ \frac{(-1)^j 4\mathcal{F}_2(j)(q)}{\Theta_4(\tau)} - \left(q \frac{d}{dq}\right)^j \frac{4\mathcal{F}_0(q)}{\Theta_4(\tau)} \right]. \end{aligned}$$

Using [Theorem 7.1](#), this can be written as

$$\frac{\eta(8\tau)^3}{(\Theta_2(\tau)\Theta_3(\tau))^{2k+2}} \sum_{j=0}^k \binom{k}{j} \frac{\Gamma(1/2)(-12)^j E_2(8\tau)^{k-j}}{\Gamma(1/2 + j)8^j} \cdot 2 \left[ (-1)^j \left( \frac{1}{2\pi i} \frac{d}{d\omega} \right)^{2j} \mu(4\tau+2\omega, 4\tau; 8\tau) - \left( \frac{1}{2\pi i} \frac{d}{d\tau} \right)^j \mu(4\tau+2\omega, 4\tau; 8\tau) \right] \Big|_{\omega=0}.$$

Using the transformation laws for  $\mu$  found in [Lemma 6.1](#), we observe that the Mordell integrals that arise as obstructions to the modular transformation of

$$(-1)^j \left( \frac{1}{2\pi i} \frac{d}{d\omega} \right)^{2j} \mu(4\tau + 2\omega, 4\tau; 8\tau) - \left( \frac{1}{2\pi i} \frac{d}{d\tau} \right)^j \mu(4\tau + 2\omega, 4\tau; 8\tau)$$

cancel directly. Therefore an argument similar to that used in the proof of [Lemma 4.10](#) shows that  $\mathcal{G}_k(q) - \mathcal{H}_k(q)$  is modular with respect to  $\Gamma_0(8)$ . It is then straightforward to verify that  $\mathcal{G}_k(q) - \mathcal{H}_k(q)$  is in  $M_0^*(\Gamma_0(8))$ , and so is a polynomial in  $\widehat{Z}_0(q)$ . This then completes the proof.

## 8.2 Examples

Here we give some numerical examples of [Theorem 4.26](#). First we give some examples when  $n = 0$ , and then we give an example when  $n = 1$ . We conclude with examples of the polynomials in  $\widehat{Z}_0(q)$  which are central to the proof of [Theorem 1.1](#).

**Example** Let  $Z_0(q) = E^*(4\tau)/\eta(8\tau)^3$  and define  $f_m(\tau)$  by

$$f_m(\tau) := \frac{\Theta_4(\tau)^9 (16\Theta_2(\tau)^4 + \Theta_3(\tau)^4)^m}{\Theta_2(\tau)^{2m+3} \Theta_3(\tau)^{2m+3}}.$$

The proof shows that the constant terms of  $Z_0(q) f_m(\tau)$  (see [\(102\)](#)) vanish for all  $m$ . The initial terms of the expansions of the first few  $f_m(\tau)$  are

$$\begin{aligned} f_0(\tau) &= q^{-3} - 24q + 273q^5 - 1976q^9 + \dots, \\ f_1(\tau) &= q^{-5} - 4q^{-1} - 269q^3 + 5188q^7 + \dots, \\ f_2(\tau) &= q^{-7} + 16q^{-3} - 411q + 272q^5 + \dots, \\ f_3(\tau) &= q^{-9} + 36q^{-5} - 153q^{-1} - \dots, \end{aligned}$$

and the initial terms of  $Z_0(q) f_m(\tau)$  are

$$\begin{aligned} Z_0(q) f_0(\tau) &= q^{-4} - 276q^4 + 4096q^8 - 33606q^{12} + \dots, \\ Z_0(q) f_1(\tau) &= q^{-6} + 20q^{-2} - 338q^2 - 1208q^6 + \dots, \\ Z_0(q) f_2(\tau) &= q^{-8} + 40q^{-4} - 8992q^4 + 65260q^8 + \dots, \\ Z_0(q) f_3(\tau) &= q^{-10} + 60q^{-6} + 738q^{-2} - 11256q^2 - \dots. \end{aligned}$$

One easily sees that the constant terms of these  $Z_0(q) f_m(\tau)$  indeed vanish.

**Example** Here we simplify the notation in [Theorem 4.26](#) by defining  $q$ -series  $\widehat{\mathbf{D}}_{m,n}^{(i)}(q)$  and  $\Lambda^{(i)}(m, n, k, j; q)$  so that

$$(107) \quad \widehat{\mathbf{D}}_{m,n}^{(i)}(q) := \mathbf{D}_{m,n}^{(i)}(q^8) = \sum_{k=0}^n \sum_{j=0}^k \Lambda^{(i)}(m, n, k, j; q).$$

By [Theorem 4.26](#), [Theorem 1.1](#) follows from the assertion that the constant term of

$$\widehat{\mathbf{D}}_{m,n}^{(1)}(q) - \widehat{\mathbf{D}}_{m,n}^{(2)}(q)$$

vanishes. The first few terms of the series  $\Lambda^{(i)}(3, 1, k, j; q)$  are

$$\begin{aligned} \Lambda^{(1)}(3, 1, 0, 0; q) &= \frac{1}{256}q^{-8} + \frac{43}{256}q^{-4} + \frac{7}{16} - \dots, \\ \Lambda^{(1)}(3, 1, 1, 0; q) &= \frac{1}{768}q^{-8} + \frac{35}{768}q^{-4} - \frac{13}{48} - \dots, \\ \Lambda^{(1)}(3, 1, 1, 1; q) &= -\frac{1}{768}q^{-8} - \frac{59}{768}q^{-4} - \frac{85}{96} + \dots, \\ \Lambda^{(2)}(3, 1, 0, 0; q) &= -\frac{1}{3072}q^{-12} - \frac{7}{256}q^{-8} - \frac{11}{16}q^{-4} - \frac{85}{96} + \dots, \\ \Lambda^{(2)}(3, 1, 1, 0; q) &= \frac{1}{3072}q^{-12} + \frac{5}{256}q^{-8} + \frac{13}{64}q^{-4} - \frac{247}{48} - \dots, \\ \Lambda^{(2)}(3, 1, 1, 1; q) &= \frac{1}{1024}q^{-12} - \frac{13}{256}q^{-8} - \frac{203}{64}q^{-4} + \frac{85}{16} + \dots. \end{aligned}$$

One sees that the constant term of  $\widehat{\mathbf{D}}_{3,1}^{(1)}(q) - \widehat{\mathbf{D}}_{3,1}^{(2)}(q)$  is

$$\frac{7}{16} - \frac{13}{48} - \frac{85}{96} + \frac{85}{96} + \frac{247}{48} - \frac{85}{16} = 0.$$

Notice that the constant terms of  $\Lambda^{(1)}(3, 1, 1, 1; q)$  and  $\Lambda^{(2)}(3, 1, 0, 0)$  agree. This equality is a special case of a general equality whose proof is equivalent to the  $n = 0$  case of the proof of [Theorem 1.1](#). In general, the constant terms of  $\Lambda^{(1)}(m, n, 0, 0; q)$  and  $\Lambda^{(2)}(m, n, n, n; q)$  agree.

**Example** The proof of [Theorem 1.1](#) relies on the fact that the expressions in [Theorem 4.26](#) can be written as (see [\(106\)](#))

$$(108) \quad \left( q \frac{d}{dq} \widehat{Z}_0(q^{1/8}) \right) \widehat{Z}_0(q^{1/8})^m P_n(\widehat{Z}_0(q^{1/8})),$$

where  $P_n(x)$  is a polynomial. In the table below we give the first few polynomials.

$n$	$P_n(x)$
0	$x$
1	$-\frac{1}{2}x^2 + 72$
2	$\frac{1}{12}x^3 + \frac{32}{3}x$
3	$-\frac{1}{120}x^4 + \frac{122}{15}x^2 - 64$
4	$\frac{1}{1680}x^5 + \frac{8}{7}x^3 + \frac{5696}{105}x$
5	$-\frac{1}{30240}x^6 + \frac{457}{1890}x^4 + \frac{9536}{315}x^2 + \frac{406016}{315}$
6	$\frac{1}{665280}x^7 + \frac{4}{135}x^5 + \frac{1712}{315}x^3 - \frac{2089984}{3465}x$

## References

- [1] **L Álvarez-Gaumé, M Mariño, F Zamora**, *Softly broken  $N = 2$  QCD with massive quark hypermultiplets, I*, Internat. J. Modern Phys. A 13 (1998) 403–430 [MR1621192](#)
- [2] **G E Andrews**, *Mordell integrals and Ramanujan’s “lost” notebook*, from: “Analytic number theory (Philadelphia, PA, 1980)”, (MI Knopp, editor), Lecture Notes in Math. 899, Springer, Berlin (1981) 10–18 [MR654518](#)
- [3] **MF Atiyah**, *The logarithm of the Dedekind  $\eta$ -function*, Math. Ann. 278 (1987) 335–380 [MR909232](#)
- [4] **MF Atiyah, IM Singer**, *Dirac operators coupled to vector potentials*, Proc. Nat. Acad. Sci. U.S.A. 81 (1984) 2597–2600 [MR742394](#)
- [5] **K Bringmann, K Ono**, *The  $f(q)$  mock theta function conjecture and partition ranks*, Invent. Math. 165 (2006) 243–266 [MR2231957](#)
- [6] **K Bringmann, K Ono**, *Dyson’s ranks and Maass forms*, Ann. of Math. 171 (2010) 419–449 [MR2630043](#)
- [7] **K Bringmann, K Ono, R C Rhoades**, *Eulerian series as modular forms*, J. Amer. Math. Soc. 21 (2008) 1085–1104 [MR2425181](#)
- [8] **J H Bruinier**, *Borchers products on  $O(2, l)$  and Chern classes of Heegner divisors*, Lecture Notes in Math. 1780, Springer, Berlin (2002) [MR1903920](#)

- [9] **J H Bruinier, J Funke**, *On two geometric theta lifts*, Duke Math. J. 125 (2004) 45–90 [MR2097357](#)
- [10] **J H Bruinier, K Ono**, *Heegner divisors,  $L$ -functions and harmonic weak Maass forms*, Ann. of Math. 172 (2010) 2135–2181 [MR2726107](#)
- [11] **J H Bruinier, G van der Geer, G Harder, D Zagier**, *The 1-2-3 of modular forms*, (K Ranestad, editor), Universitext, Springer, Berlin (2008) [MR2385372](#) Lectures from the Summer School on Modular Forms and their Applications held in Nordfjordeid, June 2004
- [12] **J H Bruinier, T Yang**, *Faltings heights of CM cycles and derivatives of  $L$ -functions*, Invent. Math. 177 (2009) 631–681 [MR2534103](#)
- [13] **Y Choie, M H Lee**, *Rankin–Cohen brackets on pseudodifferential operators*, J. Math. Anal. Appl. 326 (2007) 882–895 [MR2280950](#)
- [14] **S K Donaldson, P B Kronheimer**, *The geometry of four-manifolds*, Oxford Math. Monogr., Oxford Science Publ., The Clarendon Press, Oxford Univ. Press, New York (1990) [MR1079726](#)
- [15] **F J Dyson**, *A walk through Ramanujan’s garden*, from: “Ramanujan revisited (Urbana-Champaign, IL, 1987)”, (G E Andrews, R A Askey, B C Berndt, K G Ramanathan, R A Rankin, editors), Academic Press, Boston (1988) 7–28 [MR938957](#)
- [16] **G Ellingsrud, L Göttsche**, *Wall-crossing formulas, the Bott residue formula and the Donaldson invariants of rational surfaces*, Quart. J. Math. Oxford Ser. 49 (1998) 307–329 [MR1645556](#)
- [17] **D S Freed, K K Uhlenbeck**, *Instantons and four-manifolds*, second edition, MSRI Publ. 1, Springer, New York (1991) [MR1081321](#)
- [18] **A L Gorodenzev**, *Top Chern classes of universal bundles on the complex projective plane and correlation functions of asymptotically free SYM  $N = 2$  QFT*, J. Math. Sci. (New York) 106 (2001) 3240–3257 [MR1878046](#)
- [19] **L Göttsche**, *Modular forms and Donaldson invariants for 4-manifolds with  $b_+ = 1$* , J. Amer. Math. Soc. 9 (1996) 827–843 [MR1362873](#)
- [20] **L Göttsche**, *Donaldson invariants in algebraic geometry*, from: “School on Algebraic Geometry (Trieste, 1999)”, (L Göttsche, editor), ICTP Lect. Notes 1, Abdus Salam Int. Cent. Theoret. Phys., Trieste (2000) 101–134 [MR1795862](#)
- [21] **L Göttsche, H Nakajima, K Yoshioka**, *Instanton counting and Donaldson invariants*, J. Differential Geom. 80 (2008) 343–390 [MR2472477](#)
- [22] **L Göttsche, D Zagier**, *Jacobi forms and the structure of Donaldson invariants for 4-manifolds with  $b_+ = 1$* , Selecta Math. 4 (1998) 69–115 [MR1623706](#)
- [23] **F Hirzebruch, D Zagier**, *Intersection numbers of curves on Hilbert modular surfaces and modular forms of Nebentypus*, Invent. Math. 36 (1976) 57–113 [MR0453649](#)



- [24] **A A Klyachko**, *Moduli of vector bundles and numbers of classes*, Funktsional. Anal. i Prilozhen. 25 (1991) 81–83 [MR1113131](#)
- [25] **D Kotschick**, *SO(3)–invariants for 4–manifolds with  $b_2^+ = 1$* , Proc. London Math. Soc. 63 (1991) 426–448 [MR1114516](#)
- [26] **D Kotschick**, **P Lisca**, *Instanton invariants of  $\mathbb{C}P^2$  via topology*, Math. Ann. 303 (1995) 345–371 [MR1348804](#)
- [27] **D Kotschick**, **J W Morgan**, *SO(3)–invariants for 4–manifolds with  $b_2^+ = 1$ . II*, J. Differential Geom. 39 (1994) 433–456 [MR1267898](#)
- [28] **P B Kronheimer**, **T S Mrowka**, *Embedded surfaces and the structure of Donaldson’s polynomial invariants*, J. Differential Geom. 41 (1995) 573–734 [MR1338483](#)
- [29] **J M F Labastida**, **C Lozano**, *Duality in twisted  $\mathcal{N} = 4$  supersymmetric gauge theories in four dimensions*, Nuclear Phys. B 537 (1999) 203–242 [MR1659299](#)
- [30] **T G Leness**, *Degeneracy loci of families of Dirac operators*, Trans. Amer. Math. Soc. 364 (2012) 5995–6008 [MR2946940](#)
- [31] **A Losev**, **N Nekrasov**, **S Shatashvili**, *Issues in topological gauge theory*, Nuclear Phys. B 534 (1998) 549–611 [MR1663467](#)
- [32] **A Malmendier**, *Expressions for the generating function of the Donaldson invariants for  $\mathbb{C}P^2$* , PhD thesis, Massachusetts Institute of Technology (2007) Available at <http://hdl.handle.net/1721.1/38959>
- [33] **A Malmendier**, *The signature of the Seiberg–Witten surface*, from: “Surveys in differential geometry. Volume XV. Perspectives in mathematics and physics”, (T Mrowka, S-T Yau, editors), Surv. Differ. Geom. 15, Int. Press, Somerville, MA (2011) 255–277 [MR2815730](#)
- [34] **M Mariño**, **G Moore**, *Integrating over the Coulomb branch in  $\mathcal{N} = 2$  gauge theory*, Nuclear Phys. B Proc. Suppl. 68 (1998) 336–347 [MR1641983](#) Strings ’97 (Amsterdam, 1997)
- [35] **R Miranda**, *Persson’s list of singular fibers for a rational elliptic surface*, Math. Z. 205 (1990) 191–211 [MR1076128](#)
- [36] **R Miranda**, *An overview of algebraic surfaces*, from: “Algebraic geometry (Ankara, 1995)”, (S Sertöz, editor), Lecture Notes in Pure and Appl. Math. 193, Dekker, New York (1997) 157–217 [MR1483329](#)
- [37] **G Moore**, **E Witten**, *Integration over the  $u$ –plane in Donaldson theory*, Adv. Theor. Math. Phys. 1 (1997) 298–387 [MR1605636](#)
- [38] **W Nahm**, *On electric-magnetic duality*, Nuclear Phys. B Proc. Suppl. 58 (1997) 91–96 [MR1486333](#) Advanced quantum field theory (La Londe les Maures, 1996)
- [39] **K Oguiso**, **T Shioda**, *The Mordell–Weil lattice of a rational elliptic surface*, Comment. Math. Univ. St. Paul. 40 (1991) 83–99 [MR1104782](#)

- [40] **K Ono**, *Unearthing the visions of a master: harmonic Maass forms and number theory*, from: “Current developments in mathematics, 2008”, (D Jerison, B Mazur, T Mrowka, W Schmid, R P Stanley, S-T Yau, editors), Int. Press, Somerville, MA (2009) 347–454 [MR2555930](#)
- [41] **U Persson**, *Configurations of Kodaira fibers on rational elliptic surfaces*, Math. Z. 205 (1990) 1–47 [MR1069483](#)
- [42] **N Seiberg, E Witten**, *Electric-magnetic duality, monopole condensation, and confinement in  $N = 2$  supersymmetric Yang–Mills theory*, Nuclear Phys. B 426 (1994) 19–52 [MR1293681](#)
- [43] **N Seiberg, E Witten**, *Monopoles, duality and chiral symmetry breaking in  $N = 2$  supersymmetric QCD*, Nuclear Phys. B 431 (1994) 484–550 [MR1306869](#)
- [44] **Y Shimizu**, *Seiberg–Witten integrable systems and periods of rational elliptic surfaces*, from: “Primes and knots”, (T Kohno, M Morishita, editors), Contemp. Math. 416, Amer. Math. Soc. (2006) 237–247 [MR2276144](#)
- [45] **G Shimura**, *On modular forms of half integral weight*, Ann. of Math. 97 (1973) 440–481 [MR0332663](#)
- [46] **T Shioda**, *On elliptic modular surfaces*, J. Math. Soc. Japan 24 (1972) 20–59 [MR0429918](#)
- [47] **P F Stiller**, *Elliptic curves over function fields and the Picard number*, Amer. J. Math. 102 (1980) 565–593 [MR584462](#)
- [48] **P F Stiller**, *Monodromy and invariants of elliptic surfaces*, Pacific J. Math. 92 (1981) 433–452 [MR618076](#)
- [49] **S A Strømme**, *Ample divisors on fine moduli spaces on the projective plane*, Math. Z. 187 (1984) 405–423 [MR757480](#)
- [50] **C Vafa, E Witten**, *A strong coupling test of  $S$ -duality*, Nuclear Phys. B 431 (1994) 3–77 [MR1305096](#)
- [51] **E Witten**, *Topological quantum field theory*, Comm. Math. Phys. 117 (1988) 353–386 [MR953828](#)
- [52] **E Witten**, *Monopoles and four-manifolds*, Math. Res. Lett. 1 (1994) 769–796 [MR1306021](#)
- [53] **E Witten**, *On  $S$ -duality in abelian gauge theory*, Selecta Math. 1 (1995) 383–410 [MR1354602](#)
- [54] **K Yoshioka**, *The Betti numbers of the moduli space of stable sheaves of rank 2 on  $\mathbf{P}^2$* , J. Reine Angew. Math. 453 (1994) 193–220 [MR1285785](#)
- [55] **D Zagier**, *Ramanujan’s mock theta functions and their applications (after Zwegers and Ono–Bringmann)*, from: “Séminaire Bourbaki. Vol. 2007/2008”, Astérisque 326, Soc. Math. France (2009) vii–viii, 143–164 [MR2605321](#)

- [56] **SP Zwegers**, *Mock theta functions*, PhD thesis, Universiteit Utrecht (2002)  
Available at <http://igitur-archive.library.uu.nl/dissertations/2003-0127-094324/inhoud.htm>

*Department of Mathematics and Statistics, Colby College  
Waterville MN 04901, USA*

*Mathematics and Computer Science, Emory University  
Atlanta GA 30322, USA*

[amalmend@colby.edu](mailto:amalmend@colby.edu), [ono@mathcs.emory.edu](mailto:ono@mathcs.emory.edu)

<http://www.mathcs.emory.edu/~ono/>

Proposed: Walter Neumann

Seconded: Jim Bryan, Simon Donaldson

Received: 28 April 2010

Revised: 1 May 2012