

2–strand twisting and knots with identical quantum knot homologies

ANDREW LOBB

Given a knot, we ask how its Khovanov and Khovanov–Rozansky homologies change under the operation of introducing twists in a pair of strands. We obtain long exact sequences in homology and further algebraic structure which is then used to derive topological and computational results. Two of our applications include giving a way to generate arbitrary numbers of knots with isomorphic homologies and finding an infinite number of mutant knot pairs with isomorphic reduced homologies.

57M25

1 Introduction and results

In this paper we consider $\mathrm{sl}(n)$ Khovanov–Rozansky homology [6] (Khovanov homology appears as $n = 2$) under the operation of adding twists in a pair of strands. We observe stabilization of the homology as we add more twists and, looking a little deeper, reveal some further algebraic structure which we exploit for various structural and topological results.

We start by giving statements of and context for some results whose proofs follow in later sections (Section 2 for the algebraic proofs necessary for the topological and structural results, which are themselves proved in Section 3).

1.1 Topological and structural results on Khovanov–Rozansky homology

By relating Khovanov homology with their own instanton knot Floer homology, Kronheimer and Mrowka have shown that Khovanov homology detects the unknot [9]. It is still an open question whether the Jones polynomial, which is the graded Euler characteristic of Khovanov homology, detects the unknot. However, it is known that the Jones polynomial (and likewise the HOMFLY polynomial) does not enjoy the stronger property of being a complete invariant able to distinguish between any pair of knots. For example, the HOMFLY polynomial is unable to distinguish between mutant knots.

It has been verified by Mackaay and Vaz [20] that the mutant knot pair consisting of the Kinoshita–Terasaka and the Conway knots have isomorphic reduced Khovanov–Rozansky homologies and hence also isomorphic reduced HOMFLY homologies. Furthermore, there exist families of distinct 2–bridge knots with the same HOMFLY polynomials. Since 2–bridge knots have thin homology, these knots must also share isomorphic reduced Khovanov–Rozansky homologies; see Rasmussen [14].

Our first use of the algebraic structure we observe is to produce new families of knots with isomorphic Khovanov–Rozansky homologies.

Theorem 1.1 *Given a natural number m , there are m distinct prime knots with bridge number greater than 2, which have isomorphic $\mathrm{sl}(n)$ Khovanov–Rozansky homologies for all n .*

We note that Theorem 1.1 holds for reduced, unreduced and equivariant homology with potential $w = x^{n+1} + ax$, and consequently also for reduced HOMFLY homology. The knots undistinguished by these flavors of Khovanov–Rozansky homology that we produce are not necessarily thin nor necessarily related by mutation. For an example of two knots with isomorphic Khovanov–Rozansky homologies, see Figure 8 and the discussion in Section 3.1.

In fact, the m knots we construct for Theorem 1.1 can be distinguished by the hyperbolic structures on their complements, using Thurston’s theorem on hyperbolic Dehn surgery [19].

It remains a motivating question whether topological conclusions may be drawn from the coincidence of Khovanov–Rozansky homologies. Further consequences of Theorem 1.1 and its proof are discussed in Sections 3.1 and 3.2, where we give specific examples of interesting phenomena including an example on mutant knot pairs.

Theorem 1.2 *There exists an infinite number of mutant knot pairs with isomorphic reduced Khovanov–Rozansky homologies (and hence also isomorphic reduced HOMFLY homologies).*

From now on, all complexes are understood to be complexes of matrix factorizations, and $C(K)$ and $H(K)$ stand for the $\mathrm{sl}(n)$ Khovanov–Rozansky chain complex and homology of the knot K respectively for some fixed $n \geq 2$. Sometimes we will mean specifically the reduced, unreduced or equivariant (with potential $w = x^{n+1} - ax$) homologies (see Krasner [8]) in which case we shall make it clear. Otherwise results should be interpreted as holding for each of these three versions of Khovanov–Rozansky homology.

Our next application concerns the concordance homomorphism s_n (see Wu [21] and the author [12; 13]) arising from a perturbation of $sl(n)$ Khovanov–Rozansky homology.

Theorem 1.3 *Let the knot K_0 be obtained by changing a crossing of K_{-1} from negative to positive as in Figure 1. Then we know by [13] and Livingston [11, Corollary 3] that we must have*

$$s_n(K_0) \leq s_n(K_{-1}).$$

If in fact we have strict inequality $s_n(K_0) < s_n(K_{-1})$ then the homology group in homological degree $2p$ satisfies

$$H^{2p}(K_p) \neq 0$$

for the sequence of knots K_1, K_2, \dots shown in Figure 1.




Figure 1: Here we show a knot K_{-1} differing from a knot K_0 by a single crossing change. We have drawn local pictures of diagrams of these knots. The knots K_p for $p \geq 1$ have diagrams D_p formed by making p further positive crossing changes at the same site as shown. Alternatively, one can think of the knot K_{-1} and the knots K_p as obtained from K_0 by replacing the tangle T_0 shown in D_0 by T_{-1} or T_p respectively.

We obtain an interesting topological corollary. By the definition of the Khovanov–Rozansky chain complex it is clear that Theorem 1.3 gives the following.

Corollary 1.4 *Given the conditions of Theorem 1.3 the knot K_p must have at least $2p$ positive crossings in any diagram.* □

In other words, the crossings in K_p shown in Figure 1 are in some sense essential. We note that each s_n provides a tight bound on the unknotting number of a torus knot and, in the standard diagram of a torus knot, a single crossing change anywhere results in a knot with a smaller unknotting number. Hence Corollary 1.4 can be applied in this situation.

The exact sequences giving rise to our topological and structural results work best when we can identify one of the terms. In particular, we expect to be able to say useful things about knots with unknotting number equal to 1.

Theorem 1.5 *We consider the situation of Figure 1 where we take $K_0 = U$, the unknot. Then we have*

$$s_n(K_p) = s_n(K_1)$$

for all $p \geq 1$.

In the case where $s_n(K_1) = 0$, we can say more about the homology of the knot K_p . In fact, the homology of K_p is characterized entirely by p and the homology of the knot K_1 . We state this first for the equivariant case.

Theorem 1.6 *We consider the situation of Figure 1 where we take $K_0 = U$, the unknot, and assume that $s_n(K_1) = 0$. Taking equivariant homology with potential $w = x^{n+1} - ax$, let Δ be the bigraded $\mathbb{C}[a]$ -module isomorphic to the torsion part of $H(K_1)$. Then for $p \geq 2$ we have*

$$H(K_p) = H(K_{p-1}) \oplus \Delta[2p]\{2n(1-p)\}.$$

It is almost possible to characterize completely the homology of K_p in terms of p and the homology of K_{p-1} even if $s_n(K_1) \neq 0$. In fact, just knowing $H(K_{p-1})$ we would know $H(K_p)$ in all homological degrees apart from possibly one, and to determine $H(K_p)$ in this degree we would need one more piece of information. We discuss what piece of information this is following the proof of Theorem 1.6. Armed with Theorem 1.6, we can also consider the nonequivariant cases.

Theorem 1.7 *Suppose we are in the setup of Theorem 1.6 and let $H(K)$ stand for the standard unreduced or reduced Khovanov–Rozansky homology of K . Let Δ be the bigraded \mathbb{C} -module satisfying*

$$H(K_1) = \mathbb{C}[0]\{0\} \oplus \Delta$$

for the reduced case and

$$H(K_1) = \mathbb{C}[0]\{1-n\} \oplus \mathbb{C}[0]\{3-n\} \oplus \cdots \oplus \mathbb{C}[0]\{n-1\} \oplus \Delta$$

for the unreduced case. Then for $p \geq 2$ we have

$$H(K_p) = H(K_{p-1}) \oplus \Delta[2p]\{2n(1-p)\}.$$

1.2 Algebraic preliminaries

First we describe some chain complexes of matrix factorizations, one such for each integer, which will be the building blocks of this paper.

Definition 1.8 For $k \geq 0$, the complex T_k is the $\mathfrak{sl}(n)$ Khovanov–Rozansky chain complex of direct sums of matrix factorizations corresponding to a diagram of k full twists in two oppositely oriented strands, where the $2k$ crossings are positive (see Figures 2 and 3 for an explicit picture). When $k < 0$ we take the $-2k$ crossings to be negative. By abuse of notation sometimes we shall also refer to the corresponding tangle diagram itself by T_k .

It should be clear that there is an obvious way in which each of these complexes can be built from T_1 and T_{-1} by tensor product.

Proposition 1.9 Up to homotopy equivalence $T_k \otimes T_l = T_{k+l}$, where the tensor product of complexes of matrix factorizations is taken by concatenating in the obvious way the corresponding tangle diagrams with $|2k|$ and $|2l|$ crossings.

Proof For k and l of the same sign this is by definition, and for k and l of opposite sign it follows from the invariance up to homotopy equivalence of the Khovanov–Rozansky chain complex under Reidemeister move II. □




Figure 2: The complex T_1 is the $\mathfrak{sl}(n)$ Khovanov–Rozansky complex of direct sums of matrix factorizations corresponding to this diagram. Note that there are two positive crossings in the diagram.

In [15] Rasmussen gives a homomorphism $s: K \mapsto s(K) \in 2\mathbb{Z}$ from the smooth knot concordance group to the additive group of even integers. Furthermore, he shows that s provides a lower bound $|s(K)|/2$ on the smooth slice genus of a knot K . Rasmussen’s construction proceeds by extracting an even integer $s(K)$ from the E_∞ page of a spectral sequence which has as E_2 page the standard Khovanov homology of K . This spectral sequence is essentially due to Lee [10].




Figure 3: The complex $T_k = \bigotimes^k T_1$ is the $\mathfrak{sl}(n)$ Khovanov–Rozansky complex corresponding to the diagram above with $2k$ crossings.

Since this seminal paper, there have been generalizations of this result for other quantum knot homologies. In particular Gornik [2] has constructed a spectral sequence with E_2 page $\mathfrak{sl}(n)$ Khovanov–Rozansky homology $H(K)$. In [13], the author shows that the E_∞ page of Gornik’s spectral sequence is equivalent to an even integer $s_n(K)$ which gives a homomorphism $s_n: K \mapsto s_n(K) \in 2\mathbb{Z}$ from the smooth knot concordance group to the additive group of even integers. Earlier work by the author [12] and independently by Wu [21] implies that $|s_n(K)|/2(n-1)$ is a lower bound on the smooth slice genus of K .

In [13] it is shown that the E_∞ page of Gornik’s spectral sequence is isomorphic as a graded group to the homology of the unknot but with a shift in quantum grading $E_\infty \cong H(U)\{s_n(K)\}$, so that all the information about E_∞ is contained in the even integer $s_n(K)$.

In [15], Rasmussen asked if the concordance homomorphism s coming from Khovanov homology was the same as the concordance homomorphism τ coming Heegaard–Floer knot homology, a conjecture motivated by the observation that s and τ share many of the same properties. A negative answer to Rasmussen’s question was first provided by Hedden and Ording [3].

The homomorphisms s_n also share many properties with s and moreover we know that both s and s_n arise from the quantum world. It is an interesting open question whether the homomorphisms s_n are equivalent to the homomorphism $s = -s_2$ (see [13, Conjecture 1.5]).

We briefly discuss the relationship between s_n and equivariant Khovanov–Rozansky homology [8] with potential $w = x^{n+1} - ax$. Here, all the modules involved in the Khovanov–Rozansky complex are free $\mathbb{C}[a]$ -modules where a has quantum grading $2n$. The reason we are interested in this version of Khovanov–Rozansky homology is that the s_n invariant is then built into the homology. In fact for any knot K , we have that

the equivariant homology with this potential satisfies

$$H(K) = \text{tor} \oplus \bigoplus_{l=1}^n \mathbb{C}[a][0]\{2l - n - 1 + s_n(K)\},$$

where tor is a finitely-generated torsion $\mathbb{C}[a]$ -module.

To see this, observe that $C(K)$, as a freely-generated graded complex of $\mathbb{C}[a]$ -modules, is chain homotopy equivalent to a sum of complexes of the form:

- (1) $0 \rightarrow \mathbb{C}[a] \rightarrow 0$
- (2) $0 \rightarrow \mathbb{C}[a] \xrightarrow{a^k} \mathbb{C}[a] \rightarrow 0$

Setting $a = 0$ we recover standard Khovanov–Rozansky homology, while setting $a = 1$ destroys the quantum grading and gives us Gornik’s version of Khovanov–Rozansky homology. This also tells us that nothing is lost by considering equivariant homology since the nonequivariant unreduced homology can be obtained from the equivariant homology groups.

1.3 Stabilization and exact sequences

If we have a knot K given by a diagram D we may consider T_0 as a subtangle of D . Replacing T_0 by T_1, T_2, T_3, \dots in D we obtain a sequence of diagrams D_1, D_2, D_3, \dots and hence a sequence of knots K_1, K_2, K_3, \dots .

In the following theorems we let D be such a diagram with a subtangle of D identified with T_0 . We write c_- and c_+ for the number of negative crossings and for the number of positive crossings of D respectively.

Theorem 1.10 *For each $0 \leq i < j$ there exists a directed system of maps (to be defined)*

$$F_{i,j}: T_i \rightarrow T_j$$

that is graded of homological degree 0 and of quantum degree 0. Then for $0 \leq i < j$ (we allow $j = \infty$) we have that the induced map on homology

$$F_{i,j}: H(D_i) \rightarrow H(D_j)$$

is an isomorphism in all homological degrees $\leq 2i - c_- - 2$.

Using square brackets to denote a shift in homological grading, and curly brackets to denote a shift in quantum grading, we also have the following.

Theorem 1.11 For each $0 \leq i < j$ there exists a directed system of maps (to be defined)

$$G_{i,j}: T_i \rightarrow T_j[2(i-j)]\{2n(j-i)\}$$

that is graded of homological degree 0 and of quantum degree 0. Then for $0 \leq i < j$ we have that the induced map on homology

$$G_{i,j}: H(D_i) \rightarrow H(D_j)[2(i-j)]\{2n(j-i)\}$$

is an isomorphism in all homological degrees $\geq c_+$.

Remark To shorten our exposition, in this paper we restrict ourselves to the tangles T_k where the $2k$ crossings are positive. For each theorem we state, there is a dual theorem using negative crossings that interested readers should have no trouble in stating and proving for themselves.

If this were all that there were to say about the algebra, we would not expect to be able to prove interesting results. However, the maps $F_{i,j}$ and $G_{i,j}$ mesh well together, in a sense that we shall later make explicit.

From homology theories in different branches of mathematics we know that short exact sequences of chain complexes (and hence long exact sequences of homology groups) are useful tools when they are found in a homology theory. And even more so are morphisms of short exact sequences of chain complexes (giving natural maps between long exact sequences of homology groups). We find these relatively easily in our setup and it is these that provide the power to start proving our later topological and structural results.

The results on exact sequences are best stated in the next section, after Theorems 1.10 and 1.11 are established. For those wishing to jump ahead, these results appear as Propositions 2.6 and 2.7.

We do not expect that the topological and structural corollaries that we find represent all of that which can be proved by making use of our exact sequences. We therefore end this subsection with an encouragement for others to play with these exact sequences and see what else may drop out!

Acknowledgements This paper would not have been possible without the input of Daniel Krasner. Akio Kawachi has supplied invaluable advice concerning his paper [4]. The author would also like to thank Cameron Gordon, Marc Lackenby, Jacob Rasmussen and Scott Taylor for useful emails or discussions. Thanks to Josh Greene and Dylan Thurston for comments on earlier versions of this paper.

Much of the work in this paper was done during the Knot Homology semester at MSRI, where the author was funded as a postdoctoral fellow.

2 Algebraic structure results

In this section we shall prove Theorems 1.10 and 1.11 and derive further results enabling us to prove our more topological theorems.

2.1 Stabilization

By *stabilization* we mean, most basically, the existence of a complex T_∞ , the direct limit of a sequence of maps $T_k \rightarrow T_{k+1}$. This complex T_∞ is defined in Definition 2.2.

As before, if we have a knot K given by a diagram D we may consider T_0 as a subtangle of D . Replacing T_0 by T_1, T_2, T_3, \dots in D we obtain a sequence of diagrams D_1, D_2, D_3, \dots and hence a sequence of knots K_1, K_2, K_3, \dots .

In the chain complex $C(D_i)$, T_i appears as a tensor factor. Replacing T_i by T_∞ gives us a chain complex which we shall denote $C(D_\infty)$ and its homology by $H(D_\infty)$. We have, in effect, replaced the T_i tangle in D_i by a “tangle consisting of an infinite number of twists.”

To simplify notation we shall write V and Z (vertical and horizontal) for the matrix factorizations indicated in Figure 4.




Figure 4: We draw here the matrix factorizations V and Z . In the text of this paper, V and Z often appear with integers appended in curly and/or square parentheses to indicate quantum degree shift and homological degree respectively.

In [7], Krasner gave a compact description of the complex T_k of our Definition 1.8. As a consequence of this description, one can see that knot diagrams built up from these tangle building blocks have associated chain complexes which avoid the “thick-edged” matrix factorization, and hence much of the complication usually involved in the Khovanov–Rozansky chain complex. Understanding such knots may well be a good way to begin getting a grasp on Khovanov–Rozansky homology.

This compact description of T_k will essentially be our main ingredient. In the theorem that follows we use curly or square parentheses to indicate shift in the quantum degree and homological degree respectively and w is the *potential*. We state Krasner’s Theorem both for the standard potential $w = x^{n+1}$ and for the equivariant potential $w = x^{n+1} - ax$, although Krasner only stated it for the standard potential. Since the results that go into the proof of Krasner’s Theorem have now been established in the general equivariant setting [8], we can state the result in more generality.

Theorem 2.1 (Krasner [7]) *Up to chain homotopy equivalence, the complex T_k is isomorphic to the following chain complex of matrix factorizations:*

$$V[0]\{1 - n\} \xrightarrow{x_2 - x_4} V[1]\{-1 - n\} \xrightarrow{A} V[2]\{1 - 3n\} \xrightarrow{x_2 - x_4} \dots$$

$$\xrightarrow{x_2 - x_4} V[2k - 1]\{(1 - 2k)n - 1\} \xrightarrow{S} Z[2k]\{-2kn\},$$

where we write

$$A = x_2^{n-1} + x_2^{n-2}x_4 + x_2^{n-3}x_4^2 + \dots + x_4^{n-1}$$

and we write S for the map induced by the saddle cobordism.

Definition 2.2 Setting $k = \infty$ in Theorem 2.1 gives us a definition of a complex T_∞ .

With Krasner’s characterization, it is a quick matter to define the chain maps $F_{i,j}$ and $G_{i,j}$ of Theorems 1.10 and 1.11.

Definition 2.3 Let $0 \leq i < j$. Using the description of Theorem 2.1 of the complexes T_k , we define two maps

$$F_{i,j}: T_i \rightarrow T_j,$$

$$G_{i,j}: T_i \rightarrow T_j[2(i - j)]\{2n(j - i)\},$$

as follows. We require that $F_{i,j}$ preserves the homological grading and is the identity map on the matrix factorizations in all homological degrees less than $2i$. To the component of $F_{i,j}$ in homological degree $2i$ we assign the map $S' = (-1/n + 1)S$ where S is the map of matrix factorizations associated to the saddle cobordism. To check that $F_{i,j}$ is a chain map, it is enough to observe that

$$S^2 = -(n + 1)A \quad \text{and} \quad (x_2 - x_4) \circ S = 0.$$

The former of these identities is computed in detail by Khovanov and Rozansky in [5, Appendix A]. For the latter note that up to homotopy we have

$$\begin{aligned} (x_2 - x_4) \circ S &= x_2 \circ S - x_4 \circ S \\ &= x_2 \circ S - S \circ x_4 = x_2 \circ S - S \circ x_1 \\ &= x_2 \circ S - x_1 \circ S = x_2 \circ S - x_2 \circ S = 0. \end{aligned}$$

Clearly $F_{i,j}$ preserves the quantum grading.

We require that $G_{i,j}$ is the identity map on all homological degrees of T_i which are nonzero matrix factorizations. Certainly then $G_{i,j}$ is a chain map and we see that it is quantum graded of degree 0.

With these definitions in hand, the path to proving Theorems 1.10 and 1.11 is straightforward: in brief, we compute the cones of the maps $F_{i,j}$ and $G_{i,j}$ and show that the homology of the cones is supported well away from certain degrees in which $F_{i,j}$ and $G_{i,j}$ must therefore induce isomorphisms.

In the following propositions, we leave out quantum grading shifts and only give the leftmost and rightmost homological gradings. We do this in order to try and give an uncluttered exposition; for the reader who is making use of these propositions, we recommend having a copy of Krasner’s [7] to hand.

Proposition 2.4 *Writing $\text{Co}(F_{i,j})$ for the cone of $F_{i,j}$ we have*

$$\text{Co}(F_{i,j}) = Z[2i - 1] \xrightarrow{S'} V \xrightarrow{x_2 - x_4} V \xrightarrow{A} V \xrightarrow{x_2 - x_4} V \xrightarrow{A} V \dots \xrightarrow{x_2 - x_4} V \xrightarrow{S} Z[2j].$$

Proposition 2.5 *Writing $\text{Co}(G_{i,j})$ for the cone of $G_{i,j}$ we have*

$$\text{Co}(G_{i,j}) = V[2(i - j)] \xrightarrow{x_2 - x_4} V \xrightarrow{A} V \xrightarrow{x_2 - x_4} V \xrightarrow{A} V \dots \xrightarrow{x_2 - x_4} V[-1].$$

Proof of Propositions 2.4 and 2.5 This is a straightforward application of Gaussian elimination. Starting from the leftmost homological degree in the case of $F_{i,j}$ and the rightmost in the case of $G_{i,j}$, we cancel all the identity maps of chain factorizations appearing as components of the chain maps. □

With our precise knowledge of the cones Co of the chain maps $F_{i,j}$ and $G_{i,j}$, it is straightforward to prove our stabilization Theorems 1.10 and 1.11.

Proof of Theorems 1.10 and 1.11 There is a short exact of chain complexes

$$0 \rightarrow C(D_i) \xrightarrow{F_{i,j}} C(D_j) \rightarrow \text{Co}(F_{i,j}) \rightarrow 0$$

in which each map is graded of homological and quantum degree 0. This is clear in the unreduced and equivariant settings, and indeed holds also in the reduced setting since the map of rings $\mathbb{C}[x]/x^n \rightarrow \mathbb{C}$ is flat.

Induced by this short exact sequence is a long exact sequence of homology groups. Proposition 2.4 tells us that we must have

$$H^k(\text{Co}(F_{i,j})) = 0$$

for $k \leq 2i - c_- - 2$, so that the long exact sequence consists of isomorphisms $F_{i,j}$ in homological degrees $\leq 2i - c_- - 2$. This proves Theorem 1.10.

The proof of Theorem 1.11 follows the same argument. \square

Informally speaking, Theorem 1.10 tells us that we can generalize the class of objects for which there exists Khovanov–Rozansky homology to include *knots with infinite twist regions*, as discussed in the preamble to the statement of the theorem. More formally we could consider knot diagrams with extra singularities allowed. The concept is outlined in Figure 5. Investigating these stable homologies is an interesting project, but we shall not pursue it further in this paper. Infinite torus braids giving rise to projectors have been studied by Rozansky [16] in the context of colored Khovanov homology and Cautis [1] to categorify Reshetikhin–Turaev tangle invariants of type A .




Figure 5: We show an example of part of a knot diagram where we have allowed an extra type of singularity corresponding to an infinitely positively twisted pair of strands. From the results on stabilization in this paper it follows that such enhanced diagrams have well-defined homology groups. We also give an example of a new Reidemeister-type move for such diagrams: the infinitely twisted region donates a positive twist to the rest of the diagram. Clearly the homology groups will not change under this move. Further moves are possible of course, and we encourage the reader to investigate.

2.2 Some exact sequences

We derive some exact sequences of homology groups to use in proving the structural and topological theorems of Section 3.

Proposition 2.6 *Let knots K_0, K_1, K_2, \dots be given as in Figure 1. Then there is a commutative diagram in which the rows are exact, which has the following form:*

$$\begin{array}{ccccccccc}
 \longrightarrow & M^{-1} & \longrightarrow & H^0(K_0) & \longrightarrow & H^0(K_1) & \longrightarrow & M^0 & \longrightarrow & H^1(K_0) & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & & \text{id} \downarrow & & \downarrow & \\
 \longrightarrow & M^{-1} & \longrightarrow & H^2(K_1)\{2n\} & \longrightarrow & H^2(K_2)\{2n\} & \longrightarrow & M^0 & \longrightarrow & H^3(K_1)\{2n\} & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & & \text{id} \downarrow & & \downarrow & \\
 \longrightarrow & M^{-1} & \longrightarrow & H^4(K_2)\{4n\} & \longrightarrow & H^4(K_3)\{4n\} & \longrightarrow & M^0 & \longrightarrow & H^5(K_2)\{4n\} & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & & \text{id} \downarrow & & \downarrow &
 \end{array}$$

Here M is a bigraded, finitely-generated module (over \mathbb{C} or $\mathbb{C}[a]$ depending on the variant of homology chosen). Moreover, in the equivariant case, M is a torsion $\mathbb{C}[a]$ -module. All maps preserve the quantum grading.

Proof Each row of the commutative diagram comes about from a short exact sequence of chain complexes, and the maps between the rows are induced by morphisms of these short exact sequences. From Proposition 2.4 we first observe that

$$\text{Co}(F_{i,i+1}) = \text{Co}(F_{0,1})[2i]\{-2ni\}.$$

It is then straightforward to check that for $i \geq 0$, there is a commutative map of short exact sequences of chain complexes

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C(D_i) & \xrightarrow{F_{i,i+1}} & C(D_{i+1}) & \longrightarrow & \text{Co}(F_{0,1})[2i]\{-2ni\} \longrightarrow 0 \\
 & & \downarrow G_{i,i+1} & & \downarrow G_{i+1,i+2} & & \downarrow \text{id} \\
 0 & \longrightarrow & C(D_{i+1})[-2]\{2n\} & \xrightarrow{F_{i+1,i+2}} & C(D_{i+2})[-2]\{2n\} & \longrightarrow & \text{Co}(F_{0,1})[2i]\{-2ni\} \longrightarrow 0.
 \end{array}$$

So setting $M = H(\text{Co}(F_{0,1}: C(D_0) \rightarrow C(D_1)))$ we are almost done, it only remains to argue that M is finitely-generated and, in the equivariant case, torsion.

That M is finitely-generated follows from $H(K_0)$ and $H(K_1)$ being finitely-generated and the first row of the commutative diagram. In the equivariant case, suppose that M were not torsion, so that there is some i for which $\mathbb{C}[a]$ is a submodule of M^i . Taking a row low enough in the commutative diagram, we see that this would force $H^k(K_i)$

to be nontorsion for some $k > 0$ and some knot K_l , a contradiction. Hence M is torsion. □

Proposition 2.7 *Let knots K_0, K_1, K_2, \dots be given as in Figure 1. Then there is a commutative diagram in which the rows are exact, which has the following form:*

$$\begin{array}{ccccccccc}
 \longrightarrow & N^{-1} & \longrightarrow & H^0(K_0) & \longrightarrow & H^2(K_1)\{2n\} & \longrightarrow & N^0 & \longrightarrow & H^1(K_0) & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & & \text{id} \downarrow & & \downarrow & \\
 \longrightarrow & N^{-1} & \longrightarrow & H^0(K_1) & \longrightarrow & H^2(K_2)\{2n\} & \longrightarrow & N^0 & \longrightarrow & H^1(K_1) & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & & \text{id} \downarrow & & \downarrow & \\
 \longrightarrow & N^{-1} & \longrightarrow & H^0(K_2) & \longrightarrow & H^2(K_3)\{2n\} & \longrightarrow & N^0 & \longrightarrow & H^1(K_2) & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & & \text{id} \downarrow & & \downarrow &
 \end{array}$$

Here N is a bigraded, finitely-generated module (over \mathbb{C} or $\mathbb{C}[a]$ depending on the variant of homology chosen). Every map in the complex preserves the quantum grading.

Proof Setting $N = H(\text{Co}(G_{0,1}: C(D_0) \rightarrow C(D_1)))$, we consider the commutative map of short exact sequences

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & C(D_i) & \xrightarrow{G_{i,i+1}} & C(D_{i+1})[-2]\{2n\} & \longrightarrow & \text{Co}(G_{0,1}) & \longrightarrow & 0 \\
 & & \downarrow F_{i,i+1} & & \downarrow F_{i+1,i+2}[-2]\{2n\} & & \downarrow \text{id} & & \\
 0 & \longrightarrow & C(D_{i+1}) & \xrightarrow{G_{i+1,i+2}} & C(D_{i+2})[-2]\{2n\} & \longrightarrow & \text{Co}(G_{0,1}) & \longrightarrow & 0.
 \end{array}$$

The proof now follows as before. □

Remark Although we do not prove it in this paper, we believe that the results of Propositions 2.6 and 2.7 hold for standard Khovanov homology over the integers, allowing analogues of results such as those of the next section to be deduced in this setting.

3 Topological and structural results

With Proposition 2.6 in hand, we can now begin to prove Theorems 1.3, 1.5 and 1.6. We note that Propositions 2.6 and 2.7 seem to contain much of the same information from our point of view, but we suspect that there are some useful applications of Proposition 2.7 yet to be uncovered which make use of the fact that $\text{Co}(G_{0,1})$ is such a simple complex.

Proof of Theorem 1.3 Let us work in the equivariant setting. First note that the commutative diagram in Proposition 2.6 can in fact be extended arbitrarily upwards. This is because for any $l \geq 1$, we can add l negative full twists to K_0 forming $\tilde{K}_0 = K_{-l}$, and then make use of the short exact sequences for $C(\tilde{K}_j) = C(K_{j-l})$.

Now suppose we are in the situation of Theorem 1.3 where $s_n(K_{-1}) > s_n(K_0)$.

From Proposition 2.6 we see that we have the row-exact commutative diagram:

$$\begin{array}{ccccccc}
 \longrightarrow & H^0(K_{-1})\{-2n\} & \longrightarrow & H^0(K_0)\{-2n\} & \longrightarrow & M^2 & \longrightarrow \\
 & \downarrow & & \downarrow & & \downarrow & \\
 \longrightarrow & H^{2p}(K_{p-1})\{2np\} & \longrightarrow & H^{2p}(K_p)\{2np\} & \longrightarrow & M^2 & \longrightarrow
 \end{array}$$

Since the free parts of $H^0(K_{-1})$ and $H^0(K_0)$ do not lie in the same quantum degrees by hypothesis and M is torsion, this forces the map $H^0(K_0) \rightarrow M^2$ to be nonzero. By commutativity of the righthand square, this also forces $H^{2p}(K_p) \rightarrow M^2$ to be nonzero, and in particular we have $H^{2p}(K_p) \neq 0$. □

We note that with a little more work we could say exactly what quantum degrees of $H^{2p}(K)$ are nonzero, in terms of $s_n(K_{-1})$, $s_n(K_0)$ and p . Such exact information could be useful in investigating whether the s_n homomorphisms are equivalent. This precise knowledge is not necessary however to deduce Corollary 1.4, which follows immediately.

Proof of Theorem 1.5 Let us work in the equivariant setting. Suppose we have the hypotheses of Theorem 1.5. Let $p \geq 2$, and consider the following part of the commutative diagram of Proposition 2.6:

$$\begin{array}{ccccccc}
 \longrightarrow & H^{-2p-2}(K_0) & \longrightarrow & H^{-2p-2}(K_1) & \xrightarrow{\psi} & M^{2p-2} & \longrightarrow \\
 & \downarrow & & \downarrow & & \downarrow & \\
 \longrightarrow & H^0(K_{p-1})\{2n(p-1)\} & \longrightarrow & H^0(K_p)\{2n(p-1)\} & \xrightarrow{\varphi} & M^{2p-2} & \longrightarrow
 \end{array}$$

Observe that since by hypothesis we have that K_0 is the unknot, then we have $H^{-2p-2}(K_0) = H^{-2p-1}(K_0) = 0$ so that ψ is an isomorphism. Then the commutativity of the square involving both ψ and φ tells us that φ restricted to the torsion part of $H^0(K_p)$ is a surjection. Therefore there exists a decomposition $H^0(K_p) = \text{Fr} \oplus \text{tor}$ into free and torsion $\mathbb{C}[a]$ -modules such that $\varphi|_{\text{Fr}} = 0$. But if $s_n(K_p) \neq s_n(K_{p-1})$ then we must have $\varphi|_{\text{Fr}} \neq 0$, hence a contradiction. □

Proof of Theorem 1.6 Suppose now that we have the hypotheses of Theorem 1.6.

First of all we would like to see that $M = \Delta$, the torsion part of $H(K_1)$. This follows directly from the first row of the commutative diagram in Proposition 2.6 and the fact that the map $H^0(K_0) \rightarrow H^0(K_1)$ is onto the free part of $H^0(K_1)$. Indeed, if this map were not, we would either have a nontorsion part of M^{-1} or we would have $s_n(K_1) < 0$.

Now let $p \geq 2$ and consider the following two commutative diagrams with exact rows:

$$\begin{array}{ccccccc}
 \longrightarrow & H^{i-2(p-1)}(K_1) & \longrightarrow & M^{i-2p+2} & \xrightarrow{\varphi} & H^{i-2(p-1)+1}(K_0) & \longrightarrow \\
 & \downarrow & & \text{id} \downarrow & & \downarrow & \\
 \longrightarrow & H^i(K_p)\{2n(p-1)\} & \longrightarrow & M^{i-2p+2} & \xrightarrow{\psi} & H^{i+1}(K_{p-1})\{2n(p-1)\} & \longrightarrow , \\
 \\
 \longrightarrow & M^{i-2p+1} & \xrightarrow{\varphi'} & H^{i-2(p-1)}(K_0) & \longrightarrow & H^{i-2(p-1)}(K_1) & \longrightarrow \\
 & \text{id} \downarrow & & \downarrow & & \downarrow & \\
 \longrightarrow & M^{i-2p+1} & \xrightarrow{\psi'} & H^i(K_{p-1})\{2n(p-1)\} & \longrightarrow & H^i(K_p)\{2n(p-1)\} & \longrightarrow .
 \end{array}$$

From the first diagram, observe that $\varphi = 0$ since $H(K_0)$ is nontorsion. This implies that $\psi = 0$ by commutativity of the rightmost square. For the same reason in the second diagram we see $\varphi' = 0$, which implies that $\psi' = 0$ by the commutativity of the leftmost square. This means that each row gives rise to short exact sequences

$$0 \rightarrow H^i(K_{p-1})\{2n(p-1)\} \rightarrow H^i(K_p)\{2n(p-1)\} \rightarrow M^{i-2p+2} \rightarrow 0.$$

With this in hand, to prove the theorem it remains to see that every such short exact sequence splits to give isomorphisms

$$H^i(K_p)\{2n(p-1)\} = H^i(K_{p-1})\{2n(p-1)\} \oplus M^{i-2p+2}.$$

A splitting map is found by running anticlockwise around the square

$$\begin{array}{ccc}
 H^{i-2(p-1)}(K_1) & \longrightarrow & M^{i-2p+2} \\
 \downarrow & & \text{id} \downarrow \\
 H^i(K_p)\{2n(p-1)\} & \longrightarrow & M^{i-2p+2},
 \end{array}$$

from M^{i-2p+2} to $H^i(K_p)\{2n(p-1)\}$, which is possible since the top row of the square is an isomorphism when restricted to the torsion part of $H^{i-2(p-1)}(K_1)$. \square

Proof of Theorem 1.7 We can copy the proof of Theorem 1.6 here. In fact, this situation is simpler since there is no torsion hence every short exact sequence splits. The one almost delicate point is to deduce that the map appearing in the top row commutative diagram in Proposition 2.6,

$$F_{0,1}: H^0(U) = H^0(K_0) \rightarrow H^0(K_1),$$

is an injection. We know that it is an injection equivariantly and furthermore we have a description of the chain-homotopy type of the equivariant complex given in the discussion of Section 1.2. So it follows that $F_{0,1}$ is an injection in the unreduced case which is obtained by setting $a = 0$ in the equivariant chain complexes. The reduced case then follows from $F_{0,1}$ being an injection in the unreduced case and the generalized universal coefficients theorem for principal ideal domains. \square

Earlier we promised a discussion of the case when $s_n(K_1) \neq 0$. Notice that in this case our argument in the proof of Theorem 1.6 goes through as before for all $H^i(K_p)\{2n(p-1)\}$ except when $2p-i=2$. Hence we can determine the homology groups of K_p in terms of $H(K_{p-1})$ and p except for $H^{2p-2}(K_p)$. To fix the remaining homology group it suffices to know the image of the map $H^0(K_0) \rightarrow H^{2p-2}(K_{p-1})\{2n(p-1)\}$. We do not give a proof of this fact since it is not needed for our main application of Theorem 1.6.

To begin our proof of Theorem 1.1, we first collect a few results from the literature on hyperbolic 3–manifolds. We state the first theorem not as strongly as Thurston proved it, but strongly enough for us to use.

Theorem 3.1 (Hyperbolic Dehn surgery [19]) *Let M be a cusped hyperbolic 3–manifold with a distinguished cusp. We write $M(1/p)$ for the result of filling the distinguished cusp with filling coefficient $1/p$. Then $M(1/p)$ is hyperbolic except for a finite set of filling slopes and $M(1/p)$ converges to M in the geometric topology as $p \rightarrow \infty$.*

We shall also need a result of Kawauchi’s concerning special knots K^* in S^3 .

Theorem 3.2 (Kawauchi [4]) *For every $m > 1$ there exists an $(m + 1)$ –component link*

$$U \cup U_1 \cup U_2 \cup \dots \cup U_m \subset S^3,$$

where U is the unknot and $U_1 \cup U_2 \cup \dots \cup U_m$ is the m –component unlink, satisfying the following properties.

- (1) Each U_i bounds a disc intersecting U in two points with signed count 0.
- (2) For $i \neq j$, the link $U \cup U_i$ is distinct from the link $U \cup U_j$.
- (3) For any i , the result of $+1$ -surgery on U_i turns U into a smoothly slice knot K^* , which is independent of i .
- (4) Define the tangles $\overline{\text{Tan}}_i$ as in Figure 6. Each tangle $\overline{\text{Tan}}_i$ is hyperbolic, as is the branched double cover of each $\overline{\text{Tan}}_i$.




Figure 6: This diagram accompanies the statement of Theorem 3.2. We have drawn a tangle in a small 3-ball $\text{Tan}_i \subset B^3$ which is a subtangle of the link $(U \cup U_i) \subset S^3$. It consists of all of U_i and two strands of U which intersect a disc bounded by U_i in two points, with signed count 0. (The rest of U has been drawn schematically as a dotted line). We denote by $\overline{\text{Tan}}_i$ the complement to this tangle so that $\text{Tan}_i \cup_{\partial} \overline{\text{Tan}}_i = U \cup U_i$.

Proof of Theorem 1.1 Consider Figure 6. Tan_i is an example of a *simple* tangle (in other words prime and atoroidal). Furthermore, we know by item (4) of Theorem 3.2 that $\overline{\text{Tan}}_i$ (the complement of Tan_i) is hyperbolic.

We are in the situation where we can apply a result of Soma [18, Lemma 2]. This tells us that if we glue back Tan_i to $\overline{\text{Tan}}_i$, then the result (which is $\text{Tan}_i \cup_{\partial} \overline{\text{Tan}}_i = U \cup U_i$) is a hyperbolic link.

We now write K_N^i for the result of doing $(1/N)$ -surgery on U_i to the knot U , so that $K_0^i = U$ for each i . By item (3) of Theorem 3.2, we see that K_1^i is the knot K^* for each $i = 1, 2, \dots, m$.

Since the complement of $U \cup U_i$ is atoroidal for each i , Theorem 3.1 tells us that the complement of K_N^i is hyperbolic for large enough N and that these complements converge in the geometric topology to the complement of $U \cup U_i$ as $N \rightarrow \infty$. Since the meridians of the K_N^i converge to the meridian to U , the sequence of knots K_N^i determines the link complement to $U \cup U_i$ as well as the meridional curve to U . By filling along the meridian and taking U isotopic to any longitude relative to the meridian, we see this determines U inside the solid torus complement to U_i . Since

there is only one way to fill the boundary of this solid torus to get $U = K_0^i$ unknotted inside S^3 , we have determined the whole link $U \cup U_i$.

Hence there exists an N such that the complement to K_N^i is not diffeomorphic to the complement to K_N^j whenever $i \neq j$. Since the knot complement determines the knot, we know that for this N we have $K_N^i \neq K_N^j$ whenever $i \neq j$. This set $\{K_N^1, K_N^2, \dots, K_N^m\}$ will be the m distinct knots we are required to exhibit.

Because K^* is slice we have $s_n(K_1^i = K^*) = 0$ for all $i = 1, 2, \dots, m$. This means that we can apply Theorem 1.6 to see that $H(K_N^i) = H(K_N^j)$ for all $1 \leq i, j \leq m$.

It remains to see that each K_N^i is prime and not 2-bridge. Primeness follows from the hyperbolicity of K_N^i .

The branched double cover of K_N^i is a Dehn filling of the branched double cover of $\overline{\text{Tan}_i}$, with filling slope determined by N . Again, Theorem 3.1 implies that for N large enough, the branched double cover of K_N^i is hyperbolic. We know that branched double covers of 2-bridge knots are lens spaces, which are not hyperbolic. Hence K_N^i is not 2-bridge. □




Figure 7: Here is an example of a Brunnian pure braid: a pure braid with the property that the removal of any strand results in a trivial braid.

3.1 An example of the construction of a knot pair with isomorphic knot homologies

Kawauchi used the theory of *almost identical imitation* to create knots K^* with multiple unknotting sites [4]. In the proof of Theorem 1.1 we used these knots K^* in an essential way to create distinct knots with isomorphic Khovanov–Rozansky knot homologies. If we wished to draw a diagram of such knots it would be necessary to understand in detail the theory of almost identical imitation. However, if one is prepared to work on a more *ad hoc* basis then it is easy to create examples of knots with isomorphic knot homologies.

One such *ad hoc* construction is based on pure Brunnian braids (pure braids that become equivalent to a trivial braid when any strand is removed). We have drawn an example of such a braid (on three strands) in Figure 7.




Figure 8: Here we show a tangle determined by the braid drawn in Figure 7. There are three boundary components to this tangle, each will be filled by some tangle corresponding to the chain complex T_i as in Figure 3.

From the braid drawn in Figure 7 we obtain the tangle drawn in Figure 8. This tangle can be completed to a knot by filling the slots X, Y, Z with other tangles. We now abuse notation by referring to the tangle corresponding to the chain complex T_i itself

by T_i . We denote by K_i^X the knot obtained by filling X with T_i , Y with T_1 and Z with T_{-1} , and denote by K_i^Y the knot obtained by filling X with T_1 , Y with T_i and Z with T_{-1} .

Note that $K_0^X = K_0^Y = U$, the unknot and that $K_1^X = K_1^Y$. Furthermore since K_1^X can be transformed into the unknot both by a positive-to-negative crossing change (in place X , say) and by a negative-to-positive crossing change (in place Z), we must have $s_n(K_1^X) = s_n(K_1^Y) = 0$.

Hence it follows from Theorem 1.6 that K_i^X and K_i^Y have isomorphic homologies for all $i \geq 2$.

One can check that $K_2^X \neq K_2^Y$ using *SnapPea*. In fact, they have different hyperbolic volumes so they are not even mutant by a result of Ruberman in [17].

3.2 Pairs of mutant knots with isomorphic knot homologies

The Conway and the Kinoshita–Terasaka (KT) knots are the first (measured by crossing number) example of a pair of mutant knots. In [20], Mackaay and Vaz use techniques given by Rasmussen in [14] in order to compute that all reduced Khovanov–Rozansky homologies of the Conway and the KT knots agree. Since it is easily observed that the KT knot and the Conway knot have unknotting number equal to 1, we can build upon this computation and give an infinite number of mutant pairs.




Figure 9: This diagram shows two families of knots K_i^{KT} and K_i^C . On both sides the unknot U occurs when we put the tangle T_0 where indicated $K_0^{KT} = K_0^C = U$. When we add the tangle T_1 we get the Kinoshita–Terasaka knot K_1^{KT} on the left and the Conway knot K_1^C on the right.

Proof of Theorem 1.2 We work with reduced homology. Consider the two families of knots K_i^C and K_i^{KT} shown in Figure 9. Since the reduced homologies agree, $H(K_1^C) = H(K_1^{KT})$, $K_0^C = K_0^{KT} = U$ and $s_n(K_1^C) = s_n(K_1^{KT}) = 0$, we can apply Theorem 1.6 to see that we have isomorphic homology groups $H(K_i^C) = H(K_i^{KT})$ for all $i \geq 2$. Thurston's Theorem 3.1 tells us that $K_i^C \neq K_i^{KT}$ for large enough i , and since each is hyperbolic each must be prime. \square

References

- [1] **S Cautis**, *Clasp technology to knot homology via the affine Grassmannian* arXiv: 1207.2074
- [2] **B Gornik**, *Note on Khovanov link cohomology* arXiv:math.QA/0402266
- [3] **M Hedden, P Ording**, *The Ozsváth–Szabó and Rasmussen concordance invariants are not equal*, Amer. J. Math. 130 (2008) 441–453 MR2405163
- [4] **A Kawachi**, *Introduction to topological imitations of $(3, 1)$ -dimensional manifold pairs*, from: “Topics in knot theory”, (ME Bozhüyük, editor), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 399, Kluwer Acad. Publ., Dordrecht (1993) 69–83 MR1257906
- [5] **M Khovanov, L Rozansky**, *Virtual crossings, convolutions and a categorification of the $SO(2N)$ Kauffman polynomial*, J. Gökova Geom. Topol. GGT 1 (2007) 116–214 MR2386537
- [6] **M Khovanov, L Rozansky**, *Matrix factorizations and link homology*, Fund. Math. 199 (2008) 1–91 MR2391017
- [7] **D Krasner**, *A computation in Khovanov–Rozansky homology*, Fund. Math. 203 (2009) 75–95 MR2491784
- [8] **D Krasner**, *Equivariant $\mathfrak{sl}(n)$ -link homology*, Algebr. Geom. Topol. 10 (2010) 1–32 MR2580427
- [9] **P B Kronheimer, T S Mrowka**, *Khovanov homology is an unknot-detector*, Publ. Math. Inst. Hautes Études Sci. (2011) 97–208 MR2805599
- [10] **E S Lee**, *An endomorphism of the Khovanov invariant*, Adv. Math. 197 (2005) 554–586 MR2173845
- [11] **C Livingston**, *Computations of the Ozsváth–Szabó knot concordance invariant*, Geom. Topol. 8 (2004) 735–742 MR2057779
- [12] **A Lobb**, *A slice genus lower bound from $\mathfrak{sl}(n)$ Khovanov–Rozansky homology*, Adv. Math. 222 (2009) 1220–1276 MR2554935
- [13] **A Lobb**, *A note on Gornik's perturbation of Khovanov–Rozansky homology*, Algebr. Geom. Topol. 12 (2012) 293–305 MR2916277

- [14] **J Rasmussen**, *Some differentials on Khovanov–Rozansky homology* arXiv math/0607544
- [15] **J Rasmussen**, *Khovanov homology and the slice genus*, Invent. Math. 182 (2010) 419–447 MR2729272
- [16] **L Rozansky**, *An infinite torus braid yields a categorified Jones–Wenzl projector* arXiv:1005.3266
- [17] **D Ruberman**, *Mutation and volumes of knots in S^3* , Invent. Math. 90 (1987) 189–215 MR906585
- [18] **T Soma**, *Simple links and tangles*, Tokyo J. Math. 6 (1983) 65–73 MR707839
- [19] **W P Thurston**, *The geometry and topology of three-manifolds*, Princeton Univ. Math. Dept. Lecture Notes (1979) Available at <http://msri.org/publications/books/gt3m/>
- [20] **P Vaz**, *The reduced HOMFLY-PT homology for the Conway and the Kinoshita–Terasaka knots* arXiv:0812.1957
- [21] **H Wu**, *On the quantum filtration of the Khovanov–Rozansky cohomology*, Adv. Math. 221 (2009) 54–139 MR2509322

*Department of Mathematical Sciences, Durham University
Science Labs, South Road, Durham DH1 3LE, UK*

`andrew.lobb@durham.ac.uk`

Proposed: Tom Mrowka
Seconded: Richard Thomas, Peter Teichner

Received: 3 May 2011
Revised: 4 May 2011

