

Open book foliation

TETSUYA ITO
KEIKO KAWAMURO

We study open book foliations on surfaces in 3–manifolds and give applications to contact geometry of dimension 3. We prove a braid-theoretic formula for the self-linking number of transverse links, which reveals an unexpected connection with the Johnson–Morita homomorphism in mapping class group theory. We also give an alternative combinatorial proof of the Bennequin–Eliashberg inequality.

57M27; 57M50, 57R17, 53D35

1 Introduction

In his seminal work [1], Bennequin shows that there is an “exotic” contact structure, ξ_{ot} , on S^3 that is homotopic to the standard contact structure, ξ_{std} , as a 2–plane field but *not* contactomorphic to ξ_{std} . In other words, ξ_{std} is tight whereas ξ_{ot} is overtwisted in contemporary terminology. In order to distinguish these contact structures he studies closed braids and characteristic foliations on their Seifert surfaces induced by the contact structures. Since then, Bennequin’s method has been developed in two directions.

One direction is the theory of characteristic foliations and convex surfaces: Eliashberg uses characteristic foliations to show the Bennequin–Eliashberg inequality for tight contact 3–manifolds and generalizes the Bennequin inequality for the tight contact 3–sphere in [16]. Characteristic foliations also play important roles in Eliashberg’s classification of overtwisted contact structures [15]. In [24], Giroux extends characteristic foliation theory and initiates convex surface theory. This gives us cut-and-paste techniques to study contact structures and to classify tight contact structures for various 3–manifolds. See also Honda’s work [28] on convex surface theory.

The other direction is the theory of braid foliations studied in a series of papers by Birman and Menasco [5; 6; 7; 8; 9; 10; 11]. One of its highest achievements is the “Markov theorem without stabilization,” which states that given two closed braid representatives of any link in \mathbb{R}^3 can be transformed to each other in a very controlled manner [11]. Moreover, Birman and Menasco apply the braid foliation to contact geometry and construct examples of transversely non-simple knots in the standard

contact 3–sphere: transverse knots of the same topological type and the same self-linking number that are not transversely isotopic [12]. Their examples are closed 3–braids related by negative flypes. Analysis of a sequence of braid moves that relates one to the other (a Markov tower) reveals that the two closed braids represent distinct transverse links. See also Birman and Finkelstein [3] for a concise survey article.

We establish a foundation for *open book foliations*, which generalize braid foliations.

Our starting point is a classical theorem that dates back to Alexander: every closed, oriented 3–manifold admits an open book decomposition. An open book decomposition naturally induces a singular foliation on an embedded surface. When the foliation satisfies certain conditions, we call it an open book foliation on the surface. As shown in Theorem 2.5, any embedded surface can be isotoped to admit an open book foliation.

The idea of open book foliations has existed for some time. The project of this paper started from a conversation between John Etnyre and the authors at the conference “Braids in Seville” in 2011. Etnyre pointed out that Bennequin’s work [1] suggests that characteristic foliations and open book foliations are essentially the “same”. Also, he and Ko Honda had discussed generalizations of braid foliations. In addition, readers may find a preliminary step toward open book foliations in Pavelescu’s thesis [39].

However, a foundation for open book foliations in the general setting has not been fully developed in the literature, and an open book foliation has often been regarded as a special kind of characteristic foliation in contact geometry. In this paper we develop the basics of open book foliations in a topological and combinatorial way. It is important that open book foliation theory is independent of the theory of characteristic foliations. In fact in Remark 2.22 we list items that highlight differences between the two foliations. The most notable difference is that open book foliations are more “rigid” than characteristic foliations. For instance, the Giroux cancellation lemma [25] for characteristic foliations does not apply to open book foliations. However the two foliations have similar appearances, as Etnyre pointed out to us. We prove *the structural stability theorem* (Theorem 2.21), which states that the two foliations can be topologically conjugate to each other under certain conditions.

Hence, via the Giroux–correspondence [26], open book foliation theory gives rise to a new technique to analyze general contact 3–manifolds, just like Bennequin’s foliations and Birman and Menasco’s braid foliations have been used to study the standard tight contact 3–sphere.

Our first application of open book foliations to contact geometry is a self-linking number formula for an n –stranded braid, b , with respect to an open book (S, ϕ) :

$$\text{sl}(\widehat{b}, [\Sigma]) = -n + \widehat{\text{exp}}(b) - \phi_*(a) \cdot [b] + c([\phi], a)$$

The precise statement and definitions can be found in Theorem 3.10. Interestingly the function $c([\phi], a)$ in the formula reveals an unexpected relationship between the self-linking number, an invariant in contact geometry, and the Johnson–Morita homomorphism in mapping class group theory. We discuss this in detail in Section 3.5.

Our formula generalizes Bennequin’s self-linking number formula [1] for a braid in the open book (D^2, id) that is a usual closed braid in \mathbb{R}^3 around the z -axis. Bennequin’s formula is

$$\text{sl}(b) = -n + \exp(b),$$

where $\exp(b)$ (with no “hat” over \exp) is the exponent sum of a braid word representing b . When the page surface S is an annulus, Kawamuro and Pavelescu [33] show that

$$\text{sl}(b, [\Sigma]) = -n + \exp(b) - \phi_*(a) \cdot [b].$$

Moreover if S is planar, Kawamuro [32] shows that

$$\text{sl}(b, [\Sigma]) = -n + \exp(b) - \phi_*(a) \cdot [b] + c'(\phi, a),$$

where the function c' is a part of the function c and the gap of c and c' is essentially the Johnson–Morita homomorphism mentioned above. We can see that the formula gets more complicated as the topology of S gets complicated.

The self-linking number is not merely an invariant of knots and links in contact manifolds. By the following celebrated Bennequin–Eliashberg inequality [16], one can use the self-linking number to determine tightness or overtwistedness of a given contact structure:

Theorem 4.3 [16] *If a contact 3-manifold (M, ξ) is tight, then for any null-homologous transverse link L and its Seifert surface Σ , we have*

$$\text{sl}(L, [\Sigma]) \leq -\chi(\Sigma).$$

Our second application of open book foliations to contact geometry is to give an alternative combinatorial proof to the above Bennequin–Eliashberg inequality. Because of its rigidity, an open book foliation is effective at visualizing or constructing surfaces like overtwisted discs. In fact we define a *transverse overtwisted disc* (Definition 4.1), a notion corresponding to an overtwisted disc in contact geometry, and we use it to prove the Bennequin–Eliashberg inequality.

1.1 Origins of open book foliation

In this section we briefly review braid foliations and characteristic foliations. We generalize braid foliations to open book foliations. On the other hand, many applications of open book foliations are derived from problems in characteristic foliation theory.

1.1.1 Braid foliations In Birman and Menasco's braid foliation theory [5; 6; 7; 8; 9; 10; 11], braids are geometric objects. Let A be an oriented unknot in S^3 . We regard S^3 as $\mathbb{R}^3 \cup \{\infty\}$ and identify A with the union of the z -axis and the point $\{\infty\}$. As is well-known, A is a fibered knot. With the cylindrical coordinates (r, θ, z) of \mathbb{R}^3 , the fibration $\pi: S^3 \setminus A \rightarrow S^1$ is given by the projection $(r, \theta, z) \mapsto \theta$. An oriented link $L \subset S^3$ is called a *closed braid* with respect to A (and π) if L is disjoint from A and positively transverse to each fiber $S_\theta = \pi^{-1}(\theta)$. In other words, L winds around the z -axis in the positive direction.

Consider an incompressible Seifert surface F of a braid L , or an essential closed surface $F \subset S^3 \setminus L$. The intersection of F and the fibers $\{S_\theta \mid \theta \in S^1\}$ induces a singular foliation \mathcal{F} on F . We can put F in a position so that \mathcal{F} satisfies the following conditions:

- (i) The z -axis pierces F transversely in finitely many points, around which the foliation \mathcal{F} is radial.
- (ii) The leaves of \mathcal{F} along ∂F are transverse to ∂F .
- (iii) All but finitely many fibers S_θ meet F transversely. Each exceptional fiber is tangent to F at a single point.
- (iv) All the tangencies of F and fibers are saddles.

This \mathcal{F} is called a *braid foliation* on the surface F . (Later in Section 2.1 we borrow Birman and Menasco's axioms (i)–(iv) to define open book foliations.)

The foliation \mathcal{F} encodes both topological and algebraic information of the closed braid L . For example, let L be the closure of the braid word σ_1 in the Artin braid group B_2 and F its Bennequin surface consisting of two discs and one positively twisted band. The surface F and its braid foliation \mathcal{F} are depicted in Figure 1(a). (The meaning of \oplus will be made clear in Section 2.1.1.) We collapse the twisted band and the upper disc to get the trivial braid as in Figure 1(b). Algebraically this corresponds to *destabilization* of σ_1 and the topology of \mathcal{F} indicates that L is destabilizable.

In general if \mathcal{F} can be “simplified” then L is also “simplified”. Moreover, as the above example suggests, a simplification of \mathcal{F} can be understood as a certain braid operation. Therefore, by studying \mathcal{F} one may find a sequence of braid operations to get the “simplest” braid representative of L .

Braid foliations have numerous applications to study of knots and links in S^3 [4; 5; 6; 7; 8; 9; 10; 11]. Moreover, via the correspondence [1] between the transverse links in the standard contact S^3 and the closed braids around the z -axis, braid foliations are used to solve problems in contact geometry, in particular detecting transversely




Figure 1: Example: Braid foliation and destabilization

non-simple links [12; 13; 14; 34]. (Here a topological link type \mathcal{L} is called *transversely simple* if the transverse link representatives of \mathcal{L} can be completely classified by an invariant called the self-linking number).

In [10], Birman and Menasco study the set of 3–braids and prove that two closed 3–braids representing topologically the same link are related to each other by the so-called *flype* move. This is the key to their construction of transversely non-simple 3–braid links [12]. In [8] they prove that every closed braid representative of the unknot can be deformed into the one-stranded braid by a sequence of exchange moves and destabilizations. Based on this, Birman and Wrinkle [14] give an alternative topological proof, first proven by Eliashberg and Fraser [17], that the unknot in (S^3, ξ_{std}) is transversely simple.

It should be pointed out that the braid foliation is not too difficult to see or illustrate once we understand how a surface is embedded. This contrasts strikingly with the flexibility of the characteristic foliations, which we describe next.

1.1.2 Characteristic foliations Let (M, ξ) be a closed contact 3–manifold. Let $F \subset M$ be an oriented embedded surface, usually either closed or with Legendrian boundary. (A convex surface with transverse boundary is established by Etnyre and Van Horn-Morris [22, Section 2].) Integrating the vector field $\xi \cap TF$ on F we get

a singular foliation $\mathcal{F}_\xi(F)$ on F called the *characteristic foliation*. If two contact structures induce the same characteristic foliation on F then they are isotopic near F .

A surface F is called *convex* if there exists a vector field v whose flow preserves ξ and is transverse to F . The *dividing set* [24] on a convex surface F is a multi-curve defined by $\{p \in F \mid v_p \subset \xi_p\}$. Giroux's flexibility theorem [24; 28] states that it is the isotopy type of a dividing set (not an individual characteristic foliation compatible with the dividing set) that encodes information of the contact structure near F . If two contact structures induce isotopic dividing sets on F then they are isotopic near F .

In [28], Honda introduces *bypass attachment*, which allows us to modify dividing sets in controlled manner. With careful examination of dividing sets, one can apply topological techniques such as gluing and cutting contact 3-manifolds along convex surfaces. This leads to various results in contact geometry. For example, Etnyre and Honda prove the non-existence of tight contact structures on a Poincaré homology sphere [19]. They also prove transverse non-simplicity of the $(2, 3)$ -cable of the $(2, 3)$ -torus knot by classifying its Legendrian representatives [20]. Later, LaFountain and Menasco [34] establish Legendrian and transversal “Markov theorem without stabilization” for the above knot by using both braid foliation and convex surface techniques.

In practice, except for certain simple cases, it is not very easy to grasp the entire picture of a characteristic foliation and a dividing set. It is also not very clear how they change under isotopies of surfaces. In contrast, the structural stability theorem that we prove in Section 2.2 allows us to visualize a characteristic foliation through an open book foliation.

2 Basics of open book foliation

In this section we define open book foliations and develop basic machinery by applying (sometimes with modifications) existing notions in braid foliation theory.

Hence most of our definitions in this section can be found in Birman and Menasco's papers [5]–[12]. We also cite Birman and Finkelstein's paper [3] because it is a concise survey of braid foliation theory and conveniently contains all the basic notions we want to borrow.

2.1 Definition of open book foliation

An *open book* (S, ϕ) is a compact surface S with non-empty boundary ∂S along with a diffeomorphism $\phi \in \text{Aut}(S, \partial S)$ fixing the boundary pointwise. Given an open book

(S, ϕ) we define a closed oriented 3-manifold $M = M_{(S, \phi)}$ by

$$M_{(S, \phi)} = M_\phi \cup \left(\coprod_{|\partial S|} D^2 \times S^1 \right),$$

where M_ϕ denotes the mapping torus $S \times [0, 1]/(x, 1) \sim (\phi(x), 0)$ and the solid tori are attached so that for each point $p \in \partial S$ the circle $\{p\} \times S^1 \subset \partial M_\phi$ bounds a meridian disc of $D^2 \times S^1$. If a closed oriented manifold M is homeomorphic to $M_{(S, \phi)}$, we say that (S, ϕ) is an *open book decomposition* of the manifold M . For example, $M_{(D^2, \text{id})} \cong S^3$. The union of core circles of the attached solid tori, B , is called the *binding* of the open book. Let $\pi: M \setminus B \rightarrow S^1 = \mathbb{R}/\mathbb{Z}$ denote the fibration. The fibers $\pi^{-1}(t) =: S_t$ where $t \in [0, 1)$ are called the *pages* of the open book.

We say that an oriented link L in $M_{(S, \phi)}$ is in *braid position* with respect to the open book (S, ϕ) if L is disjoint from the binding and positively transverses each page S_t . This generalizes the familiar concept of braid position for $M_{(D^2, \text{id})} \cong S^3$.

Let F be an oriented, connected, compact surface smoothly embedded in $M_{(S, \phi)}$ whose boundary ∂F (if it exists) is in braid position with respect to the open book (S, ϕ) .

Consider the singular foliation $\mathcal{F} = \mathcal{F}(F)$ on F induced by the pages $\{S_t \mid t \in S^1\}$. That is, \mathcal{F} is obtained by integrating the singular vector field $\{T_p S_t \cap T_p F\}_{p \in F}$ on F . We call each connected component of the integral curves a *leaf*. We may regard the leaves as $F \cap S_t$. By standard general position arguments (see Hirsch [27] for example) the surface F can be perturbed while the braid isotopy class of ∂F is fixed (if ∂F is non-empty) so that F satisfies the same conditions in [7, page 23], namely:

- (\mathcal{F} i) The binding B pierces the surface F transversely in finitely many points. Moreover, $p \in B \cap F$ if and only if there exists a disc neighborhood $N_p \subset \text{Int}(F)$ of p on which the foliation $\mathcal{F}(N_p)$ is radial with the node p (see the top sketches in Figure 2). We call the singularity p an *elliptic point*.
- (\mathcal{F} ii) The leaves of \mathcal{F} along ∂F are transverse to ∂F .
- (\mathcal{F} iii) All but finitely many fibers S_t intersect F transversely. Each exceptional fiber is tangent to $\text{Int}(F)$ at a single point. In particular, \mathcal{F} has no saddle-saddle connections.
- (\mathcal{F} iv') The type of a tangency in (\mathcal{F} iii) is saddle or local extremum.

Definition 2.1 [7, page 23] We say that a page S_t is *regular* if S_t intersects F transversely and it is *singular* otherwise. Similarly, a leaf l of \mathcal{F} is called *regular* if l does not contain a tangency point.

The arguments in [3, page 272–273] imply the following:

Proposition 2.2 *Since ∂F is in braid position (if ∂F is non-empty), no regular leaf of $\mathcal{F}(F)$ has both of its endpoints on ∂F . Hence, the regular leaves of \mathcal{F} are classified into the following three types:*

a–arc: An arc where one of its endpoints lies on B and the other lies on ∂F

b–arc: An arc whose endpoints both lie on B

c–circle: A simple closed curve

Definition 2.3 We say that the singular foliation $\mathcal{F}(F)$ is an *open book foliation* if the above conditions (\mathcal{F} i), (\mathcal{F} ii), (\mathcal{F} iii) and the following condition (\mathcal{F} iv), which is stronger than (\mathcal{F} iv'), are satisfied, and we denote it by $\mathcal{F}_{\text{ob}}(F)$.

(\mathcal{F} iv) All the tangencies of F and fibers are of saddle type (see the bottom sketches of Figure 2). We call them *hyperbolic points*.

Remark 2.4 Here we list differences between the braid foliation and the open book foliation.

- (1) For braid foliations the ambient manifold M is S^3 , whereas for open book foliations M can be any closed oriented 3–manifold.
- (2) In braid foliation theory each regular leaf $l \subset S_t$ is required to be *essential* in $S_t \setminus (S_t \cap \partial F)$ [3, Theorem 1.1]. In open book foliation theory we relax this restriction, so a regular leaf can be inessential, ie F can be compressible.

We do this for the following reasons: First, we prefer to establish the basics of open book foliations under less restrictive conditions. Second, characteristic foliations, which share common properties with open book foliations, also contain inessential circles. Third, in some cases it is more convenient and natural to allow inessential leaves: For example, as we will see in Proposition 2.6, one can remove c –circles at the cost of introducing inessential leaves. (In [29] we study open book foliations where all of the b –arc leaves are essential, and give several applications to the topology of 3–manifolds.)

The open book foliation is intrinsic in the following sense:

Theorem 2.5 *If (\mathcal{F} i), (\mathcal{F} ii), (\mathcal{F} iii), (\mathcal{F} iv') are satisfied then (\mathcal{F} iv) holds. Namely, with an ambient isotopy (that fixes ∂F if it exists), every surface F admits an open book foliation $\mathcal{F}_{\text{ob}}(F)$.*

We prove Theorem 2.5 in Section 2.1.2. At a glance, this theorem is similar to [7, Lemma 2]. However we allow our pages to be of type $S_{g,r}$ (rather than D^2) and moreover we allow F to be compressible. As a result Birman and Menasco's proof (which is a refined argument of Bennequin's [1] with much more details) does not apply. For the same reason, Roussarie and Thurston's argument [41] does not work either.

As a byproduct of the proof of Theorem 2.5 we obtain:

Proposition 2.6 *Given an open book foliation $\mathcal{F}_{\text{ob}}(F)$, we can perturb F (fixing ∂F if it exists) so that the new $\mathcal{F}_{\text{ob}}(F)$ contains no c -circles.*

We prove Proposition 2.6 also in Section 2.1.2. This is a useful proposition that allows us to convert an open book foliation into Morse–Smale type. In this paper we use Proposition 2.6 many times, including in a new proof of the Bennequin–Eliashberg inequality.

2.1.1 Signs of singularities, describing arcs and orientation of leaves

Definition 2.7 [1, page 19; 3, page 280] We say that an elliptic singularity p is *positive (negative)* if the binding B is positively (negatively) transverse to F at p . The sign of the hyperbolic singularity p is *positive (negative)* whether the orientation of the tangent plane $T_p F$ does (does not) coincide with the orientation of $T_p S_t$.

See Figure 2, where we describe an elliptic point by a hollowed circle with its sign inside, a hyperbolic point by a black dot with the sign indicated nearby, and positive normals \vec{n}_F to F by dashed arrows.

With this definition, we observe that:

Claim 2.8 *The elliptic point at the end of every a -arc is positive, and the endpoints of every b -arc have opposite signs.*

Definition 2.9 (Describing arc) Consider a saddle shape subsurface of F whose leaves l_1 and l_2 (possibly $l_1 = l_2$) as in Figure 3 are sitting on a page S_t . As t increases (the page moves up) the leaves converge along a properly embedded arc $\gamma \subset S_t$ (dashed in Figure 3) joining l_1 and l_2 and switch configuration. See the passage in Figure 3.

We call γ the *describing arc* of the hyperbolic singularity. Up to isotopy, γ is uniquely determined. We also often put the sign of a hyperbolic point near its describing arc (see Figure 11).




Figure 2: Signs of singularities and normal vectors \vec{n}_F




Figure 3: The describing arc (dashed) for a hyperbolic singularity

Definition 2.10 We denote the number of positive (resp. negative) elliptic points of $\mathcal{F}_{\text{ob}}(F)$ by $e_+ = e_+(\mathcal{F}_{\text{ob}}(F))$ (resp. $e_- = e_-(\mathcal{F}_{\text{ob}}(F))$). Similarly, the number of positive (resp. negative) hyperbolic points is denoted by $h_+ = h_+(\mathcal{F}_{\text{ob}}(F))$ (resp. $h_- = h_-(\mathcal{F}_{\text{ob}}(F))$).

Proposition 2.11 The Euler characteristic of the surface F is

$$\chi(F) = (e_+ + e_-) - (h_+ + h_-).$$

To prove Proposition 2.11, we define *orientations* of leaves:

Definition 2.12 (Orientation of leaves) Both the surface F and the ambient manifold M are oriented so that the positive normal \vec{n}_F of F (in this paper we indicate \vec{n}_F by

dashed arrows like in Figure 2) is canonically defined. We orient each leaf of $\mathcal{F}_{\text{ob}}(F)$, for both regular and singular, so that if we were to stand up on the positive side of F and walk along a leaf, the positive side of the intersecting page S_t of the open book would be on our left. In other words, at a non-singular point p on a leaf $l \subset (S_t \cap F)$, let \vec{n}_S be a positive normal to S_t . Then $X_{\text{ob}} := \vec{n}_S \times \vec{n}_F$ is a positive tangent to l . As a result, positive/negative elliptic points are sources/sinks of the vector field X_{ob} .

Proof of Proposition 2.11 The orientations of the leaves give a vector field X_{ob} on F . By the axiom $(\mathcal{F} \text{ iv})$ any singularity of $\mathcal{F}_{\text{ob}}(F)$ is either elliptic or hyperbolic. The statement follows from the Poincaré–Hopf theorem. \square

2.1.2 Proofs of Theorem 2.5 and Proposition 2.6 Since we do not assume incompressibility of the surface F , we *cannot* directly apply Roussarie and Thurston’s general position theorem [41, Theorem 4] or the proof of a corresponding result in braid foliation theory [7, Lemma 2] in order to remove all the local extrema from a foliation satisfying $(\mathcal{F} \text{ i})$, $(\mathcal{F} \text{ ii})$, $(\mathcal{F} \text{ iii})$, $(\mathcal{F} \text{ iv}')$. Instead, we use a trick that we call a finger move.

Proof of Theorem 2.5 Let F be a surface in a general position such that the singular foliation $\mathcal{F} = \mathcal{F}(F)$ satisfies $(\mathcal{F} \text{ i})$, $(\mathcal{F} \text{ ii})$, $(\mathcal{F} \text{ iii})$, $(\mathcal{F} \text{ iv}')$. We show that we can isotope F so that $(\mathcal{F} \text{ iv})$ is satisfied.

Let p be a local extremal point on the page S_t . We will replace p with a pair of elliptic points and one hyperbolic point by the following isotopy, which we call a *finger move*. Repeating finger moves we can get rid of all the local extrema, ie $(\mathcal{F} \text{ iv})$ is satisfied.

Choose an arc γ in S_t that joins p and a binding component B . See Figure 4. If γ intersects other regular leaves of $\mathcal{F}(F)$, by small local perturbation, we make the intersections transverse. Take a small 3–ball neighborhood N of γ (dashed ellipses). We may assume that N contains no singularities of \mathcal{F} other than p . Push a neighborhood of p along γ so that no changes occur outside the region N . See the passage in Figure 4(a)

Call this isotopy a *finger move* supported on N . Figure 4(b) illustrates this finger move viewed from “above” the binding component B .

The finger move removes p and introduces new elliptic (black dots in Figure 4) and hyperbolic (gray dots) singularities to \mathcal{F} . But since the finger move is supported on N , no new local extrema are introduced. More precisely, if a positive normal to S_t agrees (resp. disagrees) with a positive normal to F at p , then the finger move introduces




Figure 4: A finger move supported on a small neighborhood N of γ

one negative (resp. positive) hyperbolic point and a pair of \pm elliptic points. See the top passage in Figure 5. For other part of F that is involved in the finger move, a pair of \pm elliptic points and a pair of \pm hyperbolic points are inserted. See the bottom passage of Figure 5. \square

Proof of Proposition 2.6 Let $\mathcal{F}_{\text{ob}}(F)$ be an open book foliation containing c -circles. For a c -circle c there exists a maximal annulus $c \subset A_c \subset F$ whose interior is foliated only by c -circles and whose boundary components are singular leaves. Let us call A_c a c -circle annulus. The number of c -circle annuli in $\mathcal{F}_{\text{ob}}(F)$ is finite since the number of singularities of $\mathcal{F}_{\text{ob}}(F)$ is finite.

In the following, applying finger moves introduced in the proof of Theorem 2.5 we will eliminate all the c -circle annuli. Recall that a finger move does not introduce new c -circles.

Let $A \subset F$ be a c -circle annulus whose interior consists of a smooth family of c -circles $\{c_t \subset S_t\}$ and let $c_i \subset S_{t_i} \cap \partial A$ ($i = 0, 1$) denote the limit circles of the family. There




Figure 5: (Top) Foliation change by a finger move near a local maximum. (Bottom) Non-singular region involved in a finger move.

is no restriction on the way that A may wind around the binding components. Each limit circle c_{t_i} has one (or two) hyperbolic point(s). (In the latter case the two points must be identical due to the condition $(\mathcal{F}$ iii) and the limit circle is immersed like the singular leaf in cc -pants as in Figure 7.)

Since the open book foliation $\mathcal{F}_{\text{ob}}(F)$ contains only finitely many hyperbolic points, there exists a finite family of disjoint smooth arcs and points

$$\{\text{arc } \alpha_i \subset A, \text{ point } p_i \in B \mid i = 1, \dots, k\},$$

where B is the set of binding components, such that

- every c -circle of A intersects at least one of the arcs α_i ,
- all the intersections of α_i and c -circles are transverse,
- for each α_i there exists a smooth family of arcs $\lambda_t^i \subset S_t$ from the point $\alpha_i \cap S_t$ to p_i that avoids hyperbolic points of $\mathcal{F}_{\text{ob}}(F)$ and is never tangent to leaves of $\mathcal{F}_{\text{ob}}(F)$. See Figure 6. (It is convenient to imagine a triangle $\Delta_i = \{\lambda_t^i\}$ with the bottom edge α_i and the top vertex p_i .)

We apply a finger move (see the proof of Theorem 2.5) along the triangle Δ_i . The open book foliation locally changes as in the bottom passage of Figure 5 in a neighborhood




Figure 6: Arc α_i , point p_i and triangle Δ_i

of α_i . Then all the c -circles through α_i disappear. Repeat finger moves along all $\Delta_1, \dots, \Delta_k$. As a consequence all the c -circles of A disappear. Note that the finger moves may introduce new singularities even away from A if some Δ_i intersect other parts of the surface F . We apply this procedure to every c -circle annulus. \square

2.1.3 Regions

Definition 2.13 [7, page 30] Recall the three types of regular leaves: Type a , b and c (Proposition 2.2). The hyperbolic points in $\mathcal{F}_{\text{ob}}(F)$ are classified into six types, according to types of nearby regular leaves: Type aa , ab , bb , ac , bc and cc as depicted in Figure 7. We call such model regions aa -tile, ab -tile, bb -tile, ac -annulus, bc -annulus, cc -pants, respectively. (Note that ac -annuli do not exist in braid foliation theory [3, page 279].)

For each region, the sign of the hyperbolic point can be either $+1$ or -1 , but the signs of the elliptic points are determined as depicted in Figure 7 due to Claim 2.8. For ac and bc -annuli, the hyperbolic points can be on the left parts of the annuli. The interior of a region is embedded in F as a disc, an annulus, or a pair of pants.




Figure 7: The six types of regions

Definition 2.14 (Degenerate regions) If a region R is of type aa , ac , bc or cc , some parts of ∂R are possibly identified in F . In such case we say that R is *degenerate*. For example, in Figure 8(1), two boundary a -arcs of an aa -tile are identified, and in (2) the two boundary b -arcs of a bc -annulus are identified (we have already seen this in Figure 5).

On the other hand, a region like in Figure 8(3), where two ends of the singular leaf lie on the same positive elliptic point, does not exist. This is because around an elliptic point all the leaves (both regular and singular) sit on distinct pages.




Figure 8: (1) A degenerate aa -tile (2) A degenerate bc -annulus (3) Non-existing region

We study degenerate regions in [30].

The next proposition shows one of the useful combinatorial features of open book foliations. It is originally a theorem in braid foliation theory.

Proposition 2.15 (Region decomposition [3, Theorem 1.2]) *If $\mathcal{F}_{\text{ob}}(F)$ contains a hyperbolic point, the surface F is decomposed into a union of model regions whose interiors are disjoint.*

We omit a proof and refer the readers to the proof of [3, Theorem 1.2].

The decomposition is called a *region decomposition* of F . It describes how F is embedded in $M_{(S,\phi)}$. If $\mathcal{F}_{\text{ob}}(F)$ has no c -circles then the region decomposition gives a cellular decomposition of F .

2.1.4 The graph G_{--}

Definition 2.16 The two flow lines, induced by the orientation vector field X_{ob} on F (Definition 2.12), approaching to (resp. departing from) the hyperbolic point in an aa , ab or bb -tile is called *stable* (resp. *unstable*) *separatrices*.

Definition 2.17 [11, page 471] The *graph G_{--}* is a graph embedded in F . The edges of G_{--} are the unstable separatrices for negative hyperbolic points in aa , ab and bb -tiles. See Figure 9. We regard the negative hyperbolic points as part of the edges. The vertices of G_{--} are the negative elliptic points in ab and bb -tiles and the end points of the edges of G_{--} that lie on ∂F , called the *fake vertices*.




Figure 9: The graph G_{--}

In the same way, we can define G_{++} the graph consists of positive elliptic points and stable separatrices of positive hyperbolic points.

Remark 2.18 The origin of the above definition is the graphs $G_{\pm\pm}$, $G_{\pm\mp}$ [3, page 314; 11, page 471] in braid foliation theory. In convex surface theory our G_{--} corresponds to a sub-graph of the Giroux graph [25, page 646].

We will use the graphs G_{--} and G_{++} to define a transverse overtwisted disc and to give an alternative proof to the Bennequin–Eliashberg inequality in Section 4.

2.1.5 Movie presentation A useful tool for expressing how the surface, F , is embedded in $M_{(S,\phi)}$, *movie presentations*, can be borrowed from braid foliation theory; see [7, Figure 8]. Using a movie presentation allows us to grasp the whole picture of $\mathcal{F}_{\text{ob}}(F)$.

Let $\{S_{t_i}\}_{i=1,\dots,k}$ be the set of singular pages of $\mathcal{F}_{\text{ob}}(F)$, where $0 < t_1 < t_2 < \dots < t_k < 1$. Consider the family $\{(S_t, F \cap S_t) \mid t \in [0, 1]\}$ of slices of F by the pages S_t . For $s, t \in (t_i, t_{i+1})$, the slices $(S_s, F \cap S_s)$ and $(S_t, F \cap S_t)$ are isotopic, and the isotopy type of $(S_t, F \cap S_t)$ changes only when $t = t_i$. The describing arcs (Definition 2.9) encode all the information of the configuration changes.

Choose $s_0 = 0$, $s_k = 1$ and $s_i \in (t_i, t_{i+1})$. Consider the slices

$$\{(S_{s_i}, F \cap S_{s_i}) \mid i = 0, \dots, k\}.$$

These are the slices on which we may place describing arcs. The describing arc for the singularity on S_{t_i} is found on $S_{s_{i-1}}$. The above observation shows that those are the slices that determine the embedding of F and the open book foliation $\mathcal{F}_{\text{ob}}(F)$ up to isotopy. The slice $(S_1, F \cap S_1)$ is identified with the slice $(S_0, F \cap S_0)$ under the monodromy ϕ . We call this family of slices with describing arcs a *movie presentation* of $\mathcal{F}_{\text{ob}}(F)$.

We will often use part of a movie presentation to express a local picture of a surface. Also, for the reader's convenience, some movie presentations may contain singular slices $(S_{t_i}, F \cap S_{t_i})$ like in Figure 12.

2.1.6 Examples of open book foliations

Example 2.19 First we consider the simplest open book (D^2, id) , which supports the standard tight contact structure on S^3 . This is the case that Birman and Menasco studied in their braid foliation theory. Consider a 2-sphere F embedded as shown in the left sketch of Figure 10.

Since F intersects the binding in four points, the open book foliation $\mathcal{F}_{\text{ob}}(F)$ has four elliptic points, two positive and two negative. It also has two hyperbolic points of opposite signs where F is tangent to pages of the open book (we may assume that the hyperbolic points lie on pages $S_{1/4}$ and $S_{3/4}$). The right sketch of Figure 10 depicts the whole picture of $\mathcal{F}_{\text{ob}}(F)$ and Figure 11 depicts a movie presentation of $\mathcal{F}_{\text{ob}}(F)$, where the dashed arrows indicate positive normals to F . Note that the open book foliation $\mathcal{F}_{\text{ob}}(F)$ contains inessential b -arcs so, strictly speaking, this foliation is not treated in braid foliation theory.




Figure 10: Example 2.19




Figure 11: A movie presentation of Example 2.19

Example 2.20 Next we study a more informative example. Consider the open book $(S, \phi) := (A, T_A^{-1})$ where A denotes an annulus and T_A the right-handed Dehn twist along a core circle of A . The ambient manifold is again S^3 . However in this case the binding is a negative Hopf link and (A, T_A^{-1}) supports an overtwisted contact structure.

In order to visualize an overtwisted disc, D , we cut the complement of the binding $S^3 \setminus B$ along the page S_0 . The resulting manifold is homeomorphic to $S \times [0, 1]$ and each page S_t is naturally identified with $S \times \{t\}$. The disc $D \subset M_{(S, \phi)}$ is also cut out along $D \cap S_0$ and becomes a properly embedded surface, D' , in $S \times [0, 1]$ such that $D' \cap (S \times \{0\}) = \phi(D' \cap (S \times \{1\}))$ and $D' \cap (\partial S \times [0, 1]) = (D \cap \partial S_0) \times [0, 1]$. The left sketch in Figure 12 shows how D' is embedded in $S \times [0, 1]$.

The sketch on the right depicts a movie presentation of $\mathcal{F}_{\text{ob}}(D)$. (For convenience, as we note in Section 2.1.5, redundant slices that contain hyperbolic points are added in the 2nd and 4th rows.) We see that the multi-curve in the top annulus is identified with the multi-curve in the bottom annulus under the monodromy T_A^{-1} . The movie presentation also shows that the open book foliation $\mathcal{F}_{\text{ob}}(D)$ contains two positive hyperbolic points, two positive elliptic points and one negative elliptic point. See Figure 13 for the entire picture of $\mathcal{F}_{\text{ob}}(D)$.




Figure 12: Example 2.20: An overtwisted disc in an annulus open book (A, T_A^{-1})




Figure 13: Example 2.20: The open book foliation $\mathcal{F}_{\text{ob}}(D)$

2.2 Open book foliation vs characteristic foliation

Let $\mathcal{F}_\xi(F)$ denote the characteristic foliation of a surface F embedded in (M, ξ) . In this section, we compare the open book foliation $\mathcal{F}_{\text{ob}}(F)$ and the characteristic foliation $\mathcal{F}_\xi(F)$.

Theorem 2.21 (Structural stability) *Assume that a surface F in $M_{(S, \phi)}$ admits an open book foliation $\mathcal{F}_{\text{ob}}(F)$. There exists a contact structure ξ on $M_{(S, \phi)}$ supported by the open book (S, ϕ) such that $e_\pm(\mathcal{F}_{\text{ob}}(F)) = e_\pm(\mathcal{F}_\xi(F))$ and $h_\pm(\mathcal{F}_{\text{ob}}(F)) = h_\pm(\mathcal{F}_\xi(F))$.*

Moreover, if $\mathcal{F}_{\text{ob}}(F)$ contains no c -circles, then $\mathcal{F}_{\text{ob}}(F)$ and $\mathcal{F}_\xi(F)$ are topologically conjugate, namely there exists a homeomorphism of F that takes $\mathcal{F}_{\text{ob}}(F)$ to $\mathcal{F}_\xi(F)$. In particular [22, Lemma 2.1] implies that F is a convex surface.

Proof Recall the Thurston–Winkelnkemper construction (Thurston and Winkelnkemper [42], and Geiges [23, page 151–153]) of a contact structure compatible with the open book (S, ϕ) :

Away from the binding, Thurston and Winkelnkemper’s contact 1–form is written as $\alpha = \beta_t + C dt$, where $t \in [0, 1]$ (page parameter), $C \gg 1$ is a sufficiently large constant number and $\{\beta_t\}$ is a smooth family of 1–forms on the page S_t such that $d\beta_t$ is an area form of S_t of total area 2π and $\beta_1 = \phi^*\beta_0$. Such a family $\{\beta_t\}$ is not unique, so we choose any to start with.

Near a binding component there exist cylindrical coordinates (θ, r, t) , where θ represents the positive direction of the binding and $t \in [0, 1]$ is the same t as above, such that

$$(2-1) \quad \alpha = 2 d\theta + r^2 dt.$$

Assume that F admits an open book foliation $\mathcal{F}_{\text{ob}}(F)$. In the following we use the 1–form on $M_{(S,\phi)}$, α , chosen above, and contact planes $\xi := \ker \alpha$ to study neighborhoods of singular and non-singular points.

Elliptic points Suppose $p \in \text{Int}(F)$ is an elliptic point of $\text{sgn}(p) =: \epsilon \in \{\pm 1\}$. This means that a binding component, γ , intersects F transversely at p with sign ϵ ; see Figure 14(1). Take a disc neighborhood $D \subset F$ of p whose open book foliation $\mathcal{F}_{\text{ob}}(D)$ contains no other singularities. By (2-1) we know that along γ the contact planes and γ intersect transversely with sign $+1$. We push down (or up) a very small neighborhood $D_0 \subset D$ of p along γ without touching the rest of the surface; see Figure 14(2).




Figure 14

Since this operation preserves the open book foliation, we may call the perturbed surface by the same name F . By the symmetry with respect to γ of the pushed D_0 and $\alpha = 2 d\theta + r^2 dt$, at the new $p = D_0 \cap \gamma$ the tangent plane and the contact plane satisfy $T_p F = \epsilon \cdot \xi_p$, hence the new p is an elliptic point of the characteristic foliation

$\mathcal{F}_\xi(D_0)$ of sign ϵ . (If we push *up* in Figure 14(2) instead of push *down* exactly the same argument holds.)

Hyperbolic points Let $p \in \mathcal{F}_{\text{ob}}(F)$ be a hyperbolic point of $\text{sgn}(p) = +1$. (A parallel argument holds for the negative case.) Take an open ball neighborhood $U \subset M_{(S,\phi)}$ of p in which p is the only singularity of the open book foliation $\mathcal{F}_{\text{ob}}(F)$. Let (x, y, z) be coordinates of U such that

- (i) z is a coordinate for a sub-interval of $[0, 1]$ such that $\partial_z = \partial_t$,
- (ii) (x, y) are coordinates for the open disc $U \cap S_t$,
- (iii) $p = (0, 0, 0)$.

We may assume that $F \cap U$ is a saddle surface and satisfies $z = x^2 - y^2$. The normal vector \vec{n}_F to the surface at $q = (x, y, z) \in F \cap U$ is $\vec{n}_F = (-2x, 2y, 1)$. Suppose that the contact plane $\xi_q = \ker(\alpha_q)$ at q is spanned by

$$(2-2) \quad \xi_q = \text{span}(\partial_x + f(q)\partial_z, \partial_y + g(q)\partial_z)_{\mathbb{R}}$$

for some smooth functions $f, g: U \rightarrow \mathbb{R}$. Let $\vec{n}_\xi := (-f(q), -g(q), 1)$. Then \vec{n}_ξ is a positive normal to ξ_q . We have

$$\begin{aligned} 0 &= \alpha_q(\partial_x + f(q)\partial_z) = \beta_q(\partial_x) + Cf(q), \\ 0 &= \alpha_q(\partial_y + g(q)\partial_z) = \beta_q(\partial_y) + Cg(q). \end{aligned}$$

Since C can be taken as large as we want, we have

$$(2-3) \quad |f(q)| = |\beta_q(\partial_x)|/C \ll 1 \quad \text{and} \quad |g(q)| = |\beta_q(\partial_y)|/C \ll 1.$$

Therefore, if we take U small enough there exists a unique point $p_0 \in U \cap F$ at which $\vec{n}_F = \vec{n}_\xi$, and the foliations $\mathcal{F}_{\text{ob}}(F \cap U)$ and $\mathcal{F}_\xi(F \cap U)$ are topologically conjugate. In particular, p_0 is a hyperbolic point of the characteristic foliation and $\text{sgn}(p_0) = \text{sgn}(p)$.

Non-singular points Let $p \in \text{Int}(F)$ be a non-singular point in $\mathcal{F}_{\text{ob}}(F)$. Take a small open 3-ball neighborhood $U \subset M_{(S,\phi)}$ of p so that the surface $F \cap U$ contains no singularity of $\mathcal{F}_{\text{ob}}(F)$. Let $(x, y, z) \in \mathbb{R}^3$ be coordinates of U with the above conditions (i), (ii), (iii). We may suppose that $z = ky$ is satisfied on $F \cap U$ for some $k \neq 0$. So the leaves of $\mathcal{F}_{\text{ob}}(U \cap F)$ are the integral curves of the vector field ∂_x . Given a point $q = (x, y, z) \in U$, we may assume the above (2-2) and (2-3). Hence the normal vector $\vec{n}_\xi = (-f(q), -g(q), 1)$ to ξ_q and the normal vector $\vec{n}_F = (0, -k, 1)$ to $T_q F$ are not parallel to each other, ie $\xi_q \neq \pm T_q F$, and the point q is not a singularity of $\mathcal{F}_\xi(F)$.

The above arguments conclude the first assertion of the theorem: $e_{\pm}(\mathcal{F}_{\text{ob}}(F)) = e_{\pm}(\mathcal{F}_{\xi}(F))$ and $h_{\pm}(\mathcal{F}_{\text{ob}}(F)) = h_{\pm}(\mathcal{F}_{\xi}(F))$.

To prove the second assertion, we assume that $\mathcal{F}_{\text{ob}}(F)$ contains no c -circles. By Proposition 2.15, F decomposes into type aa , ab and bb -tiles. For the stable separatrices \mathcal{S} in each tile we take a small disc neighborhood $\mathcal{D} \subset F$ of \mathcal{S} . The leaves of $\mathcal{F}_{\text{ob}}(\mathcal{D})$ are oriented outward along the boundary $\partial\mathcal{D}$. This implies that $\partial\mathcal{D}$ is a positive braid with respect to the open book (S, ϕ) , or equivalently, a positive transverse unknot in $(M_{(S, \phi)}, \xi)$, where ξ is the contact structure chosen above. Therefore, the leaves of $\mathcal{F}_{\xi}(\mathcal{D})$ are also outward along $\partial\mathcal{D}$. Moreover, the above argument shows that $\mathcal{F}_{\xi}(\mathcal{D})$ and $\mathcal{F}_{\text{ob}}(\mathcal{D})$ are topologically conjugate relative to $\partial\mathcal{D}$.

A similar argument holds for each unstable separatrices. Hence we conclude that $\mathcal{F}_{\xi}(F)$ and $\mathcal{F}_{\text{ob}}(F)$ are topologically conjugate. \square

Remark 2.22 The above proof of Theorem 2.21 shows that the open book and characteristic foliations may coincide, especially when there are no c -circles in \mathcal{F}_{ob} . Interesting contrast is found between open book foliations and characteristic foliations (on convex surfaces).

- For a given closed surface F , we can always find a convex surface F_{cv} that is isotopic and C^{∞} -close to F . However, in general, there may not exist a surface admitting an open book foliation that is even C^1 -close to F (eg when F has local extrema relative to the pages and then we apply finger moves).
- The *dividing set* Γ_F of a convex surface F encodes essential information of the contact structure near F . It yields a decomposition $F \setminus \Gamma = F_+ \sqcup F_-$ of F . If $\mathcal{F}_{\text{ob}}(F)$ has no c -circles then the region F_- is homotopy equivalent to our graph G_{--} .
- In the characteristic foliation on a convex surface, any closed leaf is either repelling or attracting, and there are no type ac , bc and cc -hyperbolic points (Figure 7) due to the Morse–Smale condition (cf [23, page 171]). On the other hand, an annular neighborhood of a c -circle in an open book foliation is foliated by parallel c -circles.
- In the theory of convex surfaces, *Giroux elimination* [25; 23, Lemma 4.6.26] allows us to remove a pair of elliptic and hyperbolic singularities of the same sign by an arbitrary C^0 -small isotopy. Morally, one thinks that Giroux elimination corresponds to elimination of a certain arrangement of a pair consisting of a local extremum and a saddle point in an open book foliation $\mathcal{F}_{\text{ob}}(F)$ by “flattening” the surface F . See Figure 15.

In a subsequent paper [30] we discuss a number of operations in open book foliation theory that allow us to remove singularities.




Figure 15: (Left) Elimination of a local extremum and a saddle point in open book foliation. (Right) Giroux elimination takes place in the shaded region of characteristic foliation.

3 The self-linking number

A *transverse knot* in a contact 3–manifold (M, ξ) is an embedding of S^1 transverse to ξ . It is known that a transverse knot is a contact submanifold of a contact 3–manifold (see [23, Remark 2.1.15] for example). In this section we study an invariant of transverse knots, called the self-linking number.

Definition 3.1 Let $L \subset (M, \xi)$ be a transverse link that bounds a surface F , ie L is 0-homologous. The rank-2 vector bundle $\xi|_F \rightarrow F$ over F is trivializable. Let s be a nowhere vanishing smooth section of the bundle. Push L into the direction of s and call the resulting link L^{+s} . The *self-linking number* of L relative to $[F] \in H_2(M, L; \mathbb{Z})$, which we denote by $\text{sl}(L, [F])$, is the algebraic intersection number of L^{+s} and F .

Using Mitsumatsu and Mori's Theorem [35, Appendix] or Pavelescu's [39; 40], we can identify a transverse link in (M, ξ) with a closed braid in any compatible open book (S, ϕ) . The goal of this section is to prove Theorem 3.10, a self-linking number formula for closed braids.

Our strategy is to construct a special Seifert surface Σ for a given closed braid and count the singularities of its open book foliation $\mathcal{F}_{\text{ob}}(\Sigma)$, then apply the following proposition:

Proposition 3.2 Suppose that $F \subset M_{(S, \phi)}$ is a surface with the open book foliation $\mathcal{F}_{\text{ob}}(F)$. In particular, ∂F is a transverse link in $(M_{(S, \phi)}, \xi_{(S, \phi)})$. Recall the integers $e_{\pm} = e_{\pm}(\mathcal{F}_{\text{ob}}(F))$, $h_{\pm} = h_{\pm}(\mathcal{F}_{\text{ob}}(F))$ defined in Definition 2.10. We have

$$\text{sl}(\partial F, [F]) = -\langle e(\xi), [F] \rangle = -(e_+ - e_-) + (h_+ - h_-).$$

Proof The self-linking number formula in characteristic foliation theory (see [23, page 203] for example) together with Theorem 2.21 yields the above formula. \square

In order to state our main theorem (Theorem 3.10), we first need to define a function $c: \mathcal{MCG}(S) \times H_1(S; \partial S) \rightarrow \mathbb{Z}$ in Section 3.1. Later in Section 3.5 we show that the function c is related to the first Johnson–Morita homomorphism, a well-studied homomorphism in mapping class group theory.

3.1 Definition of function c

Let $S = S_{g,r}$ be an oriented genus g surface with r boundary components. We divide the surface S by *walls* (dashed arcs in Figure 19) into $g + r - 1$ *chambers*, g of which are once-punctured tori and $r - 1$ of which are annuli.

Definition 3.3 (Normal form) A relative homology class $a \in H_1(S, \partial S)$ is represented by a set of properly embedded oriented simple closed curves and arcs in S . Among such multi-curve representatives, we take a special one, $N(a)$, which satisfies the following conditions:

- $N(a)$ does not intersect the walls.

- Any subset of $N(a)$ has non-trivial homology in $H(S, \partial S)$, ie the components of $N(a)$ in a torus (resp. an annulus) chamber are a torus knot or link (resp. parallel arcs joining γ_0 and γ_i in Figure 19) oriented in the same direction.

Clearly the multi-curve $N(a)$ is uniquely determined up to isotopy. We call $N(a)$ the *normal form* of the homology class $a \in H_1(S, \partial S)$.

Definition 3.4 (OB cobordism) Let A and A' be oriented, properly embedded multi-curves in S representing the same homology class $[A] = [A'] \in H_1(S, \partial S)$. An *open book foliation cobordism* (OB cobordism) between A and A' , denoted by $A \xrightarrow{\Sigma} A'$, is a properly embedded oriented compact surface Σ in $S \times [0, 1]$ such that:

- $\Sigma \cap S_0 = \partial \Sigma \cap S_0 = -A \times \{0\}$
- $\Sigma \cap S_1 = \partial \Sigma \cap S_1 = A' \times \{1\}$
- $\partial A = A \cap \partial S = A' \cap \partial S = \partial A'$
- $\partial \Sigma = (-A \times \{0\}) \cup (A' \times \{1\}) \cup (\partial A \times [0, 1])$
- The fibration $\{S_t\}_{t \in [0,1]}$ induces a foliation \mathcal{F}_Σ on Σ , all of whose singularities are of hyperbolic type

Proposition 3.5 There is an OB cobordism $A \xrightarrow{\Sigma} N(a)$ for any multi-curve representative A of $a \in H_1(S, \partial S)$. That is, if multi-curves A and A' represent the same homology class then there exists an OB cobordism $A \xrightarrow{\Sigma} A'$.

Proof We construct an oriented surface Σ embedded in $S \times [0, 1]$ with $\Sigma \cap S_0 = -A$ and $\Sigma \cap S_1 = N(a)$. Let w be one of the walls. Since $[A] = [N(a)] \in H_1(S, \partial S)$ and the normal form $N(a)$ does not intersect w , the algebraic intersection number $[A] \cdot w = 0$. We take a collar neighborhood $\nu(w) \subset S$ of w so that each component of $\nu(w) \cap A$ has geometric intersection number 1 with w . The arcs $\nu(w) \cap A$ may not all have the same orientation. As $t \in [0, 1]$ increases, we apply the configuration changes to pairs of consecutive arcs in $\nu(w) \cap A$ with opposite orientations as in the passage of Figure 16, until we remove all the arcs of $\nu(w) \cap A$. Each configuration change introduces a new hyperbolic singularity. We repeat the procedure for all the walls. The deformed multi-curve A , which we denote A' , no longer intersects the walls.

The multi-curve $A' \subset S$ may contain null-homologous sets of c -circles. We remove them by the following three steps.

Step 1 If there exist c -circles bounding concentric discs in a chamber H of S and oriented in the same direction, then we remove them from the outermost one. We can find a describing arc of a hyperbolic point (cf Figure 3) that joins the outermost



Figure 16: Configuration change of $v(w) \cap A$




Figure 17: Step 1 (top) and Step 2 (bottom)

c -circle and some curve in $A' \cap H$ and is properly embedded in $H \setminus A'$. As shown in the top row of Figure 17, one hyperbolic singularity is introduced then the c -circle disappears. The sign of the hyperbolic singularity is $+1$ if and only if the c -circle is oriented clockwise.

Step 2 If there is a pair of c -circles with opposite orientations that bounds an annulus in $S \setminus A'$, then remove the pair by introducing a hyperbolic singularity of sign ε between the two c -circles as in Figure 17. The resulting c -circle bounds a disc that can be removed by Step 1 at the expense of another hyperbolic singularity of sign $-\varepsilon$.

Step 3 Let H be a once-punctured torus chamber of S . After Steps 1 and 2, there exist $p, q, r \in \mathbb{Z}$ such that in H the multi-curve A' is the union of (p, q) torus link and r boundary parallel c -circles oriented in the same direction. As in Figure 18 we remove the c -circles by introducing r hyperbolic points of the same sign. The sign depends on the signs of p, q and the orientation of the boundary parallel c -circles.

Now A' is deformed to the normal form $N(a)$. Hence, we get a desired surface Σ . \square

Proposition 3.6 For an OB cobordism $A \xrightarrow{\Sigma} A'$, let $h_+(\mathcal{F}_\Sigma)$ (resp. $h_-(\mathcal{F}_\Sigma)$) denote the number of the positive (resp. negative) hyperbolic singularities of \mathcal{F}_Σ . The value

$$d(A \xrightarrow{\Sigma} A') := h_+(\mathcal{F}_\Sigma) - h_-(\mathcal{F}_\Sigma)$$

is independent of the choice of cobordism surface Σ and it only depends on the multi-curve representatives A and A' . Hence we may denote

$$d(A, A') := d(A \xrightarrow{\Sigma} A').$$




Figure 18: Step 3: Remove boundary parallel null-homologous c -circles by configuration changes along the dashed arcs.

Proof Suppose that $A \xrightarrow{\Sigma'} A'$ is another OB cobordism. We embed $-\Sigma'$ in $S \times [1, 2]$. We glue Σ and $-\Sigma'$ at the page S_1 and obtain a surface F' in $S \times [0, 2]$. Since $F' \cap S_2 = A = -(F' \cap S_0)$, we can further identify $F' \cap S_2$ and $F' \cap S_0$ by the identity map that defines a surface F embedded in the open book (S, id) . Since a \pm -hyperbolic singular point in $\mathcal{F}_{\Sigma'}$ turns to a \mp -hyperbolic point in $\mathcal{F}_{-\Sigma'}$ we have $d(A' \xrightarrow{-\Sigma'} A) = -d(A \xrightarrow{\Sigma'} A')$ and

$$(3-1) \quad h_+(\mathcal{F}_{\text{ob}}(F)) - h_-(\mathcal{F}_{\text{ob}}(F)) = d(A \xrightarrow{F} A) = d(A \xrightarrow{\Sigma} A') - d(A \xrightarrow{\Sigma'} A').$$

By Definition 3.4 the elliptic points in $\mathcal{F}_{\text{ob}}(F)$ correspond to the lines $\partial A \times [0, 2]$. Since the endpoints of each arc component of A correspond to two elliptic points of opposite signs, we get

$$(3-2) \quad e_+(\mathcal{F}_{\text{ob}}(F)) = e_-(\mathcal{F}_{\text{ob}}(F)).$$

Let ξ_{id} be the contact structure supported by the open book (S, id) . Since the Euler class of ξ_{id} is equal to zero, by Proposition 3.2, (3-1) and (3-2), we have

$$0 = \langle e(\xi_{\text{id}}), [F] \rangle = d(A \xrightarrow{\Sigma} A') - d(A \xrightarrow{\Sigma'} A'). \quad \square$$

We are ready to define the function $c([\phi], a)$. The following definition is geometric. Later we study algebraic properties of $c([\phi], a)$ in Propositions 3.14, 3.20 and Theorem 3.21.

Let $\mathcal{MCG}(S)$ denote the *mapping class group* of S , that is, the group of isotopy classes of orientation-preserving homeomorphisms of S fixing the boundary ∂S pointwise.

Definition 3.7 Let $[\phi] \in \mathcal{MCG}(S)$ and $a \in H_1(S, \partial S)$. Define

$$c([\phi], a) := d(\phi(N(a)), N(\phi_*a)).$$

In general, the multi-curve $\phi(N(a))$ may not be isotopic to $N(\phi_*(a))$. But if $\phi(N(a))$ is isotopic to $N(\phi_*(a))$, we can choose an OB cobordism $\phi(N(a)) \xrightarrow{\Sigma} N(\phi_*(a))$ to be the product $\Sigma \simeq \phi(N(a)) \times [0, 1]$ with no hyperbolic singularities, hence $c([\phi], a) = 0$. We call such an OB cobordism *trivial*.

3.2 A self-linking number formula for braids

Let $S = S_{g,r}$ be an oriented genus g surface with r boundary components $\gamma_1, \dots, \gamma_r$. The orientation of γ_i is induced from that of S . Let b be an n -stranded braid in $S \times [0, 1]$ with $b \cap S_1 = b \cap S_0 = \{x_1, \dots, x_n\} \subset S$. By braid isotopy we may assume that points x_1, \dots, x_n are lined up in this order on an arc parallel to and very close to γ_0 . The arc $\{x_i\} \times [0, 1]$ is called the i^{th} braid strand in $S \times [0, 1]$. We define oriented loops, $\rho_i \subset S$ ($i = 1, \dots, 2g + r - 1$) with the base point x_n as in Figure 19.




Figure 19: Surface S

Geometrically, ρ_i represents the n^{th} braid strand winding along ρ_i as $t \in [0, 1]$ increases. Let σ_i denote the positive half twist of the i^{th} and the $(i + 1)^{\text{st}}$ braid strands. As a consequence of the Birman exact sequence [2], the braid b is represented by a braid word $b_1^{\varepsilon_1} b_2^{\varepsilon_2} \dots b_l^{\varepsilon_l}$ (read from the left) where $b_i \in \{\rho_1, \dots, \rho_{2g+r-1}, \sigma_1, \dots, \sigma_{n-1}\}$ and $\varepsilon_i \in \mathbb{Z} \setminus \{0\}$.

Fix a diffeomorphism $\phi \in \text{Aut}(S, \partial S)$. Since x_i is near $\gamma_1 \subset \partial S$, we have $\phi(x_i) = x_i$ and identify $\{x_i\} \times \{1\}$ and $\{x_i\} \times \{0\}$ under ϕ , which yields a closed braid \hat{b} in $M_{(S,\phi)}$. We assume that \hat{b} is null-homologous in the rest of the section.

Claim 3.8 Put $[b] = \sum_{i=1}^l \varepsilon_i [b_i] \in H_1(S; \mathbb{Z})$, where we set $[\sigma_k] = 0$ for $k = 1, \dots, n - 1$. Then there exists a (not necessarily unique) homology class $a \in H_1(S, \partial S; \mathbb{Z})$ such that $[b] = a - \phi_*(a)$ in $H_1(S; \mathbb{Z})$.

Proof The homology group of the manifold $M_{(S,\phi)}$ is computed by Etnyre and Ozbagci [21, page 3136]:

$$H_1(M_{(S,\phi)}; \mathbb{Z}) = \langle [\rho_1], \dots, [\rho_{2g+r-1}] \mid [\rho'_i] - \phi_*[\rho'_i] = 0, i = 1, \dots, 2g+r-1 \rangle,$$

where

$$\rho'_i = \begin{cases} \text{a properly embedded arc from } \gamma_0 \text{ to } \gamma_i \text{ and dual to } \rho_i \text{ for } i = 1, \dots, r-1, \\ \rho_i \text{ for } i = r, \dots, 2g+r-1. \end{cases}$$

Though ρ'_i is an arc for $i = 1, \dots, r-1$, since $\phi = \text{id}$ on ∂S , we can view $\rho'_i \cup \phi(-\rho'_i)$ as an oriented (immersed) loop in $\text{Int}(S)$. Then we consider $[\rho'_i] - \phi_*[\rho'_i] \in H_1(S; \mathbb{Z})$ representing the loop $\rho'_i \cup \phi(-\rho'_i)$.

Since $[\widehat{b}] = 0$ in $H_1(M; \mathbb{Z})$, there exist $s_i \in \mathbb{Z}$ for $i = 1, \dots, 2g+r-1$, such that

$$[b] = \sum_{i=1}^{2g+r-1} s_i([\rho'_i] - \phi_*[\rho'_i]) \quad \text{in } H_1(S; \mathbb{Z}).$$

Hence if we put $a = \sum_{i=1}^{2g+r-1} s_i[\rho'_i]$, under the identification

$$[\rho'_i] - \phi_*[\rho'_i] = [\rho'_i \cup \phi(-\rho'_i)],$$

we have $[b] = a - \phi_*(a)$. □

Definition 3.9 For homology classes $[a_1] \in H_1(S, \partial S; \mathbb{Z})$ and $[a_2] \in H_1(S; \mathbb{Z})$, we denote the algebraic intersection number by $[a_1] \cdot [a_2] \in \mathbb{Z}$. It counts the transverse intersections of representatives a_1 and a_2 algebraically in the way described in Figure 20. For example, we have $[\rho'_1] \cdot [\rho_1] = 1$ and $[\rho_r] \cdot [\rho_{r+1}] = 1$.



Figure 20: Algebraic intersection number $[a_1] \cdot [a_2]$

Here is our main theorem of this section:

Theorem 3.10 (Self-linking number formula) *Let $[b] \in H_1(S; \mathbb{Z})$ and \widehat{b} be as above. Let $a \in H_1(S, \partial S; \mathbb{Z})$ be a homology class such that $[b] = a - \phi_*(a)$ in $H_1(S; \mathbb{Z})$ (see Claim 3.8). Recall the function $c([\phi], a)$ in Definition 3.7. For the choice of $a \in H_1(S, \partial S; \mathbb{Z})$, there exists a Seifert surface $\Sigma = \Sigma_a$ of \widehat{b} such that the self-linking number satisfies the formula*

$$(3-3) \quad \text{sl}(\widehat{b}, [\Sigma]) = -n + \widehat{\text{exp}}(b) - \phi_*(a) \cdot [b] + c([\phi], a),$$

where

$$\widehat{\text{exp}}(b) = \sum_{i=1}^l \varepsilon_i - \sum_{1 \leq j < i \leq l} \varepsilon_i \varepsilon_j [b_j] \cdot [b_i].$$

Remark 3.11 The formula (3-3) is a generalization of Bennequin’s self-linking formula of braids in the open book (D^2, id) [1], and it also covers the works in [32; 33]. When $(S, \phi) = (D^2, \text{id})$ the function $\widehat{\text{exp}}$ is equal to the usual exponent sum, $\text{exp}: B_n \rightarrow \mathbb{Z}$, for the Artin braid group B_n and $\phi_*(a) \cdot [b] = c([\phi], a) = 0$. Thus the formula (3-3) contains Bennequin’s self linking formula

$$\text{sl}(\widehat{b}) = -n + \text{exp}(b).$$

With more elaborate investigation of the function c we will deduce the self-linking number formulae of [32; 33] in Corollary 3.17 below.

Proof For each $i = 1, \dots, n$, take a point y_i on the binding component γ_0 near x_i so that y_1, \dots, y_n lined up in this order with respect to the orientation of γ_0 ; see Figure 19. Choose a properly embedded arc v_i from x_i to y_i that is contained in a small collar neighborhood of γ_0 so that $\phi(v_i) = v_i$. We require that v_1, \dots, v_n are mutually disjoint.

Construction of surface $\Sigma_* \subset S \times [0, \frac{1}{2}]$ Fix $a \in H_1(S, \partial S; \mathbb{Z})$ with $a - \phi_*(a) = [b]$. Let $N(a)$ denote the normal form of a ; see Definition 3.3. Let $A_1, A_{1/2}, A_0$ be oriented multi-curves in S defined by

$$\begin{aligned} A_1 &= v_1 \cup \dots \cup v_n \cup N(a), \\ A_{1/2} &= v_1 \cup \dots \cup v_n \cup N(\phi_*a), \\ A_0 &= \phi(A_1) = v_1 \cup \dots \cup v_n \cup \phi(N(a)). \end{aligned}$$

Unlike A_1 or $A_{1/2}$, the multi-curve A_0 possibly intersects the walls. We have

$$[A_1] = a, \quad [A_{1/2}] = [A_0] = \phi_*a.$$

Let $\phi(N(a)) \xrightarrow{\Sigma_\circ} N(\phi_*a)$ be an OB cobordism whose existence is guaranteed by Proposition 3.5. We compress Σ_\circ vertically to fit in $S \times [0, \frac{1}{2}]$ and take disjoint union

with the vertical rectangle strips $(v_1 \cup \dots \cup v_n) \times [0, \frac{1}{2}]$. We call the resulting surface Σ_* . By the construction

$$\Sigma_* \cap S_0 = -A_0, \quad \Sigma_* \cap S_{1/2} = A_{1/2},$$

where $S_0 = S \times \{0\}$ and $S_{1/2} = S \times \{\frac{1}{2}\}$ are pages of the open book, and by Definition 3.7 the algebraic count of the hyperbolic points of $\mathcal{F}_{\text{ob}}(\Sigma_*)$ is $c([\phi], a)$.

Construction of surface $\Sigma_{} \subset S \times [\frac{1}{2}, 1]$** The next goal is to construct an oriented surface Σ_{**} embedded in $S \times [\frac{1}{2}, 1]$ with $\Sigma_{**} \cap S_1 = A_1$ and $\Sigma_{**} \cap S_{1/2} = -A_{1/2}$. Recall that b is represented by the braid word $b_1^{\varepsilon_1} \dots b_l^{\varepsilon_l}$. Let $I_i = [\frac{l+i-1}{2l}, \frac{l+i}{2l}]$. Then $[\frac{1}{2}, 1] = I_1 \cup \dots \cup I_l$. We will build an oriented surface Σ_i embedded in $S \times I_i$ inductively from $i = 1$ to l such that:

- (1) $\Sigma_1 \cap S_{1/2} = -A_{1/2}$ and $\Sigma_l \cap S_1 = A_1$
- (2) $\Sigma_i \cap S_{(l+i)/2l} = -(\Sigma_{i+1} \cap S_{(l+i)/2l})$; we denote this multi-curve on the page $S_{(l+i)/2l}$ by $A_{(l+i)/2l}$
- (3) $A_{(l+i)/2l}$ does not intersect the walls
- (4) $A_{(l+i)/2l}$ contains $v_1 \cup \dots \cup v_n$ and any subset of $A_{(l+i)/2l} \setminus (v_1 \cup \dots \cup v_n)$ has non-trivial homology in $H_1(S, \partial S)$
- (5) $\partial \Sigma_i \cap (S \times \text{Int}(I_i)) = b_i^{\varepsilon_i}$, so $[A_{(l+i)/2l}] = [A_{1/2}] + \varepsilon_1[b_1] + \dots + \varepsilon_i[b_i]$ in $H_1(S, \partial S)$

Eventually we will define $\Sigma_{**} = \Sigma_1 \cup \dots \cup \Sigma_l$. Suppose that we have constructed $\Sigma_1, \dots, \Sigma_{i-1}$ satisfying the above conditions.

Case 1 If the braid word $b_i = \sigma_k$, then as $t \in I_i$ increases, apply the deformation of the graph $A_{(l+i-1)/2l}$ as in the passage of Figure 21 (where \odot and \square denote the intersection of the braid b and the page S_t) for $|\varepsilon_i|$ times in a small neighborhood of v_k and v_{k+1} . We call the surface that the graph traces out Σ_i . The surface Σ_i satisfies the above conditions (1)–(5) and the open book foliation $\mathcal{F}_{\text{ob}}(\Sigma_i)$ has $|\varepsilon_i|$ hyperbolic singularities of $\text{sgn}(\varepsilon_i)$.

Case 2-1 Suppose that $b_i = \rho_k$ and $\varepsilon_i = 1$. Let H be the chamber that ρ_k belongs to.

Assume that $r \leq k \leq 2g + r - 1$ so that H is a torus with connected boundary. For simplicity, put $u = A_{(l+i-1)/2l}$. By conditions (3), (4) above, we may assume that $u \cap H$ is some (p, q) -torus link. As $t \in I_i$ increases, move the point x_n along ρ_k . See Figure 22.

To come back to the original position x_n has to traverse u , which yields $p = [u] \cdot [\rho_k]$ many negative ($= -\text{sgn}(\varepsilon_i)$) hyperbolic points. Moreover, the last step (Sketch (4))




Figure 21: (Case 1) Deformation of graph $v_k \cup v_{k+1}$ corresponding to σ_k (top) and σ_k^{-1} (bottom)




Figure 22: (Case 2-1) Construction of surface Σ_i when $b_i^{\epsilon_i} = \rho_k$

adds one more hyperbolic singularity of positive ($= \text{sgn}(\epsilon_i)$) sign. This defines the surface Σ_i in $S \times I_i$. In summary, the value $h_+ - h_-$ increases by

$$\text{sgn}(\epsilon_i) \cdot 1 - [u] \cdot [\rho_k]$$

and the class $[u] \in H_1(S, \partial S)$ is replaced by $[u] + [\rho_k]$ (compare Sketches (1) and (4)). No circle bounding a disc in S has been created.

When $k = 1, \dots, r - 1$ (ie the chamber H is an annulus), a parallel argument holds and the value $h_+ - h_-$ increases by $\text{sgn}(\epsilon_i) \cdot 1 - [u] \cdot [\rho_k]$.

Case 2-2 If $b_i = \rho_k$ and $\varepsilon_i \neq 1, 0$, repeat the above construction $|\varepsilon_i|$ times. Since $([u] + [\rho_k]) \cdot [\rho_k] = [u] \cdot [\rho_k]$, the total change in $h_+ - h_-$ is

$$(3-4) \quad \begin{aligned} \varepsilon_i - \varepsilon_i[u] \cdot [\rho_k] &\stackrel{(5)}{=} \varepsilon_i - \varepsilon_i([A_{1/2}] + \varepsilon_1[b_1] + \cdots + \varepsilon_{i-1}[b_{i-1}]) \cdot [b_i] \\ &= \varepsilon_i - \left(\sum_{j=1}^{i-1} \varepsilon_i \varepsilon_j [b_j] \cdot [b_i] \right) - \varepsilon_i [A_{1/2}] \cdot [b_i]. \end{aligned}$$

After constructing $\Sigma_1, \dots, \Sigma_l$, we glue them and obtain a desired surface Σ_{**} in $S \times [1/2, 1]$, which increases the algebraic count of the hyperbolic singularities by

$$h_+ - h_- = \sum_{i=1}^l \varepsilon_i - \left(\sum_{i=1}^l \sum_{j=1}^{i-1} \varepsilon_i \varepsilon_j [b_j] \cdot [b_i] \right) - [A_{1/2}] \cdot [b].$$

Finally, we glue Σ_* and Σ_{**} by identifying $\Sigma_* \cap S_{1/2} = -(\Sigma_{**} \cap S_{1/2})$ and $-(\Sigma_* \cap S_0) = \phi(\Sigma_{**} \cap S_1)$, which yields a Seifert surface Σ for \hat{b} in the open book (S, ϕ) . By the construction, it is clear that $y_1, \dots, y_n \in \gamma_0$ are positive elliptic points, and the end points of arc ρ'_i are elliptic points with distinct signs. By Proposition 3.2, we obtain our self-linking formula (3-3). □

3.3 Properties of the function c

In this section we study properties of the function c in the self linking number formula (3-3). We will use the properties repeatedly in the later sections to deduce algebraic descriptions of the function c , which is originally defined geometrically.

Proposition 3.12 Take $[\phi], [\psi] \in \mathcal{MCG}(S)$ to be the mapping classes of $\phi, \psi \in \text{Aut}(S, \partial S)$, and let $a, a' \in H_1(S, \partial S)$. We have:

- (1) $c([\phi], a + a') = c([\phi], a) + c([\phi], a')$
- (2) $c([\psi\phi], a) = c([\phi], a) + c([\psi], \phi_*(a))$
- (3) Let C be a simple closed curve which does not intersect the walls, let T_C denote the right-handed Dehn twist along C , then $c([T_C], a) = 0$ for any a .
- (4) Let C be a simple closed curve in S such that $a \cdot [C] = 0$, then $c([T_C], a) = 0$.

In particular, (1) and (2) imply that the function c induces a crossed homomorphism

$$C: \mathcal{MCG}(S) \rightarrow \text{Hom}(H_1(S, \partial S), \mathbb{Z}) \simeq H^1(S; \mathbb{Z}), \quad \phi \mapsto c([\phi], -).$$

Proof First we prove (1). Let $\phi N(a) \xrightarrow{\Sigma} N(\phi_*a)$ and $\phi N(a') \xrightarrow{\Sigma'} N(\phi_*a')$ be OB cobordisms. We place the surfaces Σ and Σ' so that

- $\phi N(a)$ and $\phi N(a')$ in S_0 have the minimal geometric intersection (ie, so do $N(a)$ and $N(a')$),
- $N(\phi_*a)$ and $N(\phi_*a')$ in S_1 have the minimal geometric intersection.

Let $H \subset S$ be one of the once-punctured torus chambers. Then $H \cap N(a)$ and $H \cap N(a')$ are oriented torus links. Suppose that $H \cap N(a)$ is a (p, q) -torus link and $H \cap N(a')$ is a (p', q') -torus link. Then $H \cap N(a+a')$ is a $(p+p', q+q')$ -torus link. By isotopy, we arrange the curves $H \cap N(a)$ and $H \cap N(a')$ realizing the minimum geometric intersection, hence in particular, they transversely intersect. We resolve all the intersection points as shown in Figure 23, and call the resulting multi-curve $A_{H,a,a'}$.




Figure 23: Smoothing an intersection

Note that $[A_{H,a,a'}] = [H \cap N(a+a')]$ in $H_1(S, \partial S)$. We compare curves $A_{H,a,a'}$ and $H \cap N(a+a')$:

- (i) Suppose that

$$(\text{sgn}(p), \text{sgn}(q)) = (-\text{sgn}(p'), -\text{sgn}(q')).$$

If $n = \min\{|p|, |q|, |p'|, |q'|\}$, then $A_{H,a,a'}$ is the disjoint union of $H \cap N(a+a')$, n circles bounding concentric discs oriented counterclockwise, and n circles bounding concentric discs oriented clockwise. Removing the circles as shown in Figure 17 yields n negative and n positive hyperbolic points. Hence we obtain an OB cobordism

$$H \cap N(a+a') \xrightarrow{\Sigma_{H,a,a'}} A_{H,a,a'}$$

with $d(\Sigma_{H,a,a'}) = n - n = 0$.

- (ii) Suppose that $(\text{sgn}(p), \text{sgn}(q)) \neq (-\text{sgn}(p'), -\text{sgn}(q'))$. In this case, we have $A_{H,a,a'} = H \cap N(a+a')$. Hence we obtain a trivial OB cobordism

$$H \cap N(a+a') \xrightarrow{\Sigma_{H,a,a'}} A_{H,a,a'}$$

with $d(\Sigma_{H,a,a'}) = 0$.

Next let $H \subset S$ be the k^{th} annulus chamber. Recall the properly embedded arc $\rho'_k \subset H$ joining the boundary circles γ_0 and γ_k (cf Figure 19). Due to the definition of normal forms we may suppose that $H \cap N(a) = n\rho'_k$ and $H \cap N(a') = n'\rho'_k$. Then $H \cap N(a + a') = (n + n')\rho'_k$.

- (iii) If $\text{sgn}(n) = \text{sgn}(n')$, let $A_{H,a,a'} := (H \cap N(a)) \sqcup (H \cap N(a'))$. Then $A_{H,a,a'} = H \cap N(a + a')$. Again we obtain a trivial OB cobordism

$$H \cap N(a + a') \xrightarrow{\Sigma_{H,a,a'}} A_{H,a,a'}$$

with $d(\Sigma_{H,a,a'}) = 0$.

- (iv) If $\text{sgn}(n) \neq \text{sgn}(n')$, join $N(a)$ and $N(a')$ by describing arcs from the nearest pairs of ρ'_k and $-\rho'_k$ to introduce $m = \min\{|n|, |n'|\}$ hyperbolic singularities of the same sign, equal to ε . See Figure 24. Call the resulting set of curves $A_{H,a,a'}$. Then $A_{H,a,a'}$ is the disjoint union of $H \cap N(a + a')$ and null-homologous nested arcs. This yields an OB cobordism

$$H \cap N(a + a') \xrightarrow{\Sigma_{H,a,a'}} A_{H,a,a'}$$

with $d(\Sigma_{H,a,a'}) = \varepsilon m$.




Figure 24: Case (iv): (Left) Curves $N(a) \sqcup N(a')$. (Right) $A_{H,a,a'}$.

Let

$$A_0 = \phi \left(\bigsqcup_{H \subset S} A_{H,a,a'} \right), \quad \Sigma_0 = (\phi \times \text{id}) \left(\bigsqcup_{H \subset S} \Sigma_{H,a,a'} \right),$$

where the disjoint union is taken for all the $g + r - 1$ chambers H of S . Now we obtain an OB cobordism

$$\phi N(a + a') \xrightarrow{\Sigma_0} A_0 \quad \text{with} \quad d(\Sigma_0) = \sum_H d(\Sigma_{H,a,a'}).$$

We repeat the arguments parallel to (i)–(iv) by replacing a by $\phi_* a$ and a' by $\phi_* a'$. Namely, for each chamber H we construct a multi-curve $A_{H,\phi_* a, \phi_* a'}$ from

$$H \cap (N(\phi_* a) \cup N(\phi_* a'))$$

and obtain an OB cobordism

$$A_{H, \phi_* a, \phi_* a'} \xrightarrow{\Sigma_{H, \phi_* a, \phi_* a'}} H \cap N(\phi_*(a + a')).$$

Let $A_1 = \bigsqcup_{H \subset S} A_{H, \phi_* a, \phi_* a'}$, $\Sigma_1 = \bigsqcup_{H \subset S} \Sigma_{H, \phi_* a, \phi_* a'}$. Then we obtain an OB cobordism

$$A_1 \xrightarrow{\Sigma_1} N(\phi_*(a + a')) \quad \text{with } d(\Sigma_1) = \sum_H d(\Sigma_{H, \phi_* a, \phi_* a'}).$$

Claim 3.13 We have $d(\Sigma_1) = -d(\Sigma_0)$.

Proof For cases (i), (ii), (iii), we have $d(\Sigma_{H, \phi_* a, \phi_* a'}) = 0$. For case (iv), ie, H is the k^{th} annulus chamber, since $\phi = \text{id}$ near ∂S , we have $H \cap N(\phi_* a) = H \cap N(a) = n\rho'_k$ and $H \cap N(\phi_* a') = H \cap N(a') = n'\rho'_k$. Therefore, the OB cobordism $\Sigma_{H, \phi_* a, \phi_* a'}$ is given by the reverse direction as depicted in Figure 24. Recalling that $d(\Sigma_{H, a, a'}) = \varepsilon m$, we have $d(\Sigma_{H, \phi_* a, \phi_* a'}) = -\varepsilon m$. This concludes the claim. \square

Next we construct an OB cobordism $A_0 \xrightarrow{\Sigma^+} A_1$. Recall that the OB cobordism surfaces Σ and Σ' are obtained by a sequence of configuration changes (cf Figure 16). In general, a describing arc, δ , of a hyperbolic singularity on Σ may intersect Σ' (or vice versa) as shown in the top left sketch of Figure 25, where the black arc (resp. gray arcs) are leaves of Σ (resp. Σ'), the dashed arc is δ , and the dashed arrows indicate positive normal directions of the surfaces.




Figure 25: Modification of configuration changes

By isotopy, we make δ, Σ, Σ' have no triple intersection points and δ and Σ' attain the minimal geometric intersection. We project δ to the diagram of A_0 then replace

δ with several describing arcs for A_0 as in the vertical left passage in Figure 25 so that the diagram commutes. If the sign of original δ is ε , then the algebraic count of the replacing describing arcs is also ε . This modification of configuration changes yields an OB cobordism $A_0 \xrightarrow{\Sigma^+} A_1$. By the construction of Σ^+ , we have $d(\Sigma^+) = d(\Sigma) + d(\Sigma')$.

Finally we obtain a sequence of OB cobordisms

$$\phi N(a + a') \xrightarrow{\Sigma_0} A_0 \xrightarrow{\Sigma^+} A_1 \xrightarrow{\Sigma_1} N(\phi_*(a + a')).$$

By Claim 3.13,

$$c([\phi], a + a') = d(\Sigma_0) + d(\Sigma^+) + d(\Sigma_1) = d(\Sigma) + d(\Sigma') = c([\phi], a) + c([\phi], a').$$

We proceed to prove (2). Let $\phi N(a) \xrightarrow{\Sigma} N(\phi_*(a))$ be an OB cobordism. Extend $\psi \in \mathcal{MCG}(S, \partial S)$ to a diffeomorphism $\tilde{\psi} = \psi \times \text{id}: S \times [0, 1] \rightarrow S \times [0, 1]$ and we obtain an OB cobordism

$$\psi \phi(N(a)) \xrightarrow{\tilde{\psi} \Sigma} \psi(N(\phi_*(a))).$$

Now let us take an OB cobordism $\psi(N(\phi_*(a))) \xrightarrow{\Theta} N(\psi_*\phi_*(a))$. Gluing $\tilde{\psi} \Sigma$ and Θ , we obtain an OB cobordism

$$\psi \phi(N(a)) \xrightarrow{\tilde{\psi} \Sigma} \xrightarrow{\Theta} N(\psi_*\phi_*(a)).$$

Since $\tilde{\psi}$ preserves the signs and the number of hyperbolic singularities, $d(\Sigma) = d(\tilde{\psi} \Sigma)$. This yields the desired equation.

To see (3), we observe that if a simple closed curve C does not intersect the walls, then $T_C(N(a))$ is in the normal form for any $a \in H_1(S, \partial S)$, ie, $T_C(N(a)) = N(T_C a)$. Consider the product $\Sigma = T_C(N(a)) \times I$, which yields the trivial OB cobordism $T_C(N(a)) \xrightarrow{\Sigma} N(T_C a)$. Since the foliation is trivial, $c([T_C], a) = 0$.

Finally, we prove (4). We construct an OB cobordism $T_C N(a) \xrightarrow{\Sigma} N(a) = N(T_C a)$ with $d(T_C N(a), N(a)) = 0$ as follows. Since $[C] \cdot [T_C N(a)] = [C] \cdot a = 0$, by applying the configuration changes, described in Figure 16, to a portion of the multi-curve $T_C N(a)$ that lives in a small collar neighborhood of C , we can modify $T_C N(a)$ so that it is disjoint from C . For example, the left sketch in Figure 26 depicts the case when the geometric intersection number $i(T_C N(a), C) = 2$, where the thin dashed arcs indicate describing arcs for hyperbolic singularities. Suppose that the sign of the hyperbolic singularity corresponding to the configuration change is ε . Next, we add a describing arc of sign $-\varepsilon$ to the deformed $T_C N(a)$ (cf the middle sketch) so that the corresponding configuration change yields the multi-curve $N(a)$ (cf the

right sketch). This defines an OB cobordism $T_C N(a) \xrightarrow{\Sigma} N(a)$, which satisfies $d(T_C N(a), N(a)) = 0$.

When the geometric intersection number is greater than 2, a similar construction applies. In particular, the sum of the total algebraic count of the signs in the first operation and the second operation is 0. □




Figure 26: Untwisting multi-curve $T_C N(a)$ (left) to obtain $N(a)$ (right)

3.4 The function c : Planar surface case

In this section, we study the function c for the case of $S = S_{0,r}$, a planar surface with r boundary components. We adopt the same notation as in Section 3.2. The next proposition essentially has been proved in [32] by direct analysis of the OB cobordism (though this terminology is not explicitly used). Based on the fact that c is a crossed homomorphism, we will give a more detailed expression of c .

Recall the arcs ρ'_j and loops ρ_j ($j = 1, \dots, r - 1$) specified in Figure 19. Under Poincaré duality, $H_1(S, \partial S; \mathbb{Z}) \simeq H^1(S; \mathbb{Z})$, $[\rho'_j] \mapsto \text{PD}[\rho'_j]$, we may view $\{[\rho'_j]\}_{j=1}^{r-1}$ as a basis of $H^1(S)$. Let $\langle \cdot, \cdot \rangle$ denote the natural pairing of cohomology and homology. Then we have $\langle [\rho'_i], [\rho_j] \rangle = [\rho'_i] \cdot [\rho_j] = \delta_{i,j}$, the Kronecker delta.

Proposition 3.14 *Let $S = S_{0,r}$ be a planar surface with r boundary components. For $a \in H_1(S, \partial S; \mathbb{Z})$, the function c is formulated as*

$$(3-5) \quad c([\phi], a) = \sum_{i=1}^{r-1} \langle [\rho'_i], \phi_* a - a \rangle - \sum_{j=1}^{r-1} \langle a, [\rho_j] \rangle \langle [\rho'_j], \phi_* [\rho'_j] - [\rho'_j] \rangle$$

where $\phi_* a - a$ and $\phi_* [\rho'_i] - [\rho'_i]$ are regarded as elements of $H_1(S; \mathbb{Z})$. Moreover, let $\{t_{i,j}\}_{1 \leq i, j \leq r-1}$ be the matrix with $[\rho'_i] - \phi_* [\rho'_i] = \sum_{j=1}^{r-1} t_{i,j} [\rho_j]$ and suppose that $a = \sum_{j=1}^{r-1} x_j [\rho'_j]$. Then (3-5) can be restated as follows:

$$(3-6) \quad c([\phi], a) = - \sum_{j=1}^{r-1} x_j \sum_{\substack{1 \leq i \leq r-1 \\ i \neq j}} t_{j,i}$$

Remark 3.15 For the planar case, Proposition 3.14 shows that the crossed homomorphism \mathcal{C} or $c([\phi], -)$ is completely determined by the map $\phi_* - \text{id}: H_1(S, \partial S) \rightarrow H_1(S)$.

Proof For $j = 1, \dots, r - 1$, we have

$$(3-7) \quad c([\phi], [\rho'_j]) = \sum_{\substack{1 \leq i \leq r-1 \\ i \neq j}} \langle [\rho'_i], \phi_*[\rho'_j] - [\rho'_j] \rangle$$

for the following reasons. We recall that $c([\phi], [\rho'_j])$ algebraically counts the hyperbolic singularities produced by the configuration changes (cf Figure 16) of the multi-curve $\phi(\rho'_j)$ where it crosses the walls. We write ϕ as a product of special type of Dehn twists that are used in [32] and denoted by $A_{k,l}, A_m$ there. We observe that a \pm Dehn twist that involves the i^{th} and j^{th} binding components ($i \neq j$) contributes ± 1 hyperbolic singularity for the OB cobordism $\phi(\rho'_j) \xrightarrow{\Sigma} N(\phi_*[\rho'_j])$. But a Dehn twist around a single binding component γ_k , where $(k = 1, \dots, r - 1)$, does not contribute any hyperbolic singularity to the OB cobordism. Since the quantity $\langle [\rho'_i], \phi_*[\rho'_j] - [\rho'_j] \rangle$ algebraically counts the number of circles in $N(\phi_*[\rho'_j])$ around the binding γ_i , Equation (3-7) follows.

Recall that $\{[\rho_i] \in H_1(S)\}_{i=1}^{r-1}$ is the dual basis of $\{[\rho'_i] \in H_1(S, \partial S)\}_{i=1}^{r-1}$. We may express $a \in H_1(S, \partial S) \simeq H^1(S)$ as $a = \sum_j \langle a, [\rho_j] \rangle [\rho'_j]$. By the crossed homomorphism property of the function c (Proposition 3.12), we can deduce (3-5) as follows:

$$\begin{aligned} c([\phi], a) &= \sum_j \langle a, [\rho_j] \rangle c([\phi], [\rho'_j]) \stackrel{(3-7)}{=} \sum_j \sum_{i \neq j} \langle [\rho'_i], \langle a, [\rho_j] \rangle (\phi_*[\rho'_j] - [\rho'_j]) \rangle \\ &= \sum_i \langle [\rho'_i], \phi_* a - a \rangle - \sum_j \langle a, [\rho_j] \rangle \langle [\rho'_j], \phi_*[\rho'_j] - [\rho'_j] \rangle \end{aligned}$$

Now plugging the relation $[\rho'_j] - \phi_*[\rho'_j] = \sum_{i=1}^{r-1} t_{j,i} [\rho_i]$ to (3-7) we obtain

$$(3-8) \quad c([\phi], [\rho'_j]) = - \sum_{\substack{1 \leq i \leq r-1 \\ i \neq j}} t_{j,i}.$$

Linearly extending (3-8) to an arbitrary element $a = \sum_j x_j [\rho'_j]$, we obtain (3-6). \square

Remark 3.16 Since $\phi_*[\rho'_j] = [\rho'_j] \in H_1(S, \partial S)$, we have $c([\psi], \phi_*[\rho'_j]) = c([\psi], [\rho'_j])$. Therefore, when S is planar the property (2) in Proposition 3.12 can be restated as

$$c([\psi\phi], [\rho'_j]) = c([\phi], [\rho'_j]) + c([\psi], [\rho'_j]).$$

By using Theorem 3.10 and Proposition 3.14, now we can deduce the self-linking number formulae in [33; 32]. Let a_σ (resp. a_{ρ_j}) be the exponent sum of the braid generators $\{\sigma_i\}_{i=1}^{n-1}$ (resp. ρ_j) in the braid word $b = b_1^{\varepsilon_1} b_2^{\varepsilon_2} \cdots b_l^{\varepsilon_l}$. Let $a = \sum_{i=1}^{r-1} s_i [\rho'_i] \in H_1(S, \partial S)$, the homology class introduced in the proof of Claim 3.8 such that $[b] = a - \phi_* a$.

Corollary 3.17 (The self-linking number formula for planar open books [32]) *With the notation above, the self-linking number is given by the formula*

$$sl(\widehat{b}, [\Sigma]) = -n + a_\sigma + \sum_{j=2}^r a_{\rho_j} (1 - s_j) - \sum_{j=1}^{r-1} s_j \sum_{\substack{1 \leq i \leq r-1 \\ i \neq j}} t_{j,i}.$$

Proof Since $[b_i] \cdot [b_j] = 0$ for all $b_i, b_j \in \{\rho_1, \dots, \rho_{r-1}, \sigma_1, \dots, \sigma_{n-1}\}$, we have

$$\widehat{\text{exp}}(b) = \sum_{i=1}^l \varepsilon_i = a_\sigma + \sum_{j=1}^{r-1} a_{\rho_j},$$

and since $[\rho'_j] \cdot [\rho_k] = \delta_{j,k}$, we have

$$\phi_*(a) \cdot [b] = (a - [b]) \cdot [b] = \left(\sum_{j=1}^{r-1} s_j [\rho'_j] \right) \cdot \left(\sum_{i=1}^l \varepsilon_i [b_i] \right) = \sum_{j=1}^{r-1} a_{\rho_j} s_j.$$

Hence by Theorem 3.10 and Proposition 3.14, we have

$$\begin{aligned} sl(\widehat{b}, \Sigma) &= -n + \widehat{\text{exp}}(b) - \phi_*(a) \cdot [b] + c([\phi], a) \\ &= -n + a_\sigma + \sum_{j=1}^{r-1} a_{\rho_j} (1 - s_j) - \sum_{j=1}^{r-1} s_j \sum_{\substack{1 \leq i \leq r-1 \\ i \neq j}} t_{j,i}. \end{aligned}$$

□

3.5 The function c : surface with connected boundary

Let $S = S_{g,1}$ be a genus g surface with one boundary component. When $g = 1$, since there is no wall, Proposition 3.12(3) implies that $c([\phi], a) = 0$ for all $[\phi] \in \mathcal{MCG}(S_{1,1})$ and $a \in H_1(S, \partial S)$. Henceforth in this section we restrict our attention to the case $g \geq 2$.

We observe in the following example that, unlike the planar case discussed in Remark 3.15, the function $c([\phi], -)$ is no longer completely determined by the action of ϕ_* on homologies. In fact we see in Proposition 3.20 that $c([\phi], -)$ carries more delicate information about $[\phi] \in \mathcal{MCG}(S)$.




Figure 27: Curves C , C' and ρ in Example 3.18

Example 3.18 Let us take simple closed curves C , C' and ρ as in Figure 27.

Since C and C' cobound a subsurface, T_C and $T_{C'}$ induce the same action on the homology groups $H_1(S; \mathbb{Z})$ and $H_1(S, \partial S; \mathbb{Z})$. As shown in Figure 28, we modify the curve $T_C(\rho)$ into the normal form $N(T_C(\rho))$ by introducing three positive hyperbolic singularities and one negative hyperbolic singularity. Hence $c([T_C], [\rho]) = 3 - 1 = 2$. On the other hand, C' does not intersect the walls, so by Proposition 3.12(3), we get $c([T_{C'}], [\rho]) = 0$.




Figure 28: Configuration change of $T_C(\rho)$ to the normal form $N(T_C(\rho))$

In this section we use the following notation: Recall the circles $\rho_j, \rho'_j \subset S$ defined in Section 3.2. To distinguish elements of $H_1(S; \mathbb{Z})$ and $H_1(S, \partial S; \mathbb{Z}) \cong H^1(S; \mathbb{Z})$, we use the symbol $[\rho_j]$ ($j = 1, \dots, 2g$) to express the homology class of $H_1(S)$ represented by the circle ρ_j , and the symbol $[\rho'_j]$ for the relative homology class of

$H_1(S, \partial S) \cong H^1(S)$ represented by the circle ρ'_j . Note that since S has connected boundary, $\rho_j = \rho'_j$ as a set for all j , and as a group $H_1(S; \mathbb{Z}) \cong H_1(S, \partial S; \mathbb{Z}) \cong \mathbb{Z}^{2g}$.

Let $\langle \cdot, \cdot \rangle: H_1(S, \partial S; \mathbb{Z}) \times H_1(S; \mathbb{Z}) \rightarrow \mathbb{Z}$ denote the natural pairing of cohomology and homology, or the intersection pairing, ie, $\langle \text{PD}[\rho'_j], [\rho_k] \rangle = [\rho'_j] \cdot [\rho_k]$. For simplicity, we denote $\text{PD}[\rho'_j]$ by $[\rho'_j]$ in the following. We have

$$\langle [\rho'_j], [\rho_k] \rangle = \begin{cases} 1 & \text{if } (j, k) = (2i - 1, 2i), \\ -1 & \text{if } (j, k) = (2i, 2i - 1), \\ 0 & \text{otherwise.} \end{cases}$$

Let $\Gamma_1 := \pi_1(S)$, the fundamental group, $\Gamma_2 := [\Gamma_1, \Gamma_1]$, the commutator subgroup, and $\Gamma_{k+1} := [\Gamma_k, \Gamma_1]$, namely, $\{\Gamma_k\}_{k \geq 1}$ is the lower central series of Γ_1 . Then the natural action of $\mathcal{MCG}(S_{g,1})$ on Γ_1 induces the k^{th} Johnson–Morita representation (Morita [38, page 199]):

$$\varrho_k: \mathcal{MCG}(S_{g,1}) \rightarrow \text{Aut}(\Gamma_1/\Gamma_k), \quad k \geq 2$$

Let $\mathcal{M}(k) := \ker \varrho_k$ and $H := H_1(S; \mathbb{Z})$. Morita generalizes the Johnson homomorphism $\tau_2: \mathcal{M}(2) \rightarrow \text{Hom}(H, \Gamma_2/\Gamma_3)$ to the k^{th} Johnson–Morita homomorphism [38, page 201]:

$$\tau_k: \mathcal{M}(k) \rightarrow \text{Hom}(H, \Gamma_k/\Gamma_{k+1})$$

with $\ker \tau_k = \mathcal{M}(k + 1)$. Let \mathcal{K} be the subgroup of $\mathcal{MCG}(S_{g,1})$ generated by the Dehn twists about separating simple closed curves in S . Johnson proves in [31] that for $g \geq 3$, we can identify $\ker \tau_2 = \mathcal{K}$. Recall that by Proposition 3.12(2), (4) our crossed homomorphism $\mathcal{C}: \mathcal{MCG}(S) \rightarrow H^1(S; \mathbb{Z})$ also vanishes on \mathcal{K} . Hence it is natural to expect that the map \mathcal{C} is related to τ_2 .

Associated to the representation $\varrho_3: \mathcal{MCG}(S_{g,1}) \rightarrow \text{Aut}(\Gamma_1/\Gamma_3)$, Morita [38] finds the embedding $\mathcal{MCG}(S_{g,1})/\mathcal{M}(3) \subset \frac{1}{2} \wedge^3 H \rtimes \text{Sp}(H)$ as a finite index subgroup and the crossed homomorphism $\tilde{k}: \mathcal{MCG}(S_{g,1}) \rightarrow \frac{1}{2} \wedge^3 H$, which is the unique (modulo coboundaries for $H^1(\mathcal{MCG}(S_{g,1}), \wedge^3 H)$) extension of τ_2 . For our purpose we are interested in the composition

$$k = C \circ \tilde{k}: \mathcal{MCG}(S_{g,1}) \rightarrow H^1(S; \mathbb{Z}),$$

where $C: \frac{1}{2} \wedge^3 H \rightarrow H$ is the contraction defined by

$$C(x \wedge y \wedge z) = 2[(x \cdot y)z + (y \cdot z)x + (z \cdot x)y].$$

The associated map, which we denote by the same letter, $k: \mathcal{MCG}(S_{g,1}) \times H_1(S) \rightarrow \mathbb{Z}$, given by $k(\phi, a) = k(\phi)(a)$, is a crossed homomorphism. Since k is a generator of

the cohomology group $H^1(\mathcal{MCG}(S_{g,1}); H) \cong \mathbb{Z}$ [38, Rem 4.9], it is natural to expect that k appears in the description of $c(\phi, a)$.

Below we fix conventions and define the crossed homomorphism k following Morita's [36, Section 6] that is based on combinatorial group theory.

Definition 3.19 Let F_2 be the free group of rank two with generators α and β . Any element of F_2 is uniquely written in the form $\alpha^{\epsilon_1} \beta^{\delta_1} \dots \alpha^{\epsilon_n} \beta^{\delta_n}$, where $\epsilon_i, \delta_i \in \{-1, 0, 1\}$. With this expression, we define a function $d: F_2 \rightarrow \mathbb{Z}$ by

$$d(\alpha^{\epsilon_1} \beta^{\delta_1} \dots \alpha^{\epsilon_n} \beta^{\delta_n}) = \sum_{i=1}^n \delta_i \sum_{j=1}^i \epsilon_j.$$

Let α_i, β_i ($i = 1, \dots, g$) be generating curves of $\pi_1(S)$ as in Figure 29.




Figure 29

Let $p_i: \pi_1(S) \rightarrow F_2$ be a homomorphism defined by

$$p_i(\gamma) = \begin{cases} \alpha & \text{if } \gamma = \alpha_i, \\ \beta & \text{if } \gamma = \beta_i, \\ 1 & \text{otherwise.} \end{cases}$$

Finally we define a map $k: \mathcal{MCG}(S_{g,1}) \times H_1(S, \partial S) \rightarrow \mathbb{Z}$ by

$$k([\phi], a) = \sum_{i=1}^g d(p_i(\phi_* a)) - d(p_i(a)),$$

where $a \in \pi_1(S)$ represents $a \in H_1(S, \partial S)$. Morita proves in [36, Lemma 6.3] that $k([\phi], a)$ is a crossed homomorphism.

We give an explicit formula of the function c by using k . It provides a new geometric meaning of the classically known crossed homomorphism k : the signed count of the saddle points in an OB cobordism.

Proposition 3.20 *If $S = S_{g,1}$ has connected boundary and $g \geq 2$, then the function c is expressed as*

$$\begin{aligned}
 (3-9) \quad c([\phi], a) = & -2k([\phi], a) + \sum_{i=1}^g \langle [\rho'_{2i-1}] - [\rho'_{2i}], \phi_* a - a \rangle \\
 & - \sum_{i=1}^g \langle a, [\rho_{2i-1}] \rangle \langle [\rho'_{2i}], \phi_* [\rho'_{2i}] - [\rho'_{2i}] \rangle \\
 & - \sum_{i=1}^g \langle a, [\rho_{2i}] \rangle \langle [\rho'_{2i-1}], \phi_* [\rho'_{2i-1}] - [\rho'_{2i-1}] \rangle,
 \end{aligned}$$

where $\phi_* a - a$ and $\phi_* [\rho'_j] - [\rho'_j]$ are regarded as elements of $H_1(S; \mathbb{Z})$.

Proof Recall that the left hand side of (3-9) satisfies the crossed homomorphism properties (1), (2) in Proposition 3.12. Hence it is sufficient to verify (3-9) for a generating set of the mapping class group $\mathcal{MCG}(S_{g,1})$.

We use the Lickorish generators of $\mathcal{MCG}(S_{g,1})$. Let A_i, B_i ($i = 1, \dots, g$) and C_j ($j = 1, \dots, g - 1$) be simple closed curves as shown in Figure 30.




Figure 30: Generating curves for $\mathcal{MCG}(S_{g,1})$

Lickorish proved that the Dehn twists along these $3g - 1$ curves generate $\mathcal{MCG}(S_{g,1})$. With the orientations indicated in Figure 30, we have in $H_1(S; \mathbb{Z})$ that

$$[A_i] = [\rho_{2i-1}], \quad [B_i] = [\rho_{2i}] \quad \text{and} \quad [C_i] = -[\rho_{2i-1}] + [\rho_{2i+1}].$$

If $D \in \{A_i, B_i, C_i\}$ is disjoint from the loop ρ_j , then $c([T_D], [\rho'_j]) = k([T_D], [\rho'_j]) = 0$ and $T_{D*}[\rho'_j] - [\rho'_j] = 0$; thus the formula (3-9) holds. So we only need to consider the case where D has non-trivial intersection with ρ_j . There are four cases to study:

Case I $(\phi, a) = (T_{A_i}, [\rho'_{2i}])$ Since A_i is disjoint from the walls, Proposition 3.12(3) implies that $c([T_{A_i}], [\rho'_{2i}]) = 0$. On the other hand, $T_{A_i}(\rho_{2i}) = \rho_{2i-1}\rho_{2i} = \beta_{g-i}\alpha_{g-i}^{-1}$

in $\pi_1(S)$, hence $k([T_{A_i}], [\rho'_{2i}]) = d(\beta\alpha^{-1}) = 0$. Finally, observe that $T_{A_i*}[\rho'_{2i}] - [\rho'_{2i}] = [\rho_{2i-1}]$, hence

$$\begin{aligned}
 (\star) &:= \sum_{k=1}^g \langle [\rho'_{2k-1}] - [\rho'_{2k}], \phi_*a - a \rangle = \langle -[\rho'_{2i}], [\rho_{2i-1}] \rangle = 1, \\
 (\star\star) &:= \sum_{k=1}^g \langle a, [\rho_{2k-1}] \rangle \langle [\rho'_{2k}], \phi_*[\rho'_{2k}] - [\rho'_{2k}] \rangle \\
 &\quad + \sum_{k=1}^g \langle a, [\rho_{2k}] \rangle \langle [\rho'_{2k-1}], \phi_*[\rho'_{2k-1}] - [\rho'_{2k-1}] \rangle \\
 &= \langle [\rho'_{2i}], [\rho_{2i-1}] \rangle^2 = (-1)^2 = 1.
 \end{aligned}$$

Thus the equality (3-9) holds.

Case II $(\phi, a) = (T_{B_i}, [\rho'_{2i-1}])$ As in Case I, B_i is disjoint from the walls, so $c([T_{B_i}], [\rho'_{2i-1}]) = 0$. On the other hand, $T_{B_i}(\rho_{2i-1}) = \rho_{2i-1}\rho_{2i}^{-1} = \beta_{g-i}\alpha_{g-i}$, hence $k([T_{B_i}], [\rho'_{2i-1}]) = d(\beta\alpha) = 0$. Finally, observe that $T_{B_i*}[\rho'_{2i-1}] - [\rho'_{2i-1}] = -[\rho_{2i}]$, hence

$$\begin{aligned}
 (\star) &= \langle [\rho'_{2i-1}], -[\rho_{2i}] \rangle = -1, \\
 (\star\star) &= \langle [\rho'_{2i-1}], [\rho_{2i}] \rangle \langle [\rho'_{2i-1}], -[\rho_{2i}] \rangle = -1.
 \end{aligned}$$

Thus the equality (3-9) holds.

Case III $(\phi, a) = (T_{C_i}, [\rho'_{2i}])$ Observe that $c([T_{C_i}], [\rho'_{2i}]) = -1$. Since

$$T_{C_i}(\rho_{2i}) = \rho_{2i}\rho_{2i+1}^{-1}\rho_{2i}^{-1}\rho_{2i-1}\rho_{2i} = \alpha_{g-i}^{-1}\beta_{g-i-1}^{-1}\alpha_{g-i}\beta_{g-i}\alpha_{g-i}^{-1},$$

$k([T_{C_i}], [\rho'_{2i}]) = d(\beta\alpha^{-1}) + d(\beta^{-1}) = 0$. Finally, $T_{C_i*}[\rho'_{2i}] - [\rho'_{2i}] = [\rho_{2i-1}] - [\rho_{2i+1}]$, hence

$$\begin{aligned}
 (\star) &= 0, \\
 (\star\star) &= \langle [\rho'_{2i}], [\rho_{2i-1}] \rangle \langle [\rho'_{2i}], -[\rho_{2i+1}] + [\rho_{2i-1}] \rangle = (-1)^2 = 1.
 \end{aligned}$$

Thus the equality (3-9) holds.

Case IV $(\phi, a) = (T_{C_i}, [\rho'_{2i+2}])$ In this case, $c([T_{C_i}], [\rho'_{2i+2}]) = 1$ and

$$T_{C_i}(\rho'_{2i+2}) = \rho_{2i}^{-1}\rho_{2i-1}^{-1}\rho_{2i}\rho_{2i+1}\rho_{2i+2} = \alpha_{g-i}\beta_{g-i}\alpha_{g-i}^{-1}\beta_{g-i-1}\alpha_{g-i-1}^{-1}.$$

Hence $k([T_{C_i}], [\rho'_{2i+2}]) = d(\alpha\beta^{-1}\alpha^{-1}) + d(\beta\alpha^{-1}) = -1$. Finally,

$$T_{C_i*}[\rho'_{2i+2}] - [\rho'_{2i+2}] = -[\rho_{2i-1}] + [\rho_{2i+1}],$$

hence

$$(\star) = 0,$$

$$(\star\star) = \langle [\rho'_{2i+2}], [\rho_{2i+1}] \rangle \langle [\rho'_{2i+2}], -[\rho_{2i-1}] + [\rho_{2i+1}] \rangle = (-1)^2 = 1.$$

Thus the equality (3-9) holds. These computations complete the proof. \square

The map k appears in various contexts in the theory of mapping class groups (see Section 2 of [37] for concise overview). In particular, k can be interpreted in terms of winding numbers of curves on surfaces. Fixing a non-vanishing vector field X on S , one defines the *winding number* of an oriented simple closed curve γ on S as the rotation number of the tangent vector to γ with respect to X as γ is traversed once positively. Then $k(\phi, \gamma)$ is equal to the difference of winding numbers of $\phi(\gamma)$ and γ as stated in Definition 1.3.1 of Trapp's paper [43].

Recall that in (Step 1) near Figure 17 we have observed that a c -circle bounding a disc contributes ± 1 to the function $c([\phi], a)$. Such a disc also contributes ± 1 to the above winding number.

In addition, the self-linking number $\text{sl}(\gamma, [\Sigma])$ is the winding number of a nowhere vanishing section X of the vector bundle $\xi|_{\Sigma} \rightarrow \Sigma$ along γ relative to Σ , where Σ is a Seifert surface of γ .

Interestingly, the keyword of the above facts is “winding number”. The authors thank the anonymous referee for pointing this out.

Theorem 3.10 and Proposition 3.20 give a new relationship between the contact structures of 3-manifolds and the Johnson–Morita homomorphisms. This develops into the following question: Our result roughly says that if we choose a homology class $a \in \Gamma_1 / \Gamma_2 = H_1(S; \mathbb{Z})$ (from the geometric point of view, this choice corresponds to the choice of Seifert surface of the transverse link $L = \hat{b}$), then the Johnson–Morita representation $\varrho_3: \mathcal{MCG}(S_{g,1}) \rightarrow \text{Aut}(\Gamma_1 / \Gamma_3)$ gives the self-linking number. Now we ask whether a similar phenomenon occurs for the higher Johnson–Morita representation $\varrho_i: \mathcal{MCG}(S_{g,1}) \rightarrow \text{Aut}(\Gamma_1 / \Gamma_i)$, where $i > 3$, and provides new invariants of transverse links.

3.6 The function c : general surface case

Finally we give a complete description of the function c for general surfaces $S = S_{g,r}$. We use the same convention as in Section 3.5, that is, $[\rho'_j]$ is an element of $H_1(S, \partial S; \mathbb{Z}) \cong H^1(S; \mathbb{Z})$ and $[\rho_j]$ is an element of $H_1(S; \mathbb{Z})$. Let $S' = S_{g,1}$ be the surface obtained from $S = S_{g,r}$ by filling the boundary components $\gamma_1, \dots, \gamma_{r-1}$ by

discs and $i: S \rightarrow S'$ the canonical inclusion. Let $\pi: \mathcal{MCG}(S_{g,r}) \rightarrow \mathcal{MCG}(S_{g,1})$ be the forgetful map. Let us consider the pull-back $\pi^*k: \mathcal{MCG}(S_{g,r}) \times H_1(S, \partial S) \rightarrow \mathbb{Z}$ of the crossed homomorphism k defined by $\pi^*k: ([\phi], a) \mapsto k(\pi[\phi], i_*(a))$. For $1 \leq i \leq r - 1 + 2g$, let

$$[\zeta'_j] = \begin{cases} [\rho'_j] & \text{if } j = 1, \dots, r - 1, \\ -[\rho'_{j+1}] & \text{if } i = r, r + 2, \dots, r - 2 + 2g, \\ [\rho_{j-1}] & \text{if } i = r + 1, r + 3, \dots, r - 1 + 2g. \end{cases}$$

In particular, we have $\langle [\zeta'_i], [\rho_j] \rangle = \delta_{i,j}$. By combining Propositions 3.14 and 3.20, we get an explicit formula of the function c .

Theorem 3.21 (A formula of function c) *Let $S = S_{g,r}$ be the surface with genus g and r boundary components. The function $c: \mathcal{MCG}(S_{g,r}) \times H_1(S, \partial S; \mathbb{Z}) \rightarrow \mathbb{Z}$ has the expression*

$$c([\phi], a) = -2(\pi^*k)([\phi], a) + \sum_{j=1}^{2g+r-1} \langle [\zeta'_j], \phi_*a - a \rangle - \sum_{j=1}^{2g+r-1} \langle a, [\rho_j] \rangle \langle [\zeta'_j], \phi_*[\zeta'_j] - [\zeta'_j] \rangle,$$

where $\phi_*a - a$ and $\phi_*[\zeta'_j] - [\zeta'_j]$ are regarded as elements of $H_1(S; \mathbb{Z})$.

4 On the Bennequin–Eliashberg inequality

In this section using open book foliations we give a new proof to the Bennequin–Eliashberg inequality [16].

Recall that an *overtwisted disc* is an embedded disc whose boundary is a limit cycle in the characteristic foliation. Thus an overtwisted disc always has *Legendrian* boundary. As a corresponding notion in the framework of open book foliations, we introduce the following:

Definition 4.1 Let $D \subset M_{(S,\phi)}$ be an oriented disc whose boundary is a positively braided unknot. If the following are satisfied D is called a *transverse overtwisted disc*:

- (1) G_{--} (Definition 2.17) is a connected tree with no fake vertices.
- (2) G_{++} is homeomorphic to S^1 .
- (3) $\mathcal{F}_{\text{ob}}(D)$ contains no c -circles.

By Proposition 3.2, we observe that $\text{sl}(\partial D, [D]) = 1$ for a transverse overtwisted disc D .

Proposition 4.2 *If (S, ϕ) contains a transverse overtwisted disc then the compatible contact 3-manifold (M, ξ) contains an overtwisted disc.*

Proof By applying Theorem 2.21 and Giroux's elimination lemma (see [23, page 187]), we can convert a transverse overtwisted disc to an overtwisted disc. \square

We will prove the converse in Corollary 4.6, hence the existence of a transverse overtwisted disc is equivalent to the existence of a usual overtwisted disc.

Theorem 4.3 (The Bennequin–Eliashberg inequality [16]) *If a contact 3–manifold (M, ξ) is tight, then for any null-homologous transverse link L and its Seifert surface Σ , the following inequality holds:*

$$\text{sl}(L, [\Sigma]) \leq -\chi(\Sigma)$$

The following corollary was pointed out by John Etnyre and the proof is straightforward.

Corollary 4.4 *The following are equivalent:*

- (1) (M, ξ) is tight.
- (2) For any null-homologous transverse link L and its Seifert surface Σ we have $\text{sl}(L, [\Sigma]) \leq -\chi(\Sigma)$.
- (3) For any transverse unknot $U = \partial D$, we have $\text{sl}(U, [D]) \leq -\chi(D) = -1$.

We use the following Lemma 4.5 and Proposition 3.2 to prove Theorem 4.3.

Lemma 4.5 *Let L be a null-homologous transverse link in a contact 3–manifold (M, ξ) and Σ be a Seifert surface for L . Assume that $\text{sl}(L, [\Sigma]) > -\chi(\Sigma)$, that is, the Bennequin–Eliashberg inequality is violated. With some perturbation of Σ fixing the boundary we can make the graph G_{--} contain a contractible component with no fake vertices.*

Proof Using Propositions 2.11 and 3.2, we assume that

$$\text{sl}(L, [\Sigma]) + \chi(\Sigma) = 2(e_- - h_-) > 0,$$

ie, $e_- - h_- > 0$. Let $\Gamma_1, \dots, \Gamma_k$ denote the connected components of the graph G_{--} . Let $f(\Gamma_i)$ be the number of the fake vertices of Γ_i and $e_-(\Gamma_i)$ the number of the negative elliptic points in Γ_i . Let $h_-(\Gamma_i)$ be the number of the edges in Γ_i . By Proposition 2.6 with some perturbation of Σ fixing the boundary we may assume that $\mathcal{F}_{\text{ob}}(\Sigma)$ has no c –circles, hence the region decomposition (Proposition 2.15) does not contain type ac , bc or cc regions, so $e_- = \sum_{i=1}^k e_-(\Gamma_i)$ and $h_- = \sum_{i=1}^k h_-(\Gamma_i)$. Since Γ_i is connected, the Euler characteristic of Γ_i satisfies

$$\chi(\Gamma_i) = (f(\Gamma_i) + e_-(\Gamma_i)) - h_-(\Gamma_i) \leq 1,$$

ie $e_-(\Gamma_i) - h_-(\Gamma_i) \leq 1 - f(\Gamma_i)$. Therefore we obtain that $e_-(\Gamma_i) - h_-(\Gamma_i) = 1$ if and only if $f(\Gamma_i) = 0$ and Γ_i is contractible. Now we have

$$0 < e_- - h_- = \sum_{i=1}^k e_-(\Gamma_i) - \sum_{i=1}^k h_-(\Gamma_i) = \sum_{i=1}^k (e_-(\Gamma_i) - h_-(\Gamma_i)) \leq \sum_{i=1}^k (1 - f(\Gamma_i)).$$

Thus for some i , the equality $e_-(\Gamma_i) - h_-(\Gamma_i) = 1$ must hold, which implies that Γ_i is contractible and has no fake vertices. □

Now we are ready to prove Theorem 4.3. Eliashberg’s original proof of the Bennequin–Eliashberg inequality uses characteristic foliation theory. We give an alternative proof from the viewpoint of open book foliations.

Proof of Theorem 4.3 Suppose that there exists a null-homologous transverse link L in (M, ξ) with a Seifert surface Σ such that $sl(L, [\Sigma]) > -\chi(\Sigma)$. We will show that ξ is overtwisted.

Fix an open book (S, ϕ) that supports ξ and isotope L and Σ with the transverse link type of L , preserved until it admits an open book foliation $\mathcal{F}_{ob}(\Sigma)$. By Proposition 2.6 and Lemma 4.5, we may assume that $\mathcal{F}_{ob}(\Sigma)$ contains no c -circles and the negativity graph G_{--} contains a contractible component $\Gamma \subset G_{--}$ with no fake vertices. In particular, the induced region decomposition of Σ consists only of aa , ab , and bb -tiles.

Let $\mathcal{R} \subset \Sigma$ be the set of b -arcs that end on the vertices of Γ . Since Γ lives only in ab and bb -tiles and has no fake vertices, we have $\Gamma \subset \text{Int}(\overline{\mathcal{R}})$, where $\overline{\mathcal{R}}$ is the closure of \mathcal{R} . Let $\mathcal{P} = G_{++}(\overline{\mathcal{R}})$ be the set of positive elliptic points in $\overline{\mathcal{R}}$ and the stable separatrices approaching to the positive hyperbolic points in $\overline{\mathcal{R}}$. Since Γ is a tree with no fake vertices, $\overline{\mathcal{R}} \setminus \mathcal{P}$ is an open disc, D , embedded in Σ .

In general, $\overline{\mathcal{R}}$ may not be a disc, or $\partial D = \mathcal{P}$ may not be an embedded circle. Let $\mathcal{P}_\circ \subset \mathcal{P}$ denote the subset of \mathcal{P} where we cut out $\overline{\mathcal{R}}$ to obtain D . We have $\overline{\mathcal{R}} \setminus \mathcal{P}_\circ = \overline{D}$. A connected component λ of \mathcal{P}_\circ is either

- (i) a positive elliptic point like in Figure 31, or
- (ii) a union of stable separatrices like the thick arcs in Figures 32 and 33.

Cutting $\overline{\mathcal{R}}$ along λ produces two copies of λ , which we denote by λ_1 and λ_2 . Move λ_2 slightly away from λ_1 so that now $\partial \overline{D}$ is an embedded circle in M . We extend \overline{D} by adding a collar neighborhood along $\partial \overline{D}$ so that the resulting surface, \tilde{D} , is a disc embedded in M , its boundary $\partial \tilde{D}$ is a positive transverse unknot, and the open book foliation of the collar $\tilde{D} \setminus \overline{D}$ has no singularities. Figure 32 shows the change in open book foliation near λ and corresponding movie presentations.




Figure 31: Case (i): Construction of \tilde{D} when λ is a single positive elliptic point. The arrows depict the binding.




Figure 32: Case (ii): Construction of \tilde{D} . The thin arrows represent part of the binding.

By the construction, \tilde{D} satisfies all the requirements in Definition 4.1, so \tilde{D} is a transverse overtwisted disc. By Proposition 4.2 we conclude that ξ is overtwisted. \square

Corollary 4.6 *If a contact 3-manifold $(M_{(S,\phi)}, \xi_{(S,\phi)})$ contains an overtwisted disc, then (S, ϕ) contains a transverse overtwisted disc.*




Figure 33: Case (ii): Construction of \tilde{D}

Proof Let $\Delta \subset (M, \xi)$ be an overtwisted disc. We orient Δ so that the elliptic point of $\mathcal{F}_\xi(D)$ has negative sign. Since Δ is embedded and the boundary $L = \partial\Delta$ is a Legendrian knot, [18, page 129] implies that we can take a collar neighborhood $\nu(\Delta)$ of Δ whose characteristic foliation $\mathcal{F}_\xi(\nu(\Delta))$ is sketched in Figure 34.




Figure 34: The characteristic foliation $\mathcal{F}_\xi(\nu(\Delta))$ and a positive transverse push off L^+

Let $L^+ \subset \nu(\Delta)$ (dashed circle in Figure 34) be a positive transverse push off of L . Let $\Delta^+ \subset \nu(\Delta)$ be the disc bounded by L^+ . Then $sl(L^+, [\Delta^+]) = 1$ and the Euler characteristic has $\chi(\Delta^+) = 1$. In particular, $sl(L^+, [\Delta^+]) > -\chi(\Delta^+)$. By the proof of Theorem 4.3, we can find a transverse overtwisted disc. \square

Acknowledgements

The authors would like to thank Joan Birman, John Etnyre and the referees for numerous constructive comments. They also thank Marcos Ortiz for help with the English. The first author was supported by JSPS Research Fellowship for Young Scientists. The second author was partially supported by NSF grants DMS-0806492 and DMS-1206770.

References

- [1] **D Bennequin**, *Entrelacements et équations de Pfaff*, from “Third Schnepfenried geometry conference, Vol. 1”, Astérisque 107, Soc. Math. France, Paris (1983) 87–161 MR753131
- [2] **J S Birman**, *Braids, links, and mapping class groups*, Annals of Mathematics Studies 82, Princeton Univ. Press (1974) MR0375281
- [3] **J S Birman**, **E Finkelstein**, *Studying surfaces via closed braids*, J. Knot Theory Ramifications 7 (1998) 267–334 MR1625362
- [4] **J S Birman**, **MD Hirsch**, *A new algorithm for recognizing the unknot*, Geom. Topol. 2 (1998) 175–220 MR1658024
- [5] **J S Birman**, **W W Menasco**, *Studying links via closed braids, IV: Composite links and split links*, Invent. Math. 102 (1990) 115–139 MR1069243
- [6] **J S Birman**, **W W Menasco**, *Studying links via closed braids, II: On a theorem of Bennequin*, Topology Appl. 40 (1991) 71–82 MR1114092
- [7] **J S Birman**, **W W Menasco**, *Studying links via closed braids, I: A finiteness theorem*, Pacific J. Math. 154 (1992) 17–36 MR1154731
- [8] **J S Birman**, **W W Menasco**, *Studying links via closed braids, V: The unlink*, Trans. Amer. Math. Soc. 329 (1992) 585–606 MR1030509
- [9] **J S Birman**, **W W Menasco**, *Studying links via closed braids, VI: A nonfiniteness theorem*, Pacific J. Math. 156 (1992) 265–285 MR1186805
- [10] **J S Birman**, **W W Menasco**, *Studying links via closed braids, III: Classifying links which are closed 3–braids*, Pacific J. Math. 161 (1993) 25–113 MR1237139
- [11] **J S Birman**, **W W Menasco**, *Stabilization in the braid groups, I: MTWS*, Geom. Topol. 10 (2006) 413–540 MR2224463
- [12] **J S Birman**, **W W Menasco**, *Stabilization in the braid groups, II: Transversal simplicity of knots*, Geom. Topol. 10 (2006) 1425–1452 MR2255503
- [13] **J S Birman**, **W W Menasco**, *A note on closed 3–braids*, Commun. Contemp. Math. 10 (2008) 1033–1047 MR2468377
- [14] **J S Birman**, **N C Wrinkle**, *On transversally simple knots*, J. Differential Geom. 55 (2000) 325–354 MR1847313
- [15] **Y Eliashberg**, *Classification of overtwisted contact structures on 3–manifolds*, Invent. Math. 98 (1989) 623–637 MR1022310
- [16] **Y Eliashberg**, *Contact 3–manifolds twenty years since J Martinet’s work*, Ann. Inst. Fourier (Grenoble) 42 (1992) 165–192 MR1162559
- [17] **Y Eliashberg**, **M Fraser**, *Classification of topologically trivial Legendrian knots*, from “Geometry, topology, and dynamics” (F Lalonde, editor), CRM Proc. Lecture Notes 15, Amer. Math. Soc. (1998) 17–51 MR1619122

- [18] **J B Etnyre**, *Legendrian and transversal knots*, from “Handbook of knot theory” (W W Menasco, M Thistlethwaite, editors), Elsevier, Amsterdam (2005) 105–185 MR2179261
- [19] **J B Etnyre, K Honda**, *On the nonexistence of tight contact structures*, *Ann. of Math.* 153 (2001) 749–766 MR1836287
- [20] **J B Etnyre, K Honda**, *Cabling and transverse simplicity*, *Ann. of Math.* 162 (2005) 1305–1333 MR2179731
- [21] **J B Etnyre, B Ozbagci**, *Invariants of contact structures from open books*, *Trans. Amer. Math. Soc.* 360 (2008) 3133–3151 MR2379791
- [22] **J B Etnyre, J Van Horn-Morris**, *Fibered transverse knots and the Bennequin bound*, *Int. Math. Res. Not.* 2011 (2011) 1483–1509 MR2806512
- [23] **H Geiges**, *An introduction to contact topology*, *Cambridge Studies in Advanced Mathematics* 109, Cambridge Univ. Press (2008) MR2397738
- [24] **E Giroux**, *Convexité en topologie de contact*, *Comment. Math. Helv.* 66 (1991) 637–677 MR1129802
- [25] **E Giroux**, *Structures de contact en dimension trois et bifurcations des feuilletages de surfaces*, *Invent. Math.* 141 (2000) 615–689 MR1779622
- [26] **E Giroux**, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, from “Proceedings of the International Congress of Mathematicians, Vol. II” (T Li, editor), Higher Ed. Press, Beijing (2002) 405–414 MR1957051
- [27] **M W Hirsch**, *Differential topology*, *Graduate Texts in Mathematics* 33, Springer, New York (1976) MR0448362
- [28] **K Honda**, *On the classification of tight contact structures, I*, *Geom. Topol.* 4 (2000) 309–368 MR1786111
- [29] **T Ito, K Kawamuro**, *Essential open book foliation and fractional Dehn twist coefficient* arXiv:1208.1559
- [30] **T Ito, K Kawamuro**, *Operations on open book foliations* arXiv:1309.4486
- [31] **D Johnson**, *The structure of the Torelli group, II: A characterization of the group generated by twists on bounding curves*, *Topology* 24 (1985) 113–126 MR793178
- [32] **K Kawamuro**, *The self-linking number in planar open book decompositions*, *Math. Res. Lett.* 19 (2012) 41–58 MR2923175
- [33] **K Kawamuro, E Pavelescu**, *The self-linking number in annulus and pants open book decompositions*, *Algebr. Geom. Topol.* 11 (2011) 553–585 MR2783238
- [34] **D LaFountain, W W Menasco**, *Climbing a Legendrian mountain range without stabilization* (2009) arXiv:0801.3475v2

- [35] **Y Mitsumatsu**, *Convergence of contact structures to foliations*, from “Foliations 2005” (P Walczak, R Langevin, S Hurder, T Tsuboi, editors), World Sci. Publ., Hackensack, NJ (2006) 353–371 MR2284791
- [36] **S Morita**, *Families of Jacobian manifolds and characteristic classes of surface bundles, I*, Ann. Inst. Fourier (Grenoble) 39 (1989) 777–810 MR1030850
- [37] **S Morita**, *Mapping class groups of surfaces and three-dimensional manifolds*, from “Proceedings of the International Congress of Mathematicians, Vol. I, II” (I Satake, editor), Math. Soc. Japan, Tokyo (1991) 665–674 MR1159253
- [38] **S Morita**, *The extension of Johnson’s homomorphism from the Torelli group to the mapping class group*, Invent. Math. 111 (1993) 197–224 MR1193604
- [39] **E Pavelescu**, *Braids and open book decompositions*, PhD thesis, University of Pennsylvania (2008) Available at <http://www.math.upenn.edu/grad/dissertations/ElenaPavelescuThesis.pdf>
- [40] **E Pavelescu**, *Braiding knots in contact 3–manifolds*, Pacific J. Math. 253 (2011) 475–487 MR2878820
- [41] **W P Thurston**, *A norm for the homology of 3–manifolds*, Mem. Amer. Math. Soc. 59 (1986) i–vi and 99–130 MR823443
- [42] **W P Thurston**, **H E Winkelnkemper**, *On the existence of contact forms*, Proc. Amer. Math. Soc. 52 (1975) 345–347 MR0375366
- [43] **R Trapp**, *A linear representation of the mapping class group \mathcal{M} and the theory of winding numbers*, Topology Appl. 43 (1992) 47–64 MR1141372

Research Institute for Mathematical Sciences, Kyoto University
 Sakyo-ku, Kyoto 606-8502, Japan

Department of Mathematics, The University of Iowa
 14 McLean Hall, Iowa City, IA 52242, USA

tetitoh@kurims.kyoto-u.ac.jp, kawamuro@iowa.uiowa.edu

<http://kurims.kyoto-u.ac.jp/~tetitoh/>

Proposed: Shigeyuki Morita
 Seconded: Yasha Eliashberg, Walter Neumann

Received: 25 January 2013
 Revised: 3 September 2013