

# Geodesics and horizontal-path spaces in Carnot groups

ANDREI A AGRACHEV  
ALESSANDRO GENTILE  
ANTONIO LERARIO

We study properties of the space of horizontal paths joining the origin with a vertical point on a generic two-step Carnot group. The energy is a Morse-Bott functional on paths and its critical points (sub-Riemannian geodesics) appear in families (compact critical manifolds) with controlled topology. We study the asymptotic of the number of critical manifolds as the energy grows. The topology of the horizontal-path space is also investigated, and we find asymptotic results for the total Betti number of the sublevels of the energy as it goes to infinity. We interpret these results as local invariants of the sub-Riemannian structure.

53C17; 37J60, 58E05

## 1 Introduction

How many geodesics are there between two fixed points on a Riemannian manifold?

The classical way to answer this question, due to M Morse, is to consider the space  $\Omega$  of *all* curves (defined on the same interval) joining the two points, together with the function  $J: \Omega \rightarrow \mathbb{R}$  that associates to each curve  $\gamma$  the energy  $\frac{1}{2} \int_I \|\dot{\gamma}\|^2$ . Geodesics joining the two points are thus regarded as *critical points* of the energy function, and the celebrated *Morse inequalities* state that, in the generic case, the number of such geodesics with energy bounded by  $s$  is *at least* the number of “holes” of the topological space  $\{J \leq s\}$ .

In this Riemannian framework, if the two points  $p_0$  and  $p$  are close enough and we only allow curves with energy bounded by a small constant  $c > 0$ , there is only one geodesic joining them and the space of curves connecting  $p_0$  with  $p$  with energy bounded by  $c$  is contractible (bounding the energy ensures that the curves we consider are contained in a small neighborhood of  $p_0$ ).

In fact Morse inequalities can be used both ways: geometers study the topology of  $\{J \leq s\}$  to get information on geodesics; topologists use properties of geodesics in

order to understand the path space  $\Omega$ . This subject has been widely investigated; see for instance the classical works of Milnor [14] and Bott [6; 7].

On a *sub-Riemannian* manifold  $M$  the situation gets more complicated. Being the velocities of curves constrained to the subbundle  $\Delta \subset TM$ , the *horizontal*-path space is no longer the same as above and new phenomena can occur.

Typically, the space of horizontal curves with energy less than  $s$  joining  $p_0$  and  $p$  has finite-dimensional homology but its homological dimension and total Betti number may grow to infinity as  $p \rightarrow p_0$ . We are interested in the asymptotic behavior of these quantities and related properties of the space of sub-Riemannian geodesics joining  $p_0$  and  $p$ .

In this paper, we focus on the case of a two-step Carnot group where the situation is well controlled. A one-step Carnot group is just a Euclidean space. The general  $k$ -step Carnot group  $G$  is a simply connected  $k$ -step nilpotent Lie group equipped with a “dilation”. Such a dilation is a one-parametric group of automorphisms that generalizes homotheties of the Euclidean space. A typical example is the group of  $n \times n$  lower-triangular matrices with units on the main diagonal, zeros over the diagonal and dilation  $\delta_t: \{a_{ij}\}_{i,j=1}^n \mapsto \{t^{i-j} a_{ij}\}_{i,j=1}^n, t > 0$  (Example 1 below is the case  $n = 3$ ).

First-order elements with respect to the dilation form a vector subspace of the Lie algebra of the group and thus define a left-invariant vector distribution  $\Delta \subset TG$ . It is assumed that the first-order elements generate the whole Lie algebra (see Pansu [16] and the first author, Boscain and Barilari [2] for details and for the explanation of the role of Carnot groups in sub-Riemannian geometry).

There are no one- or two-dimensional nonabelian Carnot groups. The only three-dimensional nonabelian Carnot group is the Heisenberg group; we discuss its geometry as it will be an enlightening, leading example for the rest of the paper.

**Example 1** The *Heisenberg group*  $G$  is the smooth manifold  $\mathbb{R}^3$  with the usual coordinates  $(x, y, z)$  and the distribution

$$(1) \quad \Delta = \text{span} \left\{ \frac{\partial}{\partial x} - \frac{y}{2} \frac{\partial}{\partial z}, \frac{\partial}{\partial y} + \frac{x}{2} \frac{\partial}{\partial z} \right\}.$$

This means that one is allowed to move only along curves  $\gamma: I = [0, 2\pi] \rightarrow G$  whose velocity pointwise belongs to  $\Delta$  (these curves are called horizontal and they are all defined on the same interval; we pick this interval to be  $[0, 2\pi]$  because it will simplify notation later). In fact, for a technical reason, one considers curves whose derivative is defined almost everywhere and is square integrable, ending up with absolutely continuous curves. The sub-Riemannian structure is given by declaring the above vector fields form an orthonormal basis.

We assume one of the two points is the origin  $e = (0, 0, 0)$  and call the other  $p$ ; thus we consider

$$\Omega_p = \{\text{horizontal curves starting at } e \text{ and ending at } p\}.$$

Recall that  $\Delta_p = \text{span}\{X(p), Y(p)\}$ , where  $X, Y$  are the two vector fields given in (1); in particular for every horizontal curve  $\gamma$  we can write  $\dot{\gamma} = u_x X + u_y Y$ , where  $u_x, u_y \in L^2(I)$ . This correspondence is one-to-one: because of Cauchy’s theorem, given  $u \in L^2(I, \mathbb{R}^2)$  there is only one solution  $\gamma_u$  to the system  $\dot{\gamma} = u, \gamma(0) = e$ . Thus we can identify the space of *all* horizontal curves starting at  $e$  with  $L^2(I, \mathbb{R}^2)$  and  $\Omega_p$  coincides with the set of curves such that  $\gamma_u$  is defined for  $t = 2\pi$  and  $\gamma_u(2\pi) = p$  (thus  $u$  represents the “coordinates” of  $\gamma_u$  and it is usually called its *control*).




Figure 1: Geodesics in the Heisenberg group

The energy of an horizontal curve is by definition

$$J(\gamma_u) = \frac{1}{2} \int_0^{2\pi} \|\dot{\gamma}_u(t)\|^2 dt = \frac{1}{2} \|u\|^2.$$

Critical points of this function restricted to the various  $\Omega_p$  (ie geodesics between  $e$  and  $p$ ) are curves whose projection on the  $(x, y)$ -plane is an arc of a circle (possibly with infinite radius, ie an interval on a straight line); the signed area swept out on the circle by this projection equals the  $z$ -coordinate of the final point.

We see that if  $p$  belongs to the  $(x, y)$ -plane there is only one geodesic joining it with the origin (this is precisely the segment through  $e$  and  $p$ ), as if we were on a Riemannian manifold; if  $p$  has both nonzero components in the  $(x, y)$ -plane and the  $z$ -axis, the number of geodesics is finite; finally, if  $p$  belongs to the  $z$ -axis there

are infinitely many such geodesics (given one we obtain infinitely many others by composing it with a rotation in  $\mathbb{R}^3$  along the axis).

The behavior of geodesics ending at a point  $(0, 0, z)$  (slightly abusing notation we call this point  $z$ ) is really new compared to the classical Riemannian situation and is worth discussing in more detail.

Given a natural number  $n \in \mathbb{N}_0$  let us pick a circle in the  $(x, y)$ -plane of area  $|z|/n$  passing through the origin. Starting at  $e$  and moving clockwise around the circle  $n$  times, we obtain the projection of a geodesic ending at  $z$  and whose energy is  $2\pi n|z|$ . Rotating this curve along the  $z$ -axis of an angle  $\theta \in \text{SO}(2) \simeq S^1$  gives another such geodesic (thus we have an  $S^1$  of geodesics with the same energy and endpoints). Hence geodesics ending at  $z$  (ie critical points of  $J$  on  $\Omega_z$ ) arrange into *infinitely* many *families*, indexed by the values of their energy  $\{2\pi n|z|\}_{n \in \mathbb{N}}$ , each family being homeomorphic to an  $S^1$  in  $\Omega_z$ .

It is not possible, therefore, to study the topology of  $\Omega_z$  by using  $f = J|_{\Omega_z}$  as a Morse function, simply because its critical points are not isolated. Despite this, as we have noted these critical points appear in nice families and in fact these families are *nondegenerate* in a sense that will be specified later, allowing to use the Morse theory machinery (this generalization is usually called *Morse–Bott* theory).

It turns out that  $\Omega_z$  is homotopy equivalent to an infinite-dimensional sphere

$$\Omega_z \sim S^\infty$$

and is contractible. What kind of information can a Morse theoretical study of  $\Omega_z$  then give? As a starting point, the investigation of critical points of  $f = J|_{\Omega_z}$  gives information about the structure of geodesics, but we will see there is also more.

Let us find all the critical manifolds of  $f$  with energy bounded by  $s$ . A circle through the origin of area  $|z|/n$  is the projection of a geodesic between  $e$  and  $z$  with energy bounded by  $s$  if and only if  $n \leq s/(2\pi|z|)$ . Rotating each such geodesic we get a whole critical manifold homeomorphic to  $S^1$ . In particular,

$$(2) \quad \{\text{geodesics between } e \text{ and } z \text{ with } J \leq s\} = \underbrace{S^1 \cup \dots \cup S^1}_{\lfloor \frac{s}{2\pi|z|} \rfloor \text{ copies}}.$$

In this context the “amount” of geodesics can still be used as a measure of the topology of  $\Omega_z \cap \{J \leq s\}$ , in a form that generalizes Morse inequalities. Specifically, whenever the critical points of  $f$  arrange into smooth manifolds (with some nondegeneracy conditions), then the sum of the Betti numbers (ie the number of “holes”) of  $\{f \leq s\}$

is bounded by the sum of the Betti numbers of all the critical manifolds with  $f \leq s$ . The inequalities we get are called *Morse–Bott inequalities*, and in our case they give<sup>1</sup>

$$(3) \quad b(\Omega_z \cap \{J \leq s\}) \leq b(S^1 \cup \dots \cup S^1) \sim \frac{s}{\pi|z|}.$$

Thus, as  $s \rightarrow \infty$  there are more and more critical manifolds but, surprisingly enough, the above inequality is far from being sharp. One can in fact show that  $\Omega_z \cap \{J \leq s\}$  has the homotopy type of a finite-dimensional sphere  $S^{d_s}$  in  $\Omega_z$ :

$$\Omega_z \cap \{J \leq s\} \sim S^{d_s} \quad \text{and} \quad b(\Omega_z \cap \{J \leq s\}) = 2.$$

In fact the dimension  $d_s$  of this sphere increases linearly in  $s$  but the sum of its Betti numbers is constant: this shows that the Morse–Bott inequalities in (3) overcount the Betti numbers.

Hence the picture is the same as of a family of finite-dimensional spheres “approaching” an infinite-dimensional one; the dimension of the spheres increases with  $s$  and the topology eventually vanishes (ie  $\lim_{s \rightarrow \infty} b_i(\Omega_z \cap \{J \leq s\}) = 0$  for every  $i \in \mathbb{N}_0$ ). Nonetheless there is something that “persists” at infinity: the *sum* of all the Betti numbers of  $\Omega_z \cap \{J \leq s\}$ . What is the meaning of this invariant number?




Figure 2: The homology of  $\Omega_p \cap \{J \leq s\}$  in the Heisenberg group: the top-dimensional homology moves to infinity as  $s$  grows and eventually “disappears”.

A possible way to answer this question is to consider a local problem near  $e$ : Assuming  $d(e, p_\epsilon) = \epsilon$ , how many geodesics are there between  $e$  and  $p_\epsilon$  with *bounded* energy (say by a constant  $c$ )? As we have seen, if  $p_\epsilon$  is not a vertical point, the answer is finitely many (depending on the upper bound on the energy). On the other hand, what happens if  $p_\epsilon$  approaches  $e$  along the non-Riemannian  $z$ -axis? Let us consider the point

$$p_\epsilon = \left(0, 0, \frac{\epsilon^2}{4\pi}\right) \quad \text{such that} \quad d(e, p_\epsilon) = \epsilon.$$

<sup>1</sup>For the rest of the paper,  $b(X)$  will indicate the *sum* of the  $\mathbb{Z}_2$ -Betti numbers of  $X$ , ie  $b(X) = \sum_i \text{rank } H_i(X; \mathbb{Z}_2)$ . This sum might a priori be infinite, depending on the topological space  $X$ , but in our case it will always be finite.

Arguing as in (2), we see that the critical manifolds of  $J|_{\Omega_{p_\epsilon}}$  with energy bounded by  $c$  are  $\lfloor 2c/\epsilon^2 \rfloor$  copies of  $S^1$ : as we let  $p_\epsilon$  move closer and closer to  $e$  the number of geodesics keeps increasing. The equations for  $\Omega_{p_\epsilon} \cap \{J \leq c\}$  are

$$\Omega_{p_\epsilon} \cap \{J \leq c\} = \left\{ F_1(u) = F_2(u) = 0, q(u) = \frac{\epsilon^2}{2\pi}, \|u\|^2 \leq c \right\},$$

and the transformation  $u \mapsto u/(\epsilon\sqrt{2})$  gives a homeomorphism between  $\Omega_{p_\epsilon} \cap \{J \leq c\}$  and  $\Omega_{(0,0,1/(4\pi))} \cap \{J \leq (2c)/\epsilon^2\}$ . Hence the above discussion applies with  $z = (0, 0, \frac{1}{4\pi})$  and  $s = 2c/\epsilon^2$ , giving

$$\lim_{\epsilon \rightarrow 0} b(\Omega_{p_\epsilon} \cap \{J \leq c\}) = 2.$$

Thus, this persisting number can be interpreted as a local invariant of the Heisenberg group “in the direction” of  $z$ , as we will make more clear later.

### 1.1 A limiting procedure

The previous example shows that the sub-Riemannian case can be much richer than the Riemannian one, even at the level of the topology of the set of curves in  $\Omega_p$  when  $p$  is close to the initial point and the energy stays bounded. Let us continue along these lines and ask if we can find, in the general case, some asymptotic behavior when the distance to the final point becomes smaller and the energy stays bounded. To this end we need a procedure for taking limits.

The “time-one” picture is the following: We have two points  $e$  and  $p$  at distance  $s$  on a sub-Riemannian manifold  $M$ , and the space  $\Omega_p \cap \{J \leq c\}$  of curves reaching a neighborhood of  $p$ . Now, on a neighborhood  $U$  of  $e$  there is a well-defined family of nonisotropic “dilations” centered at  $e$  (we assume  $p$  is close enough to be in  $U$ ),

$$\delta_\lambda: U \rightarrow U, \quad \lambda > 0,$$

with the property that  $d(e, \delta_\lambda(p)) = \lambda d(e, p)$  (for example, in the Heisenberg group this family operates as  $\delta_\lambda(x, y, z) = (\lambda x, \lambda y, \lambda^2 z)$ ). We are thus led to consider the family of spaces

$$(4) \quad \Omega_{\delta_\epsilon(p)} \cap \{J \leq c\}, \quad \epsilon \rightarrow 0$$

exactly as we did in the Heisenberg case. This limiting procedure is reminiscent of the one used to construct the so called *sub-Riemannian* tangent space at  $e$ : as a vector space it is simply  $T_e M$ , but it comes endowed with much more structure that makes it into a Carnot group.

Two-step Carnot groups are “tangent spaces” to a generic sub-Riemannian manifold (the two-step condition needs some assumptions on the corank of  $\Delta$  to be generic).

Thus, on one hand our results should be interpreted as an “infinitesimal” version of a more general local theory (that the authors plan to discuss in a subsequent paper); on the other hand, Carnot groups are themselves sub-Riemannian manifolds (plenty of literature has been devoted to them) and we believe this study is of independent interest.

The questions we will be interested in are the following.

- (a) What is the structure of geodesics connecting two points  $e$  and  $p$  (ie critical points of the energy on  $\Omega_p$ ) in a two-step Carnot group?
- (b) What is the growth rate of the “number” of critical points at infinity? How does the topology of  $\Omega_p \cap \{J \leq s\}$  behave as  $s \rightarrow \infty$ ?
- (c) Is there some invariant that can be captured by  $\Omega_{\delta_\epsilon(p)} \cap \{J \leq c\}$  when we pass to the limit  $\epsilon \rightarrow 0$ ?

### 1.2 The horizontal-loop space

To start with we need to make our definition of a two-step Carnot group more precise. We consider a Lie algebra  $\mathfrak{g} = \Delta \oplus \Delta^2$  with bases  $\{e_1, \dots, e_d\}$  for  $\Delta$  and  $\{f_1, \dots, f_l\}$  for  $\Delta^2$ . We assume that the only nonzero brackets are given by

$$[e_i, e_j] = \sum_{k=1}^d a_{ij}^k f_k, \quad a_{ij}^k \in \mathbb{R}.$$

The Carnot group we will consider is the unique connected and simply connected Lie group  $G$  whose Lie algebra is  $\mathfrak{g}$  (it turns out that as a differentiable manifold,  $G$  is simply  $\mathbb{R}^{d+l}$ ). The sub-Riemannian structure is obtained by left translating  $\Delta \subset T_e G$  and declaring the vector fields obtained by left translation of the  $e_i, i = 1, \dots, d$ , to be orthonormal.

The constants  $\{a_{ij}^k\}$  come from skew symmetric matrices  $A_k = (a_{ij}^k) \in \mathfrak{so}(d), k = 1, \dots, l$ , and these matrices span an  $l$ -dimensional vector space

$$W = \text{span}\{A_1, \dots, A_l\} \subset \mathfrak{so}(d).$$

Two different choices of bases for  $W$  will produce *isomorphic* Carnot groups; for this reason we will call  $W \subset \mathfrak{so}(d)$  the *Carnot group structure*.

Horizontal curves will be absolutely continuous curves, whose velocities (almost everywhere defined) are in  $\Delta$ . As we did for the Heisenberg group,  $\Omega_p$  will denote

the set of horizontal curves joining the origin  $e$  with  $p$  and there will be a natural inclusion

$$\Omega_p \hookrightarrow L^2(I, \mathbb{R}^d).$$

In this way the *energy* of a curve  $\gamma_u$  with velocity  $u$  will simply be  $J(\gamma_u) = \frac{1}{2}\|u\|^2$ .

With this notation we have the homeomorphism

$$(5) \quad \Omega_{\delta_\epsilon(p)} \cap \{J \leq c\} = \delta_\epsilon(\Omega_p \cap \{J \leq c/\epsilon^2\}).$$

For the generic point  $p$  not in  $\exp(\Delta^2) \subset G$  (the second layer) we were able to prove under the condition  $d > l$  that the number of geodesics is finite, from which it follows that the topology of  $\Omega_p \cap \{J \leq s\}$  stabilizes for  $s \rightarrow \infty$  and the space becomes contractible in a “finite time”. We believe that this holds also without the condition  $d > l$ ; in any case what is clear is that the remarkable properties of the asymptotic of the homology are due to the second layer part of the point.

**Proposition** *If  $p$  is not a vertical point and  $d > l$ , the cohomology of  $\Omega_p \cap \{J \leq s\}$  stabilizes for  $s$  big enough.*

In particular, using the above rescaling procedure, the homology of  $\Omega_{\delta_\epsilon(p)} \cap \{J \leq c\}$  also stabilizes for  $\epsilon$  small enough if  $p$  is not vertical and  $d > l$ .

From now on we will focus on final points in  $\exp(\Delta^2)$ ; we will call such points *vertical*: they are analogous to the points in the  $z$ -axis of the Heisenberg group. By a slight abuse of notation, we can identify  $\Delta^2$  with its image under the exponential map in  $\mathbb{R}^{d+l}$ , and write the condition of being vertical simply as  $p \in \Delta^2$ .

All the above questions can be addressed by studying the Morse–Bott theory of  $J|_{\Omega_p}$ , by taking appropriate rescalings. In fact, assuming  $p$  is vertical, we have  $\delta_\epsilon(p) = \epsilon^2 p$  and the homogeneity of the endpoint map on the vertical directions gives the homeomorphism (5). In particular the limit (4) can be studied by simply fixing  $p$  and letting the energy grow (at a rate  $\epsilon^{-2}$ ).

**Theorem** (The topology of  $\Omega_p$ ) *The topological space  $\Omega_p$  is homotopy equivalent to  $\Omega_e$  and is therefore contractible. For generic  $p \in \Delta^2$ , it is a Hilbert manifold.*

In particular, analysis can be performed over  $\Omega_p$  and critical points of  $J|_{\Omega_p}$  (which is a smooth function) can be found using the Lagrange multiplier rule. As for the Heisenberg case, we can write down explicit equations: here the crucial point is that the end-point map sends a curve to  $\Delta^2$  if and only if its control has zero mean (ie  $0 = \int_I u \in \mathbb{R}^d$ ) and the restriction of this map to the space of such controls is *quadratic*.

### 1.3 The structure of geodesics

A careful investigation of the critical points of  $f$  shows that they appear in families, ie they arrange into critical manifolds. These manifolds are tori  $S^1 \times \dots \times S^1$  and are indexed by their Lagrange multipliers. The whole structure can be recovered by  $W \subset \mathfrak{so}(d)$  only, as follows.

Consider the infinite union of algebraic sets

$$\Lambda = \bigcup_{n \in \mathbb{N}} \Lambda_n, \quad \Lambda_n = \{A \in W \mid \det(A - in\mathbf{1}) = 0\}.$$

Each set  $\Lambda_n$  is an hypersurface in  $W$ : generically it is smooth, but there can be points where  $iA$  has multiple integer eigenvalues and the corresponding hypersurface is singular. In fact all these hypersurfaces (except  $\Lambda_0$ ) can be obtained by dilations of  $\Lambda_1$  (ie  $\Lambda_n = n\Lambda_1$ ). Hence  $\Lambda$  looks like an “infinite net” with two kinds of singularities (they might appear at the same time): one kind comes from the singularities of each  $\Lambda_n$  and the other from the intersections  $\Lambda_{n_1} \cap \dots \cap \Lambda_{n_\nu}$  for  $\nu \leq l$ .

As we will see  $\Lambda$  represents the set of all possible Lagrange multipliers, for all possible final points, hence to each point of  $\Lambda$  there corresponds a family of geodesics. Notice that once the basis  $\{A_1, \dots, A_l\}$  of  $W$  and coordinates are fixed the correspondence

$$\omega = (\omega_1, \dots, \omega_l) \mapsto \omega A = \omega_1 A_1 + \dots + \omega_l A_l$$

defines a linear isomorphism  $(\Delta^2)^* \simeq W$ .

The next theorem gives a detailed answer to question (a) above on the structure of geodesics.

**Theorem** (The structure of geodesics) *Let  $u$  be the control associated to a geodesic from  $e$  to a point in  $\Delta^2$  with Lagrange multiplier  $\omega$ . Then we have the following.*

- (a)  $u(t) = e^{-t\omega A} u_0$  with  $u_0 = e^{-2\pi\omega A} u_0$ .
- (b) The final point  $q(u) = (q_1(u), \dots, q_l(u)) \in \Delta^2$  of such geodesic is given by
 
$$q_i(u) = \langle u_0, L_i(\omega) u_0 \rangle, \quad L_i(\omega) = \frac{1}{2} \int_0^{2\pi} \left( \int_0^t e^{\tau\omega A} d\tau \right) A_i e^{-t\omega A} dt.$$
- (c) The energy of such geodesic is given by  $J(u) = \omega(F(u))$ .

In addition, for the generic choice of  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$ , we have the following.

- (d) The set  $\Lambda(p) \subset \Lambda$  of Lagrange multipliers for geodesics whose final point is  $p$  is a discrete set.

- (e) If  $\omega \in \Lambda(p)$  then the integer eigenvalues of the matrix  $i\omega A$  are simple and there exist  $n_1, \dots, n_\nu \in \mathbb{N}_0$  such that  $\omega$  belongs to  $\Lambda_{n_1} \cap \dots \cap \Lambda_{n_\nu}$  (the number  $\nu \leq l$  will be called the number of resonances of  $\omega$ ).
- (f) If  $\omega$  has  $\nu$  resonances, the union of all the geodesics from  $e$  to  $p$  with Lagrange multiplier  $\omega$  is a smooth manifold  $C_\omega$  homeomorphic to

$$C_\omega \simeq \underbrace{S^1 \times \dots \times S^1}_{\nu \text{ times}}.$$

Before proceeding any further, it is interesting to discuss one more example.

**Example 2** (Corank-two distributions) Let us consider the case of  $\mathbb{R}^{d+2}$  with Carnot group structure  $W = \text{span}\{A_1, A_2\} \subset \mathfrak{so}(d)$  having the above genericity property.

Let us consider the curve  $\Lambda_1 \subset W$  first: Since the set of matrices with double eigenvalues has codimension three in  $\mathfrak{so}(d)$ ,  $\Lambda_1$  is a smooth curve (with possibly many components). In this case the set  $\{\omega \in \Lambda_1 \mid \omega(p) \geq 0, p \in (T_\omega \Lambda_1)^\perp\}$  consists of Lagrange multipliers whose critical manifold  $C_\omega$  is homeomorphic to  $S^1$ ; all the other Lagrange multipliers whose associated critical manifold is a circle are obtained multiplying these covectors by a positive natural number:

$$\{\omega \in \Lambda(p) \mid C_\omega \simeq S^1\} = \bigcup_{n \in \mathbb{N}_0} \{\omega \in \Lambda_n \mid \omega(p) \geq 0, p \in (T_\omega \Lambda_1)^\perp\}.$$




Figure 3: The set  $\bigcup_{n \in \mathbb{N}_0} \{\omega \in \Lambda_n \mid \omega(p) \geq 0, p \in (T_\omega \Lambda_1)^\perp\}$

Let us now consider two natural numbers  $n_1, n_2$  such that  $\Lambda_{n_1} \cap \Lambda_{n_2} \neq \emptyset$  and let  $\omega$  be a point in this intersection such that  $\omega(p) \geq 0$ . Let  $E(\omega) = \{v \in \mathbb{R}^d \mid e^{2\pi\omega A} v = v\}$  and consider the quadratic forms  $q_i|_\omega: E(\omega) \rightarrow \mathbb{R}$  defined by

$$q_i|_\omega: v \mapsto \langle v, L_i(\omega)v \rangle, \quad i = 1, 2.$$

Let  $q|_\omega: E(\omega) \rightarrow \mathbb{R}^2$  be the quadratic map whose components are the above  $q_i|_\omega$ . The set of Lagrange multipliers whose associated critical manifold is  $S^1 \times S^1$  coincides with

$$\{\omega \in \Lambda(p) \mid C_\omega = S^1 \times S^1\} = \bigcup_{n_1, n_2 \in \mathbb{N}_0} \{\omega \in \Lambda_{n_1} \cap \Lambda_{n_2} \mid p \in \text{im}(q|_\omega)\}.$$




Figure 4: The set  $\bigcup_{n_1, n_2 \in \mathbb{N}_0} \{\omega \in \Lambda_{n_1} \cap \Lambda_{n_2} \mid p \in \text{im}(q|_\omega)\}$

### 1.4 Critical manifolds

Let us move now to question (b) above, on the growth rate of the number of critical manifolds of  $J|_{\Omega_p}$  as the energy goes to infinity. As we have seen for the Heisenberg group in (2), once  $p$  is fixed the number of critical manifolds with energy bounded by  $s$  increases linearly in  $s$ :

$$\text{Card}\{\text{critical manifolds of } J|_{\Omega_p} \text{ such that } J \leq s\} = O(s) \quad (\text{Heisenberg}).$$

The previous example shows that in the corank-two case the critical manifolds of  $J|_{\Omega_p}$  can be indexed by points on an infinite two-dimensional net and, naïvely, it is reasonable to guess that they increase quadratically in  $s$ . Indeed it is a general fact that this number increases at most as a polynomial of degree  $l = \text{corank}(\Delta)$  in  $s$  (see Figure 5).

**Theorem** (The growth rate of the number of critical manifolds) *For a generic choice of  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$  we have*

$$(6) \quad \text{Card}\{\text{critical manifolds of } J|_{\Omega_p} \text{ such that } J \leq s\} = O(s^l), \quad l = \text{dim}(\Delta^2).$$

Using the inequality in (6) and the fact that the topology of the critical manifold is uniformly bounded by a constant depending on  $d$  and  $l$  only, one is tempted to answer

the second part of question (b), concerning the topology of  $\Omega_p \cap \{J \leq s\}$ , using Morse–Bott inequalities. What we get with this strategy is, for a corank- $l$  distribution,

$$(7) \quad b(\Omega_p \cap \{J \leq s\}) = O(s^l).$$

Surprisingly enough, this estimate can be improved up to an  $O(s^{l-1})$  (as we already noticed, in the corank-one case  $b(\Omega_p \cap \{J \leq s\})$  is indeed constant). Before discussing the general case, let us continue the corank-two example.

### 1.5 A topological coarea formula

In the corank-two case not only can we show that  $b(\Omega_p \cap \{J \leq s\})$  grows as  $\tau(p)s + o(s)$ , but we can also analytically compute the leading coefficient  $\tau(p)$ .

To explain the result, consider a unit circle  $S^1 \subset W \simeq (\Delta^2)^*$ . As the parameter  $\theta$  varies on  $S^1$  let us consider the positive eigenvalues  $\alpha_1(\theta), \dots, \alpha_j(\theta)$  of the matrix  $i \cos(\theta)A_1 + i \sin(\theta)A_2$ . The genericity assumption ensures that these numbers can be taken as the value of semialgebraic functions  $\alpha_j: S^1 \rightarrow \mathbb{R}$ . Given  $p \in \Delta$  we consider the *rational* functions  $\lambda_j: S^1 \rightarrow \mathbb{R} \cup \{\infty\}$  given by

$$\lambda_j: \theta \mapsto \left| \frac{\alpha_j(\theta)}{p_1 \cos(\theta) + p_2 \sin(\theta)} \right| \quad \text{for } j = 1, \dots, d.$$

Notice that when  $\omega$  approaches  $p^\perp$  these functions might explode since they are rational in  $\theta$ ; on the other hand they are semialgebraic and differentiable almost everywhere and it makes sense to consider the integral

$$\tau(p) \doteq \frac{1}{2} \int_{S^1} \sum_{j=1}^d \left| \frac{\partial \lambda_j}{\partial \theta}(\theta) \right| - \left| \sum_{j=1}^d \frac{\partial \lambda_j}{\partial \theta}(\theta) \right| d\theta.$$

The convergence of the integral follows from the fact that where the derivatives explode, they all have the same sign and the integrand vanishes.

**Theorem** (Topological coarea formula) *If the corank  $l = 2$ , then for a generic choice of  $W \subset \mathfrak{so}(d)$  and  $p \in \Delta^2$  we have*

$$(8) \quad b(\Omega_p \cap \{J \leq s\}) = \tau(p)s + o(s).$$

The name “topological coarea” is reminiscent of integral geometry: in fact the coefficient  $\tau(p)$  is computed by considering the sum of the Betti numbers  $b = \sum_i b_i$  as an integral over the index set, regarding  $i$  as the “variable” of integration; rearranging this sum in an appropriate way in the limit gives exactly the coarea formula for functions of one variable. In fact, looking at the inequalities defining  $\Omega_p \cap \{J \leq s\}$ , one immediately

sees that they are quadratic. There is a general theory (see Appendix C or the first and third authors [3] for more details) for studying the topology of sets defined by quadratic inequalities: The idea is to consider quadratic forms obtained by taking linear combinations of the equations defining the set, ie considering quadratic forms depending on some (homogeneous) parameters (there are as many parameters as the number of inequalities minus one). The main ingredient is the function on this parameter space that counts the number of positive eigenvalues (the *positive inertia index*) of the corresponding quadratic form. Roughly, the theory says that for each “change” in the monotonicity of this function, there corresponds a “hole” in the set. Thus one can count, for each level of the positive inertia index, how many homology classes are at that level, and this gives another rearrangement of the above sum.




Figure 5: (Corank two) The number of critical manifolds with energy less than  $s$  can grow quadratically in  $s$  (this number is proportional to the volume of the shaded region; dots are Lagrange multipliers).

### 1.6 The growth rate of the topology

In the general case the upper bound we get for the growth rate of the total Betti number of  $\Omega_p \cap \{J \leq s\}$  is the following (see Figure 6).

**Theorem** (Bound on the growth rate of the topology) *For a generic choice of  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$  we have*

$$b(\Omega_p \cap \{J \leq s\}) = O(s^{l-1}), \quad l = \dim(\Delta^2).$$

Here the idea is to use the function  $-J|_{\Omega_p}$ , which is again Morse–Bott, whose critical manifolds are the same of  $J|_{\Omega_p}$  but with *infinite* index. Thus passing one of these critical manifolds amounts to attaching an infinite-dimensional cell (ie the homotopy of the sublevel set does not change). Thus one can start with  $\Omega_p \cap \{J \leq s\}$  and “push” its topology to its boundary  $\Omega_p \cap \{J = s\}$  via  $\nabla J|_{\Omega_p}$ . This observation will have important consequences for our study, including the above result.

If we look back again at the Heisenberg example, we see that  $b_i(\Omega_p \cap \{J \leq s\})$  is zero for  $i > O(s)$ , simply because  $\Omega_p \cap \{J \leq s\}$  has the homotopy type of a sphere whose dimension grows linearly in  $s$ . It is in fact a general phenomenon that the maximum nonzero Betti number increases at most linearly in  $s$  (with no restrictions on the corank).

**Theorem** (The vanishing rate of Betti numbers) *For the generic choice of  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$  we have*

$$\max\{i \mid b_i(\Omega_p \cap \{J \leq s\}) \neq 0\} = O(s).$$

## 1.7 Final remarks

At this point the study of question (b) above is complete and one can address question (c). What we know so far is that the whole  $\Omega_p$  is contractible and we have rather precise asymptotics on the behavior of the number of geodesics in  $\Omega_p \cap \{J \leq s\}$  and its total Betti number as  $s \rightarrow \infty$ . What we have to do is simply to translate the results using the homeomorphism (5):

$$\Omega_{\delta_\epsilon(p)} \cap \{J \leq c\} \simeq \Omega_p \cap \{J \leq s\} \quad \text{as } s = c/\epsilon^2 \rightarrow \infty.$$

For example one might ask whether the estimate for the number of critical points and the topology of  $\Omega_{\delta_\epsilon(p)} \cap \{J \leq c\}$  can be actually attained. In the case  $l = 2$  we have seen that the leading coefficient  $\tau(p)$  of  $b(\Omega_{\delta_\epsilon(p)})$  can be analytically computed and in fact for the generic Carnot group structure and  $p \in \Delta^2$  it is not difficult to show that  $\tau(p) \neq 0$ . In particular as we let  $\epsilon \rightarrow 0$ , the topology of  $\Omega_{\delta_\epsilon(p)}$  explodes; as a corollary the number of geodesics grows unbounded as well (this is an example of a backward use of Morse–Bott inequalities).

For the general corank  $l$  we do not know whether the limit

$$\tau(p) = \limsup_{\epsilon \rightarrow 0} b(\Omega_{\delta_\epsilon(p)} \cap \{J \leq 1\})\epsilon^{2l-2} = \limsup_{\epsilon \rightarrow 0} b(\Omega_p \cap \{J \leq 1/\epsilon\})\epsilon^{l-1}$$

is different from zero, but it is natural to guess so (we only know it is finite).




Figure 6: The Betti numbers of  $\Omega_p \cap \{J \leq s\}$ : as  $s$  goes to infinity the “wave” moves to the right and the sum of all the Betti numbers (the area below the wave) can increase as fast as  $O(s^{l-1})$ , but eventually everything “disappears”.

As a concluding remark, we believe these asymptotic Morse inequalities can be useful for generalizations and, in particular, for the next paper that we plan to devote to the general two-step sub-Riemannian structure. Of course, in general, we cannot guarantee that the functional is Morse–Bott. The quantity to estimate could be a “virtual number of geodesics”: the minimal number of critical points for a Morsification of the functional.

### 1.8 Structure of the paper

The paper is organized as follows. In Section 2 we discuss some preliminary material and give the main definitions. In Section 3 we study the structure of geodesics: the theorem on the smoothness and the topology of  $\Omega_p$  is Theorem 9 and the theorem on the structure of geodesics is a combination of Lemmas 10, 11 and 12 and Theorem 13. The growth of the number of critical manifolds is computed in Theorem 21; the bound on the growth rate of the topology of  $\Omega_p \cap \{J \leq s\}$  is proved in Theorem 26 and the exact asymptotic for the case  $l = 2$  is the content of Theorem 27; the theorem on the vanishing rate of the Betti numbers is Corollary 25. The appendices contain, for the reader’s convenience, some useful results we will use in the main body.

## 2 Preliminaries

### 2.1 Two-step Carnot groups and their geometry

Here we briefly recall the main definitions related to Carnot groups; the reader is referred to [2], Onishchick and Vinberg [15] and Semmes [17] for more details.

As a differentiable manifold a *two-step Carnot group* is a connected, simply connected Lie group  $G$  whose Lie algebra  $\mathfrak{g} = T_e G$  decomposes as

$$\mathfrak{g} = \Delta \oplus \Delta^2, \quad \text{with} \quad [\Delta, \Delta] = \Delta^2, \quad [\Delta, \Delta^2] = 0 \quad \text{and} \quad [\Delta^2, \Delta^2] = 0;$$

as vector spaces here we have  $\Delta \cong \mathbb{R}^d$  and  $\Delta^2 \cong \mathbb{R}^l$ .

Whenever a Lie algebra  $\mathfrak{g}$  as above is given, the existence (and uniqueness) of such a group  $G$  is guaranteed by Lie’s theorem. We recall that in fact under the above assumption on the structure of  $\mathfrak{g}$ , the exponential map  $\exp: \mathfrak{g} \rightarrow G$  is an analytic diffeomorphism, hence in particular  $G \simeq \mathbb{R}^{d+l}$ .

The geometric structure on  $G$  is given by fixing a scalar product  $h$  on  $\Delta$  and considering the distribution  $\Delta_q = dL_q \Delta$  together with the extension of  $h$  by left translation; in this way the triple  $(G, \Delta, h)$  defines a *sub-Riemannian manifold*. Notice that the distribution  $\Delta$  is by assumption bracket generating, hence it satisfies Hörmander’s condition, in particular meaning that  $G$  is *horizontal-path-connected* (see below).

Two such Carnot groups  $(G_1, \Delta_1, h_1)$  and  $(G_2, \Delta_2, h_2)$  are considered to be isomorphic if there exists a Lie algebra isomorphism  $L: \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$  such that  $L\Delta_1 = \Delta_2$  and  $L|_{\Delta_1}^* h_2 = h_1$ ; in fact by the simple connectedness assumption the linear map  $L$  integrates to a Lie group isomorphism  $\phi: G_1 \rightarrow G_2$  and the global geometric structures are related by  $(G_1, \Delta_1, h_1) = (G_2, \phi^{-1}\Delta_2, \phi^*h_2)$ .

If we fix now an *orthonormal* basis  $\{e_1, \dots, e_d\}$  of  $\Delta$  and a basis  $\{f_1, \dots, f_l\}$  of  $\Delta^2$ , then the bracket structure can be written as

$$(9) \quad [e_i, e_j] = \sum_{k=1}^l a_{ij}^k f_k \quad \text{for all } i, j \in \{1, \dots, d\}$$

where each matrix  $A_k = (a_{ij}^k)$  belongs to  $\mathfrak{so}(d)$ . In particular we can consider the vector space

$$W = \text{span}\{A_1, \dots, A_l\} \subset \mathfrak{so}(d).$$

Each vector space  $W$  of dimension  $l$  in  $\mathfrak{so}(d)$  defines a sub-Riemannian structure on  $G$  by fixing a basis  $\{A_1, \dots, A_l\}$  for  $W$  and declaring that the corresponding matrices define the bracket structure in an orthonormal basis  $\{e_1, \dots, e_d\}$  for  $\Delta$ . The isomorphism class of the Carnot group does not depend on the choice of the basis of  $W$ : indeed, let  $\{A'_1, \dots, A'_l\}$  be another basis of  $W$  and  $B = (b_{hk})$  the basis-change matrix, such that  $A'_h = \sum_{k=1}^l b_{hk} A_k$ . Now we can build another Carnot group by defining its Lie algebra  $\mathfrak{g}'$  with basis  $\{e'_1, \dots, e'_d, f'_1, \dots, f'_l\}$  such that  $\{e'_1, \dots, e'_d\}$  is orthonormal; the structure constants are given by the entries of the matrices  $A'_h$  as in (9). The map  $\phi$ , defined on the basis elements by  $e'_i \mapsto e_i$  and  $f'_h \mapsto \sum_{k=1}^l b_{hk} f_k$ , gives an isomorphism between  $\mathfrak{g}'$  and  $\mathfrak{g}$ .

**Remark 1** (The moduli space of Carnot groups) Once a sub-Riemannian structure is given, changing  $\{e_1, \dots, e_d\}$  to another orthonormal basis  $\{Me_1, \dots, Me_d\}$

(where  $M$  is an orthogonal matrix in  $O(d)$ ) changes  $W$  into  $W' = MWM^T$ . Thus denoting by  $G(l, \mathfrak{so}(d))$  the Grassmannian of  $l$ -planes in  $\mathfrak{so}(d)$ , the (naïve) moduli space of two-step Carnot groups is represented by the quotient  $\mathcal{M}_{l,d} = G(l, \mathfrak{so}(d))/O(d)$ . Since  $\mathcal{M}_{l,d}$  is the quotient of a manifold by a Lie group action, the quotient map is open and perturbing  $W$  defines a “genuine” perturbation of the isomorphism class of the corresponding Carnot group; in particular this means that a generic choice of  $W$  results in a generic choice of an isomorphism class of Carnot groups.

Motivated by the above remark, once  $W \in G(l, \mathfrak{so}(d))$  is fixed we consider the Carnot group given by exponentiating  $\mathfrak{g} = \mathbb{R}^d \oplus \mathbb{R}^l$ , whose Lie algebra is given as follows:  $\{e_1, \dots, e_d\}$  is the standard orthonormal basis for  $\mathbb{R}^d$ ,  $\{f_1, \dots, f_l\}$  is the standard basis for  $\mathbb{R}^l$  and fixing a basis  $\{A_1, \dots, A_l\}$  for  $W$ , the Lie brackets are given by (9); we will call  $W$  the *Carnot group structure*.

The following theorem gives a geometric realization of Carnot groups.

**Theorem 1** *Let  $\{A_1, \dots, A_l\}$  be a basis for  $W \subset \mathfrak{so}(d)$  and for  $i = 1, \dots, d$  consider the vector fields  $E_i$  on  $\mathbb{R}^{d+l}$  defined in coordinates  $(x, y)$  by*

$$E_i(x, y) = \frac{\partial}{\partial x_i}(x, y) - \frac{1}{2} \sum_{k=1}^l \sum_{j=1}^d a_{ij}^k x_j \frac{\partial}{\partial y_k}(x, y).$$

*Then the sub-Riemannian manifold  $(\mathbb{R}^{d+l}, \Delta = \text{span}\{E_1, \dots, E_l\}, g)$ , where  $g$  is the standard Euclidean metric, is isomorphic to the Carnot group defined by  $W$ .*

## 2.2 The end-point map

*Horizontal paths* on  $G$  are defined as absolutely continuous curves  $\gamma: I = [0, 2\pi] \rightarrow G$  whose derivative exists almost everywhere, is square integrable and belongs to the distribution  $\Delta$ . We denote the set of such paths by  $\Omega$ . Our choice of the interval  $I = [0, 2\pi]$  is motivated by simplicity of notation: we will later need to expand the components of an horizontal path into their Fourier series; a different choice of the interval will produce a completely equivalent theory.

As earlier noticed, the bracket generating condition implies that any two points in  $G$  can be joined by an horizontal path. The set  $\Omega$  can be given a *Hilbert manifold structure* as follows. Let  $u = (u_1, \dots, u_d) \in L^2(I, \mathbb{R}^d)$  and consider the Cauchy problem

$$\dot{\gamma}(t) = \sum_{i=1}^d u_i(t) E_i(\gamma(t)), \quad \gamma(0) = e.$$

We may assume this ODE problem is set on  $\mathbb{R}^{d+l}$  using the above geometric realization theorem; in this case the identity element  $e \in G$  corresponds to the zero of  $\mathbb{R}^{d+l}$ . By Caratheodory's theorem the above Cauchy problem has a local solution  $\gamma_u$  and we consider the set

$$\mathcal{U} = \{u \in L^2(I, \mathbb{R}^d) \mid \gamma_u \text{ is defined for } t = 2\pi \}.$$

For general sub-Riemannian manifolds the set  $\mathcal{U}$  is an open subset of  $L^2(I, \mathbb{R}^d)$  (by ODEs continuous dependence theorem) and is called the set of *controls*; in our case the estimates for the final time can be made uniform and we actually have  $\mathcal{U} = L^2(I, \mathbb{R}^d)$ . Associating to each  $u$  the corresponding path  $\gamma_u$  gives thus a local coordinate chart and by slightly abusing of notation in the sequel we will often identify  $\Omega$  with  $\mathcal{U}$ .

Once we are given the Carnot group structure  $W = \text{span}\{A_1, \dots, A_l\}$ , we can use Theorem 1 to write down the above ODE in a more explicit form:

$$\begin{cases} \dot{x} = u \\ \dot{y}_i = \frac{1}{2}x^T A_i u \end{cases} \quad \text{and} \quad \gamma(0) = 0.$$

In this framework the *end-point map* is the *smooth* map

$$F: \Omega \longrightarrow G,$$

which associates to each curve  $\gamma$  its final point  $\gamma(2\pi)$ .

If  $A := (A_1, \dots, A_l)$ , we can again use Theorem 1 and write the end-point map as

$$(10) \quad F(u) = \left( \int_0^{2\pi} u(t) dt, \frac{1}{2} \int_0^{2\pi} \left\langle \int_0^t u(\tau) d\tau, Au(t) \right\rangle dt \right);$$

(here the brackets denote the sub-Riemannian scalar product on the Lie algebra  $\mathfrak{g}$ ).

In the sequel we will mainly be interested in horizontal paths whose endpoints lie on  $\Delta^2$  (see Proposition 7 below).  $G$  being of step two, we know that  $\Delta^2$  is an abelian subalgebra of  $\mathfrak{g}$ ; therefore we can identify  $\Delta^2$  with the submanifold  $\exp(\Delta^2) \subset G$  and using Theorem 1 we can write this identification as

$$\xi_1 f_1 + \dots + \xi_l f_l \mapsto (\underbrace{0, \dots, 0}_x, \underbrace{\xi_1, \dots, \xi_l}_y),$$

(here as above  $\{f_1, \dots, f_l\}$  is a basis of  $\Delta^2$ ).

We study now the structure of the set of horizontal paths whose endpoints are on  $\Delta^2$ . It turns out that in the local coordinates given by the controls  $\mathcal{U}$  it coincides with the kernel of the differential of  $F$  at  $0 \in \mathcal{U}$ ,

$$H = \ker D_0 F,$$

as described by the following proposition.

**Proposition 2** *The following properties hold:*

- (a)  $H = \{u \in L^2(I, \mathbb{R}^d) \mid \int_I u \, dt = 0\}$ .
- (b)  $u \in H \Leftrightarrow F(u) \in \exp(\Delta^2)$ .
- (c)  $F|_H = \text{He}_0 F$ .

**Proof** For point (a) we compute the differential  $D_0 F$ : by taking a variation  $\varepsilon v$  of the constant curve  $\gamma \equiv 0$  we easily see that

$$\begin{aligned} D_0 F v &= \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \left( \varepsilon \int_0^{2\pi} v(t) \, dt, \frac{1}{2} \varepsilon^2 \int_0^{2\pi} \left\langle \int_0^t v(\tau) \, d\tau, A v(t) \right\rangle dt \right) \\ &= \left( \int_0^{2\pi} v(t) \, dt, 0 \right) \in \mathfrak{g}. \end{aligned}$$

which proves property (a).

Point (b) is a direct consequence of (10). For point (c) we notice that the Hessian  $\text{He}_0 F$  is defined on  $H = \ker D_0 F$  with values in  $\text{coker } D_0 F = \Delta^2$ , thus has the same range as  $F|_H$ . As in the proof of (a) if we consider the second derivative of a variation and we easily obtain  $\text{He}_0 F$  has the same expression of  $F$  when restricted to  $H$ .  $\square$

We denote by  $q$  the Hessian of  $F$  at zero, ie the quadratic map

$$q \doteq F|_H: H \rightarrow \mathbb{R}^l.$$

Every component  $q_i$  of  $q$  is a quadratic form on  $H$  and its explicit expression is given by

$$(11) \quad q_i(u) = \frac{1}{2} \int_0^{2\pi} \left\langle \int_0^t u(\tau) \, d\tau, A_i u(t) \right\rangle dt.$$

By polarization we obtain the expression for the associated bilinear form

$$\begin{aligned} q_i(u, v) &= \frac{1}{4} \left( \int_0^{2\pi} \left\langle \int_0^t u(\tau) + v(\tau) \, d\tau, A_i(u(t) + v(t)) \right\rangle dt \right. \\ &\quad \left. - \int_0^{2\pi} \left\langle \int_0^t u(\tau) \, d\tau, A_i u(t) \right\rangle dt - \int_0^{2\pi} \left\langle \int_0^t v(\tau) \, d\tau, A_i v(t) \right\rangle dt \right) \\ &= \frac{1}{4} \left( \int_0^{2\pi} \left\langle \int_0^t u(\tau) \, d\tau, A_i v(t) \right\rangle dt + \int_0^{2\pi} \left\langle \int_0^t v(\tau) \, d\tau, A_i u(t) \right\rangle dt \right) \\ &= \frac{1}{4} \left( \int_0^{2\pi} \left\langle \int_0^t u(\tau) \, d\tau, A_i v(t) \right\rangle dt - \int_0^{2\pi} \left\langle v(t), A_i \int_0^t u(\tau) \, d\tau \right\rangle dt \right) \\ &= \frac{1}{2} \int_0^{2\pi} \left\langle \int_0^t u(\tau) \, d\tau, A_i v(t) \right\rangle dt, \end{aligned}$$

where the fourth row follows from integration by parts.

Moreover to every  $q_i$  it corresponds a symmetric operator  $Q_i: H \rightarrow H$  defined by

$$q_i(u) = \langle u, Q_i u \rangle_H \quad \text{for all } u \in H;$$

recall that we have  $\langle u, v \rangle_H = \int_I \langle u, v \rangle dt$ . We will use the notation  $Q$  for the map  $(Q_1, \dots, Q_l): H \rightarrow H \otimes \mathbb{R}^l$ . Also, given a covector  $\omega \in \mathbb{R}^{l*}$  we will denote by  $\omega q, \omega Q, \omega A$  the compositions of  $\omega: \mathbb{R}^l \rightarrow \mathbb{R}$  respectively with  $q, Q$  and  $A$ .

In the sequel we will need to expand a control  $u \in H$  into its Fourier series: we will write  $u = \sum_{k \in \mathbb{N}_0} U_k \frac{1}{\sqrt{\pi}} \cos kt + V_k \frac{1}{\sqrt{\pi}} \sin kt$  where  $U_k, V_k \in \Delta$ ; the constant term is zero because of part (a) of Proposition 2 (mean zero condition).

**Proposition 3** *Let  $T_k$  be the subspace of  $H$  with “wave number”  $k$ , namely*

$$T_k = \Delta \otimes \text{span}\{\cos kt, \sin kt\}.$$

*Then we have the following.*

- (a)  $H = \bigoplus_{k \geq 1} T_k$ , and the sum is orthogonal with respect to the scalar product.
- (b) For every  $\omega \in \mathbb{R}^{l*}$  we have  $\omega Q T_k \subset T_k$  (ie each subspace  $T_k$  is invariant by  $\omega Q$ ).
- (c) Consider the orthonormal basis  $\{e_i \otimes \frac{1}{\sqrt{\pi}} \cos kt, e_i \otimes \frac{1}{\sqrt{\pi}} \sin kt\}_{i=1}^d$  for  $T_k$ ; in this basis the matrix associated to  $\omega Q|_{T_k}$  is

$$\frac{1}{k}(\omega P) \doteq \frac{1}{k} \begin{pmatrix} 0 & \frac{1}{2}\omega A \\ -\frac{1}{2}\omega A & 0 \end{pmatrix}.$$

**Proof** Point (a) is just Fourier decomposition theorem;  $k \geq 1$  expresses the mean zero condition. For the other two points, let us consider  $u \in T_n$  and  $v \in H$ , with Fourier series respectively  $u = U \frac{1}{\sqrt{\pi}} \cos kt + V \frac{1}{\sqrt{\pi}} \sin kt$  and  $v = \sum_{n \geq 1} U_n \frac{1}{\sqrt{\pi}} \cos nt + V_n \frac{1}{\sqrt{\pi}} \sin nt$ . By a direct computation we have

$$\begin{aligned} & \langle u, \omega Q v \rangle_H \\ &= \int_0^{2\pi} \left\langle \int_0^t U \frac{1}{\sqrt{\pi}} \cos k\tau + V \frac{1}{\sqrt{\pi}} \sin k\tau \, d\tau, \frac{1}{2}\omega A v \right\rangle dt \\ &= \sum_{n \geq 1} \int_0^{2\pi} \frac{1}{k} \langle U \frac{1}{\sqrt{\pi}} \sin kt - V \frac{1}{\sqrt{\pi}} \cos kt, \frac{1}{2}\omega A (U_n \frac{1}{\sqrt{\pi}} \cos nt + V_n \frac{1}{\sqrt{\pi}} \sin nt) \rangle dt \\ &= \frac{1}{k} \int_0^{2\pi} -\frac{1}{\pi} (\cos kt)^2 \langle V, \frac{1}{2}\omega A U_k \rangle + \frac{1}{\pi} (\sin kt)^2 \langle U, \frac{1}{2}\omega A V_k \rangle dt \\ &= \frac{1}{k} (-\langle V, \frac{1}{2}\omega A U_k \rangle + \langle U, \frac{1}{2}\omega A V_k \rangle); \end{aligned}$$

the equality between the second and third row holds because the only nonzero integrals of products of sines/cosines are  $\int_0^{2\pi} \frac{1}{\pi} (\cos kt)^2 dt = \int_0^{2\pi} \frac{1}{\pi} (\sin kt)^2 dt = 1$ .  $\square$

**Remark 2** We notice that for every  $\omega \in (\Delta^2)^*$  the operator  $\omega Q$  is compact. Indeed, it is the limit of a converging series of operators with finite-dimensional image:

$$S_n = \sum_{i=1}^n \omega Q|_{T_i}.$$

Let us prove that the operator norm of  $\omega Q - \omega S_n$  goes to zero. Given a norm-one  $v = \sum_{k \geq 1} v_k$ , we have

$$\begin{aligned} \|(\omega Q - \omega S_n)v\|^2 &= \sum_{k \geq n+1} \|\omega Q|_{T_k} v\|^2 \leq \sum_{k \geq n+1} \frac{4\|\omega P\|_{\text{op}}^2 \|v_k\|^2}{k^2} \\ &\leq \frac{4\|\omega P\|_{\text{op}}^2}{(n+1)^2} \sum_{k \geq n+1} \|v_k\|^2 \leq \frac{4\|\omega P\|_{\text{op}}^2}{(n+1)^2}. \end{aligned}$$

In particular, taking square roots,  $\|(\omega Q - \omega S_n)v\| \leq 4\|\omega P\|_{\text{op}}/(n+1)$ , ie  $\|\omega Q - \omega S_n\|_{\text{op}} \rightarrow 0$ .

We conclude this section by describing the spectrum of the operator  $\omega Q$ . Given  $\omega Q$  we consider as above the skew-symmetric matrix  $\omega A$ , which can be put in canonical form as a block matrix of the form  $\text{diag}(\alpha_1(\omega)J_2, \dots, \alpha_m(\omega)J_2, 0_n)$ , where  $J_2 \in \mathfrak{so}(2)$  is the standard symplectic matrix and  $0_n$  is the  $n \times n$  zero matrix. More precisely on  $\Delta$  we can find an orthonormal basis  $\{X_i, Y_i, Z_j, i = 1, \dots, m, j = 1, \dots, n\}$  for suitable  $m, n \in \mathbb{N}$  satisfying  $2m + n = d$  such that

$$\omega A X_i = -\alpha_i(\omega) Y_i, \quad \omega A Y_i = \alpha_i(\omega) X_i, \quad \omega A Z_j = 0.$$

Let us consider now the operator  $\omega Q$  restricted to  $T_k$ . Using the basis for  $\Delta$  defined above we get the orthogonal basis

$$\left\{ \begin{pmatrix} X_i \\ Y_i \end{pmatrix}, \begin{pmatrix} -Y_i \\ X_i \end{pmatrix}, \begin{pmatrix} X_i \\ -Y_i \end{pmatrix}, \begin{pmatrix} Y_i \\ X_i \end{pmatrix}, \begin{pmatrix} Z_j \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ Z_j \end{pmatrix} \right\},$$

$i = 1, \dots, m, j = 1, \dots, n,$

with eigenvalues  $\frac{\alpha_i(\omega)}{k}, \frac{\alpha_i(\omega)}{k}, -\frac{\alpha_i(\omega)}{k}, -\frac{\alpha_i(\omega)}{k}, 0, 0$  respectively. Thus we have proved:

**Proposition 4** *The nonzero eigenvalues of the operator  $\omega Q$  are  $\pm\alpha_i(\omega)/k$  with multiplicity two, where  $\alpha_i(\omega)$  are the coefficients of the canonical form of  $\omega A$  and  $k \in \mathbb{N}_0$ .*

### 3 Geodesics

#### 3.1 The general structure of geodesics

Here we introduce the *horizontal path space*; given a point  $p \in G$  it is defined as

$$\Omega_p = \{\text{horizontal curves starting at the origin and ending at } p\} = F^{-1}(p).$$

Since in the space  $\Omega$  of horizontal paths we are allowed to compute velocities and their lengths, we define the *energy* functional

$$J: \Omega \longrightarrow \mathbb{R},$$

by associating (as in Riemannian geometry) to each curve  $\gamma$  the integral  $\int_0^{2\pi} \|\dot{\gamma}(t)\|^2 dt$ .

In the case  $p$  is a regular value of the end-point map  $F$ , then a critical point of  $J|_{F^{-1}(p)}$  is called a *normal geodesic*. From now on the word “geodesic” will simply mean “normal geodesic”.

Theorem 9 then allows us to study the structure of geodesics whose endpoint is the generic  $p$ ; in fact they are by definition critical points of the restriction of  $J$  to  $\Omega_p$  and in the case the latter is a Hilbert manifold are defined by the Lagrange multipliers rule. Using the above coordinates  $G \simeq \mathbb{R}^d \oplus \mathbb{R}^l$  we can decompose a vector  $\lambda \in T^*G$  as  $\lambda = \eta + \omega$ , where  $\eta$  is the “horizontal” part and  $\omega \in (\Delta^2)^*$  is the “vertical” one. We have the following proposition.

**Proposition 5** *Let  $u$  be the control associated to a geodesic with Lagrange multiplier  $\lambda \in T^*G$  (ie  $\lambda d_u F = d_u J$ ). Then*

$$u(t) = e^{-(\omega A)t} u_0 \quad \text{and} \quad 2\eta = (e^{-2\pi\omega A} + \mathbf{1})u_0.$$

**Proof** A simple computation using (10) in  $\lambda d_u F = u$  gives

$$u(t) = \eta - (\omega A) \int_0^t u(s) ds + \frac{1}{2} \int_0^{2\pi} (\omega A) u(s) ds.$$

Differentiating the above equation provides  $\dot{u} = -(\omega A)u$ , which in turn implies  $u(t) = e^{-(\omega A)t} u_0$ . Substituting the explicit expression  $u(t) = e^{-(\omega A)t} u_0$  into the same equation and evaluating at zero gives  $u_0 = \eta - \frac{1}{2}u(2\pi) + \frac{1}{2}u_0$ .  $\square$

If  $p$  is not a vertical point and the rank is sufficiently big ( $d > l$ ), then the number of geodesics joining the origin to  $p$  is bounded. In order to prove this statement we need a preliminary lemma.

**Lemma 6** For the generic point  $p \in G$ , the set of normal geodesics starting at the origin and ending at  $p$  is discrete.

**Proof** We can parametrize normal geodesics with their initial covector  $\omega$  and the initial velocity  $u_0$ ; in this way we obtain a smooth map

$$f: \mathbb{R}^l \times \mathbb{R}^d \rightarrow G$$

defined by

$$(\omega, u_0) \mapsto \left( \int_I e^{-t\omega A} u_0 dt, \int_I \left\langle \int_0^t e^{-s\omega A} u_0 ds, A e^{-t\omega A} u_0 \right\rangle dt \right).$$

If  $p$  is a regular value of  $f$  (and the set of such  $p$  is a residual set) then  $f^{-1}(p)$  is a submanifold of  $\mathbb{R}^{d+l}$  of dimension zero (possibly noncompact).  $\square$

**Remark 3** (Semialgebraic geometry and genericity conditions) In the sequel we will be using frequently semialgebraic sets and their properties. The standard reference is the monography by Bochnak, Coste and Roy [4], where definitions and the most important theorems are discussed. We will mainly use the semialgebraic triviality theorem [4, Theorem 9.3.2] and the semialgebraic Sard’s lemma [4, Theorem 9.6.2].

For the notion of genericity, we will say a property is generic if it holds for all points but a set of measure zero. In fact for many statements (especially those involving semialgebraic geometry) properties are generic in a stronger sense (holding for all points but a complement of a countable union of proper algebraic sets). For example, the image  $f(X) \subset Y$  of a semialgebraic set  $X$  of dimension  $d$  under a semialgebraic map  $f: X \rightarrow Y$  has dimension smaller or equal to  $d$ ; if the target space  $Y$  has dimension  $n > d$ , then  $f(X)$  is contained in a proper algebraic set (in particular has measure zero); see [4, Chapter 2.8]. We adopted the weaker viewpoint for simplicity of exposition, leaving to the reader the check of which refined condition can be assumed.

**Proposition 7** Assume that  $d > l$ . Then for the generic  $p \notin \Delta^2$  there is a finite number of geodesics between  $e$  and  $p$ .

**Proof** We already know by the previous lemma that the set of geodesics (ie pairs  $(\omega, u_0)$  such that  $f(\omega, u_0) = p$ ) is discrete for the generic  $p$ . We will exclude that the set of possible Lagrange multipliers is unbounded: this will imply (see below) that the set of initial velocities is bounded as well, hence the set of geodesics to  $p$  is a discrete set in compact, ie it is finite.

If  $\omega$  is a Lagrange multiplier, we can always choose a basis for  $\Delta$  such that the matrix  $\omega A$  appears in canonical form: so we get  $k$  subspaces of  $\Delta$  of dimension two

on which the matrix  $\omega A$  is of the form  $\alpha_i J_2$  (where  $J_2$  is the standard  $2 \times 2$  symplectic matrix on the  $i^{\text{th}}$  eigenspace and  $\alpha_i > 0$ ) for  $i = 1, \dots, k$ ; for every subspace we take the component  $u_0^i$  of the initial velocity  $u_0$ . Since the eigenspaces are orthogonal, the computations for the endpoint of the geodesic can be performed separately: thus we split the horizontal part of the endpoint  $p_\Delta$  into the components of the eigenspaces of  $\omega A$ , say  $p_\Delta^i$ . These components are

$$p_\Delta^i = \int_0^{2\pi} e^{-t\omega A} u_0^i dt = \left[ \frac{1}{\alpha_j} J_2 e^{-t\alpha_j J_2} u_0^i \right]_0^{2\pi} = \frac{1}{\alpha_j} J_2 (e^{-2\pi\alpha_j J_2} u_{0j} - u_{0j})$$

on the eigenspaces of  $\omega A$ . The norm squared of the component  $p_\Delta^i$  is given by

$$\begin{aligned} \|p_\Delta^i\|^2 &= \frac{1}{\alpha_i^2} \langle J_2 (e^{-2\pi\alpha_j J_2} u_0^i - u_0^i), J_2 (e^{-2\pi\alpha_j J_2} u_0^i - u_0^i) \rangle \\ &= \frac{2}{\alpha_i^2} (\|u_0^i\|^2 - \langle u_0^i, e^{-2\pi\alpha_i J_2} u_0^i \rangle) = \frac{2}{\alpha_i^2} \|u_0^i\|^2 (1 - \cos(2\pi\alpha_i)). \end{aligned}$$

The vertical part is more complicated, but we only need to compute its value in the direction of the Lagrange multiplier  $\omega$ :

$$\begin{aligned} \omega(p_{\Delta^2}) &= \frac{1}{2} \int_0^{2\pi} \left\langle \int_0^t e^{-\omega A s} u_0 ds, \omega A e^{-\omega A t} u_0 \right\rangle dt \\ &= \frac{1}{2} \int_0^{2\pi} \left\langle \int_0^t -\omega A e^{-\omega A s} u_0 ds, e^{-\omega A t} u_0 \right\rangle dt \\ &= \frac{1}{2} \int_0^{2\pi} \langle e^{-\omega A t} u_0 - u_0, e^{-\omega A t} u_0 \rangle dt = \pi \|u_0\|^2 - \frac{1}{2} \langle u_0, p_\Delta \rangle. \end{aligned}$$

The equation

$$(12) \quad \omega(p_{\Delta^2}) = \pi \|u_0\|^2 - \frac{1}{2} \langle u_0, p_\Delta \rangle$$

is important at first because it tells that it is enough to prove that the set of Lagrange multipliers is bounded. Indeed, we have

$$\|\omega\| \|p_{\Delta^2}\| \geq \pi \|u_0\|^2 - \frac{1}{2} \langle u_0, p_\Delta \rangle;$$

if the set of Lagrange multipliers is bounded, the set of initial velocities cannot be unbounded otherwise the second term of the inequality would diverge while being limited by a constant.

In order to prove that the set of Lagrange multipliers of the generic endpoint is bounded, we suppose on the contrary that we have a sequence  $\omega_n$  of Lagrange multipliers; this sequence is forced to diverge since the set of the initial data  $(\omega, u_0)$  for the generic endpoint  $p$  is discrete. Up to subsequences, we may assume that the normalized Lagrange multipliers  $\hat{\omega}_n = \omega_n / \|\omega_n\|$  converge to a covector  $\lambda$ ; moreover we can

assume the rescaled eigenvalues  $\hat{\alpha}_{i,n} = \alpha_{i,n}/\|\omega_n\|$  and the corresponding eigenspaces converge. The first one is true since every eigenvalue is bounded by the norm of the matrix which is 1 by definition; the second one follows from the fact that the set of changes of basis is the orthogonal group which is compact. Hence there exists  $c > 0$  such that for every sequence of eigenvalues  $\hat{\alpha}_{i,n}$  that does not converge,  $c\|\omega_n\| \leq \alpha_{i,n}$ .

If we split the second term of the equation (12) into its components given by the eigenspaces of  $\omega A$  we obtain

$$\omega(p_{\Delta^2}) = \sum_{i=1}^k \pi \|u_0^i\|^2 - \frac{1}{2} \langle u_0^i, p_{\Delta}^i \rangle.$$

Notice that each term  $\pi \|u_0^i\|^2 - \frac{1}{2} \langle u_0^i, p_{\Delta}^i \rangle$  is nonnegative: in fact

$$\begin{aligned} \|p_{\Delta}^i\| &= \left\| \int_0^{2\pi} e^{-\omega A s} u_0^i ds \right\| \leq \int_0^{2\pi} \|e^{-\omega A s} u_0^i\| ds = 2\pi \|u_0^i\|, \\ \frac{1}{2} \langle u_0^i, p_{\Delta}^i \rangle &\leq \frac{1}{2} \|u_0^i\| \|p_{\Delta}^i\| \leq \pi \|u_0^i\|^2. \end{aligned}$$

Therefore we have

$$\|p_{\Delta^2}\| \geq \hat{\omega}_n(p_{\Delta^2}) \geq \pi \frac{\|u_{0,n}^i\|^2}{\|\omega_n\|} - \frac{\langle u_{0,n}^i, p_{\Delta} \rangle}{\|\omega_n\|}$$

and if  $\|u_{0,n}^i\|$  diverges, the last term of the previous inequality is asymptotic to its first addendum, which turns out to be bounded. Then for the corresponding component of the horizontal part of the endpoint we have that:

$$\|p_{\Delta,n}^i\|^2 = \frac{2}{\alpha_{i,n}^2} \|u_{0,n}^i\|^2 (1 - \cos(2\pi \alpha_{i,n})) \leq 2 \frac{\|u_{0,n}^i\|^2}{c^2 \|\omega_n\|^2}.$$

If  $\|u_{0,n}^i\|$  is bounded the last term of the inequality converges to 0; if it diverges we have that

$$\|p_{\Delta,n}^i\|^2 \leq 2 \frac{\|u_{0,n}^i\|^2}{c^2 \|\omega_n\|^2} \leq \text{constant} \frac{1}{\|\omega_n\|}$$

which again converges to 0. So, in the limit, the components of the horizontal part of the endpoint are orthogonal to the eigenspaces of the limit matrix  $\lambda A$  with nonzero eigenvalues.

Now, if the matrix  $\lambda A$  is not singular the horizontal part has to be zero contradicting the hypothesis; thus in order to end the argument, we will prove that the horizontal part of the generic point does not belong to the kernel of any degenerate matrix in the linear space  $W = \text{span}\{A_1, \dots, A_l\}$ .

Let us consider the semialgebraic set of the couples of degenerate matrices together with the vectors in their kernels, namely

$$X \doteq \{(\eta, x) \in S^{l-1}(W) \times \Delta \mid \eta Ax = 0, \dim \ker \eta A \geq 2\}$$

stratified in subsets with constant dimension of the kernel. Assuming that the Carnot algebra structure defined by  $W$  is transversal to every subset of the stratification described in Appendix A (condition satisfied by the generic  $W$ ), the stratum  $W_k$  of the matrices of  $k$ -dimensional kernel has codimension  $k \frac{k-1}{2}$ . Therefore the semialgebraic set  $X$  has dimension not greater than  $\max_{k \geq 2} l - 1 - k \frac{k-1}{2} + k \leq l$ . If we project  $X$  on  $\Delta$  we get the set  $\tilde{X}$  of points belonging to the kernel of a degenerate matrix of  $W$ : this is still a semialgebraic set and the projection does not increase the dimension. In particular the set

$$Y = \{(p_\Delta, p_{\Delta^2}) \in G \mid p_\Delta \in \ker \eta A \text{ for some } \eta \neq 0\}$$

has dimension less than  $2l$ , which is smaller than  $d + l = \dim(G)$  under the assumption  $d > l$ . This concludes the proof.  $\square$

**Corollary 8** *If  $p$  is a regular value of  $F$  and  $p \notin \Delta^2$ , the homology of  $\Omega_p \cap \{J \leq s\}$  stabilizes as  $s \rightarrow \infty$ .*

**Proof** If  $p$  is a regular value of  $F$ , then all geodesics to  $p$  are normal and the result follows from the previous proposition.  $\square$

### 3.2 Geodesics ending at $p \in \Delta^2$

In this section we study in more detail the case  $p \in \Delta^2$ . To start with, we prove that for the generic choice of  $p \in \Delta^2$  the set  $\Omega_p$  is a nice object; more precisely we have the following theorem.

**Theorem 9** *For the generic choice of  $W \subset \mathfrak{so}(d)$  and a generic  $p \in \Delta^2$  the topological space  $\Omega_p$  is a Hilbert manifold.*

**Proof** We will prove that the set of critical values of  $q = F|_{F^{-1}(\Delta^2)}$  is contained in a semialgebraic subset of codimension one, from this the conclusion follows.

We first notice that the condition for  $p \in \Delta^2$  to be a critical value of  $q$  is that there exist  $u \in H$  and  $\omega \in (\Delta^2)^*$  such that  $q(u) = p$  and  $dq_u = 2\omega Qu = 0$ . In particular if  $u = \sum_{k=1}^{\infty} u_k$ , where  $u_k \in T_k \simeq \mathbb{R}^{2d}$ , then  $q(u) = \sum_{k=1}^{\infty} \langle u_k, \frac{1}{k}(\omega P)u_k \rangle$  and the condition that  $\omega Qu = 0$ , by invariance of the spaces  $T_k$ , reads  $\omega P u_k = 0$  for all  $k \geq 1$ .

Let us consider the stratification  $\mathfrak{so}(d) = \coprod S_r$ , where  $S_r$  is the set of matrices with constant rank  $r$ . Each  $S_r$  is smooth and over it we have the smooth bundle  $K_r = \{(A, v) \in \mathfrak{so}(d) \times \mathbb{R}^n \mid Av = 0\}$ . Since this stratification is homogeneous, then the generic  $W$  is transversal to all strata and we have an induced stratification

$$W = \coprod_{r=0}^d W_r, \quad W_r = S_r \cap W.$$

Consider now the vector bundle  $K|_{W_r}$  over  $W_r$  (the restriction of  $K_r$ ); notice that for every  $\omega A \in W$  we have  $\ker(\omega P) = \{(x, y) \in \mathbb{R}^{2d} \mid \omega Ax = \omega Ay = 0\}$ . In particular a smooth section of  $K|_{W_r}$  produces also a smooth section of  $\{(\omega, z) \in W_r \times \mathbb{R}^{2d} \mid \omega Pz = 0\}$  over  $W_r$ .

We notice now the following interesting property: if  $p \in \Delta^2$  is a critical point for  $q$  with Lagrange multiplier  $\omega \in W_r$ , then  $p \in (T_\omega W_r)^\perp$ . In fact for every  $k \geq 1$  let us consider a smooth curve  $\omega(t)$  in  $W_r$  with  $\omega(0) = \omega$ ,  $\dot{\omega}(0) = \eta \in T_\omega W_j$  and a smooth  $z(t) \in \ker(\omega(t)\omega P)$  with  $z(0) = u_k$  such that

$$\omega(t)\omega Pz(t) = 0;$$

the existence of such a smooth  $z(t)$  follows from the above discussion. Then deriving the above equation we get  $\dot{\omega}(0)\omega Pz(0) + \omega(0)\omega P\dot{z}(0) = 0$  and considering the scalar product with  $z$  gives

$$\langle z, \dot{\omega}(0)\omega Pz \rangle = \dot{\omega}(0)(q(u_k)) = 0$$

which tells  $\eta(p) = \sum_{k=1}^\infty \eta(q(u_k))$  vanishes for every  $\eta$  in  $T_\omega W_r$ . We consider now the semialgebraic set

$$\Sigma'_1 = \bigcup_{r=0}^d \left( \bigcup_{\omega \in W_r} (T_\omega W_r)^\perp \right).$$

Because of the above argument all critical points of  $q$  are contained in  $\Sigma'_1$  and we want to show this set is of dimension strictly less than  $l$ .

We first check that  $\Sigma'_1$  is indeed semialgebraic (and as a result we compute its dimension); being a finite union, it is enough to prove that each  $\bigcup_{\omega \in W_r} (T_\omega W_r)^\perp$  is semialgebraic. To this end consider  $T_r = \{(\omega, \eta) \in W_r \times W^* \mid (\eta \in T_\omega W_r)^\perp\}$ , which is clearly semialgebraic, and the semialgebraic projection  $\pi_2: T_r \rightarrow W^*$  to the second factor. The image of  $\pi_2$  is semialgebraic and coincides with  $\bigcup_{\omega \in W_r} (T_\omega W_r)^\perp$ .

Now, each  $\bigcup_{\omega \in W_r} (T_\omega W_r)^\perp$  has dimension less than  $\dim(W_r) + l - \dim(W_r) - 1 \leq l - 1$ , where the  $-1$  comes from the fact that by homogeneity  $(T_{\omega/|\omega|} W_r)^\perp = (T_\omega W_r)^\perp$

(notice in particular that the stratum of maximal dimension is open and produces only the zero, since the orthogonal complement of its tangent space is the zero only).

In particular the dimension of  $\Sigma'_1$  is strictly less than  $l$  and the generic  $p \in \Delta^2$  is a regular value of  $q$ . □

**Remark 4** It will be useful for us to set  $\Sigma_1 = \overline{\Sigma}'_1$ . Since the Euclidean closure is smaller than the Zariski closure and dimension of a semialgebraic set is preserved after taking its Zariski closure, then  $\Sigma_1$  is *closed* and contained in a semialgebraic set of codimension one. Points in the complement of  $\Sigma_1$  are an open dense set of regular values of  $q = F|_H$ .

Following up the discussion after Proposition 5, we see that in the case the final point of  $\gamma_u$  is in  $\Delta^2$ , which we know is equivalent to  $\int_I u = 0$ , we can apply the Lagrange multiplier rule to the map  $q$ . More precisely  $u$  is the control associated to a curve which is a geodesic with endpoint  $p \in \Delta^2$  if

$$q(u) = p \quad \text{and} \quad \text{there exists } \omega \text{ such that } \omega Qu = u.$$

The covector  $\omega \in (\Delta^2)^*$  is called the *Lagrange multiplier* associated to  $u$ . Using this remark we see that  $u$  is a geodesics with Lagrange multiplier  $\omega$  if and only if

$$\langle u, v \rangle_H = \langle \omega Qu, v \rangle_H \quad \text{for all } v \in H$$

(here the final point is not specified, ie we are considering all possible geodesics with Lagrange multiplier  $\omega$ ; the final point is recovered by simply applying the expression given in (10) to  $u$ ). Thus in particular we have that for all  $v$  in  $H = \{\int v = 0\}$ ,

$$\int_0^{2\pi} \langle u(t), v(t) \rangle dt = - \int_0^{2\pi} \langle \omega AU(t), v(t) \rangle dt = 0,$$

where  $U(t) = \int_0^t u(s) ds$ . The previous condition tells that  $u + \omega AU$  is a constant function, or equivalently that  $\dot{u} = -\omega Au$ . This implies that  $u$  must be of the form

$$u(t) = e^{-t(\omega A)} u_0,$$

and since  $u \in H$ , then  $u_0$  must be in the *integer* eigenspace of  $i\omega A$ , ie

$$u_0 = e^{-2\pi\omega A} u_0.$$

In particular notice that in this case the complete Lagrange multiplier (ie the one arising by using the map  $F$ , as in Proposition 5), is  $\lambda = (u_0, \omega)$ .

We collect the result for a geodesic ending at  $p \in \Delta^2$  in a lemma.

**Lemma 10** *Let  $u$  be the control associated to a geodesic whose final point is in  $\Delta^2$  with Lagrange multiplier  $\omega$ . Then*

$$u(t) = e^{-t\omega A}u_0 \quad \text{with} \quad u_0 = e^{-2\pi\omega A}u_0.$$

Motivated by the previous lemma, for every  $\omega \in W$  we define

$$E(\omega) = \{v \in \mathbb{R}^d \mid e^{-2\pi\omega A}v = v, v \notin \ker(\omega A)\}.$$

Thus  $E(\omega)$  is the set of possible initial data for *nonconstant* geodesics with Lagrange multiplier  $\omega$ . In particular we see that in order to have a nonzero initial datum the matrix  $i\omega A$  must have nonzero integer eigenvalues, thus the set of all possible Lagrange multipliers coincides with the set

$$\Lambda = \{\omega \in (\Delta^2)^* \mid \det(\omega A - in\mathbf{1}) = 0 \text{ for some } n \in \mathbb{N}_0\}.$$

Notice that  $\Lambda$  is *not* an algebraic (or a semialgebraic set): it is indeed given by the infinite union of algebraic sets  $\Lambda_n = \{\det(\omega A - in\mathbf{1}) = 0\}$ . However  $\Lambda$  is locally algebraic: if we intersect it with a ball, then only a finite number of  $\Lambda_n$  show up.

We discuss now in more detail the structure of the set

$$E = \{(\omega, v) \in W \times \mathbb{R}^d \mid v \in E(\omega)\}.$$

As for  $\Lambda$ , this set is *not* semialgebraic, although if we take the “restriction”  $E|_B$  to a compact semialgebraic set  $B$ , ie we only allow  $\omega$  to vary on a compact semialgebraic set  $B \subset W$ , then  $E|_B$  becomes semialgebraic.

First for every  $\omega$  let us consider the canonical skew-symmetric form of  $\omega A$

$$M(\omega)^T(\omega A)M(\omega) = \text{diag}(\alpha_1(\omega)J_2, \dots, \alpha_s(\omega)J_2, 0, \dots, 0)$$

where  $M(\omega)$  is an orthogonal matrix,  $\alpha_1(\omega), \dots, \alpha_s(\omega)$  are the *positive nonzero* eigenvalues of  $i\omega A$  and  $J_2 \in \mathfrak{so}(2)$  is the canonical symplectic matrix. Thus  $\mathbb{R}^d$  decomposes as the orthogonal sum  $\mathbb{R}^d = V_1 \oplus \dots \oplus V_s \oplus K$ , where the  $V_i$  are the coordinate-two planes and  $K$  is the vector space of the last  $d - 2s$  coordinates. Using this notation we set  $V_i(\omega) = M(\omega)V_i$ : it is the invariant subspace of  $\omega A$  associated to the eigenvalue  $\alpha_i(\omega)$ . In particular we see that

$$E(\omega) = \bigoplus_{\alpha_i(\omega) \in \mathbb{N}_0} V_i(\omega).$$

Associating to each  $v \in E(\omega)$  the control  $e^{t\omega A}v$  defines a linear injection of  $E(\omega)$  into  $H = \bigoplus_{k \geq 1} T_k$ ; in particular the above curve admits the Fourier series decomposition

$$(13) \quad e^{-t\omega A}v = \sum_{k \geq 1} X_k(v) \frac{1}{\sqrt{\pi}} \cos kt - Y_k(v) \frac{1}{\sqrt{\pi}} \sin kt$$

and we denote by  $\phi_k$  the linear map  $v \mapsto (X_k(v), Y_k(v))$  (the  $k^{\text{th}}$  component of the Fourier series of  $e^{t\omega A}v$  written in coordinates  $T_k \simeq \mathbb{R}^{2d}$ ).

Let now  $v \in V_i(\omega)$  with  $\alpha_i(\omega) = k \in \mathbb{N}$ ; in order to get the expression for  $\phi_k(v)$  we compute the Taylor series of  $e^{\omega A t}v$ . We have

$$\begin{aligned} e^{-t\omega A}v &= \left( \sum_{n \geq 0} \frac{(\omega A)^n t^n}{n!} \right)v = \left( \sum_{m \geq 0} \frac{(\omega A)^{2l} t^{2l}}{(2l)!} \right)v - \left( \sum_{m \geq 0} \frac{(\omega A)^{2l+1} t^{2l+1}}{(2l+1)!} \right)v \\ &= \left( \sum_{m \geq 0} \frac{(-1)^l k^{2l} t^{2l}}{(2l)!} \right)v - \left( \sum_{m \geq 0} \frac{(-1)^l k^{2l+1} t^{2l+1}}{(2l+1)!} \right) \frac{\omega A}{k} v \\ &= v \cos kt - \frac{\omega A}{k} v \sin kt, \end{aligned}$$

where in the second line we have used the fact that  $(\omega A)^2 v = -k^2 v$  (being  $V_i(\omega)$  the space associated to the eigenvalue  $\alpha_i(\omega) = k$ ). This computation implies that

$$\phi_k(v) = \sqrt{\pi} \begin{pmatrix} v \\ -\frac{\omega A}{k} v \end{pmatrix}.$$

Notice that the same construction can be performed using the linear immersion  $v \mapsto e^{t\omega A}v$ , which gives the above control with *backward* time; the Lagrange multiplier for the corresponding geodesic is  $-\omega$  and the final point is  $-q(e^{-t\omega A}v)$ .

Slightly abusing notation, we will still denote by  $q$  the map obtained by composing the endpoint map with the linear immersion  $E(\omega) \hookrightarrow H$ .

We recall from Appendix A (see Proposition A.1) that the Lie algebra  $\mathfrak{so}(d)$  of skew-symmetric matrices of size  $d$  is stratified by the sets  $\Gamma_{k|m_1, \dots, m_r}$  consisting of those matrices  $A$  satisfying  $\dim \ker(A) = k$ , and the numbers  $m_1, \dots, m_r$  are natural nonincreasing (they are the multiplicities in the positive spectrum of  $iA$ ). Each one of these strata is smooth and has codimension  $\sum_{i=1}^r (m_i^2 - 1) + \frac{k(k-1)}{2}$ . Since (by construction) this stratification is homogeneous, then a generic choice of the Carnot structure  $W \subset \mathfrak{so}(d)$  will be transversal to all of the strata and will inherit the stratification (in particular respecting codimensions and smoothness).

To deal with integer eigenvalues we need to refine this stratification, unfortunately ending up with an infinite number of strata, but still with nice properties. More

specifically for every  $r \geq 0$  we consider  $\vec{n} = (n_1, \dots, n_r) \in \mathbb{N}^r$  with distinct nonzero components and define the *semialgebraic* set  $\Gamma_{k|m_1, \dots, m_r|\vec{n}}$  as follows: we look at the nonzero components of  $\vec{n}$ , say  $n_{j_1}, \dots, n_{j_\nu}$ , and we take those matrices in  $\Gamma_{k|m_1, \dots, m_r}$  such that the eigenvalue with multiplicity  $m_{j_1}$  equals  $n_{j_1}$ , the one with multiplicity  $m_{j_2}$  equals  $n_{j_2}$ , and so on.

For example  $(0, 2, 2, 0) \in \mathbb{N}^4$  is not an admissible  $\vec{n}$  (since there are two equal nonzero entries); on the other hand if  $\vec{n} = (0, 1, 2, 0)$  then  $\Gamma_{k|m_1, m_2, m_3, m_4|\vec{n}}$  equals the set of all matrices in  $\mathfrak{so}(d)$  with multiplicities of the spectrum  $\{m_1, m_2, m_3, m_4\}$  and with one eigenvalue equal to  $i$  (the imaginary unit) with multiplicity  $m_2$  and another equal to  $2i$  and with multiplicity  $m_3$ .

This operation of fixing some eigenvalues to some integer numbers increases the codimension by  $\nu$  (the number of nonzero components of  $\vec{n}$ ).

Using this notation we see that one can stratify  $\Lambda$  as

$$\Lambda = \coprod_r \coprod_{\{k, m_1, \dots, m_r\}} \coprod_{\{\vec{n} \in \mathbb{N}^r \text{ admissible}\}} \Gamma_{k|m_1, \dots, m_r|\vec{n}} \cap W.$$

As we already noticed, this stratification is not finite even though each stratum is semialgebraic. Nevertheless if we intersect  $\Lambda$  with a compact ball  $B \subset W$  only a finite number of the above strata appear and we are locally semialgebraic.

The next lemma says that for the generic choice of  $W \subset \mathfrak{so}(d)$  and a generic  $p \in \Delta^2$ , the Lagrange multipliers have simple integer spectrum.

**Lemma 11** *For a generic Carnot group structure  $W \subset \mathfrak{so}(d)$ , the generic  $p \in \Delta^2$  is not the final point of a geodesic with Lagrange multiplier  $\omega$  such that  $\omega A$  has multiple eigenvalues in  $i\mathbb{Z}$ .*

**Proof** First we pick the structure  $W$  to be transversal to all strata of the first one of the above stratifications, the one using only the multiplicities in the spectrum (and we know such a property is generic). We stratify now the set  $\Lambda$  by intersecting it with the different strata  $\Gamma_{k|m_1, \dots, m_r}$ ; we are interested only in those strata for which there is at least a multiple integer eigenvalue and we refine the stratification to the above infinite one, by indexing with the admissible  $\vec{n} \in \mathbb{N}^r$ .

Thus we let  $\Lambda_{\{m_j \geq 2\}}$  be one stratum  $\Gamma_{k|m_1, \dots, m_r|\vec{n}}$  such that  $m_j \geq 2$  for at least one index  $j$  with  $n_j \neq 0$ . Each  $\Lambda_{\{m_j \geq 2\}}$  obtained in this way has codimension

$$\text{codim } W \Lambda_{\{m_j \geq 2\}} = \sum_{i=1}^r (m_i^2 - 1) + \frac{k(k-1)}{2} + \nu.$$

We consider now as above the set  $E = \{(\omega, v) \in \Lambda \times \mathbb{R}^d \mid e^{2\pi\omega A}v = v, v \notin \ker(\omega A)\}$ . Over each stratum  $\Lambda_{\{m_j \geq 2\}}$  the set  $E|_{\Lambda_{\{m_j \geq 2\}}}$  is a smooth vector bundle (it is the restriction to  $\Lambda_{\{m_j \geq 2\}}$ , which is smooth, of a smooth vector bundle); moreover  $E|_{\Lambda_{\{m_j \geq 2\}}}$  is semialgebraic as well (here the vector  $\vec{n}$  is fixed).

Consider the smooth map

$$f: E|_{\Lambda_{\{m_j \geq 2\}}} \rightarrow \Delta^2$$

defined by  $(\omega, v) \mapsto q(v)$ , where  $q(v)$  is the final point of the geodesic associated to the control  $v(t) = e^{t\omega A}v$ . We compute the rank of the differential of  $f$  and show that the assumption  $m_j \geq 2$  implies this rank is less than  $l - 1$ ; in particular the image of  $f$  has measure zero. Since the set of final points of geodesics with Lagrange multipliers with multiple eigenvalues in  $i\mathbb{Z}$  is the countable union of the images of the different  $f$  obtained as  $\vec{n}$  varies over  $\mathbb{N}^r$ , the result follows.

The differential of  $f$  restricted to the base  $\Lambda_{\{m_j \geq 2\}}$  has rank smaller than the dimension of  $\Lambda_{\{m_j \geq 2\}}$ , which is  $l - \sum_{i=1}^r (m_i^2 - 1) - ((k(k - 1))/2) - \nu$ . For the rank of  $f$  restricted to the fibers we argue as follows. For every  $\omega \in \Lambda_{\{m_j \geq 2\}}$  we consider the invariant subspaces of  $\omega A$ ; for each natural nonzero eigenvalue  $\lambda_j(\omega)$  of  $i\omega A$  we find an invariant space  $V_j(\omega)$  (the real part of the  $\lambda_j(\omega)$ -eigenspace of  $i\omega A$ ) of dimension  $2\mu_j$ , twice the multiplicity of  $\lambda_j(\omega)$ ; let us call  $I \subset \{1, \dots, r\}$  the index set for such spaces  $V_j(\omega)$  (notice that  $I = \{j_1, \dots, j_\nu\}$ ).

The restriction of  $f$  to each such  $V_j(\omega)$  maps  $\mu_j$  unit circles (lying on distinct orthogonal planes) to a point, in particular the dimension of the kernel of the differential of  $f$  on each  $V_j(\omega)$  is at least  $\mu_j$ . Since the dimension of  $E(\omega)$  is  $2 \sum_{j \in I} \mu_j$ , we see that the rank of the differential of  $f$  on the fibers is at most  $\sum_{j \in I} \mu_j$ . In particular we can bound the rank of the differential of  $f$  as

$$\begin{aligned} \text{rank}(df) &\leq l - \nu - \sum_{j=1}^r (m_j^2 - 1) - \frac{k(k-1)}{2} + \sum_{j \in I} \mu_j \\ &\leq l - \nu - \sum_{j \in I} (m_j^2 - 1 - m_j) - \frac{k(k-1)}{2} \\ &\leq l - \nu - \sum_{j \in I, m_j \geq 2} (m_j^2 - 1 - m_j) - \sum_{j \in I, m_j = 1} (m_j^2 - 1 - m_j) \\ &\leq l - \nu - \sum_{j \in I, m_j \geq 2} 1 - \sum_{j \in I, m_j = 1} (-1) \\ &\leq l - \nu - 1 + (\nu - 1) < l - 1. \end{aligned}$$

This completes the proof. □

We define now the set  $\Sigma_2 \subset \Delta^2$  to be the union of the various  $f(E|_{\Lambda_{k|m_1, \dots, m_r, \vec{n}}})$  where  $\vec{n} = (n_1, \dots, n_r) \in \mathbb{N}^r$  is admissible and  $m_j \geq 2$  for at least one index  $j$  with  $n_j \neq 0$ . The above lemma says that  $\Sigma_2$  is the countable union of semialgebraic sets of codimension at least 2 (in particular, for example,  $\Sigma_2$  has measure zero).

We study now what happens for a  $p \in \Delta^2 \setminus (\Sigma_1 \cup \Sigma_2)$  (because of the above argument such  $p$  is generic). Such a  $p$  has Lagrange multipliers  $\omega$  with simple spectrum, ie  $\omega A$  belongs to a stratum  $\Lambda_{k|m_1, \dots, m_r, \vec{n}}$  with all multiplicities equal to 1; for simplicity of notation we omit the string of multiplicities and denote such stratum simply by  $\Lambda_{\vec{n}}$ . In other words  $\Lambda_{\vec{n}}$  is one of the above strata where all eigenvalues are distinct and we have fixed  $\nu$  of them to be equal to  $in_{j_1}, \dots, in_{j_\nu}$  (the nonzero entries of  $\vec{n}$ ).

**Lemma 12** *Let  $\omega \in \Lambda_{\vec{n}}$  and  $n_1, \dots, n_\nu$  be the nonzero eigenvalues of  $i\omega A$  in  $\mathbb{N}$ ; for  $j = 1, \dots, \nu$  let each  $V_j(\omega)$  be the (two-dimensional) invariant subspace of  $\omega A$  associated to  $n_j$ . Then  $E(\omega)$  splits as the direct orthogonal sum*

$$E(\omega) = \bigoplus_{j=1}^{\nu} V_j(\omega).$$

Moreover the image of  $q|_{V_j(\omega)}$  is a half-line  $l_j^+(\omega)$  and

$$\text{im}(q|_{E(\omega)}) = \text{cone}\{l_1^+(\omega), \dots, l_\nu^+(\omega)\}.$$

**Proof** Recall that the space  $E(\omega)$  is defined to be  $\{v \in \mathbb{R}^d \mid e^{-2\pi\omega A}v = v, v \notin \ker(\omega A)\}$ ; the map that associates to a vector  $v \in E(\omega)$  the curve  $e^{-t\omega A}v$  defines an embedding of  $E(\omega)$  into  $H$  and if  $v \in V_j(\omega)$  then the resulting control must be a linear combination of  $\sin(n_j t)$  and  $\cos(n_j t)$ ; in particular  $V_{n_j}(\omega) \subset T_{n_j}$ . Since the  $T_k$  are pairwise orthogonal for each operator  $Q_1, \dots, Q_l$ , then decomposing  $v \in E(\omega)$  into its pieces  $v = v_1 + \dots + v_\nu$  with  $v_j \in V_j(\omega)$ , we get

$$q(v) = q(v_1) + \dots + q(v_\nu),$$

which proves the image of  $q|_{E(\omega)}$  is the cone spanned by  $\{q(V_1(\omega)), \dots, q(V_\nu(\omega))\}$ ; the orthogonality of the  $V_j$  follows from the one of the  $T_{n_j}$ .

It remains to prove that the image of  $q|_{V_j(\omega)}$  is a half-line. By assumption  $\omega$  belongs to a smooth stratum of codimension  $\nu$  in  $W$  and recalling the definition of  $\Lambda_{n_j} = \{\det(\omega A - in_j \mathbf{1}) = 0\}$ , we have that

$$\Lambda_{\vec{n}} = \bigcap_{j=1}^r \Lambda_{n_j} \cap \Gamma_{k|1, \dots, 1}.$$

The bundle  $\coprod_{\omega \in \Lambda_{n_j}} V_j(\omega)$  is smooth (being the restriction of a smooth bundle). In particular for every  $\eta \in T_\omega \Lambda_{n_j}$  there are curves  $\omega(t) \in \Lambda_{n_j}$  and  $v(t) \in V_j(\omega(t))$  such that  $\omega(0) = \omega$ ,  $\dot{\omega}(0) = \eta$  and  $v(0) = v$ . Deriving the equation  $\omega(t)\omega P v(t) = v(t)$  and taking the inner product with  $v$  we get

$$0 = \left\langle v, \frac{\dot{\omega}(0)\omega P}{k} v \right\rangle = \eta(q(v)),$$

which tells the final point of the geodesic  $u$  associated to  $e^{\omega A t} v$  is orthogonal to  $T_\omega \Lambda_{n_j}$ , hence is it contained in a line. On the other hand since  $\omega$  is the Lagrange multiplier for the geodesic  $u$ ,  $\omega(q(u)) = J(u) > 0$ , which concludes the proof.  $\square$

Everything now is ready for the proof of the theorem that describes the structure of geodesics.

**Theorem 13** *For a generic Carnot group structure  $W \subset \mathfrak{so}(d)$  and a generic  $p \in \Delta^2$  the set  $\Lambda(p)$  of Lagrange multipliers of geodesic whose final point  $p$  is discrete. Moreover every  $\eta \in \Lambda(p)$  belongs to some  $\Lambda_{\vec{n}}$  and the set of all geodesics whose endpoint is  $p$  with Lagrange multiplier  $\eta$  is a compact manifold of dimension  $\nu \leq l$  (where  $\nu$  is the number of nonzero entries of  $\vec{n}$ ) diffeomorphic to the torus  $S^1 \times \cdots \times S^1$  ( $\nu$  times).*

**Proof** By Lemma 11 we know that for the generic choice of  $W \subset \mathfrak{so}(d)$  the generic  $p \in \Delta^2$  is a final point only of geodesics with Lagrange multipliers in  $\Lambda_{\vec{n}}$  for some  $\vec{n} \in \mathbb{N}^{\lfloor d/2 \rfloor}$ .

Moreover for every  $\eta$  in  $\Lambda(p)$  the set of all geodesics with Lagrange multipliers  $\eta$  and final point  $p$  is the preimage of  $p$  under the map  $q: E|_{\Lambda(p)} \rightarrow \Delta^2$ . For every admissible  $\vec{n} \in \mathbb{N}^{\lfloor d/2 \rfloor}$  we consider the semialgebraic set  $F_{\vec{n}}$  defined by

$$F_{\vec{n}} = \{(\omega, p) \in \Lambda_{\vec{n}} \times \Delta^2 \mid p \in \text{im}(q|_{E(\omega)})\}$$

together with the semialgebraic map  $g: F_{\vec{n}} \rightarrow \Delta^2$  defined by  $(\omega, p) \mapsto p$ . Since each point  $(\omega, p)$  in  $F_{\vec{n}}$  has  $\omega$  in  $\Lambda_{\vec{n}}$ , then by Lemma 12 the dimension of  $F_{\vec{n}}$  is at most  $l$ . In fact  $\omega$  varies on a set of dimension  $l - \nu$  and the image of  $q|_{E(\omega)}$  is a cone of dimension at most  $\nu$ . Since  $F_{\vec{n}}$  is semialgebraic we stratify it as  $F_{\vec{n}} = \coprod_{j=1}^s F_{\vec{n},j}$ , where each stratum is smooth semialgebraic of dimension at most  $l$  (in fact here the index  $s$  depends on  $\vec{n}$  as well, but we omit this dependence to simplify notation). Notice that if  $(\omega, p)$  belongs to a stratum of maximal dimension  $l$ , then the cone  $q(E(\omega))$  must have maximal dimension  $\nu$  and  $p$  must be in its interior.

The restriction  $g_{\vec{n},j} = g|_{F_{\vec{n},j}}$  is smooth semialgebraic, thus by the semialgebraic Sard's lemma (see [4]) the set  $C_{\vec{n},j}$  of its critical values is a semialgebraic set of dimension at

most  $l - 1$ . If  $p$  is not one of these critical values then  $g_{\vec{n},j}^{-1}(p)$  consists of isolated points if  $\dim(F_{\vec{n},j}) = l$ , and is empty otherwise.

We set  $\Sigma_3$  to be the union of the critical values of  $g_{\vec{n},j}$  ( $\vec{n}$  varies over  $\mathbb{N}^{\lfloor d/2 \rfloor}$  and  $j$  is the stratifying index for  $F_{\vec{n}}$  as above); such a union, being a countable union of semialgebraic set of dimension at most  $l - 1$  has measure zero, hence points belonging to its complement are generic.

On the other hand  $\Lambda(p)$  equals the union of the projections on  $\Lambda$  of the various  $g_{\vec{n},j}^{-1}(p)$ . If we intersect  $\Lambda$  with a compact ball  $B$ , we hit only a finite number of strata  $\Lambda_{\vec{n}}$  and  $\Lambda(p) \cap B$  is discrete; thus for a generic  $p$  the set  $\Lambda(p)$  is discrete set (possibly infinite).

From Lemma 12 we recall that  $q(E(\omega))$  is the cone spanned by the half-lines  $l_j^+(\omega) = q(E_j(\omega))$ : moreover  $p$  is the sum of nonzero vectors belonging to these half-lines,  $p = p_1 + \dots + p_\nu$  with  $p_j \in l_j^+(\omega)$ . Since the spaces  $E_j(\omega)$  are orthogonal with respect to the operators  $J, Q_1, \dots, Q_l$ , the condition  $q(u) = p$  with  $u \in E(\omega)$  can be split up as  $q(u_j) = p_j$  with  $u_j \in E_j(\omega)$ . The condition  $q(u_j) = p_j$  is equivalent to  $\omega q(u_j) = \omega(p_j)$  since for every covector  $\eta$  orthogonal to  $p_j$  the condition  $\eta q(u_j) = \eta(p_j)$  is automatically satisfied; on the other hand by the Lagrange multiplier condition we have that  $\omega q(u_j) = J(u_j)$ , so that the condition is a positive definite one,  $J(u_j) = \omega(p_j)$ . This implies that every component  $u_j \in E_j(\omega)$  of the geodesic  $u$  going to  $p$  is constrained on a circle  $S^1 \in E_j(\omega)$ , from which follows that the critical manifold  $C_\omega$  is a  $\nu$ -dimensional torus. □

Up to now the genericity assumptions for  $p$  come from Theorem 9, Lemma 11 and Theorem 13: specifically we require  $p \notin \Sigma_1 \cup \Sigma_2 \cup \Sigma_3$  (where  $\Sigma_3$  is defined in the proof of Theorem 13).

### 3.3 Morse–Bott theory

Since critical points of the Energy functional  $J$  are far from being isolated (they arrange themselves into compact manifolds) we cannot apply Morse theory in its standard version. What we need is a generalization of it called *Morse–Bott theory*: it allows to prove the same results as for the ordinary theory if in the definitions nondegenerate critical points are replaced by *nondegenerate critical manifolds* (see [6] and Klingenberg [11]; the basic definitions and results are recalled in Appendix B).

**Theorem 14** *For a generic Carnot Group structure  $W \subset \mathfrak{so}(d)$  and a generic point  $p \in \Delta^2$ , the Energy functional  $J$  restricted to  $\Omega_p$  is a Morse–Bott function.*

**Proof** We assume all the genericity conditions of the previous theorems to be satisfied. We need to prove that we can possibly restrict the set of “good” final points  $p$  to a smaller (but still dense) set for which  $J|_{\Omega_p}$  is Morse–Bott.

Part (a) of the definition of a Morse–Bott function given in Appendix B, immediately follows from Theorem 13.

We proceed to prove part (b). Let us take a Lagrange multiplier  $\eta$ , its corresponding critical manifold  $C_\eta$  and a point  $u \in C_\eta$ . Since we already know that the critical manifold is compact, it remains to show that the Hessian of the energy  $J$  is nondegenerate outside the tangent space to the critical manifold  $C_\eta$ . Since both  $q$  and  $J$  are quadratic, they coincide with their second derivatives. By the Lagrange multiplier rule we get the expression

$$\text{He}_u(q) = (d^2J - \eta D^2q_u)|_{T_u\Omega_p} = (J - \eta q)|_{T_u\Omega_p},$$

for the Hessian, where we have  $T_u\Omega_p = \ker D_uq$ . Notice that to the quadratic form defined by the Hessian it corresponds the self-adjoint operator  $\mathbf{1} - \eta Q$ .

The Hessian is degenerate in the direction of  $v \in T_u\Omega_p$  if and only if

$$\langle v - \eta Qv, x \rangle = 0 \quad \text{for all } x \in T_u\Omega_p,$$

meaning that  $v - \eta Qv$  is orthogonal to  $T_u\Omega_p$ . Since the tangent space  $T_u\Omega_p$  is the orthogonal space to  $\text{span}\{Q_1u, \dots, Q_lu\}$ , then  $v - \eta Qv$  is a linear combination of the vectors  $Q_iu$ , namely  $\lambda_1 Q_1u + \dots + \lambda_l Q_lu$ .

We can eventually restate the degeneracy condition by the equations

$$(14) \quad \begin{cases} v - \eta Qv = \lambda Qu, \\ \langle v, Qu \rangle = 0, \end{cases}$$

where  $\lambda = (\lambda_1, \dots, \lambda_l) \in (\Delta^2)^*$  as above.

If we take  $\lambda = 0$  we see that the degeneracy condition is satisfied by the vectors in

$$E(\eta) \cap T_u\Omega_p = T_uC_\eta,$$

and we have to prove that for the generic choice of  $p$  the degeneracy equation (14) does not admit other solutions.

Let us consider the smooth manifold  $\Lambda_{\vec{n}}$  of Lagrange multipliers containing  $\eta$  (the definition of  $\Lambda_{\vec{n}}$  is given before Lemma 12). Let us call as before  $E_{\vec{n}}$  the fiber bundle with base space  $\Lambda_{\vec{n}}$  and fiber  $E(\omega)$  with  $\omega \in \Lambda_{\vec{n}}$  (see the above discussion).

The tangent space to  $E_{\vec{n}}$  at  $(u, \eta)$  is determined as follows: take a curve  $(u(t), \eta(t))$  in  $E_{\vec{n}}$  based on  $(u, \eta)$  and compute its tangent vector in  $t = 0$ . Differentiating the

condition  $\eta(t)Qu(t) = u(t)$ , we get

$$\dot{u} - \eta Q\dot{u} = \dot{\eta}Qu,$$

which is the same condition as for the degeneracy of the Hessian (the first equation in (14)).

We consider now the smooth semialgebraic map

$$f: E_{\vec{n}} \rightarrow \Delta^2 \quad \text{given by} \quad (\omega, v) \mapsto q(v).$$

The set of regular values of  $f$  is a dense subset of  $\Delta^2$  (it is the complement of a semialgebraic set of codimension at least one): this subset is the good one we want to restrict to. In other words we consider  $\Sigma_4$  to be the union of the set of critical values of the various  $f: E_{\vec{n}} \rightarrow \Delta^2$  as  $\vec{n}$ ; the complement of  $\Sigma_4$  contains generic points.<sup>2</sup> On the preimage of a “good”  $p$  we know that the differential of  $f$  is surjective with rank  $l$ ; moreover,

$$\dim E_{\vec{n}} = \dim \Lambda_{\vec{n}} + \dim E(\eta) = l - \nu + 2\nu = l + \nu.$$

Looking at the dimensions of the domain and the range of  $d_u f$  we get that the kernel has dimension  $\nu$ . On one hand, the kernel of  $d_u f$  is the vector space satisfying both the equations (14) for the degeneracy of the Hessian; on the other hand we have the inclusion

$$E(\eta) \cap T_u \Omega_p \subset \ker d_u f.$$

Since both these spaces have dimension  $\nu$ , they must be equal. It follows that with all the above generic restrictions on  $p$ , the only directions of degeneracy for the Hessian are in  $T_u C_\eta$ .

**Remark 5** We know that the operator  $\eta Q$  is compact, and that the eigenvalues are of the form  $\pm \alpha_i/k$  with  $k \in \mathbb{N}$  nonzero and  $i = 1, \dots, s < \infty$ . Then the eigenvalues of the Hessian of the energy on a critical point are  $1 \pm \alpha_i/k$ ; it follows that the number of negative eigenvalues is always finite, so that the index of every critical manifold is finite.

It remains to prove property (c) of the definition (the Palais–Smale property). Let us consider a sequence  $\{u_k\}$  in  $\Omega_p$  with energy  $\|u_k\|^2$  bounded by  $E$  and such that  $\nabla \psi_{u_k} \rightarrow 0$ , where  $\psi \doteq J|_{\Omega_p}$ .

---

<sup>2</sup>Thus at this stage  $p \in \Delta^2 \setminus (\Sigma_1 \cup \Sigma_2 \cup \Sigma_3 \cup \Sigma_4)$ .

We have that the gradient  $\nabla\psi_u$  is the orthogonal projection of  $\nabla J_u = u$  on the space  $\text{span}\{Q_1u, \dots, Q_lu\}^\perp$ , then if we define  $\pi_u$  to be the orthogonal projection on the space  $\text{span}\{Q_1u, \dots, Q_lu\}$  we have

$$\nabla\psi_u = u - \pi_u u.$$

From  $\{u_k\}$  we can extract a subsequence (we keep calling it  $\{u_k\}$ ) such that  $\|u_k\|^2 \rightarrow L$ . Now we can compute

$$\|\nabla\psi_u\|^2 = \langle u, u \rangle - 2\langle u, \pi_u u \rangle + \langle \pi_u u, \pi_u u \rangle = \langle u, u \rangle - \langle \pi_u u, \pi_u u \rangle,$$

where the second equality follows from  $\langle u, \pi_u u \rangle = \langle \pi_u u, \pi_u u \rangle$  being  $\pi_u$  an orthogonal projection.

Since  $\langle u_k, u_k \rangle \rightarrow L$ , and  $\|\nabla\psi_{u_k}\|^2 = \langle u_k, u_k \rangle - \langle \pi_{u_k} u_k, \pi_{u_k} u_k \rangle \rightarrow 0$ , it follows that

$$\langle \pi_{u_k} u_k, \pi_{u_k} u_k \rangle \rightarrow L.$$

Now, every  $\pi_{u_k} u_k$  is a linear combination of the vectors  $Q_i u_k$ , namely

$$\pi_{u_k} u_k = \sum_{i=1}^l \eta_k^i Q_i u_k,$$

and by computing its norm we get

$$(15) \quad \sum_{i,j=1}^l \eta_k^i \eta_k^j \langle Q_i u_k, Q_j u_k \rangle \rightarrow L.$$

Since the operators  $Q_i$  are compact, they map the bounded sequence  $\{u_k\}$  to a sequence  $\{Q_i u_k\}$  with limit points, so we can iteratively extract converging subsequences (again we keep calling them  $\{u_k\}$ ) and we have  $Q_i u_k \rightarrow v_i$ . In this way (15) becomes

$$\sum_{i,j=1}^l \eta_k^i \eta_k^j \langle v_i, v_j \rangle \rightarrow L,$$

where the coefficients  $\langle v_i, v_j \rangle$  give the scalar product of the whole Hilbert space  $H$  restricted to the finite-dimensional subspace  $V = \text{span}\{v_1, \dots, v_l\}$ . Now we have a bounded sequence of vectors  $\eta_n = \sum_{i=1}^l \eta_n^i v_i$  in  $\mathbb{R}^l$  from which we can extract a converging sequence with limit  $\eta$ .

So the sequence  $\pi_{u_k} u_k = \sum_{i=1}^l \eta_k^i Q_i u_k$  tends to  $v = \sum_{i=1}^l \eta^i v_i$ , and since

$$0 = \lim_{k \rightarrow \infty} \|\nabla\psi_{u_k}\|^2 = \lim_{k \rightarrow \infty} \|u_k - \pi_{u_k} u_k\|^2 = \lim_{k \rightarrow \infty} \|u_k - v\|^2,$$

the sequence  $\{u_k\}$  also tends to  $v$ . Moreover, since  $v_i = \lim_{k \rightarrow \infty} Q_i u_k = Q_i v$  we have that  $v = \eta Q v$  which is the condition for  $v$  to be a critical point of the Energy.  $\square$

## 4 Horizontal-path space and its topology

### 4.1 Paths with bounded energy

Despite Theorem 9 shows that (a priori) only some of the  $\Omega_p$  are Hilbert manifolds, they are in fact all homotopy equivalent to each other; the argument is a simple modification of the standard one for loop spaces and appeared first by Ge in [9]; we recall it here for convenience of the reader.

**Theorem 15** *For every  $p_1, p_2 \in G$  the spaces  $\Omega_{p_1}$  and  $\Omega_{p_2}$  are homotopy equivalent.*

**Proof** It is sufficient to prove that for every  $p \in G$  the space  $\Omega_p$  is homotopy equivalent to  $\Omega_e$ . To this end let  $\gamma_0 \in \Omega_p$  be a fixed *horizontal* path and define the map

$$A: \Omega_e \rightarrow \Omega_p$$

by concatenation of loops in  $\Omega_e$  with  $\gamma$ :  $A(\gamma) = \gamma_0 \gamma$  (velocities have to be rescaled). Let also  $\hat{\gamma}_0$  be  $\gamma_0$  with backward time (it connects  $p$  to  $e$ ); then define

$$B: \Omega_p \rightarrow \Omega_e$$

by concatenation with  $\hat{\gamma}_0$ :  $B(\gamma) = \hat{\gamma}_0 \gamma$ . Let now  $\gamma_\epsilon, \epsilon \in [0, 1]$  be the paths

$$\gamma_\epsilon(t) = \gamma_0(\epsilon(1-t))$$

and  $L_\epsilon: \gamma \rightarrow \gamma_\epsilon \hat{\gamma}_0 \gamma$ . The maps  $L_\epsilon$  give a homotopy between the identity  $L_0 = \text{id}: \Omega_e \rightarrow \Omega_e$  and  $L_1 = AB$ . In a similar way  $BA$  is homotopic to the identity on  $\Omega_p$  and the two spaces are homotopy equivalent.  $\square$

As a corollary we see that  $\Omega_p$  is contractible (in particular all its nonzero Betti numbers vanish).

**Corollary 16** *For every  $p \in G$  the topological space  $\Omega_p$  is contractible.*

**Proof** By the above theorem it is enough to show that  $\Omega_e$  is contractible, and this is obvious since it is given by *homogeneous* equations.  $\square$

We start now the study of Morse–Bott theory of the function  $J|_{\Omega_p} = f$  on  $\Omega_p$ . In the sequel we will use the notation

$$\Omega_p^s \doteq \Omega_p \cap \{J \leq s\}.$$

Proposition 7 tells us that in the case  $p$  is not a vertical point the number of geodesics joining  $e$  to  $p$  is finite; in particular if  $s > 0$  is large enough  $f$  does not have critical points on  $\{f \geq s\}$  and the topology of  $\Omega_p^s$  stabilizes.

**Proposition 17** *If  $p$  is not a vertical point, then for every  $s$  large enough and  $t > 0$  the inclusion*

$$\Omega_p^s \hookrightarrow \Omega_p^{s+t}$$

*is a homotopy equivalence.*

Thus we focus on the case  $p \in \Delta^2$ . As we already mentioned, we *will not* use Morse–Bott theory to give a lower bound on  $b(\Omega_p^s)$  by counting critical manifolds; we will instead use it to reduce the problem to the study of intersection of real quadrics. In fact the following proposition shows that  $\Omega_p^s$  is homotopy equivalent to its boundary  $\partial\Omega_p^s$ ; since the map  $q$  is quadratic, the space  $\partial\Omega_p^s$  is an intersection of infinite-dimensional quadrics (it is given by the quadratic equations  $\|u\|^2 = 2s$  and  $q(u) = p$ ).

**Proposition 18** *For a generic choice of the Carnot group structure  $W \subset \mathfrak{so}(d)$  and a generic point  $p \in \Delta^2$ , for almost every  $s$  the following isomorphism holds:*

$$H_*(\Omega_p^s) \simeq H_*(\partial\Omega_p^s).$$

**Proof** We first notice that the generic  $s$  is not a critical value for the energy. Let us consider now the Morse–Bott function  $g = -J$  and let us denote by  $X^a$  the set  $\{g \leq a\}$ . The critical manifolds of  $g$  are the same as for  $J$ , except that the *index* of each one of them for  $g$  is infinite (since these manifolds have *finite* index for  $J$ , then they must have *infinite* index for  $g$ ).

After passing a critical value  $c$  with corresponding critical manifold  $C$ , the relative homology of the Lebesgue set is given by (ie “the homology changes by”)

$$H_*(X^{c+\delta}, X^{c-\delta}) \simeq H_*(D_C^-, \partial D_C^-).$$

We recall that  $D_C^-$  is the unit disk bundle in the fiber bundle over  $C$  on which the Hessian of the Morse–Bott function is negative definite; see Theorem B.3 and the subsequent discussion from Appendix B. Notice that here the choice of the coefficients field  $\mathbb{Z}_2$  prevents us from the problem of orientability of this bundle.

This relative homology is zero: since the index of  $C$  is infinite, then both  $D_c^-$  and  $\partial D_c^-$  retract on  $C$  (this follows from the fact that the infinite-dimensional sphere is contractible).

We can conclude our proof by observing that even though we pass critical values for  $-J$ , the homology remains the same of  $\partial\Omega_p^s$  until we get the whole  $\Omega_p^s$ .  $\square$

Thus we see that, since  $\Omega_p$  is contractible, each of the Betti numbers  $b_i(\Omega_p^s)$  ( $i > 0$ ) eventually vanishes as  $s \rightarrow \infty$ . Despite this their sum can still grow: the smallest  $i > 0$  for which  $b_i(\Omega_p^s) \neq 0$  will get bigger and bigger and the amount of topology can increase as well: we are interested in understanding quantitatively this phenomenon.

### 4.2 Asymptotic Morse–Bott inequalities

Before giving an explicit bound to  $b(\Omega_p^s)$ , we will see what this bound *would be* if we were to use Morse–Bott inequalities only. The Morse–Bott inequalities bound will follow from the count of the number of critical manifolds with energy less than  $s$ . It turns out that this bound is much worse than the actual one: in fact one has

$$\text{Card}\{\text{critical manifolds with energy less than } s\} = O(s^l)$$

against the actual bound  $b(\Omega_p^s) = O(s^{l-1})$  (this will be proved in the next section).

The following proposition will be fundamental for the sequel: essentially it allows us to turn the direct limits arguments into a quantitative form. Roughly it says that the wave numbers of the controls associated to the geodesics grow at most with the order of their energy.

To deal with this idea, we introduce the following useful notation: for every  $L \in \mathbb{N}$  we define

$$T^L \doteq \bigoplus_{k \leq L} T_k.$$

**Proposition 19** *For the generic choice of the Carnot group structure  $W \subset \mathfrak{so}(d)$  and the generic point  $p \in \Delta^2$  there exists a constant  $c_p > 0$  such that for every geodesic  $\gamma \in \Omega_p \cap \{J \leq s\}$ , its associated control belongs to  $T^{\lfloor sc_p \rfloor}$ .*

In order to prove the previous proposition we first need the following lemma:

**Lemma 20** *For the generic choice of the Carnot group structure  $W \subset \mathfrak{so}(d)$  and the generic point  $p \in \Delta^2$  there exists a constant  $c_p > 0$  such that for every Lagrange multiplier  $\omega$  associated to  $p$ , the following inequality holds:*

$$\frac{\langle \omega, p \rangle}{\|\omega\|} \geq \frac{1}{c_p}.$$

**Remark 6** Let the quantity  $\langle \omega, p \rangle / \|\omega\|$  be the cosine between  $\omega$  and  $p$  times the norm of  $p$ , the lemma says that the Lagrange multipliers for  $p$  are contained in a convex acute cone in  $W$ .

The norm on the space of covectors  $(\Delta^2)^* \cong W$  is the one induced by the inclusion  $W \hookrightarrow \mathfrak{so}(d)$  where  $\langle X, Y \rangle = \text{Trace}(X^T Y)$ .

Before giving the proof of the lemma we show how it implies Proposition 19.

**Proof of Proposition 19** Lemma 20 is equivalent to

$$\|\omega\| \leq c_p \langle \omega, p \rangle.$$

The norm of  $\omega A \in W$  can be written in terms of its eigenvalues  $\alpha_1, \dots, \alpha_s$  as

$$\|\omega\| = \|\omega A\| = \sqrt{2\alpha_1^2 + \dots + 2\alpha_s^2};$$

it follows that every eigenvalue of  $\omega A$  is smaller than the norm of  $\omega A$ . Since  $\omega$  is a Lagrange multiplier, if  $u = u_{k_1} + \dots + u_{k_l}$  then the  $k_j$  are integer eigenvalues of  $i\omega A$ . Since the energy of a geodesic  $u$  associated to  $\omega$  is  $J(u) = \langle \omega, p \rangle$ , we have

$$(16) \quad k_j \leq \|\omega\| \leq c_p \langle \omega, p \rangle \leq c_p s.$$

This completes the proof. □

Now we go back to the proof of Lemma 20.

**Proof of Lemma 20** Suppose on the contrary that the constant bounding  $\langle \omega, p \rangle / \|\omega\|$  from below does not exist, so that we can find a sequence of Lagrange multipliers  $\omega_n$  such that, setting  $\hat{\omega}_n \doteq \omega_n / \|\omega_n\|$ , the sequence  $\hat{\omega}_n(p) \rightarrow 0$ . Since the sequence  $\hat{\omega}_n$  is contained in  $S^{l-1}$  which is compact, we can assume (up to subsequences) that it converges, with limit  $\lambda$  such that  $\langle \lambda, p \rangle = 0$  by hypothesis. Up to subsequences we can also assume that every Lagrange multiplier  $\omega_n$  has the same number of integer eigenvalues (all distinct by Lemma 11), say  $\nu$ . For every Lagrange multiplier  $\omega_n$  we have the cone of the endpoints of the geodesics associated to  $\omega_n$ ; the Lagrange multiplier  $\omega_n$  is contained in the intersection of  $\nu$  hypersurfaces of matrices with constant eigenvalue equal the imaginary integers  $ik_1(n), \dots, ik_r(n)$ . The direct sum  $E(\omega)$  of the associated eigenvalues contains all the geodesics with Lagrange multiplier  $\omega$  and  $q(E(\omega))$  is the cone spanned by the normal vectors to each of these  $\nu$  surfaces (Lemma 12), where the normal vector has to be chosen with positive scalar product with  $\omega_n$  (since the energy is positive).

Let us call  $l_j^+$  the normal vectors to the surfaces with eigenvalue equal to  $k_j(n)$  respectively; up to subsequences again we can assume that every direction  $l_j^+(n)$  converges to some  $l_j^+$ . The point  $p$  is contained in the interior of every cone (Theorem 13) and it can be written as  $p = \sum_{j=1}^v c_j(n)l_j^+(n)$  with  $c_j(n) > 0$  for every  $j$  and every  $n$ . Since  $\langle \lambda, p \rangle = 0$ , we have

$$0 = \langle \lambda, p \rangle = \sum_{j=1}^v \langle \lambda, l_j^+ \rangle = \lim_{n \rightarrow \infty} \sum_{j=1}^v c_j(n) \langle \hat{\omega}_n, l_j^+(n) \rangle;$$

the terms of the sum above must converge to 0 one by one because they are nonnegative. Not every  $c_j(n)$  can converge to 0, otherwise  $p$  would be 0 as well. Therefore at least one of the terms  $\langle \hat{\omega}_n, l_j^+(n) \rangle$  converges to 0; in the limit  $p$  is a linear combination of directions  $l_j^+$  with  $j \in I$  for a set of indexes  $I$  such that  $\langle \lambda, l_j^+ \rangle = 0$  for every  $j \in I$ .

Now we are going to see what happens to Lagrange multipliers associated to directions  $l_j^+$  such that  $\lim_{n \rightarrow \infty} \langle \hat{\omega}_n, l_j^+(n) \rangle = 0$ . Let us take the smooth hypersurface  $S_\alpha \subset W$  of the matrices with an eigenvalue equal to a given  $i\alpha \in i\mathbb{R}$  and all the other eigenvalues different from  $i\alpha$ . Then we take a point  $\omega A \in S_\alpha$  and we compute the angle between  $\omega A$  and the normal vector to the surface  $S_\alpha$  at  $\omega A$ . The surface  $S_\alpha$  can be given as a zero locus of the real valued function  $s_\alpha(\eta) \doteq \det(\eta A - i\alpha \mathbf{1})$  in the even-dimensional case and  $i \det(\eta A - i\alpha \mathbf{1})$  in the odd-dimensional case (here for simplicity we discuss the case  $d$  is even, but the proof for the odd-dimensional case is analogous).

We may assume that the matrices  $A^1, \dots, A^l$  form an *orthonormal* basis for  $W$ ; moreover we can choose an orthonormal basis for  $\Delta$  such that the matrix  $\omega A$  is written in canonical form, ie  $\omega A = \text{diag}(\alpha J_2, \alpha_2 J_2, \dots, \alpha_{d/2} J_2)$ . Now we compute the differential of  $s_\alpha$  at  $\omega A$ :

$$(ds_\alpha)_{\omega A} = \sum_{i=1, j=1}^d \sum_{k=1}^l \frac{\partial s_\alpha}{\partial m_{ij}} \frac{\partial m_{ij}}{\partial \eta_k} d\eta_k = \sum_{i=1, j=1}^d \sum_{k=1}^l \text{adj}(\omega A - i\alpha \mathbf{1})^{ij} a_{ij}^k d\eta_k,$$

where  $m_{ij}$  are the variables for the entries of the matrices,  $\eta_k$  are the coordinates on  $W$  given by the components of the covectors in  $(\Delta^2)^*$ ,  $\text{adj}(\omega A - i\alpha \mathbf{1})^{ij}$  is the  $ij$  entry of the adjugate matrix of  $\omega A - i\alpha \mathbf{1}$  and  $a_{ij}^k$  are the entries of the matrix  $A^k$ . The matrix  $\omega A - i\alpha \mathbf{1}$  takes the form

$$\omega A - i\alpha \mathbf{1} = \text{diag} \left( \begin{pmatrix} -i\alpha & \alpha \\ -\alpha & -i\alpha \end{pmatrix}, \begin{pmatrix} -i\alpha & \alpha_2 \\ -\alpha_2 & -i\alpha \end{pmatrix}, \dots, \begin{pmatrix} -i\alpha & \alpha_{d/2} \\ -\alpha_{d/2} & -i\alpha \end{pmatrix} \right);$$

setting  $\beta \doteq \prod_{i=2}^{d/2} (a_i^2 - a^2)$ , the adjugate matrix is

$$\text{adj}(\omega A - i\alpha \mathbf{1}) = \text{diag} \left( \begin{pmatrix} -i\alpha\beta & \alpha\beta \\ -\alpha\beta & -i\alpha\beta \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \dots, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right),$$

and so we get

$$(ds_\alpha)_{\omega A} = 2\alpha\beta a_{12}^k d\eta_k.$$

Therefore we have

$$\langle (ds_\alpha)_{\omega A}, \omega A \rangle = 2\alpha\beta \omega_k a_{12}^k = 2\alpha^2 \beta.$$

Since the basis  $(A^1, \dots, A^l)$  is orthonormal for  $W$  and so is  $(d\lambda_1, \dots, d\lambda_l)$  for  $W^*$ , the norm  $\|(ds_\alpha)_{\omega A}\|$  is easily computed as

$$\|(ds_\alpha)_{\omega A}\| = 2|\alpha\beta| \sqrt{(a_{12}^1)^2 + \dots + (a_{12}^l)^2} = 2|\alpha\beta| \|a_{12}\|,$$

where  $a_{12} \doteq (a_{12}^1, \dots, a_{12}^l)$ . Now we can compute the cosine of the angle  $\theta$  between  $\omega$  and the normal to  $S_\alpha$  at  $\omega A$ :

$$\cos \theta = \frac{\langle (ds_\alpha)_{\omega A}, \omega A \rangle}{\|(ds_\alpha)_{\omega A}\| \|\omega A\|} = \frac{2\alpha^2 \beta}{2|\alpha\beta| \|a_{12}\| \|\omega\|} = \pm \frac{\alpha}{\|a_{12}\| \|\omega\|}.$$

Let us go back to the directions  $l_j^+$  such that  $\langle \lambda, l_j^+ \rangle = \lim_{n \rightarrow \infty} \langle \hat{\omega}_n, l_j^+(n) \rangle = 0$ ; recall that the imaginary integers  $ik_j(n)$  are the eigenvalue corresponding to the eigenspaces mapped on  $\mathbb{R}l_j^+(n)$ . Notice that the constant  $\|a_{12}\|$  in the previous computation depends on the space  $W$  and on the basis we choose in order to put  $\omega A$  in canonical form, but now the basis is a priori different for every  $\omega_n A$ ; indeed we have that  $\|a_{12}\| \leq \|A\|$ , where the norm  $\|A\|$  is the one induced by the norm on  $\Delta$  on its exterior algebra. It follows that

$$|\langle \hat{\omega}_n, l_j^+(n) \rangle| \geq \left| \frac{\alpha_i(n)}{\|A\| \|\omega_n\|} \right|,$$

and since the first term converges to 0, so does the second one.

Notice that the vectors  $l_j^+(c)$ , orthogonal to the surfaces with eigenvalue equal to  $k_i(n)$ , are orthogonal to the surfaces with eigenvalue equal to  $k_i(n)/\|\omega_n\|$  as well, since the surfaces are the same up to homothety. The point  $p$  is a linear combination of the directions  $l_j^+$  with  $j \in I$  with the property that  $\langle \lambda, l_j^+ \rangle = 0$ : these directions are the images of the limit eigenspaces  $E_j(\lambda)$  of the sequences  $E_j(\hat{\omega}_n)$ . Since their eigenvalues are  $k_j(n)/\|\omega_n\|$  which converge to 0, the eigenvalues of the eigenspaces  $E_j(\lambda)$  for  $j \in I$  are 0. It follows that  $p$  belongs to the image by  $q$  of the kernel of  $\omega Q$ , which means that  $p$  is a critical value for  $q$ , and this is absurd by hypothesis.  $\square$

We prove now the theorem that gives an upper bound for the number of critical manifolds with energy bounded by  $s$ .

**Theorem 21** For the generic choice of the Carnot group structure  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$ ,

$$\text{Card}\{\text{critical manifolds with energy less than } s\} = O(s^l).$$

**Proof** Let us start by considering the vector space  $P_{l,d}$  of real polynomials of degree  $d$  in  $l$  variables. For every  $\nu = 1, \dots, l$  we set  $f = (f_1, \dots, f_\nu) \in P_{l,d}^\nu$  and

$$Y_\nu = \{(f, V, \omega) \in P_{l,d}^\nu \times G(l - \nu, l) \times \mathbb{R}^l \mid \omega \in Z(f) \setminus \text{Sing}(Z(f)), V = (T_\omega Z(f))^\perp, p \in V\}.$$

Since  $Y_\nu$  is semialgebraic, we can consider the semialgebraic projection to the first factor  $\pi: Y_\nu \rightarrow P_{l,d}^\nu$  and stratify  $P_{l,d}^\nu = \coprod_{j=1}^s P_j$  such that  $\pi$  is semialgebraically trivial over each stratum (see [4]). In particular since there are finitely many strata, there exists a number  $\beta_\nu$  such that for every  $f \in P_{l,d}^\nu$ ,

$$b_0(\pi^{-1}(f)) \leq \beta_\nu$$

(only a finite number of fibers, up to semialgebraic homeomorphism, appear). In particular, in the case when there are finitely many  $\omega$  with normals  $V = (T_\omega Z(f))^\perp$  containing  $p$ , this construction implies their number is bounded by  $\beta_\nu$ .

Consider now for every  $n \in \mathbb{N}$  the polynomial  $f_n \in P_{l,d}$  defined by

$$f_n(\omega) = i^d \det(\omega A - i n \mathbf{1})$$

(the  $i^d$  factor has the only scope of turning  $f_n$  into a *real* polynomial in the case  $d$  is odd). We know from Lemma 11 that for the generic choice of  $W \subset \mathfrak{so}(d)$  and  $p \in \Delta^2$  each Lagrange multiplier belongs to some  $Z(f_{k_1}, \dots, f_{k_\nu})$ , where  $\nu$  is the number of positive integer eigenvalues of the matrix  $i\omega A$ . Now if  $u = u_1 + \dots + u_\nu$  is the control associated to a geodesic with final point  $p$  and energy less than  $s$ , Proposition 19 implies

$$k_j \leq s c_p \quad \text{for } j = 1, \dots, \nu$$

for a constant  $c_p$  depending only on  $W$  and  $p$ .

Thus the way to get all *possible* Lagrange multipliers associated to geodesics ending at  $p$  with energy less than  $s$  is by intersecting  $\nu$  of the hypersurfaces  $Z(f_k)$  for

$k = 1, \dots, \lfloor sc_p \rfloor$  and  $\nu = 1, \dots, l$  and considering the points in this intersection where the normal space contains  $p$ . There are

$$\binom{\lfloor sc_p \rfloor}{\nu} = O(s)^\nu$$

possible ways of choosing the hypersurfaces and the above argument implies each such choice can contribute by at most  $\beta_\nu$  Lagrange multipliers.

In particular the set of Lagrange multipliers for  $p$  with energy less than  $s$  is bounded by

$$\text{Card}\{\text{Lagrange multipliers } \omega \text{ such that } \omega(p) \leq s\} \leq \sum_{\nu=1}^l O(s)^\nu = O(s^l).$$

To each critical manifold with energy less than  $s$  there corresponds one and only one Lagrange multiplier whose scalar product with  $p$  is less than  $s$ , hence the conclusion follows. □

As a corollary we derive now Morse–Bott inequalities; as already stated the following bound will be improved in the next section.

**Corollary 22** (Morse–Bott inequalities) *For the generic choice of  $W \subset \mathfrak{so}(d)$  and of the point  $p \in \Delta^2$  we have*

$$b(\Omega_p^s) \leq \sum_{J(C_\omega) \leq s} b(C_\omega) = O(s^l).$$

**Proof** By Lemma 12 and Theorem 13 we know that each critical manifold is an intersection of  $l$  quadrics in  $\mathbb{R}^{2d}$  (it is the preimage of  $p$  under  $q|_{E(\omega)}$ ): in particular the possible homeomorphism types of such manifold are finite and there is a constant  $\beta$  such that  $b(C_\omega) \leq \beta$  for every critical manifold  $C_\omega$ .

Since the sum  $\sum_{J(C) \leq s} b(C)$  contains at most  $O(s^l)$  terms (by Theorem 21), the conclusion follows from Morse–Bott inequalities (see Appendix B). □

### 4.3 Asymptotic total Betti number

The aim of this section is to refine the bound for  $b(\Omega_p^s)$  given in Corollary 22.

We start by proving some technical results.

**Proposition 23** *For the generic choice of  $W \in \mathfrak{so}(d)$  and the generic point  $p \in \Delta^2$  we have: the critical manifolds of  $J$  on  $\Omega_p$  with Energy less than  $s$  coincide with the critical manifolds of  $J$  restricted to  $\Omega_p \cap T^{\lfloor sc_p \rfloor}$ ; their index is the same either if they are considered critical manifolds for  $J$  or if they are considered critical manifolds for  $J$  restricted to  $\Omega_p \cap T^{\lfloor sc_p \rfloor}$ .*

**Proof** We have already proved that there exists a constant  $c_p$  such that all the critical points with energy less than  $s$  are contained in  $\Omega_p \cap T^{\lfloor sc_p \rfloor}$  (Proposition 19); since the spaces  $T_k$  are orthogonal with respect to both the quadratic maps  $J$  and  $q$ , the critical points of  $J|_{\Omega_p \cap T^{\lfloor sc_p \rfloor}}$  are given by the same equations as for critical points of  $J|_{\Omega_p}$  using the Lagrange multipliers rule.

Let  $\omega$  be a Lagrange multiplier for  $p$  with energy less than  $s$ . The Hessian of the energy  $J$  is  $\langle \text{id} - \omega Q \cdot, \cdot \rangle$ , so its eigenvalues are  $1 - \alpha_i(\omega)/k$  with  $k \in \mathbb{N}_0$  and  $\alpha_i(\omega)$  eigenvalues of  $\omega A$  as usual. Therefore we have negative eigenvalues for every integer  $k$  such that  $k < \alpha_i(\omega)$  for at least one  $i$ ; since  $\alpha_i(\omega) \leq \|\omega\|$ , it follows from Lemma 20 (as in the proof of Proposition 19) that

$$k \leq \|\omega\| \leq c_p \langle \omega, p \rangle \leq sc_p$$

with the same constant  $c_p$  as in Proposition 19. □

The next proposition tells that if we want to compute  $b(\Omega_p^s)$  we can restrict ourselves to the intersection with a finite-dimensional subspace of the form  $T^L$  (and indeed we have a quantitative control on the dimension).

**Proposition 24** *For a generic choice of  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$  there exists a constant  $r_p > 0$  such that for every  $m \in \mathbb{N}$ ,*

$$\Omega_p^s \text{ deformation retracts to } \Omega_p^s \cap T^{\lfloor sr_p \rfloor + m}.$$

*In particular:  $H_*(\Omega_p^s) \simeq H_*(\Omega_p^s \cap T^{\lfloor sr_p \rfloor + m})$ .*

**Proof** Given  $L \in \mathbb{N}$  we can define the function  $f_L$ , “distance from  $T^L$ ” in the following way: Every  $u \in \Omega_p^s$  can be uniquely written as  $u = \bar{u} + v$ , where  $\bar{u} \in T^L$  and  $v \in (T^L)^\perp$ ; then we define  $f_L(u) \doteq \|v\|^2$ . Assume that for a suitable  $L$  there are no critical points for  $f_L$  outside  $\Omega_p^s \cap T^L$  and the function  $f_L$  satisfies the Palais–Smale condition: then we can retract the manifold  $f_L^{-1}([0, s]) = \Omega_p^s$  on the sublevel set  $f_L^{-1}(0) = \Omega_p^s \cap T^L$  by Theorem B.2 (the function  $f_L$  is bounded on  $\hat{\Omega}_{\epsilon p}$  since  $f_L \leq J \leq s$ ). The manifolds we are considering have boundary, but we can apply the argument above to  $\Omega_p \cap \{J < s + \delta\}$ : it deformation retracts to  $\Omega_p^s$  again by Theorem B.2 if we choose  $\delta$  small enough not to have new critical values for the Energy; we can indeed choose such  $\delta$  small enough not to have new critical values for  $f_L$  as well.

We are now going to prove that  $f_L$  satisfies the Palais–Smale condition and to find for which  $L$  we are sure not to have critical points for  $f_L$  in  $\Omega_p^s \setminus (\Omega_p^s \cap T^L)$ .

The gradient of the function  $f_L$  at  $u = \bar{u} + v$  is  $2v$  restricted to  $T_u\Omega_p$ : this means that there exists  $\eta \in (\Delta^2)^*$  such that  $\nabla_u F_L = 2v - 2\eta Qu$ . For every critical point  $u$  for  $f_L$  there exists  $\eta$  such that  $v - \eta Qu = v - \eta Qv - \eta Q\bar{u} = 0$ : since the space  $T^L$  and its orthogonal are invariant with respect to  $\eta Q$  we have the equivalent couple of conditions

$$v = \eta Qv, \quad \eta Q\bar{u} = 0;$$

the first condition tells that  $v$  is a geodesic going somewhere, the second one tells that  $q(\bar{u})$  is a critical value for  $q$ .

We need  $f_L$  to satisfy the Palais–Smale condition: having the explicit expression of its gradient  $\nabla f_L$ , we omit this verification whose proof is analogous to the one for  $J$  in Theorem 14.

We are going to prove that there exists a constant  $r_p$  such that if we take  $L = \lfloor sr_p \rfloor$  there are no critical points for  $f_L$  in  $\Omega_p^s \setminus (\Omega_p^s \cap T^L)$ .

Assume that such a constant does not exist: then for every term  $\rho_n$  of a diverging sequence of positive real numbers, we find  $s_n > 0$  and a critical point  $u(n)$  for the function  $f_{\lfloor s_n \rho_n \rfloor}$  outside  $T^{\lfloor \rho_n s_n \rfloor}$  with Energy less than  $s_n$ . By hypothesis  $v(n) \in (T^{\lfloor \rho_n s_n \rfloor})^\perp$  so that its Fourier expansion is  $v(n) = \sum_{k > \rho_n s_n} v_k(n)$ . Recall that  $P \doteq Q|_{T_1}$ ; then we have

$$\begin{aligned} \|q(v(n))\| &= \left\| \sum_{k > \rho_n s_n} \frac{1}{k} \langle Pv_k(n), v_k(n) \rangle \right\| \leq \sum_{k > \rho_n s_n} \frac{1}{k} \|P\| \|u_k(n)\|^2 \\ &\leq \frac{1}{\rho_n s_n} \|P\| \|v(n)\|^2 \\ &\leq \frac{1}{\rho_n s_n} \|P\| s_n = \frac{\|P\|}{\rho_n}. \end{aligned}$$

The last term of the chain of inequalities converges to zero; it follows that  $q(v(n))$  goes to zero as well. Since  $p = q(\bar{u}(n)) + q(v(n))$ , we get

$$\lim_{n \rightarrow \infty} q(\bar{u}(n)) = p,$$

and as we noticed before  $\bar{u}(n)$  is a critical point for  $q$ . This means that we have a sequence of critical values for  $q$  converging to  $p$ , which is impossible since we picked our  $p$  into an open subset of regular values (ie  $p \in \Delta^2 \setminus \Sigma_1$ ). We end the proof by noticing that the condition for having critical points for  $f_{\lfloor r_p s_n \rfloor}$  has to be verified on every  $T_k$  separately (they are orthogonal and invariant with respect to  $Q$ ): since there are no critical points of  $f_{\lfloor r_p s_n \rfloor}$  outside  $T^{\lfloor r_p s_n \rfloor}$ , there are no critical points of the same function restricted to  $T^{\lfloor r_p s_n \rfloor + m}$  (similarly to what happens to the critical points of  $J$

in Proposition 23) and it follows eventually that  $\Omega_p^s \cap T^{\lfloor r_p s n \rfloor + m}$  deformation retracts onto  $\Omega_p^s \cap T^{\lfloor r_p s n \rfloor}$  for every  $m \in \mathbb{N}$ .  $\square$

As a corollary we get the following interesting result, that controls the growth rate of the index of the highest nonzero Betti number of  $\Omega_p^s$ .

**Corollary 25** *For a generic  $W \subset \mathfrak{so}(d)$  and  $p \in \Delta^2$ , there exists a constant  $r_p > 0$  such that*

$$\max_i \{i \mid b_i(\Omega_p^s) \neq 0\} \leq 2d \lfloor r_p s \rfloor.$$

**Proof** By Proposition 24 there exists  $c_p > 0$  such that  $H_*(\Omega_p^s) \simeq H_*(\Omega_p^s \cap T^{\lfloor c_p s \rfloor})$ . In particular  $\Omega_p^s$  has the homology of a semialgebraic subset of  $\mathbb{R}^{2d \lfloor c_p s \rfloor}$  (namely  $\Omega_p^s \cap T^{\lfloor c_p s \rfloor}$ ) and its  $j^{\text{th}}$  Betti number must be zero for  $j > 2d \lfloor c_p s \rfloor$ .  $\square$

Everything is now ready for the proof of the main theorem of this section.

**Theorem 26** (Strong Morse–Bott inequalities) *For the generic choice of  $W \subset \mathfrak{so}(d)$  and of  $p \in \Delta^2$  we have*

$$b(\Omega_p^s) = O(s^{l-1}).$$

**Proof** First we know from Proposition 24 that there exists  $r_p > 0$  such that

$$b(\Omega_p^s) = b(\Omega_p^s \cap T^{\lfloor r_p s \rfloor}) \quad \text{for all } s > 0 \text{ and } m \in \mathbb{N}.$$

It means that  $\Omega_p^s$  has the same Betti numbers as

$$\{x \in T^{\lfloor r_p s \rfloor} \mid q_1(x) = p_1, \dots, q_l(x) = p_l\} \cap \{\|x\|^2 \leq 2s\}.$$

Notice that the dimension of  $T^{\lfloor r_p s \rfloor}$  is  $O(s)$ .

Let us consider the semialgebraic set (a level set of a quadratic map with  $l$  components)

$$X = \{x \in T^{\lfloor r_p s \rfloor} \mid q_1(x) = p_1, \dots, q_l(x) = p_l\}$$

and  $\epsilon > 0$  small enough such that  $X \cap \{2s - \epsilon \leq \|x\|^2 \leq 2s + \epsilon\}$  deformation retracts onto  $X$  (the existence of such  $\epsilon$  is guaranteed by semialgebraic triviality; see [4]). Let also  $A = X \cap \{\|x\|^2 \leq 2s + \epsilon\}$  (which deformation retracts onto  $X \cap \{\|x\|^2 \leq 2s\}$ ) and  $B = X \cap \{2s - \epsilon \leq \|x\|^2\}$ . The Mayer–Vietoris exact sequence of the pair  $(A, B)$  gives  $b(A) + b(B) \leq b(A \cap B) + b(A \cup B)$ , which implies

$$b(\Omega_p^s) = b(A) \leq b(A \cap B) + b(A \cup B).$$

Since  $A \cap B$  deformation retracts onto  $X \cap \{\|x\|^2 = 2s\}$ , then it is defined by  $l$  quadratic equations on a sphere of dimension  $\dim(T^{[r_p^s]}) - 1 = O(s)$ ; on the other hand  $X = A \cup B$  is given by  $l$  quadratic equations in a vector space of dimension  $\dim(T^{[r_p^s]}) = O(s)$ ; hence by Proposition C.1 the total Betti numbers of these spaces are bounded by

$$b(A \cap B) = O(s^{l-1}) \quad \text{and} \quad b(A \cup B) = O(s^{l-1})$$

and the conclusion follows. □

#### 4.4 A topological coarea formula

In this section we compute the first order asymptotic of  $b(\Omega_p^s)$  in  $s$ , for the case  $l = 2$ . It turns out that for a generic choice of the Carnot group structure  $W \subset \mathfrak{so}(d)$  and the point  $p \in \Delta^2$ , the leading term is a *real* number and can be analytically computed using only the data  $W = \text{span}\{A_1, A_2\}$ .

Consider a unit circle  $S^1 \subset W$ . For a generic  $W$  the eigenvalues of  $\omega A$  are distinct and differentiable almost everywhere (the set of matrices in  $\mathfrak{so}(d)$  with multiple eigenvalues is a cone with codimension 3, and the eigenvalues are semialgebraic functions of the parameter  $\omega \in S^1$ ). Thus there exist semialgebraic functions  $\alpha_j: S^1 \rightarrow \mathbb{R}$  such that the  $\alpha_j(\omega)$ , for  $j = 1, \dots, d$ , are the coefficients of the canonical form of  $\omega A$ . Given  $p \in \Delta$  we consider the *rational* functions  $\lambda_j: S^1 \rightarrow \mathbb{R} \cup \{\infty\}$  given by

$$\lambda_j: \omega \mapsto \left| \frac{\alpha_j(\omega)}{\langle \omega, p \rangle} \right| \quad \text{for } j = 1, \dots, d.$$

Notice that when  $\omega$  approaches  $p^\perp$  these functions might explode; that is why they are rational in  $\omega$ . On the other hand they are semialgebraic and differentiable almost everywhere and it makes sense to consider the integral

$$(17) \quad \tau(p) \doteq \frac{1}{2} \int_{S^1} \sum_{j=1}^d |\dot{\lambda}_j(\omega)| - \left| \sum_{j=1}^d \dot{\lambda}_j(\omega) \right| d\omega.$$

The convergence of the integral follows from the fact that where the derivatives of the  $\lambda_j$  explode, they all have the same sign and the integrand vanishes.

The next theorem proves that for a fixed  $p$ , as a function of  $s$ ,

$$b(\Omega_p^s) = \tau(p)s + o(s) \quad \text{as } s \rightarrow \infty.$$

**Theorem 27** *If the corank  $l = 2$ , for a generic choice of  $W \subset \mathfrak{so}(d)$  and  $p \in \Delta^2$  we have*

$$\lim_{s \rightarrow \infty} \frac{b(\Omega_p^s)}{s} = \tau(p).$$

**Proof** In order to compute the asymptotic for  $b(\Omega_p^s)$  for  $s \rightarrow \infty$  we use (4). In fact we have seen that  $\Omega_p^s$  is homotopy equivalent to  $\{v \in H \mid q(v) = p/s, \|v\|^2 = 1\}$  and the latter can be rewritten as

$$\left\{ v \in H \mid q(v) - \frac{\|v\|^2}{s} p = 0 \right\} \cap S$$

where  $S$  is the infinite-dimensional sphere  $S = \{\|v\|^2 = 1\}$ . In particular we can present our set as the intersection of two quadrics in  $H$  on the unit sphere  $S$ ; thus we let  $i_s^-$  denote the index function of the quadratic map  $q - p \cdot \|\cdot\|^2/s$  (we are using the notation of Theorem C.2). In this setting the set  $P \subset S^1$  coincides with  $\{\omega \in S^1 \mid \langle \omega, p \rangle < 0\}$ . In fact if we let  $p = (p_1, p_2)$ , here the two quadrics we are considering are  $q_1 - p_1 \|\cdot\|^2/s$  and  $q_2 - p_2 \|\cdot\|^2/s$  and for every  $\omega$  the self-adjoint operator on  $H$  corresponding to the quadratic form  $\omega q$  is  $\omega Q - \omega(p)\mathbf{1}/s$ . In particular the spectrum of  $\omega Q - \omega(p)\mathbf{1}/s$  is obtained by translating the spectrum of  $\omega Q$  by  $\langle \omega, p \rangle/s$  and since  $\omega Q$  is compact and its spectrum is symmetric with respect to the origin, we see that in order to have finitely many negative eigenvalues we need  $\langle \omega, p \rangle < 0$ .

On the other hand the subspaces  $T_k$  are invariant by both  $\omega Q$  and  $\langle \omega, p \rangle\mathbf{1}/s$ , thus the index function can be computed as

$$\begin{aligned} i_s^-(\omega) &= \sum_{k \geq 1} i^- \left( \omega Q - \frac{\langle \omega, p \rangle}{s} \mathbf{1} \Big|_{T_k} \right) = \sum_{k \geq 1} i^- \left( \frac{\omega Q_0}{k} - \frac{\langle \omega, p \rangle}{s} \mathbf{1} \right) \\ &= \sum_{j=1}^d \left\lfloor \frac{s \alpha_j(\omega)}{\langle \omega, p \rangle} \right\rfloor = \sum_{j=1}^d \lfloor s \lambda_j(\omega) \rfloor, \end{aligned}$$

where in the second line we have used the fact that the spectrum of  $(\omega Q_0)/k - (\langle \omega, p \rangle/s)\mathbf{1}$  is of the form  $(\alpha_j(\omega))/k - (\langle \omega, p \rangle/s)$ . In the sequel we also identify  $P \subset S^1$  with a subset of  $[0, 2\pi]$  in the standard way.

Denoting now by  $\mu(s)$  the number of local maxima of  $i_s^-$  on  $P$ , we see that formula (4) implies

$$(18) \quad b(\Omega_p^s) = 2\mu(s) + 1 - b_0(P_0) = 2\mu(s) + O(1).$$

In fact, using the long exact sequence of the pair  $(P_{j+1}, P_j)$ , we can rewrite

$$b_0(P_{j+1}, P_j) = b_0(P_{j+1}) - b_0(P_j) + b_1(P_{j+1}, P_j);$$

substituting these identities into  $b(\widehat{\Omega}_{\epsilon p}) = 1 + \sum_{j \geq 1} b_0(P_{j+1}, P_j) + b_1(P_{j+2}, P_{j+1})$  we get  $b(\widehat{\Omega}_{\epsilon p}) = 1 - b_0(P_0) + 2 \sum_{j \geq 1} b_1(P_{j+1}, P_j)$ . Since each local maximum of  $i_{\epsilon}^-$  contributes by 1 to one of the  $b_1(P_{j+1}, P_j)$  and  $b(P_0) \leq 1$  (since  $P_0$  is convex), then (18) follows.

To compute the asymptotic of the number of maxima of  $i_\epsilon^-$  we introduce the following auxiliary data. First we let  $\lambda = \sum_{j=1}^d \lambda_j(\omega)$  and notice that this is a semialgebraic function. In particular we can divide  $P$  into a finite number of intervals (arcs)

$$P = (\omega_0, \omega_1] \cup [\omega_1, \omega_2] \cup \dots \cup [\omega_m, \omega_{m+1}] \cup [\omega_{m+1}, \omega_{m+2})$$

such that for every  $j, k$  the functions  $\alpha_j$  as well as  $\alpha$  are monotone on  $(\omega_k, \omega_{k+1})$ . Labeling  $I_k = [\omega_k, \omega_{k+1}]$  we also see that each  $\lfloor s\lambda_j \rfloor$  is monotone on  $I_k$ . On the other hand monotonicity of  $i_s^-$  is granted only where the signs of the derivatives of the  $\lambda_j$  all agree. Since for the generic choice of  $p$  the functions  $\alpha_j$  do not vanish on  $\{\omega_0, \omega_{m+2}\}$  (the orthogonal complement of  $p$  on  $S^1$ ),  $\lambda_j$  approaches infinity when approaching  $\omega_0$  or  $\omega_{m+2}$ ; in particular  $i_s^-$  is monotone on  $I_0$  and  $I_{m+1}$  and has no local maxima on them.

For every  $j \in \{1, \dots, m\}$  let us denote respectively by  $\mu_j(s)$  and  $\sigma_j(s)$  the number of local maxima of  $i_s^-$  on  $I_j$  and the number of subintervals of  $I_j$  where  $i_s^-$  is constant (thus  $\sigma_j(s)$  equals the number of “jumps” of the integer valued function  $i_s^-$  on  $I_j$ ).

For every interval  $I_j = [\omega_j, \omega_{j+1}]$  we see that

$$|i_s^-(\omega_{j+1}) - i_s^-(\omega_j)| = \sigma_j(s) - 2\mu_j(s).$$

In particular summing all these equations and using the fact that  $i_s^-$  is monotone on  $I_0$  and  $I_{m+1}$ , combining with (18) we get

$$\begin{aligned} \frac{b(\Omega_p^s)}{s} &= 2\frac{\mu(\epsilon)}{s} + O(1/s) = 2 \sum_{j=1}^m \frac{\mu_j(s)}{s} + O(1/s) \\ &= \sum_{j=1}^m \frac{\sigma_j(s)}{s} - \sum_{j=1}^m \left| \frac{i_s^-(\omega_{j+1}) - i_s^-(\omega_j)}{s} \right| + O(1/s). \end{aligned}$$

Now we notice that as  $s \rightarrow \infty$ , the function  $i_s^-/s$  converges uniformly to  $\lambda$ ; thus,

$$\begin{aligned} (19) \quad \lim_{s \rightarrow \infty} \sum_{j=1}^m \left| \frac{i_s^-(\omega_{j+1}) - i_s^-(\omega_j)}{s} \right| + O(1/s) &= \sum_{j=1}^m |\lambda(\omega_{j+1}) - \lambda(\omega_j)| \\ &= \int_{\omega_1}^{\omega_{m+1}} |\dot{\lambda}(\omega)| d\omega. \end{aligned}$$

It remains to evaluate  $\lim_s \sum_j \sigma_j(s)/s$ . To this end, we let  $\sigma_j^i(s)$  be the number of jumps of  $\lfloor s\lambda_i \rfloor$  on the interval  $I_j$ . We notice that

$$\sigma_j^i(s) = \sum_{i=1}^d \sigma_j^i(s) + O(1);$$

in fact  $i_s^-$  jumps exactly when one of the  $\lfloor s\lambda_j \rfloor$  jumps and these function all jump at different points (except for the points where two eigenvalues are in resonance, but these are in finite number bounded independently of  $s$ ); we also notice that each function  $\lfloor s\lambda_i \rfloor/s$  converges uniformly to  $\lambda_i$ . Thus we get

$$\begin{aligned}
 (20) \quad \lim_{s \rightarrow \infty} \sum_{j=1}^m \frac{\sigma_j(s)}{s} &= \lim_{s \rightarrow \infty} \sum_{j=1}^m \left( \sum_{i=1}^d \frac{\sigma_j^i(s)}{s} \right) \\
 &= \lim_{s \rightarrow \infty} \sum_{j=1}^m \left( \sum_{i=1}^d \left| \frac{\lfloor s\lambda_i(\omega_{j+1}) \rfloor}{s} - \frac{\lfloor s\lambda_i(\omega_j) \rfloor}{s} \right| \right) \\
 &= \sum_{j=1}^m \left( \sum_{i=1}^d |\lambda_i(\omega_{j+1}) - \lambda_i(\omega_j)| \right) \\
 &= \sum_{j=1}^m \int_{\omega_j}^{\omega_{j+1}} \left( \sum_{i=1}^d |\dot{\lambda}_i(\omega)| \right) d\omega \\
 &= \int_{\omega_1}^{\omega_{m+1}} \left( \sum_{i=1}^d |\dot{\lambda}_i(\omega)| \right) d\omega.
 \end{aligned}$$

Combining (19) and (20) we finally get

$$\lim_{s \rightarrow \infty} \frac{b(\Omega_p^s)}{s} = \int_{\omega_1}^{\omega_{m+1}} \sum_{i=1}^d |\dot{\lambda}_i(\omega)| - \left| \sum_{i=1}^d \dot{\lambda}_i(\omega) \right| d\omega = \int_P \sum_{i=1}^d |\dot{\lambda}_i(\omega)| - \left| \sum_{i=1}^d \dot{\lambda}_i(\omega) \right| d\omega,$$

where the last identity follows from the fact that on  $I_0$  and  $I_{m+1}$  the two functions  $\sum_{i=1}^d |\dot{\lambda}_i(\omega)|$  and  $|\sum_{i=1}^d \dot{\lambda}_i(\omega)|$  are equal. The limit of the statement simply follows by noticing  $\alpha_i(\omega) = \alpha_i(-\omega)$  (ie the positive eigenvalues of  $i\omega A$  are  $\pi$ -periodic).  $\square$

As a corollary we get the following result: it says that the topology of the set of paths reaching the point  $\epsilon p$  with energy  $J \leq 1$  explodes; in particular the number of geodesics gets unbounded as well.

**Corollary 28** For a generic choice of  $W \subset \mathfrak{so}(d)$  and the point  $p \in \Delta^2$ ,

$$\lim_{\epsilon \rightarrow 0} b(\Omega_{\epsilon p} \cap \{J \leq 1\}) = \infty.$$

**Proof** First notice that  $\Omega_{\epsilon p} \cap \{J \leq 1\}$  is homeomorphic to  $\Omega_p \cap \{J \leq 1/\epsilon\}$ ; thus we can apply the above theorem.

In order to prove the limit it is enough to show that for the generic choice of  $W$  and  $p$  the integral  $\tau(p)$  is not zero. Since the integrand function is always nonnegative, it is enough to prove it does not vanish identically. Pick two distinct eigenvalues (functions), say  $i\alpha_1$  and  $i\alpha_2$ , for the family  $\{\omega A\}_{\omega \in S^1}$ . Since these functions are continuous semialgebraic,  $i\alpha_1$  has a maximum point  $\bar{\omega}$  and we can assume this is not a critical point for  $\alpha_2$  also (this is a generic condition). Then in a neighborhood of  $\bar{\omega}$  the derivatives of the corresponding  $\lambda_1$  and  $\lambda_2$  have different signs and the integrand is nonzero.  $\square$

We conclude the section with an example where the topological coarea formula can be computed directly.

**Example 3** (Commuting matrices, corank  $l = 2$ ) Let us fix the corank  $l = 2$ . If the matrices  $A_1$  and  $A_2$  commute, they can be written simultaneously in their canonical form

$$A_i = \text{diag}(v_1^i J_2, \dots, v_k^i, 0_h),$$

where as usual  $J_2$  is the  $2 \times 2$  symplectic matrix and  $0_h$  is the  $h \times h$  zero matrix (possibly with  $h = 0$ ). Setting  $v_j \doteq (v_j^1, v_j^2)$  and given  $\omega \in \mathbb{R}^{2*}$ , the eigenvalues of the matrix  $\omega A$  are  $\pm \langle \omega, v_j \rangle$ . Now we pick a generic  $p \in \Delta^2$ : having parametrized by  $t$  the unit circle in  $\mathbb{R}^{2*}$ , the functions we need in order to compute  $\tau(p)$  are  $\lambda_j(t) \doteq |\langle \omega(t), v_j \rangle / \langle \omega(t), p \rangle|$ , their derivatives being

$$\dot{\lambda}_j(t) = \frac{\text{sgn}(\langle \omega(t), v_j \rangle)}{\text{sgn}(\langle \omega(t), p \rangle)} \cdot \frac{\langle \dot{\omega}(t), v_j \rangle \langle \omega(t), p \rangle - \langle \omega(t), v_j \rangle \langle \dot{\omega}(t), p \rangle}{\langle \omega(t), p \rangle^2}.$$

Since the curve  $\omega(t)$  is the arc length parametrization of the unit circle, for every  $t$  the covectors  $\omega(t), \dot{\omega}(t)$  form an orthonormal basis for  $(\Delta^2)^*$ ; it follows that the term

$$m_j \doteq \langle \dot{\omega}(t), v_j \rangle \langle \omega(t), p \rangle - \langle \omega(t), v_j \rangle \langle \dot{\omega}(t), p \rangle$$

is the determinant of the matrix

$$\begin{pmatrix} p^1 & a_j^1 \\ p^2 & a_j^2 \end{pmatrix}$$

which does not depend on  $t$ . Since the functions  $\lambda_j$  are periodic by  $\pi$ , we can modify the formula (17) by integrating on the semicircle  $\{\langle \omega(t), p \rangle > 0\}$  and eliminating the coefficient of  $\frac{1}{2}$  before the integral sign.

These two remarks allow us to simplify the expression of  $\dot{\lambda}(t)$ :

$$\dot{\lambda}_j(t) = \text{sgn}(\langle \omega(t), v_j \rangle) \cdot \frac{m_j}{\langle \omega(t), p \rangle^2}.$$

Now, in order to have the term  $\tau(p) > 0$ , the integrand of

$$\begin{aligned} \tau(p) &= \int_{\langle \omega(t), p \rangle > 0} \sum_{j=1}^d |\dot{\lambda}_j(\omega)| - \left| \sum_{j=1}^d \dot{\lambda}_j(\omega) \right| d\omega \\ &= \int_{\langle \omega(t), p \rangle > 0} \frac{1}{\langle \omega(t), p \rangle^2} \cdot \left( \sum_{i=1}^d |m_i| - \left| \sum_{i=1}^d \operatorname{sgn}(\langle \omega(t), v_i \rangle) m_i \right| \right) d\omega \end{aligned}$$

must be strictly positive somewhere. This happens if and only if there exist  $i$  and  $j$  such that for some  $t_0$  the terms  $\operatorname{sgn}(\langle \omega(t_0), v_i \rangle) m_i$  and  $\operatorname{sgn}(\langle \omega(t_0), v_j \rangle) m_j$  have opposite sign. If  $m_i \cdot m_j > 0$  we must have  $t_0$  such that  $\operatorname{sgn}(\langle \omega(t), v_i \rangle) \cdot \operatorname{sgn}(\langle \omega(t), v_j \rangle) < 0$ : if such a  $t_0$  does not exist, we have  $\operatorname{sgn}(\langle \omega(t), v_i \rangle) \cdot \operatorname{sgn}(\langle \omega(t), v_j \rangle) > 0$  for all  $t$  such that  $\langle \omega(t), p \rangle > 0$ , meaning that  $v_i$  is proportional to  $v_j$ ; a similar argument holds if  $m_i \cdot m_j < 0$ .

In order to have  $\tau(p) = 0$  the only possibility is that every  $v_i$  is proportional to every other  $v_j$ , which is equivalent to saying that  $A_1$  and  $A_2$  are proportional: this implies that  $\dim \Delta^2 - \dim \Delta = 1$  which contradicts the hypothesis.

### Appendix A: Stratifications of $\mathfrak{so}(d)$

Here we construct a useful stratification of  $\mathfrak{so}(d)$ , generalizing the results of Boscaïn and Gauthier [5, Appendix]; for general results on stratifications and semialgebraic sets the reader is referred to [4].

We are interested in studying the dimensions of the semialgebraic sets with generalized eigenvalues of given multiplicities and with given dimension of the kernel. Every skew-symmetric matrix  $A$  can be written in its canonical form as a block matrix, with blocks on the diagonal of the form

$$\alpha J_2 = \begin{pmatrix} 0 & \alpha \\ -\alpha & 0 \end{pmatrix},$$

( $J_2$  being the canonical symplectic matrix in  $\mathfrak{so}(2)$ ) and a 0-block of the dimension of the kernel. By *generalized eigenvalues* we mean the entries like  $\alpha$ .

We introduce the set

$$\Gamma_{k|m_1, \dots, m_r} \subset \mathfrak{so}(d)$$

defined to be the set of skew-symmetric matrices in  $\mathfrak{so}(d)$  with dimension of the kernel equal to  $k$  and multiplicities of the generalized eigenvalues  $m_1, \dots, m_r$  (with  $m_1 \geq m_2 \geq \dots \geq m_r$ ).

By acting with  $SO(d)$  on a matrix  $A \in \Gamma_{k|m_1, \dots, m_r}$  with eigenvalues  $\alpha_1, \dots, \alpha_r$  (corresponding to the ordered multiplicities), we can put it in the form

$$\text{diag}(\alpha_1 J_{2m_1}, \dots, \alpha_r J_{2m_r}, 0_k).$$

Let us look at the stabilizer  $SO(d)_A$  of  $A$ : first of all it has to fix every eigenspace of  $A$ , since eigenspaces with different eigenvalues are orthogonal. On the kernel  $K$  the stabilizer is the restriction of  $SO(d)$  on  $K$ , ie a copy of  $SO(k)$ ; on the eigenspace with eigenvalue  $\alpha_i$  the restriction of  $SO(d)$  is a copy of  $SO(2m_i)$ , but the stabilizer has to fix the symplectic matrix; it follows that the stabilizer acts as a copy of

$$SO(2m_i) \cap Sp(2m_i) = U(m_i),$$

where  $J_{2n}$  is the symplectic matrix in  $\mathbb{R}^{2n}$  and  $0_k$  is the nullmatrix on  $\mathbb{R}^k$ .

Now it is possible to compute the codimension of the orbit  $\text{Ad}(SO(d))A$  of  $A$  by the adjoint action  $\text{Ad}$  of  $SO(d)$  on  $\mathfrak{so}(d)$  with known stabilizer:

$$\begin{aligned} \text{codim } \text{Ad}(SO(d))A &= \dim \mathfrak{so}(d) - \dim \text{Ad}(SO(d))A \\ &= \dim \mathfrak{so}(d) - \dim SO(d) + \dim SO(d)_A = \dim SO(d)_A. \end{aligned}$$

We know that the stabilizer  $SO(d)_A$  is

$$SO(d)_A = SO(k) \times U(m_1) \times \dots \times U(m_r),$$

and its dimension is

$$\dim SO(d)_A = \frac{k(k-1)}{2} + \sum_{i=1}^r m_i^2.$$

Let us now consider the eigenvalues  $\alpha_i$ : as long as they are distinct (so they preserve their multiplicities) they are smooth functions of the matrices; see Kato [10]. On the set  $\Gamma_{k|m_1, \dots, m_r}$  this condition holds true (by definition), hence we have a smooth map

$$\psi: \Gamma_{k|m_1, \dots, m_r} \rightarrow \mathbb{R}^r$$

given by  $A \mapsto (\alpha_i)$ . This map is indeed a submersion on the open subset  $\mathcal{O}$  of vectors in  $\mathbb{R}^r$  with distinct entries. The fibers of the map  $\psi$  are the orbits of the adjoint action and they are diffeomorphic to a fixed manifold  $SO(d)/SO(d)_A$ ; in particular  $\Gamma_{k|m_1, \dots, m_r}$  is a fiber bundle over  $\mathcal{O}$  with fibers diffeomorphic to  $SO(d)/SO(d)_A$ .

Now we can compute the codimension of  $\Gamma_{k|m_1, \dots, m_r}$  in  $\mathfrak{so}(d)$ :

$$\begin{aligned} \text{codim}_{\mathfrak{so}(d)} \Gamma_{k|m_1, \dots, m_r} &= \text{codim}_{\mathfrak{so}(d)} \text{Ad}(\text{SO}(d))A - r \\ &= \frac{k(k-1)}{2} - r + \sum_{i=1}^r m_i^2 = \frac{k(k-1)}{2} + \sum_{i=1}^r (m_i^2 - 1). \end{aligned}$$

Since we are interested in the matrices with integer eigenvalues, we will need to stratify  $\Gamma_{k|m_1, \dots, m_r}$  in infinite semialgebraic sets with given integer eigenvalues. Take  $\vec{n} = (n_1, \dots, n_r) \in \mathbb{N}^r$  with nonnegative entries such that all the nonzero entries are distinct. Then by  $\Gamma_{k|m_1, \dots, m_r|\vec{n}}$  we will mean the stratum in  $\Gamma_{k|m_1, \dots, m_r}$  with the eigenvalue of multiplicity  $m_i$  equal to  $in_i$  if and only if  $n_i > 0$ . The eigenvalues corresponding to zero entries of  $\vec{n}$  vary in  $\mathbb{R}$ . Since by fixing an eigenvalue we drop the dimension of the stratum by 1, we have the following:

**Proposition A.1** *Given  $\vec{n} \in \mathbb{N}^r$  with the properties described above and with  $v$  nonzero entries, the submanifold  $\Gamma_{k|m_1, \dots, m_r|\vec{n}}$  has codimension  $v$  in  $\Gamma_{k|m_1, \dots, m_r}$ , thus its codimension in the set of all matrices  $\mathfrak{so}(d)$  is*

$$\text{codim}_{\mathfrak{so}(d)} \Gamma_{k|m_1, \dots, m_r|\vec{n}} = \frac{k(k-1)}{2} + \sum_{i=1}^r (\mu_i^2 - 1) + v.$$

## Appendix B: Morse–Bott functions

In this section we give a short review of Morse–Bott theory; the interested reader is referred to the original paper by Bott [6] and to the books [11] and by Chang [8] for more details (especially for the infinite-dimensional case).

We recall that a function  $f: X \rightarrow \mathbb{R}$  on the Hilbert manifold  $X$  is called a *Morse–Bott function* if:

- (1) The critical set is the disjoint union of compact smooth manifolds.
- (2) If  $x$  is a critical point of the critical manifold  $C$  then  $\ker \text{He}_x f = T_x C$ .
- (3) For every sequence  $\{x_k\}$  on  $X$  such that  $f(x_k)$  is bounded and  $\|\nabla f_{x_k}\| \rightarrow 0$ , then the sequence  $\{x_k\}$  has limit points and every limit point is critical for  $f$ .

Condition (3) is usually referred to as *Palais–Smale condition* and is automatically satisfied in the finite-dimensional case. The smooth manifolds of critical points are called *nondegenerate critical manifolds*; notice that admitting only zero-dimensional critical manifolds we get classical Morse functions.

The second condition is equivalent to the nondegeneracy of the Hessian on the normal space  $N_x C$  for  $x \in C$ . The *index* of the critical manifold  $C$  is defined as the maximum of the dimensions of subspaces  $V \subset N_x C$  where the Hessian is negative-definite; since the Hessian is nondegenerate in the all normal bundles, this number does not depend on the point  $x \in C$  and it is denoted by  $\text{ind}(C)$ .

Under this assumption it is still possible to describe what happens to the topology of the sublevels  $X^c \doteq f^{-1}(-\infty, c)$  of the Morse–Bott function  $f$  by increasing  $c$ .

Let us consider a Morse–Bott function  $f: X \rightarrow \mathbb{R}$  together with a Riemannian metric  $g$  on  $X$  (the choice of  $g$  is only a technical convenience and in fact the following results do not depend on it). The first fundamental theorem of Morse theory describe how the sublevels (do not) change when  $c$  increases without passing critical values:

**Theorem B.1** *Let us assume that the smooth function  $f: X \rightarrow \mathbb{R}$  is a Morse–Bott function on  $X$ . If the interval  $[a, b] \subset \mathbb{R}$  does not contain critical values,  $X^b$  is diffeomorphic to  $X^a$ .*

Roughly speaking  $X^b$  is deformed to  $X^a$  along the integral curves of the gradient flow  $\nabla f$ . Moreover if we let  $a$  be a critical value for the Morse–Bott function  $f$  the sublevels are not diffeomorphic to one another anymore, but still there exists a deformation.

**Theorem B.2** *If  $f: X \rightarrow \mathbb{R}$  is a  $C^1$  function satisfying the Palais–Smale condition and  $(a, b] \subset \mathbb{R}$  does not contain critical values,  $X^a$  is a strong deformation retract of  $X^b$ .*

For the proof see [8, Lemma 3.2 in Chapter 1].

It only remains to recall what happens when we pass a critical value for a Morse–Bott function. Given a critical manifold  $C$  we restrict the tangent bundle  $TX$  to  $C$  and consider the subbundle

$$E_C^- = \{\text{directions where the Hessian of the function } f \text{ is negative-definite}\}$$

together with the unit disk bundle  $D_C^- \subset E_C^-$ . With this notation the following theorem generalizes the classical one (the statement we present here is actually the one in [11]).

**Theorem B.3** (Bott) *Let  $f: X \rightarrow \mathbb{R}$  be a Morse–Bott function, and let  $c$  be a critical value. For  $\delta > 0$  sufficiently small the sublevel  $X^{c+\delta} = X \cap \{f \leq c + \epsilon\}$  is homotopic to the sublevel  $X^{c-\epsilon}$  with the unit disk bundle  $D_C^-$  glued along the boundary.*

Since in our setting we consider homology with  $\mathbb{Z}_2$  coefficients, it follows from the Thom isomorphism that

$$H_*(X^{c+\delta}, X^{c-\delta}) \simeq H_*(D_C^-, \partial D_C^-),$$

where in the last equation we allow the critical manifold  $C$  to be nonconnected, in which case we actually have a disjoint union of different bundles (with possibly different rank, corresponding to the possibly different indexes of the components of  $C$ ).

Moreover we can state *Morse–Bott inequalities* in terms of the Poincaré polynomial of  $X^s$  and Morse (Bott) polynomial of  $f$ , which is defined by

$$M_f^s(t) = \sum_{\{C \mid f(C) \leq s\}} P_C(t) t^{\text{ind}(C)},$$

where the sum is taken amongst the critical manifolds  $C$  contained in  $\{f \leq s\}$ ; evaluations of this sum at  $t = 1$  give the following.

**Proposition B.4** (Morse–Bott inequalities) *We have*

$$b(X^s) \leq \sum_{\{C \mid f(C) \leq s\}} b(C).$$

## Appendix C: Cohomology of the intersection of real quadrics

In this section we present useful results from the authors [3; 12; 13] for the study of the Betti numbers of the intersection of real quadrics on the sphere  $S^n$ . The motivating example is the case of the zero locus  $Y$  of one single nondegenerate quadratic form  $q$  on the sphere  $S^n$ : if  $i^-(q)$  denotes the negative inertia index of  $q$ , then

$$Y \simeq S^{i^-(q)-1} \times S^{n-i^-(q)}.$$

In particular we see that the knowledge of the *index* function on the whole line spanned by  $q$  in the space of all quadratic forms determines the topology (since by nondegeneracy  $n - i^-(q) = i^-(-q) - 1$ ).

More generally if we have  $l$  quadratic forms  $q_1, \dots, q_l$  in  $n + 1$  variables, then we consider the function  $\eta \mapsto i^-(\eta q)$ ; in the generic case this function is the restriction of the negative inertia index function to the span of  $q_1, \dots, q_l$  in the space of all quadratic forms. Although the general theory is more detailed, for our purposes we need explicit computations only in the case  $l = 2$  and in the general case it will suffice to have quantitative bounds on the topology of

$$Y = \{x \in S^n \mid q_1(x) = \dots = q_l(x) = 0\}.$$

In the case  $l = 2$  we consider a unit circle  $S^1$  in  $\mathbb{R}^2$  and the restriction  $i^-|_{S^1}$ ; also for  $j \geq 0$  we let

$$P_j = \{\eta \in S^1 \mid i^-(\eta q) \leq j\}.$$

The following formula (1) is proved in [13] and relates the Betti numbers of  $Y$  to the topology of the sets  $P_j$ ; we denote by  $\tilde{b}_j(Y)$  the rank of  $\tilde{H}^j(Y; \mathbb{Z}_2)$ . The general bound (2) for the topology of  $Y$  in the case  $l \geq 2$  is proved in [12]. The reader is referred to [3; 12; 13] for more details.

**Proposition C.1** *If  $Y$  is the intersection of two quadrics on the sphere  $S^n$  and  $0 \leq j \leq n - 3$ , then*

$$(1) \quad \tilde{b}_j(Y) = \tilde{b}_{n-j-1}(S^n \setminus Y) = b_0(P_{j+1}, P_j) + b_1(P_{j+2}, P_{j+1}).$$

Moreover if  $Y$  is defined by  $l \geq 2$  quadratic equations on  $S^n$ , on  $\mathbb{R}P^n$  or in  $\mathbb{R}^n$ , then

$$(2) \quad b(Y) \leq O(n)^{l-1}.$$

It is possible to apply the above technique also in the case  $Y$  is the intersection of quadrics on the unit sphere in some (infinite-dimensional) Hilbert space  $H$ . The main differences for this infinite-dimensional case are the following:  $Y$  must be nonsingular;  $\tilde{b}_i$  denotes the rank of the  $i^{\text{th}}$  Čech cohomology group; the negative inertia index might be infinite for some  $\eta \in S^1$ , but these  $\eta$  are already excluded by the condition  $i^-(\eta) \leq j < \infty$ . With these modifications we have the following result from the first author [1]; formula (4) is the analogue of (1), but the condition that  $H$  is infinite-dimensional allows to remove the restriction on the range for  $j$ .

**Theorem C.2** *Let  $q_1, \dots, q_l$  be continuous quadratic forms on the Hilbert space  $H$  and  $Y$  be their (nondegenerate) common zero locus on the unit sphere. Then*

$$(3) \quad H_*(Y) = \varinjlim_{V \in \mathcal{F}} \{H_*(Y \cap V)\},$$

where  $\mathcal{F}$  denotes the family of all finite-dimensional subspace of  $H$ . Moreover in the case  $l = 2$ ,

$$(4) \quad \tilde{b}_j(Y) = \tilde{b}_0(P_{j+1}, P_j) + \tilde{b}_1(P_{j+2}, P_{j+1}).$$

**Sketch of proof** Since  $Y$  is assumed to be nonsingular, then it has a tubular neighborhood  $U$  in  $H$  and  $H_*(Y) \simeq H_*(U)$ . In particular every singular chain in  $U$  is homotopic to one whose image is contained in a finite-dimensional subspace and (3) follows. To prove (4) we fix a  $j \geq 0$ ; then using (1) we have

$$\tilde{b}_j(Y \cap V) = b_0(P_{j+1}(V), P_j(V)) + b_1(P_{j+2}(V), P_{j+1}(V)),$$

where  $V \subset H$  is a sufficiently big finite-dimensional subspace (the condition required on the dimension is  $\dim(V) - 3 \geq j$ ) and  $P_j(V) = \{\eta \in S^1 \mid i^-(\eta q|_V) \leq j\}$ . Now the sets  $\{P_j(V)\}_{V \in \mathcal{F}}$  are also partially ordered by inclusion: if  $V_1 \subset V_2$ , then  $P_j(V_2) \subset P_j(V_1)$ . It is not difficult to show that under the isomorphism  $\tilde{H}_j(Y \cap V) \simeq H^0(P_{j+1}(V), P_j(V)) \oplus H^1(P_{j+2}(V), P_{j+1}(V))$  the inclusion morphism on the homology  $H_j(Y \cap V_1) \rightarrow H_j(Y \cap V_2)$  is induced by the restriction morphism (see [1]):

$$\bigoplus_{i=0,1} H^i(P_{j+i+1}(V_1), P_{j+i}(V_1)) \rightarrow \bigoplus_{i=0,1} H^i(P_{j+i+1}(V_2), P_{j+i}(V_2)).$$

Since the sets  $\{P_j(V)\}_{V \in \mathcal{F}}$  are Euclidean neighborhood retracts (being semialgebraic sets), then by the continuity property of Čech cohomology,

$$\tilde{H}_*(Y) = \varprojlim_{V \in \mathcal{F}} \{H^*(P_{j+1}(V), P_j(V))\} = \check{H}^*\left(\bigcap_{V \in \mathcal{F}} P_{j+1}(V), \bigcap_{V \in \mathcal{F}} P_j(V)\right).$$

Finally  $P_j$  equals by construction  $\bigcap_{V \in \mathcal{F}} P_j(V)$  and the conclusion follows. □

**Acknowledgements** The first author is supported by Grant of the Russian Federation for the State Support of Researches (agreement number 14.B25.31.0029). The third author has received funding from the European Community’s Seventh Framework Programme ([FP7/2007-2013] and [FP7/2007-2011]) under grant agreement number [258204].

## References

- [1] **A A Agrachëv**, *The topology of quadratic mappings and Hessians of smooth mappings*, from: “Algebra, Topology, Geometry”, (R V Gamkrelidze, editor), Itogi Nauki i Tekhniki 26, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow (1988) 85–124, 162 MR978394 Translated in J. Soviet Math. 49 (1990) 990–1013
- [2] **A A Agrachëv, U Boscain, D Barilari**, *Introduction to Riemannian and sub-Riemannian geometry* (2015) Available at <https://www.imj-prg.fr/~davide.barilari/ABB-SRnotes-100415.pdf>
- [3] **A A Agrachëv, A Lerario**, *Systems of quadratic inequalities*, Proc. Lond. Math. Soc. 105 (2012) 622–660 MR2974201
- [4] **J Bochnak, M Coste, M-F Roy**, *Real algebraic geometry*, Ergeb. Math. Grenzgeb. 36, Springer, Berlin (1998) MR1659509
- [5] **U Boscain, J-P Gauthier**, *On the spherical Hausdorff measure in step 2 corank–2 sub-Riemannian geometry*, SIAM J. Control Optim. 51 (2013) 4450–4462 MR3141746

- [6] **R Bott**, *Nondegenerate critical manifolds*, Ann. of Math. 60 (1954) 248–261 MR0064399
- [7] **R Bott**, *Lectures on Morse theory, old and new*, Bull. Amer. Math. Soc. 7 (1982) 331–358 MR663786
- [8] **K-c Chang**, *Infinite-dimensional Morse theory and multiple solution problems*, Progr. Nonlinear Diff. Eq. Appl. 6, Birkhäuser, Basel (1993) MR1196690
- [9] **Z Ge**, *Horizontal path spaces and Carnot–Carathéodory metrics*, Pacific J. Math. 161 (1993) 255–286 MR1242199
- [10] **T Kato**, *Perturbation theory for linear operators*, Classics in Math., Springer, Berlin (1995) MR1335452
- [11] **W Klingenberg**, *Lectures on closed geodesics*, Ergeb. Math. Grenzgeb. 230, Springer, Berlin (1978) MR0478069
- [12] **A Lerario**, *Complexity of intersection of real quadrics and topology of symmetric determinantal varieties*, to appear in J. of the Eur. Math. Soc.
- [13] **A Lerario**, *Convex pencils of real quadratic forms*, Discrete Comput. Geom. 48 (2012) 1025–1047 MR3000573
- [14] **J Milnor**, *Morse theory*, Annals of Math. Studies 51, Princeton Univ. Press (1963) MR0163331
- [15] **A L Onishchik, È B Vinberg**, *Lie groups and algebraic groups*, Springer Series in Soviet Math., Springer, New York (1990) MR1064110
- [16] **P Pansu**, *Métriques de Carnot–Carathéodory et quasiisométries des espaces symétriques de rang un*, Ann. of Math. 129 (1989) 1–60 MR979599
- [17] **S Semmes**, *An introduction to Heisenberg groups in analysis and geometry*, Notices Amer. Math. Soc. 50 (2003) 640–646 MR1988576

SISSA & IM SO RAS I-34136 Trieste, Italy

Sissa, I-34136 Trieste, Italy

AL: Institut Camille Jordan

69622 Villeurbanne, France

agrachev@sissa.it, alemaudit@gmail.com, anto.lerario@gmail.com

Proposed: Yasha Eliashberg

Received: 22 January 2014

Seconded: Dmitri Burago, Leonid Polterovich

Revised: 16 August 2014