

Recurrent Weil–Petersson geodesic rays with non-uniquely ergodic ending laminations

JEFFREY BROCK
BABAK MODAMI

We construct Weil–Petersson geodesic rays with minimal filling non-uniquely ergodic ending lamination which are recurrent to a compact subset of the moduli space of Riemann surfaces. This construction shows that an analogue of Masur’s criterion for Teichmüller geodesics does not hold for Weil–Petersson geodesics.

30F60, 32G15; 37D40

1 Introduction

The Weil–Petersson (WP) metric on Teichmüller space provides a negatively curved, Riemannian alternative to the more familiar Teichmüller metric, a Finsler metric whose global geometry is not negatively curved in any general sense. While negative curvature allows one to harness a broad range of techniques from hyperbolic geometry, difficulties in implementing these techniques arise from the fact that the WP metric is *incomplete* and that its sectional curvatures approach both 0 and $-\infty$ asymptotically near the completion. Nevertheless, it is useful to draw analogies between these metrics and instructive to determine which of these are robust or obtainable through methods in negative curvature.

As an example, Brock, Masur and Minsky [7] introduced a notion of an *ending lamination* for WP geodesic rays, an analogue of the vertical foliation associated to a Teichmüller geodesic ray. They proved that the ending laminations parametrize the strong asymptote class of *recurrent* WP geodesic rays. Recurrent rays are the rays whose projection to the moduli space recurs to a compact set infinitely often. Brock, Masur and Minsky [8] and Modami [24] initiated a systematic study of the behavior of Weil–Petersson geodesics in terms of their ending laminations and associated *subsurface projection coefficients*. Certain diophantine-type conditions for subsurface projection coefficients give strong control over the trajectories of the corresponding geodesics.

For example, criteria on these coefficients can be given to guarantee that geodesics projected to the moduli space stay in a compact part of the moduli space [8], recur to a

compact part of the moduli space, or diverge in the moduli space [24]. A simple scenario arises from bounding the subsurface coefficients associated to the ending lamination of all proper subsurfaces from above, akin to bounded-type irrational numbers, all of whose continued fraction coefficients are bounded. In this *bounded type* case the projection of the corresponding WP geodesic to the moduli space stays in a compact subset; we say the geodesic is *co-bounded*.

In this paper we prove:

Theorem 1.1 *There are Weil–Petersson geodesic rays in the Teichmüller space with minimal, filling, non-uniquely ergodic ending lamination whose projections to the moduli space are recurrent. Moreover, these rays are not contained in any compact subset of the moduli space.*

The theorem sits in contrast with the following result of H Masur about Teichmüller geodesic rays with (minimal) non-uniquely ergodic vertical foliation. Note that a Teichmüller geodesic ray starting at a point X in the Teichmüller space is determined by a unique holomorphic quadratic differential on X . For the description of Teichmüller geodesics in terms of holomorphic quadratic differentials and the associated vertical and horizontal measured foliations, see eg Rafi [25].

Theorem 1.2 (Masur’s criterion [21, Theorem 1.1]) *Suppose that the vertical foliation of a quadratic differential determining a Teichmüller geodesic ray is not uniquely ergodic. Then the Teichmüller geodesic is divergent in the moduli space.*

Remark 1.3 Masur states and proves the theorem with the assumption that the vertical foliation is minimal. The same argument for each minimal component of the vertical foliation gives [Theorem 1.2](#).

The contrapositive of this theorem ensures that the vertical foliation (lamination) of a recurrent Teichmüller geodesic is uniquely ergodic. Comparing this fact and [Theorem 1.1](#) exhibits an essential disparity between how the behavior of a Teichmüller geodesic is encoded in its vertical foliation (lamination) and how the behavior of a Weil–Petersson geodesic is encoded in its forward ending lamination.

Remark 1.4 We remark that the methods here use explicit strong control over the family of geodesics in the Weil–Petersson metric with *bounded non-annular combinatorics* [8]. We remark that in the low-complexity cases of the five-holed sphere and two-holed torus, the more complete control over Weil–Petersson geodesics obtained

by Brock and Masur [6] allows one to apply [Theorem 2.13](#) to show that any Weil–Petersson geodesic with a filling ending lamination is recurrent. In this setting, then, the mere existence of non-uniquely ergodic filling laminations shows the failure of Masur’s Criterion. Here, we have chosen an explicit constructive approach that naturally generalizes to higher-genus cases.

Acknowledgements We would like to thank Chris Leininger and Kasra Rafi for very useful discussions. We would also like to thank Yair Minsky for continued support and encouragement. Brock was partially supported by NSF grant DMS-1207572.

2 Background

Notation 2.1 Let $f, g: X \rightarrow \mathbb{R}^{\geq 0}$ be two functions. Let $K \geq 1$ and $C \geq 0$ be two constants. We write $f \succ_{K,C} g$ if

$$\frac{1}{K}g(x) - C \leq f(x) \leq Kg(x) + C$$

holds for every $x \in X$.

2.1 The curve complex

Let $S = S_{g,n}$ be a finite-type, orientable surface with genus g and n punctures or boundary components. We define the complexity of the surface by $\xi(S) := 3g - 3 + n$. The curve complex of S , denoted by $\mathcal{C}(S)$, is a flag complex. When $\xi(S) > 1$, each vertex in the complex is the isotopy class of an essential, simple closed curve. An *essential curve* is a curve which is not isotopic to a point, a puncture or a boundary component of S . An edge corresponds to a pair of isotopy classes of simple closed curves with disjoint representatives on the surface. The curve complex is the flag complex obtained from the first skeleton, ie we have a k -dimensional simplex corresponding to any $k + 1$ vertices with an edge between any pair of them. Assigning length one to each edge makes the first skeleton of the curve complex a metric graph. When $\xi(S) = 1$, S is a four-holed sphere or a one-holed torus. The definition of the curve complex is the same, except that an edge corresponds to a pair of isotopy classes of simple closed curves with minimal intersection number 2 (in the four-holed sphere case) or 1 (in the one-holed torus case).

An *essential subsurface* of S is a connected, closed properly embedded subsurface $Y \subseteq S$ so that each boundary curve of Y is either an essential simple closed curve of S or a boundary curve of S , and Y itself is not a three-holed sphere. We frequently consider the inclusion of subcomplexes $\mathcal{C}(Y) \subseteq \mathcal{C}(S)$ induced by restriction.

For an essential annular subsurface Y with core curve α , the curve complex has a slightly more involved definition, but a simple model: it is quasi-isometric to \mathbb{Z} . Formally, let $\langle \alpha \rangle$ be the cyclic subgroup of $\pi_1(S)$ generated by α acting on the Poincaré disk \mathbb{D}^2 , the universal cover of S . Let $\tilde{Y} = \mathbb{D}^2 / \langle \alpha \rangle$ be the annular cover of S to which Y lifts homeomorphically. Let $\hat{Y} = \mathbb{D}^2 \cup \Omega_\alpha / \langle \alpha \rangle$ be the natural compactification of \tilde{Y} , where Ω_α is the complement of the fixed points of α acting on the circle at infinity of \mathbb{D}^2 . Each vertex of $\mathcal{C}(Y)$ corresponds to the homotopy class of an arc connecting the two boundaries of \hat{Y} relative to the boundary. There is an edge between any two vertices corresponding to arcs with disjoint interior. We denote $\mathcal{C}(Y)$ by $\mathcal{C}(\alpha)$ as well. For more detail see [23, Section 2].

We do not distinguish between the isotopy class of a closed curve and any curve representing the class. A *multi-curve* is a collection of pairwise disjoint simple closed curves.

Masur and Minsky [22] showed that the curve complex of a surface S is δ -hyperbolic, where δ depends only on the topological type of the surface. Indeed, it has recently been shown that δ is universal, and can be taken to be the constant 17 [15] (see also [1]).

Notation 2.2 We say that curves $\alpha, \beta \in \mathcal{C}_0(S)$ *overlap* if α and β cannot be realized by disjoint curves on S . If α and β overlap we say that $\alpha \pitchfork \beta$ holds. A curve α overlaps a subsurface Y if α cannot be realized disjoint from Y ; we denote this by $\alpha \pitchfork Y$. Multi-curves σ and σ' overlap if some $\alpha \in \sigma$ and some $\alpha' \in \sigma'$ overlap. Similarly, a multi-curve σ and a subsurface Y overlap if some $\alpha \in \sigma$ and Y overlap.

Let Y and Z be essential subsurfaces. We say that Y and Z *overlap* if $\partial Z \pitchfork Y$ and $\partial Y \pitchfork Z$ hold.

Pants decompositions and markings A *pants decomposition* P is a multi-curve with maximal number of curves $\xi(S)$. A (*partial*) *marking* μ consists of a pants decomposition P and t_α a diameter-1 subset of $\mathcal{C}_0(\alpha)$ for (some) all $\alpha \in P$. The subset of $\mathcal{C}(\alpha)$ can be represented by transversal curves to α on S . We call P the base of the marking and denote it by $\text{base}(\mu)$.

The *pants graph* of S , denoted by $P(S)$, is a graph whose vertices are the pants decompositions. There is an edge between any two pants decompositions that differ by an elementary move. An elementary move on a pants decomposition P fixes all the curves and replaces one curve α with a curve in $S \setminus \{P - \alpha\}$ whose intersection number with α is 1 if $S \setminus \{P - \alpha\}$ is a one-holed torus, and is 2 if $S \setminus \{P - \alpha\}$ is a four-holed sphere. Assigning length one to each edge we obtain a metric graph.

Laminations and foliations Fix a complete hyperbolic metric on S . A *geodesic lamination* λ is a closed subset of S consisting of disjoint, complete, simple geodesics. Each one of the geodesics is called a leaf of λ . Let $\tilde{S} = \mathbb{D}^2$ be the universal cover of S . Denote the circle at infinity of the Poincaré disk \mathbb{D}^2 by \tilde{S}_∞ . Let $M_\infty(S)$ denote $(\tilde{S}_\infty \times \tilde{S}_\infty \setminus \Delta) / \sim$, where Δ is the diagonal and \sim is the equivalence relation generated by $(x, y) \sim (y, x)$. Since the geodesics in \mathbb{D}^2 are parametrized by points of M_∞ the preimage of a geodesic lamination determines a closed subset of $M_\infty(S)$ which is invariant under the action of $\pi_1(S)$. We denote the space of geodesic laminations on S equipped with the Hausdorff topology of closed subsets of $M_\infty(S)$ by $\mathcal{GL}(S)$. The space $\mathcal{GL}(S)$ is a compact space. For more detail see [10, Section I.4]. A *transverse measure* m on λ is a measure on the set of arcs on S which is invariant under isotopies of S preserving λ . The measure of an arc a such that $a \subset \lambda$ or $a \cap \lambda = \emptyset$ is 0 and otherwise the measure of a is positive. A pair of a geodesic lamination λ and a transverse measure m on λ is a *measured (geodesic) lamination*, denoted by $\mathcal{L} = (\lambda, m)$. We say that λ is the *support* of the measured lamination. We denote the space of measured laminations of S equipped with the weak* topology by $\mathcal{ML}(S)$. The space of projective measured laminations $\mathcal{PML}(S)$ is the quotient of $\mathcal{ML}(S)$ by the natural action of \mathbb{R}^+ rescaling the measures, equipped with the quotient topology. For any $\mathcal{L} \in \mathcal{ML}(S)$, let $[\mathcal{L}]$ denote the projective class of \mathcal{L} .

A geodesic lamination λ is *minimal* if every leaf of λ is dense in λ . The geodesic lamination λ *fills the surface* S or is *filling* if $S \setminus \lambda$ is the union of topological disks and annuli with core curve isotopic to a boundary curve of S . Equivalently, if for any simple closed curve α , and any transverse measure m on λ , we have $i(\alpha, (\lambda, m)) > 0$. Here

$$i: \mathcal{ML}(S) \times \mathcal{ML}(S) \rightarrow \mathbb{R}^{\geq 0}$$

denotes the natural extension of the intersection number of curves to the space of measured geodesic laminations; see [3].

Given $[\mathcal{L}] \in \mathcal{PML}(S)$, let $|\mathcal{L}|$ be the support of \mathcal{L} . Then taking the quotient

$$\mathcal{PML}(S) / |\cdot|$$

of $\mathcal{PML}(S)$ by forgetting the measure, the *ending lamination space*

$$\mathcal{EL}(S) \subset \mathcal{PML}(S) / |\cdot|$$

is the image of projective measured laminations with minimal filling support equipped with the quotient topology of the topology of $\mathcal{PML}(S)$.

Recall that the curve complex of S is a δ -hyperbolic space. The following result of Klarreich describes the Gromov boundary of the curve complex.

Proposition 2.3 [17] *There is a homeomorphism Φ from the Gromov boundary of $\mathcal{C}(S)$ equipped with its standard topology to $\mathcal{EL}(S)$. Let $\{\alpha_i\}_{i=0}^\infty$ be a sequence of curves in $\mathcal{C}_0(S)$ that converges to a point x in the Gromov boundary of $\mathcal{C}(S)$. Regarding each α_i as a projective measured lamination, any accumulation point of the sequence $\{\alpha_i\}_{i=0}^\infty$ in $\mathcal{PML}(S)$ is supported on $\Phi(x)$.*

A singular foliation \mathcal{F} on S is a foliation of the complement of a finite set of points in S called singular points. At a regular (not singular) point, \mathcal{F} is locally modeled on an open set $U \subset \mathbb{C}$ containing the origin whose leaves are the horizontal coordinate lines. More precisely, there is a coordinate chart $x + iy$ such that the leaves of \mathcal{F} are the trajectories given by $y = \text{constant}$. At singular points the foliation is locally modeled on an open set $U \subset \mathbb{C}$ containing the origin whose leaves are the trajectories along which the real-valued 1-form $\text{Im}(\sqrt{z^k} dz^2)$ vanishes, where $k \in \mathbb{N}$. The singular point is mapped to the origin. A foliation is *minimal* if any half leaf of the foliation is dense in the surface.

A *transverse measure* on a singular foliation \mathcal{F} is a measure on the collection of arcs in the surface transversal to \mathcal{F} which is invariant under isotopies of the surface that preserve the foliation.

A pair consisting of a foliation and a transverse measure on the foliation is a *measured foliation*. Given a foliation \mathcal{F} , let $x + iy$ be a coordinate chart as above. Then $|dy|$ defines a transverse measure on \mathcal{F} .

We denote the space of measured foliations of the surface S equipped with the weak* topology by $\mathcal{MF}(S)$. For more detail, see [13, Exposé 5].

There is a one-to-one correspondence between measured laminations and measured foliations up to Whitehead moves and isotopies of foliations on a surface [19]. A lamination is minimal if and only if the corresponding foliation is minimal; see [19, Theorem 2].

Subsurface coefficients Let $Y \subseteq S$ be an essential non-annular subsurface. Masur and Minsky [22] define the *subsurface projection map*

$$\pi_Y: \mathcal{GL}(S) \rightarrow \mathcal{PC}_0(Y)$$

that assigns to $\lambda \in \mathcal{GL}(S)$ the subset of $\mathcal{C}_0(S)$ denoted by $\pi_Y(\lambda)$, as follows: Fix a complete hyperbolic metric on S and realize λ and ∂Y geodesically. If λ does not intersect Y , then define $\pi_Y(\lambda) = \emptyset$. Now suppose that λ intersects Y . Let $\lambda \cap Y$ be the intersection locus of λ and the subsurface Y . Consider isotopy classes of arcs in $\lambda \cap Y$ with endpoints on ∂Y or at cusps of the hyperbolic surface, where the endpoints

of arcs are allowed to move in ∂Y . For any arc a (up to isotopy) take the essential boundary curves of a regular neighborhood of $a \cup \partial Y$ in Y . The union of these curves where we select one arc from each isotopy class and the closed curves in $\lambda \cap Y$ is $\pi_Y(\lambda)$. Note that the diameter of $\pi_Y(\lambda)$ viewed as a subset of $\mathcal{C}(Y)$ is at most 2.

Let Y be an essential annular subsurface with core curve α . Denote the natural compactification of the annular cover of S to which Y lifts homeomorphically by \hat{Y} . Given a geodesic lamination λ , the projection of λ to Y is the set of component arcs of the lift of λ to \hat{Y} which connect the two boundaries of \hat{Y} . We denote the projection map by either π_Y or π_α . For more detail see [23, Section 2].

Note that since $\mathcal{C}_0(S) \subset \mathcal{GL}(S)$, we have in particular the subsurface projection map

$$\pi_Y: \mathcal{C}_0(S) \rightarrow \mathcal{PC}_0(Y).$$

Given a multi-curve σ and an essential subsurface Y , the projection of σ onto Y is the union of the projections $\pi_Y(\alpha)$ where $\alpha \in \sigma$. For a partial marking μ , if the subsurface Y is not an annulus with core curve in $\text{base}(\mu)$, then $\pi_Y(\mu) = \pi_Y(\text{base}(\mu))$. If Y is an annulus with core curve $\alpha \in \text{base}(\mu)$, then $\pi_Y(\mu)$ is the set of transversal curves to α in μ .

Let μ and μ' be either partial markings or laminations. Let $Y \subseteq S$ be an essential subsurface. The Y subsurface coefficient of μ and μ' is defined by

$$d_Y(\mu, \mu') := \min\{d_Y(\gamma, \gamma') : \gamma \in \pi_Y(\mu), \gamma' \in \pi_Y(\mu')\}.$$

When Y is an annular subsurface with core curve α we denote $d_Y(\cdot, \cdot)$ by $d_\alpha(\cdot, \cdot)$ as well.

Lemma 2.1 of [22] gives us the following bound on the subsurface coefficient of two curves in terms of their intersection number:

$$(2-1) \quad d_Y(\alpha, \beta) \leq 2i(\alpha, \beta) + 1.$$

Let $\alpha, \beta \in \mathcal{C}_0(S)$ and $Y \subseteq S$ be an essential subsurface. If $d_Y(\alpha, \beta) > 2$, then $\alpha \pitchfork \beta$ holds. To see this, first suppose that Y is non-annular. Recall the surgery map which assigns to any arc in Y with endpoints on ∂Y the set of curves in the boundary of a regular neighborhood of $a \cup \partial Y$. This map from the arc complex of Y to $\mathcal{PC}_0(Y)$ is 2–Lipschitz; see [23]. Let a be an arc in $\alpha \cap Y$ and b be an arc in $\beta \cap Y$ with endpoints in the boundary of Y . The assumption $d_Y(\alpha, \beta) > 2$ and the fact that the surgery map is 2–Lipschitz imply that a and b have arc complex distance at least 2. Thus the arcs a and b intersect, and therefore the curves α and β intersect. Now suppose that Y is an annular subsurface. Then $d_Y(\alpha, \beta) > 2$ implies that the interior of any two lifts of α and β to the compactified annular cover \hat{Y} that go between two boundary components of \hat{Y} intersect. Therefore, α and β intersect.

The following lemma is a consequence of [23, Lemma 2.3].

Lemma 2.4 *Let μ denote a multi-curve, (partial) marking or lamination on a surface S . Then for any essential subsurface $Y \subseteq S$ we have*

$$\text{diam}_Y(\mu) \leq 2.$$

When Y is an annulus the sharp upper bound is 1.

The reason for the second part of the lemma is that any two lifts of two disjoint curves on the surface S to the compactified annular cover corresponding to the annular subsurface $Y \subset S$ are disjoint.

Let $\alpha, \beta, \gamma \in \mathcal{C}_0(S)$. Farb, Lubotzky and Minsky [12] defined the *relative twist* of the curves β and γ with respect to the curve α by

$$\tau_\alpha(\beta, \gamma) := \{b.c : b \in \pi_\alpha(\beta), c \in \pi_\alpha(\gamma)\},$$

where $b.c$ denotes the algebraic intersection number of the arcs a and b . The arcs b and c are oriented so that they intersect the lift of α homotopic to the core of \hat{Y} in the same direction. More precisely, let $\tilde{\alpha}$ be the lift of α homotopic to the core of \hat{Y} , and fix an orientation for $\tilde{\alpha}$. Then b and c are oriented so that the tangents to $\tilde{\alpha}$ and b and the tangents to $\tilde{\alpha}$ and c at their intersection points determine the same orientation for the annulus \tilde{Y} . Note that the subset $\tau_\alpha(\beta, \gamma) \subset \mathbb{Z}$ has diameter 2.

Given arcs $b, c \in \mathcal{C}(\alpha)$, by the discussion in [23, Section 2.4],

$$d_\alpha(b, c) = |b.c| + 1.$$

Let $\beta, \gamma \in \mathcal{C}_0(S)$. Since the diameter of $\tau_\alpha(\beta, \gamma)$ is at most 2, by the above formula we have

$$(2-2) \quad |d_\alpha(\beta, \gamma) - |x|| \leq 3$$

for any $x \in \tau_\alpha(\beta, \gamma)$. Let $\gamma = \mathcal{D}_\alpha^e(\beta)$, where \mathcal{D}_α is the positive Dehn twist about α and e is a positive integer. Formula (2) in [12, Section 2] is

$$(2-3) \quad \tau_\alpha(\beta, \gamma) \subset \{e, e + 1\}.$$

The following inequality, proved by Behrstock [2], relates the subsurface coefficients of two subsurfaces that overlap.

Theorem 2.5 (Behrstock inequality) *There is a constant $B_0 > 0$ so that given a curve system μ and subsurfaces Y and Z satisfying $Y \pitchfork Z$ we have*

$$\min\{d_Y(\partial Z, \mu), d_Z(\partial Y, \mu)\} \leq B_0.$$

Remark 2.6 Chris Leininger has observed that B_0 can be taken to be the universal constant 10. However, the specific value of B_0 does not play any role in our work.

Limits of laminations Let $\mathcal{L}_i = (\lambda_i, m_i)$ ($i \in \mathbb{N}$) be a sequence of measured laminations which converges to a measured lamination $\mathcal{L} = (\lambda, m)$ in the weak* topology. Suppose that, after possibly passing to a subsequence, the laminations λ_i converge to a lamination ξ in the Hausdorff topology of $M_\infty(S)$. It is a standard fact that $\lambda \subseteq \xi$; see for example [10, Section I.4].

Lemma 2.7 *Suppose that a sequence of curves $\{\alpha_i\}_{i=0}^\infty$ converges to a lamination λ in the Hausdorff topology of $M_\infty(S)$. Let Y be an essential subsurface, so that λ intersects Y essentially. Then for any geodesic lamination λ' that intersects Y essentially we have*

$$d_Y(\alpha_i, \lambda') \asymp_{1,4} d_Y(\lambda, \lambda')$$

for all i sufficiently large.

Proof First suppose that Y is an essential non-annular subsurface. Equip S with a complete hyperbolic metric and realize ∂Y , the curves α_i and the lamination λ geodesically in this metric. Let b an arc in $\lambda \cap Y$ and $\delta > 0$ be so that the δ -neighborhood of $b \cup \partial Y$ is a regular neighborhood and at least one of the components of the boundary of the neighborhood is an essential curve in Y . Denote the neighborhood by U ; see Figure 1. Let l be the geodesic in λ so that $b \subset l$. Let \tilde{l} be a lift of l to the universal cover \mathbb{D}^2 . The convergence of the curves α_i to λ in the Hausdorff topology of $M_\infty(S)$ (see [10, Section I.4, Lemma I.4.1.11]) guarantees that, given $\delta' < \delta$ and $L > 0$, for all i sufficiently large there is a lift $\tilde{\alpha}_i$ of α_i to \mathbb{D}^2 such that $\tilde{\alpha}_i$ and \tilde{l} δ' -fellow-travel on an interval of length at least L . Then, projecting $\tilde{\alpha}_i$ and \tilde{l} to S , we can see that there is an arc a_i of $\alpha_i \cap Y$ such that the arcs b and a_i are δ' -fellow-travelers in Y . This implies that the regular neighborhood U is also a regular neighborhood of $a_i \cup \partial Y$. By the definition of the subsurface projection the essential boundary curve of this neighborhood is a curve in $\pi_Y(\alpha_i)$.

By Lemma 2.4, $\pi_Y(\lambda)$ and $\pi_Y(\alpha_i)$ are subsets of $\mathcal{C}_0(Y)$ with diameter at most 2. Moreover, as we saw in the previous paragraph, $\pi_Y(\lambda) \cap \pi_Y(\alpha_i) \neq \emptyset$. Therefore

$$\text{diam}_Y(\pi_Y(\alpha_i) \cup \pi_Y(\lambda)) \leq 4.$$

Let β be a curve in $\pi_Y(\lambda')$. Then by the above bound on the diameter we have

$$|d_Y(\beta, \alpha_i) - d_Y(\beta, \lambda)| \leq 4.$$

This completes the proof of the lemma for non-annular subsurface Y .




Figure 1: The δ -neighborhood U of an arc b in $\lambda \cap Y$ is a regular neighborhood in Y with at least one essential boundary curve. For i sufficiently large, the Hausdorff distance of α_i and λ is less than δ , so $\alpha_i \cap Y$ is contained in U .

Now suppose that Y is an essential annular subsurface with core curve γ . Let b be an arc in $\pi_Y(\lambda)$. We claim that, for all i sufficiently large, there is an arc a_i in $\pi_Y(\alpha_i)$ such that the arcs a_i and b have at most one intersection point in their interior. If a_i and b do not intersect we are done. Otherwise, after conjugation we may assume that the origin of \mathbb{D}^2 is a lift of an intersection point of a_i and b . Moreover, there are lifts \tilde{b} of b and \tilde{a}_i of a_i to $\overline{\mathbb{D}^2}$ which pass through the origin; see Figure 2. As in Figure 2, there is a lower bound for the distance of \tilde{b} and any other lift of b to \mathbb{D}^2 . Then, choosing $\delta > 0$ sufficiently small and $L > 0$ large enough, any geodesic in \mathbb{D}^2 passing through the origin which δ -fellow-travels \tilde{b} on an interval of length at least L does not intersect any of the lifts of b except \tilde{b} . The geodesic \tilde{b} is a lift of a leaf of λ to \mathbb{D}^2 and \tilde{a}_i is a lift of α_i to \mathbb{D}^2 . So the Hausdorff convergence of the curves α_i to λ implies that, given $\delta, L > 0$, for i sufficiently large, \tilde{a}_i δ fellow-travels \tilde{b} on an interval of length at least L . Therefore, as we saw above, \tilde{a}_i intersects \tilde{b} once at the origin and does not intersect any other lift of b . The number of times that the arcs a_i and b intersect is equal to the number times that \tilde{b} intersects all of the lifts of a_i to \mathbb{D}^2 . Which by the above discussion is at most 1 (see Figure 2). The proof of the claim is complete.

The fact that a_i and b intersect at most once implies that

$$d_Y(\alpha_i, \lambda) \leq 2.$$

By Lemma 2.4, $\pi_Y(\alpha_i)$ and $\pi_Y(\lambda)$ are subsets of $\mathcal{C}_0(Y)$ with diameter at most 1. Moreover, as we saw above $\pi_Y(\alpha_i)$ and $\pi_Y(\lambda)$ have distance at most 2. Therefore

$$\text{diam}_Y(\pi_Y(\alpha_i) \cup \pi_Y(\lambda)) \leq 4.$$

Let β be a curve in $\pi_Y(\lambda')$. Then by the above bound on the diameter we have

$$|d_Y(\beta, \alpha_i) - d_Y(\beta, \lambda)| \leq 4.$$




Figure 2: Left: the arc b in $\pi_Y(\lambda)$ and a_i in $\pi_Y(\alpha_i)$ in the compactified annular cover \hat{Y} . Right: the lift \tilde{b} of the arc b and \tilde{a}_i of a_i to the universal cover \mathbb{D}^2 that pass through the origin and fellow-travel for a long portion. As in the picture, \tilde{a}_i and \tilde{b} intersect once; moreover, \tilde{a}_i does not intersect any other lift of b to \mathbb{D}^2 . Thus b and a_i in \hat{Y} intersect once.

This completes the proof of the lemma for an annular subsurface Y . □

Hierarchy paths and the distance formula Hierarchy paths, introduced by Masur and Minsky [23], comprise quasi-geodesics in the pants and marking graphs of a surface, with constants depending only on the topological type of the surface. Hierarchy paths have properties encoded in their endpoints and the associated subsurface coefficients. For a list of these properties, see [8, Section 2] and [24, Section 2]. Here we only state a key feature of hierarchy paths, which is the no-backtracking property. For other properties we provide a reference wherever we use them.

Theorem 2.8 *There exists a constant $M_2 > 0$ depending only on the topological type of the surface S with the following property. Let $\rho: [m, n] \rightarrow P(S)$ be a hierarchy path. Let $i, j, k, l \in [m, n]$ with $i \leq j \leq k \leq l$. For any subsurface $Y \subseteq S$ we have*

$$d_Y(\rho(i), \rho(l)) + 2M_2 \geq d_Y(\rho(j), \rho(k)).$$

The following theorem is a straightforward consequence of the bounded geodesic image theorem [23, Theorem 3.1].

Theorem 2.9 *Given $k \geq 1$ and $c \geq 0$, there is a $G \geq 0$ with the following property. Let $\{\gamma_i\}_{i=0}^\infty$ be a sequence of curves in $\mathcal{C}_0(S)$ which form a 1-Lipschitz, (k, c) -quasi-geodesic. Let $Y \subsetneq S$ be an essential subsurface so that $\gamma_i \cap Y$ holds for all $i \geq 0$. Then*

$$\text{diam}_Y(\{\pi_Y(\gamma_i)\}_{i=0}^\infty) \leq G.$$

Here $\text{diam}_Y(\cdot)$ is the diameter of the given subset of $\mathcal{C}(Y)$.

Using the hierarchical machinery, Masur and Minsky provide the following quasi-distance formula in the pants graph of a surface [23, Theorem 6.12]. Given $A > M_1$ (M_1 is a constant depending on the topological type of S) there are constants $K \geq 1$ and $C \geq 0$ such that

$$(2-4) \quad d(\mu, \mu') \asymp_{K,C} \sum_{\substack{Y \subseteq S \\ \text{non-annular}}} \{d_Y(\mu, \mu')\}_A.$$

Here the cut-off function $\{\cdot\}_A: \mathbb{R} \rightarrow \mathbb{R}^{\geq 0}$ is defined by

$$\{a\}_A = \begin{cases} a & \text{if } a \geq A, \\ 0 & \text{if } a < A. \end{cases}$$

Bounded combinatorics A pair of laminations or partial markings (μ, μ') has non-annular R -bounded combinatorics if

$$d_Y(\mu, \mu') \leq R$$

for every proper, essential, non-annular subsurface $Y \subsetneq S$.

The following result about stability of hierarchy paths with non-annular bounded combinatorics in the pants graph is an important ingredient in the proof that bounded combinatorics of end invariants of a WP geodesic guarantees co-boundedness of the geodesic and vice versa (see [8]). We need this theorem in our study of the behavior of WP geodesics in Section 4.

Theorem 2.10 [8] *Given $R > 0$, there is a quantifier function*

$$d_R: \mathbb{R}^{\geq 1} \times \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$$

so that a hierarchy path ρ with endpoints with non-annular R -bounded combinatorics is d_R -stable in the pants graph. That is, any (K, C) -quasi-geodesic with endpoints on $|\rho|$ stays in the $d_R(K, C)$ neighborhood of $|\rho|$. Here $|\rho|$ is the union of the pants decompositions of ρ .

2.2 The Weil–Petersson metric

In this section we assemble properties of the Weil–Petersson metric we will need. For an introduction to the synthetic geometry of the Weil–Petersson metric, see [30].

The Weil–Petersson metric on the Teichmüller space $\text{Teich}(S)$ is a Riemannian metric with negative sectional curvatures. It is incomplete, but is geodesically convex: any two points are joined by a unique geodesic that lies in the interior. Its metric completion $\overline{\text{Teich}(S)}$ is a CAT(0) space. See [4, Section II.3.4] for an introduction to CAT(0)

spaces. By the work of Masur [20] the completion of the Teichmüller space with the Weil–Petersson metric is naturally identified with the *augmented Teichmüller space* obtained by adjoining nodal surfaces as limits. The completion is stratified by the data of simple closed curves on S that are pinched: each stratum $\mathcal{S}(\sigma)$ is a copy of the Teichmüller space of the surface $S \setminus \sigma$, where σ is a multicurve. Masur also gave an expansion of the metric near the completion showing that the inclusion $\mathcal{S}(\sigma) \hookrightarrow \text{Teich}(S)$ is an isometry and $\mathcal{S}(\sigma)$ is totally geodesic.

S Yamada observed that a stronger form of Masur’s expansion should hold near the completion guaranteeing that the Weil–Petersson metric is asymptotic to a metric product of strata to higher order, and work of Daskalopolous and Wentworth [11] gave the appropriate metric expansion. Their expansion showed that these completion strata have the *non-refraction property*: for any $X, Y \in \overline{\text{Teich}(S)}$, the interior of the unique geodesic connecting X and Y lies in the smallest stratum that contains X and Y . See [29] for stronger form of the asymptotic expansion of the WP metric. The Weil–Petersson metric is invariant under the action of the mapping class group of the surface $\text{Mod}(S)$ and descends under the natural orbifold cover to a metric on the moduli space of Riemann surfaces $\mathcal{M}(S)$. The completion descends to a metric on the familiar Deligne–Mumford compactification of $\mathcal{M}(S)$.

Length functions Let $X \in \text{Teich}(S)$. Let α be a closed curve on S . We denote by $\ell_\alpha(X)$ the length of the geodesic representative of α in its free homotopy class on S . The length function has a natural extension to the space of measured laminations [3]. For $\mathcal{L} \in \mathcal{ML}(S)$, we denote the length of \mathcal{L} by $\ell_{\mathcal{L}}(X)$.

Significant from the point of view of the Weil–Petersson geometry is the result of Wolpert that each length function is a strictly convex function along any WP geodesic [30].

Quasi-isometric model Let S be a surface with negative Euler characteristic. There is a constant L_S (the Bers constant) depending only on the topological type of S such that any complete hyperbolic metric on S possesses a pants decomposition (a Bers pants decomposition) with the property that the length of any curve in the pants decomposition is at most L_S ; see [9, Section 5]. Let $X \in \overline{\text{Teich}(S)}$. Suppose that $X \in \mathcal{S}(\sigma)$. A *Bers pants decomposition* of X , denoted by $Q(X)$, is the union of Bers pants decompositions of the connected components of $S \setminus \sigma$ and σ . A *Bers marking* of X , denoted by $\mu(X)$, is a partial marking obtained from a Bers pants decomposition Q of X by adding a transversal curve with minimal length for each $\alpha \in Q - \sigma$. The following result of Jeffrey Brock provides a quasi-isometric model for the Weil–Petersson metric.

Theorem 2.11 (Quasi-isometric model [5]) *There are constants $K_{WP} \geq 1$ and $C_{WP} \geq 0$ depending only on the topological type of S with the following property. The map $Q: \text{Teich}(S) \rightarrow P(S)$ assigning to each X a Bers pants decomposition of X is a (K_{WP}, C_{WP}) -quasi-isometry.*

Ending laminations Let $r: [0, a) \rightarrow \text{Teich}(S)$ be a WP geodesic ray. Any limit in the weak* topology of an infinite sequence of distinct Bers curves at $r(t_n)$, where $t_n \rightarrow a$, is an *ending measured lamination* of r . A pinching curve α along r is a curve so that $\ell_\alpha(r(t)) \rightarrow 0$ as $t \rightarrow a$. Brock, Masur and Minsky [7] showed that the union of the supports of ending measured laminations and pinching curves of r is a geodesic lamination. We call this lamination the *ending lamination* of r .

Let $g: (a, b) \rightarrow \text{Teich}(S)$ be a WP geodesic, where (a, b) is an open interval containing 0. If the forward trajectory $g|_{[0,b)}$ can be extended to b so that $g(b) \in \overline{\text{Teich}(S)}$, we define the forward end invariant of g to be a Bers partial marking of $g(b)$. If not, let the forward end invariant of g be the lamination of $g|_{[0,b)}$ we defined above. We denote the forward end invariant by $\nu^+ = \nu^+(g)$. Similarly, consider the backward trajectory $g|_{(a,0]}$ and define the backward end invariant of g , $\nu^- = \nu^-(g)$.

From [24, Section 8] we have the following result:

Lemma 2.12 (Infinite rays) *Let ν be a minimal filling lamination. There is an infinite WP geodesic ray r with forward ending lamination ν .*

The following strengthened version of Wolpert’s geodesic limit theorem (see [29] and [8]), proved in [24, Section 4], provides a limiting picture for a sequence of bounded length WP geodesic segments in the Teichmüller space.

Theorem 2.13 (Geodesic limits) *Given $T > 0$, let $\zeta_n: [0, T] \rightarrow \overline{\text{Teich}(S)}$ be a sequence of WP geodesic segments parametrized by arc-length. After possibly passing to a subsequence, there is a partition $0 = t_0 < \dots < t_{k+1} = T$ of $[0, T]$, possibly empty multi-curves $\sigma_0, \dots, \sigma_{k+1}$ and a multi-curve $\hat{\tau} \equiv \sigma_i \cap \sigma_{i+1}$ for $i = 0, 1, \dots, k$ and a piece-wise geodesic*

$$\hat{\zeta}: [0, T] \rightarrow \overline{\text{Teich}(S)},$$

with the following properties:

- (1) $\hat{\zeta}((t_i, t_{i+1})) \subset \mathcal{S}(\hat{\tau})$ for $i = 0, \dots, k$.
- (2) $\hat{\zeta}(t_i) \in \mathcal{S}(\sigma_i)$ for $i = 0, \dots, k + 1$.

- (3) Given a multi-curve σ , denote by $\text{tw}(\sigma)$ the subgroup of $\text{Mod}(S)$ generated by positive Dehn twists about the curves in σ . There are elements ψ_n of the mapping class group for each $n \in \mathbb{N}$, and $\mathcal{T}_{i,n} \in \text{tw}(\sigma_i - \hat{\tau})$ for each $1 \leq i \leq k$ and $n \in \mathbb{N}$, so that $\psi_n(\zeta_n(t)) \rightarrow \hat{\zeta}(t)$ as $n \rightarrow \infty$ for all $t \in [0, t_1]$. Let $\varphi_{i,n} = \mathcal{T}_{i,n} \circ \dots \circ \mathcal{T}_{1,n} \circ \psi_n$ for $i = 1, \dots, k$ and each $n \in \mathbb{N}$. Then $\varphi_{i,n}(\zeta_n(t)) \rightarrow \hat{\zeta}(t)$ as $n \rightarrow \infty$ for any $t \in [t_i, t_{i+1}]$.

Remark 2.14 The central difference between the above version and original versions lies in the assertion that we have one (possibly empty) multi-curve $\hat{\tau}$ rather than several multi-curves $\tau_i = \sigma_i \cap \sigma_{i+1}$, $i = 0, 1, \dots, k$, allowed in Wolpert’s geodesic limit theorem. In particular, in part (1) the geodesic segments $\hat{\zeta}((t_i, t_{i+1}))$ lie in one stratum $\mathcal{S}(\hat{\tau})$ rather than several strata $\mathcal{S}(\tau_i)$.

3 Minimal non-uniquely ergodic laminations

A (measurable) geodesic lamination λ is *non-uniquely ergodic* if there are non-proportional measures supported on λ . More precisely, λ is non-uniquely ergodic if there exist transverse measures m and m' supported on λ and curves α and β such that

$$\frac{m(\alpha)}{m'(\alpha)} \neq \frac{m(\beta)}{m'(\beta)}.$$

Gabai [14, Section 9] gave a recipe to construct minimal filling non-uniquely ergodic geodesic laminations on any surface S with $\xi(S) > 1$. In fact, Gabai outlined the construction of minimal filling laminations and measures supported on each one of them with distinct projective classes [14, Theorem 9.1]. Leininger, Lenzhen and Rafi [18, Sections 3–5] gave a detailed construction of minimal filling non-uniquely ergodic laminations on the surface $S_{0,5}$. Moreover, they studied the set of measures supported on the lamination and their projective classes.

We first recall the construction of [18]. Let $\{e_i\}_{i=1}^\infty$ be a sequence of positive integers. Let $\rho: S_{0,5} \rightarrow S_{0,5}$ be the order-five homeomorphism of $S_{0,5}$ realized as the rotation by angle $4\pi/5$ in Figure 3. Let $\mathcal{D} = \mathcal{D}_{\hat{\gamma}_2}$ be the positive Dehn twist about the curve $\hat{\gamma}_2$. Let $f_i = \mathcal{D}^{e_i} \circ \rho$, for $i \geq 1$. Define the sequence of curves $\hat{\gamma}_i = f_1 \circ f_2 \circ \dots \circ f_i(\hat{\gamma}_0)$, for $i \geq 1$. The curves $\hat{\gamma}_0, \dots, \hat{\gamma}_5$ are shown in Figure 3.

Proposition 3.1 *There exist constants $E > 0$, $k \geq 1$, $c \geq 0$ and $K \geq 1$, $C \geq 0$ with the following properties. Suppose that $\{e_i\}_{i=1}^\infty$ is a sequence of integers satisfying $e_i > E$ for all $i \geq 1$. Let $\{\hat{\gamma}_i\}_{i=0}^\infty$ be the sequence of curves described above. Then*

- (1) For any $i \geq 0$ and $j \geq i + 2$, $\hat{\gamma}_j \pitchfork \hat{\gamma}_i$ holds.
- (2) For any $i \geq 0$ and $j \geq i + 4$ the curves $\hat{\gamma}_i$ and $\hat{\gamma}_j$ fill the surface $S_{0,5}$.
- (3) The sequence of curves $\{\hat{\gamma}_i\}_{i=0}^\infty$ is a 1-Lipschitz, (k, c) -quasi-geodesic in $\mathcal{C}_0(S_{0,5})$.
- (4) $d_{\hat{\gamma}_i}(\hat{\gamma}_j, \hat{\gamma}_{j'}) \asymp_{K,C} e_{i-1}$ for any $j \geq i + 2$ and $j' \leq i - 2$.




Figure 3: The double of each pentagon in the picture is a five-times punctured 2-sphere. The curves $\hat{\gamma}_0, \hat{\gamma}_1, \dots, \hat{\gamma}_5$ are shown in the picture. Any other six consecutive curves in the sequence, after applying an appropriate element of $\text{Mod}(S_{0,5})$, are the same as the above six curves, where the last two curves have different numbers of parallel strands from $\hat{\gamma}_4$ and $\hat{\gamma}_5$, respectively.

Proof We start by proving the following lemma.

Lemma 3.2 *Let $i \geq 1$. For any $j \geq i + 2$ and $j' \leq i - 2$,*

$$(3-1) \quad \hat{\gamma}_i \pitchfork \hat{\gamma}_j \quad \text{and} \quad \gamma_i \pitchfork \hat{\gamma}_{j'}$$

hold. Furthermore, the subsurface coefficient bound

$$(3-2) \quad d_{\hat{\gamma}_i}(\hat{\gamma}_j, \hat{\gamma}_{j'}) \geq e_{i-1} - C$$

holds.

Here $C = 2B_0 + 7$ and B_0 is the constant from the Behrstock inequality (Theorem 2.5).

Proof Define the constant

$$E = C + B_0 + G_0 + 2,$$

where G_0 is the constant from [Theorem 2.9](#) for a geodesic in the curve complex of $S_{0,5}$.

We prove (3-1) and (3-2) simultaneously by induction on $j - j'$. The proof of the base of the induction breaks into the following cases:

Case $j - j' = 4, j - i = 2$ and $i - j' = 2$ Applying $(f_1 \circ \dots \circ f_{i-2})^{-1}$ to the curves $\hat{\gamma}_{j'}, \hat{\gamma}_i$ and $\hat{\gamma}_j$, we obtain the curves

$$\hat{\gamma}_0, \hat{\gamma}_2 = f_{i-1} \circ f_i(\hat{\gamma}_0) \quad \text{and} \quad f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0),$$

respectively. The curve $f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0)$ is the same as $\hat{\gamma}_4$ with a different number of parallel strands; see [Figure 3](#). Since $\hat{\gamma}_2 \pitchfork \hat{\gamma}_0$ and $\hat{\gamma}_2 \pitchfork f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0)$ hold, it follows that $\hat{\gamma}_i \pitchfork \hat{\gamma}_{i-2}$ and $\hat{\gamma}_i \pitchfork \hat{\gamma}_{i+2}$ hold. This is (3-1).

We proceed to establish (3-2). We have $f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0) = D_{\hat{\gamma}_2}^{e_{i-1}} \circ \rho(\hat{\gamma}_3)$. Then by the formula (2-3) for the relative twists we have

$$\tau_{\hat{\gamma}_2}(f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0), \rho(\hat{\gamma}_3)) \subset \{e_{i-1}, e_{i-1} + 1\}.$$

Then (2-2) implies that

$$d_{\hat{\gamma}_2}(f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0), \rho(\hat{\gamma}_3)) \geq e_{i-1} - 3.$$

Furthermore, the curves $\rho(\hat{\gamma}_3)$ and $\hat{\gamma}_0$ are disjoint and both intersect $\hat{\gamma}_2$, thus

$$d_{\hat{\gamma}_2}(\hat{\gamma}_0, \rho(\hat{\gamma}_3)) \leq 1.$$

Combining the above two subsurface coefficient bounds by the triangle inequality and using the fact that $\text{diam}_{\hat{\gamma}_2}(\rho(\hat{\gamma}_3)) \leq 1$, we have

$$(3-3) \quad d_{\hat{\gamma}_2}(\hat{\gamma}_0, f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_i)) \geq e_{i-1} - 5.$$

Applying $f_1 \circ \dots \circ f_{i-2}$ to the subsurface coefficient above and using the fact that $C > 5$, we obtain

$$d_{\hat{\gamma}_i}(\hat{\gamma}_{i-2}, \hat{\gamma}_{i+2}) \geq e_{i-1} - C.$$

This is the subsurface coefficient bound (3-2).

Case $j - j' = 5, j - i = 2$ and $i - j' = 3$ Applying $(f_1 \circ \dots \circ f_{i-2})^{-1}$ to the curves $\hat{\gamma}_{j'}, \hat{\gamma}_i$ and $\hat{\gamma}_j$, we obtain the curves

$$\hat{\gamma}_0, \hat{\gamma}_2 = f_{i-1} \circ f_i(\hat{\gamma}_0) \quad \text{and} \quad f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_0),$$

respectively. The curve $f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_0)$ is the same as $\hat{\gamma}_5$ with a different number of parallel strands; see [Figure 3](#). Since $\hat{\gamma}_0 \pitchfork \hat{\gamma}_2$ and $\hat{\gamma}_2 \pitchfork f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_0)$ hold, (3-1) holds.

We have $f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_i) = \mathcal{D}_{\hat{\gamma}_2}^{e_i-1} \circ \rho(f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i))$. Then, by (2-3),

$$\tau_{\hat{\gamma}_2}(f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_i), \rho \circ f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i)) \subset \{e_{i-1}, e_{i-1} + 1\}.$$

So (2-2) implies that

$$d_{\hat{\gamma}_2}(f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_i), \rho \circ f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i)) \geq e_{i-1} - 3.$$

Furthermore, because $\rho(\hat{\gamma}_3)$ is a curve intersecting $\hat{\gamma}_2$ and disjoint from both $\hat{\gamma}_0$ and $\rho \circ f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i)$ (to see this, note that $f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i)$ is $\hat{\gamma}_4$ with a different number of parallel strands), we have

$$d_{\hat{\gamma}_2}(\hat{\gamma}_0, \rho \circ f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i)) \leq 2.$$

Combining the above two subsurface coefficient bounds with the triangle inequality and using the fact that $\text{diam}_{\hat{\gamma}_2}(\rho \circ f_i \circ \dots \circ f_{i+3}(\hat{\gamma}_i)) \leq 1$, we have

$$d_{\hat{\gamma}_2}(\hat{\gamma}_0, f_{i-1} \circ \dots \circ f_{i+3}(\hat{\gamma}_i)) \geq e_{i-1} - 6.$$

Now applying $f_1 \circ \dots \circ f_{i-2}$ to the above subsurface coefficient and using that $C > 6$, we get

$$d_{\hat{\gamma}_i}(\hat{\gamma}_{i-2}, \hat{\gamma}_{i+3}) \geq e_{i-1} - C.$$

This is the subsurface coefficient bound (3-2).

Case $j - j' = 5, j - i = 3$ and $i - j' = 2$ Applying $(f_1 \circ \dots \circ f_{i-3})^{-1}$ to the curves $\hat{\gamma}_{j'}, \hat{\gamma}_i$ and $\hat{\gamma}_j$, we obtain the curves

$$\hat{\gamma}_0, \hat{\gamma}_3 = f_{i-2} \circ f_{i-1} \circ f_i(\hat{\gamma}_0) \quad \text{and} \quad f_{i-2} \circ \dots \circ f_{i+2}(\hat{\gamma}_0);$$

see Figure 3. The statement about the intersection of curves (3-1) holds since $\hat{\gamma}_3 \pitchfork \hat{\gamma}_0$ and $\hat{\gamma}_3 \pitchfork f_{i-2} \circ \dots \circ f_{i+2}(\hat{\gamma}_0)$.

By the triangle inequality,

$$(3-4) \quad d_{\hat{\gamma}_3}(\hat{\gamma}_0, f_{i-2} \circ \dots \circ f_{i+2}(\hat{\gamma}_0)) \geq d_{\hat{\gamma}_3}(\hat{\gamma}_1, f_{i-2} \circ \dots \circ f_{i+2}(\hat{\gamma}_0)) - d_{\hat{\gamma}_3}(\hat{\gamma}_0, \hat{\gamma}_1) - \text{diam}_{\hat{\gamma}_3}(\hat{\gamma}_1).$$

First we find a lower bound for the first term on the right-hand side of (3-4). Note that $f_{i-2}(\hat{\gamma}_0) = \hat{\gamma}_1$ and $f_{i-2}(\hat{\gamma}_2) = \hat{\gamma}_3$. Thus, applying $(f_{i-2})^{-1}$ to this term, we obtain

$$d_{\hat{\gamma}_2}(\hat{\gamma}_0, f_{i-1} \circ \dots \circ f_{i+2}(\hat{\gamma}_0)).$$

This subsurface coefficient by (3-3) is bounded below by $e_{i-1} - 5$.

The two curves $\hat{\gamma}_0$ and $\hat{\gamma}_1$ are disjoint and intersect $\hat{\gamma}_3$. So the second term on the right-hand side of (3-4) is bounded by 1.

These bounds for the two terms on the right-hand side of the inequality (3-4) and the fact that $\text{diam}_{\widehat{\gamma}_3}(\widehat{\gamma}_1) \leq 1$ (Lemma 2.4) give us

$$d_{\widehat{\gamma}_3}(\widehat{\gamma}_0, f_{i-2} \circ \cdots \circ f_{i+2}(\widehat{\gamma}_0)) \geq e_{i-1} - 7 > e_{i-1} - C.$$

Applying $f_1 \circ \cdots \circ f_{i-1}$ to the subsurface coefficient on the left-hand side of the above inequality, we obtain the bound (3-2).

We proved that (3-1) and (3-2) hold for $j - j' \leq 5$. In what follows we assume that (3-1) and (3-2) hold when $j - j' \leq n$, where $n \geq 5$, and prove that (3-1) and (3-2) hold for $j - j' = n + 1$.

If $j - i = 2$ or 3 , applying $(f_1 \circ \cdots \circ f_i)^{-1}$ to $\widehat{\gamma}_i$ and $\widehat{\gamma}_j$, we obtain $\widehat{\gamma}_0$ and $\widehat{\gamma}_{j-i}$, respectively. Then since $\widehat{\gamma}_0 \pitchfork \widehat{\gamma}_{j-i}$ (see Figure 3), we have $\widehat{\gamma}_i \pitchfork \widehat{\gamma}_j$. If $j - i = 4$ or 5 , then since $(j - 2) - i \geq 2$ and $j - (j - 2) = 2$, by the hypothesis of the induction we have

$$d_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_i, \widehat{\gamma}_j) \geq E > 2.$$

This bound implies that $\widehat{\gamma}_i \pitchfork \widehat{\gamma}_j$ holds.

Now suppose that $j - i \geq 6$. Then we have $(j - 2) - i \geq 2$. Thus, by the induction hypothesis, $\widehat{\gamma}_{j-2} \pitchfork \widehat{\gamma}_i$ holds. Moreover, $\widehat{\gamma}_j \pitchfork \widehat{\gamma}_{j+2}$. So we may write the following triangle inequality:

$$(3-5) \quad \begin{aligned} d_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_i, \widehat{\gamma}_j) &\geq d_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_{j-4}, \widehat{\gamma}_j) - d_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_{j-4}, \widehat{\gamma}_i) - \text{diam}_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_{j-4}) \\ &\geq E - C - B_0 - 1 > 2. \end{aligned}$$

To get the second inequality in (3-5), first, by the assumption of the induction, we have

$$d_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_{j-4}, \widehat{\gamma}_j) \geq e_{j-3} - C \geq E - C.$$

This gives a lower bound for the first term on the right-hand side of the first inequality of (3-5). Second, since $(j - 4) - i \geq 2$ by the assumption of the induction, we have

$$d_{\widehat{\gamma}_{j-4}}(\widehat{\gamma}_i, \widehat{\gamma}_{j-2}) \geq e_{j-5} - C > E - C > B_0,$$

where the last inequality holds because $E > C + B_0$. Then the Behrstock inequality (Theorem 2.5) implies that

$$d_{\widehat{\gamma}_{j-2}}(\widehat{\gamma}_{j-4}, \widehat{\gamma}_i) \leq B_0.$$

This is the upper bound for the second term on the right-hand side of the first inequality of (3-5). Finally, by Lemma 2.4, the last term is at most 1.

The lower bound (3-5) guarantees that $\widehat{\gamma}_i \pitchfork \widehat{\gamma}_j$ holds. The proof of that $\widehat{\gamma}_{j'} \pitchfork \widehat{\gamma}_i$ holds for each $j' \leq i - 2$ is similar. The proof of (3-1) is complete.

We proceed to establish (3-2). Let j, j' be so that $j' \leq i - 2$ and $j \geq i + 2$. By (3-1) we may write the following triangle inequality:

$$(3-6) \quad d_{\widehat{\gamma}_i}(\widehat{\gamma}_{j'}, \widehat{\gamma}_j) \geq d_{\widehat{\gamma}_i}(\widehat{\gamma}_{i-2}, \widehat{\gamma}_{i+2}) - d_{\widehat{\gamma}_i}(\widehat{\gamma}_{i-2}, \widehat{\gamma}_{j'}) \\ - d_{\widehat{\gamma}_i}(\widehat{\gamma}_{i+2}, \widehat{\gamma}_j) - \text{diam}_{\widehat{\gamma}_i}(\widehat{\gamma}_{i-2}) - \text{diam}_{\widehat{\gamma}_i}(\widehat{\gamma}_{i+2}).$$

We have that $(i - 2) - j' < i - j' < j - j'$ and $j - (i + 2) < j - i < j - j'$. Thus by the assumption of the induction, the fact that $e_i > E$ and the choice of E we have that

$$d_{\widehat{\gamma}_{i-2}}(\widehat{\gamma}_i, \widehat{\gamma}_{j'}) \geq E - C > B_0 \quad \text{and} \quad d_{\widehat{\gamma}_{i+2}}(\widehat{\gamma}_i, \widehat{\gamma}_j) \geq E - C > B_0.$$

The first lower bound above and the Behrstock inequality imply that the second term on the right-hand side of (3-6) is bounded above by B_0 . Similarly, the second bound above and the Behrstock inequality imply that the third term on the right-hand side of (3-6) is bounded above by B_0 . Moreover, by Lemma 2.4, the third and fourth terms on the right-hand side of (3-6) are less than or equal to 1. So we obtain

$$d_{\widehat{\gamma}_i}(\widehat{\gamma}_{j'}, \widehat{\gamma}_j) \geq d_{\gamma_i}(\widehat{\gamma}_{i-2}, \widehat{\gamma}_{i+2}) - 2B_0 - 2 \geq e_{i-1} - C.$$

The proof of (3-2) is complete. □

We proceed to prove the proposition. Part (1) is the statement about intersection of curves (3-1) we proved in Lemma 3.2. Note that (3-2) gives the lower bound in part (4). Part (3) is [18, Lemma 3.2]. Part (4) follows from parts (1), (3) and the bounded geodesics image theorem (Theorem 2.9).

Now we prove part (2) of the proposition. The proof is by induction on $j - i$ and is essentially the one given in [18, Lemma 3.2]. Note that here we do not assume any upper bound for the value of $j - i$.

In the rest of the proof denote the surface $S_{0,5}$ by S . Suppose that $j - i = 4$, applying $(f_1 \circ \dots \circ f_i)^{-1}$ to the curves $\widehat{\gamma}_i$ and $\widehat{\gamma}_j$, we obtain the curves $\widehat{\gamma}_0$ and $\widehat{\gamma}_4$ in Figure 3, respectively, which fill S . Thus $\widehat{\gamma}_i$ and $\widehat{\gamma}_j$ fill S .

Suppose that part (2) is true for all $j - i \leq n$, where $n \geq 5$. Let $j - i = n + 1$. To get a contradiction, suppose that the curves $\widehat{\gamma}_i$ and $\widehat{\gamma}_j$ do not fill the surface. Then $d_S(\widehat{\gamma}_i, \widehat{\gamma}_j) \leq 2$. On the other hand, by the assumption of the induction the curves $\widehat{\gamma}_i$ and $\widehat{\gamma}_{j-1}$ fill S , so $d_S(\widehat{\gamma}_i, \widehat{\gamma}_{j-1}) \geq 3$. Moreover, by the construction of the sequence of curves, $\widehat{\gamma}_j$ and $\widehat{\gamma}_{j-1}$ are disjoint, so $d_S(\widehat{\gamma}_j, \widehat{\gamma}_{j-1}) = 1$. Thus, by the triangle inequality, $d_S(\widehat{\gamma}_i, \widehat{\gamma}_j) \geq 2$. The two bounds we established for $d_S(\widehat{\gamma}_i, \widehat{\gamma}_j)$ imply that

$$d_S(\widehat{\gamma}_i, \widehat{\gamma}_j) = 2.$$

Since $j - i \geq 5$ we may choose an index h so that

$$i < h < h + 1 < j, \quad j - h - 1 \geq 2 \quad \text{and} \quad h - i \geq 2.$$

Then by (3-1) the curves $\hat{\gamma}_i$ and $\hat{\gamma}_j$ intersect $\hat{\gamma}_h$ and $\hat{\gamma}_{h+1}$. Moreover, by the bound (3-2), the fact that $e_i > E$ and the choice of E , we have that

$$d_{\hat{\gamma}_h}(\hat{\gamma}_i, \hat{\gamma}_j) \geq E - C > G_0 \quad \text{and} \quad d_{\hat{\gamma}_{h+1}}(\hat{\gamma}_i, \hat{\gamma}_j) \geq E - C > G_0,$$

where G_0 is the constant from Theorem 2.9 for a geodesic in $\mathcal{C}(S)$.

As we saw above, $d_S(\hat{\gamma}_j, \hat{\gamma}_i) = 2$, so the geodesic in $\mathcal{C}(S)$ connecting $\hat{\gamma}_i$ and $\hat{\gamma}_j$ contains three curves $\hat{\gamma}_i, \gamma'$ and $\hat{\gamma}_j$. We have that the curve γ' is disjoint from $\hat{\gamma}_h$: otherwise, the curves $\gamma_i, \gamma', \gamma_j$ which form a geodesic in $\mathcal{C}(S)$ intersect $\hat{\gamma}_h$. Then the bounded geodesic image theorem (Theorem 2.9) implies that $d_{\hat{\gamma}_h}(\hat{\gamma}_i, \hat{\gamma}_j) \leq G_0$. But this contradicts the first lower bound above. Similarly, using the second lower bound above we may show that γ' and $\hat{\gamma}_{h+1}$ are disjoint. The curves $\hat{\gamma}_h$ and $\hat{\gamma}_{h+1}$ form a pants decomposition on S . Thus the only curves disjoint from both $\hat{\gamma}_h$ and $\hat{\gamma}_{h+1}$ are themselves. So γ' is either $\hat{\gamma}_h$ or $\hat{\gamma}_{h+1}$. As we mentioned above, the curves $\hat{\gamma}_h$ and $\hat{\gamma}_{h+1}$ intersect the curves $\hat{\gamma}_i$ and $\hat{\gamma}_j$. So γ' intersects both $\hat{\gamma}_i$ and $\hat{\gamma}_j$. On the other hand, since $\hat{\gamma}_i, \gamma'$ and $\hat{\gamma}_j$ are consecutive curves on a geodesic in $\mathcal{C}(S)$, γ' is disjoint from both $\hat{\gamma}_i$ and $\hat{\gamma}_j$. This contradiction shows that in fact $\hat{\gamma}_i$ and $\hat{\gamma}_j$ fill S , and completes the proof of part (2) by induction. \square

Let the sequence of integers $\{e_i\}_{i=1}^\infty$ with $e_i > E$, and the sequence of curves $\{\hat{\gamma}_i\}_{i=0}^\infty$ be as in Proposition 3.1. Part (3) of Proposition 3.1 and hyperbolicity of the curve complex imply that the sequence of curves $\{\hat{\gamma}_i\}_{i=0}^\infty$ converges to a point in the Gromov boundary of the curve complex. By Proposition 2.3 this point determines a projective measured lamination $[\mathcal{E}]$ with minimal filling support $\hat{\nu}$ on $S_{0,5}$.

Proposition 3.3 *Let the marking $\hat{\mu}$ and the geodesic lamination $\hat{\nu}$ be as above. Then:*

- (1) *There exist $K \geq 1$ and $C \geq 0$ so that $d_{\hat{\gamma}_i}(\hat{\mu}, \hat{\nu}) \asymp_{K,C} e_{i-1}$.*

Furthermore, suppose that for some $a > 2$ we have $e_{i+1} \geq ae_i$ for each $i \geq 1$. Then:

- (2) *The geodesic lamination $\hat{\nu}$ is minimal, filling and non-uniquely ergodic.*

Part (1) follows from Proposition 3.1(4) and the bounded geodesic image theorem (Theorem 2.9). Part (2) is [18, Theorem 1.1]. Note that the growth of powers $e_{i+1} \geq ae_i$ ($a > 2$) is required to guarantee the non-unique ergodicity of the lamination $\hat{\nu}$.

The utility of the construction of Leininger, Lenzhen and Rafi lies in its control on subsurface coefficients; see Proposition 3.3(1), Theorem 3.4 and Theorem 3.5. These

are conditions similar to arithmetic conditions for coefficients of the continued fraction expansion of irrational numbers relevant to the coding of geodesics on the modular surface which is $\mathcal{M}(S_{1,1})$ as well; see [28]. Though Gabai’s construction produces a minimal filling non-uniquely ergodic lamination on any surface S with $\xi(S) > 1$, it provides no a priori control on subsurface coefficients.

Theorem 3.4 *There is a constant $\hat{R} > 0$ such that for any proper, essential, non-annular subsurface $Y \subsetneq S$ we have $d_Y(\hat{\mu}, \hat{\nu}) \leq \hat{R}$. In other words, the pair $(\hat{\mu}, \hat{\nu})$ has non-annular \hat{R} -bounded combinatorics.*

Proof By Proposition 3.1(3), $\{\hat{\gamma}_i\}_{i=0}^\infty$ is a 1-Lipschitz, (k, c) -quasi-geodesic in $\mathcal{C}(S_{0,5})$. Let G be the corresponding constant from the bounded geodesic image theorem. Let $Y \subseteq S_{0,5}$ be an essential non-annular subsurface. First, note that Y is a four-holed sphere.

If $\hat{\gamma}_i \pitchfork Y$ holds for all $i \geq 0$, then the bounded geodesic image theorem guarantees that

$$\text{diam}_Y(\{\hat{\gamma}_i\}_{i=0}^\infty) \leq G.$$

The lamination $\hat{\nu}$ is filling, so $\pi_Y(\hat{\nu}) \neq \emptyset$. Now we claim that

$$(3-7) \quad d_Y(\hat{\mu}, \hat{\nu}) \leq G + 6.$$

To see this, let $\hat{\gamma}_{j_n}$ be a convergent subsequence of $\{\hat{\gamma}_j\}_{j=i+2}^\infty$ in the $\mathcal{PML}(S)$ topology. By Proposition 2.3 the support of the limit of $\hat{\gamma}_{j_n}$ is $\hat{\nu}$. After possibly passing to a further subsequence we may assume that $\hat{\gamma}_{j_n}$ is also convergent in the Hausdorff topology of $M_\infty(S)$. Denote the limit lamination in the Hausdorff topology by ξ . Then $\hat{\nu} \subseteq \xi$ (see eg [10]). By the bound $\text{diam}_Y(\{\hat{\gamma}_i\}_{i=0}^\infty) \leq G$ we established above, we have that $d_Y(\hat{\gamma}_0, \hat{\gamma}_{j_n}) \leq G$. Then since $\hat{\gamma}_{j_n} \rightarrow \xi$ in the Hausdorff topology as $n \rightarrow \infty$, by Lemma 2.7, we obtain

$$d_Y(\xi, \hat{\gamma}_0) \leq G + 4.$$

Furthermore, we have that $\hat{\nu} \subseteq \xi$ and $\hat{\gamma}_0 \subset \hat{\mu}$. Then since $\text{diam}_Y(\xi) \leq 2$ (by Lemma 2.4), the difference of the subsurface projection distance in (3-7) and the one above is at most 2. Which gives us (3-7).

Now suppose that for some integer $i \geq 0$, $\hat{\gamma}_i \pitchfork Y$ does not hold. Then since Y is a four-holed sphere inside $S_{0,5}$ we have that $\partial Y = \hat{\gamma}_i$.

Let $j' \leq i - 2$. By Proposition 3.1(1), $\hat{\gamma}_i \pitchfork \hat{\gamma}_{j'}$. Then since $\partial Y = \hat{\gamma}_i$, we conclude that $\hat{\gamma}_{j'} \pitchfork Y$ holds. Thus the bounded geodesic image theorem guarantees that

$$\text{diam}_Y(\{\hat{\gamma}_{j'}\}_{j'=0}^{i-2}) \leq G.$$

The above bound and the fact that $\hat{\mu}$ contains $\hat{\gamma}_0$ give us the bound

$$(3-8) \quad d_Y(\hat{\mu}, \hat{\gamma}_{i-2}) \leq G.$$

Let $j \geq i + 2$. By Proposition 3.1(1), $\hat{\gamma}_i \pitchfork \hat{\gamma}_j$ holds. Then similarly to above we obtain that

$$\text{diam}_Y(\{\hat{\gamma}_j\}_{j=i+2}^\infty) \leq G.$$

Then similar to the proof of (3-7) we may obtain

$$(3-9) \quad d_Y(\hat{\nu}, \hat{\gamma}_{i+2}) \leq G + 6.$$

By Proposition 3.1(4), $\hat{\gamma}_{i-2} \pitchfork \hat{\gamma}_i$. So $\hat{\gamma}_{i-2} \pitchfork Y$ holds, because $\hat{\gamma}_i = \partial Y$. Similarly $\hat{\gamma}_{i+2} \pitchfork Y$ holds. So $d_Y(\hat{\gamma}_{i-2}, \hat{\gamma}_{i+2})$ is defined. We claim that

$$(3-10) \quad d_Y(\hat{\gamma}_{i-2}, \hat{\gamma}_{i+2}) = 1.$$

To see this, let g be the element of $\text{Mod}(S)$ given by the composition $g = f_1 \circ \dots \circ f_{i-2}$. Applying g^{-1} to the subsurface coefficient in (3-10) we get

$$d_{g^{-1}(Y)}(g^{-1}(\hat{\gamma}_{i-2}), g^{-1}(\hat{\gamma}_{i+2})).$$

Thus, to obtain the desired equality, it suffices to show that the above subsurface coefficient is equal to 1. The curves

$$g^{-1}(\hat{\gamma}_{i-2}), \dots, g^{-1}(\hat{\gamma}_{i+2})$$

are the curves $\hat{\gamma}_0, \dots, \hat{\gamma}_4$ in Figure 3, respectively, except that the twist of the curve $g^{-1}(\hat{\gamma}_{i+2})$ about $g^{-1}(\hat{\gamma}_i) = \hat{\gamma}_2$ is e_{i-1} rather than e_1 .




Figure 4: The subsurface $g^{-1}(Y) \subset S_{0,5}$ is the four-holed sphere with boundary $\partial g^{-1}(Y) = \hat{\gamma}_2$. The curves $\pi_{g^{-1}(Y)}(g^{-1}(\hat{\gamma}_{i-2}))$ and $\pi_{g^{-1}(Y)}(g^{-1}(\hat{\gamma}_{i+2}))$ are shown in the figure.

Since $\partial Y = \hat{\gamma}_i$, the subsurface $g^{-1}(Y)$ is the four-holed sphere with boundary $\hat{\gamma}_2$; see Figure 4. We have that $g^{-1}(\hat{\gamma}_{i-2}) = \hat{\gamma}_0$. Furthermore, the curve $g^{-1}(\hat{\gamma}_{i+2})$ is the curve $\hat{\gamma}_4$ in Figure 3, except that the twist of the curve $g^{-1}(\hat{\gamma}_{i+2})$ about $\hat{\gamma}_2$ is

e_{i-1} rather than e_1 . The projections of the curves $g^{-1}(\hat{\gamma}_{i-2})$ and $g^{-1}(\hat{\gamma}_{i+2})$ to the subsurface $g^{-1}(Y)$ are shown in Figure 4. The $\mathcal{C}(g^{-1}(Y))$ -distance of these two curves is 1, because these are two curves with (minimal) intersection number 2 on the four-holed sphere $g^{-1}(Y)$, yielding the desired equality.

Note that $\hat{\mu}$ is a marking and \hat{v} fills the surface. By the triangle inequality and the bounds (3-8), (3-9) and (3-10) we have

$$d_Y(\hat{\mu}, \hat{v}) \leq d_Y(\hat{\mu}, \hat{\gamma}_{i-2}) + d_Y(\hat{\gamma}_{i-2}, \hat{\gamma}_{i+2}) + d_Y(\hat{\gamma}_{i+2}, \hat{v}) + \text{diam}_Y(\hat{\gamma}_{i-2}) + \text{diam}_Y(\hat{\gamma}_{i+2}) \leq 2G + 6 + 1 + 4.$$

We conclude that the Y subsurface coefficient of $\hat{\mu}$ and \hat{v} is bounded above by $\hat{R} := 2G + 11$, as was desired. □

Theorem 3.5 *There is a constant $R' > 0$ so that $d_\beta(\hat{\mu}, \hat{v}) \leq R'$ for any curve β which is not in the sequence $\{\hat{\gamma}_i\}_{i=0}^\infty$.*

Proof By Proposition 3.1(3), $\{\hat{\gamma}_i\}_{i=0}^\infty$ is a 1-Lipschitz, (k, c) -quasi-geodesic in $\mathcal{C}(S_{0,5})$. Let G be the corresponding constant from the bounded geodesic image theorem (Theorem 2.9).

If β intersects all of the curves in the sequence then, similarly to (3-7) in the proof of Theorem 3.4, we may obtain that

$$d_\beta(\hat{\mu}, \hat{v}) \leq G + 5.$$

Now suppose that for some integer $i > 0$ the curve β is disjoint from $\hat{\gamma}_i$. By Proposition 3.1(2) for every $j \geq i + 4$ the curves $\hat{\gamma}_j$ and $\hat{\gamma}_i$ fill $S_{0,5}$. So we may deduce that $\beta \pitchfork \hat{\gamma}_j$ holds. Then Theorem 2.9 guarantees that

$$(3-11) \quad d_\beta(\hat{\mu}, \hat{\gamma}_{i-4}) \leq G.$$

Similarly, for every $j' \leq i - 4$, $\beta \pitchfork \hat{\gamma}_{j'}$ holds. Then

$$\text{diam}_\beta(\{\gamma_{j'}\}_{j'=0}^{i-4})$$

by the bounded geodesic image theorem. Then, similarly to (3-7), we may obtain

$$(3-12) \quad d_\beta(\hat{v}, \hat{\gamma}_{i+4}) \leq G + 5.$$

Let $g = f_1 \circ \dots \circ f_i$. Applying g^{-1} to the curves $\hat{\gamma}_i, \dots, \hat{\gamma}_{i+4}$, we obtain the curves $\hat{\gamma}_0, \dots, \hat{\gamma}_4$ in Figure 3, respectively. The difference is that $g^{-1}(\hat{\gamma}_{i+4})$ has e_{i+1} twists. The only curve disjoint from $\hat{\gamma}_0$ and $\hat{\gamma}_2$ is $\hat{\gamma}_1$. Therefore, the only curve disjoint

from $\widehat{\gamma}_i$ and $\widehat{\gamma}_{i+2}$ is $\widehat{\gamma}_{i+1}$. The curve β is not in the sequence $\{\widehat{\gamma}_i\}_{i=0}^\infty$; in particular, $\beta \neq \widehat{\gamma}_{i+1}$. Moreover, β is disjoint from $\widehat{\gamma}_i$. Thus $\beta \pitchfork \widehat{\gamma}_{i+2}$ holds. Furthermore, the curves $\widehat{\gamma}_0$ and $g^{-1}(\widehat{\gamma}_{i+4})$ fill S ($g^{-1}(\widehat{\gamma}_{i+4})$ is $\widehat{\gamma}_4$ with a different number of parallel strands). Thus $\widehat{\gamma}_i$ and $\widehat{\gamma}_{i+4}$ fill S . Then, since β is disjoint from $\widehat{\gamma}_i$, $\beta \pitchfork \widehat{\gamma}_{i+4}$ holds. We showed that $\beta \pitchfork \widehat{\gamma}_{i+2}$ and $\beta \pitchfork \widehat{\gamma}_{i+4}$, therefore $d_\beta(\widehat{\gamma}_{i+2}, \widehat{\gamma}_{i+4})$ is defined. Now since $i(\widehat{\gamma}_{i+2}, \widehat{\gamma}_{i+4}) = 2$, by (2-1) we obtain the bound

$$(3-13) \quad d_\beta(\widehat{\gamma}_{i+2}, \widehat{\gamma}_{i+4}) \leq 5.$$

Similarly, we may obtain the bound

$$(3-14) \quad d_\beta(\widehat{\gamma}_{i-2}, \widehat{\gamma}_{i-4}) \leq 5.$$

Let $g = f_1 \circ \dots \circ f_{i-2}$. Applying g^{-1} to the curves $\widehat{\gamma}_{i-2}, \dots, \widehat{\gamma}_{i+2}$ we obtain the first five curves in Figure 3, with the difference that the last curve has e_{i-1} twists. Let $Y \subset S_{0,5}$ be the four-holed sphere with boundary curve $\widehat{\gamma}_i$. Then $\beta \in \mathcal{C}_0(Y)$. The curves $\pi_{g^{-1}(Y)}(g^{-1}(\widehat{\gamma}_{i-2}))$ and $\pi_{g^{-1}(Y)}(g^{-1}(\widehat{\gamma}_{i+2}))$ are shown in Figure 4. These two curves intersect twice. Thus by (2-1) we have

$$(3-15) \quad d_\beta(\widehat{\gamma}_{i-2}, \widehat{\gamma}_{i+2}) \leq 5.$$

The bounds (3-11)–(3-15) for the β subsurface coefficients combined with the triangle inequality give us the bound $R' := 2G + 29$. □

We proceed to construct minimal, filling, non-uniquely ergodic laminations on any surface $S_{g,0}$ of genus $g \geq 2$ with control on the subsurface coefficients of the laminations. We construct the laminations by an appropriate lift of the lamination we described on $S_{0,5}$ using 2-dimensional orbifolds and their orbifold covers. Here, we replace each puncture with a marked point on the surface.

Let $S_{0,5}$ be the 2-sphere equipped with an orbifold structure with five orbifold points of order 2 at the five marked points of $S_{0,5}$. Let $S_{0,6}$ be the 2-sphere with an orbifold structure with orbifold points of order 2 at the marked points of $S_{0,6}$. Let $S_{g,0}$ ($g \geq 2$) be $S_{g,0}$ equipped with an orbifold structure with no orbifold points (ie a manifold structure). Let

$$f: S_{0,6} \rightarrow S_{0,5} \quad \text{and} \quad h: S_{2,0} \rightarrow S_{0,6}$$

be the orbifold covering maps shown at the top left and right of Figure 5, respectively. Given $g \geq 2$, let $\sigma_g: S_{g,0} \rightarrow S_{2,0}$ be the covering map given at the bottom of Figure 5. Let $F_g = \sigma_g \circ h \circ f$. Let ν be the lamination $\nu = F_g^{-1}(\widehat{\nu})$.

Recall the sequence of curves $\{\widehat{\gamma}_i\}_{i=0}^\infty$. Denote the surface $S_{g,0}$ by S and the covering map F_g by F .




Figure 5: Top left: $f: \mathcal{S}_{0,6} \rightarrow \mathcal{S}_{0,5}$ is the orbifold covering map realized as the rotation by angle π about the axis in the picture. Top right: $h: \mathcal{S}_{2,0} \rightarrow \mathcal{S}_{0,6}$ is the orbifold covering map realized as the rotation by angle π about the axis in the picture (hyperelliptic involution). Bottom: $\sigma_g: \mathcal{S}_{g,0} \rightarrow \mathcal{S}_{2,0}$ is the covering map realized by $g - 1$ iterations of the rotation by angle $2\pi/(g - 1)$.

Theorem 3.6 *There are constants $k \geq 1$ and $c \geq 0$ so that the sequence $\{F^{-1}(\hat{\gamma}_i)\}_{i=0}^\infty$ is a (k, c) -quasi-geodesic in $\mathcal{C}(S)$.*

Proof By [27, Theorem 8.1] we have that the set-valued map that assigns to each simple closed curve α on the orbifold $\mathcal{S}_{0,5}$ the component curves of $F^{-1}(\alpha)$ in the orbifold cover $\mathcal{S}_{g,0}$ is a (Q, Q) -quasi-isometry from $\mathcal{C}(\mathcal{S}_{0,5})$ to $\mathcal{C}(\mathcal{S}_{g,0})$, where $Q \geq 1$ is a constant depending only on the degree of the cover $4(g - 1)$. Then the theorem follows from Proposition 3.1(3). □

For each $i \geq 0$ let γ_i be a component curve of $F^{-1}(\hat{\gamma}_i)$. By Theorem 3.6 we have

$$d_S(\gamma_i, \gamma_j) \geq \frac{1}{k}|i - j| - c.$$

Let $d = 2k + kc$. If $|i - j| \geq d$, then by the above inequality we have

$$d_S(\gamma_i, \gamma_j) \geq 2,$$

which implies that

$$(3-16) \quad \gamma_j \pitchfork \gamma_i$$

holds.

Let μ be a marking that contains a curve in each $F^{-1}(\hat{\gamma}_a)$ for $a = 0, 1, 2, 3$.

Theorem 3.7 *Let the sequence of curves $\{\gamma_i\}_{i=0}^\infty$, the marking μ and the lamination ν be as above. There are constants $K \geq 1$ and $C \geq 0$ depending only on the degree of the cover such that the following hold:*

- (1) For any $i \geq d$, $j \geq i + d$ and $j' \leq i - d$ we have $d_{\gamma_i}(\gamma_j, \gamma_{j'}) \asymp_{K,C} e_{i-1}$.
- (2) $d_{\gamma_i}(\mu, \nu) \asymp_{K,C} e_{i-1}$ for all $i \geq 1$.
- (3) For any essential, non-annular subsurface W we have $d_W(\mu, \nu) \leq R$.
- (4) The lamination ν is a minimal filling non-uniquely ergodic lamination on $S_{g,0}$.

Proof First we prove part (4). Since $\hat{\nu}$ is a non-uniquely ergodic lamination there are curves $\alpha, \beta \in \mathcal{C}_0(S_{0,5})$ and measures \hat{m} and \hat{m}' supported on $\hat{\nu}$ so that

$$\frac{\hat{m}(\alpha)}{\hat{m}'(\alpha)} \neq \frac{\hat{m}(\beta)}{\hat{m}'(\beta)}.$$

Let $m = F^*(\hat{m})$ and $m' = F^*(\hat{m}')$ be the pullbacks of \hat{m} and \hat{m}' , respectively. Then m and m' are measures supported on ν . Let $\tilde{\alpha}$ be a component of $F^{-1}(\alpha)$ and $\tilde{\beta}$ be a component of $F^{-1}(\beta)$. Then $m(\tilde{\alpha}) = \hat{m}(\alpha)$ and $m(\tilde{\beta}) = \hat{m}(\beta)$. Therefore

$$\frac{m(\tilde{\alpha})}{m'(\tilde{\alpha})} \neq \frac{m(\tilde{\beta})}{m'(\tilde{\beta})},$$

and hence the lamination ν is a non-uniquely ergodic lamination.

We proceed to show that the lamination ν is minimal and filling. We use the facts stated in Section 2.1 about measured laminations and foliations and the correspondence between them. Equip $\hat{\nu}$ with a transverse measure \hat{m} and ν with the measure $m = F^*(\hat{m})$. Let $(\hat{\mathcal{F}}, \hat{m})$ and (\mathcal{F}, m) be the measured foliations corresponding to $(\hat{\nu}, \hat{m})$ and (ν, m) , respectively. Note that $\mathcal{F} = F^{-1}(\hat{\mathcal{F}})$. Since the lamination $\hat{\nu}$ is minimal, the foliation $\hat{\mathcal{F}}$ is minimal. By the result of Hubbard and Masur [16], given a complex structure on the surface $S_{0,5}$ there is a unique quadratic differential \hat{q} with vertical measured foliation $(\hat{\mathcal{F}}, \hat{m})$. Then (\mathcal{F}, m) is the vertical measured foliation of the quadratic differential $q = F^*(\hat{q})$. Since \mathcal{F} is a minimal foliation on $S_{0,5}$ any leaf of \mathcal{F} is dense in the surface. Therefore, the lift of each leaf of \mathcal{F} to $S_{g,0}$ is dense. To see this, let l be a leaf of \mathcal{F} . Suppose to the contrary that l misses an open set U in $S_{g,0}$. We may shrink U and assume that the restriction of F to U is a homeomorphism. But then $F(l)$, which is a leaf of $\hat{\mathcal{F}}$, misses $F(U)$, which is an open subset of $S_{0,5}$. This contradicts the fact that $\hat{\mathcal{F}}$ is a minimal foliation of $S_{0,5}$. Therefore \mathcal{F} is minimal, and consequently ν is as well.

To see that the lamination ν fills S , note that given $\alpha \in \mathcal{C}_0(S)$, a homotopy that realizes α and ν as disjoint subsets of $S_{g,0}$ composed with F gives us a homotopy which realizes $F(\alpha)$ (an essential closed curve on $S_{0,5}$) and $\hat{\nu}$ as disjoint subsets of $S_{0,5}$. But this contradicts the fact that $\hat{\nu}$ fills $S_{0,5}$.

Using the terminology of [27] we say that a subsurface $W \subseteq S_{g,0}$ is a symmetric subsurface if it is a component of $F^{-1}(Y)$ for some subsurface $Y \subseteq S_{0,5}$.

When the subsurface W is not symmetric, by [27, Lemma 7.2], we have

$$(3-17) \quad d_W(\mu, \nu) \leq 2T_e + 1$$

for a constant $T_e > 0$ depending only on the degree of the cover and the constant e which comes from Rafi’s characterization of short curves along Teichmüller geodesics; see [27, Section 4] and for more detail [25; 26].

When the subsurface W is an essential symmetric subsurface we have

$$d_W(\mu, \nu) \leq d_Y(\hat{\mu}, \hat{\nu})$$

(see the proof of [27, Theorem 8.1]). Furthermore, by Theorem 3.4, we know that there exists $\hat{R} > 0$ so that

$$d_Y(\hat{\mu}, \hat{\nu}) \leq \hat{R}$$

for every essential non-annular subsurface $Y \subseteq S_{0,5}$. The above two inequalities for subsurface coefficients give us

$$(3-18) \quad d_W(\mu, \nu) \leq \hat{R}.$$

Then by the subsurface coefficient bounds (3-17) and (3-18) we obtain the upper bound $R := \max\{\hat{R}, 2T_e + 1\}$ in part (3).

We proceed to prove parts (1) and (2). The fact that the subsurface coefficient $d_{\gamma_i}(\gamma_j, \gamma_{j'})$ in part (1) is defined follows from (3-16). Note that each annular subsurface with core curve γ_i is a symmetric subsurface, because γ_i is a component of $F^{-1}(\hat{\gamma}_i)$. Thus, as is shown in the proof of [27, Theorem 8.1], there exists $Q \geq 1$ so that

$$(3-19) \quad d_{\gamma_i}(\gamma_j, \gamma_{j'}) \asymp_{Q,Q} d_{\hat{\gamma}_i}(\hat{\gamma}_j, \hat{\gamma}_{j'}), \text{ and}$$

$$(3-20) \quad d_{\gamma_i}(\mu, \nu) \asymp_{Q,Q} d_{\hat{\gamma}_i}(\hat{\nu}, \hat{\mu}).$$

By Proposition 3.1(4) we have

$$d_{\hat{\gamma}_i}(\hat{\gamma}_j, \hat{\gamma}_{j'}) \asymp e_{i-1},$$

then from the quasi-equality of subsurface coefficients (3-19) the quasi-equality (1) follows.

Moreover, by Proposition 3.3(4) we have

$$d_{\hat{\gamma}_i}(\hat{\mu}, \hat{\nu}) \asymp e_{i-1},$$

then from the quasi-equality of subsurface coefficients (3-20) the quasi-equality (2) follows. □

4 Recurrence of geodesics

Let $X \in \text{Teich}(S)$ and ν^- be a Bers marking of X . Let ν^+ be a minimal filling lamination. By Lemma 2.12 there is an infinite WP geodesic ray $r: [0, \infty) \rightarrow \text{Teich}(S)$ with $r(0) = X$ and end invariant (ν^-, ν^+) . Denote the projection of r to the moduli space by \hat{r} . For $\epsilon > 0$, the ϵ -thick part of $\mathcal{M}(S)$ consists of the Riemann surfaces with injectivity radius greater than ϵ .

Theorem 4.1 *Given $R > 0$. Suppose that (ν^-, ν^+) has non-annular R -bounded combinatorics. There is an $\epsilon > 0$ such that \hat{r} is recurrent to the ϵ -thick part of the moduli space.*

Lemma 4.2 *Given $R > 0$, there are constants $T_0 > 0$ and $\epsilon > 0$ with the following property. Let $T > T_0$ and $\zeta_n: [0, T] \rightarrow \text{Teich}(S)$ be a sequence of WP geodesic segments parametrized by arc-length so that the pair $(Q(\zeta_0(0)), Q(\zeta_n(T)))$ has non-annular R -bounded combinatorics. Then there is a time $t^* \in [0, T]$ and a sequence $\{m_n\}_{n=1}^\infty$ so that for all n sufficiently large we have*

$$\text{inj}(\zeta_{m_n}(t^*)) > \epsilon.$$

Proof Consider the limiting picture of geodesic segments ζ_n as was described in Theorem 2.13. Let the partition $0 = t_0 < \dots < t_{k+1} = T$, the multi-curves σ_i , $i = 0, \dots, k + 1$, the multi-curve $\hat{\tau}$, and the piecewise geodesic

$$\hat{\zeta}: [0, T] \rightarrow \overline{\text{Teich}(S)}$$

be as in the theorem. Furthermore, recall the elements of the mapping class group ψ_n for $n \in \mathbb{N}$, and $\mathcal{T}_{i,n} \in \text{tw}(\sigma_i - \hat{\tau})$ for $i = 0, \dots, k + 1$ and $n \in \mathbb{N}$. As in the theorem set $\varphi_{i,n} = \mathcal{T}_{i,n} \circ \dots \circ \mathcal{T}_{1,n} \circ \psi_n$.

First we show that $\hat{\tau} = \emptyset$. Suppose to the contrary that $\hat{\tau} \neq \emptyset$.

From Theorem 2.13, we know that $\hat{\tau} = \sigma_0 \cap \sigma_1$, so $\hat{\tau} \subseteq \sigma_0$. Moreover, $\hat{\zeta}(0) \in \mathcal{S}(\sigma_0)$ by Theorem 2.13(2). Thus for any $\alpha \in \hat{\tau}$ we have $l_\alpha(\hat{\zeta}(0)) = 0$. Furthermore, by Theorem 2.13(3), after possibly passing to a subsequence, $\psi_n(\zeta_n(0)) \rightarrow \hat{\zeta}(0)$ as $n \rightarrow \infty$. Thus, by continuity of length functions, for all n sufficiently large and any $\alpha \in \hat{\tau}$ we have $l_\alpha(\psi_n(\zeta_n(0))) \leq L_S$. Thus there is a Bers pants decomposition $Q_{0,n}$ of $\psi_n(\zeta_n(0))$ that contains $\hat{\tau}$.

Similarly, since $\hat{\tau} = \sigma_k \cap \sigma_{k+1}$ (as in Theorem 2.13), we have $\hat{\tau} \subseteq \sigma_{k+1}$. Moreover, by Theorem 2.13(2) we know that $\hat{\zeta}(T) \in \mathcal{S}(\sigma_{k+1})$. Thus for any $\alpha \in \hat{\tau}$ we have

$\ell_\alpha(\widehat{\zeta}(T)) = 0$. Furthermore, by [Theorem 2.13\(3\)](#), after possibly passing to a subsequence, $\varphi_{k,n}(\zeta_n(T)) \rightarrow \widehat{\zeta}(T)$ as $n \rightarrow \infty$. Thus, by continuity of length functions, for all n sufficiently large and any $\alpha \in \widehat{\tau}$ we have that $\ell_\alpha(\varphi_{k,n}(\zeta_n(T))) \leq L_S$.

Now note that we have $\varphi_{k,n} = \mathcal{T}_{k,n} \circ \dots \circ \mathcal{T}_{1,n} \circ \psi_n$. The element $\mathcal{T}_{i,n}$ of the mapping class group is a composition of powers of Dehn twists about the curves in σ_i and $\widehat{\tau} \subseteq \sigma_i$. Therefore, $\mathcal{T}_{i,n}$ preserves the isotopy class and the length of every curve $\alpha \in \widehat{\tau}$. Thus, applying $(\mathcal{T}_{k,n} \circ \dots \circ \mathcal{T}_{1,n})^{-1}$ to $\ell_\alpha(\varphi_{k,n}(\zeta_n(T)))$, we obtain

$$\ell_\alpha(\varphi_{k,n}(\zeta_n(T))) = \ell_\alpha(\psi_n(\zeta_n(T))).$$

Then, by the previous paragraph, for all n sufficiently large, $\ell_\alpha(\psi_n(\zeta_n(T))) \leq L_S$. Thus there is a Bers pants decomposition $\mathcal{Q}_{k+1,n}$ of $\psi_n(\zeta_n(T))$ containing $\widehat{\tau}$.

Let the threshold constant in the distance formula [\(2-4\)](#) be $A > \max\{M_1, R, 2\}$. Then there are constants $K \geq 1$ and $C \geq 0$ such that

$$(4-1) \quad d(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n}) \asymp_{K,C} \sum_{\substack{Y \subseteq S \\ \text{non-annular}}} \{d_Y(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n})\}_A.$$

As we saw above $\widehat{\tau} \subset \mathcal{Q}_{0,n}$ and $\widehat{\tau} \subset \mathcal{Q}_{k+1,n}$. So for any essential subsurface W satisfying $\widehat{\tau} \pitchfork W$ it follows that

$$d_W(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n}) \leq 2.$$

Thus subsurfaces which overlap $\widehat{\tau}$ do not contribute to the right-hand side of [\(4-1\)](#). On the other hand, by [Theorem 2.11](#) (quasi-isometric model) there are constants $K_{WP} \geq 1, C_{WP} \geq 0$ such that

$$d(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n}) \asymp_{K_{WP}, C_{WP}} d_{WP}(\zeta_n(0), \zeta_n(T)).$$

Let $T_0 = K_{WP}(KA + KC) + K_{WP}C_{WP}$. Since $T \geq T_0$, by the above quasi-equality we have

$$d(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n}) \geq KA + KC.$$

Now [\(4-1\)](#) and the above inequality imply that for any $n \in \mathbb{N}$ there is an essential non-annular subsurface Y_n with

$$d_{Y_n}(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n}) \geq A \geq R.$$

But as we saw above Y_n can not overlap $\widehat{\tau}$ (otherwise $d_{Y_n}(\mathcal{Q}_{0,n}, \mathcal{Q}_{k+1,n}) \leq 2 < A$), therefore, $Y_n \subseteq S \setminus \widehat{\tau}$. Moreover, since $\widehat{\tau} \neq \emptyset$, Y_n is a proper subsurface. Applying ψ_n^{-1} to the subsurface coefficient above we get

$$(4-2) \quad d_{\psi_n^{-1}(Y_n)}(\psi_n^{-1}(\mathcal{Q}_{0,n}), \psi_n^{-1}(\mathcal{Q}_{k+1,n})) \geq R,$$

where $\psi_n^{-1}(Q_{0,n})$ is a Bers pants decomposition of $\zeta_n(0)$ and $\psi_n^{-1}(Q_{k+1,n})$ is a Bers pants decomposition of $\zeta_n(T)$. Moreover, $\psi_n^{-1}(Y_n)$ is a proper subsurface of S , because Y_n is a proper subsurface of S . But then the lower bound (4-2) contradicts the non-annular bounded combinatorics assumption for the two pants decompositions $Q(\zeta_n(0))$ and $Q(\zeta_n(T))$. This contradiction completes the proof of the fact that $\hat{\tau} = \emptyset$.

Let $t^* = \frac{1}{2}t_1$. By Theorem 2.13(2) and since $\hat{\tau} = \emptyset$, we have that $\hat{\zeta}(t^*) \in \text{Teich}(S)$. So $\text{inj}(\hat{\zeta}(t^*)) > 2\epsilon$ for some $\epsilon > 0$. Furthermore, by Theorem 2.13(3), there is a sequence $\{m_n\}_{n=1}^\infty$ such that $\psi_{m_n}(\zeta_{m_n}(t^*)) \rightarrow \hat{\zeta}(t^*)$ as $n \rightarrow \infty$. Therefore, $\text{inj}(\psi_{m_n}(\zeta_{m_n}(t^*))) > \epsilon$ for any n sufficiently large. Then since the action by elements of the mapping class group does not change the injectivity radius of a surface, $\text{inj}(\zeta_{m_n}(t^*)) > \epsilon$. \square

Proof of Theorem 4.1 Let $T_0 > 0$ be as in Lemma 4.2 and $T \geq T_0$. Consider the sequence of WP geodesic segments

$$\zeta_n := r|_{[nT, (n+1)T]}: [0, T] \rightarrow \text{Teich}(S).$$

Note that Theorem 2.10 guarantees that, for $D = d_R(K_{\text{WP}}, C_{\text{WP}})$, the paths $Q(r)$ and ρ D -fellow-travel in the pants graph. Let $z_n^-, z_n^+ \in [0, \infty)$ be so that

$$d(\rho(z_n^-), Q(\zeta_n(0))) \leq D, \text{ and}$$

$$d(\rho(z_n^+), Q(\zeta_n(T))) \leq D.$$

Then for every essential non-annular subsurface $Y \subsetneq S$,

$$(4-3) \quad d_Y(\rho(z_n^-), Q(\zeta_n(0))) \leq D, \text{ and}$$

$$(4-4) \quad d_Y(\rho(z_n^+), Q(\zeta_n(T))) \leq D.$$

Moreover, by the assumption that the pair (ν^-, ν^+) has non-annular R -bounded combinatorics, for any proper, essential non-annular subsurface $Y \subsetneq S$ we have $d_Y(\nu^-, \nu^+) \leq R$. Then by the no-backtracking property of hierarchy paths (Theorem 2.8) there is an $M_2 > 0$ so that

$$(4-5) \quad d_Y(\rho(z_n^-), \rho(z_n^+)) \leq R + 2M_2.$$

The subsurface coefficient bounds (4-3)–(4-5) combined with the triangle inequality imply that

$$d_Y(Q(\zeta_n(0)), Q(\zeta_n(T))) \leq 2D + R + 2M_2 + \text{diam}_Y(\rho(z_n^-)) + \text{diam}_Y(\rho(z_n^+))$$

$$\leq 2D + R + 2M_2 + 4.$$

Thus the pair $(Q(\zeta_n(0)), Q(\zeta_n(T)))$ has $R + 2D + 2M_2 + 4$ non-annular bounded combinatorics.

Then Lemma 4.2 applies to the sequence of geodesic segments $\zeta_n := r|_{[nT, (n+1)T]}$ and implies that there are $t^* \in [0, T]$, $\epsilon > 0$ and a sequence of integers $\{m_n\}_{n=1}^\infty$ such that at $a_n = m_n T + t^*$ we have

$$\text{inj}(r(a_n)) > \epsilon.$$

This implies that $\hat{r}(a_n)$ is in the ϵ -thick part of the moduli space, where \hat{r} is the projection of r to the moduli space. Furthermore, since $a_n \rightarrow \infty$, the ray is recurrent to the ϵ -thick part of the moduli space. \square

Let $r: [0, \infty) \rightarrow \text{Teich}(S)$ be the ray with end invariant (v^-, v^+) with non-annular bounded combinatorics. In Theorem 4.1 we saw that the ray \hat{r} is recurrent to a compact subset of $\mathcal{M}(S)$. In Theorem 4.4 we show that if in addition there is a sequence of curves $\{\gamma_i\}_{i=1}^\infty$ so that $d_{\gamma_i}(v^-, v^+) \rightarrow \infty$ as $i \rightarrow \infty$ then the recurrent ray \hat{r} is not contained in any compact part of the moduli space. The theorem also follows from [8, Theorem 3.1]. The proof here is different and more direct and gives some information about the excursion times. We need the following result from [24, Section 4].

Lemma 4.3 (Large twist \implies short curve) *Given T, ϵ_0 and N positive there is an $\epsilon < \epsilon_0$ with the following property. Let $\zeta: [0, T'] \rightarrow \text{Teich}(S)$ be a WP geodesic segment of length $T' \leq T$ such that*

$$\sup_{t \in [0, T']} \ell_\gamma(\zeta(t)) \geq \epsilon_0.$$

If $d_\gamma(\mu(\zeta(0)), \mu(\zeta(T'))) > N$ (where $\mu(X)$ denotes a Bers marking of the point $X \in \text{Teich}(S)$) then we have

$$\inf_{t \in [0, T']} \ell_\gamma(\zeta(t)) \leq \epsilon.$$

Moreover, $\epsilon \rightarrow 0$ as $N \rightarrow \infty$.

Theorem 4.4 *Let $r: [0, \infty) \rightarrow \text{Teich}(S)$ be a WP geodesic ray with end invariant (v^-, v^+) . Suppose (v^-, v^+) has non-annular R -bounded combinatorics. Moreover, assume that there is a sequence of curves $\{\gamma_i\}_{i=1}^\infty$ so that*

$$d_{\gamma_i}(v^-, v^+) \rightarrow \infty$$

as $i \rightarrow \infty$. Then there is a sequence of times $b_i \rightarrow \infty$ as $i \rightarrow \infty$ such that

$$\ell_{\gamma_i}(r(b_i)) \rightarrow 0$$

as $i \rightarrow \infty$.

Proof Let $\rho: [0, \infty) \rightarrow P(S)$ be a hierarchy path with endpoints v^- and v^+ . The pair (v^-, v^+) has non-annular bounded combinatorics, so [Theorem 2.10](#) implies that, for $D = d_R(K_{WP}, C_{WP})$, ρ and $Q(r)$ D -fellow-travel. Moreover, both ρ and $Q(r)$ are quasi-geodesics. Then there is a quasi-isometry $N: [0, \infty) \rightarrow [0, \infty)$ from the domain of ρ to the domain of r . The map assigns to each i in the domain of ρ any time t in the smallest interval in the domain of r which contains all t' with $d(\rho(i), Q(r(t')))) \leq D$. For more detail, see [\[24, Section 5.3\]](#). Denote the constants of the quasi-isometry N by K_1 and C_1 .

We assumed that $d_{\gamma_i}(v^-, v^+) \rightarrow \infty$ as $i \rightarrow \infty$, so for all i sufficiently large we have

$$d_{\gamma_i}(v^-, v^+) \geq M_1,$$

where M_1 is the constant from the large link lemma [\[23, Lemma 6.2\]](#); see also property (2) of hierarchy paths in [\[8, Theorem 2.6\]](#). Then the annular subsurface with core curve γ_i is a component domain of ρ . Thus there is a time $q_i \in [0, \infty)$ so that $\rho(q_i)$ contains the curve γ_i . Note that $q_i \rightarrow \infty$ as $i \rightarrow \infty$.

Since the pair (v^-, v^+) has non-annular R -bounded combinatorics, for any proper, essential non-annular subsurface $Y \subsetneq S$ we have $d_Y(v^-, v^+) \leq R$. Then by the no-backtracking property of hierarchy paths ([Theorem 2.8](#)) there is an $M_2 > 0$ so that for any $i, j \in \mathbb{N}$ we have

$$d_Y(\rho(q_i), \rho(q_j)) \leq R + 2M_2.$$

Let the threshold in the distance formula (2-4) be $\max\{M_1, R + 2M_2\}$. Then there are constants $K_R \geq 1$ and $C_R \geq 0$ corresponding to the threshold so that

$$(4-6) \quad d(\rho(q_i), \rho(q_j)) \asymp_{K_R, C_R} d_S(\rho(q_i), \rho(q_j)).$$

Let $w(D, 0, R)$ be the constant from the annular coefficient comparison lemma in [\[24, Section 6\]](#); see below. Let $w = \max\{w, 2\}$. We have that the pair (v^-, v^+) has non-annular R -bounded combinatorics. Moreover for each i sufficiently large $\gamma_i \in \rho(q_i)$. These two facts imply that for all i sufficiently large, the curve γ_i is $(w, 0)$ -isolated at q_i , where the subsurface with non-annular R -bounded combinatorics on both sides of q_i is the surface S . See [\[24, Section 6.1\]](#) for the definition of isolated curve (annular subsurface) along a hierarchy path.

Recall that ρ is a (k, c) -quasi-geodesic in $P(S)$. Let $K_2 = \max\{K_1, K_R\}$ and $C_2 = \max\{C_1, C_R\}$, where K_1, C_1 are the constants of the quasi-isometry N and K_R, C_R are the constants in the quasi-equality (4-6). For any integer $i \geq 0$, set

$$q_i^- = q_i - k(K_2(w + C_2)) - kc \quad \text{and} \quad q_i^+ = q_i + k(K_2(w + C_2)) + kc.$$

By the setup of q_i^- , for any $q \leq q_i^-$ we have $d(\rho(q_i), \rho(q)) \geq K_2(w + C_2)$, and by the setup of q_i^+ , for any $q \geq q_i^+$ we have $d(\rho(q_i), \rho(q)) \geq K_2(w + C_2)$. Then the quasi-equality (4-6) implies that

$$d_S(\rho(q_i), \rho(q)) \geq w \geq 2.$$

This inequality guarantees that the $\mathcal{C}(S)$ -distance of any curve in the pants decomposition $\rho(q_i)$ and any curve in $\rho(q)$ is at least 2. Thus any curve in the pants decomposition $\rho(q_i)$ intersects any curve in $\rho(q)$. In particular, γ_i intersects any curve in $\rho(q)$. Then there is an $M_3 > 0$ so that $d_{\gamma_i}(\rho(q_i^+), v^+) \leq M_3$ and $d_{\gamma_i}(\rho(q_i^-), v^-) \leq M_3$; see property (4) of hierarchy paths in [8, Theorem 2.6]. Therefore,

$$(4-7) \quad d_{\gamma_i}(\rho(q_i^-), \rho(q_i^+)) \asymp_{1,2M_3} d_{\gamma_i}(v^-, v^+).$$

Let $s_i^- \in N(\rho(q_i^-))$ and $s_i^+ \in N(\rho(q_i^+))$. Since $q_i^+ - q_i \geq w$ and $q_i - q_i^- \geq w$, by the annular coefficient comparison lemma in [24, Section 6] we have

- $\min\{\ell_{\gamma_i}(r(s_i^-)), \ell_{\gamma_i}(r(s_i^+))\} \geq \omega(L_S)$, where $\omega(a)$ is the width of the collar of a simple closed geodesic with length a on a complete hyperbolic surface provided by the collar lemma (see [9, Section 4.1]), and
- $d_{\gamma_i}(Q(r(s_i^-)), Q(r(s_i^+))) \asymp_{1,B} d_{\gamma_i}(\rho(q_i^-), \rho(q_i^+))$, for a constant B depending only on D .

By the setup of q_i^- and q_i^+ we have $q_i^+ - q_i^- \leq L$, where

$$L = 2k(K_2(w + C_2)) + 2kc.$$

Then since N is a (K_1, C_1) -quasi-isometry the length of the interval $[s_i^-, s_i^+]$ is bounded above by $K_1L + C_1$. This fact and the first bullet above allow us to apply Lemma 4.3 to the geodesic segment $r|_{[s_i^-, s_i^+]}$ and conclude that there exists $\epsilon_i > 0$ depending on the upper bound for the length of the interval $[s_i^-, s_i^+]$, the lower bound $\omega(L_S)$ in the first bullet above and the value of the annular coefficient $d_{\gamma_i}(Q(r(s_i^-)), Q(r(s_i^+)))$ so that

$$\inf_{t \in [s_i^-, s_i^+]} \ell_{\gamma_i}(r(t)) \leq \epsilon_i.$$

Moreover, the second bullet above, the quasi-equality (4-7) and the assumption that

$$d_{\gamma_i}(v^-, v^+) \rightarrow \infty \quad \text{as } i \rightarrow \infty$$

together imply that

$$d_{\gamma_i}(Q(r(s_i^-)), Q(r(s_i^+))) \rightarrow \infty \quad \text{as } i \rightarrow \infty.$$

Then the last statement of Lemma 4.3 guarantees that $\epsilon_i \rightarrow 0$ as $i \rightarrow \infty$.

Let $b_i \in [s_i^-, s_i^+]$ be the time that the above infimum is realized. Then $\ell_{\gamma_i}(r(b_i)) \rightarrow 0$ as $i \rightarrow \infty$. Moreover, since $q_i \rightarrow \infty$ as $i \rightarrow \infty$, we have $q_i^- \rightarrow \infty$ as $i \rightarrow \infty$. Thus $s_i^- \rightarrow \infty$ as $i \rightarrow \infty$. Then $b_i \rightarrow \infty$ as $i \rightarrow \infty$. This completes the proof of the lemma. \square

Proof of Theorem 1.1 Let $\{\gamma_i\}_{i=0}^\infty$ be a sequence of curves as in Section 3 and let ν^+ be the minimal filling non-uniquely ergodic lamination in $\mathcal{EL}(S)$ which is determined by the sequence. Let ν^- be a marking containing $\gamma_0, \dots, \gamma_3$ as in Section 3. Then by Theorem 3.7(3) the pair (ν^-, ν^+) has non-annular R -bounded combinatorics. Let $X \in \text{Teich}(S)$ be a point with a Bers marking ν^- . By Lemma 2.12 there is a geodesic ray $r: [0, \infty) \rightarrow \text{Teich}(S)$ with $r(0) = X$ and the forward ending lamination ν^+ . Then Theorem 4.1 implies that \hat{r} is recurrent to a compact subset of $\mathcal{M}(S)$. Furthermore, by Theorem 3.7(2),

$$d_{\gamma_i}(\nu^-, \nu^+) \geq \frac{1}{K} e_{i-1} - C.$$

Then since $e_i \rightarrow \infty$ as $i \rightarrow \infty$, we have $d_{\gamma_i}(\nu^-, \nu^+) \rightarrow \infty$ as $i \rightarrow \infty$. Thus by Theorem 4.4 the ray \hat{r} is not contained in any compact subset of $\mathcal{M}(S)$. \square

Remark 4.5 Masur’s criterion (Theorem 1.2) guarantees that any Teichmüller geodesic ray with vertical lamination ν^+ is divergent in $\mathcal{M}(S)$.

References

- [1] T Aougab, *Uniform hyperbolicity of the graphs of curves*, *Geom. Topol.* 17 (2013) 2855–2875 MR3190300
- [2] J A Behrstock, *Asymptotic geometry of the mapping class group and Teichmüller space*, *Geom. Topol.* 10 (2006) 1523–1578 MR2255505
- [3] F Bonahon, *Geodesic laminations on surfaces*, from: “Laminations and foliations in dynamics, geometry and topology”, (M Lyubich, J W Milnor, Y N Minsky, editors), *Contemp. Math.* 269, Amer. Math. Soc. (2001) 1–37 MR1810534
- [4] M R Bridson, A Haefliger, *Metric spaces of non-positive curvature*, *Grundlehr. Math. Wissen.* 319, Springer, Berlin (1999) MR1744486
- [5] J F Brock, *The Weil–Petersson metric and volumes of 3–dimensional hyperbolic convex cores*, *J. Amer. Math. Soc.* 16 (2003) 495–535 MR1969203
- [6] J Brock, H Masur, *Coarse and synthetic Weil–Petersson geometry: Quasi-flats, geodesics and relative hyperbolicity*, *Geom. Topol.* 12 (2008) 2453–2495 MR2443970
- [7] J Brock, H Masur, Y Minsky, *Asymptotics of Weil–Petersson geodesic, I: Ending laminations, recurrence, and flows*, *Geom. Funct. Anal.* 19 (2010) 1229–1257 MR2585573

- [8] **J Brock, H Masur, Y Minsky**, *Asymptotics of Weil–Petersson geodesics, II: Bounded geometry and unbounded entropy*, *Geom. Funct. Anal.* 21 (2011) 820–850 [MR2827011](#)
- [9] **P Buser**, *Geometry and spectra of compact Riemann surfaces*, *Progress in Mathematics* 106, Birkhäuser, Boston (1992) [MR1183224](#)
- [10] **RD Canary, DB A Epstein, PL Green**, *Notes on notes of Thurston*, from: “Fundamentals of hyperbolic geometry: Selected expositions”, (RD Canary, D Epstein, A Marden, editors), *London Math. Soc. Lecture Note Ser.* 328, Cambridge Univ. Press (2006) 1–115 [MR2235710](#)
- [11] **G Daskalopoulos, R Wentworth**, *Classification of Weil–Petersson isometries*, *Amer. J. Math.* 125 (2003) 941–975 [MR1993745](#)
- [12] **B Farb, A Lubotzky, Y Minsky**, *Rank-1 phenomena for mapping class groups*, *Duke Math. J.* 106 (2001) 581–597 [MR1813237](#)
- [13] **A Fathi, F Laudenbach, V Poenaru**, *Travaux de Thurston sur les surfaces*, *Astérisque* 66–67, Soc. Math. France, Paris (1979) [MR568308](#)
- [14] **D Gabai**, *Almost filling laminations and the connectivity of ending lamination space*, *Geom. Topol.* 13 (2009) 1017–1041 [MR2470969](#)
- [15] **S Hensel, P Przytycki, R CH Webb**, *1–slim triangles and uniform hyperbolicity for arc graphs and curve graphs*, *J. Eur. Math. Soc.* 17 (2015) 755–762 [MR3336835](#)
- [16] **J Hubbard, H Masur**, *Quadratic differentials and foliations*, *Acta Math.* 142 (1979) 221–274 [MR523212](#)
- [17] **E Klarreich**, *The boundary at infinity of the curve complex*, preprint (1999) Available at <http://www.msri.org/people/members/klarreic/curvecomplex.ps>
- [18] **C Leininger, A Lenzen, K Rafi**, *Limit sets of Teichmüller geodesics with minimal non-uniquely ergodic vertical foliation*, preprint (2015) [arXiv:1312.2305](#)
- [19] **G Levitt**, *Foliations and laminations on hyperbolic surfaces*, *Topology* 22 (1983) 119–135 [MR683752](#)
- [20] **H Masur**, *Extension of the Weil–Petersson metric to the boundary of Teichmüller space*, *Duke Math. J.* 43 (1976) 623–635 [MR0417456](#)
- [21] **H Masur**, *Hausdorff dimension of the set of nonergodic foliations of a quadratic differential*, *Duke Math. J.* 66 (1992) 387–442 [MR1167101](#)
- [22] **HA Masur, YN Minsky**, *Geometry of the complex of curves, I: Hyperbolicity*, *Invent. Math.* 138 (1999) 103–149 [MR1714338](#)
- [23] **HA Masur, YN Minsky**, *Geometry of the complex of curves, II: Hierarchical structure*, *Geom. Funct. Anal.* 10 (2000) 902–974 [MR1791145](#)
- [24] **B Modami**, *Prescribing the behavior of Weil–Petersson geodesics in the moduli space of Riemann surfaces*, *J. Topol. Anal.* 7 (2015) 543–676 [MR3400125](#)

- [25] **K Rafi**, *A characterization of short curves of a Teichmüller geodesic*, *Geom. Topol.* 9 (2005) 179–202 [MR2115672](#)
- [26] **K Rafi**, *Hyperbolicity in Teichmüller space*, *Geom. Topol.* 18 (2014) 3025–3053 [MR3285228](#)
- [27] **K Rafi, S Schleimer**, *Covers and the curve complex*, *Geom. Topol.* 13 (2009) 2141–2162 [MR2507116](#)
- [28] **C Series**, *The modular surface and continued fractions*, *J. London Math. Soc.* 31 (1985) 69–80 [MR810563](#)
- [29] **S A Wolpert**, *Geometry of the Weil–Petersson completion of Teichmüller space*, *Surv. Differ. Geom.* 8, International Press, Somerville, MA (2003) 357–393 [MR2039996](#)
- [30] **S A Wolpert**, *Families of Riemann surfaces and Weil–Petersson geometry*, *CBMS Regional Conference Series in Mathematics* 113, Amer. Math. Soc. (2010) [MR2641916](#)

*Department of Mathematics, Brown University
Box 1917, Providence, RI 02912, USA*

*Department of Mathematics, University of Illinois at Urbana-Champaign
1409 W Green St, Urbana, IL 61801, USA*

brock@math.brown.edu, babak.modami@yale.edu

Proposed: Danny Calegari

Seconded: Benson Farb, Jean-Pierre Otal

Received: 21 September 2014

Accepted: 6 April 2015

