

The metric geometry of the Hamming cube and applications

FLORENT BAUDIER
DANIEL FREEMAN
THOMAS SCHLUMPRECHT
ANDRÁS ZSÁK

The Lipschitz geometry of segments of the infinite Hamming cube is studied. Tight estimates on the distortion necessary to embed the segments into spaces of continuous functions on countable compact metric spaces are given. As an application, the first nontrivial lower bounds on the $C(K)$ -distortion of important classes of separable Banach spaces, where K is a countable compact space in the family $\{[0, \omega], [0, \omega \cdot 2], \dots, [0, \omega^2], \dots, [0, \omega^k \cdot n], \dots, [0, \omega^\omega]\}$ are obtained.

46B20, 46B80; 46B85

In memory of Luis Sánchez-González

1 Introduction

1.1 Motivation and background

Assume that one is given a Banach space Y and a class \mathcal{C} of metric spaces. Given an *arbitrary* metric space M in the class \mathcal{C} , it is natural to study the smallest distortion achievable when trying to embed M into Y through a bi-Lipschitz embedding. This quite general quantitative embedding problem is an important topic in the nonlinear geometry of Banach spaces. When Y is a Hilbert space this problem is known as estimating the Euclidean distortion of the given class. It is well recognized that being able to accurately estimate the Euclidean distortion of some specific classes of metric spaces has tremendous and far reaching applications in both mathematics and computer science. In this paper we consider the general embedding problem when Y is the Banach space $C(K)$, the space of continuous functions on a compact topological space K . We will mainly stay in the separable world and therefore consider only compact *metric* spaces K as well as classes \mathcal{C} contained in the class \mathcal{M} of separable metric spaces. The theory is clearly isometric and, although c_0 is not isometric to a $C(K)$ -space for any compact space K , embeddings into c_0 are related to those into $C(K)$ -spaces. Indeed, c_0 is a hyperplane of the space c of convergent sequences of real numbers, which can

be seen as the space $C(K)$, where $K = \gamma\mathbb{N}$ is the Alexandrov compactification (or one-point compactification) of \mathbb{N} . Moreover, it is easy to show that, whenever K is an infinite (not necessarily metrizable) compact Hausdorff space, $C(K)$ contains a subspace isometric to c_0 (see Albiac and Kalton [2, Proposition 4.3.11]). We browse briefly and chronologically through a few classical and historical embedding results into $C(K)$ -spaces and c_0 . Back in 1906, Fréchet observed [6] that every separable metric space admits an isometric embedding into the space $\ell_\infty(\mathbb{N})$. An easy application of the Hahn–Banach theorem gives a linear isometric embedding of every separable Banach space into $\ell_\infty(\mathbb{N})$. These results can actually be cast as embedding results into a $C(K)$ -space. Indeed, $\ell_\infty(\mathbb{N})$ can be identified with the space $C(\beta\mathbb{N})$, where $\beta\mathbb{N}$ denotes the Stone–Čech compactification of \mathbb{N} . Note that $\beta\mathbb{N}$ is an uncountable compact space, and since $\ell_\infty(\mathbb{N})$ is nonseparable, $\beta\mathbb{N}$ is not metrizable. The Banach–Mazur theorem [4] asserts that every separable Banach space admits a (linear) isometric embedding into the space $C[0, 1]$. Note that $[0, 1]$ equipped with its canonical distance is compact and hence $C[0, 1]$ is separable. With the help of Fréchet’s embedding, it is easily seen that every separable metric space can be isometrically embedded into $C[0, 1]$. In 1974 Aharoni proved in [1] that the c_0^+ -distortion of every separable metric space is less than 6. In that same article he also proved that the c_0 -distortion of ℓ_1 is at least 2. A few years later Assouad [3] showed that the c_0^+ -distortion of every separable metric space is at most 3. The fact that there is a bi-Lipschitz embedding with distortion exactly 3 and that this value is optimal for embeddings into c_0^+ is due to Pelant [10]. Finally, the end of the story regarding embeddings into c_0 was completed by Kalton and Lancien [8] when they constructed an embedding with distortion 2 (respectively, 1) for every separable (respectively, proper) metric space. Recall that a metric space is *proper* if all its closed balls are compact.

1.2 Notation and definitions

Let M and N be two metric spaces. Define the *distortion* of a map $f: M \rightarrow N$ to be

$$\text{dist}(f) := \|f\|_{\text{Lip}} \|f^{-1}\|_{\text{Lip}} = \left(\sup_{x \neq y \in M} \frac{d_N(f(x), f(y))}{d_M(x, y)} \right) \left(\sup_{x \neq y \in M} \frac{d_M(x, y)}{d_N(f(x), f(y))} \right).$$

If the distortion of f is finite, f is said to be a *bi-Lipschitz embedding*. The convenient notation

$$M \xrightarrow[\text{Lip}]{} N$$

means that there exists a bi-Lipschitz embedding f from M into N . We are concerned with the quantitative theory, and if $\text{dist}(f) \leq C$, we use the notation

$$M \xrightarrow[C\text{-Lip}]{} N.$$

The parameter

$$c_N(M) = \inf \left\{ C \geq 1 : M \xrightarrow[C\text{-Lip}]{} N \right\}$$

will be referred to as *the N -distortion of M* .

Let \mathcal{F} be a collection of metric spaces. We can define *the N -distortion of the class \mathcal{F}* as follows:

$$c_N(\mathcal{F}) = \sup \{ c_N(M) : M \in \mathcal{F} \}.$$

Finally, for two families \mathcal{F} and \mathcal{G} of metric spaces we define

$$c_{\mathcal{G}}(\mathcal{F}) = \sup_{M \in \mathcal{F}} \inf_{N \in \mathcal{G}} c_N(M).$$

As an application of our work on the metric geometry of the Hamming cube we will give nontrivial estimates on the parameter $c_N(\mathcal{F})$ for the following spaces and classes:

- $N = C(K)$ for some countable compact metric space K .
- \mathcal{F} is one of the following classes:
 - (1) $\mathcal{M} := \{M : M \text{ separable metric space}\}$.
 - (2) $\mathcal{SB} := \{X : X \text{ separable Banach space}\}$.
 - (3) $\mathcal{COT} := \{X : X \text{ separable Banach space with nontrivial cotype}\}$.
 - (4) $\mathcal{TCP} := \{X : X \text{ separable Banach space with nontrivial type}\}$.
 - (5) $\mathcal{SR} := \{X : X \text{ separable, reflexive Banach space}\}$.

Observe that $c_N(\mathcal{SB}) = c_N(\mathcal{M})$. Indeed, it is clear that $c_N(\mathcal{SB}) \leq c_N(\mathcal{M})$, and the reverse inequality follows from the fact that every separable metric space embeds isometrically into the separable Banach space $C[0, 1]$.

1.3 Stratification of the Hamming cube

We define a *stratification* of a metric space M to be a sequence $M_1 \subset M_2 \subset \dots$ of subsets of M such that $M = \bigcup_{k=1}^{\infty} M_k$. (More generally, it is a way of expressing M as a direct limit of metric spaces, but this generality will not be needed here.) The sets M_k are the *segments* of M , and the sets $M_k \setminus M_{k-1}$ are the *layers* of M (where we put $M_0 = \emptyset$). In this paper we are concerned with stratifications of the Hamming cube. The infinite Hamming cube H_{∞} is the set of all infinite sequences in $\{0, 1\}$ containing only finitely many 1s, equipped with the Hamming metric d_H , where $d_H(\sigma, \tau) = |\{i \in \mathbb{N} : \sigma_i \neq \tau_i\}|$. It is isometric to the metric space Δ_{∞} consisting of the set $[\mathbb{N}]^{<\omega}$ of all finite subsets of \mathbb{N} equipped with the symmetric difference metric d_{Δ} , where $d_{\Delta}(A, B) = |A \Delta B|$. The isometry between Δ_{∞} and H_{∞} is the natural one identifying a set with its indicator function.

We describe two natural stratifications of the infinite Hamming cube. For $k \in \mathbb{N}$ let $H_k = \{0, 1\}^k$ thought of as a subset of H_∞ by extending elements of H_k to infinite binary sequences with the addition of an infinite tail of 0s. The layers of the stratification $(H_k)_{k=0}^\infty$ are $\{\emptyset\}$ and families of sets of the form $\{A \subset \mathbb{N} : \max A = n\}$, $n \in \mathbb{N}$. The members of the second stratification are the families $\Delta_k = [\mathbb{N}]^{\leq k}$ of subsets of \mathbb{N} of size at most k . The set Δ_k can be identified with the rooted countably infinitely branching tree of height k . Note, however, that the metric d_Δ is not the classical graph metric of a tree.

The two stratifications share some essential metric properties despite being quite different from the combinatorial and structural standpoints. For example, Δ_k (respectively, H_k) is a $2k$ -bounded (respectively, k -bounded), 1-separated metric space. However, Δ_k is a countable non-proper metric space while H_k is a finite metric space. The two stratifications are different in the Lipschitz category in the following sense. Two families of metric spaces \mathcal{F} and \mathcal{G} shall be called *Lipschitz equivalent* if $c_{\mathcal{F}}(\mathcal{G})c_{\mathcal{G}}(\mathcal{F}) < \infty$. The stratifications $\mathcal{O} = (H_k)_{k \geq 0}$ and $\mathcal{U} = (\Delta_k)_{k \geq 0}$ are not Lipschitz equivalent. Indeed, the embedding $(\sigma_1, \dots, \sigma_k) \mapsto \{i \in \{1, \dots, k\} : \sigma_i = 1\}$ sends H_k isometrically into Δ_k (ie $c_{\mathcal{U}}(\mathcal{O}) = 1$), however, $c_{\mathcal{O}}(\mathcal{U}) = \infty$ since it is impossible to embed a single Δ_k bi-Lipschitzly into any H_i because of a cardinality obstruction (assuming $k \geq 1$ of course).

Sometimes metric information about a stratification can be used to derive metric information on the stratified space and vice versa. However, this need not be the case. As we will see H_∞ does not embed isometrically into c_0 and this will be witnessed by Δ_2 . This is in stark contrast with the fact that every H_k , as any finite metric space, embeds isometrically into c_0 . So in some sense $(\Delta_k)_{k \geq 0}$ captures more of the structure of Δ_∞ .

1.4 Organization of the article

From now on we will consider *countable* compact metric spaces and we will focus on the following nested family:

$$[0, \omega] \subset [0, \omega \cdot 2] \subset \dots \subset [0, \omega^2] \subset \dots \subset [0, \omega^\alpha \cdot n] \subset \dots \subset [0, \omega^\omega],$$

where, as usual, ω is the first infinite ordinal. It is a simple fact that if compact spaces K and L are homeomorphic then the Banach spaces $C(K)$ and $C(L)$ are isometrically isomorphic. Note that the converse is also true by the Banach–Stone theorem. Therefore the $C(K)$ -spaces arising from the nested family above are mutually non-isometric Banach spaces. However, this family has the property that $C(K)$ embeds linearly isometrically into $C(L)$ whenever $K \subset L$, since then K is in fact a clopen subset of L .

In Section 2 we estimate from above the $C(K)$ -distortion of the infinite Hamming cube and its stratification Δ_k . We will show that $c_{C([0,\omega^r])}(\Delta_k) \leq \min\{k/r, 2\}$ when $1 \leq r \leq k < \infty$. In particular, Δ_k embeds isometrically into $C([0, \omega^k])$. In Section 3 we will give lower bounds. To estimate $c_{C([0,\omega^r])}(\Delta_k)$ from below, we exhibit a connection between a topological property of the compact space K and the $C(K)$ -distortion of the metric spaces Δ_k . Roughly speaking, if the compact metric space K is small in the sense of the Cantor–Bendixson derivation, then the $C(K)$ -distortion of Δ_k cannot be too small. More precisely, we show that if the Cantor–Bendixson index of K is $k \geq 2$, then $c_{C(K)}(\Delta_k) \geq k/(k - 1)$. In Section 4 we give some applications concerning the parameters $c_{C(K)}(\mathcal{M})$, $c_{C(K)}(SB)$, $c_{C(K)}(\mathcal{COT})$ and $c_{C(K)}(SR)$. We conclude with a few open questions that arise naturally from our work.

Acknowledgements Baudier’s research was supported by ANR grant 13-PDOC-0031, project NoLiGeA. Freeman’s research was supported by NSF grant DMS-1332255. Schlumprecht’s research was supported by NSF grant DMS-1160633. Freeman and Zsák were supported by the Workshop in Analysis and Probability at Texas A&M University in 2013. Zsák was supported by Texas A&M University while he was a Visiting Scholar there in 2014.

2 Low distortion embeddings of the Hamming cube

2.1 Embeddings of the sets Δ_k

We will show, by constructing suitable bi-Lipschitz embeddings, that $c_{C([0,\omega^r])}(\Delta_k) \leq \min\{k/r, 2\}$ when $1 \leq r \leq k < \infty$. In particular, Δ_k embeds isometrically into $C([0, \omega^k])$, and hence also into $C([0, \omega^r])$ for $r \geq k$.

We will need a description of $C(K)$ -spaces as *tree spaces*, due to Bourgain [5] (see also [9]), which we now proceed to describe. Recall that a *tree* is a set T with a partial order \preceq such that $b_t = \{s \in T : s \preceq t\}$ is finite and linearly ordered by \preceq for all $t \in T$. The space $c_{00}(T)$ consists of all functions $x: T \rightarrow \mathbb{R}$ for which $\{t \in T : x(t) \neq 0\}$ is finite. The unit vector basis $(e_t)_{t \in T}$ of $c_{00}(T)$ consists of functions e_t taking the value 1 at t and 0 everywhere else. For $t \in T$ the functional β_t is defined by summing along the branch b_t :

$$\beta_t(x) = \sum_{s \in b_t} x(s) \quad (x \in c_{00}(T)).$$

We define a norm $\|\cdot\|$ on $c_{00}(T)$ by letting

$$\|x\| = \sup_{t \in T} |\beta_t(x)| \quad (x \in c_{00}(T)).$$

The tree space corresponding to T is the completion $S(T)$ of $(c_{00}(T), \|\cdot\|)$. It is easy to verify that (e_t) is a normalized, monotone basis of $S(T)$. Note that the branch functionals can be expressed in terms of the biorthogonal functional as follows: $\beta_t = \sum_{s \in b_t} e_s^*$. We now let K be the w^* -closure in $S(T)^*$ of the set $\{\beta_t : t \in T\}$. This is a compact Hausdorff space and $0 \in K$ if and only if T has infinitely many *initial nodes* (ie elements $t \in T$ for which $s \preceq t$ implies $s = t$). The restriction to K of the image of each element of $S(T)$ under the canonical embedding of $S(T)$ into $S(T)^{**}$ is an isometric isomorphism $S(T) \rightarrow C(K)$. By the Stone–Weierstrass theorem it is onto $C(K)$ if $0 \notin K$ and onto $C_0(K)$ (functions vanishing at 0) if $0 \in K$. It turns out that every $C(K)$ -space with separable dual can be represented as a tree space but we will not need this result in its full generality. We will now mention the examples relevant to us.

For each $k \in \mathbb{N}$ let T_k be the tree $([\mathbb{N}]^{\leq k}, \preceq)$, where $s \preceq t$ if and only if s is an initial segment of t . Thus T_k is the rooted, countably infinitely branching tree of height k . As usual, we identify a set $t \subset \mathbb{N}$ with the sequence i_1, i_2, \dots , where $i_1 < i_2 < \dots$ are the elements of t . So, for example, we shall write e_m for the basis element $e_{\{m\}}$ of $S(T_k)$, etc. The set $\{\beta_t : t \in T_k\}$ is homeomorphic to $(0, \omega^k]$ (and hence to $[0, \omega^k]$) via the map $\beta_\emptyset \mapsto \omega^k$ and

$$(i_1, \dots, i_r) \mapsto \sum_{j=1}^{r-1} \omega^{k-j} (i_j - i_{j-1} - 1) + \omega^{k-r} (i_r - i_{r-1}),$$

for $1 \leq r \leq k$, $i_1 < \dots < i_r$ (and with $i_0 = 0$). Thus $S(T_k) \cong C([0, \omega^k])$. Let us now denote by T the disjoint union of the trees T_k . For $s, t \in T$ we have $s \preceq t$ if and only if for some k both s and t belong to T_k and $s \preceq t$ in T_k . The tree space $S(T)$ is then isometrically isomorphic to $C_0([0, \omega^\omega])$ which of course isometrically embeds into $C([0, \omega^\omega])$. Note also that $S(T) \cong (\bigoplus_{k=1}^\infty S(T_k))_{c_0}$. For the rest of the paper we fix T_k and T to be the trees just described.

Theorem 1 *For every $1 \leq r \leq k$ there exist a map $\varphi_{k,r} : \Delta_k \rightarrow C([0, \omega^r])$ such that $\text{dist}(\varphi_{k,r}) \leq k/r$. It follows that $c_{C([0, \omega^r])}(\Delta_k) \leq \min\{k/r, 2\}$.*

Proof For each $r \in \mathbb{N}$ we define the map

$$f_r : \mathbb{N} \rightarrow S(T_r), \quad m \mapsto - \sum_{i=1}^{m-1} e_i + e_m + 2 \sum_{j=2}^r \sum_{\substack{i_1 < \dots < i_j \\ i_j = m}} e_{i_1, \dots, i_j}.$$

Then for $1 \leq r \leq k$ define

$$\varphi_{k,r} : \Delta_k \rightarrow S(T_r), \quad \sigma \mapsto \sum_{m \in \sigma} f_r(m).$$

Let $\sigma, \tau \in \Delta_k$. We will show that

$$\frac{r}{k} d_{\Delta}(\sigma, \tau) \leq \|\varphi_{k,r}(\sigma) - \varphi_{k,r}(\tau)\| \leq d_{\Delta}(\sigma, \tau).$$

Let $i_1 < \dots < i_s$ and $j_1 < \dots < j_t$ be the elements of $\sigma \setminus \tau$ and $\tau \setminus \sigma$, respectively. We need to show that

$$\frac{r}{k}(s + t) \leq \|f_r(i_1) + \dots + f_r(i_s) - f_r(j_1) - \dots - f_r(j_t)\| \leq (s + t).$$

The upper bound follows from the triangle inequality. Indeed, for each $m \in \mathbb{N}$ and for each $t \in T_r$, summing $f_r(m)$ along the branch b_t yields the values $-1, 0, 1$, and hence $f_r(m)$ is of norm 1. To see the lower bound, first note that we can assume without loss of generality that $1 \leq s$ and that either $t = 0$ or $i_1 < j_1$. We will then prove the following statement by induction on $\max\{s, t\}$. Given $s + t$ distinct positive integers $i_1 < \dots < i_s$ and $j_1 < \dots < j_t$, where $1 \leq s \leq k$ and either $t = 0$ or $1 \leq t \leq k$ and $i_1 < j_1$, setting

$$g = f_r(i_1) + \dots + f_r(i_s) - f_r(j_1) - \dots - f_r(j_t),$$

there is a branch functional $\beta_{\ell_1, \dots, \ell_u}$ with $1 \leq u \leq r$ and $i_1 \leq \ell_1$ such that

$$|\beta_{\ell_1, \dots, \ell_u}(g)| = \left| \sum_{v=1}^u e_{\ell_1, \dots, \ell_v}^*(g) \right| \geq \frac{r}{\max\{r, s, t\}}(s + t).$$

This clearly implies the lower bound of $(r/k)(s + t)$ on the norm of g .

If $s \leq r$ or $r \leq s \leq t$, then for $u = \min\{r, s\}$ we have

$$\begin{aligned} \beta_{i_1, \dots, i_u}(g) &= e_{i_1}^* \left(f_r(i_1) + \sum_{m=2}^s f_r(i_m) - \sum_{n=1}^t f_r(j_n) \right) + \sum_{v=2}^u e_{i_1, \dots, i_v}^*(f_r(i_v)) \\ &= 1 - (s - 1) + t + 2(u - 1) = -s + t + 2u. \end{aligned}$$

When $s \leq r$, then $-s + t + 2u = s + t$, and we are done. If $r \leq s \leq t$, then

$$-s + t + 2u \geq 2r = \frac{r}{(1/2)(s + t)}(s + t) \geq \frac{r}{\max\{r, s, t\}}(s + t),$$

as required. We finally deal with the case when $r < s$ and $t < s$. Set

$$h = f_r(i_2) + \dots + f_r(i_s) - f_r(j_1) - \dots - f_r(j_t).$$

If $t = 0$ or $i_2 < j_1$, then we apply the induction hypothesis to h , and if $j_1 < i_2$, then we apply the induction hypothesis to $-h$. In either case we obtain a branch functional $\beta_{\ell_1, \dots, \ell_u}$ such that $i_1 < \ell_1$ and

$$|\beta_{\ell_1, \dots, \ell_u}(h)| \geq \frac{r}{s-1}(s + t - 1).$$

Since $i_1 < \ell_1$, we have $\beta_{\ell_1, \dots, \ell_u}(f_r(i_1)) = 0$, and it follows that

$$|\beta_{\ell_1, \dots, \ell_u}(g)| = |\beta_{\ell_1, \dots, \ell_u}(h)| \geq \frac{r}{s-1}(s+t-1) \geq \frac{r}{s}(s+t).$$

This completes the proof that $\text{dist}(\varphi_{k,r}) \leq k/r$. Recall that Kalton and Lancien [8] proved that every separable metric space embeds into c_0 with distortion at most 2. It follows that $c_{C([0, \omega^r])}(\Delta_k) \leq \min\{k/r, 2\}$. □

2.2 $C([0, \omega^\omega])$ -distortion of the Hamming cube

It follows from Theorem 1 that each Δ_k embeds isometrically into $C([0, \omega^\omega])$. We now prove a stronger result. Recall that a set $A \subset \mathbb{N}$ is a *Schreier set* if $|A| \leq \min A$ (or if $A = \emptyset$). This inequality is usually referred to as the *admissibility condition*. The Schreier family, the set of all Schreier sets, is denoted by \mathcal{S}_1 . We endow \mathcal{S}_1 with the symmetric difference metric, ie we consider \mathcal{S}_1 as a subset of Δ_∞ .

Theorem 2 $(\mathcal{S}_1, d_\Delta)$ embeds isometrically into $C([0, \omega^\omega])$.

Proof Define

$$f_\omega: \mathbb{N} \rightarrow S(T), \quad m \mapsto \sum_{k=1}^m f_k(m),$$

where $f_k, k \in \mathbb{N}$, are the functions defined in Theorem 1. Here we identify $x \in S(T_k)$ with the sequence in $S(T) \cong (\bigoplus_{k=1}^\infty S(T_k))_{c_0}$ that has x in the k^{th} co-ordinate and zero everywhere else. Thus, more precisely,

$$f_\omega(m) = (f_1(m), \dots, f_m(m), 0, 0, \dots).$$

We next define

$$\varphi_\omega: \mathcal{S}_1 \rightarrow S(T), \quad \sigma \mapsto \sum_{m \in \sigma} f_\omega(m),$$

and claim that this is an isometric embedding. As before, this amounts to showing that if $\sigma, \tau \in \mathcal{S}_1$ and $i_1 < \dots < i_s$ and $j_1 < \dots < j_t$ are the elements of $\sigma \setminus \tau$ and $\tau \setminus \sigma$, respectively, then

$$\|f_\omega(i_1) + \dots + f_\omega(i_s) - f_\omega(j_1) - \dots - f_\omega(j_t)\| = s + t.$$

Setting $g = f_\omega(i_1) + \dots + f_\omega(i_s) - f_\omega(j_1) - \dots - f_\omega(j_t)$, we have $\|g\| \leq s + t$ by the triangle inequality. Indeed, for each $m \in \mathbb{N}$ we have

$$\|f_\omega(m)\| = \max_{1 \leq k \leq m} \|f_k(m)\| = 1.$$

To get the lower bound, we may assume without loss of generality that $i_1 < j_1$ (or $t = 0$) and consider the k^{th} component of g in $S(T_k)$, where $k = i_1$. We will show that

$$\|f_k(i_1) + \dots + f_k(i_s) - f_k(j_1) - \dots - f_k(j_t)\| = s + t.$$

Note that $s \leq |\sigma| \leq \min \sigma \leq i_1 = k$. It follows that we can get the lower bound $s + t$ by applying the branch functional β_{i_1, \dots, i_s} in T_k as in the proof of Theorem 1. \square

We now turn our attention to the infinite Hamming cube. With the help of Theorem 2 we are now able to embed the infinite Hamming cube into $C([0, \omega^\omega])$ with arbitrarily small distortion. We say that M embeds *almost isometrically* into N , denoted by

$$M \xrightarrow[\text{a.i.}]{} N,$$

if for every $\varepsilon > 0$ there exists a bi-Lipschitz embedding f from M into N with $\text{dist}(f) \leq 1 + \varepsilon$.

Theorem 3 *The infinite Hamming cube Δ_∞ embeds almost isometrically into $C([0, \omega^\omega])$.*

Proof As before, we will in fact embed Δ_∞ into $C_0([0, \omega^\omega])$, which is identified with $S(T) \cong (\bigoplus_{k=1}^\infty S(T_k))_{c_0}$. Fix $\varepsilon > 0$. Choose a sequence $0 = N_0 < N_1 < N_2 < \dots$ of integers satisfying

$$(1) \quad 2m \leq \varepsilon N_m \quad \text{for all } m \geq 0.$$

We next define maps f, φ similar to f_ω, Δ_ω but with a different admissibility condition. It will be clear from the definition and the proof of Theorem 2 that this new map φ will be an isometric embedding when restricted to the class of sets σ satisfying the admissibility condition $|\sigma| \leq N_{\min(\sigma)}$. We define

$$f: \mathbb{N} \rightarrow S(T), \quad m \mapsto \sum_{k=1}^{N_m} f_k(m)$$

and

$$\varphi: \Delta_\infty \rightarrow S(T), \quad \sigma \mapsto \sum_{m \in \sigma} f(m).$$

Fix $\sigma, \tau \in \Delta_\infty$. We will show that

$$(2) \quad (1 - \varepsilon)d_\Delta(\sigma, \tau) \leq \|\varphi(\sigma) - \varphi(\tau)\| \leq d_\Delta(\sigma, \tau).$$

By the triangle inequality, we have

$$\|\varphi(\sigma) - \varphi(\tau)\| = \left\| \sum_{m \in \sigma} f(m) - \sum_{m \in \tau} f(m) \right\| \leq \sum_{m \in \sigma \Delta \tau} \|f(m)\| = d_\Delta(\sigma, \tau).$$

To show the lower bound, we first observe that σ and τ can be assumed to be disjoint. Indeed, we have

$$\varphi(\sigma) - \varphi(\tau) = \varphi(\sigma \setminus \tau) - \varphi(\tau \setminus \sigma) \quad \text{and} \quad d_{\Delta}(\sigma, \tau) = d_{\Delta}(\sigma \setminus \tau, \tau \setminus \sigma),$$

and so we can replace σ and τ with $\sigma \setminus \tau$ and $\tau \setminus \sigma$ if necessary.

We next choose $m, n \in \mathbb{N}$ such that

$$(3) \quad N_{m-1} < |\sigma| \leq N_m \quad \text{and} \quad N_{n-1} < |\tau| \leq N_n.$$

Set $\sigma' = \sigma \setminus \{1, \dots, m-1\}$ and $\tau' = \tau \setminus \{1, \dots, n-1\}$. Since σ' and τ' are admissible, we have

$$(4) \quad \|\varphi(\sigma') - \varphi(\tau')\| = d_{\Delta}(\sigma', \tau').$$

Next, since σ' and τ' are small perturbations of σ and τ , respectively, we have

$$(5) \quad \left| \|\varphi(\sigma) - \varphi(\tau)\| - \|\varphi(\sigma') - \varphi(\tau')\| \right| \leq \|\varphi(\sigma) - \varphi(\sigma')\| + \|\varphi(\tau) - \varphi(\tau')\| \\ \leq d_{\Delta}(\sigma, \sigma') + d_{\Delta}(\tau, \tau') \leq (m-1) + (n-1)$$

and

$$(6) \quad |d_{\Delta}(\sigma, \tau) - d_{\Delta}(\sigma', \tau')| \leq d_{\Delta}(\sigma, \sigma') + d_{\Delta}(\tau, \tau') \leq (m-1) + (n-1).$$

It follows that

$$\begin{aligned} \|\varphi(\sigma) - \varphi(\tau)\| &\geq \|\varphi(\sigma') - \varphi(\tau')\| - (m+n-2) && \text{(by (5))} \\ &= d_{\Delta}(\sigma', \tau') - (m+n-2) && \text{(by (4))} \\ &\geq d_{\Delta}(\sigma, \tau) - 2(m+n-2) && \text{(by (6))} \\ &= |\sigma| + |\tau| - 2(m+n-2) \\ &= |\sigma| \left(1 - \frac{2(m-1)}{|\sigma|}\right) + |\tau| \left(1 - \frac{2(n-1)}{|\tau|}\right) \\ &\geq (1-\varepsilon)(|\sigma| + |\tau|) = (1-\varepsilon) d_{\Delta}(\sigma, \tau) && \text{(by (1) and (3))} \end{aligned}$$

as required. □

Remark An interesting question presents itself in light of the two theorems above. Does Δ_{∞} almost isometrically embed into \mathcal{S}_1 ? A positive answer together with Theorem 2 would provide another proof of Theorem 3.

3 Estimating the $C(K)$ -distortion from below

3.1 Aharoni’s lower bound observed with “metric lenses”

Aharoni proved that $c_{c_0}(SB) \geq 2$, and hence $c_{c_0}(\mathcal{M}) \geq 2$. Indeed, he showed that the separable Banach space ℓ_1 does not embed into c_0 with distortion strictly less than 2. A careful inspection of his proof shows that the proof and the statement of the result can be carried out and stated without using or even mentioning the linear structure of the Banach space ℓ_1 . This simple but crucial observation allows us to extend Aharoni’s proof to the much more general setting of embeddings into $C(K)$ -spaces.

Denote by $\tilde{\Delta}_2$ the subset $\{\emptyset, \{n\}, \{1, i\}, \{2, j\} : n \geq 1, i \geq 2, j \geq 3\}$ of the metric space Δ_2 . The following theorem is nothing else but Aharoni’s lower bound theorem reformulated in purely metric terms. For the sake of completeness we include the original proof using our notation in the hope that it will make the notation used in the proof of Theorem 6 more accessible.

Theorem 4 (Aharoni) *The metric space $\tilde{\Delta}_2$ does not embed into c_0 with distortion strictly less than 2.*

Proof Assume that $f: \tilde{\Delta}_2 \rightarrow c_0$ and $C < 2$ satisfy

$$d_{\Delta}(\sigma, \tau) \leq \|f(\sigma) - f(\tau)\| \leq Cd_{\Delta}(\sigma, \tau) \quad \text{for all } \sigma, \tau \in \tilde{\Delta}_2.$$

Without loss of generality one can assume that $f(\emptyset) = 0$. Let $f_n = e_n^* \circ f$, so that $f(\sigma) = (f_n(\sigma))_{n=1}^{\infty}$ for $\sigma \in \tilde{\Delta}_2$. For every $i \neq j$ in \mathbb{N} define

$$\mathcal{X}_{i,j} = \{n \in \mathbb{N} : |f_n(\{i\}) - f_n(\{j\})| \geq 4 - 2C\}.$$

Note that these are finite sets. Moreover, $\mathcal{X}_{1,2} \cap \mathcal{X}_{i,j} \neq \emptyset$ for every $i, j \geq 3, i \neq j$. Indeed, we have

$$\|f(\{1, i\}) - f(\{2, j\})\| \geq d_{\Delta}(\{1, i\}, \{2, j\}) = 4.$$

Hence there exists $n_{i,j} \in \mathbb{N}$ such that

$$|f_{n_{i,j}}(\{1, i\}) - f_{n_{i,j}}(\{2, j\})| \geq 4.$$

It follows that

$$\begin{aligned} |f_{n_{i,j}}(\{i\}) - f_{n_{i,j}}(\{j\})| &\geq |f_{n_{i,j}}(\{1, i\}) - f_{n_{i,j}}(\{2, j\})| \\ &\quad - |f_{n_{i,j}}(\{1, i\}) - f_{n_{i,j}}(\{i\})| - |f_{n_{i,j}}(\{2, j\}) - f_{n_{i,j}}(\{j\})| \\ &\geq 4 - \|f(\{1, i\}) - f(\{i\})\| - \|f(\{2, j\}) - f(\{j\})\| \\ &\geq 4 - Cd_{\Delta}(\{1, i\}, \{i\}) - Cd_{\Delta}(\{2, j\}, \{j\}) = 4 - 2C. \end{aligned}$$

This proves that $n_{i,j} \in \mathcal{X}_{i,j}$. Arguing along the same lines, one gets that $n_{i,j} \in \mathcal{X}_{1,2}$ as well. Therefore $\mathcal{X}_{1,2} \cap \mathcal{X}_{i,j} \neq \emptyset$ whenever $i \neq j$ and $i, j \geq 3$. Denote by P the canonical projection from c_0 onto the closed linear span Y of the vectors $(e_n)_{n \in \mathcal{X}_{1,2}}$. We now obtain a contradiction by observing that the sequence $(Pf(\{n\}))_{n=3}^\infty$ is a C -bounded and $(4 - 2C)$ -separated sequence in the finite-dimensional Banach space Y . Indeed, for every $n \geq 3$,

$$\|Pf(\{n\})\| \leq \|f(\{n\})\| = \|f(\{n\}) - f(\emptyset)\| \leq Cd_\Delta(\{n\}, \emptyset) = C,$$

and for every $i \neq j$ with $i, j \geq 3$, we have

$$\begin{aligned} \|Pf(\{i\}) - Pf(\{j\})\| &= \sup_{n \in \mathcal{X}_{1,2}} |f_n(\{i\}) - f_n(\{j\})| \\ &\geq |f_{n_{i,j}}(\{i\}) - f_{n_{i,j}}(\{j\})| \\ &\geq 4 - 2C > 0. \end{aligned} \quad \square$$

3.2 Estimating the $C(K)$ -distortion of Δ_k from below

A key ingredient in estimating from below the $C(K)$ -distortion of the metric space Δ_k is the Cantor–Bendixson derivation for compact spaces. We next recall the definition and a few basic properties of this derivation.

Let K be a compact topological space. The *Cantor–Bendixson derivative* K' of K is the set of all accumulation points of K , ie

$$K' = K \setminus \{x \in K : x \text{ is an isolated point}\}.$$

By transfinite induction one can define derivatives $K^{(\alpha)}$ of higher order α as follows. We set $K^{(0)} = K$. For an ordinal α we let $K^{(\alpha+1)} = (K^{(\alpha)})'$ and, finally, for a non-zero limit ordinal λ we define $K^{(\lambda)} = \bigcap_{\alpha < \lambda} K^{(\alpha)}$.

We gather in the next proposition some basic properties of the Cantor–Bendixson derivation.

Proposition 5 *Let K be a compact metric space. Then:*

- (1) K is finite $\iff K' = \emptyset \iff K$ is discrete.
- (2) K is countable \iff there exists $\alpha < \omega_1$ such that $K^{(\alpha)} = \emptyset$.
- (3) K is uncountable \iff there exists $\alpha < \omega_1$ such that $K^{(\alpha+1)} = K^{(\alpha)} \neq \emptyset$.

For a general compact topological space K the smallest ordinal α such that $K^{(\alpha)} = K^{(\alpha+1)}$ is called the *Cantor–Bendixson index* (or *rank*) of K , and we denote it by $i_{\text{CB}}(K)$. For example, consider the compact space $K = [0, \omega^\alpha \cdot n]$, where $1 \leq \alpha < \omega_1$

and $1 \leq n < \omega$. Then $i_{CB}(K) = \alpha + 1$ and $|K^{(\alpha)}| = n$. More generally, if K is a countably infinite compact metric space, then for some $1 \leq \alpha < \omega_1$ and $1 \leq n < \omega$ we have $i_{CB}(K) = \alpha + 1$, $|K^{(\alpha)}| = n$ and K is homeomorphic to $[0, \omega^\alpha \cdot n]$. Thus, the Cantor–Bendixson derivation gives rise to a topological classification of countable compact metric spaces, and hence an isometric classification of $C(K)$ –spaces with separable dual.

Inspired by the reformulation of Aharoni’s proof in terms of a metric subset of Δ_2 , we establish a link between the $C(K)$ –distortion of the sequence $(\Delta_k)_{k \geq 1}$ and the Cantor–Bendixson index of the compact space K . In Section 2 we showed that $c_{C([0, \omega^{k-1}])}(\Delta_k) \leq k/(k - 1)$ for $k \geq 2$. In the remainder of this section we will show that this upper bound is tight.

Theorem 6 *Let K be a compact topological space and k be an integer with $k \geq 2$. If Δ_k admits a bi-Lipschitz embedding into $C(K)$ with distortion strictly less than $k/(k - 1)$, then $i_{CB}(K) \geq k + 1$. It follows that $c_{C([0, \omega^{k-1}])}(\Delta_k) = k/(k - 1)$.*

Proof Assume that there is a function $f: \Delta_k \rightarrow C(K)$ and a constant $D < k/(k - 1)$ such that

$$d_\Delta(\sigma, \tau) \leq \|f(\sigma) - f(\tau)\| \leq Dd_\Delta(\sigma, \tau) \quad \text{for all } \sigma, \tau \in \Delta_k.$$

Set $\eta = 2k - 2(k - 1)D$ and observe that $\eta > 0$. For distinct $i, j \in \mathbb{N}$ define

$$\mathcal{X}_{i,j} = \{\beta \in K : |f(\{i\})(\beta) - f(\{j\})(\beta)| \geq \eta\}.$$

Consider the following statement. For each $0 \leq s \leq k$ and for any $2(k - s)$ distinct integers $i_1, i_2, \dots, i_{k-s}, j_1, j_2, \dots, j_{k-s}$, we have

$$K^{(s)} \cap \mathcal{X}_{i_1, j_1} \cap \mathcal{X}_{i_2, j_2} \cap \dots \cap \mathcal{X}_{i_{k-s}, j_{k-s}} \neq \emptyset.$$

We will now verify this statement by induction on s . The theorem will then follow by putting $s = k$.

We begin with $s = 0$. Let i_1, \dots, i_k and j_1, \dots, j_k be $2k$ distinct elements of \mathbb{N} . Set $\sigma = \{i_1, \dots, i_k\}$ and $\tau = \{j_1, \dots, j_k\}$. Since $\|f(\sigma) - f(\tau)\| \geq d_\Delta(\sigma, \tau) = 2k$, there exists $\beta \in K$ such that $|f(\sigma)(\beta) - f(\tau)(\beta)| \geq 2k$. It follows that

$$\begin{aligned} |f(\{i_r\})(\beta) - f(\{j_r\})(\beta)| &\geq |f(\sigma)(\beta) - f(\tau)(\beta)| - |f(\sigma)(\beta) - f(\{i_r\})(\beta)| \\ &\quad - |f(\{j_r\})(\beta) - f(\tau)(\beta)| \\ &\geq 2k - \|f(\sigma) - f(\{i_r\})\| - \|f(\{j_r\}) - f(\tau)\| \\ &\geq 2k - Dd_\Delta(\sigma, \{i_r\}) - Dd_\Delta(\{j_r\}, \tau) \\ &\geq 2k - 2D(k - 1) = \eta > 0 \end{aligned}$$

for each $1 \leq r \leq k$. Thus, $\beta \in \mathcal{X}_{i_1, j_1} \cap \dots \cap \mathcal{X}_{i_k, j_k}$.

Now assume that the statement holds for some $0 \leq s < k$. Let i_1, \dots, i_{k-s-1} and j_1, \dots, j_{k-s-1} be $2(k-s-1)$ distinct elements of \mathbb{N} . Let

$$L = K^{(s)} \cap \mathcal{X}_{i_1, j_1} \cap \dots \cap \mathcal{X}_{i_{k-s-1}, j_{k-s-1}}$$

and $n_0 = \max\{i_1, \dots, i_{k-s-1}, j_1, \dots, j_{k-s-1}\}$. Note that L is a closed subset of K . Let us denote by R the restriction operator $C(K) \rightarrow C(L)$. Note that for distinct $i, j > n_0$ we have $L \cap \mathcal{X}_{i, j} \neq \emptyset$ by the induction hypothesis. It follows that the functions $Rf(\{i\})$, $i > n_0$, are uniformly bounded and η -separated. Indeed, we have

$$\begin{aligned} \|Rf(\{i\})\| &\leq \|f(\{i\})\| \leq \|f(\{i\}) - f(\emptyset)\| + \|f(\emptyset)\| \\ &\leq Dd_\Delta(\{i\}, \emptyset) + \|f(\emptyset)\| \\ &\leq D + \|f(\emptyset)\|, \end{aligned}$$

and for distinct $i, j > n_0$ we can pick $\beta \in L \cap \mathcal{X}_{i, j}$ and obtain

$$\|Rf(\{i\}) - Rf(\{j\})\| \geq |f(\{i\})(\beta) - f(\{j\})(\beta)| \geq \eta.$$


We deduce that $C(L)$ must be infinite-dimensional, and hence L must be infinite. Since every infinite compact space has an accumulation point, we have

$$K^{(s+1)} \cap \mathcal{X}_{i_1, j_1} \cap \mathcal{X}_{i_2, j_2} \cap \dots \cap \mathcal{X}_{i_{k-s-1}, j_{k-s-1}} \neq \emptyset,$$

as required. □

4 Applications and open problems

Let (M, d_M) denote an arbitrary separable metric space. Let D_α be an upper bound on $c_{C([0, \omega^\alpha])}(M)$, and consider the following self-explanatory diagram:



Whereas the best distortion achievable in the two extreme cases is completely understood, essentially no estimates for the values of the parameters $c_{C([0, \omega^\alpha])}(C)$ have been hitherto known for C being any class among $\mathcal{M}, \mathcal{SB}, \mathcal{COT}, \mathcal{TYP}, \mathcal{SR}$ (besides the upper bound 2 which follows from the Kalton–Lancien embedding result [8]). It is worth noting that since any $C(K)$ -space (for K countable) is c_0 -saturated, it cannot be a linearly isometrically universal space for the class of separable Banach spaces. Moreover, it cannot be an isometrically universal space either since Godefroy

and Kalton [7] proved that if a separable Banach space X embeds isometrically into a Banach space Y , then Y contains a linear isometric copy of X . Our study of stratifications of the Hamming cube (Theorem 6) yields nontrivial lower bounds for the first time.

Corollary 7 *Let $\mathcal{C} \in \{\mathcal{M}, \mathcal{SB}, \mathcal{COT}, \mathcal{TYP}, \mathcal{SR}\}$, and let $k \in \mathbb{N}$. Then*

$$\frac{k+1}{k} \leq c_{C([0, \omega^k])}(\mathcal{C}) \leq 2.$$

Proof We first remark that the upper bound for all k is the result of Kalton and Lancien [8], and the lower bound for $k = 1$ is due to Aharoni [1]. We now consider the lower bound for $k \geq 2$.

Set $K = [0, \omega^k]$, and note that $i_{CB}(K) = k + 1$. It follows from Theorem 6 that $c_{C(K)}(\Delta_{k+1}) \geq (k + 1)/k$. Given $\varepsilon > 0$, choose p with $1 < p < \infty$ such that the function $f: \Delta_{k+1} \rightarrow \ell_p$ defined by $f(\sigma) = \sum_{i \in \sigma} e_i$ is a $(1 + \varepsilon)$ -isometric embedding. It follows that

$$c_{C(K)}(\Delta_{k+1}) \leq (1 + \varepsilon)c_{C(K)}(\ell_p).$$

Since ℓ_p belongs to the class \mathcal{C} , we have

$$c_{C(K)}(\Delta_{k+1}) \leq (1 + \varepsilon)c_{C(K)}(\mathcal{C}),$$

and the result is proved. □

The following corollary is an easy consequence of Theorem 6 and the fact that $(\Delta_k)_{k \geq 1}$ is a stratification of H_∞ .

Corollary 8 *Let K be a countable compact metric space. If*

$$H_\infty \underset{\text{a.i.}}{\hookrightarrow} C(K),$$

then $i_{CB}(K) \geq \omega + 1$. In particular, if $C(K)$ is an almost isometrically universal space for the class $\mathcal{C} \in \{\mathcal{M}, \mathcal{SB}, \mathcal{COT}, \mathcal{TYP}, \mathcal{SR}\}$, then $i_{CB}(K) \geq \omega + 1$.

Proof It follows from Theorem 6 that $i_{CB}(K) \geq k + 1$ for every $k < \omega$, and hence $K^{(\omega)} = \bigcap_{k < \omega} K^{(k)} \neq \emptyset$. The result follows from Proposition 5(2). □

Remark Procházka and Sánchez-González [11] using the technique of Section 3 exhibited a countable nonproper metric space which does not admit an embedding with distortion less than 2 into any $C(K)$ -space with K countable. Therefore for such compact spaces K we have $c_{C(K)}(\mathcal{M}) = c_{C(K)}(\mathcal{SB}) = 2$, and hence $C(K)$ cannot be an almost isometrically universal space for the classes \mathcal{M} or \mathcal{SB} .

The following theorem, of independent interest, can also be used to prove the second part of Corollary 8 in combination with either Aharoni’s original lower bound involving ℓ_1 or Corollary 7.

Theorem 9 *If $\ell_1 \xrightarrow[\text{a.i.}]{} C(K)$ then $\ell_1 \xrightarrow[\text{a.i.}]{} C(K^{(\alpha)})$ for all ordinals $\alpha < \omega$.*

Proof It is sufficient to show that

$$\ell_1 \xrightarrow[\text{a.i.}]{} C(K) \text{ implies } \ell_1 \xrightarrow[\text{a.i.}]{} C(K').$$

Fix $\varepsilon \in (0, \frac{1}{2})$ and let $f: \ell_1 \rightarrow C(K)$ be a function satisfying

$$\frac{\|x - y\|_1}{1 + \varepsilon} \leq \|f(x) - f(y)\|_\infty \leq \|x - y\|_1.$$

Define $g: \ell_1 \rightarrow C(K')$ by letting $g(x)$ be the restriction of $f(x)$ to K' ($x \in \ell_1$). We are going to show that $\text{dist}(g) \leq (1 + \varepsilon)/(1 - 2\varepsilon)$, which then completes the proof.

Fix distinct vectors $x, y \in \ell_1$ of finite support. Let

$$\delta = \|x - y\|_1 \text{ and } n_0 = \max \text{supp}(x) \cup \text{supp}(y).$$

For distinct integers $i, j > n_0$ we have

$$\|f(x + \delta e_i) - f(y + \delta e_j)\|_\infty \geq \frac{3\delta}{1 + \varepsilon}.$$

Hence there exists $\beta \in K$ such that

$$(7) \quad |f(x + \delta e_i)(\beta) - f(y + \delta e_j)(\beta)| \geq \frac{3\delta}{1 + \varepsilon}.$$

We next observe that if (7) holds, then we also have

$$(8) \quad |f(x + \delta e_i)(\beta) - f(x + \delta e_j)(\beta)| \geq \frac{(2 - \varepsilon)\delta}{1 + \varepsilon}$$

and

$$(9) \quad |f(x)(\beta) - f(y)(\beta)| \geq \frac{(1 - 2\varepsilon)\delta}{1 + \varepsilon} = \frac{(1 - 2\varepsilon)\|x - y\|_1}{1 + \varepsilon}.$$

Now let

$$L = \{\beta \in K : \exists \text{ distinct } i, j > n_0 \text{ satisfying inequality (7)}\}.$$

For $z \in \ell_1$ let $f_L(z)$ denote the restriction of $f(z)$ to L . By (8), the sequence $(f_L(x + \delta e_i))_{i > n_0}$ in $C(L)$ is bounded and $(2 - \varepsilon)\delta/(1 + \varepsilon)$ -separated. It follows that

L is infinite, and so $L \cap K' \neq \emptyset$. By (9), for any $\beta \in L \cap K'$ we have

$$|f(x)(\beta) - f(y)(\beta)| \geq \frac{(1-2\varepsilon)\|x-y\|_1}{1+\varepsilon}.$$

Thus

$$\|g(x) - g(y)\| \geq \frac{(1-2\varepsilon)\|x-y\|_1}{1+\varepsilon},$$

proving that $g: \ell_1 \rightarrow C(K')$ is a bi-Lipschitz embedding with constant $(1+\varepsilon)/(1-2\varepsilon)$, as claimed. \square

We conclude by stating some open problems. In light of the above result, it is natural to ask the following.

Question 1 Does ℓ_1 almost isometrically embed into $C([0, \omega^\omega])$?

Recall that one cannot hope for an isometric embedding because of the aforementioned result of Godefroy and Kalton [7].

Recall also that using the techniques of Theorem 6, Procházka and Sánchez-González [11] constructed a separable metric space M for which $c_{C(K)}(M) = 2$ for any (infinite) countable compact space K . However, it is not clear whether their example embeds into ℓ_1 isometrically (or with distortion less than 2). Indeed, it is not known if their example isometrically embeds into any Banach space which is not already universal for \mathcal{SB} . So the following open problems seem to be of interest.

Question 2 Is there some non-trivial class \mathcal{C} of Banach spaces and a countable compact space K such that $C(K)$ is almost isometrically universal for the class \mathcal{C} ?

The above question is deliberately vague. Examples we have in mind for non-trivial classes include \mathcal{TOP} , \mathcal{COT} and \mathcal{SR} . We conclude with a more specific quantitative question.

Question 3 Let $\alpha \in [2, \omega_1)$. What is the exact value of $c_{C([0, \omega^\alpha])}(\mathcal{C})$ for $\mathcal{C} \in \{\mathcal{TOP}, \mathcal{COT}, \mathcal{SR}\}$?

References

- [1] **I Aharoni**, *Every separable metric space is Lipschitz equivalent to a subset of c_0^+* , Israel J. Math. 19 (1974) 284–291 MR0511661
- [2] **F Albiac, N J Kalton**, *Topics in Banach space theory*, Graduate Texts in Mathematics 233, Springer, New York (2006) MR2192298

- [3] **P Assouad**, *Remarques sur un article de Israel Aharoni sur les prolongements lipschitziens dans c_0* (Israel J. Math. 19 (1974) 284–291), Israel J. Math. 31 (1978) 97–100 MR0511662
- [4] **S Banach, S Mazur**, *Zur Theorie der linearen Dimension*, Stud. Math. 4 (1933) 100–112
- [5] **J Bourgain**, *The Szlenk index and operators on $C(K)$ -spaces*, Bull. Soc. Math. Belg. Sér. B 31 (1979) 87–117 MR592664
- [6] **MM Fréchet**, *Sur quelques points du calcul fonctionnel*, Rend. Circ. Mat Palermo Math. 22 (1906) 1–72
- [7] **G Godefroy, NJ Kalton**, *Lipschitz-free Banach spaces*, Studia Math. 159 (2003) 121–141 MR2030906
- [8] **NJ Kalton, G Lancien**, *Best constants for Lipschitz embeddings of metric spaces into c_0* , Fund. Math. 199 (2008) 249–272 MR2395263
- [9] **E Odell**, *Ordinal indices in Banach spaces*, Extracta Math. 19 (2004) 93–125 MR2072539
- [10] **J Pelant**, *Embeddings into c_0* , Topology Appl. 57 (1994) 259–269 MR1278027
- [11] **A Procházka, L Sánchez-González**, *Low distortion embeddings into Asplund Banach spaces*, preprint (2014) arXiv:1311.4584

FB: Department of Mathematics, Texas A&M University
College Station, TX 77843, USA

FB: Institut de Mathématiques Jussieu-Paris Rive Gauche, Université Pierre et Marie Curie
75005 Paris, France

DF: Department of Mathematics and Computer Science, Saint Louis University
220 N. Grand Blvd., St. Louis, MO 63103, USA

TS: Department of Mathematics, Texas A&M University
College Station, TX 77843, USA

TS: Faculty of Electrical Engineering, Czech Technical University in Prague
Zikova 4, 166 27 Prague, Czech Republic

AZ: Peterhouse, University of Cambridge
Cambridge CB2 1RD, UK

florent@math.tamu.edu, dfreema7@slu.edu, schlump@math.tamu.edu,
a.zsak@dpms.cam.ac.uk

Proposed: Bruce Kleiner
Seconded: Tobias H Colding, Dmitri Burago

Received: 21 March 2014
Revised: 31 May 2015