

Cross-effects and the classification of Taylor towers

GREGORY ARONE

MICHAEL CHING

Let F be a homotopy functor with values in the category of spectra. We show that partially stabilized cross-effects of F have an action of a certain operad. For functors from based spaces to spectra, it is the Koszul dual of the little discs operad. For functors from spectra to spectra it is a desuspension of the commutative operad. It follows that the Goodwillie derivatives of F are a right module over a certain “pro-operad”. For functors from spaces to spectra, the pro-operad is a resolution of the topological Lie operad. For functors from spectra to spectra, it is a resolution of the trivial operad. We show that the Taylor tower of the functor F can be reconstructed from this structure on the derivatives.

55P65, 55P48; 18D50

Let \mathcal{C} and \mathcal{D} each be either the category of based topological spaces, or the category of spectra. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a homotopy functor. Goodwillie’s homotopy calculus provides a systematic way to decompose F into homogeneous pieces which are classified by certain spectra with Σ_n actions, denoted $\partial_n F$ and, by analogy with ordinary calculus, called the *derivatives* or *Taylor coefficients* of F .

A key problem in the homotopy calculus is to describe all the relevant structure on the symmetric sequence $\partial_* F$, and to reconstruct the original functor F (or at least its Taylor tower) from this structure. In [2] we gave a general description of this structure. In this paper we give an alternative description of the structure on $\partial_* F$ in cases when F takes values in the category of spectra. We interpret this structure as that of a “module over a pro-operad”. The structure arises from actions of certain operads on the partially stabilized cross-effects of F . Thus we connect the module structure on $\partial_* F$ with Goodwillie’s presentation of $\partial_* F$ as stabilized cross-effects of F .

Let us review briefly our previous results on the subject. Let $I_{\mathcal{T}op_*}$ be the identity functor on the category of based spaces. It was shown by the second author in [6] that $\partial_* I_{\mathcal{T}op_*}$ has an operad structure. In fact, $\partial_* I_{\mathcal{T}op_*}$ is a topological realization of the classical Lie operad, and we refer to $\partial_* I_{\mathcal{T}op_*}$ informally as the topological Lie operad. In [1] we showed that for a functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$, the symmetric sequence $\partial_* F$ is a right module over $\partial_* I_{\mathcal{T}op_*}$. We also remarked in [1] that the module structure was not sufficient for recovering the Taylor tower of F , and therefore there had to be

additional structure, that still had to be described. In [2] we gave a description of this extra information. Specifically, we constructed a comonad \mathbf{C} , defined on the category of right $\partial_* I_{\mathcal{Top}_*}$ -modules, which acts on the derivatives of a homotopy functor F . We showed that the Taylor tower of F can be recovered from this action.

The methods of [2] were quite general (applying to functors between any combinations of the categories of based spaces and spectra). The comonad \mathbf{C} has the form $\partial_* \Phi$, where ∂_* is Goodwillie differentiation, and Φ is a functor right adjoint to ∂_* . In this paper we provide a more concrete description of this comonad, for functors that take values in \mathcal{Sp} .

To understand our results, recall that the Goodwillie derivatives of F can be recovered from the cross-effects of F via “multilinearization”. Explicitly, for a functor $F: \mathcal{Top}_*^f \rightarrow \mathcal{Sp}$ we have

$$\partial_n F \simeq \operatorname{hocolim}_{L \rightarrow \infty} \Sigma^{-nL} \operatorname{cr}_n F(S^L, \dots, S^L),$$

where $\operatorname{cr}_n F$ denotes the n^{th} cross-effect of F and S^L is the topological L -sphere. We show in Proposition 3.54 that, for fixed L , the symmetric sequence of partially linearized cross-effects

$$d^L[F] := \Sigma^{-*L} \operatorname{cr}_* F(S^L, \dots, S^L)$$

naturally forms a right module over an operad KE_L that is given by the Koszul dual of the stable little L -discs operad E_L . (Strictly speaking, this is true when F is polynomial, or when F is analytic and L is sufficiently large.)

Our construction of the right modules $d^L[F]$ is quite complicated: see the sequence of steps listed just before Definition 3.11 for a summary. A key role here is played by a version of Koszul duality that relates comodules over the operad E_L to modules over the Koszul dual operad KE_L . This is described further in Propositions 3.44 and 3.67.

The Koszul dual operads KE_L form an inverse sequence

$$(0.1) \quad \cdots \rightarrow \operatorname{KE}_{L+1} \rightarrow \operatorname{KE}_L \rightarrow \cdots \rightarrow \operatorname{KE}_2 \rightarrow \operatorname{KE}_1 \rightarrow \operatorname{KE}_0.$$

We refer to such an inverse sequence as a *pro-operad*. We denote the above pro-operad by $\operatorname{KE}_\bullet$. Note that the homotopy inverse limit of the sequence $\operatorname{KE}_\bullet$ is KE_∞ , which is equivalent to the topological Lie operad $\partial_* I_{\mathcal{Top}_*}$. Therefore we refer to the sequence $\operatorname{KE}_\bullet$ as the *Lie pro-operad*.

For polynomial functors F , we construct a model for the maps

$$d^L[F] \rightarrow d^{L+1}[F]$$

between the partially linearized cross-effects that respects the operad maps

$$KE_{L+1} \rightarrow KE_L.$$

It follows that the symmetric sequence of derivatives $\partial_* F \simeq d[F] := \text{hocolim}_L d^L[F]$ inherits a limiting structure. We refer to this structure as a *right module over the Lie pro-operad*. This structure includes, but contains strictly more information than, a right module over the topological Lie operad by itself.

More precisely, we define modules over a pro-operad as coalgebras over a certain comonad, constructed as follows. First note that the category of right modules over any operad P can be identified with the category of coalgebras over a certain comonad C_P on the category of symmetric sequences. The sequence (0.1) of operads induces a directed sequence of comonads

$$C_{KE_0} \rightarrow C_{KE_1} \rightarrow C_{KE_2} \rightarrow \dots$$

If we write

$$C_{KE_\bullet} := \text{hocolim}_L C_{KE_L},$$

then C_{KE_\bullet} has a canonical comonad structure, and we define a module over the pro-operad KE_\bullet to be a coalgebra over C_{KE_\bullet} .

Remark 0.2 Our notion of a right module over a pro-operad is modelled on that of a discrete set with a continuous action of a profinite group. In particular, a module over a pro-operad is (for us, in this paper) not an object in the pro-category of symmetric sequences. Rather, it often can be presented as a filtered colimit of symmetric sequences. This is analogous to the fact that a discrete set X with a continuous action of a profinite group G is the union (ie colimit) of sets with actions of finite quotients of G .

Ultimately, we show (Proposition 3.54) that $d[F]$ has a natural KE_\bullet -module structure for all pointed simplicial functors F from \mathcal{Top}_* to \mathcal{Sp} . Our Theorem 3.64 then says that C_{KE_\bullet} is equivalent to the abstract comonad C considered in [2]. It follows that the Taylor tower of a functor F can be recovered from the KE_\bullet -module structure on $d[F]$. We show in Theorem 3.75 that there is an equivalence between the homotopy categories of polynomial functors and bounded KE_\bullet -modules. We also show (Theorem 3.92) that there is an equivalence between the homotopy categories of analytic functors (up to Taylor tower equivalence) and KE_\bullet -modules satisfying suitable connectivity conditions.

There are other approaches to the classification of polynomial functors from based spaces to spectra, most notably by Dwyer and Rezk (unpublished) in terms of functors on the category of finite sets and surjections. Polynomial functors from based spaces

to spectra are also equivalent to their left Kan extensions from the full subcategory of finite pointed sets. In Theorem 3.82 we describe how our approach is related to these others.

The story for a functor $F: \mathcal{S}p \rightarrow \mathcal{S}p$ is quite analogous. The derivatives $\partial_* F$ are again the homotopy colimit, over L , of partially linearized cross-effects

$$d^L[F] := \Sigma^{-*L} \text{cr}_* F(\Sigma^\infty S^L, \dots, \Sigma^\infty S^L)$$

and again these cross-effects, for fixed L , form a right module over a certain operad S^{-L} . The operad S^{-L} can be thought of as the L -fold desuspension of the commutative operad, and all of its terms are sphere spectra of varying dimensions. Again there is an inverse sequence

$$(0.3) \quad \dots \rightarrow S^{-(L+1)} \rightarrow S^{-L} \rightarrow \dots \rightarrow S^{-2} \rightarrow S^{-1} \rightarrow S^0 \simeq \text{Com}.$$

We refer to this sequence as the *sphere pro-operad*. Note that the homotopy inverse limit of this sequence is equivalent to the trivial operad.

We show that the derivatives $\partial_*(F) \simeq d[F] := \text{hocolim}_L d^L[F]$ form a coalgebra over the corresponding comonad

$$\mathbf{C}_{S_\bullet} := \text{hocolim}_L \mathbf{C}_{S^{-L}}.$$

We show in Theorem 4.27 that the comonad \mathbf{C}_{S_\bullet} is equivalent to that constructed in [2]. The Taylor tower of $F: \mathcal{S}p \rightarrow \mathcal{S}p$ can then be recovered from the \mathbf{C}_{S_\bullet} -coalgebra $d[F]$ and there is an equivalence between the homotopy categories of polynomial functors and bounded coalgebras, as well as between analytic functors and suitably connected coalgebras.

There is some connection and overlap between our results on functors from $\mathcal{S}p$ to $\mathcal{S}p$ and work of Randy McCarthy [16] which involves a right Com -module structure on the symmetric sequence $\text{cr}_* F(\Sigma^\infty S^L, \dots, \Sigma^\infty S^L)$. The S^{-L} -module structure that we use is equivalent to a desuspension of McCarthy’s structure. Ultimately, McCarthy classifies n -excisive endofunctors of $\mathcal{S}p$ in terms of structure on the spectrum $F(\bigvee_n S^0)$. Namely, he shows that if F is n -excisive, then the spectrum $F(\bigvee_n S^0)$ is a module over a certain ring spectrum, and that the functor F can be recovered from this module structure.

Since $F(\bigvee_n S^0)$ is equivalent to a wedge sum of copies of cross-effects $\text{cr}_1 F, \dots, \text{cr}_n F$ evaluated at S^0 , it seems likely that McCarthy’s result can be rephrased in terms of structure on the sequence of cross-effects of F . In this form, it would be analogous to the result of Dwyer and Rezk that classifies polynomial functors from $\mathcal{T}op_*$ to $\mathcal{S}p$ in terms of structure on the cross-effects.

As far as we know, until now all work on classifying polynomial functors approached the problem via describing the structure on the sequence of cross-effects of a functor. Our approach focuses instead on the derivatives of a functor. For functors from $\mathcal{T}op_*$ to $\mathcal{S}p$ we have a good understanding of the relationship between the structure on the cross-effects and the structure on derivatives: it is given by a form of Koszul duality between comodules over the commutative operad, and divided power modules over the Lie operad. It would be interesting to find a similar connection in the $\mathcal{S}p$ to $\mathcal{S}p$ case and for other classes of functors.

Some open problems, possible directions for future research

Do something similar for space-valued functors Consider, for example, functors $F: \mathcal{T}op_* \rightarrow \mathcal{T}op_*$. By results of [1], $\partial_* F$ is a bimodule over $\partial_* I_{\mathcal{T}op_*}$, and we seek to describe the additional structure on $\partial_* F$. One answer is given in [2], which says that $\partial_* I_{\mathcal{T}op_*}$ is a coalgebra over a certain comonad in the category of $\partial_* I_{\mathcal{T}op_*}$ -bimodules. But it seems desirable to have a more concrete description of the additional structure, perhaps in the form of a compatibility condition between the left and right module structures on $\partial_* F$. At the moment we are unable to provide such a description. We also do not know how to relate the module structure on $\partial_* F$ with the view of $\partial_* F$ as the stabilized cross-effects of F .

Give an explicit description of KE_L , and use it to understand the connection between the two classes of functors Let $F: \mathcal{T}op_* \rightarrow \mathcal{S}p$ be a functor. Then $F\Sigma^\infty$ is a functor from $\mathcal{S}p$ to $\mathcal{S}p$. There is an equivalence of symmetric sequences $\partial_* F \simeq \partial_* F\Sigma^\infty$. This means that there should be a forgetful functor from modules over the sphere pro-operad to modules over the Lie pro-operad. Presumably, the forgetful functor is induced by maps of operads $KE_L \rightarrow S^{-L}$. It does not seem obvious how to construct such a map of operads. This is partly because we do not have a geometric model for KE_L .

This question ties in with another conjecture about the operad KE_L . Namely, it is conjectured that there is an equivalence of operads $KE_L \simeq \Sigma^{-L}E_L$. Corresponding results on the level of homology are due to Getzler and Jones [11], and on the chain level are due to Fresse [10]. Assuming this conjecture, notice that the obvious map of operads $E_L \rightarrow \text{Com}$ induces a map $KE_L \simeq \Sigma^{-L}E_L \rightarrow S^{-L}$. We speculate that the KE_\bullet -module structure on $\partial_* F\Sigma^\infty$ is equivalent to the pullback along this hypothetical map of operads.

If this speculation is correct, we would obtain a new proof of the well-known fact that the Taylor tower of the functor $\Sigma^\infty\Omega^\infty\Sigma^\infty$ splits as a product of its layers. Indeed,

we show in Example 4.33 that the action of the pro-sphere operad on $\partial_* \Sigma^\infty \Omega^\infty$ is pulled back from an action by the single operad $\text{Com} = S^0$. It would follow that the KE_\bullet -module structure on $\partial_* \Sigma^\infty \Omega^\infty \Sigma^\infty$ is pulled back from a KE_0 -module structure. But KE_0 is the trivial operad, so it would follow that the Taylor tower of $\Sigma^\infty \Omega^\infty \Sigma^\infty$ splits.

Describe the chain rule for functors from $\mathcal{S}p$ to $\mathcal{S}p$ on the level of modules over the sphere pro-operad Let F, G be functors from spectra to spectra. By the results of [7], there is an equivalence of symmetric sequences, where the right-hand side is the composition product of $\partial_* F$ and $\partial_* G$:

$$\partial_*(FG) \simeq \partial_* F \circ \partial_* G.$$

By our present results, $\partial_* F$, $\partial_* G$ and $\partial_*(FG)$ are modules over the sphere pro-operad. This suggests that there should be a composition product in the category of such modules, refining the composition product of symmetric sequences. Moreover, we speculate that the product can be made associative (or at least A_∞ -associative), and that the chain rule can be made associative as well.

Outline

In Section 1 we review basic facts about Taylor towers and derivatives, and also about symmetric sequences, operads and modules. We also review our construction of the homotopy category of coalgebras over a comonad. The construction that we use here is a slight variation of the one in [2]. In Section 2 we introduce pro-operads, and we define the category of modules over a pro-operad as the category of coalgebras of an associated comonad. In the long Section 3 we analyze functors from \mathcal{Top}_* to $\mathcal{S}p$. In Sections 4 and 5 we do the same from functors from $\mathcal{S}p$ to $\mathcal{S}p$.

Acknowledgements

Conversations with Bill Dwyer contributed considerably to the Koszul duality between E_L -comodules and KE_L -modules that we use in this paper. The first author was supported by National Science Foundation grant DMS-0968221 and the second author by National Science Foundation grant DMS-1144149.

1 Background

1.1 Taylor towers and derivatives

Definition 1.1 (Categories) Let \mathcal{Top}_* be the category of based topological spaces and let $\mathcal{S}p$ be the category of S -modules of Elmendorf, Kriz, Mandell and May [9].

We refer to the objects of $\mathcal{S}p$ as *spectra* to avoid confusion with the other uses of the term “module” in this paper. Each of the categories $\mathcal{T}op_*$ and $\mathcal{S}p$ is enriched in simplicial sets and we write $\text{Hom}_{\mathcal{C}}(X, Y)$ for the simplicial set of maps from X to Y in the category \mathcal{C} . The category $\mathcal{S}p$ is closed symmetric monoidal and we write $\text{Map}(X, Y)$ for the internal mapping spectrum in $\mathcal{S}p$.

The suspension spectrum and zeroth space functors give us the standard adjunction:

$$\Sigma^\infty: \mathcal{T}op_* \rightleftarrows \mathcal{S}p : \Omega^\infty$$

Let $\mathcal{T}op_*^f$ denote the full subcategory of $\mathcal{T}op_*$ consisting of finite based cell complexes, and let $\mathcal{S}p^f$ denote the full subcategory of $\mathcal{S}p$ consisting of finite cell spectra (with respect to the usual generating cofibrations for the stable model structure on $\mathcal{S}p$; see [9, VII]).

Definition 1.2 (Functors) Let \mathcal{C} be either $\mathcal{T}op_*$ or $\mathcal{S}p$, and consider a functor $F: \mathcal{C} \rightarrow \mathcal{S}p$. We say that F is a *homotopy functor* if it preserves weak equivalences, is *simplicial* if it induces maps $\text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{S}p}(FX, FY)$ of simplicial sets, is *finitary* if it preserves filtered homotopy colimits, and is *pointed* if $F(*) \cong *$, where $*$ denotes either a one-point space in $\mathcal{T}op_*$, or a trivial spectrum in $\mathcal{S}p$.

Definition 1.3 (Categories of functors) For \mathcal{C} equal to either $\mathcal{T}op_*$ or $\mathcal{S}p$, let $[\mathcal{C}^f, \mathcal{S}p]_*$ denote the category whose objects are the pointed simplicial functors $F: \mathcal{C}^f \rightarrow \mathcal{S}p$, and whose morphisms are the corresponding simplicially enriched natural transformations. Since \mathcal{C}^f is skeletally small, $[\mathcal{C}^f, \mathcal{S}p]_*$ is a locally small category. Also note that any simplicial functor preserves simplicial homotopy equivalences. Since every object of \mathcal{C}^f is both cofibrant and fibrant (in the standard model structure on either $\mathcal{T}op_*$ or $\mathcal{S}p$), it follows that any object in $[\mathcal{C}^f, \mathcal{S}p]_*$ also preserves weak equivalences.

We can extend a pointed simplicial functor $F: \mathcal{C}^f \rightarrow \mathcal{S}p$ to all of \mathcal{C} by (enriched) homotopy left Kan extension along the inclusion $\mathcal{C}^f \rightarrow \mathcal{C}$. The result of this construction is a reduced finitary homotopy functor. Moreover, any reduced finitary homotopy functor $F: \mathcal{C} \rightarrow \mathcal{S}p$ arises, up to natural equivalence, in this way. We therefore view $[\mathcal{C}^f, \mathcal{S}p]_*$ as a model for the collection of reduced finitary homotopy functors $\mathcal{C} \rightarrow \mathcal{S}p$.

For a functor $F \in [\mathcal{C}^f, \mathcal{S}p]_*$ we say that F is *n-excisive* if it takes a strongly cocartesian $(n + 1)$ -cube in \mathcal{C}^f to a cartesian cube in $\mathcal{S}p$. We say F is *polynomial* if it is *n-excisive* for some n .

The category $[\mathcal{C}^f, \mathcal{S}p]_*$ has a projective model structure in which weak equivalences and fibrations are detected objectwise. We denote the associated homotopy category

as $[\mathcal{C}^f, \mathcal{S}p]_*^h$. This homotopy category has subcategories given by restricting to functors satisfying various conditions; for example, we have the homotopy categories $[\mathcal{C}^f, \mathcal{S}p]_{*,n\text{-exc}}^h$ of n -excisive functors, and $[\mathcal{C}^f, \mathcal{S}p]_{*,\text{poly}}^h$ of all polynomial functors.

Definition 1.4 (Taylor tower and derivatives) For a pointed simplicial functor $F: \mathcal{C}^f \rightarrow \mathcal{S}p$ there is a *Taylor tower* of pointed simplicial functors $P_n F: \mathcal{C}^f \rightarrow \mathcal{S}p$ of the form

$$F \rightarrow \cdots \rightarrow P_n F \rightarrow P_{n-1} F \rightarrow \cdots \rightarrow P_2 F \rightarrow P_1 F \rightarrow P_0 F = *,$$

where $P_n F$ is n -excisive in the sense of [12]. The *layers* of the Taylor tower are the functors $D_n F: \mathcal{C}^f \rightarrow \mathcal{S}p$ given by the objectwise homotopy fibres

$$D_n F := \text{hofib}(P_n F \rightarrow P_{n-1} F).$$

Goodwillie showed that for each $n \geq 1$ there is a spectrum $\partial_n F$ with Σ_n -action such that

$$D_n F(X) \simeq (\partial_n F \wedge X^{\wedge n})_{h\Sigma_n}$$

for $X \in \mathcal{C}^f$. We refer to $\partial_n F$ as the n^{th} derivative of F .

Definition 1.5 (Cross-effects and co-cross-effects) The n^{th} derivative of a functor can be calculated using cross-effects. The n^{th} *cross-effect* of $F: \mathcal{C} \rightarrow \mathcal{S}p$ is the functor of n variables $\text{cr}_n F: \mathcal{C}^{\times n} \rightarrow \mathcal{S}p$ given by

$$\text{cr}_n F(X_1, \dots, X_n) := \text{thofib}_{S \subset \underline{n}} \left\{ F \left(\bigvee_{i \notin S} X_i \right) \right\}.$$

This is the iterated (or total) homotopy fibre of an n -cube formed by applying F to the wedge sums of subsets of X_1, \dots, X_n , where the morphisms in the cube are given by the relevant collapse maps $X_i \rightarrow *$.

For spectrum-valued functors, the cross-effect can also be calculated by taking a total homotopy cofibre. The n^{th} *co-cross-effect* of $F: \mathcal{C}^f \rightarrow \mathcal{S}p$ is the functor $\text{cr}^n F: \mathcal{C}^{\times n} \rightarrow \mathcal{S}p$ given by

$$\text{cr}^n F(X_1, \dots, X_n) := \text{thocofib}_{S \subset \underline{n}} \left\{ F \left(\bigvee_{i \in S} X_i \right) \right\}.$$

The morphisms in this n -cube are given by the inclusion maps $* \rightarrow X_i$. For any $F: \mathcal{C} \rightarrow \mathcal{S}p$, there is a natural equivalence

$$\text{cr}_n F(X_1, \dots, X_n) \simeq \text{cr}^n F(X_1, \dots, X_n).$$

In particular, taking cross-effects commutes with both homotopy limits and homotopy colimits.

One of Goodwillie’s main results is that the derivatives of a functor can be recovered by multilinearizing the cross-effects. Specifically, [13, Theorem 6.1] implies that for a functor $F: \mathcal{C} \rightarrow \mathcal{S}p$, we have a natural equivalence

$$\partial_n F \simeq P_{(1,\dots,1)}(\text{cr}_n F)(S^0, \dots, S^0),$$

where the right-hand side is the multilinearization of the n^{th} cross-effect of F . The multilinearization $P_{(1,\dots,1)}(\text{cr}_n F)$ is the homotopy colimit of maps

$$\text{cr}_n F \rightarrow T_{(1,\dots,1)}(\text{cr}_n F) \rightarrow T_{(1,\dots,1)}^2(\text{cr}_n F) \rightarrow T_{(1,\dots,1)}^3 \rightarrow \dots$$

where, for a functor $G: \mathcal{C}^n \rightarrow \mathcal{S}p$ that is reduced in each variable, there are natural equivalences

$$T_{(1,\dots,1)}G(X_1, \dots, X_n) \simeq \Sigma^{-n}G(\Sigma X_1, \dots, \Sigma X_n).$$

Thus we can express Goodwillie’s result as an equivalence

$$(1.6) \quad \partial_n F \simeq \text{hocolim}_L \Sigma^{-nL} \text{cr}_n F(S^L, \dots, S^L).$$

The maps in this homotopy colimit take the form

$$(1.7) \quad \Sigma^{-nL} \text{cr}_n F(S^L, \dots, S^L) \rightarrow \Sigma^{-n(L+1)} \text{cr}_n F(S^{L+1}, \dots, S^{L+1}).$$

In building our models for the derivatives of F , we need models for these maps.

Definition 1.8 When the functor $H: \mathcal{C} \rightarrow \mathcal{S}p$ is pointed and simplicial, it determines natural maps

$$t_X: X \wedge H(Y) \rightarrow H(X \wedge Y)$$

for a based simplicial set X and $Y \in \mathcal{C}$. We refer to this as the *tensoring map* for H .

Lemma 1.9 Let $G: \mathcal{C}^n \rightarrow \mathcal{S}p$ be a model for the n^{th} cross-effect of a functor $F: \mathcal{C} \rightarrow \mathcal{S}p$ that is pointed simplicial in each variable. Then the map (1.7) is, up to equivalence, given by the tensoring map

$$\Sigma^{-n(L+1)} \Sigma^n G(S^L, \dots, S^L) \xrightarrow{t_{S^1 \circ \dots \circ S^1}} \Sigma^{-n(L+1)} G(S^{L+1}, \dots, S^{L+1}).$$

Proof It is sufficient to do the case $\mathcal{C} = \mathcal{T}op_*$ since the result for $G: \mathcal{S}p^n \rightarrow \mathcal{S}p$ follows from that for $G(\Sigma^\infty -, \dots, \Sigma^\infty -): \mathcal{T}op_*^n \rightarrow \mathcal{S}p$. We illustrate with the case $n = 1$. In this case, we have the following diagram. For $X \in \mathcal{T}op_*$, with CX denoting

the (reduced) cone on X and $\Omega X = \text{holim}(* \rightarrow X \leftarrow *) \cong \text{Hom}_{\mathcal{T}op_*}(S^1, X)$:

$$\begin{array}{ccc}
 & G(X) & \\
 \swarrow & & \searrow \\
 G(\Omega \Sigma X) & \xrightarrow{\sim} & G(\text{holim}(CX \rightarrow \Sigma X \leftarrow CX)) \\
 \downarrow & & \downarrow \\
 \Omega G(\Sigma X) & \xrightarrow{\sim} & \text{holim}(G(CX) \rightarrow G(\Sigma X) \leftarrow G(CX))
 \end{array}$$

The vertical maps come from the fact that G is pointed simplicial, and the bottom square commutes by naturality of that enrichment. The composite of the right-hand two maps is the canonical map $G(X) \rightarrow T_1 G(X)$, and the composite of the left-hand two is adjoint to the tensoring map t_{S^1} for G . It is sufficient then to show that the top triangle commutes up to homotopy. Since G is simplicial, it is therefore sufficient to show that the underlying triangle of spaces

$$\begin{array}{ccc}
 & X & \\
 \swarrow & & \searrow \\
 \Omega \Sigma X & \xrightarrow{\sim} & \text{holim}(CX \rightarrow \Sigma X \leftarrow CX)
 \end{array}$$

commutes up to homotopy. A point in this homotopy limit consists of a path $\gamma: [0, 1] \rightarrow \Sigma X$ from a point in one cone to a point in the other cone. Writing

$$\Sigma X = [0, 2]/\{0 \sim 2\} \wedge X$$

with cones $[0, 1] \wedge X$ and $[1, 2] \wedge X$, where 0 and 2 are treated as the basepoints, the required homotopy

$$H: X \times [0, 1] \rightarrow \text{holim}(CX \rightarrow \Sigma X \leftarrow CX)$$

is given by

$$(x, t) \mapsto (s \mapsto (t + 2s(1 - t), x)). \quad \square$$

Definition 1.10 We refer to the terms $\Sigma^{-nL} \text{cr}_n F(S^L, \dots, S^L)$ in the homotopy colimit (1.6) as the *partially stabilized cross-effects* of the functor F .

1.2 Symmetric sequences, operads, modules and comodules

Definition 1.11 (Symmetric sequences) We write $[\Sigma, \mathcal{S}p]$ for the category of *symmetric sequences* in $\mathcal{S}p$, that is, the category of functors $\Sigma \rightarrow \mathcal{S}p$, where Σ is the category of nonempty finite sets and bijections. Given $A \in [\Sigma, \mathcal{S}p]$ we often write

$A(n) := A(\underline{n})$, where $\underline{n} = \{1, \dots, n\}$ and consider A as the sequence of spectra $A(n)$, for $n \geq 1$, each with an action of Σ_n .

We also define a *bisymmetric sequence* to be a functor $B: \Sigma \times \Sigma \rightarrow \mathcal{S}p$, so that B consists of spectra $B(m, n)$ for $m, n \geq 1$ with commuting actions of Σ_m and Σ_n .

The category $[\Sigma, \mathcal{S}p]$ has a cofibrantly generated model structure in which weak equivalences and fibrations are detected termwise. A symmetric sequence is Σ -*cofibrant* if it is cofibrant in this model structure. The category $[\Sigma, \mathcal{S}p]$ is also enriched in $\mathcal{S}p$: the spectrum of maps between two symmetric sequences A and A' is given by

$$\text{Map}_\Sigma(A, A') := \prod_n \text{Map}(A(n), A'(n))^{\Sigma_n}.$$

The model structure respects this enrichment, making $[\Sigma, \mathcal{S}p]$ into a $\mathcal{S}p$ -enriched model category.

A symmetric sequence A is *n-truncated* if $A(m) = *$ for all $m > n$. We say that A is *bounded* if it is *n-truncated* for some n . For any symmetric sequence A we have its *n-truncation* $A_{\leq n}$, which is the symmetric sequence given by

$$A_{\leq n}(I) := \begin{cases} A(I) & \text{if } |I| \leq n, \\ * & \text{otherwise.} \end{cases}$$

Associated to A is its *truncation sequence* consisting of maps of symmetric sequences

$$A \rightarrow \dots \rightarrow A_{\leq n} \rightarrow A_{\leq(n-1)} \rightarrow \dots \rightarrow A_{\leq 1}.$$

Definition 1.12 (Operads of spectra) An *operad of spectra* is a monoid in the category $[\Sigma, \mathcal{S}p]$ with respect to the composition product of symmetric sequences (see, for example, [6, 2.11]). Explicitly, an operad P consists of a symmetric sequence together with

- a composition map

$$P_\alpha: P(k) \wedge \bigwedge_{j=1}^k P(n_j) \rightarrow P(n)$$

for each surjection $\alpha: \underline{n} \twoheadrightarrow \underline{k}$ of nonempty finite sets, where $n_j := |\alpha^{-1}(j)|$, and

- a unit map $\eta: S \rightarrow P(1)$,

that satisfy associativity, unitality and equivariance conditions.

An operad of spectra is *reduced* if the unit map $\eta: S \rightarrow P(1)$ is an isomorphism. We consider only reduced operads in this paper.

There is a cofibrantly generated model structure on the category of reduced operads of spectra, described in [1, Appendix], in which weak equivalences and fibrations are detected on the underlying symmetric sequences, ie termwise. An operad is Σ -*cofibrant*, *n-truncated* or *bounded* if its underlying symmetric sequence has the corresponding property. The truncation sequence associated to an operad is a sequence of operads.

Note that by design our operads do not include a term $P(0)$ so that they describe only non-unital structures.

Definition 1.13 (Modules over operads of spectra) Given an operad P of spectra, a (*right*) P -*module* consists of a symmetric sequence M together with a right action of P with respect to the composition product. Explicitly, such an M has a structure map

$$M_\alpha: M(k) \wedge \bigwedge_{j=1}^k P(n_j) \rightarrow M(n)$$

for each surjection $\alpha: \underline{n} \twoheadrightarrow \underline{k}$, which satisfy appropriate conditions.

Since we only consider right modules over operads in this paper, we refer to these just as P -modules and write $\text{Mod}(P)$ for the category of P -modules (whose morphisms are maps of the underlying symmetric sequences that commute with the structure maps). The category $\text{Mod}(P)$ has a cofibrantly generated stable model structure that is enriched in $\mathcal{S}p$, in which weak equivalences and fibrations are detected termwise. Note that all homotopy limits and colimits of diagrams of P -modules are also computed termwise. A P -module is Σ -*cofibrant*, *n-truncated* or *bounded* if its underlying symmetric sequence has the corresponding property. The truncation sequence of a P -module is a sequence of P -modules.

In this paper we consider also what we call a “comodule” over an operad. In order to say what this means, we recall that a module can be interpreted as a spectrally enriched functor $\underline{P} \rightarrow \mathcal{S}p$ for a particular category \underline{P} associated to the operad P .

Definition 1.14 For an operad P of spectra we define a $\mathcal{S}p$ -enriched category \underline{P} as follows:

- The objects of \underline{P} are the nonempty finite sets.

- For two nonempty finite sets I and J , the morphism spectrum $\underline{P}(I, J)$ is given by

$$\underline{P}(I, J) := \bigvee_{I \twoheadrightarrow J} \bigwedge_{j \in J} P(I_j),$$

where the wedge product is taken over all surjections $\alpha: I \twoheadrightarrow J$ and we write $I_j := \alpha^{-1}(j)$.

- The composition and identity maps for the category \underline{P} are determined by the operad multiplication and unit maps, respectively.

The category \underline{P} is also known as the PROP associated to the operad P .

Lemma 1.15 *Let P be an operad of spectra. There is an equivalence between the category $\text{Mod}(P)$ of (right) P -modules and the category of $\mathcal{S}p$ -enriched functors $\underline{P}^{\text{op}} \rightarrow \mathcal{S}p$ (with morphisms the $\mathcal{S}p$ -enriched natural transformations).*

Proof A $\mathcal{S}p$ -enriched functor $M: \underline{P}^{\text{op}} \rightarrow \mathcal{S}p$ consists of objects $M(I)$ for each finite set I , and (suitably associative and unital) maps

$$\underline{P}(I, J) \rightarrow \text{Map}(M(J), M(I)).$$

Equivalently, for each surjection $I \twoheadrightarrow J$, there is a map

$$M(J) \wedge \bigwedge_{j \in J} P(I_j) \rightarrow M(I).$$

These maps form precisely the data for a P -module. □

Lemma 1.15 inspires the following definition.

Definition 1.16 Let P be an operad of spectra. A (right) P -comodule is a $\mathcal{S}p$ -enriched functor $\underline{P} \rightarrow \mathcal{S}p$. More explicitly, a P -comodule consists of objects $N(I)$, one for each nonempty finite set I , and maps

$$N(I) \wedge \bigwedge_{j \in J} P(I_j) \rightarrow N(J),$$

one for each surjection $\alpha: I \twoheadrightarrow J$. In particular, a P -comodule N has an underlying symmetric sequence.

We write $\text{Comod}(P)$ for the category of P -comodules, with morphisms given by the $\mathcal{S}p$ -enriched natural transformations. The category $\text{Comod}(P)$ is enriched over $\mathcal{S}p$ and has a cofibrantly generated model structure in which weak equivalences and fibrations

are detected termwise, and homotopy limits and colimits are computed termwise. We say that a P -comodule is Σ -cofibrant, n -truncated or bounded if its underlying symmetric sequence has the corresponding property.

Example 1.17 A right Com-comodule can be identified with a functor $\Omega \rightarrow \mathcal{S}p$, where Ω is the category of nonempty finite sets and surjections. A right Com-module can similarly be identified with a functor $\Omega^{op} \rightarrow \mathcal{S}p$.

The dual definitions of module and comodule permit a natural “coend” construction between a module and comodule. We require a homotopy-invariant version of this construction which we now describe.

Definition 1.18 Let P be a reduced operad of spectra, N a P -comodule and M a P -module. We define the spectrum $N \tilde{\wedge}_P M$ to be the realization of the simplicial spectrum given by

$$[r] \mapsto \bigvee_{n_0, \dots, n_r} N(n_0) \wedge_{\Sigma_{n_0}} \underline{P}(n_0, n_1) \wedge_{\Sigma_{n_1}} \cdots \wedge_{\Sigma_{n_{r-1}}} \underline{P}(n_{r-1}, n_r) \wedge_{\Sigma_{n_r}} M(n_r),$$

where the wedge sum is taken over all sequences of positive integers n_0, \dots, n_r (though only the non-increasing sequences contribute non-trivial terms) and with face maps

- d_0 given by the comodule structure maps $N(n_0) \wedge_{\Sigma_{n_0}} \underline{P}(n_0, n_1) \rightarrow N(n_1)$,
- d_i for $i = 1, \dots, r - 1$ given by the operad composition maps $\underline{P}(n_{i-1}, n_i) \wedge_{\Sigma_{n_i}} \underline{P}(n_i, n_{i+1}) \rightarrow \underline{P}(n_{i-1}, n_{i+1})$,
- d_r given by the module structure maps $\underline{P}(n_{r-1}, n_r) \wedge_{\Sigma_{n_r}} M(n_r) \rightarrow M(n_{r-1})$,

and degeneracy maps

- s_j for $j = 0, \dots, r$ given by the operad unit map $S \cong \underline{P}(n_j, n_j)_{\Sigma_{n_j}}$.

Lemma 1.19 Let P be a Σ -cofibrant operad of spectra and N a Σ -cofibrant P -comodule. Let $M \xrightarrow{\sim} M'$ be a weak equivalence of P -modules. Then the induced map

$$N \tilde{\wedge}_P M \rightarrow N \tilde{\wedge}_P M'$$

is a weak equivalence of spectra. Similarly, let M be a Σ -cofibrant P -module and $N \xrightarrow{\sim} N'$ a weak equivalence of P -comodules. Then the induced map

$$N \tilde{\wedge}_P M \rightarrow N' \tilde{\wedge}_P M$$

is a weak equivalence of spectra.

Proof The simplicial spectra involved here are all proper in the sense of [9, X.2.2]. The conditions imply that the induced maps of simplicial spectra are levelwise weak equivalences. Then [9, X.2.4] implies that the given maps are weak equivalences. \square

Lemma 1.19 tells us that when P is Σ -cofibrant, the homotopy coend $N \widetilde{\wedge}_P M$ has the “correct” homotopy type if either N or M is Σ -cofibrant.

1.3 Coalgebras over comonads and their homotopy theory

In this paper we are concerned with comonads on the category $[\Sigma, \mathcal{S}p]$ of symmetric sequences of spectra.

Recall that a *comonad* C on $[\Sigma, \mathcal{S}p]$ is an endofunctor $C: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$ equipped with natural transformations $v: C \rightarrow CC$ (the *comultiplication*) and $\epsilon: C \rightarrow I_{[\Sigma, \mathcal{S}p]}$ (the *counit*) that make C into a comonoid with respect to composition of functors. For a comonad $C: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$, a *C-coalgebra* is a symmetric sequence A together with a structure map $\theta: A \rightarrow CA$ that forms a coaction of the comonoid C . Morphisms of coalgebras are maps in $[\Sigma, \mathcal{S}p]$ that commute with the structure maps, and we have a category $\text{Coalg}(C)$ of coalgebras over C . We say that a C -coalgebra is *Σ -cofibrant*, *n -truncated* or *bounded* if its underlying symmetric sequence has the corresponding property.

Our main examples of comonads arise from operads.

Definition 1.20 For an operad P of spectra, we define an endofunctor

$$C_P: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$$

by

$$C_P(A)(k) := \prod_n \left[\prod_{n \twoheadrightarrow k} \text{Map}(P(n_1) \wedge \cdots \wedge P(n_k), A(n)) \right]^{\Sigma_n},$$

where Σ_n acts on the set of surjections $\underline{n} \twoheadrightarrow \underline{k}$ by pre-composition, and Σ_k acts by post-composition.

The operad structure on P determines a comonad structure on C_P with the operad composition determining the comultiplication, and the operad unit map determining the counit. Note that the functor C_P is enriched in $\mathcal{S}p$.

Lemma 1.21 *The comonad C_P is that associated to an adjunction*

$$U_P: \text{Mod}(P) \rightleftarrows [\Sigma, \mathcal{S}p] : R_P,$$

where U_P is the forgetful functor. In particular, for any P -module M , the symmetric sequence $U_P(M)$ has a canonical C_P -coalgebra structure.

Proof The right adjoint \mathbf{R}_P is defined by the same formula as \mathbf{C}_P . The P -module structure on $\mathbf{R}_P(A)$ is determined by the operad composition map for P . It is easy to check that \mathbf{R}_P so defined is right adjoint to \mathbf{U}_P , and then $\mathbf{C}_P = \mathbf{U}_P \mathbf{R}_P$ is the associated comonad. The \mathbf{C}_P -coalgebra structure on $\mathbf{U}_P(M)$ is given by the unit map

$$\mathbf{U}_P(M) \rightarrow \mathbf{U}_P \mathbf{R}_P \mathbf{U}_P(M)$$

for the adjunction $(\mathbf{U}_P, \mathbf{R}_P)$. □

Definition 1.22 We refer to the P -module $\mathbf{R}_P(A)$ as the *cofree P -module* associated to the symmetric sequence A .

Lemma 1.23 *The functor $M \mapsto \mathbf{U}_P M$ determines an equivalence of categories*

$$\text{Mod}(P) \simeq \text{Coalg}(\mathbf{C}_P).$$

Proof Since \mathbf{U}_P is right adjoint to the free P -module functor \mathbf{F}_P , the composite $\mathbf{U}_P \mathbf{R}_P$ is right adjoint to the functor monad $\mathbf{U}_P \mathbf{F}_P$ whose algebras are the P -modules. The structure maps for a right P -module correspond precisely under this adjunction to the structure maps for a \mathbf{C}_P -coalgebra structure on the same symmetric sequence. □

Definition 1.24 Suppose that $\phi: P \rightarrow P'$ is a morphism of operads of spectra. There is an induced map of comonads

$$\mathbf{C}_\phi: \mathbf{C}_{P'} \rightarrow \mathbf{C}_P.$$

If M is a $\mathbf{C}_{P'}$ -coalgebra with structure map θ then the composite

$$M \xrightarrow{\theta} \mathbf{C}_{P'} M \xrightarrow{\mathbf{C}_\phi} \mathbf{C}_P M$$

gives M the structure of a \mathbf{C}_P -module. This corresponds to pulling back the P' -module structure on M along the map ϕ . This construction makes $P \mapsto \mathbf{C}_P$ into a contravariant functor from the category of operads in $\mathcal{S}p$ to the category of comonads on $[\Sigma, \mathcal{S}p]$.

Lemma 1.25 *Let P be a cofibrant operad (in the projective model structure on the category of reduced operads of spectra). Then the comonad \mathbf{C}_P preserves all weak equivalences.*

Proof We can rewrite $\mathbf{C}_P(A)(k)$ as

$$\prod_n \left[\prod_{n=n_1+\dots+n_k} \text{Map}(P(n_1) \wedge \dots \wedge P(n_k), A(n))^{\Sigma_{n_1} \times \dots \times \Sigma_{n_k}} \right].$$

Since P is cofibrant, each $P(n_i)$ is a cofibrant Σ_{n_i} -spectrum, and hence $P(n_1) \wedge \dots \wedge P(n_k)$ is a cofibrant $\Sigma_{n_1} \times \dots \times \Sigma_{n_k}$ -spectrum. It follows that \mathbf{C}_P preserves weak equivalences (since all objects in $[\Sigma, \mathcal{S}p]$ are fibrant). □

We now turn to the homotopy theory of coalgebras over comonads. This topic was studied in detail in [2, Section 1] and we first recall some definitions and results from there, though with one change. In this paper we work with comonads on the category of symmetric sequences that respect the spectral enrichment of that category. In this case we are able to define mapping spectra for coalgebras as well as mapping spaces.

Definition 1.26 Let $\mathbf{C}: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$ be a $\mathcal{S}p$ -enriched comonad on the category of symmetric sequences. We define the *derived mapping spectrum* for two \mathbf{C} -coalgebras A, A' to be the spectrum

$$\widetilde{\text{Map}}_{\mathbf{C}}(A, A') := \text{Tot}[\text{Map}_{\Sigma}(A, A') \rightrightarrows \text{Map}_{\Sigma}(A, \mathbf{C}(A')) \rightrightarrows \cdots].$$

This is the (fat) totalization of a cosimplicial spectrum constructed using the comonad structure on \mathbf{C} , the coalgebra structures on A and A' , and the spectral enrichment of \mathbf{C} .

As in [2], though with mapping spectra instead of mapping spaces, the derived mapping spectra $\widetilde{\text{Map}}_{\mathbf{C}}(A, A')$ determine an A_{∞} - $\mathcal{S}p$ -enriched category with composition maps parametrized by a certain A_{∞} -operad \mathcal{A} . We obtain the *homotopy category of \mathbf{C} -coalgebras*, which we denote $\text{Coalg}(\mathbf{C})^h$, with objects the Σ -cofibrant \mathbf{C} -coalgebras, and morphism sets

$$[A, A']_{\mathbf{C}} := \pi_0 \widetilde{\text{Map}}_{\mathbf{C}}(A, A').$$

A morphism $f: A \rightarrow A'$ in this homotopy category is determined by a *derived morphism of \mathbf{C} -coalgebras* which consists of compatible maps of symmetric sequences

$$f_k: \Delta_+^k \wedge A \rightarrow \mathbf{C}^k(A').$$

By [2, Proposition 1.16], such a morphism induces an isomorphism in the homotopy category if and only if the map $f_0: A \rightarrow A'$ is an equivalence of symmetric sequences.

In our classifications of analytic functors, we require a slight modification of the homotopy category constructed in Definition 1.26.

Definition 1.27 Let $\mathbf{C}: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$ be a $\mathcal{S}p$ -enriched comonad on the category of symmetric sequences. We say that \mathbf{C} *preserves truncations* if whenever A is n -truncated, then $\mathbf{C}(A)$ is also n -truncated. In this case, for every \mathbf{C} -coalgebra A and each n , there is a unique \mathbf{C} -coalgebra structure on the truncations $A_{\leq n}$ such that the truncation sequence for A consists of (strict) maps of \mathbf{C} -coalgebras.

We define the *pro-truncated mapping spectrum* for two \mathbf{C} -coalgebras A, A' to be the spectrum

$$\widetilde{\text{Map}}_{\mathbf{C}}^t(A, A') := \text{Tot holim}_n \text{Map}_{\Sigma}(A, \mathbf{C}^{\bullet}A'_{\leq n}).$$

The homotopy limit is taken over the maps of cosimplicial spectra induced by the truncation maps $A'_{\leq n} \rightarrow A'_{\leq (n-1)}$ which, we note, are strict morphisms of \mathbf{C} -coalgebras. Since totalization commutes with homotopy limits, this definition is weakly equivalent to $\text{holim}_n \widetilde{\text{Map}}_{\mathbf{C}}(A, A'_{\leq n})$, the homotopy limit of the corresponding derived mapping spectra of Definition 1.26.

Proposition 1.28 *Suppose that $\mathbf{C}: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$ is a $\mathcal{S}p$ -enriched comonad that preserves truncations and weak equivalences between Σ -cofibrant objects. Then the pro-truncated mapping spectra of Definition 1.27 are the mapping spectra in an A_∞ - $\mathcal{S}p$ -enriched category whose objects are the Σ -cofibrant \mathbf{C} -coalgebras.*

Proof Since \mathbf{C} preserves truncations, we have isomorphisms

$$\text{Map}_\Sigma(A, \mathbf{C}^k(A'_{\leq n})) \cong \text{Map}_\Sigma(A_{\leq n}, \mathbf{C}^k(A'_{\leq n})).$$

We therefore have maps

$$\text{Map}_\Sigma(A, \mathbf{C}^\bullet(A'_{\leq n})) \square \text{Map}_\Sigma(A', \mathbf{C}^\bullet(A''_{\leq n})) \rightarrow \text{Map}_\Sigma(A, \mathbf{C}^\bullet(A''_{\leq n})),$$

where \square denotes the box-product of [18] applied to cosimplicial spectra. The box-product is enriched over spaces in each variable and so we have canonical, and suitably associated maps of the form

$$(\text{holim}_n X_n^\bullet) \square (\text{holim}_n Y_n^\bullet) \rightarrow \text{holim}_n (X_n^\bullet \square Y_n^\bullet).$$

Combining these with those above we get

$$\begin{aligned} \text{holim}_n \text{Map}_\Sigma(A, \mathbf{C}^\bullet(A'_{\leq n})) \square \text{holim}_n \text{Map}_\Sigma(A', \mathbf{C}^\bullet(A''_{\leq n})) \\ \rightarrow \text{holim}_n \text{Map}_\Sigma(A, \mathbf{C}^\bullet(A''_{\leq n})). \end{aligned}$$

Taking totalizations, and applying the arguments of [2, Proposition 1.14], we get composition maps for the desired spectral A_∞ -category. \square

Definition 1.29 With \mathbf{C} as above, we define the *pro-truncated homotopy category of \mathbf{C} -coalgebras*, which we denote as $\text{Coalg}^t(\mathbf{C})^h$, to have objects the Σ -cofibrant \mathbf{C} -coalgebras and morphism sets

$$[A, A']_{\mathbf{C}}^t := \pi_0 \widetilde{\text{Map}}_{\mathbf{C}}^t(A, A').$$

A morphism $f: A \rightarrow A'$ in this homotopy category is determined by what we call a *pro-truncated derived morphism of \mathbf{C} -coalgebras*. This consists of maps of symmetric sequences

$$f_{k,n}: \Delta_+^k \wedge A \rightarrow \mathbf{C}^k(A'_{\leq n})$$

that are compatible both with the truncation sequence for A' , and with various cosimplicial structure maps.

In particular, the maps $f_{0,n}$ together make up a morphism of symmetric sequences

$$f_0: A \rightarrow A',$$

and the maps $f_{1,n}$ ensure that the large rectangle in the diagram

$$\begin{array}{ccccc}
 A & \xrightarrow{f_0} & A' & \xrightarrow{\sim} & \operatorname{holim}_n A'_{\leq n} \\
 \downarrow & & \downarrow & & \downarrow \\
 C(A) & \xrightarrow{C(f_0)} & C(A') & \longrightarrow & \operatorname{holim}_n C(A'_{\leq n})
 \end{array}$$

commutes up to homotopy, though this does not in general imply that the left-hand square commutes up to homotopy.

Remark 1.30 The pro-truncated homotopy category is in general coarser than the homotopy category of C -coalgebras defined in [2]. The subcategories of bounded coalgebras are equivalent in both homotopy categories, but, when C does not commute with homotopy limits, the pro-truncated category can have more equivalences between unbounded coalgebras.

Proposition 1.31 A pro-truncated derived morphism of Σ -cofibrant C -coalgebras $f: A \rightarrow A'$, in the sense of Definition 1.29, induces an isomorphism in the pro-truncated homotopy category if and only if the underlying map

$$f_0: A \rightarrow A'$$

is a weak equivalence of symmetric sequences.

Proof The argument is almost identical to that of [2, Proposition 1.16] using the fact that a map $f_0: A \rightarrow A'$ of Σ -cofibrant symmetric sequences is a weak equivalence if and only if for every bounded symmetric sequence X , the induced map

$$f_0^*: \operatorname{Map}_\Sigma(A', X) \rightarrow \operatorname{Map}_\Sigma(A, X)$$

is a weak equivalence. □

Lemma 1.32 Let P be a Σ -cofibrant operad of spectra. Then the comonad C_P preserves truncations and the following categories are equivalent:

- (1) The homotopy category associated to the projective model structure on $\operatorname{Mod}(P)$.

- (2) The homotopy category of C_P -coalgebras of Definition 1.26.
- (3) The pro-truncated homotopy category of C_P -coalgebras of Definition 1.29.

Proof Let M, M' be Σ -cofibrant P -modules (ie C_P -coalgebras). Then the homotopy category associated to the projective model structure on $\text{Mod}(P)$ is determined by the mapping spectra

$$\text{Map}_P(M, M') \simeq \text{Tot Map}_\Sigma((U_P F_P)^\bullet(M), M'),$$

where $F_P: [\Sigma, \mathcal{S}P] \rightarrow \text{Mod}(P)$ is the free P -module functor. Applying the adjunctions (U_P, R_P) and (F_P, U_P) , this is equivalent to the derived mapping spectra of [2, Definition 1.10]:

$$\widetilde{\text{Map}}_{C_P}(M, M') = \text{Tot Map}_\Sigma(M, (U_P R_P)^\bullet M').$$

Finally, notice that C_P commutes with homotopy limits, from which it follows that

$$\widetilde{\text{Map}}_{C_P}(M, M') \simeq \widetilde{\text{Map}}^t_{C_P}(M, M').$$

The homotopy categories associated to these three mapping spectrum constructions are therefore equivalent. □

2 Pro-operads and their modules

We now consider inverse sequences of operads and sequences of modules over them. In this section we develop some theory for these objects and see how they are related to coalgebras over associated comonads.

Definition 2.1 (Modules over pro-operads) A *pro-operad* is a cofiltered diagram in the category of operads of spectra. The only pro-operads that we consider in this paper are straightforward inverse sequences of operads so we focus on these. Let P_\bullet denote a sequence of operads of the form

$$\dots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0.$$

A *module over the pro-operad* P_\bullet is a direct sequence M^\bullet of symmetric sequences

$$M^0 \rightarrow M^1 \rightarrow M^2 \rightarrow \dots$$

together with a P_L -module structure on M^L for each L , such that the map

$$M^L \rightarrow M^{L+1}$$

is a morphism of P_{L+1} -modules. Here, M^L inherits a P_{L+1} -module structure via the operad morphism $P_{L+1} \rightarrow P_L$.

A morphism of P_\bullet -modules $f: M_1^\bullet \rightarrow M_2^\bullet$ consists of maps $f^L: M_1^L \rightarrow M_2^L$ for each L such that each diagram

$$\begin{array}{ccc} M_1^L & \longrightarrow & M_1^{L+1} \\ \downarrow & & \downarrow \\ M_2^L & \longrightarrow & M_2^{L+1} \end{array}$$

commutes. There is a category $\text{Mod}(P_\bullet)$ whose objects are the P_\bullet -modules, with morphisms as defined above. By recent work of Greenlees and Shipley [14], the category $\text{Mod}(P_\bullet)$ can be given a *strict projective model structure* arising from the projective model structures on the categories $\text{Mod}(P_L)$, in which a morphism f is a weak equivalence (or a fibration) if and only if each map f^L is a weak equivalence (or, respectively, a fibration) of P_L -modules.

Remark 2.2 More generally, given a pro-operad P_\bullet indexed by an arbitrary (small) cofiltered category \mathbb{I} , we can define a P_\bullet -module to be a diagram of symmetric sequences M^\bullet indexed by \mathbb{I}^{op} such that each M^i is a P_i -module, and, for each morphism $i \rightarrow i'$ in \mathbb{I} , the map $M^{i'} \rightarrow M^i$ is a morphism of P_i -modules.

We now consider the structure that is inherited by the homotopy colimit of a module over a pro-operad. The main result we need for this is that the homotopy colimit of a diagram of comonads (calculated objectwise) inherits a canonical comonad structure.

Definition 2.3 (Colimits of comonads) Let \mathbf{C}_\bullet be a diagram in the category of comonads on $[\Sigma, \mathcal{S}p]$ indexed by a small category \mathbb{I} . We define the colimit $\text{colim}_i \mathbf{C}_i$ objectwise, that is,

$$(\text{colim}_i \mathbf{C}_i)(A) := \text{colim}_i (\mathbf{C}_i(A)).$$

Then $\text{colim}_i \mathbf{C}_i$ has a canonical comonad structure with comultiplication given by the composite

$$\text{colim}_i \mathbf{C}_i(A) \rightarrow \text{colim}_i \mathbf{C}_i(\mathbf{C}_i(A)) \rightarrow \text{colim}_i \mathbf{C}_i(\text{colim}_{i'} \mathbf{C}_{i'}(A)),$$

in which the first map is built from the comultiplication maps for the comonads \mathbf{C}_i and the second is induced by the natural maps $\mathbf{C}_i \rightarrow \text{colim}_{i'} \mathbf{C}_{i'}$, and counit given by

$$\text{colim}_i \mathbf{C}_i(A) \rightarrow A$$

is built from the counit maps $\mathbf{C}_i(A) \rightarrow A$ for the individual comonads \mathbf{C}_i .

Definition 2.4 (Tensoring of comonads) Let \mathbf{C} be a simplicially enriched comonad on the category $[\Sigma, \mathcal{S}p]$ and let X be a simplicial set. We define the objectwise tensoring of \mathbf{C} by X to be the functor $X \otimes \mathbf{C}$ given by

$$(X \otimes \mathbf{C})(A) := X_+ \wedge \mathbf{C}(A).$$

Then $X \otimes \mathbf{C}$ inherits a canonical comonad structure with comultiplication given by the composite

$$X_+ \wedge \mathbf{C}(A) \rightarrow X_+ \wedge X_+ \wedge \mathbf{C}(\mathbf{C}(A)) \rightarrow X_+ \wedge \mathbf{C}(X_+ \wedge \mathbf{C}(A)),$$

in which the first map is built from the diagonal on X and the comultiplication for \mathbf{C} , and the second is given by the simplicial enrichment of \mathbf{C} , and counit map given by

$$X_+ \wedge \mathbf{C}(A) \rightarrow \mathbf{C}(A) \rightarrow A,$$

where the first map is induced by the collapse map $X \rightarrow *$ and the second is the counit for \mathbf{C} .

Definition 2.5 (Homotopy colimits of comonads) Let \mathbf{C}_\bullet be a diagram of simplicially enriched comonads on $[\Sigma, \mathcal{S}p]$ indexed by a small category \mathbb{I} . We define the homotopy colimit $\text{hocolim}_i \mathbf{C}_i$ objectwise by the coend

$$(\text{hocolim}_i \mathbf{C}_i)(A) := \text{hocolim}_i (\mathbf{C}_i(A)) = \Delta^r \otimes_{r \in \mathbf{\Delta}} \left[\bigvee_{i_0 \rightarrow \dots \rightarrow i_r} \mathbf{C}_{i_r}(A) \right],$$

where we use the standard model for the homotopy colimit of a diagram of spectra as the geometric realization of a simplicial object: $\mathbf{\Delta}$ is the simplicial indexing category and this is a coend over $\mathbf{\Delta}$. Notice that the homotopy colimit is defined by a combination of tensoring with the simplicial sets Δ^r and taking colimits. Using the constructions of Definitions 2.3 and 2.4 we thus obtain a canonical comonad structure on the functor $\text{hocolim}_i \mathbf{C}_i$.

Definition 2.6 Let \mathbf{P}_\bullet be an inverse sequence of operads in $\mathcal{S}p$ as before. By the construction in Definition 1.24 we obtain a corresponding sequence of comonads

$$\mathbf{C}_{\mathbf{P}_0} \rightarrow \mathbf{C}_{\mathbf{P}_1} \rightarrow \mathbf{C}_{\mathbf{P}_2} \rightarrow \dots$$

We define a new comonad

$$\mathbf{C}_{\mathbf{P}_\bullet}: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p], \quad \mathbf{C}_{\mathbf{P}_\bullet}(A) := \text{hocolim}_L \mathbf{C}_{\mathbf{P}_L}(A),$$

with comonad structure given by the construction of Definition 2.5. Note that the functor $\mathbf{C}_{\mathbf{P}_\bullet}$ is enriched in $\mathcal{S}p$ since each individual $\mathbf{C}_{\mathbf{P}_L}$ is.

We now show that when M^\bullet is a module over the pro-operad P_\bullet , the homotopy colimit of the sequence M^\bullet becomes a coalgebra over the comonad C_{P_\bullet} of Definition 2.6. The coalgebra structure map can be built from corresponding maps for tensors and strict colimits, just as is the comonad structure on C_{P_\bullet} .

Definition 2.7 (Colimits of coalgebras) Let C_\bullet be a diagram of comonads on $[\Sigma, \mathcal{S}p]$ indexed by a small category \mathbb{I} , and let M^\bullet be a diagram of symmetric sequences indexed by \mathbb{I}^{op} such that, for each morphism $i \rightarrow i'$ in \mathbb{I} , the map

$$M^{i'} \rightarrow M^i$$

is a morphism of $C_{i'}$ -coalgebras, where the coalgebra structure on M^i is given by the composite $M^i \rightarrow C_i(M^i) \rightarrow C_{i'}(M^i)$. So M^\bullet is a “module” over the diagram C_\bullet in the sense of Remark 2.2.

Then we define a $(\text{colim}_i C_i)$ -coalgebra structure on the symmetric sequence $\text{colim}_i M^i$ with structure map

$$\text{colim}_i M^i \rightarrow \text{colim}_i C_i(M^i) \rightarrow \text{colim}_i C_i(\text{colim}_{i'} M^{i'}).$$

Definition 2.8 (Tensoring of coalgebras) Let C be a simplicially enriched comonad on $[\Sigma, \mathcal{S}p]$, M a C -coalgebra, and X a simplicial set. Then we make $X \otimes M := X_+ \wedge M$ into a $(X \otimes C)$ -coalgebra via the structure map

$$X_+ \wedge M \rightarrow X_+ \wedge X_+ \wedge C(M) \rightarrow X_+ \wedge C(X_+ \wedge M).$$

Definition 2.9 (Homotopy colimits of coalgebras) Suppose that C_\bullet and M^\bullet are as in Definition 2.7. We then get a $(\text{hocolim}_i C_i)$ -coalgebra structure on the homotopy colimit

$$\text{hocolim}_i M^i = \Delta_+^r \wedge_{r \in \Delta} \left[\bigvee_{i_0 \rightarrow \dots \rightarrow i_r} M^{i_r} \right]$$

by combining the constructions of Definitions 2.7 and 2.8, as in Definition 2.5.

Definition 2.10 Let P_\bullet be an inverse sequence of operads and M^\bullet a P_\bullet -module in the sense of Definition 2.1. Let M be the homotopy colimit symmetric sequence

$$M := \text{hocolim}_L M^L.$$

By the construction of Definition 2.9, M inherits the structure of a C_{P_\bullet} -coalgebra.

Remark 2.11 If M^\bullet is a P_\bullet -module, then each term M^L is a module over the operad $\lim P_\bullet$ given by the inverse limit (in the category of operads) of the sequence P_\bullet . It follows that the homotopy colimit M also has a canonical structure of a module over $\lim P_\bullet$. In general, the structure of a C_{P_\bullet} -coalgebra includes and extends that of a $\lim P_\bullet$ -module. Another way to see this is via the existence of a canonical map of comonads

$$C_{P_\bullet} \rightarrow C_{\lim P_\bullet},$$

which is typically not an equivalence.

There is one technical construction that we need for modules over inverse sequences of operads. That is, we need to be able to construct a replacement for a given module in which all terms are Σ -cofibrant. We show how to do this now.

Definition 2.12 Let P_\bullet be an inverse sequence of reduced operads in $\mathcal{S}p$ such that each P_L is cofibrant in the projective model structure on reduced operads of spectra. Let

$$\underline{M}^\bullet : \underline{M}^0 \rightarrow \underline{M}^1 \rightarrow \underline{M}^2 \rightarrow \dots$$

be a P_\bullet -module. We recursively construct a commutative diagram of the form

$$(2.13) \quad \begin{array}{ccccccc} M^0 & \longrightarrow & M^1 & \longrightarrow & M^2 & \longrightarrow & \dots \\ \sim \downarrow & & \sim \downarrow & & \sim \downarrow & & \\ \underline{M}^0 & \longrightarrow & \underline{M}^1 & \longrightarrow & \underline{M}^2 & \longrightarrow & \dots \end{array}$$

in which $M^L \xrightarrow{\sim} \underline{M}^L$ is a weak equivalence of P_L -modules, and each M^L is Σ -cofibrant.

First, let $M^0 \xrightarrow{\sim} \underline{M}^0$ be a cofibrant replacement for \underline{M}^0 as a P_0 -module. Note that M^0 is then Σ -cofibrant because a cofibrant module over the cofibrant operad P_0 is Σ -cofibrant by [1, Proposition 2.3.14]. Now, suppose that we have built the diagram as far as the weak equivalence $M^L \xrightarrow{\sim} \underline{M}^L$. We can factor the composite

$$M^L \xrightarrow{\sim} \underline{M}^L \rightarrow \underline{M}^{L+1}$$

in the category of P_{L+1} -modules as a cofibration followed by a trivial fibration which we write as

$$M^L \twoheadrightarrow M^{L+1} \twoheadrightarrow \underline{M}^{L+1}.$$

Then $M^L \twoheadrightarrow M^{L+1}$ is a cofibration of P_{L+1} -modules with Σ -cofibrant domain, hence is a Σ -cofibration by [1, Lemma A.0.11]. It follows that M^{L+1} is Σ -cofibrant as required. Recursively this defines the diagram (2.13).

We refer to the \mathbf{P}_\bullet -module M^\bullet as a Σ -cofibrant replacement for \underline{M}^\bullet . Notice that because each M^L is Σ -cofibrant, it follows that the homotopy colimit

$$M := \operatorname{hocolim}_L M^L$$

is a Σ -cofibrant $\mathbf{C}_{\mathbf{P}_\bullet}$ -coalgebra. In particular M determines an object in the homotopy category of $\mathbf{C}_{\mathbf{P}_\bullet}$ -coalgebras described in Definition 1.26, as well as in the pro-truncated homotopy category of Definition 1.29.

3 Functors from based spaces to spectra

We now take up the study of homotopy functors from based spaces to spectra. Recall that we are writing $\mathcal{T}op_*^f$ for the full subcategory of $\mathcal{T}op_*$ consisting of the finite cell complexes, and that $[\mathcal{T}op_*^f, \mathcal{S}p]_*$ is the category of pointed simplicially enriched functors $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$.

In [2] we proved that the Taylor tower (expanded at the one-point space $*$) of a functor $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$ is determined by the action of a certain comonad \mathbf{C} on the symmetric sequence $\partial_* F$ formed by the derivatives of F (at $*$). The comonad $\mathbf{C} = \partial_* \Phi$ arises from an adjunction

$$(3.1) \quad \partial_*: [\mathcal{T}op_*^f, \mathcal{S}p]_* \rightleftarrows [\Sigma, \mathcal{S}p] : \Phi$$

in which the left adjoint ∂_* is a model for the functor taking a (cofibrant) F to its symmetric sequence of derivatives. The main result of this section is a separate description of \mathbf{C} as the comonad associated to an inverse sequence of operads via the constructions of the previous section. We start by describing that sequence.

3.1 Koszul duals of the stable little disc operads

The *little disc operads* of Boardman and Vogt [5] were introduced to classify iterated loop spaces, but have since been seen to arise in a variety of other contexts in algebraic topology: for example, Ayala and Francis show in [4] that algebras over the little n -discs operad classify certain types of homology theory on n -dimensional manifolds; the Deligne conjecture (proved by McClure and Smith [17] and others) shows that the Hochschild complex on an associative algebra forms an algebra over the chain model for the little 2-discs operad.

In this paper we show that the sequence of operads of spectra given by the suspension spectra of the topological little disc operads captures the information needed to recover the Taylor tower of a functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ from its derivatives. More precisely, the

Koszul duals of this direct sequence form an inverse sequence \mathbf{KE}_\bullet whose associated comonad $\mathbf{C}_{\mathbf{KE}_\bullet}$ coacts on the derivatives of any such F .

Throughout this paper we use the Fulton–MacPherson models for the little disc operads, as described by Getzler and Jones [11]. The only place where the precise model matters is in our proof of Lemma 3.31.

Definition 3.2 For a fixed non-negative integer L , we write \mathbb{E}_L for the operad of unbased spaces in which

$$\mathbb{E}_L(r)$$

is given by the Fulton–MacPherson compactified configuration space of r points in \mathbb{R}^L . Recall that a point in $\mathbb{E}_L(r)$ includes an r -tuple $y = [y_1, \dots, y_r]$ of points in \mathbb{R}^L defined up to translation and positive scalar multiplication. When two or more of the points y_i are equal, the point y also includes information about the relative directions and distances between these “equal” points. More details on the definition of \mathbb{E}_L can be found in [21].

There is a sequence of operads of the form

$$\mathbb{E}_0 \rightarrow \mathbb{E}_1 \rightarrow \dots \rightarrow \mathbb{E}_L \rightarrow \mathbb{E}_{L+1} \rightarrow \dots,$$

where the map

$$\mathbb{E}_L(r) \rightarrow \mathbb{E}_{L+1}(r)$$

extends points in \mathbb{R}^L to \mathbb{R}^{L+1} via the standard inclusion. Note that \mathbb{E}_0 is the trivial operad of unbased spaces which we also denote by $\mathbb{1}$.

We write \mathbf{Com} for the commutative operad of unbased spaces with $\mathbf{Com}(r) = *$ for all $r \geq 1$. Since \mathbf{Com} is terminal among operads of unbased spaces, there are operad maps $\mathbb{E}_L \rightarrow \mathbf{Com}$ that are compatible with the maps in the above sequence.

Definition 3.3 The *stable little L -discs operad* is the reduced operad E_L of spectra given by

$$E_L(n) := \Sigma^\infty \mathbb{E}_L(n)_+,$$

with operad structure maps induced by those of \mathbb{E}_L . There is a corresponding sequence of operads of spectra of the form

$$(3.4) \quad E_0 \rightarrow E_1 \rightarrow \dots \rightarrow E_L \rightarrow E_{L+1} \rightarrow \dots.$$

Note that E_0 is the trivial operad of spectra which we also denote by $\mathbb{1}$.

Let \mathbf{Com} be the commutative operad of spectra, given by

$$\mathbf{Com}(n) := \Sigma^\infty \mathbf{Com}(n)_+ \cong S,$$

the sphere spectrum. There are then operad maps $E_L \rightarrow \text{Com}$ that are compatible with the maps in the sequence (3.4). The induced map

$$E_\infty := \text{colim } E_L \rightarrow \text{Com}$$

is a weak equivalence of operads.

Definition 3.5 (Koszul dual) Let P be a reduced operad of spectra. Then the *Koszul dual* of P is a cofibrant replacement (in the projective model category of reduced operads of spectra) for the Spanier–Whitehead dual of the reduced bar construction on P . We denote such a replacement by KP . The construction K is a contravariant functor from the category of reduced operads of spectra to itself, and for operads formed from finite spectra, there is an equivalence of operads $KKP \simeq P$ described in [8].

Definition 3.6 Applying the Koszul duality functor to the sequence (3.4), we obtain an inverse sequence of operads of spectra

$$(3.7) \quad KE_\bullet : \dots \rightarrow KE_2 \rightarrow KE_1 \rightarrow KE_0.$$

Associated to the pro-operad KE_\bullet we have a comonad C_{KE_\bullet} given as in Definition 2.6.

Explicitly, for a symmetric sequence A and positive integer k , we have an equivalence

$$(3.8) \quad C_{KE_\bullet}(A)(k) \simeq \text{hocolim}_L \prod_{n \geq k} \left[\prod_{n \rightarrow k} \text{Map}(KE_L(n_1) \wedge \dots \wedge KE_L(n_k), A(n)) \right]^{h\Sigma_n}.$$

Remark 3.9 It is conjectured that the operad KE_L is equivalent to an L -fold desuspension of E_L itself. Corresponding results on the level of homology are due to Getzler and Jones [11], and on the chain level are due to Fresse [10].

Remark 3.10 Recall from Remark 2.11 that a C_{KE_\bullet} -coalgebra forms, in particular, a module over the operad $\lim KE_L$ which is equivalent to $K\text{Com}$, and hence to $\partial_* I_{\mathcal{T}op_*}$, the operad formed by the derivatives of the identity on the category of based spaces, described in [6]. We therefore have a forgetful functor

$$\text{Coalg}(C_{KE_\bullet}) \rightarrow \text{Mod}(\partial_* I_{\mathcal{T}op_*})$$

associated to the comonad map $C_{KE_\bullet} \rightarrow C_{K\text{Com}}$.

The main goal of the rest of this section is to prove that the comonad C_{KE_\bullet} acts on (a suitable choice of model of) the derivatives of a functor $F: \mathcal{T}op_* \rightarrow \mathcal{S}p$, that C_{KE_\bullet} is equivalent to the comonad C constructed from the adjunction (3.1), and hence that

the Taylor tower of F can be recovered from the derivatives of F together with their structure as a $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebra.

We should remark here that the construction of models for the derivatives of a functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ on which $\mathbf{C}_{\mathbf{KE}_\bullet}$ acts is rather involved (see steps (1)–(5) described below), as is the proof of Theorem 3.64 that $\mathbf{C}_{\mathbf{KE}_\bullet}$ is equivalent to \mathbf{C} . The reader who wishes to skip some of the technical constructions is advised to turn to Theorem 3.75 and following, where the main results of this section are laid out.

3.2 Models for the derivatives of functors from based spaces to spectra

Let $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ be a pointed simplicial functor. In this section we construct a module $d^\bullet[F]$ over the sequence of operads \mathbf{KE}_\bullet , such that the homotopy colimit $d[F] = \text{hocolim } d^L[F]$ is equivalent to the symmetric sequence $\partial_* F$ of Goodwillie derivatives of F . It then follows that these derivatives form a $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebra by the construction of Definition 2.10.

When F is a polynomial functor, the term $d^L[F]$ in the module $d^\bullet[F]$ (which is a \mathbf{KE}_L -module) is equivalent to the symmetric sequence of partially stabilized cross-effects of F , that is,

$$d^L[F](n) \simeq \Sigma^{-nL} \text{cr}_n F(S^L, \dots, S^L).$$

More generally, when F is ρ -analytic in the sense of [12], this equivalence holds for $L \geq \rho$.

Here is an outline of the construction of $d[F]$ for a given $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$:

- (1) Construct a Com-comodule $\mathbf{N}[F]$ (and hence, by pulling back, a \mathbf{E}_L -comodule for each L) that models the cross-effects of F evaluated at S^0 , and show that a polynomial functor F can be recovered from $\mathbf{N}[F]$ (Proposition 3.15).
- (2) Show that the “derived indecomposables” of $\mathbf{N}[F]$ as a \mathbf{E}_L -comodule (which can be described as a homotopy coend of the form $\mathbf{N}[F] \widetilde{\wedge}_{\mathbf{E}_L} \underline{1}$) recover the partially stabilized cross-effects of a polynomial functor F (Proposition 3.24).
- (3) Construct a model for the derived indecomposables of an \mathbf{E}_L -comodule that has a \mathbf{KE}_L -module structure (Proposition 3.44) and thus obtain the required construction for polynomial functors (Proposition 3.47).
- (4) Generalize the construction of these models to analytic functors F using the Taylor tower (Lemma 3.49).
- (5) Produce the necessary models for any functor by left Kan extension from representables (Proposition 3.54).

We start then with (1).

Definition 3.11 For each positive integer n , the construction $Y \mapsto \Sigma^\infty Y^{\wedge n}$ defines a pointed simplicial functor $\mathcal{T}op_*^f \rightarrow \mathcal{S}p$. Let $\widetilde{\Sigma^\infty Y^{\wedge n}}$ denote a (functorial) cofibrant replacement of this in the category $[\mathcal{T}op_*^f, \mathcal{S}p]_*$. A surjection $\underline{n} \twoheadrightarrow \underline{k}$ determines a natural transformation

$$\Sigma^\infty Y^{\wedge k} \rightarrow \Sigma^\infty Y^{\wedge n}$$

by way of the diagonal on a based space Y , and hence, via naturality of the cofibrant replacement, a natural map

$$\widetilde{\Sigma^\infty Y^{\wedge k}} \rightarrow \widetilde{\Sigma^\infty Y^{\wedge n}}.$$

For each Y , these maps make $\widetilde{\Sigma^\infty Y^{\wedge *}}$ into a Com–module, naturally in $Y \in \mathcal{T}op_*^f$.

Definition 3.12 For $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$ and $n \in \mathbb{N}$ we define

$$N[F](n) := \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge n}}, FY).$$

This is the spectrum of natural transformations between two functors in $[\mathcal{T}op_*^f, \mathcal{S}p]_*$, that is, the enrichment of $[\mathcal{T}op_*^f, \mathcal{S}p]_*$ over $\mathcal{S}p$. The Com–module structure maps on $\widetilde{\Sigma^\infty Y^{\wedge *}}$ make the symmetric sequence $N[F]$ into a Com–comodule (naturally in F).

Lemma 3.13 For any $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$, there is a natural equivalence

$$N[F](n) \simeq \text{cr}_n F(S^0, \dots, S^0),$$

where the right-hand side is the n^{th} cross-effect of F , evaluated at the zero-sphere in each variable.

Proof We can write $\Sigma^\infty Y^{\wedge n}$ as the total homotopy cofibre of the n –cube of spectra given by

$$J \mapsto \Sigma^\infty \text{Hom}_{\mathcal{T}op_*^f}(J_+, Y)$$

for subsets $J \subseteq \underline{n}$, and whose edges are determined by the maps $J_+ \rightarrow I_+$ for $I \subseteq J$ that collapse $J - I$ to the basepoint. Applying $\text{Nat}(-, F)$ to this total homotopy cofibre n –cube, and using the Yoneda lemma we get the total homotopy fibre of the n –cube that defines the given cross-effect. \square

Corollary 3.14 The construction $F \mapsto N[F]$ preserves homotopy colimits (defined objectwise on both sides).

Proof The cross-effects in Lemma 3.13 are equivalent to the corresponding co-cross-effects. It is easy to see that taking co-cross-effects preserves homotopy colimits. \square

The next result is unpublished work of Bill Dwyer and Charles Rezk. It says that the value $F(X)$ of a polynomial functor F at a based space X can be recovered from the Com–comodule $N(F)$ by way of a homotopy coend with the Com–module $\Sigma^\infty X^{\wedge*}$.

Proposition 3.15 *Let $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ be a polynomial functor and let $\tilde{N}[F]$ denote a Σ –cofibrant replacement for the Com–comodule $N[F]$. Then the canonical evaluation map*

$$\epsilon: \tilde{N}[F] \tilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge*}} \longrightarrow F(X)$$

is a weak equivalence for each $X \in \mathcal{T}op_*^f$.

Proof Since both sides preserve objectwise (co)fibration sequences in F , we can use the Taylor tower to reduce to the case that F is n –homogeneous, that is, of the form

$$F(X) = (E \wedge X^{\wedge n})_{h\Sigma_n}$$

for some spectrum E with Σ_n –action. Both sides also commute with the homotopy orbit construction, the left-hand side by Corollary 3.14, and with smashing with a fixed spectrum so we can reduce to the case that

$$F(X) = \Sigma^\infty X^{\wedge n}.$$

For this case, we first claim that there is an equivalence of Com–comodules

$$(3.16) \quad \phi: \underline{\text{Com}}(n, *) \xrightarrow{\sim} \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge*}}, \Sigma^\infty Y^{\wedge n}).$$

The map ϕ is constructed as follows. Notice that $\underline{\text{Com}}(n, k) = \bigvee_{n \twoheadrightarrow k} S$. Then for each surjection $\alpha: \underline{n} \twoheadrightarrow \underline{k}$, we have a corresponding natural transformation

$$\widetilde{\Sigma^\infty Y^{\wedge k}} \xrightarrow{\sim} \Sigma^\infty Y^{\wedge k} \xrightarrow{\Delta_\alpha} \Sigma^\infty Y^{\wedge n},$$

where Δ_α is the diagonal map $X^{\wedge k} \rightarrow X^{\wedge n}$ associated to α . It is clear from the definition that ϕ is a morphism of Com–comodules, where $\underline{\text{Com}}(n, *)$ has the free comodule structure built from the composition maps in the category $\underline{\text{Com}}$.

To see that ϕ is an equivalence, we consider the following commutative diagram:

$$\begin{array}{ccc} \bigvee_{n \twoheadrightarrow k} S & \xrightarrow{\phi} & \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge k}}, \Sigma^\infty Y^{\wedge n}) \\ \downarrow \sim & & \downarrow \sim \\ \text{thofib}_{J \subseteq k} \Sigma^\infty J_+^n & \xrightarrow{\sim} & \text{thofib}_{J \subseteq k} \text{Nat}_{Y \in \mathcal{T}op_*^f}(\Sigma^\infty \text{Hom}_{\mathcal{T}op_*}(J_+, Y), \Sigma^\infty Y^{\wedge n}) \end{array}$$

The right-hand vertical map is the equivalence described in the proof of Lemma 3.13. The left-hand vertical map is given as follows: a surjection $\underline{n} \twoheadrightarrow \underline{k}$ determines an element of \underline{k}_+^n that maps to the basepoint in J_+^n for any proper subset $J \subset \underline{k}$. There is therefore an induced map into the total homotopy fibre. It is easy to check that this map is an equivalence. The bottom horizontal map is induced by a levelwise equivalence of k -cubes by the enriched Yoneda Lemma, so is an equivalence. It follows that ϕ is an equivalence.

Now consider the diagram

$$(3.17) \quad \begin{array}{ccc} \underline{\text{Com}}(n, *) \widetilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge *}} & \longrightarrow & \widetilde{\Sigma^\infty X^{\wedge n}} \\ \phi \downarrow \sim & & \downarrow \sim \\ \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge *}}, \Sigma^\infty Y^{\wedge n}) \widetilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge *}} & \xrightarrow{\epsilon} & \Sigma^\infty X^{\wedge n} \end{array}$$

where the bottom map is the evaluation map that we want to show is an equivalence, the left vertical map is induced by ϕ and so is an equivalence, the right-hand map is the fixed equivalence used in the construction of ϕ , and the top horizontal map is induced by the augmentation of the simplicial object underlying

$$\underline{\text{Com}}(n, *) \widetilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge *}},$$

that is built from the composition maps

$$\bigvee_{n_0, \dots, n_r} \underline{\text{Com}}(n, n_0) \wedge_{\Sigma_{n_0}} \cdots \wedge_{\Sigma_{n_{r-1}}} \underline{\text{Com}}(n_{r-1}, n_r) \wedge_{\Sigma_{n_r}} \widetilde{\Sigma^\infty X^{\wedge n_r}} \rightarrow \widetilde{\Sigma^\infty X^{\wedge n}}.$$

This simplicial object has extra degeneracies and so the augmentation map is an equivalence. It can be checked that the diagram (3.17) is commutative and so it follows that the evaluation map is an equivalence as required. \square

Remark 3.18 The equivalence of Proposition 3.15 is part of a classification of reduced polynomial functors $\mathcal{T}op_*^f \rightarrow \mathcal{S}p$ in terms of Com-comodules (also from unpublished work of Dwyer and Rezk) to which we return in Theorem 3.82.

We now turn to part (2) of our construction of $d[F]$. We start by taking derivatives of each side of the equivalence in Proposition 3.15 to get models for the derivatives of a polynomial functor F . To state this result we use the following definition.

Definition 3.19 Let $\underline{1}$ be the bisymmetric sequence given by

$$\underline{1}(I, J) := \bigvee_{I \cong J} S,$$

where S is the sphere spectrum and the wedge sum is taken over the set of bijections from I to J . Thus $\underline{1}(I, J)$ is trivial if $|I| \neq |J|$. Notice also that this notation is consistent with that of Definition 1.14 applied to the trivial operad $\mathbf{1}$.

Corollary 3.20 *Let $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ be a polynomial functor. Then there are natural Σ_n -equivariant equivalences*

$$N[F] \widetilde{\wedge}_{\text{Com}} \underline{1}(*, n) \xrightarrow{\sim} \partial_n(F),$$

where, for each n , the symmetric sequence $\underline{1}(*, n)$ is given a trivial Com -module structure.

Proof This follows from Proposition 3.15 by applying ∂_* to each side, using the fact that taking derivatives commutes with arbitrary homotopy colimits (for $\mathcal{S}p$ -valued functors) and that there are equivalences of Com -modules

$$\partial_n(\Sigma^\infty X^{\wedge *}) \simeq \underline{1}(*, n). \quad \square$$

Notation 3.21 Let P be an operad, N a P -comodule, and \underline{B} a bisymmetric sequence that has a P -module structure on its first variable. We then write $N \widetilde{\wedge}_P \underline{B}$ for the symmetric sequence given by

$$(N \widetilde{\wedge}_P \underline{B})(n) := N(*) \widetilde{\wedge}_P \underline{B}(*, n).$$

Remark 3.22 For a P -comodule N , the symmetric sequence

$$N \widetilde{\wedge}_P \underline{1}$$

is a model for the “derived indecomposables” of N , that is, for the left derived functor of the adjoint to the trivial comodule structure functor $[\Sigma, \mathcal{S}p] \rightarrow \text{Comod}(\text{Com})$. We can rephrase Corollary 3.20 as saying that, for a polynomial functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$, the derived indecomposables of the Com -comodule $N[F]$ recover the derivatives of F .

Now recall from Definition 3.2 that we have a sequence of operads

$$E_0 \rightarrow E_1 \rightarrow E_2 \rightarrow \dots$$

and an equivalence of operads $\text{hocolim}_L E_L \xrightarrow{\sim} \text{Com}$. Using Definition 1.18 (and the fact that homotopy colimits commute with each other), one can show that there is a corresponding equivalence

$$(3.23) \quad \text{hocolim}_L N[F] \widetilde{\wedge}_{E_L} \underline{1} \xrightarrow{\sim} N[F] \widetilde{\wedge}_{\text{Com}} \underline{1} \simeq \partial_* F,$$

where $N[F]$ is given an E_L -module structure by pulling back the Com -module structure along the operad map $E_L \rightarrow \text{Com}$. This expresses the derivatives of F as a homotopy colimit of the derived indecomposables of $N[F]$ as an E_L -comodule, as $L \rightarrow \infty$.

We now prove that these derived indecomposables recover the partially stabilized cross-effects of F , thus completing part (2) of our construction.

Proposition 3.24 *Let F be a polynomial functor. Then there are equivalences of symmetric sequences*

$$N[F] \widetilde{\wedge}_{E_L} \underline{1} \simeq \Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L)$$

that are natural in F and make the following diagrams commute in the homotopy category of symmetric sequences:

$$(3.25) \quad \begin{array}{ccc} N[F] \widetilde{\wedge}_{E_L} \underline{1} & \xrightarrow{\sim} & \Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L) \\ \downarrow & & \downarrow \\ N[F] \widetilde{\wedge}_{E_{L+1}} \underline{1} & \xrightarrow{\sim} & \Sigma^{-*(L+1)} \text{cr}_* F(S^{L+1}, \dots, S^{L+1}) \end{array}$$

Here the left-hand vertical map is induced by the operad map $E_L \rightarrow E_{L+1}$, and the right-hand map is that of (1.7).

Proof Our strategy is to use Proposition 3.15 to reduce to the case that F is one of the functors $\Sigma^\infty X^{\wedge n}$. Proving the result in those cases seems to require some significant geometric input about the relationship of the operad E_L to the sphere S^L . For us, this input is contained in the construction of the equivalence in Lemma 3.31 below.

By Proposition 3.15, we can write

$$F(X) \simeq N[F] \widetilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge *}}$$

Since taking cross-effects commutes with homotopy colimits, this gives us

$$(3.26) \quad \Sigma^{-kL} \text{cr}_k F(S^L, \dots, S^L) \simeq N[F] \widetilde{\wedge}_{\text{Com}} \Sigma^{-kL} \text{cr}_k (\Sigma^\infty X^{\wedge *})(S^L, \dots, S^L),$$

where the Com -module structure on the right-hand side comes from that on $\Sigma^\infty X^{\wedge *}$. The cross-effects of the homogeneous functors $\Sigma^\infty X^{\wedge *}$ are easily calculated: we have

$$\text{cr}_k (\Sigma^\infty X^{\wedge n})(X_1, \dots, X_k) \simeq \bigvee_{\alpha: \underline{n} \twoheadrightarrow \underline{k}} \Sigma^\infty X_1^{n_1} \wedge \dots \wedge \Sigma^\infty X_k^{n_k},$$

where, as usual, we write $n_i := |\alpha^{-1}(i)|$ for a surjection $\alpha: \underline{n} \twoheadrightarrow \underline{k}$.

This model for $\text{cr}_k(\Sigma^\infty X^{\wedge n})$ is pointed simplicial in each variable with tensoring maps

$$S^k \wedge \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^\infty X_1^{n_1} \wedge \cdots \wedge \Sigma^\infty X_k^{n_k} \rightarrow \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^\infty (\Sigma X_1)^{\wedge n_1} \wedge \cdots \wedge \Sigma^\infty (\Sigma X_k)^{\wedge n_k}$$

induced by the diagonal maps $S^k = (S^1)^{\wedge k} \xrightarrow{\Delta_\alpha} (S^1)^{\wedge n}$ associated to the surjections $\underline{n} \twoheadrightarrow \underline{k}$. By Lemma 1.9, the map (1.7) for the functor $\Sigma^\infty X^{\wedge n}$ is modelled by the map

$$\bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^{-kL} \Sigma^\infty (S^L)^{\wedge n} \rightarrow \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^{-k(L+1)} \Sigma^\infty (S^{L+1})^{\wedge n}$$

induced also by the diagonal maps Δ_α . For each fixed k , we therefore have a commutative diagram in the homotopy category of Com-modules (with variable n) of the form

$$(3.27) \quad \begin{array}{ccc} \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^{-kL} (\Sigma^\infty S^L)^{\wedge n} & \xrightarrow{\sim} & \Sigma^{-kL} \text{cr}_k(\Sigma^\infty X^{\wedge n})(S^L, \dots, S^L) \\ \downarrow & & \downarrow \\ \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^{-k(L+1)} (\Sigma^\infty S^{L+1})^{\wedge n} & \xrightarrow{\sim} & \Sigma^{-k(L+1)} \text{cr}_k(\Sigma^\infty X^{\wedge n})(S^{L+1}, \dots, S^{L+1}) \end{array}$$

The Com-module structure on the left-hand side terms in (3.27) is given by composition of surjections combined with the diagonal maps for S^L and S^{L+1} .

The next step is to construct a diagram (in the homotopy category of Com-modules) of the form

$$(3.28) \quad \begin{array}{ccc} \underline{\text{Com}}(n, *) \tilde{\wedge}_{E_L} \underline{1}(*, k) & \xrightarrow{\sim} & \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^{-kL} (\Sigma^\infty S^L)^{\wedge n} \\ \downarrow & & \downarrow \\ \underline{\text{Com}}(n, *) \tilde{\wedge}_{E_{L+1}} \underline{1}(*, k) & \xrightarrow{\sim} & \bigvee_{\underline{n} \twoheadrightarrow \underline{k}} \Sigma^{-k(L+1)} (\Sigma^\infty S^{L+1})^{\wedge n} \end{array}$$

Combining (3.28) with (3.27) and applying $N[F] \tilde{\wedge}_{\text{Com}} -$ to the resulting diagram, we get, by (3.26), the required diagram (3.25). Note that there are equivalences $N[\Sigma^\infty X^{\wedge n}] \simeq \underline{\text{Com}}(n, *)$, which preserve both the Com-comodule structure and the

Com-module structure on the variable n , so the construction of diagram (3.28) amounts to proving the proposition in the case $F = \Sigma^\infty X^{\wedge n}$.

To get (3.28) we first note that there are natural isomorphisms of Com-modules

$$(3.29) \quad \underline{\text{Com}}(n, *) \widetilde{\wedge}_{\mathbb{E}_L} \underline{\mathbb{1}}(*, k) \cong \bigvee_{n \rightarrow k} \bigwedge_{i=1}^k \text{B}(1, \mathbb{E}_L, \text{Com})(n_i),$$

where the right-hand side is the two-sided operadic bar construction, as described in [6]. The above isomorphism can be identified by comparing the structure of the simplicial spectra underlying the homotopy coend and the bar construction.

The most significant part of the proof now concerns the construction of equivalences of Com-modules

$$(3.30) \quad \text{B}(1, \mathbb{E}_L, \text{Com}) \xrightarrow{\sim} \Sigma^{-L}(\Sigma^\infty S^L)^{\wedge*}.$$

We make this construction at the unstable level. For an operad \mathbb{P} of unbased spaces, let us write \mathbb{P}_+ for the corresponding operad of based spaces given by adding a disjoint basepoint to each term. The following lemma is the key calculation underlying the proof of Proposition 3.24.

Lemma 3.31 *There is a weak equivalence of Com_+ -modules (in the category of based spaces)*

$$f_L: \text{B}(\mathbb{1}_+, \mathbb{E}_{L+}, \text{Com}_+) \xrightarrow{\sim} S^{L(*-1)},$$

where we identify $S^{L(n-1)}$ with the one-point compactification of $(\mathbb{R}^L)^n/\mathbb{R}^L$ (the space of n -tuples $[y_1, \dots, y_n]$ of vectors in \mathbb{R}^L defined up to translation) and give the collection $S^{L(*-1)}$ a Com_+ -module structure via the diagonal map $\mathbb{R}^L \rightarrow \mathbb{R}^L \times \mathbb{R}^L$. These maps make the following diagrams commute:

$$\begin{array}{ccc} \text{B}(\mathbb{1}_+, \mathbb{E}_{L+}, \text{Com}_+) & \xrightarrow{f_L} & S^{L(*-1)} \\ \downarrow & & \downarrow \\ \text{B}(\mathbb{1}_+, \mathbb{E}_{L+1+}, \text{Com}_+) & \xrightarrow{f_{L+1}} & S^{(L+1)(*-1)} \end{array}$$

where the right-hand vertical map is induced by the fixed inclusion $\mathbb{R}^L \rightarrow \mathbb{R}^{L+1}$, which induces the operad map $\mathbb{E}_L \rightarrow \mathbb{E}_{L+1}$.

Proof Write $\check{\mathbb{E}}_L$ for the symmetric sequence (of unbased spaces) given by

$$\check{\mathbb{E}}_L(r) := \begin{cases} \mathbb{E}_L(r) & \text{if } r \geq 2, \\ \emptyset & \text{if } r = 1. \end{cases}$$

Note that $\check{\mathbb{E}}_L$ has a right \mathbb{E}_L -module structure coming from the operad structure on \mathbb{E}_L and that there is a homotopy cofibre sequence

$$\check{\mathbb{E}}_{L+} \rightarrow \mathbb{E}_{L+} \rightarrow \mathbb{1}_+$$

of right \mathbb{E}_{L+} -modules. Applying the bar construction, we obtain a homotopy cofibre sequence

$$(3.32) \quad B(\check{\mathbb{E}}_{L+}, \mathbb{E}_{L+}, \mathbb{C}om_+) \rightarrow B(\mathbb{E}_{L+}, \mathbb{E}_{L+}, \mathbb{C}om_+) \rightarrow B(\mathbb{1}_+, \mathbb{E}_{L+}, \mathbb{C}om_+)$$

of right $\mathbb{C}om_+$ -modules.

Now let $S^{L(n-1)-1}$ denote the sphere in the vector space $(\mathbb{R}^L)^n / \mathbb{R}^L$, or equivalently, the space of n -tuples $[y_1, \dots, y_n]$ of vectors in \mathbb{R}^L , not all equal, defined up to translation and positive scalar multiplication. Notice that $S^{L(n-1)}$ is the unreduced suspension of $S^{L(n-1)-1}$ so that we have a homotopy cofibre sequence of right $\mathbb{C}om_+$ -modules

$$(3.33) \quad S_+^{L(*-1)-1} \rightarrow \mathbb{C}om_+ \rightarrow S^{L(*-1)}.$$

The second map here sends the non-basepoint of $\mathbb{C}om_+(n)$ to the point in $S^{L(n-1)}$ that represents the n -tuple $[0, \dots, 0]$.

Our goal now is to construct an equivalence between the sequences (3.32) and (3.33) of which the required equivalence f_L will be part.

Notice first that

$$B(\check{\mathbb{E}}_{L+}, \mathbb{E}_{L+}, \mathbb{C}om_+) = B(\check{\mathbb{E}}_L, \mathbb{E}_L, \mathbb{C}om)_+.$$

We now claim that $\check{\mathbb{E}}_L$ is cofibrant as an \mathbb{E}_L -module (in the projective model structure on right modules over an operad in the category of compactly generated spaces). To see this, suppose we are given a diagram of \mathbb{E}_L -modules

$$(3.34) \quad \begin{array}{ccc} & & \mathbb{M} \\ & & \downarrow \sim \\ \check{\mathbb{E}}_L & \longrightarrow & \mathbb{M}' \end{array}$$

where the right-hand vertical map is a trivial fibration in the projective model structure on modules over the operad \mathbb{E}_L (ie each map $\mathbb{M}(n) \rightarrow \mathbb{M}'(n)$ is a Serre fibration and weak homotopy equivalence). We recursively construct a lifting $l: \check{\mathbb{E}}_L \rightarrow \mathbb{M}$ as follows. Suppose we have constructed maps

$$l_r: \check{\mathbb{E}}_L(r) \rightarrow \mathbb{M}(r),$$

for each $r < k$, that commute with the relevant \mathbb{E}_L -module structure maps. Together the l_r determine the top horizontal map in a commutative diagram

$$(3.35) \quad \begin{array}{ccc} \partial\check{\mathbb{E}}_L(k) & \longrightarrow & \mathbb{M}(k) \\ \downarrow & & \downarrow \sim \\ \check{\mathbb{E}}_L(k) & \longrightarrow & \mathbb{M}'(k) \end{array}$$

where $\partial\check{\mathbb{E}}_L(k)$ denotes the boundary of the manifold with corners $\check{\mathbb{E}}_L(k)$. (Recall that this boundary is identical to the union of the images of the non-trivial module structure maps

$$\check{\mathbb{E}}_L(r) \times \mathbb{E}_L(k_1) \times \cdots \times \mathbb{E}_L(k_r) \rightarrow \check{\mathbb{E}}_L(k).)$$

Since the inclusion $\partial\check{\mathbb{E}}_L(k) \rightarrow \check{\mathbb{E}}_L(k)$ is a relative cell complex, we can choose a lift

$$l_k: \check{\mathbb{E}}_L(k) \rightarrow \mathbb{M}(k)$$

for the diagram (3.35) which continues the recursion. Together the maps l_k determine the required lift for the diagram (3.34). It follows that $\check{\mathbb{E}}_L$ is a cofibrant \mathbb{E}_L -module, as claimed.

It then follows that there is an equivalence of $\mathbb{C}om$ -modules

$$\gamma_L: \mathbb{B}(\check{\mathbb{E}}_L, \mathbb{E}_L, \mathbb{C}om) \xrightarrow{\sim} \check{\mathbb{E}}_L \circ_{\mathbb{E}_L} \mathbb{C}om.$$

We claim that there is then an isomorphism of $\mathbb{C}om$ -modules

$$(3.36) \quad \check{\mathbb{E}}_L \circ_{\mathbb{E}_L} \mathbb{C}om \cong S^{L(*-1)-1}.$$

To see this we first define a map of $\mathbb{C}om$ -modules

$$g: \check{\mathbb{E}}_L \circ \mathbb{C}om \rightarrow S^{L(*-1)-1}.$$

The n^{th} term on the left-hand side can be written as

$$\coprod_{k \geq 2} \left[\coprod_{n \twoheadrightarrow k} \mathbb{E}_L(k) \right]_{\Sigma_k}.$$

Given an integer k and surjection $\alpha: \underline{n} \twoheadrightarrow \underline{k}$, we can define a map

$$g_\alpha: \check{\mathbb{E}}_L(k) \rightarrow S^{L(n-1)-1}$$

as follows. A point x in $\check{\mathbb{E}}_L(k)$ consists of a k -tuple $[x_1, \dots, x_k]$ of vectors in \mathbb{R}^L defined up to translation and scalar multiplication, together with extra information if some (but not all) of the x_i are equal. We set

$$g_\alpha(x) := [x_{\alpha(1)}, \dots, x_{\alpha(n)}] \in S^{L(n-1)-1}.$$

Together the maps g_α determine the required map g . We now show that g induces an isomorphism of the form (3.36).

First note that the two composites in the following diagram are equal:

$$\check{\mathbb{E}}_L \circ \mathbb{E}_L \circ \mathbb{C}om \rightrightarrows \check{\mathbb{E}}_L \circ \mathbb{C}om \rightarrow S^{L(*-1)-1},$$

so that g determines a map

$$\bar{g}: \check{\mathbb{E}}_L \circ_{\mathbb{E}_L} \mathbb{C}om \rightarrow S^{L(*-1)-1}.$$

To see that each \bar{g}_n is a bijection, we construct its inverse which we denote \bar{h}_n .

A point $y \in S^{L(n-1)-1}$ is represented by an n -tuple $[y_1, \dots, y_n]$ of vectors in \mathbb{R}^L , not all equal, defined up to translation and positive scalar multiplication. We can write $y_i = x_{\alpha(i)}$ for a k -tuple $x = [x_1, \dots, x_k]$ of *distinct* vectors, also defined up to translation and positive scalar multiplication, where $k \geq 2$ is a positive integer and $\alpha: \underline{n} \twoheadrightarrow \underline{k}$ is a surjection. Note that k and α are uniquely determined, up to action of Σ_k , by the condition that all x_i are distinct. This data determines a point in $\check{\mathbb{E}}_L \circ \mathbb{C}om$ (where, in particular, the k -tuple x represents a point in the *interior* of the space $\check{\mathbb{E}}_L(k)$), and hence a point in $(\check{\mathbb{E}}_L \circ_{\mathbb{E}_L} \mathbb{C}om)(n)$ which we take to be $\bar{h}_n(y)$.

It is now relatively easy to see that \bar{h}_n is inverse to \bar{g}_n and so that \bar{g}_n is a continuous bijection. The space $(\check{\mathbb{E}}_L \circ_{\mathbb{E}_L} \mathbb{C}om)(n)$ is compact (since each $\mathbb{E}_L(k)$ is for $2 \leq k \leq n$) and $S^{L(n-1)-1}$ is Hausdorff. Therefore \bar{g}_n is a homeomorphism, and we obtain the required isomorphism (3.36). We then have a commutative diagram of $\mathbb{C}om_+$ -modules

$$\begin{array}{ccc} B(\check{\mathbb{E}}_{L+}, \mathbb{E}_{L+}, \mathbb{C}om_+) & \longrightarrow & B(\mathbb{E}_{L+}, \mathbb{E}_{L+}, \mathbb{C}om_+) \\ \gamma_L \downarrow \sim & & \downarrow \sim \\ S_+^{L(*-1)-1} & \longrightarrow & \mathbb{C}om_+ \end{array}$$

where the right-hand map is the standard bar resolution map. We therefore get an induced equivalence between the homotopy cofibres which provides the required equivalence

$$f_L: B(\mathbb{1}_+, \mathbb{E}_{L+}, \mathbb{C}om_+) \xrightarrow{\sim} S^{L(*-1)}.$$

It is easy to see that the maps g_L commute with the maps induced by the inclusion $\mathbb{R}^L \rightarrow \mathbb{R}^{L+1}$, and hence the maps f_L do too, completing the proof of Lemma 3.31. \square

Remark 3.37 Low-dimensional calculations suggest that the based space

$$B(\mathbb{1}_+, \mathbb{E}_{L+}, \mathbb{C}om_+)(n)$$

is actually homeomorphic to $S^{L(n-1)}$ though we do not need that claim here.

Proof of Proposition 3.24 (continued) We now get the required stable equivalences (3.30) from the maps f_L of Lemma 3.31 by the composite

$$B(1, E_L, Com) \cong \Sigma^\infty B(\mathbb{1}_+, \mathbb{E}_{L+}, \mathbb{C}om_+) \xrightarrow{\sim} \Sigma^\infty S^{L(*-1)} \xrightarrow{\sim} \Sigma^{-L}(\Sigma^\infty S^L)^{\wedge*}.$$

The final equivalence is adjoint to the map

$$S^{L(k-1)} \wedge S^L \rightarrow (S^L)^{\wedge k}$$

that is the one-point compactification of the map of $\mathbb{C}om$ -modules (in the k variables)

$$(\mathbb{R}^L)^k / \mathbb{R}^L \times \mathbb{R}^L \rightarrow (\mathbb{R}^L)^k$$

given by

$$([y_1, \dots, y_k], u) \mapsto (y_1 - m + u, \dots, y_k - m + u),$$

where m is the coordinate-wise minimum of the vectors y_1, \dots, y_k . (That is, $m_i := \min_{1 \leq j \leq k} (y_j)_i$.)

Using (3.29) we then get the required diagram (3.28). This completes the proof of Proposition 3.24. \square

We now turn to part (3) of the construction of $d[F]$: showing that the derived indecomposables $N[F] \widetilde{\wedge}_{E_L} \underline{1}$ can be given the structure of a KE_L -module. Behind this construction is a form of derived Koszul duality for comodules and modules over an operad P and its Koszul dual KP . For any operad P of spectra and P -comodule N , we construct a model for the symmetric sequence

$$N \widetilde{\wedge}_P \underline{1}$$

that is a module over the Koszul dual operad KP . At the heart of our construction is a particular bisymmetric sequence \underline{B}_P , equivalent to $\underline{1}$, that has a P -module structure in one variable, and a KP -module structure in the other. We define this now.

Definition 3.38 For an operad P of spectra, let \underline{B}_P be the bisymmetric sequence given on finite nonempty sets I, J by

$$\underline{B}_P(I, J) := \prod_{I \twoheadrightarrow J} \bigwedge_{j \in J} B(1, P, P)(I_j),$$

where the product is indexed by the set of surjections from I to J . Recall from [6] that the symmetric sequence $B(1, P, P)$ has both a right P -module structure and a left BP -comodule structure (where $BP = B(1, P, 1)$ is the cooperad given by the reduced bar construction on P). We use these structures to produce the required operad actions on \underline{B}_P .

We make \underline{B}_P into a right P -module in its first variable as follows. For each surjection $\alpha: I \twoheadrightarrow I'$ we have to describe a map

$$r_\alpha: \underline{B}_P(I', J) \wedge \bigwedge_{i' \in I'} P(I_{i'}) \rightarrow \underline{B}_P(I, J).$$

The target here is a product indexed by surjections $\beta: I \twoheadrightarrow J$ and we define each component individually. There are two cases. If β does not factor as $\gamma\alpha$ for some surjection $\gamma: I' \twoheadrightarrow J$ then we take the relevant component of r_α to be the trivial map. If β does factor in this way then γ is uniquely determined. We then build the relevant component of r_α from the projection map

$$p_\gamma: \underline{B}_P(I', J) \rightarrow \bigwedge_{j \in J} B(1, P, P)(I'_j)$$

and by taking the smash product, over $j \in J$, of the maps

$$B(1, P, P)(I'_j) \wedge \bigwedge_{i' \in I'_j} P(I_{i'}) \rightarrow B(1, P, P)(I_j)$$

associated to the right P -action on $B(1, P, P)$ and the surjections $I_j \twoheadrightarrow I'_j$ given by restricting α .

We also make \underline{B}_P into a right KP -module in its second variable as follows. For each surjection $\alpha: J \twoheadrightarrow J'$ we have to describe a map

$$(3.39) \quad s_\alpha: \underline{B}_P(I, J') \wedge \bigwedge_{j' \in J'} KP(J_{j'}) \rightarrow \underline{B}_P(I, J).$$

The target is still indexed by surjections $\beta: I \twoheadrightarrow J$ and again we define each component individually. Let $\gamma = \alpha\beta: I \twoheadrightarrow J'$. Then we build the relevant component of s_α from the projection

$$p_\gamma: \underline{B}_P(I, J') \rightarrow \bigwedge_{j' \in J'} B(1, P, P)(I_{j'})$$

and by taking the smash product, over $j' \in J'$, of maps

$$B(1, P, P)(I_{j'}) \wedge KP(J_{j'}) \rightarrow \bigwedge_{j \in J_{j'}} B(1, P, P)(I_j)$$

associated to the *left* coaction by the cooperad BP on $B(1, P, P)$. Recall that the Koszul dual KP is the Spanier–Whitehead dual of BP and the above map is adjoint to the coaction map associated to the surjection $I_{j'} \twoheadrightarrow J_{j'}$ given by restricting β .

Lemma 3.40 *The constructions of Definition 3.38 make \underline{B}_P into a spectrally enriched functor $\underline{P}^{op} \times \underline{KP}^{op} \rightarrow \mathcal{S}p$, that is, a bisymmetric sequence with commuting right module structures for the operads P and KP on the first and second variables respectively.*

Proof We have to check associativity and unit conditions for each of the actions, and then a commutativity condition between them. For the P–action on the first variable, the unit condition is that the map r_α associated to the identity $\alpha: I \twoheadrightarrow I$ be the identity on $\underline{B}_P(I, J)$, which it is.

The associativity condition concerns the maps $r_\alpha, r_{\alpha'}, r_{\alpha'\alpha}$ for surjections $\alpha: I \twoheadrightarrow I'$ and $\alpha': I' \twoheadrightarrow I''$. The key point is then the following: a surjection $\beta: I \twoheadrightarrow J$ factors via $\alpha'\alpha$ if and only if β factors as $\gamma\alpha$, and then γ factors via α' . If these factorizations do not exist, then the components of $r_{\alpha'\alpha}$ and of $r_\alpha r_{\alpha'}$ corresponding to β are both trivial. If the factorizations do exist, then those components are equal by the associativity of the right P–action on $B(1, P, P)$.

For the KP–action on the second variable, the corresponding checks are similar, but they are easier since there are no separate cases in the construction of the maps s_α . The associativity condition follows from the coassociativity of the coaction of BP on $B(1, P, P)$.

To see that the two actions commute, suppose we have surjections $\alpha: I \twoheadrightarrow I'$ and $\delta: J \twoheadrightarrow J'$. We have to show that the following square commutes:

$$(3.41) \quad \begin{array}{ccc} \bigwedge_{J' \in J'} KP(J_{j'}) \wedge \underline{B}_P(I', J') \wedge \bigwedge_{i' \in I'} P(I_{i'}) & \xrightarrow{r_\alpha} & \bigwedge_{j' \in J'} KP(J_{j'}) \wedge \underline{B}_P(I, J') \\ \downarrow s_\delta & & \downarrow s_\delta \\ \underline{B}_P(I', J) \wedge \bigwedge_{i' \in I'} P(I_{i'}) & \xrightarrow{r_\alpha} & \underline{B}_P(I, J) \end{array}$$

Since the target is a product, we can check commutativity by considering the components of the composite maps corresponding to each surjection $\beta: I \twoheadrightarrow J$ in turn. There are three cases.

If $\beta = \gamma\alpha$ for some $\gamma: I' \twoheadrightarrow J$ then also $\delta\beta = (\delta\gamma)\alpha$. In this case, the relevant components of the horizontal maps in (3.41) are given by the P-action on $B(1, P, P)$. The required commutativity then follows from the commutativity of the corresponding diagram

$$\begin{array}{ccc}
 B(1, P, P)(I'_{j'}) \wedge \bigwedge_{i' \in I'_{j'}} P(I_{i'}) & \longrightarrow & B(1, P, P)(I_{j'}) \\
 \downarrow & & \downarrow \\
 BP(J_{j'}) \wedge \bigwedge_{j \in J_{j'}} B(1, P, P)(I'_j) \wedge \bigwedge_{i' \in I'_{j'}} P(I_{i'}) & \longrightarrow & BP(J_{j'}) \wedge \bigwedge_{j \in J_{j'}} B(1, P, P)(I_j)
 \end{array}$$

where the vertical maps are given by the BP-coaction maps on $B(1, P, P)$ associated to the restrictions of β to $I_{j'}$ for $j' \in J'$.

Secondly, suppose that β does not factor via α , but that $\delta\beta = \gamma'\alpha$ for some $\gamma': I' \twoheadrightarrow J'$. Then the relevant component of the bottom horizontal map in (3.41) is trivial. We need to show that the other composite in (3.41) is trivial. This composite is given by the smash product, over $j' \in J'$, of the following part of the previous diagram:

$$\begin{array}{ccc}
 B(1, P, P)(I'_{j'}) \wedge \bigwedge_{i' \in I'_{j'}} P(I_{i'}) & \longrightarrow & B(1, P, P)(I_{j'}) \\
 & & \downarrow \\
 & & BP(J_{j'}) \wedge \bigwedge_{j \in J_{j'}} B(1, P, P)(I_j)
 \end{array}$$

The only obstruction to factoring β via α is that there must be some two elements $i_1, i_2 \in I$ with $\alpha(i_1) = \alpha(i_2)$ but $\beta(i_1) \neq \beta(i_2)$. In this case, let $j' = \gamma'\alpha(i_1) = \gamma'\alpha(i_2)$ and consider the above diagram for this particular j' . To see that this composite is trivial, we recall from [6, Definition 7.8] the structure of the bar construction $B(1, P, P)$. In particular, $B(1, P, P)(I'_{j'})$ is stratified by trees with leaves labelled surjectively by the set $I'_{j'}$. Since $\alpha(i_1) = \alpha(i_2)$, these two elements must label the same leaf of such a tree. It follows that the image of the top horizontal map is contained in the strata corresponding to trees (with leaves labelled by $I_{j'}$) where i_1 and i_2 label the same leaf. But then according to the definition of the vertical map, since $\beta(i_1) \neq \beta(i_2)$, each

such stratum is mapped to the basepoint in the bottom-right corner. It follows that the above composite is trivial as required.

Finally, suppose that neither β nor $\delta\beta$ factors via α . In this case, the relevant components of both horizontal maps in (3.41) are trivial and the square commutes. \square

Lemma 3.42 *There is an equivalence of bisymmetric sequences*

$$\underline{\mathbb{B}}_P \simeq \underline{\mathbb{1}}$$

that preserves the P -module structure on the first variable.

Proof For finite sets I, J , the equivalence of right P -modules $\mathbb{B}(1, P, P) \xrightarrow{\sim} \underline{\mathbb{1}}$ determines a natural equivalence

$$\prod_{I \rightarrow J} \bigwedge_{j \in J} \mathbb{B}(1, P, P)(I_j) \rightarrow \prod_{I \rightarrow J} \bigwedge_{j \in J} \underline{\mathbb{1}}(I_j),$$

and this preserves the right P -module structures in the I variable (with that on the right-hand side the trivial structure). The right-hand side is the same as $\underline{\mathbb{1}}(I, J)$ but with $\bigvee_{\Sigma_n} S$ replaced by the equivalent $\prod_{\Sigma_n} S$ (when $n = |I| = |J|$). \square

Remark 3.43 Lemma 3.42 tells us that the P -module structure on each symmetric sequence $\underline{\mathbb{B}}_P(-, J)$ is equivalent to the trivial structure. The same is true about the KP -module structure on each $\underline{\mathbb{B}}_P(I, -)$. This is proved via equivalences

$$\underline{\mathbb{1}} \simeq \underline{\mathbb{B}}_P$$

that preserve the KP -module structure on the second variables. The essential point, however, is that these two structures are *not* equivalent when taken together. That is, the bisymmetric sequence $\underline{\mathbb{B}}_P$ is *not* equivalent to $\underline{\mathbb{1}}$ in the category of functors $P^{op} \times KP^{op} \rightarrow \mathcal{S}p$. There is no zigzag of equivalences between $\underline{\mathbb{B}}_P$ and $\underline{\mathbb{1}}$ that preserves both structures simultaneously.

Proposition 3.44 *Let P be an operad of spectra and \tilde{N} a Σ -cofibrant P -comodule. Then the symmetric sequence*

$$\tilde{N} \tilde{\wedge}_P \underline{\mathbb{B}}_P$$

is equivalent to $\tilde{N} \tilde{\wedge}_P \underline{\mathbb{1}}$, and has a canonical KP -module structure.

Proof The only part to prove is the equivalence to $\tilde{N} \tilde{\wedge}_P \underline{\mathbb{1}}$, which follows from Lemmas 3.42 and 1.19. \square

We now consider the naturality of the construction of $\underline{\mathbb{B}}_P$ in the operad P .

Lemma 3.45 *Let $\phi: P \rightarrow P'$ be a morphism of operads in $\mathcal{S}p$. The induced natural transformation of bisymmetric sequences*

$$\underline{B}_\phi: \underline{B}_P \rightarrow \underline{B}_{P'}$$

is a morphism of P -modules (in the first variables, with the P -action on $\underline{B}_{P'}(-, J)$ given by pulling back along ϕ), and of KP' -modules (in the second variables, with the KP' -action on $\underline{B}_P(I, -)$ given by pulling back the KP -action along the operad map $K\phi: KP' \rightarrow KP$).

Proof This is an easy but lengthy diagram chase. □

We are now in position to describe models for the partially stabilized cross-effects of a polynomial functor that have the desired KE_L -module structure.

Definition 3.46 For a polynomial functor $F \in [\mathcal{T}op^f_*, \mathcal{S}p]_*$, write $\tilde{N}[F]$ for a (functorial) Σ -cofibrant replacement of the Com-comodule $N[F]$ of Definition 3.12. We then define a KE_L -module $\underline{d}^L[F]$ by

$$\underline{d}^L[F] := \tilde{N}[F] \tilde{\wedge}_{E_L} \underline{B}_{E_L},$$

with KE_L -module structure arising as in Proposition 3.44. We also have, for each L , a map of symmetric sequences

$$\underline{m}^L: \underline{d}^L[F] \rightarrow \underline{d}^{L+1}[F]$$

given by the composite

$$\tilde{N}[F] \tilde{\wedge}_{E_L} \underline{B}_{E_L} \rightarrow \tilde{N}[F] \tilde{\wedge}_{E_L} \underline{B}_{E_{L+1}} \rightarrow \tilde{N}[F] \tilde{\wedge}_{E_{L+1}} \underline{B}_{E_{L+1}},$$

in which the second map is induced by the operad map $E_L \rightarrow E_{L+1}$ and the first is the corresponding map of E_L -modules $\underline{B}_{E_L} \rightarrow \underline{B}_{E_{L+1}}$ from Lemma 3.45. The map \underline{m}^L is also a morphism of KE_{L+1} -modules by Lemma 3.45.

The sequence

$$\underline{d}^\bullet[F]: \underline{d}^0[F] \rightarrow \underline{d}^1[F] \rightarrow \underline{d}^2[F] \rightarrow \dots$$

is therefore a module over the pro-operad KE_\bullet , in the sense of Definition 2.1. It follows that the homotopy colimit

$$\underline{d}[F] := \text{hocolim}_L \underline{d}^L[F]$$

is a coalgebra over the comonad C_{KE_\bullet} .

Proposition 3.47 *Let $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$ be polynomial. Then we have natural equivalences of symmetric sequences*

$$\Sigma^{-*L} cr_* F(S^L, \dots, S^L) \simeq \underline{d}^L[F]$$

such that the following diagrams commute in the homotopy category of symmetric sequences, where the left-hand maps are those of (1.7):

$$\begin{array}{ccc} \Sigma^{-*L} cr_* F(S^L, \dots, S^L) & \xrightarrow{\sim} & \underline{d}^L[F] \\ \downarrow & & \downarrow \underline{m}^L \\ \Sigma^{-*(L+1)} cr_* F(S^{L+1}, \dots, S^{L+1}) & \xrightarrow{\sim} & \underline{d}^{L+1}[F] \end{array}$$

Taking homotopy colimits as $L \rightarrow \infty$ we get an equivalence

$$\partial_*(F) \simeq \underline{d}[F].$$

Proof This follows from Propositions 3.44 and 3.24. □

We have therefore constructed models for the partially stabilized cross-effects of a polynomial functor as a module over the sequence of operads KE_\bullet , and for the Goodwillie derivatives as a coalgebra over the associated comonad C_{KE_\bullet} . We now extend these constructions to analytic functors by taking the inverse limit over the Taylor tower (part (4) of our overall construction).

Recall that a functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ is ρ -analytic if it satisfies Goodwillie’s condition for being stably n -excisive in a way that depends in a controlled manner on ρ . Specifically, there should exist a number q such that F satisfies condition $E_n(n\rho - q, \rho + 1)$ of [12, Definition 4.1]. We say that F is analytic if it is ρ -analytic for some ρ .

Definition 3.48 For an analytic functor $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$, we define

$$\underline{d}^L[F] := \text{holim}_n \underline{d}^L[P_n F],$$

where the homotopy inverse limit is formed in the category of KE_L -modules (or, equivalently, in the underlying category of symmetric sequences). The maps in the inverse diagram for the homotopy limit are those induced by the maps in the Taylor tower of F . The maps \underline{m}^L for each $P_n F$ induce corresponding maps (of KE_{L+1} -modules)

$$\underline{m}^L: \underline{d}^L[F] \rightarrow \underline{d}^{L+1}[F],$$

and hence we obtain a KE_\bullet -module $\underline{d}^*[F]$ and hence a C_{KE_\bullet} -coalgebra

$$\underline{d}[F] := \text{hocolim}_L \underline{d}^L[F].$$

(If F is polynomial, this new definition of $\underline{d}^\bullet[F]$ is termwise equivalent to that of Definition 3.46.)

Lemma 3.49 For a ρ -analytic functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ and positive integer L , there are natural maps in the homotopy category of symmetric sequences of the form

$$\Sigma^{-*L} cr_* F(S^L, \dots, S^L) \rightarrow \underline{d}^L[F]$$

that are equivalences when $L \geq \rho + 1$, and such that the following diagrams commute, where the left-hand maps are those of (1.7):

$$\begin{array}{ccc} \Sigma^{-*L} cr_* F(S^L, \dots, S^L) & \longrightarrow & \underline{d}^L[F] \\ \downarrow & & \downarrow \underline{m}^L \\ \Sigma^{-*(L+1)} cr_* F(S^{L+1}, \dots, S^{L+1}) & \longrightarrow & \underline{d}^{L+1}[F] \end{array}$$

Taking the homotopy colimit as $L \rightarrow \infty$, we obtain an equivalence of symmetric sequences

$$\partial_*(F) \simeq \underline{d}[F].$$

Proof By Proposition 3.47 we have

$$\underline{d}[P_n F] \simeq \Sigma^{-*L} cr_*(P_n F)(S^L, \dots, S^L).$$

Therefore

$$\begin{aligned} \underline{d}^L[F] &\simeq \text{holim } \Sigma^{-*L} cr_*(P_n F)(S^L, \dots, S^L) \\ &\simeq \Sigma^{-*L} cr_*(\text{holim } P_n F)(S^L, \dots, S^L), \end{aligned}$$

since taking cross-effects commutes with taking homotopy limits. The required map is then induced by the natural transformation

$$F \rightarrow \text{holim } P_n F.$$

Since F is ρ -analytic, the natural map $F \rightarrow \text{holim } P_n F$ is an equivalence on ρ -connected spaces. It follows that we have an equivalence

$$\Sigma^{-*L} cr_* F(S^L, \dots, S^L) \xrightarrow{\sim} \Sigma^{-*L} cr_*(\text{holim } P_n F)(S^L, \dots, S^L)$$

when $L - 1 \geq \rho$, which yields the first claim. Taking homotopy colimits over the L variable on each side then gives the second claim. \square

The final goal of this section (part (5)) is to extend our constructions to arbitrary (ie non-analytic) functors from based spaces to spectra. For this, we recall from [2, Proposition 4.3] that the derivatives of an arbitrary functor F can be obtained by forming the coend between the values $F(X)$ and the derivatives of the representable functors $\Sigma^\infty \text{Hom}_{\mathcal{T}op_*}(X, -)$. Since the functor represented by a finite spectrum X is analytic, we can use the models for derivatives from Definition 3.48, and since the comonad \mathbf{C}_{KE_\bullet} is enriched in spectra, its coaction on those derivatives extends to the coend.

Definition 3.50 For $X \in \mathcal{T}op_*^f$, we write

$$R_X: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$$

for the representable functor given by

$$R_X(Y) := \Sigma^\infty \text{Hom}_{\mathcal{T}op_*}(X, Y).$$

Definition 3.51 For $X \in \mathcal{T}op_*^f$, we now apply the construction of Definition 2.12 to the module $\underline{d}^\bullet[R_X]$ of Definition 3.48 to get a \mathbf{KE}_\bullet -module $d^\bullet[R_X]$ and a morphism of \mathbf{KE}_\bullet -modules

$$d^\bullet[R_X] \rightarrow \underline{d}^\bullet[R_X]$$

formed from weak equivalences

$$d^L[R_X] \xrightarrow{\sim} \underline{d}^L[R_X]$$

and such that each $d^L[R_X]$ is Σ -cofibrant. In particular, the homotopy colimit

$$d[R_X] := \text{hocolim } d^L[R_X]$$

is a Σ -cofibrant \mathbf{C}_{KE_\bullet} -coalgebra that is equivalent, as a symmetric sequence, to $\partial_*(R_X)$.

Remark 3.52 It is important that the functor

$$(\mathcal{T}op_*^f)^{\text{op}} \rightarrow [\Sigma, \mathcal{S}p]; \quad X \mapsto d[R_X],$$

be simplicially enriched in order to make Definition 3.53 below. The reader can check that we have built $d[R_X]$ from a succession of simplicially enriched constructions, including: applying Goodwillie’s construction P_n ; forming the natural transformation objects $\mathbb{N}[F]$; taking cofibrant replacements (which can be done simplicially by work of Rezk, Schwede and Shipley [20]); forming homotopy coends over E_L ; taking homotopy limits; applying functorial factorizations as in the construction of Definition 2.12; and taking homotopy colimits.

Finally, then, we obtain models for the derivatives of any pointed simplicial functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ by enriched left Kan extension from our models for the representable functors.

Definition 3.53 For any $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$ and each L , define a \mathbf{KE}_L -module $d^L[F]$ by taking the enriched coend

$$d^L[F] := F(X) \wedge_{X \in \mathcal{T}op_*^f} d^L[R_X]$$

over the simplicial category $\mathcal{T}op_*^f$. We then obtain a $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebra $d[F]$ given by

$$d[F] := \text{hocolim}_L d^L[F] \cong F(X) \wedge_{X \in \mathcal{T}op_*^f} d[R_X].$$

Notice that, by the dual Yoneda Lemma, we have $R_Y(X) \wedge_X d[R_X] \cong d[R_Y]$ so that this definition of $d[F]$ is consistent with that already made for the functors R_X themselves.

Note that in Definition 3.53 we use a strict (not homotopy) coend which typically only has the correct homotopy type when F is a cofibrant object in the projective model structure on $[\mathcal{T}op_*^f, \mathcal{S}p]_*$.

Proposition 3.54 Let $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ be pointed simplicial and let \mathbf{c} denote an arbitrary cofibrant replacement functor for the projective model structure on $[\mathcal{T}op_*^f, \mathcal{S}p]_*$. There are natural maps (in the homotopy category of symmetric sequences)

$$\eta_L(F): \Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L) \rightarrow d^L[\mathbf{c}F]$$

such that the diagrams

$$\begin{array}{ccc} \Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L) & \xrightarrow{\eta_L(F)} & d^L[\mathbf{c}F] \\ \downarrow & & \downarrow \\ \Sigma^{-*(L+1)} \text{cr}_* F(S^{L+1}, \dots, S^{L+1}) & \xrightarrow{\eta_{L+1}(F)} & d^{L+1}[\mathbf{c}F] \end{array}$$

commute, where the left-hand maps are those of (1.7) and the right-hand maps are those that form the \mathbf{KE}_\bullet -module $d^\bullet[\mathbf{c}F]$. Taking the homotopy colimit as $L \rightarrow \infty$ we therefore get a map

$$\eta(F): \partial_*(F) \xrightarrow{\sim} d[\mathbf{c}F],$$

and this is an equivalence for all F .

Proof From Lemma 3.49 we have a zigzag of maps (in which all the backwards maps are equivalences)

$$(3.55) \quad \Sigma^{-*L} \text{cr}_*(R_X)(S^L, \dots, S^L) \dashrightarrow d^L[R_X].$$

Since $\mathbf{c}F$ is a cofibrant functor, the coend construction $\mathbf{c}F(X) \wedge_X -$ preserves weak equivalences between objectwise Σ -cofibrant diagrams $(\text{Top}_*^f)^{\text{op}} \rightarrow [\Sigma, \mathcal{S}p]$. So applying this coend construction to the zigzag of maps (3.55), we obtain a corresponding zigzag

$$(3.56) \quad \mathbf{c}F(X) \wedge_X \Sigma^{-*L} \widetilde{\text{cr}}_*(R_X)(S^L, \dots, S^L) \dashrightarrow d^L[\mathbf{c}F],$$

where $\widetilde{\text{cr}}_*(R_X)$ denotes a Σ -cofibrant replacement of the cross-effects of R_X . The strict coend on the left-hand side is equivalent to the corresponding homotopy coend since $\mathbf{c}F$ is cofibrant. Therefore, because taking cross-effects commutes with homotopy colimits, the left-hand term in (3.56) is equivalent to

$$\Sigma^{-*L} \text{cr}_*(\mathbf{c}F(X) \wedge_X R_X)(S^L, \dots, S^L),$$

which is equivalent to

$$\Sigma^{-*L} \text{cr}_*(F)(S^L, \dots, S^L)$$

by the dual Yoneda Lemma. It follows that the zigzag (3.56) represents the required map $\eta_L(F)$ in the homotopy category of symmetric sequences.

It remains to show that $\eta(F) = \text{hocolim } \eta_L(F)$ is an equivalence. We know from Lemma 3.49 that the homotopy colimit of the maps (3.55), as $L \rightarrow \infty$, is an equivalence for each X . Since this homotopy colimit commutes with the homotopy coend, we deduce that $\eta(F)$ is an equivalence as required. \square

This completes the construction (started in Definition 3.11) of a model for the derivatives of a pointed simplicial functor $F: \text{Top}_*^f \rightarrow \mathcal{S}p$ that form a $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebra.

Note that for an analytic functor $F \in [\text{Top}_*^f, \mathcal{S}p]_*$ we have now defined two different $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebra models for the derivatives of F : the object $\underline{d}[F]$ of Definition 3.48 and the object $d[F]$ of Definition 3.53. It is important to know that these objects are actually equivalent as $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebras.

There is a small technical obstacle to constructing an equivalence of $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebras between $d[F]$ and $\underline{d}[F]$, which is that the construction $F \mapsto \underline{d}[F]$ is not enriched in spectra. (In particular, we do not have a natural map (of $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebras) of the form $X \wedge \underline{d}[F] \rightarrow \underline{d}[X \wedge F]$ for a spectrum X .) The underlying reason for this is that the cofibrant replacement in the category of Com -comodules used to form the object $\widetilde{N}[F]$ is not enriched in spectra. The following lemma shows that there is at least a zigzag of coalgebra maps (which are, in fact, equivalences).

Lemma 3.57 For ρ -analytic $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$, cofibrant $X \in \mathcal{S}p$, and $L \geq \rho + 1$, there are, naturally in L , zigzags of equivalences of KE_L -modules

$$\underline{d}^L[X \wedge F] \simeq X \wedge \underline{d}^L[F]$$

and a zigzag of equivalences of C_{KE_\bullet} -coalgebras

$$\underline{d}[X \wedge F] \simeq X \wedge \underline{d}[F].$$

The C_{KE_\bullet} -coalgebra structure on the right-hand side relies on the fact that the comonad C_{KE_\bullet} itself is enriched over spectra.

Proof Consider first the case that F is polynomial. Then there is a zigzag of equivalences of Σ -cofibrant Com-comodules

$$(3.58) \quad X \wedge \tilde{N}[F] \xleftarrow{\simeq} \widetilde{[X \wedge \tilde{N}[F]]} \xrightarrow{\simeq} \widetilde{[X \wedge N[F]]} \xrightarrow{\simeq} \tilde{N}[X \wedge F],$$

where the tildes denote a fixed cofibrant replacement functor in the category of Com-comodules. The final equivalence is induced by the natural map of Com-comodules

$$X \wedge \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge *}}, FY) \rightarrow \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge *}}, X \wedge FY),$$

and is an equivalence by Lemma 3.13.

We get from (3.58) the required chain of equivalences of KE_L -modules

$$X \wedge \underline{d}^L[F] := X \wedge \tilde{N}[F] \tilde{\lambda}_{E_L} \underline{B}_{E_L} \simeq \tilde{N}[X \wedge F] \tilde{\lambda}_{E_L} \underline{B}_{E_L} = \underline{d}^L[X \wedge F].$$

It is important for this that each term in the chain (3.58) is Σ -cofibrant. Note that these equivalences are compatible with the operad maps $KE_{L+1} \rightarrow KE_L$ and thus induce a corresponding chain of equivalences of C_{KE_\bullet} -coalgebras

$$X \wedge \underline{d}[F] \simeq \underline{d}[X \wedge F].$$

Now suppose that F is ρ -analytic with $L \geq \rho + 1$. Then we have a sequence of maps/equivalences of KE_L -modules of the form

$$(3.59) \quad \begin{aligned} X \wedge \underline{d}^L[F] &= X \wedge \text{holim}_n \underline{d}^L[P_n F] \rightarrow \text{holim}_n X \wedge \underline{d}^L[P_n F] \\ &\simeq \text{holim}_n \underline{d}^L[X \wedge P_n F] \\ &\simeq \text{holim}_n \underline{d}^L[P_n(X \wedge F)] = \underline{d}^L[X \wedge F]. \end{aligned}$$

By Lemma 3.49 the composite can be identified (in the homotopy category of symmetric sequences) with the map

$$X \wedge \Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L) \rightarrow \Sigma^{-*L} \text{cr}_*(X \wedge F)(S^L, \dots, S^L),$$

which is an equivalence since cross-effects are equivalent to co-cross-effects, and hence commute with smashing by X . Taking the homotopy colimit of the equivalences (3.59) as $L \rightarrow \infty$ we get the required equivalence

$$X \wedge \underline{d}[F] \simeq \underline{d}[X \wedge F]. \quad \square$$

Lemma 3.60 *For a cofibrant ρ -analytic functor $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$ and $L \geq \rho + 1$, there are, naturally in L , zigzags of equivalences of \mathbf{KE}_L -modules*

$$\underline{d}^L[F] \simeq d^L[F]$$

and a zigzag of equivalences of $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebras

$$\underline{d}[F] \simeq d[F].$$

Proof In the case $F = R_X$, the required equivalence follows immediately from the construction of $d^\bullet[R_X]$ from $\underline{d}^\bullet[R_X]$ in Definition 3.51. For arbitrary F , we have

$$d[F] = F(X) \wedge_X d[R_X] \simeq F(X) \wedge_X \underline{d}[R_X] \simeq \underline{d}[F(X) \wedge_X R_X] \cong \underline{d}[F],$$

and similarly for d^L instead of d , with the penultimate equivalence arising from Lemma 3.57. □

Corollary 3.61 *If F is ρ -analytic and $L \geq \rho + 1$, then the map*

$$\eta_L(F): \Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L) \rightarrow d^L[\mathbf{c}F]$$

of Proposition 3.54 is an equivalence.

Proof The map $\eta_L(F)$ is equivalent to the composite

$$\Sigma^{-*L} \text{cr}_*(\mathbf{c}F)(S^L, \dots, S^L) \xrightarrow{\sim} \underline{d}^L[\mathbf{c}F] \xrightarrow{\sim} d^L[\mathbf{c}F]$$

given by the equivalences of Lemmas 3.49 and 3.60. □

3.3 Classification of polynomial functors from based spaces to spectra

The main goal of this section is to prove Theorem 3.75, which provides an equivalence between the homotopy theory of polynomial functors from based spaces to spectra and the homotopy theory of bounded $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebras. Our approach is to make use of the general theory of [2] in which we have already constructed another comonad \mathbf{C} whose coalgebras classify polynomial functors.

Definition 3.62 The functor

$$d: [\mathcal{T}op_*^f, \mathcal{S}p]_* \rightarrow [\Sigma, \mathcal{S}p]$$

constructed in Definition 3.53 has a right adjoint Φ given by

$$\Phi(A)(X) := \text{Map}_\Sigma(d[R_X], A).$$

Since $d[R_X]$ is a Σ -cofibrant symmetric sequence, the right adjoint Φ preserves fibrations and acyclic fibrations, so that (d, Φ) is a Quillen adjunction (with respect to the projective model structures on both sides).

We can now apply the work of [2] to the adjunction (d, Φ) . This gives us a comonad \mathbf{C} on the category of symmetric sequences that acts on $d[F]$ for any $F \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$. According to [2, Theorem 3.13], the Taylor tower of a functor F can then be recovered from $d[F]$ by a cobar construction, and the homotopy theory of pointed simplicial polynomial functors $\mathcal{T}op_*^f \rightarrow \mathcal{S}p$ is equivalent to the homotopy theory of bounded \mathbf{C} -coalgebras. Here we show that \mathbf{C} is equivalent, as a comonad, to \mathbf{C}_{KE_\bullet} , and we deduce that the Taylor tower of F can be recovered from the \mathbf{C}_{KE_\bullet} -coalgebra structure on $d[F]$ constructed in Definition 3.53.

Definition 3.63 The comonad $\mathbf{C}: [\Sigma, \mathcal{S}p] \rightarrow [\Sigma, \mathcal{S}p]$ can be written as

$$\mathbf{C} := d\mathbf{c}\Phi,$$

where \mathbf{c} is a cofibrant replacement comonad for the category $[\mathcal{T}op_*^f, \mathcal{S}p]_*$. See [2, Section 3] for more details.¹

The functor \mathbf{C} takes values in \mathbf{C}_{KE_\bullet} -coalgebras (since d does) and so we get a morphism of comonads

$$\theta: \mathbf{C} \rightarrow \mathbf{C}_{KE_\bullet}$$

given by the composite

$$\mathbf{C}(A) \rightarrow \mathbf{C}_{KE_\bullet}(\mathbf{C}(A)) \rightarrow \mathbf{C}_{KE_\bullet}(A),$$

where the first map is given by the \mathbf{C}_{KE_\bullet} -coalgebra structure on $\mathbf{C}(A)$ and the second by the counit for the comonad \mathbf{C} . The map θ respects the comonad structures because the comultiplication for \mathbf{C} is a natural map of \mathbf{C}_{KE_\bullet} -coalgebras.

¹More precisely, we constructed \mathbf{C} in [2] using a Quillen equivalence $u: [\mathcal{T}op_*^f, \mathcal{S}p]_c \rightleftarrows [\mathcal{T}op_*^f, \mathcal{S}p]_* : c$, where $[\mathcal{T}op_*^f, \mathcal{S}p]_c$ is a model category in which every object is cofibrant. The comonad \mathbf{C} is then given by $duc\Phi$. In this case the composite uc provides a cofibrant replacement comonad for the category $[\mathcal{T}op_*^f, \mathcal{S}p]_*$ which we denote by \mathbf{c} .

Theorem 3.64 *The comonad map*

$$\theta_A: \mathbf{C}(A) \rightarrow \mathbf{C}_{\mathbf{KE}_\bullet}(A)$$

is an equivalence for any symmetric sequence A.

Proof Our proof is based on the construction, for bounded symmetric sequences A, of natural zigzags of equivalences of \mathbf{KE}_L -modules

$$(3.65) \quad d^L[\mathbf{c}\Phi A] \simeq \mathbf{C}_{\mathbf{KE}_L}(A)$$

that commute with the \mathbf{KE}_\bullet -module structure maps

$$d^L[\mathbf{c}\Phi A] \rightarrow d^{L+1}[\mathbf{c}\Phi A]$$

on the left-hand side, and the comonad maps

$$\mathbf{C}_{\mathbf{KE}_L}(A) \rightarrow \mathbf{C}_{\mathbf{KE}_{L+1}}(A)$$

on the right-hand side.

We start by noting that there are equivalences of Com-comodules

$$\begin{aligned} N[\Phi A] &= \text{Nat}_{Y \in \mathcal{J}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge *}}, \Phi A(Y)) \\ &\cong \text{Map}_\Sigma(d[\widetilde{\Sigma^\infty Y^{\wedge *}}], A) \\ &\simeq \text{Map}_\Sigma(\underline{1}(-, *), A(-)) \\ &\simeq A, \end{aligned}$$

where A has the trivial Com-comodule structure. We therefore get, using Lemma 3.60, an equivalence of \mathbf{KE}_L -modules

$$(3.66) \quad d^L[\mathbf{c}\Phi A] \simeq \tilde{N}[\Phi A] \tilde{\wedge}_{E_L} \underline{B}_{E_L} \simeq A \tilde{\wedge}_{E_L} \underline{B}_{E_L},$$

where A is given the trivial E_L -comodule structure. Note that ΦA is analytic because A is bounded and so Lemma 3.60 applies here.

We next claim that when A is bounded, $A \tilde{\wedge}_{E_L} \underline{B}_{E_L}$ is equivalent to the cofree \mathbf{KE}_L -module generated by A. This is part of a version of Koszul duality between E_L -comodules and \mathbf{KE}_L -modules under which trivial E_L -comodules correspond to cofree \mathbf{KE}_L -modules. We record this claim as a separate proposition.

Proposition 3.67 *Let A be a bounded trivial Σ -cofibrant E_L -comodule. Then there is a zigzag of equivalences of \mathbf{KE}_L -modules*

$$A \tilde{\wedge}_{E_L} \underline{B}_{E_L} \simeq \mathbf{C}_{\mathbf{KE}_L}(A).$$

In other words, the derived indecomposables of a bounded trivial E_L -comodule form a cofree KE_L -module.

Proof To define the required zigzag, we introduce a symmetric sequence equivalent to A , but that accepts a natural map from $A \widetilde{\wedge}_{E_L} \underline{B}_{E_L}$. We define

$$\widehat{A}(n) := A(n) \wedge_{\Sigma_n} \prod_{\Sigma_n} S.$$

Note that there is an isomorphism $A(n) \cong A(n) \wedge_{\Sigma_n} \bigvee_{\Sigma_n} S$ and that, since A is Σ -cofibrant, the map

$$A \rightarrow \widehat{A},$$

induced by the map from coproduct to product, is an equivalence.

We can now define a map of symmetric sequences $A \widetilde{\wedge}_{E_L} \underline{B}_{E_L} \rightarrow \widehat{A}$ as follows. For given n , we have a map

$$(3.68) \quad [A \widetilde{\wedge}_{E_L} \underline{B}_{E_L}](n) \rightarrow [A_{=n} \widetilde{\wedge}_{E_L} \underline{B}_{E_L}](n)$$

induced by the projection $A \rightarrow A_{=n}$, where $A_{=n}$ is the symmetric sequence with $A(n)$ in the n^{th} term and trivial otherwise. The right-hand side in (3.68) is the geometric realization of a simplicial spectrum that is constant with terms

$$A(n) \wedge_{\Sigma_n} \underline{B}_{E_L}(n, n) \cong \widehat{A}(n)$$

since $\underline{B}_{E_L}(n, n) \cong \prod_{\Sigma_n} S$ by Definition 3.38. It follows that we have isomorphisms

$$(3.69) \quad [A_{=n} \widetilde{\wedge}_{E_L} \underline{B}_{E_L}](n) \cong \widehat{A}(n).$$

Composing (3.68) and (3.69) we get the required map of symmetric sequences

$$\epsilon: A \widetilde{\wedge}_{E_L} \underline{B}_{E_L} \rightarrow \widehat{A}.$$

We can now define the zigzag of maps required by the proposition. We have maps of KE_L -modules

$$(3.70) \quad A \widetilde{\wedge}_{E_L} \underline{B}_{E_L} \rightarrow C_{KE_L}(A \widetilde{\wedge}_{E_L} \underline{B}_{E_L}) \xrightarrow{\epsilon} C_{KE_L}(\widehat{A}) \xleftarrow{\sim} C_{KE_L}(A),$$

where the first map is given by the KE_L -module structure on \underline{B}_{E_L} . It is now sufficient to check that this composite is an equivalence.

We start by reducing to the case that $A = \underline{1}(n, *)$, ie A consists of only a free Σ_n -spectrum concentrated in arity n . Since A is a trivial E_L -comodule, we have an equivalence of E_L -comodules

$$A \simeq \bigvee_n A(n) \wedge_{h\Sigma_n} \underline{1}(n, *).$$

To reduce our claim to the case $A = \underline{1}(n, *)$ it is sufficient to check that the functors $-\widetilde{\wedge}_{E_L} \underline{B}_{E_L}$ and C_{KE_L} commute with: (1) smashing with a cofibrant spectrum; (2) taking homotopy orbits with respect to Σ_n -action; and (3) finite coproducts. For $-\widetilde{\wedge}_{E_L} \underline{B}_{E_L}$ these are clear and for C_{KE_L} they follow from the next lemma.

Lemma 3.71 *For bounded symmetric sequences A , there is a natural equivalence of spectra*

$$C_{KE_L}(A)(k) \simeq \bigvee_{m=m_1+\dots+m_k} [BE_L(m_1) \wedge \dots \wedge BE_L(m_k) \wedge A(m)]_{h\Sigma_{m_1} \times \dots \times \Sigma_{m_k}},$$

where the coproduct is taken over all ordered partitions of a positive integer m as a sum of positive integers m_1, \dots, m_k .

Proof From Definition 1.20 we have

$$C_{KE_L}(A)(k) = \prod_m \left[\prod_{\underline{m} \rightarrow \underline{k}} \text{Map}(KE_L(m_1) \wedge \dots \wedge KE_L(m_k), A(m)) \right]^{\Sigma_m}.$$

First note that the terms in the outer product can be written as

$$\prod_{m=m_1+\dots+m_k} [\text{Map}(KE_L(m_1) \wedge \dots \wedge KE_L(m_k), A(m))]^{\Sigma_{m_1} \times \dots \times \Sigma_{m_k}},$$

where the product is taken over all ordered partitions as in the statement of the lemma. Since each $KE_L(m_j)$ is cofibrant as a Σ_{m_j} -spectrum, the strict fixed points here are equivalent to the corresponding homotopy fixed points. Furthermore, we have equivariant equivalences

$$(3.72) \quad \text{Map}(KE_L(m_1) \wedge \dots \wedge KE_L(m_k), A(m)) \simeq BE_L(m_1) \wedge \dots \wedge BE_L(m_k) \wedge A(m),$$

and each of the products involved here is finite (since A is bounded), so altogether we can write

$$C_{KE_L}(A)(k) \simeq \bigvee_{m=m_1+\dots+m_k} [BE_L(m_1) \wedge \dots \wedge BE_L(m_k) \wedge A(m)]^{h\Sigma_{m_1} \times \dots \times \Sigma_{m_k}}.$$

All that remains is to show that the homotopy fixed points here are equivalent to the corresponding homotopy orbits. To see this we note that the Σ_m -spectrum $BE_L(m)$ can be built from finitely many Σ_m -free cell spectra (because each space $\mathbb{E}_L(k)$ is a finite free Σ_k -cell complex). Therefore the norm map from homotopy orbits to homotopy fixed points is an equivalence. \square

Continuing with the proof of Proposition 3.67, this completes the reduction to the case $A = \underline{1}(n, *)$, so we now consider this case. Recall that we have to prove that the composite

$$(3.73) \quad \underline{1}(n, *) \widetilde{\wedge}_{E_L} \underline{B}_{E_L} \rightarrow \mathbf{C}_{KE_L}(\underline{1}(n, *) \widetilde{\wedge}_{E_L} \underline{B}_{E_L}) \rightarrow \mathbf{C}_{KE_L}(\widehat{\underline{1}}(n, *))$$

is an equivalence. For this we really need to get our hands on the definition of \underline{B}_{E_L} in Definition 3.38, including the E_L and KE_L -module structures described there.

The k^{th} term in this map of symmetric sequences takes the form

$$\begin{aligned} & \underline{1}(n, *) \widetilde{\wedge}_{E_L} \underline{B}_{E_L}(*, k) \\ & \rightarrow \left[\prod_{n \twoheadrightarrow k} \text{Map}(KE_L(n_1) \wedge \cdots \wedge KE_L(n_k), \underline{1}(n, *) \widetilde{\wedge}_{E_L} \underline{B}_{E_L}(*, n)) \right]^{\Sigma_n} \\ & \cong \left[\prod_{n \twoheadrightarrow k} \text{Map}(KE_L(n_1) \wedge \cdots \wedge KE_L(n_k), \widehat{\underline{1}}(n, n)) \right]^{\Sigma_n}, \end{aligned}$$

where the first map comes from the right KE_L -module structure on \underline{B}_{E_L} . The only relevant structure map is that of the form

$$\underline{B}_{E_L}(n, k) \wedge \bigwedge_{i=1}^k KE_L(n_i) \rightarrow \underline{B}_{E_L}(n, n),$$

which, by (3.39), is built from the smash product of the maps

$$B(1, E_L, E_L)(n_i) \wedge KE_L(n_i) \rightarrow B(1, E_L, E_L)(1) \wedge \cdots \wedge B(1, E_L, E_L)(1) \cong S$$

that are dual to the left BE_L -comodule structure maps for $B(1, E_L, E_L)$.

Now notice that by analyzing the homotopy coend we get equivalences

$$\underline{1}(n, *) \widetilde{\wedge}_{E_L} \underline{B}_{E_L}(*, k) \xrightarrow{\sim} \prod_{n \twoheadrightarrow k} \bigwedge_{i=1}^k B(1, E_L, 1)(n_i).$$

(This amounts to the fact that $B(1, E_L, 1)$ is equivalent to the derived indecomposables of the E_L -module structure on $B(1, E_L, E_L)$.) Putting this together with the KE_L -module structure from above, we have to show that the maps

$$\begin{aligned} & B(1, E_L, 1)(n_i) \rightarrow \text{Map}(KE_L(n_i), B(1, E_L, 1)(1) \wedge \cdots \wedge B(1, E_L, 1)(1)) \\ & \cong \text{Map}(KE_L(n_i), S), \end{aligned}$$

given by the cooperad structure on BE_L , are equivalences. But these are just evaluation maps for the Spanier–Whitehead duals of the finite spectra $BE_L(n_i)$ and hence are

indeed equivalences. It follows that (3.73) too is an equivalence as required. This completes the proof of Proposition 3.67. \square

We return now to the rest of the proof of Theorem 3.64. Together with (3.66), the equivalence of Proposition 3.67 forms the required equivalences (3.65) of the form

$$\zeta^L: d^L[\mathbf{c}\Phi A] \simeq \mathbf{C}_{KE_L}(A)$$

for bounded symmetric sequences A . Note that taking homotopy colimits over L gives us an equivalence of \mathbf{C}_{KE_\bullet} -coalgebras $\mathbf{C}(A) \simeq \mathbf{C}_{KE_\bullet}(A)$.

Suppose now that A is an arbitrary symmetric sequence. We then have a zigzag of equivalences of \mathbf{C}_{KE_\bullet} -coalgebras

$$\zeta: \text{hocolim}_L \text{holim}_n d^L[\mathbf{c}\Phi A_{\leq n}] \simeq \text{hocolim}_L \text{holim}_n \mathbf{C}_{KE_L}(A_{\leq n}),$$

where we take the homotopy limit along the truncation maps $A_{\leq n} \rightarrow A_{\leq(n-1)}$ and the homotopy colimit as $L \rightarrow \infty$. Now the right-hand side above is equivalent to $\mathbf{C}_{KE_\bullet}(A)$ (because the homotopy limit commutes with \mathbf{C}_{KE_L} and $\text{holim}_n A_{\leq n} \simeq A$). For the left-hand side, we have a diagram (in the homotopy category of symmetric sequences)

$$\begin{array}{ccc} \text{hocolim}_L \Sigma^{-*L} \text{cr}_*(\Phi A)(S^L, \dots) & \xrightarrow{\sim} & \text{hocolim}_L \text{holim}_n \Sigma^{-*L} \text{cr}_*(\Phi A_{\leq n})(S^L, \dots) \\ \sim \downarrow & & \downarrow \sim \\ \text{hocolim}_L d^L[\mathbf{c}\Phi A] & \xrightarrow{\quad\quad\quad} & \text{hocolim}_L \text{holim}_n d^L[\mathbf{c}\Phi A_{\leq n}] \end{array}$$

in which the vertical maps are the equivalences of Proposition 3.54. The top horizontal map is an equivalence because cross-effects commute with homotopy limits, so we deduce that the bottom horizontal map is an equivalence. It therefore follows that ζ is a zigzag of equivalences of \mathbf{C}_{KE_\bullet} -coalgebras

$$\zeta: \mathbf{C}(A) \simeq \mathbf{C}_{KE_\bullet}(A).$$

Finally, we consider the following diagram (in the homotopy category)

$$\begin{array}{ccccc} \mathbf{C}(A) & \longrightarrow & \mathbf{C}_{KE_\bullet}(\mathbf{C}(A)) & \xrightarrow{\epsilon_{\mathbf{C}}} & \mathbf{C}_{KE_\bullet}(A) \\ \downarrow \zeta \sim & & \downarrow \zeta \sim & & \downarrow \zeta \sim \\ \mathbf{C}_{KE_\bullet}(A) & \longrightarrow & \mathbf{C}_{KE_\bullet}(\mathbf{C}_{KE_\bullet}(A)) & \xrightarrow{\epsilon_{\mathbf{C}_{KE_\bullet}}} & \mathbf{C}_{KE_\bullet}(A) \end{array}$$

where the left-hand horizontal maps are given by the \mathbf{C}_{KE_\bullet} -coalgebra structures on \mathbf{C} and \mathbf{C}_{KE_\bullet} , respectively.

The composite of the bottom row is the identity, and the composite of the top row is θ . To show that θ is an equivalence, it is sufficient then to show that this diagram commutes in the homotopy category. The left-hand square commutes because ζ is a zigzag of equivalences of $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebras. To see that the right-hand square commutes (in the homotopy category) it is sufficient to show that

$$\begin{array}{ccc}
 \mathbf{C}(A) & & \\
 \downarrow \zeta & \searrow \epsilon_{\mathbf{C}} & \\
 \mathbf{C}_{\mathbf{KE}_\bullet}(A) & \nearrow \epsilon_{\mathbf{C}_{\mathbf{KE}_\bullet}} & A
 \end{array}$$

commutes up to homotopy. This follows by comparing the description of $\epsilon_{\mathbf{C}}$ given in [2, Lemma 4.10] with that of the counit for $\mathbf{C}_{\mathbf{KE}_\bullet}$ in Definition 1.20: in each case the counit map is, up to homotopy, a projection on the summand of the respective comonad. This completes the proof of Theorem 3.64. \square

Remark 3.74 Theorem 3.64 provides an alternative proof of the result of [2, Proposition 6.1] that describes the comonad \mathbf{C} up to homotopy in terms of “divided power” module structures for the operad $\partial_* I$ formed from the derivatives of the identity on based spaces. For a bounded symmetric sequence A we have, using Lemma 3.71,

$$\begin{aligned}
 \mathbf{C}(A)(k) &\simeq \mathbf{C}_{\mathbf{KE}_\bullet}(A)(k) \\
 &\simeq \operatorname{hocolim}_L \bigvee_{m=m_1+\dots+m_k} [\mathbf{BE}_L(m_1) \wedge \dots \wedge \mathbf{BE}_L(m_k) \wedge A(m)]_{h\Sigma_{m_1} \times \dots \times \Sigma_{m_k}} \\
 &\simeq \bigvee_{m=m_1+\dots+m_k} [\operatorname{hocolim}_L (\mathbf{BE}_L(m_1) \wedge \dots \wedge \mathbf{BE}_L(m_k)) \wedge A(m)]_{h\Sigma_{m_1} \times \dots \times \Sigma_{m_k}} \\
 &\simeq \bigvee_{m=m_1+\dots+m_k} [\mathbf{BCom}(m_1) \wedge \dots \wedge \mathbf{BCom}(m_k) \wedge A(m)]_{h\Sigma_{m_1} \times \dots \times \Sigma_{m_k}} \\
 &\simeq \prod_{m=m_1+\dots+m_k} [\operatorname{Map}(\partial_{m_1} I \wedge \dots \wedge \partial_{m_k} I, A(m))]_{h\Sigma_{m_1} \times \dots \times \Sigma_{m_k}},
 \end{aligned}$$

where the last equivalence is given by the equivalence $\partial_* I \simeq \mathbf{KCom}$ of [6].

We showed in [2] that the functor d induces an equivalence between the homotopy theory of polynomial functors in $[\mathcal{Top}_*^f, \mathcal{Sp}]_*$ and the homotopy theory of bounded \mathbf{C} -coalgebras. Using Theorem 3.64 we can now replace \mathbf{C} with $\mathbf{C}_{\mathbf{KE}_\bullet}$ in that statement.

Theorem 3.75 *The functor d induces an equivalence*

$$[\mathcal{T}op_*^f, \mathcal{S}p]_{*, \text{poly}}^h \xrightarrow{\sim} \text{Coalg}_b(\mathbf{C}_{\text{KE}_\bullet})^h$$

between the homotopy category of polynomial pointed simplicial functors $\mathcal{T}op_*^f \rightarrow \mathcal{S}p$ and the homotopy category of bounded KE_\bullet -coalgebras of Definition 1.26. Moreover, for cofibrant polynomial functors $F, G \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$, we have equivalences of mapping spectra

$$(3.76) \quad \text{Nat}_{\mathcal{T}op_*^f}(F, G) \simeq \widetilde{\text{Map}}_{\mathbf{C}_{\text{KE}_\bullet}}(d[F], d[G]).$$

Proof We prove (3.76) first. This is made more difficult by the fact that the comonad \mathbf{C} is not enriched in $\mathcal{S}p$ and so we are unable to define mapping spectra for \mathbf{C} -coalgebras. Instead we have only mapping spaces $\widetilde{\text{Hom}}_{\mathbf{C}}(-, -)$ defined as in [2, Definition 1.10]. We do then have a sequence

$$\Omega^\infty \text{Nat}_{\mathcal{T}op_*^f}(F, G) \xrightarrow{\sim} \widetilde{\text{Hom}}_{\mathbf{C}}(d[F], d[G]) \xrightarrow{\sim} \Omega^\infty \widetilde{\text{Map}}_{\mathbf{C}_{\text{KE}_\bullet}}(d[F], d[G]),$$

where the first equivalence is [2, Corollary 3.15] and the second equivalence follows from Theorem 3.64. This implies that the map

$$\text{Nat}_{\mathcal{T}op_*^f}(F, G) \rightarrow \widetilde{\text{Map}}_{\mathbf{C}_{\text{KE}_\bullet}}(d[F], d[G])$$

induces an isomorphism on homotopy groups π_k for $k \geq 0$. For $k < 0$, we can apply the result just proved to see that

$$\text{Nat}_{\mathcal{T}op_*^f}(F, \Sigma^{-k}G) \rightarrow \widetilde{\text{Map}}_{\mathbf{C}_{\text{KE}_\bullet}}(d[F], d[\Sigma^{-k}G])$$

is an equivalence on π_0 and hence so is the equivalent map

$$\Sigma^{-k} \text{Nat}_{\mathcal{T}op_*^f}(F, G) \rightarrow \Sigma^{-k} \widetilde{\text{Map}}_{\mathbf{C}_{\text{KE}_\bullet}}(d[F], d[G]).$$

Therefore the map (3.76) induces an isomorphism on π_k for all k , so is an equivalence of spectra.

It now follows that d determines a fully faithful embedding of the homotopy theory of polynomial functors in $[\mathcal{T}op_*^f, \mathcal{S}p]_*$ into the homotopy theory of bounded $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebras. To complete the proof, we must show that every bounded $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebra A is in the image of this embedding.

Given a bounded $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebra A , we define a functor $F_A: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ by

$$F_A(X) := \widetilde{\text{Map}}_{\mathbf{C}_{\text{KE}_\bullet}}(d[R_X], A) \cong \text{Tot}(\Phi \mathbf{C}_{\text{KE}_\bullet}^* A),$$

with the asterisk denoting the cosimplicial variable. We then claim that F_A is polynomial, and that $d[\mathbf{c}F_A]$ is equivalent to A in the homotopy category of $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebras.

If A is N -truncated, then so is $\mathbf{C}_{\mathbf{KE}_\bullet}^r(A)$ for any r . Therefore, each term in the cosimplicial object $\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^*A$ is N -excisive, by [2, Remark 3.10]. It follows that the (fat) totalization $\text{Tot}(\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^*A) = F_A$ is also N -excisive.

Since d is a simplicial functor we have natural maps

$$f_r: \Delta_+^r \wedge d[\mathbf{c}\text{Tot}(\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^*A)] \rightarrow d\mathbf{c}\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^rA \rightarrow \mathbf{C}_{\mathbf{KE}_\bullet}^rA,$$

where the first map is the projection from the totalization, and the second is the counit associated to the comonad $\mathbf{C} = d\mathbf{c}\Phi$. These commute with coface maps in the relevant way and determine a derived $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebra map $f: d[\mathbf{c}F_A] \rightarrow A$ (in the sense of Definition 1.26).

By [2, Proposition 1.16], it is sufficient to show that f_0 is an equivalence of symmetric sequences. We can write f_0 , up to equivalence, as a composite

$$d[\mathbf{c}\text{Tot}(\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^*A)] \rightarrow \text{Tot}(d\mathbf{c}\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^*A) \xrightarrow{\sim} \text{Tot}(\mathbf{C}_{\mathbf{KE}_\bullet}^{**+1}A) \xrightarrow{\sim} A,$$

where the second map is induced by the equivalence of Theorem 3.64 and the third is the coaugmentation equivalence associated to the extra codegeneracies in the cosimplicial object $\mathbf{C}_{\mathbf{KE}_\bullet}^{**+1}A$. It remains then to show that the first map is an equivalence; that is, d commutes with the totalization of the cosimplicial object $\Phi\mathbf{C}_{\mathbf{KE}_\bullet}^*A$. The proof of this is virtually identical to that of [2, Corollary 3.16], using the equivalence $\theta: d\mathbf{c}\Phi \xrightarrow{\sim} \mathbf{C}_{\mathbf{KE}_\bullet}$ of Theorem 3.64. □

3.4 Comparison of classifications of polynomial functors

We take this opportunity to describe how the result of Theorem 3.75 is related to other approaches to the classification of polynomial functors from based spaces to spectra. We have already mentioned the work of Dwyer and Rezk in which a polynomial functor $F: \mathcal{Top}_*^f \rightarrow \mathcal{Sp}$ corresponds to the bounded Com-comodule $N[F]$. One half of this equivalence was observed in Proposition 3.15 where we showed that F can be recovered from $N[F]$ via a homotopy coend with $\Sigma^\infty X^{\wedge*}$. In Theorem 3.82 below we show the other half: that any bounded Com-comodule is equivalent to $N[F]$ for some polynomial functor F .

Another classification of polynomial functors from based spaces to spectra follows from the observation that such a functor F is determined, by left Kan extension, by its values on the full subcategory of finite pointed sets. In fact, an n -excisive F is determined by its values on sets of cardinality at most n (not including the basepoint). This fact can be related to the classification described in the previous paragraph via a homotopical version of an equivalence of Pirashvili [19].

Definition 3.77 Let $\Omega_{\leq n}$ denote the category whose objects are the non-empty finite sets of cardinality at most n , and whose morphisms are the surjections. Notice that an n -truncated Com-comodule can be identified with a functor $\Omega_{\leq n} \rightarrow \mathcal{S}p$. Also, let $\Gamma_{\leq n}$ denote the category whose objects are the pointed finite sets of cardinality at most n (not including the basepoint), and whose morphisms are functions that preserve the basepoint. We write Ω and Γ for the corresponding categories of finite sets without restriction on the cardinality.

In [19] Pirashvili showed, among other things, that there is an equivalence of categories

$$[\Omega_{\leq n}, \mathcal{A}b] \simeq [\Gamma_{\leq n}, \mathcal{A}b]_*$$

between the category of functors from $\Omega_{\leq n}$ to the category of abelian groups $\mathcal{A}b$, and the category of pointed functors from $\Gamma_{\leq n}$ to $\mathcal{A}b$ (ie those that send the one-point set to the zero group). In fact, if one includes the empty set as an object of $\Omega_{\leq n}$, then one can remove the pointedness restriction on the right-hand side.

A homotopical version of this equivalence for *contravariant* functors was constructed by Helmstutler in [15] but the covariant version does not appear to be in the literature, so we prove it now. This result includes the case $n = \infty$ where there is no restriction, beyond finiteness, on the cardinality of the sets involved.

Theorem 3.78 For each $1 \leq n \leq \infty$, there is a Quillen equivalence of the form

$$(3.79) \quad \mathbf{L}: \text{Comod}_{\leq n}(\text{Com}) = [\Omega_{\leq n}, \mathcal{S}p] \rightleftarrows [\Gamma_{\leq n}, \mathcal{S}p]_* : \mathbf{R},$$

where $[\Omega_{\leq n}, \mathcal{S}p]$ is the category of functors $\Omega_{\leq n} \rightarrow \mathcal{S}p$, and $[\Gamma_{\leq n}, \mathcal{S}p]_*$ is the category of pointed functors $\Gamma_{\leq n} \rightarrow \mathcal{S}p$ (ie those which take the one-element set to the trivial spectrum). Each of these functor categories has the projective model structure in which weak equivalences and fibrations are detected objectwise, and the Quillen functors are given by

$$\mathbf{L}(\mathbf{N})(J_+) := \mathbf{N} \wedge_{\text{Com}} \widetilde{\Sigma^\infty(J_+)^{\wedge *}}$$

and

$$\mathbf{R}(G) := \text{Map}_{J_+ \in \Gamma_{\leq n}}(\widetilde{\Sigma^\infty(J_+)^{\wedge *}}, G(J_+)).$$

Here $\widetilde{\Sigma^\infty(J_+)^{\wedge *}}$ is as in Definition 3.11.

Proof The same proof works for all n ; we describe it for $n = \infty$.

First note that for each k , the functor $\Gamma \rightarrow \mathcal{S}p$ given by

$$J_+ \mapsto \widetilde{\Sigma^\infty(J_+)^{\wedge k}}$$

is cofibrant in the projective model structure. It follows that \mathbf{R} preserves fibrations and trivial fibrations, so (\mathbf{L}, \mathbf{R}) is a Quillen adjunction.

We now identify the right adjoint \mathbf{R} in terms of taking the cross-effects of a functor $G: \Gamma \rightarrow \mathcal{S}p$, just as in Pirashvili’s original result. First observe that there is an equivalence of spaces, natural in $J_+ \in \Gamma$, of the form

$$(3.80) \quad \text{thocofib}_{I \subseteq K} \Gamma(I_+, J_+) \xrightarrow{\sim} (J_+)^K,$$

given by identifying a pointed function $I_+ \rightarrow J_+$ with an I -tuple in J_+ . (The left-hand side is the total homotopy cofibre of a cube of discrete spaces indexed by subsets $I \subseteq K$ where, for $I \subseteq I'$, the map $\Gamma(I_+, J_+) \rightarrow \Gamma(I'_+, J_+)$ is given by extending a function by the basepoint on $I' - I$. The total homotopy cofibre of this cube is equivalent to the strict total cofibre, ie $(J_+)^K$, since, for each face, the map from its pushout to terminal vertex is an inclusion of discrete spaces, hence a cofibration.)

Applying $\text{Map}_{J_+ \in \Gamma}(-, G(J_+))$ to (3.80) we get an equivalence

$$(3.81) \quad \mathbf{R}(G)(K) \xrightarrow{\sim} \text{thofib}_{I \subseteq K} \text{Map}_{J_+ \in \Gamma}(\Gamma(I_+, J_+), G(J_+)) \cong \text{thofib}_{I \subseteq K} G(I_+),$$

and the right-hand side here is the definition of the K^{th} cross-effect of G evaluated at $(1_+, \dots, 1_+)$.

An important consequence of (3.81) is that the right derived functor of \mathbf{R} preserves homotopy colimits. This follows from the fact that the total homotopy fibre of a cube of spectra can be calculated via the corresponding total homotopy cofibre. Another consequence is that a natural equivalence between functors $\Gamma \rightarrow \mathcal{S}p$ can be detected by \mathbf{R} , in the following sense.

Let $G \rightarrow G'$ be a natural transformation between pointed functors $\Gamma \rightarrow \mathcal{S}p$ such that the induced map $\mathbf{R}(G) \rightarrow \mathbf{R}(G')$ is a weak equivalence of Com-comodules. We prove by induction on $|K|$ that $G(K_+) \rightarrow G'(K_+)$ is a weak equivalence of spectra for any finite set K . For $|K| = 0$ this follows from the fact that G and G' are pointed. For arbitrary K , consider the following diagram of spectra:

$$\begin{array}{ccc} \mathbf{R}(G)(K) & \xrightarrow{\sim} & \text{thofib}_{I \subseteq K} G(I_+) \\ \sim \downarrow & & \downarrow \\ \mathbf{R}(G')(K) & \xrightarrow{\sim} & \text{thofib}_{I \subseteq K} G'(I_+) \end{array}$$

The horizontal maps are the equivalences of (3.81) and the left-hand vertical map is an equivalence by hypothesis. It follows that the right-hand vertical map is an equivalence.

Now note that we also have the following diagram of spectra in which the rows are fibration sequences:

$$\begin{array}{ccccc}
 \text{thofib}_{I \subseteq K} G(I_+) & \longrightarrow & G(K_+) & \longrightarrow & \text{holim}_{I \subsetneq K} G(I_+) \\
 \downarrow \sim & & \downarrow & & \downarrow \sim \\
 \text{thofib}_{I \subseteq K} G'(I_+) & \longrightarrow & G'(K_+) & \longrightarrow & \text{holim}_{I \subsetneq K} G'(I_+)
 \end{array}$$

where the right-hand vertical map is an equivalence by the induction hypothesis. It follows that $G(K_+) \rightarrow G'(K_+)$ is also an equivalence, as desired.

Using these preliminary results, we now turn to showing that the adjunction (\mathbf{L}, \mathbf{R}) is a Quillen equivalence, which we do by showing that the derived unit and counit are both equivalences.

Consider first the derived unit map $\eta_N: N \rightarrow \mathbf{R}N$ for a Com -comodule N . We prove that this is an equivalence by using the cofibrantly generated model structure on $\text{Comod}(\text{Com})$ to reduce to the case where N is the source or target of one of the generating cofibrations, ie is of the form

$$N(I) := E \wedge \underline{\text{Com}}(n, I)$$

for some finite spectrum E and some $n \geq 1$. In this case, we have

$$\mathbf{L}N(J_+) = E \wedge \underline{\text{Com}}(n, I) \wedge_{\text{Com}} \widetilde{\Sigma^\infty(J_+)^{\wedge *}} \cong E \wedge \widetilde{\Sigma^\infty(J_+)^{\wedge n}}$$

and hence

$$\mathbf{R}N(k) = \text{Map}_\Gamma(\widetilde{\Sigma^\infty(J_+)^{\wedge k}}, E \wedge \widetilde{\Sigma^\infty(J_+)^{\wedge n}}).$$

Since E is finite, it is sufficient to show the canonical map

$$\underline{\text{Com}}(n, k) \rightarrow \text{Map}_\Gamma(\widetilde{\Sigma^\infty(J_+)^{\wedge k}}, \widetilde{\Sigma^\infty(J_+)^{\wedge n}}),$$

that takes a surjection $\underline{n} \twoheadrightarrow \underline{k}$ to the map induced by the diagonals on the pointed finite sets (J_+) , is an equivalence. This follows by the argument used to prove that (3.16) is an equivalence, relying on the equivalence (3.80) from above.

Now recall that any Com -comodule N is a retract of a cell complex, which is in turn formed by taking (homotopy) pushouts and sequential colimits, starting from the generating cofibrations. Since these homotopy colimits commute with both \mathbf{L} and \mathbf{R} , we deduce that η_N is an equivalence for any N .

Finally, consider the derived counit map $\epsilon_G: \mathbf{LRG} \rightarrow G$ for pointed $G: \Gamma \rightarrow \mathcal{S}p$. We know that the map

$$\eta_{\mathbf{R}G}: \mathbf{R}G \rightarrow \mathbf{RLRG}$$

is an equivalence, and so by the triangle identity, we deduce that

$$\mathbf{R}(\epsilon_G): \mathbf{RLRG} \rightarrow \mathbf{R}G$$

is an equivalence. But since we have shown that \mathbf{R} reflects equivalences, it follows that ϵ_G is an equivalence too. This completes the proof that (\mathbf{L}, \mathbf{R}) is a Quillen equivalence. \square

Theorem 3.82 *There is a diagram of functors, as follows, which commutes up to natural equivalence and in which each functor determines an equivalence of homotopy categories:*

$$(3.83) \quad \begin{array}{ccc} & \text{Coalg}_{\leq n}(\mathbf{C}_{\text{KE}\bullet}) & \\ \text{hocolim}_L \tilde{\wedge}_{E_L} \mathbb{B}_{E_L} \nearrow & & \nwarrow d \\ \text{Comod}_{\leq n}(\text{Com}) & \xrightarrow{-\wedge_{\text{Com}} \widetilde{\Sigma^{\infty} X^{\wedge *}}} & [\mathcal{T}op_*^f, \mathcal{S}p]_{*,n\text{-exc}} \\ \searrow \mathbf{L} & & \nearrow \text{LKan} \\ & [\Gamma_{\leq n}, \mathcal{S}p]_* & \end{array}$$

In this diagram,

- $\text{Coalg}_{\leq n}(\mathbf{C}_{\text{KE}\bullet})$ is the category of n -truncated $\mathbf{C}_{\text{KE}\bullet}$ -coalgebras;
- $\text{Comod}_{\leq n}(\text{Com})$ is the category of n -truncated Com -comodules;
- $[\mathcal{T}op_*^f, \mathcal{S}p]_{*,n\text{-exc}}$ is the category of n -excisive pointed simplicial functors $\mathcal{T}op_*^f \rightarrow \mathcal{S}p$;
- $[\Gamma_{\leq n}, \mathcal{S}p]_*$ is as in Theorem 3.78;

and

- \mathbf{L} is the left adjoint of the Quillen equivalence in Theorem 3.78;
- LKan denotes left Kan extension along the inclusion $\Gamma_{\leq n} \rightarrow \mathcal{T}op_*^f$;
- d is as in Definition 3.53.

Proof Theorem 3.75 implies that d determines an equivalence of homotopy theories. The top triangle commutes up to natural equivalence by the construction of d . We turn then to the horizontal functor in diagram (3.83).

We have already shown in Proposition 3.15 that the evaluation map

$$\text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge *}}, FY) \widetilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge *}} \rightarrow F(X)$$

is an equivalence when F is polynomial. We now show that the unit map

$$N(r) \rightarrow \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge r}}, N \widetilde{\wedge}_{\text{Com}} \widetilde{\Sigma^\infty Y^{\wedge *}})$$

is an equivalence when N is a cofibrant Com -comodule. To see this, note that by Corollary 3.14, it is sufficient to show that, for each r , the map

$$N(r) \rightarrow N \widetilde{\wedge}_{\text{Com}} \text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge r}}, \widetilde{\Sigma^\infty Y^{\wedge *}})$$

induced by the identity on $\widetilde{\Sigma^\infty Y^{\wedge *}}$ is an equivalence. But by (3.16) there is an equivalence

$$\text{Nat}_{Y \in \mathcal{T}op_*^f}(\widetilde{\Sigma^\infty Y^{\wedge r}}, \widetilde{\Sigma^\infty Y^{\wedge *}}) \simeq \underline{\text{Com}}(*, r),$$

and the map

$$N(r) \rightarrow N \widetilde{\wedge}_{\text{Com}} \underline{\text{Com}}(*, r)$$

is an equivalence because the right-hand side is a simplicial object with extra degeneracies. It now follows that the horizontal map in (3.83) induces an equivalence of homotopy theories. It then also follows that the top-left map induces an equivalence.

By Theorem 3.78, \mathbf{L} also induces an equivalence, so it only remains to show that the bottom triangle commutes up to natural equivalence. Notice that we have a commutative diagram

$$\begin{array}{ccc} \text{Comod}_{\leq n}(\text{Com}) & \xrightarrow{-\wedge_{\text{Com}} \widetilde{\Sigma^\infty X^{\wedge *}}} & [\mathcal{T}op_*^f, \mathcal{S}p]_{*,n-\text{exc}} \\ & \searrow & \swarrow \text{res} \\ & & [\Gamma_{\leq n}, \mathcal{S}p]_* \\ & \swarrow & \nwarrow \\ & & \end{array}$$

$-\wedge_{\text{Com}} \widetilde{\Sigma^\infty (J_+)^{\wedge *}}$

where res is restriction to the subcategory $\Gamma_{\leq n} \subseteq \mathcal{T}op_*^f$. Since the other two functors induce equivalences, it follows that res does too. Since $\mathbf{L}\text{Kan}$ is left adjoint to res , it induces the inverse equivalence to res on the homotopy category and hence the bottom triangle in the original diagram commutes up to natural equivalence. \square

3.5 Classification of analytic functors from based spaces to spectra

One of the main advantages of using $\mathbf{C}_{\text{KE}_\bullet}$ -coalgebras to classify polynomial functors from based spaces to spectra (over the other categories appearing in Theorem 3.82) is that this approach generalizes, to some extent, to non-polynomial functors.

Here we consider functors that are “analytic at the one-point space $*$ ” in the sense described below. This condition is weaker than Goodwillie’s notion of analyticity since it only concerns the values of a functor in a “neighbourhood” of $*$, that is, only on highly connected spaces.

Definition 3.84 We say that a homotopy functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ satisfies condition $E_n^*(c, \kappa)$ if for any strongly cocartesian $(n + 1)$ -cube $\mathcal{X}: \mathcal{P}(S) \rightarrow \mathcal{T}op_*^f$ such that $\mathcal{X}(\emptyset) \rightarrow \mathcal{X}(s)$ is a k_s -connected map between κ -connected spaces for any $s \in S$, the cube $F(\mathcal{X})$ is $(-c + \sum k_s)$ -cartesian.

We say that $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ is ρ -analytic at $*$ if there is a constant q such that F satisfies

$$E_n^*(n\rho - q, \rho + 1)$$

for all $n \geq 1$. We say that F is analytic at $*$ if it is ρ -analytic at $*$ for some number ρ .

Remark 3.85 These conditions should be compared to those of Goodwillie [12, Definition 4.1]. Since a map between κ -connected spaces is κ -connected, it is easy to see that Goodwillie’s condition $E_n(c, \kappa)$ implies our $E_n^*(c, \kappa)$. It follows that a functor that is ρ -analytic in the sense of Goodwillie is, in particular, ρ -analytic at $*$. More generally, we can say that F is ρ -analytic at X , for some based space X , if it satisfies the analyticity condition on cubes of spaces that are $(\rho + 1)$ -connected over X .

Definition 3.86 Let A be a symmetric sequence of spectra. We say that A is ρ -analytic if there is a constant c such that $A(n)$ is $(-\rho n + c)$ -connected for all n . We say A is analytic if it is ρ -analytic for some ρ .

Lemma 3.87 Let $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ be a functor that is ρ -analytic at $*$. Then the symmetric sequence $\partial_* F$ is ρ -analytic in the sense of Definition 3.86.

Proof For each $n \geq 2$, we apply the condition $E_{n-1}^*((n-1)\rho - q, \rho + 1)$ to the strongly cocartesian n -cube with initial maps $* \rightarrow S^L$ for $L \geq \rho + 2$. These maps are $(L - 1)$ -connected and so we deduce that the cube $T \mapsto F(\bigvee_T S^L)$ is $(q - (n-1)\rho + n(L-1))$ -cartesian. It follows that the total homotopy cofibre of this cube of spectra, which is

precisely the n^{th} co-cross-effect $\text{cr}^n F(S^L, \dots, S^L)$, is $(q - (n - 1)\rho + n(L - 1) + n - 1)$ -connected.

Since the cross-effect is equivalent to the co-cross-effect, we deduce that the desuspended cross-effect

$$\Sigma^{-nL} \text{cr}_n F(S^L, \dots, S^L)$$

is $(-n\rho + \rho + q - 1)$ -connected. Taking the homotopy colimit as $L \rightarrow \infty$, we deduce that $\partial_n F$ has this same connectivity, and so the symmetric sequence $\partial_* F$ is ρ -analytic, with the constant c in Definition 3.86 equal to $\rho + q - 1$. \square

Lemma 3.88 *Consider a sequence of functors*

$$(3.89) \quad \dots \rightarrow F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1$$

that looks like a Taylor tower in the sense that it induces equivalences $P_n(F_{n+1}) \xrightarrow{\sim} P_n(F_n) \simeq F_n$ for each n . Suppose that the symmetric sequence $\{\partial_n(F_n)\}$ is ρ -analytic for some $\rho \geq 0$. Then

$$F := \text{holim}_n F_n$$

is ρ -analytic at $*$, and the sequence (3.89) is equivalent to the Taylor tower of F .

Proof Let us write D_n for the homotopy fibre of the natural transformation $F_n \rightarrow F_{n-1}$. Then we have

$$D_n(X) \simeq (A_n \wedge X^{\wedge n})_{h\Sigma_n},$$

where $A_n := \partial_n(F_n)$. Choose a constant c such that A_n is $(-\rho n + c)$ -connected for all n .

To show that F is ρ -analytic at $*$, consider a strongly cocartesian $(n + 1)$ -cube \mathcal{X} of $(\rho + 1)$ -connected based spaces with each initial map $\mathcal{X}(\emptyset) \rightarrow \mathcal{X}(s)$ being k_s -connected. In particular, then, the cube $(\Sigma^{-\rho} \Sigma^\infty \mathcal{X})$ is a strongly cocartesian cube of 1-connected spectra with each initial map being $(-\rho + k_s)$ -connected. We can apply a version of [12, Example 4.4] for cubes of spectra to deduce that, for each m , the cube $(\Sigma^{-\rho m} \Sigma^\infty X^{\wedge m})$ is $(-\rho(n + 1) + \sum k_s)$ -cartesian.

Since A_m is $(-\rho m + c)$ -connected, it therefore follows that the cube $D_m(\mathcal{X})$ is

$$\left(\rho m - \rho m + c - \rho(n + 1) + \sum k_s\right) = \left(-\rho n - \rho + c + \sum k_s\right)$$

-cartesian. By induction on m , using the fibre sequences $D_m \rightarrow F_m \rightarrow F_{m-1}$ we deduce that each cube $F_m(\mathcal{X})$ is $(-\rho n - \rho + c + \sum k_s)$ -cartesian, and so the homotopy limit $F(\mathcal{X})$ is $(-\rho n - \rho + c - 1 + \sum k_s)$ -cartesian. This verifies that F is ρ -analytic at $*$ with the number q in Definition 3.84 equal to $c - \rho - 1$.

Now suppose that X is a k -connected based space. Then $D_n(X) = (A_n \wedge X^{\wedge n})_{h\Sigma_n}$ is $[(k - \rho)n + c]$ -connected. As long as $k \geq \rho$, it follows that the map $F(X) \rightarrow F_n(X)$ is $[(k - \rho)(n + 1) + c - 1]$ -connected, for each n . This means that F and F_n agree to order n in the sense of Goodwillie [13, Definition 1.2] from which it follows by [13, Proposition 1.6] that $P_n F \simeq P_n F_n \simeq F_n$. \square

Corollary 3.90 *If $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ is analytic at $*$, then $P_\infty F := \text{holim}_n P_n F$ is analytic at $*$ and the map $F \rightarrow P_\infty F$ induces an equivalence of Taylor towers.*

Definition 3.91 Let us say that a functor $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ is *strongly analytic at $*$* if F is analytic at $*$ and the map $F \rightarrow P_\infty F := \text{holim } P_n F$ is an equivalence. It follows from Corollary 3.90 that, for any $F: \mathcal{T}op_*^f \rightarrow \mathcal{S}p$ that is analytic at $*$, the functor $P_\infty F$ is strongly analytic at $*$, and the map $F \rightarrow P_\infty F$ determines an equivalence of Taylor towers. We can therefore think of the category of functors strongly analytic at $*$ as a model for the category of all functors that are analytic at $*$ up to equivalence of Taylor towers.

We write $[\mathcal{T}op_*^f, \mathcal{S}p]_{*,\text{an}(*)}^h$ for the subcategory of the homotopy category $[\mathcal{T}op_*^f, \mathcal{S}p]_*^h$ consisting of the functors that are strongly analytic at $*$.

Now consider the restriction of $d: [\mathcal{T}op_*^f, \mathcal{S}p]_* \rightarrow \text{Coalg}(\mathbf{C}_{KE_\bullet})$ to strongly analytic functors. Firstly, the argument of Theorem 3.75 implies that d determines weak equivalences

$$\text{Nat}_{\mathcal{T}op_*^f}(F, G) \xrightarrow{\sim} \widetilde{\text{Map}}_{\mathbf{C}_{KE_\bullet}}(d[F], d[G])$$

for any $F, G \in [\mathcal{T}op_*^f, \mathcal{S}p]_*$ such that $G \simeq P_\infty G$. It follows that d is a fully faithful embedding of $[\mathcal{T}op_*^f, \mathcal{S}p]_{*,\text{an}(*)}^h$ into the homotopy category of \mathbf{C}_{KE_\bullet} -coalgebras. We have not been able to explicitly identify the image of this embedding. Instead, however, by considering the “pro-truncated” version of the homotopy theory of \mathbf{C}_{KE_\bullet} -coalgebras, as constructed in Definition 1.29, we have the following result.

Theorem 3.92 *The functor d sets up an equivalence*

$$[\mathcal{T}op_*^f, \mathcal{S}p]_{*,\text{an}(*)}^h \simeq \text{Coalg}_{\text{an}}^t(\mathbf{C}_{KE_\bullet})^h$$

between the homotopy category of functors $\mathcal{T}op_^f \rightarrow \mathcal{S}p$ that are strongly analytic at $*$, and the pro-truncated homotopy category of analytic \mathbf{C}_{KE_\bullet} -coalgebras. Moreover, for cofibrant $F, G \in [\mathcal{T}op_*^f, \mathcal{S}p]_{*,\text{an}(*)}$, we have equivalences*

$$\text{Nat}_{\mathcal{T}op_*^f}(F, G) \xrightarrow{\sim} \widetilde{\text{Map}}_{\mathbf{C}_{KE_\bullet}}(d[F], d[G]) \simeq \widetilde{\text{Map}}^t_{\mathbf{C}_{KE_\bullet}}(d[F], d[G]).$$

Proof For two functors F, G strongly analytic at $*$ we have already noted that there is an equivalence of spectra

$$\text{Nat}_{\mathcal{J}op_*^f}(F, G) \xrightarrow{\sim} \widetilde{\text{Map}}_{\mathbf{C}_{KE_\bullet}}(d[F], d[G]).$$

Since the map

$$\text{Nat}_{\mathcal{J}op_*^f}(F, G) \rightarrow \text{holim}_n \text{Nat}_{\mathcal{J}op_*^f}(F, P_n G)$$

is an equivalence, so too is the map

$$\widetilde{\text{Map}}_{\mathbf{C}_{KE_\bullet}}(d[F], d[G]) \rightarrow \text{holim}_n \widetilde{\text{Map}}_{\mathbf{C}_{KE_\bullet}}(d[F], d[G]_{\leq n}) \simeq \widetilde{\text{Map}}^t_{\mathbf{C}_{KE_\bullet}}(d[F], d[G]).$$

So, between \mathbf{C}_{KE_\bullet} -coalgebras of the form $d[G]$ where G is strongly analytic at $*$, pro-truncated mapping spectra are equivalent to ordinary mapping spectra. In particular, then, the functor d is an embedding of the homotopy category of strongly analytic functors into the pro-truncated homotopy category of analytic \mathbf{C}_{KE_\bullet} -coalgebras.

To see that this embedding is an equivalence, let A be an arbitrary analytic \mathbf{C}_{KE_\bullet} -coalgebra. Let F be the functor given by

$$F := \text{holim}_n F_{A_{\leq n}},$$

where $F_{A_{\leq n}}$ is as in the proof of Theorem 3.64. Then F is strongly analytic at $*$ by Lemma 3.88.

We claim that there is a derived equivalence, in the pro-truncated sense of Definition 1.27, of \mathbf{C}_{KE_\bullet} -coalgebras $f: d[\mathbf{c}F] \rightarrow A$. This consists of the maps

$$f_{k,n}: \Delta_+^k \wedge d[\mathbf{c} \text{holim}_n F_{A_{\leq n}}] \rightarrow \Delta_+^k \wedge d[\mathbf{c}F_{A_{\leq n}}] \rightarrow \mathbf{C}_{KE_\bullet}^k(A_{\leq n})$$

constructed by projecting from the homotopy limit, together with the corresponding maps from the proof of Theorem 3.64. It is sufficient then to show that the composite

$$f_0: d[\mathbf{c} \text{holim}_n F_{A_{\leq n}}] \rightarrow \text{holim}_n d[\mathbf{c}F_{A_{\leq n}}] \rightarrow \text{holim}_n A_{\leq n} \simeq A$$

is an equivalence. The first map is an equivalence by Lemma 3.88, and the second map is an equivalence by Theorem 3.64, so the required result follows. \square

Examples 3.93 Under the equivalence of Theorem 3.92, we can identify the analytic functors with split Taylor tower with those \mathbf{C}_{KE_\bullet} -coalgebras A whose structure is induced via the map of comonads (of symmetric sequences)

$$I_{[\Sigma, \mathcal{J}p]} = \mathbf{C}_1 = \mathbf{C}_{KE_0} \rightarrow \text{hocolim } \mathbf{C}_{KE_L} = \mathbf{C}_{KE_\bullet}.$$

More generally, it is possible to characterize those functors whose corresponding $\mathbf{C}_{\mathbf{KE}_\bullet}$ -coalgebra is induced via the map of comonads

$$\mathbf{C}_{\mathbf{KE}_L} \rightarrow \mathbf{C}_{\mathbf{KE}_\bullet}$$

for some fixed L , ie those functors whose derivatives possess a \mathbf{KE}_L -module structure from which the Taylor tower can be constructed. We leave this analysis to a future paper.

4 Functors from spectra to spectra

We now turn to pointed simplicial functors $\mathcal{S}p^f \rightarrow \mathcal{S}p$, where $\mathcal{S}p^f$ is the category of finite cell spectra. We denote the category of such functors by $[\mathcal{S}p^f, \mathcal{S}p]_*$. As with the spaces to spectra case there is an inverse sequence of operads that acts on the sequence of partially stabilized cross-effects of a functor $F \in [\mathcal{S}p^f, \mathcal{S}p]_*$. We start by constructing this sequence.

4.1 Desuspensions of the commutative operad

Underlying the sequence of operads we are interested in is a notion of “desuspension” for operads of spectra. This uses a certain cooperad \mathbb{S} , in the category of based spaces, whose terms are homeomorphic to spheres. The structure maps for this cooperad are homeomorphisms and so \mathbb{S} is also an operad in a canonical way. This operad is isomorphic to the operad S_∞ of [3].

Definition 4.1 For a nonempty finite set I we set

$$\mathbb{R}_0^I := \{t \in \mathbb{R}^I \mid \min_{i \in I} t_i = 0\}.$$

We then write

$$\mathbb{S}(I) := (\mathbb{R}_0^I)^+.$$

This is the based space given by the one-point compactification of \mathbb{R}_0^I . Note that $\mathbb{S}(I)$ is homeomorphic to $S^{|I|-1}$. The permutation action of the symmetric group Σ_I on \mathbb{R}^I restricts to an action on \mathbb{R}_0^I and hence on $\mathbb{S}(I)$. This makes \mathbb{S} into a symmetric sequence of based spaces.

Definition 4.2 We construct a cooperad structure on the symmetric sequence \mathbb{S} as follows. It arises from a cooperad structure on the unbased spaces \mathbb{R}_0^n of Definition 4.1.

For a surjection of nonempty finite sets $\alpha: I \twoheadrightarrow J$, writing $I_j := \alpha^{-1}(j)$, there is a continuous map

$$d_\alpha: \mathbb{R}_0^I \rightarrow \mathbb{R}_0^J \times \prod_{j \in J} \mathbb{R}_0^{I_j}$$

given by

$$d_\alpha((t_i)_{i \in I}) := ((v_j)_{j \in J}, \{(u_{j,i})_{i \in I_j}\}_{j \in J}),$$

where

$$v_j := \min_{i \in I_j} t_i$$

and

$$u_{j,i} := t_i - v_j \quad \text{for } i \in I_j.$$

The condition that the minimum of the t_i is zero implies that the minimum of the v_j is zero. For each $j \in J$, we clearly have $\min_{i \in I_j} u_{j,i} = 0$.

The functions d_α extend to the one-point compactifications to give maps

$$d_\alpha: \mathbb{S}(I) \rightarrow \mathbb{S}(J) \wedge \bigwedge_{j \in J} \mathbb{S}(I_j).$$

Example 4.3 If $\alpha: \{1, 2, 3\} \rightarrow \{1, 2\}$ is the map given by $\alpha(1) = 1$, $\alpha(2) = 2$, $\alpha(3) = 2$, then

$$d_\alpha(t_1, t_2, t_3) = [(t_1, m), (0), (t_2 - m, t_3 - m)],$$

where $m = \min\{t_2, t_3\}$.

Lemma 4.4 Each map d_α of Definition 4.2 is a homeomorphism, and together they make \mathbb{S} into a reduced cooperad of based spaces. (The inverses d_α^{-1} thus make \mathbb{S} into a reduced operad.)

Proof We define an inverse to d_α on non-basepoints by

$$d_\alpha^{-1}((v_j)_{j \in J}, \{(u_{j,i})_{i \in I_j}\}_{j \in J}) := (t_i)_{i \in I},$$

where

$$t_i := u_{\alpha(i),i} + v_{\alpha(i)}.$$

The map d_α^{-1} is continuous and inverse to d_α . We leave the reader to check the unit and associativity conditions for \mathbb{S} to be a cooperad. \square

Definition 4.5 Let A be a symmetric sequence of spectra. We define the *desuspension* of A to be the symmetric sequence $S^{-1}A$ given by

$$(S^{-1}A)(n) := \text{Map}(\mathbb{S}(n), A(n))$$

with the diagonal Σ_n -action. Here $\text{Map}(-, -)$ denotes the cotensoring of $\mathcal{S}p$ over based spaces.

Now suppose that P is an operad of spectra. Then we make $S^{-1}P$ into an operad of spectra by convolving the operad structure on P with the cooperad structure on \mathbb{S} in the following way. For a surjection $\alpha: I \twoheadrightarrow J$ we have structure map

$$\begin{aligned} \text{Map}(\mathbb{S}(J), P(J)) \wedge \bigwedge_{j \in J} \text{Map}(\mathbb{S}(I_j), P(I_j)) & \rightarrow \text{Map}(\mathbb{S}(J) \wedge \bigwedge_{j \in J} \mathbb{S}(I_j), P(J) \wedge \bigwedge_{j \in J} P(I_j)) \\ & \rightarrow \text{Map}(\mathbb{S}(I), P(I)), \end{aligned}$$

where the first map is the canonical smash product of mapping spectra and the second employs the cooperad structure on \mathbb{S} and the operad structure on P . We refer to $S^{-1}P$ with this structure as the (*operadic*) *desuspension* of P .

We can iterate the desuspension process to get operads $S^{-L}P$ given by

$$S^{-L}P(I) \cong \text{Map}(\mathbb{S}(I)^{\wedge L}, P(I)).$$

Definition 4.6 For each nonempty finite set I , we define a map

$$\eta_I: S^0 \rightarrow \mathbb{S}(I)$$

by sending the non-basepoint in S^0 to the point in $\mathbb{S}(I)$ given by the origin $0 \in \mathbb{R}_0^I$.

Lemma 4.7 *The maps η_I together form a morphism of cooperads (of based spaces)*

$$\eta: \text{Com}_+^c \rightarrow \mathbb{S},$$

where Com_+^c is the commutative cooperad in based spaces (given by S^0 in each term with trivial Σ_n -actions and homeomorphisms as structure maps).

Proof It is sufficient to check that for each surjection $\alpha: I \twoheadrightarrow J$, we have

$$d_\alpha(0) = (0, \{0\}_{j \in J}),$$

which follows from the definition of d_α . □

Definition 4.8 For any operad P of spectra, the map η of Lemma 4.7 induces a natural map of operads

$$\eta: S^{-1}P \rightarrow P.$$

Iterating this construction, we get, for each operad P of spectra, an inverse sequence

$$S^{-\bullet}P: \dots \rightarrow S^{-2}P \rightarrow S^{-1}P \rightarrow P.$$

The example we are concerned with in this paper is the sequence of desuspensions of $P = \text{Com}$, the commutative operad of spectra.

Definition 4.9 We write S^{-L} for a (functorial) cofibrant replacement (in the projective model structure on operads) of $S^{-L} \text{Com}$. Applying this cofibrant replacement to the sequence in Definition 4.8 gives us an inverse sequence of operads

$$S_{\bullet}: \dots \rightarrow S^{-2} \rightarrow S^{-1} \rightarrow S^0 \xrightarrow{\simeq} \text{Com}.$$

Associated to this sequence, by the construction of Definition 2.6, is a comonad $C_{S_{\bullet}}$ on the category of symmetric sequences given by

$$\begin{aligned} C_{S_{\bullet}}(A)(k) &:= \text{hocolim}_L C_{S^{-L}}(A)(k) \\ &= \text{hocolim}_L \prod_n \left[\prod_{\underline{n} \rightarrow \underline{k}} \text{Map}(S^{-L}(n_1) \wedge \dots \wedge S^{-L}(n_k), A(n)) \right]^{\Sigma_n}. \end{aligned}$$

4.2 Models for the derivatives of a functor from spectra to spectra

Our next task is to construct models for the partially stabilized cross-effects of a pointed simplicial functor $F \in [\mathcal{S}p^f, \mathcal{S}p]_*$ that admit the structure of a S^{-L} -module. From these we obtain models for the derivatives of F that form a $C_{S_{\bullet}}$ -coalgebra. We focus first on the representable functors.

Definition 4.10 Let $X \in \mathcal{S}p^f$. The functor represented by X is the pointed simplicial functor $R_X: \mathcal{S}p^f \rightarrow \mathcal{S}p$ given by

$$R_X(Y) := \Sigma^{\infty} \text{Hom}_{\mathcal{S}p}(X, Y).$$

A significant difference from the case $[\mathcal{T}op_*^f, \mathcal{S}p]_*$ is that there are simple models for the cross-effects of the representable functors from spectra to spectra.

Lemma 4.11 The n^{th} cross-effect of R_X is given by

$$\text{cr}_n R_X(Y_1, \dots, Y_n) \simeq \Sigma^{\infty} \text{Hom}_{\mathcal{S}p}(X, Y_1) \wedge \dots \wedge \text{Hom}_{\mathcal{S}p}(X, Y_n).$$

Proof Finite coproducts of spectra are equivalent to the corresponding products. Therefore, the co-cross-effect (and hence the desired cross-effect) is given by the total homotopy cofibre

$$\text{thocofib}_{S \subseteq \underline{n}} \left\{ \Sigma^\infty \text{Hom}_{\mathcal{G}p} \left(X, \prod_{i \in S} Y_i \right) \right\},$$

which is equivalent to

$$\Sigma^\infty \text{thocofib}_{S \subseteq \underline{n}} \left\{ \prod_{i \in S} \text{Hom}_{\mathcal{G}p}(X, Y_i) \right\}.$$

The total homotopy cofibre of the cube of simplicial sets whose terms are $\prod_{i \in S} Z_i$ is equivalent to the smash product $Z_1 \wedge \cdots \wedge Z_n$. We thus obtain the required formula for the cross-effect. \square

Corollary 4.12 *The partially stabilized cross-effects of R_X are therefore given by*

$$\Sigma^{-nL} \text{cr}_n(R_X)(S^L, \dots, S^L) \simeq \Sigma^{-nL} \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge n},$$

with the stabilization maps of (1.7) given by the maps

$$\text{Map}(S^{nL}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge n}) \rightarrow \text{Map}(S^{n(L+1)}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^{L+1})^{\wedge n})$$

induced by the n -fold smash power of the canonical map

$$S^1 \wedge \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L) \rightarrow \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^{L+1}).$$

Proof The models for $\text{cr}_n R_X$ in Lemma 4.11 are pointed simplicial in each variable and so, by Lemma 1.9, the maps (1.7) are given by the relevant tensoring maps, which are as shown. \square

We now construct models for these partially stabilized cross-effects that admit actions of the operads S^{-L} . The key is to identify the desuspension spheres with terms of the cooperad S .

Definition 4.13 For $X \in \mathcal{G}p^f$ and a non-negative integer L , we define a symmetric sequence $\underline{d}^L[R_X]$ by

$$\underline{d}^L[R_X](I) := \text{Map}((S^1 \wedge \mathbb{S}(I))^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge I}).$$

We then make the symmetric sequence $\underline{d}^L[R_X]$ into a right S^{-L} Com-module (and hence a S^{-L} -module via pullback) by combining the cooperad structure on \mathbb{S} with the right Com-module structure on $\Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge *}$ described in Definition 3.11

(ie that given by the diagonal on the space $\text{Hom}_{\mathcal{G}p}(X, S^L)$). Explicitly, for a surjection $\alpha: I \twoheadrightarrow J$, we have a map

$$\nu_\alpha: \underline{d}^L[R_X](J) \wedge \bigwedge_{j \in J} S^{-L}(I_j) \rightarrow \underline{d}^L[R_X](I)$$

given by the composite

$$\begin{aligned} & \text{Map}((S^1)^{\wedge L} \wedge \mathbb{S}(J)^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge J}) \wedge \bigwedge_{j \in J} \text{Map}(\mathbb{S}(I_j)^{\wedge L}, \text{Com}(I_j)) \\ & \rightarrow \text{Map}\left((S^1)^{\wedge L} \wedge \mathbb{S}(J)^{\wedge L} \wedge \bigwedge_{j \in J} \mathbb{S}(I_j)^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge J} \wedge \bigwedge_{j \in J} \text{Com}(I_j)\right) \\ & \rightarrow \text{Map}((S^1)^{\wedge L} \wedge \mathbb{S}(I)^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge I}), \end{aligned}$$

where the first map is the canonical smash product of mapping spectra, and the second is induced by the L -fold smash power of the cooperad structure map

$$d_\alpha: \mathbb{S}(I) \rightarrow \mathbb{S}(J) \wedge \bigwedge_{j \in J} \mathbb{S}(I_j)$$

and the right Com-module structure map

$$\Delta_\alpha: \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge J} \rightarrow \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge I}$$

associated to α .

Lemma 4.14 *We have equivalences of symmetric sequences (natural in X)*

$$\Sigma^{-*L} \text{cr}_*(R_X)(S^L, \dots, S^L) \simeq \underline{d}^L[R_X].$$

Proof This follows from Corollary 4.12. □

Our next task is to provide models for the maps between the desuspended cross-effects that respect these operad actions.

Definition 4.15 There are homeomorphisms

$$h_I: \mathbb{R} \times \mathbb{R}_0^I \xrightarrow{\cong} \mathbb{R}^I$$

given by

$$(t, (t_i)_{i \in I}) \mapsto (t + t_i)_{i \in I},$$

which extend to homeomorphisms (which we also call h_I)

$$S^1 \wedge \mathbb{S}(I) \xrightarrow{\cong} (S^1)^{\wedge I},$$

where we identify $(S^1)^{\wedge I}$ with the one-point compactification of \mathbb{R}^I . The following diagram of homeomorphisms then commutes:

$$(4.16) \quad \begin{array}{ccc} \mathbb{R} \times \mathbb{R}_0^J \times \prod_{j \in J} \mathbb{R}_0^{I_j} & \xrightarrow{d_\alpha^{-1}} & \mathbb{R} \times \mathbb{R}_0^I \\ \downarrow h_J & & \downarrow h_I \\ \mathbb{R}^J \times \prod_{j \in J} \mathbb{R}_0^{I_j} & \xrightarrow{\prod h_{I_j}} & \mathbb{R}^I \end{array}$$

Definition 4.17 We construct a map of symmetric sequences

$$\underline{m}_L: \underline{d}^L[R_X] \rightarrow \underline{d}^{L+1}[R_X]$$

as follows. For a positive integer n the required map $\underline{m}_L(n)$,

$$\begin{aligned} \text{Map}((S^1 \wedge \mathbb{S}(n))^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge n}) \\ \rightarrow \text{Map}((S^1 \wedge \mathbb{S}(n))^{\wedge L+1}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^{L+1})^{\wedge n}), \end{aligned}$$

is adjoint to a composite of the homeomorphism given by applying h_n to the final copy of $S^1 \wedge \mathbb{S}(n)$,

$$(S^1 \wedge \mathbb{S}(n))^{\wedge(L+1)} \xrightarrow{\cong} (S^1 \wedge \mathbb{S}(n))^{\wedge L} \wedge S^n,$$

and the n -fold smash power of the canonical map

$$S^1 \wedge \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L) \rightarrow \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^{L+1}).$$

Lemma 4.18 *The homotopy colimit of the sequence*

$$\underline{d}^0[R_X] \xrightarrow{\underline{m}_0} \underline{d}^1[R_X] \xrightarrow{\underline{m}_1} \underline{d}^2[R_X] \xrightarrow{\underline{m}_2} \dots$$

is naturally (in X) equivalent to the symmetric sequence $\partial_*(R_X)$ of derivatives of the representable functor R_X .

Proof It follows from the definition of \underline{m}_L , and Corollary 4.12, that \underline{m}_L models the stabilization map

$$\Sigma^{-*L} \text{cr}_*(R_X)(S^L, \dots, S^L) \rightarrow \Sigma^{-*(L+1)} \text{cr}_*(R_X)(S^{L+1}, \dots, S^{L+1})$$

of (1.7). The claim follows. □

Unfortunately the map m_L is not a morphism of $S^{-(L+1)}$ -modules. (We explain why not in Section 5; see diagram (5.2) and after.) We therefore cannot apply our general theory to produce a \mathbf{C}_{S_\bullet} -coalgebra structure on the homotopy colimit in Lemma 4.18.

However, we show in Section 5 that it is possible to extend m_L to a morphism of $S^{-(L+1)}$ -modules of the form

$$m'_L: B(\underline{d}^L[R_X], S^{-(L+1)}, S^{-(L+1)}) \rightarrow \underline{d}^{L+1}[R_X],$$

where the left-hand side is the standard bar resolution of $\underline{d}^L[R_X]$ as a $S^{-(L+1)}$ -module.

Recall that for any operad P and P -module M , the bar resolution is a weak equivalence of P -modules

$$p: B(M, P, P) \xrightarrow{\sim} M.$$

The map p has a section

$$s: M \xrightarrow{\sim} B(M, P, P)$$

that is a morphism of symmetric sequences, but not of P -modules. The map s is given by the composite

$$M \xrightarrow{\eta} M \circ P \rightarrow B(M, P, P)$$

of the unit map for the operad P with the inclusion of the 0-simplices into the bar construction. We then have the following proposition.

Proposition 4.19 *There is a natural morphism of $S^{-(L+1)}$ -modules*

$$m'_L: B(\underline{d}^L[R_X], S^{-(L+1)}, S^{-(L+1)}) \rightarrow \underline{d}^{L+1}[R_X]$$

such that the following diagram commutes:

$$\begin{array}{ccc} B(\underline{d}^L[R_X], S^{-(L+1)}, S^{-(L+1)}) & & \\ \uparrow s \sim & \searrow m'_L & \\ \underline{d}^L[R_X] & \xrightarrow{m_L} & \underline{d}^{L+1}[R_X] \end{array}$$

The construction of the map m'_L is quite involved; we defer the proof of Proposition 4.19 to Section 5.

We now use the maps m'_L to construct a sequence of S^{-L} -modules whose actions do commute with the operad maps $S^{-(L+1)} \rightarrow S^{-L}$.

Definition 4.20 For $X \in \mathcal{S}p^f$, we construct the diagram

$$\begin{array}{ccccccc}
 & & & & & & B(\underline{d}^2[R_X], S^{-3}, S^{-3}) \xrightarrow{\underline{m}'_2} \dots \\
 & & & & & & \downarrow \sim p \\
 & & & & & & B(\underline{d}^1[R_X], S^{-2}, S^{-2}) \xrightarrow{\underline{m}'_1} \underline{d}^2[R_X] \\
 & & & & & & \downarrow \sim p \\
 & & & & & & B(\underline{d}^0[R_X], S^{-1}, S^{-1}) \xrightarrow{\underline{m}'_0} \underline{d}^1[R_X] \\
 & & & & & & \downarrow \sim p \\
 & & & & & & \underline{d}^0[R_X] \xrightarrow{\underline{m}_0} \underline{d}^1[R_X] \xrightarrow{\underline{m}_1} \underline{d}^2[R_X] \xrightarrow{\underline{m}_2} \dots
 \end{array}$$

so that each rectangle is a homotopy pushout in the category of S^{-L} -modules for respective values of L . We thus obtain a sequence

$$\underline{d}^0[R_X] = \underline{d}^0[R_X] \xrightarrow{\underline{m}_0} \underline{d}^1[R_X] \xrightarrow{\underline{m}_1} \underline{d}^2[R_X] \xrightarrow{\underline{m}_2} \dots$$

such that $\underline{d}^L[R_X]$ is an S^{-L} -module, and such that the map

$$\underline{m}_L: \underline{d}^L[R_X] \rightarrow \underline{d}^{L+1}[R_X]$$

is a morphism of $S^{-(L+1)}$ -modules that is equivalent (in the homotopy category) to \underline{m}_L . In other words $\underline{d}^\bullet[R_X]$ is a module over the pro-operad S_\bullet .

Finally, by applying the construction of Definition 2.12, we obtain a diagram

$$(4.21) \quad \begin{array}{ccccccc}
 \underline{d}^0[R_X] & \xrightarrow{\underline{m}_0} & \underline{d}^1[R_X] & \xrightarrow{\underline{m}_1} & \underline{d}^2[R_X] & \xrightarrow{\underline{m}_2} & \dots \\
 \downarrow \sim & & \downarrow \sim & & \downarrow \sim & & \\
 \underline{d}^0[R_X] & \xrightarrow{\underline{m}_0} & \underline{d}^1[R_X] & \xrightarrow{\underline{m}_1} & \underline{d}^2[R_X] & \xrightarrow{\underline{m}_2} & \dots
 \end{array}$$

in which each vertical map is an equivalence of S^{-L} -modules for the appropriate L , and each $\underline{d}^L[R_X]$ is Σ -cofibrant.

We are now in a position to choose models for the derivatives of representable functors as C_{S_\bullet} -coalgebras.

Definition 4.22 For $X \in \mathcal{S}p^f$, we define

$$d[R_X] := \text{hocolim}_L d^L[R_X].$$

As in Definition 2.10, this symmetric sequence has the structure of a \mathbf{C}_{S_\bullet} -coalgebra, where \mathbf{C}_{S_\bullet} is the comonad of Definition 4.9. Moreover, $d[R_X]$ is Σ -cofibrant and the functor $X \mapsto d[R_X]$ is simplicially enriched.

Lemma 4.23 The \mathbf{C}_{S_\bullet} -coalgebra $d[R_X]$ is naturally equivalent to the symmetric sequence $\partial_*(R_X)$ of derivatives of the representable functor $R_X = \Sigma^\infty \text{Hom}_{\mathcal{S}p}(X, -)$.

Proof By Proposition 4.19, the homotopy colimits of the horizontal sequences in (4.21) are equivalent to the homotopy colimit in Lemma 4.18. \square

Having established a \mathbf{C}_{S_\bullet} -coalgebra structure on the derivatives of the representable functors, we left Kan extend to all pointed simplicial $F: \mathcal{S}p^f \rightarrow \mathcal{S}p$.

Definition 4.24 For an arbitrary pointed simplicial functor $F: \mathcal{S}p^f \rightarrow \mathcal{S}p$, we define $d^L[F]$ by the enriched coend

$$d^L[F] := F(X) \wedge_{X \in \mathcal{S}p^f} d^L[R_X]$$

over the simplicial category $\mathcal{S}p^f$, and

$$d[F] := F(X) \wedge_{X \in \mathcal{S}p^f} d[R_X] \cong \text{hocolim}_L d^L[F].$$

The diagram $d^\bullet[F]$ is a module over the pro-operad S_\bullet and its homotopy colimit $d[F]$ is therefore a \mathbf{C}_{S_\bullet} -coalgebra.

Proposition 4.25 For $F \in [\mathcal{S}p^f, \mathcal{S}p]_*$ cofibrant, we have natural equivalences of symmetric sequences

$$\Sigma^{-*L} \text{cr}_* F(S^L, \dots, S^L) \xrightarrow{\sim} d^L[F]$$

and, taking the homotopy colimit as $L \rightarrow \infty$, an equivalence

$$\partial_*(F) \xrightarrow{\sim} d[F].$$

Proof Since taking cross-effects commutes with homotopy colimits, this follows from Lemmas 4.14 and 4.23. \square

Note that we have a stronger result than in the case of functors from spaces to spectra: the terms $d^L[F]$ are equivalent to the partially stabilized cross-effects without any analyticity condition on F .

4.3 Classification of polynomial functors from spectra to spectra

We now turn to the classification of polynomial functors from spectra to spectra. The structure of this section is similar to the corresponding section for functors from spaces to spectra. The main result is that the comonad \mathbf{C}_{S_\bullet} is equivalent to the comonad \mathbf{C} constructed in [2] which is known to act on the derivatives of such a functor, and from which the Taylor tower can be recovered.

Definition 4.26 The functor $d: [\mathcal{S}p^f, \mathcal{S}p]_* \rightarrow [\Sigma, \mathcal{S}p]$ is simplicially enriched and has a simplicial right adjoint $\Phi: [\Sigma, \mathcal{S}p] \rightarrow [\mathcal{S}p^f, \mathcal{S}p]_*$ given by

$$\Phi(A)(X) := \text{Map}_\Sigma(d[R_X], A).$$

We then define a comonad \mathbf{C} on $[\Sigma, \mathcal{S}p]$ by

$$\mathbf{C} = d\mathbf{c}\Phi,$$

where \mathbf{c} is a comonad cofibrant replacement functor for $[\mathcal{S}p^f, \mathcal{S}p]_*$. Since d takes values in \mathbf{C}_{S_\bullet} -coalgebras, so too does \mathbf{C} and the comonad structure map $\mathbf{C} \rightarrow \mathbf{C}\mathbf{C}$ is a map of \mathbf{C}_{S_\bullet} -coalgebras. Therefore we obtain a map of comonads $\theta: \mathbf{C} \rightarrow \mathbf{C}_{S_\bullet}$ given by the composite

$$\theta_A: \mathbf{C}(A) \rightarrow \mathbf{C}_{S_\bullet}(\mathbf{C}(A)) \rightarrow \mathbf{C}_{S_\bullet}(A),$$

in which the first map is given by the \mathbf{C}_{S_\bullet} -coalgebra structure on $d[\mathbf{c}\Phi A]$, and the second by the counit for \mathbf{C} .

Theorem 4.27 For any symmetric sequence A , the comonad map

$$\theta_A: \mathbf{C}(A) \rightarrow \mathbf{C}_{S_\bullet}(A)$$

is a weak equivalence.

Proof We follow a similar approach to the proof of Theorem 3.64. The main step is to construct, when A is bounded, equivalences of S^{-L} -modules of the form

$$\zeta^L: d^L[\mathbf{c}\Phi A] \simeq \mathbf{C}_{S^{-L}}(A)$$

that commute with the maps $d^L \rightarrow d^{L+1}$ on the left-hand side, and those induced by the operad morphisms $S^{-(L+1)} \rightarrow S^{-L}$ on the right-hand side. The remainder of the proof of Theorem 3.64, the part starting after the proof of Lemma 3.71, then applies in the same way to deduce Theorem 4.27.

Since both sides of the required equivalence commute with products, we can reduce to the case that A is concentrated in a single term, say the n^{th} . Recall that by [2, Corollary 5.11] there is then a natural equivalence

$$\Phi A(X) \simeq (A(n) \wedge X^{\wedge n})^{h\Sigma_n}.$$

Then there is a Σ_n -equivariant equivalence

$$\begin{aligned} d^L[\mathbf{c}\Phi A](n) &\simeq \Sigma^{-nL} \text{cr}_n(\Phi A)(S^L, \dots, S^L) \\ &\simeq \Sigma^{-nL} \left[\bigvee_{\underline{n} \twoheadrightarrow \underline{n}} A(n) \wedge (S^L)^{\wedge n} \right]^{h\Sigma_n} \simeq A(n). \end{aligned}$$

It is therefore enough to show that there is an equivalence of S^{-L} -modules

$$d^L[\mathbf{c}\Phi A] \simeq \mathbf{C}_{S^{-L}}(d^L[\mathbf{c}\Phi A(n)]),$$

or alternatively that the S^{-L} -module structure map

$$d^L[\mathbf{c}\Phi A](k) \rightarrow \left[\prod_{\underline{n} \twoheadrightarrow \underline{k}} \text{Map}(S^{-L}(n_1) \wedge \dots \wedge S^{-L}(n_k), d^L[\mathbf{c}\Phi A](n)) \right]^{\Sigma_n}$$

is an equivalence for each k .

Since the functor d^L (which computes partially stabilized cross-effects) commutes both with taking homotopy fixed points and with smashing with a fixed spectrum, it is then sufficient to show that the S^{-L} -action map

$$d^L[\mathbf{c}(X^{\wedge n})](k) \rightarrow \left[\prod_{\underline{n} \twoheadrightarrow \underline{k}} \text{Map}(S^{-L}(n_1) \wedge \dots \wedge S^{-L}(n_k), d^L[\mathbf{c}(X^{\wedge n})](n)) \right]^{\Sigma_n}$$

is an equivalence.

Recall that we have

$$d^L[\mathbf{c}(X^{\wedge n})](k) \simeq \mathbf{c}(X^{\wedge n}) \wedge_{X \in \mathcal{G}p^f} \text{Map}((S^1 \wedge S(k))^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge k}).$$

Considering the definition of the S^{-L} -module structure here, and consolidating the sphere factors on one side, it is sufficient to show that there is an equivalence

$$(4.28) \quad \mathbf{c}(X^{\wedge n}) \wedge_{X \in \mathcal{G}p^f} \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge k} \xrightarrow{\simeq} \prod_{\underline{n} \twoheadrightarrow \underline{k}} S^{nL},$$

where the component corresponding to a surjection $\alpha: \underline{n} \twoheadrightarrow \underline{k}$ is given by the composite

$$\begin{aligned} \mathbf{c}(X^{\wedge n}) \wedge_{X \in \mathcal{G}p^f} \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge k} &\xrightarrow{\Delta_\alpha} X^{\wedge n} \wedge_{X \in \mathcal{G}p^f} \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge n} \\ &\rightarrow (S^L)^{\wedge n} \cong S^{nL}, \end{aligned}$$

where the first map is given by the diagonal map associated to the surjection α , and the second is the canonical evaluation map. To see that (4.28) is an equivalence, consider the commutative diagram

$$\begin{array}{ccc}
 \text{thocofib}_{I \subseteq \underline{k}} X^{\wedge n} \wedge_{X \in \mathcal{S}p^f} \Sigma^\infty \text{Hom}_{\mathcal{S}p} \left(X, \prod_I S^L \right) & \xrightarrow{\sim} & \text{thocofib}_{I \subseteq \underline{k}} \left(\prod_I S^L \right)^{\wedge n} \\
 \downarrow \sim & & \downarrow \sim \\
 X^{\wedge n} \wedge_{X \in \mathcal{S}p^f} \Sigma^\infty \text{Hom}_{\mathcal{S}p} (X, S^L)^{\wedge k} & \longrightarrow & \prod_{n \rightarrow k} S^{nL}
 \end{array}$$

where the top map is an equivalence by the enriched co-Yoneda Lemma, the left-hand map is an equivalence because for any based space Y , the smash power $Y^{\wedge k}$ is equivalent to the total homotopy cofibre of the k -cube with terms $\prod_I Y$ for $I \subseteq \underline{k}$, and the right-hand map is an equivalence by direct calculation. \square

Theorem 4.29 *The functor d induces an equivalence*

$$[\mathcal{S}p^f, \mathcal{S}p]_{*, \text{poly}}^h \simeq \text{Coalg}_{\mathbf{C}_{S_\bullet}}(S_\bullet)^h$$

between the homotopy category of polynomial pointed simplicial functors $\mathcal{S}p^f \rightarrow \mathcal{S}p$ and the homotopy category of bounded \mathbf{C}_{S_\bullet} -coalgebras of Definition 1.26. Moreover, for cofibrant polynomial functors $F, G \in [\mathcal{S}p^f, \mathcal{S}p]_*$, we have equivalences of simplicial mapping objects

$$(4.30) \quad \text{Nat}_{\mathcal{S}p^f}(F, G) \simeq \widetilde{\text{Map}}_{\mathbf{C}_{S_\bullet}}(d[F], d[G]).$$

Proof The proof is entirely analogous to that of Theorem 3.75, using Theorem 4.27 instead of Theorem 3.64. \square

4.4 Classification of analytic functors from spectra to spectra

As in the case of functors from spaces to spectra, the classification result of the previous section extends naturally to analytic functors in the following way.

Definition 4.31 The notions of *analytic at $*$* and *strongly analytic at $*$* from Definitions 3.84 and 3.91 generalize immediately to functors from spectra to spectra. We therefore have a category $[\mathcal{S}p^f, \mathcal{S}p]_{*, \text{an}(\ast)}$ of pointed simplicial functors $F: \mathcal{S}p^f \rightarrow \mathcal{S}p$ that are strongly analytic at $*$, and an associated homotopy category $[\mathcal{S}p^f, \mathcal{S}p]_{*, \text{an}(\ast)}^h$.

Theorem 4.32 *The functor d sets up an equivalence*

$$[\mathcal{S}p^f, \mathcal{S}p]_{*, \text{an}(*)}^h \simeq \text{Coalg}_{\text{an}}^t(\mathbf{C}_{S_\bullet})^h$$

between the homotopy category of pointed simplicial functors $\mathcal{S}p^f \rightarrow \mathcal{S}p$ that are strongly analytic at $*$ and the pro-truncated homotopy category of analytic \mathbf{C}_{S_\bullet} -coalgebras. Moreover, for cofibrant $F, G \in [\mathcal{S}p^f, \mathcal{S}p]_{*, \text{an}(*)}$ we have equivalences

$$\text{Nat}_{\mathcal{S}p^f}(F, G) \xrightarrow{\sim} \widetilde{\text{Map}}_{\mathbf{C}_{S_\bullet}}(d[F], d[G]) \xrightarrow{\sim} \widetilde{\text{Map}}_{\mathbf{C}_{S_\bullet}}^t(d[F], d[G]).$$

Proof This is analogous to the proof of Theorem 3.92. □

Example 4.33 The derivatives of the functor $\Sigma^\infty \Omega^\infty: \mathcal{S}p \rightarrow \mathcal{S}p$ are given by

$$\partial_*(\Sigma^\infty \Omega^\infty) \simeq \text{Com}.$$

The symmetric sequence Com has a canonical right Com -module structure coming directly from the operad structure, and hence Com is a $\mathbf{C}_{\text{Com}} = \mathbf{C}_{S^0}$ -coalgebra. The comonad map

$$\mathbf{C}_{S^0} \rightarrow \text{hocolim}_{L \rightarrow \infty} \mathbf{C}_{S^{-L}} = \mathbf{C}_{S_\bullet}$$

then endows Com with a \mathbf{C}_{S_\bullet} -coalgebra structure. It follows from [2, Lemma 5.19] that this structure encodes the Taylor tower of $\Sigma^\infty \Omega^\infty$.

5 Proof of Proposition 4.19

The goal of this section is to prove Proposition 4.19; that is, to construct a map

$$(5.1) \quad \underline{m}'_L: \mathbb{B}(\underline{d}^L[R_X], S^{-(L+1)}, S^{-(L+1)}) \rightarrow \underline{d}^{L+1}[R_X],$$

where the source is the standard simplicial bar resolution of $\underline{d}^L[R_X]$ as a $S^{-(L+1)}$ -module.

First recall that S^{-L} is a cofibrant replacement for the desuspension $S^{-L} \text{Com}$ and the S^{-L} -module structure on $\underline{d}^L[R_X]$ is pulled back from a $S^{-L} \text{Com}$ -module structure along the equivalence $S^{-L} \rightarrow S^{-L} \text{Com}$. It is therefore sufficient to construct maps of the form \underline{m}'_L with $S^{-(L+1)}$ replaced with $S^{-(L+1)} \text{Com}$. To keep the notation under control we will just write $S^{-(L+1)}$ in this section, but we mean $S^{-(L+1)} \text{Com}$.

We construct the map \underline{m}'_L by defining, for each non-negative integer N , a map of $S^{-(L+1)}$ -modules

$$\underline{m}'_L{}^N: \Delta_+^N \wedge \underline{d}^L[R_X] \circ \underbrace{S^{-(L+1)} \circ \dots \circ S^{-(L+1)}}_{N+1 \text{ terms}} \rightarrow \underline{d}^{L+1}[R_X].$$

We then show that these maps commute with the face and degeneracy maps in the simplicial bar construction and the coface and codegeneracy maps on the simplexes Δ^N .

For example, \underline{m}_L^0 is the $S^{-(L+1)}$ -module map

$$\underline{m}_L^0: \underline{d}^L[R_X] \circ S^{-(L+1)} \rightarrow \underline{d}^{L+1}[R_X]$$

which is induced by the map of symmetric sequences \underline{m}_L of Definition 4.17. At the next stage, \underline{m}_L^1 is an $S^{-(L+1)}$ -module map

$$\underline{m}_L^1: \Delta_+^1 \wedge \underline{d}^L[R_X] \circ S^{-(L+1)} \circ S^{-(L+1)} \rightarrow \underline{d}^{L+1}[R_X]$$

which is induced by a map of symmetric sequences

$$\Delta_+^1 \wedge \underline{d}^L[R_X] \circ S^{-(L+1)} \rightarrow \underline{d}^{L+1}[R_X]$$

that forms a homotopy which measures the failure of \underline{m}_L to be a map of $S^{-(L+1)}$ -modules on the nose. For larger N , the maps \underline{m}_L^N form a set of higher coherent homotopies.

To illustrate the situation, we can give an explicit description of the two composites in the following (non-commutative) diagram

$$(5.2) \quad \begin{array}{ccc} \underline{d}^L[R_X] \circ S^{-(L+1)} & \xrightarrow{\underline{m}_L} & \underline{d}^{L+1}[R_X] \circ S^{-(L+1)} \\ \eta \downarrow & & \downarrow \mu_{L+1} \\ \underline{d}^L[R_X] \circ S^{-L} & & \\ \mu_L \downarrow & & \downarrow \\ \underline{d}^L[R_X] & \xrightarrow{\underline{m}_L} & \underline{d}^{L+1}[R_X] \end{array}$$

where η is the map of Definition 4.8, μ_L denotes the S^{-L} -module structure on $\underline{d}^L[R_X]$ and \underline{m}_L is the map of Definition 4.17. Each of these composites is built from maps indexed by surjections $\alpha: I \twoheadrightarrow J$ of finite sets, with the component corresponding to α taking the form

$$(5.3) \quad \begin{array}{c} \text{Map}((S^1 \wedge \mathbb{S}(J))^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge J}) \wedge \bigwedge_{j \in J} \text{Map}(\mathbb{S}(I_j)^{L+1}, S) \\ \downarrow \\ \text{Map}((S^1 \wedge \mathbb{S}(I))^{\wedge L+1}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^{L+1})^{\wedge I}) \end{array}$$

The components of the two composites in (5.2) share some features: both can be expressed using

- (i) the diagonal map $\text{Hom}_{\mathcal{G}P}(X, S^L)^{\wedge J} \rightarrow \text{Hom}_{\mathcal{G}P}(X, S^L)^{\wedge I}$, and
- (ii) a homeomorphism

$$(S^1 \wedge \mathbb{S}(J))^{\wedge L} \wedge \bigwedge_{j \in J} \mathbb{S}(I_j)^{\wedge L} \cong (S^1 \wedge \mathbb{S}(I))^{\wedge L}$$

coming from the L -fold smash product of the one-point compactification of the homeomorphism d_α^{-1} of Lemma 4.4. The map (5.3) then arises by combining the maps (i) and (ii) with

- (iii) a map

$$(5.4) \quad S^1 \wedge \mathbb{S}(I) \rightarrow (S^1)^{\wedge I} \wedge \bigwedge_{j \in J} \mathbb{S}(I_j),$$

and

- (iv) the I -fold assembly map

$$(S^1)^{\wedge I} \wedge \text{Hom}_{\mathcal{G}P}(X, S^L)^{\wedge I} \rightarrow \text{Hom}_{\mathcal{G}P}(X, S^{L+1})^{\wedge I}.$$

It is the map (5.4) in which the two composites of (5.2) differ:

- the composite $\underline{m}_L \mu_L \eta$ is determined by the smash product (over $j \in J$) of the maps

$$\eta: S^0 \rightarrow \mathbb{S}(I_j)$$

of Definition 4.6, combined with the homeomorphism $h_I: S^1 \wedge \mathbb{S}(I) \cong (S^1)^{\wedge I}$ of Definition 4.15;

- the composite $\mu_{L+1} \underline{m}_L$ is determined by the composite

$$\begin{aligned} S^1 \wedge \mathbb{S}(I) &\xrightarrow{d_\alpha} S^1 \wedge \mathbb{S}(J) \wedge \bigwedge_{j \in J} \mathbb{S}(I_j) \\ &\xrightarrow{h_J} (S^1)^{\wedge J} \wedge \bigwedge_{j \in J} \mathbb{S}(I_j) \\ &\xrightarrow{\Delta_{S^1}} (S^1)^{\wedge I} \wedge \bigwedge_{j \in J} \mathbb{S}(I_j). \end{aligned}$$

It is the difference between these two maps which implies that (5.2) does not commute and hence that \underline{m}_L is not a map of $S^{-(L+1)}$ -modules. However, the two maps are homotopic and we use a homotopy

$$\tilde{r}^1: \Delta_+^1 \wedge S^1 \wedge \mathbb{S}(I) \rightarrow (S^1)^{\wedge I} \wedge \bigwedge_{j \in J} \mathbb{S}(I_j)$$

to construct the map \underline{m}'_L which forms the first stage in the construction of the map \underline{m}'_L . The map \tilde{r}^1 is the map induced on one-point compactifications by an underlying homotopy of the form

$$r^1: \Delta^1 \times U \rightarrow U \times U; \quad (t, u) \mapsto (tu, (1-t)u),$$

where $U = \prod_{j \in J} \mathbb{R}_0^{I_j}$.

In order to build the map \underline{m}'_L on all levels of the bar construction, we need “higher coherent” versions of the homotopy r^1 . These take the following form.

Definition 5.5 Let V_0, \dots, V_N be finite-dimensional real vector spaces, and for each $i = 0, \dots, N$ suppose $U_i \subseteq V_i$ is a subset that is closed under non-negative scalar multiplication. Then we define continuous functions

$$r^N: \Delta^N \times U_0 \times \dots \times U_N \rightarrow U_0 \times \dots \times U_N \times U_0 \times \dots \times U_N$$

by identifying a point in the simplex Δ^N as a sequence $t = (t_0, \dots, t_N)$ with all $t_j \geq 0$ and $\sum_{j=0}^N t_j = 1$, and then by setting

$$r^N(t, u_0, \dots, u_N)$$

to be equal to

$$((t_1 + \dots + t_N)u_0, \dots, (t_{j+1} + \dots + t_N)u_j, \dots, 0, t_0u_0, \dots, (t_0 + \dots + t_j)u_j, \dots, u_N).$$

Note that the U_N term does not really feature in the homotopy. This term corresponds to the “free” copy of $S^{-(L+1)}$ that appears when we take a map of symmetric sequences with target $d^{L+1}[R_X]$ and extend it to a map of $S^{-(L+1)}$ -modules. Including this term in the homotopy also makes the interactions with coface and codegeneracy maps more clear. Those interactions are described by the following result.

Lemma 5.6 (a) Let U_0, \dots, U_N be as in Definition 5.5. For an integer $1 \leq j \leq N$, the following diagram commutes:

$$\begin{array}{ccc} \Delta^{N-1} \times U_0 \times \dots \times (U_{j-1} \times U_j) \times \dots \times U_N & \xrightarrow{r^{N-1}} & U_0 \times \dots \times U_N \times U_0 \times \dots \times U_N \\ \delta^j \downarrow & & \downarrow \cong \\ \Delta^N \times U_0 \times \dots \times U_N & \xrightarrow{r^N} & U_0 \times \dots \times U_N \times U_0 \times \dots \times U_N \end{array}$$

where $\delta^j: \Delta^{N-1} \rightarrow \Delta^N$ is the coface map given by

$$(t_0, \dots, t_{N-1}) \mapsto (t_0, \dots, t_{j-1}, 0, t_j, \dots, t_{N-1}).$$

(b) For $j = 0$, the following diagram commutes:

$$\begin{array}{ccc} \Delta^{N-1} \times U_1 \times \cdots \times U_N & \xrightarrow{r^{N-1}} & U_1 \times \cdots \times U_N \times U_1 \times \cdots \times U_N \\ \delta^0 \times \iota \downarrow & & \downarrow \iota \times \iota \\ \Delta^N \times U_0 \times U_1 \times \cdots \times U_N & \xrightarrow{r^N} & U_0 \times U_1 \times \cdots \times U_N \times U_0 \times U_1 \times \cdots \times U_N \end{array}$$

where $\iota: U_1 \times \cdots \times U_N \rightarrow U_0 \times U_1 \times \cdots \times U_N$ is the inclusion $(u_1, \dots, u_N) \mapsto (0, u_1, \dots, u_N)$.

(c) For an integer $0 \leq i \leq N$, the following diagram commutes:

$$\begin{array}{ccc} \Delta^{N+1} \times U_0 \times \cdots & & U_0 \times \cdots \times \mathbf{0} \times \cdots \\ \cdots \times U_{i-1} \times \mathbf{0} \times U_i \times \cdots \times U_N & \xrightarrow{r^{N+1}} & \cdots \times U_N \times U_0 \times \cdots \times U_N \\ \sigma^i \downarrow & & \downarrow \cong \\ \Delta^N \times U_0 \times \cdots \times U_N & \xrightarrow{r^N} & U_0 \times \cdots \times U_N \times U_0 \times \cdots \times U_N \end{array}$$

where $\mathbf{0}$ is a zero-dimensional vector space and $\sigma^i: \Delta^{N+1} \rightarrow \Delta^N$ is the codegeneracy map given by

$$(t_0, \dots, t_{N+1}) \mapsto (t_0, \dots, t_{i-1}, t_i + t_{i+1}, t_{i+2}, \dots, t_{N+1}).$$

Proof Each of these is easily checked directly from the definitions. □

We now show how to construct the maps \underline{m}_L^N that generalize the map \underline{m}_L^1 described before Definition 5.5, and from which the desired map \underline{m}'_L is built.

Definition 5.7 Now let I, J be nonempty finite sets and suppose that

$$\lambda: I = I^{(N+1)} \xrightarrow{\alpha_N} I^{(N)} \xrightarrow{\alpha_{N-1}} \cdots \xrightarrow{\alpha_1} I^{(1)} \xrightarrow{\alpha_0} I^{(0)} = J$$

is a sequence of surjections (which determines a sequence of partitions of I). We apply the construction of Definition 5.5 to the sets

$$U_k := \prod_{j \in I^{(k)}} \mathbb{R}_0^{I_j^{(k+1)}},$$

which are subsets of the vector spaces $V_k = \prod_{j \in I^{(k)}} \mathbb{R}^{I_j^{(k+1)}}$ that are closed under non-negative scalar multiplication. This gives us a map

$$r^\lambda: \Delta^N \times U_0 \times \cdots \times U_N \rightarrow U_0 \times \cdots \times U_N \times U_0 \times \cdots \times U_N.$$

We can now compose r^λ with various homeomorphisms as follows:

$$\begin{aligned} \Delta^N \times \mathbb{R} \times \mathbb{R}_0^I &\xrightarrow{\prod d_\alpha} \Delta^N \times \mathbb{R} \times \mathbb{R}_0^J \times U_0 \times \cdots \times U_N \\ &\xrightarrow{r^\lambda} \mathbb{R} \times \mathbb{R}_0^J \times U_0 \times \cdots \times U_N \times U_0 \times \cdots \times U_N \\ &\xrightarrow{\prod d_\alpha^{-1}} \mathbb{R} \times \mathbb{R}_0^I \times U_0 \times \cdots \times U_N \\ &\xrightarrow{h_I} \mathbb{R}^I \times U_0 \times \cdots \times U_N. \end{aligned}$$

Taking one-point compactifications, we get

$$\tilde{r}^\lambda: \Delta_+^N \wedge S^1 \wedge \mathbb{S}(I) \rightarrow (S^1)^{\wedge I} \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} \mathbb{S}(I_j^{(k+1)}).$$

We now combine (ie smash/compose as appropriate) this with

- (i) the diagonal map $\text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge J} \rightarrow \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge I}$,
- (ii) the homeomorphisms

$$(S^1 \wedge \mathbb{S}(J))^{\wedge L} \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} \mathbb{S}(I_j^{(k+1)})^{\wedge L} \xrightarrow{\prod d_\alpha^{-1}} (S^1 \wedge \mathbb{S}(I))^{\wedge L},$$

- (iv) the I -fold assembly map $(S^1)^{\wedge I} \wedge \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge I} \rightarrow \text{Hom}_{\mathcal{G}p}(X, S^{L+1})^{\wedge I}$,

to get a map

$$\begin{aligned} \text{Map}((S^1 \wedge \mathbb{S}(J))^{\wedge L}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^L)^{\wedge J}) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} \text{Map}(\mathbb{S}(I_j)^{\wedge L+1}, S) \\ \downarrow \underline{m}_L^\lambda \\ \text{Map}((S^1 \wedge \mathbb{S}(I))^{\wedge L+1}, \Sigma^\infty \text{Hom}_{\mathcal{G}p}(X, S^{L+1})^{\wedge I}). \end{aligned}$$

That is, we have a map

$$\underline{m}_L^\lambda: \Delta_+^N \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) \rightarrow \underline{d}^{L+1}[R_X](I).$$

Putting these together for all sequences of surjections λ we get the desired map

$$\underline{m}_L^N: \Delta_+^N \wedge \underline{d}^L[R_X] \circ S^{-(L+1)} \circ \cdots \circ S^{-(L+1)} \rightarrow \underline{d}^{L+1}[R_X].$$

In order to show that the maps \underline{m}_L^N fit together to the desired map (5.1), we have to check that the maps \underline{m}_L^λ respect the face and degeneracy maps in the bar construction.

Lemma 5.8 *Let I, J be nonempty finite sets and suppose that as before λ is a sequence of surjections*

$$I = I^{(N+1)} \xrightarrow{\alpha_N} \dots \xrightarrow{\alpha_0} I^{(0)} = J.$$

(a) *For an integer $1 \leq r \leq N$, let $d_r(\lambda)$ denote the sequence of surjections*

$$(\alpha_0, \dots, \alpha_{r-1} \circ \alpha_r, \dots, \alpha_N).$$

Also write

$$\tilde{I}_j^{(r)} := (\alpha_{r-1} \circ \alpha_r)^{-1}(j).$$

Then the following diagram commutes:

$$\begin{array}{ccc}
 \Delta_+^{N-1} \wedge \underline{d}^L[R_X](J) \wedge \dots \wedge \bigwedge_{j \in I^{(r-1)}} S^{-(L+1)}(\tilde{I}_j^{(r)}) \wedge \dots & & \\
 \uparrow \mu & \searrow \underline{m}_L^{d_r(\lambda)} & \\
 \Delta_+^{N-1} \wedge \underline{d}^L[R_X](J) \wedge \dots \wedge \bigwedge_{j \in I^{(r-1)}} S^{-(L+1)}(I_j^{(r)}) & & \\
 \wedge \bigwedge_{j \in I^{(r)}} S^{-(L+1)}(I_j^{(r+1)}) \wedge \dots & \searrow & \underline{d}^{L+1}[R_X](I) \\
 \downarrow \delta^j & \nearrow \underline{m}_L^\lambda & \\
 \Delta_+^N \wedge \underline{d}^L[R_X](J) \wedge \dots \wedge \bigwedge_{j \in I^{(r-1)}} S^{-(L+1)}(I_j^{(r)}) \wedge \bigwedge_{j \in I^{(r)}} S^{-(L+1)}(I_j^{(r+1)}) \wedge \dots & &
 \end{array}$$

where

$$\mu: \bigwedge_{j \in I^{(r-1)}} S^{-(L+1)}(I_j^{(r)}) \wedge \bigwedge_{j \in I^{(r)}} S^{-(L+1)}(I_j^{(r+1)}) \rightarrow \bigwedge_{j \in I^{(r-1)}} S^{-(L+1)}(\tilde{I}_j^{(r)})$$

is the smash product of operad composition maps for $S^{-(L+1)}$, and δ^j is induced by the corresponding coface map $\Delta^{N-1} \rightarrow \Delta^N$.

(b) For the case $r = 0$, let $d_0(\lambda)$ denote the sequence $(\alpha_1, \dots, \alpha_N)$. Then the following diagram commutes:

$$\begin{array}{ccc}
 \Delta_+^{N-1} \wedge \underline{d}^L[R_X](I^{(1)}) \wedge \bigwedge_{k=1}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) & & \\
 \uparrow \nu & \searrow m_L^{d_0(\lambda)} & \\
 \Delta_+^{N-1} \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) & & \underline{d}^{L+1}[R_X](I) \\
 \downarrow \delta^0 & \nearrow m_L^\lambda & \\
 \Delta_+^N \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) & &
 \end{array}$$

where

$$\nu: \underline{d}^L[R_X](J) \wedge \bigwedge_{j \in I^{(0)}} S^{-(L+1)}(I_j^{(1)}) \rightarrow \underline{d}^L[R_X](I^{(1)})$$

is the $S^{-(L+1)}$ -module structure map on $\underline{d}^L[R_X]$, pulled back from the S^{-L} -module structure of Definition 4.13 along the operad map $\eta: S^{-(L+1)} \rightarrow S^{-L}$.

(c) For an integer $0 \leq r \leq N$, denote the sequence $(\alpha_0, \dots, \alpha_{r-1}, 1_{I^{(r)}}, \alpha_r, \dots, \alpha_N)$ by $s_r(\lambda)$. Then the following diagram commutes:

$$\begin{array}{ccc}
 \Delta_+^{N+1} \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) \wedge \bigwedge_{j \in I^{(r)}} S^{-(L+1)}(1) & & \\
 \uparrow \eta & \searrow m_L^{s_i(\lambda)} & \\
 \Delta_+^{N+1} \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) & & \underline{d}^{L+1}[R_X](I) \\
 \downarrow \sigma^r & \nearrow m_L^\lambda & \\
 \Delta_+^N \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) & &
 \end{array}$$

where $\eta: S^0 \xrightarrow{\sim} S^{-(L+1)}(1)$ is the unit map for the operad $S^{-(L+1)}$, and σ^r is induced by the corresponding codegeneracy map $\Delta^{N+1} \rightarrow \Delta^N$.

Proof Working through the definitions of the maps \underline{m}_L^λ , and the relevant operad and module structure maps, each of these claims follows from the corresponding part of Lemma 5.6. \square

The following result is then a restatement of the required Proposition 4.19.

Proposition 5.9 *The maps \underline{m}_L^λ of Definition 5.7 together define a morphism of $S^{-(L+1)}$ -modules*

$$\underline{m}'_L: B(\underline{d}^L[R_X], S^{-(L+1)}, S^{-(L+1)}) \rightarrow \underline{d}^{L+1}[R_X]$$

such that the composite

$$\underline{d}^L[R_X] \xrightarrow{s} B(\underline{d}^L[R_X], S^{-(L+1)}, S^{-(L+1)}) \xrightarrow{\underline{m}'_L} \underline{d}^{L+1}[R_X]$$

is equal to the map \underline{m}_L of Definition 4.17.

Proof The existence of a map \underline{m}'_L follows from Lemma 5.8. To see this is a morphism of $S^{-(L+1)}$ -modules, it is sufficient to show that each map m_L^N is a morphism of $S^{-(L+1)}$ -modules. Let $\lambda = (\alpha_0, \dots, \alpha_N)$ be a sequence of surjections of finite sets

$$I = I^{(N+1)} \xrightarrow{\alpha_N} \dots \xrightarrow{\alpha_0} I^{(0)} = J$$

and let $\lambda' = (\alpha_0, \dots, \alpha_{N-1})$. Then it is sufficient to show that the following diagram commutes:

$$\begin{array}{ccc} & \underline{d}^{L+1}[R_X](I^{(N)}) \wedge \bigwedge_{j \in I^{(N)}} S^{-(L+1)}(I_j^{(N+1)}) & \\ & \nearrow \underline{m}'_L & \downarrow \nu_{L+1} \\ \Delta_+^N \wedge \underline{d}^L[R_X](J) \wedge \bigwedge_{k=0}^N \bigwedge_{j \in I^{(k)}} S^{-(L+1)}(I_j^{(k+1)}) & \xrightarrow{\underline{m}_L^\lambda} & \underline{d}^{L+1}[R_X](I) \end{array}$$

where ν_{L+1} is the $S^{-(L+1)}$ -module structure map for $\underline{d}^{L+1}[R_X]$ associated to the surjection α_N (from Definition 4.13). This follows from the fact that in the map r^N of Definition 5.5, the term U_N is mapped by the identity into the first copy of U_N in the target, and by zero into the second copy.

Finally, the composite $\underline{m}'_L \circ s$ is equal to the composite

$$\underline{d}^L[R_X](I) \xrightarrow{\eta} \underline{d}^L[R_X](I) \wedge \bigwedge_{i \in I} S^{-(L+1)}(1) \xrightarrow{\underline{m}_L^\lambda} \underline{d}^{L+1}[R_X](I),$$

where λ is the sequence consisting just of the identity map $I \xrightarrow{=} I$. Following through the definitions we see that this is precisely \underline{m}_L . \square

References

- [1] **G Arone, M Ching**, *Operads and chain rules for the calculus of functors*, Astérisque 338, Société Mathématique de France, Paris (2011) MR2840569
- [2] **G Arone, M Ching**, *A classification of Taylor towers of functors of spaces and spectra*, Adv. Math. 272 (2015) 471–552 MR3303239
- [3] **G Arone, M Kankaanrinta**, *The sphere operad*, Bull. Lond. Math. Soc. 46 (2014) 126–132 MR3161768
- [4] **D Ayala, J Francis**, *Factorization homology of topological manifolds*, J. Topol. 8 (2015) 1045–1084 MR3431668
- [5] **J M Boardman, R M Vogt**, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics 347, Springer, Berlin (1973) MR0420609
- [6] **M Ching**, *Bar constructions for topological operads and the Goodwillie derivatives of the identity*, Geom. Topol. 9 (2005) 833–933 MR2140994
- [7] **M Ching**, *A chain rule for Goodwillie derivatives of functors from spectra to spectra*, Trans. Amer. Math. Soc. 362 (2010) 399–426 MR2550157
- [8] **M Ching**, *Bar-cobar duality for operads in stable homotopy theory*, J. Topol. 5 (2012) 39–80 MR2897049
- [9] **A D Elmendorf, I Kriz, M A Mandell, J P May**, *Rings, modules, and algebras in stable homotopy theory*, Mathematical Surveys and Monographs 47, Amer. Math. Soc., Providence, RI (1997) MR1417719
- [10] **B Fresse**, *Koszul duality of E_n -operads*, Selecta Math. 17 (2011) 363–434 MR2803847
- [11] **E Getzler, J D S Jones**, *Operads, homotopy algebra and iterated integrals for double loop spaces* arXiv:hep-th/9403055
- [12] **T G Goodwillie**, *Calculus, II: Analytic functors*, K-Theory 5 (1991/92) 295–332 MR1162445
- [13] **T G Goodwillie**, *Calculus, III: Taylor series*, Geom. Topol. 7 (2003) 645–711 MR2026544
- [14] **J P C Greenlees, B Shipley**, *Homotopy theory of modules over diagrams of rings*, Proc. Amer. Math. Soc. Ser. B 1 (2014) 89–104 MR3254575
- [15] **R Helmstutler**, *Conjugate pairs of categories and Quillen equivalent stable model categories of functors*, J. Pure Appl. Algebra 218 (2014) 1302–1323 MR3168497
- [16] **R McCarthy**, *On n -excisive functors of module categories*, preprint (1999) Available at <http://www.math.uiuc.edu/~randy/Vita/>

- [17] **J E McClure, J H Smith**, *A solution of Deligne’s Hochschild cohomology conjecture*, from: “Recent progress in homotopy theory”, (D M Davis, J Morava, G Nishida, W S Wilson, N Yagita, editors), Contemp. Math. 293, Amer. Math. Soc. (2002) 153–193 MR1890736
- [18] **J E McClure, J H Smith**, *Cosimplicial objects and little n -cubes, I*, Amer. J. Math. 126 (2004) 1109–1153 MR2089084
- [19] **T Pirashvili**, *Dold–Kan type theorem for Γ -groups*, Math. Ann. 318 (2000) 277–298 MR1795563
- [20] **C Rezk, S Schwede, B Shipley**, *Simplicial structures on model categories and functors*, Amer. J. Math. 123 (2001) 551–575 MR1833153
- [21] **D P Sinha**, *Manifold-theoretic compactifications of configuration spaces*, Selecta Math. 10 (2004) 391–428 MR2099074

Department of Mathematics, Stockholm University
SE-106 91 Stockholm, Sweden

Department of Mathematics and Statistics, Amherst College
Amherst, MA 01002, United States

gregory.arone@math.su.se, mching@amherst.edu

Proposed: Mark Behrens
Seconded: Haynes Miller, Bill Dwyer

Received: 14 April 2014
Revised: 30 March 2015

