

Vanishing of cohomology and parameter rigidity of actions of solvable Lie groups

HIROKAZU MARUHASHI

We give a sufficient condition for parameter rigidity of actions of solvable Lie groups, by vanishing of (uncountably many) first cohomologies of the orbit foliations. In some cases, we can prove that vanishing of finitely many cohomologies is sufficient. For this purpose we use a rigidity property of quasiisometry.

As an application we prove some actions of 2-step solvable Lie groups on mapping tori are parameter rigid. Special cases of these actions are considered in a paper of Matsumoto and Mitsumatsu.

We also remark on the relation between transitive locally free actions of solvable Lie groups and lattices in solvable Lie groups, and apply results in rigidity theory of lattices in solvable Lie groups to construct transitive locally free actions with some properties.

37A20; 37C15, 37C85

1 Introduction

Let ρ_0 be a C^∞ right action of a connected, simply connected, solvable Lie group S on a closed C^∞ manifold M . We always assume ρ_0 is locally free, that is, every isotropy subgroup is discrete in S . Then we have a foliation \mathcal{F} of M by the orbits of ρ_0 , which is called the orbit foliation of ρ_0 . We say that ρ_0 is *parameter rigid* if any C^∞ locally free action ρ of S on M whose orbit foliation coincides with \mathcal{F} is *parameter equivalent* to ρ_0 . Here parameter equivalence means the following: there are an automorphism Φ of S and a diffeomorphism F of M such that $F(\rho_0(x, s)) = \rho(F(x), \Phi(s))$ for all $x \in M$ and $s \in S$, and F preserves each leaf of \mathcal{F} and is homotopic to the identity via C^∞ maps preserving each leaf.

First we review a criterion for parameter rigidity when S is nilpotent. Instead of using S , let N denote the acting group. We have the *leafwise cohomology* $H^*(\mathcal{F})$ of the foliation \mathcal{F} , which is defined in a similar way to the usual de Rham cohomology. Using the action ρ_0 we can define a canonical injection $H^*(\mathfrak{n}) \hookrightarrow H^*(\mathcal{F})$. We will always use fraktur for the corresponding Lie algebras. The author of this article proved in [5; 4] the following:

1.0.1 Theorem ρ_0 is parameter rigid if and only if $H^1(\mathcal{F}) = H^1(\mathfrak{n})$.

This theorem reduces proving parameter rigidity — which looks like a nonlinear problem at first sight — to a linear one, that is, calculation of first cohomology.

However this criterion is no longer true when the acting group is a general solvable Lie group. In fact there are a solvable Lie group S and a lattice Γ in S for which the natural transitive action $\Gamma \backslash S \curvearrowright S$ is not parameter rigid while $H^1(\Gamma \backslash S) = H^1(\mathfrak{s})$. We will see it in Section 6, together with an example of a locally parameter rigid action of a contractible group which is not parameter rigid. These are obtained by looking at the relations between transitive locally free actions and lattices in solvable Lie groups.

Therefore, the formulation should be changed and we will do it by using twisted leafwise cohomologies. Let us return to the first notations. For the action $M \overset{\rho_0}{\curvearrowright} S$ and a representation π of \mathfrak{s} on a finite-dimensional real vector space V , we will define the leafwise cohomology $H^*(\mathcal{F}; \mathfrak{s} \overset{\pi}{\curvearrowright} V)$ of \mathcal{F} with coefficient π as follows. Let ω_0 denote the canonical 1-form of ρ_0 , ie for any $x \in M$, $(\omega_0)_x: T_x \mathcal{F} \rightarrow \mathfrak{s}$ is the inverse of the derivative at the identity of the map $S \rightarrow M$ which sends g to $\rho_0(x, g)$. This is a leafwise \mathfrak{s} -valued 1-form satisfying $d_{\mathcal{F}} \omega_0 + [\omega_0, \omega_0] = 0$, where $d_{\mathcal{F}}$ is the leafwise exterior derivative of \mathcal{F} . By composing with π we get an $\text{End}(V)$ -valued leafwise 1-form $\pi \omega_0$ satisfying $d_{\mathcal{F}} \pi \omega_0 + [\pi \omega_0, \pi \omega_0] = 0$. Therefore, the trivial vector bundle $M \times V \rightarrow M$ carries the flat leafwise connection whose connection form is $\pi \omega_0$. Here, this connection form is relative to any global frame of the bundle which has constant V components. The square of the exterior derivative with respect to this connection on the space of leafwise V -valued forms $\Omega^*(\mathcal{F}; V)$ is zero by flatness of the connection. So we obtain the cohomology $H^*(\mathcal{F}; \mathfrak{s} \overset{\pi}{\curvearrowright} V)$. On the other hand, the cohomology $H^*(\mathfrak{s}; \mathfrak{s} \overset{\pi}{\curvearrowright} V)$ of the Lie algebra \mathfrak{s} with coefficient π is defined from the complex $\text{Hom}(\wedge^* \mathfrak{s}, V)$. Consider the map $\text{Hom}(\wedge^* \mathfrak{s}, V) \hookrightarrow \Omega^*(\mathcal{F}; V)$ mapping φ to $\omega_0^* \varphi$, where ω_0^* means pullback by ω_0 . This is a cochain map, so that it induces a map between cohomologies.

1.0.2 Proposition The induced map $H^*(\mathfrak{s}; \mathfrak{s} \overset{\pi}{\curvearrowright} V) \rightarrow H^*(\mathcal{F}; \mathfrak{s} \overset{\pi}{\curvearrowright} V)$ is injective.

We will prove this in Section 2.1. Hereafter we shall regard $H^*(\mathfrak{s}; \mathfrak{s} \overset{\pi}{\curvearrowright} V)$ as a subspace of $H^*(\mathcal{F}; \mathfrak{s} \overset{\pi}{\curvearrowright} V)$.

Next we will specify which representations are needed for our sufficient condition. Let \mathfrak{n} be the nilradical of \mathfrak{s} , that is, the largest nilpotent ideal of \mathfrak{s} , which contains $[\mathfrak{s}, \mathfrak{s}]$ since $[\mathfrak{s}, \mathfrak{s}]$ is a nilpotent ideal. Take any subspace \mathfrak{h} satisfying $[\mathfrak{s}, \mathfrak{s}] \subset \mathfrak{h} \subset \mathfrak{n}$. Since \mathfrak{h} is a nilpotent ideal, the descending central series of \mathfrak{h} terminates at some term: $\mathfrak{h} \supset \mathfrak{h}^2 \supset \cdots \supset \mathfrak{h}^d \supset 0$. The adjoint representation ad of \mathfrak{s} has the following invariant

filtration: $\mathfrak{s} \xrightarrow{\text{ad}} \mathfrak{s} \supset \mathfrak{h} \supset \mathfrak{h}^2 \supset \dots \supset \mathfrak{h}^d \supset 0$. We take the associated graded quotient of this filtration: $\mathfrak{s} \xrightarrow{\text{ad}} \text{Gr}_{\mathfrak{h}}(\mathfrak{s}) = \mathfrak{s}/\mathfrak{h} \oplus \bigoplus_{i=1}^d \mathfrak{h}^i/\mathfrak{h}^{i+1}$. Note that $\mathfrak{s}/\mathfrak{h}$ is a direct sum of trivial representations. On this associated graded quotient \mathfrak{h} acts trivially, so that we actually have a representation $\mathfrak{s}/\mathfrak{h} \xrightarrow{\text{ad}} \text{Gr}_{\mathfrak{h}}(\mathfrak{s})$. Let \mathcal{X} denote the set of all surjective homomorphisms $\varphi: \mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{h}$ of Lie algebras. An element $\varphi \in \mathcal{X}$ is just a surjective linear map, which vanishes on $[\mathfrak{s}, \mathfrak{s}]$ since $\mathfrak{s}/\mathfrak{h}$ is abelian. For any $\varphi \in \mathcal{X}$ we form a representation $\mathfrak{s} \xrightarrow{\text{ad} \circ \varphi} \text{Gr}_{\mathfrak{h}}(\mathfrak{s})$.

1.0.3 Theorem (sufficient condition for parameter rigidity) *If*

$$H^1(\mathcal{F}; \mathfrak{s} \xrightarrow{\text{ad} \circ \varphi} \text{Gr}_{\mathfrak{h}}(\mathfrak{s})) = H^1(\mathfrak{s}; \mathfrak{s} \xrightarrow{\text{ad} \circ \varphi} \text{Gr}_{\mathfrak{h}}(\mathfrak{s}))$$

for some \mathfrak{h} and any $\varphi \in \mathcal{X}$, then ρ_0 is parameter rigid.

In general, $[\mathfrak{s}, \mathfrak{s}] \neq \mathfrak{n}$ and we choose \mathfrak{h} between them depending on the situations. For instance, we take $\mathfrak{h} = \mathfrak{n}$ in Theorem 1.0.4 while we take $\mathfrak{h} = [\mathfrak{s}, \mathfrak{s}]$ in Theorem 1.0.5.

The assumption requires vanishing of the cohomologies for every $\varphi \in \mathcal{X}$ and this sometimes causes a difficulty when applying the theorem.¹ But it is very likely that, for most cases, vanishing of the cohomologies for almost all $\varphi \in \mathcal{X}$ is unnecessary, that is, vanishing for finitely many φ is sufficient. To prove such results we have two approaches. First is the method appearing in Matsumoto and Mitsumatsu [6, Section 6], which uses a volume form of the manifold M . Here we generalize this method to obtain the next theorem:

1.0.4 Theorem *Assume the following four conditions:*

- \mathfrak{s} is nonunimodular.
- $\dim \mathfrak{s}/\mathfrak{n} = 1$.
- M is orientable.
- $H^1(\mathcal{F}; \mathfrak{s} \xrightarrow{\text{ad}} \text{Gr}_{\mathfrak{n}}(\mathfrak{s})) = H^1(\mathfrak{s}; \mathfrak{s} \xrightarrow{\text{ad}} \text{Gr}_{\mathfrak{n}}(\mathfrak{s}))$.

Then ρ_0 is parameter rigid.

Recall that \mathfrak{s} is unimodular if and only if $\text{tr ad } X = 0$ for every $X \in \mathfrak{s}$. Note that we have used \mathfrak{n} for \mathfrak{h} . In this theorem, only vanishing for the natural projection $\mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{n} \in \mathcal{X}$ is required by assuming the first three conditions.

The next approach is a new one, in which we use large scale geometry of solvable Lie groups. This can also be applied to unimodular groups. We can obtain several theorems

¹We will see such a situation in Maruhashi [5].

using this method. Here we appeal to a theorem of Ogasawara [8]. For $i = 1, \dots, k$, let

$$A_i = \begin{pmatrix} \alpha_1^{(i)} & & \\ & \ddots & \\ & & \alpha_n^{(i)} \end{pmatrix}$$

be a diagonal matrix with positive diagonal entries and put $A(t) = A_1^{t_1} \cdots A_k^{t_k}$ for $t = (t_1, \dots, t_k) \in \mathbb{R}^k$. Here $A_i^{t_i}$ means

$$\begin{pmatrix} (\alpha_1^{(i)})^{t_i} & & \\ & \ddots & \\ & & (\alpha_n^{(i)})^{t_i} \end{pmatrix}.$$

Consider $S_A = \mathbb{R}^n \rtimes_{A(t)} \mathbb{R}^k$. Let W_S denote the image of the natural map $\text{Aut}(S) \rightarrow \text{GL}(S/[S, S])$. We take $\mathfrak{h} = [\mathfrak{s}, \mathfrak{s}]$ in the next theorem:

1.0.5 Theorem *Assume $S = S_A$ satisfies the condition: for any j there exists i such that $\alpha_j^{(i)} \neq 1$. If*

$$H^1(\mathcal{F}; \mathfrak{s} \overset{\text{ad}\circ\varphi}{\curvearrowright} \text{Gr}_{[\mathfrak{s}, \mathfrak{s}]}(\mathfrak{s})) = H^1(\mathfrak{s}; \mathfrak{s} \overset{\text{ad}\circ\varphi}{\curvearrowright} \text{Gr}_{[\mathfrak{s}, \mathfrak{s}]}(\mathfrak{s}))$$

for all $\varphi \in W_S$, then ρ_0 is parameter rigid.

First $\varphi \in W_S$ is regarded as an element of \mathcal{X} in the following way. By pulling back through $\text{exp}: \mathfrak{s}/[\mathfrak{s}, \mathfrak{s}] \simeq S/[S, S]$, $\varphi: S/[S, S] \rightarrow S/[S, S]$ is regarded as $\varphi: \mathfrak{s}/[\mathfrak{s}, \mathfrak{s}] \rightarrow \mathfrak{s}/[\mathfrak{s}, \mathfrak{s}]$. Composing with the natural projection $\mathfrak{s} \twoheadrightarrow \mathfrak{s}/[\mathfrak{s}, \mathfrak{s}]$ we get $\varphi: \mathfrak{s} \twoheadrightarrow \mathfrak{s}/[\mathfrak{s}, \mathfrak{s}] \in \mathcal{X}$. The second remark is about the condition that for any j there exists i such that $\alpha_j^{(i)} \neq 1$. This is equivalent to $[S, S] = \mathbb{R}^n$. In this theorem, if we put some genericity condition on A , the parameter set W_S becomes finite.

Here we have used a theorem of Ogasawara, but we can also use other theorems treating rigidity of quasiisometry. We will give other applications in a forthcoming paper.

Finally we give an application of this method to get parameter rigid actions. We describe a somewhat generalized version of the usual construction of suspensions of actions. Let $M_0 \overset{\rho_0}{\curvearrowright} H$ be a smooth, locally free action of a connected Lie group H on a closed C^∞ -manifold M_0 , let $G \overset{\Phi}{\curvearrowright} H$ be a smooth action of a connected Lie group G on H by automorphisms and let $\Gamma \overset{\lambda}{\curvearrowright} M_0$ be a smooth action of a cocompact lattice Γ of G on M_0 . Assume these three actions satisfy the compatibility condition

$$\lambda(\gamma, \rho_0(x, h)) = \rho_0(\lambda(\gamma, x), \Phi_\gamma(h))$$

for any $\gamma \in \Gamma$, $x \in M_0$ and $h \in H$. Let $H \rtimes_\Phi G$ be the semidirect product whose multiplication is defined by $(h_1, g_1)(h_2, g_2) = (h_1 \Phi_{g_1}(h_2), g_1 g_2)$ for $h_1, h_2 \in H$

and $g_1, g_2 \in G$. We define two actions $\Gamma \curvearrowright M_0 \times G \curvearrowleft H \rtimes_{\Phi} G$. The action of Γ is defined diagonally: $\gamma(x, g) = (\lambda(\gamma, x), \gamma g)$ for $\gamma \in \Gamma, x \in M_0$ and $g \in G$. The action of $H \rtimes_{\Phi} G$ is defined like the multiplication rule of a semidirect product: $(x, g)(h, g') = (\rho_0(x, \Phi_g(h)), gg')$ for $x \in M_0, g, g' \in G$ and $h \in H$. Then these two actions commute by the compatibility condition. So we get an action

$$\Gamma \backslash (M_0 \times G) \overset{\rho}{\curvearrowleft} H \rtimes_{\Phi} G.$$

This is locally free and the fiber bundle $\Gamma \backslash (M_0 \times G) \rightarrow \Gamma \backslash G$ with a typical fiber M_0 is $(H \rtimes_{\Phi} G \rightarrow G)$ -equivariant. The case in which H is trivial is the usual construction of suspensions.

We deal with a special case of the above construction. Consider $A \in GL(n, \mathbb{Z})$ and an A -invariant subspace V of \mathbb{R}^n . Take a one-parameter subgroup $\Phi: \mathbb{R} \rightarrow GL(V)$ satisfying $\Phi_1 = A|_V$. With respect to the above notation, we let $M_0 = \mathbb{T}^n = \mathbb{Z}^n \backslash \mathbb{R}^n, H = V, G = \mathbb{R}$ and $\Gamma = \mathbb{Z}$, and the given three actions are $\mathbb{Z}^n \backslash \mathbb{R}^n \curvearrowleft V$ by translations, $\mathbb{Z} \curvearrowright \mathbb{T}^n$ by $1 \in \mathbb{Z}$ acting as A , and $\mathbb{R} \overset{\Phi}{\curvearrowright} V$, which are compatible in the above sense. The acting group $S = V \rtimes_{\Phi} \mathbb{R}$ is solvable and two actions $\mathbb{Z} \curvearrowright \mathbb{T}^n \times \mathbb{R} \curvearrowleft S$ are defined by $1(x, t) = (Ax, t + 1)$ and $(x, t)(v, s) = (x + \Phi_t(v), t + s)$. The resulting action is $M = \mathbb{Z} \backslash (\mathbb{T}^n \times \mathbb{R}) \curvearrowleft S$, where M is the mapping torus of A . Note that the images $\Phi_{\mathbb{Z}}$ and $\Phi_{\mathbb{R}}$ of \mathbb{Z} and \mathbb{R} by Φ lie in a real algebraic group $GL(V)$.

1.0.6 Theorem *We assume the following four conditions:*

- V is Diophantine in \mathbb{R}^n .
- $\Phi_{\mathbb{Z}}$ is Zariski dense in $\Phi_{\mathbb{R}}$, meaning $\overline{\Phi_{\mathbb{Z}}} = \overline{\Phi_{\mathbb{R}}}$.
- 1 is not an eigenvalue of $A|_V$.
- $A|_V$ has an eigenvalue whose absolute value is not 1.

Then $M \curvearrowleft S$ is parameter rigid.

The Diophantus condition means that there are a basis v_1, \dots, v_p of V and positive constants C and α satisfying $\max_i |m \cdot v_i| \geq C / \|m\|^\alpha$ for all $m \in \mathbb{Z}^n \setminus \{0\}$.

This theorem² is a generalization of a theorem of Matsumoto and Mitsumatsu [6]. They deal with the case when A is hyperbolic with its characteristic polynomial irreducible over \mathbb{Q} and without eigenvalues on the interval $(-1, 0)$, V is the intersection with \mathbb{R}^n of the direct sum of all eigenspaces of $A: \mathbb{C}^n \rightarrow \mathbb{C}^n$ with eigenvalues of modulus less than 1, and Φ is the most naturally defined one. In this case A is diagonalizable, since its characteristic polynomial has no multiple roots. So they consider the suspension

²In a future work, I plan to generalize this and prove it with a simpler, different calculation.

Anosov flow of the Anosov diffeomorphism A on \mathbb{T}^n and its weak stable foliation, which will be the orbit foliation of the action of $V \rtimes_{\Phi} \mathbb{R}$. In this setting the group $V \rtimes_{\Phi} \mathbb{R}$ is always nonunimodular and Theorem 1.0.4 can be applied.

In our situation A may have nontrivial Jordan blocks, may have a reducible characteristic polynomial such as $A = \begin{pmatrix} A_1 & \\ & A_2 \end{pmatrix}$ for some $A_1 \in \mathrm{GL}(k, \mathbb{Z})$ and $A_2 \in \mathrm{GL}(n - k, \mathbb{Z})$, or V may be some smaller part of stable directions or a mixture of some stable and unstable directions; in particular $V \rtimes_{\Phi} \mathbb{R}$ can be unimodular. The method we use to prove Theorem 1.0.6 is that using large scale geometry instead of that by Matsumoto and Mitsumatsu.

Acknowledgements This is a part of the PhD Thesis of the author. Most of the paper was written when the author was a Research Fellow of the Japan Society for the Promotion of Science. I would like to thank my advisor, Masayuki Asaoka, and an anonymous referee, who pointed out the surjectivity in Proposition 6.1.2.

2 General sufficient condition for parameter rigidity

2.1 Proof of Proposition 1.0.2

As in Section 1, let $M \overset{\rho_0}{\curvearrowright} S$ be an action and $\mathfrak{s} \overset{\pi}{\curvearrowright} V$ a representation. Let $S \overset{\Pi}{\curvearrowright} V$ denote the representation whose derivative is π . We define an action $M \times V \curvearrowright S$ by $(x, v)s = (\rho_0(x, s), \Pi(s^{-1})v)$ for $(x, v) \in M \times V$ and $s \in S$ and this turns $M \times V \rightarrow M$ into an S -equivariant vector bundle. We equip V with a norm which comes from an inner product. Then the space $\Gamma_{\mathrm{cont}}(V)$ of all continuous sections of the trivial vector bundle $M \times V \rightarrow M$ is a Banach space with the supremum norm, and on it we have a natural representation $S \curvearrowright \Gamma_{\mathrm{cont}}(V)$, defined by $(s\xi)(x) = \Pi(s)\xi(\rho_0(x, s))$ for $s \in S$, $\xi \in \Gamma_{\mathrm{cont}}(V)$ and $x \in M$. We regard V as a subspace of $\Gamma_{\mathrm{cont}}(V)$ consisting of constant sections.

2.1.1 Lemma *There is an S -equivariant continuous linear map $\mu: \Gamma_{\mathrm{cont}}(V) \rightarrow V$ which is the identity on V .*

Proof Since S is amenable and M is compact, there exists a ρ_0 -invariant Borel probability measure μ on M . We define $\mu: \Gamma_{\mathrm{cont}}(V) \rightarrow V$ by $\xi \mapsto \int_M \xi d\mu$. Then it is easy to show $\|\int_M \xi d\mu\| \leq \dim V \|\xi\|_{\infty}$ and $\mu(s\xi) = \Pi(s)\mu(\xi)$ for all $\xi \in \Gamma_{\mathrm{cont}}(V)$ and $s \in S$. \square

Using the map μ , we define a map $r: \Omega^*(\mathcal{F}; V) \rightarrow \mathrm{Hom}(\wedge^* \mathfrak{s}, V)$ which, on the p^{th} degree, takes η to $r(\eta)$ defined by $r(\eta)(X_1, \dots, X_p) = \mu(\eta(X_1, \dots, X_p))$ for

$X_1, \dots, X_p \in \mathfrak{s}$. Here X_1, \dots, X_p are regarded as vector fields on M using ρ_0 . Namely, X_i is regarded as a vector field $x \mapsto (\omega_0)_x^{-1}(X_i)$ tangent to the foliation \mathcal{F} , where ω_0 is the canonical 1-form of ρ_0 . We will always make this identification during the paper.

Before proving that r is a cochain map, let us look at our connection more closely. Denote by ∇ the covariant derivative with respect to the flat leafwise connection of the trivial bundle $M \times V \rightarrow M$, defined in Section 1. Since the connection form is $\pi\omega_0$, we have $\nabla\xi = d_{\mathcal{F}}\xi + \pi\omega_0\xi$ for a section $\xi \in \Omega^0(\mathcal{F}; V)$. Let $D: \Omega^p(\mathcal{F}; V) \rightarrow \Omega^{p+1}(\mathcal{F}; V)$ denote the covariant exterior derivative arising from ∇ . Then it is easy to check that $D\eta = d_{\mathcal{F}}\eta + \pi\omega_0 \wedge \eta$ for $\eta \in \Omega^p(\mathcal{F}; V)$.

Next let us see which sections are parallel, that is, sections $\xi \in \Omega^0(\mathcal{F}; V)$ satisfying $\nabla\xi = 0$. Fix a point $x_0 \in M$ and a vector $v \in V$. Define ξ_0 locally along the leaf passing through x_0 by $\xi_0(\rho_0(x_0, s)) = \Pi(s^{-1})v$ for $s \in S$ close to the identity. Then, for any $y = \rho_0(x_0, s_0)$ with small $s_0 \in S$ and any $Y \in \mathfrak{s}$, we have

$$\begin{aligned} \nabla_{d\rho_0(y, e^{tY})/dt|_{t=0}} \xi_0 &= d_{\mathcal{F}}\xi_0 \left(\frac{d}{dt} \rho_0(y, e^{tY}) \Big|_{t=0} \right) + \pi(Y)\xi_0(y) \\ &= \frac{d}{dt} \Pi(e^{-tY} s_0^{-1})v \Big|_{t=0} + \pi(Y)\Pi(s_0^{-1})v \\ &= 0. \end{aligned}$$

Therefore, $\nabla\xi_0 = 0$ and this means the directions of orbits of the action $M \times V \curvearrowright S$ is horizontal for the leafwise connection. So we have

$$(\nabla_X \xi)(x) = \lim_{t \rightarrow 0} \frac{\Pi(e^{tX})\xi(\rho_0(x, e^{tX})) - \xi(x)}{t} = \lim_{t \rightarrow 0} \frac{(e^{tX}\xi)(x) - \xi(x)}{t}$$

for any $\xi \in \Omega^0(\mathcal{F}; V)$, $X \in \mathfrak{s}$ and $x \in M$.

2.1.2 Lemma $(e^{tX}\xi - \xi)/t$ converges uniformly to $\nabla_X \xi$ as $t \rightarrow 0$.

Proof Take a basis v_1, \dots, v_l of V and write $(e^{tX}\xi)(x) = \sum_{i=1}^l f_i(t, x)v_i$ for some real-valued functions f_i . Then $(\nabla_X \xi)(x) = \sum_{i=1}^l f'_i(0, x)v_i$. The function $f_i(t, x)$ has the Taylor expansion $f_i(t, x) = f_i(0, x) + t f'_i(0, x) + \frac{1}{2} t^2 f''_i(\theta_{i,x,t}, x)$, where $\theta_{i,x,t}$ is a number between 0 and t . Since

$$\frac{(e^{tX}\xi)(x) - \xi(x)}{t} - (\nabla_X \xi)(x) = \frac{t}{2} \sum_{i=1}^l f''_i(\theta_{i,x,t}, x)v_i$$

and $f''_i(\theta, x)$ is bounded for $-1 \leq \theta \leq 1$ and $x \in M$, we get the conclusion. □

Recall that we have a cochain map $\omega_0^*: \text{Hom}(\wedge^* \mathfrak{s}, V) \hookrightarrow \Omega^*(\mathcal{F}; V)$.

2.1.3 Lemma *The map r is a cochain map and $r \circ \omega_0^*$ is the identity. Therefore, ω_0^* induces the injective map between cohomologies.*

Proof Using the definitions we verify easily that $r \circ \omega_0^*$ is the identity.

By Lemma 2.1.2, we have

$$\mu(\nabla_X \xi) = \lim_{t \rightarrow 0} \mu\left(\frac{e^{tX}\xi - \xi}{t}\right) = \lim_{t \rightarrow 0} \frac{\Pi(e^{tX})\mu(\xi) - \mu(\xi)}{t} = \pi(X)\mu(\xi)$$

for $X \in \mathfrak{s}$ and $\xi \in \Omega^0(\mathcal{F}; V)$. By this property of μ and a direct calculation, we see that r is a cochain map. □

2.2 Proof of Theorem 1.0.3

Let $M \overset{\rho_0}{\curvearrowright} S$ be an action, \mathcal{F} its orbit foliation and ω_0 the canonical 1-form of ρ_0 . The set of all smooth actions $M \curvearrowright S$ with the orbit foliation \mathcal{F} is denoted by $A(\mathcal{F}, S)$. We will prove that any $\rho \in A(\mathcal{F}, S)$ is parameter equivalent to ρ_0 under the assumption of Theorem 1.0.3. Let ω be the canonical 1-form of ρ . To show that ρ is parameter equivalent to ρ_0 , it is sufficient to prove the existence of some C^∞ map $P: M \rightarrow S$ and some endomorphism $\Phi: S \rightarrow S$ satisfying $\omega = \text{Ad}(P^{-1})\Phi_*\omega_0 + P^*\Theta$, where $\Theta \in \Omega^1(S; \mathfrak{s})$ is the left Maurer–Cartan form of S . See, for instance, Asaoka [1, Proposition 1.4.4]. In other words, we will show that the \mathfrak{s} -valued cocycle ω is cohomologous to a constant \mathfrak{s} -valued cocycle $\Phi_*\omega_0$. We call ω an \mathfrak{s} -valued cocycle because it is the infinitesimal version of a usual S -valued cocycle over ρ_0 . In our situation, the usual K -valued cocycles over ρ_0 for some connected, simply connected Lie group K are in one-to-one correspondence with elements $\eta \in \Omega^1(\mathcal{F}; \mathfrak{k})$ satisfying $d_{\mathcal{F}}\eta + [\eta, \eta] = 0$. Two \mathfrak{k} -valued cocycles η_1 and η_2 are cohomologous if and only if $\eta_1 = \text{Ad}(P^{-1})\eta_2 + P^*\Theta_K$ for some smooth $P: M \rightarrow K$, where Θ_K denotes the left Maurer–Cartan form on K . Also, η is a constant cocycle if and only if $\eta = \Phi_*\omega_0$ for some homomorphism $\Phi: S \rightarrow K$. See Asaoka [1, Section 1.4.1] for more details.

Now recall that we have a filtration $\mathfrak{s} \supset \mathfrak{h} \supset \mathfrak{h}^2 \supset \dots \supset \mathfrak{h}^d \supset 0$. Fix complementary subspaces V_i for $i = 0, \dots, d$, so that $\mathfrak{s} = V_0 \oplus \mathfrak{h}$ and $\mathfrak{h}^i = V_i \oplus \mathfrak{h}^{i+1}$. The adjoint representations $\mathfrak{s} \curvearrowright \mathfrak{s}/\mathfrak{h}$ and $\mathfrak{s} \curvearrowright \mathfrak{h}^i/\mathfrak{h}^{i+1}$ are canonically identified with representations³ of \mathfrak{s} on V_i , which we call π_i , and π_0 is a multiple of the trivial representation. For any element $X \in \mathfrak{s}$, let X^i be the V_i -component with respect to the decomposition $\mathfrak{s} = \bigoplus_{i=0}^d V_i$, so that we have $X = X^0 + X^1 + \dots + X^d$. In this section we use this superscript i to indicate the projection operator onto V_i , or to indicate that the element belongs to V_i . Accordingly, ω is decomposed as $\omega = \omega^0 + \dots + \omega^d$.

³Here we do not assume V_i are invariant under ad .

By looking at the V_0 -component of the equation $d_{\mathcal{F}}\omega + [\omega, \omega] = 0$, we get $d_{\mathcal{F}}\omega^0 = 0$. Our assumption $H^1(\mathcal{F}; V_0) = H^1(\mathfrak{s}; V_0)$ implies that there are a linear map $\varphi: \mathfrak{s} \rightarrow V_0$ vanishing on $[\mathfrak{s}, \mathfrak{s}]$ and a smooth map $h: M \rightarrow V_0$ satisfying

$$(1) \quad \omega^0 = \varphi\omega_0 + d_{\mathcal{F}}h.$$

We also write φ_ρ for φ . This φ_ρ plays an important role (or causes trouble) in our problem.⁴ Unexpectedly, to determine φ_ρ is not so easy. If we can show φ_ρ has some restricted form, we can weaken the assumption of vanishing of cohomologies to a smaller subset of \mathcal{X} . This is what we will do in Sections 3 and 4. Here let us see some properties of φ_ρ .

Let $a: M \times S \rightarrow S$ be the unique smooth map satisfying $\rho_0(x, s) = \rho(x, a(x, s))$ and $a(x, 1) = 1$. This is defined since ρ_0 and ρ have the same orbit foliation. The map a is a cocycle over ρ_0 and is important in our problem.

2.2.1 Lemma *For any $x \in M$ and $s \in S$,*

$$\int_1^s \varphi_\rho \Theta + h(\rho_0(x, s)) - h(x) = \int_1^{a(x,s)} \Theta^0.$$

Proof Since Θ satisfies $d\Theta + [\Theta, \Theta] = 0$, we see that $\varphi_\rho \Theta$ and Θ^0 are V_0 -valued, closed 1-forms on S . Fix $x \in M$ and $s \in S$. Then, by (1),

$$\int_x^{\rho_0(x,s)} (\varphi_\rho \omega_0 + d_{\mathcal{F}}h) = \int_x^{\rho(x,a(x,s))} \omega^0.$$

These integrals are along a curve contained in a leaf. Take a curve $\gamma(t)$, $0 \leq t \leq 1$, on S connecting 1 and s . Then the left-hand side of the above equation is

$$\begin{aligned} \int_0^1 \left(\varphi_\rho \omega_0 \left(\frac{d}{dt} \rho_0(x, \gamma(t)) \right) + d_{\mathcal{F}}h \left(\frac{d}{dt} \rho_0(x, \gamma(t)) \right) \right) dt \\ = \int_0^1 \varphi_\rho \Theta \left(\frac{d}{dt} \gamma(t) \right) dt + h(\rho_0(x, s)) - h(x) \end{aligned}$$

and, to compute the right-hand side, take a curve $\gamma_1(t)$ connecting 1 and $a(x, s)$ and then

$$\int_0^1 \omega^0 \left(\frac{d}{dt} \rho(x, \gamma_1(t)) \right) dt = \int_0^1 \Theta^0 \left(\frac{d}{dt} \gamma_1(t) \right) dt. \quad \square$$

2.2.2 Lemma *The map $\varphi: \mathfrak{s} \rightarrow V_0$ is surjective.*

⁴By (1), $\varphi_\rho = r(\omega^0)$ for r as defined in the previous section. By this formula, we can define φ_ρ without any assumption on the cohomology. But in this case the definition might depend on the choice of a S -invariant Borel probability measure μ .

Proof Let $q: \mathfrak{s} \rightarrow \mathfrak{s}/\ker \varphi_\rho$ be the natural projection and let $\bar{\varphi}_\rho: \mathfrak{s}/\ker \varphi_\rho \rightarrow V_0$ be the map induced from φ_ρ . The map $S \rightarrow V_0$ mapping s to $\int_1^s \Theta^0$ is surjective because $\int_1^{e^X} \Theta^0 = X^0$ for every $X \in \mathfrak{s}$. Fix a point $x \in M$. The map $S \rightarrow S$ defined by $s \mapsto a(x, s)$ is bijective. See Asaoka [1, Lemma 1.4.6]. By the above lemma, we have $\bar{\varphi}_\rho(\int_1^s q\Theta) = \int_1^{a(x,s)} \Theta^0 - h(\rho_0(x, s)) + h(x)$. This and the boundedness of h show $\bar{\varphi}_\rho$ is surjective. \square

Therefore we can regard φ as an element of \mathcal{X} .

2.2.3 Lemma Set $P = e^h: M \rightarrow S$. Then

$$\text{Ad}(P)(\omega - P^*\Theta) = \varphi\omega_0 + \bar{\omega}^1 + \dots + \bar{\omega}^d$$

for some leafwise 1-forms $\bar{\omega}^i$ with values in V_i . So we have

$$(2) \quad \omega = \text{Ad}(P^{-1})(\varphi\omega_0 + \bar{\omega}^1 + \dots + \bar{\omega}^d) + P^*\Theta.$$

Proof First we show

$$P^*\Theta = \sum_{j=0}^{\infty} (-1)^j \frac{(\text{ad } h)^j}{(j+1)!} d_{\mathcal{F}}h = d_{\mathcal{F}}h - \frac{1}{2}(\text{ad } h)d_{\mathcal{F}}h + \dots .$$

This is because, for any point $x \in M$ and $X \in T_x\mathcal{F}$, we have

$$\begin{aligned} (P^*\Theta)(X) &= \Theta \left. \frac{d}{dt} e^{h(x(t))} \right|_{t=0} = (L_{e^{-h(x)}})_* \left. \frac{d}{dt} e^{h(x(t))} \right|_{t=0} \\ &= \left. \frac{d}{dt} e^{-h(x)} e^{h(x)+h(x(t))-h(x)} \right|_{t=0} \\ &= \sum_{j=0}^{\infty} (-1)^j \frac{(\text{ad } h(x))^j}{(j+1)!} Xh, \end{aligned}$$

where $x(t)$ is a curve satisfying $dx(t)/dt|_{t=0} = X$.

So $\omega - P^*\Theta = \varphi\omega_0 + \bar{\omega}^1 + \dots + \bar{\omega}^d$ for some $\bar{\omega}^i$ taking values in V_i . Since $\mathfrak{s}/\mathfrak{h}$ is an abelian Lie algebra, $S \overset{\text{Ad}}{\curvearrowright} \mathfrak{s}/\mathfrak{h}$ is trivial. Therefore, $\text{Ad}(P)(\omega - P^*\Theta) = \varphi\omega_0 + \bar{\omega}^1 + \dots + \bar{\omega}^d$ for some $\bar{\omega}^i$. \square

By this lemma we can replace ω by a cohomologous cocycle whose V_0 -component is constant. We say an element of $\Omega^*(\mathcal{F}; W)$ for some vector space W is constant if it lies in the image of $\omega_0^*: \text{Hom}(\wedge^* \mathfrak{s}, W) \hookrightarrow \Omega^*(\mathcal{F}; W)$. Replacing by cohomologous cocycles, we will gradually make components constant, and finally get a constant cocycle. So now we may assume $\omega = \varphi\omega_0 + \omega^1 + \dots + \omega^d$ and proceed to the next step.

2.2.4 Lemma Assume that

$$\omega = \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^{k-1}\omega_0 + \omega^k + \dots + \omega^d$$

for some linear maps $\varphi^i: \mathfrak{s} \rightarrow V_i$, that is, ω is already constant up to the V_{k-1} -component. Then we can choose some smooth $P: M \rightarrow S$ so that

$$\text{Ad}(P)(\omega - P^*\Theta) = \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^{k-1}\omega_0 + \varphi^k\omega_0 + \bar{\omega}^{k+1} + \dots + \bar{\omega}^d$$

for some linear map $\varphi^k: \mathfrak{s} \rightarrow V_k$ and $\bar{\omega}^i$.

Proof Looking at the V_k -component of the equation $d_{\mathcal{F}}\omega + [\omega, \omega] = 0$, we obtain

$$\begin{aligned} 0 &= d_{\mathcal{F}}\omega^k + [\varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^{k-1}\omega_0 + \omega^k, \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^{k-1}\omega_0 + \omega^k]^k \\ &= d_{\mathcal{F}}\omega^k + \pi_k\varphi\omega_0 \wedge \omega^k + \text{constant form.} \end{aligned}$$

The k appearing in $[\dots, \dots]^k$ in the first line of the above denotes the projection onto V_k . Let $D: \Omega^p(\mathcal{F}; V_k) \rightarrow \Omega^{p+1}(\mathcal{F}; V_k)$ be the covariant exterior derivative arising from the leafwise connection defined by the connection form $\pi_k\varphi\omega_0$. We saw $D = d_{\mathcal{F}} + \pi_k\varphi\omega_0 \wedge$ in the previous section, so that

$$(3) \quad D\omega^k = \omega_0^*\psi$$

for some $\psi \in \text{Hom}(\wedge^2\mathfrak{s}, V_k)$ by the above computation. Recall r , which is defined in the previous section. We set $\theta = r(\omega^k): \mathfrak{s} \rightarrow V_k$. Then

$$(4) \quad \psi = r(\omega_0^*\psi) = r(D\omega^k) = Dr(\omega^k) = D\theta.$$

Here D also denotes the differential of $\text{Hom}(\wedge^*\mathfrak{s}, V_k)$. By (3) and (4), we get

$$D(\omega^k - \omega_0^*\theta) = 0.$$

By Lemma 2.2.2 and by our assumption, we have $H^1(\mathcal{F}; \mathfrak{s} \underset{\mathcal{V}}{\overset{\pi_k\varphi}{\curvearrowright}} V_k) = H^1(\mathfrak{s}; \mathfrak{s} \underset{\mathcal{V}}{\overset{\pi_k\varphi}{\curvearrowright}} V_k)$. Therefore there exist a linear map $\theta': \mathfrak{s} \rightarrow V_k$ and a smooth map $h: M \rightarrow V_k$ satisfying

$$\omega^k = \theta\omega_0 + \theta'\omega_0 + d_{\mathcal{F}}h + \pi_k\varphi\omega_0h = \varphi^k\omega_0 + d_{\mathcal{F}}h + \pi_k\varphi\omega_0h.$$

Here we set $\varphi^k = \theta + \theta'$. As before we let $P = e^h$ and then

$$\omega - P^*\Theta = \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^{k-1}\omega_0 + (\varphi^k\omega_0 + \pi_k\varphi\omega_0h) + \bar{\omega}^{k+1} + \dots + \bar{\omega}^d$$

for some $\bar{\omega}^i$. Finally we compute as follows:

$$\begin{aligned} \text{Ad}(P)(\omega - P^*\Theta) &= e^{\text{ad}h}(\varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^k\omega_0 + \pi_k\varphi\omega_0h) + \text{higher terms} \\ &= \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^k\omega_0 + \pi_k\varphi\omega_0h - \pi_k\varphi\omega_0h + \text{higher terms} \\ &= \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^k\omega_0 + \text{higher terms.} \end{aligned} \quad \square$$

Applying this lemma repeatedly, we see the given ω is cohomologous to a cocycle of the form $\omega' = \varphi\omega_0 + \varphi^1\omega_0 + \dots + \varphi^d\omega_0$. Set $\Phi_* = \varphi + \varphi^1 + \dots + \varphi^d: \mathfrak{s} \rightarrow \mathfrak{s}$. Then $\omega' = \Phi_*\omega_0$ is a constant cocycle, because the equation $d_{\mathcal{F}}\omega' + [\omega', \omega'] = 0$ implies that Φ_* is an endomorphism of the Lie algebra \mathfrak{s} . This completes the proof of Theorem 1.0.3.

Let \mathcal{X}_{ρ_0} be the set of all $\varphi \in \mathcal{X}$ which can be written as φ_ρ for some $\rho \in A(\mathcal{F}, S)$ (using the isomorphism $\mathfrak{s}/\mathfrak{h} \simeq V_0$). What we actually proved in this section is:

2.2.5 Theorem Assume $H^1(\mathcal{F}) = H^1(\mathfrak{s})$, so that we can define \mathcal{X}_{ρ_0} as a subset of \mathcal{X} . If

$$H^1(\mathcal{F}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}\circ\varphi}{\curvearrowright}} \text{Gr}_{\mathfrak{h}}(\mathfrak{s})) = H^1(\mathfrak{s}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}\circ\varphi}{\curvearrowright}} \text{Gr}_{\mathfrak{h}}(\mathfrak{s}))$$

for all $\varphi \in \mathcal{X}_{\rho_0}$, then ρ_0 is parameter rigid.

The next task is to prove the set \mathcal{X}_{ρ_0} is small.

Remark Although we deal with only \mathfrak{s} -valued cocycles arising from actions ρ in $A(\mathcal{F}, S)$, what we do in this section is actually valid for any \mathfrak{k} -valued cocycles over ρ_0 for any connected, simply connected, solvable Lie group K and a subspace \mathfrak{h} between $[\mathfrak{k}, \mathfrak{k}]$ and the nilradical of \mathfrak{k} . Also the acting group S need not be solvable; we need only assume that the action has an invariant Borel probability measure. Solvability is used only for the value group K . This might be useful for purposes other than parameter rigidity.

Remark For actions of semisimple Lie groups or groups with property (T), \mathbb{R} -valued cocycle rigidity implies K -valued cocycle rigidity for any connected simply connected solvable Lie groups. This is shown by an obvious argument and valid for actions in broader categories.

Remark As explained in Asaoka [1, Section 1.4.4], $H^1(\mathcal{F}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}}{\curvearrowright}} \mathfrak{s})/H^1(\mathfrak{s}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}}{\curvearrowright}} \mathfrak{s})$ can be viewed as the formal tangent space at ρ_0 in $A(\mathcal{F}, S)/(\text{parameter equivalence})$. We can show that $H^1(\mathcal{F}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}}{\curvearrowright}} \text{Gr}_{\mathfrak{h}}(\mathfrak{s})) = H^1(\mathfrak{s}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}}{\curvearrowright}} \text{Gr}_{\mathfrak{h}}(\mathfrak{s}))$ implies $H^1(\mathcal{F}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}}{\curvearrowright}} \mathfrak{s})/H^1(\mathfrak{s}; \mathfrak{s} \underset{\curvearrowright}{\overset{\text{ad}}{\curvearrowright}} \mathfrak{s}) = 0$ by an argument using spectral sequences. But the converse seems to be false.

3 Sufficient condition by the method of Matsumoto and Mitsumatsu

We prove Theorem 1.0.4 here. Let $M \overset{\rho_0}{\curvearrowright} S$ be an action which we consider and take any $\rho \in A(\mathcal{F}, S)$. According to Theorem 2.2.5, what we need to show is that φ_ρ ,

defined in Section 2.2, coincides with the natural projection $\mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{n}$. In this section we also use the notation $\rho_0^s(x) = \rho_0(x, s)$ and $\rho^s(x) = \rho(x, s)$. As in Section 2.2 we have a S -valued cocycle $a: M \times S \rightarrow S$ over ρ_0 satisfying $\rho_0^s(x) = \rho^{a(x,s)}(x)$ for all $x \in M$ and $s \in S$. For any $X \in \mathfrak{s}$, $s \in S$ and $x \in M$,

$$(\rho_0^s)_* X_x = \left. \frac{d}{dt} \rho_0(x, e^{tX}s) \right|_{t=0} = \left. \frac{d}{dt} \rho_0(\rho_0^s(x), e^{t \text{Ad}(s^{-1})X}) \right|_{t=0} = (\text{Ad}(s^{-1})X)_{\rho_0^s(x)}.$$

Take a basis of \mathfrak{s} and its dual basis of \mathfrak{s}^* . The dual basis of \mathfrak{s}^* , after being pulled back by the canonical 1-form ω_0 of ρ_0 , is regarded as a global frame of the bundle $T^*\mathcal{F}$. Let $\Omega_0 \in \Omega^{\dim S}(\mathcal{F})$ be the wedge product of the global frame, which is a leafwise volume form. By the above computation, we see

$$(\rho_0^s)^* \Omega_0 = \det \text{Ad}(s^{-1}) \Omega_0$$

for all $s \in S$. We do the same thing for ρ , getting another leafwise volume form $\Omega \in \Omega^{\dim S}(\mathcal{F})$ which satisfies

$$(\rho^s)^* \Omega = \det \text{Ad}(s^{-1}) \Omega$$

for all $s \in S$. Here we must use the canonical 1-form ω of ρ rather than ω_0 . Fix a complementary subbundle E to $T\mathcal{F}$ in TM , that is, $TM = T\mathcal{F} \oplus E$. Since M is orientable, we can choose a nowhere-vanishing smooth section Ω_{tr} of $\wedge^{\dim E} (TM/T\mathcal{F})^*$. We have natural projections $T\mathcal{F} \leftarrow TM \rightarrow TM/T\mathcal{F}$ defined by E . Let $\bar{\Omega}_0, \bar{\Omega} \in \Omega^{\dim S}(M)$ and $\bar{\Omega}_{\text{tr}} \in \Omega^{\dim E}(M)$ be the pullbacks of Ω_0, Ω and Ω_{tr} by the projections. In this section, bars written over something stand for pulling back something by the projections. Both $\bar{\Omega}_0 \wedge \bar{\Omega}_{\text{tr}}, \bar{\Omega} \wedge \bar{\Omega}_{\text{tr}} \in \Omega^{\dim M}(M)$ are volume forms of M . So there is a smooth map $c: M \times S \rightarrow \mathbb{R}_{>0}$ satisfying $(\rho^s)^*(\bar{\Omega} \wedge \bar{\Omega}_{\text{tr}}) = c(\cdot, s)(\bar{\Omega} \wedge \bar{\Omega}_{\text{tr}})$ for all $s \in S$. It is easy to see that c is a cocycle over ρ , that is, $c(x, ss') = c(x, s)c(\rho(x, s), s')$ for all $x \in M$ and $s, s' \in S$. Our assumption $H^1(\mathcal{F}) = H^1(\mathfrak{s})$ is equivalent to the \mathbb{R} -valued cocycle rigidity of ρ . See Maruhashi [4, Section 2], for instance. Thus we can find a homomorphism $\alpha: S \rightarrow \mathbb{R}_{>0}$ and a smooth map $P: M \rightarrow \mathbb{R}_{>0}$ such that $c(x, s) = P(x)^{-1} \alpha(s) P(\rho(x, s))$ holds for all $x \in M$ and $s \in S$. Then, for any $s \in S$,

$$(\rho^s)^*(P^{-1}(\bar{\Omega} \wedge \bar{\Omega}_{\text{tr}})) = P(\rho^s(\cdot))^{-1} c(\cdot, s)(\bar{\Omega} \wedge \bar{\Omega}_{\text{tr}}) = \alpha(s) P^{-1}(\bar{\Omega} \wedge \bar{\Omega}_{\text{tr}}).$$

By integrating over M , we get $\alpha(s) = 1$ for all $s \in S$. By replacing $P^{-1} \Omega_{\text{tr}}$ by Ω_{tr} we may assume that

$$(5) \quad (\rho^s)^*(\bar{\Omega} \wedge \bar{\Omega}_{\text{tr}}) = \bar{\Omega} \wedge \bar{\Omega}_{\text{tr}}$$

for all $s \in S$. For any $x \in M$ and $s \in S$, two maps

$$(\rho_0^s)_*, (\rho^{a(x,s)})_*: (TM/T\mathcal{F})_x \rightarrow (TM/T\mathcal{F})_{\rho_0^s(x) = \rho^{a(x,s)}(x)}$$

coincide. This is because ρ_0 and ρ have the same orbit foliation and it is easy to see this for small $s \in S$. For two small $s, s' \in S$, we have

$$\begin{array}{ccccc} (TM/T\mathcal{F})_x & \xrightarrow{(\rho_0^s)_*} & (TM/T\mathcal{F})_{\rho_0^s(x)} & \xrightarrow{(\rho_0^{s'})_*} & (TM/T\mathcal{F})_{\rho_0^{s'}(x)} \\ \parallel & \wr & \parallel & \wr & \parallel \\ (TM/T\mathcal{F})_x & \xrightarrow{(\rho^{a(x,s)})_*} & (TM/T\mathcal{F})_{\rho_0^s(x)} & \xrightarrow{(\rho^{a(\rho_0^s(x),s')})_*} & (TM/T\mathcal{F})_{\rho_0^{s'}(x)} \end{array}$$

so that $(\rho_0^{s'})_* = (\rho_0^{s'})_* \circ (\rho_0^s)_* = (\rho^{a(\rho_0^s(x),s')})_* \circ (\rho^{a(x,s)})_* = (\rho^{a(x,s)a(\rho_0^s(x),s')})_* = (\rho^{a(x,s,s')})_*$. Using this we can prove the same for general $s \in S$. Therefore,

$$(6) \quad ((\rho_0^s)^* \Omega_{tr})_x = ((\rho^{a(x,s)})^* \Omega_{tr})_x$$

for all $x \in M$ and $s \in S$. Let $\beta: M \times S \rightarrow \mathbb{R}_{>0}$ be the smooth map satisfying $(\rho^s)^* \Omega_{tr} = \beta(\cdot, s) \Omega_{tr}$ for all $s \in S$.

3.0.1 Lemma We have $\beta(\cdot, s) = \det \text{Ad}(s)$ for all $s \in S$. Therefore,

$$(7) \quad (\rho^s)^* \Omega_{tr} = \det \text{Ad}(s) \Omega_{tr}.$$

Proof By (5),

$$\overline{\Omega} \wedge \overline{\Omega}_{tr} = (\rho^s)^* (\overline{\Omega} \wedge \overline{\Omega}_{tr}) = (\rho^s)^* \overline{\Omega} \wedge (\rho^s)^* \overline{\Omega}_{tr}.$$

Since

$$\begin{array}{ccc} T_x M & \xrightarrow{(\rho^s)_*} & T_{\rho^s(x)} M \\ \downarrow & \wr & \downarrow \\ (TM/T\mathcal{F})_x & \xrightarrow{(\rho^s)_*} & (TM/T\mathcal{F})_{\rho^s(x)} \end{array}$$

we have $(\rho^s)^* \overline{\Omega}_{tr} = \overline{(\rho^s)^* \Omega}_{tr}$. On the other hand, the diagram

$$\begin{array}{ccc} T_x M & \xrightarrow{(\rho^s)_*} & T_{\rho^s(x)} M \\ \downarrow & & \downarrow \\ T_x \mathcal{F} & \xrightarrow{(\rho^s)_*} & T_{\rho^s(x)} \mathcal{F} \end{array}$$

does not commute unless E is ρ -invariant, so usually $(\rho^s)^* \overline{\Omega} \neq \overline{(\rho^s)^* \Omega}$. But

$$(\rho^s)^* \overline{\Omega} \wedge (\rho^s)^* \overline{\Omega}_{tr} = \overline{(\rho^s)^* \Omega} \wedge \overline{(\rho^s)^* \Omega}_{tr}$$

holds, because $(\rho^s)^* \overline{\Omega}$ and $\overline{(\rho^s)^* \Omega}$ coincide when only vectors in $T\mathcal{F}$ are substituted, and $(\rho^s)^* \overline{\Omega}_{tr}$ vanishes if we substitute a vector in $T\mathcal{F}$. Therefore, $\overline{\Omega} \wedge \overline{\Omega}_{tr} =$

$\overline{(\rho^s)^*\Omega} \wedge \overline{(\rho^s)^*\Omega_{\text{tr}}} = \overline{\det \text{Ad}(s^{-1})\Omega} \wedge \overline{\beta(\cdot, s)\Omega_{\text{tr}}} = \det \text{Ad}(s^{-1})\overline{\Omega} \wedge \overline{\Omega_{\text{tr}}}$ and the claim follows. □

Combining (6) and (7), we get

$$(\rho_0^s)^*\Omega_{\text{tr}} = \det \text{Ad}(a(\cdot, s))\Omega_{\text{tr}}.$$

Then, by the same argument as in the proof of the above lemma,

$$\begin{aligned} (8) \quad (\rho_0^s)^*(\overline{\Omega}_0 \wedge \overline{\Omega}_{\text{tr}}) &= (\rho_0^s)^*\overline{\Omega}_0 \wedge (\rho_0^s)^*\overline{\Omega}_{\text{tr}} \\ &= \overline{(\rho_0^s)^*\Omega_0} \wedge \overline{(\rho_0^s)^*\Omega_{\text{tr}}} \\ &= \overline{\det \text{Ad}(s^{-1})\Omega_0} \wedge \overline{\det \text{Ad}(a(x, s))\Omega_{\text{tr}}} \\ &= \det \text{Ad}(s^{-1}) \det \text{Ad}(a(x, s))\overline{\Omega}_0 \wedge \overline{\Omega}_{\text{tr}}. \end{aligned}$$

This formula will be the key to proving Theorem 1.0.4.

3.0.2 Lemma For any $s \in S$,

$$\det \text{Ad}(s) = e^{\text{tr ad } \int_1^s p^\Theta},$$

where $p: \mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{n}$ denotes the natural projection and ad is also used for $\mathfrak{s}/\mathfrak{n} \curvearrowright \text{Gr}_{\mathfrak{n}}(\mathfrak{s})$.

Proof Any $s \in S$ can be written as $s = e^{X_1} \dots e^{X_k}$ for some $X_1, \dots, X_k \in \mathfrak{s}$. It is easy to see

$$\int_1^s p^\Theta = p(X_1) + \dots + p(X_k).$$

Since $\text{tr ad } p(X) = \text{tr ad } X$ for $X \in \mathfrak{s}$, we have

$$\det \text{Ad}(s) = e^{\text{tr ad } X_1} \dots e^{\text{tr ad } X_k} = e^{\text{tr ad}(p(X_1) + \dots + p(X_k))} = e^{\text{tr ad } \int_1^s p^\Theta}. \quad \square$$

Since we have $\int_1^s \varphi_\rho^\Theta + h(\rho_0(x, s)) - h(x) = \int_1^{a(x, s)} p^\Theta$ as in Lemma 2.2.1, the factor appearing in the formula (8) will be

$$\begin{aligned} \det \text{Ad}(s^{-1}) \det \text{Ad}(a(x, s)) &= e^{-\text{tr ad } \int_1^s p^\Theta} e^{\text{tr ad } \int_1^{a(x, s)} p^\Theta} \\ &= \exp\left(\text{tr ad} \int_1^s (\varphi_\rho - p)^\Theta + \text{tr ad}(h(\rho_0(x, s)) - h(x))\right). \end{aligned}$$

Assume $\varphi_\rho \neq p$ and we will get a contradiction as follows. We have $X \in \mathfrak{s}$, so that $A := (\varphi_\rho - p)(X) \neq 0$. Since $\dim \mathfrak{s}/\mathfrak{n} = 1$ and \mathfrak{s} is not unimodular, $\text{tr ad } A \neq 0$. We may assume $\text{tr ad } A > 0$ by replacing X . Then, for $s = e^{TX}$ with $T > 0$ large, the function

$$\det \text{Ad}(s^{-1}) \det \text{Ad}(a(x, s)) = e^{T \text{tr ad } A} e^{\text{tr ad}(h(\rho_0(x, s)) - h(x))}$$

on M must be large because h is bounded. This contradicts the formula (8) by integrating over M . Therefore, $\varphi_\rho = p$ and this completes the proof of Theorem 1.0.4.

Remark The assumptions $\dim \mathfrak{s}/\mathfrak{n} = 1$ and nonunimodularity of \mathfrak{s} are used only in the final part of the proof, so we have proved the following: if M is orientable, $\text{tr ad}(\varphi_\rho - p)(X) = 0$ for all $X \in \mathfrak{s}$, where ad denotes $\mathfrak{s}/\mathfrak{n} \curvearrowright^{\text{ad}} \text{Gr}_n(\mathfrak{s})$.

4 Sufficient condition given by large scale geometry of solvable Lie groups

4.1 Key proposition to the method

Let X and B be metric spaces. We say a surjective map $p: X \rightarrow B$ is a *distance-respecting projection* if $d(b, b') = d(p^{-1}(b), p^{-1}(b')) = d_{\mathcal{H}}(p^{-1}(b), p^{-1}(b'))$ holds for any two points $b, b' \in B$. Here, $d_{\mathcal{H}}$ denotes the Hausdorff distance. Let $p: X \rightarrow B$ and $p': X' \rightarrow B'$ be distance-respecting projections. For a given diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & X' \\ p \downarrow & & \downarrow p' \\ B & \xrightarrow{\varphi} & B' \end{array}$$

we say f is *fiber-respecting over φ* if there is $C > 0$ such that

$$d_{\mathcal{H}}(f(p^{-1}(b)), (p')^{-1}(\varphi(b))) < C$$

for all $b \in B$.

4.1.1 Lemma *Let G be a Lie group and H be a closed subgroup of G .*

- (1) *Left-invariant Riemannian metrics on G/H are in one-to-one correspondence with inner products on $\mathfrak{g}/\mathfrak{h}$ invariant under $H \curvearrowright^{\text{Ad}} \mathfrak{g}/\mathfrak{h}$ by the canonical identification $\mathfrak{g}/\mathfrak{h} \simeq T_H G/H$.*
- (2) *Assume that there exists an invariant inner product for $H \curvearrowright^{\text{Ad}} \mathfrak{g}/\mathfrak{h}$. Take an inner product of \mathfrak{g} for which the restriction $\mathfrak{h}^\perp \xrightarrow{\sim} \mathfrak{g}/\mathfrak{h}$ of the projection $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h}$ is an isometry. Endow G and G/H with left invariant Riemannian metrics defined by these inner products. Let $\pi: G \rightarrow G/H$ be the projection. Then, for every $g \in G$, the restriction $(L_g)_* \mathfrak{h}^\perp \xrightarrow{\sim} T_g G/H$ of $\pi_*: T_g G \rightarrow T_g G/H$ is an isometry, and the kernel of $\pi_*: T_g G \rightarrow T_g G/H$ is $(L_g)_* \mathfrak{h}$.*
- (3) *Assume G is connected. Under the assumption of (2), the map $\pi: G \rightarrow G/H$ is a distance-respecting projection.*

Proof It is easy to check (1) and (2).

(3) Take arbitrary two points $b, b' \in G/H$.

Claim $d(\pi^{-1}(b), \pi^{-1}(b')) \geq d(b, b')$.

For any $g \in \pi^{-1}(b)$ and $g' \in \pi^{-1}(b')$, take a minimal geodesic $\gamma: [0, 1] \rightarrow G$ connecting g and g' . Since $\|(\pi \circ \gamma)'(t)\| = \|\pi_* \gamma'(t)\| \leq \|\gamma'(t)\|$ by the property of our metric, $d(g, g') = \int_0^1 \|\gamma'(t)\| dt \geq \int_0^1 \|(\pi \circ \gamma)'(t)\| dt \geq d(b, b')$.

Claim $d(\pi^{-1}(b), \pi^{-1}(b')) \leq d(b, b')$.

Take a minimal geodesic $\gamma: [0, 1] \rightarrow G/H$ connecting b and b' . Fix a point $g \in \pi^{-1}(b)$. Then there exists a curve $\tilde{\gamma}: [0, 1] \rightarrow G$ starting from g such that $\pi \circ \tilde{\gamma} = \gamma$ and $\tilde{\gamma}'(t) \in (L\tilde{\gamma}(t))_* \mathfrak{h}^\perp$ for all $t \in [0, 1]$, by a standard argument. For this curve, $d(b, b') = \int_0^1 \|\gamma'(t)\| dt = \int_0^1 \|\tilde{\gamma}'(t)\| dt \geq d(\pi^{-1}(b), \pi^{-1}(b'))$.

Claim $d(b, b') = d_{\mathcal{H}}(\pi^{-1}(b), \pi^{-1}(b'))$.

We have $B(\pi^{-1}(b), C) = \pi^{-1}(B(b, C))$ and $B(\pi^{-1}(b'), C) = \pi^{-1}(B(b', C))$ by the above discussion; this is obvious. □

4.1.2 Corollary *Let G be a connected Lie group and H a connected, normal, closed subgroup of G . Take an inner product of \mathfrak{g} . Endow $\mathfrak{g}/\mathfrak{h}$ with the inner product for which the restriction $\mathfrak{h}^\perp \xrightarrow{\sim} \mathfrak{g}/\mathfrak{h}$ of the projection $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h}$ is an isometry. Consider left-invariant Riemannian metrics on G and G/H corresponding to these inner products. Then the projection $\pi: G \rightarrow G/H$ is a distance-respecting projection.*

Proof This is because $H \overset{\text{Ad}}{\curvearrowright} \mathfrak{g}/\mathfrak{h}$ is trivial. □

Let us return to our previous setting $M \overset{\rho_0}{\curvearrowright} S$. Take an action $\rho \in A(\mathcal{F}, S)$ and let $a: M \times S \rightarrow S$ be the cocycle over ρ_0 defined by ρ , that is, $\rho_0(x, s) = \rho(x, a(x, s))$.

4.1.3 Lemma *For any $x \in M$, the map $S \rightarrow S$ taking s to $a(x, s)$ is a bi-Lipschitz diffeomorphism for any left-invariant Riemannian metric on S .*

Proof See Asaoka [1, Lemma 1.4.6] for a proof that the map is a diffeomorphism.

To see that the map is bi-Lipschitz, equip $T\mathcal{F}$ with the metric $\langle \cdot, \cdot \rangle_{\rho_0}$ for which $(\omega_0)_x: T_x\mathcal{F} \xrightarrow{\sim} \mathfrak{s}$ is isometric for all $x \in M$. We can easily verify that the metric obtained by pulling back $\langle \cdot, \cdot \rangle_{\rho_0}$ through the map $S \rightarrow M$ taking s to $\rho_0(x, s)$ is the original Riemannian metric on S . Similarly, we also have the metric $\langle \cdot, \cdot \rangle_\rho$ on $T\mathcal{F}$ defined by using ρ . Since M is compact, there is a constant $C > 1$ such that

$C^{-1} \|\cdot\|_{\rho_0} \leq \|\cdot\|_{\rho} \leq C \|\cdot\|_{\rho_0}$. For any $s, s' \in S$, take a minimal geodesic $\gamma: [0, 1] \rightarrow S$ connecting s and s' . Then

$$\begin{aligned} d(s, s') &= \int_0^1 \|\gamma'(t)\| dt = \int_0^1 \left\| \frac{d}{dt} \rho_0(x, \gamma(t)) \right\|_{\rho_0} dt \\ &= \int_0^1 \left\| \frac{d}{dt} \rho(x, a(x, \gamma(t))) \right\|_{\rho_0} dt \\ &\geq \frac{1}{C} \int_0^1 \left\| \frac{d}{dt} \rho(x, a(x, \gamma(t))) \right\|_{\rho} dt \\ &= \frac{1}{C} \int_0^1 \left\| \frac{d}{dt} a(x, \gamma(t)) \right\| dt \\ &\geq \frac{1}{C} d(a(x, s), a(x, s')). \end{aligned}$$

The inverse is also Lipschitz by the same argument. □

Recall that \mathfrak{h} is a subspace between $[\mathfrak{s}, \mathfrak{s}]$ and \mathfrak{n} . Let K_{ρ} and H be the Lie subgroups corresponding to $\ker \varphi_{\rho}$ and \mathfrak{h} and $\tilde{\varphi}_{\rho}: S/K_{\rho} \rightarrow S/H$ be the map induced by $\varphi_{\rho}: \mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{h}$. Both S/K_{ρ} and S/H are vector groups and $\tilde{\varphi}_{\rho}$ is a linear isomorphism. The following proposition is the key for our method to reduce the set \mathcal{X}_{ρ_0} :

4.1.4 Proposition For any $\rho \in A(\mathcal{F}, S)$ and $x \in M$, consider the diagram

$$\begin{array}{ccc} S & \xrightarrow{a(x, \cdot)} & S \\ \tilde{q} \downarrow & & \downarrow \tilde{p} \\ S/K_{\rho} & \xrightarrow{\tilde{\varphi}_{\rho}} & S/H \end{array}$$

where the vertical maps are the natural projections. We give S a left-invariant Riemannian metric, and S/K_{ρ} and S/H the left-invariant Riemannian metrics considered in Corollary 4.1.2. Then $a(x, \cdot)$ is a fiber-respecting bi-Lipschitz diffeomorphism over $\tilde{\varphi}_{\rho}$.

Proof Let $q: \mathfrak{s} \rightarrow \mathfrak{s}/\ker \varphi_{\rho}$ and $p: \mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{h}$ be the natural projections. Then we have

$$e^{\int_1^s q^{\Theta}} = \tilde{q}(s) \quad \text{and} \quad e^{\int_1^s p^{\Theta}} = \tilde{p}(s)$$

for all $s \in S$. Indeed, writing $s = e^{X_1} \dots e^{X_k}$ for some $X_1, \dots, X_k \in \mathfrak{s}$,

$$e^{\int_1^s q^{\Theta}} = \exp\left(\int_1^{e^{X_1}} q^{\Theta} + \int_{e^{X_1}}^{e^{X_1}e^{X_2}} q^{\Theta} + \dots + \int_{e^{X_1} \dots e^{X_{k-1}}}^s q^{\Theta}\right) = e^{q(X_1) + \dots + q(X_k)}.$$

Let $\bar{\varphi}_\rho: \mathfrak{s}/\ker \varphi_\rho \rightarrow \mathfrak{s}/\mathfrak{h}$ be the map induced by φ_ρ . By exponentiating the equation $\bar{\varphi}_\rho(\int_1^s q^\Theta) + h(\rho_0(x, s)) - h(x) = \int_1^{a(x, s)} p^\Theta$ from Lemma 2.2.1, we get

$$\tilde{\varphi}_\rho \tilde{q}(s) e^{h(\rho_0(x, s)) - h(x)} = \tilde{p}(a(x, s)).$$

Since M is compact, there is a constant $C > 0$ such that

$$(9) \quad d(\tilde{\varphi}_\rho \tilde{q}(s), \tilde{p}(a(x, s))) = d(1, e^{h(\rho_0(x, s)) - h(x)}) < C$$

for any $s \in S$. This means that the diagram commutes up to bounded distance. Let f be the inverse map of $a(x, \cdot)$. By Lemma 2.2.1,

$$\int_1^{f(s)} q^\Theta + \bar{\varphi}_\rho^{-1}(h(\rho_0(x, f(s))) - h(x)) = \bar{\varphi}_\rho^{-1}\left(\int_1^s p^\Theta\right).$$

Exponentiating both sides of the equation,

$$\tilde{q} f(s) e^{\bar{\varphi}_\rho^{-1}(h(\rho_0(x, f(s))) - h(x))} = \tilde{\varphi}_\rho^{-1} \tilde{p}(s).$$

As before there is a constant $C' > 0$ such that

$$(10) \quad d(\tilde{q} f(s), \tilde{\varphi}_\rho^{-1} \tilde{p}(s)) < C'.$$

Let $C'' > 0$ be a constant such that $d(a(x, s_1), a(x, s_2)) \leq C'' d(s_1, s_2)$ for all $s_1, s_2 \in S$. Take any $b \in S/K_\rho$. Then we can show that

$$d_{\mathcal{H}}(a(x, \tilde{q}^{-1}(b)), \tilde{p}^{-1}(\tilde{\varphi}_\rho(b))) \leq \max\{C, C' C''\}$$

in the following way:

Claim $a(x, \tilde{q}^{-1}(b)) \in B(\tilde{p}^{-1}(\tilde{\varphi}_\rho(b)), C)$.

By (9), $d(\tilde{\varphi}_\rho(b), \tilde{p}(a(x, s))) < C$ for any $s \in \tilde{q}^{-1}(b)$. Therefore, $a(x, s)$ is in $B(\tilde{p}^{-1}(\tilde{\varphi}_\rho(b)), C)$.

Claim $\tilde{p}^{-1}(\tilde{\varphi}_\rho(b)) \in B(a(x, \tilde{q}^{-1}(b)), C' C'')$.

By (10), $d(\tilde{q} f(s), b) < C'$ for any $s \in \tilde{p}^{-1}(\tilde{\varphi}_\rho(b))$, so that we can find $s' \in \tilde{q}^{-1}(b)$ satisfying $d(s', f(s)) \leq C'$. Then $d(a(x, s'), s) \leq C'' d(s', f(s)) \leq C'' C'$. □

This proposition puts a strong restriction on the map $\tilde{\varphi}_\rho$ which we want to know, as in the next section or Section 5.2.

4.2 Proof of Theorem 1.0.5

As in Section 1, let

$$A_i = \begin{pmatrix} \alpha_1^{(i)} & & \\ & \ddots & \\ & & \alpha_n^{(i)} \end{pmatrix}$$

for $i = 1, \dots, k$ be diagonal matrices with positive diagonal entries and set $S_A = \mathbb{R}^n \rtimes_{A(t)} \mathbb{R}^k$, where $A(t) = A_1^{t_1} \cdots A_k^{t_k}$ for $t = (t_1, \dots, t_k) \in \mathbb{R}^k$. Ogasawara [8] proved the following theorem to classify up to quasiisometry groups obtained from \mathbb{Z}^n by performing certain HNN extensions several times, which is a generalization of a result of Farb and Mosher [3] on quasiisometric rigidity of abelian-by-cyclic groups.

4.2.1 Theorem *If*

$$\begin{array}{ccc} S_A & \xrightarrow{f} & S_{A'} \\ \downarrow & & \downarrow \\ \mathbb{R}^k & \xrightarrow{\varphi} & \mathbb{R}^k \end{array}$$

is a diagram in which vertical maps are natural projections and f is a fiber-respecting quasiisometry over some linear map φ , then there is a permutation matrix $P \in GL(n, \mathbb{R})$ such that $PA(t) = A'(\varphi(t))P$ for every $t \in \mathbb{R}^k$. In particular, there exists a diagram

$$\begin{array}{ccc} S_A & \xrightarrow{\sim} & S_{A'} \\ \downarrow & \wr & \downarrow \\ \mathbb{R}^k & \xrightarrow{\varphi} & \mathbb{R}^k \end{array}$$

where the upper horizontal map is an isomorphism of Lie groups taking (v, t) to $(Pv, \varphi(t))$.

Consider an action $M \overset{\rho_0}{\curvearrowright} S$ for $S = S_A$ and assume that for any j there is some i for which $\alpha_j^{(i)} \neq 1$. It is easy to check that this is equivalent to $[\mathfrak{s}, \mathfrak{s}] = \mathbb{R}^n$. We let $\mathfrak{h} = [\mathfrak{s}, \mathfrak{s}]$. For any $\rho \in A(\mathcal{F}, S)$, we have $\ker \varphi_\rho = [\mathfrak{s}, \mathfrak{s}]$ and then $K_\rho = \mathbb{R}^n$. Therefore, by Proposition 4.1.4, $a(x, \cdot)$ is a fiber-respecting bi-Lipschitz diffeomorphism over $\tilde{\varphi}_\rho$ in:

$$\begin{array}{ccc} S & \xrightarrow{a(x, \cdot)} & S \\ \downarrow & & \downarrow \\ \mathbb{R}^k & \xrightarrow{\tilde{\varphi}_\rho} & \mathbb{R}^k \end{array}$$

Hence, by the above theorem we get $\tilde{\varphi}_\rho \in W_S$. Using Theorem 2.2.5, this completes the proof of Theorem 1.0.5.

Remark There are solvable Lie groups for which this method does not work. Consider the solvable Lie group $S = \mathbb{R}^2 \rtimes_{R(t)} \mathbb{R}$ for

$$R(t) = \begin{pmatrix} \cos 2\pi t & -\sin 2\pi t \\ \sin 2\pi t & \cos 2\pi t \end{pmatrix}.$$

Then the standard Euclidean metric on \mathbb{R}^3 identified with S is a left-invariant Riemannian metric on S . Hence, for any $c \in \mathbb{R} \setminus \{0\}$ there is a bi-Lipschitz diffeomorphism $f: S \rightarrow S$ such that:

$$\begin{array}{ccc} S & \xrightarrow{f} & S \\ \downarrow & \circlearrowleft & \downarrow \\ \mathbb{R} & \xrightarrow{\times c} & \mathbb{R} \end{array}$$

So we do not have any restriction on c .

5 Parameter rigidity of suspension-like actions on mapping tori

We prove Theorem 1.0.6 here. The proof is based on Theorem 2.2.5, Proposition 4.1.4 and a theorem of Farb and Mosher [3]. To calculate cohomologies we use Mayer–Vietoris exact sequences.

The action which we consider is $M = \mathbb{Z} \backslash (\mathbb{T}^n \times \mathbb{R}) \curvearrowright S$, where $S = V \rtimes_{\Phi} \mathbb{R}$. Let \mathcal{F} be its orbit foliation.

5.1 Cohomology

The bracket operation of the Lie algebra $\mathfrak{s} = V \rtimes_{\Phi_*} \mathbb{R}$ is

$$\begin{aligned} [(v_1, t_1), (v_2, t_2)] &= ([v_1, v_2] + \Phi_*(t_1)v_2 - \Phi_*(t_2)v_1, [t_1, t_2]) \\ &= (\Phi_*(t_1)v_2 - \Phi_*(t_2)v_1, 0). \end{aligned}$$

Let $T = (0, 1) \in \mathfrak{s}$. We will use the notation $\text{ad}^0 X = \text{ad } X|_V$ for $X \in \mathfrak{s}$. Then $\text{ad}^0 T = \Phi_*(1)$, so that $\Phi_t = e^{t \text{ad}^0 T}$ and $A|_V = e^{\text{ad}^0 T}$. Since we assume 1 is not an eigenvalue of $A|_V$, $\text{ad}^0 T: V \rightarrow V$ does not have 0 as an eigenvalue, hence it is invertible. So we have $[\mathfrak{s}, \mathfrak{s}] = V$.

Let us examine which cohomologies to compute. We take $[\mathfrak{s}, \mathfrak{s}] = V$ as a subspace \mathfrak{h} appearing in Theorem 1.0.3. We consider representations $\mathfrak{s} \begin{smallmatrix} \text{ad}^0 \varphi \\ \curvearrowright \end{smallmatrix} \text{Gr}_V(\mathfrak{s}) = \mathfrak{s}/V \oplus V$ for $\varphi = \varphi_\rho$ for some $\rho \in A(\mathcal{F}, S)$. The first component $\mathfrak{s} \begin{smallmatrix} \text{ad}^0 \varphi \\ \curvearrowright \end{smallmatrix} \mathfrak{s}/V$ is just the trivial

representation. Let \bar{T} be the image of T under the natural projection $\mathfrak{s} \twoheadrightarrow \mathfrak{s}/V$ and put $\varphi(T) = c\bar{T}$. Then the second component $\mathfrak{s} \overset{\text{ad}^0\varphi}{\curvearrowright} V$ is given by $(v, t)v' = ct(\text{ad}^0 T)v' = c(\text{ad}^0(v, t))v'$, so that it is just $\mathfrak{s} \overset{c\text{ad}^0}{\curvearrowright} V$. Therefore we must compute

$$H^1(\mathcal{F}) \quad \text{and} \quad H^1(\mathcal{F}; \mathfrak{s} \overset{c\text{ad}^0}{\curvearrowright} V)$$

for all $c \in \mathbb{R}$. But we will see that vanishing of $H^1(\mathcal{F})$ and $H^1(\mathcal{F}; \mathfrak{s} \overset{\pm\text{ad}^0}{\curvearrowright} V)$ suffices by the method of large scale geometry and this reduction fits our assumption of Zariski density nicely.

5.2 Application of large scale geometry

Here we use rigidity of quasiisometries between solvable Lie groups found by Farb and Mosher. They prepare the following theorem as a tool to prove quasiisometric rigidity of abelian-by-cyclic groups. In [3, Theorem 5.2] a slightly different statement is given, but the following is also proved:

5.2.1 Theorem *Let $\Phi_t^{(1)}$ and $\Phi_t^{(2)}$ be one-parameter subgroups of $\text{GL}(p, \mathbb{R})$ and $S_i = \mathbb{R}^p \rtimes_{\Phi^{(i)}} \mathbb{R}$. If*

$$\begin{array}{ccc} S_1 & \xrightarrow{\phi} & S_2 \\ \downarrow & & \downarrow \\ \mathbb{R} & \xrightarrow{\text{id}} & \mathbb{R} \end{array}$$

is a diagram in which vertical maps are natural projections and ϕ is a fiber-respecting quasiisometry over id . Then absolute Jordan forms of $\Phi_1^{(1)}$ and $\Phi_1^{(2)}$ coincide up to permutation of Jordan blocks. (Here an absolute Jordan form refers to a matrix obtained by replacing the diagonal entries of a Jordan normal form over the complex field with their absolute values.)

Returning to our situation, we have

$$\begin{array}{ccc} S & \xrightarrow{a(x, \cdot)} & S \\ \downarrow & & \downarrow \\ \mathbb{R} & \xrightarrow{\times c} & \mathbb{R} \end{array}$$

where $a(x, \cdot)$ is a fiber-respecting bi-Lipschitz diffeomorphism by Proposition 4.1.4. Composing with

$$\begin{array}{ccc} S & \xrightarrow{\sim} & V \rtimes_{\Phi_{ct}} \mathbb{R} \\ \downarrow & \wr & \downarrow \\ \mathbb{R} & \xrightarrow{\times 1/c} & \mathbb{R} \end{array}$$

where the above horizontal map takes (v, t) to $(v, t/c)$, we get

$$\begin{array}{ccc} S & \longrightarrow & V \rtimes_{\Phi_{ct}} \mathbb{R} \\ \downarrow & & \downarrow \\ \mathbb{R} & \xrightarrow{\text{id}} & \mathbb{R} \end{array}$$

in which the above horizontal map is a fiber-respecting bi-Lipschitz diffeomorphism over id . Then, by Theorem 5.2.1, absolute Jordan forms of Φ_1 and Φ_c coincide. Let $\alpha_1, \dots, \alpha_p$ be the eigenvalues of $\text{ad}^0 T$, so that eigenvalues of $\Phi_t = e^{t \text{ad}^0 T}$ are $e^{t\alpha_1}, \dots, e^{t\alpha_p}$. Therefore the sets $\{|e^{\alpha_1}|, \dots, |e^{\alpha_p}|\}$ and $\{|e^{\alpha_1}|^c, \dots, |e^{\alpha_p}|^c\}$ coincide with multiplicity. Since Φ_1 has an eigenvalue of absolute value not equal to 1, we must have $c = \pm 1$. So we need to verify only the vanishing of $H^1(\mathcal{F})$ and $H^1(\mathcal{F}; \mathfrak{s}^{\pm \text{ad}^0 V})$ to show parameter rigidity of the action.

5.3 Mayer–Vietoris argument

Let $\mathfrak{s} \curvearrowright W$ be a representation on a finite-dimensional real vector space satisfying $\pi(v) = 0$ for all $v \in V$. We will try to calculate the first cohomology $H^1(\mathcal{F}; \mathfrak{s} \curvearrowright W)$.

Let U_1 and U_2 be the projections to M of $\mathbb{T}^n \times (0, 1)$ and $\mathbb{T}^n \times (-\frac{1}{2}, \frac{1}{2})$, so that $M = U_1 \cup U_2$. Then we have a short exact sequence of cochain complexes

$$0 \rightarrow \Omega^*(\mathcal{F}; W) \rightarrow \Omega^*(\mathcal{F}|_{U_1}; W) \oplus \Omega^*(\mathcal{F}|_{U_2}; W) \rightarrow \Omega^*(\mathcal{F}|_{U_1 \cap U_2}; W) \rightarrow 0.$$

The second map is $\xi \mapsto (\xi|_{U_1}, \xi|_{U_2})$ and the third is $(\xi_1, \xi_2) \mapsto \xi_2|_{U_1 \cap U_2} - \xi_1|_{U_1 \cap U_2}$. Hence, we obtain an exact sequence of cohomology

$$\begin{aligned} H^0(\mathcal{F}|_{U_1}; \pi) \oplus H^0(\mathcal{F}|_{U_2}; \pi) &\xrightarrow{P} H^0(\mathcal{F}|_{U_1 \cap U_2}; \pi) \\ &\rightarrow H^1(\mathcal{F}; \pi) \rightarrow H^1(\mathcal{F}|_{U_1}; \pi) \oplus H^1(\mathcal{F}|_{U_2}; \pi) \xrightarrow{Q} H^1(\mathcal{F}|_{U_1 \cap U_2}; \pi), \end{aligned}$$

so that we have

$$(11) \quad H^1(\mathcal{F}; \pi) \simeq \text{coker } P \oplus \text{ker } Q.$$

To compute $H^1(\mathcal{F}; \pi)$, we first compute $H^1(\mathcal{F}|_{U_1}; \pi)$. Let \mathcal{G} be the orbit foliation of the translation action $\mathbb{T}^n \curvearrowright V$. Define $\iota: \mathbb{T}^n \hookrightarrow U_1$ by $\iota(x) = (x, \frac{1}{2})$. The map $\iota^*: \Omega^*(\mathcal{F}|_{U_1}; W) \rightarrow \Omega^*(\mathcal{G}; W)$ obtained by pullback is a cochain map⁵ between $(\Omega^*(\mathcal{F}|_{U_1}; W), d_{\mathcal{F}} + \pi\omega_0 \wedge)$ and $(\Omega^*(\mathcal{G}; W), d_{\mathcal{G}})$, where ω_0 is the canonical 1-form of the action $M \curvearrowright S$. In fact, $\iota^*(d_{\mathcal{F}} + \pi\omega_0 \wedge)\xi = d_{\mathcal{G}}\iota^*\xi + \iota^*(\pi\omega_0 \wedge \xi) = d_{\mathcal{G}}\iota^*\xi$

⁵However, $p^*: \Omega^*(\mathcal{G}; W) \rightarrow \Omega^*(\mathcal{F}|_{U_1}; W)$ is not a cochain map, where $p: U_1 \rightarrow \mathbb{T}^n$ maps (x, t) to x .

for $\xi \in \Omega^k(\mathcal{F}|_{U_1}; W)$ due to our assumption on π . Thus we get the induced map on cohomologies

$$\iota^*: H^*(\mathcal{F}|_{U_1}; \pi) \rightarrow H^*(\mathcal{G}; W).$$

Note that W in $H^*(\mathcal{G}; W)$ is just a direct sum of the trivial coefficient \mathbb{R} .

5.3.1 Lemma $\iota^*: H^1(\mathcal{F}|_{U_1}; \pi) \rightarrow H^1(\mathcal{G}; W)$ is an isomorphism.

Proof We first show injectivity:

Take $[\eta] \in H^1(\mathcal{F}|_{U_1}; \pi)$ satisfying $\iota^*[\eta] = 0$. We must show the existence of a smooth $\alpha: U_1 \rightarrow W$ such that

$$(12) \quad \eta = d_{\mathcal{F}}\alpha + \pi\omega_0\alpha.$$

Such an α must satisfy $\eta(T) = T\alpha + \pi(T)\alpha$, where T is regarded as a vector field using the action. Note that T is just the vector field $\partial/\partial t$, where $(x, t) \in U_1$. We have a smooth map $h: \mathbb{T}^n \rightarrow W$ satisfying $\iota^*\eta = d_{\mathcal{G}}h$. Using this we let

$$\alpha(x, t) = e^{-(t-1/2)\pi(T)} \left(\int_{1/2}^t e^{(s-1/2)\pi(T)} \eta(T)(x, s) ds + h(x) \right)$$

for $x \in \mathbb{T}^n$ and $t \in (0, 1)$. This α satisfies $\partial\alpha(x, t)/\partial t = -\pi(T)\alpha(x, t) + \eta(T)(x, t)$, ie $\eta(T) = T\alpha + \pi(T)\alpha$ and $\alpha(x, \frac{1}{2}) = h(x)$. To show (12) we have to prove

$$\eta(X) - X\alpha = 0$$

for all $X \in V$. Note that

$$(13) \quad (\eta(X) - X\alpha)(x, \frac{1}{2}) = 0$$

for $X \in V$. Since $0 = (d_{\mathcal{F}}\eta + \pi\omega_0 \wedge \eta)(T, X) = T\eta(X) - X\eta(T) - \eta([T, X]) + \pi(T)\eta(X)$, we have

$$(14) \quad \begin{aligned} T(\eta(X) - X\alpha) &= X\eta(T) + \eta([T, X]) - \pi(T)\eta(X) - X(\eta(T) - \pi(T)\alpha) - [T, X]\alpha \\ &= -\pi(T)(\eta(X) - X\alpha) + \eta([T, X]) - [T, X]\alpha. \end{aligned}$$

Choose a basis of V which turns $\text{ad}^0 T$ into a real Jordan normal form

$$\bigoplus \left(\begin{array}{ccc} a & 1 & \\ & \ddots & \ddots \\ & & 1 & \\ & & & a \end{array} \right) \oplus \bigoplus \left(\begin{array}{ccccccc} b & -c & 1 & & & & \\ c & b & & 1 & & & \\ & & \ddots & & \ddots & & \\ & & & \ddots & & \ddots & \\ & & & & \ddots & & 1 \\ & & & & & & & 1 \\ & & & & & & & b & -c \\ & & & & & & & c & b \end{array} \right),$$

where $a, b, c \in \mathbb{R}$ and $c \neq 0$. Let X_1, \dots, X_k be the basis of the subspace corresponding to the matrix

$$\begin{pmatrix} a & 1 & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ & & & & a \end{pmatrix},$$

so that the relations

$$[T, X_1] = aX_1, \quad [T, X_2] = aX_2 + X_1, \quad \dots \quad [T, X_k] = aX_k + X_{k-1}$$

hold. By (14), $T(\eta(X_1) - X_1\alpha) = (a - \pi(T))(\eta(X_1) - X_1\alpha)$. The solution is

$$(\eta(X_1) - X_1\alpha)(x, t) = e^{(t-1/2)(a-\pi(T))}(\eta(X_1) - X_1\alpha)(x, \frac{1}{2}).$$

So we get $\eta(X_1) - X_1\alpha = 0$ by (13). For X_2 we also have $T(\eta(X_2) - X_2\alpha) = (a - \pi(T))(\eta(X_2) - X_2\alpha)$, so $\eta(X_2) - X_2\alpha = 0$. Repeating this we get $\eta(X_i) - X_i\alpha = 0$ for $i = 1, \dots, k$.

Let $X_1, Y_1, \dots, X_k, Y_k$ be the basis of the subspace corresponding to

$$\begin{pmatrix} b & -c & 1 & & & \\ c & b & & 1 & & \\ & & \ddots & & \ddots & \\ & & & \ddots & & \ddots \\ & & & & \ddots & 1 \\ & & & & & c & b \end{pmatrix}.$$

We proceed similarly. The relations are

$$\begin{aligned} [T, X_1] &= bX_1 + cY_1, & [T, Y_1] &= -cX_1 + bY_1, \\ [T, X_2] &= bX_2 + cY_2 + X_1, & [T, Y_2] &= -cX_2 + bY_2 + Y_1, \\ &\vdots & &\vdots \end{aligned}$$

The first equation to solve is

$$T \begin{pmatrix} \eta(X_1) - X_1\alpha \\ \eta(Y_1) - Y_1\alpha \end{pmatrix} = \begin{pmatrix} b - \pi(T) & c \\ -c & b - \pi(T) \end{pmatrix} \begin{pmatrix} \eta(X_1) - X_1\alpha \\ \eta(Y_1) - Y_1\alpha \end{pmatrix},$$

and the solution is

$$\begin{pmatrix} \eta(X_1) - X_1\alpha \\ \eta(Y_1) - Y_1\alpha \end{pmatrix}(x, t) = \exp\left(\left(t - \frac{1}{2}\right) \begin{pmatrix} b - \pi(T) & c \\ -c & b - \pi(T) \end{pmatrix}\right) \begin{pmatrix} \eta(X_1) - X_1\alpha \\ \eta(Y_1) - Y_1\alpha \end{pmatrix}(x, \frac{1}{2}).$$

So we have $\eta(X_1) - X_1\alpha = 0$ and $\eta(Y_1) - Y_1\alpha = 0$. Repeating, we eventually get $\eta(X_i) - X_i\alpha = 0$ and $\eta(Y_i) - Y_i\alpha = 0$ for $i = 1, \dots, k$. This proves the injectivity.

We now prove surjectivity:

Take any $[\xi] \in H^1(\mathcal{G}; W)$. We must construct $\eta \in \Omega^1(\mathcal{F}|_{U_1}; W)$ satisfying

$$(15) \quad d_{\mathcal{F}}\eta + \pi\omega_0 \wedge \eta = 0$$

and $\iota^*\eta = \xi$. We will construct η , requiring the additional property $\eta(T) = 0$. In order to satisfy (15), for any $X \in V$ we should construct $\eta(X)$ such that

$$(16) \quad T\eta(X) - \eta([T, X]) + \pi(T)\eta(X) = 0$$

holds. Fix a basis X_1, \dots, X_p of V and let (a_{ij}) be the matrix representing $\text{ad}^0 T$ with respect to X_1, \dots, X_p ; then $[T, X_j] = \sum_{i=1}^p a_{ij} X_i$. The $\eta(X_j)$ should satisfy $T\eta(X_j) - \sum_{i=1}^p a_{ij}\eta(X_i) + \pi(T)\eta(X_j) = 0$ and $\eta(X_j)(x, \frac{1}{2}) = \xi(X_j^0)$. Here, for $X \in V$, we denote by X^0 the section of $T\mathcal{G}$ satisfying $\iota_* X^0 = X|_{t=1/2}$. So we must solve

$$T \begin{pmatrix} \eta(X_1) \\ \vdots \\ \eta(X_p) \end{pmatrix} = \left((a_{ji}) - \begin{pmatrix} \pi(T) & & \\ & \ddots & \\ & & \pi(T) \end{pmatrix} \right) \begin{pmatrix} \eta(X_1) \\ \vdots \\ \eta(X_p) \end{pmatrix}.$$

The solution is

$$\begin{pmatrix} \eta(X_1) \\ \vdots \\ \eta(X_p) \end{pmatrix} (x, t) = \exp \left(\left(t - \frac{1}{2} \right) \begin{pmatrix} \pi(T) & & \\ & \ddots & \\ & & \pi(T) \end{pmatrix} \right) \begin{pmatrix} \xi(X_1^0) \\ \vdots \\ \xi(X_p^0) \end{pmatrix} (x).$$

We define $\eta(X_i)$ by this, then $\eta(X)$ for any $X \in V$ is defined by linearity, so that we have defined $\eta \in \Omega^1(\mathcal{F}|_{U_1}; W)$. Then (16) and $\iota^*\eta = \xi$ are satisfied. To see that (15) holds, we only have to show $(d_{\mathcal{F}}\eta + \pi\omega_0 \wedge \eta)(X, Y) = 0$ for all $X, Y \in V$. Put $\theta = d_{\mathcal{F}}\eta + \pi\omega_0 \wedge \eta$. Then

$$\begin{aligned} T(\theta(X, Y)) &= TX\eta(Y) - TY\eta(X) \\ &= X(\eta([T, Y]) - \pi(T)\eta(Y)) + [T, X]\eta(Y) \\ &\quad - Y(\eta([T, X]) - \pi(T)\eta(X)) - [T, Y]\eta(X) \\ &= -\pi(T)\theta(X, Y) + \theta(X, [T, Y]) + \theta([T, X], Y) \end{aligned}$$

and $\theta(X, Y)(x, \frac{1}{2}) = X^0\xi(Y^0) - Y^0\xi(X^0) = d_{\mathcal{G}}\xi(X^0, Y^0) = 0$. As in the proof of injectivity we can show $\theta(X_i, X_j) = 0$, so that (15) is satisfied. This completes the proof. □

By the assumption that V is a Diophantine subspace of \mathbb{R}^n , the leafwise cohomology $H^1(\mathcal{G}; W)$ vanishes, ie $H^1(\mathcal{G}; W) = H^1(V; W) = V^* \otimes W$. Thus $H^1(\mathcal{F}|_{U_1}; \pi) \simeq V^* \otimes W$. Let $q: \mathfrak{s} \rightarrow V$ be the natural projection.

5.3.2 Lemma *An element $\varphi \otimes w \in V^* \otimes W$ corresponds to*

$$(e^{(t-1/2)(\text{ad}^0 T)^*} \varphi)q\omega_0 \otimes e^{-(t-1/2)\pi(T)}w \in \Omega^1(\mathcal{F}|_{U_1}; W)$$

by the above isomorphism, where $(\text{ad}^0 T)^*: V^* \rightarrow V^*$ is the adjoint of $\text{ad}^0 T$.

Proof Put $\eta = (e^{(t-1/2)(\text{ad}^0 T)^*} \varphi)q\omega_0 \otimes e^{-(t-1/2)\pi(T)}w$. Then $\iota^* \eta = \varphi \otimes w$ and $\eta(T) = 0$. For $X \in V$, $\eta(X) = (e^{(t-1/2)(\text{ad}^0 T)^*} \varphi)(X)e^{-(t-1/2)\pi(T)}w$ satisfies $T\eta(X) = \eta([T, X]) - \pi(T)\eta(X)$. So we see $d_{\mathcal{F}}\eta + \pi\omega_0 \wedge \eta = 0$, as in the proof of Lemma 5.3.1. □

Now, $U_1 \cap U_2$ is the disjoint union of the projections to M of $\mathbb{T}^n \times (0, \frac{1}{2})$ and $\mathbb{T}^n \times (\frac{1}{2}, 1)$. We define the maps

$$\begin{aligned} \iota_{1/2}: \mathbb{T}^n &\hookrightarrow U_1, & \iota_{1/2}(x) &= (x, \frac{1}{2}), \\ \iota_0: \mathbb{T}^n &\hookrightarrow U_2, & \iota_0(x) &= (x, 0), \\ \iota_{1/4}: \mathbb{T}^n &\hookrightarrow U_1 \cap U_2, & \iota_{1/4}(x) &= (x, \frac{1}{4}), \\ \iota_{3/4}: \mathbb{T}^n &\hookrightarrow U_1 \cap U_2, & \iota_{3/4}(x) &= (x, \frac{3}{4}). \end{aligned}$$

We will calculate the bottom map of the next commutative diagram, in which the vertical maps are isomorphisms:

$$\begin{array}{ccc} H^1(\mathcal{F}|_{U_1}; \pi) \oplus H^1(\mathcal{F}|_{U_2}; \pi) & \xrightarrow{\mathcal{Q}} & H^1(\mathcal{F}|_{U_1 \cap U_2}; \pi) \\ \downarrow \iota_{1/2}^* \oplus \iota_0^* & \wr & \downarrow \iota_{1/4}^* \oplus \iota_{3/4}^* \\ (V^* \otimes W) \oplus (V^* \otimes W) & \longrightarrow & (V^* \otimes W) \oplus (V^* \otimes W) \end{array}$$

5.3.3 Lemma *The bottom map of the above diagram is*

$$\begin{pmatrix} -e^{-(\text{ad}^0 T)^*/4} \otimes e^{\pi(T)/4} & e^{(\text{ad}^0 T)^*/4} \otimes e^{-\pi(T)/4} \\ -e^{(\text{ad}^0 T)^*/4} \otimes e^{-\pi(T)/4} & e^{-(\text{ad}^0 T)^*/4} \otimes e^{\pi(T)/4} \end{pmatrix}.$$

Proof We define $F: \mathbb{T}^n \times \mathbb{R} \rightarrow \mathbb{T}^n \times \mathbb{R}$ by $F(x, t) = (A^{-1}x, t - 1)$. Take an element $(\varphi_1 \otimes w_1, \varphi_2 \otimes w_2)$ of $(V^* \otimes W) \oplus (V^* \otimes W)$. This is mapped by the vertical map to

$$((e^{(t-1/2)(\text{ad}^0 T)^*} \varphi_1)q\omega_0 \otimes e^{-(t-1/2)\pi(T)}w_1, (e^{t(\text{ad}^0 T)^*} \varphi_2)q\omega_0 \otimes e^{-t\pi(T)}w_2),$$

which is in turn mapped by Q to

$$\left((e^{t(\text{ad}^0 T)^*} \varphi_2) q \omega_0 \otimes e^{-t\pi(T)} w_2 - (e^{(t-1/2)(\text{ad}^0 T)^*} \varphi_1) q \omega_0 \otimes e^{-(t-1/2)\pi(T)} w_1, \right. \\ \left. F^* \left((e^{t(\text{ad}^0 T)^*} \varphi_2) q \omega_0 \otimes e^{-t\pi(T)} w_2 \right) - (e^{(t-1/2)(\text{ad}^0 T)^*} \varphi_1) q \omega_0 \otimes e^{-(t-1/2)\pi(T)} w_1 \right).$$

Since $F^* \left((e^{t(\text{ad}^0 T)^*} \varphi_2) q \omega_0 \otimes e^{-t\pi(T)} w_2 \right) = (e^{(t-1)(\text{ad}^0 T)^*} \varphi_2) q \omega_0 \otimes e^{-(t-1)\pi(T)} w_2$, the above element equals

$$\left((e^{(t-1/4)(\text{ad}^0 T)^*} e^{(\text{ad}^0 T)^*/4} \varphi_2) q \omega_0 \otimes e^{-(t-1/4)\pi(T)} e^{-\pi(T)/4} w_2 \right. \\ \left. - (e^{(t-1/4)(\text{ad}^0 T)^*} e^{-(\text{ad}^0 T)^*/4} \varphi_1) q \omega_0 \otimes e^{-(t-1/4)\pi(T)} e^{\pi(T)/4} w_1, \right. \\ (e^{(t-3/4)(\text{ad}^0 T)^*} e^{-(\text{ad}^0 T)^*/4} \varphi_2) q \omega_0 \otimes e^{-(t-3/4)\pi(T)} e^{\pi(T)/4} w_2 \\ \left. - (e^{(t-3/4)(\text{ad}^0 T)^*} e^{(\text{ad}^0 T)^*/4} \varphi_1) q \omega_0 \otimes e^{-(t-3/4)\pi(T)} e^{-\pi(T)/4} w_1 \right)$$

and finally this is mapped by the vertical arrow to

$$\left(e^{(\text{ad}^0 T)^*/4} \varphi_2 \otimes e^{-\pi(T)/4} w_2 - e^{-(\text{ad}^0 T)^*/4} \varphi_1 \otimes e^{\pi(T)/4} w_1, \right. \\ \left. e^{-(\text{ad}^0 T)^*/4} \varphi_2 \otimes e^{\pi(T)/4} w_2 - e^{(\text{ad}^0 T)^*/4} \varphi_1 \otimes e^{-\pi(T)/4} w_1 \right). \quad \square$$

By using the canonical isomorphism $V^* \otimes W \simeq \text{Hom}(V, W)$, the map in the above lemma becomes a map $\bar{Q}: \text{Hom}(V, W) \oplus \text{Hom}(V, W) \rightarrow \text{Hom}(V, W) \oplus \text{Hom}(V, W)$ which sends $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ to

$$\begin{pmatrix} -e^{\pi(T)/4} \circ \alpha \circ e^{-\text{ad}^0 T/4} + e^{-\pi(T)/4} \circ \beta \circ e^{\text{ad}^0 T/4} \\ -e^{-\pi(T)/4} \circ \alpha \circ e^{\text{ad}^0 T/4} + e^{\pi(T)/4} \circ \beta \circ e^{-\text{ad}^0 T/4} \end{pmatrix}.$$

5.3.4 Lemma If $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ is in the kernel of \bar{Q} , then α and β satisfy $e^{\pi(T)} \circ \alpha = \alpha \circ e^{\text{ad}^0 T}$ and $e^{\pi(T)} \circ \beta = \beta \circ e^{\text{ad}^0 T}$.

Proof The pair $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ is in the kernel if and only if $-e^{\pi(T)/2} \circ \alpha + \beta \circ e^{\text{ad}^0 T/2} = 0$ and $-\alpha \circ e^{\text{ad}^0 T/2} + e^{\pi(T)/2} \circ \beta = 0$. By eliminating β or α we get the conclusion. \square

Next we will calculate the map $H^0(\mathcal{F}|_{U_1}; \pi) \oplus H^0(\mathcal{F}|_{U_2}; \pi) \xrightarrow{P} H^0(\mathcal{F}|_{U_1 \cap U_2}; \pi)$. The group $H^0(\mathcal{F}|_{U_1}; \pi)$ consists of all smooth functions $f: U_1 \rightarrow W$ satisfying $d_{\mathcal{F}} f + \pi \omega_0 \wedge f = 0$, which is equivalent to the equations $Tf + \pi(T)f = 0$ and $Xf = 0$ for all $X \in V$. Since the action $\mathbb{T}^n \curvearrowright V$ has a dense orbit, such an f must be constant along the directions of tori; write $f(x, t) = f(t)$. Solving the differential equation, we

get $f(t) = e^{-(t-1/2)\pi(T)} f(\frac{1}{2})$. Thus we have an isomorphism $H^0(\mathcal{F}|_{U_1}; \pi) \simeq W$ which sends f to $f(\frac{1}{2})$. The bottom arrow of the diagram

$$\begin{array}{ccc}
 H^0(\mathcal{F}|_{U_1}; \pi) \oplus H^0(\mathcal{F}|_{U_2}; \pi) & \xrightarrow{P} & H^0(\mathcal{F}|_{U_1 \cap U_2}; \pi) \\
 \downarrow \iota_{1/2}^* \oplus \iota_0^* & \wr & \downarrow \iota_{1/4}^* \oplus \iota_{3/4}^* \\
 W \oplus W & \xrightarrow{\bar{P}} & W \oplus W
 \end{array}$$

sends an element (w_1, w_2) as follows:

$$\begin{aligned}
 (w_1, w_2) &\mapsto (e^{-(t-1/2)\pi(T)} w_1, e^{-t\pi(T)} w_2) \\
 &\mapsto (e^{-t\pi(T)} w_2 - e^{-(t-1/2)\pi(T)} w_1, F^*(e^{-t\pi(T)} w_2) - e^{-(t-1/2)\pi(T)} w_1) \\
 &= (e^{-t\pi(T)} w_2 - e^{-(t-1/2)\pi(T)} w_1, e^{-(t-1)\pi(T)} w_2 - e^{-(t-1/2)\pi(T)} w_1) \\
 &\mapsto (e^{-\pi(T)/4} w_2 - e^{\pi(T)/4} w_1, e^{\pi(T)/4} w_2 - e^{-\pi(T)/4} w_1).
 \end{aligned}$$

An element (w_1, w_2) is in $\ker \bar{P}$ if and only if $w_2 = e^{\pi(T)/2} w_1$ and $w_1 = e^{\pi(T)/2} w_2$. So,

$$(17) \quad \ker \bar{P} \simeq \ker(\text{id} - e^{\pi(T)}) \subset W$$

by $(w_1, w_2) \mapsto w_2$.

Finally we calculate the cohomology $H^1(\mathfrak{s}; \pi)$ of the Lie algebra. A 1-cocycle is a linear map $\varphi: \mathfrak{s} \rightarrow W$ satisfying $\pi(T)\varphi(X) - \varphi([T, X]) = 0$ for all $X \in V$. So the space of 1-cocycles is isomorphic to $\text{Hom}_T(V, W) \oplus W$ by the isomorphism which sends φ to $(\varphi|_V, \varphi(T))$, where $\text{Hom}_T(V, W)$ denotes the space of $(\text{ad}^0 T, \pi(T))$ -equivariant linear maps from V to W . On the other hand, a 1-coboundary maps $Z \in \mathfrak{s}$ to $\pi(Z)c \in W$ for some $c \in W$. So a linear map $\varphi: \mathfrak{s} \rightarrow W$ is a 1-coboundary if and only if $\varphi(X) = 0$ for all $X \in V$ and $\varphi(T) \in \text{im } \pi(T)$. The space of 1-coboundaries is isomorphic to $0 \oplus \text{im } \pi(T)$ by the above isomorphism. Therefore, we have

$$(18) \quad H^1(\mathfrak{s}; \pi) \simeq \text{Hom}_T(V, W) \oplus (W / \text{im } \pi(T)).$$

5.4 Vanishing of the cohomology

As we saw in Section 5.2 we must show vanishing of the following cohomologies:

- (a) The trivial representation $\mathfrak{s} \curvearrowright \mathbb{R}$.
- (b) The restriction of ad to V and its negative, $\mathfrak{s} \curvearrowright^{\pm \text{ad}^0} V$.

(a) By (18) in last section, $H^1(\mathfrak{s}) \simeq \{\varphi: V \rightarrow \mathbb{R} \mid \varphi \circ \text{ad}^0 T = 0\} \oplus \mathbb{R}$. By (11) and (17), $H^1(\mathcal{F}) \simeq \mathbb{R} \oplus \ker \bar{Q}$. If (α, β) is in $\ker \bar{Q}$, then, by Lemma 5.3.4, $\alpha = \alpha \circ \Phi_k$ for all $k \in \mathbb{Z}$. By the assumption of Zariski density, $\alpha = \alpha \circ \Phi_t$ for all $t \in \mathbb{R}$. Thus, $\alpha \circ \text{ad}^0 T = 0$ by differentiation. The same thing holds for β and then we have $\alpha = \beta$. So $\ker \bar{Q} \simeq \{\varphi: V \rightarrow \mathbb{R} \mid \varphi \circ \text{ad}^0 T = 0\}$ and $H^1(\mathcal{F}) = H^1(\mathfrak{s})$.

(b) By (18), $H^1(\mathfrak{s}; \pm \text{ad}^0) \simeq \{\varphi: V \rightarrow V \mid \varphi \circ \text{ad}^0 T = \pm \text{ad}^0 T \circ \varphi\}$. By (11) and (17), $H^1(\mathcal{F}; \pm \text{ad}^0) \simeq \ker \bar{Q}$. If (α, β) is in $\ker \bar{Q}$, then, by Lemma 5.3.4, $\Phi_{\pm k} \circ \alpha = \alpha \circ \Phi_k$ for all $k \in \mathbb{Z}$. By the assumption of Zariski density, $\Phi_{\pm t} \circ \alpha = \alpha \circ \Phi_t$ for all $t \in \mathbb{R}$. Thus, $\pm \text{ad}^0 T \circ \alpha = \alpha \circ \text{ad}^0 T$ by differentiation. Note that here we use the benefit of large scale geometry. Since the map $\ker \bar{Q} \rightarrow \{\varphi: V \rightarrow V \mid \varphi \circ \text{ad}^0 T = \pm \text{ad}^0 T \circ \varphi\}$ mapping (α, β) to α is injective, we have $H^1(\mathcal{F}; \pm \text{ad}^0) = H^1(\mathfrak{s}; \pm \text{ad}^0)$.

This completes the proof of Theorem 1.0.6.

6 Parameter rigidity of transitive locally free actions and rigidity of lattices

6.1 Relations between transitive locally free actions and lattices

Let S be a connected, simply connected solvable Lie group, Γ_0 be a cocompact lattice in S and $M = \Gamma_0 \backslash S \xrightarrow{\rho_0} S$ be the action by right multiplication.

We put

$$\mathcal{A}(\Gamma_0, S) := \{M \xrightarrow{\rho} S \mid \rho \text{ is transitive locally free}\},$$

$$\mathcal{H}(\Gamma_0, S) := \{\alpha: \Gamma_0 \rightarrow S \mid \alpha \text{ is an injective homomorphism and } \alpha(\Gamma_0) \text{ is a lattice}\}.$$

Then

$$\mathcal{H}(\Gamma_0, S) / \text{Aut}(\Gamma_0) = \{\Gamma \subset S \mid \Gamma \text{ is a cocompact lattice isomorphic to } \Gamma_0\}.$$

6.1.1 Proposition *There is a one-to-one correspondence between*

$$\mathcal{A}(\Gamma_0, S) / (C^\infty\text{-conjugacy} + \text{Aut}(S))$$

and

$$\text{Aut}(S) \backslash \mathcal{H}(\Gamma_0, S) / \text{Aut}(\Gamma_0)$$

$$= \text{Aut}(S) \backslash \{\Gamma \subset S \mid \Gamma \text{ is a cocompact lattice isomorphic to } \Gamma_0\}$$

taking ρ to the isotropy subgroup of ρ at the point $x_0 = \Gamma_0$.

Proof Well-definedness and injectivity are easy. For surjectivity, take any cocompact lattice Γ in S isomorphic to Γ_0 . Then, by a theorem of Mostow (see for example Raghunathan [9, Theorem 3.6]), $\Gamma \backslash S$ and $\Gamma_0 \backslash S = M$ are diffeomorphic. So the map is also surjective. \square

6.1.2 Proposition *There is a one-to-one correspondence between*

$$\mathcal{A}(\Gamma_0, S)/\text{parameter equivalence}$$

and

$$\text{Aut}(S) \backslash \mathcal{H}(\Gamma_0, S).$$

Proof The definition of the map is as follows: For a transitive locally free action $M \overset{\rho}{\curvearrowright} S$, let $a_\rho: M \times S \rightarrow S$ be the smooth map uniquely defined by $\rho_0(x, s) = \rho(x, a_\rho(x, s))$ and $a_\rho(x, 1) = 1$. Then a_ρ is a cocycle over ρ_0 :

$$a_\rho(x, ss') = a_\rho(x, s)a_\rho(\rho_0(x, s), s').$$

Let Γ_ρ be the isotropy subgroup of ρ at x_0 . Then we have $a_\rho(x_0, \cdot): \Gamma_0 \xrightarrow{\sim} \Gamma_\rho \subset S$. So we will define the map by $\rho \mapsto a_\rho(x_0, \cdot)$.

Let us see the well-definedness of this map. Take two transitive locally free actions $M \overset{\rho_i}{\curvearrowright} S$ for $i = 1, 2$ which are parameter equivalent. So there are $\Phi \in \text{Aut}(S)$ and a diffeomorphism $F: M \rightarrow M$ which is homotopic to the identity such that

$$(19) \quad F(\rho_1(x, s)) = \rho_2(F(x), \Phi(s)).$$

Let $b: M \times S \rightarrow S$ be the cocycle over ρ_1 defined by $\rho_1(x, s) = \rho_0(x, b(x, s))$. Note that $s = a_{\rho_1}(x, b(x, s))$. We have $\rho_1(x, s) = \rho_2(x, a_{\rho_2}(x, b(x, s)))$ and then $a_{\rho_2}(x, b(x, s))$ is a cocycle over ρ_1 . Since F is homotopic to the identity, we can define a smooth map $P: M \rightarrow S$ by $F(x) = \rho_2(x, P(x)^{-1})$. By (19), we see

$$\begin{aligned} a_{\rho_2}(x, b(x, s)) &= P(x)^{-1} \Phi(s) P(\rho_1(x, s)) \\ &= P(x)^{-1} \Phi(a_{\rho_1}(x, b(x, s))) P(\rho_0(x, b(x, s))), \end{aligned}$$

so that

$$a_{\rho_2}(x, s) = P(x)^{-1} \Phi(a_{\rho_1}(x, s)) P(\rho_0(x, s))$$

for all $s \in S$, since $b(x, \cdot): S \rightarrow S$ is invertible. Taking $x = x_0$ and $s = \gamma \in \Gamma_0$, we have

$$a_{\rho_2}(x_0, \gamma) = P(x_0)^{-1} \Phi(a_{\rho_1}(x_0, \gamma)) P(x_0).$$

So the map is well-defined.

Next let us show the injectivity. Take two transitive locally free actions $M \overset{\rho_i}{\curvearrowright} S$ for $i = 1, 2$ for which there exists $\Phi \in \text{Aut}(S)$ such that $a_{\rho_2}(x_0, \gamma) = \Phi(a_{\rho_1}(x_0, \gamma))$ for every $\gamma \in \Gamma_0$. Take $b: M \times S \rightarrow S$ satisfying $\rho_2(x, s) = \rho_0(x, b(x, s))$, so that $s = b(x, a_{\rho_2}(x, s))$. We have S -equivariant diffeomorphisms $\Gamma_{\rho_i} \backslash S \xrightarrow{\sim} M$ mapping $\Gamma_{\rho_i} s$ to $\rho_i(x_0, s)$. Using these, define a diffeomorphism F by:

$$\begin{array}{ccc} M & \xleftarrow{\sim} & \Gamma_{\rho_1} \backslash S \\ F \downarrow & \circlearrowleft & \downarrow \Phi \\ M & \xleftarrow{\sim} & \Gamma_{\rho_2} \backslash S \end{array}$$

Here Φ denotes the map $\Gamma_{\rho_1} s \mapsto \Gamma_{\rho_2} \Phi(s)$. Obviously $F(\rho_1(x, s)) = \rho_2(F(x), \Phi(s))$. To see that F is homotopic to the identity, take any $\gamma \in \Gamma_0 = \pi_1(M)$ and a curve $c: [0, 1] \rightarrow S$ connecting 1 and γ and consider the curve $\rho_0(x_0, c(\cdot)): [0, 1] \rightarrow M$. Since $\rho_0(x_0, c(t)) = \rho_1(x_0, a_{\rho_1}(x_0, c(t)))$, we have

$$F(\rho_0(x_0, c(t))) = \rho_2(x_0, \Phi(a_{\rho_1}(x_0, c(t)))) = \rho_0(x_0, b(x_0, \Phi(a_{\rho_1}(x_0, c(t))))).$$

Therefore, $F_*: \pi_1(M) \rightarrow \pi_1(M)$ maps γ to

$$b(x_0, \Phi(a_{\rho_1}(x_0, \gamma))) = b(x_0, a_{\rho_2}(x_0, \gamma)) = \gamma.$$

So F must be homotopic to the identity.

Finally, we show the surjectivity. Take any $\alpha: \Gamma_0 \rightarrow S$. Put $\Gamma := \alpha(\Gamma_0)$ and $y_0 := \Gamma \in \Gamma \backslash S$. We have an isomorphism $\alpha: \Gamma_0 \xrightarrow{\sim} \Gamma$ of the fundamental groups of (M, x_0) and $(\Gamma \backslash S, y_0)$. By Witte [10, Theorem 7.4], which is a refinement of Mostow’s theorem used in the proof of the previous proposition, we can find a diffeomorphism $F: (M, x_0) \rightarrow (\Gamma \backslash S, y_0)$, which induces $\alpha: \Gamma_0 \xrightarrow{\sim} \Gamma$ on the fundamental groups. Define $M \overset{\rho}{\curvearrowright} S$ by $\rho(x, s) = F^{-1}(F(x)s)$. Then ρ is a transitive, locally free action. We have $F(\rho_0(x_0, s)) = y_0 a_\rho(x_0, s)$ for all $s \in S$. Take any $\gamma_0 \in \Gamma_0$. Choose a curve $s: [0, 1] \rightarrow S$ connecting 1 and γ_0 . Then the curve $\rho_0(x_0, s(t))$ represents γ_0 in $\pi_1(M, x_0)$. Thus the curve $y_0 a_\rho(x_0, s(t))$ represents $F_*(\gamma_0) = \alpha(\gamma_0)$ in $\pi_1(\Gamma \backslash S, y_0)$. Since the curve $a_\rho(x_0, s(t))$ in S is the lift of the curve $y_0 a_\rho(x_0, s(t))$ starting from 1, we get $a_\rho(x_0, \gamma_0) = \alpha(\gamma_0)$. This proves the surjectivity. \square

6.1.3 Proposition *The map*

$$\mathcal{A}(\Gamma_0, S) \rightarrow \mathcal{H}(\Gamma_0, S)$$

defined in Proposition 6.1.2 is continuous. Here $\mathcal{A}(\Gamma_0, S)$ is endowed with the topology induced from the C^∞ compact–open topology of $C^\infty(M \times S, M)$ and $\mathcal{H}(\Gamma_0, S)$ has the topology of pointwise convergence.

Proof Take any $\gamma_0 \in \Gamma_0$. We must show that $a_\rho(x_0, \gamma_0) \in S$ is continuous with respect to ρ . The map $\mathcal{A}(\Gamma_0, S) \rightarrow C^\infty(S, M)$ which sends ρ to $\rho(x_0, \cdot)$ is continuous. Take a C^∞ -curve $c: [0, 1] \rightarrow S$ connecting 1 and γ_0 . Let $c_\rho: [0, 1] \rightarrow S$ be the lift of the curve $\rho_0(x_0, c(t))$ with respect to the covering $\rho(x_0, \cdot): S \rightarrow M$ starting at 1. Then $a_\rho(x_0, \gamma_0) = c_\rho(1)$. The map $\mathcal{A}(\Gamma_0, S) \rightarrow C^\infty([0, 1], S)$ taking ρ to c_ρ is continuous. In particular, $\rho \mapsto c_\rho(1) = a_\rho(x_0, \gamma_0)$ is continuous. □

6.1.4 Proposition We have a commutative diagram

$$\begin{array}{ccc}
 \mathcal{A}(\Gamma_0, S)/\text{parameter equivalence} & \xrightarrow{\sim} & \text{Aut}(S) \backslash \mathcal{H}(\Gamma_0, S) \\
 \downarrow & \circlearrowleft & \downarrow \\
 \mathcal{A}(\Gamma_0, S)/(C^\infty\text{-conjugacy} + \text{Aut}(S)) & \xrightarrow{\sim} & \text{Aut}(S) \backslash \mathcal{H}(\Gamma_0, S) / \text{Aut}(\Gamma_0)
 \end{array}$$

where the horizontal maps are as defined above and the vertical surjective maps are defined obviously.

Proof This is obvious. □

By combining Proposition 6.1.2 with Theorem 1.0.3, we get the following:

6.1.5 Corollary Let S be a connected, simply connected solvable Lie group, Γ_0 be a lattice in S and \mathfrak{h} be a subspace between $[\mathfrak{s}, \mathfrak{s}]$ and \mathfrak{n} . If

$$H^1(\Gamma_0 \backslash S; \mathfrak{s} \xrightarrow{\text{ad} \circ \varphi} \text{Gr}_{\mathfrak{h}}(\mathfrak{s})) = H^1(\mathfrak{s}; \mathfrak{s} \xrightarrow{\text{ad} \circ \varphi} \text{Gr}_{\mathfrak{h}}(\mathfrak{s}))$$

for all surjective Lie algebra homomorphisms $\varphi: \mathfrak{s} \rightarrow \mathfrak{s}/\mathfrak{h}$, then any injective homomorphism $\alpha: \Gamma_0 \rightarrow S$ whose image is a lattice is transformed into the inclusion $\Gamma_0 \hookrightarrow S$ by an element of $\text{Aut}(S)$. In particular, if Γ is a lattice in S isomorphic to Γ_0 , there is an isomorphism of S which transforms Γ into Γ_0 .

6.2 A counterexample to Theorem 1.0.1 for solvable Lie groups

Now we show a counterexample to Theorem 1.0.1 for solvable Lie groups. The next example is taken from Baues and Klopsch [2, Example 2.2], which is due to Milovanov [7]. Consider $A_0 = \begin{pmatrix} 0 & -1 \\ 1 & 3 \end{pmatrix} \in \text{SL}(2, \mathbb{Z})$, which has $\lambda = \frac{1}{2}(3 + \sqrt{5})$ and $\lambda^{-1} = \frac{1}{2}(3 - \sqrt{5})$ as eigenvalues, and let

$$X(t) = \begin{pmatrix} \lambda^t \cos 2\pi t & -\lambda^t \sin 2\pi t & & \\ \lambda^t \sin 2\pi t & \lambda^t \cos 2\pi t & & \\ & & \lambda^{-t} & \\ & & & \lambda^{-t} \end{pmatrix}$$

for $t \in \mathbb{R}$. Since $X(1)$ and $A = \begin{pmatrix} A_0 & \\ & A_0 \end{pmatrix} \in \mathrm{SL}(4, \mathbb{Z})$ are conjugate, the solvable Lie group $S = \mathbb{R}^4 \rtimes_{X(t)} \mathbb{R}$ contains a lattice Γ_0 isomorphic to $\mathbb{Z}^4 \rtimes_A \mathbb{Z}$. Then the set

$$\mathcal{H}(\Gamma_0, S)/\mathrm{Aut}(S)$$

is uncountable, as in Baues and Klopsch [2]. Therefore, by Proposition 6.1.2 the set

$$\mathcal{A}(\Gamma_0, S)/\text{parameter equivalence}$$

is uncountable, so that the natural action $\Gamma_0 \backslash S \curvearrowright S$ is not parameter rigid. However, by computing $\Gamma_0/[\Gamma_0, \Gamma_0]$ using $\Gamma_0 \simeq \mathbb{Z}^4 \rtimes_A \mathbb{Z}$, we can show $H^1(\Gamma_0 \backslash S) = H^1(\mathfrak{s})$.

6.3 A locally parameter rigid action of a contractible group which is not parameter rigid

Let $M \overset{\rho_0}{\curvearrowright} S$ be an action with the orbit foliation \mathcal{F} . The set of all smooth actions $M \curvearrowright S$ with the orbit foliation \mathcal{F} is denoted by $\mathcal{A}(\mathcal{F}, S)$, which is endowed with the topology induced from the C^∞ compact–open topology of $C^\infty(M \times S, M)$. An action $M \overset{\rho_0}{\curvearrowright} S$ is called *locally parameter rigid* if any $\rho \in \mathcal{A}(\mathcal{F}, S)$ which is close enough to ρ_0 is parameter equivalent to ρ_0 . In [1, Section 1.1.2], Asaoka comments that there is no known locally parameter rigid action of a contractible group which is not parameter rigid. Here we give an example of such an action. In [2, Example 2.5], Baues and Klopsch give an example of a connected, simply connected, solvable Lie group S which has a Zariski dense lattice Γ_0 such that $\mathrm{Aut}(S) \backslash \mathcal{H}(\Gamma_0, S)$ is countably infinite. Here Zariski density means that the Zariski closures of $\mathrm{Ad}(\Gamma_0)$ and $\mathrm{Ad}(S)$ in $\mathrm{GL}(\mathfrak{s})$ coincide. If we take such S and Γ_0 then the action $\Gamma_0 \backslash S \overset{\rho_0}{\curvearrowright} S$ by right multiplication is not parameter rigid, by Proposition 6.1.2. Let us show that this action is locally parameter rigid. Since Γ_0 is a Zariski dense lattice in S , the inclusion $\iota: \Gamma_0 \hookrightarrow S$ is *locally rigid*, that is, the $\mathrm{Aut}(S)$ –orbit of ι in $\mathcal{H}(\Gamma_0, S)$ is a neighborhood of ι . See for example Baues and Klopsch [2, Theorem 1.9]. Take an open neighborhood U of ι in $\mathcal{H}(\Gamma_0, S)$ which is contained in the $\mathrm{Aut}(S)$ –orbit of ι . Let V be the inverse image of U by the map $\mathcal{A}(\Gamma_0, S) \rightarrow \mathcal{H}(\Gamma_0, S)$; then V is an open neighborhood of ρ_0 in $\mathcal{A}(\Gamma_0, S)$ by continuity. Since U projects to a one-point set in $\mathrm{Aut}(S) \backslash \mathcal{H}(\Gamma_0, S)$, V also projects to a one-point set in $\mathcal{A}(\Gamma_0, S)/\text{parameter equivalence}$. Therefore, ρ_0 is locally parameter rigid.

References

- [1] **M Asaoka**, *Deformation of locally free actions and leafwise cohomology*, from “Foliations: dynamics, geometry and topology” (JA López, M Nicolau, editors), Springer, Basel (2014) 1–40 MR

- [2] **O Baues, B Klopsch**, *Deformations and rigidity of lattices in solvable Lie groups*, J. Topol. 6 (2013) 823–856 MR
- [3] **B Farb, L Mosher**, *On the asymptotic geometry of abelian-by-cyclic groups*, Acta Math. 184 (2000) 145–202 MR
- [4] **H Maruhashi**, *Parameter rigid actions of simply connected nilpotent Lie groups*, Ergodic Theory Dynam. Systems 33 (2013) 1864–1875 MR
- [5] **H Maruhashi**, *Vanishing of cohomology and parameter rigidity of actions of solvable Lie groups II*, preprint (2016)
- [6] **S Matsumoto, Y Mitsumatsu**, *Leafwise cohomology and rigidity of certain Lie group actions*, Ergodic Theory Dynam. Systems 23 (2003) 1839–1866 MR
- [7] **M V Milovanov**, *The extension of automorphisms of uniform discrete subgroups of solvable Lie groups*, Dokl. Akad. Nauk BSSR 17 (1973) 892–895 MR In Russian
- [8] **N Ogasawara**, *Quasiisometric classification of groups obtained from \mathbb{Z}^n by HNN extension performed several times*, Masters thesis, Kyoto University (2012) In Japanese
- [9] **MS Raghunathan**, *Discrete subgroups of Lie groups*, Ergeb. Math. Grenzgeb. 68, Springer, Berlin (1972) MR
- [10] **D Witte**, *Superrigidity of lattices in solvable Lie groups*, Invent. Math. 122 (1995) 147–193 MR

Max Planck Institute for Mathematics
Vivatsgasse 7, D-53111 Bonn, Germany
h-maruha@math.kyoto-u.ac.jp

Proposed: Martin Bridson
Seconded: Walter Neumann, Benson Farb

Received: 5 May 2014
Revised: 7 August 2015

