

Analytic nonabelian Hodge theory

JONATHAN PRIDHAM

The proalgebraic fundamental group can be understood as a completion with respect to finite-dimensional noncommutative algebras. We introduce finer invariants by looking at completions with respect to Banach and C^* -algebras, from which we can recover analytic and topological representation spaces, respectively. For a compact Kähler manifold, the C^* -completion also gives the natural setting for nonabelian Hodge theory; it has a pure Hodge structure, in the form of a pro- C^* -dynamical system. Its representations are pluriharmonic local systems in Hilbert spaces, and we study their cohomology, giving a principle of two types, and splittings of the Hodge and twistor structures.

32G13, 32G20

Introduction

Simpson [24; 25] defined the coarse Betti, de Rham and Dolbeault moduli spaces of a smooth projective complex variety. These are all algebraic spaces, with a complex analytic isomorphism between the Betti and de Rham moduli spaces. The nonabelian Hodge theorem of [25, Theorem 7.18] is a homeomorphism between the de Rham and Dolbeault moduli spaces, the key to which was the correspondence between semisimple local systems, pluriharmonic bundles and Higgs bundles.

The reductive proalgebraic fundamental group $\pi_1(X, x)^{\text{red}}$ introduced by Simpson [23] encapsulates, in a single object, all the information about the category of semisimple local systems. When X is a compact Kähler manifold, the group scheme $\pi_1(X, x)^{\text{red}}$ also has a pure Hodge structure in the form of a discrete circle action, and a description in terms of Higgs bundles.

However, it has long been realised that the reductive proalgebraic fundamental group is slightly inadequate. From it we can recover the points of the Betti moduli space, and from the full proalgebraic fundamental group we can even recover their infinitesimal neighbourhoods, but in general these groups convey no information about how the neighbourhoods glue together. A further source of dissatisfaction is the discontinuity of the circle action on $\pi_1(X, x)^{\text{red}}$, since it is continuous on moduli spaces.

The key idea behind this paper is that we can produce finer and more satisfactory invariants by looking at representations with analytic structure, an approach previously considered by Sullivan [29, Section 9]. The group scheme $\pi_1(X, x)^{\text{red}}$ can be recovered from representations in finite-dimensional matrix algebras, but the Riemann–Hilbert correspondence between Betti and de Rham moduli holds with coefficients in any Banach algebra. We accordingly construct Betti, de Rham and Dolbeault moduli functors on Banach algebras, and recover the analytic moduli spaces from these functors. The framed Betti and de Rham functors are represented by a Fréchet algebra which we regard as the analytic completion of $\mathbb{R}[\pi_1(X, x)]$.

To understand the topological structure underlying these analytic spaces, we then restrict to C^* -algebras rather than Banach algebras. There are notions of unitary and pluriharmonic representations with coefficients in any C^* -algebra, and the homeomorphism of moduli spaces above extends to an isomorphism between the semisimple de Rham functor and the polystable Dolbeault functor on polynormal C^* -algebras, via isomorphisms with the pluriharmonic functor. The C^* -algebra of bounded operators gives us a notion of pluriharmonic local systems in Hilbert spaces, and there is a form of Hodge decomposition for these local systems.

Lurking behind these comparisons is the twistor moduli functor on multiplicatively convex Fréchet $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ -algebras. Its fibres at $\pm i \in \mathbb{P}^1$ are the Dolbeault functor and its conjugate, while all other fibres are isomorphic to the de Rham functor. The Deligne–Hitchin twistor space can be recovered as an analytic space from the twistor moduli functor, and pluriharmonic torsors give a splitting of the twistor moduli functor on C^* -algebras over $C(\mathbb{P}^1(\mathbb{C}))$. Twistor cochains then admit a Hodge decomposition on pulling back along the Hopf fibration $\text{SU}_2 \rightarrow \mathbb{P}^1(\mathbb{C})$, and a continuous circle action serves to promote twistor structures to Hodge structures.

The structure of the paper is as follows. In Section 1, we cover some background material on prorepresentability. Proposition 1.19 then establishes a topological analogue of Tannaka duality for polynormal C^* -algebras and unitary representations, while Lemma 1.28 gives a similar result for nonunitary representations.

In Section 2, we introduce the framed Betti and de Rham functors $\mathbf{R}_{X,x}^B$ and $\mathbf{R}_{X,x}^{\text{dR}}$ on Banach algebras for any manifold X . In Proposition 2.3, we establish an isomorphism $\mathbf{R}_{X,x}^B(A) \cong \mathbf{R}_{X,x}^{\text{dR}}(A)$ for any pro-Banach algebra A . We can even recover the analytic structure of moduli spaces of G -bundles from these symmetric monoidal functors (Remark 2.6). Proposition 2.10 then shows that $\mathbf{R}_{X,x}^B$ is represented by a Fréchet algebra completion $E_{X,x}^B$ of $\mathbb{R}[\pi_1(X, x)]$. This is a Fréchet bialgebra, from which $\pi_1(X, x)$ can be recovered (Lemma 2.11).

In Definition 2.18, we introduce a symmetric monoidal functor $R_{X,x}^J$ on C^* -algebras, parametrising pluriharmonic bundles on a compact Kähler manifold X . This is representable by a pro- C^* -bialgebra $E_{X,x}^J$ (Proposition 2.22 and Lemma 2.24). There are also a symmetric monoidal functor $R_{X,x}^{\text{Dol}}$ on Banach algebras associated to Dolbeault moduli, which is seldom representable, and a harmonic functor extending the definition of $R_{X,x}^J$ to all Riemannian manifolds X , but with substantial loss of functoriality.

In Section 3 we establish relations between the various functors. We can recover the topology on moduli spaces of semisimple representations from $E_{X,x}^J$ (Theorem 3.6). Proposition 3.8 then gives a Tannakian description of the polynormal completion of $E_{X,x}^J$, while Corollary 3.11 gives a simple characterisation of continuous morphisms from $E_{X,x}^J$ to polynormal C^* -algebras. Section 3.3 then gives similar results for $R_{X,x}^{\text{Dol}}$. Lemma 3.19 shows that grouplike elements $G((E_{X,x}^J)^{\text{ab}})$ of the abelianisation of $E_{X,x}^J$ are just $H_1(X, \mathbb{Z} \oplus \mathbb{R})$, with consequences for complex tori.

There is a continuous circle action on $E_{X,x}^J$, so it is a pro- C^* -dynamical system (Proposition 3.31). This allows us to regard $E_{X,x}^J$ as an analytic nonabelian Hodge structure of weight 0 (Remark 3.32). In Example 3.33, we see that the circle action on $G((E_{X,x}^J)^{\text{ab}})$ is just given by the Hodge structure on $H^1(X, \mathbb{R})$. Proposition 3.38 then characterises pure Hilbert variations of Hodge structure as representations of $E_{X,x}^J \rtimes S^1$.

Section 4 is concerned with Hilbert space representations of $E_{X,x}^J$, which correspond to pluriharmonic local systems \mathbb{V} in Hilbert spaces. We can identify reduced cohomology $\bar{H}^*(X, \mathbb{V})$ with the space of smooth \mathbb{V} -valued harmonic forms, as well as establishing the principle of two types and a formality results (Section 4.2.1). There are analogous, but weaker, results for nonreduced cohomology (Section 4.2.2). The same is true of direct limits of Hilbert space representations, and Corollary 4.22 shows that the universal such is the continuous dual $(E_{X,x}^J)'$ (which can be regarded as the predual of the W^* -envelope of $E_{X,x}^J$).

In Section 5, these results are extended to show that the natural twistor structure on $\bar{H}^n(X, \mathbb{V})$ is pure of weight n (Corollary 5.5), with a weaker result for nonreduced cohomology (Proposition 5.6). If \mathbb{E} is the local system associated to the $\pi_1(X, x)$ -representation $E_{X,x}^J$, then Proposition 5.10 shows that this twistor structure can be enhanced to a form of analytic Hodge filtration on the de Rham algebra $A^\bullet(X, \mathbb{E}')$. In Section 5.4, we reinterpret splittings of twistor structures and Archimedean monodromy in terms of the Hopf fibration.

Finally, in Section 6, we introduce a whole twistor family of framed moduli functors on multiplicatively convex Fréchet algebras over $\mathbb{P}^1(\mathbb{C})$, from which we can recover both de Rham and Dolbeault functors. The coarse quotient of the framed twistor space

is just the Deligne–Hitchin twistor space (Remark 6.17). The twistor family carries a natural involution σ , and we show in Proposition 6.13 that σ -equivariant sections of the framed twistor space are just framed pluriharmonic bundles. Theorem 6.20, Proposition 6.22 and Corollary 6.24 then give analogues of Theorem 3.6, Proposition 3.8 and Corollary 3.11 in the twistor setting, describing twistors with C^* -algebra coefficients and the topology of the twistor space.

Notation

We will use k to denote either of the fields \mathbb{R} or \mathbb{C} .

Definition 0.1 Given a k -Hilbert space H , write $L(H)$ for the space of k -linear bounded operators on H , with the norm topology.

Definition 0.2 Given topological spaces X and Y , we write $C(X, Y)$ for the set of continuous maps from X to Y .

Definition 0.3 Given a group G acting on a set X , write $[X/G]$ for the groupoid with objects X and morphisms $X \times G$, where the source of (x, g) is x and the target is xg . Composition of morphisms is given by $(xg, h) \circ (x, g) = (x, gh)$.

Definition 0.4 Given a group G acting on sets S and T , write $S \times_G T$ for the quotient of $S \times T$ by the G -action $g(s, t) = (gs, g^{-1}t)$.

Acknowledgements I would like to thank Carlos Simpson for originally posing the problem of finding finer invariants and for helpful discussions. I would also like to thank the anonymous referee for spotting erroneous statements about Fréchet algebras.

This work was supported by the Engineering and Physical Sciences Research Council (grant numbers EP/F043570/1 and EP/I004130/1).

1 Prorepresentability of functors on unital Banach algebras and C^* -algebras

Definition 1.1 Given a functor $F: \mathcal{C} \rightarrow \text{Set}$, an object $A \in \mathcal{C}$ and an element $\xi \in F(A)$, we follow [13, Section A.3] in saying that the pair (A, ξ) is *minimal* if, for any pair (A', ξ') , any strict monomorphism $f: A' \rightarrow A$ with $F(f)(\xi') = \xi$ must be an isomorphism.

We say that a pair (A'', ξ'') is *dominated* by a minimal pair if there exists a minimal pair (A, ξ) and a morphism $g: A \rightarrow A''$ with $F(g)(\xi) = \xi''$.

Definition 1.2 As in [13, Section A.3], we say that a functor $F: \mathcal{C} \rightarrow \text{Set}$ on a category \mathcal{C} containing all finite limits is *left exact* if it preserves all finite limits. This is equivalent to preserving finite products and equalisers, or to preserving fibre products and the final object.

Lemma 1.3 Let \mathcal{C} be a category containing finite limits and take a left exact functor $F: \mathcal{C} \rightarrow \text{Set}$. Assume that, for any cofiltered inverse system $\{A_i\}_i$ of strict subobjects of any object $A \in \mathcal{C}$, the limit $\varprojlim_i A_i$ exists, and that the map

$$F(\varprojlim_i A_i) \rightarrow \varprojlim_i F(A_i)$$

is an isomorphism.

Then every pair $(A, \xi \in F(A))$ is dominated by a minimal pair.

Proof Given the pair (A, ξ) , let I be the full subcategory of the overcategory $\mathcal{C} \downarrow A$ consisting of strict monomorphisms $B \rightarrow A$ for which ξ lifts to $F(B)$. Note that this lift must be unique by the monomorphism property. If $f \circ g$ is a strict monomorphism, then so is g , which implies that all morphisms in I must be strict monomorphisms in \mathcal{C} . Moreover, left-exactness of F guarantees that I is closed under the fibre product \times_A . the monomorphism properties imply that parallel arrows in I are equal, so I is a cofiltered category.

By hypothesis, the limit $L := \varprojlim_{B \in I} B$ exists in \mathcal{C} . It is necessarily a strict subobject of A , since it is the limit of all parallel maps sourced at A and equalised by some $B \in I$. The unique lifts of ξ to each $F(B)$ define an element of

$$\varprojlim_{B \in I} F(B),$$

so by hypothesis we have a corresponding element $\eta \in F(L)$. Therefore L is an object of I and is in fact the initial object of I . The pair (L, η) is therefore minimal, and dominates (A, ξ) as required. □

Definition 1.4 Recall from [13, Section A.2] that a proobject $X \in \text{pro}(\mathcal{C})$ is said to be *strict* if it is isomorphic to a proobject of the form $\{X_i\}$, where each map $X_i \rightarrow X_j$ is an epimorphism.

A functor $F: \mathcal{C} \rightarrow \text{Set}$ is said to be *strictly prorepresentable* if there exists a strict proobject X with $F \cong \text{Hom}_{\text{pro}(\mathcal{C})}(X, -)$.

Proposition 1.5 Let \mathcal{C} be a category containing finite limits and limits of cofiltered inverse systems of strict subobjects. Then a functor $F: \mathcal{C} \rightarrow \text{Set}$ is strictly prorepresentable if and only if

- (1) F is left exact;
- (2) F preserves limits of cofiltered inverse systems of strict subobjects.

Proof If F satisfies the conditions, then it is left-exact, and by Lemma 1.3 every pair is dominated by a minimal pair. It therefore satisfies the conditions of [13, Proposition A.3.1], so is strictly prorepresentable.

Conversely, every prorepresentable functor F is left-exact, so we need only show that the second condition holds. Write $F = \varprojlim_{\alpha} \text{Hom}(R_{\alpha}, -)$ for a strict inverse system $\{R_{\alpha}\}_{\alpha}$ and take a cofiltered inverse system $\{A_i\}_i$ of strict subobjects of some object $A \in \mathcal{C}$.

Given an element $x \in \varprojlim_i F(A_i)$ with image x_i in $F(A_i)$, by definition there exist objects R_{α_i} and maps $y_i: R_{\alpha_i} \rightarrow A_i$ lifting x_i . Now fix i ; the liftings are compatible in the sense that for $j > i$ (increasing α_j if necessary), there is a commutative diagram:

$$\begin{array}{ccc} R_{\alpha_j} & \xrightarrow{y_j} & A_j \\ \downarrow & & \downarrow \\ R_{\alpha_i} & \xrightarrow{y_i} & A_i \end{array}$$

Since $A_j \rightarrow A_i$ is a strict monomorphism and $R_{\alpha_j} \rightarrow R_{\alpha_i}$ an epimorphism, y_i must lift to a map $y_{ij}: R_{\alpha_i} \rightarrow A_j$. Since $A_j \rightarrow A_i$ is a monomorphism, the lifting y_{ij} is unique. Considering all $j > i$ together, this gives us a unique map $\tilde{x}: R_{\alpha_i} \rightarrow \varprojlim_j A_j$, which gives rise to a unique preimage $\tilde{x} \in F(\varprojlim_j A_j)$, as required. \square

1.1 Banach algebras

Definition 1.6 Write BanAlg_k for the category of unital (not necessarily commutative) Banach algebras over k , with bounded morphisms.

Proposition 1.7 Take a functor $F: \text{BanAlg}_k \rightarrow \text{Set}$ such that

- (1) F preserves all finite limits (equivalently preserves fibre products and the final object);
- (2) F preserves monomorphisms (ie maps closed subalgebras to subsets); and
- (3) for all inverse systems S of closed subalgebras, the map

$$F\left(\bigcap_{s \in S} A_s\right) \rightarrow \bigcap_{s \in S} F(A_s)$$

is an isomorphism.

Then F is strictly prorepresentable.

Proof Given a Banach algebra B , every strict subobject of B is a closed subalgebra. For any cofiltered inverse system $\{A_i\}_i$ of strict subobjects of any object B , the limit $\varprojlim_i A_i$ exists in BanAlg_k and is given by $\bigcap_i A_i$. The result now follows from Proposition 1.5. \square

1.2 C^* -algebras

Definition 1.8 Write $C^*\text{Alg}_k$ for the category of unital (not necessarily commutative) C^* -algebras over k , with bounded involutive morphisms.

Explicitly, a complex C^* -algebra is a complex Banach algebra equipped with an antilinear involution $*$ satisfying $(ab)^* = b^*a^*$ and $\|a^*a\| = \|a\|^2$ for all $a \in A$.

A real C^* -algebra is a real Banach algebra equipped with a linear involution $*$ satisfying the conditions above and having the additional property that $1 + a^*a$ is invertible for all $a \in A$.

A Banach $*$ -algebra over k is a C^* -algebra if and only if it is isometrically $*$ -isomorphic to a self-adjoint norm-closed algebra of bounded operators on a Hilbert k -space; for $k = \mathbb{R}$, this is Engelstam’s theorem [11, Proposition 8.2 and Theorem 15.3].

Proposition 1.9 Take a functor $F: C^*\text{Alg}_k \rightarrow \text{Set}$ such that

- (1) F preserves all finite limits (equivalently preserves fibre products and the final object);
- (2) F preserves monomorphisms (ie maps C^* -subalgebras to subsets); and
- (3) for all inverse systems S of nested C^* -subalgebras, the map

$$F\left(\bigcap_{s \in S} A_s\right) \rightarrow \bigcap_{s \in S} F(A_s)$$

is an isomorphism.

Then F is strictly prorepresentable.

Proof The proof of Proposition 1.7 carries over. \square

Lemma 1.10 Every complex C^* -algebra becomes a real C^* -algebra by forgetting the multiplication by \mathbb{C} .

Proof It suffices to show that $1 + a^*a$ is invertible for all $a \in A$. Now, $x \mapsto (1 + |x|)^{-1}$ is a continuous function on \mathbb{R} and a^*a is positive self-adjoint, so the continuous functional calculus implies that $(1 + a^*a)^{-1} \in A$, as required. \square

Lemma 1.11 *The category $C^* \text{Alg}_{\mathbb{R}}$ is equivalent to the category of pairs (A, τ) for $A \in C^* \text{Alg}_{\mathbb{C}}$ and an involution $\tau: A \rightarrow A$ satisfying*

- (1) $\tau(ab) = \tau(a)\tau(b)$;
- (2) $\tau(a)^* = \tau(a^*)$; and
- (3) $\tau(\lambda) = \bar{\lambda}$ for $\lambda \in \mathbb{C}$.

Proof Given $B \in C^* \text{Alg}_{\mathbb{R}}$, set $A := B \otimes_{\mathbb{R}} \mathbb{C}$; this is a complex C^* -algebra, with involution $(b \otimes \lambda)^* = b^* \otimes \bar{\lambda}$. The involution τ is then given by complex conjugation, with $\tau(b \otimes \lambda) = b \otimes \bar{\lambda}$. For the quasi-inverse construction, we send a pair (A, τ) to the algebra A^τ of τ -invariants. That this is a real C^* -algebra follows from Lemma 1.10.

To see that these are quasi-inverse functors, first note that $(B \otimes_{\mathbb{R}} \mathbb{C})^\tau = B$. Next, observe that because τ is antilinear, we can write $A = A^\tau \oplus iA^\tau \cong A^\tau \otimes_{\mathbb{R}} \mathbb{C}$ for all pairs (A, τ) as above. □

Definition 1.12 For a complex (resp. real) $*$ -algebra A , write $U(A)$ for the group of unitary (resp. orthogonal) elements

$$\{a \in A \mid a^*a = aa^* = 1\}.$$

Write $\mathfrak{u}(A)$ for the Lie algebra of anti-self-adjoint elements

$$\{a \in A \mid a^* + a = 0\}$$

and write $S(A)$ for the self-adjoint elements of A , noting that $\mathfrak{u}(A) = iS(A)$ when A is complex.

1.3 Representations and polynomial C^* -algebras

1.3.1 Representation spaces Fix a real unital C^* -algebra A .

Definition 1.13 Write $\text{Rep}_n^*(A)$ for the space of unital continuous $*$ -homomorphisms $\rho: A \rightarrow \text{Mat}_n(\mathbb{C})$ equipped with the topology of pointwise convergence. Write $\text{Irr}_n^*(A) \subset \text{Rep}_n^*(A)$ for the subspace of irreducible representations.

Definition 1.14 Define FDHilb to be the groupoid of complex finite-dimensional Hilbert spaces and unitary isomorphisms.

Definition 1.15 Write $\text{FD}^* \text{Rep}(A)$ for the groupoid of pairs (V, ρ) for $V \in \text{FDHilb}$ and $\rho: A \rightarrow \text{End}(V)$ a unital continuous $*$ -homomorphism. Morphisms are given by unitary isomorphisms intertwining representations. The set of objects of $\text{FD}^* \text{Rep}(A)$ is given the topology of pointwise convergence.

Definition 1.16 Define A_{PN} to be the ring of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant continuous additive endomorphisms of the fibre functor from $\text{FD}^*\text{Rep}(A)$ to vector spaces. Explicitly, $a \in A_{\text{PN}}$ associates to each pair (V, ρ) an element $a(V, \rho) \in \text{End}(V)$, subject to the conditions:

- (1) For any unitary isomorphism $u: V \rightarrow W$, we have $a(W, u\rho u^{-1}) = ua(V, \rho)u^{-1}$.
- (2) For any $(V_1, \rho_1), (V_2, \rho_2) \in \text{FD}^*\text{Rep}(A)$, we have $a(V_1 \oplus V_2, \rho_1 \oplus \rho_2) = a(V_1, \rho_1) \oplus a(V_2, \rho_2)$.
- (3) The maps $a: \text{Rep}_n^*(A) \rightarrow \text{Mat}_n(\mathbb{C})$ are continuous and $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant.

Lemma 1.17 The ring A_{PN} has the structure of a pro- C^* -algebra over A .

Proof We can describe A_{PN} as the categorical limit of a diagram of $*$ -homomorphisms between the C^* -algebras $C(\text{Rep}_n^*(A), \text{Mat}_n(\mathbb{R}))$, thus making it into a pro- C^* -algebra. The $*$ -homomorphism $A \rightarrow A_{\text{PN}}$ is given by mapping a to the transformation $a(V, \rho) = \rho(a)$. □

1.3.2 Polynormal C^* -algebras

Definition 1.18 Recall from [17] that an algebra A is said to be n -normal if, for all $a_1, \dots, a_{2n} \in A$, we have

$$\sum_{\sigma \in S_{2n}} \text{sgn}(\sigma) a_{\sigma(1)} a_{\sigma(2)} \cdots a_{\sigma(2n)} = 0.$$

Call an algebra *polynormal* if it is n -normal for some n .

By the Amitsur–Levitzki theorem, the algebra of $n \times n$ -matrices over a commutative ring is n -normal. Also note that an n -normal algebra is k -normal for all $k \geq n$.

Note that by restricting to n -dimensional representations we get that, for any real C^* -algebra A , the ring A_{PN} of Definition 1.16 is an inverse limit $A_{\text{PN}} = \varprojlim_n A_{\text{PN},n}$ of n -normal C^* -algebras.

We now have a result combining aspects of Tannaka and Takesaki duality:

Proposition 1.19 If A is a polynormal unital C^* -algebra then the morphism $A \rightarrow A_{\text{PN}}$ of Lemma 1.17 is an isomorphism.

Proof Since A is N -normal for some integer N , [17, Section 3] implies that A is of type I , with all complex irreducible representations of A having dimension at most N .

For a sufficiently large cardinal α , Bichteler [3] characterises the W^* -envelope $A'' \otimes \mathbb{C}$ of $A \otimes \mathbb{C}$ as the ring defined analogously to A_{PN} by replacing “continuous” with “bounded” and “finite-dimensional” with “of dimension at most α ”. Since all irreducible representations are at most N -dimensional, the direct integral decomposition of A -representations gives us an injective map $A_{\text{PN}} \rightarrow A''$, with boundedness following because any $a \in A_{\text{PN}}$ is bounded on $\coprod_{k \leq N} \text{Rep}_k^*(A)$.

Now, Akemann and Shultz [1] define a ring $A_c \subset A''$ to consist of those b for which the functions b , b^*b and bb^* are weakly $*$ -continuous on the space $P(A)$ of pure states of A . Since all irreducible representations arise as subrepresentations of N -dimensional representations, continuity on $\text{Rep}_N^*(A)$ suffices to give continuity on $P(A)$, so the inclusion $A_{\text{PN}} \rightarrow A_c$ is an isomorphism.

By [4], the spectrum of A is Hausdorff, then [1] observes, since A is type I , that A is perfect, which means that the inclusion $A \rightarrow A_c$ is in fact an isomorphism. Thus the map $A \rightarrow A_{\text{PN}}$ is an isomorphism. \square

1.4 The category of C^* -algebras with completely bounded morphisms

1.4.1 Basic properties

Lemma 1.20 *If A is a C^* -algebra and $f: A \rightarrow B$ a morphism of Banach algebras, then the image of f has the natural structure of a C^* -algebra, with $f: A \rightarrow \text{Im}(f)$ becoming a C^* -homomorphism.*

Proof The kernel of f is a closed two-sided ideal. Thus, by [22, Theorem 3], $A/\ker f$ is a C^* -algebra, as required. \square

Definition 1.21 Recall that a homomorphism $\pi: A \rightarrow B$ of Banach algebras is said to be *completely bounded* if

$$\sup_{n \in \mathbb{N}} \|M_n(\pi)\| < \infty,$$

where $M_n(\pi): M_n(A) \rightarrow M_n(B)$ is the morphism on $n \times n$ matrices given by π .

Given a pro-Banach algebra $A = \varprojlim_i A_i$ with completely bounded structure maps $A_j \rightarrow A_i$, and a Banach algebra B , any morphism $\pi: A \rightarrow B$ factors through some A_i , and we say that π is *completely bounded* if the map $A_j \rightarrow B$ is so for some sufficiently large $j \geq i$.

Write $\text{Hom}(A, B)_{\text{cb}}$ for the set of completely bounded homomorphisms from A to B .

Lemma 1.22 *If A is a C^* -algebra, then any completely bounded homomorphism $f: A \rightarrow L(H)$ is conjugate to a $*$ -homomorphism of C^* -algebras.*

Proof This is the main result of [16]. □

Remark 1.23 Kadison’s similarity problem asks whether all bounded (noninvolutive) homomorphisms between C^* -algebras are in fact completely bounded. The answer is affirmative in a wide range of cases, but the general problem remains open. Note that Gardner showed [9, Theorem A] that all Banach isomorphisms of C^* -algebras are conjugate to C^* -homomorphisms (and hence completely bounded).

Also note that by [22, Theorem 3], every closed two-sided ideal of a C^* -algebra is a $*$ -ideal; combined with Gardner’s result, this implies that any bounded surjective map $A \rightarrow B$ between C^* -algebras must be conjugate to a C^* -homomorphism, hence completely bounded. The same is true of any bounded map between C^* -algebras whose image is a C^* -subalgebra.

Definition 1.24 Let $C^*BA\text{Alg}_k$ be the category of unital C^* -algebras over k , with completely bounded morphisms (which need not preserve $*$).

Lemma 1.25 *For complex C^* -algebras A, B , giving a $U(B)$ -equivariant function $f: \text{Hom}_{C^*\text{Alg}_{\mathbb{C}}}(A, B) \rightarrow B$ (for the adjoint action on B) is equivalent to giving a B^\times -equivariant function $\tilde{f}: \text{Hom}_{C^*BA\text{Alg}_{\mathbb{C}}}(A, B) \rightarrow B$.*

Proof There is a canonical inclusion $\iota: \text{Hom}_{C^*\text{Alg}_{\mathbb{C}}}(A, B) \rightarrow \text{Hom}_{C^*BA\text{Alg}_{\mathbb{C}}}(A, B)$, so given \tilde{f} , we just set f to be $\tilde{f} \circ \iota$.

The polar decomposition allows us to write $B^\times = B_{++}U(B)$, where $B_{++} \subset S(B)$ is the subset of strictly positive self-adjoint elements. Given f , there is thus an associated B^\times -equivariant function $\tilde{f}: \text{Hom}_{C^*\text{Alg}_{\mathbb{C}}}(A, B) \times B_{++} \rightarrow B$ given by $\tilde{f}(p, g) = g^{-1}f(p)g$. By Lemma 1.22, the map $\text{Hom}_{C^*\text{Alg}_{\mathbb{C}}}(A, B) \times B_{++} \rightarrow \text{Hom}_{C^*BA\text{Alg}_{\mathbb{C}}}(A, B)$ is surjective, and we need to check that \tilde{f} descends.

Now, if $\xi \in S(B)$ has the property that $\exp(\xi)$ fixes $p(A)$ under conjugation, then $\exp(i\xi t)$ commutes with $f(p)$ for all t , so $i\xi$ must also. Thus ξ and hence $\exp(\xi)$ commute with $f(p)$, so the map \tilde{f} does indeed descend. □

1.4.2 Representations We now fix a real unital C^* -algebra A .

Definition 1.26 Define FDVect to be the category of complex finite-dimensional vector spaces and linear maps.

Definition 1.27 Write $\text{FDRep}(A)$ for the category of pairs (V, ρ) for $V \in \text{FDVect}$ and $\rho: A \rightarrow \text{End}(V)$ a unital continuous morphism of Banach algebras. Morphisms are given by linear maps intertwining representations. The set of objects of $\text{FDRep}(A)$ is given the topology of pointwise convergence.

Note that the objects of $\text{FDRep}(A)$ decompose into direct sums of irreducibles.

Lemma 1.28 *The ring A_{PN} of Lemma 1.17 is isomorphic to $A_{\text{PN}'}$, the ring of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant continuous endomorphisms of the fibre functor $\eta: \text{FDRep}(A) \rightarrow \text{FDVect}$. Explicitly, $A_{\text{PN}'}$ consists of elements a such that:*

- (1) *For any linear map $f: (V, \rho_1) \rightarrow (W, \rho_2)$, we have $a(W, \rho_2)f = fa(V, \rho_1)$.*
- (2) *The maps $a: \text{Rep}_n^*(A) \rightarrow \text{Mat}_n(\mathbb{C})$ are continuous and $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant.*

Proof First note that condition (1) applied to the two projections $V_1 \oplus V_2 \rightarrow V_i$ ensures that, for any $(V_1, \rho_1), (V_2, \rho_2) \in \text{FD}^*\text{Rep}(A)$, we have $a(V_1 \oplus V_2, \rho_1 \oplus \rho_2) = a(V_1, \rho_1) \oplus a(V_2, \rho_2)$.

Restriction to $*$ -representations then gives us a map $\psi: A_{\text{PN}'} \rightarrow A_{\text{PN}}$. For a commutative C^* -algebra C , the C^* -algebra $\text{Mat}_k(C)$ is of type I . This means that any bounded map $A \rightarrow \text{Mat}_k(C)$ is completely bounded, so taking $B = \text{Mat}_k(C)$ in Lemma 1.25 for all k ensures that ψ is an isomorphism. \square

Definition 1.29 Given a commutative unital real C^* -algebra A and a $*$ -homomorphism $A \rightarrow B$ of real C^* -algebras, write $\text{FDRep}_{\hat{A}}(B)$ for the category of triples (f, V, ρ) for $f \in \hat{A}$ (the spectrum of A), $V \in \text{FDVect}$ and $\rho: B \rightarrow \text{End}(V)$ a unital continuous morphism of Banach algebras for which $\rho(a) = f(a) \text{id}$ for all $a \in A$. Morphisms are given by linear maps intertwining representations. The set of objects of $\text{FDRep}_{\hat{A}}(B)$ is given the topology of pointwise convergence.

The category $\text{FDRep}_{\hat{A}}(B)$ has an additive structure over \hat{A} , given by

$$(f, V_1, \rho_1) \oplus (f, V_2, \rho_2) = (f, V_1 \oplus V_2, \rho_1 \oplus \rho_2).$$

Lemma 1.30 *Given a commutative unital C^* -algebra A and a $*$ -homomorphism $A \rightarrow B$ of real C^* -algebras, the ring B_{PN} of Lemma 1.17 is isomorphic to the ring of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant continuous endomorphisms of the fibre functor*

$$\eta: \text{FDRep}_{\hat{A}}(B) \rightarrow \text{FDVect}.$$

Proof This just combines the proofs of Lemma 1.28 and Proposition 1.19. The only modification is to observe that $A = C(\hat{A}, \mathbb{C})^{\text{Gal}(\mathbb{C}/\mathbb{R})}$ and that, for any irreducible representation $\rho: B \rightarrow \text{End}(V)$, we necessarily have $\rho|_A = f \text{id}$ for some $f \in \hat{A}$. \square

2 The Betti, de Rham and harmonic functors on Banach algebras

2.1 The Riemann–Hilbert correspondence

Definition 2.1 Given a path-connected topological space X with basepoint x and a unital \mathbb{R} –algebra B , define the Betti representation space $\mathbf{R}_{X,x}^B(B)$ by

$$\mathbf{R}_{X,x}^B(B) := \text{Hom}_{\text{Gp}}(\pi_1(X, x), B^\times),$$

where B^\times is the multiplicative group of units in B .

Define the representation groupoid $\mathcal{R}_X^B(B)$ by $\mathcal{R}_X^B(B) := [\mathbf{R}_{X,x}^B(B)/B^\times]$, where B^\times acts by conjugation. Note that this is independent of the choice of basepoint (being equivalent to the groupoid of B^\times –torsors on X).

Definition 2.2 Given a connected manifold X with basepoint x and a Banach algebra B , define the de Rham groupoid $\mathcal{R}_X^{\text{dR}}(B)$ to be the groupoid of smooth B^\times –bundles with flat connections. Thus $\mathcal{R}_X^{\text{dR}}(B)$ consists of pairs (\mathcal{T}, D) , where \mathcal{T} is a right $\mathcal{A}_X^0(B^\times)$ –torsor, and D is a flat connection on \mathcal{T} .

Explicitly, write $\mathcal{A}_X^n(\text{ad } \mathcal{T}) := \mathcal{T} \times_{\mathcal{A}_X^0(B^\times)} \mathcal{A}_X^n(B)$, for the adjoint action of B^\times on B . Then a flat connection on \mathcal{T} is

$$D: \mathcal{T} \rightarrow \mathcal{A}^1(\text{ad } \mathcal{T})$$

satisfying

- (1) D is a d –connection: $D(pg) = \text{ad}_g D(p) + g^{-1}dg$ for $g \in \mathcal{A}_X^0(B^\times)$;
- (2) D is flat: $(\text{ad } D) \circ D = 0$.

Define $\mathbf{R}_{X,x}^{\text{dR}}(B)$ to be the groupoid of triples (\mathcal{T}, D, f) , where $(\mathcal{T}, D) \in \mathcal{R}_X^{\text{dR}}(B)$ and $f \in x^*\mathcal{T}$ is a distinguished element. Since $\mathbf{R}_{X,x}^{\text{dR}}(B)$ has no nontrivial automorphisms, we will regard it as a set-valued functor (given by its set of isomorphism classes).

Note that B^\times acts on $\mathbf{R}_{X,x}^{\text{dR}}(B)$ by changing the framing, whence the quotient groupoid is equivalent to $\mathcal{R}_X^{\text{dR}}(B)$.

Proposition 2.3 For any pointed connected manifold (X, x) and any Banach algebra B there are canonical equivalences

$$\mathcal{R}_X^{\text{dR}}(B) \simeq \mathcal{R}_X^B(B), \quad \mathbf{R}_{X,x}^{\text{dR}}(B) \cong \mathbf{R}_{X,x}^B(B)$$

functorial in X, x and B .

Proof When B is a finite-dimensional matrix algebra, this is [10, Theorem 5.10]. The same proof carries over to Banach algebras, noting that the argument for existence of parallel transport [15, Section II.3] holds in this generality, since $\exp(b) = \sum_{n \geq 0} b^n/n!$ converges and is invertible for all $b \in B$. \square

Remark 2.4 The functor $R_{X,x}^{\text{dR}}$ naturally extends to a functor on the category of pro-Banach algebras $\text{pro}(\text{BanAlg})$, by sending any cofiltered inverse system $\{A_i\}_i$ to $\varprojlim_i R_{X,x}^{\text{dR}}(A_i)$. Since the functor $R_{X,x}^B$ commutes with all limits, the equivalences of Proposition 2.3 then extend to pro-Banach algebras. The category $\text{pro}(\text{BanAlg})$ contains all multiplicatively convex Fréchet algebras (since they are countable inverse limits of Banach algebras) and indeed all complete LMC algebras via the Arens–Michael decomposition.

For any open subset $U \subset \mathbb{C}^n$, the ring $O(U) := \Gamma(U, \mathcal{O}_U)$ of holomorphic functions on U can be realised as a pro-Banach algebra by looking at the system of sup norms on compact subspaces. Taking quotients by finitely generated ideals I then gives local models $Y := (V(I), \mathcal{O}_U/I)$ for complex analytic spaces, and realises $O(Y)$ as the pro-Banach algebra $O(U)/I$. A complex analytic morphism from Y to the variety $\text{Hom}(\pi_1(X, x), \text{GL}_n(\mathbb{C}))$ is then just an element of $\text{Hom}(\pi_1(X, x), \text{GL}_n(O(Y))) = R_{X,x}^B(\text{Mat}_n(O(Y)))$ for the pro-Banach algebra $\text{Mat}_n(O(Y))$ of morphisms $Y \rightarrow \text{Mat}_n(\mathbb{C})$.

We can therefore recover the analytic structure of the variety $\text{Hom}(\pi_1(X, x), \text{GL}_n(\mathbb{C}))$ from the set-valued functor $R_{X,x}^B$ on Banach algebras and hence (by Proposition 2.3) from the set-valued functor $R_{X,x}^{\text{dR}}$.

In [25], the varieties $R_{X,x}^B(\text{Mat}_n(-))$ and $R_{X,x}^{\text{dR}}(\text{Mat}_n(-))$ are denoted by $R_B(X, x, n)$, and $R_{\text{DR}}(X, x, n)$.

Lemma 2.5 *For any real Banach algebras B, C , there is a canonical map*

$$m: R_{X,x}^B(B) \times R_{X,x}^B(C) \rightarrow R_{X,x}^B(B \otimes_{\mathbb{R}}^{\pi} C),$$

where \otimes^{π} is the projective tensor product. This makes $R_{X,x}^B$ into a symmetric monoidal functor, with unit corresponding to the trivial representation in each $R_{X,x}^B(B)$.

Proof Given representations $\rho_1: \pi_1(X, x) \rightarrow B^{\times}$ and $\rho_2: \pi_1(X, x) \rightarrow C^{\times}$, we obtain $\rho_1 \otimes \rho_2: \pi_1(X, x) \rightarrow (B \otimes C)^{\times}$. Taking the completion with respect to the projective cross norm gives the required result. \square

Remark 2.6 Given any complex affine group scheme G , we may use the tensor structure on $R_{X,x}^B$ to recover the affine analytic variety $\text{Hom}(\pi_1(X, x), G(\mathbb{C}))$ in the same spirit as Remark 2.4. Explicitly, $O(G)$ is a coalgebra, so can be written as a nested union of finite-dimensional coalgebras. Therefore $O(G)^{\vee}$ is a pro-finite-dimensional algebra, and hence a pro-Banach algebra.

Multiplication on G gives us a comultiplication $\mu: O(G)^\vee \rightarrow O(G \times G)^\vee$. For any complex analytic space Y , we may then characterise $\text{Hom}(\pi_1(X, x), G(Y))$ as

$$\{\rho \in \mathbf{R}_{X,x}^B(O(G)^\vee \widehat{\otimes} O(Y)) \mid \mu(\rho) = m(\rho, \rho) \in \mathbf{R}_{X,x}^B(O(G \times G)^\vee \widehat{\otimes} O(Y))\}.$$

2.2 Representability of the de Rham functor

Lemma 2.7 *Given a free group $\Gamma = F(X)$, the functor*

$$A \mapsto \text{Hom}_{\text{Gp}}(\Gamma, A^\times)$$

on the category of real Banach algebras is prorepresentable.

Proof Given a function $v: X \rightarrow [1, \infty)$, let $\bar{v}: \Gamma \rightarrow [1, \infty)$ be the largest function subject to the conditions

- (1) $\bar{v}(1) = 1$;
- (2) $\bar{v}(x) = \bar{v}(x^{-1}) = v(x)$ for all $x \in X$;
- (3) $\bar{v}(gh) \leq \bar{v}(g)\bar{v}(h)$.

Explicitly, we write any $g \in \Gamma$ as a reduced word $g = x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$, then set $\bar{v}(g) := \prod_{i=1}^k v(x_i)^{|n_i|}$. We now define a norm $\|-\|_{1,v}$ on $k[\Gamma]$ by setting

$$\left\| \sum_{\gamma \in \Gamma} \lambda_\gamma \gamma \right\|_{1,v} := \sum_{\gamma \in \Gamma} |\lambda_\gamma| \cdot \bar{v}(\gamma).$$

Now, given any representation $\rho: \Gamma \rightarrow A^\times$, we may define $v: X \rightarrow [1, \infty)$ by

$$v(x) := \max\{\|\rho(x)\|, \|\rho(x^{-1})\|\};$$

this at least 1 because $1 = \rho(x)\rho(x^{-1})$, so $1 \leq \|\rho(x)\| \cdot \|\rho(x^{-1})\|$. It follows that for all $v \in k[\Gamma]$ we have $\|\rho(v)\| \leq \|v\|_{1,v}$, so ρ determines a map

$$k[\Gamma]^{\wedge v} \rightarrow A,$$

where $k[\Gamma]^{\wedge v}$ denotes the Banach algebra obtained by completing $k[\Gamma]$ with respect to the norm $\|-\|_{1,v}$.

Next, give $[1, \infty)^X$ the structure of a poset by saying $v_1 \leq v_2$ provided $v_1(x) \leq v_2(x)$ for all $x \in X$. This is in fact a directed set, since we can define $\max\{v_1, v_2\}$ pointwise. There is a canonical morphism

$$k[\Gamma]^{\wedge v_2} \rightarrow k[\Gamma]^{\wedge v_1}$$

whenever $v_1 \leq v_2$, which gives us an inverse system $k[\Gamma]^{\text{an}} := \{k[\Gamma]^{\wedge v}\}_v$ of Banach algebras, indexed by the directed set $([1, \infty)^X)$.

Thus we have shown that

$$\text{Hom}_{\text{Gp}}(\Gamma, A^\times) \cong \varinjlim_{\nu \in [1, \infty)^X} \text{Hom}_{k\text{BanAlg}}(k[\Gamma]^{\wedge \nu}, A),$$

functorial in Banach k -algebras A . In other words, our prorepresenting object is the inverse system $k[\Gamma]^{\text{an}}$. □

Example 2.8 Take $X = \{z\}$, so $\Gamma = \mathbb{Z}$, and let $\nu(z) = R$. Then elements of $k[\mathbb{Z}]^{\wedge \nu}$ are $\sum_{i \in \mathbb{Z}} \lambda_i z^i$ such that

$$\sum_{i \geq 0} |\lambda_i| R^i < \infty, \quad \sum_{i \leq 0} |\lambda_i| R^{-i} < \infty.$$

Thus $\mathbb{C}[\mathbb{Z}]^{\wedge R}$ is the ring of analytic functions converging on the annulus $R^{-1} \leq |z| \leq R$. Hence $\varprojlim_R \mathbb{C}[\mathbb{Z}]^{\wedge R}$ is the ring of analytic functions on \mathbb{C}^* , while $\varprojlim_R \mathbb{R}[\mathbb{Z}]^{\wedge R}$ is the subring consisting of functions f with $\overline{f(z)} = f(\bar{z})$

Contrast this with the isometric Banach completion of $\mathbb{C}[\mathbb{Z}]$, which just gives us $\mathbb{C}[\mathbb{Z}]^{\wedge 1} = \ell^1(\mathbb{Z})$.

Lemma 2.9 *Given a finitely generated free group $\Gamma = F(X)$, the functor*

$$A \mapsto \text{Hom}_{\text{Gp}}(\Gamma, A^\times)$$

on the category of multiplicatively convex Fréchet k -algebras is representable.

Proof We may embed \mathbb{N}_1 in $[1, \infty)^X$ as a subset of the constant functions. Since X is finite, \mathbb{N}_1 is a cofinal subset of $[1, \infty)^X$, giving us an isomorphism

$$\{k[\Gamma]^{\wedge \nu}\}_{\nu \in [1, \infty)^X} \cong \{k[\Gamma]^{\wedge n}\}_{n \in \mathbb{N}_1}$$

in the category of pro-Banach k -algebras. Since \mathbb{N}_1 is countable, $k[\Gamma]^{\text{an}} := \varprojlim_n k[\Gamma]^{\wedge n}$ is a Fréchet algebra.

Applying the proof of Lemma 2.7, we have shown that

$$\text{Hom}(\Gamma, A^\times) \cong \text{Hom}_{k\text{FrAlg}}(k[\Gamma]^{\text{an}}, A)$$

for all Banach algebras A . Since any m -convex Fréchet algebra A can be expressed as an inverse limit $A = \varprojlim_i A_i$ of Banach algebras, it follows that the same isomorphism holds for all such algebras, so the functor is representable in m -convex Fréchet algebras. □

Proposition 2.10 *Given a finitely generated group Γ , the functor*

$$A \mapsto \text{Hom}_{\text{Gp}}(\Gamma, A^\times)$$

on the category of multiplicatively convex Fréchet k -algebras is representable.

Proof Choose generators X for Γ , so $\Gamma = F(X)/K$ for some normal subgroup K . Lemma 2.9 gives a Fréchet k -algebra $k[F(X)]^{\text{an}}$ governing representations of $F(X)$. Since

$$\text{Hom}_{\text{Gp}}(\Gamma, A^\times) = \{f \in \text{Hom}_{\text{Gp}}(F(X), A^\times) \mid f(K) = \{1\}\},$$

our functor will be represented by a quotient of $k[F(X)]^{\text{an}}$. Specifically, let I be the closed ideal of $k[F(X)]^{\text{an}}$ generated by $\{k-1 \mid k \in K\}$ and set $k[\Gamma]^{\text{an}} := k[F(X)]^{\text{an}}/I$. This is an m -convex Fréchet algebra and

$$\text{Hom}_{\text{Gp}}(\Gamma, A^\times) \cong \text{Hom}_{k\text{FrAlg}}(k[\Gamma]^{\text{an}}, A)$$

for all such algebras A .

For an explicit description of $k[\Gamma]^{\text{an}}$, note that the system of norms is given by

$$\left\| \sum \lambda_\gamma \gamma \right\|_{1,n} = \sum |\lambda_\gamma| \cdot n^{w(\gamma)},$$

where $w(\gamma)$ is the minimal word length of γ in terms of X . □

When combined with its tensor structure, this implies that the functor of Proposition 2.10 is a very strong invariant indeed:

Lemma 2.11 *The tensor structure of Lemma 2.5 gives $k[\Gamma]^{\text{an}}$ the structure of an m -convex Fréchet bialgebra. The group*

$$G(k[\Gamma]^{\text{an}}) = \{a \in k[\Gamma]^{\text{an}} \mid \mu(a) = a \otimes a \in k[\Gamma]^{\text{an}} \otimes^\pi k[\Gamma]^{\text{an}}, \varepsilon(a) = 1 \in k\}$$

of grouplike elements of $k[\Gamma]^{\text{an}}$ is then Γ .

Proof Applying the map m of Lemma 2.5 to (ξ, ξ) for the canonical element $\xi \in \text{Hom}_{\text{Gp}}(\Gamma, k[\Gamma]^{\text{an}})$ gives us a comultiplication $\mu: k[\Gamma]^{\text{an}} \rightarrow k[\Gamma]^{\text{an}} \otimes^\pi k[\Gamma]^{\text{an}} = k[\Gamma \times \Gamma]^{\text{an}}$ and a counit $\varepsilon: k[\Gamma]^{\text{an}} \rightarrow k$. On the topological basis Γ , we must have $\mu(\gamma) = (\gamma, \gamma)$ and $\varepsilon(\gamma) = 1$.

Expressing $a \in G(k[\Gamma]^{\text{an}})$ as $\sum_{\gamma \in \Gamma} a_\gamma \gamma$, note that the conditions become $a_\gamma a_\delta = 0$ for $\gamma \neq \delta$, and $\sum a_\gamma = 1$; thus $a = \gamma$ for some $\gamma \in \Gamma$. □

Example 2.12 Arguing as in Example 2.8, for Γ abelian and finitely generated, $\mathbb{C}[\Gamma]^{\text{an}}$ is isomorphic to the ring of complex analytic functions on $\text{Hom}_{\text{Gp}}(\Gamma, \mathbb{C}^*)$, while $\mathbb{R}[\Gamma]^{\text{an}} \subset \mathbb{C}[\Gamma]^{\text{an}}$ consists of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant functions. The multiplicative analytic functions are of course just Γ itself.

Proposition 2.3 then implies:

Corollary 2.13 *The functors $R_{X,x}^{\text{dR}}$ and $R_{X,x}^B$ on real multiplicatively convex Fréchet algebras are representable.*

Remark 2.14 Adapting the ideas of [19], the functor $R_{X,x}^B$ has a natural extension to those simplicial Banach algebras B_\bullet for which $B_n \rightarrow \pi_0 B$ is a nilpotent extension for each n . Explicitly, we could set $R_{X,x}^B(B)$ to be the set of homotopy classes of maps $G(\text{Sing}(X, x)) \rightarrow B_\bullet^\times$ of simplicial groups, where G is Kan’s loop group. This functor admits a tensor structure extending Lemma 2.5,

The functor $R_{X,x}^{\text{dR}}$ has a natural extension to those differential graded Banach algebras B_\bullet for which $B_0 \rightarrow H_0 B$ is a nilpotent extension. Explicitly, $R_{X,x}^{\text{dR}}(B)$ would consist of pairs (\mathcal{T}_0, D) , where \mathcal{T}_0 is a $\mathcal{A}_X^0(B_0^\times)$ -torsor and $D: \mathcal{T}_0 \rightarrow \prod_n \mathcal{A}_X^{n+1} \otimes_{\mathcal{A}_X^0} \text{ad } \mathcal{T}_n(n+1)$ is a flat hyperconnection, where $\text{ad } \mathcal{T}_n := \mathcal{T} \times_{\mathcal{A}_X^0(B_0^\times)} \mathcal{A}_X^0(B_n)$.

It then seems likely that [19, Corollary 4.41] should adapt to give natural isomorphisms $R_{X,x}^B(B) \cong R_{X,x}^{\text{dR}}(NB)$, where N is Dold–Kan normalisation.

2.3 The pluriharmonic functor

Fix a compact connected Kähler manifold X , with basepoint $x \in X$.

Definition 2.15 Given a real Banach space B , denote the sheaf of B -valued C^∞ n -forms on X by $\mathcal{A}_X^n(B)$, and let \mathcal{A}_X^\bullet be the resulting complex. Write $A^\bullet(X, B) := \Gamma(X, \mathcal{A}_X^\bullet(B))$. We also write $\mathcal{A}_X^\bullet := \mathcal{A}_X^\bullet(\mathbb{R})$ and $A^\bullet(X) := A^\bullet(X, \mathbb{R})$.

Definition 2.16 Define S to be the real algebraic group $\prod_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$ obtained as in [6, Section 2.1.2] from $\mathbb{G}_{m,\mathbb{C}}$ by restriction of scalars. Note that there is a canonical inclusion $\mathbb{G}_m \hookrightarrow S$.

The following is a slight generalisation of [20, Definition 2.49]:

Definition 2.17 For any real Banach space B , there is an action of S on $\mathcal{A}_X^*(B)$, which we will denote by $a \mapsto \lambda \diamond a$, for $\lambda \in \mathbb{C}^* = S(\mathbb{R})$. For $a \in (A^*(X) \otimes \mathbb{C})^{pq}$, it is given by

$$\lambda \diamond a := \lambda^p \bar{\lambda}^q a.$$

Definition 2.18 Given a real C^* -algebra B , define $\mathcal{R}_X^J(B)$ to be the groupoid of pairs $(U(\mathcal{P}), D)$, where $U(\mathcal{P})$ is a right $\mathcal{A}_X^0(U(B))$ -torsor, and D is a pluriharmonic connection on $U(\mathcal{P})$.

Explicitly, write $\mathcal{P} := U(\mathcal{P}) \times_{\mathcal{A}_X^0(U(B))} \mathcal{A}_X^0(B^\times)$, and

$$\begin{aligned} \text{ad } \mathcal{P} &:= \mathcal{P} \times_{\mathcal{A}_X^0(B^\times)} \mathcal{A}_X^0(B) \\ &= U(\mathcal{P}) \times_{\mathcal{A}_X^0(U(B))} \mathcal{A}_X^0(B) \\ &= [U(\mathcal{P}) \times_{\mathcal{A}_X^0(U(B))} \mathcal{A}_X^0(u(B))] \oplus [U(\mathcal{P}) \times_{\mathcal{A}_X^0(U(B))} \mathcal{A}_X^0(S(B))], \end{aligned}$$

where $U(B)$ and B^\times act on B by the adjoint action. Then a pluriharmonic connection on \mathcal{P} is

$$D: U(\mathcal{P}) \rightarrow \text{ad } \mathcal{P}$$

satisfying

- (1) D is a d -connection: $D(pu) = \text{ad}_u D(p) + u^{-1} du$ for $u \in \mathcal{A}_X^0(U(B))$;
- (2) D is flat: $(\text{ad } D) \circ D = 0$;
- (3) D is pluriharmonic: $(\text{ad } D) \circ D^c + (\text{ad } D^c) \circ D = 0$.

Here, $D = d^+ + \vartheta$ comes from the decomposition of $\text{ad } \mathcal{P}$ into anti-self-adjoint and self-adjoint parts, and $D^c = i \diamond d^+ - i \diamond \vartheta$.

Define $\mathbf{R}_{X,x}^J(B)$ to be the groupoid of triples $(U(\mathcal{P}), D, f)$, where $(U(\mathcal{P}), D)$ is in $\mathcal{R}_X^J(B)$ and $f \in x^*U(\mathcal{P})$ is a distinguished element. Since $\mathbf{R}_{X,x}^J(B)$ has no nontrivial automorphisms, we will regard it as a set-valued functor (given by its set of isomorphism classes).

Remark 2.19 There is a natural action of $U(B)$ on $\mathbf{R}_{X,x}^J(B)$, given by changing the framing. The quotient groupoid $[\mathbf{R}_{X,x}^J(B)/U(B)]$ is thus equivalent to $\mathcal{R}_X^J(B)$. In [25, Lemma 7.13], the set $\mathbf{R}_{X,x}^J(\text{Mat}_n(\mathbb{C}))$ is denoted by $\mathbf{R}_{\text{DR}}^J(X, x, n)$.

Also note that the definition of $\mathbf{R}_{X,x}^J(B)$ can be extended to any real Banach $*$ -algebra B . However, this will not be true of the harmonic functor of Section 2.5.

Example 2.20 When V is a real Hilbert space, the algebra $L(V)$ of bounded operators on V is a real C^* -algebra. Then $\mathcal{R}_X^J(B)$ is equivalent to the groupoid of pluriharmonic local systems \mathbb{V} in Hilbert spaces on X , fibrewise isometric to V . The connection $D: \mathcal{A}^0(\mathbb{V}) \rightarrow \mathcal{A}^1(\mathbb{V})$ must satisfy the pluriharmonic condition that $DD^c + D^cD = 0$ for D^c defined with respect to the smooth inner product $\mathbb{V} \times \mathbb{V} \rightarrow \mathcal{A}_X^0$. Isomorphisms in $\mathcal{R}_X^J(B)$ preserve the inner product.

Definition 2.21 Define the de Rham projection

$$\pi_{\text{dR}}: \mathbf{R}_{X,x}^J(B) \rightarrow \mathbf{R}_{X,x}^{\text{dR}}(B)$$

by mapping $(U(\mathcal{P}), D, f)$ to the framed flat torsor

$$(\mathcal{P}, D, f) = (U(\mathcal{P}) \times_{\mathcal{A}_X^0(U(B))} \mathcal{A}_X^0(B^\times), D, f \times_{U(B)} B^\times).$$

Proposition 2.22 The functor $\mathbf{R}_{X,x}^J: C^*\text{Alg} \rightarrow \text{Set}$ is strictly prorepresentable, by an object $E_{X,x}^J \in \text{pro}(C^*\text{Alg})$.

Proof The final object in $C^*\text{Alg}$ is 0, and $\mathbf{R}_{X,x}^J(0)$ is the one-point set, so $\mathbf{R}_{X,x}^J$ preserves the final object.

Given maps $A \rightarrow B \leftarrow C$ in $C^*\text{Alg}$ and $(p_A, p_B) \in \mathbf{R}_{X,x}^J(A) \times_{\mathbf{R}_{X,x}^J(B)} \mathbf{R}_{X,x}^J(C)$, we get

$$\begin{aligned} \pi_{\text{dR}}(p_A, p_C) &\in \mathbf{R}_{X,x}^{\text{dR}}(A) \times_{\mathbf{R}_{X,x}^{\text{dR}}(B)} \mathbf{R}_{X,x}^{\text{dR}}(C) \cong \mathbf{R}_{X,x}^B(A) \times_{\mathbf{R}_{X,x}^B(B)} \mathbf{R}_{X,x}^B(C) \\ &\cong \mathbf{R}_{X,x}^B(A \times_B C). \end{aligned}$$

Thus we have a flat torsor $(\mathcal{P}, D) \in \mathbf{R}_{X,x}^{\text{dR}}(A \times_B C)$.

It follows that $p_A \cong (U(\mathcal{P}_A), D)$ for some orthogonal form

$$U(\mathcal{P}_A) \subset \mathcal{P}_A = \mathcal{P} \times_{\mathcal{A}_X^0((A \times_B C)^\times)} \mathcal{A}_X^0(A^\times),$$

and similarly for p_C . Since the images of p_A and p_C are equal in $\mathbf{R}_{X,x}^{\text{dR}}(B)$, there is a framed orthogonal isomorphism

$$\alpha: U(\mathcal{P}_A) \times_{\mathcal{A}_X^0(U(A))} \mathcal{A}_X^0(U(B)) \rightarrow U(\mathcal{P}_C) \times_{\mathcal{A}_X^0(U(C))} \mathcal{A}_X^0(U(B)),$$

inducing the identity on \mathcal{P}_B . Hence α must itself be the identity, so both $U(\mathcal{P}_A)$ and $U(\mathcal{P}_C)$ give the same unitary form $U(\mathcal{P}_B)$ for \mathcal{P}_B . It is easy to check the pluriharmonic conditions, giving an element

$$(U(\mathcal{P}_A) \times_{U(\mathcal{P}_B)} U(\mathcal{P}_C), D) \in \mathbf{R}_{X,x}^J(A \times_B C)$$

over (p_A, p_C) . This is essentially unique, so $\mathbf{R}_{X,x}^J$ preserves fibre products, and hence finite limits.

Now, given a C^* -subalgebra $A \subset B$, the map $\mathbf{R}_{X,x}^J(A) \rightarrow \mathbf{R}_{X,x}^J(B)$ is injective. This follows because if two framed pluriharmonic bundles $\mathcal{P}_1, \mathcal{P}_2 \in \mathbf{R}_{X,x}^J(A)$ become isomorphic in $\mathbf{R}_{X,x}^J(B)$, compatibility of framings ensures that the isomorphism f maps $x^*\mathcal{P}_1$ to $x^*\mathcal{P}_2$. Since f is compatible with the connections, it thus gives an isomorphism $f: \mathcal{P}_1 \rightarrow \mathcal{P}_2$, by considering the associated local systems.

Finally, given an inverse system $\{A_i\}_i$ of nested C^* -subalgebras of a C^* -algebra B and an element of $\bigcap_i R_{X,x}^J(A_i)$, we have a compatible system $\{(\mathcal{P}_i, D_i, f_i)\}_i$. Set $\mathcal{P} := \varprojlim_i \mathcal{P}_i$, with connection D and framing f induced by the D_i and f_i . This defines a unique element of $R_{X,x}^J(\bigcap_i A_i)$, showing that

$$F\left(\bigcap_i A_i\right) \cong \bigcap_i F(A_i).$$

Thus all the conditions of Proposition 1.9 are satisfied, so $R_{X,x}^J$ is strictly prorepresentable. \square

Definition 2.23 Given pro- C^* -algebras B and C over k , define $B \widehat{\otimes}_k C$ to be the maximal k -tensor product of B and C , as defined in [18, Definition 3.1]; this is again a pro- C^* -algebra.

Lemma 2.24 For any real pro- C^* -algebras B and C , there is a canonical map

$$m: R_{X,x}^J(B) \times R_{X,x}^J(C) \rightarrow R_{X,x}^J(B \widehat{\otimes}_{\mathbb{R}} C),$$

making $R_{X,x}^J$ into a symmetric monoidal functor, with unit corresponding to the trivial torsor in each $R_{X,x}^J(B)$.

Proof Given $(U(\mathcal{P}), D, f, U(\mathcal{Q}), E, \beta)$ on the left-hand side, we first form the $\mathcal{A}_X^0(U(B \widehat{\otimes} C))$ -torsor $U(\mathcal{R})$ given by

$$U(\mathcal{R}) := (U(\mathcal{P}) \times U(\mathcal{Q}))_{\mathcal{A}_X^0(U(B) \times U(C))} \mathcal{A}_X^0(U(B \widehat{\otimes} C)).$$

We then define a connection F on $U(\mathcal{R})$ determined by

$$\begin{aligned} F(p, q, 1) &= (Dp, q) + (p, Dq) \in \mathcal{A}_X^1(\text{ad } \mathcal{R}) \\ &= (U(\mathcal{P}) \times U(\mathcal{Q}))_{\mathcal{A}_X^0(U(B) \times U(C))} \mathcal{A}_X^1(B \widehat{\otimes} C) \end{aligned}$$

for $p \in U(\mathcal{P})$ and $q \in U(\mathcal{Q})$. This is clearly flat and pluriharmonic, and the construction is also symmetric monoidal. \square

Note that this gives $E_{X,x}^J$ the structure of a pro- C^* -bialgebra, with comultiplication $\mu: E_{X,x}^J \rightarrow E_{X,x}^J \widehat{\otimes} E_{X,x}^J$ coming from m , and counit $\varepsilon: E_{X,x}^J \rightarrow k$ coming from the trivial torsor.

The following is immediate:

Lemma 2.25 For any morphism $f: X \rightarrow Y$ of compact connected Kähler manifolds, there is a natural transformation

$$f^*: R_{Y, f_x}^J \rightarrow R_{X, x}^J$$

of functors.

2.4 Higgs bundles

Definition 2.26 Given a complex Banach algebra B , write $\mathcal{O}_X(B)$ for the sheaf on X given locally by holomorphic functions $X \rightarrow B$.

Definition 2.27 For a complex Banach algebra B , a Higgs B -torsor on X consists of an $\mathcal{O}_X(B)^\times$ -torsor \mathcal{T} , together with a Higgs form $\theta \in \text{ad } \mathcal{T} \otimes_{\mathcal{O}_X} \Omega_X^1$, where $\text{ad } \mathcal{T} := \mathcal{T} \times_{\mathcal{O}_X(B)^\times, \text{ad}} \mathcal{O}_X(B)$ satisfying

$$\theta \wedge \theta = 0 \in \mathcal{T} \otimes_{\mathcal{O}_X} \Omega_X^2.$$

Definition 2.28 Let $\mathcal{R}_X^{\text{Dol}}(B)$ be the groupoid of Higgs B -torsors, and $\mathbf{R}_{X, x}^{\text{Dol}}(B)$ the groupoid of framed Higgs bundles (\mathcal{T}, θ, f) , where $f: B^\times \rightarrow x^* \mathcal{T}$ is a B^\times -equivariant isomorphism. Alternatively, we may think of f as a distinguished element of $x^* \mathcal{T}$.

Note that $\mathbf{R}_{X, x}^{\text{Dol}}(B)$ is a discrete groupoid, so we will usually identify it with its set of isomorphism classes. Also note that there is a canonical action of B^\times on $\mathbf{R}_{X, x}^{\text{Dol}}(B)$ given by the action on the framings. This gives an equivalence

$$\mathcal{R}_X^{\text{Dol}}(B) \simeq [\mathbf{R}_{X, x}^{\text{Dol}}(B) / B^\times]$$

of groupoids.

The following is immediate:

Lemma 2.29 Giving a Higgs B -torsor X is equivalent to giving an $\mathcal{A}_X^0(B^\times)$ torsor \mathcal{Q} equipped with a flat $\bar{\partial}$ -connection, ie a map

$$D'': \mathcal{Q} \rightarrow \text{ad } \mathcal{Q} := \mathcal{Q} \times_{\mathcal{A}_X^0(B^\times), \text{ad}} \mathcal{A}_X^1(B)$$

satisfying

- (1) $D''(pg) = \text{ad}_g D''(p) + g^{-1} \bar{\partial} g$ for $g \in \mathcal{A}_X^0(B^\times)$;
- (2) $(\text{ad } D'') \circ D'' = 0$.

Remark 2.30 Unlike the Betti, de Rham and harmonic functors, the Dolbeault functor cannot be prorepresentable in general. This is for the simple reason that a left-exact scheme must be affine, but the Dolbeault moduli space is seldom so, since it contains the Picard scheme.

Definition 2.31 Given $(U(\mathcal{P}), D) \in \mathcal{R}_X^J(B)$, decompose d^+ and ϑ into $(1, 0)$ and $(0, 1)$ types as $d^+ = \partial + \bar{\partial}$ and $\vartheta = \theta + \bar{\theta}$. Now set $D' = \partial + \bar{\theta}$ and $D'' = \bar{\partial} + \theta$. Note that $D = D' + D''$ and $D^c = iD' - iD''$.

Definition 2.32 For a complex C^* -algebra B , define the Dolbeault projection map $\pi_{\text{Dol}}: \mathcal{R}_X^J(B) \rightarrow \mathcal{R}_X^{\text{Dol}}(B)$ by sending $(U(\mathcal{P}), D)$ to $(\mathcal{P} \times_{\mathcal{A}_X^0(B^\times)} \mathcal{A}_X^0(B^\times), D'')$.

2.5 The harmonic functor

We now let X be any compact Riemannian real manifold.

Definition 2.33 Given a compact Riemannian manifold X , a real C^* -algebra B , a right $\mathcal{A}_X^0(U(B))$ -torsor $U(\mathcal{P})$ and a flat connection

$$D: U(\mathcal{P}) \rightarrow \text{ad } \mathcal{P},$$

say that D is a harmonic connection if $(d^+)^*\vartheta = 0 \in \Gamma(X, \text{ad } \mathcal{P})$ for d^+ and ϑ defined as in Definition 2.18 and the adjoint $*$ given by combining the involution $*$ on $\text{ad } \mathcal{P}$ with the adjoint on \mathcal{A}_X^* given by the Kähler form.

Lemma 2.34 A flat connection D as above on a compact Kähler manifold is harmonic if and only if it is pluriharmonic.

Proof The proof of [23, Lemma 1.1] carries over to this generality. □

Definition 2.35 The lemma allows us to extend Definitions 2.18 and 2.21 to any compact Riemannian manifold X , replacing pluriharmonic with harmonic in the definition of $\mathcal{R}_X^J(B)$ and $\mathbf{R}_{X,x}^J(B)$.

Proposition 2.36 The functor $\mathbf{R}_{X,x}^J: C^*\text{Alg}_{\mathbb{R}} \rightarrow \text{Set}$ is strictly prorepresentable, by an object $E_{X,x}^J \in \text{pro}(C^*\text{Alg}_{\mathbb{R}})$.

Proof The proof of Proposition 2.22 carries over. □

Note that Lemma 2.24 carries over to the functor $\mathbf{R}_{X,x}^J$ for any compact Riemannian manifold X .

The following is immediate:

Lemma 2.37 *For any local isometry $f: X \rightarrow Y$ of compact connected real Riemannian manifolds, there is a natural transformation*

$$f^*: \mathbf{R}_{Y,fx}^J \rightarrow \mathbf{R}_{X,x}^J$$

of functors.

Note that this is much weaker than Lemma 2.25, the pluriharmonic functor being a priori functorial with respect to all morphisms.

3 Analytic nonabelian Hodge theorems

3.1 The de Rham projection

Fix a compact connected real Riemannian manifold X , with basepoint $x \in X$.

The argument of [25, Lemma 7.17] (which is only stated for X Kähler) shows that π_{dR} gives a homeomorphism

$$\mathbf{R}_{X,x}^J(\text{Mat}_n(\mathbb{C}))/U(n) \rightarrow \text{Hom}(\pi_1(X, x), \text{GL}_n(\mathbb{C}))/\text{GL}_n(\mathbb{C}),$$

where $//$ denotes the coarse quotient (in this case, the Hausdorff completion of the topological quotient).

As an immediate consequence, note that

$$\mathbf{R}_{X,x}^J(\mathbb{C}) \rightarrow \text{Hom}_{\text{Gp}}(\pi_1(X, x), \mathbb{C}^*)$$

is a homeomorphism. Thus the abelianisation of $E_{X,x}^J \otimes \mathbb{C}$ is isomorphic to the commutative C^* -algebra $C(\text{Hom}(\pi_1(X, x), \mathbb{C}^*), \mathbb{C})$, with $\mathbf{R}_{X,x}^J \subset \mathbf{R}_{X,x}^J \otimes \mathbb{C}$ consisting of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant functions.

We now adapt these results to recover a finer comparison between the respective functors.

3.1.1 Harmonic representations

Proposition 3.1 *For a compact Riemannian manifold X and all real C^* -algebras B , the de Rham projection*

$$\pi_{\text{dR}}: \mathbf{R}_{X,x}^J(B) \rightarrow \mathbf{R}_{X,x}^{\text{dR}}(B)$$

has the property that if $p_1, p_2 \in \mathbf{R}_{X,x}^J(B)$ and $\text{ad}_b \pi_{\text{dR}}(p_1) = \pi_{\text{dR}}(p_2)$ for some strictly positive self-adjoint element $b \in B$, then $p_1 = p_2$.

Thus

$$\pi_{\text{dR}}: \mathbf{R}_{X,x}^J(B)/U(B) \rightarrow \mathbf{R}_{X,x}^{\text{dR}}(B)/B^\times$$

is injective.

Proof The first statement above implies the second: it suffices to show that, for any $(U(\mathcal{P}), D, f) \in \mathbf{R}_{X,x}^J(B)$, there are no other harmonic representations in the B_{++} -orbit of $\pi_{\text{dR}}(U(\mathcal{P}), D, f)$. Since $B_{++} = \exp(S(B))$ (by the continuous functional calculus), we can equivalently look at the orbit under the exponential action of the set $S(B)$.

We adapt the proof of [5, Proposition 2.3]. The harmonic condition $(d^+)^*\vartheta = 0$ is equivalent to saying that, for all $\xi \in A^0(X, i \text{ ad } \mathcal{P})$,

$$\langle \vartheta, d^+\xi \rangle = 0 \in A^0(X, B),$$

where $\langle -, - \rangle$ is defined using the Riemannian metric.

Now, the set of flat connections on \mathcal{P} admits a gauge action \star of the smooth automorphism group of \mathcal{P} , and hence via exponentiation an action of the additive group $\Gamma(X, \text{ad } \mathcal{P})$. An isomorphism in B_{++} between two flat connections corresponds to an element of $\exp(\Gamma(X, S(\text{ad } \mathcal{P})))$ for $S(\text{ad } \mathcal{P}) \subset \text{ad } \mathcal{P}$ consisting of symmetric elements, giving a gauge between the respective connections. Thus the $S(B)$ -orbit above is given by looking at the orbit of D under $\Gamma(X, S(\text{ad } \mathcal{P}))$.

By analogy with [5, Proposition 2.3], we fix $\xi \in A^0(X, S(\text{ad } \mathcal{P}))$, let d_t^+ and ϑ_t be the anti-self-adjoint and self-adjoint parts of $\exp(\xi t) \star D$, and set

$$f(t) := \langle \vartheta_t, \vartheta_t \rangle \in A^0(X, B).$$

Now, $d(\exp(\xi t) \star D)/dt = (\exp(\xi t) \star D)\xi = d_t^+\xi + \vartheta_t \wedge \xi$, so $d\vartheta_t/dt = d_t^+\xi$ and

$$f'(t) = 2\langle \xi, (d_t^+)^*\vartheta_t \rangle \in A^0(X, B).$$

In other words, D_t is harmonic if and only if $f'(t) = 0$ for all ξ .

Now, if we set $\widehat{D}_t := d_t^+ - \vartheta_t$, the calculations of [5, Proposition 2.3] adapt to give

$$2f''(t) = \|D_t\xi + \widehat{D}_t\xi\|^2 + \|D_t\xi - \widehat{D}_t\xi\|^2 \in A^0(X, B),$$

where $\|v\|^2 := \langle v, v \rangle$; unlike Corlette, we are only taking inner product with respect to the Kähler metric, not imposing an additional inner product on B .

Note that $f''(t)$ an element of $A^0(X, B_+)$, which lies in $A^0(X, B_{++})$ unless $D_t\xi = 0$. If we start with a harmonic connection D , this implies that $\exp(\xi) \star D$ is harmonic if and only if $D\xi = 0$. However, when $D\xi = 0$ we have $\exp(\xi) \star D = D$, showing that D is the unique harmonic connection in its B_{++} -orbit. \square

Corollary 3.2 For a complex C^* -algebra B and an element $p \in \mathbf{R}_{X,x}^J(B^\times)$, the centraliser $Z(\pi_{\text{dR}}(p), B^\times)$ of $\pi_{\text{dR}}(p)$ under the adjoint action of B is given by

$$Z(\pi_{\text{dR}}(p), B^\times) = \exp(\{b \in S(B) \mid e^{ibt} \in Z(p, U(B)) \text{ for all } t \in \mathbb{R}\}) \rtimes Z(p, U(B));$$

beware that this is the semidirect product of a set with a group.

Proof Take $g \in Z(\pi_{\text{dR}}(p), B^\times)$, and observe that the polar decomposition allows us to write $g = \exp(b)u$ for $u \in U(B)$ and $b \in S(B)$. Since π_{dR} is $U(B)$ -equivariant, we have

$$\text{ad}_{\exp(b)} \pi_{\text{dR}}(\text{ad}_u p) = \text{ad}_g \pi_{\text{dR}}(p) = \pi_{\text{dR}}(p).$$

Thus Proposition 3.1 implies that $\text{ad}_u p = p$, so $u \in Z(p, U(B))$.

Since $Z(\pi_{\text{dR}}(p), B^\times)$ is a group, this implies that $\exp(b) \in Z(\pi_{\text{dR}}(p), B^\times)$, and hence that $\exp(b)$ commutes with the image of $\pi_{\text{dR}}(p)$. We may apply the continuous functional calculus to take logarithms, showing that b itself commutes with the image of $\pi_{\text{dR}}(p)$, so ibt does also. But then $\exp(ibt) \in Z(p, U(B))$ for all t , as required.

Conversely, if $\exp(ibt) \in Z(p, U(B))$ for all t , then $\exp(-ibt)\pi_{\text{dR}}(p)\exp(ibt) = \pi_{\text{dR}}(p)$ and, differentiating in t for each element of $\pi_1(X, x)$, we see that ib commutes with $\pi_{\text{dR}}(p)$. Thus $\exp(b) \in Z(\pi_{\text{dR}}(p), B^\times)$. □

3.1.2 Topological representation spaces and completely bounded maps

Lemma 3.3 *For the real pro- C^* -algebra $E_{X,x}^J$ of Proposition 2.22, there is a canonical map $\pi_{\text{dR}}: \text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, B) \rightarrow \mathbf{R}_{X,x}^{\text{dR}}(B)$, functorial in real Banach algebras B .*

Proof Given $f: E_{X,x}^J \rightarrow B$, Lemma 1.20 factors f as the composition of a surjective C^* -homomorphism $g: E_{X,x}^J \rightarrow C$ and a continuous embedding $C \hookrightarrow B$. The de Rham projection of Definition 2.21 then gives us an element $\pi_{\text{dR}}(g) \in \mathbf{R}_{X,x}^{\text{dR}}(C)$. Combining this with the embedding $C^\times \rightarrow B^\times$ then provides the required element of $\mathbf{R}_{X,x}^{\text{dR}}(B)$. □

Proposition 3.4 *For any real C^* -algebra B , the map of Lemma 3.3 induces an injection*

$$\pi_{\text{dR}}: \text{Hom}(E_{X,x}^J, B)_{\text{cb}} \hookrightarrow \mathbf{R}_{X,x}^{\text{dR}}(B)$$

for the completely bounded morphisms of Definition 1.21.

Proof Since B can be embedded as a closed C^* -subalgebra of $L(H)$ for some complex Hilbert space H , we may replace B with $L(H)$. By Lemma 1.22, any completely bounded homomorphism $f: E_{X,x}^J \rightarrow L(H)$ is conjugate to a $*$ -morphism, since $E_{X,x}^J$ is a pro- C^* -algebra. Therefore Proposition 3.1 shows that

$$\text{Hom}(E_{X,x}^J, L(H))_{\text{cb}}/\text{GL}(H) \hookrightarrow \mathbf{R}_{X,x}^{\text{dR}}(L(H))/\text{GL}(H).$$

Take a homomorphism $f: E_{X,x}^J \rightarrow L(H)$ of C^* -algebras; it suffices to show that the centraliser of f and of $\pi_{\text{dR}}(f)$ are equal. By Corollary 3.2, we know that

$$\begin{aligned} &Z(\pi_{\text{dR}}(f), \text{GL}(H)) \\ &= \exp(\{b \in S(L(H)) \mid e^{ibt} \in Z(f, U(H)) \text{ for all } t \in \mathbb{R}\}) \rtimes Z(f, U(H)). \end{aligned}$$

If e^{ibt} commutes with f for all f , then $e^{ibt}fe^{-ibt} = f$, and differentiating in t shows that b commutes with f . Therefore $\exp(b)$ commutes with f , showing that

$$Z(\pi_{\text{dR}}(f), \text{GL}(H)) \subset Z(f, \text{GL}(H)).$$

The reverse inclusion is automatic, giving the required result. □

Remark 3.5 In [23; 20], the proreductive fundamental group $\pi_1(X, x)_k^{\text{red}}$ is studied — this is an affine group scheme over k . By Tannakian duality [7, Chapter II], we can interpret the dual $O(\pi_1(X, x)_{\mathbb{R}}^{\text{red}})^{\vee}$ of the ring of functions as the ring of discontinuous $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant endomorphisms of $\eta_x^{\text{dR,ss}}$.

The group scheme $\pi_1(X, x)_k^{\text{red}}$ encodes all the information about the sets of finite-dimensional representations of $\pi_1(X, x)$. As we will now see, $(E_{X,x}^J)_{\text{PN}}$ encodes all the information about their topologies as well.

Theorem 3.6 For any positive integer n , π_{dR} gives a homeomorphism $\pi_{\text{dR,ss}}$ between the space $\text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, \text{Mat}_n(\mathbb{C}))$ with the topology of pointwise convergence and the subspace of $\mathbf{R}_{X,x}^{\text{dR}}(\text{Mat}_n(\mathbb{C}))$ whose points correspond to semisimple local systems.

Proof The isomorphism $\pi_{\text{dR,ss}}$ is given on points by the proof of [5, Theorem 3.3], since completely bounded algebra homomorphisms $E_{X,x}^J \rightarrow \text{Mat}_n(\mathbb{C})$ are those conjugate to $*$ -homomorphisms, which in turn correspond to harmonic local systems. We need to show that this is a homeomorphism.

Consider the map $\pi_{\text{dR}}^{\sharp}: E_{X,x}^{\text{dR}} \rightarrow E_{X,x}^J$ of pro-Banach algebras. If $T_i \rightarrow T$ is a convergent net in $\text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, \text{Mat}_n(\mathbb{C}))$, then $T_i(\pi_{\text{dR}}^{\sharp}(\gamma)) \rightarrow T(\pi_{\text{dR}}^{\sharp}(\gamma))$, so $\pi_{\text{dR,ss}}$ is continuous.

Now, the de Rham projection $\pi_{\text{dR}}: \mathbf{R}_{X,x}^J(B) \rightarrow \mathbf{R}_{X,x}^{\text{dR}}(B)$ is automatically injective for C^* -algebras B . Thus the inclusion in $E_{X,x}^J$ of the pro- C^* -subalgebra generated by $\pi_{\text{dR}}^{\sharp}(\pi_1(X, x))$ must be an epimorphism, since $\mathbb{R}[\pi_1(X, x)]$ is dense in $E_{X,x}^{\text{dR}}$. By [21, Proposition 2], an epimorphism of C^* -algebras is surjective, so $E_{X,x}^J$ must be generated as a pro- C^* -algebra by $\pi_{\text{dR}}^{\sharp}(\pi_1(X, x))$, and as a pro-Banach algebra by $\pi_{\text{dR}}^{\sharp}(\pi_1(X, x)) \cup \pi_{\text{dR}}^{\sharp}(\pi_1(X, x))^*$.

Now, if we have a convergent sequence $\pi_{\text{dR}}(T_i) \rightarrow \pi_{\text{dR}}(T)$, then $T_i(\pi_{\text{dR}}^\# \gamma) \rightarrow T(\pi_{\text{dR}}^\# \gamma)$ for all $\gamma \in \pi_1(X, x)$, so it suffices to show that the same holds for $(\pi_{\text{dR}}^\# \gamma)^*$. Given $T \in \text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, \text{Mat}_n(\mathbb{C}))$, define T^* by $T^*(e) := T(e^*)^*$. We wish to show that $T_i^*(\pi_{\text{dR}}^\# \gamma) \rightarrow T^*(\pi_{\text{dR}}^\# \gamma)$, which will follow if $\pi_{\text{dR}}(T_i^*) \rightarrow \pi_{\text{dR}}(T^*)$.

As in the proof of Proposition 3.4, we can write $T = \text{ad}_g S$ for $S: E_{X,x}^J, \text{Mat}_n(\mathbb{C})$ a $*$ -homomorphism and $g \in \text{GL}_n(\mathbb{C})$. Then $T^* = \text{ad}_{(g^*)^{-1}} S = \text{ad}_{(gg^*)^{-1}} T$; this means that if we write $\pi_{\text{dR}}(T) = (\mathcal{V}, D, f)$, then $\pi_{\text{dR}}(T^*) = (\mathcal{V}, D, (f^*)^{-1})$, where $f^*: x^* \mathcal{V} \rightarrow \mathbb{C}^n$ is defined using the harmonic metric on \mathcal{V} and the standard inner product on \mathbb{C}^n .

Explicitly, this means that we can describe the involution $*$ on semisimple elements of $\mathbf{R}_{X,x}^B(\text{Mat}_n(\mathbb{C}))$ by $\rho^* = (C(\rho)^{-1})^\dagger$, where C is the Cartan involution of [23]. If $D = d^+ + \vartheta$ is the decomposition into antihermitian and hermitian parts with respect to the harmonic metric, then $C(\mathcal{V}, d^+ + \vartheta, f) = (\mathcal{V}, d^+ - \vartheta, f)$. The proof of [5, Theorem 3.3] ensures that the decomposition $D \mapsto (d^+, \vartheta)$ is continuous in D , so C is continuous. Hence $\pi_{\text{dR}}(T) \mapsto \pi_{\text{dR}}(T^*)$ is also continuous, which gives the convergence required. □

3.1.3 The polynormal completion and Tannaka duality

Definition 3.7 Let $\text{FDR}_{X,x}^{\text{dR}}$ be the category of pairs (V, p) for $V \in \text{FDVect}$ and $p \in \mathbf{R}_{X,x}^{\text{dR}}(\text{End}(V))$. Morphisms $f: (V_1, p_1) \rightarrow (V_2, p_2)$ are given by linear maps $f: V_1 \rightarrow V_2$ for which the adjoint action of

$$\begin{pmatrix} \text{id} & 0 \\ f & \text{id} \end{pmatrix} \in \begin{pmatrix} \text{End}(V_1) & 0 \\ \text{Hom}(V_1, V_2) & \text{End}(V_2) \end{pmatrix} \quad \text{on } \text{FDR}_{X,x}^{\text{dR}} \begin{pmatrix} \text{End}(V_1) & 0 \\ \text{Hom}(V_1, V_2) & \text{End}(V_2) \end{pmatrix}$$

fixes $p_1 \oplus p_2$.

Write $\eta_x^{\text{dR}}: \text{FDR}_{X,x}^{\text{dR}} \rightarrow \text{FDVect}$ for the fibre functor $(V, p) \mapsto V$. Let $\text{FDR}_{X,x}^{\text{dR,ss}} \subset \text{FDR}_{X,x}^{\text{dR}}$ be the full subcategory in which objects correspond to semisimple local systems, with fibre functor $\eta_x^{\text{dR,ss}}$.

Proposition 3.8 *The ring $(E_{X,x}^J)_{\text{PN}}$ is isomorphic to the ring of continuous $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant endomorphisms of $\eta_x^{\text{dR,ss}}$.*

Proof This just combines Lemma 1.28 and Theorem 3.6. □

Remark 3.9 This leads us to contemplate the structure of the ring of continuous endomorphisms f of η_x^{dR} . Any finite-dimensional \mathbb{C} -algebra arises as a subalgebra of some matrix algebra, so any such f induces continuous maps $\text{Hom}_{\text{Gp}}(\pi_1(X, x), B^\times) \rightarrow B$

for all finite-dimensional algebras B . In particular, this holds when $B = \text{Mat}_n(A)$ for some Artinian \mathbb{C} -algebra A , from which it follows that the maps

$$f_V: \mathbf{R}_{X,x}^{\text{dR}}(\text{End}(V)) \rightarrow \text{End}(V)$$

are all analytic. In other words, any continuous endomorphism of η_x^{dR} is automatically analytic.

When $\pi_1(X, x)$ is abelian, this ensures that the ring $(E_{X,x}^{\text{dR}})_{\text{FD}} \otimes \mathbb{C}$ of such endomorphisms is the ring of complex analytic functions on $\text{Hom}_{\text{Gp}}(\pi_1(X, x), \mathbb{C}^*)$, which by Example 2.12 is just $\mathbb{C}[\pi_1(X, x)]^{\text{an}}$.

In general, the ring $(E_{X,x}^{\text{dR}})_{\text{FD}}$ is an inverse limit of polynormal Banach algebras, but it is not clear to the author whether it is the propolynormal completion of the Fréchet algebra $E_{X,x}^{\text{dR}}$. In general, the map $\pi_1(X, x) \rightarrow (E_{X,x}^{\text{dR}})_{\text{FD}}$ need not be injective: Section 6.5 of [2] gives examples of Kähler groups with no faithful linear representations.

Definition 3.10 Given a k -normal real C^* -algebra B , define $\mathbf{R}_{X,x}^{\text{dR,ss}}(B) \subset \mathbf{R}_{X,x}^{\text{dR}}(B)$ to be the subspace consisting of those p for which $\psi(p)$ corresponds to a semisimple local system for all $\psi: B \rightarrow \text{Mat}_k(\mathbb{C})$.

Corollary 3.11 For any k -normal real C^* -algebra B , the space $\mathbf{R}_{X,x}^{\text{dR,ss}}(B)$ is isomorphic to the set of continuous algebra homomorphisms $E_{X,x}^J \rightarrow B$.

Proof Since B is k -normal, any such morphism $E_{X,x}^J \rightarrow B$ factors uniquely through $(E_{X,x}^J)_{\text{PN}}$. By Proposition 3.8, a homomorphism $(E_{X,x}^J)_{\text{PN}} \rightarrow B$ corresponds to a continuous $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant functor $p^*: \text{FDRep}(B) \rightarrow \text{FDR}_{X,x}^{\text{dR,ss}}$ of topological categories fibred over FDVect . An element $p \in \mathbf{R}_{X,x}^{\text{dR}}(B)$ satisfies this condition provided p^* maps to $\text{FDR}_{X,x}^{\text{dR,ss}} \subset \text{FDR}_{X,x}^{\text{dR}}$. \square

Remark 3.12 It is natural to ask whether the nonabelian Hodge theorem of [25] extends from finite-dimensional matrix algebras to more general C^* -algebras B . Proposition 3.8 can be thought of as an extension of the correspondence to polynormal C^* -algebras, but it seems unlikely to adapt much further, because the arguments of [25; 5] rely on sequential compactness of U_n . However, it is conceivable that suitable solutions of the heat equation might give rise to asymptotic C^* -homomorphisms.

3.2 Residually finite-dimensional completion, products and complex tori

Definition 3.13 A pro- C^* -algebra A is said to be *residually finite-dimensional* if it has a separating family of finite-dimensional $*$ -representations.

Given a pro- C^* -algebra A , define the pro- C^* -algebra A_{RFD} to be the universal residually finite-dimensional quotient of A . Explicitly, A_{RFD} is the quotient of A with respect to the proideal given by the system of kernels of finite-dimensional $*$ -representations of A .

Note that polynormal C^* -algebras are residually finite-dimensional, so we have completions $A \rightarrow A_{\text{RFD}} \rightarrow A_{\text{PN}}$ for general A .

Proposition 3.14 *Given compact connected Kähler manifolds X and Y , there is an isomorphism $(E_{X \times Y, (x,y)}^J)_{\text{RFD}} \cong (E_{X,x}^J)_{\text{RFD}} \widehat{\otimes} (E_{Y,y}^J)_{\text{RFD}}$.*

Proof The projections give canonical elements of the spaces $R_{X \times Y, (x,y)}^J(E_{X,x}^J)$ and $R_{X \times Y, (x,y)}^J(E_{Y,y}^J)$, which by Lemma 2.24 give rise to a canonical map

$$f: E_{X \times Y, (x,y)}^J \rightarrow E_{X,x}^J \widehat{\otimes} E_{Y,y}^J.$$

By [5], every finite-dimensional representation of $E_{X \times Y, (x,y)}^J$ corresponds to a semi-simple representation of $\pi_1(X \times Y, (x, y)) = \pi_1(X, x) \times \pi_1(Y, y)$, so factors through $E_X^J \widehat{\otimes} E_Y^J$. Since $(E_{X \times Y, (x,y)}^J)_{\text{RFD}} \subset \prod_i \text{End}(V_i)$ where V_i ranges over finite-dimensional irreducible representations, this implies that

$$f_{\text{RFD}}: (E_{X \times Y, (x,y)}^J)_{\text{RFD}} \rightarrow (E_{X,x}^J)_{\text{RFD}} \widehat{\otimes} (E_{Y,y}^J)_{\text{RFD}}$$

is injective. However, the basepoint y gives us a map $X \rightarrow X \times Y$, and hence $E_{X,x}^J \rightarrow E_{X \times Y, (x,y)}^J$, ensuring that $(E_X^J)_{\text{RFD}}$ lies in the image of f_{RFD} ; a similar argument applies to Y . Thus f_{RFD} is surjective, and hence an isomorphism. \square

Remark 3.15 We have only imposed the hypothesis that the manifolds be Kähler in order to use the functoriality properties of Lemma 2.25, since Lemma 2.37 is too weak to apply to the maps between $X \times Y$ and X .

Lemma 3.16 *Given a compact connected Kähler manifold X , the commutative quotient $(E_{X,x}^J)^{\text{ab}}$ is given by*

$$\begin{aligned} (E_{X,x}^J)^{\text{ab}} \otimes \mathbb{C} &= C(\mathbb{H}^1(X, \mathbb{C}^*), \mathbb{C}), \\ (E_{X,x}^J)^{\text{ab}} &= \{f \in C(\mathbb{H}^1(X, \mathbb{C}^*), \mathbb{C}) \mid f(\bar{\rho}) = \overline{f(\rho)}\}. \end{aligned}$$

Proof Since $(E_{X,x}^J)^{\text{ab}}$ is a commutative C^* -algebra, the Gelfand–Naimark theorem gives $(E_{X,x}^J)^{\text{ab}} \otimes \mathbb{C} \cong C(\text{Hom}(E_{X,x}^J, \mathbb{C}), \mathbb{C})$, with $(E_{X,x}^J)^{\text{ab}}$ given by $\text{Gal}(\mathbb{C}/\mathbb{R})$ -invariants. By Section 3.1.3, we have a homeomorphism $\text{Hom}(E_{X,e}^J, \mathbb{C}) \cong \mathbb{H}^1(X, \mathbb{C}^*)$, which completes the proof. \square

Corollary 3.17 For X a complex torus with identity e and a fixed Riemannian metric, we have

$$(E_{X,e}^J)_{\text{RFD}} \otimes \mathbb{C} = C(H^1(X, \mathbb{C}^*), \mathbb{C}),$$

$$(E_{X,e}^J)_{\text{RFD}} = \{f \in C(H^1(X, \mathbb{C}^*), \mathbb{C}) \mid f(\bar{\rho}) = \overline{f(\rho)}\}.$$

Proof Multiplication on X gives a pointed morphism $X \times X \rightarrow X$ and hence, by functoriality of P in X and Proposition 3.14, we have a morphism

$$(E_{X,e}^J)_{\text{RFD}} \widehat{\otimes} (E_{X,e}^J)_{\text{RFD}} \rightarrow (E_{X,e}^J)_{\text{RFD}}$$

of real pro- C^* -algebras, and we may apply Lemma 3.16. □

Remark 3.18 If it were the case that all irreducible representations of $\pi_1(X, x)$ were harmonic and similarly for $\pi_1(Y, y)$, then the proof of Proposition 3.14 would adapt to show that $E_{X \times Y, (x,y)}^J \cong E_{X,x}^J \widehat{\otimes} E_{Y,y}^J$. As in the proof of Corollary 3.17, that would then imply commutativity of $E_{X,e}^J$ for complex tori (X, e) , giving $E_{X,e}^J \otimes \mathbb{C} = C(\text{Hom}(\pi_1(X, e), \mathbb{C}^*))$.

Lemma 3.19 Given a compact connected Kähler manifold X , the grouplike elements $G((E_{X,x}^J)^{\text{ab}})$ (see Lemma 2.11) of the commutative quotient $(E_{X,x}^J)^{\text{ab}}$ are given by

$$G((E_{X,x}^J)^{\text{ab}} \otimes \mathbb{C}) \cong H_1(X, \mathbb{Z} \oplus \mathbb{C}),$$

$$G((E_{X,x}^J)^{\text{ab}}) \cong H_1(X, \mathbb{Z} \oplus \mathbb{R}),$$

with the map $\pi_1(X, x)^{\text{ab}} \rightarrow G((E_{X,x}^J)^{\text{ab}})$ given by the diagonal map $\mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{R}$.

Proof The coalgebra structure on $(E_{X,x}^J)^{\text{ab}}$ corresponds under Lemma 3.16 to the group structure on $H^1(X, \mathbb{C}^*)$. Thus $G((E_{X,x}^J)^{\text{ab}})$ consists of continuous functions $f: H^1(X, \mathbb{C}^*) \rightarrow \mathbb{C}$ with $f(1) = 1$ and $f(ab) = f(a)f(b)$.

We have an isomorphism $\mathbb{C}^* \cong S^1 \times \mathbb{R}$, given by $re^{i\phi} \mapsto (\phi, r)$. Thus $H^1(X, \mathbb{C}^*) \cong H^1(X, S^1 \times H^1(X, \mathbb{R}))$. By Pontrjagin duality, a continuous group homomorphism $H^1(X, S^1) \rightarrow \mathbb{C}^*$ is just an element of $H_1(X, \mathbb{Z})$, and a continuous group homomorphism $H^1(X, \mathbb{R}) \rightarrow \mathbb{C}^*$ is an element of $H_1(X, \mathbb{C})$. □

3.3 The Dolbeault projection

Now let X be a compact connected Kähler manifold with basepoint $x \in X$.

Proposition 3.20 For all complex C^* -algebras B , the Dolbeault projection

$$\pi_{\text{Dol}}: \mathbf{R}_{X,x}^J(B) \rightarrow \mathbf{R}_{X,x}^{\text{Dol}}(B)$$

has the property that, if $p_1, p_2 \in \mathbf{R}_{X,x}^J(B)$ and $\text{ad}_b \pi_{\text{Dol}}(p_1) = \pi_{\text{Dol}}(p_2)$ for some strictly positive self-adjoint element $b \in B$, then $p_1 = p_2$.

Thus

$$\pi_{\text{Dol}}: \mathbf{R}_{X,x}^J(B)/U(B) \rightarrow \mathbf{R}_{X,x}^{\text{Dol}}(B)/B^\times$$

is injective.

Proof The proof of Proposition 3.1 adapts, replacing D with D'' . □

Corollary 3.21 For an element $p \in \mathbf{R}_{X,x}^J(B^\times)$, the centraliser $Z(\pi_{\text{Dol}}(p), B^\times)$ of $\pi_{\text{Dol}}(p)$ under the adjoint action of B is given by

$$Z(\pi_{\text{Dol}}(p), B^\times) = \exp(\{b \in S(B) \mid e^{ibt} \in Z(p, U(B)) \text{ for all } t \in \mathbb{R}\}) \rtimes Z(p, U(B));$$

beware that this is the semidirect product of a set with a group.

Proof The proof of Corollary 3.2 carries over. □

Proposition 3.22 For the real pro- C^* -algebra $E_{X,x}^J$ of Proposition 2.22, there is a canonical map $\text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, B) \rightarrow \mathbf{R}_{X,x}^{\text{Dol}}(B)$, functorial in complex Banach algebras B . This induces an injection

$$\text{Hom}(E_{X,x}^J, B)_{\text{cb}} \hookrightarrow \mathbf{R}_{X,x}^{\text{Dol}}(B)$$

whenever B is a C^* -algebra.

Proof The proofs of Lemma 3.3 and Proposition 3.4 carry over to this context, replacing Proposition 3.1 and Corollary 3.2 with Proposition 3.20 and Corollary 3.21. □

Theorem 3.23 For any positive integer n , there is a homeomorphism $\pi_{\text{Dol,st}}$ between the space $\text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, \text{Mat}_n(\mathbb{C}))$ with the topology of pointwise convergence and the subspace of $\mathbf{R}_{X,x}^{\text{Dol}}(\text{Mat}_n(\mathbb{C}))$ consisting of polystable Higgs bundles E with $\text{ch}_1(E) \cdot [\omega]^{\dim X - 1} = 0$ and $\text{ch}_2(E) \cdot [\omega]^{\dim X - 2} = 0$.

Proof The isomorphism of points is given by [23, Theorem 1]. Replacing Proposition 3.4 with Proposition 3.22, the argument from the proof of Theorem 3.6 shows that the map $\pi_{\text{Dol}}: \text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, \text{Mat}_n(\mathbb{C})) \rightarrow \mathbf{R}_{X,x}^{\text{Dol}}(\text{Mat}_n(\mathbb{C}))$ is continuous, so we just need to show that it is open.

Now, Proposition 7.9 of [25] implies that the isomorphism $\pi_{\text{dR,ss}} \circ \pi_{\text{Dol,st}}^{-1}$ is continuous. Since $\pi_{\text{dR,ss}}$ is a homeomorphism by Theorem 3.6, $\pi_{\text{Dol,st}}$ must also be a homeomorphism. □

Definition 3.24 Let $\text{FDR}_{X,x}^{\text{Dol}}$ be the category of pairs (V, p) for $V \in \text{FDVect}$ and $p \in \mathbf{R}_{X,x}^{\text{Dol}}(\text{End}(V))$, with morphisms defined by adapting the formulae of Definition 3.7.

Let $\text{FDR}_{X,x}^{\text{Dol, st}} \subset \text{FDR}_{X,x}^{\text{Dol}}$ be the full subcategory in which objects correspond to those of Theorem 3.23. Write $\eta_x^{\text{Dol}}: \text{FDR}_{X,x}^{\text{Dol}} \rightarrow \text{FDVect}$, $\eta_x^{\text{Dol, st}}: \text{FDR}_{X,x}^{\text{Dol, st}} \rightarrow \text{FDVect}$ for the fibre functors $(V, p) \mapsto V$.

Proposition 3.25 *The ring $(E_{X,x}^J)_{\text{PN}} \otimes \mathbb{C}$ is isomorphic to the ring of continuous endomorphisms of $\eta_x^{\text{Dol, st}}$.*

Proof The proof of Proposition 3.8 carries over, replacing Theorem 3.6 with Theorem 3.23. □

Definition 3.26 Given a k -normal complex C^* -algebra B , define $\mathbf{R}_{X,x}^{\text{Dol, st}}(B) \subset \mathbf{R}_{X,x}^{\text{Dol}}(B)$ to be the subspace consisting of those p for which $\psi(p) \in \text{FDR}_{X,x}^{\text{Dol, st}}$ for all $\psi: B \rightarrow \text{Mat}_k(\mathbb{C})$.

Corollary 3.27 *For any k -normal complex C^* -algebra B , the space $\mathbf{R}_{X,x}^{\text{Dol, st}}(B)$ is isomorphic to the set of continuous algebra homomorphisms $E_{X,x}^J \rightarrow B$.*

Proof The proof of Corollary 3.11 carries over, replacing Proposition 3.8 with Proposition 3.25. □

3.4 Circle actions and C^* -dynamical systems

Definition 3.28 Define a circle action on a (real or complex) pro- C^* -algebra A to be a continuous group homomorphism from S^1 to $\text{Aut}_{\text{pro}(C^*\text{Alg})}(A)$. Here, the topology on $\text{Aut}_{\text{pro}(C^*\text{Alg})}(A)$ is defined pointwise, so a net f_i converges to f if and only if $f_i(a) \rightarrow f(a)$ for all $a \in A$.

The following is immediate:

Lemma 3.29 *Giving a circle action on a C^* -algebra A is equivalent to giving a pro- C^* -algebra homomorphism $f: A \rightarrow C(S^1, A)$ that satisfies*

- (1) $1^* \circ f = \text{id}_A: A \rightarrow C(\{1\}, A) = A$;
- (2) the diagram

$$\begin{array}{ccc}
 A & \xrightarrow{f} & C(S^1, A) \\
 f \downarrow & & \downarrow C(S^1, f) \\
 C(S^1, A) & \xrightarrow{m^*} & C(S^1 \times S^1, A)
 \end{array}$$

commutes, where $m: S^1 \times S^1 \rightarrow S^1$ is the multiplication.

Lemma 3.30 *If a functor $F: C^* \text{Alg}_k \rightarrow \text{Set}$ is represented by a pro- C^* -algebra A , then to give a circle action on A is equivalent to giving maps*

$$\alpha_B: F(B) \rightarrow F(C(S^1, B)),$$

functorial in B , such that

- (1) $F(1^*) \circ \alpha_B = \text{id}_{F(B)}: F(B) \rightarrow F(B)$;
- (2) *the diagram*

$$\begin{array}{ccc} F(B) & \xrightarrow{\alpha_B} & F(C(S^1, B)) \\ \alpha_B \downarrow & & \downarrow \alpha_{C(S^1, B)} \\ F(C(S^1, B)) & \xrightarrow{F(m^*)} & F(C(S^1 \times S^1, B)) \end{array}$$

commutes, where $m: S^1 \times S^1 \rightarrow S^1$ is the multiplication.

Proof If A has a circle action α , then a homomorphism $h: A \rightarrow B$ gives rise to $C(S^1, h): C(S^1, A) \rightarrow C(S^1, B)$, and we define $\alpha_B(h) := C(S^1, h) \circ \alpha$. This clearly satisfies the required properties.

Conversely, given maps α_B as above, write $A = \varprojlim_i A_i$ as an inverse limit of C^* -algebras, and let $h_i: A \rightarrow A_i$ be the structure map. Then $\alpha_{A_i}(h_i) \in F(C(S^1, A_i))$ is a map $A \rightarrow C(S^1, A_i)$. Since the $\alpha_{A_i}(h_i)$ are compatible, we may take the inverse limit, giving a map

$$\alpha: A \rightarrow C(S^1, A)$$

To see that this is a group homomorphism, just observe that the conditions above ensure that $h_i \circ 1^* \circ \alpha = h_i$ and

$$C(S^1 \times S^1, h_i) \circ C(S^1, \alpha) \circ \alpha = C(S^1 \times S^1, h_i) \circ m^* \alpha$$

for all i . Taking the inverse limit over i shows that this satisfies the conditions of Lemma 3.29.

Finally, note that these two constructions are clearly inverse to each other. □

Proposition 3.31 *For every compact Kähler manifold X , there is a canonical continuous circle action on $E_{X,x}^J$.*

Proof Given $(U(\mathcal{P}), D, f) \in \mathbf{R}_{X,x}^J(B)$, define $\alpha(U(\mathcal{P}), D, f) \in \mathbf{R}_{X,x}^J(C(S^1, B))$ as follows.

Decompose $D = d^+ + \vartheta$ into anti-self-adjoint and self-adjoint parts. Set $\alpha(U(\mathcal{P})) := C(S^1, U(\mathcal{P})) = U(\mathcal{P}) \times_{\mathcal{A}_X^0(U(B))} \mathcal{A}_X^0(C(S^1, U(B)))$, then define $\alpha(D) := d^+ + t \diamond \vartheta$,

where $t \in C(S^1, \mathbb{C})$ is the canonical embedding and \diamond is from Definition 2.17. Thus we have constructed $\alpha(U(\mathcal{P}), D, f) := (C(S^1, U(\mathcal{P}), d^+ + t \diamond \vartheta, C(S^1, f))$, and it is easy to check that this satisfies the conditions of Lemma 3.30. \square

Remark 3.32 By considering finite dimensional quotients of $E_{X,x}^J$, the circle action induces a continuous map

$$S^1 \times E_{X,x}^J \rightarrow O(\pi_1(X, x)_{\mathbb{R}}^{\text{red}})'$$

for $O(\pi_1(X, x)_{\mathbb{R}}^{\text{red}})^\vee$ as in Remark 3.5. This descends to a discontinuous action of S^1 on $O(\pi_1(X, x)_{\mathbb{R}}^{\text{red}})^\vee$, as in [23] (made explicit in the real case as [20, Lemma 6.5]).

Note, however, that the circle action descends to continuous actions on $(E_{X,x}^J)_{\text{RFD}}$ and $(E_{X,x}^J)_{\text{PN}}$ (which are subalgebras of $O(\pi_1(X, x)_{\mathbb{R}}^{\text{red}})^\vee$, though not closed).

Continuity of the circle action ensures that the map

$$S^1 \times \pi_1(X, x) \rightarrow E_{X,x}^J$$

is continuous, and hence that the induced map $S^1 \times \pi_1(X, x) \rightarrow \pi_1(X, x)_{\mathbb{R}}^{\text{red}}(\mathbb{R}) \subset O(\pi_1(X, x)_{\mathbb{R}}^{\text{red}})^\vee$ is continuous. Thus a continuous circle action on $E_{X,x}^J$ gives rise to a pure Hodge structure on $\pi_1(X, x)^{\text{red}}$ in the sense of [23, Section 5], but without needing to refer to $\pi_1(X, x)$ itself. This suggests that the most natural definition of a pure nonabelian Hodge structure is a continuous circle action on a pro- C^* -bialgebra.

Example 3.33 Lemma 3.16 gives an isomorphism

$$(E_{X,x}^J)^{\text{ab}} = \{f \in C(\mathbb{H}^1(X, \mathbb{C}^*), \mathbb{C}) \mid f(\bar{\rho}) = \overline{f(\rho)}\},$$

then Lemma 3.19 shows that the grouplike elements are $G((E_{X,x}^J)^{\text{ab}}) \cong \mathbb{H}_1(X, \mathbb{Z} \oplus \mathbb{R})$. To describe the circle action on $(E_{X,x}^J)^{\text{ab}}$, it thus suffices to describe it on the space $\mathbb{H}^1(X, \mathbb{C}^*)$ of one-dimensional complex representations.

Taking the decomposition $D = d^+ + \vartheta$ of a flat connection D into antihermitian and hermitian parts, note that we must have $(d^+)^2 = \vartheta^2 = 0$, because commutativity of \mathbb{C}^* ensures that commutators vanish, everything else vanishing by hypothesis. This decomposition therefore corresponds to the isomorphism $\mathbb{H}^1(X, \mathbb{C}^*) \cong \mathbb{H}^1(X, \mathbb{S}^1) \times \mathbb{H}^1(X, \mathbb{R})$. Since the action is given by $\vartheta \mapsto t \diamond \vartheta$ for $t \in S^1$, it follows that the S^1 -action is just the \diamond -action on $\mathbb{H}^1(X, \mathbb{R})$.

On $G((E_{X,x}^J)^{\text{ab}}) \cong \mathbb{H}_1(X, \mathbb{Z}) \oplus \mathbb{H}_1(X, \mathbb{R})$, this means that the circle action fixes $\mathbb{H}_1(X, \mathbb{Z})$ and acts with the \diamond -action on $\mathbb{H}_1(X, \mathbb{R}) = \mathbb{H}_1(X, \mathbb{R})^\vee$.

Definition 3.34 Recall from [31, Definition 2.6] that a C^* -dynamical system is a triple (A, G, α) for G a locally compact topological group, A a C^* -algebra and α a continuous action of G on a A .

Lemma 3.35 The circle action α of Proposition 3.31 gives rise to a pro- C^* -dynamical system $(E_{X,x}^J, S^1, \alpha)$, ie an inverse system of C^* -dynamical systems.

Proof Since $E_{X,x}^J$ is a pro- C^* -algebra, we may write it as an inverse system $E_{X,x}^J = \varprojlim_i E_i$ for C^* -algebras E_i . The circle action then sends the structure map $h_i: E_{X,x}^J \rightarrow E_i$ to the map $C(S^1, h_i) \circ \alpha: E_{X,x}^J \rightarrow C(S^1, E_i)$, and evaluation at $1 \in S^1$ recovers h_i . We may therefore set $E_{\alpha(i)}$ to be the closure of the image of $E_{X,x}^J \rightarrow C(S^1, E_i)$, and observe that $E_{\alpha(i)}$ is S^1 -equivariant, with $E_{X,x}^J = \varprojlim_i E_{\alpha(i)}$.

Thus $(E_{X,x}^J, S^1, \alpha) = \varprojlim_i (E_{\alpha(i)}, S^1, \alpha)$ is a pro- C^* -dynamical system. □

The following is taken from [31, Lemma 2.27]:

Definition 3.36 Given a C^* -dynamical system (A, G, α) and $f \in C_c(G, A)$, define

$$\|f\| := \sup\{\|\pi \rtimes U(f)\| : (\pi, U) \text{ is a covariant representation of } (A, G, \alpha)\}.$$

Then $\|-\|$ is called the universal norm, and dominated by $\|-\|_1$.

The completion of $C_c(G, A)$ with respect to $\|-\|$ is the crossed product of A by G , denoted by $A \rtimes_{\alpha} G$.

Definition 3.37 Define a polarised real Hilbert variation of Hodge structures of weight n on X to be a real local system \mathbb{V} , with a pluriharmonic metric on $\mathcal{A}_X^0(\mathbb{V})$, equipped with a Hilbert space decomposition

$$\mathcal{A}_X^0(\mathbb{V}) \otimes \mathbb{C} = \bigoplus_{p+q=n} \widehat{\gamma}^{pq}$$

(where $\widehat{\oplus}$ denotes Hilbert space direct sum), with $\overline{\widehat{\gamma}^{pq}} = \widehat{\gamma}^{qp}$ and satisfying the conditions

$$\partial: \widehat{\gamma}^{pq} \rightarrow \widehat{\gamma}^{pq} \otimes_{\mathcal{A}_X^0(\mathbb{C})} \mathcal{A}_X^{1,0}, \quad \bar{\partial}: \widehat{\gamma}^{pq} \rightarrow \widehat{\gamma}^{p+1, q-1} \otimes_{\mathcal{A}_X^0(\mathbb{C})} \mathcal{A}_X^{0,1}$$

for the decomposition $D = \partial + \bar{\partial} + \theta + \bar{\theta}$ of Definition 2.31.

Proposition 3.38 Real Hilbert space representations of the nonunital pro- C^* -algebra $E_{X,x}^J \rtimes_{\alpha} S^1$ correspond to framed weight 0 polarised real Hilbert variations of Hodge structure.

Proof By [31, Proposition 2.29], a $*$ -representation $E_{X,x}^J \rtimes_{\alpha} S^1 \rightarrow L(H)$ for a Hilbert space H consists of

- (1) a $*$ -representation $\rho: E_{X,x}^J \rightarrow L(H)$, and
- (2) a continuous representation $u: S^1 \rightarrow U(H)$

such that

$$\rho(\alpha(t, a)) = u(t)\rho(a)u(t)^{-1}$$

for all $a \in E_{X,x}^J, t \in S^1$.

In other words, in $R_{X,x}^J(C(S^1, L(H)))$, we have $\alpha(\rho) = u\rho u^{-1}$, so $\alpha(\rho)$ and ρ are isomorphic in the groupoid $\mathcal{R}_X^J(C(S^1, L(H)))$.

Now, by definition of $E_{X,x}^J$, the representation ρ corresponds to a real local system \mathbb{V} , with a pluriharmonic metric on $\mathcal{A}_X^0(\mathbb{V})$ and a Hilbert space isomorphism $f: \mathbb{V}_x \rightarrow H$. The representation $\alpha(\rho)$ corresponds to the connection $\alpha(D) := d^+ + t \diamond \vartheta$ on $\mathcal{A}_X^0(C(S^1, \mathbb{V}))$ for the standard coordinate $t: S^1 \rightarrow \mathbb{C}$, together with framing f .

The condition that $\alpha(\rho)$ and ρ are isomorphic then gives us a unitary gauge transformation g between them. In other words, we have a continuous representation $g: S^1 \rightarrow \Gamma(X, U(\mathcal{A}_X^0(\mathbb{V})))$ with $\alpha(D) \circ g = g \circ D$. We must also have $g_x = u$.

Thus g gives us a Hilbert space decomposition

$$\mathcal{A}_X^0(\mathbb{V}) \otimes \mathbb{C} = \widehat{\bigoplus}_{p+q=0} \mathcal{V}^{pq},$$

with $\overline{\mathcal{V}^{pq}} = \mathcal{V}^{qp}$ and $g(t)$ acting on \mathcal{V}^{pq} as multiplication by t^{p-q} . The condition $\alpha(D) \circ g = g \circ D$ then forces the conditions

$$\partial: \mathcal{V}^{pq} \rightarrow \mathcal{V}^{pq} \otimes_{\mathcal{A}_X^0(\mathbb{C})} \mathcal{A}_X^{10}, \quad \bar{\partial}: \mathcal{V}^{pq} \rightarrow \mathcal{V}^{p+1,q-1} \otimes_{\mathcal{A}_X^0(\mathbb{C})} \mathcal{A}_X^{01X},$$

as required. □

Remark 3.39 For any $E_{X,x}^J$ -representation V , [31, Example 2.14] gives an $E_{X,x}^J \rtimes_{\alpha} S^1$ -representation $\text{Ind}_e^{S^1} V$. Its underlying Hilbert space is just the space $L^2(S^1, V)$ of L^2 -measurable V -valued forms on the circle with respect to Haar measure. For the pluriharmonic local system \mathbb{V} associated to V , this therefore gives a weight 0 variation $\text{Ind}_e^{S^1} \mathbb{V}$ of Hodge structures on X , with $\mathcal{A}_X^0(\text{Ind}_e^{S^1} \mathbb{V}) = \mathcal{A}_X^0(L^2(S^1, \mathbb{V}))$.

4 Hodge decompositions on cohomology

Fix a compact Kähler manifold X .

Definition 4.1 Given a pluriharmonic local system \mathbb{V} in real Hilbert spaces on X (as in Example 2.20), the inner product on \mathbb{V} combines with the Kähler metric on X to give inner products $\langle -, - \rangle$ on the spaces $A^n(X, \mathbb{V})$ for all n . Given an operator F on $A^*(X, \mathbb{V})$, we denote the adjoint operator by F^* . Let $\Delta = DD^* + D^*D$.

4.1 Sobolev spaces

Note that, in general, the Laplacian Δ is not a bounded operator in the L^2 norm. We therefore introduce a system of Sobolev norms:

Definition 4.2 Define $L^n_{(2),s}(X, \mathbb{V})$ to be the completion of $A^n(X, \mathbb{V})$ with respect to the inner product $\langle v, w \rangle_s := \langle v, (I + \Delta)^s w \rangle$.

Note that we then have bounded operators $D, D^c: L^n_{(2),s}(X, \mathbb{V}) \rightarrow L^{n+1}_{(2),s-1}(X, \mathbb{V})$, $D^*, D^{c*}: L^n_{(2),s}(X, \mathbb{V}) \rightarrow L^{n-1}_{(2),s-1}(X, \mathbb{V})$ and $\Delta: L^n_{(2),s}(X, \mathbb{V}) \rightarrow L^n_{(2),s-2}(X, \mathbb{V})$.

Proposition 4.3 The maps $(I + \Delta)^k: L^n_{(2),s}(X, \mathbb{V}) \rightarrow L^n_{(2),s-2k}(X, \mathbb{V})$ are Hilbert space isomorphisms, and there are canonical inclusions $L^n_{(2),s}(X, \mathbb{V}) \subset L^n_{(2),s-1}(X, \mathbb{V})$.

Proof The proofs of [8, Proposition 2.3 and Lemma 2.4] carry over to this generality. □

Definition 4.4 Define $\mathcal{H}^n(X, \mathbb{V}) \subset A^n(X, \mathbb{V})$ to consist of those forms with $\Delta\alpha = 0$. Regard this as a pre-Hilbert space with the inner product $\langle -, - \rangle$.

The following implies that $\mathcal{H}^n(X, \mathbb{V})$ is in fact a Hilbert space:

Lemma 4.5 *The inclusions*

$$\mathcal{H}^n(X, \mathbb{V}) \rightarrow \{\alpha \in L^n_{(2),0} \mid D\alpha = D^*\alpha = 0\} \rightarrow \{\alpha \in L^n_{(2),0} \mid \Delta\alpha = 0\}$$

are Hilbert space isomorphisms.

Proof This is [8, Lemma 2.5] when \mathbb{V} is the local system associated to the $\pi_1(X)$ -representation $\ell^2(\pi_1(X, x))$, but the same proof carries over. □

4.1.1 Decomposition into eigenspaces

Definition 4.6 Define T to be the composition

$$L^n_{(2),0}(X, \mathbb{V}) \xrightarrow{(I+\Delta)^{-1}} L^n_{(2),2}(X, \mathbb{V}) \hookrightarrow L^n_{(2),0}(X, \mathbb{V}).$$

This is bounded and self-adjoint, with spectrum $\sigma(T) \subset (0, 1]$. Thus the spectral decomposition gives $T = \int_{(0,1]} \lambda \, d\pi_\lambda$ for some projection-valued measure π on $(0, 1]$.

For $S \subset [0, \infty)$ measurable, write

$$\nu(S) := \pi_{\{(1+\rho)^{-1} | \rho \in S\}}.$$

Thus

$$T = \int_{\rho \in [0, \infty)} (1 + \rho)^{-1} \, d\nu_\rho,$$

and for $v \in \mathcal{L}_{(2),s+2}(X, \mathbb{V})$ we have

$$\Delta v = \int_{\rho \in [0, \infty)} \rho \, d\nu_\rho v \in \mathcal{L}_{(2),s}(X, \mathbb{V}).$$

If we set $E^n(S) := \nu(S) \mathcal{L}^n_{(2),0}(X, \mathbb{V})$, observe that E^n defines a measurable family of Hilbert spaces on $[0, \infty)$, and that we have direct integral decompositions

$$\mathcal{L}^n_{(2),s}(X, \mathbb{V}) = \int_{\rho \in [0, \infty)}^\oplus (1 + \rho)^{-s/2} E^n_\rho.$$

4.1.2 Harmonic decomposition of eigenspaces Since the operators D, D^c, D^* and D^{c*} commute with Δ , they descend to each graded Hilbert space $E^n(S)$ provided S is bounded above.

If S also has a strictly positive lower bound, then Δ is invertible on $E^n(S)$, so

$$E^n(S) = \Delta E^n(S).$$

As $\Delta = DD^* + D^*D = D^c D^{c*} + D^{c*} D^c$, with $[D, D^c] = [D^*, D^c] = [D^{c*}, D] = 0$, this implies that

$$\begin{aligned} E^n(S) &= DE^{n-1}(S) \oplus D^*E^{n+1}(S) = D^c E^{n-1}(S) \oplus D^{c*} E^{n+1}(S) \\ &= DD^c E^{n-2}(S) \oplus D^* D^c E^n(S) \oplus DD^{c*} E^n(S) \oplus D^* D^{c*} E^{n+2}(S). \end{aligned}$$

Furthermore, $D: D^*E^n(S) \rightarrow DE^{n-1}(S)$ and $D^*: DE^n(S) \rightarrow D^*E^{n+1}(S)$ are isomorphisms, with similar statements for D^c .

If S is just bounded above and does not contain 0, then the statements above still hold if we replace subspaces with their closures:

Proposition 4.7 *There are Hilbert space decompositions*

$$\begin{aligned} \mathcal{L}_{(2),s}^n(X, \mathbb{V}) &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{\Delta \mathcal{L}_{(2),s+2}^n(X, \mathbb{V})} \\ &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{D \mathcal{L}_{(2),s+1}^{n-1}(X, \mathbb{V})} \oplus \overline{D^* \mathcal{L}_{(2),s+1}^{n+1}(X, \mathbb{V})} \\ &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{D^c \mathcal{L}_{(2),s+1}^{n-1}(X, \mathbb{V})} \oplus \overline{D^{c*} \mathcal{L}_{(2),s+1}^{n+1}(X, \mathbb{V})} \\ &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{DD^c \mathcal{L}_{(2),s+2}^{n-2}(X, \mathbb{V})} \oplus \overline{D^* D^c \mathcal{L}_{(2),s+2}^n(X, \mathbb{V})} \\ &\quad \oplus \overline{DD^{c*} \mathcal{L}_{(2),s+2}^n(X, \mathbb{V})} \oplus \overline{D^* D^{c*} \mathcal{L}_{(2),s+2}^{n+2}(X, \mathbb{V})} \end{aligned}$$

for all s .

Proof This is essentially the Hodge theorem, and we can construct the decomposition by a slight modification of [12, pages 94–96].

We can define an approximate Green’s function by

$$G_\epsilon := \int_{(0,1-\epsilon]} \frac{\lambda}{1-\lambda} d\pi_\lambda.$$

Now, note that $(I + \Delta)G_\epsilon = \int_{(0,1-\epsilon]} 1/(1-\lambda) d\pi_\lambda$, which is bounded, so G_ϵ is the composition of the inclusion $L_{(2),s+2}^n(X, \mathbb{V}) \hookrightarrow L_{(2),s}^n(X, \mathbb{V})$ with a map

$$G_\epsilon: L_{(2),s}^n(X, \mathbb{V}) \rightarrow L_{(2),s+2}^n(X, \mathbb{V}).$$

Also note that G_ϵ commutes with Δ , and that $\Delta G_\epsilon = \pi(0, 1 - \epsilon] = I - \pi[1 - \epsilon, 1]$. As $\epsilon \rightarrow 0$, this means that $\pi(1) + \Delta G_\epsilon$ converges weakly to I . Since $\pi(1)$ is projection onto $\mathcal{H}^n(X, \mathbb{V})$, this gives the decomposition required (noting that norm closure and weak closure of a subspace are the same, by the Hahn–Banach theorem). \square

Now, for $v \in D^* E^n(S)$, we have $\langle Dv, Dv \rangle = \langle \Delta v, v \rangle$, so $\|Dv\|^2/\|v\|^2$ lies in S .

Definition 4.8 Define $\Delta^{1/2}: \mathcal{L}_{(2),s+1}^n \rightarrow \mathcal{L}_{(2),s}^n$ by

$$\Delta^{1/2} := \int_{\rho \in [0, \infty)} \rho^{1/2} d\nu_\rho.$$

This gives

$$\begin{aligned} D \mathcal{L}_{(2),s+1}^{n-1}(X, \mathbb{V}) &= \Delta^{1/2} \overline{D \mathcal{L}_{(2),s+2}^{n-1}(X, \mathbb{V})}, \\ \ker D \cap \mathcal{L}_{(2),s}^n(X, \mathbb{V}) &= \overline{D \mathcal{L}_{(2),s+1}^{n-1}(X, \mathbb{V})} \oplus \mathcal{H}^n(X, \mathbb{V}). \end{aligned}$$

Thus

$$H^n \mathcal{L}_{(2),s}^\bullet(X, \mathbb{V}) \cong \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{(D \mathcal{L}_{(2),s-1}^{n-1}(X, \mathbb{V}) / \Delta^{1/2} \overline{D \mathcal{L}_{(2),s-2}^{n-1}(X, \mathbb{V})})}.$$

There are similar statements for the operators D^c , D^* and D^{c*} .

Definition 4.9 Define $\Delta^{-1/2} D: \overline{D^* \mathcal{L}_{(2),s+1}^n(X, \mathbb{V})} \rightarrow \mathcal{L}_{(2),s}^n(X, \mathbb{V})$ by

$$\Delta^{-1/2} := \int_{\rho \in (0, \infty)} \rho^{-1/2} d\nu_\rho \circ D,$$

and define $\Delta^{-1/2} D^c: \overline{D^{c*} \mathcal{L}_{(2),s+1}^n(X, \mathbb{V})} \rightarrow \mathcal{L}_{(2),s}^n(X, \mathbb{V})$ similarly.

Proposition 4.10 *The operator $\Delta^{-1/2} D$ is a Hilbert space isomorphism from the closed subspace $\overline{D^* \mathcal{L}_{(2),s+1}^n(X, \mathbb{V})}$ of $\mathcal{L}_{(2),s}^{n-1}(X, \mathbb{V})$ to the closed subspace $\overline{D \mathcal{L}_{(2),s+1}^{n-1}(X, \mathbb{V})}$ of $\mathcal{L}_{(2),s}^n(X, \mathbb{V})$.*

Likewise, $\Delta^{-1/2} D^c: \overline{D^{c} \mathcal{L}_{(2),s+1}^n(X, \mathbb{V})} \rightarrow \overline{D^c \mathcal{L}_{(2),s+1}^{n-1}(X, \mathbb{V})}$ is a Hilbert space isomorphism.*

Proof We prove this for the first case, the second being entirely similar. Given $a, b \in D^* A^n(X, \mathbb{V})$, we have

$$\begin{aligned} \langle \Delta^{-1/2} Da, \Delta^{-1/2} Db \rangle_s &= \langle (I + \Delta)^s \Delta^{-1/2} Da, \Delta^{-1/2} Db \rangle \\ &= \langle (I + \Delta)^s \Delta^{-1} D^* Da, b \rangle \\ &= \langle (I + \Delta)^s a, b \rangle = \langle a, b \rangle_s, \end{aligned}$$

since $D^* a = 0$ gives $D^* Da = \Delta a$. Taking Hilbert space completions with respect to $\langle -, - \rangle_s$ then gives the required result. \square

4.2 The Hodge decomposition and cohomology

Proposition 4.11 *The nested intersection $\bigcap_s L_{(2),s}^p(X, \mathbb{V})$ is the space $A^p(X, \mathbb{V})$ of C^∞ \mathbb{V} -valued p -forms.*

Proof When \mathbb{V} is finite-dimensional, this is the global Sobolev lemma, but the same proof applies for Hilbert space coefficients. \square

Theorem 4.12 *There are pre-Hilbert space decompositions*

$$\begin{aligned} A^n(X, \mathbb{V}) &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{\Delta A^n(X, \mathbb{V})} \\ &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{DA^{n-1}(X, \mathbb{V})} \oplus \overline{D^* A^{n+1}(X, \mathbb{V})} \\ &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{D^c A^{n-1}(X, \mathbb{V})} \oplus \overline{D^{c*} A^{n+1}(X, \mathbb{V})} \\ &= \mathcal{H}^n(X, \mathbb{V}) \oplus \overline{DD^c A^{n-2}(X, \mathbb{V})} \oplus \overline{D^* D^c A^n(X, \mathbb{V})} \oplus \overline{DD^{c*} A^n(X, \mathbb{V})} \\ &\quad \oplus \overline{D^* D^{c*} A^{n+2}(X, \mathbb{V})} \end{aligned}$$

for all n .

Proof We just take the inverse limit \varprojlim_s of the decomposition in Proposition 4.7, and then make the substitution of Proposition 4.11. \square

4.2.1 Reduced cohomology

Definition 4.13 Given a cochain complex C^\bullet in topological vector spaces, write

$$\overline{H}^n(C^\bullet) := H^n(C^\bullet) / \overline{\{0\}},$$

where $\overline{\{0\}}$ is the closure of 0. Note that we could equivalently define \overline{H}^* as the quotient of the space of cocycles by the *closure* of the space of coboundaries.

Given a local system \mathbb{V} in topological vector spaces on X , define

$$\overline{H}^n(X, \mathbb{V}) := \overline{H}^n(A^\bullet(X, \mathbb{V})).$$

Corollary 4.14 *The maps*

$$\mathcal{H}^n(X, \mathbb{V}) \rightarrow \overline{H}^n(X, \mathbb{V})$$

are all topological isomorphisms.

Corollary 4.15 (principle of two types) *As subspaces of $A^n(X, \mathbb{V})$,*

$$\ker D \cap \ker D^c \cap (\overline{DA^{n-1}(X, \mathbb{V})} + \overline{D^cA^{n-1}(X, \mathbb{V})}) = \overline{DD^cA^{n-2}(X, \mathbb{V})}.$$

Lemma 4.16 (formality) *The morphisms*

$$(\overline{H}_{D^c}^*(X, \mathbb{V}), 0) \leftarrow (Z_{D^c}^*(X, \mathbb{V}), D) \rightarrow (A^*(X, \mathbb{V}), D)$$

induce isomorphisms on reduced cohomology.

Proof The proof of [23, Lemma 2.2] carries over to this generality. □

Remark 4.17 Usually, formality statements such as Lemma 4.16 lead to isomorphisms on deformation functors (see [10] for the original case and [19, Proposition 5.3] for the case closest to our setting).

However, there does not appear to be a natural deformation functor associated to topological DGLAs L with obstruction space $\overline{H}^2(L)$. Thus, in contrast to the proalgebraic case, it is not clear whether there are natural completions of the homotopy groups which can be described in terms of the reduced cohomology ring.

The description of the Archimedean monodromy in [20, Theorem 9.13] is even less likely to adapt, since it features the Green’s operator G , which we have had to replace with a nonconvergent sequence of operators.

4.2.2 Nonreduced cohomology Taking the inverse limit \varprojlim_s of the decompositions of Section 4.1.2, we obtain

$$DA^{n-1}(X, \mathbb{V}) = \Delta^{1/2} \overline{DA^{n-1}(X, \mathbb{V})},$$

$$D^*A^{n+1}(X, \mathbb{V}) = \Delta^{1/2} \overline{D^*A^{n+1}(X, \mathbb{V})},$$

with similar statements for D^c and D^{c*} .

Thus:

Proposition 4.18

$$H^n A^\bullet(X, \mathbb{V}) \cong \mathcal{H}^n(X, \mathbb{V}) \oplus \left(\overline{DA^{n-1}(X, \mathbb{V})} / \Delta^{1/2} \overline{DA^{n-1}(X, \mathbb{V})} \right).$$

Applying the operator $*$ then gives

$$H^n A^\bullet(X, \mathbb{V}) \oplus_{\mathcal{H}^n(X, \mathbb{V})} H^{2d-n} A^\bullet(X, \mathbb{V}') \cong A^n(X, \mathbb{V}) / \Delta^{1/2} A^n(X, \mathbb{V}).$$

Moreover, we have topological isomorphisms

$$\Delta^{-1/2} D: \overline{D^*A^n(X, \mathbb{V})} \rightarrow \overline{DA^{n-1}(X, \mathbb{V})},$$

$$\Delta^{-1/2} D^c: \overline{D^{c*}A^n(X, \mathbb{V})} \rightarrow \overline{D^cA^{n-1}(X, \mathbb{V})},$$

for $\Delta^{-1/2} D$ and $\Delta^{-1/2} D^c$ as in Definition 4.9.

4.3 The W^* -enveloping algebra

4.3.1 $E^J(X, x)'$

Definition 4.19 Given a C^* -algebra B and a positive linear functional f , define B_f to be the Hilbert space completion of B with respect to the bilinear form $\langle a, b \rangle_f := f(a^*b)$. We define π_f to be the representation of B on B_f by left multiplication. Note that this is a cyclic representation, generated by $1 \in B_f$.

Lemma 4.20 Given a C^* -algebra B , the topological dual is given by $B' = \varinjlim_f B'_f$, where f ranges over the filtered inverse system of positive linear functionals on B .

Proof This amounts to showing that $B^{\vee\vee} = \varinjlim_f B_f$. Since the system is filtered (with $f + g \geq f$, g and $B_g \rightarrow B_f$ for $g \geq f$), we see that $\widehat{B} := \varinjlim_f B_f$ is the completion of B with respect to the seminorms $\|b\|_f := f(b^*b)^{1/2}$. The space B_f is the strong closure of B in the cyclic representation π_f , which is just the image of $B^{\vee\vee}$, by the von Neumann bicommutant theorem. Since $B^{\vee\vee}$ is the completion of B with respect to the system of weak seminorms for all representations, this implies that the map $\widehat{B} \rightarrow B^{\vee\vee}$ is an equivalence. \square

Lemma 4.21 For a C^* -algebra B and a B -representation V in Hilbert spaces,

$$\text{Hom}_B(V, B') \cong V'.$$

Proof The space $\text{Hom}_B(V, B')$ consists of continuous B -linear maps $V \rightarrow B'$, and hence to continuous (A, k) -bilinear maps $A \times V \rightarrow k$. These correspond to continuous linear maps $V \rightarrow k$, as required \square

Considering smooth morphisms from X then gives:

Corollary 4.22 For any $E := E^J(X, x)$ -representation V in real Hilbert spaces, with corresponding local system \mathbb{V} on X , there is a canonical topological isomorphism

$$A^\bullet(X, \mathbb{V}) \cong \text{Hom}_E(V', A^\bullet(X, \mathbb{E}')),$$

where \mathbb{E}' is the direct system of pluriharmonic local systems corresponding to the ind- E -representation E' given by left multiplication.

Of course, all the cohomological decomposition results of this section extend to direct limits, so apply to \mathbb{E}' . Conversely, Corollary 4.22 all such results for local systems \mathbb{V} can be inferred from the corresponding results with \mathbb{E}' -coefficients.

Remark 4.23 The comultiplication $E^J(X, x) \rightarrow E^J(X, x) \widehat{\otimes} E^J(X, x)$ of Lemma 2.24 induces a multiplication

$$E^J(X, x)' \widehat{\otimes} E^J(X, x)' \rightarrow E^J(X, x)'$$

on continuous duals, where $(\varprojlim_i E_i)' \widehat{\otimes} (\varprojlim_j E_j)' := \varinjlim_{i,j} E_i \widehat{\otimes} E_j$ for C^* -algebras E_i and the dual tensor product $\widehat{\otimes}$ of [30, page 210]. In particular, $\widehat{\otimes}$ is a crossnorm, so we have a jointly continuous multiplication on $E^J(X, x)'$.

Thus $A^\bullet(X, \mathbb{E}')$ is also equipped with a jointly continuous (graded) multiplication, so has the structure of a differential graded topological algebra.

4.3.2 Failure of continuity Since direct integrals of harmonic representations must be harmonic, Corollary 4.14 and Proposition 4.18 provide us with information about the behaviour of cohomology in measurable families. In particular, they allow us to recover space of measures on the topological spaces of cohomology groups fibred over the moduli spaces of local systems. Thus $E^J_{X,x}$ is a much finer invariant than the proalgebraic completion of $\pi_1(X, x)$.

It is natural to ask whether we can strengthen the Hodge decomposition to incorporate finer topological data. The following example indicates that it does not hold for coefficients in \mathbb{E} itself:

Example 4.24 Let X be a complex torus, so $\pi_1(X, e) \cong \mathbb{Z}^{2g}$. By Corollary 3.17, $(E_{X,e}^J)_{\text{RFD}} \otimes \mathbb{C} = C(\text{Hom}(\pi_1(X, e), \mathbb{C}^*), \mathbb{C}) \cong C((\mathbb{C}^*)^{2g}, \mathbb{C})$.

The complex $A^\bullet(X, \mathbb{E}_{\text{RFD}} \otimes \mathbb{C})$ is then quasi-isomorphic to $H^*(\mathbb{Z}^{2g}, C((\mathbb{C}^*)^{2g}, \mathbb{C}))$. This is given by taking the completed tensor product of $2g$ copies of the complex F given by $C(\mathbb{C}^*, \mathbb{C}) \xrightarrow{z-1} C(\mathbb{C}^*, \mathbb{C})$, so

$$\overline{H}^*(X, \mathbb{E}_{\text{RFD}} \otimes \mathbb{C}) \cong \mathbb{C}[-2g].$$

However, $D^*D + DD^* = |z - 1|^2$ on the complex F , so harmonic forms are given by

$$\mathcal{H}^*(X, \mathbb{E}_{\text{RFD}} \otimes \mathbb{C}) \cong 0.$$

5 Twistor and Hodge structures on cochains, and SU_2

5.1 Preliminaries on nonabelian twistor and Hodge filtrations

The following is [20, Definition 2.3]:

Definition 5.1 Define C to be the real affine scheme $\prod_{\mathbb{C}/\mathbb{R}} \mathbb{A}^1$ obtained from $\mathbb{A}_{\mathbb{C}}^1$ by restriction of scalars, so $C(A) = \mathbb{A}_{\mathbb{C}}^1(A \otimes_{\mathbb{R}} \mathbb{C}) \cong A \otimes_{\mathbb{R}} \mathbb{C}$ for any real algebra A . Choosing $i \in \mathbb{C}$ gives an isomorphism $C \cong \mathbb{A}_{\mathbb{R}}^2$, and we let C^* be the quasiaffine scheme $C - \{0\}$.

We let the real algebraic group $S = \prod_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$ of Definition 2.16 act on C and C^* by inverse multiplication, ie

$$S \times C \rightarrow C, \quad (\lambda, w) \mapsto (\lambda^{-1}w).$$

Fix an isomorphism $C \cong \mathbb{A}^2$, with coordinates u, v on C such that the isomorphism $C(\mathbb{R}) \cong \mathbb{C}$ is given by $(u, v) \mapsto u + iv$. Thus the algebra $O(C)$ associated to C is the polynomial ring $\mathbb{R}[u, v]$. The group scheme S is isomorphic as a scheme to $\mathbb{A}_{\mathbb{R}}^2 - \{(\alpha, \beta) \mid \alpha^2 + \beta^2 = 0\}$, with the isomorphism $S(\mathbb{R}) \cong C^*$ given by $(\alpha, \beta) \mapsto \alpha + i\beta$ and the isomorphism $S(\mathbb{C}) \cong (C^*)^2$ given by $(\alpha, \beta) \mapsto (\alpha + i\beta, \alpha - i\beta)$.

By [20, Corollary 2.6 and Proposition 2.28], real Hodge filtrations (resp. real twistor structures) correspond to S -equivariant (resp. \mathbb{G}_m -equivariant) flat vector bundles on C^* . The latter arises because $[C^*/\mathbb{G}_m] \simeq \mathbb{P}_{\mathbb{R}}^1$, so \mathbb{G}_m -equivariant sheaves on C^* correspond to sheaves on \mathbb{P}^1 .

The following is [20, Definition 2.15]:

Definition 5.2 Define an S -action on the real affine scheme SL_2 by

$$(\alpha, \beta, A) \mapsto \begin{pmatrix} 1 & 0 \\ 0 & \alpha^2 + \beta^2 \end{pmatrix} A \begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix}^{-1} = \begin{pmatrix} \alpha^2 + \beta^2 & 0 \\ 0 & 1 \end{pmatrix}^{-1} A \begin{pmatrix} \alpha & -\beta \\ \beta & \alpha \end{pmatrix}.$$

Let $\text{row}_1: SL_2 \rightarrow C^*$ be the S -equivariant map given by projection onto the first row.

The subgroup scheme $\mathbb{G}_m \subset S$ is given by $\beta = 0$ in the coordinates above, and there is a subgroup scheme $S^1 \subset S$ given by $\alpha^2 + \beta^2 = 1$. These induce an isomorphism $(\mathbb{G} \times S^1)/(-1, -1) \cong S$. On these subgroups, the action on SL_2 simplifies as follows:

Lemma 5.3 *The action of $\mathbb{G}_m \subset S$ on SL_2 is given by*

$$(\alpha, A) \mapsto \begin{pmatrix} \alpha^{-1} & 0 \\ 0 & \alpha \end{pmatrix} A$$

and the action of $S^1 \subset \mathbb{G}_m$ is given by

$$(\alpha, \beta, A) \mapsto A \begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix}^{-1}.$$

The action of $S^1 \subset S$ descends via the maps above to an action on $\mathbb{P}_{\mathbb{R}}^1$, which is just given by identifying S^1 with the real group scheme SO_2 .

5.2 The twistor structure on cochains

Fix a compact Kähler manifold X .

Definition 5.4 Define $\tilde{D}: A^n(X, \mathbb{V}) \otimes \mathcal{O}_{C^*} \rightarrow A^{n+1}(X, \mathbb{V}) \otimes \mathcal{O}_{C^*}$ by

$$\tilde{D} = uD + vD^c,$$

and write $\tilde{A}^\bullet(X, \mathbb{V})$ for the resulting complex. Put a \mathbb{G}_m -action on $\tilde{A}^\bullet(X, \mathbb{V})$ by letting $A^n(X, \mathbb{V})$ have weight n , and giving \mathcal{O}_{C^*} the action of $\mathbb{G}_m \subset S$ from Definition 5.1.

Define $\tilde{Z}^n(X, \mathbb{V}) := \ker(\tilde{D}: A^n(X, \mathbb{V}) \otimes \mathcal{O}_{C^*} \rightarrow A^{n+1}(X, \mathbb{V}) \otimes \mathcal{O}_{C^*})$, $\tilde{B}^n(X, \mathbb{V}) := \text{Im}(\tilde{D}: A^{n-1}(X, \mathbb{V}) \otimes \mathcal{O}_{C^*} \rightarrow A^n(X, \mathbb{V}) \otimes \mathcal{O}_{C^*})$ and

$$\bar{H}^n(X, \mathbb{V}) := \tilde{Z}^n(X, \mathbb{V}) / \overline{\tilde{B}^n(X, \mathbb{V})}.$$

By analogy with [20, Proposition 2.28 and Theorem 7.1], we regard the \mathbb{G}_m -equivariant complex $\tilde{A}^\bullet(X, \mathbb{V})$ over C^* as a twistor filtration on $A^n(X, \mathbb{V})$.

Corollary 5.5 *The canonical inclusion $\mathcal{H}^n(X, \mathbb{V})(n) \otimes \mathcal{O}_{C^*} \rightarrow \widetilde{\mathcal{H}}^n(X, \mathbb{V})$ is a \mathbb{G}_m -equivariant topological isomorphism.*

Proof It suffices to prove this on pulling back along the flat cover $\text{row}_1: \text{SL}_2 \rightarrow C^*$. We define $\widetilde{D}^*: A^n(X, \mathbb{V}) \otimes \mathcal{O}(\text{SL}_2) \rightarrow A^{n-1}(X, \mathbb{V}) \otimes \mathcal{O}(\text{SL}_2)$ by $\widetilde{D}^* = yD^* - xD^{c*}$. Then $[\widetilde{D}, \widetilde{D}^*] = \Delta$, and since Δ commutes with D and D^c it also commutes with \widetilde{D} .

The result now follows with the same proof as that of Corollary 4.14, replacing D and D^* with \widetilde{D} and \widetilde{D}^* . □

Proposition 5.6 *If we write $\mathcal{H}^n = \mathcal{H}^n(X, \mathbb{V})$ and $M^m = \overline{DD^cA^{m-2}(X, \mathbb{V})}$, then there is a \mathbb{G}_m -equivariant isomorphism*

$$\widetilde{\mathcal{H}}^n(X, \mathbb{V}) \cong [(\mathcal{H}^n \oplus M/\Delta^{1/2}M^n)(n) \oplus (M^{n+1}/\Delta^{1/2}M^{n+1})(n-1)] \otimes \mathcal{O}_{C^*}$$

of quasicoherent sheaves on C^ .*

Proof Writing $A^m := A^m(X, \mathbb{V})$, we have a commutative diagram

$$\begin{array}{ccc} & \overline{DD^cA^{n-1}}(n+1) \otimes \mathcal{O}_{C^*} & \\ \nearrow^{vD^c} & & \nwarrow^{-uD} \\ \overline{DD^cA^n}(n) \otimes \mathcal{O}_{C^*} & & \overline{D^*D^cA^n}(n) \otimes \mathcal{O}_{C^*} \\ \nwarrow^{uD} & & \nearrow^{vD^c} \\ & \overline{D^*D^cA^{n+1}}(n-1) \otimes \mathcal{O}_{C^*} & \end{array}$$

which we may regard as a bicomplex. By Theorem 4.12, the complex $\widetilde{A}^\bullet(X, \mathbb{V})$ decomposes into a direct sum of \mathcal{H}^n 's and total complexes of the bicomplexes above.

Arguing as in Proposition 4.10, we have topological isomorphisms

$$\begin{aligned} \Delta^{-1}DD^c: \overline{D^*D^cA^{n+1}} &\rightarrow \overline{DD^cA^{n-1}}, \\ \Delta^{-1/2}D: \overline{D^*D^cA^n} &\rightarrow \overline{DD^cA^{n-1}}, \\ \Delta^{-1/2}D^c: \overline{DD^cA^n} &\rightarrow \overline{DD^cA^{n-1}}, \end{aligned}$$

so the bicomplex above is linearly isomorphic to:

$$\begin{array}{ccc} & \overline{DD^cA^{n-1}}(n+1) \otimes \mathcal{O}_{C^*} & \\ \nearrow^{v\Delta^{1/2}} & & \nwarrow^{-u\Delta^{1/2}} \\ \overline{DD^cA^{n-1}}(n) \otimes \mathcal{O}_{C^*} & & \overline{DD^cA^{n-1}}(n) \otimes \mathcal{O}_{C^*} \\ \nwarrow^{u\Delta^{1/2}} & & \nearrow^{v\Delta^{1/2}} \\ & \overline{DD^cA^{n-1}}(n-1) \otimes \mathcal{O}_{C^*} & \end{array}$$

Since the ideal (u, v) generates \mathcal{O}_{C^*} , cohomology of the top level of the associated total complex is just

$$(\overline{DD^c A^{n-1}} / \Delta^{1/2} \overline{DD^c A^{n-1}})(n+1) \otimes \mathcal{O}_{C^*},$$

while cohomology of the bottom level is 0. Moreover, the map

$$\overline{DD^c A^{n-1}}(n-1) \otimes \mathcal{O}_{C^*} \xrightarrow{(u,v)} \overline{DD^c A^{n-1}}(n)^2 \otimes \mathcal{O}_{C^*}$$

is an isomorphism to the kernel of $(v, -u)$, so cohomology of the middle level is isomorphic to

$$(\overline{DD^c A^{n-1}}(n-1) / \Delta^{1/2} \overline{DD^c A^{n-1}}(n-1)) \otimes \mathcal{O}_{C^*} \otimes \mathcal{O}_{C^*},$$

which completes the proof. □

Remark 5.7 In particular, $\tilde{H}^n(X, \mathbb{V})$ is of weights $n, n-1$ in general, unlike $\tilde{H}^n(X, \mathbb{V})$, which is pure of weight n . This means that the weight filtration given good truncation cannot define a mixed twistor structure on $\tilde{H}^n(X, \mathbb{V})$.

We now have the following generalisation of the principle of two types:

Lemma 5.8 *As subspaces of $A^n(X, \mathbb{V}) \otimes O(SL_2)$,*

$$\begin{aligned} \ker \tilde{D} \cap \ker \tilde{D}^c \cap (\overline{\tilde{D}(A^{n-1}(X, \mathbb{V}) \otimes O(SL_2))} + \overline{\tilde{D}^c(A^{n-1}(X, \mathbb{V}) \otimes O(SL_2))}) \\ = \overline{\tilde{D}\tilde{D}^c(A^{n-2}(X, \mathbb{V}) \otimes O(SL_2))}. \end{aligned}$$

Proof This follows from Corollary 4.15, with the same reasoning as [20, Lemma 1.3]. □

Lemma 5.9 (formality) *The morphisms*

$$\begin{aligned} (\overline{H}_{\tilde{D}^c}^*(A^*(X, \mathbb{V}) \otimes O(SL_2)), 0) \leftarrow (Z_{\tilde{D}^c}^*(A(X, \mathbb{V}) \otimes O(SL_2)), \tilde{D}) \\ \rightarrow (A^*(X, \mathbb{V}) \otimes O(SL_2), \tilde{D}) \end{aligned}$$

induce isomorphisms on reduced cohomology.

Proof The proof of [23, Lemma 2.2] carries over to this generality, using Lemma 5.8. □

Following Corollary 4.22 and Remark 4.23, the results above can all be regarded as statements about the topological differential graded algebra $\tilde{A}^\bullet(X, \mathbb{E}')$.

5.3 The analytic Hodge filtration on cochains

Recall from Section 4.3.1 that the local system \mathbb{E}' on X is defined to correspond to the $\pi_1(X, x)$ -representation given by left multiplication on $E^J(X, x)'$.

Proposition 5.10 *The topological cochain complex $\tilde{A}^\bullet(X, \mathbb{E}')$ is equipped with a continuous circle action, satisfying*

- (1) *the S^1 -action and \mathbb{G}_m -actions on $\tilde{A}^\bullet(X, \mathbb{E}')$ commute;*
- (2) *the action of $S^1 \subset \mathbb{C}^* = S(\mathbb{R})$ on C^* makes $\tilde{A}^\bullet(X, \mathbb{E}')$ into an S^1 -equivariant sheaf on C^* ; and*
- (3) *$-1 \in S^1$ acts as $-1 \in \mathbb{G}_m$.*

Proof Since S^1 acts on $E^J(X, x)$, it acts on \mathbb{E}' , and we denote this action by $v \mapsto t \otimes v$ for $t \in S^1$. We may now adapt the proof of [20, Theorem 6.12], defining an S^1 -action on $\mathcal{A}^*(X, \mathbb{R}) \otimes_{\mathbb{R}} \mathbb{E}'$ by setting $t \boxtimes (a \otimes v) := (t \diamond a) \otimes (t^2 \otimes v)$ for $t \in S^1$ and \diamond as in Definition 2.17. Passing to the completion $\mathcal{A}^*(X, \mathbb{E}')$ completes the proof, with continuity following from Proposition 3.31. □

Remark 5.11 If the circle action of Proposition 5.10 were algebraic, then by [20, Lemma 2.23] it would correspond to a Hodge filtration on $A^\bullet(X, \mathbb{E}')$. Since finite-dimensional circle representations are algebraic, we may regard Proposition 5.10 as the natural structure of an infinite-dimensional Hodge filtration.

In Proposition 5.10, note that we can of course replace $E^J(X, x)$ with any inverse system B of C^* -algebra quotients of $E^J(X, x)$ to which the S^1 -action descends, provided we replace \mathbb{E} with the local system associated to B .

As observed in Section 4.3.1, we may substitute $\mathbb{V} = \mathbb{E}$ in Proposition 5.6 and Lemma 5.9. Note that the resulting isomorphisms are then equivariant with respect to the circle action of Proposition 5.10.

5.4 SU_2

As we saw in Corollary 5.5, in order to define the adjoint operator \tilde{D}^* to \tilde{D} , it is necessary to pull $\tilde{A}^\bullet(X, \mathbb{V})$ back along the morphism $\text{row}_1: \text{SL}_2 \rightarrow C^*$. This gives us the complex

$$\text{row}_1^* \tilde{A}^\bullet(X, \mathbb{V}) = (A^*(X, \mathbb{V}) \otimes O(\text{SL}_2), \tilde{D}),$$

where $\tilde{D} = uD + vD^c$, with adjoint $\tilde{D}^* = yD^* - xD^{c*}$.

This leads us to consider the $*$ -structure on $O(\mathrm{SL}_2)$ determined by $u^* = y$ and $v^* = -x$. This implies $x^* = -v$ and $y^* = u$, so

$$\begin{pmatrix} u & v \\ x & y \end{pmatrix}^* = \begin{pmatrix} y & -x \\ -v & u \end{pmatrix},$$

or $A^* = (A^{-1})^t$.

Lemma 5.12 *The real C^* -enveloping algebra $C^*(O(\mathrm{SL}_2))$ of the real $*$ -algebra $O(\mathrm{SL}_2)$ is the ring of continuous complex functions f on SU_2 for which*

$$f(\bar{A}) = \overline{f(A)}.$$

Proof A $*$ -morphism $O(\mathrm{SL}_2) \rightarrow \mathbb{C}$ is a matrix $A \in \mathrm{SL}_2(\mathbb{C})$ with $\bar{A} = A^* = (A^{-1})^t$, so $A \in \mathrm{SU}_2$. Thus the Gelfand representation gives $C^*(O(\mathrm{SL}_2)) \otimes \mathbb{C} \cong C(\mathrm{SU}_2, \mathbb{C})$.

Now, writing $\mathrm{Gal}(\mathbb{C}/\mathbb{R}) = \langle \tau \rangle$, and taking $f \in O(\mathrm{SL}_2) \otimes \mathbb{C}$ and $A \in \mathrm{SU}_2$, we have

$$\tau(f)(A) = \overline{f(\bar{A})}.$$

This formula extends to give a $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ -action on $C(\mathrm{SU}_2, \mathbb{C})$, and Lemma 1.11 then gives

$$C^*(O(\mathrm{SL}_2)) = C(\mathrm{SU}_2, \mathbb{C})^\tau. \quad \square$$

Note that complex conjugation on SU_2 is equivalent to conjugation by the matrix $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

5.4.1 The Hopf fibration The action of $S(\mathbb{C})$ on $\mathrm{SL}_2(\mathbb{C})$ from Definition 5.2 does not preserve SU_2 . However, Lemma 5.3 ensures that, for the subgroup schemes $\mathbb{G}_m, S^1 \subset S$, the groups $S^1 = S^1(\mathbb{R}) \subset S^1(\mathbb{C}) \cong \mathbb{C}^*$ and $S^1 \subset \mathbb{C}^* \cong \mathbb{G}_m(\mathbb{C})$ both preserve SU_2 .

Thus, in the C^* setting, the S -action becomes an action of $(S^1 \times S^1)/(-1, -1)$ on SU_2 , given by

$$(s, t, A) \mapsto \begin{pmatrix} s^{-1} & 0 \\ 0 & s \end{pmatrix} A \begin{pmatrix} \Re t & \Im t \\ -\Im t & \Re t \end{pmatrix}^{-1}.$$

Moreover, there is a $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ -action on this copy of $S^1 \times S^1$, with the nontrivial automorphism τ given by $\tau(s, t) = (s^{-1}, t)$. The action of $(S^1 \times S^1)/(-1, -1)$ is then τ -equivariant. Alternatively, we may characterise our group as $S^1 \times S^1 \subset \mathbb{C}^* \times \mathbb{C}^* \cong S(\mathbb{C})$, by sending (s, t) to (st, st^{-1}) . On this group $S^1 \times S^1$, the $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ -action is then given by $\tau(w', w'') = (\overline{w''}, \overline{w'})$.

Now, consider the composition

$$\mathrm{SL}_2 \xrightarrow{\mathrm{row}_1} C^* \rightarrow [C^*/\mathbb{G}_m] \cong \mathbb{P}^1.$$

On taking Gelfand representations of C^* -enveloping algebras, this gives rise to the map

$$\mathrm{SU}_2 \rightarrow \mathbb{P}^1(\mathbb{C}),$$

which is just the Hopf fibration $p: S^3 \rightarrow S^2$, corresponding to the quotient by the action of $S^1 \subset \mathbb{G}_m(\mathbb{C})$ by diagonal matrices. The action of τ on SU_2 and on $\mathbb{P}^1(\mathbb{C})$ is just given by complex conjugation.

5.4.2 Smooth sections If we write ρ_n for the weight n action of \mathbb{G}_m on \mathbb{A}^1 , then we may consider the topological vector bundle

$$\mathrm{SU}_2 \times_{S^1, \rho_n} \mathbb{C}$$

on $\mathbb{P}^1(\mathbb{C})$ for the action of $S^1 \subset \mathbb{G}_m(\mathbb{C})$ above.

Definition 5.13 Let $\mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n)$ be the sheaf of smooth sections of $\mathrm{SU}_2 \times_{S^1, \rho_n} \mathbb{C} \rightarrow \mathbb{P}^1(\mathbb{C})$ and write $A^0(\mathbb{P}^1, \mathbb{C}(n)) := \Gamma(\mathbb{P}^1(\mathbb{C}), \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n))$. Beware that, for $n \neq 0$, there is no local system generating $\mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n)$.

For $U \subset \mathbb{P}^1(\mathbb{C})$, observe that $\Gamma(U, \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n))$ consists of smooth maps

$$f: p^{-1}(U) \rightarrow \mathbb{C} \quad \text{satisfying } f\left(\begin{pmatrix} s^{-1} & 0 \\ 0 & s \end{pmatrix} A\right) = s^n f(A)$$

for all $s \in S^1$.

For the quotient map $q: C^* \rightarrow \mathbb{P}^1$, we may characterise $\Gamma(U, \mathcal{O}_{\mathbb{P}^1}^{\mathrm{hol}}(n))$ as the space of holomorphic maps

$$f: q^{-1}(U) \rightarrow \mathbb{C}$$

satisfying $f(su, sv) = s^n f(u, v)$ for all $s \in \mathbb{C}^*$. The embedding $S^3 \subset C^*(\mathbb{C})$ thus yields

$$\mathcal{O}_{\mathbb{P}^1}^{\mathrm{hol}}(n) \subset \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n),$$

and indeed $\mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n) = \mathcal{O}_{\mathbb{P}^1}^{\mathrm{hol}}(n) \otimes_{\mathcal{O}_{\mathbb{P}^1}^{\mathrm{hol}}} \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}$.

Now, for the conjugate sheaf $\overline{\mathcal{O}_{\mathbb{P}^1}^{\mathrm{hol}}(n)}$, note that $\Gamma(U, \overline{\mathcal{O}_{\mathbb{P}^1}^{\mathrm{hol}}(n)})$ is the space of antiholomorphic maps

$$f: q^{-1}(U) \rightarrow \mathbb{C}$$

satisfying $f(su, sv) = \bar{s}^n f(u, v)$ for all $s \in \mathbb{C}^*$. Thus we have a canonical embedding

$$\overline{\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(n)} \in \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(-n),$$

with $\mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(-n) = \overline{\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(n)} \otimes_{\overline{\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}}} \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}$.

Note that the inclusion $O(\text{SL}_2) \subset C(\text{SU}_2, \mathbb{C})$ gives

$$u, v \in A^0(\mathbb{P}^1, \mathbb{C}(1))^\tau, \quad \bar{u}, \bar{v} \in A^0(\mathbb{P}^1, \mathbb{C}(-1))^\tau,$$

and

$$O(\text{SL}_2) \subset \bigoplus_{n \in \mathbb{Z}} A^0(\mathbb{P}^1, \mathbb{C}(n))^\tau.$$

Definition 5.14 By [20, Definition 2.22], there is a derivation N on $O(\text{SL}_2)$ given by $Nx = u$, $Ny = v$ and $Nu = Nv = 0$ for coordinates $\begin{pmatrix} u & v \\ x & y \end{pmatrix}$ on SL_2 . Since this annihilates u and v , it is equivalent to the $O(\text{SL}_2)$ -linear map

$$\Omega(\text{SL}_2/C^*) \rightarrow O(\text{SL}_2)$$

given by $dx \mapsto u$ and $dy \mapsto v$.

Note that N has weight 2, and extends (by completeness) to give τ -equivariant differentials

$$N: \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n) \rightarrow \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n + 2).$$

Also, N is the composition of the antiholomorphic differential

$$\bar{\partial}_{\mathbb{P}^1}: \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n) \rightarrow \mathcal{A}_{\mathbb{P}^1}^0 \mathbb{C}(n) \otimes_{\overline{\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}}} \overline{\Omega_{\mathbb{P}^1}}$$

with the canonical isomorphism $\Omega_{\mathbb{P}^1} \cong \mathcal{O}(-2)$.

5.4.3 Splittings of the twistor structure As in [20, Remark 2.16], we can characterise the map $\text{row}_1: \text{SL}_2 \rightarrow C^*$ as the quotient $C^* = [\text{SL}_2/\mathbb{G}_a]$, where \mathbb{G}_a acts on SL_2 as left multiplication by $\begin{pmatrix} 1 & 0 \\ \mathbb{G}_a & 1 \end{pmatrix}$. Here, the S -action on \mathbb{G}_a has λ acting as multiplication by $\lambda\bar{\lambda}$.

Therefore the map $q \circ \text{row}_1: \text{SL}_2 \rightarrow \mathbb{P}^1$ is given by taking the quotient of SL_2 by the Borel subgroup $B = \mathbb{G}_m \times \mathbb{G}_a$, for the action above. The action of \mathbb{G}_m corresponds to weights, while the action of \mathbb{G}_a corresponds to the derivation N above, which we regard as the Archimedean monodromy operator as in [20, Section 3.4].

Definition 5.15 Given a pluriharmonic local system \mathbb{V} , define $\check{A}^\bullet(X, \mathbb{V})$ to be the sheaf of $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ -modules associated to the \mathbb{G}_m -equivariant sheaf $\tilde{A}^\bullet(X, \mathbb{E}^\mathbb{V})$ on C^* . Explicitly,

$$\check{A}^\bullet(X, \mathbb{V}) = \left(\bigoplus_n (q_* \tilde{A}^\bullet(X, \mathbb{V})) \otimes_{\mathcal{O}_{\mathbb{P}^1}^{\text{alg}}} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(n) \right)^{\mathbb{G}_m},$$

so

$$\check{A}^n(X, \mathbb{V}) = A^n(X, \mathbb{V}) \otimes_{\mathbb{R}} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(n),$$

with differential $\check{D} = uD + vD^c$ for $u, v \in \Gamma(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$.

Definition 5.16 Write $\mathcal{A}^\bullet(X, \mathbb{V}) := \mathcal{A}_{\mathbb{P}^1}^0 \otimes_{\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}} \check{A}^\bullet(X, \mathbb{V})$, and observe that this admits an operator

$$\check{D}^c := -\bar{v}D + uD^c: \mathcal{A}^n(X, \mathbb{V}) \rightarrow \mathcal{A}^{n+1}(X, \mathbb{V})(-2).$$

Now, applying the map $O(\text{SL}_2) \hookrightarrow \bigoplus_{n \in \mathbb{Z}} A^0(\mathbb{P}^1, \mathbb{C}(n))^\tau$ to Lemmas 5.8 and 5.9 yields the following:

Lemma 5.17 As subspaces of $\mathcal{A}^n(X, \mathbb{V})$,

$$\ker \check{D} \cap \ker \check{D}^c \cap \overline{(\check{D} \mathcal{A}^{n-1}(X, \mathbb{V}) + \check{D}^c \mathcal{A}^{n-1}(X, \mathbb{V})(2))} = \overline{\check{D} \check{D}^c (\mathcal{A}^{n-2}(X, \mathbb{V})(2))}.$$

Thus the morphisms

$$(\overline{\mathcal{H}}_{\check{D}^c}^* (\mathcal{A}^\bullet(X, \mathbb{V})), 0) \leftarrow (\mathcal{A}_{\check{D}^c}^* (\mathcal{A}^\bullet(X, \mathbb{V})), \check{D}) \rightarrow \mathcal{A}^n(X, \mathbb{V})$$

induce isomorphisms on reduced cohomology sheaves.

Now, $\tilde{A}^\bullet(X, \mathbb{E}')$ can be recovered from $\text{row}_1^* \tilde{A}^\bullet(X, \mathbb{E}')$ and its nilpotent monodromy operator N , and, by Lemma 4.16, $\text{row}_1^* \tilde{A}^\bullet(X, \mathbb{E}')$ is equivalent to $\mathcal{H}^*(X, \mathbb{E}') \otimes O(\text{SL}_2)$ up to reduced quasi-isomorphism.

Under the base change above, we have $N = \bar{\partial}_{\mathbb{P}^1}$, giving an exact sequence

$$0 \rightarrow \check{A}^\bullet(X, \mathbb{V}) \rightarrow \mathcal{A}^\bullet(X, \mathbb{V}) \xrightarrow{N} \mathcal{A}^\bullet(X, \mathbb{V})(2) \rightarrow 0.$$

In other words, we can recover the topological DGA $\check{A}^\bullet(X, \mathbb{E}')$ from the differential $\bar{\partial} = N$ on the topological DGA $\mathcal{A}^\bullet(X, \mathbb{E}')$ and the latter is just $\bigoplus_n \mathcal{A}_{\mathbb{P}^1}(\mathcal{H}^n(X, \mathbb{E}')(n))$ up to reduced quasi-isomorphism.

Also note that, when we substitute $\mathbb{V} := \mathbb{E}'$ in Lemma 5.17, the morphisms all become equivariant with respect to the circle action of Proposition 5.10. This action makes $\mathcal{A}^n(X, \mathbb{E}')$ into an S^1 -equivariant sheaf over \mathbb{P}^1 , where the action on \mathbb{P}^1 is given by $t^2 \in S^1$ sending $(u : v)$ to $t \diamond (u : v) = (au - bv : av + bu)$ for $t = a + ib \in S^1$.

6 The twistor family of moduli functors

6.1 Pro-Banach algebras on projective space

On the complex manifold $\mathbb{P}^1(\mathbb{C})$, we have a sheaf $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ of holomorphic functions, which we may regard as a sheaf of pro-Banach algebras. As a topological space, $\mathbb{P}^1(\mathbb{C})$ is equipped with a $\text{Gal}(\mathbb{C}/\mathbb{R})$ -action, the nontrivial element τ acting on points by complex conjugation. There is also an isomorphism

$$\tau_{\mathcal{O}}^{\#}: \tau^{-1} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}} \rightarrow \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$$

given by

$$\tau_{\mathcal{O}}^{\#}(f)(z) = \overline{f(\bar{z})}$$

and satisfying

$$\tau_{\mathcal{O}}^{\#} \circ \tau^{-1}(\tau_{\mathcal{O}}^{\#}) = \text{id}: \mathcal{O}_{\mathbb{P}^1}^{\text{hol}} \rightarrow \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}.$$

Definition 6.1 Define $\text{FrAlg}_{\mathbb{P}^1, \mathbb{C}}$ to be the category of sheaves \mathcal{F} of unital multiplicatively convex Fréchet $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ -algebras (ie countable pro-Banach algebras equipped with a morphism from $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$), quasicohherent in the sense that the maps

$$\mathcal{F}(U) \otimes_{\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(U)}^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(V) \rightarrow \mathcal{F}(V)$$

are isomorphisms for all open subspaces $V \subset U$, where \otimes^{π} here denotes projective tensor product.

Define $\text{FrAlg}_{\mathbb{P}^1, \mathbb{R}}$ to be the category of pairs $(\mathcal{F}, \tau_{\mathcal{F}}^{\#})$ for $\mathcal{F} \in \text{FrAlg}_{\mathbb{P}^1, \mathbb{C}}$ and

$$\tau_{\mathcal{F}}^{\#}: \tau^{-1} \mathcal{F} \rightarrow \mathcal{F}$$

an $(\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}, \tau_{\mathcal{O}}^{\#})$ -linear isomorphism satisfying

$$\tau_{\mathcal{F}}^{\#} \circ \tau^{-1}(\tau_{\mathcal{F}}^{\#}) = \text{id}_{\mathcal{F}}.$$

Note that for any m -convex Fréchet k -algebra B the sheaf $B \otimes_k^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ lies in $\text{FrAlg}_{\mathbb{P}^1, \mathbb{C}}$. When $k = \mathbb{R}$, the involution $\text{id}_B \otimes \tau_{\mathcal{O}}^{\#}$ makes $B \otimes_k^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ an object of $\text{FrAlg}_{\mathbb{P}^1, \mathbb{R}}$.

The forgetful functor $\text{FrAlg}_{\mathbb{P}^1, \mathbb{R}} \rightarrow \text{FrAlg}_{\mathbb{P}^1, \mathbb{C}}$ has a right adjoint, given by $\mathcal{F} \mapsto \mathcal{F} \oplus \tau^{-1} \mathcal{F}$, with the involution $\tau^{\#}$ given by swapping summands and the $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ -structure on $\tau^{-1} \mathcal{F}$ defined using $\tau_{\mathcal{O}}$.

6.2 The twistor functors

Definition 6.2 Define the groupoid-valued functor $\mathcal{R}_X^{\mathbb{T},\mathbb{C}}$ on $\text{FrAlg}_{\mathbb{P}^1,\mathbb{C}}$ by letting $\mathcal{R}_X^{\mathbb{T},\mathbb{C}}(\mathcal{B})$ consist of pairs (\mathcal{T}, \tilde{D}) for $\mathcal{A}_X^0(\text{pr}_2^{-1}\mathcal{B}^\times)$ -torsors \mathcal{T} on $X \times \mathbb{P}^1(\mathbb{C})$ with $x^*\mathcal{T}$ trivial as a \mathcal{B}^\times -torsor on $\mathbb{P}^1(\mathbb{C})$ and flat $ud+vd^c$ -connections

$$\tilde{D}: \mathcal{T} \rightarrow \mathcal{A}_X^1 \otimes_{\mathcal{A}_X^0} \text{ad } \mathcal{T}(1).$$

Here u and v are the basis of $\Gamma(\mathbb{P}^1_{\mathbb{R}}, \mathcal{O}(1))$ given by the coordinates u and v of C^* and the canonical map $C^* \rightarrow \mathbb{P}^1$.

Define the set-valued functor $\mathbf{R}_{X,x}^{\mathbb{T},\mathbb{C}}$ on $\text{FrAlg}_{\mathbb{P}^1,\mathbb{C}}$ by letting $\mathbf{R}_{X,x}^{\mathbb{T},\mathbb{C}}(\mathcal{B})$ be the groupoid of triples $(\mathcal{T}, \tilde{D}, f)$, with $(\mathcal{T}, \tilde{D}) \in \mathcal{R}_X^{\mathbb{T},\mathbb{C}}(\mathcal{B})$ and framing $f \in \Gamma(\mathbb{P}^1(\mathbb{C}), x^*\mathcal{T})$.

Definition 6.3 Define the groupoid-valued functor $\mathcal{R}_X^{\mathbb{T}}$ on $\text{FrAlg}_{\mathbb{P}^1,\mathbb{R}}$ by letting $\mathcal{R}_X^{\mathbb{T}}(\mathcal{B})$ consist of triples $(\mathcal{T}, \tilde{D}, \tau_{\mathcal{T}}^\#)$ for $(\mathcal{T}, \tilde{D}) \in \mathcal{R}_X^{\mathbb{T},\mathbb{C}}(\mathcal{B})$ and isomorphism $\tau_{\mathcal{T}}^\#: (\text{id}_X \times \tau)^{-1}\mathcal{T} \rightarrow \mathcal{T}$ satisfying the following conditions. The isomorphism

$$(\tau_{\mathcal{B}}^\#, \tau_{\mathcal{T}}^\#): \mathcal{A}_X^1(\text{pr}_2^{-1}\mathcal{B}^\times) \times_{\tau_{\mathcal{B}}^\#, \mathcal{A}_X^1(\text{pr}_2^{-1}\tau^{-1}\mathcal{B}^\times)} (\text{id}_X \times \tau)^{-1}\mathcal{T} \rightarrow \mathcal{T}$$

must be a morphism of $\mathcal{A}_X^1(\text{pr}_2^{-1}\mathcal{B}^\times)$ -torsors, and the diagram

$$\begin{array}{ccc} \tau^{-1}\mathcal{T} & \xrightarrow{\tau^{-1}\tilde{D}} & \mathcal{A}_X^1 \otimes_{\mathcal{A}_X^0} \tau^{-1}\mathcal{T}(1) \\ \tau_{\mathcal{T}}^\# \downarrow & & \downarrow \tau_{\mathcal{T}}^\# \\ \mathcal{T} & \xrightarrow{\tilde{D}} & \mathcal{A}_X^1 \otimes_{\mathcal{A}_X^0} \mathcal{T}(1) \end{array}$$

must commute.

Define the functor $\mathbf{R}_{X,x}^{\mathbb{T}}$ on $\text{FrAlg}_{\mathbb{P}^1,\mathbb{R}}$ by letting $\mathbf{R}_{X,x}^{\mathbb{T}}(\mathcal{B})$ be the groupoid of quadruples $(\mathcal{T}, \tilde{D}, \tau_{\mathcal{T}}^\#, f)$ for $(\mathcal{T}, \tilde{D}, \tau_{\mathcal{T}}^\#)$ in $\mathcal{R}_X^{\mathbb{T}}(\mathcal{B})$ and

$$f \in \Gamma(\mathbb{P}^1(\mathbb{C}), x^*\mathcal{T})^{\tau_{\mathcal{T}}^\#}$$

a framing.

Remark 6.4 The groupoids $\mathbf{R}_{X,x}^{\mathbb{T},\mathbb{C}}(\mathcal{B}), \mathbf{R}_{X,x}^{\mathbb{T}}(\mathcal{B})$ are equivalent to discrete groupoids, so we will regard them as set-valued functors (given by isomorphism classes of objects). Also, $\mathcal{R}_X^{\mathbb{T},\mathbb{C}}(\mathcal{B})$ and $\mathcal{R}_X^{\mathbb{T}}(\mathcal{B})$ are equivalent to the groupoid quotients $[\mathbf{R}_{X,x}^{\mathbb{T},\mathbb{C}}(\mathcal{B})/\Gamma(\mathbb{P}^1(\mathbb{C}), \mathcal{B}^\times)]$ and $[\mathbf{R}_{X,x}^{\mathbb{T}}(\mathcal{B})/\Gamma(\mathbb{P}^1(\mathbb{C}), \mathcal{B}^\times)^{\tau_{\mathcal{T}}^\#}]$, respectively, with the action given by changing the framing.

The following is straightforward:

Lemma 6.5 *The functors $R_{X,x}^{\mathbb{T},\mathbb{C}}$ and $\mathcal{R}_X^{\mathbb{T},\mathbb{C}}$ can be recovered from $R_{X,x}^{\mathbb{T}}$ and $\mathcal{R}_X^{\mathbb{T}}$, respectively, via isomorphisms*

$$R_{X,x}^{\mathbb{T},\mathbb{C}}(\mathcal{B}) \cong R_{X,x}^{\mathbb{T}}(\mathcal{B} \oplus \tau^{-1}\mathcal{B}), \quad \mathcal{R}_X^{\mathbb{T},\mathbb{C}}(\mathcal{B}) \cong \mathcal{R}_X^{\mathbb{T}}(\mathcal{B} \oplus \tau^{-1}\mathcal{B}).$$

Note that the proof of Lemma 2.5 carries over to give the canonical comultiplication

$$R_{X,x}^{\mathbb{T}}(\mathcal{B}_1) \times R_{X,x}^{\mathbb{T}}(\mathcal{B}_2) \rightarrow R_{X,x}^{\mathbb{T}}(\mathcal{B}_1 \otimes^{\pi} \mathcal{B}_2).$$

6.3 Universality and σ -invariant sections

Proposition 6.6 *The functors $\mathcal{R}_X^{\text{dR}}$ and $\mathcal{R}_X^{\text{Dol}}$ can be recovered from $\mathcal{R}_X^{\mathbb{T}}$ and $\mathcal{R}_X^{\mathbb{T},\mathbb{C}}$, respectively. Likewise, $R_{X,x}^{\text{dR}}$ and $R_{X,x}^{\text{Dol}}$ can be recovered from $R_{X,x}^{\mathbb{T}}$ and $R_{X,x}^{\mathbb{T},\mathbb{C}}$.*

Proof Given a point $p \in \mathbb{P}^1(\mathbb{C})$ and a complex m -convex Fréchet algebra B , we may regard the skyscraper sheaf p_*B as an object of $\text{FrAlg}_{\mathbb{P}^1,\mathbb{C}}$. For a real m -convex Fréchet algebra B and $p \in \mathbb{P}^1(\mathbb{R})$, we may regard $p_*B \otimes \mathbb{C}$ as an object of $\text{FrAlg}_{\mathbb{P}^1,\mathbb{R}}$, with $\tau^{\#}$ given by complex conjugation.

Now, just observe that on pulling back to $(1 : 0) \in \mathbb{P}^1(\mathbb{R})$, the differential $ud + vd^c$ is just d . At $(1 : -i) \in \mathbb{P}^1(\mathbb{R})$, we have $ud + vd^c = d - id^c = 2\partial$. Uncoiling the definitions, this gives

$$R_{X,x}^{\text{dR}}(B) = R_{X,x}^{\mathbb{T}}((1 : 0)_*B), \quad R_{X,x}^{\text{Dol}}(B) = R_{X,x}^{\mathbb{T},\mathbb{C}}((1 : -i)_*B),$$

and similarly for \mathcal{R} . □

Lemma 6.7 *For a real m -convex Fréchet algebra B , the groupoid $\mathcal{R}_X^{\mathbb{T}}(B \otimes_{\mathbb{R}}^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}})$ is equivalent to the groupoid of triples (\mathcal{P}, D, E) , where \mathcal{P} is an $\mathcal{A}_X^0(B^{\times})$ -torsor on X , the maps $D, E: \mathcal{P} \rightarrow \mathcal{A}_X^1 \otimes_{\mathcal{A}_X^0} \mathcal{P}$ are flat d - and d^c -connections, respectively, and $DE + ED = 0$.*

Proof Take an object $(\mathcal{S}, \tilde{D}) \in \mathcal{R}_X^{\mathbb{T}}(B \otimes_{\mathbb{R}} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}})$. Triviality of the $(B \otimes_{\mathbb{R}}^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}})^{\times}$ -torsors $x^*\mathcal{S}$ for $x \in X$ ensures that \mathcal{S} must be of the form $\mathcal{A}_X^0(B \otimes_{\mathbb{R}}^{\pi} B)^{\times} \times_{\mathcal{A}_X^0(B^{\times})} \mathcal{P}$ for some \mathcal{P} as above.

Now, since $\Gamma(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(1))^{\tau^{\#}} = \mathbb{R}u \oplus \mathbb{R}v$, the connection \tilde{D} can be regarded as a $ud + vd^c$ -connection

$$\tilde{D}: \mathcal{P} \rightarrow (\mathcal{A}_X^1 \otimes_{\mathcal{A}_X^0} \mathcal{P}) \otimes_{\mathbb{R}} (\mathbb{R}u \oplus \mathbb{R}v),$$

which we write as $uD + vE$.

Flatness of \tilde{D} is now a statement about $\Gamma(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(2))^{\tau^{\#}} = \mathbb{R}u^2 \oplus \mathbb{R}uv \oplus \mathbb{R}v^2$, with the u^2 and v^2 terms giving flatness of D and E and the uv term giving the condition $[D, E] = 0$. □

Definition 6.8 Define the involution $\tau\sigma_{\mathbb{P}^1}$ of the polarised scheme $(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ to be the map induced by the action of $i \in S(\mathbb{R})$ on C^* from Definition 5.1. In particular, $\tau\sigma_{\mathbb{P}^1}(u : v) = (v : -u)$.

Remark 6.9 On [27, page 12], the coordinate system $(u + iv : u - iv)$ on $\mathbb{P}^1(\mathbb{C})$ is used, and antiholomorphic involutions σ and τ are defined. In our coordinates, these become $\sigma(u : v) = (\bar{v}, -\bar{u})$ and $\tau(u : v) = (\bar{u}, \bar{v})$. This justifies the notation $\tau\sigma$ used above. Also note that the $\mathbb{G}_{m,\mathbb{C}}$ -action on $\mathbb{P}^1_{\mathbb{C}}$ given in [27, page 4] is just the complex form of our circle action on $\mathbb{P}^1_{\mathbb{R}}$ from Sections 5.3 and 5.4.3.

Definition 6.10 Given a real Banach algebra B , define the involution $\tau\sigma'$ of the space $\mathcal{R}_X^{\mathbb{T}}(B \otimes_{\mathbb{R}}^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}})$ by sending the pair (\mathcal{S}, \tilde{D}) to $(\tau\sigma_{\mathbb{P}^1}^{-1}\mathcal{S}, J\tau\sigma_{\mathbb{P}^1}^{-1}\tilde{D})$. Note that this is well-defined because $\tau\sigma_{\mathbb{P}^1}^{-1}\tilde{D}$ is a $\tau\sigma_{\mathbb{P}^1}(ud + vd^c) = (vd - ud^c)$ -connection, and $Jd = d^c$ and $Jd^c = -d$, so $J\tau\sigma_{\mathbb{P}^1}^{-1}\tilde{D}$ is a $ud + vd^c$ -connection.

Definition 6.11 Given a C^* -algebra B , define the Cartan involution C of B^{\times} to be given by $C(g) = (g^{-1})^*$. Note that this induces a Lie algebra involution $\text{ad } C : b \mapsto -b^*$ on the tangent space B of B^{\times} .

If $B = A \otimes \mathbb{C}$ for a real C^* -algebra A , we write τ for complex conjugation, so $C\tau$ is the involution $C\tau(g) = (\bar{g}^{-1})^*$. Note that $\text{ad } C\tau$ is the \mathbb{C} -linear extension of $\text{ad } C$ on A .

Since $\mathcal{R}_X^{\mathbb{T}}(\mathcal{B})$ only depends on the group of units \mathcal{B}^{\times} of \mathcal{B} and its tangent Lie algebra \mathcal{B} , the Cartan involution induces an involution $C\tau$ of $\mathcal{R}_X^{\mathbb{T}}(B \otimes_{\mathbb{R}}^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}})$.

Definition 6.12 Given a real Banach algebra B , define the involution σ of the space $\mathcal{R}_X^{\mathbb{T}}(B \otimes_{\mathbb{R}}^{\pi} \mathcal{O}_{\mathbb{P}^1}^{\text{hol}})$ by $\sigma := (C\tau)(\tau\sigma')$.

Proposition 6.13 For a real C^* -algebra B , there is a canonical isomorphism

$$\mathbf{R}_{X,x}^{\mathbb{T}}(\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(B))^{\sigma} \cong \mathbf{R}_{X,x}^J(B).$$

Proof Since $\mathbf{R}_{X,x}^{\mathbb{T}}$ is the groupoid fibre of $\mathcal{R}_X^{\mathbb{T}} \rightarrow \mathcal{R}_{\{x\}}^{\mathbb{T}}$ over the trivial torsor, Lemma 6.7 shows that an object $(\mathcal{S}, \tilde{D}, f)$ of $\mathbf{R}_{X,x}^{\mathbb{T}}(\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(B))$ is a quadruple (\mathcal{P}, D, E, f) for (\mathcal{P}, D, E) as in that lemma and f our framing. We therefore begin by describing the σ -action on such data. For (\mathcal{P}, D, E, f) to be σ -invariant, we must have an isomorphism $\alpha : (\mathcal{P}, D, E, f) \rightarrow \sigma(\mathcal{P}, D, E, f)$.

The torsor \mathcal{P} maps under σ to $C(\mathcal{P})$, with $f \in x^*\mathcal{P}$ mapping to $C(f)$ (since σ' and τ affect neither). The isomorphism α then gives $\alpha : \mathcal{P} \rightarrow C(\mathcal{P})$ such that

$\alpha(f) = C(f) \in x^*C(\mathcal{P})$. Let $U(\mathcal{P}) \subset \mathcal{P}$ consist of sections q with $\alpha(q) = C(q)$. This is nonempty (since its fibre at x contains f), so it must be an $\mathcal{A}_X^0(U(B))$ -torsor, noting that $U(B)$ is the group of C -invariants in B^\times . Moreover, $\mathcal{P} = \mathcal{A}_X^0(B^\times) \times_{\mathcal{A}_X^0(U(B))} U(\mathcal{P})$.

Meanwhile, $\tau\sigma'(uD + vE) = vJD - uJE$, so $\tau\sigma'(D, E) = (-JE, JD)$. Thus the isomorphism α gives

$$D|_{\mathcal{Q}} = -JCE, \quad E|_{\mathcal{Q}} = JCD.$$

In other words, $E = D^c$ and $E^c = -D$ (which are equivalent conditions).

Thus $\mathbf{R}_{X,x}^{\mathbb{T}}(\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}(B))^\sigma$ is equivalent to the groupoid of triples $(U(\mathcal{P}), D, f)$, with D flat and $[D, D^c] = 0$. □

Remark 6.14 When $B = \text{Mat}_n(\mathbb{C})$, this shows that framed pluriharmonic local systems correspond to framed σ -invariant sections of the twistor functor. Without the framings, this will not be true in general, since a σ -invariant section of the coarse moduli space will give a nondegenerate bilinear form which need not be positive definite. Note that for $U \subset \mathbb{P}^1$, the set of isomorphism classes of $\mathcal{R}_X^{\mathbb{T},\mathbb{C}}(\text{Mat}_n(\mathcal{O}_U^{\text{hol}}))$ is the set of sections over U of the twistor space $TW \rightarrow \mathbb{P}^1$ of [26, Section 3].

Remark 6.15 Although we have seen that $\mathcal{R}_X^{\mathbb{T}}$ together with its comultiplication encodes all the available information about twistor structures on moduli spaces of local systems, it does not carry information about higher homotopy and cohomology groups. There is, however, a natural extension of $\mathcal{R}_X^{\mathbb{T}}$ to differential graded m -convex Fréchet algebras, by analogy with [19; 20]. This would involve taking \tilde{D} to be a hyperconnection $\tilde{D}: \mathcal{T}_0 \rightarrow \prod_n \mathcal{A}_X^{n+1} \otimes_{\mathcal{A}_X^0} \text{ad } \mathcal{T}_n(n+1)$. The structures of Section 4 can all be recovered from this functor.

6.4 Topological twistor representation spaces

In this section, we will show that, by considering continuous homomorphisms rather than $*$ -homomorphisms, we can describe the entire semisimple locus of the twistor family from $(E_{X,x}^J)_{\mathbb{P}N}$, rather than just the σ -equivariant sections.

Given a point $(a : b) \in \mathbb{P}^1(\mathbb{C})$ and a complex Banach algebra B , we can generalise the construction of Proposition 6.6 and consider the set $\mathbf{R}_{X,x}^{\mathbb{T},\mathbb{C}}((a : b)_*B)$. This consists of torsors with flat $ad + bd^c$ -connections.

Definition 6.16 Define $T_{X,x,n} := \coprod_{(a:b) \in \mathbb{P}^1(\mathbb{C})} \mathbf{R}_{X,x}^{\mathbb{T},\mathbb{C}}((a : b)_*\text{Mat}_n(\mathbb{C}))$.

Note that $T_{X,x,n}$ inherits a $\text{Gal}(\mathbb{C}/\mathbb{R})$ -action from $R_{X,x}^{\mathbb{T},\mathbb{C}}$. We can also give $T_{X,x,n}$ a complex analytic structure, by saying that a map $f: U \rightarrow T_{X,x}(B)$ from an analytic space U consists of an analytic map $f: U \rightarrow \mathbb{P}^1(\mathbb{C})$ together with an element of $R_{X,x}^{\mathbb{T},\mathbb{C}}(f_*(\text{Mat}_n \mathcal{O}_U))$. We will now investigate the underlying topological structure.

Remark 6.17 The adjoint action of $\text{GL}_n(\mathbb{C})$ on $T_{X,x,n}$ is continuous and, indeed, compatible with the complex analytic structure. This allows us to consider the coarse quotient $T_{X,x,n} // \text{GL}_n(\mathbb{C})$, which is the Hausdorff completion of the topological quotient, equipped with a natural complex analytic structure over $\mathbb{P}^1(\mathbb{C})$. A straightforward calculation shows that this coarse moduli space is precisely the Deligne–Hitchin twistor space, as constructed in [14] and described in [28, Section 3].

Now, Proposition 6.13 induces a map $\pi_{\mathbb{T}}: R_{X,x}^J(\text{Mat}_n(\mathbb{C})) \times \mathbb{P}^1(\mathbb{C}) \rightarrow T_{X,x,n}$. For an explicit characterisation, note that an $ad+bd^c$ -connection \tilde{D} lies in $R_{X,x}^J(\text{Mat}_n(\mathbb{C}))$ if and only if

$$[\tilde{D}, JC \tilde{D}] = 0,$$

and that $JC \tilde{D}$ is a $-\bar{b}d + \bar{a}d^c = \sigma_{\mathbb{P}^1}^{-1}(ad+bd^c)$ -connection.

Definition 6.18 Given a flat $ad+bd^c$ -connection \tilde{D} on a finite-dimensional \mathcal{C}^∞ vector bundle \mathcal{V} on X for $(a : b) \in \mathbb{P}^1(\mathbb{C}) - \{\pm i\}$, we say that (\mathcal{V}, \tilde{D}) is *semisimple* if the local system $\ker \tilde{D}$ is so.

Definition 6.19 Define $T_{X,x,n}^{\text{st}} \subset T_{X,x,n}$ by requiring that the fibre over any point of $\mathbb{P}^1(\mathbb{C}) - \{\pm i\}$ consist of the semisimple objects, the fibre over i be $R_{X,x}^{\text{Dol,st}}(\text{Mat}_n(\mathbb{C}))$ (Section 3.3), and the fibre over $-i$ be its conjugate.

We give $T_{X,x,n}^{\text{st}}$ the subspace topology, so a map $K \rightarrow T_{X,x,n}^{\text{st}}$ is continuous if the projection $f: K \rightarrow \mathbb{P}^1(\mathbb{C})$ is so, and the map lifts to an element of $R_{X,x}^{\mathbb{T},\mathbb{C}}(f_*C(K, \text{Mat}_n(\mathbb{C})))$.

Theorem 6.20 For any positive integer n , there is a natural homeomorphism $\pi_{\mathbb{T},\text{st}}$ over $\mathbb{P}^1(\mathbb{C})$ between the space $\text{Hom}_{\text{pro}(\text{BanAlg})}(E_{X,x}^J, \text{Mat}_n(\mathbb{C})) \times \mathbb{P}^1(\mathbb{C})$ with the topology of pointwise convergence and the space $T_{X,x,n}^{\text{st}}$.

Proof The homeomorphism is given on the fibre over $(a : b) \in \mathbb{P}^1(\mathbb{C})$ by

$$\pi_{\mathbb{T},\text{st}}(U(\mathcal{P}), D, f) = (\mathcal{P}, aD + bD^c, f).$$

The proofs of Theorems 3.6 and 3.23 adapt to show that $\pi_{\mathbb{T},\text{st}}$ induces homeomorphisms on fibres over $\mathbb{P}^1(\mathbb{C})$ and, in particular, is an isomorphism on points. The same arguments also show that $\pi_{\mathbb{T},\text{st}}$ and $\pi_{\text{dR}} \circ \pi_{\mathbb{T},\text{st}}^{-1}: \pi_{\mathbb{T},\text{st}} \rightarrow R_{X,x}^{\text{dR,ss}}(\text{Mat}_n(\mathbb{C}))$ are continuous, so the result follows from Theorem 3.6. \square

Definition 6.21 Let $\text{FDT}_{X,x}^{\text{st}}$ be the category of pairs $(V, p, (a : b))$ for $V \in \text{FDVect}$ and $(p, (a : b)) \in \mathbf{T}_{X,x,n}^{\text{st}}$ where $n = \dim V$. Morphisms are defined by adapting the formulae of Definition 3.7. Write $\eta_x^{\mathbb{T},\text{st}}: \text{FDT}_{X,x}^{\text{st}} \rightarrow \text{FDVect} \times \mathbb{P}^1(\mathbb{C})$ for the fibre functors $(V, p, (a : b)) \mapsto (V, (a : b))$.

Proposition 6.22 The C^* -algebra $(E_{X,x}^J)_{\text{PN}} \widehat{\otimes}_{\mathbb{R}} C(\mathbb{P}^1(\mathbb{C}), \mathbb{C})$ is isomorphic, via a $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant isomorphism, to the ring of continuous endomorphisms of $\eta_x^{\mathbb{T},\text{st}}$.

Proof The proof of Proposition 3.8 carries over, replacing Theorem 3.6 with Theorem 6.20 and Lemma 1.28 with Lemma 1.30. □

Definition 6.23 Given a k -normal real C^* -algebra B over $C(\mathbb{P}^1(\mathbb{C}), \mathbb{C})^{\text{Gal}(\mathbb{C}/\mathbb{R})}$, we may regard B as an $\mathcal{O}_{\mathbb{P}^1}^{\text{hol}}$ -algebra via the inclusion of holomorphic functions in continuous functions. Then define $\mathbf{R}_{X,x}^{\mathbb{T},\text{st}}(B) \subset \mathbf{R}_{X,x}^{\mathbb{T}}(B)$ to be the subspace consisting of those p for which

$$(\psi(p), (a : b)) \in \mathbf{T}_{X,x,k}^{\text{st}}$$

for all $(a : b) \in \mathbb{P}^1(\mathbb{C})$ and $\psi: B \rightarrow \text{Mat}_k(\mathbb{C})$ with $\psi|_{C(\mathbb{P}^1(\mathbb{C}), \mathbb{C})^{\text{Gal}(\mathbb{C}/\mathbb{R})}} = \text{ev}_{(a:b)} \text{id}$.

Corollary 6.24 For any k -normal real C^* -algebra B over $C(\mathbb{P}^1(\mathbb{C}), \mathbb{C})^{\text{Gal}(\mathbb{C}/\mathbb{R})}$, there is a natural isomorphism between $\mathbf{R}_{X,x}^{\mathbb{T},\text{st}}(B)^{\text{Gal}(\mathbb{C}/\mathbb{R})}$ and the set of continuous algebra homomorphisms $E_{X,x}^J \rightarrow B$.

Proof The proof of Corollary 3.11 carries over, replacing Proposition 3.8 with Proposition 6.22. □

References

- [1] C A Akemann, F W Shultz, *Perfect C^* -algebras*, Mem. Amer. Math. Soc. 326, Amer. Math. Soc., Providence, RI (1985) MR
- [2] J Amorós, M Burger, K Corlette, D Kotschick, D Toledo, *Fundamental groups of compact Kähler manifolds*, Mathematical Surveys and Monographs 44, Amer. Math. Soc., Providence, RI (1996) MR
- [3] K Bichteler, *A generalization to the non-separable case of Takesaki’s duality theorem for C^* -algebras*, Invent. Math. 9 (1969) 89–98 MR
- [4] J W Bunce, J A Deddens, *C^* -algebras with Hausdorff spectrum*, Trans. Amer. Math. Soc. 212 (1975) 199–217 MR
- [5] K Corlette, *Flat G -bundles with canonical metrics*, J. Differential Geom. 28 (1988) 361–382 MR

- [6] **P Deligne**, *Théorie de Hodge, II*, Inst. Hautes Études Sci. Publ. Math. 40 (1971) 5–57 MR
- [7] **P Deligne, J S Milne, A Ogus, K-y Shih**, *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics 900, Springer, Berlin (1982) MR
- [8] **J Dodziuk**, *De Rham–Hodge theory for L^2 -cohomology of infinite coverings*, Topology 16 (1977) 157–165 MR
- [9] **L T Gardner**, *On isomorphisms of C^* -algebras*, Amer. J. Math. 87 (1965) 384–396 MR
- [10] **W M Goldman, J J Millson**, *The deformation theory of representations of fundamental groups of compact Kähler manifolds*, Inst. Hautes Études Sci. Publ. Math. 67 (1988) 43–96 MR
- [11] **K R Goodearl**, *Notes on real and complex C^* -algebras*, Shiva Mathematics Series 5, Shiva, Nantwich (1982) MR
- [12] **P Griffiths, J Harris**, *Principles of algebraic geometry*, Interscience, New York (1978) MR
- [13] **A Grothendieck**, *Technique de descente et théorèmes d’existence en géométrie algébrique, II: Le théorème d’existence en théorie formelle des modules*, from “Séminaire Bourbaki 1959/1960”, W A Benjamin, Amsterdam (1966) Exposé 195 MR Reprinted in Séminaire Bourbaki 5 (1958–1960) 369–390, Soc. Math. France, Paris, 1995
- [14] **N J Hitchin**, *The self-duality equations on a Riemann surface*, Proc. London Math. Soc. 55 (1987) 59–126 MR
- [15] **S Kobayashi, K Nomizu**, *Foundations of differential geometry, I*, Interscience, New York (1963) MR
- [16] **V I Paulsen**, *Completely bounded homomorphisms of operator algebras*, Proc. Amer. Math. Soc. 92 (1984) 225–228 MR
- [17] **C Pearcy**, *A complete set of unitary invariants for operators generating finite W^* -algebras of type I*, Pacific J. Math. 12 (1962) 1405–1416 MR
- [18] **N C Phillips**, *Inverse limits of C^* -algebras*, J. Operator Theory 19 (1988) 159–195 MR
- [19] **J P Pridham**, *Pro-algebraic homotopy types*, Proc. Lond. Math. Soc. 97 (2008) 273–338 MR
- [20] **J P Pridham**, *Real non-abelian mixed Hodge structures for quasi-projective varieties: formality and splitting*, Mem. Amer. Math. Soc. 1150, Amer. Math. Soc., Providence, RI (2016) MR
- [21] **G A Reid**, *Epimorphisms and surjectivity*, Invent. Math. 9 (1969/1970) 295–307 MR
- [22] **I E Segal**, *Two-sided ideals in operator algebras*, Ann. of Math. 50 (1949) 856–865 MR

- [23] **C T Simpson**, *Higgs bundles and local systems*, Inst. Hautes Études Sci. Publ. Math. 75 (1992) 5–95 MR
- [24] **C T Simpson**, *Moduli of representations of the fundamental group of a smooth projective variety, I*, Inst. Hautes Études Sci. Publ. Math. 79 (1994) 47–129 MR
- [25] **C T Simpson**, *Moduli of representations of the fundamental group of a smooth projective variety, II*, Inst. Hautes Études Sci. Publ. Math. 80 (1994) 5–79 MR
- [26] **C Simpson**, *The Hodge filtration on nonabelian cohomology*, from “Algebraic geometry” (J Kollár, R Lazarsfeld, D R Morrison, editors), Proc. Sympos. Pure Math. 62, Amer. Math. Soc., Providence, RI (1997) 217–281 MR
- [27] **C Simpson**, *Mixed twistor structures*, preprint (1997) arXiv
- [28] **C Simpson**, *A weight two phenomenon for the moduli of rank one local systems on open varieties*, from “From Hodge theory to integrability and TQFT tt*-geometry” (R Y Donagi, K Wendland, editors), Proc. Sympos. Pure Math. 78, Amer. Math. Soc., Providence, RI (2008) 175–214 MR
- [29] **D Sullivan**, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. 47 (1977) 269–331 MR
- [30] **M Takesaki**, *Theory of operator algebras, I*, Springer, New York (1979) MR Reprinted in Encyclopaedia of Mathematical Sciences 124
- [31] **D P Williams**, *Crossed products of C^* -algebras*, Mathematical Surveys and Monographs 134, Amer. Math. Soc., Providence, RI (2007) MR

*School of Mathematics and Maxwell Institute, The University of Edinburgh
James Clerk Maxwell Building, The King’s Buildings Mayfield Road
Edinburgh, EH9 3FD, United Kingdom*

`j.pridham@ed.ac.uk`

Proposed: Richard Thomas
Seconded: Peter Teichner, Jim Bryan

Received: 7 July 2014
Revised: 4 January 2016