



Geometry & Topology

Volume 27 (2023)

On self-shrinkers of medium entropy in \mathbb{R}^4

ALEXANDER MRAMOR



On self-shrinkers of medium entropy in \mathbb{R}^4

ALEXANDER MRAMOR

We study smooth asymptotically conical self-shrinkers in \mathbb{R}^4 with Colding–Minicozzi entropy bounded above by Λ_1 .

53A10, 53E10

1 Introduction

Self-shrinkers are basic singularity models for the mean curvature flow, and, in the noncompact case, nongeneric ones (generic ones being generalized round cylinders $S^k(\sqrt{2k}) \times \mathbb{R}^{n-k}$) are expected to often be asymptotically conical. Our purpose is to understand the topology of smooth self-shrinkers $M^3 \subset \mathbb{R}^4$ with Colding–Minicozzi entropy $\lambda(M)$, discussed in [Section 2](#), bounded above by Λ_1 , the entropy of the round circle.

Our main result is, in part, inspired by arguments of Bernstein and L Wang [3], Hershkovits and White [21], Ilmanen and White [24], Mramor [28], Mramor and S Wang [29] and White [39]. The basic idea is that by considering renormalized mean curvature flows out of (appropriate perturbations of) asymptotically conical self-shrinkers, we may use the entropy assumption to constrain which types of singularities may occur. This has strong implications for how topology may change under the flow. This is useful because topology can, in a sense, be used to “trap” the flow. On the other hand, the flow must clear out; these two principles can then be combined to constrain the topology of the self-shrinker in question.

Theorem 1.1 *Suppose $M^3 \subset \mathbb{R}^4$ is a smooth 2-sided asymptotically conical self-shrinker with entropy less than Λ_1 and k ends. Then it is diffeomorphic to S^3 with k 3-balls removed and replaced with k copies of $S^2 \times \mathbb{R}_+$ attached along their respective boundaries. If $k = 1$ then $M \simeq \mathbb{R}^3$, and in particular this is the case when $\lambda(M) \leq \Lambda_2$.*

This extends to the noncompact case joint work of the author and S Wang [29] on compact self-shrinkers $M^n \subset \mathbb{R}^{n+1}$ when $n = 3$, where they showed that, for each $n \geq 3$, closed self-shrinkers M^n with $\Lambda(M) < \Lambda_{n-2}$ are diffeomorphic to S^n . This in turn extends a result of Colding, Ilmanen, Minicozzi, and White [9], which says closed self-shrinkers with entropy less than $\Lambda_{n-1} \lesssim \Lambda_{n-2}$ are diffeomorphic to S^n , hence weakening the assumed entropy bound. In a similar manner, the result above extends (in a weaker sense than the compact case) a result of Bernstein and L Wang [4] for noncompact shrinkers in \mathbb{R}^4 , where they showed (amongst other results; see Corollary 1.4 therein), for asymptotically conical self-shrinkers $M^3 \subset \mathbb{R}^4$ satisfying $\lambda(M) \leq \Lambda_2$, the stronger conclusion that they are diffeomorphic to \mathbb{R}^3 . Our argument does at least recover their statement, as discussed at the end of the proof. With the round cylinder in mind, our conclusion seems likely to be sharp in this sense, although it could be possible that a shrinker in \mathbb{R}^4 with this entropy bound has more than one end precisely when it is a cylinder.

In this dimension and under this entropy bound, we remark that generic mean curvature flow through neck-pinch singularities has been established by Chodosh, Choi, Mantoulidis and Schulze [7; 8], so for some applications of the flow (see for instance Daniels-Holgate [12]) the study of self-shrinkers in this regime is unnecessary. However, besides its intrinsic interest, this result might still be of use in understanding singularity along nongeneric flows, which could imaginably occur, for instance, in problems involving families of flows (although to the author's knowledge, potential fattening is a more serious concern). It also paints an explicit picture of how a perturbation of a nongeneric flow might only develop neck-pinch singularities, by some copies of the $S^2 \times \mathbb{R}_+$ in the statement above pinching off before, roughly speaking, the S^3 factor collapses to a point (as opposed to handles prematurely pinching off a more complicated model).

An important extra difficulty to consider in the noncompact case versus the closed case is that, a priori, nontrivial topology may be "lost" to spatial infinity under the flow without being properly understood. To illustrate this concern by an admittedly crude thought experiment, a hypothetical translator asymptotically modeled on $T^2 \times \mathbb{R}$ would never develop a singularity, and hence its topology would never be "encountered" as a high curvature region in the flow. In particular it seems, for $n = 3$, asymptotically conical self-shrinkers could a priori have a complicated link. Our first task, and really most of the work of this paper, will be to show that in fact the link is simple.

Theorem 1.2 *Suppose $M^3 \subset \mathbb{R}^4$ is a smooth 2-sided asymptotically conical self-shrinker with entropy less than Λ_1 . Then its link L is homeomorphic to a union of S^2 .*

As an indication of why one might argue this is reasonable, consider that, in general, the link L of a shrinker $M^n \subset \mathbb{R}^{n+1}$ is of dimension $n - 1$, so an entropy bound of Λ_{n-2} on M^n implies morally that its link is low-entropy; for a submanifold $N^k \subset \mathbb{R}^{k+1}$, we say N is low entropy if $\lambda(N) < \Lambda_{k-1}$ (hence the title of paper, since $\Lambda_{k-2} > \Lambda_{k-1}$, as discussed in the next section). These compact surfaces with this entropy bound, at least in low dimensions ($n = 2, 3$), are known to be spheres.

The dimension bound assumption is for topological reasons¹ that perhaps indicate a deficit in knowledge and finesse more than any true difficulty. Potentially providing a sliver of hope that this is the case, Ilmanen and White [24] showed lower bounds for the densities of area-minimizing cones in terms of the topology of their link in every dimension. The area-minimizing property there is employed by using a foliation near the cone by minimal surfaces, which are used as barriers in a mean curvature flow argument (at a high level our argument is similar to theirs). Since cones are noncompact, this is clearly the same sort of result as ours.

For instance, one might hope to directly modify our argument in the next higher dimension ($n = 4$) because simply connected 3-manifolds are spherical by the resolution of the 3D Poincaré conjecture by Perelman [30; 31; 32]—we use the corresponding (much easier) fact for surfaces below to classify the link. As an example of why this simple criterion alone doesn't seem to immediately lead to a proof of the corresponding statement for $n = 4$, a potential issue (to the author's understanding) in this dimension is that the link could be a nontrivial homology sphere—below we use that nonspherical oriented surfaces have nontrivial homology in a seemingly essential way. For higher dimensions, of course, there are higher-dimensional versions of the Poincaré conjecture, as verified by Freedman [14] and Smale [33]; this naturally seems even more complicated, for a number of reasons, than the $n = 4$ case just discussed.

Acknowledgements The author is supported by an AMS–Simons travel grant and thanks them for their generosity, as well as the referee for their careful reading and critique.

2 Preliminaries

Let $X: M \rightarrow N^{n+1}$ be an embedding of M realizing it as a smooth closed hypersurface of N , which by abuse of notation we also refer to as M . Then the mean curvature flow

¹In the argument we use the classification of surfaces, Alexander's theorem, and Dehn's lemma, which are dimension-dependent. In a probably less essential way, the 3D Poincaré conjecture is also used.

M_t of M is given by (the image of) $X: M \times [0, T) \rightarrow N^{n+1}$ satisfying the following, where ν is the outward normal:

$$(2-1) \quad \frac{dX}{dt} = \vec{H} = -H\nu, \quad X(M, 0) = X(M).$$

By the comparison principle, singularities occur often, which makes their study important. To study these singularities, one may parabolically rescale about the developing high curvature region to obtain an ancient flow defined for times $(-\infty, T]$; when the basepoint is fixed, this is called a *tangent flow blowup* which will be modeled on self-shrinkers. By Huisken monotonicity [22] these are surfaces equivalently defined as

- (1) $M^n \subset \mathbb{R}^{n+1}$ satisfying $H - \frac{1}{2}\langle X, \nu \rangle = 0$, where X is the position vector;
- (2) minimal surfaces in the Gaussian metric $G_{ij} = e^{-|x|^2/(2n)}\delta_{ij}$; or
- (3) surfaces M which give rise to ancient flows M_t that move by dilations by setting $M_t = \sqrt{-t}M$.

(These notions all make sense at least when the shrinker is smooth, but some definitions apply in the varifold sense as well.) As is well known, the second variation formula for area shows there are no stable minimal surfaces in Ricci positive manifolds; see, for instance, Chapter 1 of [10]. This turns out to also be true for minimal surfaces of polynomial volume growth in \mathbb{R}^n endowed with the Gaussian metric as discussed in [11]. To see why this is so, the Jacobi operator for the Gaussian metric is given by

$$(2-2) \quad L = \Delta + |A|^2 - \frac{1}{2}\langle X, \nabla(\cdot) \rangle + \frac{1}{2}.$$

The extra $\frac{1}{2}$ term is essentially the reason such self-shrinkers are unstable in the Gaussian metric. For example, owing to the constant term it's clear in the compact case that one could simply plug in the function "1" to get a variation with $Lu > 0$ which doesn't change sign, implying that the first eigenvalue is negative. In fact, every properly embedded shrinker has polynomial volume growth by Q Ding and YL Xin:

Theorem 2.1 [13, Theorem 1.1] *Any complete noncompact properly immersed self-shrinker M^n in \mathbb{R}^{n+m} has Euclidean volume growth at most.*

We combine these facts below to conclude that the self-shrinker we find in some cases must in fact be unstable.

The mean curvature flow is best understood in the mean convex case because it turns out, under quite weak assumptions, the only possible shrinkers are generalized cylinders $S^k \times \mathbb{R}^{n-k}$. This is especially so for 2-convex surfaces ($\lambda_1 + \lambda_2 > 0$), and a surgery

theory with this convexity condition similar to the Ricci flow with surgery has been carried out. For the mean curvature flow with surgery, one finds, for a 2-convex surface M , curvature scales $H_{\text{th}} < H_{\text{neck}} < H_{\text{trig}}$ such that when $H = H_{\text{trig}}$ at some point p and time t the flow is stopped, and suitable points where $H \sim H_{\text{neck}}$ are found to do surgery where “necks” (at these points the surface will be approximately cylindrical) are cut and caps are glued in. The high curvature regions are topologically identified as S^n or $S^{n-1} \times S^1$ and discarded, and the low curvature regions will have curvature bounded on the order of H_{th} . The flow is then restarted and the process repeated.

It was initially established for compact 2-convex hypersurfaces in \mathbb{R}^{n+1} where $n \geq 3$ by Huisken and Sinestrari in [23], and their approach was later extended to the case $n = 2$ by Brendle and Huisken in [5], where 2-convexity is mean convexity. A somewhat different approach covering all dimensions simultaneously was given later by Haslhofer and Kleiner in [17] shortly afterwards. Haslhofer and Ketover then showed several years later in Section 8 of [15], en route to proving their main result, that the mean curvature flow with surgery can be applied to compact mean convex hypersurfaces in general ambient manifolds. Important to this article, the author with S Wang established it for (compact) mean convex hypersurfaces with entropy less than Λ_{n-2} in the sense of Colding and Minicozzi.

In [10], Colding and Minicozzi introduced their important notion of entropy, which is defined as the supremum of translated and rescaled Gaussian densities; indeed, consider a hypersurface $\Sigma^k \subset \mathbb{R}^\ell$. Then, given $x_0 \in \mathbb{R}^\ell$ and $r > 0$, define the functional $F_{x_0,r}$ by

$$(2-3) \quad F_{x_0,r}(\Sigma) = \frac{1}{(4\pi r)^{k/2}} \int_{\Sigma} e^{-|x-x_0|^2/(4r)} d\mu.$$

(When $x_0 = \vec{0}$ and $r = 1$, this is just a normalization of area in the Gaussian metric.) Colding and Minicozzi then define the entropy $\lambda(\Sigma)$ of a submanifold to be the supremum over all $F_{x_0,r}$ functionals:

$$(2-4) \quad \lambda(\Sigma) = \sup_{x_0,r} F_{x_0,r}(\Sigma).$$

The aforementioned Huisken monotonicity [22] implies that this quantity is in fact monotone under the flow, and because it is defined as a supremum over rescalings and recenterings, it also controls the nature of singularities encountered along the flow; see [9; 2; 3; 4] for instance. Note that surfaces of polynomial volume growth have finite entropy.

The current state of knowledge of mean curvature flow singularities approached from an entropy perspective seems to be “quantized” by the entropy Λ_k of round spheres as we now discuss. By a calculation of Stone [34] we have

$$\Lambda_1 > \frac{3}{2} > \Lambda_2 > \cdots > \Lambda_n \rightarrow \sqrt{2}.$$

So far in the literature, many results using an entropy condition assume that the submanifold M under consideration satisfies $\lambda(M) < \Lambda_{n-1}$, which seems to most often be referred to as a low or small entropy condition. The next natural entropy condition to consider then is a bound by Λ_{n-2} , which we refer to as a medium entropy bound; one might expect studying surfaces with this entropy bound to be tractable because morally it implies that mean convex singularities encountered will be 2-convex, which as implied above in the discussion on surgery are the easiest to consider/flow through (after convex ones). Indeed this philosophy was carried out in the compact case in the joint work with S Wang [29] (“low” in its title refers to what we define as medium). An important observation for our argument is that this philosophy can be extended to the noncompact setting, but there are significant new issues to consider. For instance, in the noncompact case the asymptotics of the submanifold in question matter.

Throughout this article we will say an end E of a self-shrinker is *asymptotically conical* if E satisfies $\lim_{\tau \rightarrow \infty} \tau^{-1} E = C(E)$ in $C_{\text{loc}}^\infty(\mathbb{R}^{n+1} \setminus 0)$ for $C(E)$ a regular cone in \mathbb{R}^{n+1} . A similar definition can be made for asymptotically cylindrical ends, and by results of L Wang [35], for $n = 2$ every end of a self-shrinker of finite topology is either asymptotically conical or cylindrical (with multiplicity one). Naturally, one says a self-shrinker is asymptotically conical if every end is. Considering singular/GMT extensions of shrinkers and asymptotically conical ends in a natural way, under suitable entropy assumptions and assumptions on the underlying measure, the support of the shrinker and asymptotic cone can be shown to be smooth (so asymptotically conical as in the sense above); see [4, Propositions 3.2 and 3.3; 8, Lemma 2.1]. In particular, our theorem applies to asymptotically conical (in the weak sense) shrinkers which arise as blowups under our entropy assumption in \mathbb{R}^4 . Note, since $\Lambda_1 < 2$, that the convergence will be with multiplicity one.

In the same paper where they introduced entropy, Colding and Minicozzi showed the only singularities which morally shouldn’t be able to be perturbed away are the mean convex ones, the generalized round cylinders $S^k(\sqrt{2k}) \times \mathbb{R}^{n-k}$, called round because the spherical factor is a standard round sphere of a radius appropriate to satisfy the shrinker equation. In particular, other singularity models should be able to be perturbed

away, so round cylinders are called generic singularity models. Their numbers are few (only n of them), whereas for instance in \mathbb{R}^3 there are self-shrinkers of arbitrarily large genus by [25], so one could say most self-shrinkers are nongeneric.

Concerning nongeneric singularity models, the no-cylinder conjecture of Ilmanen says that the types of ends shouldn't be "mixed" in that if there is a single cylindrical end then M is a cylinder, so one expects that "most" self-shrinkers in \mathbb{R}^3 are asymptotically conical (see [36] for a partial result confirming this). Extending this conjecture to the next higher dimension, this provides our justification with the above paragraph in mind for the claim that self-shrinkers are "often" asymptotically conical — it is also quite convenient for analytical reasons.

Returning to flows through singularities, an important advantage of the mean curvature flow with surgery is that the topological change across discontinuous times, when necks are cut and high curvature regions discarded, is easy to understand. A disadvantage is that it isn't quite a Brakke flow (a geometric measure theory formulation of the mean curvature flow) and so does not immediately inherit some of the consequences thereof, but at least it is closely related to the level set flow by results of Lauer [26] and Head [19; 20] which, in the nonfattening case, is (modulo some technicalities). In their work they show that surgery converges to the level set flow in Hausdorff distance (and in fact in the varifold sense, as Head shows) as the surgery parameters degenerate (ie as one lets $H_{\text{th}} \rightarrow \infty$). This connection is useful for us because deep results of White [38] show that a mean convex LSF will converge to a (possibly empty) stable minimal surface long-term.

As mentioned above, mean curvature flow with surgery in a curved ambient setting (at least for 3-manifolds and bounded geometry) has been already accomplished by Haslhofer and Ketover, but some extra care is needed for the Gaussian metric, especially in the noncompact case. This is because the metric is poorly behaved at infinity (as one sees from the calculation of its scalar curvature), which introduces some analytic difficulties for using the flow, so instead we consider the renormalized mean curvature flow (which we'll abbreviate RMCF) defined by

$$(2-5) \quad \frac{dX}{dt} = \vec{H} + \frac{1}{2}X.$$

Here, as before, X is the position vector on M . It is related to the regular mean curvature flow by the following reparametrization; this allows one to transfer many deep theorems on the MCF to the RMCF. Suppose that M_t is a mean curvature flow on

$[-1, T)$ for $-1 < T \leq 0$ ($T = 0$ is the case for a self-shrinker). Then the renormalized flow \widehat{M}_τ of M_t , defined on $[0, -\log(-T))$, is given by

$$(2-6) \quad \widehat{X}_\tau = e^{\tau/2} X_{-e^{-\tau}}, \quad \tau = -\log(-t).$$

Up to any finite time the reparametrization is bounded and preserves many properties of the regular MCF, like the avoidance principle and that entropy is monotone under the RMCF. With this in mind, the author showed in his previous article [28] that one can then construct a flow with surgery using the RMCF on suitable perturbations of noncompact self-shrinkers, and that as one lets the surgery parameters degenerate, indeed the surgery converges to the level set flow when $n = 2$. This can be readily combined with the aforementioned joint work with S Wang [29] to show the following:

Theorem 2.2 *Let $M^n \subset \mathbb{R}^{n+1}$ be a smoothly asymptotically conical hypersurface such that $H - \frac{1}{2}\langle X, \nu \rangle \geq c(1 + |X|^2)^{-\alpha}$ for some constants $c, \alpha > 0$ and choice of normal such that $\lambda(M) < \Lambda_{n-2}$. Then denoting by K the region bounded by M whose outward normal corresponds to the choice of normal on M , the level set flow M_t of M with respect to the renormalized mean curvature flow satisfies:*

- (1) *The flow is inward, in that $K_{t_1} \subset K_{t_2}$ for any $t_1 > t_2$, considering the corresponding motion of K .*
- (2) *M_t is the Hausdorff limit of surgery flows S_t^k with initial data M .*
- (3) *M_t is a forced Brakke flow (with forcing term given by position vector).*

Here α -noncollapsedness means there are inner and outer osculating balls of radius proportional to the shrinker mean curvature, and this has many consequences; see [1; 16]. The assumption on the asymptotics are conditions for which shrinker mean convexity is preserved and existence of an entropy-decreasing perturbation of a self-shrinker smoothly asymptotic to a cone can always be assumed to satisfy this by work of Bernstein and L Wang in [3]. We use this theorem (often implicitly) below with $n = 3$ when we discuss the flow of M .

H_G , the mean curvature in the Gaussian metric, is related to the renormalized mean curvature by $H_G = e^{|x|^2/4}(H - \frac{1}{2}\langle X, \nu \rangle)$, and as a result the time limit of the flow defined in the theorem above by White's theory for mean convex MCF (in particular [38]) will be a stable self-shrinker if nonempty. It will also have finite entropy by Huisken monotonicity, and hence have polynomial volume growth. As a result, either by the instability results mentioned above or by the Frenkel theorem for self-shrinkers given in the appendix of [7], we have the following:

Lemma 2.3 *Let M_t be the flow defined in Theorem 2.2. Then $\lim_{t \rightarrow \infty} M_t = \emptyset$.*

Lastly, note that by switching our choice of normal and using minimality (in the Gaussian metric) of the original surface we may shrinker mean convex perturb either inward or outward (for a 2-sided surface, of course, the distinction is somewhat arbitrary) to study its topology as observed in [3; 21]—this idea is critical to our argument and we will make our choice of perturbation depending on which case we are considering in the argument below.

3 Proof of Theorem 1.2

Note, by the entropy assumption (in particular that $\Lambda_1 < 2$), that M is embedded, and hence its link is too. Without loss of generality for this section, the link L is connected. Supposing L is not diffeomorphic to S^2 , there exists some $R_1 \gg 0$ such that $M \cap S(0, R) := L_R$ is not diffeomorphic to S^2 for $R > R_1$ and that, by the asymptotically conical assumption, $L_R \simeq L_{R_1}$ for all $R > R_1$. By the classification of surfaces (note that L_R is orientable, which can be seen by projecting the normal of $M \cap S(0, R)$ onto $TS(0, R)$, giving a section of the normal bundle of L_R , which has no kernel because the sphere intersects M transversely) L_R is topologically a connect sum of tori which bounds domains (not necessarily handlebodies) $K_R, K_R^c \subset S(0, R)$.

Fixing a choice of $R > R_1$, consider a standard generator $\gamma \subset L_R$ of $H_1(L_R)$; that is, writing L_R as a connect sum of tori, γ is homotopic to one of the two generators of a single one of the tori. Note that γ is also homotopically nontrivial in L_R . We consider two cases: either γ is homotopically trivial in M or not. Without loss of generality, γ is embedded and smooth as well.

3.1 Case 1: γ is nullhomotopic in M

Since γ is homotopically trivial in M , it bounds a disc D in M ; suppose $D \subset M \cap B(0, R_2)$. Hence, for any embedded curve $\gamma' \subset L_{R'}$ isotopic (in M) to γ for $R' > R_2$, γ' is nullhomotopic in $M \cap B(0, R')$ and hence bounds an embedded disc $D' \subset M \cap B(0, R')$ by Dehn's lemma (see [18])—Dehn's lemma gives a PL embedded disc, but when γ' is smooth note that D' can be taken to be smooth as well by the Whitney approximation theorem [27]. The idea is, morally, such discs serve as barriers in a sense to keep the flow of (a perturbation of) M “propped” up. The following indicates which domain M bounds to perturb and flow into:




Figure 1: This figure illustrates toy examples one might imagine for Case 1 (left) and Case 2 (right), considered when L_R is a standardly embedded torus. In the first case, $\gamma \in \partial(M \cap B(0, R))$ is nullhomotopic in M , and hence bounds an embedded disc $D \subset M \cap B(0, R)$ by Dehn’s lemma.

Lemma 3.1 *In one of K_R or K_R^c the curve γ is not homotopically trivial.*

Proof It seems one could probably use the Mayer–Vietoris sequence and Hurewicz isomorphism here as in the proof of Theorem 1 in [21], but we present a more geometric argument. Suppose, for the sake of contradiction, it were homotopically trivial in both simultaneously. By Dehn’s lemma, γ bounds PL embedded (of course, in fact, smooth) discs $D_1 \subset K_R$ and $D_2 \subset K_R^c$ which intersect along γ , giving an embedded $S^2 \subset S^3$. Since γ is smooth their union gives a PL embedded S^2 , and so by Alexander’s theorem (see [18]) $D_1 \cup D_2$ then bounds a (PL embedded) 3–ball $B \subset S^3$. From its construction, L_R intersects B in one boundary component, namely γ . In particular, γ is homologically trivial in L_R , giving a contradiction. \square

Of course, the lemma applies equally for γ' homotopic to γ in $L_{R'}$ for $R' > R > R_1$. After potentially relabeling, γ is homotopically nontrivial in K_R . In this case, consider a shrinker mean convex perturbation of M , as constructed in [3], which descends (ie intersecting with $S(0, R)$) to a perturbation of L_R into K_R , and consider the corresponding renormalized flow M_t (recalling that we can choose which direction to flow into, as discussed in the preliminaries). This flow likewise descends to a flow $(L_R)_t$ of L_R . Note though that, although M_t is an RMCF, $(L_R)_t$ isn’t necessarily (to the author’s knowledge) an easily described flow in $S(0, R)$, but we will still find it profitable to consider.

By Lemma 2.3, M_t must leave every bounded set in some finite time, and hence $(L_R)_t$ must eventually become empty. Denote this time by T . We will play it off against the next two lemmas, the first essentially that the disc we find by Dehn’s lemma persists:

Lemma 3.2 *For any $\tau > 0$, one can pick \bar{R} sufficiently large that, for $t \in [0, \tau]$, there will be a smoothly embedded curve $\bar{\gamma}_t \subset (L_{\bar{R}})_t$ isotopic to γ which bounds a smoothly embedded disc $\bar{D}_t \subset B(0, \bar{R}) \cap M_t$.*

Proof With the construction of the flow by surgery flows given in [Theorem 2.2](#) in mind, we first show, for exposition, that this holds for an approximating surgery flow S_t of M . Since the $L_{R'}$ are all diffeomorphic for R' large, there is clearly an initial choice of curve $\bar{\gamma}$ which is isotopic to γ . Up to the first surgery time and in between surgery times when the surgery flow is smooth, this curve is just given by restricting the motion of $(L_{\bar{R}})_t$ along $\bar{\gamma}$ because, for \bar{R} large enough, $(L_{\bar{R}})_t$ will be a graph over $(L_{\bar{R}})_0$ by pseudolocality and [Theorem 2.2\(3\)](#) on $[0, \tau]$ (in particular, $\bar{\gamma}_t$ can be taken to be embedded and smoothly vary) for times in $[0, \tau]$. Concerning the bounded disc for smooth times, the flow is an isotopy which restricts to an isotopy of the disc (modding out tangential components of the flow). Now consider a surgery time $t_s < \tau$; we must check that after surgery $\bar{\gamma}_{t_s}$ still bounds a disc. Again by pseudolocality for all surgery necks N , $N \cap S(0, \bar{R})$ is empty, and similarly all N must be within $B(0, \bar{R})$. Considering a cap C in the surgery procedure, since it is topologically a ball, the intersection of D_{t_s} with $\partial C \simeq S^2$ is a disjoint union of closed curves which bound discs by the Schoenflies theorem (without loss of generality D_t enters all caps transversely). Surgering along these discs gives a union of S^2 along with a new disc whose boundary is $\bar{\gamma}_{t_s}$ (essentially filling in the part of the disc between the end and the “closest” surgery necks). In particular, $\bar{\gamma}_{t_s}$ continues to bound a disc after surgery. Note that it’s conceivable at this stage that the discarded copies of S^2 bound nontrivial topology of M_t , so the D_t do not necessarily form an isotopic family of discs, a priori. Now we discuss how to show the curve $\bar{\gamma}$ from the previous paragraph always bounds a disc in the limiting flow. What one might first wish for is to take a limit (by compactness) of the discs as the surgery parameters degenerate, but if the limiting disc enters a singular region of the flow it could potentially complicate things, so it’s best if the disc is taken to avoid it completely. There is also the matter of boundedness along this sequence of discs needed to apply a compactness theorem, which suggests it’s best, in terms of the disc, to work only within the context of the level set flow.

To begin, we consider high curvature points we might encounter as we travel sufficiently deep within a high curvature region (loosely speaking) from a low curvature region, as in our situation of a disc starting from an end (where $\bar{\gamma}$ is) approaching a singularity in the interior of $B(0, \bar{R})$. At points where, say, $H \sim H_{\text{can}}$, referring to parameters in the canonical neighborhood theorem (see [\[17\]](#), noting that here we suppress some notation),

one can find nearby “neck-like” points (see Proposition 3.2 in [17]) in any corresponding ancient model that could appear irrespective of surgery parameters. Intuitively a surgery flow near such a point is modeled locally by a neck or a cap bordered by a neck facing towards the low curvature region. With the Hausdorff convergence in mind then, one can use Arzelà–Ascoli to pass to the limit on these bounded curvature regions for the surgery flows to see that the level set flow always has necks where $H \sim H_{\text{can}}$ as one approaches a singular region of the level set flow from a low curvature one (of course, these are smooth points as well). These necks on the level set flow give points to surger the disc as we did for the surgery flows; in this case, we perform the surgery on the disc whenever one of its points is in a region of the level set flow where, say, $H = 10H_{\text{can}}$. Note that between these times the disc varies continuously since it is within a region of the level set flow of bounded curvature, and that the disc can be taken to be smooth at all times since it is surgered on along cross sections of necks of bounded curvature. \square

Without using Dehn’s lemma (and, in particular, with repeating the argument in higher dimensions in mind), it seems the intersection of the disc with the boundary of the cap could be much more complicated, although naively it seems likely that γ would remain homotopically trivial. We will pit this lemma against the definition of the time T using the following lemma, which says the discs must “leave” the end no matter what:

Lemma 3.3 *With $\bar{\gamma}$ and τ as in Lemma 3.2, after potentially taking R larger, there is an $\epsilon > 0$ such that, in $B(0, R - \epsilon)^c \cap M_t$, the curve $\bar{\gamma}_t$ isn’t nullhomotopic and so doesn’t bound a disc $B(0, R - \epsilon)^c \cap M_t$. In particular, the disc D_t from the previous lemma satisfies that $D_t \cap S(0, R - \epsilon)$ is nonempty on $[0, \tau]$.*

Proof Denote by K the region M bounds which includes K_R . Note then that, for R large enough, $B(0, R - \epsilon)^c \cap K \simeq K_R \times [R - \epsilon, \infty)$, and in particular $\bar{\gamma}$ is homotopically nontrivial in this domain since it is homotopically nontrivial in K_R . If, for some time $t \in [0, \tau]$, $\bar{\gamma}_t$ is nullhomotopic in $B(0, R - \epsilon)^c \cap M_t$, then in particular $\bar{\gamma}_t$ bounds (the image of) a disc in $B(0, R - \epsilon)^c \cap K_t$. By the set monotonicity of the flow, ie that $K_t \subset K$, we get in fact that $\bar{\gamma}_t$ and hence $\bar{\gamma}$ are nullhomotopic in $B(0, R - \epsilon)^c \cap K$, giving a contradiction. \square

Applying the above lemmas with $\tau = T + 1$, we see that we arrive at a contradiction. Considering a time $t \in (T, T + 1)$ and the disc D_t given from Lemma 3.2, the disc, by Lemma 3.3, must have nonempty intersection with $S(0, R - \epsilon)$. On the other hand, it cannot pass through $S(0, R)$ because $(L_R)_t = \emptyset$ for $t > T$. This completes the argument in this case.

3.2 Case 2: γ is homotopically nontrivial in M

This case is easier in a sense, because we may directly apply the deep ideas of White [39], in particular Theorems 1.1 and 5.2 therein. Specialized to our setting, a consequence is that if K is a smooth mean convex set (and compact as stated in Theorem 1.1, but this can also apply in the noncompact case as long as singularities occur only in a bounded ball for the time it is applied, by Theorem 5.2) in a Riemannian manifold N of dimension 4 (in particular, less than 7), then if a curve in K^c is initially homotopically nontrivial and later becomes contractible in $(K^c)_t$, a singularity of the form $S^1 \times S^2$ must have occurred, contradicting the entropy bound. Here we will consider N to be a subset of \mathbb{R}^4 (possibly all of \mathbb{R}^4 , depending on which case we are in below) endowed with the Gaussian metric, so the flow constructed is more precisely a mean convex foliation. However, the flow is monotone, satisfies the Brakke regularity theorem and the singular set dimension results of White [37], and all the singularities are modeled on round cylinders so the results of the paper apply — this is essentially the upshot of Hershkovits and White [21] (although they phrase things entirely in terms of the RMCF), where they study the interplay of entropy and topology for compact self-shrinkers.

There are two possible cases for γ : that γ is homotopically nontrivial in one of the components K or K^c of \mathbb{R}^4 it bounds, or not. First suppose that γ is homotopically nontrivial in (at least) one of K or K^c , say K^c to align with White's terminology. Consider then a nontrivial curve γ in K^c . Since γ is contractible in \mathbb{R}^4 , the corresponding homotopy gives that it bounds a (continuous image of, perhaps not embedded) disc D — note that this disc must intersect K . Perturbing and flowing into K by Lemma 2.3, eventually we must have $D \subset K_t^c$, say, by T , implying by this time that γ' is nullhomotopic in K_t^c . By pseudolocality [6] there is an $R \gg 0$ such that, near $S(0, R)$, M_t is a smooth flow which intersects the sphere transversely, so defining $N = B(0, R)$, Section 5 of White [39] implies a singularity modeled on $S^1 \times \mathbb{R}^2$ formed, contradicting the entropy bound.

Now we consider the possibility that γ is homotopically trivial in both K and K^c ; this naively seems to be a more exotic case than above, but we are unsure it can be ruled out a priori by purely topological reasoning. Then $\gamma \in M$ bounds a disc in both K and K^c . Picking essentially arbitrarily (only to align with White's notation), we define \tilde{N} to be the union of K and $K^c \cap M \times [0, \epsilon) \nu$ (ie a collar of M), where ν is the normal pointing away from K and $\epsilon > 0$ is some number small enough that the ϵ level set of the collar is also embedded in \mathbb{R}^4 . Note that this collar region retracts onto M ; the utility of this is that now γ is a homotopically nontrivial curve in $K^c \cap \tilde{N} \subset \tilde{N} \subsetneq \mathbb{R}^4$. Consider, as in the

previous paragraph, a disc $D \subset K$ bounding γ and flow out of K^c into K (that the disc can be taken to be contained in a single component, and hence in N , is why we split this up into cases). As above, [Lemma 2.3](#) (this still applies since the flow of M_t is the same considered in \mathbb{R}^4 or \tilde{N}) gives that eventually $D \subset K_t^c \cap \tilde{N}$ — call this time T . Let $R \gg 0$ be large enough that M_t intersects $S(0, R)$ only transversely and as a smooth flow; again such an R exists by pseudolocality. Defining $N = \tilde{N} \cap B(0, R)$ and noting that γ is still homologically nontrivial in $K^c \cap \tilde{N} \cap B(0, R)$, Section 5 of [\[39\]](#) gives that a singularity modeled on $S^1 \times \mathbb{R}^2$ must have formed (in fact, by time T), giving a contradiction.

4 Proof of [Theorem 1.1](#)

By the Frenkel property for self-shrinkers, M must be connected. By [Theorem 1.2](#) there exists R sufficiently large that $M \cap B(0, R)$ is diffeomorphic to a connected 2-sided hypersurface N^3 whose boundary consists of a number of 2-spheres along each of which an end homeomorphic to $S^2 \times \mathbb{R}_+$ is attached, where by ends here we mean, for an appropriate choice of R , disjoint connected components of $M \setminus B(0, R)$ which are diffeomorphic to half cylinders over distinct (for distinct ends) connected components of the link; such an R exists since M is asymptotically conical, and the convergence is multiplicity one. The point is to confirm that N is simply connected. Then, by capping off each component of N (considering N as an intrinsically defined manifold, as a hypersurface in \mathbb{R}^4 it seems some ends could be “parallel”, which would preclude doing this at least in an embedded way) with a 3-ball, we obtain a closed connected simply connected 3-manifold \tilde{N} which, by the resolution of the 3D Poincaré conjecture, is diffeomorphic to S^3 . If there is a homotopically nontrivial curve on N and hence M , by the Seifert–Van Kampen theorem and [Theorem 1.2](#) we can proceed directly as we do in the second case of the proof above using [\[39\]](#), giving the first part of [Theorem 1.1](#). Note that, with surgery for compact manifolds in mind, one should be able to argue directly with a bit more work that \tilde{N} is diffeomorphic to either S^3 or a connect sum of $S^2 \times S^1$, the latter of which could subsequently be ruled out, avoiding the use of the Poincaré conjecture — this seems to be naturally a more robust line of reasoning for considering higher-dimensional versions of our statement.

When the number of ends is equal to one, M is diffeomorphic to \mathbb{R}^3 as a consequence of Alexander’s theorem, as noted in [\[4\]](#). Now suppose that $\lambda(M) < \Lambda_2$ (we will discuss the case of equality afterwards) and M had (at least) two ends, labeled E_1 and E_2 . Fixing an R in our definition of end given in the paragraph above, consider a curve $\gamma: \mathbb{R} \rightarrow M$

such that for s sufficiently negative $\gamma(s) \in E_1$ and for s sufficiently positive $\gamma(s) \in E_2$. With this in mind, intersect M with an embedded hypersurface $P \simeq \mathbb{R}^3$ such that

- (i) E_1 lies on one side of P and E_2 lies on the other side,
- (ii) P intersects M transversely, and
- (iii) $P \cap M$ is compact.

This is always possible by the asymptotically conical assumption. Denote by $P \cap M$ the surface S ; note that S is closed since its compact and M is boundaryless. Similarly, denote the bounded portion of P that S bounds by K_S . By perturbing and flowing M so that, restricted to N , the flow is into K_S (using (ii)) we see, as above, by [Lemma 2.3](#) that S_t is eventually empty. Because of this, one may argue that a singularity of M_t must occur which disconnects E_1 from E_2 along γ ; note that by using large spheres as barriers far along the ends toward spatial infinity (or, alternatively, pseudolocality), for any given finite time there will be points originating from E_1 and E_2 on one side of P and the other, respectively. In other words, one end can't flow from one side of P to the other in finite time, so a singularity which disconnects M must indeed occur. Clearly such a singularity must be modeled on $S^2 \times \mathbb{R}$, which has entropy Λ_2 , contradicting $\lambda(M) < \Lambda_2$. In the case $\lambda(M) = \Lambda_2$, we note that the perturbation of Bernstein and Wang we used strictly decreases entropy placing us in the case of strict inequality.

References

- [1] **B Andrews**, *Noncollapsing in mean-convex mean curvature flow*, *Geom. Topol.* 16 (2012) 1413–1418 [MR](#) [Zbl](#)
- [2] **J Bernstein, L Wang**, *A sharp lower bound for the entropy of closed hypersurfaces up to dimension six*, *Invent. Math.* 206 (2016) 601–627 [MR](#) [Zbl](#)
- [3] **J Bernstein, L Wang**, *A topological property of asymptotically conical self-shrinkers of small entropy*, *Duke Math. J.* 166 (2017) 403–435 [MR](#) [Zbl](#)
- [4] **J Bernstein, L Wang**, *Topology of closed hypersurfaces of small entropy*, *Geom. Topol.* 22 (2018) 1109–1141 [MR](#) [Zbl](#)
- [5] **S Brendle, G Huisken**, *Mean curvature flow with surgery of mean convex surfaces in \mathbb{R}^3* , *Invent. Math.* 203 (2016) 615–654 [MR](#) [Zbl](#)
- [6] **B-L Chen, L Yin**, *Uniqueness and pseudolocality theorems of the mean curvature flow*, *Comm. Anal. Geom.* 15 (2007) 435–490 [MR](#) [Zbl](#)
- [7] **O Chodosh, K Choi, C Mantoulidis, F Schulze**, *Mean curvature flow with generic initial data*, preprint (2020) [arXiv 2003.14344](#)

- [8] **O Chodosh, K Choi, C Mantoulidis, F Schulze**, *Mean curvature flow with generic low-entropy initial data*, preprint (2021) [arXiv 2102.11978](https://arxiv.org/abs/2102.11978)
- [9] **T H Colding, T Ilmanen, W P Minicozzi, II, B White**, *The round sphere minimizes entropy among closed self-shrinkers*, *J. Differential Geom.* 95 (2013) 53–69 [MR](#) [Zbl](#)
- [10] **T H Colding, W P Minicozzi, II**, *Generic mean curvature flow, I: Generic singularities*, *Ann. of Math.* 175 (2012) 755–833 [MR](#) [Zbl](#)
- [11] **T H Colding, W P Minicozzi, II**, *Smooth compactness of self-shrinkers*, *Comment. Math. Helv.* 87 (2012) 463–475 [MR](#) [Zbl](#)
- [12] **J M Daniels-Holgate**, *Approximation of mean curvature flow with generic singularities by smooth flows with surgery*, *Adv. Math.* 410 (2022) art. id. 108715 [MR](#) [Zbl](#)
- [13] **Q Ding, Y L Xin**, *Volume growth, eigenvalue and compactness for self-shrinkers*, *Asian J. Math.* 17 (2013) 443–456 [MR](#) [Zbl](#)
- [14] **M H Freedman**, *The topology of four-dimensional manifolds*, *J. Differential Geometry* 17 (1982) 357–453 [MR](#) [Zbl](#)
- [15] **R Haslhofer, D Ketover**, *Minimal 2–spheres in 3–spheres*, *Duke Math. J.* 168 (2019) 1929–1975 [MR](#) [Zbl](#)
- [16] **R Haslhofer, B Kleiner**, *Mean curvature flow of mean convex hypersurfaces*, *Comm. Pure Appl. Math.* 70 (2017) 511–546 [MR](#) [Zbl](#)
- [17] **R Haslhofer, B Kleiner**, *Mean curvature flow with surgery*, *Duke Math. J.* 166 (2017) 1591–1626 [MR](#) [Zbl](#)
- [18] **A Hatcher**, *Notes on basic 3–manifold topology*, unpublished manuscript (2007) Available at <https://pi.math.cornell.edu/~hatcher/3M/3Mdownloads.html>
- [19] **J Head**, *The surgery and level-set approaches to mean curvature flow*, PhD thesis, Max Planck Institute for Gravitational Physics (2011) Available at <https://d-nb.info/1025939417/34>
- [20] **J Head**, *On the mean curvature evolution of two-convex hypersurfaces*, *J. Differential Geom.* 94 (2013) 241–266 [MR](#) [Zbl](#)
- [21] **O Hershkovits, B White**, *Sharp entropy bounds for self-shrinkers in mean curvature flow*, *Geom. Topol.* 23 (2019) 1611–1619 [MR](#) [Zbl](#)
- [22] **G Huisken**, *Asymptotic behavior for singularities of the mean curvature flow*, *J. Differential Geom.* 31 (1990) 285–299 [MR](#) [Zbl](#)
- [23] **G Huisken, C Sinestrari**, *Mean curvature flow with surgeries of two-convex hypersurfaces*, *Invent. Math.* 175 (2009) 137–221 [MR](#) [Zbl](#)
- [24] **T Ilmanen, B White**, *Sharp lower bounds on density for area-minimizing cones*, *Camb. J. Math.* 3 (2015) 1–18 [MR](#) [Zbl](#)
- [25] **N Kapouleas, S J Kleene, N M Møller**, *Mean curvature self-shrinkers of high genus: non-compact examples*, *J. Reine Angew. Math.* 739 (2018) 1–39 [MR](#) [Zbl](#)

- [26] **J Lauer**, *Convergence of mean curvature flows with surgery*, *Comm. Anal. Geom.* 21 (2013) 355–363 [MR](#) [Zbl](#)
- [27] **JM Lee**, *Introduction to smooth manifolds*, 2nd edition, *Graduate Texts in Math.* 218, Springer (2013) [MR](#) [Zbl](#)
- [28] **A Mramor**, *An unknottedness result for noncompact self shrinkers*, preprint (2020) [arXiv 2005.01688](#)
- [29] **A Mramor, S Wang**, *Low entropy and the mean curvature flow with surgery*, *Calc. Var. Partial Differential Equations* 60 (2021) art. id. 96 [MR](#) [Zbl](#)
- [30] **G Perelman**, *The entropy formula for the Ricci flow and its geometric applications*, preprint (2002) [Zbl](#) [arXiv math/0211159](#)
- [31] **G Perelman**, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, preprint (2003) [Zbl](#) [arXiv math/0307245](#)
- [32] **G Perelman**, *Ricci flow with surgery on three-manifolds*, preprint (2003) [Zbl](#) [arXiv math/0303109](#)
- [33] **S Smale**, *Generalized Poincaré’s conjecture in dimensions greater than four*, *Ann. of Math.* 74 (1961) 391–406 [MR](#) [Zbl](#)
- [34] **A Stone**, *A density function and the structure of singularities of the mean curvature flow*, *Calc. Var. Partial Differential Equations* 2 (1994) 443–480 [MR](#) [Zbl](#)
- [35] **L Wang**, *Asymptotic structure of self-shrinkers*, preprint (2016) [arXiv 1610.04904](#)
- [36] **L Wang**, *Uniqueness of self-similar shrinkers with asymptotically cylindrical ends*, *J. Reine Angew. Math.* 715 (2016) 207–230 [MR](#) [Zbl](#)
- [37] **B White**, *Stratification of minimal surfaces, mean curvature flows, and harmonic maps*, *J. Reine Angew. Math.* 488 (1997) 1–35 [MR](#) [Zbl](#)
- [38] **B White**, *The size of the singular set in mean curvature flow of mean-convex sets*, *J. Amer. Math. Soc.* 13 (2000) 665–695 [MR](#) [Zbl](#)
- [39] **B White**, *Topological change in mean convex mean curvature flow*, *Invent. Math.* 191 (2013) 501–525 [MR](#) [Zbl](#)

Department of Mathematics, Johns Hopkins University
Baltimore, MD, United States

amramor1@jhu.edu

Proposed: Tobias H Colding
Seconded: Gang Tian, Bruce Kleiner

Received: 30 June 2021
Revised: 22 March 2022

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Frances Kirwan	University of Oxford frances.kirwan@balliol.oxford.ac.uk
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	John Lott	University of California, Berkeley lott@math.berkeley.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Haynes Miller	Massachusetts Institute of Technology harm@math.mit.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Walter Neumann	Columbia University neumann@math.columbia.edu
Benson Farb	University of Chicago farb@math.uchicago.edu	Jean-Pierre Otal	Université Paul Sabatier, Toulouse otal@math.univ-toulouse.fr
Steve Ferry	Rutgers University sferry@math.rutgers.edu	Peter Ozsváth	Princeton University petero@math.princeton.edu
David M Fisher	Rice University davidfisher@rice.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Mike Freedman	Microsoft Research michaelf@microsoft.com	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David Gabai	Princeton University gabai@princeton.edu	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Peter Teichner	Max Planck Institut für Mathematik teichner@mac.com
Cameron Gordon	University of Texas gordon@math.utexas.edu	Richard P Thomas	Imperial College, London richard.thomas@imperial.ac.uk
Lothar Götsche	Abdus Salam Int. Centre for Th. Physics gotsche@ictp.trieste.it	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Mark Gross	University of Cambridge mgross@dpms.cam.ac.uk	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2023 is US \$740/year for the electronic version, and \$1030/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2023 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 27 Issue 9 (pages 3387–3831) 2023

- Hyperbolic groups acting improperly 3387
DANIEL GROVES and JASON FOX MANNING
- Cyclic homology, S^1 -equivariant Floer cohomology and Calabi–Yau structures 3461
SHEEL GANATRA
- Congruences on K-theoretic Gromov–Witten invariants 3585
JÉRÉMY GUÉRE
- Moduli of spherical tori with one conical point 3619
ALEXANDRE EREMENKO, GABRIELE MONDELLO and
DMITRI PANOV
- The derivative map for diffeomorphism of disks: an example 3699
DIARMUID CROWLEY, THOMAS SCHICK and
WOLFGANG STEIMLE
- On self-shrinkers of medium entropy in \mathbb{R}^4 3715
ALEXANDER MRAMOR
- The Gromov–Hausdorff distance between spheres 3733
SUNHYUK LIM, FACUNDO MÉMOLI and ZANE SMITH
- Contact three-manifolds with exactly two simple Reeb orbits 3801
DANIEL CRISTOFARO-GARDINER, UMBERTO
HRYNIEWICZ, MICHAEL HUTCHINGS and HUI LIU