



# *Geometry & Topology*

Volume 28 (2024)

**Algebraic uniqueness of Kähler–Ricci flow limits  
and optimal degenerations of Fano varieties**

JIYUAN HAN

CHI LI

# Algebraic uniqueness of Kähler–Ricci flow limits and optimal degenerations of Fano varieties

JIYUAN HAN

CHI LI

We prove that for any Fano manifold  $X$ , the special  $\mathbb{R}$ –test configuration that minimizes the  $H^{\text{NA}}$ –functional is unique and has a  $\mathbb{K}$ –semistable  $\mathbb{Q}$ –Fano central fiber  $(W, \xi)$ . Moreover there is a unique  $\mathbb{K}$ –polystable degeneration of  $(W, \xi)$ . As an application, we confirm the conjecture of Chen, Sun and Wang about the algebraic uniqueness for Kähler–Ricci flow limits on Fano manifolds, which implies that the Gromov–Hausdorff limit of the flow does not depend on the choice of initial Kähler metrics. The results are achieved by studying algebraic optimal degeneration problems via new functionals for real valuations over  $\mathbb{Q}$ –Fano varieties, which are analogous to the minimization problem for normalized volumes.

14J45, 32Q26, 53E30

1. Introduction	539
2. Preliminaries	544
3. $H^{\text{NA}}$ invariant and MMP	564
4. A minimization problem for real valuations	570
5. Initial term degeneration of filtrations	575
6. Uniqueness of minimizing special $\mathbb{R}$ –test configurations	578
7. Cone construction and $g$ –normalized volume	580
8. Uniqueness of polystable degeneration	583
Appendix. Properties of $\tilde{\mathcal{S}}(v)$	585
References	589

## 1 Introduction

Let  $X$  be a smooth Fano manifold. It is now known that  $X$  admits a Kähler–Einstein metric if and only if  $X$  is  $\mathbb{K}$ –polystable; see Berman [5], Chen, Donaldson and Sun [25; 26; 27] and Tian [67; 68]. In this paper, we are interested in the case when  $X$  is not  $\mathbb{K}$ –polystable. If  $X$  is strictly  $\mathbb{K}$ –semistable,

then  $X$  admits a unique  $K$ -polystable degeneration by Li, Wang and Xu [55]. If  $X$  is  $K$ -unstable (ie not  $K$ -semistable), several kinds of optimal degenerations have been studied which are related to continuity methods or geometric flows in the analytic study of canonical metrics. For example, related to Aubin's continuity method, there is a (not necessarily unique) special degeneration whose associated valuation minimizes the  $\delta$  invariant; see Blum, Liu and Zhou [16] and Székelyhidi [65]. There is also a unique destabilizing geodesic ray which arises in the study of inverse Monge–Ampère flow (resp. Calabi flow) and whose associated non-Archimedean metric minimizes an  $L^2$ -normalized non-Archimedean Ding invariant (resp.  $L^2$ -normalized radial Calabi functional); see Donaldson [33], Hisamoto [42] and Xia [77]. In this paper we are interested in optimal degenerations that arise in the study of Hamilton–Tian conjecture about the long time behavior of Kähler–Ricci flows. The latter conjecture states that starting from any Kähler metric  $\omega \in c_1(X)$ , the normalized Kähler–Ricci converges in the Gromov–Hausdorff sense to a Kähler–Ricci soliton on a  $\mathbb{Q}$ -Fano variety  $X_\infty$ . The Hamilton–Tian conjecture has been solved (see Bamler [4], Chen and Wang [29] and Tian and Zhang [70]) and applied to give a proof of the Yau–Tian–Donaldson conjecture in Chen, Sun and Wang [28].

It is known that  $X_\infty$  coincides with  $X$  if and only if there is already a Kähler–Ricci soliton on  $X$ ; see Dervan and Székelyhidi [31] and Tian and Zhu [72]. In general, Chen, Sun and Wang [28] proved the following phenomenon. The metric degeneration from  $X$  to  $X_\infty$  induces a finitely generated filtration  $\mathcal{F}$  on  $R = \bigoplus_m H^0(X, -mK_X)$ , and there is a two-step degeneration:

- (i) The filtration  $\mathcal{F}$  as an  $\mathbb{R}$ -test configuration (see Definition 2.8) degenerates  $X$  to a normal Fano variety  $W$  with a torus  $\mathbb{T}$ -action generated by a holomorphic vector field  $\xi$ . For simplicity, we call this step the semistable degeneration.
- (ii) There is a  $\mathbb{T}$ -equivariant test configuration of  $(W, \xi)$  to  $(X_\infty, \xi)$ . We call this step the polystable degeneration.

As explained in Chen, Sun and Wang [28], this picture is a global analogue of the picture in Donaldson and Sun's study [34] of metric tangent cones on Gromov–Hausdorff limits of Fano Kähler–Einstein manifolds. In [34], Donaldson and Sun conjectured that metric tangent cones depend only on the algebraic structure near the singularity. This conjecture has been confirmed in a series of works of the second author with his collaborators (see Li [51], Li and Xu [58; 57] and Li, Wang and Xu [55]), which depends on the study of the minimization problem of a normalized volume functional over the space of valuations centered at the singularity; see Li, Liu and Xu [54] for a survey. Analogous to this conjecture on metric tangent cones, the following conjecture was proposed in [28]:

**Conjecture 1.1** *The data  $\mathcal{F}$ ,  $W$  and  $X_\infty$  depend only on the algebraic structure of  $X$  but not on the initial metric for the Kähler–Ricci flow.*

In this paper we will confirm Conjecture 1.1. The idea and method to prove this conjecture are in some sense parallel to the study of minimizing normalized volumes. However, the correct framework for achieving this goal has not been established until now. So the second purpose of this paper is to study

an analogous minimization problem in the global setting, which can be studied for all  $\mathbb{Q}$ –Fano varieties possibly singular, and prove various results about it.

The functional we want to minimize is called the  $H^{\text{NA}}$ –functional of  $\mathbb{R}$ –test configurations.<sup>1</sup> Tian, Zhang, Zhang and Zhu [69, Proposition 5.1] first introduced the  $H^{\text{NA}}$ –functional for holomorphic vector fields in their study of Kähler–Ricci flow on Fano manifolds. This invariant was generalized to any *special*  $\mathbb{R}$ –test configuration by Dervan and Székelyhidi [31], who then used the results of Chen and Wang [29], Chen, Sun and Wang [28] and He [40] to prove that the semistable degeneration mentioned above minimizes the  $H^{\text{NA}}$ –functional among all special  $\mathbb{R}$ –test configurations; see Remark 2.44. For general test configurations, such an  $H^{\text{NA}}$ –functional is a nonlinear version of the non-Archimedean Berman–Ding functional, and was first explicitly used by Hisamoto in [43] to reprove Dervan and Székelyhidi’s result using pluripotential theory. Note that in this paper, for the convenience of our argument and comparison with the case of the  $\delta$  invariant (or with the  $\beta$  invariant, see equation (107)), we will use the negative of the sign convention in these previous works.

Conjecture 1.1 follows from two purely algebrogeometric statements for each step of the semistable and polystable degenerations.

**Theorem 1.2** *For any  $\mathbb{Q}$ –Fano variety, the special  $\mathbb{R}$ –test configuration that minimizes  $H^{\text{NA}}$  is unique and its central fiber  $(W, \xi)$  is  $K$ –semistable (Definition 2.49).*

**Theorem 1.3** *If  $(X, \xi)$  is  $K$ –semistable, then there exists a unique  $K$ –polystable degeneration.*

**Corollary 1.4** *Conjecture 1.1 is true for any smooth Fano manifold. In particular, the Gromov–Hausdorff limit  $X_\infty$  for the Kähler–Ricci flow does not depend on the initial metric of the flow.*

To prepare for the proof of such results, we will first carry out an algebraic study of the  $H^{\text{NA}}$ –functional, which is analogous to the study of the minimization problem for normalized volume or the  $\delta$  invariant. We will prove a new interesting fact in Theorem 3.5, that the MMP process devised in [56] decreases the  $H^{\text{NA}}$  invariant of test configurations. This requires us to derive new intersection formulas (see (121)) and derivative formulas for the  $H^{\text{NA}}$  invariant. The proof of such formulas depends on a fibration technique in the study of equivariant cohomology. This technique is partly motivated by some construction from our previous work [39], although there are key differences which require more concrete calculations; see Remark 3.2.

We will then introduce the following  $\tilde{\beta}$ –functional on  $\text{Val}(X)$  the space of valuations on  $X$ : for any  $v \in \text{Val}(X)$  with  $A_X(v) < +\infty$ , we define

$$(1) \quad \tilde{\beta}(v) = A_X(v) + \log \left( \frac{1}{(-K_X)^n} \int_{\mathbb{R}} e^{-\lambda} (-d\text{vol}(\mathcal{F}_v^{(\lambda)})) \right).$$

<sup>1</sup>We will mostly use the notation of non-Archimedean functionals, as advocated in Boucksom, Hisamoto and Jonsson [19]. However, note that  $H^{\text{NA}}$  here is not the non-Archimedean entropy functional used in [19]. We will not use the non-Archimedean entropy in this article.

See Section 4 for details. If  $A_X(v) = +\infty$ , then we define  $\tilde{\beta}(v) = +\infty$ . The transition from  $H^{\text{NA}}$  to the  $\tilde{\beta}$ -functional is similar to the transition from Ding–Tian’s generalized Futaki invariant to the  $\beta$ -functional in the literature of  $K$ -stability (as first appeared in Li [50], where it was called  $\Theta$ , and in Fujita [37]). In other words,  $\tilde{\beta}$  is a nonlinear version of  $\beta$  and could also be considered as a global analogue of the normalized volume. Unlike the case of normalized volume functional, the  $\tilde{\beta}$  invariant is not invariant under rescaling of valuations. Indeed, we find the following new phenomenon: when restricted to the ray of multiples of a fixed valuation  $v \in \text{Val}(X)$  with  $A(v) < +\infty$ , it is strictly convex and proper and its derivative at the origin is exactly  $\beta(v)$ . As a consequence there is a unique minimizer along the ray, which is nontrivial if and only if  $\beta(v) < 0$  (Proposition 4.6). The above MMP result implies that the minimum can be approached by a sequence of special divisorial valuations. As a consequence, one can adapt the method developed in [14] to show that there is minimizing valuation which is quasimonomial; see Theorem 4.10. On the other hand, the  $H^{\text{NA}}$  invariant for special test configurations is expressed as the  $\tilde{\beta}$  invariant; see Lemma 4.2. Combining these discussions, we will prove (see Sections 2.2 and 2.5 for relevant notation):

**Theorem 1.5** For any  $\mathbb{Q}$ -Fano variety  $X$ , we have the identity

$$(2) \quad \inf_{\mathcal{F} \text{ filtration}} H^{\text{NA}}(\mathcal{F}) = \inf_{(\mathcal{X}, \mathcal{L}, a\eta) \text{ special}} (H^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta)) = \inf_{v \in \text{Val}(X)} \tilde{\beta}(v).$$

Moreover, the last infimum is achieved by a quasimonomial valuation.

As in the cases of normalized volume, we conjecture that the minimizer is unique and induces a special  $\mathbb{R}$ -test configuration (see Conjecture 4.11)<sup>2</sup> whose central fiber (with the induced vector field) must then be  $K$ -semistable by the following result. When  $X$  is smooth, by the result of Dervan and Székelyhidi [31] the existence of such special minimizing valuation is implied by the work of Chen and Wang [29] and Chen, Sun and Wang [28]. We also note that optimal degenerations (of various kinds) in the toric case are well studied; see Wang and Zhu [75] for the toric result for Kähler–Ricci flow.

**Theorem 1.6** (Theorem 5.2) A special  $\mathbb{R}$ -test configuration minimizes  $H^{\text{NA}}$  if and only if its central fiber is  $K$ -semistable.

The uniqueness in Theorem 1.2 about the semistable degeneration is nothing but the result on the uniqueness of the minimizer of  $\tilde{\beta}$  among all quasimonomial valuations associated to special  $\mathbb{R}$ -test configurations. The proof of this fact uses the technique of initial term degeneration, again motivated by study of normalized volumes; see Li [50] and Li and Xu [58; 57]. This process essentially reduces the question to the uniqueness of minimizer of  $H^{\text{NA}}$  (actually a variant of  $H^{\text{NA}}$  after the work of Xu and Zhuang [79]) along an interpolation between a fixed filtration and a weight filtration (induced by a holomorphic vector field) on the central fiber. The interpolation is constructed by using the rescaling of

<sup>2</sup>This has recently been confirmed by Blum, Liu, Xu and Zhuang [15].

twist of the fixed filtration (the twist of filtration is in the sense of Li [52] generalizing Hisamoto [41]), which we can deal with using the technique of Newton–Okounkov bodies and Boucksom and Jonsson’s work on the characterization of asymptotically equivalent filtrations. Our valuative formulation is useful because filtrations associated to valuations are asymptotically equivalent if and only if they are the same (Corollary 2.28 and Lemma 2.29). Again unlike the case of normalized volumes or the case of the  $\delta$  invariant in Blum, Liu and Zhou [16], the minimizing valuation in the current global setting is expected to be absolutely unique, not just up to rescaling or twisting. This is because of a strict convex property of the  $H^{\text{NA}}$ -functional, which goes back to Tian and Zhu’s work in [71] on the uniqueness of Kähler–Ricci vector fields from the Lie algebra of a torus.

To deal with the polystable step, we first introduce the equivariant version of normalized volumes. Most results about normalized volumes can be generalized for the equivariant version. Finally we complete the proof of Theorem 1.3 by adapting the argument in Li, Wang and Xu [55] about uniqueness of K-polystable degeneration of K-semistable Fano varieties.

To end this introduction, the following table summarizes the quantities used in each of the two steps:

degenerations	semistable	polystable
valuations	$\text{Val}(X)$	$\text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}}$
antiderivative	$H^{\text{NA}}, \tilde{\beta}$	$\widehat{\text{vol}}_g$
derivative	$D_\xi^{\text{NA}}, \text{Fut}_\xi$	$D_\xi^{\text{NA}}, \beta_g$
derivative formula	(172)	(191)

**Postscript** After we finished the paper, we were informed by F Wang and X Zhu that they use analytic methods to prove related uniqueness results for Kähler–Ricci flow limits based on their recent work on the Hamilton–Tian conjecture; see [73; 74].

After this paper appeared, there have been several important developments. In Li and Li [59], based on the minimization setup and uniqueness results in this paper, the authors calculated nontrivial examples of limits of Kähler–Ricci flows on some unstable Fano varieties which are compactifications of homogeneous varieties under some complex reductive group. Very recently, the paper of Blum, Liu, Xu and Zhuang [15] continues and completes the algebraic study of minimization problem proposed in this paper, based on the recent breakthrough on the high-rank finite generation conjecture. In particular, Conjecture 4.11 in our paper is now confirmed in [15].

**Acknowledgements** Han would like to thank Jeff Viaclovsky for his teaching and support over many years. Li is partially supported by NSF (grant DMS-1810867) and an Alfred P Sloan research fellowship. We would like to thank G Tian and X Zhu for their interest and comments, and F Wang and X Zhu for informing us of their work. We also thank M Jonsson and C Xu for helpful comments. We would like to thank a referee for the careful reading and useful suggestions for improving the paper.

## 2 Preliminaries

### 2.1 Some notation

Let  $X$  be a  $\mathbb{Q}$ -Fano variety. In this paper for the simplicity of notation, we assume that  $-K_X$  is Cartier. The modification to the general  $\mathbb{Q}$ -Cartier case is straightforward; see eg [52]. For any  $m, \ell \in \mathbb{N}$ , set

$$(3) \quad R_m := H^0(X, -mK_X), \quad R = \bigoplus_{m=0}^{+\infty} R_m,$$

$$(4) \quad N_m = \dim R_m, \quad V = (-K_X)^n = \lim_{m \rightarrow +\infty} \frac{N_m}{m^n/n!},$$

$$(5) \quad R_m^{(\ell)} := H^0(X, -m\ell K_X), \quad R^{(\ell)} = \bigoplus_{m=0}^{+\infty} R_m^{(\ell)}.$$

We will denote by  $\text{Val}(X)$  the space of real valuations on  $\mathbb{C}(X)$ , by  $\overset{\circ}{\text{Val}}(X)$  the set of real valuations  $v$  with  $A_X(v) < +\infty$ , and by  $X_{\mathbb{Q}}^{\text{div}}$  the set of divisorial valuations, ie the valuations of the form  $a \cdot \text{ord}_E$  with  $a \geq 0$  and  $E$  a prime divisor over  $X$ . A valuation  $v \in \text{Val}(X)$  is quasimonomial if there exist a birational morphism  $Y \rightarrow X$  and a simple normal crossing divisors  $E = \bigcup_{i=1}^d E_i \subset Y$  such that  $v$  is a monomial valuation on  $Y$  with respect to the local coordinates defining  $E_i$ , whose center of  $v$  over  $Y$  is an irreducible component of  $\bigcap_{i \in J} E_i$ , where  $J \subseteq \{1, \dots, d\}$  is a subset. We denote by  $\text{QM}(Y, E)$  the set of such quasimonomial valuations. We refer to [35; 44] for more details about such quasimonomial (or equivalently the Abhyankar) valuations.

In this paper,  $\mathbb{T}$  denotes a complex torus  $(\mathbb{C}^*)^r = ((S^1)^r)_{\mathbb{C}}$  that acts effectively on a  $\mathbb{Q}$ -Fano variety  $X$ . There is a canonical action of  $\mathbb{T}$  on (any multiple of)  $-K_X$ . Set

$$(6) \quad N_{\mathbb{Z}} = \text{Hom}(\mathbb{C}^*, \mathbb{T}), \quad N_{\mathbb{R}} = N_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{R}, \quad M_{\mathbb{Z}} = \text{Hom}(\mathbb{T}, \mathbb{C}^*), \quad M_{\mathbb{R}} = M_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{R}.$$

For any  $\xi \in N_{\mathbb{R}}$ , we have a valuation  $\text{wt}_{\xi} \in \text{Val}(X)$  as follows. For any  $f \in \mathbb{C}(X) = \bigoplus_{\alpha \in M_{\mathbb{Z}}} \mathbb{C}(X)_{\alpha}$ ,

$$(7) \quad \text{wt}_{\xi}(f) = \min \left\{ \langle \alpha, \xi \rangle \mid f = \sum_{\alpha} f_{\alpha}, f_{\alpha} \neq 0 \right\}.$$

Moreover, for any  $m \in \mathbb{N}$ , we have a weight decomposition induced by the canonical  $\mathbb{T}$ -action on  $(X, -mK_X)$ :

$$(8) \quad R_m = \bigoplus_{\alpha \in M_{\mathbb{Z}}} (R_m)_{\alpha} = (R_m)_{\alpha_1^{(m)}} \oplus \cdots \oplus (R_m)_{\alpha_{N_m}^{(m)}}.$$

Moreover, we will use the following notation for any  $\mathbb{Q}$ -Fano variety. Let  $e^{-\tilde{\varphi}}$  be an  $(S^1)^r$ -invariant smooth positively curved Hermitian metric on  $-K_X$  (eg as the restriction of a Fubini–Study metric under

an equivariant embedding of  $X$  into projective space). We identify any  $\eta \in N_{\mathbb{R}}$  with the corresponding holomorphic vector field on  $X$ . Because  $\mathbb{T}$ –action canonically lifts to an action on  $-K_X$ , we can set

$$(9) \quad \theta_{\tilde{\varphi}}(\eta) = -\frac{\mathfrak{L}_{\eta}e^{-\tilde{\varphi}}}{e^{-\tilde{\varphi}}}.$$

Then  $\theta_{\tilde{\varphi}}(\eta)$  is a Hamiltonian function of  $\eta$  with respect to  $dd^c\tilde{\varphi} = (\sqrt{-1}/2\pi)\partial\bar{\partial}\tilde{\varphi} \geq 0$ :

$$(10) \quad \iota_{\eta}dd^c\tilde{\varphi} = \frac{\sqrt{-1}}{2\pi}\partial\theta_{\tilde{\varphi}}(\eta).$$

Moreover,  $(z, \eta) \mapsto \theta_{\tilde{\varphi}}(\eta)(z)$  is equivalent to the moment map  $\mathbf{m}_{\tilde{\varphi}}: X \rightarrow M_{\mathbb{R}}$  whose image is the moment polytope  $P$  of  $\mathbb{T}$ –action on  $(X, -K_X)$  which does not depend on the choice of  $\tilde{\varphi}$ . It is known that the measure

$$(11) \quad \frac{n!}{m^n} \sum_i \dim(R_m)_{\alpha_i^{(m)}} \cdot \delta_{\alpha_i^{(m)}/m}$$

converges weakly to the Duistermaat–Heckman measure  $(\mathbf{m}_{\tilde{\varphi}})_*(dd^c\tilde{\varphi})^n$ ; see [23] or [8, Proposition 4.1].

For any subset  $S \subseteq \mathbb{R}^n$ , we will use  $dy_S$  or just  $dy$  to denote the Lebesgue measure of  $S$ .

## 2.2 $\mathbb{R}$ –test configuration and filtrations

We will use extensively the language of filtrations:

**Definition 2.1** [17] A filtration  $\mathcal{F} := \mathcal{F}R_{\bullet}$  of the graded  $\mathbb{C}$ –algebra  $R = \bigoplus_{m=0}^{+\infty} R_m$  consists of a family of subspaces  $\{\mathcal{F}^{\lambda}R_m\}_x$  of  $R_m$  for each  $m \geq 0$  with the following properties:

- **Decreasing**  $\mathcal{F}^{\lambda}R_m \subseteq \mathcal{F}^{\lambda'}R_m$  if  $\lambda \geq \lambda'$ .
- **Left continuous**  $\mathcal{F}^{\lambda}R_m = \bigcap_{\lambda' < \lambda} \mathcal{F}^{\lambda'}R_m$ .
- **Multiplicative**  $\mathcal{F}^{\lambda}R_m \cdot \mathcal{F}^{\lambda'}R_{m'} \subseteq \mathcal{F}^{\lambda+\lambda'}R_{m+m'}$  for any  $\lambda, \lambda' \in \mathbb{R}$  and  $m, m' \in \mathbb{Z}_{\geq 0}$ .
- **Linearly bounded** There exist  $e_-, e_+ \in \mathbb{Z}$  such that  $\mathcal{F}^{m e_-}R_m = R_m$  and  $\mathcal{F}^{m e_+}R_m = 0$  for all  $m \in \mathbb{Z}_{\geq 0}$ .

Similarly one defines filtration on  $R^{(\ell)}$  for any  $\ell \geq 1 \in \mathbb{N}$ .

**Example 2.2** Given any valuation  $v \in \mathring{\text{Val}}(X)$ , we have an associated filtration  $\mathcal{F} = \mathcal{F}_v$ :

$$(12) \quad \mathcal{F}_v^{\lambda}R_m := \{s \in R_m \mid v(s) \geq \lambda\}.$$

In particular, if there is a  $\mathbb{T}$ –action on  $X$ , for any  $\xi \in N_{\mathbb{R}}$ , we have a filtration  $\mathcal{F}_{\text{wt}_{\xi}}$  associated to the valuation  $\text{wt}_{\xi}$  in (7).

The trivial filtration  $\mathcal{F}_{\text{triv}}$  is the filtration associated to the trivial valuation:  $\mathcal{F}_{\text{triv}}^x R_m$  is equal to  $R_m$  if  $x \leq 0$ , and is equal to 0 if  $x > 0$ .



**Example 2.3** For any filtration  $\mathcal{F}$ , we will denote by  $\mathcal{F}_{\mathbb{Z}}$  the filtration defined by  $\mathcal{F}_{\mathbb{Z}}^\lambda R_m = \mathcal{F}^{\lceil \lambda \rceil} R_m$ .

**Definition 2.4** [45; 46; 49] We say a valuation  $\mathfrak{v} : \mathbb{C}(X) \rightarrow \mathbb{Z}^n$  (where  $\mathbb{Z}^n$  is ordered lexicographically) is a faithful valuation if  $\mathfrak{v}(\mathbb{C}(X)) \cong \mathbb{Z}^n$ . Note that such a valuation always has at most one-dimensional leaves (in the sense of [45]): if  $\mathfrak{v}(f) = \mathfrak{v}(g)$  for  $f, g \in \mathbb{C}(X)$ , then there exists  $c \in \mathbb{C}^*$  satisfying  $\mathfrak{v}(f + cg) > \mathfrak{v}(f)$ .

Fix such a faithful valuation  $\mathfrak{v}$ . For any  $t \in \mathbb{R}$ , define the Newton–Okounkov body of the graded linear series

$$(13) \quad \mathcal{F}^{(t)} := \mathcal{F}^{(t)} R_\bullet := \{\mathcal{F}^{tm} R_m\}$$

as the closed convex hull of unions of rescaled values of elements from  $\mathcal{F}^{(t)}$ :

$$(14) \quad \Delta(\mathcal{F}^{(t)}) = \overline{\bigcup_{m=1}^{+\infty} \frac{1}{m} \mathfrak{v}(\mathcal{F}^{tm} R_m)}.$$

By the theory of Newton–Okounkov bodies [62; 49; 46], we know that

$$(15) \quad n! \cdot \text{vol}(\Delta(\mathcal{F}^{(t)})) = \text{vol}(\mathcal{F}^{(t)} R_\bullet) = \lim_{m \rightarrow +\infty} \frac{\dim_{\mathbb{C}} \mathcal{F}^{mt} R_m}{m^n/n!}.$$

When  $t \ll 0$ ,

$$(16) \quad \Delta(\mathcal{F}^{(t)}) =: \Delta_{\mathfrak{v}}(X, -K_X) = \Delta(X)$$

is associated to the complete graded linear series  $\{R_m\}_m$ . Following [17], define the concave transform

$$(17) \quad G^{\mathcal{F}} : \Delta(X) \rightarrow \mathbb{R}, \quad G^{\mathcal{F}}(y) = \sup\{t \mid y \in \Delta(\mathcal{F}^{(t)})\}.$$

Given any filtration  $\mathcal{F} = \{\mathcal{F}^\lambda R_m\}_{\lambda \in \mathbb{R}}$  and  $m \in \mathbb{Z}_{\geq 0}$ , the successive minima on  $R_m$  is the decreasing sequence

$$\lambda_{\max}^{(m)} = \lambda_1^{(m)} \geq \dots \geq \lambda_{N_m}^{(m)} = \lambda_{\min}^{(m)}$$

defined by

$$\lambda_j^{(m)} = \max\{\lambda \in \mathbb{R} \mid \dim_{\mathbb{C}} \mathcal{F}^\lambda R_m \geq j\}.$$

**Theorem 2.5** [17] (i) The function  $x \mapsto \text{vol}(\mathcal{F}^{(x)} R_\bullet)^{1/n}$  is concave on  $(-\infty, \lambda_{\max})$  and vanishes on  $(\lambda_{\max}, +\infty)$ .

(ii) As  $m \rightarrow +\infty$ , the Dirac-type measure

$$(18) \quad \nu_m = \frac{n!}{m^n} \sum_i \delta_{\lambda_i^{(m)}/m} = -\frac{d}{dt} \frac{\dim_{\mathbb{C}} \mathcal{F}^{mt} H^0(Z, m\ell_0 L)}{m^n/n!}$$

converges weakly to a measure with total mass  $V = (-K_X)^n$ :

$$(19) \quad \text{DH}(\mathcal{F}) := n! \cdot (G^{\mathcal{F}})_* dy = -d\text{vol}(\mathcal{F}^{(t)}),$$

where  $dy$  is the Lebesgue measure on  $\Delta(X)$ .

(iii) The support of the measure  $\text{DH}(\mathcal{F})$  is given by  $\text{supp}(\text{DH}(\mathcal{F})) = [\lambda_{\min}, \lambda_{\max}]$ , with

$$(20) \quad \lambda_{\min} := \lambda_{\min}(\mathcal{F}) := \inf\{t \in \mathbb{R} \mid \text{vol}(\mathcal{F}^{(t)}) < V\},$$

$$(21) \quad \lambda_{\max} := \lambda_{\max}(\mathcal{F}) := \lim_{m \rightarrow +\infty} \frac{\lambda_{\max}^{(m)}}{m} = \sup_{m \geq 1} \frac{\lambda_{\max}^{(m)}}{m}.$$

Moreover,  $\text{DH}(\mathcal{F})$  is absolutely continuous with respect to the Lebesgue measure, except perhaps for a point mass at  $\lambda_{\max}$ .

**Example 2.6** If  $v \in \text{Val}(X)$  is quasimonomial, it is shown in [22] that  $\text{DH}$  is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}$ , ie there is no Dirac mass at  $\lambda_{\max}(\mathcal{F}_v)$ .

**Definition 2.7** Let  $\mathcal{F}$  be any filtration. For any  $a > 0$  the  $a$ -rescaling of  $\mathcal{F}$  is given by

$$(22) \quad (a\mathcal{F})^\lambda R_m = \mathcal{F}^{\lambda/a} R_m.$$

For any  $b \in \mathbb{R}$ , the  $b$ -shift is given by

$$(23) \quad \mathcal{F}(b)^\lambda R_m = \mathcal{F}^{\lambda-bm} R_m.$$

Set

$$(24) \quad a\mathcal{F}(b) = (a\mathcal{F})(b) = a(\mathcal{F}(b/a)), \quad \text{ie } a\mathcal{F}(b)^x R_m = \mathcal{F}^{(x-bm)/a} R_m.$$

We have the easy identities

$$(25) \quad \Delta(a\mathcal{F}(b)^{(t)}) = \Delta(\mathcal{F}^{((t-b)/a)}), \quad G_{a\mathcal{F}(b)} = aG_{\mathcal{F}} + b, \quad \text{vol}(a\mathcal{F}(b)^{(t)}) = \text{vol}(\mathcal{F}^{((t-b)/a)}).$$

For any  $f_m \in R_m$ , set

$$(26) \quad \bar{v}_{\mathcal{F}}(f_m) = \sup\{\lambda \mid f_m \in \mathcal{F}^\lambda R_m\} = \max\{\lambda; f_m \in \mathcal{F}^\lambda R_m\},$$

and for any  $f = \sum_m f_m \in R = \bigoplus_m R_m$  with  $f_m \in R_m$ , set

$$(27) \quad \bar{v}_{\mathcal{F}}\left(\sum_m f_m\right) = \min\{\bar{v}_{\mathcal{F}}(f_m) \mid f_m \neq 0 \in R_m\}.$$

Then  $\bar{v}_{\mathcal{F}}$  is a semivaluation on  $R = \bigoplus_m R_m$ , satisfying

$$(28) \quad \bar{v}_{\mathcal{F}}(f + g) \geq \min\{\bar{v}_{\mathcal{F}}(f), \bar{v}_{\mathcal{F}}(g)\} \quad \text{and} \quad \bar{v}_{\mathcal{F}}(fg) \geq \bar{v}_{\mathcal{F}}(f) \cdot \bar{v}_{\mathcal{F}}(g).$$

Set

$$(29) \quad \Gamma^+(\mathcal{F}) := \{\lambda_i^{(m)} \mid m \geq 0, 1 \leq i \leq N_m\}.$$

Denote by  $\Gamma(\mathcal{F})$  the group of  $\mathbb{R}$  generated by  $\Gamma^+(\mathcal{F})$ .

**Definition 2.8** • The extended Rees algebra and associated graded algebra of a filtration  $\mathcal{F}$  are defined as

$$(30) \quad \mathcal{R}(\mathcal{F}) = \bigoplus_{m \geq 0} \bigoplus_{\lambda \in \Gamma(\mathcal{F})} t^{-\lambda} \mathcal{F}^\lambda R_m,$$

$$(31) \quad \text{Gr}(\mathcal{F}) = \bigoplus_{m \geq 0} \bigoplus_{\lambda \in \Gamma(\mathcal{F})} t^{-\lambda} \mathcal{F}^\lambda R_m / \mathcal{F}^{>\lambda} R_m,$$

where  $\mathcal{F}^{>\lambda} R_m = \{f \in R_m \mid v_{\mathcal{F}}(f) > \lambda\}$ .

- If  $\mathcal{R}(\mathcal{F})$  is finitely generated, we say that  $\mathcal{F}$  is finitely generated and call  $\mathcal{F}$  an  $\mathbb{R}$ -test configuration. In this case,  $\Gamma(\mathcal{F})$  is a finitely generated free Abelian group:  $\Gamma(\mathcal{F}) \cong \mathbb{Z}^{\text{rk}(\mathcal{F})}$  for some positive integer  $\text{rk}(\mathcal{F}) \in \mathbb{Z}_{>0}$ , and we will call  $\text{rk}(\mathcal{F})$  the rank of  $\mathcal{F}$ . Moreover,  $\text{Gr}(\mathcal{F})$  is also finitely generated, and we call the projective scheme  $\text{Proj}(\text{Gr}(\mathcal{F})) =: X_{\mathcal{F},0}$  the central fiber of  $\mathcal{F}$ .

There is an induced filtration  $\mathcal{F}|_{X_{\mathcal{F},0}} := \mathcal{F}'R' = \{\mathcal{F}'R'_m\}$  on  $R' := \text{Gr}(\mathcal{F})$ , the homogeneous coordinate ring of the central fiber:

$$(32) \quad \mathcal{F}'^\lambda R'_m = \bigoplus_{\lambda_i^{(m)} \geq \lambda} \mathcal{F}^{\lambda_i^{(m)}} R_m / \mathcal{F}^{>\lambda_i^{(m)}} R_m.$$

The  $\Gamma(\mathcal{F})$  grading of  $\text{Gr}(\mathcal{F})$  corresponds to a holomorphic vector field  $\eta = \eta_{\mathcal{F}}$  on the central fiber, which generates an action by a complex torus of dimension  $\text{rk}(\mathcal{F})$ .

- We say an  $\mathbb{R}$ -test configuration  $\mathcal{F}$  is special if its central fiber  $X_{\mathcal{F},0}$  is a  $\mathbb{Q}$ -Fano variety and there is an isomorphism  $\text{Gr}(\mathcal{F}) \cong R(X_{\mathcal{F},0}, -K_{X_{\mathcal{F},0}}) =: R'$ . In this case, there is a  $\sigma \in \mathbb{R}$  such that

$$(33) \quad \mathcal{F}'R' = \mathcal{F}'_{\text{wt}_\eta} R'(-\sigma).$$

**Remark 2.9** We can naturally extend the above definition to filtrations on  $R^{(\ell)}$  for any  $\ell \in \mathbb{N}_{\geq 1}$ . Indeed we will actually identify two filtrations if they induce the same non-Archimedean metric on  $(X^{\text{NA}}, L^{\text{NA}})$  with  $L = -K_X$ . See [Definition 2.17](#).

There are two equivalent geometric descriptions of  $\mathbb{R}$ -test configurations, which we now explain.

- (I) **Geometric  $\mathbb{R}$ -TC I** Let  $t: X \rightarrow \mathbb{P}^{N_\ell}$  be a Kodaira embedding by a basis of  $R_\ell = H^0(X, \ell(-K_X))$  for some  $\ell > 0$ , and let  $\eta$  be a holomorphic vector field on  $\mathbb{P}^{N_\ell-1} = \mathbb{P}(H^0(X, \ell(-K_X))^*)$  that generates an effective holomorphic action on  $\mathbb{P}^{N_\ell-1}$  by a torus  $\mathbb{T}$  of rank  $r$ . Then we get a weight decomposition  $R_\ell = \bigoplus_{\alpha \in \mathbb{Z}^r} R_{\ell,\alpha}$  and a filtration on  $R_\ell$  by setting

$$(34) \quad \mathcal{F}^\lambda R_\ell = \bigoplus_{\langle \alpha, \eta \rangle \geq \lambda} R_{\ell,\alpha}.$$

The filtration  $\mathcal{F}R_\ell$  generates a filtration on  $\mathcal{F}R^{(\ell)}$ , which is an  $\mathbb{R}$ -test configuration  $\mathcal{F}$ . The following lemma generalizes the well-known fact for test configurations; see [\[33; 76; 64\]](#).

**Lemma 2.10** Any  $\mathbb{R}$ –test configuration, which by definition is a finitely generated filtration, is obtained in this way.

**Proof** To see this, we assume again that  $\mathcal{F}$  is generated by  $\mathcal{F}R_\ell$ . For simplicity of notation, set  $V = R_\ell$  and  $\lambda_i = \lambda_i^{(\ell)}$ . By shifting the filtration, we can normalize  $\lambda_{N_\ell} = 0$  and assume that we have the relation

$$\begin{aligned} \lambda_1 = \dots = \lambda_{i_1} &=: w_1 > \lambda_{i_1+1} = \dots = \lambda_{i_2} &=: w_2 \\ &\vdots \\ &> \lambda_{i_{k-2}+1} = \dots = \lambda_{i_{k-1}} &=: w_{k-1} \\ &> \lambda_{i_{k-1}+1} = \dots = \lambda_{N_\ell} &=: w_k = 0. \end{aligned}$$

In other words,  $\{w_1, \dots, w_k\}$  is the set of distinct values of successive minima and we have a usual filtration,

$$(35) \quad \{0\} \subsetneq \mathcal{F}^{w_1} V \subsetneq \mathcal{F}^{w_2} V \subsetneq \dots \subsetneq \mathcal{F}^{w_k} = V.$$

In other words, we can equivalently describe an  $\mathbb{R}$ –filtration by the language of weighted flags. Fixing a reference Hermitian inner product  $H_0$  on  $V = R_\ell$ , we can assign to the flag (35) a decomposition

$$(36) \quad V = V_1 \oplus V_2 \oplus \dots \oplus V_k,$$

where  $V_1 = \mathcal{F}^{w_1} V$  and  $V_j$  is the  $H_0$ –orthogonal complement of  $\mathcal{F}^{w_{j-1}} V$  inside  $\mathcal{F}^{w_j} V$ , which has dimension  $i_j - i_{j-1} =: d_j$ .

Fix a maximal  $\mathbb{Q}$ –linearly independent subset of  $\{w_1, \dots, w_k\}$  to be

$$(37) \quad 0 > w_2 =: \zeta_1 > \dots > w_{p_r} =: \zeta_r.$$

So for each  $w_j$  we can find a vector of rational numbers  $\vec{r}_j = (r_{j1}, \dots, r_{jr}) \in \mathbb{Q}$  such that  $w_j = \sum_{p=1}^r r_{jp} \zeta_p$ . Finding a common multiple  $D$  of the denominators of  $\{r_{jp} \mid 1 \leq j \leq k, 1 \leq p \leq r\}$ , we set  $\eta = \zeta/D$  and  $\alpha_j = D\vec{r}_j$ , so that

$$(38) \quad w_j = \sum_{p=1}^r \alpha_{jp} \zeta_p = \langle \alpha_j, \eta \rangle.$$

In this way we get a  $(\mathbb{C}^*)^r$  representation  $V$ , whose weight decomposition is given by (36), where  $V_j$  consists of elements of weight  $\alpha_j$ , and

$$\mathcal{F}^\lambda V = \bigoplus_{\langle \alpha_j, \eta \rangle \geq \lambda} V_j = \left\{ v = \sum_{j=1}^k v_j \mid \min\{\langle \alpha_j, \eta \rangle \mid v_j \neq 0\} \geq \lambda \right\}. \quad \square$$

From another point of view, let  $\mathcal{I}_X \subset \mathbb{C}[Z_1, \dots, Z_{N_\ell}] = S$  be the homogeneous ideal of  $X$ . For each  $d \in \mathbb{N}$ , the  $\mathbb{T}$ –action induces a representation of  $\mathbb{T}$  on  $S_d$ , the set of degree- $d$  homogeneous polynomials. The holomorphic vector field  $\eta$  induces an order on the weights of these  $\mathbb{T}$ –representations. Choosing a set of homogeneous generators of  $\mathcal{I}_X$ , the initial term with respect to this order generates the ideal

of  $X_{\mathcal{F},0}$ . If  $\sigma_\eta$  denotes the one-parameter  $\mathbb{R}$ -group generated by  $\eta$ , we have the convergence of algebraic cycles (or schemes)

$$(39) \quad \lim_{s \rightarrow +\infty} \sigma_\eta(s) \circ [X] = [X_{\mathcal{F},0}].$$

So we say that the  $\mathbb{R}$ -action generated by  $\eta$  degenerates  $X$  into a projective scheme  $X_{\mathcal{F},0}$ .

By perturbing  $\eta \in N_{\mathbb{R}}$ , we can find a sequence of rational vector  $\eta_k \in N_{\mathbb{Q}}$  converging to  $\eta$ . For  $k \gg 1$ ,  $\eta_k$  induces an  $\mathbb{R}$ -test configuration of rank one with the same central fiber  $X_{\mathcal{F},0}$ .

(II) **Geometric  $\mathbb{R}$ -TC II** This description is essentially contained in [66, Section 2]. For any  $\mathbb{R}$ -test configuration, we set  $B = \text{Spec}(\mathbb{C}(\Gamma^+(\mathcal{F})) \cong \mathbb{C}^r$ . Then there is a flat family

$$(40) \quad \mathcal{X} = \text{Proj}_{\mathbb{C}^r}(\mathcal{R}(\mathcal{F})) \rightarrow B$$

such that the generic fiber is isomorphic to  $X$  and a special fiber isomorphic to  $X_{\mathcal{F},0}$ . Set  $\mathcal{L}$  to be the relative ample line bundle  $\mathcal{O}_{\mathcal{X}/\mathbb{C}^r}(1)$ . Fix  $m \geq 0$ . For any  $\lambda \in \mathbb{R}$ , we set  $\lceil \lambda \rceil = \min\{\lambda_i^{(m)} \mid \lambda_i^{(m)} \geq \lambda\} = \langle \alpha, \eta_{\mathcal{F}} \rangle$  for  $\alpha \in M_{\mathbb{Z}}$ . Then for any  $\tau = (\tau_1, \dots, \tau_r) \in \mathbb{C}^r$ , we set  $\tau^{-\lceil \lambda \rceil} = \prod_{i=1}^r \tau_i^{\alpha_i}$ , to get

$$(41) \quad \mathcal{F}^\lambda R_m = \{s \in R_m \mid \tau^{-\lceil \lambda \rceil} \bar{s} \text{ extends to a holomorphic section of } m\mathcal{L} \rightarrow \mathcal{X}\},$$

where  $\bar{s}$  is the meromorphic section of  $m\mathcal{L}$  defined as the pullback of  $s$  via the projection  $(\mathcal{X}, \mathcal{L}) \times_B (\mathbb{C}^*)^r \cong (X, -K_X) \times (\mathbb{C}^*)^r \rightarrow X$ .

**Lemma 2.11** *If  $\text{Gr}(\mathcal{F})$  is an integral domain, then the semivaluation  $\bar{v}_{\mathcal{F}}$  in (26) defines a valuation on the quotient field of  $R$ . Denote by  $v_{\mathcal{F}}$  the restriction of  $\bar{v}_{\mathcal{F}}$  to  $\mathbb{C}(X)$ : for  $f = s_1/s_2 \in \mathbb{C}(X)$  with  $s_1, s_2 \in R_m$ , set*

$$(42) \quad v_{\mathcal{F}}(f) = \bar{v}_{\mathcal{F}}(s_1) - \bar{v}_{\mathcal{F}}(s_2).$$

*Then there exists  $\sigma > 0$  such that  $\mathcal{F} = \mathcal{F}_{v_{\mathcal{F}}}(-\sigma)$ . In particular, this statement applies to any special  $\mathbb{R}$ -test configuration.*

**Proof** Fix any two homogeneous elements  $s_i \in R_{m_i}$  for  $i = 1, 2$ . Assume that  $\bar{v}_{\mathcal{F}}(f_i) = s_i$ . Then  $s'_i \in R'_{m_i, x_i}$ . Because  $\text{Gr}(\mathcal{F})$  is integral,  $s'_1 s'_2 \neq 0 \in R'_{m_1+m_2, x_1+x_2}$ , which implies that  $\bar{v}_{\mathcal{F}}(s_1 s_2) = x_1 + x_2 = \bar{v}_{\mathcal{F}}(s_1) + \bar{v}_{\mathcal{F}}(s_2)$ . From this, we easily see that  $\bar{v}_{\mathcal{F}}$  is a real valuation.

Assume  $f = s_1/s_2 = \tilde{s}_1/\tilde{s}_2$ . Then  $s_1 \cdot \tilde{s}_2 = s_2 \cdot \tilde{s}_1$  and hence  $\bar{v}_{\mathcal{F}}(s_1) - \bar{v}_{\mathcal{F}}(s_2) = \bar{v}_{\mathcal{F}}(\tilde{s}_1) - \bar{v}_{\mathcal{F}}(\tilde{s}_2)$ . So  $v_{\mathcal{F}}$  in (42) is well defined.

For any  $s_i \neq 0 \in R_m$  with  $i = 1, 2$ , by construction  $\bar{v}_{\mathcal{F}}(s_1) - v_{\mathcal{F}}(s_1) = \bar{v}_{\mathcal{F}}(s_2) - \bar{v}_{\mathcal{F}}(s_2)$ . This means  $b_m := v_{\mathcal{F}} - \bar{v}_{\mathcal{F}}$  is constant on  $R_m \setminus \{0\}$ . It is easy to see that  $\sigma_{m_1} \sigma_{m_2} = \sigma_{m_1+m_2}$ . So we can set  $\sigma = \sigma_m/m$  to get the conclusion. □

An  $\mathbb{R}$ -test configuration with  $\text{rk}(\mathcal{F}) = 1$  is, up to rescaling, associated to the usual test configuration, a notion that plays a basic role in the subject of  $\mathbb{K}$ -stability.

**Definition 2.12** [67; 32; 56] A test configuration of  $(X, L)$  is a triple  $(\mathcal{X}, \mathcal{L}, \eta)$ , sometimes just denoted by  $(\mathcal{X}, \mathcal{L})$ , that consists of

- a variety  $\mathcal{X}$  admitting a  $\mathbb{C}^*$ -action generated by a holomorphic vector field  $\eta$  and a  $\mathbb{C}^*$ -equivariant morphism  $\pi : \mathcal{X} \rightarrow \mathbb{C}$ , where the action of  $\mathbb{C}^*$  on  $\mathbb{C}$  is given by the standard multiplication generated by  $-t\partial_t$ , and
- a  $\mathbb{C}^*$ -equivariant  $\pi$ -semiample  $\mathbb{Q}$ -Cartier  $\mathbb{Q}$ -divisor  $\mathcal{L}$  on  $\mathcal{X}$  such that there is an  $\mathbb{C}^*$ -equivariant isomorphism  $i_\eta : (\mathcal{X}, \mathcal{L})|_{\pi^{-1}(\mathbb{C} \setminus \{0\})} \cong (X, L) \times \mathbb{C}^*$ .

We denote by  $(\overline{\mathcal{X}}, \overline{\mathcal{L}})$  the natural compactification of  $(\mathcal{X}, \mathcal{L})$  obtained by adding a trivial fiber at infinity using the isomorphism  $i_\eta$ .

$(X_{\mathbb{C}}, (-K_X)_{\mathbb{C}}, \eta_{\text{triv}}) := (X \times \mathbb{C}, -K_X \times \mathbb{C}, -t\partial_t)$  is called the trivial test configuration.  $(\mathcal{X}, \mathcal{L}, \eta)$  is a normal test configuration if  $\mathcal{X}$  is a normal variety.

A normal test configuration  $(\mathcal{X}, \mathcal{L}, \eta)$  is a special test configuration (resp. weakly special) if  $(\mathcal{X}, \mathcal{X}_0)$  is plt (resp. if  $(\mathcal{X}, \mathcal{X}_0)$  is log canonical) and  $\mathcal{L} = -K_{\mathcal{X}} + c\mathcal{X}_0$  for some  $c \in \mathbb{Q}$ . By inversion of adjunction,  $(\mathcal{X}, \mathcal{L}, \eta)$  being special is equivalent to the condition that  $(\mathcal{X}_0, -K_{\mathcal{X}_0})$  is  $\mathbb{Q}$ -Fano.

Two test configurations  $(\mathcal{X}_i, \mathcal{L}_i)$  for  $i = 1, 2$  are equivalent if there exists a test configuration  $(\mathcal{X}', \mathcal{L}')$  and two  $\mathbb{C}^*$ -equivariant birational morphisms  $\rho_i : \mathcal{X}' \rightarrow \mathcal{X}_i$  such that  $\rho_1^* \mathcal{L}_1 = \mathcal{L}' = \rho_2^* \mathcal{L}_2$ .

Assume that  $\mathbb{G}$  is a reductive complex Lie group acting on  $(X, L)$ . A  $\mathbb{G}$ -equivariant test configuration of  $(X, L)$  is a test configuration  $(\mathcal{X}, \mathcal{L}, \eta)$  with the following property:

- There is a  $\mathbb{G}$ -action on  $(\mathcal{X}, \mathcal{L})$  that commutes with the  $\mathbb{C}^*$ -action generated by  $\eta$  and the action of  $\mathbb{G}$  on  $(\mathcal{X}, \mathcal{L}) \times_{\mathbb{C}} \mathbb{C}^* \cong_{i_\eta} (X, L) \times \mathbb{C}^*$  coincides with the fiberwise action of  $\mathbb{G}$  on (the first factor of)  $(X, L) \times \mathbb{C}^*$ .

As mentioned above, by the work of [76; 65; 19], for any  $\mathbb{R}$ -test configuration  $\mathcal{F}$  with  $\text{rk}(\mathcal{F}) = 1$ , there exists a test configuration  $(\mathcal{X}, \mathcal{L}, \eta)$  and  $a > 0$ , such that  $\Gamma(\mathcal{F}) \cong a\mathbb{Z}$  and  $\mathcal{F} = a\mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)}$ . In this case, we will also denote the  $\mathbb{R}$ -test configuration  $\mathcal{F}$  by  $(\mathcal{X}, \mathcal{L}, a\eta)$  and set

$$(43) \quad \mathcal{F}_{(\mathcal{X}, \mathcal{L}, a\eta)} := a\mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)}.$$

The identity (40) becomes

$$(44) \quad \mathcal{X} = \text{Proj}_{\mathbb{C}[t]} \left( \bigoplus_{m \geq 0} \bigoplus_{j \in \mathbb{Z}} t^{-aj} \mathcal{F}^j R_m \right).$$

Conversely, assume  $(\mathcal{X}, \mathcal{L})$  is a test configuration of  $(X, L := -K_X)$ . Then we associate to it a filtration  $\mathcal{F} = \mathcal{F}_{(\mathcal{X}, \mathcal{L})}$  as in (41); so  $s \in \mathcal{F}^\lambda R_m$  if and only if  $t^{-\lceil \lambda \rceil} \bar{s}$  extends to a holomorphic section of  $m\mathcal{L}$ . In particular, such a construction sets up a one-to-one correspondence between test configurations  $(\mathcal{X}, \mathcal{L})$  with ample  $\mathcal{L}$ , and  $\mathbb{R}$ -test configurations  $\mathcal{F}$  with  $\Gamma(\mathcal{F}) \subseteq \mathbb{Z}$ ; see [19, Proposition 2.15].

Now assume that  $(\mathcal{X}, \mathcal{L})$  is normal and there is a  $\mathbb{C}^*$ -equivariant birational morphism  $\rho: \mathcal{X} \rightarrow X_{\mathbb{C}} := X \times \mathbb{C}$ . Write  $\mathcal{L} = \rho^* L_{\mathbb{C}} + D$ , where  $L_{\mathbb{C}} = p_1^* L$ . Then by [19, Lemma 5.17], the filtration  $\mathcal{F}$  has the more explicit description

$$(45) \quad \mathcal{F}^\lambda R_m = \bigcap_E \{s \in H^0(X, mL) \mid r(\text{ord}_E)(s) + m\ell_0 \text{ord}_E(D) \geq xb_E\},$$

where  $E$  runs over the irreducible components of the central fiber  $\mathcal{X}_0$ , and  $b_E = \text{ord}_E(\mathcal{X}_0) = \text{ord}_E(t)$  while  $r(\text{ord}_E)$  denotes the restriction of  $\text{ord}_E$  to  $\mathbb{C}(Z)$  under the inclusion  $\mathbb{C}(Z) \subset \mathbb{C}(X \times \mathbb{C}^*) = \mathbb{C}(\mathcal{X})$ .

When  $\mathcal{F} = \mathcal{F}_{(\mathcal{X}, -K_{\mathcal{X}}, \eta)}$  is associated to a special test configuration, Lemma 2.11 applies. In fact, by [19],  $v_{\mathcal{F}} = v_{\mathcal{X}_0} = r(\text{ord}_{\mathcal{X}_0})$  and by [50],  $\sigma = A_X(v_{\mathcal{X}_0})$ , so  $\mathcal{F}_{(\mathcal{X}, -K_{\mathcal{X}}, \eta)} = \mathcal{F}_{v_{\mathcal{X}_0}}(-A(v_{\mathcal{X}_0}))$ . As a consequence, for any  $a > 0$ , by (24) we have the identity

$$(46) \quad \mathcal{F}_{(\mathcal{X}, -K_{\mathcal{X}}, a\eta)} = \mathcal{F}_{av_{\mathcal{X}_0}}(-A(av_{\mathcal{X}_0})).$$

Note that following Definition 2.8, for any  $a > 0$  we say that  $(\mathcal{X}, \mathcal{L}, a\eta)$  is a special (resp. normal)  $\mathbb{R}$ -test configuration if  $(\mathcal{X}, \mathcal{L}, \eta)$  is a special (resp. normal) test configuration.

Note that we use the negative sign  $-t\partial_t$  in our Definition 2.12. This sign convention will be convenient for our subsequent computations, as illustrated in the following simple example.

**Example 2.13** Consider the product test configuration  $(\mathcal{X}, \mathcal{L})$  of  $(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$  induced by the  $\mathbb{C}^*$ -action

$$t \circ [Z_0, Z_1] = [Z_0, tZ_1].$$

Let  $s_i$  for  $i = 0, 1$  be two holomorphic sections of  $H^0(\mathbb{P}^1, \mathcal{O}(1))$  corresponding to the homogeneous coordinates  $Z_i$  for  $i = 0, 1$ . Then  $t$  acts on the holomorphic sections by  $t \cdot s_0 = s_0$  and  $t \circ s_1 = t^{-1}s_1$ . The corresponding filtration is given by

$$(47) \quad \mathcal{F}^\lambda R_m = \text{Span}\{s_0^{m-i} s_1^i \mid 0 \geq -i \geq \lambda\};$$

cf (34). The natural compactification  $\bar{\mathcal{X}}$  can be identified with the Hirzebruch surface  $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1})$ , and  $\bar{\mathcal{L}}$  is given by  $\mathcal{O}_{\bar{\mathcal{X}}}(D_\infty)$ , where  $D_\infty$  is the divisor at infinity; see [56, Example 3]. The successive minima are given by  $\{\lambda_i^{(m)}\} = \{-m, -m + 1, \dots, 0\}$ . In particular, we have

$$(48) \quad \sum_i \lambda_i^{(m)} = -\frac{1}{2}m^2 - \frac{1}{2}m = \frac{1}{2}\bar{\mathcal{L}}^2 m^2 + \left(\frac{1}{2}K_{\bar{\mathcal{X}}}^{-1} \cdot \bar{\mathcal{L}} - 1\right)m.$$

Moreover,  $\eta = -z \partial/\partial z$ , whose Hamiltonian function is given by  $\theta(\eta) = -|Z_1|^2/(|Z_1|^2 + |Z_2|^2)$ . Note that  $\theta(\eta)_* \omega_{\text{FS}} = dy_{[-1,0]} = \text{DH}(\mathcal{F})$ .

**Example 2.14** If  $\mathcal{F}$  is an  $\mathbb{R}$ -test configuration, then  $a\mathcal{F}(b)$  is an  $\mathbb{R}$ -test configurations for any  $(a, b) \in \mathbb{R}_{>0} \times \mathbb{R}$ .

Assume  $\mathcal{F} = \mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)}$  for a test configuration  $(\mathcal{X}, \mathcal{L}, \eta)$ . Then as mentioned above, for simplicity of notation we will identify  $a\mathcal{F}(b)$  with the data  $(\mathcal{X}, \mathcal{L} + b\mathcal{X}_0, a\eta)$ .

For any  $d > 0 \in \mathbb{N}$ , we can consider the normalization of the base change,

$$(49) \quad (\mathcal{X}, \mathcal{L}, \eta)^{(d)} := ((\mathcal{X}, \mathcal{L}, \eta) \times_{\mathbb{C}, t \rightarrow t^d} \mathbb{C})^{\text{norm}} =: (\mathcal{X}^{(d)}, \mathcal{L}^{(d)}, \eta^{(d)}).$$

On the other hand,  $\mathbb{Z}_d = \langle e^{2\pi\sqrt{-1}/d} \rangle \hookrightarrow \mathbb{C}^*$  naturally acts on the  $(\mathcal{X}, \mathcal{L})$  and we can take a quotient

$$(50) \quad (\mathcal{X}, \mathcal{L}, \eta)/\mathbb{Z}_d = (\mathcal{X}^{(1/d)}, \mathcal{L}^{(1/d)}, \eta^{(1/d)})$$

to get a test configuration with a nonreduced central fiber in general.

With this notation, for any  $a > 0 \in \mathbb{Q}$  we then have the natural identification

$$(51) \quad \mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)^{(a)}} = a \cdot \mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)} = \mathcal{F}_{(\mathcal{X}, \mathcal{L}, a\eta)}.$$

For a filtration  $\mathcal{F}R_\bullet$ , choose  $e_-$  and  $e_+$  as in Definition 2.1. For convenience, we can choose  $e_+ = \lceil \lambda_{\max}(\mathcal{F}R) \rceil \in \mathbb{Z}$ . Set  $e = e_+ - e_-$  and define (fractional) ideals

$$(52) \quad I_{m, \lambda} := I_{m, \lambda}^{\mathcal{F}} := \text{Image}(\mathcal{F}^\lambda R_m \otimes \mathcal{O}_X(-mL) \rightarrow \mathcal{O}_X),$$

$$(53) \quad \tilde{\mathcal{I}}_m := \tilde{\mathcal{I}}_m^{\mathcal{F}} := I_{(m, me_+)}^{\mathcal{F}} t^{-me_+} + I_{(m, me_+-1)}^{\mathcal{F}} t^{1-me_+} + \dots + I_{(m, me_+-1)}^{\mathcal{F}} t^{-me_+-1} + \mathcal{O}_X \cdot t^{-me_-},$$

$$(54) \quad \mathcal{I}_m := \mathcal{I}_m^{\mathcal{F}(e_+)} = \tilde{\mathcal{I}}_m^{\mathcal{F}} \cdot t^{me_+} = I_{(m, me_+)}^{\mathcal{F}} + I_{(m, me_+-1)}^{\mathcal{F}} t^1 + \dots + I_{(m, me_+-1)}^{\mathcal{F}} t^{me_+-1} + (t^{me}) \subseteq \mathcal{O}_{X_{\mathbb{C}}}.$$

**Definition–Proposition 2.15** [36, Lemma 4.6] *With the above notation, for  $m$  sufficiently divisible, define the  $m^{\text{th}}$  approximating test configuration  $(\check{\mathcal{X}}_m^{\mathcal{F}}, \check{\mathcal{L}}_m^{\mathcal{F}})$  as follows:*

(i)  $\check{\mathcal{X}}_m^{\mathcal{F}}$  is the normalization of blowup of  $X \times \mathbb{C}$  along the ideal sheaf  $\mathcal{I}_m^{\mathcal{F}(e_+)}$ .

(ii) The semiample  $\mathbb{Q}$ –divisor is given by

$$(55) \quad \check{\mathcal{L}}_m^{\mathcal{F}} = \pi^*((-K_X) \times \mathbb{C}) - \frac{1}{m} E_m + e_+ \check{\mathcal{X}}_0,$$

where  $E_m$  is the exceptional divisor of the normalized blowup.

For simplicity of notation, we also denote the data by  $(\check{\mathcal{X}}_m, \check{\mathcal{L}}_m)$  if the filtration is clear.

It is easy to see that the filtration  $\mathcal{F}_{(\check{\mathcal{X}}_m, \check{\mathcal{L}}_m)}$  on  $R^{(m)}$  is induced by  $\mathcal{F}_{\mathbb{Z}} R_m$  under the canonical map  $S^k R_m \rightarrow R_{km}$ . By [20, Proof of Theorem 4.13], we have the following approximation result.

**Proposition 2.16** [20, Proof of Theorem 4.13] *With notation as in Definition–Proposition 2.15, the Duistermaat–Heckmann measures  $\text{DH}(\check{\mathcal{X}}_m, \check{\mathcal{L}}_m)$  converge weakly to  $\text{DH}(\mathcal{F})$  as  $m \rightarrow +\infty$ .*

Following Boucksom and Jonsson, it is very convenient to use the non-Archimedean metric defined by filtrations. Any filtration (in the sense of Definition 2.1) defines a non-Archimedean metric on  $L^{\text{NA}} \rightarrow X^{\text{NA}}$ . If we denote by  $\phi_{\text{triv}}$  the non-Archimedean metric associated to the trivial filtration, then any non-Archimedean metric  $\phi$  on  $L^{\text{NA}}$  is represented by the real valued function  $\phi - \phi_{\text{triv}}$  on  $X_{\mathbb{Q}}^{\text{div}}$ .



**Definition 2.17** Let  $\mathcal{F} = \mathcal{F}R_\bullet$  be a filtration. For any  $w \in \mathring{\text{Val}}(X)$ , define the non-Archimedean metric associated to  $\mathcal{F}$  by

$$(56) \quad (\phi_m^{\mathcal{F}} - \phi_{\text{triv}})(w) = -\frac{1}{m}G(w)(\tilde{\mathcal{I}}_m^{\mathcal{F}}) = -\frac{1}{m}G(w)(\mathcal{I}_m^{\mathcal{F}(e_+)}t^{-me_+}) = -\frac{1}{m}G(w)(\mathcal{I}_m^{\mathcal{F}(e_+)}) + e_+,$$

$$(57) \quad (\phi^{\mathcal{F}} - \phi_{\text{triv}})(w) = -G(w)(\tilde{\mathcal{I}}_\bullet^{\mathcal{F}}) = \lim_{m \rightarrow +\infty} \phi_m^{\mathcal{F}}(w).$$

In particular, if  $v \in \mathring{\text{Val}}(Z)$  and  $\mathcal{F} = \mathcal{F}_v$ , then we write  $\phi_v = \phi^{\mathcal{F}_v}$ .

Note that  $\phi_m^{\mathcal{F}} = \phi_{\mathcal{F}(\check{\mathcal{X}}_m, \check{\mathcal{L}}_m)}$  converges to  $\phi$  as  $m \rightarrow +\infty$ . Moreover, if (for simplicity) we assume that  $S^k R_m \rightarrow R_{km}$  is surjective for all  $k, m \geq 1$ , then it is an increasing sequence in the sense that if  $m_1 \mid m_2$ , then  $\phi_{m_1}^{\mathcal{F}} \leq \phi_{m_2}^{\mathcal{F}}$ . If  $\phi^{\mathcal{F}}$  is continuous, then  $\phi_m$  converges to  $\phi$  uniformly by Dini's theorem.

The following transformation rule can be easily verified.

**Lemma 2.18** For any filtration  $\mathcal{F}$  and any  $(a, b) \in \mathbb{R}_{>0} \times \mathbb{R}$  and  $v \in X_{\mathbb{Q}}^{\text{div}}$ ,

$$(58) \quad (\phi_{a\mathcal{F}(b)} - \phi_{\text{triv}})(v) = a(\phi_{\mathcal{F}} - \phi_{\text{triv}})\left(\frac{v}{a}\right) + b.$$

### 2.3 Twist of filtrations

Let  $\mathcal{F} = \mathcal{F}R_\bullet$  be a  $\mathbb{T}$ -equivariant filtration, which means that  $\mathcal{F}^\lambda R_m$  is a  $\mathbb{T}$ -invariant subspace of  $R_m$  for any  $x \in \mathbb{R}$ . For  $\alpha \in M_{\mathbb{Z}} = N_{\mathbb{Z}}^\vee$ , denote the weight space by

$$(59) \quad (R_m)_\alpha = \{s \in R_m \mid \tau \circ s = \tau^\alpha s \text{ for all } \tau \in (\mathbb{C}^*)^r\}.$$

Then we have

$$(60) \quad (\mathcal{F}^\lambda R_m)_\alpha := \{s \in \mathcal{F}^\lambda R_m \mid \tau \circ s = \tau^\alpha s\} = \mathcal{F}^\lambda R_m \cap (R_m)_\alpha,$$

and the decomposition

$$(61) \quad \mathcal{F}^\lambda R_m = \bigoplus_{\alpha \in M_{\mathbb{Z}}} (\mathcal{F}^\lambda R_m)_\alpha.$$

**Definition 2.19** [52] For any  $\xi \in N_{\mathbb{R}}$ , the  $\xi$ -twist of  $\mathcal{F}$  is the filtration  $\mathcal{F}_\xi R_\bullet$  defined by

$$(62) \quad \mathcal{F}_\xi^\lambda R_m = \bigoplus_{\alpha \in M_{\mathbb{Z}}} (\mathcal{F}_\xi^\lambda R_m)_\alpha, \quad \text{where } (\mathcal{F}_\xi^\lambda R_m)_\alpha := (\mathcal{F}^{\lambda - \langle \alpha, \xi \rangle} R_m)_\alpha.$$

**Example 2.20** If  $\mathcal{F}$  is a  $\mathbb{T}$ -equivariant  $\mathbb{R}$ -test configuration, then  $\mathcal{F}_\xi$  is also an  $\mathbb{R}$ -test configuration.

If  $\mathcal{F} = \mathcal{F}_{(\mathcal{X}, \mathcal{L}, a\eta)}$  for a test configuration, then we can identify the data  $\mathcal{F}_\xi$  with the data  $(\mathcal{X}, \mathcal{L}, a\eta + \xi)$ ; see [41]. If  $\xi \in N_{\mathbb{Z}}$ , then  $(\mathcal{X}, \mathcal{L}, a\eta + \xi)$  is equivalent to the birational image of the  $(\mathcal{X}, \mathcal{L})$  via the birational transform  $\sigma_\xi: \mathcal{X} \dashrightarrow \mathcal{X}, (z, t) \rightarrow (\sigma_\xi(t) \cdot z, t)$ ; see [52].

Moreover, if we start with the trivial filtration  $\mathcal{F}_{\text{triv}} = \mathcal{F}_{(X_{\mathbb{C}}, (-K_X)_{\mathbb{C}}, -t\partial_t)}$ , then  $(\mathcal{F}_{\text{triv}})_\xi$  is equal to  $\mathcal{F}_{\text{wt}_\xi}$ .

**Definition 2.21** We say that a faithful valuation  $\mathfrak{v}$  in the sense of Definition 2.4 is adapted to the torus action if for any  $f \in \mathbb{C}(X)_\alpha$  we have  $\mathfrak{v}(f) = (\alpha, \mathfrak{v}^{r+1}(f), \dots, \mathfrak{v}^n(f)) \in \mathbb{Z}^r \times \mathbb{Z}^{n-r}$ .

There always exists a faithful valuation that is adapted to the torus action. This can be constructed as follows. First we choose a  $\mathbb{T}$ -invariant Zariski-open set  $U$  of  $X$  as in [3]. Then by the theory of affine  $T$ -varieties as developed in [2], there exists a variety  $Y$  of dimension  $n - r$  and a polyhedral divisor  $\mathcal{D}$  such that

$$(63) \quad U = \text{Spec}_{\alpha \in M_{\mathbb{Z}}} H^0(Y, \mathcal{O}(\mathcal{D}(\alpha))).$$

We can choose a faithful valuation  $\mathfrak{v}_Y$  on  $Y$  (for example via a flag of varieties as in [49]) and define, for any  $f \in H^0(Y, \mathcal{O}(\mathcal{D}(\alpha)))$ ,

$$(64) \quad \mathfrak{v}(f) = (\alpha, \mathfrak{v}_Y(f)).$$

Let  $\mathfrak{v}$  be such a valuation and  $\Delta = \Delta_{\mathfrak{v}}(X, -K_X) \subset \mathbb{R}^n$  be the associated Newton–Okounkov body. If  $p: \mathbb{R}^n = \mathbb{R}^r \times \mathbb{R}^{n-r} \rightarrow \mathbb{R}^r$  denotes the natural projection, then we have

$$(65) \quad p(\Delta) = P = \text{moment map of the } \mathbb{T}\text{-action on } (X, -K_X).$$

The following lemma was already observed in [81], in which a faithful valuation adapted to the torus action was constructed using equivariant infinitesimal flags in the sense of [49]. Here we give a different and direct proof for the reader’s convenience.

For simplicity of notation, we write  $y = (y_1, \dots, y_n) = (y', y'') \in \mathbb{R}^r \times \mathbb{R}^{n-r}$  and set

$$(66) \quad \langle y', \xi \rangle = \sum_{i=1}^r y'_i \xi^i =: \langle y, \xi \rangle.$$

In the last identity, we identify  $\xi \in N_{\mathbb{R}} = \mathbb{R}^r$  with  $(\xi, 0) \in \mathbb{R}^n$ .

**Lemma 2.22** [81] *If  $\mathfrak{v}$  is a  $\mathbb{Z}^n$ -valued valuation adapted to the torus action, then for any  $y \in \Delta(-K_X)$ ,*

$$(67) \quad G_{\mathcal{F}_\xi}(y) = G_{\mathcal{F}}(y) + \langle y', \xi \rangle.$$

**Proof** For any  $t > G_{\mathcal{F}}(y) = \lambda$ , there exists  $\epsilon > 0$  such that  $y \notin \Delta(\mathcal{F}^{(t-\epsilon)})$ . Let  $\delta_1 = \text{dist}(y, \Delta(\mathcal{F}^{(t-\epsilon)}))$ .

Choose any  $f \in \mathcal{F}_\xi^{(t+\langle y', \xi \rangle)m} R_{m,\alpha} = \mathcal{F}^{(t+\langle y', \xi \rangle)m - \langle \alpha, \xi \rangle} R_{m,\alpha}$ . Consider two cases:

- (i)  $\langle \alpha/m, \xi \rangle - \langle y', \xi \rangle < \epsilon$ . Then  $\mathfrak{v}(f) \in \Delta^{(t-\epsilon)}$ , so  $|\mathfrak{v}(f)/m - y| \geq \delta_1$ .
- (ii)  $\langle \alpha/m, \xi \rangle - \langle y', \xi \rangle \geq \epsilon$ . Then  $|\mathfrak{v}(f)/m - y| \geq |\alpha/m - y'| \geq \epsilon/|\xi|$ .

The two cases together imply that  $y \notin \Delta(\mathcal{F}_\xi^{(t+\langle y', \xi \rangle)})$ . So we get the inequality  $G_{\mathcal{F}_\xi} \leq G_{\mathcal{F}} + \langle y', \xi \rangle$ .

On the other hand, since  $\mathcal{F} = (\mathcal{F}_\xi)_{-\xi}$ , we also get  $G_{\mathcal{F}} \leq G_{\mathcal{F}_\xi} - \langle y', \xi \rangle$ . So we get the desired identity.  $\square$

## 2.4 Asymptotically equivalent filtrations

In this section we recall Boucksom and Jonsson’s characterization in [20; 21] of asymptotically equivalent filtrations; see also [1].

For a filtration  $\mathcal{F}R_m$  of  $R_m$ , we say that a basis  $\mathcal{B} = \{s_1, \dots, s_{N_m}\}$  of  $R_m$  is compatible with  $\mathcal{F}R_m$  if for any  $\lambda \in \mathbb{R}$  there exists a subset of  $\mathcal{B}$  that spans  $\mathcal{F}^\lambda R_m$ .

Let  $\mathcal{F}_i = \{\mathcal{F}_i R_m\}$  for  $i = 0, 1$  be two filtrations. For each  $m$ , we can find a basis  $\mathcal{B} := \{s_1, \dots, s_{N_m}\}$  of  $R_m$  that is compatible with both  $\mathcal{F}_i R_m$  with  $i = 0, 1$ . We refer to [1; 18] and the discussion in Section 5 for more details. Assume that for each  $i = 0, 1$ , we have  $s_k \in \mathcal{F}_i^{\mu_{k,i}} \setminus \mathcal{F}^{>\mu_{k,i}}$ . Then  $\mathcal{B}$  is an orthogonal basis for the non-Archimedean norm  $\|\cdot\|_{m,i}$  corresponding to  $\mathcal{F}_i$ : for any  $s = \sum_k a_k s_k \in R_m$ ,

$$(68) \quad \|s\|_{m,i} = e^{-\max\{\lambda | s \in \mathcal{F}^\lambda R_m\}} = \max_k |a_k|_0 e^{-\mu_{k,i}},$$

where  $|\cdot|_0$  is the trivial norm on  $\mathbb{C}$ .

Following [24; 20], we define the set of successive minima of  $\mathcal{F}_1$  with respect to  $\mathcal{F}_0$  to be the set  $\{\mu_{k,1} - \mu_{k,0}\}$ . The following result was proved in [18; 24].

**Theorem 2.23** [18; 24] *As  $m \rightarrow +\infty$ , the measures*

$$(69) \quad \frac{n!}{m^n} \sum_{k=1}^{N_m} \delta_{(\mu_{k,1} - \mu_{k,0})/m}$$

*converge weakly as  $m \rightarrow +\infty$  to a relative limit measure, denoted by  $d\nu := d\nu(\mathcal{F}_0, \mathcal{F}_1)$ .*

**Corollary 2.24** *For any  $p \in [1, \infty)$ , the limit*

$$(70) \quad d_p(\mathcal{F}_0, \mathcal{F}_1) := \lim_{m \rightarrow +\infty} \left( \frac{n!}{m^n} \sum_{k=1}^{N_m} m^{-1} |\mu_{k,1} - \mu_{k,0}|^p \right)^{1/p}$$

*exists and is given by*

$$(71) \quad d_p(\mathcal{F}_0, \mathcal{F}_1) = \left( \int_{\mathbb{R}} |\lambda|^p d\nu(\lambda) \right)^{1/p}.$$

**Definition 2.25** [20, Section 3.6]  $\mathcal{F}_0$  and  $\mathcal{F}_1$  are asymptotically equivalent if  $d_2(\mathcal{F}_0, \mathcal{F}_1) = 0$ .

In fact, by [20] the  $d_p$  are comparable to each other for all  $p \in [1, \infty)$ , and the above equivalence can be defined by using any  $p \in [1, +\infty)$ .

**Theorem 2.26** [20, Theorem 4.16] *Assume that  $X$  is smooth. Let  $\mathcal{F}_0$  and  $\mathcal{F}_1$  be two filtrations on  $R$ . Then  $\mathcal{F}_0$  and  $\mathcal{F}_1$  are asymptotically equivalent if and only if  $\phi_{\mathcal{F}_1} = \phi_{\mathcal{F}_2}$ .*

We also need:

**Proposition 2.27** *If  $\mathcal{F}_{v_i}$  for  $i = 0, 1$  are two  $\mathbb{R}$ -test configurations associated to two valuations  $v_i \in \mathring{\text{Val}}(X)$  for  $i = 0, 1$ , then  $\phi_{\mathcal{F}_{v_1}} = \phi_{\mathcal{F}_{v_2}} + c$  for a constant  $c \in \mathbb{R}$  if and only if  $v_1 = v_2$  (and hence  $\mathcal{F}_{v_1} = \mathcal{F}_{v_2}$ ).*

**Proof** Recall that  $\phi_{\mathcal{F}_{v_i}} = \lim_{m \rightarrow +\infty} \phi_m^{\mathcal{F}_{v_i}}$  is an increasing limit along the subsequence  $m = 2^k$ , where for any  $w \in \text{Val}(X)$ ,

$$(72) \quad \phi_m^{\mathcal{F}_{v_i}}(w) = -\frac{1}{m} G(w) \left( \sum_{\lambda \in \mathbb{N}} I_{m,\lambda}^{\mathcal{F}_{v_i}} t^{-\lambda} \right),$$

where  $I_{m,\lambda}^{\mathcal{F}_{v_i}}$  is the base ideal of the sublinear system  $\mathcal{F}_{v_i}^\lambda R_m$ . Note that it is easy to see that  $v_1 = v_2$  if and only if  $\alpha_\lambda(v_1) = \alpha_\lambda(v_2)$  for any  $\lambda \in \mathbb{N}$ , where  $\alpha_\lambda(v_i) = \{f \in \mathcal{O}_X \mid v_i(f) \geq \lambda\}$ .

For any  $d \in \mathbb{N}$ , by choosing  $m \gg 1$  we can assume that  $mL \otimes \alpha_d(v_1)$  is globally generated. Then we get  $I_{m,d}^{\mathcal{F}_{v_1}} = \alpha_d(v_1)$ . From this it is also clear that  $\phi_{\mathcal{F}_{v_i}}(v_i) = 0$ . So we get

$$-c = -\phi_{\mathcal{F}_{v_1}}(v_2) \leq -\phi_{2^k}^{\mathcal{F}_{v_1}}(v_2) = \frac{1}{2^k} G(v_2) \left( \sum_{\lambda} I_{2^k,\lambda}^{\mathcal{F}_{v_1}} t^{-\lambda} \right) \leq \frac{1}{2^k} (v_2(I_{2^k,d}^{\mathcal{F}_{v_1}}) - d) = \frac{1}{2^k} (v_2(\alpha_d(v_1)) - d).$$

Since  $k$  can be arbitrarily large, we get  $-c \leq 0$ , ie  $c \geq 0$ . Switching  $v_1$  and  $v_2$  in the above argument, we get  $c \leq 0$ . So  $c = 0$ . We then have the inequality  $v_2(\alpha_d(v_1)) \geq d$  for any  $d \in \mathbb{N}$ . This easily implies  $v_2 \geq v_1$ . Switching  $v_1$  and  $v_2$ , we get  $v_1 \leq v_2$ . Hence  $v_1 = v_2$ , as required.  $\square$

**Corollary 2.28** Assume that  $X$  is smooth. With the same notation as above, if  $\mathcal{F}_{v_1}$  is asymptotically equivalent to  $\mathcal{F}_{v_2}$ , then  $v_1 = v_2$ .

More recently, this result has been proved for any  $\mathbb{Q}$ -Fano variety:

**Lemma 2.29** [15, Lemma 3.16; 21, Theorem C] For any  $\mathbb{Q}$ -Fano variety, if  $v_i$  for  $i = 1, 2$  are two valuations in  $\text{Val}(X)$  such that  $\mathcal{F}_{v_1}$  is asymptotically equivalent to  $\mathcal{F}_{v_2}$ , then  $v_1 = v_2$ .

**Remark 2.30** In the first version of this paper, Corollary 2.28 was stated for any  $\mathbb{Q}$ -Fano variety. However, it has been pointed out by experts that the validity of Theorem 2.26 from [20] for singular  $\mathbb{Q}$ -Fano varieties depends on a still conjectural property called *continuity of envelopes*. Fortunately, recently, in [15; 21], the result in Lemma 2.29 has been given a direct proof without using the continuity of envelopes.

### 2.5 Non-Archimedean invariants of filtrations

For any filtration  $\mathcal{F}$  on  $R = R(X, -K_X)$ , we set

$$(73) \quad L^{\text{NA}}(\phi^{\mathcal{F}}) = L^{\text{NA}}(\mathcal{F}) = L_X^{\text{NA}}(\mathcal{F}) = \inf_{v \in X_{\mathbb{Q}}^{\text{div}}} (A_X(v) + (\phi_{\mathcal{F}} - \phi_{\text{triv}})(v)),$$

$$(74) \quad \tilde{\mathcal{S}}^{\text{NA}}(\phi^{\mathcal{F}}) = \tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}) = \tilde{\mathcal{S}}_X^{\text{NA}}(\mathcal{F}) = -\log \left( \frac{1}{V} \int_{\mathbb{R}} e^{-\lambda} \text{DH}(\mathcal{F}) \right) = -\log \left( \frac{n!}{V} \int_{\Delta} e^{-G_{\mathcal{F}}(y)} dy \right),$$

$$(75) \quad E^{\text{NA}}(\phi^{\mathcal{F}}) = E^{\text{NA}}(\mathcal{F}) = E_X^{\text{NA}}(\mathcal{F}) = \frac{1}{V} \int_{\mathbb{R}} \lambda \cdot \text{DH}(\mathcal{F}) = \frac{n!}{V} \int_{\Delta} G_{\mathcal{F}}(y) dy,$$

$$(76) \quad H^{\text{NA}}(\phi^{\mathcal{F}}) = H^{\text{NA}}(\mathcal{F}) = H_X^{\text{NA}}(\mathcal{F}) = L^{\text{NA}}(\mathcal{F}) - \tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}),$$

$$(77) \quad D^{\text{NA}}(\phi^{\mathcal{F}}) = D^{\text{NA}}(\mathcal{F}) = D_X^{\text{NA}}(\mathcal{F}) = L^{\text{NA}}(\mathcal{F}) - E^{\text{NA}}(\mathcal{F}).$$

The above functionals are by now well known, and we use notation following that in [19; 43]. The formula involving  $G_{\mathcal{F}}$  follows from Theorem 2.5(ii).

**Proposition 2.31** (see [37; 20; 52]) *For a filtration  $\mathcal{F}$ , with the notation from Definition 2.17, we have the following convergence: the sequence from Definition–Proposition 2.15 satisfies, for any  $F \in \{\tilde{\mathcal{S}}, E\}$ ,*

$$(78) \quad \lim_{m \rightarrow +\infty} F^{\text{NA}}(\phi_m^{\mathcal{F}}) = F^{\text{NA}}(\phi^{\mathcal{F}}).$$

Moreover, we have

$$(79) \quad \lim_{m \rightarrow +\infty} L^{\text{NA}}(\phi_m^{\mathcal{F}}) \leq L^{\text{NA}}(\phi^{\mathcal{F}}).$$

**Proof** By Proposition 2.16 we know that  $\text{DH}(\mathcal{F}_{(\mathcal{X}_m, \mathcal{L}_m, \eta_m)})$  converges weakly to  $\text{DH}(\mathcal{F})$  as  $m \rightarrow +\infty$ , from this we easily get the convergence of  $\tilde{\mathcal{S}}^{\text{NA}}$  and  $E^{\text{NA}}$ .

The inequality (79) follows easily from the inequality  $\phi_m^{\mathcal{F}} \leq \phi^{\mathcal{F}}$ . □

For our later argument, we will use a different formulation of the  $L^{\text{NA}}$ -functional studied in [80; 15]. For any filtration  $\mathcal{F}$ , denote by  $I_{\bullet}^{\mathcal{F}(x)} = \{I_{m, mx}^{\mathcal{F}}\}$  the graded sequence of base ideals defined in (52). In [79], Xu and Zhuang introduced the functional

$$(80) \quad \hat{L}^{\text{NA}}(\mathcal{F}) = \sup\{x \in \mathbb{R} \mid \text{lct}(X; I_{\bullet}^{\mathcal{F}(x)}) \geq 1\},$$

and proved that  $\hat{L}^{\text{NA}}(\mathcal{F}) \geq L^{\text{NA}}(\mathcal{F})$ . More recently it has been shown that in fact the two functionals are identical to each other. More specifically, we will need the following comparison results.

**Proposition 2.32** [79, Proposition 4.2 and Theorem 4.3; 15, Lemma 3.8] *For any filtration  $\mathcal{F}$ , we have:*

- (i)  $A_X(E) \geq \hat{L}^{\text{NA}}(\mathcal{F}_{\text{ord}_E})$  for any prime divisor  $E$  over  $X$ , with equality holding if  $\text{ord}_E$  induces a weakly special test configuration.
- (ii)  $\hat{L}^{\text{NA}}(\mathcal{F}) = L^{\text{NA}}(\mathcal{F})$  for any filtration  $\mathcal{F}$ .

For later purposes, we also introduce, for any  $a > 0$ ,

$$(81) \quad E_k^{\text{NA}}(\mathcal{F}) = \frac{1}{V} \int_{\mathbb{R}} x^k \text{DH}(\mathcal{F}) = \lim_{m \rightarrow +\infty} \frac{1}{N_m} \sum_i \left(\frac{\lambda_i^{(m)}}{m}\right)^k,$$

$$(82) \quad Q^{(a)}(\mathcal{F}) = \frac{1}{V} \int_{\mathbb{R}} e^{-ax} \text{DH}(\mathcal{F}) = \frac{1}{V} \sum_{k=0}^{+\infty} \frac{(-1)^k}{k!} a^k E_k^{\text{NA}}(\mathcal{F}),$$

$$(83) \quad Q(\mathcal{F}) := Q^{(1)}(\mathcal{F}).$$

Note that  $E_1^{\text{NA}}(\mathcal{F}) = E^{\text{NA}}(\mathcal{F})$  and  $\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}) = -\log Q(\mathcal{F})$ .

For any  $v \in \text{Val}(X)$  (resp. test configuration  $(\mathcal{X}, \mathcal{L}, a\eta)$ ), we will often write  $F^{\text{NA}}(v)$  (resp.  $F^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta)$ ) for the above various functionals  $F^{\text{NA}}(\mathcal{F})$  with  $\mathcal{F}$  being the corresponding filtration.

**Example 2.33** If  $(\mathcal{X}, \mathcal{L}, a\eta)$  is a normal  $\mathbb{R}$ –test configuration, then we have

$$(84) \quad E^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = a \cdot \frac{\bar{\mathcal{L}}^{n+1}}{(n+1)V},$$

$$(85) \quad L^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = a \cdot (\text{lct}(\mathcal{X}, -K_{\mathcal{X}} - \mathcal{L}; \mathcal{X}_0) - 1).$$

If  $(\mathcal{X}, \mathcal{X}_0)$  has log canonical singularities and  $K_{\mathcal{X}} + \mathcal{L} = \sum_i e_i E_i$  (which is centered at  $\mathcal{X}_0$ ), then

$$(86) \quad L^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = a \cdot \min_i e_i.$$

**Example 2.34** Let  $\mathcal{F}$  be a special  $\mathbb{R}$ –test configuration and let  $(X_0, \eta) = (X_{\mathcal{F},0}, \eta_{\mathcal{F}})$  be the corresponding central fiber. Assume that  $\mathcal{F}|_{X_0} = \mathcal{F}'_{w_{\eta}} R'(-\sigma)$ ; see (33). Let  $\tilde{\varphi}$  be any  $(S^1)^r$ –invariant smooth positively curved Hermitian metric on  $-K_X$ . Then with the notation as in the paragraph containing (9), we have

$$(87) \quad L_X^{\text{NA}}(\mathcal{F}) = L_{X_0}^{\text{NA}}(\mathcal{F}|_{X_0}) = \frac{\int_{X_0} \theta_{\tilde{\varphi}}(\eta) e^{-\tilde{\varphi}}}{\int_{X_0} e^{-\tilde{\varphi}}} - \sigma = -\sigma,$$

$$(88) \quad E_X^{\text{NA}}(\mathcal{F}) = E_{X_0}^{\text{NA}}(\mathcal{F}|_{X_0}) = \frac{1}{V} \int_{X_0} \theta_{\tilde{\varphi}}(\eta) (\text{dd}^c \tilde{\varphi})^n - \sigma,$$

$$(89) \quad \tilde{S}_X^{\text{NA}}(\mathcal{F}) = \tilde{S}_{X_0}^{\text{NA}}(\mathcal{F}|_{X_0}) = -\log\left(\frac{1}{V} \int_{X_0} e^{-\theta_{\eta}} (\text{dd}^c \tilde{\varphi})^n\right) - \sigma.$$

The above identity is well known if  $\mathcal{F}$  comes from a special test configuration. For more general  $\mathcal{F}$ , one can use a sequence of special test configuration to approximate and get the above formula.

Corresponding to (58), we have the following simple transformation rule, which can be checked easily from the defining expressions of the functionals.

**Lemma 2.35** For any  $(a, b) \in \mathbb{R}_{>0} \times \mathbb{R}$ , we have

$$(90) \quad L^{\text{NA}}(a\mathcal{F}(b)) = \hat{L}^{\text{NA}}(a\mathcal{F}(b)) = aL^{\text{NA}}(\mathcal{F}) + b,$$

$$(91) \quad \tilde{S}^{\text{NA}}(\mathcal{F}(b)) = \tilde{S}^{\text{NA}}(\mathcal{F}) + b,$$

$$(92) \quad H^{\text{NA}}(\mathcal{F}(b)) = H^{\text{NA}}(\mathcal{F}).$$

We also note:

**Lemma 2.36** The function  $a \mapsto H^{\text{NA}}(a\mathcal{F})$  is a convex function on  $\mathbb{R}_{\geq 0}$ .

**Proof** Since  $L^{\text{NA}}(a\mathcal{F})$  is linear in  $a$  by (90), we just need to show that  $f(a) := -\tilde{S}^{\text{NA}}(a\mathcal{F})$  is convex in  $a \in \mathbb{R}_{\geq 0}$ . By (25) and (74), we get

$$f(a) = \log\left(\frac{n!}{V} \int_{\Delta} e^{-aG(y)} dy\right),$$

where  $G = G(y) = G_{\mathcal{F}}(y)$ . So we can calculate

$$f''(a) = \frac{\int_{\Delta} G^2 e^{-aG} dy}{\int_{\Delta} e^{-aG} dy} - \frac{\left(\int_{\Delta} G e^{-aG} dy\right)^2}{\left(\int_{\Delta} e^{-aG} dy\right)^2} \geq 0$$

by Hölder’s inequality. □

Note the first identity in (90) comes from Proposition 2.32. Moreover, combining [52, Lemma 3.10] and Proposition 2.32, we have the following invariance property under the twisting.

**Lemma 2.37** *Let  $\mathcal{F}$  be a  $\mathbb{T}$ -equivariant filtration. For any  $\xi \in N_{\mathbb{R}}$ , we have*

$$(93) \quad L^{\text{NA}}(\mathcal{F}_{\xi}) = \widehat{L}^{\text{NA}}(\mathcal{F}_{\xi}) = L^{\text{NA}}(\mathcal{F}).$$

As a consequence, we have

$$(94) \quad L^{\text{NA}}(\mathcal{F}_{\text{wt}_{\xi}}) = \widehat{L}^{\text{NA}}(\mathcal{F}_{\text{wt}_{\xi}}) = 0.$$

The following lemma is a prototype uniqueness result in this paper, and can be seen as a generalization of the uniqueness of Kähler–Ricci soliton vector fields shown by Tian and Zhu [71] (the case when  $\mathcal{F} = \mathcal{F}_{\text{triv}}$ ). See Section 2.6 for more discussion.

**Lemma 2.38** *Let  $\mathcal{F}$  be a  $\mathbb{T}$ -equivariant filtration. Then the function  $\xi \mapsto H^{\text{NA}}(\mathcal{F}_{\xi})$  on  $N_{\mathbb{R}}$  admits a unique minimizer.*

**Proof** By (93),  $L^{\text{NA}}(\mathcal{F}_{\xi})$  is constant in  $\xi$ . Using the identity (67) and (74),

$$-\widetilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_{\xi}) = \log\left(\frac{n!}{V} \int_{\Delta} e^{-G_{\mathcal{F}_{\xi}}(y)} dy\right) = \log\left(\frac{n!}{V} \int_{\Delta} e^{-G_{\mathcal{F}}(y) - \langle y, \xi \rangle} dy\right).$$

It is easy to use this expression to show that  $f(\xi) := -\widetilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_{\xi})$  is strictly convex in  $\xi \in N_{\mathbb{R}}$ , which implies the uniqueness of minimizer. To prove the existence of minimizer, we need to show that  $f(\xi)$  is proper, ie  $\lim_{|\xi| \rightarrow +\infty} f(\xi) = +\infty$ . To see this, recall that we have the vanishing

$$(95) \quad \int_X \theta_{\widetilde{\varphi}}(\xi) e^{-\widetilde{\varphi}} = - \int_X \mathfrak{L}_{\eta} e^{-\widetilde{\varphi}} = 0.$$

This implies that  $0 > \inf_X \theta_{\widetilde{\varphi}}(\xi) = \inf_{\Delta} \langle y, \xi \rangle$  if  $\xi \neq 0$ , which indeed implies the properness. □

**Definition 2.39** We say that a filtration  $\mathcal{F}$  is normalized if

$$(96) \quad L^{\text{NA}}(\mathcal{F}) = 0.$$

A test configuration  $(\mathcal{X}, \mathcal{L}, a\eta)$  is normalized if  $\mathcal{F}_{(\mathcal{X}, \mathcal{L}, a\eta)}$  is normalized.

With the above discussion, the following lemma is easy to prove.

**Lemma 2.40** (i) *Any special test configuration  $(\mathcal{X}, -K_{\mathcal{X}})$  is normalized. More generally, a special  $\mathbb{R}$ -test configuration  $\mathcal{F}$  (see [Definition 2.8](#)) if and only if  $\sigma = 0$  in [\(33\)](#).*

(ii) *For any filtration  $\mathcal{F}$ , the shift  $\mathcal{F}(-L^{\text{NA}}(\mathcal{F}))$  is normalized. If  $\mathcal{F}$  is normalized, then so are  $a\mathcal{F}$  for any  $a > 0$ , and any twist  $\mathcal{F}_{\xi}$ .*

As a consequence of this approximation result in [Proposition 2.31](#), it is convenient for us to introduce:

**Definition–Proposition 2.41** *For any  $\mathbb{Q}$ -Fano variety  $X$ , we define*

$$(97) \quad h(X) = \inf_{(\mathcal{X}, \mathcal{L}, a\eta)} \mathbf{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = \inf_{\mathcal{F}} \mathbf{H}^{\text{NA}}(\mathcal{F}),$$

where  $(\mathcal{X}, \mathcal{L}, a\eta)$  ranges over all test configurations, and  $\mathcal{F}$  ranges over all filtrations or  $\mathbb{R}$ -test configurations.

The following lemma is similar to [\[31, Lemma 2.5\]](#).

**Lemma 2.42** *For any filtration  $\mathcal{F}$ , we have*

$$(98) \quad \tilde{\mathbf{S}}^{\text{NA}}(\mathcal{F}) \leq \mathbf{E}^{\text{NA}}(\mathcal{F}) \quad \text{and} \quad \mathbf{H}^{\text{NA}}(\mathcal{F}) \geq \mathbf{D}^{\text{NA}}(\mathcal{F}).$$

The identities hold true if and only if  $\mathcal{F}(c)$  is asymptotically equivalent to the trivial filtration for some  $c \in \mathbb{R}$ ; see [Definition 2.25](#).

**Proof** The first inequality, which implies the second, follows from the concavity of the logarithmic function. When the identity holds, the DH measure  $\text{DH}(\mathcal{F})$  is a Dirac measure  $V \cdot \delta_c$ . Then  $d_2(\mathcal{F}(c), \mathcal{F}_{\text{triv}}) = 0$ , which by [Definition 2.25](#) means that  $\mathcal{F}(c)$  is asymptotically equivalent to the trivial filtration.  $\square$

Based on the work in [\[29; 28; 40\]](#), Dervan and Székelyhidi proved:

**Theorem 2.43** [\[31\]](#) *Assume that  $X$  is a smooth Fano manifold. There is an identity*

$$(99) \quad h(X) := \inf_{(\mathcal{X}, \mathcal{L}, a\eta) \text{ special}} \mathbf{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = - \inf_{\omega \in c_1(X)} \int_X h_{\omega} e^{h_{\omega}} \omega^n,$$

where  $\omega$  ranges over smooth Kähler metrics from  $c_1(X)$  and  $h_{\omega}$  is the normalized Ricci potential of  $\omega$ . Moreover, the infimum is achieved by a special test configuration constructed via the Gromov–Hausdorff limit Kähler–Ricci soliton from [\[29; 28\]](#).

More recently, Hisamoto [\[43\]](#) gave a different proof of [\(99\)](#) based on the destabilizing geodesic rays constructed from [\[30\]](#).

**Remark 2.44** Our sign convention differs from that of Dervan–Székelyhidi and Hisamoto by a minus. Dervan and Székelyhidi defined a non-Archimedean functional for general  $\mathbb{R}$ -test configuration by mimicking Tian’s CM weight (or the so-called Donaldson–Futaki invariant). But in such generality, their



normalization seems imprecise. Differently from their definition, for any test configuration  $(\mathcal{X}, \mathcal{L}, a\eta)$  one could define

$$(100) \quad \tilde{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = \frac{a}{V} (K_{\bar{\mathcal{X}}/\mathbb{P}^1} \cdot \bar{\mathcal{L}}^n + \bar{\mathcal{L}}^{n+1}) - \tilde{S}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta).$$

By the same argument as [19, Proposition 7.32], we have

$$(101) \quad H^{\text{NA}}(\mathcal{X}, \mathcal{L}, \eta) \leq \tilde{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, \eta),$$

with strict inequality if  $(\mathcal{X}, \mathcal{L}, \eta)$  is anticanonical. Moreover, by (98) we also get

$$(102) \quad \tilde{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, \eta) \geq \text{CM}(\mathcal{X}, \mathcal{L}, \eta) = \frac{1}{V} \left( K_{\bar{\mathcal{X}}/\mathbb{P}^1} \cdot \bar{\mathcal{L}}^n + \frac{n}{n+1} \bar{\mathcal{L}}^{n+1} \right),$$

with the identity being true only if  $(\mathcal{X}, \mathcal{L}, \eta)$  is trivial. One advantage of  $H^{\text{NA}}$  over  $\tilde{H}^{\text{NA}}$  is that the former can be defined for any filtration, not necessarily finitely generated. Due to this reason, we will not use  $\tilde{H}^{\text{NA}}$  in this paper.

### 2.6 $g$ -Ding-stability and Kähler-Ricci solitons

Let  $\mathcal{F}$  be a  $\mathbb{T}$ -equivariant filtration. For any  $\lambda \in \mathbb{R}$ , we have a (finite) decomposition

$$(103) \quad \mathcal{F}^\lambda R_m = \bigoplus_{\alpha \in M_{\mathbb{Z}}} \mathcal{F}^\lambda R_{m,\alpha}.$$

Let  $P$  be the moment polytope of  $(X, -K_X)$  with respect to the  $\mathbb{T}$ -action. Let  $g$  be a smooth positive function on  $P$ . Fix a faithful  $\mathbb{Z}^n$ -valuation that is adapted to the torus action (see Definition 2.21) and let  $\Delta \subset \mathbb{R}^n$  be the Okounkov body that satisfies (65):  $p(\Delta) = P$ , where  $p: \mathbb{R}^n \rightarrow \mathbb{R}^r$  is the natural projection. Still denote by  $g$  the function  $p^*g$  on  $\Delta$ . Define the  $g$ -volume of graded linear series  $\{\mathcal{F}^{(t)} R_m\}$  as

$$\text{vol}_g(\mathcal{F}^{(t)}) := \lim_{m \rightarrow +\infty} \sum_{\alpha} g\left(\frac{\alpha}{m}\right) \frac{\dim \mathcal{F}^{mt} R_{m,\alpha}}{m^n/n!} = n! \cdot \int_{\Delta(\mathcal{F}^{(t)})} g(y) dy_{\text{Leb}} =: n! \cdot \text{vol}_g(\Delta(\mathcal{F}^{(t)})).$$

Then, as in the  $g \equiv 1$  case, we have the convergence

$$\text{DH}_g(\mathcal{F}) := \lim_{m \rightarrow +\infty} \sum_{\alpha} g\left(\frac{\alpha}{m}\right) \delta_{\lambda_i^{(m,\alpha)}/m} = -d \text{vol}_g(\mathcal{F}^{(t)}) = n! \cdot (G_{\mathcal{F}})_*(g(y) dy_{\text{Leb}}).$$

We also set

$$(104) \quad V_g := n! \cdot \text{vol}_g(\Delta) = n! \cdot \int_{\Delta} g(y) dy_{\text{Leb}} = \int_{\mathbb{R}} \text{DH}_g(\mathcal{F}),$$

$$(105) \quad E_g^{\text{NA}}(\mathcal{F}) := \frac{n!}{V_g} \int_{\Delta} G_{\mathcal{F}}(y) g(y) dy_{\text{Leb}} = \frac{1}{V_g} \int_{\mathbb{R}} \lambda \cdot \text{DH}_g(\mathcal{F}),$$

$$(106) \quad D_g^{\text{NA}}(\mathcal{F}) := L^{\text{NA}}(\mathcal{F}) - E_g^{\text{NA}}(\mathcal{F}).$$

If  $(\mathcal{X}, \mathcal{L}, \eta)$  is a test configuration, then we set  $D_g^{\text{NA}}(\mathcal{X}, \mathcal{L}, \eta) = D_g^{\text{NA}}(\mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)})$ .

**Definition 2.45**  $(X, \xi)$  is  $g$ –Ding-semistable if  $D_g^{\text{NA}}(\mathcal{X}, \mathcal{L}, \eta) \geq 0$  for any  $\mathbb{T}$ –equivariant test configuration  $(\mathcal{X}, \mathcal{L}, \eta)$  of  $(X, -K_X)$ .

$(X, \xi)$  is  $g$ –Ding-polystable if it is  $g$ –Ding-semistable, and  $D_g^{\text{NA}}(\mathcal{X}, \mathcal{L}, \eta) = 0$  for a  $\mathbb{T}$ –equivariant weakly special test configuration (see Definition 2.12) only if  $(\mathcal{X}, \mathcal{L}, \eta)$  is a product test configuration.

The following result was proved by adapting the techniques of MMP from [56; 37; 7].

**Theorem 2.46** (see [39]) *To test the  $g$ –Ding-semistability, or the  $g$ –Ding-polystability, of  $(X, \xi)$ , it suffices to test over all special test configurations.*

We have the following valuative criterion:

**Theorem 2.47** [39]  *$X$  is  $g$ –Ding-semistable if and only if for any  $v \in (X_{\mathbb{Q}}^{\text{div}})^{\mathbb{T}}$ , we have*

$$(107) \quad \beta_g(v) := A_X(v) - \frac{1}{V_g} \int_0^{+\infty} \text{vol}_g(\mathcal{F}_v^{(t)}) dt \geq 0.$$

Now we use our notation to reformulate the holomorphic invariants of Tian and Zhu [71] in the study of Kähler–Ricci solitons. We refer to [71; 8; 39] for more details and references. Let  $X$  be a  $\mathbb{Q}$ –Fano variety with an effective  $\mathbb{T}$ –action. We use the same notation, such as an  $(S^1)^r$ –invariant smooth Hermitian metric  $\tilde{\varphi}$  on  $-K_X$ , a moment polytope  $P \subset M_{\mathbb{R}}$ , a function  $\theta_{\tilde{\varphi}}(\eta) = \mathfrak{L}_{\eta} e^{-\tilde{\varphi}} / e^{-\tilde{\varphi}}$ , etc. We identify any  $\eta \in N_{\mathbb{R}}$  with the corresponding holomorphic vector field on  $X$ .

A Kähler–Ricci soliton on  $(X, \xi)$  is a positively curved bounded Hermitian metric  $e^{-\varphi}$  on  $-K_X$  that satisfies the equation

$$(108) \quad e^{\varphi} (\text{dd}^c \varphi)^n = e^{\theta_{\varphi}(\xi)},$$

where  $\theta_{\varphi}(\xi) = \theta_{\tilde{\varphi}}(\xi) + \xi(\varphi - \tilde{\varphi})$ . Over  $X^{\text{reg}}$ ,  $\varphi$  is smooth [6; 39] and satisfies the identity

$$(109) \quad \text{Ric}(\text{dd}^c \varphi) - \text{dd}^c \varphi = -\text{dd}^c \theta_{\varphi}(\xi).$$

As a consequence, the family of metrics  $\varphi(s) := \sigma_{\xi}(s)^* \varphi$  satisfies the normalized Kähler–Ricci flow,

$$(110) \quad \frac{d}{ds} \text{dd}^c \varphi(s) = -\text{Ric}(\text{dd}^c \varphi(s)) + \text{dd}^c \varphi(s).$$

For any  $\xi \in N_{\mathbb{R}}$ , we set  $g_{\xi}(x) = e^{-\langle x, \xi \rangle} = e^{-\sum_{i=1}^r \xi^i x_i}$ , which is a smooth positive function on  $P$ , and write  $F_{g_{\xi}}$  as  $F_{\xi}$  for  $F \in \{\mathbf{L}, \mathbf{D}\}$  etc, and  $V_{\xi} := V_{g_{\xi}}$ . Tian and Zhu [71] defined a modified Futaki invariant as an obstruction to the existence of Kähler–Ricci solitons on  $(X, \xi)$ : for any  $\eta \in N_{\mathbb{R}}$ ,

$$(111) \quad \text{Fut}_{\xi}(\eta) := -\frac{1}{V_{\xi}} \int_X \theta_{\tilde{\varphi}}(\eta) e^{-\theta_{\tilde{\varphi}}(\xi)} (\text{dd}^c \tilde{\varphi})^n = D_{\xi}^{\text{NA}}(\text{wt}_{\eta}),$$

where  $V_{\xi} = \int_X e^{-\theta_{\tilde{\varphi}}(\xi)} (\text{dd}^c \tilde{\varphi})^n$ . The second identity follows by noting that  $D_{\xi}^{\text{NA}}(\text{wt}_{\eta}) = -E_{\xi}^{\text{NA}}(\text{wt}_{\eta})$  because of the vanishing  $L^{\text{NA}}(\text{wt}_{\eta}) = 0$ .

**Remark 2.48** Again, here we have used the negative sign convention compared to [71].

$\text{Fut}_\xi$  does not depend on the choice of  $\tilde{\varphi}$  and  $(X, \xi)$  admits a KR soliton only if  $\text{Fut}_\xi \equiv 0$  on  $N_{\mathbb{R}}$ . Moreover, by [71, Lemma 2.2] the soliton vector field is a priori uniquely determined by minimizing the strictly convex functional (Tian and Zhu didn't use the logarithm) on  $N_{\mathbb{R}}$  (see Lemma 2.38), which is the antiderivative of  $\eta \mapsto \text{Fut}_\xi(\eta)$ ,

$$(112) \quad \xi \mapsto \log\left(\frac{1}{V} \int_X e^{-\theta_{\tilde{\varphi}}(\xi)} (\text{dd}^c \tilde{\varphi})^n\right) = \log\left(\frac{1}{V} \int_{\mathbb{R}} e^{-\lambda} \text{DH}(\mathcal{F}_{\text{wt}_\xi})\right) = -\tilde{\mathcal{S}}^{\text{NA}}(\text{wt}_\xi).$$

Recall also that  $L^{\text{NA}}(\text{wt}_\eta) = \hat{L}^{\text{NA}}(\text{wt}_\eta) \equiv 0$  on  $N_{\mathbb{R}}$ ; see (94). Combining these discussions we get the derivative identity

$$(113) \quad \frac{d}{ds} H^{\text{NA}}(\text{wt}_{\xi+s\eta}) = \frac{d}{ds} \mathbf{H}^{\text{NA}}(\text{wt}_{\xi+s\eta}) = \mathbf{D}_\xi^{\text{NA}}(\text{wt}_\eta) = \text{Fut}_\xi(\eta).$$

For simplicity of notation, we introduce:

**Definition 2.49** We say that  $(X, \xi)$  is K-semistable (resp. K-polystable) if  $X$  is  $g_\xi$ -Ding-semistable (resp.  $g_\xi$ -Ding-polystable).

**Remark 2.50** Since by Theorem 2.46 it is enough to test the stability on special test configurations, this definition coincides with the original modified K-(poly)stability adopted by Tian as well as Berman, Witt and Nyström, and others. To respect the original notation, we will just call  $(X, \xi)$  K-(poly)stable, although we will also freely use the notion of Ding-(poly)stability.

By [8; 31], when  $X$  is smooth, the Yau–Tian–Donaldson conjecture is true, ie K-polystability is equivalent to the existence of Kähler–Ricci solitons. For singular  $X$ , we proved in [39] a version of the Yau–Tian–Donaldson conjecture involving  $\text{Aut}(X, \xi)_0$ -uniform Ding-stability.

### 3 $H^{\text{NA}}$ invariant and MMP

#### 3.1 An intersection formula for higher moments

Let  $(\mathcal{X}, \mathcal{L}, \eta)$  be any normal ample test configuration. Choose a smooth (semipositive) curvature form  $\omega$  in  $c_1(\mathcal{L}|_{\mathcal{X}_0})$ . Let  $\theta$  be the Hamiltonian function for  $\eta$  with respect to  $\omega$ , so  $\iota_\eta \omega = (\sqrt{-1}/2\pi) \bar{\partial} \theta$ . By the equivariant Riemann–Roch formula, we get

$$E_k^{\text{NA}}(\mathcal{X}, \mathcal{L}) := E_k^{\text{NA}}(\mathcal{F}_{(\mathcal{X}, \mathcal{L})}) = \lim_{m \rightarrow +\infty} \frac{1}{N_m} \sum_i \binom{\lambda_i^{(m)}}{m}^k = \frac{1}{V} \int_{\mathcal{X}_0} \theta^k \omega^n.$$

To motivate our calculations, we will first give a direct proof of two identities which can already be derived from the above discussion.

**Lemma 3.1** We have

$$(114) \quad E_k^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \frac{1}{V} \int_{\mathbb{R}} x^k \text{DH}(\mathcal{F}^{(x)}),$$

$$(115) \quad E^{\text{NA}}(\mathcal{X}, \mathcal{L}) = E_1^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \frac{1}{V} \frac{\bar{\mathcal{L}}^{n+1}}{n+1}.$$

**Proof** When we change  $\mathcal{L}$  to  $\mathcal{L} + d\mathcal{X}_0$ ,  $\mathcal{F}$  is changed to  $\mathcal{F}(d)$ , and both sides of the above identities have  $d$  added to them. So we can assume that  $\bar{\mathcal{L}}$  is very ample over  $\bar{\mathcal{X}}$ . Then we have

$$(116) \quad \bar{\mathcal{X}} = \text{Proj} \left( \bigoplus_{m \geq 0} \bigoplus_{j=0}^{+\infty} t^{-j} \mathcal{F}^j R_m \right)$$

and  $\bar{\mathcal{L}}_d = \mathcal{O}_{\bar{\mathcal{X}}}(1)$ .

For simplicity of notation, we write

$$(117) \quad \begin{aligned} f_k(m) &= \sum_{i=1}^{N_m} (\lambda_i^{(m)})^k = \sum_{j=0} j^k (\dim \mathcal{F}^j R_m - \dim \mathcal{F}^{j+1} R_m) \\ &= \sum_{j=1} (j^k - (j-1)^k) \dim \mathcal{F}^j R_m = \sum_{j=1} (kj^{k-1} + O(j^{k-2})) \dim \mathcal{F}^j R_m. \end{aligned}$$

We easily get the identity

$$(118) \quad E_k^{\text{NA}} = \frac{1}{V} \lim_{m \rightarrow +\infty} \frac{n!}{m^{n+k}} f_k(m) = \frac{1}{V} \int_0^{+\infty} kx^{k-1} \text{vol}(\mathcal{F}^{(x)} R_\bullet) dx = \frac{1}{V} \int_{\mathbb{R}} x^k (-d \text{vol}(\mathcal{F}^{(x)})).$$

Moreover, we have the dimension formula

$$N_m := h^0(\bar{\mathcal{X}}, m\bar{\mathcal{L}}) = \sum_{j=0}^{+\infty} \dim \mathcal{F}^j R_m = \frac{m^{n+1}}{n!} \int_0^{+\infty} \text{vol}(\mathcal{F}^{(x)} R_\bullet) dx + O(m^n),$$

which, by the Riemann–Roch formula, gives the identity

$$(119) \quad \frac{1}{V} \frac{\bar{\mathcal{L}}^{n+1}}{n+1} = \frac{1}{V} \int_0^{+\infty} \text{vol}(\mathcal{F}^{(x)} R_\bullet) dx = \frac{1}{V} \int_0^{+\infty} x \text{DH}(\mathcal{F}). \quad \square$$

The formula (115) goes back to Mumford’s study of GIT [61], and has also been used in the study of K–stability. The following result is a generalization of it to higher moments. We will use the following notation as in [39]. Let  $\mathbb{C}^* \rightarrow \mathbb{C}^{k+1} \setminus \{0\} \rightarrow \mathbb{P}^k$  be the principal  $\mathbb{C}^*$ –bundle and set

$$(120) \quad (\bar{\mathcal{X}}^{[k]}, \bar{\mathcal{L}}^{[k]}) := ((\bar{\mathcal{X}}, \bar{\mathcal{L}}) \times (\mathbb{C}^{k+1} \setminus \{0\})) / \mathbb{C}^*.$$

**Remark 3.2** Since the  $\mathbb{C}^*$ –action on  $\bar{\mathcal{X}}$  moves the fiber  $\bar{\mathcal{X}} \rightarrow \mathbb{P}^1$ , the situation here is different from the situation in [19, Corollary 3.4] or [39], where a similar fiber construction with respect to a vertical torus action is used.

**Proposition 3.3** *Let  $(\mathcal{X}, \mathcal{L})$  be a normal ample test configuration. For any  $k \geq 1$ , we have the intersection formula*

$$(121) \quad E_k^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \frac{1}{V} \frac{k!n!}{(n+1)!} (\bar{\mathcal{L}}^{[k-1]})^{n+k}.$$

**Proof** We use the notation from the above proof and without loss of generality assume that  $\bar{\mathcal{L}}$  is very ample over  $\bar{\mathcal{X}}$ .

The weights  $\{\mu_\alpha \mid \alpha = 1, \dots, \mathcal{N}_m\}$  and multiplicities of  $\mathbb{C}^*$ -action on  $H^0(\bar{\mathcal{X}}, \bar{\mathcal{L}})$  are given according to the isomorphism (44). By the identity (117), the weight of  $\mathbb{C}^*$  on  $\det H^0(\bar{\mathcal{X}}, m\bar{\mathcal{L}})$  is given by

$$(122) \quad \sum_{\alpha=1}^{\mathcal{N}_m} \mu_\alpha^{k-1} = \sum_{j=0}^{+\infty} j^{k-1} \dim \mathcal{F}^j R_m = k^{-1} f_k(m) + O(m^{n+k-1}).$$

Choose a smooth Kähler metric  $\Omega \in c_1(\bar{\mathcal{L}})$  on  $\bar{\mathcal{X}}$  and let  $\Theta$  be the Hamiltonian function for  $\eta$ . Then by the equivariant Riemann–Roch formula, we get

$$(123) \quad \lim_{m \rightarrow +\infty} \frac{(n+1)!}{m^{n+1}} \sum_{\alpha} \left(\frac{\mu_\alpha}{m}\right)^{k-1} = \int_{\bar{\mathcal{X}}} \Theta^{k-1} \Omega^{n+1} = \frac{(k-1)!(n+1)!}{(k+n)!} \int_{\bar{\mathcal{X}}^{[k-1]}} (\Omega + \Theta t)^{n+k} \\ = \frac{(k-1)!(n+1)!}{(k+n)!} (\bar{\mathcal{L}}^{[k-1]})^{n+k}.$$

Combining (118), (122) and (123), we get

$$E_k^{\text{NA}} = \frac{1}{V} \lim_{k \rightarrow \infty} \frac{n!}{m^{n+k}} k \sum_{\alpha} \mu_\alpha^{k-1} = \frac{1}{V} \frac{k}{n+1} \frac{(k-1)!(n+1)!}{(k+n)!} (\bar{\mathcal{L}}^{[k-1]})^{n+k} \\ = \frac{1}{V} \frac{k!n!}{(k+n)!} (\bar{\mathcal{L}}^{[k-1]})^{n+k}. \quad \square$$

Recall from (82) that  $\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = -\log \mathcal{Q}^{(a)}$ , where

$$\mathcal{Q}^{(a)} = \frac{1}{V} \int_{\mathcal{X}_0} e^{-a\theta} \omega^n = \sum_{k=0}^{\infty} (-1)^k a^k \frac{1}{V} \int_{\mathcal{X}_0} \frac{\theta^k}{k!} \omega^n = \sum_k (-1)^k \frac{a^k}{k!} E_k^{\text{NA}}.$$

**Proposition 3.4** Let  $(\mathcal{X}, \mathcal{L}_\lambda, a\eta)_{\lambda \in (-\epsilon, \epsilon)}$  be a family of normal test configurations of  $(X, -K_X)$ , with a fixed total space and varying polarization. Assume that  $\mathcal{X}_0 = \sum_i b_i E_i$  for irreducible components  $E_i$ , and that  $\mathcal{L}_\lambda$  is differentiable with respect to  $\lambda$ . Then we have the derivative formula

$$(124) \quad \frac{d}{d\lambda} \tilde{\mathcal{S}}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) = a \frac{\sum_i e_i \mathcal{Q}_i^{(a)}}{\mathcal{Q}^{(a)}},$$

where  $\mathcal{Q}_i^{(a)} = (1/V) \int_{E_i} e^{-a\theta} \omega^n$ .

**Proof** We use the intersection formula (121) to get

$$V \cdot \frac{d}{d\lambda} E_k^{\text{NA}} = \frac{d}{d\lambda} \frac{k!n!}{(k+n)!} (\bar{\mathcal{L}}^{[k-1]})^{n+k} = \frac{k!n!}{(k+n-1)!} (\bar{\mathcal{L}}^{[k-1]})^{n+k-1} \cdot \dot{\bar{\mathcal{L}}}^{[k-1]} \\ = \frac{k!n!}{(k+n-1)!} \sum_i e_i \int_{E_i^{[k-1]}} (\Omega + \Theta t)^{n+k-1} = k \sum_i e_i \int_{E_i} \theta^{k-1} \omega^n,$$

where  $E_i^{[k-1]} = (E_i \times \mathbb{C}^{k-1} \setminus \{0\})/\mathbb{C}^*$ . So we get the desired formula

$$\begin{aligned} \frac{d}{d\lambda} \mathbf{Q}^{(a)} &= \sum_k (-1)^k \frac{a^k}{k!} \frac{d}{d\lambda} \mathbf{E}_k^{\text{NA}} = \sum_{k=1} (-1)^k \frac{a^k}{(k-1)!} \sum_i e_i \frac{1}{V} \int_{E_i} \theta^{k-1} \omega^n \\ &= -a \sum_i e_i \frac{1}{V} \int_{E_i} \sum_{j=0} \frac{(-1)^j}{j!} (a\theta)^j \omega^n = -a \sum_i e_i \frac{1}{V} \int_{E_i} e^{-a\theta} \omega^n = -a \sum_i \mathbf{Q}_i^{(a)}. \end{aligned}$$

The termwise differentiation and the change of summation are valid because of absolute convergence.  $\square$

### 3.2 Decreasing of $H^{\text{NA}}$ along MMP

**Theorem 3.5** *Let  $\mathbb{G}$  be a reductive group and  $(\mathcal{X}, \mathcal{L}, a\eta)$  be a  $\mathbb{G}$ –equivariant normal test configuration. There exists a  $\mathbb{G}$ –equivariant special test configuration  $(\mathcal{X}^s, \mathcal{L}^s, a^s \eta^s)$  such that*

$$(125) \quad H^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) \geq H^{\text{NA}}(\mathcal{X}^s, \mathcal{L}^s, a^s \eta^s).$$

Moreover, if  $\mathcal{X}_0$  is reduced, then the identity holds true if  $\mathcal{X}$  is already a special test configuration.

**Proof** For simplicity of notation, we assume  $\mathbb{G}$  is trivial. The general case is obtained by running the  $\mathbb{G}$ –equivariant MMP in the following arguments.

**Step 1** Choose a semistable reduction of  $\mathcal{X} \rightarrow \mathbb{C}$ . By this, we mean that there is an integer  $d$  and a  $\mathbb{G}$ –equivariant log resolution of singularities  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}^{(d_1)} := (\mathcal{X} \times_{\mathbb{C}, t \rightarrow t^{d_1}} \mathbb{C})^{\text{norm}}$  (see (49)) such that  $(\tilde{\mathcal{X}}, \tilde{\mathcal{X}}_0)$  is simple normal crossing. In particular,  $\mathcal{X}_0^{(d_1)}$  is reduced. By using the identity (51) and Lemma 2.35 we easily get

$$(126) \quad H^{\text{NA}}(\mathcal{X}^{(d_1)}, \mathcal{L}^{(d_1)}, a\eta^{(d_1)}/d_1) = H^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta).$$

**Step 2** In this step, we show that there exist  $d_1 \in \mathbb{Z}_{>0}$ , a projective birational  $\mathbb{C}^*$ –equivariant morphism  $\pi: \mathcal{X}^{\text{lc}} \rightarrow \mathcal{X}^{(d_1)}$  and a normal, ample test configuration  $(\mathcal{X}^{\text{lc}}, \mathcal{L}^{\text{lc}})/\mathbb{C}$  for  $(X, L)$ , such that

$$(127) \quad H^{\text{NA}}(\mathcal{X}^{(d_1)}, \mathcal{L}^{(d_1)}, a\eta^{(d_1)}/d_1) \geq H^{\text{NA}}(\mathcal{X}^{\text{lc}}, \mathcal{L}^{\text{lc}}, a\eta^{\text{lc}}/d_1).$$

Moreover, if the equality holds, then  $(\mathcal{X}^{(d_1)}, \mathcal{L}^{(d_1)})$  is isomorphic to  $(\mathcal{X}^{\text{lc}}, \mathcal{L}^{\text{lc}})$ , and hence  $(\mathcal{X}, \mathcal{X}_0)$  is already log canonical.

We run a  $\mathbb{C}^*$ –equivariant MMP to get a log canonical modification  $\pi^{\text{lc}}: \mathcal{X}^{\text{lc}} \rightarrow \mathcal{X}^{(d_1)}$  such that  $(\mathcal{X}^{\text{lc}}, \mathcal{X}_0^{\text{lc}})$  is log canonical and  $K_{\mathcal{X}^{\text{lc}}}$  is relatively ample over  $\mathcal{X}^{(d_1)}$ . Set  $E = K_{\mathcal{X}^{\text{lc}}} + (\pi^{\text{lc}})^* \mathcal{L} = \sum_{i=1}^k e_i \mathcal{X}_{0,i}$  with  $e_1 \leq e_2 \leq \dots \leq e_k$  and  $\mathcal{L}_\lambda^{\text{lc}} = (\pi^{\text{lc}})^* \mathcal{L}^{(d_1)} + \lambda E$ . Then since  $E$  is relatively ample over  $\mathcal{X}^{(d_1)}$ ,  $\mathcal{L}_\lambda$  is ample over  $\mathcal{X}^{\text{lc}}$  for  $0 < \lambda \ll 1$ . So

$$\mathbf{L}^{\text{NA}}(\mathcal{X}^{\text{lc}}, \mathcal{L}_\lambda^{\text{lc}}, a\eta^{\text{lc}}/d_1) = \frac{a}{d_1} \mathbf{L}^{\text{NA}}(\mathcal{X}^{\text{lc}}, \mathcal{L}_\lambda^{\text{lc}}, \eta^{\text{lc}}) = \frac{a}{d_1} (1 + \lambda)e_1.$$

By definition (76), we have

$$\tilde{\mathbf{S}}^{\text{NA}}(\mathcal{X}^{\text{lc}}, \mathcal{L}_\lambda^{\text{lc}}, a\eta^{\text{lc}}/d_1) = -\log \mathbf{Q}^{(ad_1^{-1})}, \quad \mathbf{H}^{\text{NA}}(\mathcal{X}^{\text{lc}}, \mathcal{L}_\lambda^{\text{lc}}, a\eta^{\text{lc}}/d_1) = \frac{a(1 + \lambda)e_1}{d_1} + \log \mathbf{Q}^{(ad_1^{-1})}.$$

We then use (124) to calculate

$$\frac{d}{d\lambda} \mathbf{H}^{\text{NA}}(\mathcal{X}^{\text{lc}}, \mathcal{L}_\lambda^{\text{lc}}, a\eta^{\text{lc}}/d_1) = \frac{ae_1}{d_1} - \frac{a}{d_1} \frac{\sum_i e_i \mathcal{Q}_i^{(ad_1^{-1})}}{\sum_i \mathcal{Q}_i^{(ad_1^{-1})}} = -\frac{a}{d_1} \frac{\sum_i (e_i - e_1) \mathcal{Q}_i^{(ad_1^{-1})}}{\mathcal{Q}^{(ad_1^{-1})}} \leq 0.$$

The last identity holds if and only if  $e_i \equiv e_1$ , and hence  $(\mathcal{X}^{(d_1)}, \mathcal{L}^{(d_1)}) \cong (\mathcal{X}^{\text{lc}}, \mathcal{L}^{\text{lc}})$ . In this case,  $(\mathcal{X}^{(d_1)}, \mathcal{X}_0^{(d_1)})$  is log canonical, which implies that  $(\mathcal{X}, \mathcal{X}_0)$  is already log canonical, by the pullback formula for the log differential; see [56, page 210].

**Step 3** With the  $(\mathcal{X}^{\text{lc}}, \mathcal{L}^{\text{lc}})$  obtained from the first step, we run a relative MMP with scaling to get a normal, ample test configuration  $(\mathcal{X}^{\text{ac}}, \mathcal{L}^{\text{ac}})/\mathbb{P}^1$  for  $(X, -K_X)$  with  $(\mathcal{X}^{\text{ac}}, \mathcal{X}_0^{\text{ac}})$  log canonical such that  $-K_{\mathcal{X}^{\text{ac}}} \sim_{\mathbb{Q}, \mathbb{C}} \mathcal{L}^{\text{ac}}$ . More concretely, we take  $q \gg 1$  such that  $\mathcal{H}^{\text{lc}} = \mathcal{L}^{\text{lc}} - (q + 1)^{-1}(\mathcal{L}^{\text{lc}} + K_{\mathcal{X}^{\text{lc}}})$  is relatively ample. Set  $\mathcal{X}^0 = \mathcal{X}^{\text{lc}}, \mathcal{L}^0 = \mathcal{L}^{\text{lc}}, \mathcal{H}^0 = \mathcal{H}^{\text{lc}}$  and  $\lambda_0 = q + 1$ . Then  $K_{\mathcal{X}^0} + \lambda_0 \mathcal{H}^0 = q\mathcal{L}^0$ . We run a sequence of  $K_{\mathcal{X}^0}$ -MMP over  $\mathbb{C}$  with scaling of  $\mathcal{H}^0$ . Then we obtain a sequence of models

$$\mathcal{X}^0 \dashrightarrow \mathcal{X}^1 \dashrightarrow \dots \dashrightarrow \mathcal{X}^k$$

and a sequence of critical values

$$\lambda_{i+1} = \min\{\lambda \mid K_{\mathcal{X}^i} + \lambda \mathcal{H}^i \text{ is nef over } \mathbb{C}\}$$

with  $q + 1 = \lambda_0 \geq \lambda_1 \geq \dots \geq \lambda_k > \lambda_{k+1} = 1$ . For any  $\lambda_i \geq \lambda \geq \lambda_{i+1}$ , we let  $\mathcal{H}^i$  be the pushforward of  $\mathcal{H}$  to  $\mathcal{X}^i$  and set

$$(128) \quad \mathcal{L}_\lambda^i = \frac{1}{\lambda - 1} (K_{\mathcal{X}^i} + \lambda \mathcal{H}^i) = \frac{1}{\lambda - 1} (K_{\mathcal{X}^i} + \mathcal{H}^i) + \mathcal{H}^i =: \frac{1}{\lambda - 1} E + \mathcal{H}^i.$$

Write  $E = \sum_{j=1}^k e_j \mathcal{X}_{0,j}^i$  with  $e_1 \leq e_2 \leq \dots \leq e_k$ . Then we have  $(d/d\lambda)\mathcal{L}_\lambda^i = -(1/(\lambda - 1)^2)E$  and

$$\mathbf{L}^{\text{NA}}(\mathcal{X}^i, \mathcal{L}^i, a\eta^i/d_1) = \frac{a\lambda}{\lambda - 1} e_1.$$

So we can again use (124) to calculate

$$\begin{aligned} \frac{d}{d\lambda} \mathbf{H}^{\text{NA}}\left(\mathcal{X}^i, \mathcal{L}_\lambda^i, \frac{a\eta^i}{d_1}\right) &= -\frac{a}{d_1(\lambda - 1)^2} e_1 + \frac{a}{d_1(\lambda - 1)^2} \frac{\sum_i e_i \mathcal{Q}_i^{(ad_1^{-1})}}{\mathcal{Q}^{(ad_1^{-1})}} \\ &= \frac{a}{d_1(\lambda - 1)^2} \frac{\sum_i (e_i - e_1) \mathcal{Q}_i^{(a)}}{\mathcal{Q}^{(a)}} \geq 0. \end{aligned}$$

The last identity holds only if  $e_i \equiv e_1$ , which implies  $(\mathcal{X}^{\text{lc}}, \mathcal{L}^{\text{lc}}) \cong (\mathcal{X}^{\text{ac}}, \mathcal{L}^{\text{ac}} + e_1 \mathcal{X}_0^{\text{ac}})$ .

**Step 4** With the test configuration  $(\mathcal{X}^{\text{ac}}, \mathcal{L}^{\text{ac}})$  obtained from Step 2, there exists a  $d_2 \in \mathbb{Z}_{>0}$  and a projective birational  $T_{\mathbb{C}} \times \mathbb{C}^*$ -equivariant birational map  $(\mathcal{X}^{\text{ac}})^{(d_2)} \dashrightarrow \mathcal{X}^s$  over  $\mathbb{P}^1$  such that  $(\mathcal{X}^s, -K_{\mathcal{X}^s})$  is a special test configuration and

$$(129) \quad \mathbf{H}^{\text{NA}}\left(\mathcal{X}^{\text{ac}}, \mathcal{L}^{\text{ac}}, \frac{a\eta}{d_1 d_2}\right) \geq \mathbf{H}^{\text{NA}}\left(\mathcal{X}^s, \mathcal{L}^s, \frac{a\eta^s}{d_1 d_2}\right).$$

As in [56], this is achieved by doing a base change and running an MMP. Let  $E = -K_{\mathcal{X}^s/\mathbb{P}^1} - (-K_{\mathcal{X}'/\mathbb{P}^1})$ . Then  $E \geq 0$  by the negativity lemma. So  $\mathcal{L}'_\lambda = -K_{\mathcal{X}'/\mathbb{P}^1} + \lambda E$ , and

$$(130) \quad \text{lct}\left(\mathcal{X}', \mathcal{L}'_\lambda, \frac{a\eta'}{d_1 d_2}\right) = \frac{a}{d_1 d_2} \lambda e_1.$$

So, as before, we get

$$\frac{d}{d\lambda} \mathbf{H}^{\text{NA}}\left(\mathcal{X}', \mathcal{L}'_\lambda, \frac{a\eta'}{d_1 d_2}\right) = \frac{a}{d_1 d_2} e_1 - \frac{a}{d_1 d_2} \frac{\sum_i e_i \mathcal{Q}_i^{(ad_1^{-1})}}{\mathcal{Q}^{(ad_1^{-1})}} = -\frac{a}{d_1 d_2} \frac{\sum_i (e_i - e_1) \mathcal{Q}_i^{(ad_1^{-1})}}{\mathcal{Q}^{(ad_1^{-1})}} \leq 0.$$

The last identity holds only if  $e_i \equiv e_1$  which implies  $(\mathcal{X}^{\text{ac}}, \mathcal{L}^{\text{ac}}) \cong (\mathcal{X}^s, \mathcal{L}^s)$ . □

**Corollary 3.6** *We have the identity*

$$(131) \quad h(X) = \inf_{(\mathcal{X}, \mathcal{L}, a\eta) \text{ special}} \mathbf{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta).$$

**Lemma 3.7** *For any normal test configuration  $(\mathcal{X}, \mathcal{L}, \eta)$ , there exists a unique  $a_* > 0$  such that*

$$(132) \quad \mathbf{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a_*\eta) = \inf_{c>0} \mathbf{H}^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) =: \mathbf{H}_*^{\text{NA}}(\mathcal{X}, \mathcal{L}).$$

As a consequence, we have

$$(133) \quad h(X) = \inf_{(\mathcal{X}, -K_{\mathcal{X}}) \text{ special}} \mathbf{H}_*^{\text{NA}}(\mathcal{X}, \mathcal{L}).$$

**Proof** By taking normalization of a fiber product, without loss of generality we can assume that  $\mathcal{X}$  dominates  $X_{\mathbb{C}} := X \times \mathbb{C}$  by a  $\mathbb{C}^*$ -equivariant birational morphism  $\rho: \mathcal{X} \rightarrow X_{\mathbb{C}}$ .

Choose a  $\mathbb{C}^*$ -equivariant resolution of singularities  $\mu: \tilde{\mathcal{X}} \rightarrow \mathcal{X}$  such that the pair  $(\tilde{\mathcal{X}}, \tilde{\mathcal{X}}_0^{\text{red}})$  has simple normal crossing singularities. Set  $\tilde{\rho} = \rho \circ \mu$ . Then we can write

$$K_{\tilde{\mathcal{X}}} = \tilde{\rho}^* K_{X_{\mathbb{C}}} + \sum_i a_i E_i + \sum_j a'_j E'_j, \quad \pi^* \mathcal{X}_0 = \sum_i b_i E_i, \quad \mu^* \mathcal{L} = \tilde{\rho}^* (-K_{X_{\mathbb{C}}}) + \sum_i c_i E_i,$$

where  $\{E_i\}$  are irreducible components of  $\tilde{\mathcal{X}}_0$  and  $\{E'_j\}$  are horizontal exceptional divisors. Then we have the identity (see [19, Proposition 7.29])

$$\begin{aligned} L^{\text{NA}}(\mathcal{X}, \mathcal{L}) &= \text{lct}(\mathcal{X}, -(K_{\mathcal{X}} + \mathcal{L}); \mathcal{X}_0) - 1 \\ &= \min_i \left( b_i^{-1} A_{(X \times \mathbb{C}, X \times \{0\})}(E_i) + b_i^{-1} \text{ord}_{E_i} \left( \sum_k c_k E_k \right) \right) = \min_i \frac{1 + a_i + c_i}{b_i} - 1. \end{aligned}$$

Because  $\mathbf{H}^{\text{NA}}$  is translation-invariant, by adding a multiple of  $\mathcal{X}_0$  to  $\mathcal{L}$  we can normalize  $\phi = \phi_{\mathcal{F}}$  to satisfy  $L^{\text{NA}}(\phi) = 0$ . So we get

$$(134) \quad c_i \geq b_i - 1 - a_i$$

and, without loss of generality,  $c_1 = b_1 - 1 - a_1$ . So

$$(135) \quad \lambda_{\min} = \min_i \frac{c_i}{b_i} \leq \frac{c_1}{b_1} = 1 - \frac{a_1 + 1}{b_1} = 1 - A_{X_{\mathbb{C}}}(b_1^{-1} \text{ord}_{E_1}) = -A_X(v_{E_1}) \leq 0,$$



where  $v_{E_1} := r(b_1^{-1} \text{ord}_{E_1})$  is the restriction of the valuation  $\text{ord}_{E_1}$  to the function field  $\mathbb{C}(X)$ . Here we used the identity between log discrepancies from [19, Proposition 4.11] and the assumption that  $X$  has log terminal singularities.

Set  $\mathcal{F} = \mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)}$ . Then according to (43), we have  $\mathcal{F}_{(\mathcal{X}, \mathcal{L}, a\eta)} = a\mathcal{F}_{(\mathcal{X}, \mathcal{L}, \eta)}$ . Moreover, by (74), (76) and (90), we get the expression

$$f(a) := H^{\text{NA}}(a\mathcal{F}) = aL^{\text{NA}}(\mathcal{F}) + \log\left(\frac{1}{V} \int_{\lambda_{\min}}^{+\infty} e^{-a\lambda} \text{DH}(\mathcal{F})\right).$$

By Lemma 2.36,  $f(a)$  is convex in  $a \in [0, +\infty)$ . If  $\lambda_{\min} < 0$ , then  $f(a)$  diverges to  $+\infty$  as  $a \rightarrow +\infty$  by the above expression. So  $f(a)$  admits a unique minimum over  $[0, +\infty)$ .

If  $\lambda_{\min} = 0$ , then by (135)  $A_X(v_{E_1}) = 0$ , which implies that  $v_{E_1}$  is trivial so that

$$\lambda_{\max} = \text{ord}_{E_1}\left(\sum_k c_k E_k\right) = c_1 = 0 = \lambda_{\min}.$$

See [19, Theorem 5.16]. This implies that the normal test configuration  $(\mathcal{X}, \mathcal{L})$  is equivalent to a trivial test configuration and hence  $H^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) \equiv 0$ . □

### 4 A minimization problem for real valuations

In this section, we will introduce a minimization problem for valuations analogous to the normalized volume functional in the local setting [51].

**Definition 4.1** For any  $v \in \text{Val}(X)$ , define

$$(136) \quad \tilde{\beta}(v) = \begin{cases} A_X(v) - \tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_v) & \text{if } A_X(v) < +\infty, \\ +\infty & \text{otherwise.} \end{cases}$$

Note that by integration by parts we have

$$(137) \quad \begin{aligned} e^{-\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_v)} &= \frac{1}{V} \int_{\mathbb{R}} e^{-x} \text{DH}(\mathcal{F}_v) = \frac{1}{V} \int_0^{+\infty} e^{-x} (-d\text{vol}(\mathcal{F}_v^{(x)})) \\ &= 1 - \frac{1}{V} \int_0^{+\infty} \text{vol}(\mathcal{F}_v^{(x)} R_{\bullet}) e^{-x} dx \leq 1, \end{aligned}$$

with identity if and only if  $v$  is trivial. So we can rewrite  $\tilde{\beta}(v)$  as

$$(138) \quad \tilde{\beta}(v) = A_X(v) + \log\left(1 - \frac{1}{V} \int_0^{+\infty} e^{-x} \text{vol}(\mathcal{F}_v^{(x)} R_{\bullet}) dx\right).$$

**Lemma 4.2** For any  $v \in X_{\mathbb{Q}}^{\text{div}}$ , we have the inequality

$$(139) \quad H^{\text{NA}}(\mathcal{F}_v) \leq \tilde{\beta}(v).$$

Moreover, if  $(\mathcal{X}, -K_{\mathcal{X}}, a\eta)$  is a special test configuration, then equality holds for  $v = av_{\mathcal{X}_0} = a \cdot r(\text{ord}_{\mathcal{X}_0})$ . (See Definition 2.12.)

**Proof** The inequality follows immediately from

$$(140) \quad \inf_w (A(w) + \phi_w(w)) \leq A(v) + \phi_v(v) = A(v).$$

When  $(\mathcal{X}, -K_{\mathcal{X}}, a\eta)$  is a special test configuration and  $v = av_{\mathcal{X}_0}$ , then

$$(141) \quad L^{\text{NA}}(\mathcal{X}, -K_{\mathcal{X}}, a\eta) = aL^{\text{NA}}(\mathcal{X}, -K_{\mathcal{X}}, \eta) = a(\text{lct}(\mathcal{X}; \mathcal{X}_0) - 1) = 0.$$

On the other hand, by (46),

$$(142) \quad L^{\text{NA}}(\mathcal{X}, -K_{\mathcal{X}}, a\eta) = L^{\text{NA}}(\mathcal{F}_v(-A(v))) = L^{\text{NA}}(\mathcal{F}_v) - A(v).$$

So we get

$$(143) \quad H^{\text{NA}}(\mathcal{F}_v) = L^{\text{NA}}(\mathcal{F}_v) - \tilde{S}^{\text{NA}}(\mathcal{F}_v) = A(v) - \tilde{S}^{\text{NA}}(v) = \tilde{\beta}(v). \quad \square$$

**Lemma 4.3** For any  $\phi = \phi_{\mathcal{F}}$  and  $v \in X_{\mathbb{Q}}^{\text{div}}$ , we have the inequality

$$(144) \quad \tilde{S}(v) + \phi(v) \geq \tilde{S}^{\text{NA}}(\phi).$$

**Proof** We use the same argument as in [52, Section 4.1]. Set  $\gamma = \phi(v)$ . Then by the argument there, we have  $\lambda_{\min} = \lambda_{\min}(\mathcal{F}) \leq \gamma$  and we can then estimate

$$\begin{aligned} e^{-\tilde{S}^{\text{NA}}(\phi)} &= \mathcal{Q}(\phi) = \frac{1}{V} \int_{\mathbb{R}} e^{-x} (-d\text{vol}(\mathcal{F}^{(x)})) = e^{-\lambda_{\min}} - \frac{1}{V} \int_{\lambda_{\min}} e^{-x} \text{vol}(\mathcal{F}^{(x)} R_{\bullet}) dx \\ &\geq e^{-\gamma} - \frac{1}{V} \int_{\gamma}^{+\infty} e^{-x} \text{vol}(\mathcal{F}^{(x)} R_{\bullet}) dx \geq e^{-\gamma} - \frac{1}{V} \int_{\gamma}^{+\infty} e^{-x} \text{vol}(\mathcal{F}_v^{(x-a)}) dx \\ &= e^{-\gamma} - e^{-\gamma} \frac{1}{V} \int_0^{+\infty} e^{-x} \text{vol}(\mathcal{F}_v^{(t)}) dt = e^{-\gamma} \frac{1}{V} \int_0^{+\infty} e^{-x} (-d\text{vol}(\mathcal{F}_v^{(t)})) \\ &= e^{-\phi(v)} e^{-\tilde{S}^{\text{NA}}(v)}. \end{aligned} \quad \square$$

**Proposition 4.4** For any  $\mathbb{Q}$ -Fano variety, we have the identity

$$(145) \quad h(X) = \inf_{v \in X_{\mathbb{Q}}^{\text{div}}} \tilde{\beta}(v).$$

**Proof** For any test configuration  $(\mathcal{X}, \mathcal{L}, a\eta)$ , by Theorem 3.5 there exists a special test configuration  $(\mathcal{X}^s, \mathcal{L}^s, a^s \eta^s)$  such that

$$(146) \quad H^{\text{NA}}(\mathcal{X}, \mathcal{L}, a\eta) \geq H^{\text{NA}}(\mathcal{X}^s, \mathcal{L}^s, a^s \eta^s) = \tilde{\beta}(a^s v_{\mathcal{X}_0^s}).$$

The last identity is from Lemma 4.2. This together with Corollary 3.6 implies identity (145).

Alternatively, recall that  $L^{\text{NA}}(\phi) = \inf_{v \in X_{\mathbb{Q}}^{\text{div}}} (A_X(v) + \phi(v))$ . So for any  $\epsilon > 0$  we can choose  $v$  such that  $A_X(v) + \phi(v) < L^{\text{NA}}(\phi) + \epsilon$ . We can then use  $v$  in (144) to get

$$L^{\text{NA}}(\phi) - \tilde{S}^{\text{NA}}(\phi) \geq A_X(v) + \phi(v) - \epsilon - (\phi(v) + \tilde{S}(v)) = \tilde{\beta}(v) - \epsilon.$$

Since  $\epsilon$  is arbitrary, we can use (97) to get the identity (145). □

With the identity (145) and Proposition 2.32, we get:

**Corollary 4.5** For any  $\mathbb{Q}$ -Fano variety, we have the equality

$$(147) \quad h(X) = \inf_{\mathcal{F}} H^{\text{NA}}(\mathcal{F}).$$

The next result should be compared to Lemma 3.7.

**Proposition 4.6** For any  $v \in \mathring{\text{Val}}(X)$ , there exists a unique  $a_* = a_*(v) \geq 0$  such that

$$(148) \quad \tilde{\beta}(a_*v) = \inf_{a>0} \tilde{\beta}(av) =: \tilde{\beta}_*(v).$$

When  $\beta(v) \geq 0$ , then  $a_* = 0$ , so that  $a_*v$  is the trivial valuation and  $\tilde{\beta}_*(v) = 0$ . Otherwise,  $a_*(v) > 0$  and  $\tilde{\beta}_*(v) < 0$ .

**Proof** Consider the function defined on  $\mathbb{R}_{\geq 0}$  by

$$(149) \quad \begin{aligned} f(a) &= A(av) - \tilde{\mathcal{S}}^{\text{NA}}(av) = aA(v) + \log\left(\frac{1}{V} \int_0^{+\infty} e^{-x} \text{DH}(\mathcal{F}_{av})\right) \\ &= aA(v) + \log\left(\frac{1}{V} \int_0^{+\infty} e^{-ax} \text{DH}(\mathcal{F}_v)\right). \end{aligned}$$

We will show that  $a \mapsto f(a)$  is convex and goes to  $+\infty$  as  $a \rightarrow +\infty$ . Now

$$\begin{aligned} f'(a) &= A(v) - \frac{\int_0^{+\infty} x e^{-ax} \text{DH}(\mathcal{F}_v)}{\int_0^{+\infty} e^{-ax} \text{DH}(\mathcal{F}_v)}, \\ f''(a) &= \frac{\int x^2 e^{-ax} \text{DH}}{\int e^{-ax} \text{DH}} - \frac{(\int x e^{-ax} \text{DH})^2}{(\int e^{-ax} \text{DH})^2} = \|x - \bar{x}\|_{L^2(dv)}^2 \geq 0, \end{aligned}$$

where

$$(150) \quad dv = \frac{e^{-ax} \text{DH}}{\int e^{-ax} \text{DH}} \quad \text{and} \quad \bar{x} = \int x \, dv.$$

So  $f''(a) = 0$  if and only if  $av$  is trivial. Moreover,  $f'(0) = A(v) - (1/V) \int_0^{+\infty} x \text{DH}(\mathcal{F}_v) = \beta(v)$ .

On the other hand,  $f(0) = 0$  and we claim that  $\lim_{a \rightarrow +\infty} f(a) = +\infty$ , which then implies the statement. To prove this divergence, we set  $g(x) = V^{-1/n} \text{vol}(\mathcal{F}^{(x)} R_{\bullet})^{1/n}$ . Then  $g(x)$  is decreasing, and concave on  $[0, \lambda_{\max}]$  by Theorem 2.5. As a consequence, the subset  $\{x \in \mathbb{R}_{\geq 0} \mid g'(x) \text{ exists}\}$  is dense in  $\mathbb{R}_{\geq 0}$ , by Aleksandrov's differentiability theorem for concave functions. Fix  $0 < \epsilon \ll \lambda_{\max}$  such that  $g'(\epsilon)$  exists and  $g(\epsilon) < g(0) = 1$ . Setting  $C = -g'(\epsilon) > 0$  and  $T = (1 + C\epsilon)/C$ , define a function

$$(151) \quad \hat{g}(x) = \begin{cases} 1 & \text{if } x \in [0, \epsilon], \\ 1 + C\epsilon - Cx & \text{if } x \in (\epsilon, T], \\ 0 & \text{if } x \in (T, +\infty). \end{cases}$$

Then  $\hat{g}(x) \geq g(x)$  over  $[0, +\infty)$ , by concavity. Then we calculate to get

$$(152) \quad a \int_0^{+\infty} \hat{g}^n(x) e^{-ax} \, dx = 1 - nCm_{n-1},$$

where  $m_k = \int_{\epsilon}^T (1 + C\epsilon - Cx)^k e^{-ax} dx$  satisfies

$$m_k = \frac{1}{a} e^{-a\epsilon} - \frac{kC}{a} m_{k-1} = \frac{1}{a} e^{-a\epsilon} - \frac{kC}{a} \left( \frac{1}{a} e^{-a\epsilon} - \frac{(k-1)C}{a} m_{k-2} \right).$$

Using induction we get  $m_{n-1} = a^{-1} e^{-a\epsilon} (1 + O(a^{-1}))$ . So

$$\begin{aligned} e^{-\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_{av})} &= 1 - a \int_0^{+\infty} g^n(x) e^{-ax} dx \geq 1 - a \int_0^{+\infty} \hat{g}^n(x) e^{-ax} dx \\ &= nCm_{n-1} = nCa^{-1} e^{-a\epsilon} (1 + O(a^{-1})). \end{aligned}$$

So we get  $-\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_{av}) \geq -\log a - a\epsilon + O(1)$ , giving

$$(153) \quad f(a) = \tilde{\beta}(av) \geq (A(v) - \epsilon)a - \log a + O(1),$$

which approaches  $+\infty$  as  $a \rightarrow +\infty$  if we choose  $0 < \epsilon < A(v)$ . □

**Remark 4.7** By the above proof, we get an estimate: for any  $C_1 > 0$ , there exists a  $C_2 = C_2(C_1, v) > 0$  such that for any  $w \in \text{Val}(X)$  with  $w \leq C_1 v$ , we have

$$(154) \quad a_*(w) \leq \frac{C_2}{A(v)}.$$

**Corollary 4.8** We always have  $h(X) \leq 0$ , with equality holding if and only if  $h(X) = 0$ .

**Proof** By [37; 50],  $X$  is  $\mathbb{K}$ -semistable if and only if  $\beta(v) \geq 0$ , which implies  $\tilde{\beta}_*(v) = 0$ . If  $X$  is not  $\mathbb{K}$ -semistable then there exists  $v'$  such that  $\beta(v') < 0$ . By Proposition 4.6, we then have  $\tilde{\beta}_*(v') < 0$ , which implies  $h(X) < 0$ . □

**Lemma 4.9** If  $v$  computes  $h(X)$ , then  $v$  is the unique valuation, up to rescaling, that computes  $\text{lct}(\mathfrak{a}_\bullet(v))$ .

**Proof** Recall that  $\text{lct}(\mathfrak{a}_\bullet) = \inf_w A(w)/w(\mathfrak{a}_\bullet(w))$ . For any  $w \in \text{Val}(X)$ , assume that  $w(\mathfrak{a}_\bullet(v)) = a > 0$ . Then  $a^{-1}w \geq v$ . By Proposition A.1, the function

$$w \mapsto \tilde{\mathcal{S}}^{\text{NA}}(w) = -\log \frac{1}{V} \int_{\mathbb{R}} e^{-\lambda} \text{DH}(\mathcal{F}_w)$$

is strictly increasing on  $\text{Val}(X)$ . So we use the assumption to get

$$\begin{aligned} \frac{A(w)}{w(\mathfrak{a}_\bullet(v))} &= A(a^{-1}w) = A(a^{-1}w) - \tilde{\mathcal{S}}^{\text{NA}}(a^{-1}w) + \tilde{\mathcal{S}}^{\text{NA}}(a^{-1}w) \\ &\geq A(v) - \tilde{\mathcal{S}}^{\text{NA}}(v) + \tilde{\mathcal{S}}^{\text{NA}}(v) = A(v) \\ &= \frac{A(v)}{v(\mathfrak{a}_\bullet(v))}. \end{aligned}$$

When the equality holds, then  $a^{-1}w = v$ . □

We now observe that the method developed in [14] can be used to prove:

**Theorem 4.10** For any  $\mathbb{Q}$ -Fano variety, there exists a minimizing valuation of  $\tilde{\beta}$  which is quasi-monomial.

Since the argument is almost verbatim to [14] except for the continuity property of  $\tilde{\beta}$ , we just give a sketch of key points and explain the required continuity of  $\tilde{\beta}$  in Section 8. Without the properties of  $\tilde{\beta}(S)$  explained in Section 8, the existence of a valuation calculating  $h(X)$  (but without the quasimonomial property) can also be obtained using the argument in [11, Section 6].

**Proof** By Corollary 3.6,  $h(X) = \inf_E \tilde{\beta}_*(E)$ , where  $E$  ranges over prime divisors over  $X$  that induce special test configurations of  $(X, -K_X)$ . By [14, Theorem A.2], we know that such an  $E$  is an lc place of an  $N$ -complement  $D$  of  $X$ , where  $N$  depends only on the dimension  $n$  (this depends on the deep result of Birkar about the boundedness of  $\mathbb{Q}$ -complements). So we have

$$(155) \quad h(X) = \inf_v \tilde{\beta}_*(v),$$

where  $v$  ranges over all divisorial valuations that are lc places of an  $N$ -complement. For such a valuation  $v$ , there exists a  $D \in (1/N)|-NK_X|$  such that  $(X, D)$  is lc and  $A_{(X,D)}(v) = 0$ . We then parametrize such  $\mathbb{Q}$ -divisors as in [14, Proof of Theorem 4.5]. Set  $W = \mathbb{P}(H^0(X, \mathcal{O}_X(-NK_X)))$  and denote by  $H$  the universal divisor on  $X \times W$  parametrizing divisors in  $|-NK_X|$  and set  $D := (1/N)H$ . By the lower semicontinuity of log canonical thresholds, the locus  $Z = \{w \in W \mid \text{lct}(X_w; D_w) = 1\}$  is locally closed in  $W$ . For each  $z \in Z$ , set  $b_z := \inf_v \tilde{\beta}(v)$ , where  $v$  ranges over all  $v \in \text{Val}(X)$  with  $A_{(X,D_z)}(v) = 0$ .

Let  $g: Y_z \rightarrow X$  be a log resolution of  $(X, D_z)$ . Write  $K_Y + D_{Y_z} = g^*(K_X + D_z)$ . Consider the section of the simplicial cone,  $\mathcal{S} := \text{QM}(Y_z, D_{Y_z}) \cap \{v \in \text{Val}(X) \mid A(v) = 1\}$ . By Proposition 4.6, we know that for each  $v \in \mathcal{S}$  there exists  $a_*(v)$  such that  $\inf_{a>0} \tilde{\beta}(av) = \tilde{\beta}(a_*(v)v) =: \tilde{\beta}_*(v)$ . By Izumi's estimate (see [48, Example 11.3.9; 44, Proposition 5.10] for the smooth case, and [51, Section 3] in the klt case), we know that there exists  $C_1 > 0$  such that for any  $v \in \mathcal{S}$  we have  $v \leq C_1 \cdot \text{ord}_F$ , where  $F = \bigcap_i D_{Y_z, i}$ . Now by the proof of Proposition 4.6 (see Remark 4.7), we know that  $a_*(v)$  is uniformly bounded for any  $v \in \mathcal{S}$ . By Proposition A.2, we know that  $v \mapsto \tilde{\beta}(v)$  is continuous on  $\text{QM}(Y_z, D_{Y_z})$  and hence is uniformly continuous over compact subsets. We then get the continuity of  $v \mapsto \tilde{\beta}_*(v)$  over the compact set  $\mathcal{S}$ . So we know that there exists  $v_z^* \in \mathcal{S}$  such that  $\tilde{\beta}_*(v_z^*) = \min_{v \in \mathcal{S}} \tilde{\beta}_*(v)$  and  $a_*(v_z^*) \cdot v_z^*$  is then a minimizer of  $\tilde{\beta}$  over  $\text{QM}(Y_z, D_{Y_z})$ .

Then as [14, Proof of Theorem 4.5], choose a locally closed decomposition  $Z = \bigcup_{i=1}^r Z_i$  so that  $Z_i$  is smooth and there is an étale map  $Z'_i \rightarrow Z_i$  such that  $(X_{Z'_i}, D_{Z'_i})$  admits fiberwise log resolutions. By the same arguments as [14, Proof of Propositions 4.1 and 4.2], which depend on the deformation invariance of log plurigenera in the work of Hacon, McKernan and Xu, we know that  $b_z$  is independent of  $z \in Z_i$ . So  $b_z$  takes finitely many values and there is a  $z_0 \in Z$  such that  $h(X) = \min_{z \in Z} b_z = b_{z_0}$  is computed by  $v_{z_0}^*$ .  $\square$

As in the case of normalized volume, we expect the following:

**Conjecture 4.11** *The minimizer  $v_*$  is unique, and is special, which means that  $\mathcal{F}_{v_*}$  is a special  $\mathbb{R}$ -test configuration.*

**Remark 4.12** As [11, Proposition 4.11], using Lemma 4.9 one can show that any divisorial (ie rational rank one) minimizing valuation is primitive and plt.

Besides the case of stability threshold treated in [14], in the local setting of normalized volumes, the existence of quasimonomial minimizers is also known thanks to the work of Blum [10] and Xu [78]. Moreover, one might also be able to adapt the techniques in Xu and Zhuang [80] to the current global setting to prove the uniqueness of minimizing valuations.<sup>3</sup> We will prove in Section 6 the uniqueness of special minimizers (in a similar spirit to the work in [58; 57; 55]).

### 5 Initial term degeneration of filtrations

Let  $\mathcal{F}_0$  be a special  $\mathbb{R}$ -test configuration of  $(X, -K_X)$  with central fiber  $(W := \text{Proj}(\text{Gr}(\mathcal{F}_0)), \xi_0 := \xi_{\mathcal{F}_0})$ . Let  $\mathcal{F}_1$  be another filtration of  $R$ . We define a filtration on

$$(156) \quad R' := R(W, -K_W) = \bigoplus_{m \geq 0} \bigoplus_{\lambda \in \Gamma(\mathcal{F}_0)} t^{-\lambda} \mathcal{F}_0^\lambda R_m / \mathcal{F}_0^{>\lambda} R_m =: \bigoplus_{m \geq 0} R'_m$$

in the following way. Recall that we can write

$$(157) \quad R'_m = \bigoplus_{\alpha \in M_{\mathbb{Z}}} t^{-\langle \alpha, \xi_0 \rangle} \mathcal{F}_0^{\langle \alpha, \xi_0 \rangle} R_m / \mathcal{F}_0^{>\langle \alpha, \xi_0 \rangle} R_m.$$

For any  $f \in R_m$ , set

$$(158) \quad \mathbf{in}_{\mathcal{F}_0}(f) = (t^{-\langle \alpha, \xi \rangle} \bar{f})(0) =: f' \in \mathcal{F}_0^{\langle \alpha, \xi_0 \rangle} R_m / \mathcal{F}_0^{>\langle \alpha, \xi_0 \rangle} R_m, \quad \text{where } \langle \alpha, \xi_0 \rangle = v_{\mathcal{F}_0}(f).$$

For any  $\lambda \in \mathbb{R}$ , take the Gröbner base-type degeneration

$$(159) \quad \mathcal{F}_1^\lambda R'_m = \text{Span}_{\mathbb{C}}\{\mathbf{in}_{\mathcal{F}_0}(f) \mid f \in \mathcal{F}_1^\lambda R_m\} \subseteq R'_m.$$

Note that because  $R'$  is integral,  $\mathbf{in}_{\mathcal{F}_0}(fg) = \mathbf{in}_{\mathcal{F}_0}(f) \cdot \mathbf{in}_{\mathcal{F}_0}(g)$  if  $f \in R_{m_1}$  and  $g \in R_{m_2}$ . So in this way, we get a  $\mathbb{T}_0$ -equivariant filtration

$$(160) \quad \mathcal{F}_1^\lambda R'_m = \bigoplus_{\alpha \in \mathbb{Z}^{r_0}} \mathcal{F}_1^\lambda R'_{m,\alpha}.$$

There is an equivalent way to describe  $\mathcal{F}_1^\lambda R'_m$ , as follows. For any  $f' \in R'_{m,\alpha}$ , we choose  $f \in R_m$  such that  $f' = t^{-\langle \alpha, \xi_0 \rangle} \bar{f}(0)$ . Then we have

$$(161) \quad \mathcal{F}_1^\lambda R'_{m,\alpha} = \{f' \in R'_{m,\alpha} \mid f + h \in \mathcal{F}_1^\lambda R_m \text{ for some } h \in \mathcal{F}_0^{>\langle \alpha, \xi_0 \rangle} R_m\}.$$

This is well defined since  $f$  is determined up to addition by elements from  $\mathcal{F}_0^{>\langle \alpha, \xi_0 \rangle} R_m$ .

Note that this construction allows us to find a basis  $\mathcal{B} = \{s_1, \dots, s_{N_m}\}$  of  $R_m$  that is compatible with both  $\mathcal{F}_0 R_m$  and  $\mathcal{F}_1 R_m$ . Recall that this means that for any  $\lambda \in \mathbb{R}$  and  $i = 0, 1$ , there exists a subset of  $\mathcal{B}$  which depends on  $\lambda$  and  $i$  and is a basis of  $\mathcal{F}_i^\lambda R_m$ . To find such a basis, we can first find a basis  $\mathcal{B}'_\alpha$  of  $R'_{m,\alpha}$  which is compatible with  $\mathcal{F}_1 R'_{m,\alpha}$ . Then  $\mathcal{B} = \bigcup_\alpha \mathcal{B}'_\alpha =: \{f'_1, \dots, f'_{N_m}\}$  is a basis compatible with

<sup>3</sup>This has indeed been recently achieved in [15].

both  $\mathcal{F}'_1 R_m$  and  $\mathcal{F}'_{\text{wt}_{\xi_0}} R'_m$ . For each  $f'_k \in R'_{m,\alpha_k}$ , there exists  $\lambda_k \in \mathbb{R}$  such that  $f'_k \in \mathcal{F}'_1^{\lambda_k} R'_m \setminus \mathcal{F}'_1^{>\lambda_k} R'_m$ . Then by (161), there exists  $h_k \in \mathcal{F}_0^{>(\alpha_k, \xi)} R_m$  such that  $s_k := f_k + h_k \in \mathcal{F}_1^{\lambda_k} R_m$ . Moreover, we have  $s_k \notin \mathcal{F}_1^{>\lambda_k} R_m$  since otherwise  $\mathbf{in}(s_k) = \mathbf{in}(f_k) = f'_k \in \mathcal{F}'_1^{>\lambda_k} R'_m$ . It is easy to verify that  $\{s_k\}$  is the desired basis. So the relative successive minima of  $\mathcal{F}_1$  with respect to  $\mathcal{F}_0$  (see [20]) is given by the set  $\{\lambda_k - \langle \alpha_k, \xi_0 \rangle\}$ , which is the same as the relative successive minima of  $\mathcal{F}'_1 := \mathcal{F}'_1 R'$  with respect to  $\mathcal{F}'_0 := \mathcal{F}'_{\text{wt}_{\xi_0}} R'$ . This immediately proves a useful fact:

**Lemma 5.1** *With the above constructions and notation, we have the identity*

$$(162) \quad d_2^X(\mathcal{F}_0, \mathcal{F}_1) = d_2^W(\mathcal{F}'_0, \mathcal{F}'_1).$$

Since the initial term degeneration does not change the dimension of vector spaces, it is clear that the successive minima of  $\mathcal{F}_1$  and  $\mathcal{F}'_1$  coincide. As a consequence, we get

$$(163) \quad \tilde{\mathbf{S}}_X^{\text{NA}}(\mathcal{F}_1) = \tilde{\mathbf{S}}_W^{\text{NA}}(\mathcal{F}'_1).$$

On the other hand, consider the  $\mathbb{T}_0$ -equivariant graded filtration of the Rees algebra  $\mathcal{R}' := \mathcal{R}(\mathcal{F}_0)$  (see (30)) given by

$$(164) \quad \mathcal{F}'^\lambda \mathcal{R}'_{m,\alpha} = \{s = t^{-\langle \alpha, \xi \rangle} \bar{f} \in \mathcal{R}'_{m,\alpha} \mid t^{-\lambda} \bar{f} \in \mathcal{R}(\mathcal{F}_1)\}.$$

Then  $\mathcal{F}'^\lambda \mathcal{R}'$  coincides with  $\mathcal{F}R$  on the generic fiber and coincides with  $\mathcal{F}'R'$  on the central fiber. By the lower semicontinuity of  $\text{lct}$  for a family, it is easy to see that  $\hat{\mathbf{L}}^{\text{NA}}$  in (80) is also lower semicontinuous for a family. This is standard if  $\mathcal{F}_0$  has rank one, which corresponds to a special test configuration; see [47, Lemma 8.1; 13, Proof of Lemma 6.5]. In general, one can restrict to a generic curve passing through 0 in the family in Teissier’s construction in the paragraph above Lemma 2.11; alternatively, see Remark 6.2. So we can get

$$(165) \quad \mathbf{L}^{\text{NA}}(\mathcal{F}_1) = \hat{\mathbf{L}}_X^{\text{NA}}(\mathcal{F}_1) \geq \hat{\mathbf{L}}_W^{\text{NA}}(\mathcal{F}'_1) = \mathbf{L}^{\text{NA}}(\mathcal{F}'_1),$$

where the first and the last identity come from Proposition 2.32. Combining the above discussion, we get the inequality

$$(166) \quad \mathbf{H}_X^{\text{NA}}(\mathcal{F}_1) \geq \mathbf{H}_W^{\text{NA}}(\mathcal{F}'_1).$$

**Theorem 5.2** *Assume  $v$  induces a special  $\mathbb{R}$ -test configuration  $\mathcal{F}_v$  of  $X$ . Then  $v$  is a minimizer of  $\tilde{\beta}$  over  $\text{Val}(X)$  if and only if  $v$  is Ding-semistable (or equivalently  $K$ -semistable).*

**Proof** For simplicity of notation, set  $\mathcal{F}_0 = \mathcal{F}_v$  and  $(W, \xi_0) := (X_{\mathcal{F}_v, 0}, \xi_{\mathcal{F}_0})$  and let  $\mathbb{T}_0$  be the torus generated by  $\xi_0$ .

We first prove that minimizer is Ding-semistable. Suppose  $(W, \xi_0)$  is not Ding-semistable. Then by Theorem 2.46 from [39], there exists a  $\mathbb{T}$ -equivariant special test configuration  $(\mathcal{W}, -K_{\mathcal{W}})$  of  $(W, -K_W)$  with central fiber  $Y := \mathcal{W}_0$  such that

$$(167) \quad \mathbf{D}_g^{\text{NA}}(\mathcal{W}, -K_{\mathcal{W}}) = \text{Fut}_{Y, \xi}(\eta) < 0.$$

We can now construct a family of valuations  $\{v_\epsilon\}$  such that  $v_\epsilon$  induces special test configurations with central fiber  $Y$  and corresponds to a vector field  $\xi_\epsilon = \xi_0 + \epsilon\eta$  on  $Y$ . This can be done by using the cone construction to reduce to the situation in [58, Section 6] or [57, Proof of Theorem 2.64]. Alternatively, one can use an argument involving the Hilbert scheme as in [55, Proof of Lemma 3.1].

Here we will use the Chow variety to explain this construction. Recall that the Chow point of a cycle  $Z \subset \mathbb{P}^{N-1}$  of degree  $d$  and dimension  $n$  corresponds to a divisor in the Grassmannian  $\text{Gr}(N - n - 1, \mathbb{C}^N)$  which is the zero scheme of a section:

$$\text{CH}(Z) \in H^0(\text{Gr}(N - n - 1, \mathbb{C}^N), \mathcal{O}(d)) =: \mathbb{M}.$$

$\text{CH}(Z)$  is determined up to rescaling and we call it the Chow coordinate of  $Z$ . Let  $\text{CH}(X)$ ,  $\text{CH}(W)$  and  $\text{CH}(Y)$  be the Chow coordinates of  $X$ ,  $W$  and  $Y$ , respectively. Denote by  $[\text{CH}(X)]$  the Chow point of  $X$  in the projectivization  $\mathbb{P}(\mathbb{M})$ , and similarly for  $Y$  and  $W$ . Since the  $\mathbb{T}$ -action on  $\mathbb{P}^{N-1}$  induces a weight decomposition  $\mathbb{M} = \bigoplus_\alpha \mathbb{M}_\alpha$ , we have

$$(168) \quad \lim_{s \rightarrow +\infty} \sigma_\xi(s) \circ [\text{CH}(X)] = [\text{CH}(W)] \quad \text{and} \quad \lim_{s \rightarrow +\infty} \sigma_\eta(s) \circ [\text{CH}(W)] = [\text{CH}(Y)].$$

If we set

$$(169) \quad \text{CW}_\xi(X) = \min\{\langle \alpha, \xi \rangle \mid \text{CH}(X)_\alpha \neq 0\} \quad \text{and} \quad \text{CW}_\eta(W) = \min\{\langle \alpha, \eta \rangle \mid \text{CH}(W)_\alpha \neq 0\},$$

then

$$[\text{CH}(W)] = \left[ \sum_{\alpha \in I_W} \text{CH}(X)_\alpha \right], \quad \text{where } I_W = \{\alpha \mid \text{CH}(X)_\alpha \neq 0, \langle \alpha, \xi \rangle = \text{CW}_\xi(X)\},$$

$$[\text{CH}(Y)] = \left[ \sum_{\alpha \in I_Y} \text{CH}(W)_\alpha \right], \quad \text{where } I_Y = \{\alpha \mid \text{CH}(W)_\alpha \neq 0, \langle \alpha, \eta \rangle = \text{CW}_\eta(W)\}.$$

Note that  $I_Y \subseteq I_W$ . For any  $\alpha \in M_{\mathbb{Z}}$  with  $\text{CH}(X)_\alpha \neq 0$ , we have that  $\langle \alpha, \xi \rangle \geq \text{CW}_\xi(X)$ , with equality if and only if  $\alpha \in I_W$ . Similarly, for any  $\alpha \in M_{\mathbb{Z}}$  with  $\text{CH}(W)_\alpha \neq 0$  (and hence  $\text{CH}(X)_\alpha \neq 0$ ), we have that  $\langle \alpha, \eta \rangle \geq \text{CW}_\eta(W)$ , with equality if and only if  $\alpha \in I_Y$ . So when  $0 < \epsilon \ll 1$  and for any  $\text{CH}(X)_\alpha \neq 0$ , we have that  $\langle \alpha, \xi + \epsilon\eta \rangle \geq \text{CW}_\xi(X) + \epsilon\text{CW}_\eta(W)$ , with equality if and only if  $\alpha \in I_Y$ . So we get

$$\lim_{t \rightarrow 0} \sigma_{\xi + \epsilon\eta}(t) \circ [\text{CH}(X)] = \lim_{t \rightarrow 0} \left[ \sum_{\alpha} t^{\langle \alpha, \xi + \epsilon\eta \rangle} \text{CH}(X)_\alpha \right] = [\text{CH}(Y)].$$

So for  $0 < \epsilon \ll 1$ ,  $\xi + \epsilon\eta$  induces an  $\mathbb{R}$ -test configuration that degenerates  $X$  to  $Y$ . By Lemma 2.11, we get the corresponding valuations  $v_\epsilon$ .

Now we use the identity (113) to get

$$(170) \quad \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \tilde{\beta}(v_\epsilon) = \frac{d}{d\epsilon} \mathbf{H}_Y^{\text{NA}}(\mathcal{F}_{\text{wt}_{\xi + \epsilon\eta}}) = \text{Fut}_{Y, \xi}(\eta) < 0.$$

But this contradicts the assumption that  $v_0 = v$  is the minimizer of  $\tilde{\beta}$ .



Conversely, we need to show that a Ding-semistable valuation is a minimizer. Let  $(X, \mathcal{L}, a\eta)$  be any special test configuration of  $(X, -K_X)$  and  $\mathcal{F}_1 = \mathcal{F}_{(X, \mathcal{L}, a\eta)}$  be the associated filtration. We consider the initial term degeneration of  $\mathcal{F}_1$  with respect to  $\mathcal{F}_0$  defined as above. Then we can use (166) to get

$$H_X^{\text{NA}}(\mathcal{F}_1) \geq H_W^{\text{NA}}(\mathcal{F}'_1) \geq H_W^{\text{NA}}(\mathcal{F}_{\text{wt}_{\xi_0}}) = H^{\text{NA}}(\mathcal{F}_0) = \tilde{\beta}(v),$$

where the second inequality follows from the results in Lemma 6.1 in the next section and the assumption that  $(W, \xi_0)$  is Ding-semistable. □

## 6 Uniqueness of minimizing special $\mathbb{R}$ -test configurations

We prove Theorem 1.2 in this section. We first generalize the formula (113). Let  $(X, -K_X, \mathbb{T}, \xi)$  be the data as before and  $\mathcal{F}$  be a  $\mathbb{T}$ -equivariant filtration. We consider a family of  $\mathbb{T}$ -equivariant filtrations

$$(171) \quad \mathcal{F}_s = s\mathcal{F}_{((1-s)/s)\xi} \quad \text{for } s \in (0, 1], \text{ with } \mathcal{F}_0 = \mathcal{F}_{\text{wt}_{\xi}} \text{ and } \mathcal{F}_1 = \mathcal{F},$$

which interpolates  $\mathcal{F}_{\text{wt}_{\xi}}$  and  $\mathcal{F}$ .

**Lemma 6.1** *For the family of filtrations (171), the following statements hold true:*

- (i) *The map  $s \mapsto H^{\text{NA}}(\mathcal{F}_s)$  is smooth and convex. It is affine if and only if  $G_{\mathcal{F}}$  is a multiple of  $\langle x, \xi \rangle$ .*
- (ii) *We have the derivative formula*

$$(172) \quad \left. \frac{d}{ds} \right|_{s=0} H^{\text{NA}}(\mathcal{F}_s) = \beta_{\xi}(\mathcal{F}_{-\xi}).$$

To get (113) from (172), we just need to set  $\mathcal{F} = \mathcal{F}_{\text{wt}_{\xi} + \eta}$  so that  $\mathcal{F}_s = \mathcal{F}_{\xi + s\eta}$ . Moreover, we fix a faithful valuation that is adapted to the torus action (see Definition 2.21) and will freely use the associated Newton–Okounkov body  $\Delta = \Delta(-K_X)$  of  $(X, -K_X)$ .

**Proof** By Lemma 2.22 and (25), as functions on  $\Delta = \Delta(-K_X)$ , we have

$$(173) \quad G(s, y) := G_{\mathcal{F}_s}(y) = (1-s)\langle y, \xi \rangle + sG_{\mathcal{F}}(y).$$

So, by using Lemma 2.35, we get

$$(174) \quad L^{\text{NA}}(\mathcal{F}_s) = sL^{\text{NA}}(\mathcal{F}),$$

$$(175) \quad -\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_s) = \log\left(\frac{n!}{V} \int_{\Delta} e^{-G(s,y)} dy\right).$$

$L^{\text{NA}}(\mathcal{F}_s)$  is linear in  $s$  and  $-\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_s)$  is smooth in  $s$ . By Hölder’s inequality,  $-\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_s)$  is strictly convex in  $s$  unless  $G_{\mathcal{F}}$  is a multiple of  $\langle x, \xi \rangle$ . This implies that  $H^{\text{NA}}(\mathcal{F}_s) = L^{\text{NA}} - \mathcal{S}^{\text{NA}}$  is convex in  $s \in [0, 1]$ .

To see (172), we calculate

$$\begin{aligned} \left. \frac{d}{ds} \right|_{s=0} H^{\text{NA}}(\mathcal{F}_s) &= L^{\text{NA}}(\mathcal{F}) + \frac{\int_{\Delta} (\langle y, \xi \rangle - G_{\mathcal{F}}(y)) e^{-G(0,y)} dx}{\int_{\Delta} e^{-G(0,y)} dy} \\ &= L^{\text{NA}}(\mathcal{F}_{-\xi}) - \frac{n!}{V_{\xi}} \int_{\Delta} G_{\mathcal{F}_{-\xi}}(y) e^{-\langle y, \xi \rangle} dy = \beta_{\xi}(\mathcal{F}_{-\xi}). \end{aligned} \quad \square$$

Assume that there are two special  $\mathbb{R}$ –test configurations  $\mathcal{F}_i = \{\mathcal{F}_i R_m\}$  for  $i = 0, 1$  of  $(X, -K_X)$  that minimize  $\mathbf{H}^{\text{NA}}$ . By [Theorem 5.2](#), the central fibers  $(W^{(i)} := \text{Proj}(\text{Gr}_{\mathcal{F}_i}), \xi_i = \xi_{\mathcal{F}_i})$  are both Ding-semistable. Now consider the initial term degeneration of  $\mathcal{F}_1$  with respect to  $\mathcal{F}_0$  as in the above section. We get a  $\mathbb{T}_0$ –equivariant filtration  $\mathcal{F}'_1$  on  $R' = R(W^{(0)}, -K_{W^{(0)}})$  and by [\(166\)](#),  $\mathbf{H}_X^{\text{NA}}(\mathcal{F}_1) \geq \mathbf{H}_{W^{(0)}}^{\text{NA}}(\mathcal{F}'_1)$ .

Now, as at the beginning of this section, consider the family of filtrations that interpolates  $\mathcal{F}'_1$  and  $\mathcal{F}_{\text{wt}_{\xi_0}} R' =: \mathcal{F}'_{\text{wt}_{\xi_0}}$ ,

$$(176) \quad \mathcal{F}'_s := s\mathcal{F}'_{((1-s)/s)\xi_0} R'.$$

Applying [Lemma 6.1](#) to  $(W^{(0)}, \xi_0, \mathcal{F}'_s)$ , we know that  $\mathbf{D}(s) := \mathbf{H}^{\text{NA}}(\mathcal{F}'_s)$  is convex in  $s \in [0, 1]$ . Moreover we have the relation

$$(177) \quad \mathbf{D}(0) = \mathbf{H}_{W^{(0)}}^{\text{NA}}(\mathcal{F}_{\text{wt}_{\xi_0}}) = \mathbf{H}_X^{\text{NA}}(\mathcal{F}_0) = \mathbf{H}_X^{\text{NA}}(\mathcal{F}_1) \geq \mathbf{H}^{\text{NA}}(\mathcal{F}'_1) = \mathbf{D}(1).$$

The 3<sup>rd</sup> identity is by [Theorem 5.2](#), that the  $\mathcal{F}_i$  for  $i = 0, 1$  both obtain the minimum of  $\mathbf{H}^{\text{NA}}$ .

On the other hand, by [\(172\)](#),

$$\frac{d}{ds} \Big|_{s=0} \mathbf{H}^{\text{NA}}(\mathcal{F}'_s) = \beta_{\xi_0}(\mathcal{F}'_{-\xi_0}) \geq 0.$$

The last inequality is because  $(W^{(0)}, \xi_0)$  is Ding-semistable.

By convexity of  $\mathbf{D}(s)$ , we conclude that  $\mathbf{D}(s)$  is constant in  $s$  and by [Lemma 6.1](#) that  $G_{\mathcal{F}'_1}(y) \equiv \langle y, \xi_0 \rangle$  for any  $y \in \Delta' = \Delta(W^{(0)}, -K_{W^{(0)}})$  (the Okounkov body of  $(W^{(0)}, -K_{W^{(0)}})$ ).

By the discussion in previous section, we know that the relative successive minima of  $\mathcal{F}_1$  with respect to  $\mathcal{F}_0$  is the same as the relative successive minima of  $\mathcal{F}'_1$  with respect to  $\mathcal{F}'_{\text{wt}_{\xi_0}}$ , which is the same as the successive minima of  $\mathcal{F}'_{-\xi_0}$  and is given by the difference  $\lambda_k - \langle \alpha_k, \xi_0 \rangle$  with the notation there. So we get by [Lemma 5.1](#) that

$$\begin{aligned} d_2(\mathcal{F}_0, \mathcal{F}_1)^2 &= d_2(\mathcal{F}'_0, \mathcal{F}'_1) = \lim_{m \rightarrow +\infty} \sum_k \frac{(\lambda_k - \langle \alpha_k, \xi_0 \rangle)^2}{m^2} = \lim_{m \rightarrow +\infty} \sum_i \frac{\lambda_i^{(m)}(\mathcal{F}'_{-\xi_0})^2}{m^2} \\ &= \int_{\mathbb{R}} \lambda^2 \text{DH}(\mathcal{F}'_{-\xi_0})^2 = \int_{\Delta'} G_{\mathcal{F}'_{-\xi_0}}^2 dy = \int_{\Delta'} (G_{\mathcal{F}'_{-\xi_0}} - \langle y, \xi_0 \rangle)^2 dy = 0. \end{aligned}$$

By [\[20\]](#), we know that  $\mathcal{F}_0$  is asymptotically equivalent to  $\mathcal{F}_1$ . By [Lemmas 2.11](#) and [2.29](#) (see also [Proposition 2.27](#)), we get  $\mathcal{F}_0 = \mathcal{F}_1$ .

**Remark 6.2** Although here we are dealing with filtration of arbitrary ranks, the unique result in this section (and minimization result in previous section) can also be proved by using  $r := \text{rk}(\mathcal{F}_0)$ –step degenerations to reduce to the rank-one case. To see this, we first choose  $\{\eta_1, \dots, \eta_r\} \in N_{\mathbb{Q}} \cong \mathbb{Q}^r$  (where  $N = \text{Hom}(\mathbb{C}^*, \mathbb{T}_0)$  as before) such that:

- $\text{Span}_{\mathbb{R}}\{\eta_1, \dots, \eta_r\} = N_{\mathbb{R}}$ .
- For any  $1 \leq k \leq r$ ,  $\eta_k$  induces a special test configuration whose central fiber is the same as  $W^{(0)}$ . This is achieved by choosing  $\eta_k$  satisfying  $|\eta_k - \xi_0| \ll 1$ .

By abuse of notation, we denote by  $\mathcal{F}'_{\xi_0}$  (resp.  $\mathcal{F}'_{\eta_1}$ ) the filtration on  $R = R(X, -K_X)$  corresponding to the  $\mathbb{R}$ -test configuration induced by  $\xi_0$  (resp.  $\eta_1$ ), and also the filtration on  $R' = R(W^{(0)}, -K_{W_0^{(0)}})$  corresponding to the weight filtration induced by  $\xi_0$  (resp.  $\eta_k$  for  $2 \leq k \leq r$ ). Set  $\mathcal{F}'_1{}^{(0)} = \mathcal{F}_1$  and inductively define  $\mathcal{F}'_1{}^{(k)}$  to be the initial term degeneration of  $\mathcal{F}'_1{}^{(k-1)}$  with respect to  $\mathcal{F}'_{\eta_k}$  for  $1 \leq k \leq r$ . By (166) for the rank-one case, we have

$$(178) \quad H_X^{\text{NA}}(\mathcal{F}'_1{}^{(0)}) \geq H_{W^{(0)}}^{\text{NA}}(\mathcal{F}'_1{}^{(1)}) \quad \text{and} \quad H_{W^{(0)}}^{\text{NA}}(\mathcal{F}'_1{}^{(k-1)}) \geq H_{W^{(0)}}^{\text{NA}}(\mathcal{F}'_1{}^{(k)}) \quad \text{for } 2 \leq k \leq r.$$

So if  $\mathcal{F}_1 = \mathcal{F}'_1{}^{(0)}$  obtains the minimum of  $H_X^{\text{NA}}$ , then  $\mathcal{F}'_1{}^{(k)}$  for any  $1 \leq k \leq r$  also obtains the minimum of  $H_{W^{(0)}}^{\text{NA}}$ . Now because  $\mathcal{F}'^{(r)}$  is  $\mathbb{T}_0$ -invariant and  $\mathcal{F}'_{\xi_0} = \mathcal{F}'_{\text{wt}_{\xi_0}}$  also obtains the minimum of  $H_{W^{(0)}}^{\text{NA}}$ , we can use Lemma 2.38 to conclude that  $\mathcal{F}'^{(r)} = \mathcal{F}'_{\xi_0}$ .

On the other hand, by Lemma 5.1, we get for  $2 \leq k \leq r$  that

$$d_2^X(\mathcal{F}'_1{}^{(0)}, \mathcal{F}'_{\eta_1}) = d_2^{W^{(0)}}(\mathcal{F}'_1{}^{(1)}, \mathcal{F}'_{\eta_1}) \quad \text{and} \quad d_2^{W^{(0)}}(\mathcal{F}'_1{}^{(k-1)}, \mathcal{F}'_{\eta_k}) = d_2^{W^{(0)}}(\mathcal{F}'_1{}^{(k)}, \mathcal{F}'_{\eta_k}).$$

So for any  $1 \leq k \leq r$ , we get, by omitting the superscripts and using the triangle inequality,

$$\begin{aligned} d_2(\mathcal{F}'_1{}^{(k-1)}, \mathcal{F}'_{\xi_0}) &\leq d_2(\mathcal{F}'_1{}^{(k-1)}, \mathcal{F}'_{\eta_k}) + d_2(\mathcal{F}'_{\eta_k}, \mathcal{F}'_{\xi_0}) \\ &= d_2(\mathcal{F}'_1{}^{(k)}, \mathcal{F}'_{\eta_k}) + d_2(\mathcal{F}'_{\eta_k}, \mathcal{F}'_{\xi_0}) \\ &\leq d_2(\mathcal{F}'_1{}^{(k)}, \mathcal{F}'_{\xi_0}) + 2d_2(\mathcal{F}'_{\eta_k}, \mathcal{F}'_{\xi_0}). \end{aligned}$$

So we can inductively estimate

$$\begin{aligned} d_2(\mathcal{F}_1, \mathcal{F}_0) &= d_2(\mathcal{F}'_1{}^{(0)}, \mathcal{F}'_{\xi_0}) \leq d_2(\mathcal{F}'_1{}^{(1)}, \mathcal{F}'_{\xi_0}) + 2d_2(\mathcal{F}'_{\eta_1}, \mathcal{F}'_{\xi_0}) \\ &\leq d_2(\mathcal{F}'_1{}^{(2)}, \mathcal{F}'_{\xi_0}) + 2(d_2(\mathcal{F}'_{\eta_2}, \mathcal{F}'_{\xi_0}) + d_2(\mathcal{F}'_{\eta_1}, \mathcal{F}'_{\xi_0})) \\ &\vdots \\ &\leq d_2(\mathcal{F}'_1{}^{(r)}, \mathcal{F}'_{\xi_0}) + 2 \sum_{k=1}^r d_2(\mathcal{F}'_{\eta_k}, \mathcal{F}'_{\xi_0}) \\ &= 2 \sum_{k=1}^r d_2(\mathcal{F}'_{\eta_k}, \mathcal{F}'_{\xi_0}). \end{aligned}$$

Now we can choose  $\eta_k$  so that  $d_2(\mathcal{F}'_{\eta_k}, \mathcal{F}'_{\xi_0})$  is arbitrarily small for all  $1 \leq k \leq r$ . So we indeed get  $d_2(\mathcal{F}_1, \mathcal{F}_0) = 0$ , as desired.

## 7 Cone construction and $g$ -normalized volume

Let  $X$  be an  $n$ -dimensional  $\mathbb{Q}$ -Fano variety and for simplicity of notation, assume that  $-K_X$  is Cartier. Recall that  $R = \bigoplus_m R_m = \bigoplus H^0(X, m(-K_X))$ . We define the cone

$$(179) \quad C = C(X, -K_X) = \text{Spec}_{\mathbb{C}} R, \quad \mathfrak{o} = \mathfrak{m} = \bigoplus_{m>0} R_m.$$

Then  $(C, \mathfrak{o})$  is a klt cone singularity.

Since  $X$  admits a  $\mathbb{C}^* \times \mathbb{T}$ -action, we have a decomposition of the coordinate ring of  $R$ ,

$$(180) \quad R = \bigoplus_{m \geq 0} \bigoplus_{\alpha \in \mathbb{Z}^r} R_{m,\alpha}.$$

For any  $\mathbb{T}$ -invariant homogeneous primary ideal  $\mathfrak{a} = \bigoplus_m \bigoplus_{\alpha} \mathfrak{a}_{m,\alpha} \subset R$ , define the  $g$ -colength and  $g$ -multiplicity of  $\mathfrak{a}$  by

$$(181) \quad \text{colen}_g(\mathfrak{a}) = \sum_{m \geq 0} \sum_{\alpha} g\left(\frac{\alpha}{m}\right) \dim R_{m,\alpha} / \mathfrak{a}_{m,\alpha},$$

$$(182) \quad \text{mult}_g(\mathfrak{a}) = \lim_{k \rightarrow +\infty} \frac{\text{colen}_g(\mathfrak{a}^k)}{k^{n+1} / (n+1)!}.$$

See [63] for the study of such equivariant multiplicity. More generally, let  $\mathfrak{a}_\bullet = \{\mathfrak{a}_k\}_{k \in \mathbb{N}}$  be a graded sequence of  $\mathbb{C}^* \times \mathbb{T}$ -invariant ideals. We define

$$(183) \quad \text{mult}_g(\mathfrak{a}_\bullet) = \lim_{k \rightarrow +\infty} \frac{\text{colen}_g(\mathfrak{a}_k)}{k^{n+1} / (n+1)!}.$$

One can use the techniques of Newton–Okounkov bodies to show that the limit exists. To see this, we can adapt the argument in the work in [46] as follows. First choose a valuation  $\mathfrak{v}$  adapted to the  $\mathbb{T}$ -action on  $X$  (in the sense of Definition 2.21). We can construct a  $\mathbb{C}^* \times \mathbb{T}$ -invariant  $\mathbb{Z}^{n+1}$ -valuation on  $C$  by

$$(184) \quad \mathfrak{V}(f) = (m, \mathfrak{v}(f)) \quad \text{for any } f \in R_m.$$

Denote by  $\mathfrak{C}$  the strongly convex cone which is the closure of the convex hull of the value semigroup  $\mathfrak{V}(R)$ . To each graded sequence of  $\mathbb{C}^* \times \mathbb{T}$ -invariant ideals  $\mathfrak{a}_\bullet$ , one can associate a convex region  $\bar{P} := \bar{P}(\mathfrak{a}_\bullet) \subset \mathfrak{C}$  such that  $\bar{P}^c := \mathfrak{C} \setminus \bar{P}$  is bounded. If we still denote by  $g(y)$  the pullback of function  $g$  by the projection  $\mathbb{R}^{n+1} = \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ , then  $\text{mult}_g$  is given by the weighted volume of the co-convex set  $\bar{P}^c$ ,

$$(185) \quad \text{mult}_g(\mathfrak{a}_\bullet) = (n+1)! \int_{\bar{P}^c} g(y) dy.$$

Let  $\text{Val}_{C,o}$  be the space of real valuations that are centered at  $o$ , and by  $\text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}}$  the subset of  $\mathbb{C}^* \times \mathbb{T}$ -invariant real valuations in  $\text{Val}_{C,o}$ . If  $\tilde{C} \rightarrow C$  is the blowup of the vertex  $o \in X$ , then the exceptional divisor on  $\tilde{C}$  is isomorphic to  $X$ , and we will denote by  $\text{ord}_X$  the associated divisorial valuation contained in  $\text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}}$ .

Let  $\bar{\mathfrak{v}} \in \text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}}$  be any  $\mathbb{C}^* \times \mathbb{T}$ -invariant valuation. Then for any  $\lambda \in \mathbb{R}$ ,  $\mathfrak{a}_\lambda(\bar{\mathfrak{v}}) = \{f \in R \mid \bar{\mathfrak{v}}(f) > m\}$  is a  $\mathbb{T}$ -invariant homogeneous primary ideal. Set  $\mathfrak{a}_\bullet(\bar{\mathfrak{v}}) = \mathfrak{a}_m(\bar{\mathfrak{v}})$  and define (see [35] for the  $g = 1$  case)

$$\text{vol}_g(\bar{\mathfrak{v}}) := \text{mult}_g(\mathfrak{a}_\bullet(\bar{\mathfrak{v}})) = \lim_{m \rightarrow +\infty} \frac{\text{colen}_g(\mathfrak{a}_m(\bar{\mathfrak{v}}))}{\lambda^{n+1} / (n+1)!}.$$

We define the following equivariant version of normalized volume [51]:

$$\widehat{\text{vol}}_g : \text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}} \rightarrow \mathbb{R}_{>0} \cup \{+\infty\}, \quad \widehat{\text{vol}}_g(\bar{\mathfrak{v}}) = \begin{cases} A_C(\bar{\mathfrak{v}})^{n+1} \cdot \text{vol}_g(\bar{\mathfrak{v}}) & \text{when } A_C(\bar{\mathfrak{v}}) < +\infty, \\ +\infty & \text{otherwise.} \end{cases}$$

By using the same argument as in the study of normalized volumes, one can generalize almost all the results about normalized volume to work for the  $g$ -normalized volume functional. Here we just write down a few results that we need in the next section. We have the following equivariant version of an identity from [60].

**Lemma 7.1** *With the above notation, we have the identity*

$$(186) \quad \inf_{\bar{v}} \widehat{\text{vol}}_g(\bar{v}) = \inf_{\mathfrak{a}} \text{lct}(\mathfrak{a})^n \cdot \text{mult}_g(\mathfrak{a}) = \inf_{\mathfrak{a}_\bullet} \text{lct}(\mathfrak{a}_\bullet)^n \cdot \text{mult}_g(\mathfrak{a}_\bullet),$$

where  $\bar{v}$  ranges over  $\mathbb{C}^* \times \mathbb{T}$ -invariant valuations, and  $\mathfrak{a}$  (resp.  $\mathfrak{a}_\bullet$ ) ranges over  $\mathbb{C}^* \times \mathbb{T}$ -invariant ideals (resp. graded sequences of  $\mathbb{C}^* \times \mathbb{T}$ -invariant ideals).

This is proved by using exactly the same argument. For the reader's convenience, we give the short proof.

**Proof** For any  $\bar{v} \in \mathring{\text{Val}}_{C,o}$ , we have

$$(187) \quad \text{lct}(\mathfrak{a}_\bullet(\bar{v}))^n \cdot \text{mult}_g(\mathfrak{a}_\bullet(\bar{v})) \leq \left( \frac{A_C(\bar{v})}{\bar{v}(\mathfrak{a}_\bullet)} \right)^n \text{vol}_g(\bar{v}) = A_C(\bar{v})^n \text{vol}_g(\bar{v}).$$

Conversely, for any graded sequence of ideals  $\mathfrak{a}_\bullet$ , let  $\bar{w} \in \mathring{\text{Val}}_{C,o}$  be the valuation that calculates  $\text{lct}(\mathfrak{a}_\bullet)$ , which exists by [44]. By multiplying by a constant, we can assume  $1 = \bar{w}(\mathfrak{a}_\bullet) = \inf_m \bar{w}(\mathfrak{a}_m)/m$ . So  $\mathfrak{a}_m \subseteq \mathfrak{a}_m(\bar{w})$ , which implies  $\text{mult}_g(\mathfrak{a}_\bullet) \geq \text{mult}_g(\mathfrak{a}_\bullet(\bar{w})) = \text{vol}_g(\bar{w})$ . Then we get

$$(188) \quad \text{lct}(\mathfrak{a}_\bullet)^n \cdot \text{mult}_g(\mathfrak{a}_\bullet) = \left( \frac{A_C(\bar{w})}{\bar{w}(\mathfrak{a}_\bullet)} \right)^n \cdot \text{mult}_g(\mathfrak{a}_\bullet) \geq A_C(\bar{w})^n \cdot \text{vol}_g(\bar{w}) = \widehat{\text{vol}}_g(\bar{w}). \quad \square$$

For any  $v \in X_{\mathbb{Q}}^{\text{div}}$  and  $\tau > 0$ , we denote by  $\bar{v}_\tau$  the  $\mathbb{C}^*$ -invariant valuation on  $C$  given by

$$(189) \quad \bar{v}_\tau \left( \sum_i f_i t^i \right) = \min_i (v(f_i) + \tau i).$$

By using the same calculation as in [50], we get:

**Theorem 7.2** *The  $g$ -volume of  $\bar{v}_\tau$  is given by the formula*

$$(190) \quad \text{vol}_g(\bar{v}_\tau) = \frac{1}{\tau^{n+1}} V_g - (n+1) \int_0^{+\infty} \text{vol}_g(\mathcal{F}_v R^{(x)}) \frac{dx}{(x+\tau)^{n+2}}.$$

We have the following criterion for  $g$ -Ding-semistability, which generalizes the results in [50; 53; 58] about normalized volumes.

**Theorem 7.3** *The pair  $(X, \eta)$  is  $g$ -Ding-semistable if and only if  $\text{ord}_X$  obtains the minimum of  $\widehat{\text{vol}}_g$  over  $\text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}}$ .*

**Proof** For any  $v \in (X_{\mathbb{Q}}^{\text{div}})^{\mathbb{T}}$ , consider  $w_s := (\overline{sv})_{(1-s)A_X(v)} \in \text{Val}_{C,o}^{\mathbb{C}^* \times \mathbb{T}}$ . Then  $w_0 = A_X(v)\bar{v}_0$  and  $w_1 = v$ . We also have  $A_C(w_s) \equiv A_X(v)$ . Set

$$\begin{aligned} f(s) &= \widehat{\text{vol}}(w_s) = A_C(w_s)^{n+1} \text{vol}_g(w_s) \\ &= A_X(v)^{n+1} \left( \frac{V_g}{(1-s)^{n+1} A_X(v)^{n+1}} - (n+1) \int_0^{+\infty} \text{vol}_g(\mathcal{F}_v R^{(x)}) \frac{s \, dx}{(sx + (1-s)A_X(v))^{n+2}} \right) \\ &= A_X(v)^{n+1} \int_0^{+\infty} \frac{-d \text{vol}_g(\mathcal{F}_v R^{(x)})}{(sx + (1-s)A_X(v))^{n+1}}. \end{aligned}$$

Then  $f(s)$  is a convex function in  $s \in [0, 1]$ . Its derivative at  $s = 0$  is given by

$$\begin{aligned} (191) \quad f'(0) &= A_X(v)^{n+1} \left( (n+1) \frac{V_g}{A_X(v)^{n+1}} - (n+1) \int_0^{+\infty} \text{vol}_g(\mathcal{F}_v R^{(x)}) \, dx \frac{1}{A_X(v)^{n+2}} \right) \\ &= \frac{n+1}{A_X(v)V_g} \left( A_X(v) - \frac{1}{V_g} \int_0^{+\infty} \text{vol}_g(\mathcal{F}_v R^{(x)}) \, dx \right) \\ &= \frac{n+1}{A_X(v)V_g} \cdot \beta_g(v). \end{aligned}$$

With this and [Theorem 2.47](#), we can easily derive the conclusion as in [\[50\]](#). □

**Remark 7.4** By the same argument as in the case of normalized volume [\[12; 78\]](#), one shows that  $g$ –Ding-semistability is Zariski-open for a  $\mathbb{T}$ –equivariant family of Fano varieties.

## 8 Uniqueness of polystable degeneration

In this section, we prove [Theorem 1.3](#). The proof is verbatim the same as the proof of the existence and uniqueness of  $K$ –polystable degenerations for any  $K$ –semistable  $\mathbb{Q}$ –Fano varieties, as proved in [\[55\]](#); see also [\[16\]](#). Indeed, we just need to carry out the same argument by using the equivariant version of normalized volume and the modified Futaki invariant  $\text{Fut}_{\xi}$ , etc. To avoid redundancy, we only sketch the key steps and refer to [\[55; 16\]](#) for more details.

Assume that  $(X, \xi)$  is semistable and admits two polystable degenerations via two special test configurations  $(\mathcal{X}^{(i)}, -K_{\mathcal{X}^{(i)}})$  for  $i = 0, 1$ . Take cones fiberwise to get a special test configuration of Fano cones  $(\mathcal{C}^{(i)}, \zeta^{(i)})$ , where  $\zeta^{(i)}$  is the radial vector field.

Let  $E_k$  be the Kollár component (see [\[58\]](#) for the definition) obtained by blowing up the vertex of  $\mathcal{C}^{(0)}$  with weight  $(k, 1)$ . Then we have

$$\widehat{\text{vol}}_g(E_k) = \widehat{\text{vol}}_g(\text{ord}_X) + O(k^{-2}).$$

Set  $\mathbf{a}_{\bullet} = \{\mathbf{a}_{\ell}(E_k)\}$ . Then

$$\text{lct}(\mathbf{a}_{\bullet}) = \frac{A(E_k)}{\text{ord}_{E_k}(\mathbf{a}_{\bullet})} = A(E_k) =: c_k = O(k), \quad \text{lct}(X, \mathbf{a}_{\bullet})^n \cdot \text{mult}_g(\mathbf{a}_{\bullet}) = \widehat{\text{vol}}_g(E_k).$$

Consider the initial degeneration of  $\mathfrak{a}_\bullet$  with respect to  $\mathcal{C}^{(1)}$ ,

$$(192) \quad \mathbf{in}(\mathfrak{a}_\ell) = \text{span}_{\mathbb{C}}\{\mathbf{in}(f), f \in \mathfrak{a}_\ell(E_k)\}.$$

Using the preservation of co-length under initial term degeneration, we get

$$\begin{aligned} \text{lct}^n(\mathcal{C}_0^{(1)}, \mathbf{in}(\mathfrak{a}_\bullet)) &\geq \frac{\widehat{\text{vol}}_g(\text{ord}_{\mathcal{X}_0^{(1)}})}{\text{mult}_g(\mathbf{in}(\mathfrak{a}_\bullet))} = \frac{V_g}{\text{mult}_g(\mathfrak{a}_\bullet)} \\ &= \frac{V_g}{\widehat{\text{vol}}_g(E_k)} \text{lct}(\mathfrak{a}_\bullet)^n = \frac{V_g}{V_g + O(k^{-2})} \text{lct}(\mathfrak{a}_\bullet)^n \\ &= (1 + O(k^{-2}))c_k = c_k + O(k^{-1}). \end{aligned}$$

Let  $Z_k \rightarrow \mathcal{C}^{(0)}$  be the extraction of  $E_k$ , and let  $Z_k \times \mathbb{C}^*$  be the product along  $\mathcal{C}^{(1)} \setminus \mathcal{C}_0^{(1)} \cong C \times \mathbb{C}^*$  with exceptional divisor  $\mathcal{E}_k$ . Let  $\mathfrak{B}_\bullet = \{\mathfrak{B}_\ell\}$  be ideal on the total space  $\mathcal{C}^{(1)}$  obtained by the above degenerating  $\mathfrak{a}_\ell$ . Then we have

$$(193) \quad A(\mathcal{C}^{(1)}, c_k(1 - \epsilon k^{-1})\mathfrak{B}_\bullet, \mathcal{E}_k) = A(C, c_k(1 - \epsilon k^{-1})\mathfrak{a}_\bullet, E_k) = \epsilon k^{-1} c_k = \epsilon O(1),$$

$$(194) \quad \text{lct}(\mathcal{C}_0^{(1)}, c_k(1 - \epsilon k^{-1})\mathbf{in}(\mathfrak{a}_\bullet)) \geq c_k^{-1}(1 - \epsilon k^{-1})(c_k + O(k^{-1})) = 1 - \epsilon k^{-1} + O(k^{-2}).$$

By inversion of adjunction,

$$(195) \quad \text{lct}(\mathcal{C}^{(1)}, c_k(1 - \epsilon k^{-1})\mathfrak{B}_\bullet) \geq 1 - \epsilon k^{-1} + O(k^{-2}).$$

When  $0 < \epsilon \ll 1$ , by [9], we can extract the divisor  $\mathcal{E}_k$  over  $\mathcal{C}^{(1)}$ . By the same argument as [55], we get the commutative diagram

$$(196) \quad \begin{array}{ccccc} & & \mathcal{C}^{(1)} \xleftarrow{\mathcal{Z}_k^{(1)}} \xleftarrow{\mathcal{E}_k^{(1)}} & C & \xleftarrow{Z_k} \xleftarrow{E_k} \\ & \nearrow & \text{---} & \text{---} & \text{---} \\ & \mathcal{X}_0^{(1)} & \xleftarrow{\mathcal{X}'^{(1)}} & X & \\ & \downarrow \mathcal{X}'^{(1)} & & \downarrow \mathcal{X}^{(0)} & \\ & \mathcal{X}'_0 & \xleftarrow{\mathcal{X}'^{(0)}} & X_0^{(0)} & \\ & \searrow & \text{---} & \text{---} & \text{---} \\ & & \mathcal{C}'_0 & \xleftarrow{Z_{k,0}} \xleftarrow{E_k} & C_0^{(0)} \end{array}$$

$\mathcal{C}'^{(1)} \leftarrow \mathcal{C}'_0 \leftarrow \mathcal{C}'_0^{(0)}$  (left side),  $\mathcal{C}^{(0)} \leftarrow \mathcal{Z}_k \leftarrow \mathcal{E}_k = E_k \times \mathbb{A}^1$  (right side),  $\mathcal{C}^{(0)} \leftarrow \mathcal{Z}_{k,0} \leftarrow E_k$  (bottom right)

By the same argument as in [55], we know that both test configurations  $\mathcal{X}'^{(i)}$  for  $i = 0, 1$  are weakly special and have vanishing  $\text{Fut}_\xi$  invariant. By [39], we know that both of them are special and hence  $X_0^{(1)} \cong X'_0 \cong X_0^{(0)}$  by the polystability of  $X_0^{(i)}$ .

The existence part can again be proved by the similar arguments as in [55], which deals with the case when  $\xi = 0$ . We just sketch the arguments. If  $(X, \xi)$  is K-polystable, then we are done. Otherwise, we can find a nontrivial  $\mathbb{T}$ -equivariant special test configuration such that the central fiber (with the vector

field  $\xi$ ) has a vanishing  $\text{Fut}_\xi$  invariant. By [55, Proof of Lemma 3.1], we know that the central fiber is  $\mathbb{K}$ –semistable, and has an effective action by a larger torus. If the central fiber is  $\mathbb{K}$ –polystable, then we are done again. Otherwise, we can continue this process, which must stop since the dimension of the torus is bounded by the dimension of  $X$ .

**Proof of Corollary 1.4** By the work of Chen, Sun and Wang in [28], which is based on the resolution of the Hamilton–Tian conjecture [29], we get a special  $\mathbb{R}$ –test configuration  $\mathcal{F}^{\text{ss}}$  with central fiber  $(W, \xi)$ , and a special test configuration of  $(W, \xi)$  with central fiber  $(X_\infty, \xi)$ , which admits a Kähler–Ricci soliton and hence is  $\mathbb{K}$ –polystable. By the work of Dervan and Székelyhidi [31],  $\mathcal{F}^{\text{ss}}$  obtains the minimum  $h(X)$ . The statement follows directly from Theorems 1.2 and 1.3.  $\square$

**Remark 8.1** The fact that  $\mathcal{F}^{\text{ss}}$  obtains the minimum also follows from the  $\mathbb{K}$ –semistability of  $(W, \xi)$  and Theorem 5.2. The  $\mathbb{K}$ –semistability of  $(W, \xi)$  follows from the same degeneration argument as used in [58], or the Zariski–openness of  $\mathbb{K}$ –semistability as pointed out in Remark 7.4.

**Remark 8.2** As in the more general setting of [55] or [52], the algebraic results in this paper can be generalized to the log Fano case in a straightforward way.

## Appendix Properties of $\tilde{\mathcal{S}}(v)$

Recall that by (137), for any valuation  $v \in \mathring{\text{Val}}(X)$  we have

$$(197) \quad \mathcal{Q}(v) := \mathcal{Q}(\mathcal{F}_v) = e^{-\tilde{\mathcal{S}}^{\text{NA}}(\mathcal{F}_v)} = 1 - \frac{1}{V} \int_0^{T(v)} e^{-x} \text{vol}(\mathcal{F}_v^{(x)} R_\bullet) dx =: 1 - \Psi(v),$$

where, for simplicity of notation, we have written

$$(198) \quad T(v) = \lambda_{\max}(\mathcal{F}_v) \quad \text{and} \quad \Psi(v) = \frac{1}{V} \int_0^{T(v)} e^{-x} \text{vol}(\mathcal{F}_v^{(x)} R_\bullet) dx.$$

**Proposition A.1** *The function  $v \mapsto \Psi(v)$  is strictly increasing on  $\mathring{\text{Val}}(X)$ . In other words, if  $v \leq w$ , then  $\Psi(v) \leq \Psi(w)$ , with the identity true only if  $v = w$ . As a consequence,  $v \mapsto \tilde{\mathcal{S}}(v)$  is strictly increasing on  $\text{Val}(X)$ .*

This is proved as in [11, Proof of Proposition 3.15] (which is based on an argument in the local case from [58]). We sketch the argument for the reader’s convenience.

**Proof** First, by using Theorem 2.5, we can show that

$$(199) \quad \Psi(v) = \lim_{m \rightarrow +\infty} \frac{1}{mN_m} \sum_{j \geq 1} e^{-j/m} \dim \mathcal{F}_v^j R_m.$$

Suppose that  $v \leq w$  but  $v \neq w$ . Then by rescaling  $v, w$  and  $L = -K_X$ , we can assume that there exists  $s \in H^0(X, L)$  with  $w(s) = p \in \mathbb{N}^*$  and  $v(s) \leq p - 1$ . Then, arguing as in [11, Proof of Proposition 3.15],



we have

$$(200) \quad \dim(\mathcal{F}_w^j R_m / \mathcal{F}_v^j R_m) \geq \sum_{1 \leq i \leq \min\{j/p, m\}} \dim(\mathcal{F}_v^{j-ip} R_m / \mathcal{F}_v^{j-ip+1} R_{m-i}).$$

One the other hand, with  $C = \max\{T(v), T(w)\}$ , we get

$$\begin{aligned} \sum_{j \geq 1} \dim e^{-j/m} (\mathcal{F}_w^j R_m - \mathcal{F}_v^j R_m) &\geq e^{-C} \sum_{j \geq 1} (\mathcal{F}_w^j R_m - \mathcal{F}_v^j R_m) \\ &\geq e^{-C} \sum_{1 \leq i \leq m} \sum_{j \geq pi} (\dim \mathcal{F}_v^{j-ip} R_{m-i} - \mathcal{F}_v^{j-ip+1} R_{m-i}) \\ &= e^{-C} \sum_{1 \leq i \leq m} \dim R_{m-i}. \end{aligned}$$

So we conclude

$$\Psi(v) - \Psi(w) \geq e^{-C} \lim_{m \rightarrow +\infty} \frac{1}{mN_m} \sum_{1 \leq i \leq m} \dim R_{m-i} > 0. \quad \square$$

Let  $\pi: Y \rightarrow X$  be a proper birational morphism with  $Y$  a regular and  $E = \sum_i E_i$  a reduced simple normal crossing divisor.

**Proposition A.2** *The function  $v \mapsto \mathcal{Q}(v)$  is continuous on  $\text{QM}(Y, E)$ .*

We use the same strategy as [14, Proposition 2.4]. As noted in [38], for any  $v \in \text{Val}(X)$ , we have  $A(v)/T(v) \geq \alpha(X) > 0$ , which implies, with  $C = \alpha(X)^{-1}$ ,

$$(201) \quad T(v) \leq CA(v).$$

**Lemma A.3** *For any  $v \in \text{Val}(X)$ , we have the inequality*

$$(202) \quad \Psi(v) \leq CA(v).$$

**Proof** Since  $\text{vol}(\mathcal{F}^{(x)} R_\bullet) \leq V$ , we immediately get

$$\Psi(v) \leq \int_0^{T(v)} e^{-x} dx = 1 - e^{-T(v)} \leq T(v) \leq CA(v),$$

where we used the inequality  $1 - e^{-x} \leq x$  for any  $x \in \mathbb{R}_{\geq 0}$ , and the inequality (201). □

Similarly to [38; 11], we introduce the approximation

$$(203) \quad \mathcal{Q}_m(\mathcal{F}) = \frac{1}{N_m} \sum_i e^{-\lambda_i^{(m)}/m} = \frac{1}{N_m} \int_0^{+\infty} e^{-x/m} d(-\dim \mathcal{F}^x R_m)$$

$$(204) \quad = 1 - \frac{1}{N_m} \int_0^{\lambda_{\max}^{(m)}(\mathcal{F})/m} e^{-x} \dim \mathcal{F}^{xm} R_m dx =: 1 - \Psi_m(\mathcal{F}),$$

where we set

$$(205) \quad \Psi_m(v) = \frac{1}{N_m} \int_0^{\lambda_{\max}^{(m)}} e^{-x} \dim \mathcal{F}^{xm} R_m dx = \frac{1}{N_m} \int_0^{T(v)} e^{-x} \dim \mathcal{F}^{xm} R_m dx.$$

Similarly to [38; 11], for any valuation  $v \in \text{Val}(X)$  we have the identity

$$(206) \quad \mathbf{Q}_m(v) = \mathbf{Q}_m(\mathcal{F}_v) = \min_{\{s_j\}} \frac{1}{N_m} \sum_{j=1}^{N_m} e^{-v(s_j)/m},$$

where the minimum is taken over all bases  $s_1, \dots, s_{N_m}$  of  $H^0(X, -mK_X)$ .

For any  $\mathfrak{s} := \{s_1, \dots, s_{N_m}\} \in H^0(X, -mK_X)^{N_m}$ , define a function

$$(207) \quad \varphi_{\mathfrak{s}}(v) := \sum_{j=1}^{N_m} e^{-v(s_j)/m}.$$

By the same argument as in [14, Proof of Lemma 2.5], the set of functions  $\{\varphi_{\mathfrak{s}}(v) \mid \mathfrak{s} \in R_m^{N_m}\}$  is finite. So  $\mathbf{Q}_m$  is continuous on  $\text{QM}(Y, E)$ .

As in [14, Proof of Proposition 2.4], the continuity of  $\Psi$  and hence  $\mathbf{Q}$  follows easily from the following proposition, which we prove by using the techniques developed in [11; 13].

**Lemma A.4** (i) For any  $v \in \text{Val}(X)$  with  $A(v) < +\infty$ , we have the convergence

$$(208) \quad \lim_{m \rightarrow +\infty} \Psi_m(v) = \Psi(v).$$

(ii) For any  $\epsilon > 0$  and any  $C_1 > 0$ , there exists  $C_2 > 0$  and  $m_0 > 0$  such that if  $v \in \text{Val}(X)$  satisfies  $A(v) < C_1$ , we have

$$(209) \quad |\Psi_m(\mathcal{F}_v) - \Psi(\mathcal{F}_v)| \leq \epsilon$$

for all  $m$  divisible by  $m_0$ .

**Proof** The first statement follows from Theorem 2.5(ii). We focus on the second statement.

Note that  $e^{-G}$  is convex and  $0 \leq e^{-G} \leq 1$ . By [11, Lemma 2.2], for any  $\epsilon' > 0$  there exists  $m_0(\epsilon')$  such that for any  $m \geq m_0$ ,

$$(210) \quad \int_{\Delta} e^{-G} d\rho_m \geq \int_{\Delta} e^{-G} dy - \epsilon'.$$

By the same argument as [11, Proof of Lemma 2.9], we get

$$(211) \quad \mathbf{Q}_m(\mathcal{F}_v) \geq \frac{m^n}{N_m} \int_{\Delta} e^{-G} d\rho_m.$$

Note that  $\lim_{m \rightarrow +\infty} m^n/N_m = V$ . So for any  $\epsilon > 0$  there exists  $m_0$  such that for any  $m \geq m_0$ ,

$$(212) \quad \mathbf{Q}_m(\mathcal{F}_v) \geq \frac{n!}{V} \int_{\Delta} e^{-G} dy - \epsilon = \mathbf{Q}(\mathcal{F}_v) - \epsilon.$$

We need to prove the other direction of inequality. Following [11], define a graded linear series

$$(213) \quad \tilde{\mathcal{F}}_{m,p}^{(t)} R_{mp} := H^0(X, mpL \otimes \overline{\mathfrak{b}(|\mathcal{F}^{mt} R_m|^p)}),$$

where  $\mathfrak{b}(|\mathcal{F}^{mt} R_m|)$  is the base ideal of the sublinear system  $\mathcal{F}^{tm} R_m$ . Set

$$(214) \quad \tilde{\Psi}_m(\mathcal{F}) = \int_0^{T(v)} e^{-t} \text{vol}(\tilde{\mathcal{F}}_{m,\bullet}^{(t)}) dt.$$

By [11, Proposition 5.13], there exists  $a = a(X, -K_X) > 0$  such that for all  $t \in \mathbb{Q}_{>0}$  with  $mt > A(v)$ , we have, with  $t' = t - m^{-1}(at + A(v))$ ,

$$(215) \quad \left(\frac{m-a}{m}\right)^{n+1} \text{vol}(\mathcal{F}_{\bullet,\bullet}^{(t)}) \leq \frac{1}{m^n} \text{vol}(\tilde{\mathcal{F}}_{m,\bullet}^{(t')}).$$

So we can estimate as in [11, Proof of Proposition 5.15] to get

$$\begin{aligned} \tilde{\Psi}_m(v) &\geq \left(\frac{m-a}{m}\right)^{n+1} \left( \Psi(v) - e^{(aT(v)+A(v))/m} \int_0^{A(v)/(m-a)} \frac{\text{vol}(\mathcal{F}^{(t)})}{V} e^{-t} dt \right) \\ &\geq \left(\frac{m-a}{m}\right)^{n+1} \left( \Psi(v) - e^{CA(v)/m} \frac{A(v)}{m-a} \right). \end{aligned}$$

From this it is easy to get

$$\Psi(v) - \tilde{\Psi}_m(v) \leq C \frac{A(v)}{m}.$$

To compare with  $\Psi_m$ , we further set

$$(216) \quad \mathcal{F}_{m,p}^{(x)} = \text{Im}(S^p \mathcal{F}^{mx} R_m \rightarrow H^0(X, pmL)).$$

By [13, Propositions 5.14 and 3.2], there exists a positive constant  $C > 0$  independent of  $v$  such that for all  $x \leq T(v) - CA(v)/m$ , we have  $\text{vol}(\mathcal{F}_{m,\bullet}^{(x)}) = \text{vol}(\tilde{\mathcal{F}}_{m,\bullet}^{(x)})$ . So as in [13, Proof of Proposition 5.15], we get

$$\begin{aligned} \Psi(v) &\leq \tilde{\Psi}_m(v) + C \frac{A(v)}{m} = \frac{1}{V} \int_0^{T(v)} \frac{\text{vol}(\tilde{\mathcal{F}}_{m,\bullet}^{(x)})}{m^n} e^{-x} dx + \frac{CA(v)}{m} \\ &\leq \frac{1}{V} \int_0^{T(v)-CA(v)/m} \frac{\text{vol}(\mathcal{F}_{m,\bullet}^{(x)})}{m^n} e^{-x} dx + \frac{CA(v)}{m} \\ &\leq \frac{1}{V} \int_0^{T(v)} \frac{\text{vol}(\mathcal{F}_{m,\bullet}^{(x)})}{m^n} e^{-x} dx + \frac{CA(v)}{m}. \end{aligned}$$

For the second inequality we used the estimate that, as  $m \rightarrow +\infty$ ,

$$\int_{T(v)-CA(v)/m}^{T(v)} e^{-x} dx = e^{-(T(v)-CA(v)/m)} - e^{-T(v)} \leq e^{CA(v)/m} - 1 = O\left(\frac{A(v)}{m}\right).$$

Fixing any  $\epsilon > 0$ , by choosing  $m \gg 1$  and  $p \gg 1$  we have (see [13, equation (5.6)]):

$$(217) \quad \left| \frac{\text{vol}(\mathcal{F}_{m,\bullet}^{(x)})}{m^n V} - \frac{\dim \mathcal{F}_{m,p}^{(x)}}{N_{mp}} \right| < \epsilon.$$

Finally we can estimate as in [13, Proof of Theorem 5.13]: for  $m \gg 1$ ,

$$\begin{aligned} \Psi(v) &\leq \frac{1}{V} \int_0^{T(v)} \frac{\text{vol}(\mathcal{F}_{m,\bullet}^{(x)})}{m^n} e^{-x} dx + \frac{CA(v)}{m} \leq \int_0^{+\infty} \frac{\dim \mathcal{F}_{m,p}^{(x)}}{N_{mp}} e^{-x} dx + \epsilon T(v) + \frac{CA(v)}{m} \\ &\leq \int_0^{+\infty} \frac{\dim \mathcal{F}^{pmx} R_m}{N_{mp}} e^{-x} dx + 2\epsilon A(v) = \Psi(v) + 2\epsilon A(v). \end{aligned}$$

In the third inequality, we used again the inequality  $1 - e^{-T(v)} \leq T(v)$ . Since  $\epsilon > 0$  is arbitrary, we get the conclusion.  $\square$

## References

- [1] **H Abban, Z Zhuang**, *K-stability of Fano varieties via admissible flags*, Forum Math. Pi 10 (2022) art. id. e15 [MR](#) [Zbl](#)
- [2] **K Altmann, J Hausen**, *Polyhedral divisors and algebraic torus actions*, Math. Ann. 334 (2006) 557–607 [MR](#) [Zbl](#)
- [3] **K Altmann, J Hausen, H Süß**, *Gluing affine torus actions via divisorial fans*, Transform. Groups 13 (2008) 215–242 [MR](#) [Zbl](#)
- [4] **R Bamler**, *Convergence of Ricci flows with bounded scalar curvature*, Ann. of Math. 188 (2018) 753–831 [MR](#) [Zbl](#)
- [5] **R J Berman**, *K-polystability of  $\mathbb{Q}$ -Fano varieties admitting Kähler–Einstein metrics*, Invent. Math. 203 (2016) 973–1025 [MR](#) [Zbl](#)
- [6] **R J Berman, S Boucksom, P Eyssidieux, V Guedj, A Zeriahi**, *Kähler–Einstein metrics and the Kähler–Ricci flow on log Fano varieties*, J. Reine Angew. Math. 751 (2019) 27–89 [MR](#) [Zbl](#)
- [7] **R J Berman, S Boucksom, M Jonsson**, *A variational approach to the Yau–Tian–Donaldson conjecture*, J. Amer. Math. Soc. 34 (2021) 605–652 [MR](#) [Zbl](#)
- [8] **R J Berman, D W Nystrom**, *Complex optimal transport and the pluripotential theory of Kähler–Ricci solitons*, preprint (2014) [arXiv 1401.8264](#)
- [9] **C Birkar, P Cascini, C D Hacon, J McKernan**, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. 23 (2010) 405–468 [MR](#) [Zbl](#)
- [10] **H Blum**, *Existence of valuations with smallest normalized volume*, Compos. Math. 154 (2018) 820–849 [MR](#) [Zbl](#)
- [11] **H Blum, M Jonsson**, *Thresholds, valuations, and K-stability*, Adv. Math. 365 (2020) art. id. 107062 [MR](#) [Zbl](#)
- [12] **H Blum, Y Liu**, *The normalized volume of a singularity is lower semicontinuous*, J. Eur. Math. Soc. 23 (2021) 1225–1256 [MR](#) [Zbl](#)
- [13] **H Blum, Y Liu**, *Openness of uniform K-stability in families of  $\mathbb{Q}$ -Fano varieties*, Ann. Sci. Éc. Norm. Supér. 55 (2022) 1–41 [MR](#) [Zbl](#)
- [14] **H Blum, Y Liu, C Xu**, *Openness of K-semistability for Fano varieties*, Duke Math. J. 171 (2022) 2753–2797 [MR](#) [Zbl](#)
- [15] **H Blum, Y Liu, C Xu, Z Zhuang**, *The existence of the Kähler–Ricci soliton degeneration*, Forum Math. Pi 11 (2023) art. id. e9 [MR](#) [Zbl](#)
- [16] **H Blum, Y Liu, C Zhou**, *Optimal destabilization of K-unstable Fano varieties via stability thresholds*, Geom. Topol. 26 (2022) 2507–2564 [MR](#) [Zbl](#)
- [17] **S Boucksom, H Chen**, *Okounkov bodies of filtered linear series*, Compos. Math. 147 (2011) 1205–1229 [MR](#) [Zbl](#)
- [18] **S Boucksom, D Eriksson**, *Spaces of norms, determinant of cohomology and Fekete points in non-Archimedean geometry*, Adv. Math. 378 (2021) art. id. 107501 [MR](#) [Zbl](#)

- [19] **S Boucksom, T Hisamoto, M Jonsson**, *Uniform  $K$ -stability, Duistermaat–Heckman measures and singularities of pairs*, Ann. Inst. Fourier (Grenoble) 67 (2017) 743–841 [MR](#) [Zbl](#)
- [20] **S Boucksom, M Jonsson**, *A non-Archimedean approach to  $K$ -stability*, preprint (2018) [arXiv 1805.11160v1](#)
- [21] **S Boucksom, M Jonsson**, *A non-Archimedean approach to  $K$ -stability, I: Metric geometry of spaces of test configurations and valuations*, preprint (2021) [arXiv 2107.11221](#)
- [22] **S Boucksom, A Küronya, C Maclean, T Szemberg**, *Vanishing sequences and Okounkov bodies*, Math. Ann. 361 (2015) 811–834 [MR](#) [Zbl](#)
- [23] **M Brion**, *Sur l’image de l’application moment*, from “Séminaire d’algèbre Paul Dubreil et Marie-Paule Malliavin” (M-P Malliavin, editor), Lecture Notes in Math. 1296, Springer (1987) 177–192 [MR](#) [Zbl](#)
- [24] **H Chen, C Maclean**, *Distribution of logarithmic spectra of the equilibrium energy*, Manuscripta Math. 146 (2015) 365–394 [MR](#) [Zbl](#)
- [25] **X Chen, S Donaldson, S Sun**, *Kähler–Einstein metrics on Fano manifolds, I: Approximation of metrics with cone singularities*, J. Amer. Math. Soc. 28 (2015) 183–197 [MR](#) [Zbl](#)
- [26] **X Chen, S Donaldson, S Sun**, *Kähler–Einstein metrics on Fano manifolds, II: Limits with cone angle less than  $2\pi$* , J. Amer. Math. Soc. 28 (2015) 199–234 [MR](#) [Zbl](#)
- [27] **X Chen, S Donaldson, S Sun**, *Kähler–Einstein metrics on Fano manifolds, III: Limits as cone angle approaches  $2\pi$  and completion of the main proof*, J. Amer. Math. Soc. 28 (2015) 235–278 [MR](#) [Zbl](#)
- [28] **X Chen, S Sun, B Wang**, *Kähler–Ricci flow, Kähler–Einstein metric, and  $K$ -stability*, Geom. Topol. 22 (2018) 3145–3173 [MR](#) [Zbl](#)
- [29] **X Chen, B Wang**, *Space of Ricci flows, II: Part B: weak compactness of the flows*, J. Differential Geom. 116 (2020) 1–123 [MR](#) [Zbl](#)
- [30] **T Darvas, W He**, *Geodesic rays and Kähler–Ricci trajectories on Fano manifolds*, Trans. Amer. Math. Soc. 369 (2017) 5069–5085 [MR](#) [Zbl](#)
- [31] **R Dervan, G Székelyhidi**, *The Kähler–Ricci flow and optimal degenerations*, J. Differential Geom. 116 (2020) 187–203 [MR](#) [Zbl](#)
- [32] **S K Donaldson**, *Scalar curvature and stability of toric varieties*, J. Differential Geom. 62 (2002) 289–349 [MR](#) [Zbl](#)
- [33] **S K Donaldson**, *Lower bounds on the Calabi functional*, J. Differential Geom. 70 (2005) 453–472 [MR](#) [Zbl](#)
- [34] **S Donaldson, S Sun**, *Gromov–Hausdorff limits of Kähler manifolds and algebraic geometry, II*, J. Differential Geom. 107 (2017) 327–371 [MR](#) [Zbl](#)
- [35] **L Ein, R Lazarsfeld, K E Smith**, *Uniform approximation of Abhyankar valuation ideals in smooth function fields*, Amer. J. Math. 125 (2003) 409–440 [MR](#) [Zbl](#)
- [36] **K Fujita**, *Optimal bounds for the volumes of Kähler–Einstein Fano manifolds*, Amer. J. Math. 140 (2018) 391–414 [MR](#) [Zbl](#)
- [37] **K Fujita**, *A valuative criterion for uniform  $K$ -stability of  $\mathbb{Q}$ -Fano varieties*, J. Reine Angew. Math. 751 (2019) 309–338 [MR](#) [Zbl](#)
- [38] **K Fujita, Y Odaka**, *On the  $K$ -stability of Fano varieties and anticanonical divisors*, Tohoku Math. J. 70 (2018) 511–521 [MR](#) [Zbl](#)
- [39] **J Han, C Li**, *On the Yau–Tian–Donaldson conjecture for generalized Kähler–Ricci soliton equations*, Comm. Pure Appl. Math. 76 (2023) 1793–1867 [MR](#) [Zbl](#)
- [40] **W He**, *Kähler–Ricci soliton and  $H$ -functional*, Asian J. Math. 20 (2016) 645–663 [MR](#) [Zbl](#)

- [41] **T Hisamoto**, *Stability and coercivity for toric polarizations*, preprint (2016) [arXiv 1610.07998](#)
- [42] **T Hisamoto**, *Mabuchi’s soliton metric and relative  $D$ -stability*, preprint (2019) [arXiv 1905.05948](#)
- [43] **T Hisamoto**, *Geometric flow, multiplier ideal sheaves and optimal destabilizer for a Fano manifold*, *J. Geom. Anal.* 33 (2023) art. id. 265 [MR](#) [Zbl](#)
- [44] **M Jonsson**, **M Mustața**, *Valuations and asymptotic invariants for sequences of ideals*, *Ann. Inst. Fourier (Grenoble)* 62 (2012) 2145–2209 [MR](#) [Zbl](#)
- [45] **K Kaveh**, **A G Khovanskii**, *Newton–Okounkov bodies, semigroups of integral points, graded algebras and intersection theory*, *Ann. of Math.* 176 (2012) 925–978 [MR](#) [Zbl](#)
- [46] **K Kaveh**, **A Khovanskii**, *Convex bodies and multiplicities of ideals*, *Proc. Steklov Inst. Math.* 286 (2014) 268–284 [MR](#) [Zbl](#)
- [47] **S J Kovács**, **Z Patakfalvi**, *Projectivity of the moduli space of stable log-varieties and subadditivity of log-Kodaira dimension*, *J. Amer. Math. Soc.* 30 (2017) 959–1021 [MR](#) [Zbl](#)
- [48] **R Lazarsfeld**, *Positivity in algebraic geometry, II: Positivity for vector bundles, and multiplier ideals*, *Ergebnisse der Math. (3)* 49, Springer (2004) [MR](#) [Zbl](#)
- [49] **R Lazarsfeld**, **M Mustața**, *Convex bodies associated to linear series*, *Ann. Sci. Éc. Norm. Supér.* 42 (2009) 783–835 [MR](#) [Zbl](#)
- [50] **C Li**,  *$K$ -semistability is equivariant volume minimization*, *Duke Math. J.* 166 (2017) 3147–3218 [MR](#) [Zbl](#)
- [51] **C Li**, *Minimizing normalized volumes of valuations*, *Math. Z.* 289 (2018) 491–513 [MR](#) [Zbl](#)
- [52] **C Li**,  *$G$ -uniform stability and Kähler–Einstein metrics on Fano varieties*, *Invent. Math.* 227 (2022) 661–744 [MR](#) [Zbl](#)
- [53] **C Li**, **Y Liu**, *Kähler–Einstein metrics and volume minimization*, *Adv. Math.* 341 (2019) 440–492 [MR](#) [Zbl](#)
- [54] **C Li**, **Y Liu**, **C Xu**, *A guided tour to normalized volume*, from “Geometric analysis: in honor of Gang Tian’s 60th birthday” (J Chen, P Lu, Z Lu, Z Zhang, editors), *Progr. Math.* 333, Springer (2020) 167–219 [MR](#) [Zbl](#)
- [55] **C Li**, **X Wang**, **C Xu**, *Algebraicity of the metric tangent cones and equivariant  $K$ -stability*, *J. Amer. Math. Soc.* 34 (2021) 1175–1214 [MR](#) [Zbl](#)
- [56] **C Li**, **C Xu**, *Special test configuration and  $K$ -stability of Fano varieties*, *Ann. of Math.* 180 (2014) 197–232 [MR](#) [Zbl](#)
- [57] **C Li**, **C Xu**, *Stability of valuations: higher rational rank*, *Peking Math. J.* 1 (2018) 1–79 [MR](#) [Zbl](#)
- [58] **C Li**, **C Xu**, *Stability of valuations and Kollár components*, *J. Eur. Math. Soc.* 22 (2020) 2573–2627 [MR](#) [Zbl](#)
- [59] **Y Li**, **Z Li**, *Equivariant  $\mathbb{R}$ -test configurations and semistable limits of  $\mathbb{Q}$ -Fano group compactifications*, preprint (2021) [arXiv 2103.06439](#)
- [60] **Y Liu**, *The volume of singular Kähler–Einstein Fano varieties*, *Compos. Math.* 154 (2018) 1131–1158 [MR](#) [Zbl](#)
- [61] **D Mumford**, *Stability of projective varieties*, *Enseign. Math.* 23 (1977) 39–110 [MR](#) [Zbl](#)
- [62] **A Okounkov**, *Brunn–Minkowski inequality for multiplicities*, *Invent. Math.* 125 (1996) 405–411 [MR](#) [Zbl](#)
- [63] **W Rossmann**, *Equivariant multiplicities on complex varieties*, from “Orbites unipotentes et représentations, III” (M Andler, editor), *Astérisque* 173–174, Soc. Math. France, Paris (1989) 313–330 [MR](#) [Zbl](#)
- [64] **G Székelyhidi**, *Filtrations and test-configurations*, *Math. Ann.* 362 (2015) 451–484 [MR](#) [Zbl](#)

- [65] **G Székelyhidi**, *The partial  $C^0$ -estimate along the continuity method*, J. Amer. Math. Soc. 29 (2016) 537–560 [MR](#)
- [66] **B Teissier**, *Valuations, deformations, and toric geometry*, from “Valuation theory and its applications, II” (F-V Kuhlmann, S Kuhlmann, M Marshall, editors), Fields Inst. Commun. 33, Amer. Math. Soc., Providence, RI (2003) 361–459 [MR](#) [Zbl](#)
- [67] **G Tian**, *Kähler–Einstein metrics with positive scalar curvature*, Invent. Math. 130 (1997) 1–37 [MR](#) [Zbl](#)
- [68] **G Tian**, *K-stability and Kähler–Einstein metrics*, Comm. Pure Appl. Math. 68 (2015) 1085–1156 [MR](#) [Zbl](#)
- [69] **G Tian, S Zhang, Z Zhang, X Zhu**, *Perelman’s entropy and Kähler–Ricci flow on a Fano manifold*, Trans. Amer. Math. Soc. 365 (2013) 6669–6695 [MR](#) [Zbl](#)
- [70] **G Tian, Z Zhang**, *Regularity of Kähler–Ricci flows on Fano manifolds*, Acta Math. 216 (2016) 127–176 [MR](#) [Zbl](#)
- [71] **G Tian, X Zhu**, *A new holomorphic invariant and uniqueness of Kähler–Ricci solitons*, Comment. Math. Helv. 77 (2002) 297–325 [MR](#) [Zbl](#)
- [72] **G Tian, X Zhu**, *Convergence of Kähler–Ricci flow*, J. Amer. Math. Soc. 20 (2007) 675–699 [MR](#) [Zbl](#)
- [73] **F Wang, X Zhu**, *Tian’s partial  $C^0$ -estimate implies Hamilton–Tian’s conjecture*, Adv. Math. 381 (2021) art. id. 107619 [MR](#) [Zbl](#)
- [74] **F Wang, X Zhu**, *Uniformly strong convergence of Kähler–Ricci flows on a Fano manifold*, Sci. China Math. 65 (2022) 2337–2370 [MR](#) [Zbl](#)
- [75] **X-J Wang, X Zhu**, *Kähler–Ricci solitons on toric manifolds with positive first Chern class*, Adv. Math. 188 (2004) 87–103 [MR](#) [Zbl](#)
- [76] **D Witt Nyström**, *Test configurations and Okounkov bodies*, Compos. Math. 148 (2012) 1736–1756 [MR](#) [Zbl](#)
- [77] **M Xia**, *On sharp lower bounds for Calabi-type functionals and destabilizing properties of gradient flows*, Anal. PDE 14 (2021) 1951–1976 [MR](#) [Zbl](#)
- [78] **C Xu**, *A minimizing valuation is quasi-monomial*, Ann. of Math. 191 (2020) 1003–1030 [MR](#) [Zbl](#)
- [79] **C Xu, Z Zhuang**, *On positivity of the CM line bundle on K-moduli spaces*, Ann. of Math. 192 (2020) 1005–1068 [MR](#) [Zbl](#)
- [80] **C Xu, Z Zhuang**, *Uniqueness of the minimizer of the normalized volume function*, Camb. J. Math. 9 (2021) 149–176 [MR](#) [Zbl](#)
- [81] **Y Yao**, *Relative Ding stability and an obstruction to the existence of Mabuchi solitons*, J. Geom. Anal. 32 (2022) art. id. 105 [MR](#) [Zbl](#)

Westlake University  
Hangzhou, China

Department of Mathematics, Rutgers University  
Piscataway, NJ, United States

[hanjiyuan@westlake.edu.cn](mailto:hanjiyuan@westlake.edu.cn), [chi.li@rutgers.edu](mailto:chi.li@rutgers.edu)

Proposed: Gang Tian

Seconded: Simon Donaldson, John Lott

Received: 14 April 2021

Revised: 1 June 2022

# GEOMETRY & TOPOLOGY

[msp.org/gt](http://msp.org/gt)

## MANAGING EDITOR

András I Stipsicz    Alfréd Rényi Institute of Mathematics  
[stipsicz@renyi.hu](mailto:stipsicz@renyi.hu)

## BOARD OF EDITORS

Mohammed Abouzaid	Stanford University <a href="mailto:abouzaid@stanford.edu">abouzaid@stanford.edu</a>	Mark Gross	University of Cambridge <a href="mailto:mgross@dpms.cam.ac.uk">mgross@dpms.cam.ac.uk</a>
Dan Abramovich	Brown University <a href="mailto:dan_abramovich@brown.edu">dan_abramovich@brown.edu</a>	Rob Kirby	University of California, Berkeley <a href="mailto:kirby@math.berkeley.edu">kirby@math.berkeley.edu</a>
Ian Agol	University of California, Berkeley <a href="mailto:ianagol@math.berkeley.edu">ianagol@math.berkeley.edu</a>	Bruce Kleiner	NYU, Courant Institute <a href="mailto:bkleiner@cims.nyu.edu">bkleiner@cims.nyu.edu</a>
Arend Bayer	University of Edinburgh <a href="mailto:arend.bayer@ed.ac.uk">arend.bayer@ed.ac.uk</a>	Sándor Kovács	University of Washington <a href="mailto:skovacs@uw.edu">skovacs@uw.edu</a>
Mark Behrens	University of Notre Dame <a href="mailto:mbehrens1@nd.edu">mbehrens1@nd.edu</a>	Urs Lang	ETH Zürich <a href="mailto:urs.lang@math.ethz.ch">urs.lang@math.ethz.ch</a>
Mladen Bestvina	University of Utah <a href="mailto:bestvina@math.utah.edu">bestvina@math.utah.edu</a>	Marc Levine	Universität Duisburg-Essen <a href="mailto:marc.levine@uni-due.de">marc.levine@uni-due.de</a>
Martin R Bridson	University of Oxford <a href="mailto:bridson@maths.ox.ac.uk">bridson@maths.ox.ac.uk</a>	Ciprian Manolescu	University of California, Los Angeles <a href="mailto:cm@math.ucla.edu">cm@math.ucla.edu</a>
Jim Bryan	University of British Columbia <a href="mailto:jbryan@math.ubc.ca">jbryan@math.ubc.ca</a>	Haynes Miller	Massachusetts Institute of Technology <a href="mailto:hmr@math.mit.edu">hmr@math.mit.edu</a>
Dmitri Burago	Pennsylvania State University <a href="mailto:burago@math.psu.edu">burago@math.psu.edu</a>	Tomasz Mrowka	Massachusetts Institute of Technology <a href="mailto:mrowka@math.mit.edu">mrowka@math.mit.edu</a>
Tobias H Colding	Massachusetts Institute of Technology <a href="mailto:colding@math.mit.edu">colding@math.mit.edu</a>	Aaron Naber	Northwestern University <a href="mailto:anaber@math.northwestern.edu">anaber@math.northwestern.edu</a>
Simon Donaldson	Imperial College, London <a href="mailto:s.donaldson@ic.ac.uk">s.donaldson@ic.ac.uk</a>	Peter Ozsváth	Princeton University <a href="mailto:petero@math.princeton.edu">petero@math.princeton.edu</a>
Yasha Eliashberg	Stanford University <a href="mailto:eliash-gt@math.stanford.edu">eliash-gt@math.stanford.edu</a>	Leonid Polterovich	Tel Aviv University <a href="mailto:polterov@post.tau.ac.il">polterov@post.tau.ac.il</a>
Benson Farb	University of Chicago <a href="mailto:farb@math.uchicago.edu">farb@math.uchicago.edu</a>	Colin Rourke	University of Warwick <a href="mailto:gt@maths.warwick.ac.uk">gt@maths.warwick.ac.uk</a>
David M Fisher	Rice University <a href="mailto:davidfisher@rice.edu">davidfisher@rice.edu</a>	Roman Sauer	Karlsruhe Institute of Technology <a href="mailto:roman.sauer@kit.edu">roman.sauer@kit.edu</a>
Mike Freedman	Microsoft Research <a href="mailto:michaelf@microsoft.com">michaelf@microsoft.com</a>	Stefan Schwede	Universität Bonn <a href="mailto:schwede@math.uni-bonn.de">schwede@math.uni-bonn.de</a>
David Gabai	Princeton University <a href="mailto:gabai@princeton.edu">gabai@princeton.edu</a>	Natasa Sesum	Rutgers University <a href="mailto:natasas@math.rutgers.edu">natasas@math.rutgers.edu</a>
Stavros Garoufalidis	Southern U. of Sci. and Tech., China <a href="mailto:stavros@mpim-bonn.mpg.de">stavros@mpim-bonn.mpg.de</a>	Gang Tian	Massachusetts Institute of Technology <a href="mailto:tian@math.mit.edu">tian@math.mit.edu</a>
Cameron Gordon	University of Texas <a href="mailto:gordon@math.utexas.edu">gordon@math.utexas.edu</a>	Ulrike Tillmann	Oxford University <a href="mailto:tillmann@maths.ox.ac.uk">tillmann@maths.ox.ac.uk</a>
Jesper Grodal	University of Copenhagen <a href="mailto:jg@math.ku.dk">jg@math.ku.dk</a>	Nathalie Wahl	University of Copenhagen <a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>
Misha Gromov	IHÉS and NYU, Courant Institute <a href="mailto:gromov@ihes.fr">gromov@ihes.fr</a>	Anna Wienhard	Universität Heidelberg <a href="mailto:wienhard@mathi.uni-heidelberg.de">wienhard@mathi.uni-heidelberg.de</a>

---

See inside back cover or [msp.org/gt](http://msp.org/gt) for submission instructions.

---

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

---

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

GT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>

© 2024 Mathematical Sciences Publishers



# GEOMETRY & TOPOLOGY

Volume 28    Issue 2 (pages 497–1003)    2024

---

On the top-weight rational cohomology of $\mathcal{A}_g$	497
MADELINE BRANDT, JULIETTE BRUCE, MELODY CHAN, MARGARIDA MELO, GWYNETH MORELAND and COREY WOLFE	
Algebraic uniqueness of Kähler–Ricci flow limits and optimal degenerations of Fano varieties	539
JIYUAN HAN and CHI LI	
Valuations on the character variety: Newton polytopes and residual Poisson bracket	593
JULIEN MARCHÉ and CHRISTOPHER-LLOYD SIMON	
The local (co)homology theorems for equivariant bordism	627
MARCO LA VECCHIA	
Configuration spaces of disks in a strip, twisted algebras, persistence, and other stories	641
HANNAH ALPERT and FEDOR MANIN	
Closed geodesics with prescribed intersection numbers	701
YANN CHAUBET	
On endomorphisms of the de Rham cohomology functor	759
SHIZHANG LI and SHUBHODIP MONDAL	
The nonabelian Brill–Noether divisor on $\overline{\mathcal{M}}_{13}$ and the Kodaira dimension of $\overline{\mathcal{R}}_{13}$	803
GAVRIL FARKAS, DAVID JENSEN and SAM PAYNE	
Orbit equivalences of $\mathbb{R}$ –covered Anosov flows and hyperbolic-like actions on the line	867
THOMAS BARTHELMÉ and KATHRYN MANN	
Microlocal theory of Legendrian links and cluster algebras	901
ROGER CASALS and DAPING WENG	
Correction to the article Bimodules in bordered Heegaard Floer homology	1001
ROBERT LIPSHITZ, PETER OZSVÁTH and DYLAN P THURSTON	