



Geometry & Topology

Volume 28 (2024)

**Orbit equivalences of \mathbb{R} -covered Anosov flows
and hyperbolic-like actions on the line**

THOMAS BARTHELMÉ

KATHRYN MANN

APPENDIX WRITTEN JOINTLY WITH JONATHAN BOWDEN

Orbit equivalences of \mathbb{R} -covered Anosov flows and hyperbolic-like actions on the line

THOMAS BARTHELMÉ

KATHRYN MANN

APPENDIX WRITTEN JOINTLY WITH JONATHAN BOWDEN

We prove a rigidity result for group actions on the line whose elements have what we call “hyperbolic-like” dynamics. Using this, we give a rigidity theorem for \mathbb{R} -covered Anosov flows on 3-manifolds, characterizing orbit equivalent flows in terms of the elements of the fundamental group represented by periodic orbits. As consequences of this, we give an efficient criterion to determine the isotopy classes of self-orbit equivalences of \mathbb{R} -covered Anosov flows, and prove finiteness of contact Anosov flows on any given manifold.

In the appendix, with Jonathan Bowden, we prove that orbit equivalences of contact Anosov flows correspond exactly to isomorphisms of the associated contact structures. This gives a powerful tool to translate results on Anosov flows to contact geometry and vice versa. We illustrate its use by giving two new results in contact geometry: the existence of manifolds with arbitrarily many distinct Anosov contact structures, answering a question of Foulon, Hasselblatt and Vaugon, and a virtual description of the group of contact transformations of a Anosov contact structure, generalizing a result of Giroux and Massot.

37D20, 57M60

1 Introduction

1.1 Hyperbolic-like actions

A well-known theorem of Hölder states that any group acting freely by homeomorphisms of the line is abelian and conjugate to a group of translations. This was generalized in unpublished work of Solodov (see eg [Farb and Franks 2003; Kovačević 1999; Barbot 1995a]) to the statement that a group action on the line where each nontrivial element has at most *one* fixed point is either semiconjugate to an action by affine transformations, or abelian with a global fixed point. Later, the proof of the convergence group theorem [Gabai 1992; Casson and Jungreis 1994] established that a group action on the circle where each element has at most *two* fixed points is, under some additional technical dynamical hypotheses, conjugate to a subgroup of $\mathrm{PSL}(2, \mathbb{R})$ acting on $\mathbb{R}P^1$ by Möbius transformations. This important result was the last step in the proof of the Seifert fiber space conjecture.

While one cannot reasonably expect further generalizations in this vein,¹ it is a natural question to ask what other fixed-point data might determine an action. In this spirit, we show that, under suitable hypotheses, an action of a group on the line is determined up to conjugacy by the set of elements acting with fixed points. Like the statement of the convergence group theorem, our hypotheses are motivated by an application to a classification problem, in our case the classification of \mathbb{R} -covered Anosov flows on 3-manifolds. Say that an action on the line is *hyperbolic-like* if it commutes with integer translation and each nontrivial element either acts freely or has exactly two fixed points in $[0, 1)$, one attracting and one repelling. We prove the following rigidity result for such actions:

Theorem 1.1 (rigidity of hyperbolic-like actions) *A minimal, hyperbolic-like action of a nonabelian group G on \mathbb{R} is determined up to conjugacy by the set of elements of G that act with fixed points.*

Minimal and hyperbolic-like are both properties of the actions of 3-manifold fundamental groups on \mathbb{R} induced by \mathbb{R} -covered Anosov flows. This allows us to use Theorem 1.1 to classify such flows up to orbit equivalence.

1.2 Orbit equivalence of Anosov flows

Recall that two flows on a manifold M are *orbit equivalent* if there is a homeomorphism $f: M \rightarrow M$ taking orbits of one to orbits of the other, and *isotopically equivalent* if this homeomorphism can be taken to be isotopic to the identity.

An Anosov flow is called \mathbb{R} -covered if the leaf space of its weak-stable foliation is homeomorphic to \mathbb{R} . (In the case of 3-manifolds, it is equivalent that the weak-unstable foliation has leaf space \mathbb{R}). On 3-manifolds, there are many constructions of \mathbb{R} -covered flows, and examples of manifolds admitting arbitrarily many inequivalent \mathbb{R} -covered flows. Here, we give a characterization of orbit and isotopy equivalent \mathbb{R} -covered flows on 3-manifolds by their *free homotopy classes of periodic orbits*. For a flow φ on a manifold M , let $\mathcal{P}(\varphi)$ denote the set of conjugacy classes of elements in $\pi_1(M)$ represented by the free homotopy classes of periodic orbits of φ . We show the following:

Theorem 1.2 (classification of \mathbb{R} -covered Anosov flows) *Let φ and ψ be \mathbb{R} -covered Anosov flows on a closed 3 manifold M .*

- (1) *φ and ψ are isotopically equivalent if and only if $\mathcal{P}(\varphi) = \mathcal{P}(\psi)$.*
- (2) *φ and ψ are orbit equivalent² if and only if there exists a homeomorphism $f: M \rightarrow M$ such that $f_*(\mathcal{P}(\varphi)) = \mathcal{P}(\psi)$. Moreover, the orbit equivalence can be taken to be in the isotopy class of f .*

¹One reason for this is that the target groups \mathbb{R} , $\text{Aff}_+(\mathbb{R})$ and $\text{PSL}(2, \mathbb{R})$ are essentially the only Lie groups acting transitively on 1-manifolds, so the only natural candidates for such targets.

²In our definition of orbit equivalence, we do not require the homeomorphism to match *oriented* orbits to oriented orbits. If one wants to consider only orbit equivalences that preserve orbit orientation, then the conclusion of Theorem 1.2 will be that, if one flow is transversally orientable, then the other is also, and the orbit equivalence can be upgraded to an orientation-preserving orbit equivalence (using [Barbot 1995a, théorème C]). If one (and hence both) of the flows are not transversally orientable, then φ is orbit equivalent to either ψ or ψ^{-1} .

The \mathbb{R} -covered Anosov flows on closed 3-manifolds form a rich class of examples, including geodesic flows on closed surfaces, all contact Anosov flows, and diverse examples on many hyperbolic 3-manifolds and on manifolds with nontrivial JSJ decomposition (see [Fenley 1994; Foulon and Hasselblatt 2013; Barthelmé and Fenley 2017; Bonatti and Iakovoglou 2023; Bowden and Mann 2022]). Among other applications, our main theorem allows us to prove finiteness of contact Anosov flows. We describe and motivate the main applications now.

1.3 Application 1: classifying self-orbit equivalences

Describing the centralizer of a given diffeomorphism is a classical question in discrete-time dynamical systems; notably, Smale's conjecture [1998] is that the centralizer of a generic diffeomorphism should be trivial. By contrast, diffeomorphisms which embed in a flow have an \mathbb{R} -subgroup in their centralizer given by the flow, so the right analog of Smale's conjecture in this case is to ask whether the flow agrees (virtually) with the centralizer of the diffeomorphism. This motivates the general program to classify all symmetries of a given flow, and, more generally, classify the symmetries of the foliation given by orbits of a flow, ie classify the self-orbit equivalences. As with Smale's conjecture, this question is quite sensitive to regularity — for instance, 3-dimensional Anosov flows often have many self-orbit equivalences, while the set of those which may be realized by C^1 diffeomorphisms was shown to be virtually trivial by Barthelmé, Fenley and Potrie [Barthelmé et al. 2023].

Theorem 1.2 gives the following immediate characterization of isotopy classes of self-orbit equivalences:

Corollary 1.3 *A map $f: M \rightarrow M$ is in the isotopy class of a self-orbit equivalence of an \mathbb{R} -covered Anosov flow φ if and only if $f_*: \pi_1(M) \rightarrow \pi_1(M)$ preserves the set of conjugacy classes realized by periodic orbits of φ .*

With more work, we improve this to give a criterion to explicitly describe such classes, as follows. The case of interest here is for *skew* flows, those which are not orbit equivalent to a suspension of an Anosov diffeomorphism on the torus, as the orbit equivalences of suspension flows are essentially trivial.

If M is a 3-manifold with a skew Anosov flow φ , then M is orientable and irreducible, so admits a JSJ decomposition along tori into Seifert and atoroidal pieces. By Mostow rigidity and the structure of mapping class groups of Seifert spaces, the group $\text{Dehn}(M)$ of isotopy classes of diffeomorphisms generated by Dehn twists along embedded tori in M has finite index in $\text{MCG}(M)$ [Johannson 1979]. We give a criterion for when maps generated by certain Dehn twists represent a self-orbit equivalence of a flow. Given a flow φ and Dehn twist D_β , we may define an *orbit displacement* function, as follows. For each periodic orbit c in M , we set

$$d_\varphi(c, D_\beta) = \sum_{i=1}^k 2\epsilon_i t_\varphi(\beta_i),$$

where k is the number of transverse intersections of c with T (when T is in quasitransverse position with respect to the flow, $t_\varphi(\beta)$ is the *translation number* of the action of β on $\Lambda^s(\varphi)$, and $\epsilon_i = \pm 1$ is

an orientation term for each intersection. When f is a composition of Dehn twists D_{β_i} on disjoint nonisotopic tori, we define $d_\varphi(c, f)$ to be the sum of the displacements $d_\varphi(c, D_{\beta_i})$. Formal definitions are given in Section 3.

Theorem 1.4 (criterion for self-orbit equivalence) *Let φ be a transversally oriented skew Anosov flow. A map f which is a composition of Dehn twists on disjoint nonisotopic tori is isotopic to a self-orbit equivalence of φ if and only if, for all periodic orbits c of φ , we have $d_\varphi(c, f) = 0$.*

There are many situations in which this criterion is easy to check. As two sample applications, we have the following results for any transversally oriented skew Anosov flow φ :

Corollary 1.5 *Let D_{per} be the subgroup of the mapping class group of M generated by Dehn twists along curves represented by periodic orbits. Then any isotopy class $[h] \in D_{\text{per}}$ is represented by a self-orbit equivalence of φ .*

Corollary 1.6 *Let T be an embedded torus and D_β a Dehn twist on T with nonzero translation number. Then D_β is isotopic to a self-orbit equivalence of φ if and only if T is separating in M .*

In Section 3.2, we discuss a number of other special cases where one may use the topology of M to reduce Theorem 1.4 to a simpler statement. We also discuss the complementary case to Theorem 1.4 for maps generated by Dehn twists in tori which cannot be realized disjointly, namely tori inside a single Seifert piece. See Theorems 3.17 and 3.18.

Remark 1.7 One can easily describe all self-orbit equivalences of a given skew Anosov flow in a fixed isotopy class. Such a flow comes with the data of a homeomorphism $\eta: M \rightarrow M$ realizing the *half-step-up* map on the orbit space. See Section 2.1 for details. It follows from [Barthelmé and Gogolev 2019, Theorem 1.1] that any two self-orbit equivalences h_1 and h_2 in the same isotopy class differ, up to isotopy along the flow lines, by some power of η .

Theorem 1.4 identifies which elements of a large subgroup of the mapping class group of M are represented by self-orbit equivalences. However, passing to the full mapping class group requires a different approach. In particular, we do not know the answer to the following:

Question 1.8 Does there exist an (\mathbb{R} -covered or not) Anosov flow on a hyperbolic 3-manifold M such that the only self-orbit equivalences are isotopic to the identity?

Question 1.9 Does there exist an (\mathbb{R} -covered or not) Anosov flow on a hyperbolic 3-manifold M such that every element of the mapping class group of M is represented by a self-orbit equivalence?

Remark 1.10 The case of most interest for both questions is when the manifold considered has nontrivial mapping class group. However, there are, as yet, no constructions of Anosov flows on a 3-manifold that has a trivial mapping class group. So even the trivial case for these questions is not yet known.

Remark 1.11 Barthelmé et al. [2023] introduced a class of partially hyperbolic diffeomorphisms, called “collapsed Anosov flows”, which are semiconjugate to self-orbit equivalences of Anosov flows. Hence, the criterion of Theorem 1.4 (and its applications for certain manifolds, as in Theorems 3.17 and 3.18) describes the possible isotopy classes of collapsed Anosov flows associated with \mathbb{R} -covered Anosov flows.

1.4 Application 2: contact Anosov flows

An Anosov flow is said to be *contact* if it is the Reeb flow of a contact 1-form α . Notice that, with this definition, the contact structure $\xi = \ker \alpha$ is automatically transversely orientable since it is given as the kernel of a (globally defined) contact form. The contact Anosov flows are an important and well-studied class of examples, as they can be thought of as a generalization of the geodesic flow on manifolds of negative curvature, and many dynamical results on Anosov flows, for instance exponential decay of correlations [Liverani 2004], are known only for the contact case. In the context of 3-manifolds, Barbot [2001] proved that contact Anosov flows on 3-manifolds are necessarily \mathbb{R} -covered and skew, while Foulon–Hasselblatt surgery [Foulon et al. 2021] produces many examples. In fact, it is currently an open question whether every \mathbb{R} -covered skew flow is orbit equivalent to a contact flow. As progress towards a better understanding of these flows, we show that isomorphism of the associated contact structures is the same as orbit equivalence of flows, giving a powerful tool to use the machinery of flows to answer questions in contact geometry and vice versa.

Theorem 1.12 *Two contact Anosov flows on a 3-manifold are orbit equivalent if and only if their respective contact structures are contactomorphic. They are isotopically equivalent if and only if the contact structures are isotopic.*

Recall that two contact structures ξ_1 and ξ_2 on a manifold M are *contactomorphic* if there exists a diffeomorphism $g: M \rightarrow M$ such that $g_*\xi_1 = \xi_2$, and they are *isotopic* if g can be taken to be isotopic to the identity.

We prove the reverse of Theorem 1.12 in Section 4, and the forward direction in Theorem A.2. Using the coarse classification of tight contact structures of Colin, Giroux and Honda [Colin et al. 2009] and the result (proved in the appendix) that Anosov contact structures have zero torsion, we also obtain the following:

Theorem 1.13 (finiteness for contact Anosov flows) *On any given 3-manifold M , there are only finitely many contact Anosov flows on M up to orbit equivalence.*

Thanks to Theorem 1.12, one can now fully translate results about contact Anosov flows to results about Anosov contact structures and vice versa. We illustrate this principle in the appendix with two

examples: First, we show (Theorem A.7) that there exists hyperbolic 3-manifolds with arbitrarily many noncontactomorphic Anosov contact structures, answering a question raised in [Foulon et al. 2021]. Second, we give a virtual description of the group of contact transformations of a Anosov contact structure up to isotopy on some manifolds (Theorem A.8). This generalizes a result by Giroux and Massot [2017].

Outline of the article

Section 2 gives a brief introduction to the structure of \mathbb{R} -covered flows on 3-manifolds, followed by the proof of Theorem 1.2. Section 3 contains the proof of Theorem 1.4, and the applications to contact flows are given in Section 4 and the appendix.

Acknowledgements

Mann was partially supported by NSF CAREER grant DMS 1844516 and a Sloan Fellowship. Barthelmé was partially supported by the NSERC (funding reference number RGPIN-2017-04592). Bowden was partially supported by the Special Priority Programme SPP 2026 *Geometry at infinity* funded by the DFG. The authors thank Anne Vaugon and Vincent Colin for very helpful discussions. We also thank Thierry Barbot and Sergio Fenley for their detailed comments and suggestions on an earlier version of the article.

2 Proofs of Theorems 1.1 and 1.2

The first statement of Theorem 1.2 is a special case of the second, so we in fact only need to prove that assertion. Recall this is the statement that two flows φ and ψ are orbit equivalent if and only if there exists a homeomorphism $f: M \rightarrow M$ such that $f_*(\mathcal{P}(\varphi)) = \mathcal{P}(\psi)$, and, if this holds, the orbit equivalence can be taken to be in the isotopy class of f . The forward direction is immediate; we now set up the proof of the reverse direction. This will lead us to the statement and proof of Theorem 1.1; we finish the proof of Theorem 1.2 at the end of this section.

Throughout the work, we assume the reader has basic familiarity with Anosov flows. We recall below the essential structure theory of \mathbb{R} -covered flows on 3-manifolds that is used in the proof. Further background can be found in [Fisher and Hasselblatt 2019], and results specific to the topological theory of Anosov flows in dimension 3 can be found in [Barbot 2005].

By work of Fenley [1994] and Barbot [1995a], an \mathbb{R} -covered Anosov flow on a closed 3-manifold is either conjugate to the suspension of an Anosov diffeomorphism of T^2 or is *skew*, meaning that the orbit space of the lift of the flow to \tilde{M} is homeomorphic to the infinite diagonal strip

$$\mathbb{C} = \{(x, y) \in \mathbb{R}^2 : |x - y| < 1\}$$

via a homeomorphism taking the stable leaves of the flows to the horizontal cross-sections of the strip, and unstable leaves to the vertical cross-sections.




Figure 1: The orbit space \mathcal{O} .

Theorem 1.2 follows from purely topological considerations in the suspension case, as follows. Suppose that M is a 3-manifold that fibers as the mapping torus of an Anosov diffeomorphism A on the torus. By a theorem of Plante [1981], any Anosov flow on M is necessarily of suspension type. Any other suspension flow comes from a fibering of M as the mapping torus of an Anosov diffeomorphism. It is a “folklore” result that such a diffeomorphism must be conjugate to either A or A^{-1} — a detailed proof can be found in [Funar 2013]. Since our definition of orbit equivalence allows a flow to be conjugate to its inverse, we conclude that M admits only one Anosov flow up to orbit equivalence.

The case of skew flows is much more interesting. For instance, examples of (closed, hyperbolic) manifolds that admit arbitrarily many inequivalent skew Anosov flows were constructed in [Bowden and Mann 2022]. As a first step towards the proof of Theorem 1.2 in the skew case, we need to recall some general structure theory due to Barbot and Fenley that will allow us to essentially reduce the theorem to a statement about actions of $\pi_1(M)$ on S^1 .

2.1 Skew Anosov flows

Consider again the infinite diagonal strip model for the orbit space as shown in Figure 1.

In this model, each point $o \in \mathcal{O}$ can be assigned a point o_u on the upper boundary of the strip by following the unstable leaf through o , and a point o_l on the lower boundary by following the stable leaf. Taking the intersection of the stable leaf through o_u and unstable through o_l defines a continuous, fixed-point-free map $\eta: \mathcal{O} \rightarrow \mathcal{O}$, which we call the *half-step-up* map.³ This map exchanges stable leaves and unstable leaves, so $\tau = \eta^2$ induces a map on the leaf space Λ^s of the weak stable foliation. We call τ the *one-step-up* map.

If the weak foliations are transversely orientable, then τ commutes with the action of $\pi_1(M)$ on Λ^s . Identifying $\Lambda^s \cong \mathbb{R}$ so that τ is identified with the translation $x \mapsto x + 1$ realizes $\pi_1(M)$ as a subgroup

³This map and its square have both been referred to as the one-step-up map elsewhere in the literature. We choose to call the square the one-step-up and η the half-step-up.

of $\text{Homeo}^{\mathbb{Z}}(\mathbb{R})$, the group of orientation-preserving homeomorphisms of \mathbb{R} commuting with integer translations. In the nonorientable case, τ is twisted π_1 -equivariant: for any $\gamma \in \pi_1(M)$ that reverses the orientation of Λ^s , we have $\tau(\gamma x) = \gamma \tau^{-1}(x)$.

Our perspective going forward will be to study the flows through the action of $\pi_1(M)$ on the leaf space. For simplicity, in Section 2.3 we assume first that the weak foliations are transversely orientable, and then state the necessary modifications for the nonorientable case. In the orientable case, the dynamics of the action of $\pi_1(M)$ on Λ^s is what Thurston [1997] calls an *extended convergence group action*. However, the only dynamical property that we will need is the fact that, in such a group, any orientation-preserving homeomorphism with fixed points has exactly two fixed points in $[0, 1)$, one attracting and one repelling. This can be seen directly from the Anosov dynamics of the flow.

Another dynamical property that will be of use comes from [Barbot 1995a, Theorem 2.5], which states that skew Anosov flows are transitive and the action of the group generated by $\pi_1(M)$ and τ on \mathbb{R} is *minimal*, meaning that all orbits are dense. Since $\mathbb{R}/\tau \cong S^1$ and the action of $\pi_1(M)$ descends to this circle, this latter statement is equivalent to the statement that the action of $\pi_1(M)$ on \mathbb{R}/τ is minimal. The reader may find it useful to visualize the action on \mathbb{R} by thinking of it on the circle \mathbb{R}/τ , but for simplicity we will work in $\text{Homeo}^{\mathbb{Z}}(\mathbb{R})$. The next subsection establishes some general results about such subgroups of $\text{Homeo}^{\mathbb{Z}}(\mathbb{R})$, concluding with the proof of Theorem 1.1.

2.2 Hyperbolic-like homeomorphisms: proof of Theorem 1.1

Definition 2.1 We say an element $f \in \text{Homeo}^{\mathbb{Z}}(\mathbb{R})$ is *hyperbolic-like* if it has exactly two fixed points in $[0, 1)$, one attracting and one repelling, and a group action $\rho: G \rightarrow \text{Homeo}(\mathbb{R})$ is hyperbolic-like if its image lies in $\text{Homeo}^{\mathbb{Z}}(\mathbb{R})$ and every element with fixed points is hyperbolic-like.

In the context of the action of the fundamental group of a 3-manifold with a skew Anosov flow on the stable leaf space (which is what we have in mind), Thurston [1997] calls hyperbolic-like elements *space-like* homeomorphisms. Since this notation is not commonplace, we have chosen the terminology “hyperbolic-like” since the induced action of such elements on \mathbb{R}/\mathbb{Z} are topologically conjugate to hyperbolic Möbius transformations.

If a is a hyperbolic-like element, we will use the notation a_+ and a_- to respectively denote an attracting and a repelling fixed point for a . For two hyperbolic-like elements a and b , we say the fixed sets of a and b are *linked* if each connected component of $\mathbb{R} \setminus \text{Fix}(a)$ contains a fixed point for b , or if a and b have fixed points in common. We say they are *unlinked* otherwise. By convention, when we speak about the order of fixed points on \mathbb{R} , we use the notation

$$a_+ < b_- < a_- < b_+$$

to mean that there exist four *consecutive* elements of $\text{Fix}(a) \cup \text{Fix}(b)$, ordered as indicated by the inequality. In this example, the fixed sets of a and b are linked. Since both a and b commute with integer translation,

the next element of $\text{Fix}(a) \cup \text{Fix}(b)$ to the right of b_+ is another attracting fixed point for a , equal to $a_+ + 1$.

The next series of lemmas shows that the configuration of fixed points of a pair or triple of elements can be detected by the set of words in those elements which act with fixed points.

Lemma 2.2 *Let a and b be hyperbolic-like elements of $\text{Homeo}^{\mathbb{Z}}(\mathbb{R})$. The fixed sets of a and b are linked if and only if $\text{Fix}(a^n b^m) \neq \emptyset$ for all $n, m \in \mathbb{Z}$.*

Proof Suppose first that the fixed sets of a and b are linked, and let n and m be given. Since the property of having linked fixed sets does not change after passing to inverses, up to replacing a or b with their inverses we can assume that $n, m \geq 0$. Since the fixed sets of a and b are linked, either a and b have a common fixed point (in which case we are done) or there exists some connected component of $\mathbb{R} \setminus (\text{Fix}(a) \cup \text{Fix}(b))$ bounded on one side by an attracting fixed point for a , and on the other by an attracting fixed point for b . Let I denote the closure of this component. Then $a^n b^m(I) \subset I$, so $a^n b^m$ has a fixed point in I .

To prove the converse, suppose now that the sets are unlinked. Up to passing to inverses, we may find consecutive attracting and repelling fixed points for a and b that lie in the order

$$a_+ < a_- < b_+ < b_-$$

with no other fixed points between a_+ and b_- . For m large enough, $b^m(a_+)$ will lie in the open interval (a_-, b_+) . For n large enough, $a^n b^m(a_+)$ will therefore lie in the open interval $(b_-, a_+ + 1)$. Similarly, $b^m a^n b^m(a_+)$ will lie to the right of $a_+ + 1$, as will $a^n b^m a^n b^m(a_+)$. Thus, $(a^n b^m)^2$ translates some point a distance at least 1. Any such element of $\text{Homeo}^{\mathbb{Z}}(\mathbb{R})$ is fixed-point-free; hence, its root $a^n b^m$ is fixed-point-free as well. \square

The next lemma says that, if a and b have unlinked fixed sets, then we can detect the cyclic order of attracting and repelling fixed points by understanding which words in a and b have fixed points.

Lemma 2.3 *Suppose a and b have unlinked fixed sets. The word $b^N a^N$ has a fixed point for every $N > 0$ if and only if one may find a set of consecutive fixed points either in the order*

$$a_+ < a_- < b_- < b_+$$

or that obtained from the above by replacing a and b simultaneously with their inverses.

Proof If the ordering shown above occurs, then the interval $[b_+, a_+ + 1]$ is mapped into itself by $b^N a^N$, so the map has a fixed point. Note that the ordering obtained by replacing a and b with their inverses can also be obtained simply by reversing the orientation of \mathbb{R} and considering a sequence of consecutive fixed points starting with a_- . Since reversing orientation of \mathbb{R} obviously does not change the property of

a map having fixed points, we have already proved one direction of the lemma. For the converse, if the ordering above does not occur even after passing to inverses, the order is either

$$a_+ < a_- < b_+ < b_- \quad \text{or} \quad a_- < a_+ < b_- < b_+.$$

In the first case, for sufficiently large N we have $(b^N a^N)^2(b_+) > b_+ + 1$, so the map is fixed-point-free, and in the second case we have $(b^N a^N)^2(b_+) < b_+ - 1$, so the map is again fixed-point-free. \square

Our next goal is to use this information to reconstruct a minimal action, up to conjugacy, from the data of the ordering of fixed-point sets. For this we need an elementary lemma:

Lemma 2.4 *Suppose $G \subset \text{Homeo}^{\mathbb{Z}}(\mathbb{R})$ is a nonabelian group whose action on \mathbb{R} is minimal and hyperbolic-like. Then, for any point $x \in \mathbb{R}$ and any $\epsilon > 0$, there exists $a \in G$ such that a has two fixed points in the ϵ -neighborhood of x .*

Proof If no nontrivial element of G acted with fixed points, then G would be abelian by Hölder's theorem, so by assumption this case does not occur. Thus, there exist hyperbolic-like elements and, by minimality of the action of G , the set of their attracting fixed points is dense.

Let x and $\epsilon > 0$ be given. Fix any hyperbolic-like element g with an attracting fixed point in the $\frac{\epsilon}{2}$ -neighborhood of x . Observe that, if f and g are hyperbolic-like and f does not fix a repelling point g_- for g , then all fixed points of the conjugate $g^N f g^{-N}$ approach the attracting fixed points of g as $N \rightarrow \infty$. Thus, it suffices to find a hyperbolic-like f that does not fix g_- . By minimality, there exists $h \in G$ such that $h(g_-)$ lies strictly between g_- and g_+ , where $g_- < g_+$ are consecutive fixed points. If $h(g_+ - 1) \neq g_-$, then hgh^{-1} is hyperbolic-like and has fixed points distinct from g_- , since they are the images of the fixed points of g under h , and we are done. If instead $h(g_+ - 1) = g_-$, or equivalently $h(g_+) = g_- + 1$, then we have

$$g_- < h^{-1}(g_+) < g_+ < g_- + 1$$

and therefore $h^{-1}gh$ has fixed points g_+ and $h^{-1}(g_+) \neq g_-$. \square

Although not strictly needed in our proof, Lemma 2.4 can be strengthened to the following density for pairs of fixed points:

Lemma 2.5 *Suppose $G \subset \text{Homeo}^{\mathbb{Z}}(\mathbb{R})$ is a nonabelian group whose action on \mathbb{R} is minimal and hyperbolic-like. Given $x, y \in [0, 1)$, and $\epsilon > 0$, there exists a hyperbolic-like element $g \in G$ with fixed points satisfying $|g_- - x| < \epsilon$ and $|g_+ - y| < \epsilon$.*

Proof Let $x, y \in [0, 1)$ and $\epsilon > 0$ be given. Without loss of generality, assume $x < y$ and assume that ϵ is small enough that the ϵ -neighborhoods of x and y and all of their integer translates are pairwise disjoint.

By Lemma 2.4, we may find hyperbolic-like elements a and b with fixed points in the $\frac{\epsilon}{2}$ -neighborhoods of x and y , respectively. Replacing a or b with their inverses if needed, we can assume these fixed points are ordered

$$a_+ < a_- < b_- < b_+.$$

By Lemma 2.3, this implies that $b^N a^N$ has a fixed point for every $N > 0$. Furthermore, if N is sufficiently large, an attracting fixed point for $b^N a^N$ will lie within the $\frac{\epsilon}{2}$ -neighborhood of b_+ , and a repelling fixed point within the $\frac{\epsilon}{2}$ -neighborhood of a_- ; this is simply because $b^N a^N$ takes a complement of the $\frac{\epsilon}{2}$ -neighborhood of the union of translates of a_- to a neighborhood of the attracting fixed points for b . \square

We can now finish the proof of our first main theorem; first we recall the statement:

Theorem 1.1 (rigidity of hyperbolic-like actions) *A minimal, hyperbolic-like action of a nonabelian group G on \mathbb{R} is determined up to conjugacy by the set of elements of G that act with fixed points.*

Proof Let G be nonabelian, hyperbolic-like and acting minimally on \mathbb{R} , and $\rho(G)$ another such faithful action of G on \mathbb{R} , with the same set of hyperbolic-like elements as G . Let $g \in G$ be a hyperbolic-like element (recall that such an element exists by Hölder's theorem since G is nonabelian). Choose coordinates on \mathbb{R} so that the attracting fixed points of g and of $\rho(g)$ are precisely the integers. We need to choose an orientation on the line \mathbb{R} on which ρ acts. To do this, take some $f \in G$ with an axis unlinked with g . Such a map f exists by Lemma 2.4. Replacing f with its inverse if needed, we can find such a map for which consecutive fixed points are ordered

$$g_+ < g_- < f_- < f_+.$$

By Lemma 2.2 and the fact the action of $\rho(G)$ has the same elements with fixed points as the original action of G , we conclude that the fixed sets of $\rho(g)$ and $\rho(f)$ are unlinked. By the same reasoning, using Lemma 2.3, the map $f^N g^N$ has a fixed point for all $N > 0$, so by hypothesis the same is true for $\rho(f)^N \rho(g)^N$. Applying the other direction of Lemma 2.3, we can now fix an orientation on \mathbb{R} so that consecutive fixed points of $\rho(f)$ and $\rho(g)$ are ordered

$$\rho(g)_+ < \rho(g)_- < \rho(f)_- < \rho(f)_+.$$

Our next goal is to show that this determines the ordering of the set of all attracting fixed points of hyperbolic-like elements of $\rho(G)$. We then define a map Θ on the (dense) subset of \mathbb{R} consisting of attracting fixed points of other elements by sending the unique attracting fixed point of an element h that lies in $[m, m + 1)$ to the unique attracting fixed point of $\rho(h)$ in $[m, m + 1)$. This order-preserving property is sufficient to show that our map Θ is continuous, from which it will easily follow that it can be extended continuously to a homeomorphism of \mathbb{R} that conjugates the action of G and $\rho(G)$.

Consider first an element h with fixed set that is unlinked with both f and g . Up to switching h with h^{-1} , there are three cases to consider.

Case 1 Suppose first that we have the ordering

$$g_+ < g_- < f_- < h_- < h_+ < f_+.$$

Applying Lemma 2.3, we have that $f^N h^N$ and $g^N h^N$ have fixed points for all positive N , and hence so do $\rho(f)^N \rho(h)^N$ and $\rho(g)^N \rho(h)^N$. Applying the lemma again implies that one of these three orderings occur:

$$(*) \quad \rho(g)_+ < \rho(h)_+ < \rho(h)_- < \rho(g)_- < \rho(f)_- < \rho(f)_+,$$

$$(**) \quad \rho(h)_+ < \rho(g)_+ < \rho(g)_- < \rho(h)_- < \rho(f)_- < \rho(f)_+,$$

$$(***) \quad \rho(g)_+ < \rho(g)_- < \rho(f)_- < \rho(h)_- < \rho(h)_+ < \rho(f)_+.$$

We want to show that only case (***) can occur. We will show that (*) does not occur. Eliminating the possibility of (**) is done by exactly the same argument, switching the roles of g and h ; we omit the details.

Since $\text{Fix}(aba^{-1}) = a \text{Fix}(b)$, for any $n > 0$ we also have

$$g_+ < g_- < f_- < (f^n h f^{-n})_- < (f^n h f^{-n})_+ < f_+$$

and so $g^N (f^n h f^{-n})^N$ has a fixed point for all $N > 0$, as does its image under ρ . If ordering (*) were to occur, then for sufficiently large n we would have

$$\rho(g)_+ < \rho(g)_- < \rho(f)_- < \rho(f)_+ < \rho(f^n h f^{-n})_+ < \rho(f^n h f^{-n})_-,$$

contradicting Lemma 2.3 applied to $\rho(g)^N \rho(f^n h f^{-n})^N$. Thus, the ordering of consecutive fixed points of f , g and h under ρ agrees with that for the original action.

Case 2 The ordering

$$g_+ < h_- < h_+ < g_- < f_- < f_+.$$

is handled exactly as above, exchanging the roles of f and g .

Case 3 Now suppose instead that the ordering of the fixed points of f , g and h is

$$g_+ < g_- < h_+ < h_- < f_- < f_+.$$

Consider the elements $a = g^n f^n$ and $b = f^n h^n$ for some large positive n . As $n \rightarrow \infty$, the attracting fixed point a_+ approaches f_+ , and similarly a_- approaches g_- , b_+ approaches h_+ , and b_- approaches f_- . Thus, provided n is chosen large enough, we have

$$g_+ < g_- < a_- < b_+ < h_+ < h_- < b_- < f_- < f_+ < a_+.$$

We can then apply the previous cases to the triples (g, a, h) , (g, a, f) , (g, a, b) and (g, b, h) to show the ordering of their fixed points is preserved by ρ . We deduce that the ordering of the fixed points of $\rho(f)$, $\rho(g)$ and $\rho(h)$ matches that of the fixed points of f , g , and h .

We can now quickly finish the proof of the theorem. Suppose we have some hyperbolic-like a and b with

$$0 \leq a_+ < b_+ < 1.$$

Rather than consider cases depending on whether a and b are linked or not, we can instead use Lemma 2.4 to choose c and d in G with fixed sets very close to a_+ and b_+ such that c and d have fixed sets unlinked with a and b . Suppose for simplicity that $0 < a_+$ (otherwise, simply replace 0 in what follows with some very small $-\epsilon$ and 1 with $1 - \epsilon$, and repeat the proof). By Lemma 2.5, we can choose elements c and d with fixed points so that we have the ordering

$$0 < c_- < c_+ < a_+ < d_- < d_+ < b_+ < 1$$

and additionally have that a does not have a repelling fixed point between c_- and a_+ , and b does not have a repelling fixed point between d_- and d_+ . We can also choose such c and d whose fixed sets are each unlinked with respect to f and to g , so that we may apply the observation above to c and d and determine the relative order of their fixed-point sets. Any choice of c and d with fixed sets sufficiently close to a_+ and b_+ will have this property. Thus, by our convention on $\rho(g)$, we conclude that

$$\rho(g)_+ = 0 < \rho(c)_- < \rho(c)_+ < \rho(d)_- < \rho(d)_+ < 1.$$

Since c and d had unlinked fixed points with respect to a , b and g , we can apply the observation again and conclude that the ordering of fixed sets is preserved, namely

$$\rho(g)_+ = 0 < \rho(c)_- < \rho(c)_+ < \rho(a)_+ < \rho(d)_- < \rho(d)_+ < \rho(b)_+ < 1$$

and, in particular, $\rho(g)_+ = 0 < \rho(a)_+ < \rho(b)_+ < 1$, as we needed to show.

Thus, we have defined an order-preserving (and hence continuous) injective map between two dense subsets of \mathbb{R} . This extends uniquely to a continuous map $\mathbb{R} \rightarrow \mathbb{R}$ with continuous inverse, which we denote by Θ . It remains to see that Θ conjugates the actions of G and $\rho(G)$. Let $g \in G$ be given. Note that, if a_+ is a fixed point of a hyperbolic-like element of $a \in G$, then $g(a_+)$ is an attracting fixed point of (gag^{-1}) ; thus, $\Theta(ga_+)$ is some attracting fixed point of $\rho(gag^{-1})$, ie the image of an attracting fixed point of $\rho(a)$ under $\rho(g)$. In other words, $\Theta(ga_+) = \rho(g)\Theta(a_+) + n$ for some $n \in \mathbb{Z}$. Since Θ is continuous and fixed points of hyperbolic-like elements are dense in the source and the target, we conclude that $n = 0$, so g -equivariance holds on a dense set, and hence everywhere. \square

Remark 2.6 If G is not assumed to act minimally (but $\rho(G)$ is), the same proof strategy can be used to produce a *semiconjugacy* between the actions of G and $\rho(G)$, defined on the closure of the G -invariant set consisting of hyperbolic fixed points.

2.3 Conclusion of the proof of Theorem 1.2

Returning to the setup of Theorem 1.2, suppose that φ and ψ are two skew Anosov flows with $f_*\mathcal{P}(\varphi) = \mathcal{P}(\psi)$ for some homeomorphism $f: M \rightarrow M$. Replacing φ with its conjugate under f , we obtain a

flow φ satisfying $\mathcal{P}(\varphi) = \mathcal{P}(\psi)$. What we need to show is that φ and ψ are orbit equivalent by some homeomorphism of M that is isotopic to the identity.

Orientable case Assume first that $\Lambda^s(\varphi)$ and $\Lambda^s(\psi)$ are both transversely orientable. We then will apply Theorem 1.1 to the group $G := \pi_1(M)$. Recall the action of this group on the leaf space $\Lambda^s(\varphi) \cong \mathbb{R}$ is faithful, minimal and has the property that all elements with fixed points are hyperbolic-like, under the parametrization of \mathbb{R} where $\tau(\varphi)$ acts as translation by 1, and the same holds for ψ .

That $\mathcal{P}(\varphi) = \mathcal{P}(\psi)$ means precisely that these two representations have the same elements with fixed points. Thus, Theorem 1.1 implies that these two actions are conjugate. This also gives a conjugacy between the actions on the unstable leaf spaces $\Lambda^u(\varphi)$ and $\Lambda^u(\psi)$ via further conjugation by the half-step-up map η . Considering intersections of stable and unstable leaves, we can promote this to a conjugacy of the actions of $\pi_1(M)$ on the orbit space \mathcal{O} . Following an argument of Ghys using Haefliger's theory of classifying spaces of foliations, Barbot [1995a, Theorem 3.4] showed using an averaging trick that such a conjugacy on the orbit space can always be realized by a homeomorphism of M giving an orbit equivalence of the flows. In our case, this homeomorphism is easily seen to be isotopic to the identity by considering the action on π_1 . This concludes the proof in the transversely orientable case.

General case For the general case, consider again the action of $\pi_1(M)$ on $\Lambda^s(\varphi) \cong \mathbb{R}$ and on $\Lambda^s(\psi)$. Let $G \subset \pi_1(M)$ be the normal subgroup generated by all squares of elements. Since each element of G is a product of squares, its action on $\Lambda^s(\varphi)$ and $\Lambda^s(\psi)$ is by orientation-preserving homeomorphisms. The proof of [Barbot 1995a, théorème 2.5] shows directly that the action of G on $\Lambda^s(\varphi)$ and $\Lambda^s(\psi)$ is also minimal. Thus, we may apply Theorem 1.1 and conclude that the actions of G on the respective leaf spaces are conjugate. We wish to show that this conjugacy extends to a conjugacy of the actions of $\pi_1(M)$.

Apply a conjugacy so that the actions of G on $\Lambda^s(\varphi)$ and $\Lambda^s(\psi)$ agree. Now our goal is to show that, after possibly further conjugating by an integer translation (which commutes with the action of G), the actions of $\pi_1(M)$ agree. Let ρ_φ and ρ_ψ denote the actions on $\Lambda^s(\varphi)$ and $\Lambda^s(\psi)$, respectively, assumed to agree on the restrictions to G .

Note that, if $\gamma \in \pi_1(M)$, and $x \in \mathbb{R}$ is an attracting fixed point of $\rho_\varphi(g)$ for some element $g \in G$, then $\rho_\varphi(\gamma)(x)$ is an attracting fixed point of $\rho_\varphi(\gamma g \gamma^{-1})$, where we have $\gamma g \gamma^{-1} \in G$. The same applies to $\rho_\psi(\gamma)$. The set of all attracting fixed points for elements of G is dense, and each element with fixed points has a \mathbb{Z} -invariant set of attracting fixed points with exactly one in $[0, 1)$.

First we verify that $\rho_\varphi(\gamma)$ preserves orientation if and only if $\rho_\psi(\gamma)$ does. Suppose $\rho_\varphi(\gamma)$ reverses orientation. Let g_1, g_2 and g_3 in G be elements with attracting fixed points satisfying

$$\rho_\varphi(g_1)_+ < \rho_\varphi(g_2)_+ < \rho_\varphi(g_3)_+ < \rho_\varphi(g_1)_+ + 1.$$

Since $\rho_\varphi(\gamma)$ reverses orientation, we have

$$\rho_\varphi(\gamma g_1 \gamma^{-1})_+ > \rho_\varphi(\gamma g_2 \gamma^{-1})_+ > \rho_\varphi(\gamma g_3 \gamma^{-1})_+ > \rho_\varphi(\gamma g_1 \gamma^{-1})_+ - 1$$

whereas, if $\rho_\varphi(\gamma)$ preserved the orientation, the original order would not be affected by conjugation. Since $\gamma g \gamma^{-1} \in G$, this ordering holds also for the action under ρ_ψ , showing that $\rho_\psi(\gamma)$ necessarily reverses orientation as well. The situation being symmetric, we have proved the claimed “if and only if” statement.

Now consider the subgroup P of elements of $\pi_1(M)$ whose action preserves orientation on $\Lambda^s(\varphi)$, or, as we have just shown, equivalently preserves orientation of $\Lambda^s(\psi)$. Our description above also implies that, for any $\gamma \in P$, we have $\rho_\varphi(\gamma) = \rho_\psi(\gamma) \circ T_\gamma$ for some integer translation T_γ . Since orientation-preserving elements commute with integer translation, the map $\gamma \in P \mapsto T_\gamma$ is a group homomorphism. Now, when $\gamma \in G$, T_γ is the identity; thus, T_γ is the identity for any $\gamma \in P$. So, in particular, the actions of ρ_φ and ρ_ψ are identical on P .

If instead we consider γ an element reversing the orientation on the leaf spaces, then $\rho_\psi(\gamma)$ and $\rho_\varphi(\gamma)$ each have a unique fixed point. Their action being determined modulo integer translations means that there exists an integer translation T'_γ such that $\rho_\varphi(\gamma) = T'_\gamma \rho_\psi(\gamma) T'^{-1}_\gamma$. Fix some orientation-reversing element γ_0 . Up to conjugating the action of ρ_ψ by an integer translation, we can assume that T'_{γ_0} is the identity, ie that $\rho_\psi(\gamma_0) = \rho_\varphi(\gamma_0)$. Notice that this conjugation does not affect the fact that ρ_φ and ρ_ψ are identical on P , since the action of elements in P commutes with integer translations. We wish to show now that $T'_\gamma = 0$ for all $\gamma \in \pi_1(M) \setminus P$, so the actions agree.

Let γ be any orientation-reversing element. Then $\gamma_0 \gamma$ preserves the orientation so $\rho_\psi(\gamma_0 \gamma) = \rho_\varphi(\gamma_0 \gamma)$. As $\rho_\psi(\gamma_0) = \rho_\varphi(\gamma_0)$, we deduce directly that $\rho_\psi(\gamma) = \rho_\varphi(\gamma)$. Hence, we proved that the actions ρ_φ and ρ_ψ are the same, ending the proof of Theorem 1.2.

3 Classifying self-orbit equivalences

This section gives the proof of Theorem 1.4. We start by introducing some additional necessary background material on the orbit space of the flow.

Returning to the picture from Section 2.1, recall that the orbit space of a skew Anosov flow is homeomorphic to a diagonal strip in \mathbb{R}^2 foliated by Λ^s and Λ^u in the two coordinate directions. For each orbit $o \in \mathbb{C}$, the ideal quadrilateral in \mathbb{C} with corners o and $\eta(o)$ and sides the stable and unstable (half-)leaves of o and $\eta(o)$ is called a *lozenge*. The union of lozenges associated with the orbits $\eta^n(o)$ for $n \in \mathbb{Z}$ is called a *string of lozenges*. The reader may consult [Barthelmé and Fenley 2017, Section 2] for more background about lozenges in general Anosov flows.

Recall [Barbot 1995b] that an (immersed) incompressible torus T is *quasitransverse* to an Anosov flow φ if it is transverse everywhere except along finitely many periodic orbits of φ . Barbot [1995b, théorème C] (see also [Barbot and Fenley 2013, Theorem 6.10]) showed that any incompressible embedded torus in M can be isotoped to a quasitransverse torus unless it is the boundary of a tubular neighborhood of an embedded one-sided Klein bottle. Such embeddings of a Klein bottle do not actually arise if the flow is




Figure 2: The orbit space \mathbb{O} with (part of) a string of lozenges.

transversally orientable and \mathbb{R} -covered. We were not able to locate a proof of this fact in the literature, so add it as a lemma below. We thank Sergio Fenley for discussing this fact with us.

Lemma 3.1 *Let φ be a transversally orientable skew \mathbb{R} -covered Anosov flow on M . Then M does not contain a π_1 -injective immersion of the Klein bottle.*

Proof Suppose there is a subgroup G of $\pi_1(M)$ isomorphic to the fundamental group of the Klein bottle, ie we have nontrivial elements $a, b \in \pi_1(M)$ such that $aba^{-1} = b^{-1}$ and $G = \langle a, b \rangle$. Since φ is transversely orientable, G acts faithfully on $\Lambda^s(\varphi) \simeq \mathbb{R}$ by orientation-preserving homeomorphisms commuting with integer translations. Consider the induced action of G on \mathbb{R}/η^2 . For this induced action, both a and b have a single attracting and single repelling fixed point. Since $aba^{-1} = b^{-1}$, we must have that a interchanges the attracting and repelling points for b . But this gives a an orbit of period two, contradicting that it acts with fixed points. \square

The main result that we need is the following:

Lemma 3.2 [Barbot 1995b] *If T is a quasitransverse torus in M for a skew Anosov flow, then each lift \tilde{T} of T to the universal cover projects to a unique string of lozenges \mathcal{C} in \mathbb{O} . That string of lozenges is the unique string of lozenges left invariant by the \mathbb{Z}^2 -subgroup of $\pi_1(M)$ that fixes \tilde{T} . The interior of each lozenge in the string corresponds to a maximal annulus in T transverse to the flow; its corners are the periodic orbits bounding the annulus.*

Conversely, if G is a subgroup of $\pi_1(M)$ isomorphic to \mathbb{Z}^2 , then there exists a unique string of lozenges in \mathbb{O} fixed by G . Moreover that string of lozenge is the projection of an (a priori only immersed) torus T .

Remark 3.3 We have stated this lemma in the special case of skew Anosov flows. For general Anosov flows on 3-manifolds, one needs to replace “string of lozenges” by “chain of lozenges” — see [Barthelmé and Fenley 2017, Section 2] — and some \mathbb{Z}^2 -subgroups may fix two distinct chains of lozenges.

We also need the following easy proposition, giving another condition for an element $\gamma \in \pi_1(M)$ to be represented by a periodic orbit of a skew Anosov flow φ . Recall that $\Lambda^s \cong \mathbb{R}$ denotes the leaf space of the weak stable foliation of the lifted flow $\tilde{\varphi}$. We denote the natural projection by $\pi^s: \tilde{M} \rightarrow \Lambda^s$.

Proposition 3.4 *Let φ be a skew Anosov flow on a 3-manifold M . Let $\gamma \in \pi_1(M)$. The following are equivalent:*

- (i) *The element γ is represented by a periodic orbit of φ ;*
- (ii) *There exists a γ -invariant curve $c \subset \tilde{M}$ such that $\pi^s(c)$ is bounded (either above or below) in Λ^s ;*
- (iii) *For any γ -invariant curve $c \subset \tilde{M}$, its image $\pi^s(c)$ is bounded in Λ^s .*

Proof Item (iii) trivially implies (ii), we will show that (ii) implies (i) and that (i) implies (iii).

(ii) \implies (i) Assume that there exists a γ -invariant curve $c \subset \tilde{M}$ such that $\pi^s(c)$ is bounded above in Λ^s , and let L^+ denote its upper bound. (The case where $\pi^s(c)$ is bounded below is analogous.) If the action of γ reverses the orientation of Λ^s , then it fixes a leaf which necessarily contains a periodic orbit represented by γ , and we are done. If γ instead preserves the orientation, then the upper bound $L^+ \in \Lambda^s$ is a γ -invariant point since $\pi^s(c)$ is γ -invariant, and thus the leaf corresponding to this point contains a unique γ -invariant orbit. This shows (i).

(i) \implies (iii) Suppose γ is represented by a periodic orbit of the flow and $c \subset \tilde{M}$ is a γ -invariant curve. Supposing first that γ preserves orientation on \mathbb{R} , the action has an attracting fixed point $x \in \Lambda^s$. Moreover, the images of this point under powers of the half-step-up map are alternately attracting and repelling fixed points of γ ; if i is even, then $\eta^i(x)$ is an attracting fixed point, if i is odd, then $\eta^i(x)$ is a repelling fixed point, as described in Section 2.1. In the case where γ reverses orientation, the same argument above applies if we replace γ with γ^2 , which preserves orientation.

In either case, since the action of γ^2 on c is free, we may take a compact fundamental domain I for the action. The projection $\pi^s(I)$ is contained in some bounded interval $[\eta^{-k}(x), \eta^k(x)]$. Therefore, $\pi^s(c) = \bigcup_{n \in \mathbb{Z}} \gamma^{2n} \cdot \pi^s(I) \subset [\eta^{-k}(x), \eta^k(x)]$, so it is bounded, proving (iii). \square

Combining Theorem 1.2 with Proposition 3.4 gives the following as a direct consequence:

Theorem 3.5 *Let φ and ψ be two skew \mathbb{R} -covered Anosov flows on a 3-manifold M . The flows φ and ψ are isotopically equivalent if and only if, for any periodic orbit α of ψ (resp. φ) with lift $\tilde{\alpha} \subset \tilde{M}$, the projection $\pi_1^s(\tilde{\alpha}) \subset \Lambda^s(\varphi)$ (resp. $\pi_1^s(\tilde{\alpha}) \subset \Lambda^s(\psi)$) is bounded.*

3.1 Proof of Theorem 1.4

To set up for the proof, we begin by giving precise definitions of translation number with respect to φ , and the displacement of a curve by a Dehn twist.

Definition 3.6 (translation number) Let $\beta \in \pi_1(M)$, and consider its action on $\Lambda^s(\varphi)$. Fix $x \in \Lambda^s(\varphi)$. For each $q \in \mathbb{Z}$, there exists a unique $p_q \in \mathbb{Z}$ such that $\tau^{p_q}(x) \leq \beta^q(x) < \tau^{p_q+1}(x)$. We define

$$t_\varphi(\beta) := \lim_{q \rightarrow \infty} \frac{p_q}{q}.$$

Since τ and β commute, it is a standard exercise to show that this limit exists and is in fact independent of the choice of x ; indeed, $t_\varphi(\beta)$ is simply the classical *translation number* for the action of β on Λ_s with respect to any parametrization of Λ_s where τ acts as translation by 1.

Remark 3.7 We will consider below only the case when β is freely homotopic to a curve in a quasitransverse torus T . Thus, β fixes a (unique) string of lozenges in the orbit space. In this case we therefore have $t_\varphi(\beta) = k \in \mathbb{Z}$, where k is such that, if x is any of the corner of the string of lozenges, then $\beta x = \tau^k(x)$.

If a skew Anosov flow φ on M is transversally oriented, we saw (Lemma 3.1) that M cannot admit an incompressible embedding of the Klein bottle; thus, by [Barbot 1995b, théorème C], any incompressible embedded torus in M can be put in quasitransverse position with respect to the flow. We further have, thanks to [Barbot 1995b, théorème E], that any collection of pairwise disjoint, nonisotopic incompressible embedded tori can be simultaneously isotoped to a collection of still disjoint (and obviously still nonisotopic) quasitransverse tori. It is thus no loss of generality to adopt the following convention for transversely orientable flows:

Convention For the remainder of this section, we restrict our attention to transversely orientable flows, and we will always assume that the tori we consider are in quasitransverse position.

Next we will define the “displacement” of an orbit by a Dehn twist. We first recall the definition of Dehn twists to emphasize that they come with a specification of a transverse orientation on the torus.

Definition 3.8 Let T be an embedded torus, and $\beta \in \pi_1(T)$. A *Dehn twist along β* is the mapping class of a map D_β defined as follows: Take a small product region $T \times [-1, 1] \in M$, and fix a basis $\{\alpha, \nu\}$ for $\pi_1(T)$ giving an identification of T with $S^1 \times S^1 = \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$, where $\beta = \alpha^p \nu^q$ for some p and q . For $(x, y, z) \in T \times [-1, 1]$, we define $D_\beta(x, y, t) = (x + ph(t), y + qh(t), t)$, where $h: [-1, 1] \rightarrow [0, 1]$ is a smooth bump function with $h(-1) = 0$ and $h(1) = 1$, and extend D_β to be the identity elsewhere on M .

Remark 3.9 Reversing the transverse orientation of T and applying the same construction results in a map isotopic to the *inverse* of that defined above. Thus, the notation D_β , while standard, is somewhat misleading because the mapping class does not depend on β alone. There is no intrinsic way to distinguish D_β from D_β^{-1} . Thus, by convention, we say that a map D_β *comes with* the data of a choice of transverse orientation. When we speak of a Dehn twist supported on a torus neighborhood $T \times [-1, 1]$, we always assume the orientation is as given by the interval $[-1, 1]$.

It will be convenient for us to choose homeomorphisms representing Dehn twists which are in a particularly nice form with respect to the flow, as follows:

Convention 3.10 (good Dehn twist coordinates) Given a quasitransverse torus T , we choose the coordinates $(x, y, z) \in T \times [-1, 1]$ in the following way: Let $\alpha_1, \dots, \alpha_{2n}$ be the periodic orbits of φ on T . Then we assume that the local stable leaves of the orbits α_i in $T \times [-1, 1]$ are given by the equation $x = i/2n$.

With this convention, a Dehn twist D_α on T , where $\alpha \in \pi_1(T)$ represents any power of the periodic orbits, will preserve the local stable leaves of the periodic orbits α_i . More generally, given any Dehn twist D_β on T , and any segment $c(t) := (x_0, y_0, t)$ for $t \in [-1, 1]$ through $T \times [-1, 1]$, the number of times its image $D_\beta(c(t))$ intersects the union of stable leaves of the α_i is the minimal intersection number of β with α_i .

Definition 3.11 (sign of an intersection) Given a Dehn twist D_β supported on $T \times [-1, 1]$ as above, and an orbit φ_t intersecting T transversely at $t = 0$, we say this intersection is *positive* if $\varphi^t(z) \in T \times [0, 1]$ for small positive t , and *negative* if $\varphi^t(z) \in T \times [-1, 0]$ for small positive t .

Before giving the formal definition of the displacement $d_\varphi(c, f)$ of a closed orbit c under a product of Dehn twists f , we motivate this with the following lemma, which describes how a Dehn twist on a torus T affects a segment of a periodic orbit transverse to T , from the perspective of the leaf space of φ .

For the statement, we fix a quasitransverse torus T in M , a point z with orbit $\varphi^t(z)$ transverse to T , and a small product neighborhood $T \times [-1, 1]$ so that the orbit $\varphi^t(z) \cap T \times [-1, 1]$ is, locally near $t = 0$, a segment J between some point $z_- \in T \times \{-1\}$ and $z_+ \in T \times \{1\}$. Let D_β be a Dehn twist supported on $T \times [-1, 1]$.

Lemma 3.12 Let $\tilde{T} \times [-1, 1]$ be a lift of $T \times [-1, 1]$ to \tilde{M} , let \tilde{J} be the lift of J in $\tilde{T} \times [-1, 1]$ with endpoints \tilde{z}_- and \tilde{z}_+ , and let \tilde{D}_β be the lift of D_β fixing \tilde{z}_- . Finally, let $\{L_i\}_{i \in \mathbb{Z}}$ be the string of lozenges associated with \tilde{T} .

- (1) If \tilde{J} projects to a point in the lozenge L_i , then $\tilde{D}_\beta(\tilde{z}_+)$ projects into L_k , where $k = i + 2t_\varphi(\beta)$ if the intersection of J and T is positive, and $k = i - 2t_\varphi(\beta)$ if the intersection is negative.
- (2) The stable saturation of $\tilde{D}_\beta(\tilde{J})$ to the orbit space stays inside the stable saturation of the lozenges between L_i and L_k .

Proof Without loss of generality, we assume that \tilde{J} projects to a point in L_0 , and that the sign of the intersection is positive (the negative case is analogous). Recall that the lozenges L_i are the projections of strips A_i inside \tilde{T} , bounded by periodic orbits of $\tilde{\varphi}$.

By definition of D_β , the image $\tilde{D}_\beta(\tilde{z}_+)$ projects in the orbit space to the lozenge $\beta(L_0)$. Now, if x_i are the corners of the lozenge L_i (enumerated so that $x_i < x_{i+1}$ in $\Lambda_s(\varphi)$), then $\eta^k(x_0) = x_{2k}$. Thus, by definition of translation number (and Remark 3.7), $\beta(L_0) = L_{2t_\varphi(\beta)}$, proving the first part of the lemma.

The second statement follows immediately from Convention 3.10 and the structure of lozenges (Lemma 3.2). \square




Figure 3: The configuration on the right is impossible if the shaded lozenge is L'_j .

Definition 3.13 (displacement) Let c be a periodic orbit of the skew flow φ and D_β a Dehn twist on a quasitransverse torus T . Let x_1, \dots, x_n be the intersection points of c with T , and let $\epsilon_i = \{\pm 1\}$ be the sign of the intersection at x_i . The *displacement* of c by D_β is defined by

$$d_\varphi(c, D_\beta) = \sum_{i=1}^n 2\epsilon_i t_\varphi(\beta).$$

If $f = D_{\beta_1} \circ \dots \circ D_{\beta_k}$ is a composition of Dehn twists on *pairwise disjoint* quasitransverse tori, then we set

$$d_\varphi(c, f) = \sum_{i=1}^k d_\varphi(c, D_{\beta_i}).$$

We will need the following observation for the proof of Theorem 1.4:

Observation 3.14 Let \tilde{T} and \tilde{T}' be two **disjoint** lifts of (the same or distinct) quasitransverse tori. Let \tilde{J} be a segment of a periodic orbit that first crosses \tilde{T} and then crosses \tilde{T}' . Suppose \tilde{T} projects to a string of lozenges $\{L_i\}$, and \tilde{T}' to a string $\{L'_j\}$. Let i and j be such that \tilde{J} is contained in $L_i \cap L'_j$.

Then L'_j is contained in the saturation of L_i by stable leaves; equivalently, L_i is contained in the saturation of L'_j by unstable leaves.

Proof Since the lifts \tilde{T} and \tilde{T}' are disjoint, observe that either $L_i \subset \tilde{\mathcal{F}}^s(L'_j)$ and $L'_j \subset \tilde{\mathcal{F}}^u(L_i)$, or $L_i \subset \tilde{\mathcal{F}}^u(L'_j)$ and $L'_j \subset \tilde{\mathcal{F}}^s(L_i)$. Both configurations are shown in Figure 3.

We will show that the second case cannot occur because orbits converge in the positive direction of the flow along \tilde{J} . Precisely, consider the strong unstable leaf through the starting point \tilde{x} of the orbit segment \tilde{J} . Denote this leaf by λ^u , and let $l_i \subset \lambda^u$ denote the set of points whose forward orbit intersects the strip A_i in \tilde{T} corresponding to L_i . Similarly, define $l'_j \subset \lambda^u$ to be the set of points whose forward

orbit intersects the strip A'_j in \tilde{T}' corresponding to L'_j . Hyperbolicity of the flow means that $l'_j \subset l_i$. Hence, $L'_j \subset \tilde{\mathcal{F}}^s(L_i)$, and equality happens if and only if $L'_j = L_i$, which in turn implies that the orbits bounding A_i and A'_j are the same, which contradicts the assumption that \tilde{T} and \tilde{T}' are quasitransverse and disjoint. \square

We next reduce the proof of Theorem 1.4 to the following proposition:

Proposition 3.15 *Suppose $f = D_{\beta_1} \circ \cdots \circ D_{\beta_n}$ is a composition of Dehn twists on pairwise disjoint quasitransverse tori T_1, \dots, T_n , and let $\tilde{c} \subset \tilde{M}$ be a lift of a periodic orbit c of φ . Then $d_\varphi(c, f) = 0$ if and only if, for some lift (equivalently, for all lifts) \tilde{f} of f , the image $\tilde{f}(\tilde{c})$ has bounded projection to $\Lambda^s(\varphi)$.*

Proof of Theorem 1.4 given Proposition 3.15 Indeed, this is an easy consequence of Proposition 3.4 and Theorem 1.2. Theorem 1.2 states that f is a self-orbit equivalence if and only if $\mathcal{P}(\varphi) = f_*\mathcal{P}(\varphi) = \mathcal{P}(f\varphi f^{-1})$. By Proposition 3.4, $\mathcal{P}(\varphi)$ can be characterized as the set of $\gamma \in \pi_1(M)$ such that every γ -invariant curve $\tilde{c} \subset \tilde{M}$ has bounded projection to $\Lambda^s(\varphi)$. Thus, f is isotopic to a self-orbit equivalence if and only if f_* preserves this set. Proposition 3.15 characterizes this set in terms of the vanishing of $d_\varphi(c, f)$. \square

Proof of Proposition 3.15 Fix $f = D_{\beta_1} \circ \cdots \circ D_{\beta_k}$, where D_{β_i} is a Dehn twist on T_i , and let \tilde{f} be a lift of f to \tilde{M} chosen so that \tilde{f} fixes some point \tilde{x} on \tilde{c} . Recall that we assumed, without loss of generality, that all the T_i are quasitransverse.

Let $\gamma \in \pi_1(M)$ represent a periodic orbit c of φ . As a first trivial case, suppose c has no transverse intersections with any torus T_i , so it is either contained in a single torus or disjoint from all of them. Let \tilde{c} be a lift of c . Note that c is isotopic to a curve c' disjoint from the support of f ; lifting this isotopy means the lift \tilde{c} is isotopic to a lift \tilde{c}' such that, after applying some deck transformation g of the cover, this is disjoint from the support of \tilde{f} . Thus, $\tilde{f}(g\tilde{c})$ is uniformly bounded distance away from $\tilde{f}(g\tilde{c}') = g\tilde{c}'$, and this is uniformly bounded distance away from $g\tilde{c}$, showing that $g\tilde{c}$ (and hence also \tilde{c}) has bounded projection.

For the case where c intersects some T_i transversely, we may without loss of generality choose the product neighborhoods of the tori T_i in the definition of D_{β_i} small enough that, every time c crosses a torus T_i , it enters one side $T_i \times \{\pm 1\}$ and leaves the other, $T_i \times \{\mp 1\}$. Consider the positive-time ray $r = \{\varphi^t(\tilde{x}) : t \geq 0\} \subset \tilde{c}$, where \tilde{x} is as before a point fixed by \tilde{f} . We will show that the projection of $\tilde{f}(r)$ to $\Lambda^s(\varphi)$ is bounded. Reversing the argument (using unstable leaves instead of stable) will show that the projection of $\tilde{f}(\tilde{c})$ to $\Lambda^s(\varphi)$ is also bounded, so we do only the forward case.

Between \tilde{x} and $\gamma\tilde{x}$, the ray r intersects a finite number of lifts of the tori T_i on which the Dehn twists D_{β_i} are supported. Let $\mathcal{T}_1, \dots, \mathcal{T}_n$ denote these lifts, indexed along the path of r so that r first intersects \mathcal{T}_1

after \tilde{x} . Note that two distinct lifts \mathcal{T}_i and \mathcal{T}_j may project to the same torus in M ; this will happen whenever the orbit c crosses the same torus twice. Since the T_i are quasitransverse tori, each \mathcal{T}_j projects to a string of lozenges \mathcal{C}_j . Let $L_0^{(1)}$ be the lozenge in \mathcal{C}_1 containing the projection of \tilde{c} . The main technical part of the proof is the following claim:

Claim 3.16 *Let r_0 denote the segment of r between \tilde{x} and $\gamma\tilde{x}$.*

- (1) *The stable leaf of $\tilde{f}(\gamma\tilde{x})$ intersects the lozenge $\eta^{d_\varphi(c,f)}(L_0^{(1)})$.*
- (2) *There exists $N > 0$, depending only on f , such that the stable leaf saturation of $\tilde{f}(r_0)$ intersects the chain of lozenges \mathcal{C}_1 only between $\eta^{-N}(L_0^{(1)})$ and $\eta^N(L_0^{(1)})$.*

Given this claim, the proof of Proposition 3.15 can be finished quickly, by considering the positive iterates of r_0 under γ . Let us assume the claim for the moment and use it to derive the conclusion of Proposition 3.15.

We use the fact that $r = \bigcup_{i=1}^\infty \gamma^i(r_0)$. For the first direction, suppose that $d_\varphi(c, f) = 0$. Fix a segment $r_i = \gamma^i(r_0)$. Then $\gamma^{-i}(r_i) = r_0$, so, by Claim 3.16, the stable leaf of $\tilde{f}(r_i) = f_*(\gamma^i)\tilde{f}(r_0)$ intersects the lozenge $f_*(\gamma^i)(L_0^{(1)})$, and $\tilde{f}(r_i)$ intersects the chain of lozenges \mathcal{C}_1 only between $f_*(\gamma^i)\eta^{-N}(L_0^{(1)})$ and $f_*(\gamma^i)\eta^N(L_0^{(1)})$. But Observation 3.14 implies that $f_*(\gamma^i)(L_0^{(1)})$ is contained in the stable saturation of $L_0^{(1)}$; thus, $\tilde{f}(r_i)$ is contained in the union of the stable saturation of leaves between $\eta^{-N}(L_0^{(1)})$ and $\eta^N(L_0^{(1)})$ for all i ; hence, r has bounded projection to $\Lambda^s(\varphi)$. As remarked above, applying the same argument to unstable leaves and using the negative time ray shows that c has bounded projection.

Conversely, if $\iota = i_\varphi(c, f) \neq 0$, then the argument above shows that $\tilde{\mathcal{F}}^s(\tilde{f}(\gamma^n\tilde{x}))$ intersects $\eta^{n\iota}(L_1^{(1)})$; thus, the projection of $\tilde{f}(r)$ to $\Lambda^s(\varphi)$ is unbounded. □

So, to finish the proof of the proposition, we only need to prove Claim 3.16.

Proof of Claim 3.16 Recall that f has support in disjoint small neighborhoods of the quasitransverse tori in M which lift to disjoint neighborhoods of \mathcal{T}_j in \tilde{M} , and we use $L_k^{(j)}$ to denote the string of lozenges associated to \mathcal{T}_j . Let \tilde{N}_j denote the lifted neighborhood containing \mathcal{T}_j . We use these neighborhoods to split the ray r_0 into a union of intervals I_j and J_j , where $J_j := \tilde{N}_j \cap r_0$, and I_j denotes the connected component of r_0 between \tilde{N}_{j-1} and \tilde{N}_j . The interval I_0 is the segment of r_0 from \tilde{x} to \tilde{N}_1 .

For each \mathcal{T}_i , let t_i denote the translation number of the Dehn twist on the corresponding torus. Notice that, since f is supported on the union of the projection of the neighborhoods \tilde{N}_i to M , it follows that, for any j , there exists $h_j \in \pi_1(M)$ such that $\tilde{f}(I_j) = h_j(I_j) \subset h_j\tilde{c}$. Our choice of lift implies that h_0 is the identity.

To demonstrate the first statement of the claim, we will first establish the fact that, if $\tilde{f}(I_{i-1}) = h_{i-1}(I_{i-1})$ intersects a lozenge $h_{i-1}L_k^{(i)}$, then the stable leaf of $\tilde{f}(I_i)$ intersects the translate of $h_{i-1}L_k^{(i)}$ by $\eta^{\epsilon_i 2t_i}$.




Figure 4: The action of the lift of a Dehn twist on a ray, shown in \tilde{M} on the left and schematically indicating intersections with lifted neighborhoods of tori on the right.

Here is the proof of the fact: Fix some i . The segment $\tilde{f}(J_i)$ is obtained from $h_{i-1}(J_i)$ by applying the (unique) lift of the Dehn twist supported on the projection of \mathcal{T}_i that preserves $h_{i-1}\mathcal{T}_i$ and fixes $h_{i-1}(I_{i-1})$. The (projection to the orbit space of the) endpoint shared by $\tilde{f}(I_{i-1}) = h_{i-1}(I_{i-1})$ and $\tilde{f}(J_i)$ lies in some lozenge $L_k^{(i-1)}$ associated to $h_{i-2}(\mathcal{T}_{i-1})$. By Observation 3.14, the lozenge $h_{i-1}(L_{k'}^{(i)})$ for $h_{i-1}(\mathcal{T}_i)$ that also contains this point is such that $h_{i-1}(L_{k'}^{(i)})$ is contained in the saturation by stable leaves of $h_{i-2}(L_k^{(i-1)})$. Applying Lemma 3.12, we conclude that after applying the lifted Dehn twist to $h_{i-1}(J_i)$ (see Figure 4), the image of its other endpoint, which is also an endpoint of $\tilde{f}(I_i)$, lies in $\eta^{\epsilon_i 2t_i} h_{i-1}(L_{k'}^{(i)})$. This is contained in the saturation by stable leaves of $\eta^{\epsilon_i 2t_i} h_{i-2}(L_k^{(i-1)})$, which proves the fact.

To deduce (1) using the fact, we apply it iteratively, observing first that the stable leaf of $\tilde{f}(I_1)$ intersects $\eta^{\epsilon_1 2t_1}(L_0^{(1)})$ and some lozenge $h_1(L_k^{(2)})$, and the stable saturation of $L_k^{(2)}$ is contained in $\eta^{\epsilon_1 2t_1} L_0^{(1)}$. Thus, $\tilde{f}(I_2)$ has stable leaf intersecting $\eta^{\epsilon_2 2t_2} h_1(L_k^{(2)})$, and thus $\eta^{\epsilon_1 2t_1} \eta^{\epsilon_2 2t_2}(L_0^{(1)})$. Continuing iteratively, we conclude that $\tilde{f}(\gamma\tilde{x})$ has stable leaf intersecting the translate of $L_0^{(1)}$ by $\eta^{d_\varphi(c,f)}$.

Now the second part of the claim follows directly from the application of the second part of Lemma 3.12 in the proof above by choosing

$$N = \sum_{i=1}^k 2|t_i|. \quad \square$$

3.2 Application: classification of self-orbit equivalences in special cases

We will now use the criterion given by Theorem 1.4 to describe the isotopy class of self-orbit equivalences in some special cases, starting with the proof of Corollaries 1.5 and 1.6.

Proof of Corollary 1.5 Suppose that D_α is a Dehn twist on a torus T in a direction α that is represented by a periodic orbit of φ . Then the translation number of α is zero, and thus, for any periodic orbit c of φ ,

the displacement of c by D_α is zero. Therefore, the criterion of Theorem 1.4 implies that any such Dehn twist is isotopic to a self-orbit equivalence of φ . \square

Proof of Corollary 1.6 Let T be a quasitransverse torus in M and D_β a Dehn twist on T with nonzero translation number. Suppose first that T is separating. Consider a periodic orbit c of φ . Then either c does not intersect T or it intersects T an even number of times with alternating signs. In either case, $d_\varphi(c, D_\beta) = 0$. Thus, D_β is isotopic to a self-orbit equivalence by Theorem 1.4.

Now, if we assume instead that T is nonseparating, we claim that we can find a periodic orbit c of φ such that its intersections with T are always in the same transverse direction, and therefore are assigned the same sign. To do this, take a closed oriented loop based in T in M intersecting T exactly once, transversely, at the basepoint (and therefore inducing a transverse orientation of T). Lift this loop to a path in \tilde{M} with endpoints in lifts \tilde{T}_0 and \tilde{T}_1 of T , and iteratively choose successive lifts \tilde{T}_i . Then, for each i , \tilde{T}_i separates \tilde{T}_{i-1} from \tilde{T}_{i+1} , and no lift of T separates \tilde{T}_i from \tilde{T}_{i+1} .

Fix n large, and let \tilde{d} be a segment of an orbit in \tilde{M} with endpoints in \tilde{T}_0 and \tilde{T}_n . Such an orbit always exists because the flow is \mathbb{R} -covered and skew. If n is chosen large enough, \tilde{d} projects down to an orbit segment in M that will contain points close enough to satisfy the conditions of the Anosov closing lemma. Hence, we obtain a periodic orbit c such that all its intersections with T have the same transverse orientation. We conclude that $d_\varphi(c, D_\beta)$ is nonzero, which, by the criterion, implies that D_β is not isotopic to a self-orbit equivalence. \square

The proof of this corollary gives an example of a general strategy to identify self-orbit equivalences by understanding the intersection of periodic orbits with quasitransverse tori. One could broaden this to give a characterization of self-orbit equivalences of a given flow in terms of the configuration of tori in M (the JSJ graph and geometry of the pieces) and translation numbers of periodic orbits in tori. Working through the details of this is beyond the scope of this article; instead we illustrate our criterion with some examples where we can virtually describe the group of self-orbit equivalences in the mapping class group.

For both of the following statements, we assume φ is a transversally orientable skew Anosov flow on M .

Theorem 3.17 *Suppose that the JSJ decomposition of M has a single piece which is atoroidal, glued to itself along n boundary tori T_1, \dots, T_n . Let D_{per} be the group generated by Dehn twists on the T_i in the directions of the periodic orbits of φ .*

Let D be the finite-index subgroup of the mapping class group generated by Dehn twists on tori. Then D_{per} is the set of self-orbit equivalences in D .

Theorem 3.18 *Suppose that each torus of the JSJ decomposition of M is separating. Let D be the finite-index subgroup of the mapping class generated by Dehn twists on tori. Then the set of self-orbit equivalences in D is equal to the group generated by Dehn twists in the directions of periodic orbits inside Seifert pieces, together with any Dehn twists on the JSJ tori.*

Remark 3.19 Using Handel–Thurston surgery and Foulon–Hasselblatt Dehn surgeries on periodic orbits of geodesic flows, one can easily construct many examples of skew Anosov flows on manifolds as in the statements of Theorems 3.17 and 3.18.

The proofs of both statements use the fact that the group generated by Dehn twists on tori has finite index in the mapping class group of any orientable 3-manifold. (See [Johannson 1979] for an explanation and history of the proof.)

Proof of Theorem 3.17 By Corollary 1.5, any isotopy class $[f] \in D_{\text{per}}$ is represented by a self-orbit equivalence. For the other containment, suppose that f is a self-orbit equivalence contained in D . We assume for a contradiction that f cannot be written as a product of elements in D_{per} . Since elements of D_{per} are self-orbit equivalences, up to composition with such a map we can assume that $f = D_{\beta_1} \circ \cdots \circ D_{\beta_k}$, where $t_\varphi(\beta_i) \neq 0$ for all i . In fact, we will only use that one of these has nonzero translation number.

We now produce a periodic c such that $d(c, f) \neq 0$, leading to a contradiction by Theorem 1.4. Reindexing if needed, suppose T_1 is a torus in the support of f , with associated Dehn twist D_{β_1} . By hypothesis, T_1 is nonseparating in $M \setminus (T_2 \cup \cdots \cup T_n)$. By considering lifts of curves as in the proof of Corollary 1.6, one can find a periodic orbit c of φ that intersects only T_1 , and each intersection point has the same orientation. Thus, $d(c, f) = d(c, D_{\beta_1}) = \pm t_\varphi(\beta_1) \neq 0$, contradicting Theorem 1.4. \square

Proof of Theorem 3.18 Given the assumption, all JSJ tori are separating, so Corollary 1.6 shows that any product of Dehn twists along them is isotopic to a self-orbit equivalence. This, together with Corollary 1.5, shows that any element of the group generated by Dehn twists in the direction of periodic orbits together with Dehn twists on the JSJ tori is isotopic to a self-orbit equivalence.

We are left to show that any element of the finite-index subgroup generated by Dehn twists that is a self-orbit equivalence is of this form. Let f be a self-orbit equivalence generated by Dehn twists; thus, f preserves each Seifert piece. Fix a Seifert piece S and let Σ denote the base orbifold of S , and g the restriction of f to Σ .

Up to composing f with some Dehn twists in the direction of periodic orbits on S and Dehn twists on the boundary of S , we can assume that g projects to the identity in the mapping class group of the base orbifold Σ (see [Johannson 1979] or [Bonatti et al. 2020, Section 4.7]). Thus, for any $\gamma \in \pi_1(S)$ that is obtained as a lift of an element of $\pi_1(\Sigma)$ that is not homotopic to a boundary component of Σ , there exists $k \in \mathbb{Z}$ such that $g_*(\gamma) = \gamma s^k$, where s is the element of $\pi_1(S)$ representing a regular fiber of the Seifert fibration of S .

For any such γ , the group $H = \langle \gamma, s \rangle$ is a subgroup of $\pi_1(M)$ homeomorphic to \mathbb{Z}^2 , and hence there exists a (unique) string of lozenges \mathcal{C} in \mathbb{C} that is preserved by H . The permutation action of H on this string gives a homomorphism $H \rightarrow \mathbb{Z}$; thus, there exists some nontrivial $\gamma' \in H$ whose action on \mathbb{C} fixes

each corner of \mathcal{C} ; equivalently, the conjugacy class of γ' is represented by the periodic orbits of φ that project to the corners of \mathcal{C} .

Since the weak foliations are transverse to the Seifert fibration, s acts as a translation on the corners of \mathcal{C} (see [Barbot and Fenley 2013]). In particular, if $k \neq 0$, then $g_*(\gamma') = \gamma' s^k$ does not fix a point of \mathcal{C} , and hence is not represented by a periodic orbit of φ . This contradicts the fact that g was the restriction of a self-orbit equivalence. Thus, $k = 0$, and, since the choice of γ was arbitrary, we conclude that g is isotopic to the identity.

Applying this argument to the restriction of f to each Seifert piece, we conclude that f is a product of Dehn twists in the direction of periodic orbits and Dehn twists on the JSJ tori, as claimed. \square

Remark 3.20 The proof of Theorem 3.18 gives a virtual characterization of the self-orbit equivalences of a given flow that fix a Seifert piece and are isotopic to identity on its boundary: they are exactly the isotopy classes of the group generated by Dehn twists in the direction of periodic orbits in the Seifert piece.

4 Application to contact flows

We will now apply some results of contact geometry to obtain the reverse direction of Theorem 1.12, as well as its corollaries. Recall that a (coorientable) contact structure ξ is *Anosov* if ξ admits an Anosov Reeb flow, ie there exists a 1-form α such that $\xi = \ker \alpha$ and the Reeb flow of α is Anosov.

We will use as a black box the theory of cylindrical contact homology. Introduced in a more general context by Eliashberg, Givental and Hofer [Eliashberg et al. 2000], it has been proven to be well defined for dynamically convex contact structures by Hutchings and Nelson [2016]. This context includes the case of Anosov contact forms: if the Reeb flow of a contact 1-form α is Anosov, then it is automatically nondegenerate and, since Anosov flows have no contractible orbits, α is dynamically convex (see also eg [Macarini and Paternain 2012]). Some of the fundamental results in this theory can be summarized as follows:

Theorem 4.1 *If α_1 and α_2 are nondegenerate dynamically convex contact forms on a closed, hypertight, contact 3-manifold (M, ξ) , then $C\mathbb{H}_{\text{cyl}}^\Lambda(\alpha_1) \cong C\mathbb{H}_{\text{cyl}}^\Lambda(\alpha_2)$ for any set Λ of free homotopy classes in M . The space $C\mathbb{H}_{\text{cyl}}^\Lambda(\alpha_i)$ is a \mathbb{Q} -vector space, it is the homology of a complex generated by the periodic orbits of the Reeb flow of α_i in the free homotopy classes belonging to Λ .*

Having the cylindrical contact homology groups associated to α_1 and α_2 being isomorphic is not quite enough for our purpose: what we will need in order to apply Theorem 1.2 is to obtain that the chain complexes themselves are equal, when α_1 and α_2 are Anosov contact forms. This follows from the fact that the differential in the chain complex is trivial. That result was proven by Macarini and Paternain [2012, Theorem 2.1] for any Anosov contact structure (and independently by Vaugon — see [Foulon et al. 2021] — with the additional assumption that the foliations are orientable).

Proof of Theorem 1.12, reverse implication Let φ_1 and φ_2 be two contact Anosov flows with respective contact forms α_1 and α_2 and contact structures $\xi_1 = \ker \alpha_1$ and $\xi_2 = \ker \alpha_2$. Suppose that ξ_1 and ξ_2 are contactomorphic. Then there exists a diffeomorphism $g: M \rightarrow M$ such that $g_*\xi_1 = \xi_2$.

In our situation, the 1-forms $g^*\alpha_1$ and α_2 are two contact forms of (M, ξ_2) with Anosov Reeb flows. Applying the theorem on cylindrical contact homology above, where Λ runs over all possible free homotopy classes in $\pi_1(M)$, we deduce that each homotopy class represented by a periodic orbit of the Reeb flow of $g^*\alpha_1$ is also represented by a periodic orbit of the Reeb flow φ_2 .

The Reeb flow of $g^*\alpha_1$ is $g^{-1} \circ \varphi_1^t \circ g$. Thus, by Theorem 1.2, $g^{-1} \circ \varphi_1 \circ g$ and φ_2 are isotopically equivalent. Equivalently, φ_1 and φ_2 are orbit equivalent. If additionally g is isotopic to the identity, we conclude that φ_1 and φ_2 are isotopically equivalent. \square

Using this direction of Theorem 1.12, together with the coarse classification of contact structures in dimension 3 of Colin–Giroux–Honda, we can quickly deduce Theorem 1.13:

Proof of Theorem 1.13 By [Colin et al. 2009, théorème 6], on any irreducible 3-manifold, there exist at most finitely many noncontactomorphic contact structures with bounded Giroux torsion. Now Proposition A.1 shows that the contact structure of any contact Anosov flow has zero Giroux torsion. Thus, Theorem 1.12 implies the result.⁴ \square

Appendix Further applications to contact topology joint with Jonathan Bowden

In this appendix, we show that any Anosov contact structure has zero Giroux torsion, prove the converse to Theorem 1.12, and then use this to obtain new results about contact structures.

Proposition A.1 *Let ξ be an Anosov contact structure on a 3-manifold M . Then ξ has zero Giroux torsion. In fact, a double cover of M will be strongly symplectically fillable.*

Proof Let α be a contact 1-form such that $\ker \alpha = \xi$ and such that the Reeb flow φ of α is Anosov. Let X be the Reeb vector field. Consider the 4-manifold $M \times [-1, 1]$ with symplectic form $\omega = d(e^t \alpha)$, where t is the coordinate on $[-1, 1]$. Then $\xi = \ker \alpha$ is a contact structure that we can assume without loss of generality to be positive and, since $\omega|_{M \times \{1\}} = e^1 d\alpha$, the form ω is nonzero on ξ .

Up to taking a double cover, we can assume that φ is transversally orientable, ie its Anosov splitting is orientable.

Let X^{ss} and X^{uu} be vector fields in, respectively, the stable and unstable direction of the Anosov splitting of X , with orientations chosen so that the plane spanned by $X^{\text{ss}} + X^{\text{uu}}$ and X defines a (coorientable) contact structure ξ_- with *negative* orientation (see [Mitsumatsu 1995]). Put this contact structure on $M \times \{-1\}$.

⁴When M is hyperbolic, one does not need Bowden's result, thanks to [Colin et al. 2009, théorème 2].

Since the defining property of a contact form is open in the C^1 topology, we can take a C^1 -small approximation $\tilde{\xi}_-$ of ξ_- such that X is transverse to $\tilde{\xi}_-$. Then ω is nonzero on $\tilde{\xi}_-$ (since $\omega|_{M \times \{-1\}} = e^{-1} d\alpha$, so its kernel is spanned by X).

This gives us a weak semifilling of $(M \times \{1\}, \xi)$. By [Eliashberg 2004, Corollary 1.4], a weak semifilling can be capped off to give a weak filling. Since ω is exact, by [Eliashberg 2004, Proposition 4.1] this weak filling can be modified to give a strong filling (this result was independently obtained by Etnyre [2004]). Now, Gay [2006, Corollary 3] proved that any contact plane field that is strongly fillable has zero Giroux torsion. This proves the proposition for possibly a double cover of M .

Now the Giroux torsion of a contact structure cannot decrease under finite covers, since any component of a finite cover of a Giroux torsion domain is again a Giroux torsion domain. So, in any case, a contact structure ξ has zero Giroux torsion if and only if any finite lift of it has zero Giroux torsion. This proves the proposition for the original ξ . \square

Thanks to Giroux's correspondence [2002] between open books and contact structures, we can prove the following, giving the second implication needed for the statement of Theorem 1.12:

Theorem A.2 *Let φ_1 and φ_2 be two contact Anosov flows with respective contact structures ξ_1 and ξ_2 . If φ_1 and φ_2 are orbit equivalent (resp. isotopically equivalent) then ξ_1 and ξ_2 are contactomorphic (resp. isotopic).*

To apply Giroux's result in our proof, we first need to recall a result about Birkhoff sections, starting with their definition:

Definition A.3 Let φ be a flow on a 3-manifold M . A surface S is called a *topological Birkhoff section* of φ if

- (i) S is topologically immersed in M , and its interior is topologically embedded;
- (ii) the flow φ is topologically transverse to the interior of S ;
- (iii) each connected component of the boundary of S consists of a periodic orbit of φ ;
- (iv) every orbit of φ intersects S ; and
- (v) the return time of φ to the interior of S is uniformly bounded above and away from zero.

A topological Birkhoff section that is smoothly immersed is called a (smooth) Birkhoff section.

A Birkhoff section S in an orientable manifold is called *positive* if the orientations of all the boundary orbits correspond to the orientation induced by the flow on the interior of S .

Fried showed that any transitive Anosov flow has a Birkhoff section, which can in fact be taken to be embedded on the boundary as well. Moreover, Bonatti and Guelman [2010] show that this section can be assumed to be *tame*, meaning that, after an isotopy along flow lines, one can assume that the restriction of the Birkhoff section to a small tubular neighborhood of any component of its boundary is a smooth

helicoid. Finally, Marty [2021] showed that any \mathbb{R} -covered Anosov flow admits a *positive* tame Birkhoff section. Moreover, an adaptation of Marty's proof can be seen to yield a section that is also embedded on the boundary (T Marty, personal communication, 2021).⁵ Thus, one obtains an open book supporting the contact structure. We recall:

Definition A.4 (open book) Let $B \subset M$ be an oriented link in a connected oriented manifold. Then an open book (B, θ) with binding B is a fibration with connected oriented (noncompact) fibers $\theta: M \setminus B \rightarrow S^1$ such that, in a neighborhood of each binding component, the map is equivalent to the map given by projecting to the angular polar coordinate on $(D^2 \setminus \{0\}) \times S^1$, and the boundary of any fiber agrees with B as an oriented link.

The fibers of an open book are called *pages*. Note that the tameness condition in [Bonatti and Guelman 2010] corresponds precisely to a Birkhoff section S inducing an open book with binding the (oriented) periodic orbits ∂S .

Definition A.5 (supporting open book) An open book (B, θ) supports a (cooriented) contact structure (M, ξ) if there is a contact form α for ξ such that:

- The form $d\alpha$ is positive on the pages of θ , which are oriented to be compatible with the binding.
- The form α is positive on (each component of) B .

The fundamental fact due to Giroux [2002] is that any two contact structures supported by a *fixed* open book are isotopic through contact structures supported by the open book. This is essentially due to the fact that the above condition is convex, although some care is needed near the binding.

In order to apply the above, we will use the following result of [Bonatti and Guelman 2010]:

Lemma A.6 [Bonatti and Guelman 2010, Lemma 4.16] *Let S be a topological Birkhoff section of a flow φ ; then S can be isotoped along the orbits of φ to a smooth tame Birkhoff section S' .*

Proof of Theorem A.2 Let h be an orbit equivalence between φ_1 and φ_2 . Up to conjugating φ_1 by a diffeomorphism in the isotopy class of h , we can assume that φ_1 and φ_2 are isotopically equivalent and h is isotopic to the identity. Showing that ξ_1 and ξ_2 are isotopic for this new flow will imply that ξ_1 and ξ_2 are contactomorphic for the original one.

Let S be a smooth Birkhoff section for φ_1 . Then $h(S)$ is a topological Birkhoff section for φ_2 . By the lemma above, we can isotope $h(S)$ to a smooth tame Birkhoff section S_2 of φ_2 . Thus, the open book (B_2, θ_2) induced by S_2 supports ξ_2 . Now we isotope h to a smooth map g relative to the boundary of the

⁵One could also run the proof below using *rational* open books, which correspond to positive Birkhoff sections with immersed boundaries; see [Baker et al. 2012].

original Birkhoff section S . Then the open book (B_2, θ_2) pulls back to an open book (B', θ') with page $S' = g^{-1}S_2$ that supports $\xi'_1 = g^*\xi_2$. This smoothing can be arranged so that preimages of pages agree with those of an open book (B, θ) coming from the original Birkhoff section S near the binding. Then one notes that the open book (B', θ') is isotopic to the original one and we deduce that the corresponding contact structures are contactomorphic. Since h (and hence g) is isotopic to the identity, they are in fact isotopic. \square

A.1 Applications to contact topology

Now we can use Theorem 1.12 to translate to the language of contact topology some known results about Anosov flows. As a first example, Theorem 1.12 implies that the examples of skew Anosov flows on hyperbolic 3-manifolds built in [Bowden and Mann 2022] (which are all contact flows when done using Foulon–Hasselblatt contact surgery) have noncontactomorphic contact structures. Thus, we immediately obtain:

Theorem A.7 *For any $N \in \mathbb{N}$ there exists an hyperbolic 3-manifold with at least N noncontactomorphic Anosov contact structures.*

This result answers affirmatively a question raised in [Foulon et al. 2021]; see also [Bowden and Mann 2022, Question 7.4].

We can also translate Theorems 3.17 and 3.18 to a description of contact transformation groups, as follows. For a 3-manifold M with Anosov contact structure ξ , we follow [Giroux and Massot 2017] and denote by $\mathcal{D}(M, \xi)$ the group of contact transformations of (M, ξ) . Thus, there is a natural inclusion of $\pi_0\mathcal{D}(M, \xi)$ in $\text{MCG}(M)$.

Theorem A.8 *Let ξ be a Anosov contact structure on M . Suppose that either*

- (1) *M has a unique JSJ piece which is atoroidal, or*
- (2) *each torus of the JSJ decomposition of the manifold M is separating.*

In the first case, let D_ξ denote the subgroup of $\text{MCG}(M)$ generated by Dehn twists on the JSJ tori. In the second case, let D_ξ be the subgroup of $\text{MCG}(M)$ generated by all Dehn twists in the directions of periodic orbits together with any Dehn twists on the JSJ tori.

Then any class $[f] \in D_\xi$ admits a representative $f \in \mathcal{D}(M, \xi)$, and, conversely, there exists $n \in \mathbb{N}$ such that, for any class $[f] \in \pi_0\mathcal{D}(M, \xi)$, we have $[f]^n \in D_\xi$.

Remark A.9 This result partially extends a theorem of Giroux and Massot [2017], who obtained this for the case of Seifert fibered manifolds. Note that the result of Giroux and Massot is more precise, as ours only gives a description of $\pi_0\mathcal{D}(M, \xi)$ up to finite powers.

Remark A.10 It is not necessary to know the Anosov Reeb flow in order to detect which Dehn twist is in a direction of a periodic orbit, by the following observation: Let T be an embedded torus that is quasitransverse to the Anosov flow φ . Up to an arbitrarily small perturbation, one can put T in a convex position with respect to the contact structure ξ . Then an element $\alpha \in \pi_1(T)$ corresponding to a periodic orbit of the flow φ also corresponds to the free homotopy class of a connected component of the dividing set of the characteristic foliation of ξ on T . Therefore, one can use Theorem A.8 (or the translation of Corollary 1.5, which can be obtained in the same way) directly in contact geometry without having to go through the Anosov side.

Proof of Theorem A.8 We start by proving the converse implication. Let $f \in \mathcal{D}(M, \xi)$ and let φ be the (Anosov) Reeb flow. Then $f^{-1} \circ \varphi^t \circ f$ is a contact Anosov flow with contact structure $f_*\xi = \xi$. Thus, by Theorem 1.12, $f^{-1} \circ \varphi^t \circ f$ and φ^t are isotopically equivalent. Let $h: M \rightarrow M$ be an orbit equivalence isotopic to the identity between $f^{-1} \circ \varphi^t \circ f$ and φ^t ; then $h \circ f$ is a self-orbit equivalence of φ^t . Hence, by Theorems 3.17 or 3.18 depending on the case, there exists n such that $[f^n] = [(h \circ f)^n] \in D_\xi$.

Now, for the second part, let $[f]$ be a class in D_ξ . Then, by Theorems 3.17 or 3.18, $\psi = f^{-1} \circ \varphi \circ f$ is isotopically equivalent to φ . Moreover, the contact structure of ψ is $f_*\xi$, where ξ is the contact structure of φ . By Theorem 1.12, $f_*\xi$ and ξ are isotopic, so there exists g in the same isotopy class as f such that $g_*\xi = \xi$. That is, $g \in \mathcal{D}(M, \xi)$. \square

References

- [Baker et al. 2012] **K L Baker, J B Etnyre, J Van Horn-Morris**, *Cabling, contact structures and mapping class monoids*, J. Differential Geom. 90 (2012) 1–80 MR Zbl
- [Barbot 1995a] **T Barbot**, *Caractérisation des flots d’Anosov en dimension 3 par leurs feuilletages faibles*, Ergodic Theory Dynam. Systems 15 (1995) 247–270 MR Zbl
- [Barbot 1995b] **T Barbot**, *Mise en position optimale de tores par rapport à un flot d’Anosov*, Comment. Math. Helv. 70 (1995) 113–160 MR Zbl
- [Barbot 2001] **T Barbot**, *Plane affine geometry and Anosov flows*, Ann. Sci. École Norm. Sup. 34 (2001) 871–889 MR Zbl
- [Barbot 2005] **T Barbot**, *De l’hyperbolique au globalement hyperbolique*, Habilitation à diriger des recherches, Université Claude Bernard Lyon 1 (2005) Available at <https://theses.hal.science/tel-00011278>
- [Barbot and Fenley 2013] **T Barbot, S R Fenley**, *Pseudo-Anosov flows in toroidal manifolds*, Geom. Topol. 17 (2013) 1877–1954 MR Zbl
- [Barthelmé and Fenley 2017] **T Barthelmé, S R Fenley**, *Counting periodic orbits of Anosov flows in free homotopy classes*, Comment. Math. Helv. 92 (2017) 641–714 MR Zbl
- [Barthelmé and Gogolev 2019] **T Barthelmé, A Gogolev**, *A note on self orbit equivalences of Anosov flows and bundles with fiberwise Anosov flows*, Math. Res. Lett. 26 (2019) 711–728 MR Zbl
- [Barthelmé et al. 2023] **T Barthelmé, S R Fenley, R Potrie**, *Collapsed Anosov flows and self orbit equivalences*, Comment. Math. Helv. 98 (2023) 771–875 MR Zbl

- [Bonatti and Guelman 2010] **C Bonatti, N Guelman**, *Axiom A diffeomorphisms derived from Anosov flows*, *J. Mod. Dyn.* 4 (2010) 1–63 MR Zbl
- [Bonatti and Iakovoglou 2023] **C Bonatti, I Iakovoglou**, *Anosov flows on 3–manifolds: the surgeries and the foliations*, *Ergodic Theory Dynam. Systems* 43 (2023) 1129–1188 MR Zbl
- [Bonatti et al. 2020] **C Bonatti, A Gogolev, A Hammerlindl, R Potrie**, *Anomalous partially hyperbolic diffeomorphisms, III: Abundance and incoherence*, *Geom. Topol.* 24 (2020) 1751–1790 MR Zbl
- [Bowden and Mann 2022] **J Bowden, K Mann**, *C^0 stability of boundary actions and inequivalent Anosov flows*, *Ann. Sci. École Norm. Sup.* 55 (2022) 1003–1046 MR Zbl
- [Casson and Jungreis 1994] **A Casson, D Jungreis**, *Convergence groups and Seifert fibered 3–manifolds*, *Invent. Math.* 118 (1994) 441–456 MR Zbl
- [Colin et al. 2009] **V Colin, E Giroux, K Honda**, *Finitude homotopique et isotopique des structures de contact tendues*, *Publ. Math. Inst. Hautes Études Sci.* 109 (2009) 245–293 MR Zbl
- [Eliashberg 2004] **Y Eliashberg**, *A few remarks about symplectic filling*, *Geom. Topol.* 8 (2004) 277–293 MR Zbl
- [Eliashberg et al. 2000] **Y Eliashberg, A Givental, H Hofer**, *Introduction to symplectic field theory*, *Geom. Funct. Anal.* special volume (2000) 560–673 MR Zbl
- [Etnyre 2004] **J B Etnyre**, *On symplectic fillings*, *Algebr. Geom. Topol.* 4 (2004) 73–80 MR Zbl
- [Farb and Franks 2003] **B Farb, J Franks**, *Group actions on one-manifolds, II: Extensions of Hölder’s theorem*, *Trans. Amer. Math. Soc.* 355 (2003) 4385–4396 MR Zbl
- [Fenley 1994] **S R Fenley**, *Anosov flows in 3–manifolds*, *Ann. of Math.* 139 (1994) 79–115 MR Zbl
- [Fisher and Hasselblatt 2019] **T Fisher, B Hasselblatt**, *Hyperbolic flows*, *Eur. Math. Soc.*, Berlin (2019) MR Zbl
- [Foulon and Hasselblatt 2013] **P Foulon, B Hasselblatt**, *Contact Anosov flows on hyperbolic 3–manifolds*, *Geom. Topol.* 17 (2013) 1225–1252 MR Zbl
- [Foulon et al. 2021] **P Foulon, B Hasselblatt, A Vaugon**, *Orbit growth of contact structures after surgery*, *Ann. H. Lebesgue* 4 (2021) 1103–1141 MR Zbl
- [Funar 2013] **L Funar**, *Torus bundles not distinguished by TQFT invariants*, *Geom. Topol.* 17 (2013) 2289–2344 MR Zbl
- [Gabai 1992] **D Gabai**, *Convergence groups are Fuchsian groups*, *Ann. of Math.* 136 (1992) 447–510 MR Zbl
- [Gay 2006] **D T Gay**, *Four-dimensional symplectic cobordisms containing three-handles*, *Geom. Topol.* 10 (2006) 1749–1759 MR Zbl
- [Giroux 2002] **E Giroux**, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, from “Proceedings of the International Congress of Mathematicians, II” (T Li, editor), Higher Ed. Press, Beijing (2002) 405–414 MR Zbl
- [Giroux and Massot 2017] **E Giroux, P Massot**, *On the contact mapping class group of Legendrian circle bundles*, *Compos. Math.* 153 (2017) 294–312 MR Zbl
- [Hutchings and Nelson 2016] **M Hutchings, J Nelson**, *Cylindrical contact homology for dynamically convex contact forms in three dimensions*, *J. Symplectic Geom.* 14 (2016) 983–1012 MR Zbl
- [Johannson 1979] **K Johannson**, *Homotopy equivalences of 3–manifolds with boundaries*, *Lecture Notes in Math.* 761, Springer (1979) MR Zbl

- [Kovačević 1999] **N Kovačević**, *Möbius-like groups of homeomorphisms of the circle*, Trans. Amer. Math. Soc. 351 (1999) 4791–4822 MR Zbl
- [Liverani 2004] **C Liverani**, *On contact Anosov flows*, Ann. of Math. 159 (2004) 1275–1312 MR Zbl
- [Macarini and Paternain 2012] **L Macarini, G P Paternain**, *Equivariant symplectic homology of Anosov contact structures*, Bull. Braz. Math. Soc. 43 (2012) 513–527 MR Zbl
- [Marty 2021] **T Marty**, *Flots d’Anosov et sections de Birkhoff*, PhD thesis, Université Grenoble Alpes (2021) Available at <https://theses.hal.science/tel-03510071>
- [Mitsumatsu 1995] **Y Mitsumatsu**, *Anosov flows and non-Stein symplectic manifolds*, Ann. Inst. Fourier (Grenoble) 45 (1995) 1407–1421 MR Zbl
- [Plante 1981] **J F Plante**, *Anosov flows, transversely affine foliations, and a conjecture of Verjovsky*, J. Lond. Math. Soc. 23 (1981) 359–362 MR Zbl
- [Smale 1998] **S Smale**, *Mathematical problems for the next century*, Math. Intelligencer 20 (1998) 7–15 MR Zbl
- [Thurston 1997] **W P Thurston**, *Three-manifolds, foliations and circles, I*, preprint (1997) arXiv math/9712268 Reprinted in “Collected works with commentary, I: Foliations, surfaces and differential geometry” (B Farb, D Gabai, S P Kerckhoff, editors), Amer. Math. Soc., Providence, RI (2022) 353–412

*Department of Mathematics and Statistics, Queen’s University
Kingston ON, Canada*

*Department of Mathematics, Cornell University
Ithaca, NY, United States*

*Fakultät für Mathematik, Universität Regensburg
Regensburg, Germany*

`thomas.barthelme@queensu.ca, k.mann@cornell.edu,
jonathan.bowden@mathematik.uni-regensburg.de`

`sites.google.com/site/thomasbarthelme, https://e.math.cornell.edu/people/mann`

Proposed: David Fisher

Received: 13 January 2022

Seconded: Mladen Bestvina, Leonid Polterovich

Revised: 20 September 2022

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Mohammed Abouzaid	Stanford University abouzaid@stanford.edu	Mark Gross	University of Cambridge mgross@dpmms.cam.ac.uk
Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Arend Bayer	University of Edinburgh arend.bayer@ed.ac.uk	Sándor Kovács	University of Washington skovacs@uw.edu
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	Haynes Miller	Massachusetts Institute of Technology hrm@math.mit.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Aaron Naber	Northwestern University anaber@math.northwestern.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Peter Ozsváth	Princeton University petero@math.princeton.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Benson Farb	University of Chicago farb@math.uchicago.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David M Fisher	Rice University davidfisher@rice.edu	Roman Sauer	Karlsruhe Institute of Technology roman.sauer@kit.edu
Mike Freedman	Microsoft Research michaelf@microsoft.com	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
David Gabai	Princeton University gabai@princeton.edu	Natasa Sesum	Rutgers University natasas@math.rutgers.edu
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Cameron Gordon	University of Texas gordon@math.utexas.edu	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2024 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 28 Issue 2 (pages 497–1003) 2024

On the top-weight rational cohomology of \mathcal{A}_g	497
MADELINE BRANDT, JULIETTE BRUCE, MELODY CHAN, MARGARIDA MELO, GWYNETH MORELAND and COREY WOLFE	
Algebraic uniqueness of Kähler–Ricci flow limits and optimal degenerations of Fano varieties	539
JIYUAN HAN and CHI LI	
Valuations on the character variety: Newton polytopes and residual Poisson bracket	593
JULIEN MARCHÉ and CHRISTOPHER-LLOYD SIMON	
The local (co)homology theorems for equivariant bordism	627
MARCO LA VECCHIA	
Configuration spaces of disks in a strip, twisted algebras, persistence, and other stories	641
HANNAH ALPERT and FEDOR MANIN	
Closed geodesics with prescribed intersection numbers	701
YANN CHAUBET	
On endomorphisms of the de Rham cohomology functor	759
SHIZHANG LI and SHUBHODIP MONDAL	
The nonabelian Brill–Noether divisor on $\overline{\mathcal{M}}_{13}$ and the Kodaira dimension of $\overline{\mathcal{R}}_{13}$	803
GAVRIL FARKAS, DAVID JENSEN and SAM PAYNE	
Orbit equivalences of \mathbb{R} –covered Anosov flows and hyperbolic-like actions on the line	867
THOMAS BARTHELMÉ and KATHRYN MANN	
Microlocal theory of Legendrian links and cluster algebras	901
ROGER CASALS and DAPING WENG	
Correction to the article Bimodules in bordered Heegaard Floer homology	1001
ROBERT LIPSHITZ, PETER OZSVÁTH and DYLAN P THURSTON	