

# involve

a journal of mathematics

Vertex polygons

Candice Nielsen



# Vertex polygons

Candice Nielsen

(Communicated by Colin Adams)

We look at hexagons whose vertex triangles have equal area, and identify necessary conditions for these hexagons to also have vertex quadrilaterals with equal area. We discover a method for creating a hexagon whose vertex quadrilaterals have equal area without necessarily having vertex triangles of equal area. Finally, we generalize the process to build any polygon with an even number of sides to have certain vertex polygons with equal area.

## 1. Introduction

In the article “Polygons whose vertex triangles have equal area,” [Harel and Rabin \[2003\]](#) discuss the properties of polygons with the very special characteristic described in the title. To clarify, the authors offer the following definitions:

**Definition 1.** A triangle formed using three adjacent vertices of any polygon is called a *vertex triangle*.

**Definition 2.** A polygon  $V_1 V_2 \cdots V_n$  for which all vertex triangles have the same nonzero area is called an *equal-area polygon*.

Harel and Rabin take an algebraic approach, assigning direction and magnitude to each side of the polygon. In this article, we take a geometric approach, using area formulas and triangle congruencies to identify properties of certain polygons.

To extend from triangles, we offer the following definitions:

**Definition 3.** A polygon of  $n$  sides, formed using  $n$  adjacent vertices of any  $m$ -sided polygon (with  $m \geq n$ ), is called a *vertex  $n$ -gon*.


**Definition 4.** A polygon  $V_1 V_2 \cdots V_m$  for which all vertex  $n$ -gons have the same nonzero area is called an *equal- $n$ -gon polygon*.

It is clear that every equal-area quadrilateral is also an equal-quadrilateral polygon, since any vertex quadrilateral is the whole quadrilateral. Furthermore, every equal-area pentagon is an equal-quadrilateral polygon because, for  $P$  equal to the area of


---

*MSC2010:* primary 51N20; secondary 00A99.

*Keywords:* vertex polygons, vertex triangles, vertex quadrilaterals, geometry, equal area, hexagon, proof by contradiction.



**Figure 1.**  $\triangle BCD$  and  $\triangle DEF$  are vertex triangles of hexagon  $ABCDEF$ , but  $\triangle BEA$  is not.




**Figure 2.** An equal-area pentagon is always an equal-quadrilateral pentagon.

the pentagon, and  $T$  equal to the area of any vertex triangle, the area of every vertex quadrilateral is equal to  $P - T$  (Figure 2). This means every equal-quadrilateral pentagon is also an equal-area pentagon. For this reason, we begin with hexagons.

### 2. Equal-area hexagons

The first nontrivial case of the equal-area and equal-quadrilateral polygon is the hexagon. The first task is to construct an equal-area hexagon. We can show that, for any equal-area polygon  $V_1 V_2 \cdots V_n$ , the line  $V_i V_{i+1}$  is parallel to the line  $V_{i-1} V_{i+2}$ . In other words, each side is parallel to the line formed by the surrounding two vertices.

*Proof.* Let  $V_1 V_2 \cdots V_n$  be an equal-area polygon. Then  $\text{Area}(\triangle V_{i-1} V_i V_{i+1}) = \text{Area}(\triangle V_i V_{i+1} V_{i+2})$ . Let  $b = V_i V_{i+1}$ ,  $h_1 = d(V_{i-1}, V_i V_{i+1})$ ,  $h_2 = d(V_{i+2}, V_i V_{i+1})$ . So  $\text{Area}(\triangle V_{i-1} V_i V_{i+1}) = \frac{1}{2}bh_1 = \frac{1}{2}bh_2 = \text{Area}(\triangle V_i V_{i+1} V_{i+2})$ . Therefore,  $h_1 = h_2$  and  $V_{i-1} V_{i+2}$  is parallel to  $V_i V_{i+1}$ .  $\square$




**Figure 3.** Every equal-area hexagon enjoys parallelism between opposite sides and corresponding main diagonals.


With this property, an equal-area hexagon can be uniquely determined by any trapezoid. As we build an equal-area hexagon, it is important to note that the diagonals of the hexagon need not intersect at a single point. This is a key observation as we transform the equal-area hexagon into an equal-quadrilateral hexagon.

Using the hexagon  $CDEFGH$  in Figure 3, certain geometric properties arise. First, the sides of the hexagon, along with the diagonals, divide the hexagon into four triangles and three trapezoids. Let us define these as follows:

**Definition 5.** Let  $ABCDEF$  be any hexagon, with  $\overline{AD} \cap \overline{BE} = J$ ,  $\overline{BE} \cap \overline{CF} = L$ ,  $\overline{CF} \cap \overline{AD} = K$ . The triangle  $\triangle JKL$  is called the *center triangle*. The triangles  $\triangle ABJ$ ,  $\triangle CKD$ , and  $\triangle ELF$  are called *interior triangles*, and  $BCKJ$ ,  $DELK$ , and  $FAJL$  are called *interior trapezoids* (see Figure 4).



**Figure 4.** Equal-area hexagon  $ABCDEF$  and two of the associated trapezoids.



**Figure 5.** Equal-area, equal-quadrilateral hexagon  $ABCDEF$ .


**Lemma.** For an equal-area hexagon, each interior trapezoid has the same nonzero area and each interior triangle has the same nonzero area.

*Proof.* Let  $ABCDEF$  be an equal-area hexagon, with  $\overline{AD} \cap \overline{BE} = J$ ,  $\overline{BE} \cap \overline{CF} = L$ ,  $\overline{CF} \cap \overline{AD} = K$  (see Figure 5). From the previous proof,  $\overline{AF} \parallel \overline{BE}$  and  $\overline{AB} \parallel \overline{FL}$ , so  $ABLF$  is a parallelogram. Likewise,  $\overline{AK} \parallel \overline{BC}$  and  $\overline{AB} \parallel \overline{CK}$ , so  $ABCK$  is a parallelogram, and the parallelograms share a base,  $\overline{AB}$ . Let  $b_1 = AB$ ,  $h_1 = \text{height}(ABLF) = \text{height}(\triangle BAF)$ , and  $h_2 = \text{height}(ABCK) = \text{height}(\triangle ABC)$ . Then  $\text{Area}(ABLF) = b_1 h_1 = 2\text{Area}(\triangle BAF)$  and  $\text{Area}(ABCK) = b_1 h_2 = 2\text{Area}(\triangle ABC)$ . Since  $ABCDEF$  is an equal-area hexagon, we have  $\text{Area}(\triangle BAF) = \text{Area}(\triangle ABC)$ , so  $\text{Area}(ABLF) = \text{Area}(ABCK)$ . Let  $A_1 = \text{Area}(AJLF)$ ,  $A_2 = \text{Area}(\triangle ABJ)$ , and  $A_3 = \text{Area}(BCKJ)$ . Then  $\text{Area}(ABLF) = A_1 + A_2$  and  $\text{Area}(ABCK) = A_2 + A_3$ . This implies that  $A_1 = A_3$ . Similar argument supports that all interior trapezoids have the same nonzero area, as do all interior triangles.  $\square$

**Definition 6.** For any integer  $n > 1$  and any polygon having  $n$  sides with vertices  $V_1, V_2, \dots, V_{2n}$ , a true diagonal has endpoints  $V_i$  and  $V_{i+n}$ , where  $i \in \{1, 2, \dots, n\}$ .

**Theorem 1.** An equal-area hexagon is equal-quadrilateral if and only if all its true diagonals intersect at a single point.

*Proof.* Let  $ABCDEF$  be an equal-area, equal-quadrilateral hexagon, with the following properties:  $\overline{AD} \cap \overline{BE} = J$ ,  $\overline{BE} \cap \overline{CF} = L$ ,  $\overline{CF} \cap \overline{AD} = K$ . Suppose, for the sake of



**Figure 6.** Equal-area hexagon  $ABCDEF$  with main diagonals intersecting.

contradiction, that  $J$ ,  $K$ , and  $L$  are three distinct points. Let  $A_1$  be the area of the interior triangles,  $A_2$  be the area of the interior trapezoids, and  $A_c$  be the area of the center triangle. Consider the vertex quadrilaterals  $ABCD$  and  $BCDE$ . Since  $ABCDEF$  is an equal-quadrilateral hexagon, the areas of the vertex quadrilaterals are equal to each other. Thus,  $\text{Area}(ABCD) = 2A_1 + A_2 = A_1 + 2A_2 + A_c = \text{Area}(BCDE)$ .

Let  $b_1 = DE$ , and let  $h_1$  equal the height of trapezoid  $ELKD$ , which is equal to the height of vertex triangle  $EDC$ .

Let  $b_2 = LK$ , and let  $h_2$  equal the height of center triangle  $JKL$ .

Let  $b_3 = AB$  and let  $h_3$  equal the height of vertex triangle  $ABC$ , so the height of interior triangle  $ABJ$  is  $h_3 - h_2$ . Then  $2A_1 + A_2 = A_1 + 2A_2 + A_c$  implies

$$2\left(\frac{1}{2}b_3(h_3 - h_2)\right) + \frac{1}{2}(b_1 + b_2)h_1 = \frac{1}{2}b_3(h_3 - h_2) + 2\left(\frac{1}{2}(b_1 + b_2)h_1\right) + \frac{1}{2}b_2h_2$$

This simplifies to

$$b_2h_2 + b_1h_1 + b_2h_1 = b_3h_3 - b_3h_2. \tag{1}$$

Since  $ABCDEF$  is equal-area, the vertex triangles have the same area, and  $\frac{1}{2}b_1h_1 = \frac{1}{2}b_3h_3$ , so  $b_1h_1 = b_3h_3$  and (1) becomes

$$b_2h_2 + b_2h_1 = -b_3h_2. \tag{2}$$

Since  $b_2h_2$ ,  $b_2h_1$ ,  $b_3h_2$  are all positive values, this is a contradiction. Therefore,  $J = K = L$ , and the diagonals of  $ABCDEF$  intersect at a single point.

For the other direction, let  $ABCDEF$  be an equal-area hexagon, satisfying  $\overline{AD} \cap \overline{BE} \cap \overline{CF} = X$ .

Without loss of generality, consider  $\triangle ABX$ . Since the height of  $\triangle ABX$  is equal to the height of  $\triangle ABC$ , their areas are equal. Thus, the area of each interior triangle


is the area of a vertex triangle. Since all vertex triangles share an equal area, so do the interior triangles. Each vertex quadrilateral is made up of three interior triangles, so each vertex quadrilateral shares an equal area. Therefore,  $ABCDEF$  is an equal-quadrilateral hexagon.  $\square$

### 3. Equal-quadrilateral hexagons


While constructing an equal-quadrilateral hexagon out of an equal-area hexagon is helpful, the question arose: if a hexagon is equal-quadrilateral, is it necessarily equal-area? We are able to observe, through interior triangle congruencies, that the intersection of the three diagonals is the midpoint of each diagonal. Since the three diagonals are diameters of three concentric circles, we have a new way to construct the equal-quadrilateral hexagon.

**Theorem 2.** *A hexagon whose true diagonals are diameters of concentric circles is an equal-quadrilateral hexagon.*

*Proof.* Let  $AD$ ,  $BE$ ,  $CF$  be diameters of three concentric circles with center  $X$  and also be diagonals of hexagon  $ABCDEF$  (see Figure 7). Without loss of generality, consider  $ABCD$  and  $BCDE$ . We have  $ABCD \cap BCDE = BCDX$ . We also have  $EX = XB$  and  $AX = XD$  because they are radii of the same respective circles. Furthermore,  $\angle EXD \cong \angle BXA$  because they are vertical angles. Thus, by the side-angle-side condition,  $\triangle EXD \cong \triangle BXA$ . Since  $BCDX$  is congruent to itself,  $ABCD$  and  $BCDE$  are congruent, and therefore share an equal, nonzero area. With this argument, every vertex quadrilateral of  $ABCDEF$  shares the same, nonzero area. Thus,  $ABCDEF$  is an equal-quadrilateral hexagon.  $\square$



**Figure 7.** Equal-quadrilateral hexagon  $ABCDEF$ .



**Figure 8.**  $ABCDEF$  is an equal-quadrilateral hexagon, but it is not an equal-area hexagon:  $\text{Area}(\triangle ABC)$ ,  $\text{Area}(\triangle BCD)$ , and  $\text{Area}(\triangle CDE)$  are all different (and each is equal to the area of the symmetrically placed triangle).

To answer the question posed at the beginning of this section, [Figure 8](#) offers a counterexample. All vertex quadrilaterals share an equal area, while the vertex triangles have varying areas.


#### 4. Equal-( $n+1$ )-gon polygons

**Corollary.** *For any integer  $n > 1$ , a polygon with  $2n$  sides is an equal-( $n+1$ )-gon polygon if its true diagonals are diameters of  $n$  concentric circles (see [Figure 9](#)).*

*Proof.* Let  $n \in \mathbb{Z}$ , with  $n > 1$ . Let  $P_0$  be a  $2n$ -sided polygon constructed using the endpoints of diameters of  $n$  concentric circles. Call the center of the circles  $B$ , and denote the vertices of  $P_0$  by  $V_1, V_2, V_3, \dots, V_{2n}$ .

Let  $P_1$  be a polygon with vertices  $V_i, V_{i+1}, \dots, V_{i+n}$ , and let  $\text{Area}(P_1) = A_1$ . Let  $P_2$  be the polygon with vertices  $V_{i+1}, V_{i+2}, \dots, V_{i+n+1}$  and let  $\text{Area}(P_2) = A_2$ . We have  $P_1 \cap P_2 = \text{polygon}(V_{i+1}, V_{i+2}, \dots, V_{i+n}) \cup \triangle V_{i+1}V_{n+1}B$ , which we will call  $Q_0$ . Note that  $\text{Area}(Q_0)$  is equal to itself, so we need only to prove that  $\text{Area}(P_1 - Q_0) = \text{Area}(P_2 - Q_0)$ . Since  $BV_{i+n}$  and  $BV_i$  are radii of the same circle, they are congruent, and likewise for  $BV_{i+1}$  and  $BV_{i+n+1}$ . Angles  $V_iBV_{i+1}$  and  $V_{i+n}BV_{i+n+1}$  are congruent because they are vertical angles. Thus, by the side-angle-side formula, the triangles are congruent and therefore have equal area.





**Figure 9.**  $P_1$  constructed from diameters of concentric circles.

So  $\text{Area}(P_1 - Q_0) = \text{Area}(P_2 - Q_0)$ , and we finally have

$$\begin{aligned} \text{Area}(P_1) &= \text{Area}(Q_0) + \text{Area}(P_1 - Q_0) \\ &= \text{Area}(P_2 - Q_0) + \text{Area}(Q_0) = \text{Area}(P_2). \end{aligned}$$

Therefore, the areas of all vertex  $(n+1)$ -gons are equal to each other. □

### 5. Results and open questions

Using known properties of equal-area polygons, we discovered properties of the equal-quadrilateral hexagon. We stated and proved a result that gives necessary conditions for an equal-area hexagon to also be equal-quadrilateral. Finally, we were able to generalize the process of constructing an equal-quadrilateral hexagon to allow construction of any equal- $(n+1)$ -gon polygon.

An additional observation on the equal-area hexagon, whether convex or non-convex, is that the area of the hexagon is equal to the sum of the areas of the vertex triangles. Likewise, the area of any equal-quadrilateral hexagon is twice the area of the vertex quadrilaterals. While this is immediately clear for a convex hexagon, it is not so when the hexagon is nonconvex. Since it is likely the proofs for these observations are simple, they were omitted from this article.

Some questions to consider in extending the idea of equal- $n$ -gon polygons are:

- (1) Given an equal-area heptagon, what are the necessary conditions to imply an equal-quadrilateral heptagon? Does equal-quadrilateral imply equal-area in heptagons? If not, how can we construct an equal-quadrilateral heptagon?

- (2) Our corollary applies only to polygons with an even number of sides. Given a polygon with an odd number of sides, are there sufficient conditions to ensure vertex polygons of equal area?

### Acknowledgements

The author wishes to thank her advisor, Dr. Jon Johnson, for his guidance and support throughout the semester.

### References

[Harel and Rabin 2003] G. Harel and J. M. Rabin, “Polygons whose vertex triangles have equal area”, *Amer. Math. Monthly* **110**:7 (2003), 606–619. [MR 2004e:51021](#) [Zbl 1046.51008](#)

Received: 2012-09-18      Revised: 2012-11-10      Accepted: 2012-12-15

[nielsenc@net.elmhurst.edu](mailto:nielsenc@net.elmhurst.edu)

*Mathematics Department, Elmhurst College,  
Elmhurst, IL 60126, United States*

## EDITORS

### MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, [berenhks@wfu.edu](mailto:berenhks@wfu.edu)

### BOARD OF EDITORS

|                      |   |                        |  |
|----------------------|---|------------------------|--|
| Colin Adams          | Williams College, USA<br><a href="mailto:colin.c.adams@williams.edu">colin.c.adams@williams.edu</a>         | David Larson           | Texas A&M University, USA<br><a href="mailto:larson@math.tamu.edu">larson@math.tamu.edu</a>                          |
| John V. Baxley       | Wake Forest University, NC, USA<br><a href="mailto:baxley@wfu.edu">baxley@wfu.edu</a>                       | Suzanne Lenhart        | University of Tennessee, USA<br><a href="mailto:lenhart@math.utk.edu">lenhart@math.utk.edu</a>                       |
| Arthur T. Benjamin   | Harvey Mudd College, USA<br><a href="mailto:benjamin@hmc.edu">benjamin@hmc.edu</a>                          | Chi-Kwong Li           | College of William and Mary, USA<br><a href="mailto:ckli@math.wm.edu">ckli@math.wm.edu</a>                           |
| Martin Bohner        | Missouri U of Science and Technology, USA<br><a href="mailto:bohner@mst.edu">bohner@mst.edu</a>             | Robert B. Lund         | Clemson University, USA<br><a href="mailto:lund@clemson.edu">lund@clemson.edu</a>                                    |
| Nigel Boston         | University of Wisconsin, USA<br><a href="mailto:boston@math.wisc.edu">boston@math.wisc.edu</a>              | Gaven J. Martin        | Massey University, New Zealand<br><a href="mailto:g.j.martin@massey.ac.nz">g.j.martin@massey.ac.nz</a>               |
| Amarjit S. Budhiraja | U of North Carolina, Chapel Hill, USA<br><a href="mailto:budhiraj@email.unc.edu">budhiraj@email.unc.edu</a> | Mary Meyer             | Colorado State University, USA<br><a href="mailto:meyer@stat.colostate.edu">meyer@stat.colostate.edu</a>             |
| Pietro Cerone        | Victoria University, Australia<br><a href="mailto:pietro.cerone@vu.edu.au">pietro.cerone@vu.edu.au</a>      | Emil Minchev           | Ruse, Bulgaria<br><a href="mailto:eminchev@hotmail.com">eminchev@hotmail.com</a>                                     |
| Scott Chapman        | Sam Houston State University, USA<br><a href="mailto:scott.chapman@shsu.edu">scott.chapman@shsu.edu</a>     | Frank Morgan           | Williams College, USA<br><a href="mailto:frank.morgan@williams.edu">frank.morgan@williams.edu</a>                    |
| Joshua N. Cooper     | University of South Carolina, USA<br><a href="mailto:cooper@math.sc.edu">cooper@math.sc.edu</a>             | Mohammad Sal Moslehian | Ferdowsi University of Mashhad, Iran<br><a href="mailto:moslehian@ferdowsi.um.ac.ir">moslehian@ferdowsi.um.ac.ir</a> |
| Jem N. Corcoran      | University of Colorado, USA<br><a href="mailto:corcoran@colorado.edu">corcoran@colorado.edu</a>             | Zuhair Nashed          | University of Central Florida, USA<br><a href="mailto:znashed@mail.ucf.edu">znashed@mail.ucf.edu</a>                 |
| Toka Diagana         | Howard University, USA<br><a href="mailto:tdiagana@howard.edu">tdiagana@howard.edu</a>                      | Ken Ono                | Emory University, USA<br><a href="mailto:ono@mathcs.emory.edu">ono@mathcs.emory.edu</a>                              |
| Michael Dorff        | Brigham Young University, USA<br><a href="mailto:mdorff@math.byu.edu">mdorff@math.byu.edu</a>               | Timothy E. O'Brien     | Loyola University Chicago, USA<br><a href="mailto:tbriell@luc.edu">tbriell@luc.edu</a>                               |
| Sever S. Dragomir    | Victoria University, Australia<br><a href="mailto:sever@matilda.vu.edu.au">sever@matilda.vu.edu.au</a>      | Joseph O'Rourke        | Smith College, USA<br><a href="mailto:orourke@cs.smith.edu">orourke@cs.smith.edu</a>                                 |
| Behrouz Emamizadeh   | The Petroleum Institute, UAE<br><a href="mailto:bemamizadeh@pi.ac.ae">bemamizadeh@pi.ac.ae</a>              | Yuval Peres            | Microsoft Research, USA<br><a href="mailto:peres@microsoft.com">peres@microsoft.com</a>                              |
| Joel Foisy           | SUNY Potsdam<br><a href="mailto:foisyjs@potsdam.edu">foisyjs@potsdam.edu</a>                                | Y.-F. S. Pétermann     | Université de Genève, Switzerland<br><a href="mailto:petermann@math.unige.ch">petermann@math.unige.ch</a>            |
| Errin W. Fulp        | Wake Forest University, USA<br><a href="mailto:fulp@wfu.edu">fulp@wfu.edu</a>                               | Robert J. Plemmons     | Wake Forest University, USA<br><a href="mailto:plemmons@wfu.edu">plemmons@wfu.edu</a>                                |
| Joseph Gallian       | University of Minnesota Duluth, USA<br><a href="mailto:kgallian@d.umn.edu">kgallian@d.umn.edu</a>           | Carl B. Pomerance      | Dartmouth College, USA<br><a href="mailto:carl.pomerance@dartmouth.edu">carl.pomerance@dartmouth.edu</a>             |
| Stephan R. Garcia    | Pomona College, USA<br><a href="mailto:stephan.garcia@pomona.edu">stephan.garcia@pomona.edu</a>             | Vadim Ponomarenko      | San Diego State University, USA<br><a href="mailto:vadim@sciences.sdsu.edu">vadim@sciences.sdsu.edu</a>              |
| Anant Godbole        | East Tennessee State University, USA<br><a href="mailto:godbole@etsu.edu">godbole@etsu.edu</a>              | Bjorn Poonen           | UC Berkeley, USA<br><a href="mailto:poonen@math.berkeley.edu">poonen@math.berkeley.edu</a>                           |
| Ron Gould            | Emory University, USA<br><a href="mailto:rg@mathcs.emory.edu">rg@mathcs.emory.edu</a>                       | James Propp            | U Mass Lowell, USA<br><a href="mailto:jpropp@cs.uml.edu">jpropp@cs.uml.edu</a>                                       |
| Andrew Granville     | Université Montréal, Canada<br><a href="mailto:andrew@dms.umontreal.ca">andrew@dms.umontreal.ca</a>         | József H. Przytycki    | George Washington University, USA<br><a href="mailto:przytyck@gwu.edu">przytyck@gwu.edu</a>                          |
| Jerrold Griggs       | University of South Carolina, USA<br><a href="mailto:griggs@math.sc.edu">griggs@math.sc.edu</a>             | Richard Rebarber       | University of Nebraska, USA<br><a href="mailto:rrebarbe@math.unl.edu">rrebarbe@math.unl.edu</a>                      |
| Sat Gupta            | U of North Carolina, Greensboro, USA<br><a href="mailto:sgupta@uncg.edu">sgupta@uncg.edu</a>                | Robert W. Robinson     | University of Georgia, USA<br><a href="mailto:rwr@cs.uga.edu">rwr@cs.uga.edu</a>                                     |
| Jim Haglund          | University of Pennsylvania, USA<br><a href="mailto:jhaglund@math.upenn.edu">jhaglund@math.upenn.edu</a>     | Filip Saidak           | U of North Carolina, Greensboro, USA<br><a href="mailto:f_saidak@uncg.edu">f_saidak@uncg.edu</a>                     |
| Johnny Henderson     | Baylor University, USA<br><a href="mailto:johnny_henderson@baylor.edu">johnny_henderson@baylor.edu</a>      | James A. Sellers       | Penn State University, USA<br><a href="mailto:sellersj@math.psu.edu">sellersj@math.psu.edu</a>                       |
| Jim Hoste            | Pitzer College<br><a href="mailto:jhoste@pitzer.edu">jhoste@pitzer.edu</a>                                  | Andrew J. Sterge       | Honorary Editor<br><a href="mailto:andy@ajsterge.com">andy@ajsterge.com</a>  |
| Natalia Hritonenko   | Prairie View A&M University, USA<br><a href="mailto:nhritonenko@pvamu.edu">nhritonenko@pvamu.edu</a>        | Ann Trenk              | Wellesley College, USA<br><a href="mailto:atrenk@wellesley.edu">atrenk@wellesley.edu</a>                             |
| Glenn H. Hurlbert    | Arizona State University, USA<br><a href="mailto:hurlbert@asu.edu">hurlbert@asu.edu</a>                     | Ravi Vakil             | Stanford University, USA<br><a href="mailto:vakil@math.stanford.edu">vakil@math.stanford.edu</a>                     |
| Charles R. Johnson   | College of William and Mary, USA<br><a href="mailto:crjohnso@math.wm.edu">crjohnso@math.wm.edu</a>          | Antonia Vecchio        | Consiglio Nazionale delle Ricerche, Italy<br><a href="mailto:antonia.vecchio@cnr.it">antonia.vecchio@cnr.it</a>      |
| K. B. Kulasekera     | Clemson University, USA<br><a href="mailto:kk@ces.clemson.edu">kk@ces.clemson.edu</a>                       | Ram U. Verma           | University of Toledo, USA<br><a href="mailto:verma99@msn.com">verma99@msn.com</a>                                    |
| Gerry Ladas          | University of Rhode Island, USA<br><a href="mailto:gladas@math.uri.edu">gladas@math.uri.edu</a>             | John C. Wierman        | Johns Hopkins University, USA<br><a href="mailto:wierman@jhu.edu">wierman@jhu.edu</a>                                |
|                      |   | Michael E. Zieve       | University of Michigan, USA<br><a href="mailto:zieve@umich.edu">zieve@umich.edu</a>                                  |

## PRODUCTION


Silvio Levy, Scientific Editor

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2012 is US \$105/year for the electronic version, and \$145/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2012 Mathematical Sciences Publishers

# involve

2012

vol. 5

no. 3

|  |     |
|--|-----|
| Analysis of the steady states of a mathematical model for Chagas disease                             | 237 |
| MARY CLAUSON, ALBERT HARRISON, LAURA SHUMAN, MEIR SHILLOR AND ANNA MARIA SPAGNUOLO                   |     |
| Bounds on the artificial phase transition for perfect simulation of hard core Gibbs processes        | 247 |
| MARK L. HUBER, ELISE VILLELLA, DANIEL ROZENFELD AND JASON XU   |     |
| A nonextendable Diophantine quadruple arising from a triple of Lucas numbers                         | 257 |
| A. M. S. RAMASAMY AND D. SARASWATHY  |     |
| Alhazen's hyperbolic billiard problem  | 273 |
| NATHAN POIRIER AND MICHAEL MCDANIEL  |     |
| Bochner $(p, Y)$ -operator frames  | 283 |
| MOHAMMAD HASAN FAROUGH, REZA AHMADI AND MORTEZA RAHMANI  |     |
| $k$ -furus semigroups  | 295 |
| NICHOLAS R. BAETH AND KAITLYN CASSITY  |     |
| Studying the impacts of changing climate on the Finger Lakes wine industry                           | 303 |
| BRIAN MCGAUVRAN AND THOMAS J. PFAFF  |     |
| A graph-theoretical approach to solving Scramble Squares puzzles                                     | 313 |
| SARAH MASON AND MALI ZHANG   |     |
| The $n$ -diameter of planar sets of constant width   | 327 |
| ZAIR IBRAGIMOV AND TUAN LE   |     |
| Boolean elements in the Bruhat order on twisted involutions  | 339 |
| DELONG MENG  |     |
| Statistical analysis of diagnostic accuracy with applications to cricket                             | 349 |
| LAUREN MONDIN, COURTNEY WEBER, SCOTT CLARK, JESSICA WINBORN, MELINDA M. HOLT AND ANANDA B. W. MANAGE |     |
| Vertex polygons  | 361 |
| CANDICE NIELSEN  |     |
| Optimal trees for functions of internal distance   | 371 |
| ALEX COLLINS, FEDELIS MUTISO AND HUA WANG  |     |