

involve

a journal of mathematics

Polygonal bicycle paths
and the Darboux transformation

Ian Alevy and Emmanuel Tsukerman



Polygonal bicycle paths and the Darboux transformation

Ian Alevy and Emmanuel Tsukerman

(Communicated by Kenneth S. Berenhaut)

A bicycle (n, k) -gon is an equilateral n -gon whose k diagonals are of equal length. In this paper we introduce periodic bicycle (n, k) -paths, which are a natural variation in which the polygon is replaced with a periodic polygonal path, and study their rigidity and integrals of motion.

1. Background

Our motivation comes from three seemingly unrelated problems. The first is the problem of *floating bodies of equilibrium in two dimensions*. From 1935 to 1941, mathematicians at the University of Lviv, among them Stefan Banach and Mark Kac, collected mathematical problems in a book, which became known as “the Scottish book”, since they often met in the Scottish Coffee House. Stanislaw Ulam posed problem 19 of this book: “Is a sphere the only solid of uniform density which will float in water in any position?” The answer in the two-dimensional case, as it turns out, depends on the density of the solid.

The second problem, known as the *tire track problem*, originated in the story, “The adventure of the priory school” by Arthur Conan Doyle, where Sherlock Holmes and Dr. Watson discuss in view of the two tire tracks of a bicycle which way the bicycle went. The problem is: “Is it possible that tire tracks other than circles or straight lines are created by bicyclists going in both directions?” As shown in Figure 1, the answer to this subtle question is affirmative.

The third problem is that of describing the trajectories of *electrons in a parabolic magnetic field*. All three problems turn out to be equivalent [Wegner 2007].

Often in mathematics it is fruitful to discretize a problem. As such, S. Tabachnikov [2006] proposed a “discrete bicycle curve” (also known as a “bicycle polygon”), which is a polygon satisfying discrete analogs of the properties of a bicycle track. The main requirement turns out to be that, in the language of discrete differential geometry, the polygon is “self-Darboux”. That is, the discrete differential

MSC2010: 37J35, 52C25.

Keywords: bicycle polygons, tire track problem, floating bodies in equilibrium.




Figure 1. Ambiguous bicycle tracks. The rear-wheel track is the inner curve and the front-wheel track is the outer curve. One cannot tell which way the bicycle went because a bicycle could have followed either one of two trajectories [Wegner 2007].

geometric notion of a discrete Darboux transformation [Bobenko and Suris 2008; Tsuruga 2010], which relates one polygon to another, relates a discrete bicycle curve to itself.

The topic of bicycle curves and polygons belongs to a number of active areas of research. On the one hand, it is part of rigidity theory. As an illustration, R. Connelly and B. Csikós [2009] consider the problem of classifying first-order flexible regular bicycle polygons. Other work on the rigidity theory of bicycle curves and polygons can be found in [Csikós 2007; Cyr 2012; Tabachnikov 2006].

The topic is also part of the subject of discrete integrable systems. This point of view is taken in [Tabachnikov and Tsukerman 2013], where the authors find integrals of motion (i.e., quantities which are conserved) of bicycle curves and polygons under the Darboux transformation and recutting of polygons [Adler 1993; 1995].

In this paper, in analogy with bicycle polygons, we introduce a new concept called a periodic discrete bicycle path and study both its rigidity and integrals.

2. Bicycle (n, k) -paths

A bicycle (n, k) -gon is an equilateral n -gon whose k diagonals are of equal length [Tabachnikov 2006]. We consider the following analog.

Definition 1. Define $P = \{V_i \in \mathbb{R}^2 : i \in \mathbb{Z}\}$ (for brevity, $V_0 V_1 \cdots V_{n-1}$) to be a *discrete periodic bicycle (n, k) -path* (or discrete (n, k) -path) if the following conditions hold:

- (i) $V_{n+i} = V_i + e_1$ for all i , where $e_1 = (1, 0)$ and $V_0 = (0, 0)$ (periodicity condition).
- (ii) $|V_i V_{i+1}| = |V_j V_{j+1}|$ for all i, j (equilateralness).
- (iii) $|V_i V_{i+k}| = |V_j V_{j+k}|$ for all i, j (equality of k -diagonals).

Definition 1 is meant to model the motion of a bicycle whose trajectory is spatially periodic. The condition that $|V_j V_{j+1}|$ is independent of j prescribes a constant speed for the motion of the bike. The condition that $|V_j V_{j+k}|$ is independent of j

represents the ambiguity of the direction in which the bicycle went (see [Tabachnikov 2006] for details).

Some natural questions regarding periodic (n, k) -paths are for which pairs (n, k) they exist, how many there are and whether they are rigid or flexible. We consider these questions in Section 3. A simple example of a bicycle (n, k) -path, analogous to the regular (n, k) -polygon, is $V_i = (i/n, 0)$, i.e., when all vertices lie at equal intervals on the line. We call this the *regular path*. Since bicycle (n, k) -paths are discretized bicycle paths, it is also interesting to see if there are any integrals of motion. We show that this is indeed the case in Section 4, by showing that area is an integral of motion.

3. Rigidity

The following two lemmas will be helpful when analyzing the rigidity of discrete bicycle paths.

Lemma 2. *Let $n \in \mathbb{N}$, $\chi_i \in \{-1, 1\}$ for every $i \in \mathbb{Z}/n\mathbb{Z}$ and let*

$S = \{(x_0, x_1, \dots, x_{n-1}) \in \mathbb{R}^n : (x_{i+1} - x_i)^2 = (x_{j+1} - x_j)^2 \text{ for all } i, j \in \mathbb{Z}/n\mathbb{Z}\}$.
Then

$$S = \left\{ (x_0, x_1, \dots, x_{n-1}) : x_{i+1} = x_i + \chi_i r \text{ for } i \in \mathbb{Z}/n\mathbb{Z}, \sum_{i=0}^{n-1} r \chi_i = 0 \text{ and } r \geq 0 \right\}.$$

In particular, if n is odd, then $S = \{(x_0, x_1, \dots, x_{n-1}) : x_i = x_j \text{ for all } i, j \in \mathbb{Z}/n\mathbb{Z}\}$.

Proof. First note that the candidate set is well-defined since

$$x_{j+n} = x_j + \sum_{i=j}^{j+n-1} r \chi_i = x_j + \sum_{i=0}^{n-1} r \chi_i = x_j.$$

Let $(x_0, x_1, \dots, x_{n-1}) \in S$. Recall that

$$\text{sgn}(x) = \begin{cases} 1 & \text{if } x > 0, \\ 0 & \text{if } x = 0, \\ -1 & \text{if } x < 0, \end{cases}$$

and that $\text{sgn}(x)|x| = x$. Set $r := |x_{i+1} - x_i|$ and $\chi_i = \text{sgn}(x_{i+1} - x_i) + (1 - \text{sgn}(r))$. Then

$$x_{i+1} = x_i + \chi_i r,$$

and

$$\sum_{i=0}^{n-1} r \text{sgn}(x_{i+1} - x_i) = 0.$$

It follows that any n -tuple in S satisfies the conditions $x_{i+1} = x_i + \chi_i r$, $\sum_{i=0}^{n-1} r \chi_i = 0$ and $r \geq 0$. The opposite inclusion is clear. \square

Lemma 3. *Let $x_i \in \mathbb{R}$ for every $i \in \mathbb{Z}$ with $x_0 = 0$ and let k and n be coprime integers. Assume that $x_{i+k} - x_i = x_i - x_{i-k}$ for each i and that $x_{i+n} = 1 + x_i$. Then $x_i = i/n$ for each i .*

Proof. Define z_i via $x_i = z_i + i/n$. The hypothesis $x_{i+n} = 1 + x_i$ implies that

$$z_{i+n} = z_i.$$

The difference

$$\Delta z := z_{i+k} - z_i$$

is independent of i due to the assumption that $x_{i+k} - x_i = x_i - x_{i-k}$ and because k and n are coprime. This implies that

$$0 = z_{i+n} - z_i = z_{i+nk} - z_i = n\Delta z.$$

It follows that $z_i = 0$ for every i . □

The following theorem gives a classification of a family of periodic (n, k) -paths.

Theorem 4. *The discrete $(n, dn - 1)$ -paths $V_i = (x_i, y_i)$, $i \in \mathbb{N}$ with $d \neq 0$ are exactly those paths which satisfy*

$$x_j = \frac{j}{n}$$

and

$$y_{j+1} = y_j + \chi_{jr} \quad \text{for } j \in \mathbb{Z}/n\mathbb{Z} \text{ with } \sum_{i=0}^{n-1} r \chi_i = 0 \text{ and } r \geq 0$$

for each j . In particular, if n is odd then a discrete $(n, dn - 1)$ -path must be regular.

Proof. For every i ,

$$\begin{aligned} |V_i V_{i+1}| &= |V_{i+dn-1} V_{i+dn}|, \\ |V_i V_{i+dn-1}| &= |V_{i+1} V_{i+dn}|. \end{aligned}$$

Therefore

$$\begin{aligned} (x_{i+1} - x_i)^2 + (y_{i+1} - y_i)^2 &= (x_i - x_{i-1})^2 + (y_i - y_{i-1})^2, \\ (d + x_{i-1} - x_i)^2 + (y_{i-1} - y_i)^2 &= (d + x_i - x_{i+1})^2 + (y_i - y_{i+1})^2. \end{aligned}$$

It follows that

$$d(x_{i+1} - x_i) = d(x_i - x_{i-1}).$$

Since $d \neq 0$,

$$x_{i+1} - x_i = x_i - x_{i-1}.$$

By Lemma 3, $x_j = j/n$ for each j . Now equation $|V_i V_{i+1}| = |V_j V_{j+1}|$ for all i, j implies that

$$(x_{i+1} - x_i)^2 + (y_{i+1} - y_i)^2 = (x_{j+1} - x_j)^2 + (y_{j+1} - y_j)^2 \quad \text{for all } i, j,$$




Figure 2. An example of a discrete $(6, 5)$ -path.

so that

$$(y_{i+1} - y_i)^2 = (y_{j+1} - y_j)^2.$$

By Lemma 2, we are done. □

Theorem 5. *The discrete $(n, dn + 1)$ -paths $V_i = (x_i, y_i)$, $i \in \mathbb{N}$ with $d \neq 0$ are exactly those paths which satisfy*

$$x_j = \frac{j}{n}$$

and

$$y_{j+1} = y_j + \chi_{jr} \quad \text{for } j \in \mathbb{Z}/n\mathbb{Z} \text{ with } \sum_{i=0}^{n-1} r \chi_i = 0 \text{ and } r \geq 0$$

for each j . In particular, if n is odd then a discrete $(n, dn + 1)$ -path must be regular.

Proof. Set $C_1 = |V_i V_{i+dn+1}|^2$ and $C_2 = |V_i V_{i+1}|^2$. Then

$$\begin{aligned} (d + x_{i+1} - x_i)^2 + (y_{i+1} - y_i)^2 &= C_1, \\ (x_{i+1} - x_i)^2 + (y_{i+1} - y_i)^2 &= C_2. \end{aligned}$$

Substituting, we get

$$d^2 + 2d(x_{i+1} - x_i) + C_2 = C_1,$$

so that $x_{i+1} - x_i$ is constant. By Lemma 3, $x_i = i/n$. It follows that $(y_{i+1} - y_i)^2$ is constant, so by Lemma 3 we are done. □

Corollary 6. *Any $(n, dn + 1)$ -path is an $(n, dn - 1)$ -path and vice versa.*

For an example, see Figure 2.

4. Darboux transformation and integrals

It is important to make a distinction between infinitesimal “trapezoidal” movement and infinitesimal “parallelogram” movement of the bicycle. Consider a pair of conjoined bikes, sharing a back wheel and facing in opposite directions. Since

this bicycle moves in such a way that the distance between the turnable wheels is constant, at each moment of time the turnable wheels must enclose equal angles with the line of the frame. When the two turnable wheels are parallel, the trike is gliding, but then the common back wheel of the bikes is slipping, which is not allowed. That is why we exclude parallelogram movements from consideration for the remainder of this paper.

Definition 7 (trapezoidal condition). We will say that a discrete (n, k) -path satisfies the trapezoidal condition if $V_i V_{i+k+1}$ and $V_{i+1} V_{i+k}$ are parallel for each $i \in \mathbb{Z}$.

As an illustration of these concepts, consider Figure 2: $V_0 V_1 V_5 V_6$ is a trapezoidal motion, while $V_1 V_2 V_6 V_7$ is a parallelogram motion. Consequently, the bicycle path in the figure does not satisfy the trapezoidal condition.

Assuming the trapezoidal condition, we may view bicycle paths in terms of an important construction in discrete differential geometry called the Darboux transformation [Bobenko and Suris 2008; Tsuruga 2010].

Definition 8 (Darboux transform). We say that two polygons $P = P_1 P_2 \cdots$ and $Q = Q_1 Q_2 \cdots$ are in Darboux correspondence if for each $i = 1, 2, \dots$, we have that Q_{i+1} is the reflection of P_i in the perpendicular bisector of the segment $P_{i+1} Q_i$.

If segment $P_1 Q_1$ is of length ℓ then for each i , $P_i Q_i$ is of length ℓ . We then say that P and Q are in Darboux correspondence with parameter ℓ . We also note that each quadrilateral $P_i Q_i P_{i+1} Q_{i+1}$ is an isosceles trapezoid.

We denote the map taking vertex P_i to Q_i by \mathfrak{D} . We will also refer to the map of polygons $\mathfrak{D}(P) = Q$ by the same letter, since no confusion ought to occur.

Consider a polygonal line P with vertices V_0, V_1, \dots, V_{n-1} . Let v_0 be a vector with its origin at V_0 . Having a vector v_i at vertex V_i , we obtain a vertex v_{i+1} of the same length at V_{i+1} via the trapezoidal condition. For example, in Figure 3, $v_1 = P_1 Q_1$ and $v_2 = P_2 Q_2$. For a fixed length of v_0 , we may view the map taking v_0 to v_j as a self-map of the circle of radius $|v_0| = |v_j|$ by identifying the circle at V_0 with circle at V_j via parallel translation.

Definition 9 (monodromy map of the Darboux transformation). The monodromy map is the map acting on the identified circles at V_0 and V_n which takes v_0 to v_n .

It is known that the monodromy map is a cross-ratio preserving transformation (in terms of affine coordinates, a fractional linear transformation) on a circle of fixed radius after we identify the circle with the real projective line via stereographic projection [Tabachnikov and Tsukerman 2013]. We will assume throughout, unless otherwise stated, that the monodromy map is acting on a fixed point; in other words, we will assume that the Darboux transform has been chosen so that the initial vector v_0 is equal to the vector v_n , where n is the period. This is analogous to applying the Darboux transform to a closed polygon and requiring that its image is closed also.




Figure 3. Two polygons in Darboux correspondence.

We mention in passing that in the case of closed polygons, Darboux correspondence implies that the monodromies of the two polygons are conjugated to each other. The invariants of the conjugacy class of the monodromy, viewed as functions of the length parameter, are consequently integrals of the Darboux correspondence [Tabachnikov and Tsukerman 2013].

Connection between Darboux transformation and discrete (n, k) -paths. A discrete (n, k) -path satisfying the trapezoidal condition may be interpreted in terms of the Darboux transform. Indeed, given such a path, we consider the periodic equilateral linkages $L_i = \cdots V_{0+i} V_{k+i} V_{2k+i} \cdots$ for $i = 0, 1, \dots, k - 1$. The trapezoidal condition implies that there is a Darboux correspondence $\mathfrak{D}(L_i) = L_{i+1}$ of the same parameter (since the (n, k) -path is equilateral) for consecutive linkages (see Figure 4).

The Darboux transformation also preserves the area of periodic paths. More precisely, let $y = -c$ for $c > 0$ sufficiently large so that the periodic path P and its Darboux transformation P' lie completely above $y = -c$. We define an area function as follows. Let $\check{V}_i = (x(V_i), -c)$. We define the area of P to be the signed area of the polygon $\check{V}_0 V_0 V_1 \cdots V_n \check{V}_n$ and denote it by $|P|$. We show that this area is preserved under Darboux transformation (see Figure 5). In particular, it will follow that the area of $V_0 V_k \cdots V_{nk}$ is equal to the area of $V_m V_{k+m} \cdots V_{nk+m}$ for every $m \in \mathbb{Z}$.

Theorem 10. *The Darboux transformation is area-preserving on periodic polygonal paths.*

Proof. Let P and P' be two periodic polygonal paths in Darboux correspondence. We show that the difference of the areas of P and P' is zero. We denote the vertex of P' which corresponds via the Darboux transformation to the vertex V_i in P by V'_i for each i . We have

$$|P| = \sum_{i=0}^{n-1} |\check{V}_i V_i V_{i+1} \check{V}_{i+1}|,$$




Figure 4. Viewing a discrete (n, k) -path satisfying the trapezoidal condition (top) in terms of the Darboux transformation. The path is decomposed into equilateral linkages (middle). Any two consecutive linkages are in Darboux correspondence (bottom).

and similarly for P' . Therefore

$$|P| - |P'| = \sum_{i=0}^{n-1} |\check{V}_i V_i V_{i+1} \check{V}_{i+1}| - |\check{V}'_i V'_i V'_{i+1} \check{V}'_{i+1}|.$$

From the isosceles trapezoids,

$$|V_i V_{i+1} V'_{i+1}| = |V'_i V'_{i+1} V_i|. \quad (4-1)$$

Also,

$$|\check{V}_i V_i V_{i+1} \check{V}_{i+1}| = |\check{V}_i V_i V'_{i+1} \check{V}'_{i+1}| + |\check{V}'_{i+1} V'_{i+1} V_{i+1} \check{V}_{i+1}| + |V_i V_{i+1} V'_{i+1}|.$$

Similarly,

$$|\check{V}'_i V'_i V'_{i+1} \check{V}'_{i+1}| = |\check{V}'_i V'_i V_i \check{V}_i| + |\check{V}_i V_i V'_{i+1} \check{V}'_{i+1}| + |V'_i V'_{i+1} V_i|.$$

Using (4-1),

$$|\check{V}_i V_i V_{i+1} \check{V}_{i+1}| - |\check{V}'_i V'_i V'_{i+1} \check{V}'_{i+1}| = |\check{V}'_{i+1} V'_{i+1} V_{i+1} \check{V}_{i+1}| - |\check{V}'_i V'_i V_i \check{V}_i|.$$

It follows that

$$|P| - |P'| = \sum_{i=0}^{n-1} |\check{V}'_{i+1} V'_{i+1} V_{i+1} \check{V}_{i+1}| - |\check{V}'_i V'_i V_i \check{V}_i|,$$




Figure 5. Two periodic paths P and P' in Darboux correspondence. By Theorem 10, the two paths have equal areas under the curve.

which telescopes to

$$|P| - |P'| = |\check{V}'_n V'_n \check{V}_n \check{V}'_n| - |\check{V}'_0 V'_0 \check{V}_0 \check{V}'_0|.$$

Since $V'_n = V'_0 + e_1$ and $V_n = V_0 + e_1$, it follows that $\overrightarrow{V_n V'_n} = \overrightarrow{V_0 V'_0}$ and $|\check{V}'_n V'_n \check{V}_n \check{V}'_n| = |\check{V}'_0 V'_0 \check{V}_0 \check{V}'_0|$, so that $|P| = |P'|$. \square

5. Questions

We end our discussion with some research topics and questions of interest concerning bicycle (n, k) -paths.

- (1) Construct interesting families of bicycle (n, k) -paths. For example, ones for which the condition $x_j = j/n$ does not hold.
- (2) What is the m -th order ($m \in \mathbb{N}$) infinitesimal rigidity theory of bicycle (n, k) -paths like?
- (3) For closed bicycle polygons, there are many integrals of motion [Tabachnikov and Tsukerman 2013]. For example, a geometric center called the circumcenter of mass [Tabachnikov and Tsukerman 2014] is invariant under Darboux transformation for closed polygons. Are there other integrals of motion for bicycle (n, k) -paths?

Acknowledgments

It is a pleasure to acknowledge the valuable help and discussion with S. Tabachnikov. This project originated during the program Summer@ICERM 2012; we are grateful to ICERM for the support. We are also thankful to the anonymous referee for valuable suggestions.

References

- [Adler 1993] V. È. Adler, “Recuttings of polygons”, *Funktsional. Anal. i Prilozhen.* **27**:2 (1993), 79–82. In Russian; translated in *Funct. Anal. Appl.* **27**:2 (1993), 141–143. MR 94j:58072 Zbl 0812.58072
- [Adler 1995] V. È. Adler, “Integrable deformations of a polygon”, *Phys. D* **87**:1–4 (1995), 52–57. MR 96m:58100 Zbl 1194.35353
- [Bobenko and Suris 2008] A. I. Bobenko and Y. B. Suris, *Discrete differential geometry: Integrable structure*, Graduate Studies in Mathematics **98**, American Mathematical Society, Providence, RI, 2008. MR 2010f:37125 Zbl 1158.53001
- [Connelly and Csikós 2009] R. Connelly and B. Csikós, “Classification of first-order flexible regular bicycle polygons”, *Studia Sci. Math. Hungar.* **46**:1 (2009), 37–46. MR 2011d:52042 Zbl 1240.11057
- [Csikós 2007] B. Csikós, “On the rigidity of regular bicycle (n, k) -gons”, *Contrib. Discrete Math.* **2**:1 (2007), 93–106. MR 2007m:37144 Zbl 1191.52001
- [Cyr 2012] V. Cyr, “A number theoretic question arising in the geometry of plane curves and in billiard dynamics”, *Proc. Amer. Math. Soc.* **140**:9 (2012), 3035–3040. MR 2917076 Zbl 1282.37023
- [Tabachnikov 2006] S. Tabachnikov, “Tire track geometry: variations on a theme”, *Israel J. Math.* **151** (2006), 1–28. MR 2007d:37091 Zbl 1124.52005
- [Tabachnikov and Tsukerman 2013] S. Tabachnikov and E. Tsukerman, “On the discrete bicycle transformation”, *Publ. Mat. Urug.* **14** (2013), 201–219. MR 3235356 Zbl 1317.51027
- [Tabachnikov and Tsukerman 2014] S. Tabachnikov and E. Tsukerman, “Circumcenter of mass and generalized Euler line”, *Discrete Comput. Geom.* **51** (2014), 815–836. MR 3216665 Zbl 1301.51023
- [Tsuruga 2010] M. Tsuruga, “Discrete Differential Geometry”, Lecture notes, Freie Universität, 2010, available at <http://boolesrings.org/matsguru/files/2012/11/DDG1.pdf>.
- [Wegner 2007] F. Wegner, “Floating bodies of equilibrium in 2D, the tire track problem and electrons in a parabolic magnetic field”, preprint, 2007. arXiv physics/0701241

Received: 2013-09-07 Revised: 2014-09-01 Accepted: 2014-12-08

ian_alevy@brown.edu

*Division of Applied Mathematics, Brown University,
Providence, RI 02912, United States*

e.tsukerman@berkeley.edu

*Department of Mathematics, University of California, Berkeley,
Berkeley, CA 94720, United States*

MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	La Trobe University, Australia P.Cerone@latrobe.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moselehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyjs@potsdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemmons	Wake Forest University, USA rplemmons@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA jgallian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sngupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2016 is US \$160/year for the electronic version, and \$215/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

involve

2016

vol. 9

no. 1

Using ciliate operations to construct chromosome phylogenies JACOB L. HERLIN, ANNA NELSON AND MARION SCHEEPERS	1
On the distribution of the greatest common divisor of Gaussian integers TAI-DANAE BRADLEY, YIN CHOI CHENG AND YAN FEI LUO	27
Proving the pressing game conjecture on linear graphs ELIOT BIXBY, TOBY FLINT AND ISTVÁN MIKLÓS	41
Polygonal bicycle paths and the Darboux transformation IAN ALEVY AND EMMANUEL TSUKERMAN	57
Local well-posedness of a nonlocal Burgers' equation SAM GOODCHILD AND HANG YANG	67
Investigating cholera using an SIR model with age-class structure and optimal control K. RENEE FISTER, HOLLY GAFF, ELSA SCHAEFER, GLENNA BUFORD AND BRYCE C. NORRIS	83
Completions of reduced local rings with prescribed minimal prime ideals SUSAN LOEPP AND BYRON PERPETUA	101
Global regularity of chemotaxis equations with advection SAAD KHAN, JAY JOHNSON, ELLIOT CARTEE AND YAO YAO	119
On the ribbon graphs of links in real projective space IAIN MOFFATT AND JOHANNA STRÖMBERG	133
Depths and Stanley depths of path ideals of spines DANIEL CAMPOS, RYAN GUNDERSON, SUSAN MOREY, CHELSEY PAULSEN AND THOMAS POLSTRA	155
Combinatorics of linked systems of quartet trees EMILI MOAN AND JOSEPH RUSINKO	171

