

# involve

a journal of mathematics

On weak lattice point visibility

Neil R. Nicholson and Rebecca Rachan





# On weak lattice point visibility

Neil R. Nicholson and Rebecca Rachan

(Communicated by John C. Wierman)

We say that a point  $Q$  in a specific rectangular array of lattice points is weakly visible from a lattice point  $P$  not in the array if no point in the array other than  $Q$  lies on the line connecting the external point  $P$  to  $Q$ . A necessary and sufficient condition for determining if a point in the array is weakly viewable by the external point, as well as the number of points that are weakly visible from the external point, is determined.

## 1. Introduction

Imagine a photographer attempting to capture a picture in which every member of a band in a rectangular array formation is visible, with all persons, including the photographer, standing on lattice points. The photographer must stand at a fixed position, and each band member must have a straight-line view of the photographer, unobstructed by all other band members. Laison and Schick [2007] describe this situation and prove that there are positions for the photographer to stand but these may be quite far away from the marching band. If the photographer decides to stand closer, how can we determine which band members she can see?

These are examples of the questions arising in lattice point visibility that have been investigated for decades. For example, another question that has gotten much attention considers two sets  $A$  and  $B$  of lattice points. When is every point in  $A$  visible from every point in  $B$ ? Are there relationships between the sizes of the sets when this is the case, and if so, as the set  $B$  grows does its size act predictably? Much work has been done looking at questions such as these [Adhikari and Granville 2009; Adhikari and Balasubramanian 1996; Adhikari and Chen 1999; 2002; Chen and Cheng 2003; Herzog and Stewart 1971].

Here, we fix this second set to be a single point  $P$ , playing the role of the photographer trying to see the members of the marching band, those points in set  $A$ . In [Nicholson and Sharp 2010], a lower bound was placed on the distance  $P$  must be from a rectangular array of lattice points to weakly view every point in the array.

---

*MSC2010:* 11H06.

*Keywords:* weak visibility, lattice point.

Our main result can be used to prove this result in an alternate manner and provides another tool to address one of the primary questions in weak visibility: is there a formula for this minimum distance only dependent upon the dimensions of the lattice points?

## 2. Definitions

In this paper, all points are assumed to be lattice points in the first quadrant. Let  $m, n \in \mathbb{Z}^+$  with  $n \leq m$ . Define  $\Delta_{m,n} = \{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$ . We say  $Q \in \Delta_{m,n}$  is *weakly visible* from a point  $P \notin \Delta_{m,n}$  if no other point in  $\Delta_{m,n}$  lies on the segment  $PQ$ .

It was proven in [Nicholson and Sharp 2010] that the  $m \times m$  square of lattice points with its lower left-hand corner on  $(m, n)$  (that is, the square of lattice points with corners  $(m, n)$ ,  $(m, n + m - 1)$ ,  $(2m - 1, n)$ , and  $(2m - 1, n + m - 1)$ ), called the *adjacency square to  $\Delta_{m,n}$*  and denoted  $\text{Adj}_{m,n}$ , contains no point that weakly views every point in  $\Delta_{m,n}$ . As a corollary to this, there is a lower bound that can be placed on how close a point viewing every point in  $\Delta_{m,n}$  can be to  $\Delta_{m,n}$ :

**Theorem 2.1** [Nicholson and Sharp 2010]. *If a point  $P$  weakly views every point in  $\Delta_{m,n}$ , then  $P$  is at least  $\sqrt{m^2 + 1}$  units from  $(m, n)$ .*

What follows is a complete classification of which points in  $\Delta_{m,n}$  are weakly visible from a general point  $P \notin \Delta_{m,n}$  as well as two corollaries that follow from that classification.

## 3. Determining weak visibility

We begin this section with our main result: necessary and sufficient conditions for a point  $Q \in \Delta_{m,n}$  to be weakly visible from a point  $P \notin \Delta_{m,n}$ .

**Theorem 3.1.** *The point  $Q = (x_0, y_0) \in \Delta_{m,n}$  is not weakly visible by the point  $P = (a, b)$  if and only if all of the following conditions hold:*

- (1)  $\gcd(a - x_0, b - y_0) > 1$ .
- (2)  $m - x_0 \geq (a - x_0) / \gcd(a - x_0, b - y_0)$ .
- (3)  $n - y_0 \geq (b - y_0) / \gcd(a - x_0, b - y_0)$ .

*Proof.* Suppose  $Q$  is not weakly viewable by  $P$ . Then, there exist  $t \geq 1$  points on the interior of the segment  $PQ$ , and let  $R = (x_1, y_1)$  be the first of these points to the right of  $Q$ . Thus,

$$\begin{aligned} b - y_0 &= (y_1 - y_0)(t + 1), \\ a - x_0 &= (x_1 - x_0)(t + 1), \end{aligned} \tag{1}$$

implying that

$$\begin{aligned} \gcd(a - x_0, b - y_0) &\geq t + 1 \\ &> 1. \end{aligned} \tag{2}$$

Moreover, in order to have  $(x_1, y_1) \in \Delta_{m,n}$ , we have

$$\begin{aligned} m - x_0 &\geq x_1 - x_0 \\ &= \frac{a - x_0}{t + 1} \\ &\geq \frac{a - x_0}{\gcd(a - x_0, b - y_0)}, \end{aligned} \tag{3}$$

with the third property following similarly.

Now, assume the three properties hold and  $d = \gcd(a - x_0, b - y_0) > 1$ , with

$$\begin{aligned} a - x_0 &= dp, \\ b - y_0 &= dq. \end{aligned} \tag{4}$$

We claim that  $(x_0 + p, y_0 + q) \in \Delta_{m,n}$  lies on the segment  $PQ$ . To see this, note that the slope of  $PQ$  is  $q/p$ , so that  $PQ$  has equation

$$y - y_0 = \frac{q}{p}(x - x_0). \tag{5}$$

The point  $(x_0 + p, y_0 + q)$  satisfies this equation, and

$$\begin{aligned} x_0 + p &= x_0 + \frac{a - x_0}{d} \\ &\leq x_0 + m - x_0 \\ &= m. \end{aligned} \tag{6}$$

Similarly,  $y_0 + q \leq n$ , showing  $(x_0 + p, y_0 + q) \in \Delta_{m,n}$  and consequently that  $Q$  is not weakly viewable by  $P$ .  $\square$

What points then can a particularly chosen external point  $P$  weakly view? It is only natural to insist  $P$  lies strictly above the line  $y = n$  and to the right of the line  $x = m$ . Thus, the closest such point (with distance measured to the nearest point,  $(m, n)$ , in  $\Delta_{m,n}$ ) would be  $P = (m + 1, n + 1)$ . The aforementioned result from [Nicholson and Sharp 2010] guarantees  $P$  cannot weakly view every point in  $\Delta_{m,n}$  (for sufficiently large values of  $m$  and  $n$ ). Which points then can  $P$  weakly view? Corollary 3.2 follows immediately from Theorem 3.1.

**Corollary 3.2.** *The point  $(x, 1) \in \Delta_{m,n}$  is weakly viewable by the point  $P = (m + 1, n + 1)$  if and only if  $\gcd((m + 1) - x, n) = 1$ .*

This corollary states that the number of points in the first row of  $\Delta_{m,n}$  that are weakly visible by  $P$  is the number of positive integers less than or equal to  $m$  that

are relatively prime to  $n$ . This is a variation of the *Euler totient function* (or *Euler phi function*,  $\phi(m)$ ), defined on positive integers  $m$  as the number of positive integers less than or equal to  $m$  that are relatively prime to  $m$ ). For  $n \leq m$ , call this  $\phi(n, m)$ , precisely the number of points in the first row of  $\Delta_{m,n}$  weakly viewable by this particular point  $P$ . This allows us to count the total number of points in  $\Delta_{m,n}$  that  $P$  weakly views:

**Corollary 3.3.** *The number of points of  $\Delta_{m,n}$  weakly viewable by the point  $P = (m + 1, n + 1)$  is  $\sum_{i=1}^n \phi(i, m)$ .*

*Proof.* The number of points in the  $j$ -th row of  $\Delta_{m,n}$  that are weakly viewable by  $P$  is  $\phi(n - j + 1, m)$ .  $\square$

We conclude by noting that the main question, amongst numerous other interesting questions, related to the results here remains open. Is there a formula dependent only upon  $m$  and  $n$  for the point closest to  $\Delta_{m,n}$  that weakly views every point of  $\Delta_{m,n}$ ? Such a formula would lend itself not only to deeper development in other lattice point visibility questions and graph theory but would potentially have applications in a multitude of fields [Ghosh and Goswami 2013].

## References

- [Adhikari and Balasubramanian 1996] S. D. Adhikari and R. Balasubramanian, “On a question regarding visibility of lattice points”, *Mathematika* **43**:1 (1996), 155–158. MR 97k:11105 Zbl 0855.11009
- [Adhikari and Chen 1999] S. D. Adhikari and Y.-G. Chen, “On a question regarding visibility of lattice points, II”, *Acta Arith.* **89**:3 (1999), 279–282. MR 2000i:11152 Zbl 0936.11039
- [Adhikari and Chen 2002] S. D. Adhikari and Y.-G. Chen, “On a question regarding visibility of lattice points, III”, *Discrete Math.* **259**:1-3 (2002), 251–256. MR 2004a:11107 Zbl 1033.11049
- [Adhikari and Granville 2009] S. D. Adhikari and A. Granville, “Visibility in the plane”, *J. Number Theory* **129**:10 (2009), 2335–2345. MR 2010m:11117 Zbl 1176.11027
- [Chen and Cheng 2003] Y.-G. Chen and L.-F. Cheng, “Visibility of lattice points”, *Acta Arith.* **107**:3 (2003), 203–207. MR 2004g:11053 Zbl 1116.11048
- [Ghosh and Goswami 2013] S. K. Ghosh and P. P. Goswami, “Unsolved problems in visibility graphs of points, segments and polygons”, *ACM Comput. Surv.* **46**:2 (2013), 22:1–22:29. Zbl 1288.05056
- [Herzog and Stewart 1971] F. Herzog and B. M. Stewart, “Patterns of visible and nonvisible lattice points”, *Amer. Math. Monthly* **78** (1971), 487–496. MR 44 #1630 Zbl 0217.03501
- [Laison and Schick 2007] J. D. Laison and M. Schick, “Seeing dots: visibility of lattice points”, *Math. Mag.* **80**:4 (2007), 274–282. MR 2008j:11079 Zbl 1208.11082
- [Nicholson and Sharp 2010] N. Nicholson and R. Sharp, “Weakly viewing lattice points”, *Involve J. of Math.* **3**:1 (2010), 9–16. Zbl 1269.11064

Received: 2014-12-01      Revised: 2015-03-20      Accepted: 2015-04-05

nrnicholson@noctrl.edu

*Department of Mathematics, North Central College,  
30 North Brainard Street, Naperville, IL 60540, United States*

rarachan@noctrl.edu

*Department of Mathematics, North Central College,  
30 North Brainard Street, Naperville, IL 60540, United States*

# involve

msp.org/involve

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

|                      |   |                        |   |
|----------------------|---|------------------------|---|
| Colin Adams          | Williams College, USA                     | Suzanne Lenhart        | University of Tennessee, USA              |
| John V. Baxley       | Wake Forest University, NC, USA           | Chi-Kwong Li           | College of William and Mary, USA          |
| Arthur T. Benjamin   | Harvey Mudd College, USA                  | Robert B. Lund         | Clemson University, USA                   |
| Martin Bohner        | Missouri U of Science and Technology, USA | Gaven J. Martin        | Massey University, New Zealand            |
| Nigel Boston         | University of Wisconsin, USA              | Mary Meyer             | Colorado State University, USA            |
| Amarjit S. Budhiraja | U of North Carolina, Chapel Hill, USA     | Emil Minchev           | Ruse, Bulgaria                            |
| Pietro Cerone        | La Trobe University, Australia            | Frank Morgan           | Williams College, USA                     |
| Scott Chapman        | Sam Houston State University, USA         | Mohammad Sal Moslehian | Ferdowsi University of Mashhad, Iran      |
| Joshua N. Cooper     | University of South Carolina, USA         | Zuhair Nashed          | University of Central Florida, USA        |
| Jem N. Corcoran      | University of Colorado, USA               | Ken Ono                | Emory University, USA                     |
| Toka Diagana         | Howard University, USA                    | Timothy E. O'Brien     | Loyola University Chicago, USA            |
| Michael Dorff        | Brigham Young University, USA             | Joseph O'Rourke        | Smith College, USA                        |
| Sever S. Dragomir    | Victoria University, Australia            | Yuval Peres            | Microsoft Research, USA                   |
| Behrouz Emamizadeh   | The Petroleum Institute, UAE              | Y.-F. S. Pétermann     | Université de Genève, Switzerland         |
| Joel Foisy           | SUNY Potsdam, USA                         | Robert J. Plemmons     | Wake Forest University, USA               |
| Errin W. Fulp        | Wake Forest University, USA               | Carl B. Pomerance      | Dartmouth College, USA                    |
| Joseph Gallian       | University of Minnesota Duluth, USA       | Vadim Ponomarenko      | San Diego State University, USA           |
| Stephan R. Garcia    | Pomona College, USA                       | Bjorn Poonen           | UC Berkeley, USA                          |
| Anant Godbole        | East Tennessee State University, USA      | James Propp            | U Mass Lowell, USA                        |
| Ron Gould            | Emory University, USA                     | József H. Przytycki    | George Washington University, USA         |
| Andrew Granville     | Université Montréal, Canada               | Richard Rebarber       | University of Nebraska, USA               |
| Jerrold Griggs       | University of South Carolina, USA         | Robert W. Robinson     | University of Georgia, USA                |
| Sat Gupta            | U of North Carolina, Greensboro, USA      | Filip Saidak           | U of North Carolina, Greensboro, USA      |
| Jim Haglund          | University of Pennsylvania, USA           | James A. Sellers       | Penn State University, USA                |
| Johnny Henderson     | Baylor University, USA                    | Andrew J. Sterge       | Honorary Editor                           |
| Jim Hoste            | Pitzer College, USA                       | Ann Trenk              | Wellesley College, USA                    |
| Natalia Hritonenko   | Prairie View A&M University, USA          | Ravi Vakil             | Stanford University, USA                  |
| Glenn H. Hurlbert    | Arizona State University, USA             | Antonia Vecchio        | Consiglio Nazionale delle Ricerche, Italy |
| Charles R. Johnson   | College of William and Mary, USA          | Ram U. Verma           | University of Toledo, USA                 |
| K. B. Kulasekera     | Clemson University, USA                   | John C. Wierman        | Johns Hopkins University, USA             |
| Gerry Ladas          | University of Rhode Island, USA           | Michael E. Zieve       | University of Michigan, USA               |

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2016 is US \$160/year for the electronic version, and \$215/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

*Involve* (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# involve

2016

vol. 9

no. 3

|  |     |
|--|-----|
| A combinatorial proof of a decomposition property of reduced residue systems         | 361 |
| YOTSANAN MEEMARK AND THANAKORN PRINYASART  |     |
| Strong depth and quasigeodesics in finitely generated groups                         | 367 |
| BRIAN GAPINSKI, MATTHEW HORAK AND TYLER WEBER  |     |
| Generalized factorization in $\mathbb{Z}/m\mathbb{Z}$                                | 379 |
| AUSTIN MAHLUM AND CHRISTOPHER PARK MOONEY  |     |
| Cocircular relative equilibria of four vortices                                      | 395 |
| JONATHAN GOMEZ, ALEXANDER GUTIERREZ, JOHN LITTLE,<br>ROBERTO PELAYO AND JESSE ROBERT |     |
| On weak lattice point visibility   | 411 |
| NEIL R. NICHOLSON AND REBECCA RACHAN   |     |
| Connectivity of the zero-divisor graph for finite rings                              | 415 |
| REZA AKHTAR AND LUCAS LEE  |     |
| Enumeration of $m$ -endomorphisms  | 423 |
| LOUIS RUBIN AND BRIAN RUSHTON  |     |
| Quantum Schubert polynomials for the $G_2$ flag manifold                             | 437 |
| RACHEL E. ELLIOTT, MARK E. LEWERS AND LEONARDO C.<br>MIHALCEA                        |     |
| The irreducibility of polynomials related to a question of Schur                     | 453 |
| LENNY JONES AND ALICIA LAMARCHE  |     |
| Oscillation of solutions to nonlinear first-order delay differential equations       | 465 |
| JAMES P. DIX AND JULIO G. DIX  |     |
| A variational approach to a generalized elastica problem                             | 483 |
| C. ALEX SAFSTEN AND LOGAN C. TATHAM  |     |
| When is a subgroup of a ring an ideal?   | 503 |
| SUNIL K. CHEBOLU AND CHRISTINA L. HENRY  |     |
| Explicit bounds for the pseudospectra of various classes of matrices and operators   | 517 |
| FEIXUE GONG, OLIVIA MEYERSON, JEREMY MEZA, MIHAI<br>STOICIU AND ABIGAIL WARD         |     |

