

# involve

a journal of mathematics

## Prime vertex labelings of several families of graphs

Nathan Diefenderfer, Dana C. Ernst, Michael G. Hastings,  
Levi N. Heath, Hannah Prawzinsky, Briahna Preston,  
Jeff Rushall, Emily White and Alyssa Whittemore





# Prime vertex labelings of several families of graphs

Nathan Diefenderfer, Dana C. Ernst, Michael G. Hastings,  
Levi N. Heath, Hannah Prawzinsky, Briahna Preston,  
Jeff Rushall, Emily White and Alyssa Whittlemore

(Communicated by Joseph A. Gallian)

A simple and connected  $n$ -vertex graph has a prime vertex labeling if the vertices can be injectively labeled with the integers  $1, 2, 3, \dots, n$  such that adjacent vertices have relatively prime labels. We will present previously unknown prime vertex labelings for new families of graphs, including cycle pendant stars, cycle chains, prisms, and generalized books.

## 1. Introduction

The focus of this paper is prime vertex labelings, wherein adjacent vertices of simple, connected graphs are assigned integer labels that are relatively prime. Currently, the two most prominent open conjectures involving prime vertex labelings are:

- All tree graphs have a prime vertex labeling (Entringer–Tout conjecture).
- All unicyclic graphs have a prime vertex labeling [Seoud and Youssef 1999].

While we will address one infinite family that is unicyclic, our primary concern will be nonunicyclic graphs.

A *graph*  $G$  is a set of vertices,  $V(G)$ , together with a set of edges,  $E(G)$ , connecting some subset, possibly empty, of the vertices. If  $u, v \in V(G)$  are connected by an edge, we say  $u$  and  $v$  are *adjacent*. The *degree* of a vertex  $u$  is the number of edges incident to  $u$ . A *subgraph*  $H$  of a graph  $G$  is a graph whose vertex set is a subset of that of  $G$ , and whose adjacency relation is a subset of that of  $G$  restricted to this subset.


We will restrict our attention to graphs that are simple (i.e., graphs that do not have multiple edges between pairs of vertices nor edges that connect a vertex to itself) and connected (i.e., graphs that do not consist of two or more disjoint “pieces”). For the remainder of this paper, all graphs are assumed to be simple and connected.

---

*MSC2010:* 05C78.

*Keywords:* graph labeling, prime vertex labeling, prime graphs.

This research was supported by the National Science Foundation grant #DMS-1148695 through the Center for Undergraduate Research in Mathematics (CURM).



**Figure 1.** The star  $S_4$ .

Next, we define a few important families of graphs. For  $n \geq 2$ , an  $n$ -*path* (or simply *path*), denoted  $P_n$ , is a connected graph consisting of two vertices of degree 1 and  $n - 2$  vertices of degree 2. For  $n \geq 3$ , an  $n$ -*cycle* (or simply *cycle*), denoted  $C_n$ , is a connected graph consisting of  $n$  vertices, each of degree 2. Note that both  $P_n$  and  $C_n$  have  $n$  vertices, while  $P_n$  has  $n - 1$  edges and  $C_n$  has  $n$  edges. An  $n$ -*star* (or simply *star*), denoted  $S_n$ , is a graph consisting of one vertex of degree  $n$ , called the *center*, and  $n$  vertices of degree 1. Note that  $S_n$  consists of  $n + 1$  vertices and  $n$  edges. The star  $S_4$  is shown in Figure 1.


A simple and connected  $n$ -vertex graph is said to have a *prime vertex labeling* if the vertices can be injectively labeled with the integers  $1, 2, 3, \dots, n$  such that adjacent vertices have relatively prime labels. For brevity, if a graph has a prime vertex labeling, we will say that the graph is *prime*. The many familiar families of graphs that are known to be prime include paths, cycles, and stars.

In this paper, we will present previously unknown prime vertex labelings for several infinite families of graphs, including cycle pendant stars (Section 2), cycle chains (Section 3), prisms (Section 4), and generalized books (Section 5). Finally, some conjectures and potential future work will be described in Section 6.

## 2. Cycle pendant stars


The focus of this section is on a type of unicyclic graph (i.e., a graph containing exactly one cycle as a subgraph), which was inspired by Seoud and Youssef's conjecture [1999] that all unicyclic graphs are prime. Instead of attempting to prove the conjecture outright, which we anticipate would require advanced machinery from linear algebra, most authors concentrate on finding prime labelings for specific families of unicyclic graphs. In particular, this was our endeavor in [Diefenderfer et al. 2015].

Every vertex lying on the cycle of a unicyclic graph will be referred to as a *cycle vertex*. In a unicyclic graph, a *spur* is an edge with exactly one vertex on the cycle. The noncycle vertex of a spur is called a *spur vertex*. For example, the graph shown in Figure 2 is a unicyclic graph with five spurs. In this case, the vertices labeled by  $c_1, c_2, c_3$ , and  $c_4$  are cycle vertices, while the vertices labeled by  $p_1, p_2, p_3, p_4$ , and  $p_5$  are spur vertices.




**Figure 2.** Example of a unicyclic graph with five spurs.

Seoud and Youssef [1999] investigated cycles with identical complete binary trees attached to each cycle vertex. These unicyclic graphs can also be viewed as cycles with various levels of the star  $S_2$  attached to one another. This observation led to the following generalization, which features a single “level” of stars, but with differing star sizes. A *cycle pendant star*, denoted  $C_n \star P_2 \star S_m$ , is the graph that results from attaching the path  $P_2$  to each vertex of  $C_n$  followed by attaching the star  $S_m$  at its center to each spur vertex. For example, the graph  $C_5 \star P_2 \star S_6$  is shown in Figure 3. Note that our  $\star$  notation is not a construction typically found in the literature and refers to “selectively gluing” copies of one graph to another.



**Figure 3.** The graph  $C_5 \star P_2 \star S_6$ .



**Figure 4.** The generalized prime vertex labeling of  $C_n \star P_2 \star S_4$ .

**Theorem 2.1.** All  $C_n \star P_2 \star S_m$  with  $0 \leq m \leq 8$  are prime.

*Proof.* The cases involving  $m = 0, 1, 2, 3$  correspond to familiar graphs with known prime vertex labelings. Seoud and Youssef [1999] showed that pendant graphs ( $m = 0$ ), double pendant graphs ( $m = 1$ ), and graphs with identical complete binary trees attached to the spur vertices of a pendant graph are prime (which includes the case  $m = 2$ ). Diefenderfer et al. [2015] showed that  $C_n \star P_2 \star S_3$  has a prime vertex labeling.

Now, consider  $C_n \star P_2 \star S_m$ , with  $4 \leq m \leq 8$ . Let  $c_1, c_2, \dots, c_n$  denote the consecutive cycle vertices of  $C_n$ , let  $p_i$  denote the spur vertex adjacent to  $c_i$ , and let  $o_{i,k}$ , with  $1 \leq k \leq m$ , denote the outer vertices adjacent to  $p_i$ . For example, see the identification of vertices depicted in Figure 3. For each case, it is straightforward and routine to verify that all adjacent vertices have relatively prime labels. However, for the reader’s benefit, we will describe the case involving  $m = 6$  in detail. Similar reasoning is required for each of the remaining cases.

For the  $m = 4$  case, the labeling function  $f : V \rightarrow \{1, 2, \dots, 6n\}$  is given by

$$\begin{aligned}
 f(c_i) &= 6i - 5 && \text{if } 1 \leq i \leq n, \\
 f(p_i) &= 6i - 1 && \text{if } 1 \leq i \leq n, \\
 f(o_{i,1}) &= 6i - 2 && \text{if } 1 \leq i \leq n, \\
 f(o_{i,2}) &= 6i - 3 && \text{if } 1 \leq i \leq n, \\
 f(o_{i,3}) &= 6i - 4 && \text{if } 1 \leq i \leq n, \\
 f(o_{i,4}) &= 6i && \text{if } 1 \leq i \leq n.
 \end{aligned}$$

Figure 4 depicts the generalized labeling function in this case for  $C_n \star P_2 \star S_4$ . The prime vertex labeling of  $C_4 \star P_2 \star S_4$  using this labeling appears in Figure 5.


For the  $m = 5$  case, the labeling function  $f : V \rightarrow \{1, 2, \dots, 7n\}$  is given by

$$\begin{aligned}
 f(c_i) &= 7i - 6 \quad \text{if } 1 \leq i \leq n, \\
 f(p_i) &= \begin{cases} 7i - 2 & \text{if } i \equiv_6 1, 3, \\ 7i - 3 & \text{if } i \equiv_6 2, 4, \\ 7i - 4 & \text{if } i \equiv_6 5, \\ 7i - 5 & \text{if } i \equiv_6 0, i \not\equiv_{30} 0, \\ 7i - 1 & \text{if } i \equiv_{30} 0, \end{cases} \\
 f(o_{i,1}) &= \begin{cases} 7i - 5 & \text{if } i \not\equiv_6 0 \text{ or } i \equiv_{30} 0, \\ 7i - 4 & \text{if } i \equiv_6 0, i \not\equiv_{30} 0, \end{cases} \\
 f(o_{i,2}) &= \begin{cases} 7i - 4 & \text{if } i \not\equiv_6 0, 5 \text{ or } i \equiv_{30} 0, \\ 7i - 3 & \text{if } i \equiv_6 5 \text{ or } i \equiv_6 0, i \not\equiv_{30} 0, \end{cases} \\
 f(o_{i,3}) &= \begin{cases} 7i - 3 & \text{if } i \not\equiv_6 0, 2, 4, 5 \text{ or } i \equiv_{30} 0, \\ 7i - 2 & \text{if } i \not\equiv_6 1, 3, \end{cases} \\
 f(o_{i,4}) &= \begin{cases} 7i - 2 & \text{if } i \equiv_{30} 0, \\ 7i - 1 & \text{if } i \not\equiv_{30} 0, \end{cases} \\
 f(o_{i,5}) &= 7i \quad \text{if } 1 \leq i \leq n.
 \end{aligned}$$

As an example, the prime vertex labeling of  $C_3 \star P_2 \star S_5$  using this labeling appears in Figure 6.

For  $m = 6$ , the labeling function  $f : V \rightarrow \{1, 2, \dots, 8n\}$  is given by

$$\begin{aligned}
 f(c_i) &= 8i - 7 \quad \text{if } 1 \leq i \leq n, \\
 f(p_i) &= \begin{cases} 8i - 3 & \text{if } i \not\equiv_3 0, \\ 8i - 5 & \text{if } i \equiv_3 0, i \not\equiv_{15} 0, \\ 8i - 1 & \text{if } i \equiv_{15} 0, \end{cases} \\
 f(o_{i,1}) &= 8i - 6 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,2}) &= \begin{cases} 8i - 5 & \text{if } i \not\equiv_3 0 \text{ or } i \equiv_{15} 0, \\ 8i - 3 & \text{if } i \equiv_3 0, i \not\equiv_{15} 0, \end{cases} \\
 f(o_{i,3}) &= 8i - 4 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,4}) &= \begin{cases} 8i - 3 & \text{if } i \equiv_{15} 0, \\ 8i - 1 & \text{if } i \not\equiv_{15} 0, \end{cases} \\
 f(o_{i,5}) &= 8i - 2 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,6}) &= 8i \quad \text{if } 1 \leq i \leq n.
 \end{aligned}$$



**Figure 5.** A prime vertex labeling of  $C_4 \star P_2 \star S_4$ .

As an example, the prime vertex labeling of  $C_5 \star P_2 \star S_6$  using this labeling appears in Figure 7. When  $m = 6$ , the cycle vertex labels will be consecutive integers congruent to 1 modulo 8. So, each pair of adjacent labels are odd integers that differ by 8 and hence are relatively prime. The remaining adjacencies to consider involve three cases. These cases refer to the three possible labelings for the spur vertices and within each case there are seven adjacencies to check.

**Case 1.** Assume  $i \not\equiv_3 0$ , so that  $f(p_i) = 8i - 3$ , and refer to Figure 8 for the corresponding labeling. Then

- (1)  $8i - 7$  and  $8i - 3$  are odd integers that differ by 4;
- (2)  $8i - 3$  and  $8i - 6$  are not multiples of 3 and differ by 3;
- (3)  $8i - 3$  and  $8i - 5$  are consecutive odd integers;
- (4)  $8i - 3$  and  $8i - 4$  are consecutive integers;
- (5)  $8i - 3$  and  $8i - 2$  are consecutive integers;
- (6)  $8i - 3$  and  $8i - 1$  are consecutive odd integers;
- (7)  $8i - 3$  and  $8i$  are not multiples of 3 and differ by 3.

**Case 2.** Next, assume that  $i \equiv_3 0$  and  $i \not\equiv_{15} 0$ , so that  $f(p_i) = 8i - 5$ , and refer to Figure 9 for the corresponding labeling. Then

- (1)  $8i - 7$  and  $8i - 5$  are consecutive odd integers;
- (2)  $8i - 5$  and  $8i - 6$  are consecutive integers;
- (3)  $8i - 5$  and  $8i - 4$  are consecutive integers;
- (4)  $8i - 5$  and  $8i - 3$  are consecutive odd integers;
- (5)  $8i - 5$  and  $8i - 2$  are not multiples of 3 and differ by 3;
- (6)  $8i - 5$  and  $8i - 1$  are odd integers that differ by 4;
- (7)  $8i - 5$  and  $8i$  are not multiples of 5 and differ by 5.






Figure 6. A prime vertex labeling of  $C_3 \star P_2 \star S_5$ .





Figure 7. A prime vertex labeling of  $C_5 \star P_2 \star S_6$ .


**Case 3.** Lastly, assume  $i \equiv_{15} 0$ , so that  $f(p_i) = 8i - 1$ , and refer to Figure 10 for the corresponding labeling. Then

- (1)  $8i - 7$  and  $8i - 1$  are odd integers that are not multiples of 3 and differ by 6;
- (2)  $8i - 1$  and  $8i - 6$  are not multiples of 5 and differ by 5;
- (3)  $8i - 1$  and  $8i - 5$  are odd integers that differ by 4;
- (4)  $8i - 1$  and  $8i - 4$  are not multiples of 3 and differ by 3;
- (5)  $8i - 1$  and  $8i - 3$  are consecutive odd integers;
- (6)  $8i - 1$  and  $8i - 2$  are consecutive integers;
- (7)  $8i - 1$  and  $8i$  are consecutive integers.


In all cases, adjacent labels are relatively prime, showing that  $C_n \star P_2 \star S_6$  is prime.




**Figure 8.** The generalized prime vertex labeling of  $C_n \star P_2 \star S_6$  for  $i \not\equiv_3 0$  (Case 1).



**Figure 9.** The generalized prime vertex labeling of  $C_n \star P_2 \star S_6$  for  $i \equiv_3 0$  and  $i \not\equiv_{15} 0$  (Case 2).



**Figure 10.** The generalized prime vertex labeling of  $C_n \star P_2 \star S_6$  for  $i \equiv_{15} 0$  (Case 3).




**Figure 11.** A prime vertex labeling of  $C_4 \star P_2 \star S_7$ .

Next, for  $m = 7$ , the labeling function  $f : V \rightarrow \{1, 2, \dots, 9n\}$  is given by

$$\begin{aligned}
 f(c_i) &= 9i - 8 \quad \text{if } 1 \leq i \leq n, \\
 f(p_i) &= \begin{cases} 9i - 4 & \text{if } i \equiv_2 1, \\ 9i - 5 & \text{if } i \equiv_2 0, i \not\equiv_{10} 0, \\ 9i - 7 & \text{if } i \equiv_{10} 0, i \not\equiv_{70} 0, \\ 9i - 1 & \text{if } i \equiv_{70} 0, \end{cases} \\
 f(o_{i,1}) &= \begin{cases} 9i - 7 & \text{if } i \not\equiv_{10} 0 \text{ or } i \equiv_{70} 0, \\ 9i - 5 & \text{if } i \equiv_{10} 0, i \not\equiv_{70} 0, \end{cases} \\
 f(o_{i,2}) &= 9i - 6 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,3}) &= \begin{cases} 9i - 5 & \text{if } i \equiv_2 1 \text{ or } i \equiv_{70} 0, \\ 9i - 4 & \text{if } i \equiv_2 0, i \not\equiv_{70} 0, \end{cases} \\
 f(o_{i,4}) &= \begin{cases} 9i - 4 & \text{if } i \equiv_{70} 0, \\ 9i - 1 & \text{if } i \not\equiv_{70} 0, \end{cases} \\
 f(o_{i,5}) &= 9i - 3 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,6}) &= 9i - 2 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,7}) &= 9i \quad \text{if } 1 \leq i \leq n.
 \end{aligned}$$

Figure 11 shows the prime vertex labeling of  $C_4 \star P_2 \star S_7$  using this labeling.




**Figure 12.** A prime vertex labeling of  $C_3 \star P_2 \star S_8$ .

Finally, for the  $m = 8$  case, the labeling function  $f : V \rightarrow \{1, 2, \dots, 10n\}$  is given by

$$\begin{aligned}
 f(c_i) &= 10i - 9 \quad \text{if } 1 \leq i \leq n, \\
 f(p_i) &= \begin{cases} 10i - 3 & \text{if } i \not\equiv_3 0, \\ 10i - 7 & \text{if } i \equiv_3 0, i \not\equiv_{21} 0, \\ 10i - 1 & \text{if } i \equiv_{21} 0, \end{cases} \\
 f(o_{i,1}) &= 10i - 8 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,2}) &= \begin{cases} 10i - 7 & \text{if } i \equiv_{21} 0 \text{ or } i \equiv_3 1, 2, \\ 10i - 3 & \text{if } i \equiv_3 0, i \not\equiv_{21} 0, \end{cases} \\
 f(o_{i,3}) &= 10i - 6 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,4}) &= 10i - 5 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,5}) &= 10i - 4 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,6}) &= \begin{cases} 10i - 3 & \text{if } i \equiv_{21} 0, \\ 10i - 1 & \text{if } i \not\equiv_{21} 0, \end{cases} \\
 f(o_{i,7}) &= 10i - 2 \quad \text{if } 1 \leq i \leq n, \\
 f(o_{i,8}) &= 10i \quad \text{if } 1 \leq i \leq n.
 \end{aligned}$$

The prime vertex labeling of  $C_3 \star P_2 \star S_8$  using this labeling appears in Figure 12.  $\square$

Given the relative simplicity of the labelings for the graphs in Theorem 2.1, it might be surprising that determining prime vertex labelings for  $C_n \star P_2 \star S_m$  with  $m \geq 9$  appears to be more difficult. Consistent with Seoud and Youssef’s conjecture, we expect that all cycle pendant stars are prime.



**Figure 13.** An example of the cycle chain  $C_6^5$ .

### 3. Cycle chains

The “gluing together” of identical cycles appears in various guises in the literature. But the construction of chains of cycles, with adjacent cycles sharing a single common vertex, is not prevalent. For this reason, we require the following definition, where we assume that  $n$  is even. The graph  $C_n^2$  results from attaching two  $n$ -cycles together at a single shared vertex. Continuing in this manner, we define  $C_n^3$  by attaching a third  $n$ -cycle to one of the  $n$ -cycles of  $C_n^2$  in a similar fashion so that the cycle containing two shared vertices consists of two identical  $(n/2)$ -paths. Recursively, the graph  $C_n^m$  consists of a “chain” of  $m$  consecutive  $n$ -cycles. We refer to each of the graphs in this family as a *cycle chain*. For example, the cycle chain  $C_6^5$  consisting of five consecutive 6-cycles is shown in Figure 13.

In what follows, we will show that each  $C_n^m$  is prime for  $n = 4, 6, 8$  and all  $m$ . The labeling functions involved are all similar and relatively simple, which is a consequence of the vertex identification employed on each family. We also show that one of these labeling schemes generalizes in an obvious way, providing a prime vertex labeling for a specific family of cycle chains associated with Mersenne primes.

**Theorem 3.1.** *All  $C_4^m$  are prime.*


*Proof.* The vertices of  $C_4^m$  are identified as follows. First, the vertices of  $C_1$  are identified clockwise as  $c_{1,1}, c_{1,2}, c_{1,3}, c_{1,4}$ , where  $c_{1,4}$  is the vertex also belonging to  $C_2$ . The remaining vertex identifications are based on the parity of  $i$ .

If  $i$  is even, the vertices  $c_{i,1}, c_{i,2}, c_{i,3}, c_{i,4}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . If  $i$  is odd and greater than 1, the vertices  $c_{i,1}$  and  $c_{i,2}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ , and the vertices  $c_{i,3}$  and  $c_{i,4}$  are identified counterclockwise from the vertex that is counterclockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . Note that in both cases,  $c_{i,4}$  is the common vertex of  $C_i$  and  $C_{i+1}$ .

The graph  $C_4^m$  can now be prime labeled via

$$f(c_{i,k}) = \begin{cases} k + 1 & \text{if } i = 1, 1 \leq k \leq 4, \\ 3i + k - 1 & \text{if } 2 \leq i \leq m, 1 \leq k \leq 3, \text{ except } i = m \text{ and } k = 3, \\ 1 & \text{if } i = m, k = 3. \end{cases}$$

It is simple to verify that all adjacent vertices receive relatively prime labels.  $\square$



**Figure 14.** Prime vertex labelings of  $C_4^4$  and  $C_4^5$ .

In Figure 14, we see two labeled examples of  $C_4^m$ , one for each of the possible locations of the integer 1, the last vertex label to be assigned.

**Theorem 3.2.** All  $C_6^m$  are prime.

*Proof.* The vertices of  $C_6^m$  are identified as follows. First, the vertices of  $C_1$  are identified counterclockwise as  $c_{1,1}, c_{1,2}, \dots, c_{1,6}$ , where  $c_{1,1}$  is the vertex also belonging to  $C_2$ . The remaining vertex identifications are based on the congruence class modulo 3 to which  $i$  belongs.


If  $i \equiv_3 0, 2$ , the vertices  $c_{i,1}$  and  $c_{i,2}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ , and the vertices  $c_{i,3}, c_{i,4}, c_{i,5}$  are identified counterclockwise from the vertex that is counterclockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . If  $i \equiv_3 1$ , the vertices  $c_{i,1}, c_{i,2}, \dots, c_{i,5}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . Note that in both cases,  $c_{i,5}$  is the common vertex of  $C_i$  and  $C_{i+1}$ .

The graph of  $C_6^m$  is then labeled using the function given by


$$f(c_{i,k}) = \begin{cases} k & \text{if } i = 1, 1 \leq k \leq 6, \\ 5i + k - 4 & \text{if } i > 1, 1 \leq k \leq 6. \end{cases}$$

Again, it is straightforward to verify that all adjacent vertices receive relatively prime labels, and the result follows.  $\square$

A prime vertex labeling of  $C_6^5$  using Theorem 3.2 appears in Figure 15.



**Figure 15.** A prime vertex labeling of  $C_6^5$ .



**Figure 16.** A prime vertex labeling of  $C_8^5$ .

**Theorem 3.3.** All  $C_8^m$  are prime.

*Proof.* The vertices of  $C_8^m$  are identified as follows. The vertices of  $C_1$  are identified counterclockwise as  $c_{1,1}, c_{1,2}, \dots, c_{1,8}$ , where  $c_{1,1}$  is the vertex also belonging to  $C_2$ . The remaining vertex identifications are based on the parity of  $i$ .

If  $i$  is even, the vertices  $c_{i,1}, c_{i,2}, c_{i,3}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ , and the vertices  $c_{i,4}, c_{i,5}, c_{i,6}, c_{i,7}$  are identified counterclockwise from the vertex that is counterclockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . If  $i$  is odd and greater than 1, the vertices  $c_{i,1}, c_{i,2}, \dots, c_{i,7}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . Note that in both cases,  $c_{i,4}$  is the common vertex of  $C_i$  and  $C_{i+1}$ .

The graph of  $C_8^m$  is then labeled using the function given by

$$f(c_{i,k}) = \begin{cases} k & \text{if } i = 1, 1 \leq k \leq 8, \\ 7i + k - 6 & \text{if } i > 1, 1 \leq k \leq 7. \end{cases}$$

It is simple to verify that all adjacent vertices receive relatively prime labels.  $\square$


An example of a prime vertex labeling of  $C_8^5$  using the labeling in Theorem 3.3 appears in Figure 16.

For any positive integer  $k$ , a Mersenne number is an integer of the form  $M_k = 2^k - 1$ . If  $M_k = 2^k - 1$  is a prime number, then  $M_k$  is called a *Mersenne prime*. The first few Mersenne primes are  $M_2 = 2^2 - 1 = 3$ ,  $M_3 = 2^3 - 1 = 7$ , and  $M_5 = 2^5 - 1 = 31$ . There are 49 known Mersenne primes.

**Theorem 3.4.** Let  $k \in \mathbb{N}$ ,  $k \geq 3$ , and let  $n = 2^k$ . If  $2^k - 1$  is a Mersenne prime, then  $C_n^m$  has a prime vertex labeling.

*Proof.* Both the vertex identification and the labeling function follow from what appeared in the proof of Theorem 3.3. It should be noted that these are different from the machinery used to show that  $C_4^m$  are prime. The vertices of  $C_1$  are identified clockwise as  $c_{1,1}, c_{1,2}, \dots, c_{1,2^k}$ , where  $c_{1,1}$  is the vertex also belonging to  $C_2$ . The remaining vertex identifications are based on the parity of  $i$ .

If  $i$  is even, the vertices  $c_{i,1}, c_{i,2}, \dots, c_{i,2^{k-1}-1}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ , and



**Figure 17.** A prime vertex labeling of the Fibonacci chain  $C_F^5$ .

the vertices  $c_{i,2^{k-1}}, c_{i,2^{k-1}+1}, \dots, c_{i,2^k-1}$  are identified counterclockwise from the vertex that is counterclockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . If  $i$  is odd and greater than 1, the vertices  $c_{i,1}, c_{i,2}, \dots, c_{i,2^k-1}$  are identified clockwise from the vertex that is clockwise adjacent to the common vertex of  $C_i$  and  $C_{i-1}$ . Note that  $c_{i,2^k-1}$  is the common vertex of  $C_i$  and  $C_{i+1}$  in both cases.

The graph  $C_n^m$  can be prime labeled using the function given by

$$f(c_{i,k}) = \begin{cases} k & \text{if } i = 1, 1 \leq k \leq n, \\ (n-1)i + k - (n-2) & \text{if } i > 1, 1 \leq k \leq n-1. \end{cases}$$

It is again relatively straightforward to verify that all adjacent vertices receive relatively prime labels. □

Another historically significant sequence of integers that can be used to generate prime chains of cycles is the sequence of Fibonacci numbers. However, in this case, we will attach cycles of increasing size determined by the terms of the Fibonacci sequence. Recall that the Fibonacci numbers are defined by the recurrence relation  $F_i = F_{i-1} + F_{i-2}$ , where  $F_1 = 1$  and  $F_2 = 1$ . The first several Fibonacci numbers are 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144. To construct our graph, begin with a single path consisting of  $m + 2$  vertices. Starting from one end of the path, identify the vertices as  $p_1, p_2, \dots, p_{m+2}$ . First, add an edge between  $p_1$  and  $p_3$ . Next, for  $i \geq 3$ , build out a cycle by adding an additional  $F_i$  edges between  $p_i$  and  $p_{i+1}$ . The resulting graph consists of  $m$  consecutive cycles, where the first cycle will consist of three vertices and, for  $j \geq 2$ , the  $j$ -th cycle will consist of  $F_{j+1} + 1$  vertices. We denote this graph by  $C_F^m$  and call it a *Fibonacci cycle chain*.

The well-known fact that consecutive Fibonacci numbers are relatively prime leads directly to the following result.

**Theorem 3.5.** *All Fibonacci cycle chains  $C_F^m$  are prime.*



*Proof.* To label a Fibonacci cycle chain, begin by consecutively labeling the path initially used to construct the graph with the Fibonacci numbers starting with the second 1 of the sequence. The remaining vertices can be labeled using the complement of the Fibonacci sequence in the obvious way.  $\square$

Figure 17 shows a Fibonacci cycle chain with a prime vertex labeling. Note that the colored vertices are those that are assigned Fibonacci numbers as labels, which form a path within the graph.

#### 4. Prisms

A *prism graph* is a graph of the form  $C_n \times P_2$ , which consists of an inner and an outer  $n$ -cycle connected with spurs. Prajapati and Gajjar [2014] showed that if  $n + 1$  is prime, then  $C_n \times P_2$  has a prime vertex labeling. Moreover, they also showed that if  $n \geq 3$  is odd, then  $C_n \times P_2$  does not have a prime vertex labeling. In this section, we will prove that if  $n - 1$  is prime, then  $C_n \times P_2$  has a prime vertex labeling. The remaining cases involving prism graphs are currently open.

When  $n$  is even, the initial strategy for labeling prism graphs is to divide the set  $\{1, 2, \dots, 2n\}$  into two subsets, namely  $\{1, 2, \dots, n\}$  and  $\{n + 1, n + 2, \dots, 2n\}$ . We then attempt to label the inner-cycle vertices clockwise using the consecutive integers from 1 to  $n$  and label the outer-cycle vertices with consecutive integers from  $n + 1$  to  $2n$  in the same direction so that the vertex labeled  $n + 1$  is adjacent to the vertex labeled 2. In this case, the difference between the labels assigned to adjacent pairs of inner and outer vertices will be  $n - 1$  except between 1 and  $2n$ . Since  $n - 1$  is prime, most pairs of labels for inner and outer vertices will be guaranteed to be relatively prime. However, we have a problem with the labels  $n - 1$  and  $2(n - 1)$  that the labeling in the next theorem will address by swapping the labels 1 and  $n$  with  $n - 1$  and  $2n$ , respectively. It should be noted that our technique does not generalize to arbitrary  $C_n \times P_2$  (for  $n$  even).


**Theorem 4.1.** *If  $n - 1$  is a prime number and  $n \geq 4$ , then  $C_n \times P_2$  is prime.*

*Proof.* Denote the vertices on the inner cycle by  $c_{1,1}, c_{1,2}, \dots, c_{1,n}$  and their corresponding vertices on the outer cycle by  $c_{2,1}, c_{2,2}, \dots, c_{2,n}$ . The labeling function  $f : V \rightarrow \{1, 2, \dots, 2n\}$  is given by

$$f(c_{1,i}) = \begin{cases} i & \text{if } i = 2, 3, \dots, n - 2, \\ n - 1 & \text{if } i = 1, \\ 1 & \text{if } i = n - 1, \\ 2n & \text{if } i = n, \end{cases}$$

and

$$f(c_{2,i}) = \begin{cases} i + n - 1 & \text{if } i = 2, 3, \dots, n, \\ n & \text{if } i = 1. \end{cases}$$



**Figure 18.** The generalized labeling for the prism  $C_n \times P_2$  when  $n - 1$  is prime.

Recall that consecutive integers are relatively prime, which forces most pairs of adjacent inner-cycle vertices and most pairs of adjacent outer-cycle vertices to receive relatively prime labels. To verify the relative primeness of the labels on the remaining pairs of adjacent vertices, first note that, for  $i = 2, 3, \dots, n - 2$ , we have

$$(f(c_{1,i}), f(c_{2,i})) = (i, i + n - 1) = 1.$$

Next, for  $i = 1$ , we see that

$$(f(c_{1,1}), f(c_{2,1})) = (n - 1, n) = 1.$$

Finally, observe that

$$(f(c_{1,1}), f(c_{1,2})) = (n - 1, 2) = 1,$$


$$(f(c_{1,1}), f(c_{1,n})) = (n - 1, 2n) = 1,$$

$$(f(c_{1,1}), f(c_{1,2})) = (n - 1, 2) = 1,$$


$$(f(c_{1,n-2}), f(c_{2,n-2})) = (n - 2, 2n - 3) = 1.$$

In Figure 18, we see a portion of the labeling of  $C_n \times P_2$  ( $n - 1$  prime). Note that vertices absent from the figure are labeled clockwise using consecutive integers. Thus,  $C_n \times P_2$  has a prime vertex labeling when  $n - 1$  is a prime number.  $\square$

As an example, Figure 19 depicts the labeling of Theorem 4.1 for  $C_6 \times P_2$ .



**Figure 19.** A prime vertex labeling of  $C_6 \times P_2$ .



**Figure 20.** The generalized book  $S_6 \times P_3$ .

### 5. Generalized books


A *book* is a graph of the form  $S_n \times P_2$ , where  $S_n$  is the star with  $n$  spur vertices and  $P_2$  is the path with two vertices. Using a simple parity argument, Seoud and Youssef [1999] showed that all books have a prime vertex labeling. In this section, we extend their work by providing a prime vertex labeling for some *generalized books*, which are graphs of the form  $S_n \times P_m$ . Observe that  $S_n \times P_m$  looks like  $m - 1$  books glued together. The generalized book  $S_6 \times P_3$  is shown in Figure 20.

**Theorem 5.1.** *All  $S_n \times P_m$  with  $3 \leq m \leq 7$  are prime.*


*Proof.* We will handle each of the cases involving  $3 \leq m \leq 7$  separately. For each  $3 \leq m \leq 7$ , let  $c_1, c_3, \dots, c_m$  denote the vertices on the path through the center of each star  $S_n$ . Next, let  $v_{i,1}, v_{i,2}, \dots, v_{i,n}$  denote the vertices of degree 1 on the  $i$ -th star so that  $v_{i,k}$  in the  $i$ -th star is adjacent to  $v_{i+1,k}$  in the  $(i+1)$ -th star. For an example, see the identification of vertices depicted in Figure 20.

First, we consider the generalized book  $S_n \times P_3$ . We define our labeling function  $f : V \rightarrow \{1, 2, \dots, 3n + 3\}$  as follows. Let  $f(c_j) = j$ . For  $k$  odd, define  $f(v_{i,k}) = 3k - i + 4$ . For  $k$  even, let  $f(v_{1,k}) = 3k + 2$ ,  $f(v_{2,k}) = 3k + 3$ , and  $f(v_{3,k}) = 3k + 1$ . Using this labeling, one can quickly see that  $S_n \times P_3$  has a prime vertex labeling. For an example, see the labeling of  $S_6 \times P_3$  given in Figure 21.

For  $S_n \times P_4$ , we define our labeling function  $f : V \rightarrow \{1, 2, \dots, 4n + 4\}$  as follows. Let  $f(c_j) = j$ . For  $k \equiv_3 1$ , let  $f(v_{1,k}) = 4k + 2$ ,  $f(v_{2,k}) = 4k + 3$ ,  $f(v_{3,k}) = 4k + 4$ , and  $f(v_{4,k}) = 4k + 1$ . For  $k \not\equiv_3 1$ , define  $f(v_{i,k}) = 4k - i + 5$ . It then follows that each  $S_n \times P_4$  has a prime vertex labeling. An example of this labeling for  $S_6 \times P_4$  appears in Figure 22.



**Figure 21.** A prime vertex labeling of  $S_6 \times P_3$ .




**Figure 22.** A prime vertex labeling of  $S_6 \times P_4$ .

Next, for  $S_n \times P_5$ , we define our labeling function  $f : V \rightarrow \{1, 2, \dots, 5n + 5\}$  in the following manner. Let  $f(c_j) = j$ . The rest of the labeling function is given by

$$f(v_{i,k}) = \begin{cases} 11 - i & \text{if } k = 1, \\ 5k + 1 + i & \text{if } 1 \leq i \leq 4, k > 1, k - 1 \equiv_6 1, 5, \\ 5k + 1 & \text{if } i = 5, k > 1, k - 1 \equiv_6 1, 5, \\ 5k + 2 + i & \text{if } 1 \leq i \leq 3, k > 1, k - 1 \equiv_6 0, 2, \\ 5k + 6 - i & \text{if } i = 4, 5, k > 1, k - 1 \equiv_6 0, 2, \\ 5k + 5 - i & \text{if } 1 \leq i \leq 3, k > 1, k - 1 \equiv_6 3, \\ 5k + 5 & \text{if } i = 4, k > 1, k - 1 \equiv_6 3, \\ 5k + 1 & \text{if } i = 5, k > 1, k - 1 \equiv_6 3, \\ 5k + 6 - i & \text{if } k > 1, k - 1 \equiv_6 4. \end{cases}$$

It then follows that each  $S_n \times P_5$  has a prime vertex labeling. For an example, see the labeling of  $S_7 \times P_5$  given in Figure 23.




**Figure 23.** A prime vertex labeling of  $S_7 \times P_5$ .

For  $S_n \times P_6$ , we define our labeling function  $f : V \rightarrow \{1, 2, \dots, 6n + 6\}$  as follows. Let  $f(c_j) = j$  and let

$$f(v_{i,k}) = \begin{cases} 6(k+1) + 1 - i & \text{if } k = 1, 2, \\ 6k + 1 + i & \text{if } 1 \leq i \leq 5, k = 3, \\ 6k + 1 & \text{if } i = 6, k = 3, \\ 6k + 1 - i & \text{if } k > 3, k - 3 \equiv_5 1, 2, 3, 4, \\ 6k + 1 + i & \text{if } 1 \leq i \leq 5, k > 3, k - 3 \equiv_5 0, \\ 6k + 1 & \text{if } i = 6, k > 3, k - 3 \equiv_5 0. \end{cases}$$

Using this labeling, we see that each  $S_n \times P_6$  has a prime vertex labeling. An example of this labeling for  $S_8 \times P_6$  appears in Figure 24.



**Figure 24.** A prime vertex labeling of  $S_8 \times P_6$ .

Lastly, for  $S_n \times P_7$ , we define our labeling function  $f : V \rightarrow \{1, 2, \dots, 7n+7\}$  as follows. Let  $f(c_j) = j$ . The remaining portion of the labeling function will involve 10 ordered “row” (i.e., set of corresponding positions in each star) permutation patterns, which we will denote by  $A, B, C, D, E, F, G, H, I$ , and  $J$ . Each ordered row permutation pattern takes the seven consecutive integers that will be used to label the vertices in the  $k$ -th row ( $7k+1, \dots, 7k+7$ ) and assigns them the labels  $w_1 = 7k+1, w_2 = 7k+2, \dots, w_7 = 7k+7$ . For instance, if permutation  $A$  is applied to the  $k$ -th row, then the first vertex in the  $k$ -th row is given the label  $w_2 = 7k+2$ , the second vertex in the  $k$ -th row is given the label  $w_3 = 7k+3$ , etc. Here are the 10 row permutations written in 1-line notation:

$$\begin{aligned} A &= [w_2, w_3, w_4, w_5, w_6, w_7, w_1], \\ B &= [w_2, w_3, w_6, w_7, w_4, w_5, w_1], \\ C &= [w_3, w_2, w_7, w_6, w_5, w_4, w_1], \\ D &= [w_5, w_6, w_7, w_2, w_3, w_4, w_1], \\ E &= [w_4, w_5, w_6, w_7, w_2, w_3, w_1], \\ F &= [w_3, w_4, w_5, w_6, w_7, w_2, w_1], \\ G &= [w_6, w_7, w_2, w_3, w_4, w_5, w_1], \\ H &= [w_2, w_1, w_3, w_7, w_6, w_5, w_4], \\ I &= [w_7, w_6, w_1, w_2, w_3, w_4, w_5], \\ J &= [w_7, w_6, w_5, w_4, w_3, w_2, w_1]. \end{aligned}$$

After assigning  $f(c_j) = j$  to the center of each star, the labeling for  $S_n \times P_7$  has the following 30-row repeating pattern:

$C, E, J, A, J, A, D, E, J, A, F, G, C, H, J, A, J, A, I, E, F, E, J, A, D, E, J, A, J, A.$

It is straightforward to check that this “block” of 30 rows provides a prime vertex labeling for  $S_{30} \times P_7$ . To label the next block of 30 rows in the generalized book, simply add 210 to each of the labels from the first block of 30 rows. Since any of the vertices in this second block can only be adjacent to the center of its star and any neighbors in its row, and since congruence classes modulo 2, 3, 4, 5, 6, and 7 are invariant under addition of 210, all adjacent vertices in this second block must still have relatively prime labels. Thus, this second block has a prime vertex labeling. It then follows that the entire generalized book has a prime vertex labeling.  $\square$

We conjecture that all generalized books  $S_n \times P_m$  have a prime vertex labeling; however, it appears that extending our results using the current approach becomes increasingly difficult as one increases the size of the path  $P_m$ .

## 6. Conclusion

The prime vertex labeling functions included in this paper for cycle pendant stars may be extended to somewhat larger sizes of stars. Specifically, we conjecture that similar processes will work for cycle pendant stars up to stars of size 15. The reasoning for this restriction on stars of size 15 is the following result of Pillai [1941].

**Proposition 6.1.** *When  $k \geq 17$ , we can find  $k$  consecutive integers such that no integer in the set is relatively prime to all other integers in the set.*

This implies that a new labeling scheme must be devised for finding prime vertex labelings of  $C_n \star P_2 \star S_m$  for  $m \geq 15$ .

In Section 3, we constructed prime vertex labelings for cycle chains having cycles of sizes 4, 6, and 8, respectively. We conjecture that all cycle chains are prime. Using the fact that consecutive Fibonacci numbers are relatively prime, we constructed a prime graph. One can likely construct similar graphs using other well-known sequences that exhibit traits of relative primeness.

We conjecture that all prisms constructed from even-order cycles have prime vertex labelings. We also have preliminary results implying that some families of generalized prisms of the form  $C_n \times P_m$  for  $m > 2$  have prime vertex labelings.

In addition, we conjecture that every generalized book has a prime vertex labeling. But our pairwise matching approach would need to be replaced in order for this conjecture to be realized. For the interested reader, additional information can be found in Gallian's dynamic survey on graph labelings [1998].

## References

- [Diefenderfer et al. 2015] N. Diefenderfer, M. G. Hastings, L. N. Heath, H. Prawzinsky, B. Preston, E. White, and A. Whittemore, "Prime vertex labelings of families of unicyclic graphs", *Rose-Hulman Undergrad. Math J.* **16**:1 (2015), 253–269. MR 3367500
- [Gallian 1998] J. A. Gallian, "A dynamic survey of graph labeling", *Electron. J. Combin. Dynamic Surveys*:6 (1998). 18th ed., December 7, 2015. MR 1668059 Zbl 0953.05067
- [Pillai 1941] S. S. Pillai, "On  $m$  consecutive integers—III", *Proc. Indian Acad. Sci. (A)* **13**:6 (1941), 530–533. MR 0004824 Zbl 0063.06228
- [Prajapati and Gajjar 2014] U. M. Prajapati and S. J. Gajjar, "Some results on prime labeling", *Open J. Discret. Math.* **4**:3 (2014), 60–66.
- [Seoud and Youssef 1999] M. A. Seoud and M. Z. Youssef, "On prime labeling of graphs", *Congr. Numer.* **141** (1999), 203–215. MR 1745237 Zbl 0968.05068

Received: 2015-03-21      Revised: 2015-08-12      Accepted: 2015-08-17

ned32@nau.edu

Northern Arizona University, Flagstaff, AZ 86011,  
United States

dana.ernst@nau.edu

Northern Arizona University, Flagstaff, AZ 86001,  
United States

mgh64@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>
lnh57@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>
hpp3@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>
bep38@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>
jeffrey.rushall@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>
ekw49@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>
anw99@nau.edu	<i>Northern Arizona University, Flagstaff, AZ 86011, United States</i>



# involve

msp.org/involve

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Errin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2016 is US \$160/year for the electronic version, and \$215/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

*Involve* (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# involve

2016

vol. 9

no. 4

Affine hyperbolic toral automorphisms COLIN THOMSON AND DONNA K. MOLINEK	541
Rings of invariants for the three-dimensional modular representations of elementary abelian $p$ -groups of rank four THÉO PIERRON AND R. JAMES SHANK	551
Bootstrap techniques for measures of center for three-dimensional rotation data L. KATIE WILL AND MELISSA A. BINGHAM	583
Graphs on 21 edges that are not 2-apex JAMISON BARSOTTI AND THOMAS W. MATTMAN	591
Mathematical modeling of a surface morphological instability of a thin monocrystal film in a strong electric field AARON WINGO, SELAHITTIN CINAR, KURT WOODS AND MIKHAIL KHENNER	623
Jacobian varieties of Hurwitz curves with automorphism group $\mathrm{PSL}(2, q)$ ALLISON FISCHER, MOUCHEN LIU AND JENNIFER PAULHUS	639
Avoiding approximate repetitions with respect to the longest common subsequence distance SERINA CAMUNGOL AND NARAD RAMPERSAD	657
Prime vertex labelings of several families of graphs NATHAN DIEFENDERFER, DANA C. ERNST, MICHAEL G. HASTINGS, LEVI N. HEATH, HANNAH PRAWZINSKY, BRIAHNA PRESTON, JEFF RUSHALL, EMILY WHITE AND ALYSSA WHITTEMORE	667
Presentations of Roger and Yang's Kauffman bracket arc algebra MARTIN BOBB, DYLAN PEIFER, STEPHEN KENNEDY AND HELEN WONG	689
Arranging kings $k$ -dependently on hexagonal chessboards ROBERT DOUGHTY, JESSICA GONDA, ADRIANA MORALES, BERKELEY REISWIG, JOSIAH REISWIG, KATHERINE SLYMAN AND DANIEL PRITIKIN	699
Gonality of random graphs ANDREW DEVEAU, DAVID JENSEN, JENNA KAINIC AND DAN MITROPOLSKY	715



1944-4176(2016)9:4;1-1