

involve

a journal of mathematics

Discrete dynamics of contractions on graphs

Olena Ostapyuk and Mark Ronnenberg



Discrete dynamics of contractions on graphs

Olena Ostapyuk and Mark Ronnenberg

(Communicated by Martin Bohner)

We study the dynamical behavior of functions on vertices of a graph that are contractions in the graph metric. We show that the fixed point set of such functions must be convex. If a function has no fixed points and the graph is a tree, we prove that every dynamical cycle must have an even period and the function behaves eventually like a symmetry.

1. Introduction

This work was inspired by dynamics of analytic functions on the unit disk. The key property of such functions is the point-invariant Schwarz lemma, i.e., that analytic functions are contractions in the hyperbolic metric of the disk. This property allows the proof of various results about iteration of analytic functions; see, for example, the survey paper [Poggi-Corradini 2011].

Our purpose is to study dynamics of contractions in a discrete setting. In particular, we study dynamics on finite graphs (in most cases, trees). A connected graph can be considered as a discrete metric space of vertices with the *graph metric*. Let $G = (V, E)$ be a finite, connected, simple graph with the set of vertices V and the set of edges E . Then for all vertices $x, y \in V$, we say the distance between x and y , denoted $d(x, y)$, is the number of edges in the shortest path connecting x to y . Such path is called a *geodesic*. Note that trees as metric spaces are 0-hyperbolic [Anderson 1999], so we expect them to have some similar properties to the unit disk with hyperbolic metric.

We wish to study contractions (in the graph metric) on the vertices of a graph. Let f be a function on the vertices of G to the vertices of G . We say f is a *contraction* if, for all vertices $x, y \in V$, we have $d(f(x), f(y)) \leq d(x, y)$. We will need some terminology from dynamics. Let f be a function. We denote by $f^{on}(x) = f \circ f \circ f \circ \dots \circ f(x)$ (n terms) the n -th *iterate* of f . If for some point x and some positive integer n , we have $f^{on}(x) = x$, then we say x is a periodic point, x lies on a *dynamical cycle* of f of period n , or that x lies on a *dynamical n -cycle*

MSC2010: primary 39B12, 54H20; secondary 05C05.

Keywords: discrete dynamics, dynamics of contractions, graphs.

of f . If $f(x) = x$, we say x is a *fixed point* of f . We use the term *dynamical cycle* to distinguish these cycles from the graph cycles.

It is easy to show by induction that, given a contraction f , the map f^{on} is also a contraction for any positive integer n . Dynamical cycles and fixed points will be the main focus of our study.

2. Fixed point sets

Our goal is to characterize the set of fixed points of a contraction on graph vertices. Note that in the general case, the fixed point set can be empty:

Example 2.1. Let G_1 be a graph with four vertices x, y, z, w . Let f be a function on the vertices of G_1 defined by $f(x) = y, f(y) = z, f(z) = w$, and $f(w) = x$. Then $\{x, y, z, w\}$ forms a dynamical 4-cycle of f (see Figure 1). The map f is clearly a contraction since for all $a, b \in \{x, y, w, z\}$, we have $d(f(a), f(b)) = d(a, b)$. In this case, the set of fixed points of f is empty.

Example 2.2. Let G_2 be a graph with vertices $x_0, x_1, x_2, y_0, y_1, z_0$ and z_1 as shown in Figure 2. Let f be a contraction on the vertices of G_2 such that x_0, x_1, x_2 are fixed by f , and $\{y_0, y_1\}$ and $\{z_0, z_1\}$ are dynamical 2-cycles of f .

Note that one main difference between the two examples is that for any two vertices in G_2 , the geodesic connecting them is unique, whereas this is not the case with G_1 . Notice also that for any two fixed points in G_2 , the geodesic connecting them contains only fixed points.




Figure 1. Dynamical 4-cycle.




Figure 2. Cycles and fixed points.




Figure 3. The graph G_3 with nonunique geodesics.

Definition 2.3. Let $G = (V, E)$ be a graph and let $H \subset V$. We say H is *convex* if for any two vertices in H , the geodesic connecting them contains only vertices in H . (See, for example, [Gross and Yellen 2006].)

Thus in [Example 2.2](#), the set of fixed points of f is convex. In fact, this is true in general:

Theorem 2.4. *Let $G = (V, E)$ be a graph such that the geodesic between any two vertices is unique. Let f be a contraction on the vertices of G . Then the set of fixed points of f is convex.*

Proof. Let $x, y \in V$ be fixed by f . Let L be the unique geodesic connecting them. Let $z \in L$. We need to show that $f(z) = z$. We will first show that $f(z) \in L$ and it will follow that $f(z) = z$.

By way of contradiction, suppose $f(z) \notin L$. Then there exist unique geodesics connecting x to $f(z)$ and y to $f(z)$, respectively. We can concatenate these geodesics to construct a walk K connecting x to y , so the length of K is $d(x, f(z)) + d(f(z), y)$ and the length of L is $d(x, z) + d(z, y)$. Since f is a contraction and x and y are fixed points, we have $d(f(z), x) \leq d(z, x)$ and $d(f(z), y) \leq d(z, y)$. Then it follows that $d(x, f(z)) + d(f(z), y) \leq d(x, z) + d(z, y)$. If $d(x, f(z)) + d(f(z), y) = d(x, z) + d(z, y)$, then L is not a unique geodesic between x and y , a contradiction. If $d(x, f(z)) + d(f(z), y) < d(x, z) + d(z, y)$, then K is shorter than L , which is also a contradiction. Thus it must be that $z \in L$.

Now we will show $f(z) = z$. Suppose $f(z) \neq z$. Since $f(z)$ lies on the geodesic L connecting x to y , we have $d(x, z) + d(z, y) = d(x, f(z)) + d(f(z), y) = d(x, y)$. We can assume without loss of generality that $d(x, f(z)) < d(x, z)$, in which case we obtain $d(y, f(z)) = d(x, y) - d(x, f(z)) > d(x, y) - d(x, z) = d(y, z)$, contradicting the fact that f is a contraction. Thus we conclude that $f(z) = z$. \square

Note that if for any two points in G the geodesic connecting them is not unique then the conclusion of [Theorem 2.4](#) does not necessarily hold, as can be seen in the following counterexample.

Example 2.5. Let G_3 be a graph with vertices x_0, x_1, y and z as shown in [Figure 3](#).

Let f be a contraction such that the vertices z and y are fixed and the points x_0 and x_1 form a dynamical 2-cycle. Note that the geodesic connecting z to y is

not unique, since the path from z to y through x_0 is the same length as the path through x_1 . Despite the fact that z and y are fixed and that x_0, x_1 lie on the geodesics connecting them, x_0 and x_1 are clearly not fixed. Thus the conclusion of [Theorem 2.4](#) does not hold in this case.

Corollary 2.6. *Let $G = (V, E)$ be a graph such that for any two vertices in G the geodesic connecting them is unique. Let f be a contraction on V . Suppose f has a dynamical cycle J of period k . Let z be a point which lies on the geodesic connecting two consecutive points in J . Then z lies on a dynamical cycle whose period divides k .*

Proof. Let $x, y \in J$. Let $z \in V$ such that z lies on the geodesic between x and y . Since J is a dynamical cycle of period k , we know $f^{ok}(x) = x$ and $f^{ok}(y) = y$. Thus x and y are fixed by the k -th iterate of f . Since f is a contraction, any iterate of f is also a contraction. Thus [Theorem 2.4](#) applied to f^{ok} implies $f^{ok}(z) = z$. So z must lie on a dynamical cycle whose period divides k . \square

Now we will consider a particular case when the graph is a tree. For any tree, a path connecting any two points is unique, hence geodesics are unique, so [Theorem 2.4](#) holds. But the converse is also true for trees: any convex set of vertices will be a fixed point set for some contraction.

We will need the following property of a tree structure: in a tree, a concatenation of two geodesics from x to y and from y to z is either a geodesic from x to z or a walk that follows the geodesic connecting x to y until the first common point of the two geodesics, y' , then follows the geodesic from y' to y , then goes back to y' along the same geodesic and finally follows the geodesic from y' to z . Note that concatenation of geodesics from x to y' and from y' to z will form a geodesic that connects x to z .

Proposition 2.7. *Let $T = (V, E)$ be a tree and $H \subset V$ be convex set. Then there exists a contraction f such that H is the fixed point set of f .*

Proof. Given H , we define the desired contraction f as follows: for all $x \in V$, $f(x) = y$, where $y \in H$ is the closest vertex to x in H ; see [Figure 4](#). Note that such a y is unique. Indeed, suppose $y_1, y_2 \in H$ are at the same shortest distance from $x \notin H$. Apply the property mentioned above the proposition to the concatenation of geodesics connecting y_1 to x and x to y_2 . If it is a geodesic, then $x \in H$, which is a contradiction. If instead there is a common point y' , then $y' \in H$ and it is closer to x than y_1 and y_2 are, again a contradiction. Thus the point y is unique and the function f is well-defined. Also, H is clearly fixed point set of f .

Now we need to show that f is a contraction. Let $f(x_1) = y_1$ and $f(x_2) = y_2$. Consider a walk following the geodesic from x_1 to y_1 , then from y_1 to y_2 . If there is a common point of these geodesics other than y_1 , then this point is in H and within a shorter distance to x_1 than y_1 , which contradicts the construction of y_1 . So the




Figure 4. Constructing the contraction f for a given convex subset of vertices H .

concatenation of these two geodesics is the geodesic from x_1 to y_2 . Similarly, the geodesic from y_1 to x_2 passes through y_2 , and finally, the geodesic from x_1 to x_2 is just a concatenation of those from x_1 to y_1 , y_1 to y_2 and y_2 to x_1 . So we have

$$d(x_1, x_2) = d(x_1, y_1) + d(y_1, y_2) + d(y_2, x_2) \geq d(y_1, y_2),$$

and f is a contraction. □

3. Contractions with no fixed points

In the previous section, we characterized the set of fixed points of a contraction on the vertices of a graph with unique geodesics, in particular a tree. Next we want to consider the case when a contraction has no fixed points. Then there must exist a dynamical cycle. We will use the following property of periodic points:

Lemma 3.1. *Let G be a finite graph, and f be a contraction on vertices of G . If x and y are two periodic points of f (not necessarily from the same dynamical cycle), then $d(f(x), f(y)) = d(x, y)$.*

Proof. Assume x belongs to a dynamical m -cycle and y belongs to a dynamical n -cycle. Let K be a common multiple of m and n . Then we have

$$d(x, y) \geq d(f(x), f(y)) \geq \dots \geq d(f^{\circ K}(x), f^{\circ K}(y)) = d(x, y).$$

So all inequalities must be, in fact, equalities and in particular, $d(f(x), f(y)) = d(x, y)$. □

Now let us introduce some notation. Let $G = (V, E)$ be a graph and f a contraction on V . Let $J \subset V$ be a dynamical cycle of f . Then we denote by J' the set of all vertices which lie on geodesics connecting consecutive points in J , together with the vertices in J .

Theorem 3.2. *Let T be a finite tree. Let f be a contraction on the vertices of T . If f has no fixed points, then f has a dynamical 2-cycle such that the points in the cycle are connected by an edge. Moreover, such a cycle is unique.*

Proof. Suppose f has no fixed points. Since the number of vertices of T is finite, every vertex of T either lies on a dynamical cycle of period greater than 1 or is eventually mapped into one. Let k be the smallest period of all dynamical cycles of f . Let J be a dynamical cycle of period k such that the distance between consecutive points in J is least among all dynamical cycles of f of period k . We want to show that $k = 2$.

We claim that for $k > 2$ there must exist two geodesics connecting consecutive points in J that intersect at a point other than their endpoints. If not, the points in J would form a graph cycle, which is a contradiction since T is a tree. Thus there must exist two geodesics which intersect at a point which is not one of their endpoints.

Suppose two nonconsecutive geodesics intersect at some point y . Then we claim that there must exist two consecutive geodesics which intersect at point z which is not one of their endpoints. Indeed, if we start from the point y of intersection of two nonconsecutive geodesics and follow one of the geodesics to the point x_j on the cycle J , then follow the next geodesic to the point $x_{j+1} = f(x_j)$, and so on, we will eventually return to the point y . Since the graph is a tree, the walk constructed this way must go over each edge in this walk at least twice. In particular, there must exist a vertex w which is farthest away from y on this walk and an edge $\{w, z\}$ such that our walk will follow the edge from z to w and then immediately return to z through the same edge. Note that w must be an endpoint of two consecutive geodesics, because one geodesic cannot follow the same edge twice. Then z lies on the intersection of two consecutive geodesics.

Without loss of generality, let x_0, x_1, x_2 be the endpoints of the two consecutive geodesics constructed above. By [Corollary 2.6](#), z must lie on a dynamical cycle whose period divides k , but since k is the smallest possible cycle length, z must lie on a dynamical k -cycle.

Since f is a contraction and x_0, x_1, x_2 are points on a dynamical cycle, f must map the geodesic from x_0 to x_1 bijectively to the geodesic from x_1 to x_2 . Since z lies on the geodesic from x_0 to x_1 , the point $f(z)$ must lie on the geodesic from x_1 to x_2 . Thus both z and $f(z)$ lie on the geodesic from x_1 to x_2 and we have $d(z, f(z)) < d(x_1, x_2) = d(x_0, x_1)$. So we have found a dynamical k -cycle $\{z, f(z), \dots, f^{\circ(k-1)}(z)\}$ such that the distance between two consecutive points in this cycle is less than $d(x_0, x_1)$.

This contradicts the way we selected J , so k must be equal to 2 and the geodesic from x_0 to x_1 , which is the same as the geodesic from x_1 to x_0 , must contain no other points. This means there is a dynamical 2-cycle $\{x_0, x_1\}$ and x_0 and x_1 are connected by an edge.

Now we need prove that such a dynamical 2-cycle is unique. Let $\{y_0, y_1\}$ be another such cycle. Without loss of generality assume that the distance a between x_0 and y_0 is the shortest among all distances from a point in $\{x_0, x_1\}$ to a point in $\{y_0, y_1\}$. Now consider x_1 ; it is connected to x_0 by an edge. If x_1 lies on the geodesic from x_0




Figure 5. Unique 2-cycle $\{x_0, x_1\}$ and sets X_0 and X_1 .

to y_0 , then $d(x_1, y_0) < d(x_0, y_0)$, which contradicts the choice of x_0, y_0 . Otherwise, the geodesic from y_0 to x_1 follows the geodesic from y_0 to x_0 and then the edge connecting x_0 to x_1 , so $d(y_0, x_1) = a + 1$. Similarly, $d(x_0, y_1) = a + 1$, and finally, $d(x_1, y_1) = a + 2$. But then $d(x_1, y_1) = d(f(x_0), f(y_0)) > d(x_0, y_0)$, which contradicts the assumption that f is a contraction. \square

It will in fact turn out that every dynamical cycle of a contraction with no fixed points has even period. To prove this, we will need the following corollary to [Theorem 3.2](#). Let us introduce the following notation. Let $\{x_0, x_1\}$ be the points in the 2-cycle constructed in [Theorem 3.2](#). We let X_0 denote the set of all points which are within shorter distance to x_0 than to x_1 . Similarly we let X_1 denote the set of all points which are within shorter distance to x_1 than to x_0 ; see [Figure 5](#).

Corollary 3.3. *Let T be a finite tree and f a contraction on the vertices of T such that f has no fixed points. Let $\{x_0, x_1\}$ be the unique dynamical 2-cycle, where x_0 and x_1 are connected by an edge. Then for all vertices z that lie on any dynamical cycle, if $z \in X_0$ (respectively X_1), then $f(z) \in X_1$ (respectively X_0).*

Proof. Let z lie on a dynamical cycle and $z \in X_0$. By way of contradiction, suppose $f(z) \in X_0$. Let $a = d(z, x_0)$; then $d(z, x_1) = a + 1$. By [Lemma 3.1](#), $d(f(z), x_1) = a$, and since $f(z) \in X_0$, we must have $d(f(z), x_0) < d(f(z), x_1) = a$. But by [Lemma 3.1](#) again, $d(f(z), x_0) = d(f(z), f(x_1)) = d(z, x_1) = a + 1$, which is a contradiction. So $f(z) \in X_1$. \square

Note that if z is not a periodic point, then the above claim does not hold.

Example 3.4. Let T be a tree with vertices x_0, x_1 and z such that there are edges between x_0 and x_1 and between x_0 and z , and f be a contraction such that $\{x_0, x_1\}$ forms a dynamical 2-cycle and $f(z) = x_0$ (see [Figure 6](#)). Then f has no fixed points, and x_0 and x_1 form the unique 2-cycle connected by an edge. Since $f(z) = x_0$, we have $z \in X_0$ and also $f(z) \in X_0$. Thus we see that if a point z is in X_0 but does not lie on a dynamical cycle, it is not necessarily true that $f(z) \in X_1$.




Figure 6

Now we are ready to prove the following:

Theorem 3.5. *Let T be a finite tree and f a contraction on the vertices of T such that f has no fixed points. Then every dynamical cycle of f has even period.*

Proof. Since f has no fixed points, f has a dynamical 2-cycle $\{x_0, x_1\}$ whose points are connected by an edge and sets of vertices X_0 and X_1 as defined above. Let $\{y_0, y_1, \dots, y_{n-1}\}$ be a dynamical n -cycle of f . Without loss of generality, suppose $y_0 \in X_0$. Then by [Corollary 3.3](#), we have $y_1 \in X_1$, and in general, $y_{2k} \in X_0$ and $y_{2k+1} \in X_1$. If n is odd, then $y_0 = f(y_{n-1}) \in X_1$, which is a contradiction to $y_0 \in X_0$. Hence every dynamical cycle of f has even period. \square

If a contraction f on the vertices of a tree T has no fixed points, then f eventually behaves like a symmetry. More precisely:

Theorem 3.6. *Let $T = (V, E)$ be a finite tree and f a contraction on V without fixed points. Then there exists a subset H of V and a nonnegative integer N such that $f^{\circ N}(V) = H$ and f is a symmetry on the connected subgraph induced by H . In particular, there is an edge in the subgraph such that two connected components obtained by removing this edge are isomorphic graphs and f is an isomorphism.*

Proof. Since T is finite and has no fixed points, each vertex of T will be mapped eventually to a point on a dynamical cycle. Thus there exists N such that $f^{\circ N}(V) = H$ contains only periodic points of f . Note that by [Corollary 2.6](#), the subgraph induced by H is connected. Let $\{x_0, x_1\}$ be the unique dynamical 2-cycle whose points are connected by an edge. Then by [Corollary 3.3](#), for all $z \in H \cap X_0$, we have $f(z) \in H \cap X_1$ and for all $z \in H \cap X_1$, we have $f(z) \in H \cap X_0$. Moreover, since all points in H are periodic, f bijectively maps $H \cap X_0$ to $H \cap X_1$. Now we need to show that any two vertices y, z in $H \cap X_0$ are connected by an edge if and only if $f(y)$ and $f(z)$ are connected by an edge. But being connected by an edge is equivalent to $d(y, z) = 1$, and since by [Lemma 3.1](#), $d(y, z) = d(f(y), f(z))$, the required conclusion follows. \square

4. Conclusion

Note that in the classical case of the unit disk in the complex plane, any analytic self-map of the disk always has a fixed point in the closed disk. This is the consequence of the classical Denjoy–Wolff theorem (see, for example, [\[Abate 1989\]](#)

and references therein). In our study, a contraction without fixed points must behave like a symmetry. Symmetries are contractions in the unit disk, but they are not analytic (in fact, they are anticonformal, i.e., they preserve the value of angles, but change their orientation). So we can say that our result agrees with the classical case.

References

- [Abate 1989] M. Abate, *Iteration theory of holomorphic maps on taut manifolds*, Mediterranean Press, Rende, 1989. [MR](#) [Zbl](#)
- [Anderson 1999] J. W. Anderson, *Hyperbolic geometry*, Springer, London, 1999. [MR](#) [Zbl](#)
- [Gross and Yellen 2006] J. L. Gross and J. Yellen, *Graph theory and its applications*, 2nd ed., Chapman & Hall/CRC, Boca Raton, FL, 2006. [MR](#) [Zbl](#)
- [Poggi-Corradini 2011] P. Poggi-Corradini, “Iteration in the disk and the ball: a survey of the role of hyperbolic geometry”, *Anal. Math. Phys.* **1**:4 (2011), 289–310. [MR](#) [Zbl](#)

Received: 2016-02-14

Revised: 2016-04-12

Accepted: 2016-04-14

ostapyuk@math.uni.edu

*Department of Mathematics, University of Northern Iowa,
Cedar Falls, IA 50614-0506, United States*

ronnema@uni.edu

*Department of Mathematics, University of Northern Iowa,
Cedar Falls, IA 50614-0506, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

involve

2017

vol. 10

no. 3

Dynamics of vertical real rhombic Weierstrass elliptic functions	361
LORELEI KOSS AND KATIE ROY	
Pattern avoidance in double lists	379
CHARLES CRATTY, SAMUEL ERICKSON, FREHIWET NEGASSI AND LARA PUDWELL	
On a randomly accelerated particle	399
MICHELLE NUNO AND JUHI JANG	
Reeb dynamics of the link of the A_n singularity	417
LEONARDO ABBRESCIA, IRIT HUQ-KURUVILLA, JO NELSON AND NAWAZ SULTANI	
The vibration spectrum of two Euler–Bernoulli beams coupled via a dissipative joint	443
CHRIS ABRIOLA, MATTHEW P. COLEMAN, AGLIKA DARAKCHIEVA AND TYLER WALES	
Loxodromes on hypersurfaces of revolution	465
JACOB BLACKWOOD, ADAM DUKEHART AND MOHAMMAD JAVAHERI	
Existence of positive solutions for an approximation of stationary mean-field games	473
NOJOD ALMAYOUF, ELENA BACHINI, ANDREIA CHAPOUTO, RITA FERREIRA, DIOGO GOMES, DANIELA JORDÃO, DAVID EVANGELISTA JUNIOR, AVETIK KARAGULYAN, JUAN MONASTERIO, LEVON NURBEKYAN, GIORGIA PAGLIAR, MARCO PICCIRILLI, SAGAR PRATAPSI, MARIANA PRAZERES, JOÃO REIS, ANDRÉ RODRIGUES, ORLANDO ROMERO, MARIA SARGSYAN, TOMMASO SENECCI, CHULIANG SONG, KENGO TERAJ, RYOTA TOMISAKI, HECTOR VELASCO-PEREZ, VARDAN VOSKANYAN AND XIANJIN YANG	
Discrete dynamics of contractions on graphs	495
OLENA OSTAPYUK AND MARK RONNENBERG	
Tiling annular regions with skew and T-tetrominoes	505
AMANDA BRIGHT, GREGORY J. CLARK, CHARLES LUNDON, KYLE EVITTS, MICHAEL P. HITCHMAN, BRIAN KEATING AND BRIAN WHETTER	
A bijective proof of a q -analogue of the sum of cubes using overpartitions	523
JACOB FORSTER, KRISTINA GARRETT, LUKE JACOBSEN AND ADAM WOOD	
Ulrich partitions for two-step flag varieties	531
IZZET COSKUN AND LUKE JASKOWIAK	